

Analytical and computational study of the variable inverse sum deg index

Walter Carballosa · J. A.
Méndez-Bermúdez · José M. Rodríguez ·
José M. Sigarreta

Received: date / Accepted: date

Abstract A large number of graph invariants of the form $\sum_{uv \in E(G)} F(d_u, d_v)$ are studied in mathematical chemistry, where uv denotes the edge of the graph G connecting the vertices u and v , and d_u is the degree of the vertex u . Among them the variable inverse sum deg index ISD_a , with $F(d_u, d_v) = 1/(d_u^a + d_v^a)$, was found to have applicative properties. The aim of this paper is to obtain new inequalities for the variable inverse sum deg index, and to characterize graphs extremal with respect to them. Some of these inequalities generalize and improve previous results for the inverse sum indeg index. In addition, we computationally validate some of the obtained inequalities on ensembles of random graphs and show that the ratio $\langle ISD_a(G) \rangle / n$ (n being the order of the graph) depends only on the average degree $\langle d \rangle$.

Keywords variable inverse sum deg index · inverse sum indeg index · optimization on graphs · degree-based topological index

Walter Carballosa
Department of Mathematics and Statistics, Florida International University, 11200 SW 8th Street Miami, FL 33199, USA
E-mail: waltercarb@gmail.com

J. A. Méndez-Bermúdez
Instituto de Física, Benemérita Universidad Autónoma de Puebla, Apartado Postal J-48, Puebla 72570, Mexico
E-mail: jmendezb@ifuap.buap.mx

José M. Rodríguez
Departamento de Matemáticas, Universidad Carlos III de Madrid, Avenida de la Universidad 30, 28911 Leganés, Madrid, Spain
E-mail: jomaro@math.uc3m.es

José M. Sigarreta
Facultad de Matemáticas, Universidad Autónoma de Guerrero, Carlos E. Adame No.54 Col. Garita, 39650 Acapulco Gro., Mexico
E-mail: josemariasigarretaalmira@hotmail.com

Mathematics Subject Classification (2010) 05C09 · 05C92

1 Introduction

Topological indices are parameters associated with chemical compounds that associate the chemical structure with several physical, chemical or biological properties.

A family of degree-based topological indices, named *Adriatic indices*, was put forward in [37, 38]. Twenty of them were selected as significant predictors. One of them, the *inverse sum indeg* index, *ISI*, was singled out in [37, 38] as a significant predictor of total surface area of octane isomers. This index is defined as

$$ISI(G) = \sum_{uv \in E(G)} \frac{d_u d_v}{d_u + d_v} = \sum_{uv \in E(G)} \frac{1}{\frac{1}{d_u} + \frac{1}{d_v}},$$

where uv denotes the edge of the graph G connecting the vertices u and v , and d_u is the degree of the vertex u . In the last years there is an increasing interest in the mathematical properties of this index (see, e.g., [6, 11, 13, 14, 21, 26, 30]).

We study here the properties of the *variable inverse sum deg index* defined, for each $a \in \mathbb{R}$, as

$$ISD_a(G) = \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a}.$$

Note that ISD_{-1} is the inverse sum indeg index *ISI*.

The variable inverse sum deg index $ISD_{-1.950}$ was selected in [39] as a significant predictor of standard enthalpy of formation.

The idea behind the variable molecular descriptors is that the variables are determined during the regression so that the standard error of estimate for a particular studied property is as small as possible (see, e.g., [20]).

The aim of this paper is to obtain new inequalities for the variable inverse sum deg index, and to characterize graphs extremal with respect to them. Some of these inequalities generalize and improve previous results for the inverse sum indeg index. Also, we want to remark that many previous results are proved for connected graphs, but our inequalities hold for both connected and non-connected graphs.

Throughout this paper, $G = (V(G), E(G))$ denotes an undirected finite simple (without multiple edges and loops) graph without isolated vertices. We denote by n , m , Δ and δ the cardinality of the set of vertices of G , the cardinality of the set of edges of G , its maximum degree and its minimum degree, respectively. Thus, we have $1 \leq \delta \leq \Delta < n$. We denote by $N(u)$ the set of neighbors of the vertex $u \in V(G)$.

2 Inequalities for the ISD_a index

Proposition 1 *If G is a graph with minimum degree δ , maximum degree Δ and m edges, and $a \in \mathbb{R}$, then*

$$\begin{aligned} \frac{m}{2\Delta^a} &\leq ISD_a(G) \leq \frac{m}{2\delta^a}, & \text{if } a > 0, \\ \frac{m}{2\delta^a} &\leq ISD_a(G) \leq \frac{m}{2\Delta^a}, & \text{if } a < 0. \end{aligned}$$

The equality in each bound is attained if and only if G is regular.

Proof If $a > 0$, then $2\delta^a \leq d_u^a + d_v^a \leq 2\Delta^a$ and

$$\begin{aligned} ISD_a(G) &= \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \leq \sum_{uv \in E(G)} \frac{1}{2\delta^a} = \frac{m}{2\delta^a}, \\ ISD_a(G) &= \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \geq \sum_{uv \in E(G)} \frac{1}{2\Delta^a} = \frac{m}{2\Delta^a}. \end{aligned}$$

If $a < 0$, then the previous argument gives the converse inequalities.

If G is a regular graph, then the lower and upper bounds are the same, and they are equal to $ISD_a(G)$.

Assume now that the equality in some bound is attained. Thus, by the previous argument we have either $d_u = d_v = \delta$ for every $uv \in E(G)$, or $d_u = d_v = \Delta$ for every $uv \in E(G)$. Hence, G is regular.

In 1998 Bollobás and Erdős [3] generalized the Randić index by replacing $1/2$ by any real number. Thus, for $a \in \mathbb{R} \setminus \{0\}$, the *general Randić index* of a graph G is defined as

$$R_a(G) = \sum_{uv \in E(G)} (d_u d_v)^a.$$

The general Randić index, also called *variable Zagreb index* in 2004 by Milicević and Nikolić [20], has been extensively studied [15]. Note that $R_{-1/2}$ is the usual Randić index, R_1 is the second Zagreb index M_2 , R_{-1} is the modified Zagreb index [24], etc. In Randić's original paper [27], in addition to the particular case $a = -1/2$, also the index with $a = -1$ was briefly considered.

The next result relates the ISD_a and R_{-a} indices.

Theorem 1 *If G is a graph with minimum degree δ and maximum degree Δ , and $a \in \mathbb{R}$, then*

$$\begin{aligned} \frac{1}{2} \delta^a R_{-a}(G) &\leq ISD_a(G) \leq \frac{1}{2} \Delta^a R_{-a}(G), & \text{if } a > 0, \\ \frac{1}{2} \Delta^a R_{-a}(G) &\leq ISD_a(G) \leq \frac{1}{2} \delta^a R_{-a}(G), & \text{if } a < 0. \end{aligned}$$

The equality in each bound is attained if and only if G is regular.

Proof We have

$$ISD_a(G) = \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} = \sum_{uv \in E(G)} \frac{(d_u d_v)^{-a}}{d_u^{-a} + d_v^{-a}}.$$

If $a > 0$, then $2\Delta^{-a} \leq d_u^{-a} + d_v^{-a} \leq 2\delta^{-a}$, and

$$\begin{aligned} \frac{1}{2} \delta^a R_{-a}(G) &= \sum_{uv \in E(G)} \frac{(d_u d_v)^{-a}}{2\delta^{-a}} \leq \sum_{uv \in E(G)} \frac{(d_u d_v)^{-a}}{d_u^{-a} + d_v^{-a}} \\ &\leq \sum_{uv \in E(G)} \frac{(d_u d_v)^{-a}}{2\Delta^{-a}} = \frac{1}{2} \Delta^a R_{-a}(G). \end{aligned}$$

If $a < 0$, then the previous argument gives the converse inequalities.

If G is a regular graph, then the lower and upper bounds are the same, and they are equal to $ISD_a(G)$.

Assume now that the equality in some bound is attained. Thus, by the previous argument we have either $d_u = d_v = \delta$ for every $uv \in E(G)$, or $d_u = d_v = \Delta$ for every $uv \in E(G)$. Hence, G is regular.

The following result relates the ISD_a and ISD_{-a} indices.

Theorem 2 *If G is a graph with minimum degree δ and maximum degree Δ , and $a \in \mathbb{R}$, then*

$$\begin{aligned} \Delta^{-2a} ISD_{-a}(G) &\leq ISD_a(G) \leq \delta^{-2a} ISD_{-a}(G), & \text{if } a > 0, \\ \delta^{-2a} ISD_{-a}(G) &\leq ISD_a(G) \leq \Delta^{-2a} ISD_{-a}(G), & \text{if } a < 0. \end{aligned}$$

The equality in each bound is attained if and only if G is regular.

Proof We have

$$ISD_a(G) = \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} = \sum_{uv \in E(G)} \frac{(d_u d_v)^{-a}}{d_u^{-a} + d_v^{-a}}.$$

Similarly, we obtain the result if $a > 0$.

$$\begin{aligned} \Delta^{-2a} ISD_{-a}(G) &= \sum_{uv \in E(G)} \frac{\Delta^{-2a}}{d_u^{-a} + d_v^{-a}} \leq ISD_a(G) \\ &\leq \sum_{uv \in E(G)} \frac{\delta^{-2a}}{d_u^{-a} + d_v^{-a}} = \delta^{-2a} ISD_{-a}(G). \end{aligned}$$

If $a < 0$, then the previous argument gives the converse inequalities.

If G is a regular graph, then the lower and upper bounds are the same, and they are equal to $ISD_a(G)$.

If the equality in some bound is attained, by the previous argument we have either $d_u = d_v = \delta$ for every $uv \in E(G)$, or $d_u = d_v = \Delta$ for every $uv \in E(G)$. Therefore, G is regular.

The *general sum-connectivity index* was defined in [42] as

$$\chi_a(G) = \sum_{uv \in E(G)} (d_u + d_v)^a.$$

Note that χ_1 is the first Zagreb index M_1 , $2\chi_{-1}$ is the harmonic index H , $\chi_{-1/2}$ is the sum-connectivity index χ , etc.

The following result relates the variable inverse sum deg and the general sum-connectivity indices.

Theorem 3 *If G is a graph and $a \in \mathbb{R} \setminus \{0, 1\}$, then*

$$\chi_{-a}(G) < ISD_a(G) \leq 2^{a-1} \chi_{-a}(G), \quad \text{if } a > 1, \quad (1)$$

$$2^{a-1} \chi_{-a}(G) \leq ISD_a(G) < \chi_{-a}(G), \quad \text{if } 0 < a < 1, \quad (2)$$

$$ISD_a(G) \leq 2^{a-1} \chi_{-a}(G), \quad \text{if } a < 0. \quad (3)$$

The equality in the first or third upper bound or in the second lower bound is attained if and only if each connected component of G is regular.

Proof We want to compute the minimum and maximum values of the function $f : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ given by

$$f(x, y) = \frac{(x + y)^a}{x^a + y^a}.$$

In order to do that, we are going to compute the extremal values of $g(x, y) = (x + y)^a$ with the restrictions $h(x, y) = x^a + y^a = 1$, $x, y > 0$. If (x, y) is a critical point, then there exists $\lambda \in \mathbb{R}$ such that

$$\begin{aligned} a(x + y)^{a-1} &= \lambda a x^{a-1}, \\ a(x + y)^{a-1} &= \lambda a y^{a-1}, \end{aligned}$$

and so, $x = y$; this fact and the equality $x^a + y^a = 1$ give $x = y = 2^{-1/a}$ and $g(2^{-1/a}, 2^{-1/a}) = 2^{a-1}$.

If $a > 0$ and $x \rightarrow 0^+$ (respectively, $y \rightarrow 0^+$), then $y \rightarrow 1$ (respectively, $x \rightarrow 1$) and $g(x, y) \rightarrow 1$.

If $a > 1$, then $1 < g(x, y) \leq 2^{a-1}$ and the upper bound is attained if and only if $x = y$. By homogeneity, we have $1 < f(x, y) \leq 2^{a-1}$ for every $x, y > 0$ and the upper bound is attained if and only if $x = y$.

If $0 < a < 1$, then $2^{a-1} \leq g(x, y) < 1$ and the lower bound is attained if and only if $x = y$. Thus, $2^{a-1} \leq f(x, y) < 1$ for every $x, y > 0$ and the lower bound is attained if and only if $x = y$.

If $a < 0$, then $x, y > 1$. If $x \rightarrow 1^+$ (respectively, $y \rightarrow 1^+$), then $y \rightarrow \infty$ (respectively, $x \rightarrow \infty$) and $g(x, y) \rightarrow 0$. Hence, $0 < g(x, y) \leq 2^{a-1}$ and the upper bound is attained if and only if $x = y$. Thus, $0 < f(x, y) \leq 2^{a-1}$ for every $x, y > 0$ and the upper bound is attained if and only if $x = y$.

Note that if $c_a \leq f(x, y) \leq C_a$, then

$$c_a \frac{1}{(d_u + d_v)^a} \leq \frac{1}{d_u^a + d_v^a} \leq C_a \frac{1}{(d_u + d_v)^a}$$

for every $uv \in E(G)$ and, consequently, $c_a \chi_{-a}(G) \leq ISD_a(G) \leq C_a \chi_{-a}(G)$. These facts give the inequalities.

If G is a connected δ -regular graph with m edges, then

$$2^{a-1} \chi_{-a}(G) = 2^{a-1} (2\delta)^{-a} m = \frac{m}{2\delta^a} = ISD_a(G).$$

By linearity, the equality $2^{a-1} \chi_{-a}(G) = ISD_a(G)$ also holds if each connected component of G is regular.

Assume now that the equality in the first or third upper bound or in the second lower bound is attained. Thus, the previous argument gives that $d_u = d_v$ for every $uv \in E(G)$ and, consequently, each connected component of G is regular.

Note that Theorem 3, with $a = -1$, gives $ISI(G) \leq M_1(G)/4$, a known inequality (see [30, Theorem 4]). Hence, Theorem 3 generalizes [30, Theorem 4].

Remark 1 Note that if we take limits as $a \rightarrow 1$ in Theorem 3, then we obtain by continuity the trivial equality $ISD_1(G) = \chi_{-1}(G)$.

The geometric-arithmetic index was introduced in [40] as

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v}.$$

Although it was introduced in 2009, there are many papers dealing with this index (see, e.g., [9], [10], [18], [22], [25], [28], [32], [33], [34], [40] and the references therein). The predicting ability of the GA index compared with Randić index is reasonably better (see [9, Table 1]). The graphic in [9, Fig.7] (from [9, Table 2], [35]) shows that there exists a good linear correlation between GA and the heat of formation of benzenoid hydrocarbons (the correlation coefficient is equal to 0.972). Furthermore, the improvement in prediction with GA index comparing to Randić index in the case of standard enthalpy of vaporization is more than 9%. That is why one can think that GA_1 index should be considered in the QSPR/QSAR researches.

The following result relates the variable inverse sum deg and the geometric-arithmetic indices.

Theorem 4 *If G is a graph and $a \in \mathbb{R}$, then*

$$\begin{aligned} ISD_a(G) &\geq \frac{1}{2} \Delta^{-a} GA(G), & \text{if } a > 0, \\ ISD_a(G) &\geq \frac{1}{2} \delta^{-a} GA(G), & \text{if } a < 0. \end{aligned}$$

The equality in each bound is attained if and only if G is a regular graph.

Proof We are going to compute the minimum and maximum values of the function $V : [\delta, \Delta] \times [\delta, \Delta] \rightarrow \mathbb{R}^+$ given by

$$V(x, y) = \frac{x + y}{2\sqrt{xy}(x^a + y^a)}.$$

We have

$$\begin{aligned} \frac{\partial V}{\partial x}(x, y) &= \frac{1}{2\sqrt{y}} \frac{x^{1/2}(x^a + y^a) - (x + y)(\frac{1}{2}x^{-1/2}(x^a + y^a) + x^{1/2}ax^{a-1})}{x(x^a + y^a)^2} \\ &= \frac{2x(x^a + y^a) - (x + y)(x^a + y^a + 2ax^a)}{4x^{3/2}y^{1/2}(x^a + y^a)^2} \\ &= \frac{(x - y)(x^a + y^a) - 2a(x + y)x^a}{4x^{3/2}y^{1/2}(x^a + y^a)^2}. \end{aligned}$$

Assume first that $a > 0$. By symmetry, we can assume that $x \leq y$. Thus, $\partial V / \partial x(x, y) < 0$ for $\delta \leq x \leq y \leq \Delta$, and so,

$$V(x, y) \geq V(y, y) = \frac{1}{2y^a} \geq \frac{1}{2} \Delta^{-a},$$

and the equality in the bound is attained if and only if $x = y = \Delta$. Hence,

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} &\geq \frac{1}{2} \Delta^{-a} \frac{2\sqrt{d_u d_v}}{d_u + d_v}, \\ ISD_a(G) &= \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \geq \frac{1}{2} \Delta^{-a} \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v} = \frac{1}{2} \Delta^{-a} GA(G), \end{aligned}$$

and the equality in the bound is attained if and only if $d_u = d_v = \Delta$ for every $uv \in E(G)$, i.e., G is a regular graph.

Assume now that $a < 0$. We can assume that $y \leq x$. Thus, $\partial V / \partial x(x, y) > 0$ for $\delta \leq y \leq x \leq \Delta$, and so,

$$V(x, y) \geq V(y, y) = \frac{1}{2y^a} \geq \frac{1}{2} \delta^{-a},$$

and the equality in the bound is attained if and only if $x = y = \Delta$. Hence,

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} &\geq \frac{1}{2} \delta^{-a} \frac{2\sqrt{d_u d_v}}{d_u + d_v}, \\ ISD_a(G) &= \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \geq \frac{1}{2} \delta^{-a} \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v} = \frac{1}{2} \delta^{-a} AG(G), \end{aligned}$$

and the equality in the bound is attained if and only if $d_u = d_v = \delta$ for every $uv \in E(G)$, i.e., G is a regular graph.

As an inverse variant of the geometric-arithmetic index, in 2015, the arithmetic-geometric index was introduced in [31] as

$$AG(G) = \sum_{uv \in E(G)} \frac{d_u + d_v}{2\sqrt{d_u d_v}}.$$

In [23] it is shown that the arithmetic-geometric index has a good predictive power for entropy of octane isomers. The paper [41] studied spectrum and energy of arithmetic-geometric matrix, in which the sum of all elements is equal to $2AG$. Other bounds of the arithmetic-geometric energy of graphs appeared in [12], [8]. The paper [36] studies optimal AG -graphs for several classes of graphs. In [4], [7], [23] and [29] there are more bounds on the AG index.

The following result relates the variable inverse sum deg and the arithmetic-geometric indices.

Theorem 5 *If G is a graph and $a \in \mathbb{R}$, then*

$$\begin{aligned} ISD_a(G) &\leq \frac{1}{2} \delta^{-a} AG(G), & \text{if } a > 0, \\ ISD_a(G) &\leq \frac{1}{2} \Delta^{-a} AG(G), & \text{if } a < 0. \end{aligned}$$

The equality in each bound is attained if and only if G is a regular graph.

Proof We are going to compute the minimum and maximum values of the function $U : [\delta, \Delta] \times [\delta, \Delta] \rightarrow \mathbb{R}^+$ given by

$$U(x, y) = \frac{2\sqrt{xy}}{(x+y)(x^a + y^a)}.$$

We have

$$\begin{aligned} \frac{\partial U}{\partial x}(x, y) &= \sqrt{y} \frac{x^{-1/2}(x+y)(x^a + y^a) - 2x^{1/2}(x^a + y^a + (x+y)ax^{a-1})}{(x+y)^2(x^a + y^a)^2} \\ &= \sqrt{y} \frac{(x+y)(x^a + y^a) - 2(x(x^a + y^a) + (x+y)ax^a)}{\sqrt{x}(x+y)^2(x^a + y^a)^2} \\ &= \sqrt{y} \frac{(y-x)(x^a + y^a) - 2(x+y)ax^a}{\sqrt{x}(x+y)^2(x^a + y^a)^2}. \end{aligned}$$

Assume first that $a > 0$. By symmetry, we can assume that $x \geq y$. Thus, $\partial U / \partial x(x, y) < 0$ for $\delta \leq y \leq x \leq \Delta$, and so,

$$U(x, y) \leq U(y, y) = \frac{1}{2y^a} \leq \frac{1}{2} \delta^{-a},$$

and the equality in the bound is attained if and only if $x = y = \delta$. Hence,

$$\frac{1}{d_u^a + d_v^a} \leq \frac{1}{2} \delta^{-a} \frac{d_u + d_v}{2\sqrt{d_u d_v}},$$

$$ISD_a(G) = \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \leq \frac{1}{2} \delta^{-a} \sum_{uv \in E(G)} \frac{d_u + d_v}{2\sqrt{d_u d_v}} = \frac{1}{2} \delta^{-a} AG(G),$$

and the equality in the bound is attained if and only if $d_u = d_v = \delta$ for every $uv \in E(G)$, i.e., G is a regular graph.

Assume now that $a < 0$. We can assume that $x \leq y$. Thus, $\partial U / \partial x(x, y) > 0$ for $\delta \leq x \leq y \leq \Delta$, and so,

$$U(x, y) \leq U(y, y) = \frac{1}{2y^a} \leq \frac{1}{2} \Delta^{-a},$$

and the equality in the bound is attained if and only if $x = y = \Delta$. Hence,

$$\frac{1}{d_u^a + d_v^a} \leq \frac{1}{2} \Delta^{-a} \frac{d_u + d_v}{2\sqrt{d_u d_v}},$$

$$ISD_a(G) = \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \leq \frac{1}{2} \Delta^{-a} \sum_{uv \in E(G)} \frac{d_u + d_v}{2\sqrt{d_u d_v}} = \frac{1}{2} \Delta^{-a} AG(G),$$

and the equality in the bound is attained if and only if $d_u = d_v = \Delta$ for every $uv \in E(G)$, i.e., G is a regular graph.

Miličević and Nikolić defined in [20] the *variable first Zagreb index* as

$$M_1^a(G) = \sum_{u \in V(G)} d_u^a,$$

with $a \in \mathbb{R}$. Note that M_1^2 is the first Zagreb index M_1 , M_1^{-1} is the the inverse index ID , $M_1^{-1/2}$ is the zeroth-order Randić index, M_1^3 is the forgotten index F , etc.

Theorem 6 *If G is a graph with m edges, and $a \in \mathbb{R}$, then*

$$ISD_a(G) + M_1^{a+1}(G) \geq \frac{5}{2} m, \quad \text{if } a > 0, \quad (4)$$

$$ISD_a(G) + M_1^{a+1}(G) \geq 2m, \quad \text{if } a < 0. \quad (5)$$

The equality in the first bound is attained if and only if G is a union of path graphs P_2 .

Proof Recall that we have for any function h

$$\sum_{uv \in E(G)} (h(d_u) + h(d_v)) = \sum_{u \in V(G)} d_u h(d_u).$$

In particular,

$$\sum_{uv \in E(G)} (d_u^a + d_v^a) = \sum_{u \in V(G)} d_u^{a+1} = M_1^{a+1}(G).$$

The function $f(x) = x + 1/x$ is strictly decreasing on $(0, 1]$ and strictly increasing on $[1, \infty)$, and so, $f(x) \geq f(1) = 2$ for every $x > 0$. Hence,

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} + d_u^a + d_v^a &\geq 2, \\ ISD_a(G) + M_1^{a+1}(G) &\geq 2m. \end{aligned}$$

If $a > 0$, then $d_u^a + d_v^a \geq 2$ and

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} + d_u^a + d_v^a &\geq f(2) = \frac{5}{2}, \\ ISD_a(G) + M_1^{a+1}(G) &\geq \frac{5}{2}m. \end{aligned}$$

The previous argument gives that the equality in this bound is attained if and only if $d_u = d_v = 1$ for every $uv \in E(G)$, i.e., G is a union of path graphs P_2 .

Theorem 7 *Let G be a graph with minimum degree δ and m edges, and $a \in \mathbb{R}$.*

(1) *If $a > 0$, then*

$$ISD_a(G) + M_1^{a+1}(G) \geq \left(2\delta^a + \frac{1}{2\delta^a}\right)m.$$

(2) *If $\delta > 1$ and $a \leq -\log 2 / \log \delta$, then*

$$ISD_a(G) + M_1^{a+1}(G) \geq \left(2\delta^a + \frac{1}{2\delta^a}\right)m.$$

The equality in each bound is attained if and only if G is regular.

Proof If $a > 0$, then $d_u^a + d_v^a \geq 2\delta^a \geq 2 > 1$. The argument in the proof of Theorem 6 gives

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} + d_u^a + d_v^a &\geq f(2\delta^a) = 2\delta^a + \frac{1}{2\delta^a}, \\ ISD_a(G) + M_1^{a+1}(G) &\geq \left(2\delta^a + \frac{1}{2\delta^a}\right)m. \end{aligned}$$

If $\delta > 1$ and $a \leq -\log 2 / \log \delta < 0$, then $2\delta^a \leq 1$ and $d_u^a + d_v^a \leq 2\delta^a \leq 1$. Thus,

$$\begin{aligned} \frac{1}{d_u^a + d_v^a} + d_u^a + d_v^a &\geq f(2\delta^a) = 2\delta^a + \frac{1}{2\delta^a}, \\ ISD_a(G) + M_1^{a+1}(G) &\geq \left(2\delta^a + \frac{1}{2\delta^a}\right)m. \end{aligned}$$

The previous argument gives that the equality in each bound is attained if and only if $d_u^a + d_v^a = 2\delta^a$ for every $uv \in E(G)$, i.e., $d_u = d_v = \delta$ for every $uv \in E(G)$; and this holds if and only if G is regular.

Theorem 8 *Let G be a graph with maximum degree Δ and m edges, and $a > 0$. Then*

$$ISD_a(G) + M_1^{a+1}(G) \leq \left(2\Delta^a + \frac{1}{2\Delta^a}\right)m,$$

and the equality in the bound is attained if and only if G is regular.

Proof If $a > 0$, then $1 < 2 \leq d_u^a + d_v^a \leq 2\Delta^a$. The argument in the proof of Theorem 6 gives

$$\frac{1}{d_u^a + d_v^a} + d_u^a + d_v^a \leq f(2\Delta^a) = 2\Delta^a + \frac{1}{2\Delta^a},$$

$$ISD_a(G) + M_1^{a+1}(G) \leq \left(2\Delta^a + \frac{1}{2\Delta^a}\right)m.$$

The previous argument gives that the equality in the bound is attained if and only if $d_u^a + d_v^a = 2\Delta^a$ for every $uv \in E(G)$, i.e., $d_u = d_v = \Delta$ for every $uv \in E(G)$; and this holds if and only if G is regular.

We need the following well known result, that provides a converse of the Cauchy-Schwarz inequality (see, e.g., [19, Lemma 3.4]).

Lemma 1 *If $a_j, b_j \geq 0$ and $\omega b_j \leq a_j \leq \Omega b_j$ for $1 \leq j \leq k$, then*

$$\left(\sum_{j=1}^k a_j^2\right)^{1/2} \left(\sum_{j=1}^k b_j^2\right)^{1/2} \leq \frac{1}{2} \left(\sqrt{\frac{\Omega}{\omega}} + \sqrt{\frac{\omega}{\Omega}}\right) \sum_{j=1}^k a_j b_j.$$

If $a_j > 0$ for some $1 \leq j \leq k$, then the equality holds if and only if $\omega = \Omega$ and $a_j = \omega b_j$ for every $1 \leq j \leq k$.

Recall that a (Δ, δ) -biregular graph is a bipartite graph for which any vertex in one side of the given bipartition has degree Δ and any vertex in the other side of the bipartition has degree δ .

Theorem 9 *If G is a graph with m edges, maximum degree Δ and minimum degree δ , and $a \in \mathbb{R} \setminus \{0\}$, then*

$$m^2 \leq ISD_a(G)M_1^{a+1}(G) \leq \frac{(\Delta^a + \delta^a)^2}{4\Delta^a\delta^a} m^2. \quad (6)$$

The equality in the upper bound is attained if and only if G is regular. The equality in the lower bound is attained if G is regular or biregular. Furthermore, if G is a connected graph, then the equality in the lower bound is attained if and only if G is a regular or biregular graph.

Proof Cauchy-Schwarz inequality gives

$$m^2 = \left(\sum_{uv \in E(G)} \frac{1}{\sqrt{d_u^a + d_v^a}} \sqrt{d_u^a + d_v^a} \right)^2 \leq \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \sum_{uv \in E(G)} (d_u^a + d_v^a)$$

$$= ISD_a(G)M_1^{a+1}(G).$$

If $a > 0$, then

$$2\delta^a \leq d_u^a + d_v^a = \frac{\sqrt{d_u^a + d_v^a}}{\frac{1}{\sqrt{d_u^a + d_v^a}}} \leq 2\Delta^a.$$

If $a < 0$, then

$$2\Delta^a \leq d_u^a + d_v^a = \frac{\sqrt{d_u^a + d_v^a}}{\frac{1}{\sqrt{d_u^a + d_v^a}}} \leq 2\delta^a.$$

Lemma 1 gives for every $a \neq 0$

$$\begin{aligned} m^2 &= \left(\sum_{uv \in E(G)} \frac{1}{\sqrt{d_u^a + d_v^a}} \sqrt{d_u^a + d_v^a} \right)^2 \\ &\geq \frac{\sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \sum_{uv \in E(G)} (d_u^a + d_v^a)}{\frac{1}{4} \left(\frac{\Delta^{a/2}}{\delta^{a/2}} + \frac{\delta^{a/2}}{\Delta^{a/2}} \right)^2} \\ &= \frac{4\Delta^a \delta^a}{(\Delta^a + \delta^a)^2} ISD_a(G) M_1^{a+1}(G). \end{aligned}$$

If G is a regular graph, then the lower and upper bounds are the same, and they are equal to $ISD_a(G) M_1^{a+1}(G)$.

Assume now that the equality in the upper bound is attained. Lemma 1 gives $2\Delta^a = 2\delta^a$ and so, $\Delta = \delta$ and G is regular.

If G is a regular or biregular graph, then

$$ISD_a(G) M_1^{a+1}(G) = \frac{m}{\Delta^a + \delta^a} (\Delta^a + \delta^a) m = m^2, \quad (7)$$

and the lower bound is attained.

Assume now that G is a connected graph. By Cauchy-Schwarz inequality, the equality in the lower bound is attained if and only if there exists a constant η such that, for every $uv \in E(G)$,

$$\frac{1}{\sqrt{d_u^a + d_v^a}} = \eta \sqrt{d_u^a + d_v^a}, \quad d_u^a + d_v^a = \eta^{-1}. \quad (8)$$

If $uv, uw \in E(G)$, then

$$\eta^{-1} = d_u^a + d_v^a = d_u^a + d_w^a,$$

and $d_w = d_v$, since $h(t) = t^a$ is a one to one function. Thus, we conclude that (8) is equivalent to the following: for each vertex $u \in V(G)$, every neighbor of u has the same degree. Since G is connected, this holds if and only if G is regular or biregular.

3 Computational study of the ISD_a index on random graphs

Here we follow a recently introduced approach under which topological indices are applied to ensembles of random graphs. Thus instead of computing the index of a single graph, the index average value over a large number of random graphs is measured as a function of the random graph parameters; see the application of this approach to Erdős-Rényi graphs and random regular graphs in [16, 2, 1, 17].

We consider random graphs G from the standard Erdős-Rényi (ER) model $G(n, p)$, i.e., G has n vertices and each edge appears independently with probability $p \in (0, 1)$. The computational study of the ISD_a index we perform below is justified by the random nature of the ER model: since a given parameter pair (n, p) represents an infinite-size ensemble of ER graphs, the computation of a ISD_a index on a single ER graph is irrelevant. In contrast, the computation of $\langle ISD_a \rangle$ over a large ensemble of ER graphs, all characterized by the same parameter pair (n, p) , may provide useful average information about the ensemble. Also, we extend some of the inequalities derived in the previous Section to index average values.

3.1 Scaling of the average ISD_a index on random graphs

In Fig. 1(a) we plot the average variable inverse sum deg index $\langle ISD_a(G) \rangle$ as a function of the probability p of ER graphs of size $n = 1000$. There, we show curves for $a \in [-2, 2]$. As a reference we plot in different colors the curves corresponding to $a = -1$ (blue), $a = 0$ (red), and $a = 1$ (green). Recall that $ISD_{-1}(G) = ISI(G)$ and $ISD_1(G) = \chi_{-1}(G)$. While for $a = 0$, $\langle ISD_0(G) \rangle$ gives half of the average number of edges of the ER graph; that is,

$$\langle ISD_0(G) \rangle = \sum_{uv \in E(G)} \frac{1}{d_u^0 + d_v^0} = \frac{1}{2} |E(G)| = \frac{1}{4} n(n-1)p. \quad (9)$$

Also, from Fig. 1(a) we observe that the curves of $\langle ISD_a(G) \rangle$ show three different behaviors as a function of p depending on the value of a : For $a < a_0$, they grow for small p , approach a maximum value and then decrease when p is further increased. For $a > a_0$, they are monotonically increasing functions of p . For $a = a_0$ the curves saturate above a given value of p . Here $a_0 = 1$.

Moreover, when $np \gg 1$, we can write $d_u \approx d_v \approx \langle d \rangle$, with

$$\langle d \rangle \approx (n-1)p. \quad (10)$$

Therefore, for $np \gg 1$, $\langle ISD_a(G) \rangle$ is well approximated by:

$$\langle ISD_a(G) \rangle \approx \sum_{uv \in E(G)} \frac{1}{\langle d \rangle^a + \langle d \rangle^a} = |E(G)| \frac{1}{2 \langle d \rangle^a} \approx \frac{n}{4} [(n-1)p]^{1-a}. \quad (11)$$

In Fig. 1(b), we show that Eq. (11) (red-dashed lines) indeed describes well the data (thick black curves) for $np \geq 10$.




Fig. 1 (a,b) Average variable inverse sum deg index $\langle ISD_a(G) \rangle$ as a function of the probability p of Erdős-Rényi graphs of size $n = 1000$. Here we show curves for $\alpha \in [-2, 2]$ in steps of 0.2 (from top to bottom). Blue, red and green curves in (a) correspond to $a = -1$, $a = 0$ and $a = 1$, respectively. The red dashed lines in (b) are Eq. (11). The blue dashed line in (b) marks $\langle d \rangle = 10$. (c) $\langle ISD_a(G) \rangle$ as a function of the probability p of ER graphs of four different sizes n . (d) $\langle ISD_a(G) \rangle / n$ as a function of the average degree $\langle d \rangle$. Same curves as in panel (c). The inset in (d) is the enlargement of the cyan rectangle. All averages are computed over $10^7/n$ random graphs.

Now in Fig. 1(c) we show $\langle ISD_a(G) \rangle$ as a function of the probability p of ER random graphs of four different sizes n . It is quite clear from this figure that the blocks of curves, characterized by the different graph sizes, display similar curves but displaced on both axes. Thus, our next goal is to find the scaling parameters that make the blocks of curves to coincide.

First, we recall that the average degree $\langle d \rangle$, see Eq. (10), is known to scale both topological and spectral measures applied to ER graphs. In particular, $\langle d \rangle$ was shown to scale the normalized Randic index [16], the normalized Harmonic [17] index, as well as several variable degree-based indices [1] on ER graphs. Thus, we expect $\langle ISD_a(G) \rangle \propto f(\langle d \rangle)$. Second, we observe in Fig. 1(c) that the effect of increasing the graph size is to displace the blocks of curves $\langle ISD_a(G) \rangle$ vs. p , characterized by the different graph sizes, upwards in the y -axis. Moreover, the fact that these blocks of curves, plotted in semi-log scale, are shifted the same amount on the y -axis when doubling n is a clear signature of scalings of the form $\langle ISD_a(G) \rangle \propto n^\beta$. By plotting $\langle ISD_a(G) \rangle$ vs. n for given values of p (not shown here) we conclude that $\beta = 1$ for all a .

Therefore, in Fig. 1(d) we plot $\langle ISD_a(G) \rangle / n$ as a function of $\langle d \rangle$ showing that all curves are now properly scaled; i.e. the blocks of curves painted in different colors for different graph sizes fall on top of each other (see a detailed view in the inset of this figure). Moreover, following Eq. (11), we obtain

$$\frac{\langle ISD_a(G) \rangle}{n} \approx \frac{1}{4} \langle d \rangle^{1-a}. \quad (12)$$

We have verified that Eq. (12) is valid when $\langle d \rangle \geq 10$.

3.2 Inequalities of the average ISD_a index on random graphs

Most inequalities obtained in the previous Section are not restricted to any particular type of graph. Thus, they should also be valid for random graphs and, moreover, can be extended to index average values, as needed in computational studies of random graphs.

Now, in order to ease the computational validation of some of the inequalities derived in the previous Section, we:

- (i) write the right inequality of Eq. (1) in Theorem 3 as

$$0 \leq \langle 2^{a-1} \chi_{-a}(G) - ISD_a(G) \rangle, \quad \text{if } a > 1, \quad (13)$$

- (ii) write the left inequality of Eq. (2) in Theorem 3 as

$$0 \leq \langle ISD_a(G) - 2^{a-1} \chi_{-a}(G) \rangle, \quad \text{if } 0 < a < 1, \quad (14)$$

- (iii) write the inequality of Eq. (3) in Theorem 3 as

$$0 \leq \langle 2^{a-1} \chi_{-a}(G) - ISD_a(G) \rangle, \quad \text{if } a < 0, \quad (15)$$

- (iv) write the inequality of Eq. (4) in Theorem 6 as

$$\frac{5}{2} \langle m \rangle \leq \langle ISD_a(G) + M_1^{a+1}(G) \rangle, \quad \text{if } a > 0, \quad (16)$$

- (v) write the inequality of Eq. (5) in Theorem 6 as

$$2 \langle m \rangle \leq \langle ISD_a(G) + M_1^{a+1}(G) \rangle, \quad \text{if } a < 0, \quad (17)$$

and

- (vi) write the left inequality of Eq. (6) in Theorem 9 as

$$\langle m^2 \rangle \leq \langle ISD_a(G) M_1^{a+1}(G) \rangle. \quad (18)$$

Therefore, in Figs. 2(a-f) we plot the r.h.s. of the inequalities (13-18), respectively, as a function of the probability p of ER graphs of size $n = 100$.

In particular, since the curves in Figs. 2(a-c) are all positive, the inequalities (13-15) are easily validated. Now, in order to validate inequalities (16,17) we include (as dashed lines) in both Fig. 2(d) and Fig. 2(e) the functions $(5/2) \langle m \rangle$ vs. p (red) and $2 \langle m \rangle$ vs. p (blue); where we used $\langle m \rangle = n(n-1)p/2$. Then,




Fig. 2 In panels (a-f) we plot the r.h.s. of relations (13-18), respectively, as a function of the probability p of Erdős-Rényi graphs of size $n = 100$. In (a) $a \in [1.1, 2]$ in steps of 0.1, in (b) $a \in [0.1, 0.9]$ in steps of 0.1, in (c) $a \in [-2, -0.1]$ in steps of 0.1, in (d) $a \in [0.1, 2]$ in steps of 0.1, in (e) $a \in [-2, -0.1]$ in steps of 0.1, and in (f) $a \in [-2, 2]$ in steps of 0.2. Dashed lines in (d,e) are $(5/2) \langle m \rangle$ (red) and $2 \langle m \rangle$ (blue). The red dashed line in (f) is $\langle m^2 \rangle$. Here we used $\langle m \rangle = n(n-1)p/2$. Insets in (d-f) are enlargements of the cyan rectangles of the corresponding main panels. Blue arrows in panels (a,c-e) indicate increasing a . All averages are computed over $10^7/n$ random graphs.

we can clearly see that all curves $\langle \text{ISD}_a(G) + M_1^{a+1}(G) \rangle$ vs. p in Fig. 2(d) lie above the red dashed line, corresponding to $(5/2) \langle m \rangle$; while all curves $\langle \text{ISD}_a(G) + M_1^{a+1}(G) \rangle$ vs. p in Fig. 2(e) lie above the blue dot-dashed line, corresponding to $2 \langle m \rangle$. This can be better appreciated in the enlargements shown in the panel insets. Finally, in Fig. 2(f) we include, as a red dashed line, the function $\langle m^2 \rangle$ vs. p to clearly show that all curves $\langle \text{ISD}_a(G) M_1^{a+1}(G) \rangle$ vs. p lie above it, as stated in inequality (18). Moreover, note that the equality in (18) is attained for $p \rightarrow 1$. This is indeed expected since for $np \gg 1$ we can

write

$$\begin{aligned}\langle ISD_a(G)M_1^{a+1}(G) \rangle &= \left\langle \sum_{uv \in E(G)} \frac{1}{d_u^a + d_v^a} \sum_{uv \in E(G)} d_u^a + d_v^a \right\rangle \\ &\approx \left\langle \sum_{uv \in E(G)} \frac{1}{\langle d \rangle^a + \langle d \rangle^a} \sum_{uv \in E(G)} \langle d \rangle^a + \langle d \rangle^a \right\rangle \\ &= \left\langle \frac{m}{\langle d \rangle^a + \langle d \rangle^a} m(\langle d \rangle^a + \langle d \rangle^a) \right\rangle = \langle m^2 \rangle,\end{aligned}$$

which we have observed to be valid for several graph sizes when $\langle d \rangle \geq 10$.

4 Summary

In this work we performed analytical and computational studies of the variable inverse sum deg index $ISD_a(G)$. First, we analytically obtained new inequalities connecting $ISD_a(G)$ with other well-known topological indices such as the Randić index, the general sum-connectivity index, the geometric-arithmetic index, the arithmetic-geometric index, as well as the variable first Zagreb index. Then, we computationally validated some of the obtained inequalities on ensembles of Erdős-Rényi graphs $G(n, p)$ characterized by n vertices connected independently with probability $p \in (0, 1)$. Additionally, we showed that the ratio $\langle ISD_a(G) \rangle / n$ depends only on the average degree $\langle d \rangle = (n - 1)p$.

Acknowledgements The research of W.C., J.M.R. and J.M.S. was supported by a grant from Agencia Estatal de Investigación (PID2019-106433GB-I00/AEI/10.13039/501100011033), Spain. J.M.R. was supported by the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with UC3M in the line of Excellence of University Professors (EPUC3M23), and in the context of the V PRICIT (Regional Programme of Research and Technological Innovation).

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. R. Aguilar-Sánchez, I. F. Herrera-González, J. A. Méndez-Bermúdez, J. M. Sigarreta, Computational properties of general indices on random networks, *Symmetry* **12** (2020) 1341.
2. R. Aguilar-Sánchez, J. A. Méndez-Bermúdez, F. A. Rodrigues, J. M. Sigarreta-Almira, Topological versus spectral properties of random geometric graphs, *Phys. Rev. E* **102** (2020) 042306.
3. B. Bollobás, P. Erdős, Graphs of extremal weights, *Ars Combin.* **50** (1998) 225.
4. W. Carballosa, A. Granados, J. A. Méndez-Bermúdez, D. Pestana, A. Portilla, Computational properties of the arithmetic-geometric index, Submitted.

5. W. Carballosa, J. M. Rodríguez, J. M. Sigarreta, Extremal problems on the variable sum exdeg index, *MATCH Commun. Math. Comput. Chem.* **84:3** (2020) 753.
6. H. Chen, H. Deng, The inverse sum indeg index of graphs with some given parameters, *Discr. Math. Algor. Appl.* **10:01**, 1850006 (2018).
7. S.-Y. Cui, W. Wang, G.-X. Tian, B. Wu, On the arithmetic-geometric index of graphs, *MATCH Commun. Math. Comput. Chem.* **85:1** (2021) 87.
8. K. C. Das, I. Gutman, Degree-based energies of graphs, *Linear Algebra Appl.* **554** (2018) 185.
9. K. C. Das, I. Gutman, B. Furtula, Survey on Geometric-Arithmetic Indices of Graphs, *MATCH Commun. Math. Comput. Chem.* **65** (2011) 595.
10. K. C. Das, I. Gutman, B. Furtula, On first geometric-arithmetic index of graphs, *Discrete Appl. Math.* **159** (2011) 2030.
11. F. Falahati-Nezhad, M. Azari, T. Došlić, T., Sharp bounds on the inverse sum indeg index, *Discrete Appl. Math.* **217**, (2017) 185.
12. X. Guo, Y. Gao, Arithmetic-geometric spectral radius and energy of graphs, *MATCH Commun. Math. Comput. Chem.* **83** (2020) 651.
13. I. Gutman, M. Matejić, E. Milovanović, I. Milovanović, Lower bounds for inverse sum indeg index of graphs, *Kragujevac J. Math.* **44:4** (2020) 551.
14. I. Gutman, J. M. Rodríguez, J. M. Sigarreta, Linear and non-linear inequalities on the inverse sum indeg index, *Discrete Appl. Math.* **258** (2019) 123.
15. X. Li, I. Gutman, Mathematical Aspects of Randić Type Molecular Structure Descriptors, Univ. Kragujevac, Kragujevac, 2006.
16. C. T. Martínez-Martínez, J. A. Méndez-Bermúdez, J. M. Rodríguez, J. M. Sigarreta, Computational and analytical studies of the Randić index in Erdős-Rényi models. *Appl. Math. Comput.* **377** (2020) 125137.
17. C. T. Martínez-Martínez, J. A. Méndez-Bermúdez, J. M. Rodríguez, J. M. Sigarreta, Computational and analytical studies of the harmonic index in Erdős-Rényi models, *MATCH Commun. Math. Comput. Chem.* **85** (2021) 395.
18. A. Martínez-Pérez, J. M. Rodríguez, New lower bounds for the Geometric-Arithmetic index, *MATCH Commun. Math. Comput. Chem.* **79** (2018) 451.
19. A. Martínez-Pérez, J. M. Rodríguez, J. M. Sigarreta, A new approximation to the geometric-arithmetic index, *J. Math. Chem.* **56:7** (2018) 1865.
20. A. Miličević, S. Nikolić, On variable Zagreb indices, *Croat. Chem. Acta* **77** (2004) 97.
21. A. Mingqiang, X. Liming, Some results on the inverse sum indeg index of a graph, *Inf. Process. Lett.* **134**, (2018) 42.
22. M. Mogharrab, G. H. Fath-Tabar, Some bounds on GA_1 index of graphs, *MATCH Commun. Math. Comput. Chem.* **65** (2010) 33.
23. E. D. Molina, J. M. Rodríguez, J. L. Sánchez, J. M. Sigarreta, Some Properties of the Arithmetic-Geometric Index, *Symmetry* (2021) 13, 857.
24. S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, The Zagreb Indices 30 years after, *Croat. Chem. Acta* **76** (2003) 113.
25. D. Pestana, J. M. Sigarreta, E. Tourís, Geometric-arithmetic index and line graph, *J. Math. Chem.* **57:5** (2019) 1427.
26. J. Rada, J. M. Rodríguez, J. M. Sigarreta, Optimization problems for the variable inverse sum deg index, Submitted.
27. M. Randić, On characterization of molecular branching, *J. Am. Chem. Soc.* **97** (1975) 6609.
28. J. M. Rodríguez, J. A. Rodríguez-Velázquez, J. M. Sigarreta, New inequalities involving the Geometric-Arithmetic Index, *MATCH Commun. Math. Comput. Chem.* **78:2** (2017) 361.
29. J. M. Rodríguez, J. L. Sánchez, J. M. Sigarreta, E. Tourís, Bounds on the arithmetic-geometric index, *Symmetry* 2021, 13(4), 689.
30. J. Sedlar, D. Stevanović, A. Vasilyev, On the inverse sum indeg index, *Discrete Appl. Math.* **184** (2015) 202.
31. V. S. Shegehall, R. Kanabur, Arithmetic-geometric indices of path graph, *J. Math. Comput. Sci.* **16** (2015) 19.
32. J. M. Rodríguez, J. M. Sigarreta, On the Geometric-Arithmetic Index, *MATCH Commun. Math. Comput. Chem.* **74** (2015) 103.

-
33. J. M. Rodríguez, J. M. Sigarreta, Spectral properties of geometric-arithmetic index, *Appl. Math. Comput.* **277** (2016) 142.
 34. J. M. Sigarreta, Bounds for the geometric-arithmetic index of a graph, *Miskolc Math. Notes* **16** (2015) 1199.
 35. TRC Thermodynamic Tables. Hydrocarbons; Thermodynamic Research Center, The Texas A & M University System: College Station, TX, 1987.
 36. S. Vujošević, G. Popivoda, Ž. Kovijanić Vukičević, B. Furtula, R. Škrekovski, Arithmetic-geometric index and its relations with geometric-arithmetic index, *Appl. Math. Comput.* **391** (2021) #125706.
 37. D. Vukičević, M. Gašperov, Bond additive modeling 1. Adriatic indices, *Croat. Chem. Acta* **83** (2010) 243.
 38. D. Vukičević, Bond additive modeling 2. Mathematical properties of max-min rodeg index, *Croat. Chem. Acta* **83** (2010) 261.
 39. D. Vukičević, Bond additive modeling 4. QSPR and QSAR studies of the variable Adriatic indices, *Croat. Chem. Acta* **84** (2011) 87.
 40. D. Vukičević, B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, *J. Math. Chem.* **46** (2009) 1369.
 41. L. Zheng, G.-X. Tian, S.-Y. Cui, On spectral radius and energy of arithmetic-geometric matrix of graphs, *MACTH Commun. Math. Comput. Chem.* **83** (2020) 635.
 42. B. Zhou, N. Trinajstić, On general sum-connectivity index, *J. Math. Chem.* **47** (2010) 210.