ON THE NUMBER OF ERGODIC MEASURES OF MAXIMAL ENTROPY FOR PARTIALLY HYPERBOLIC DIFFEOMORPHISMS WITH COMPACT CENTER LEAVES

JOAS ELIAS ROCHA AND ALI TAHZIBI

ABSTRACT. In this paper, we study the number of ergodic measures of maximal entropy for partially hyperbolic diffeomorphisms defined on 3-torus with compact center leaves. Assuming the existence of a periodic leaf with Morse-Smale dynamics we prove a sharp upper bound for the number of maximal measures in terms of the number of sources and sinks of Morse-Smale dynamics. A well-known class of examples for which our results apply are the so-called Kan-type diffeomorphisms admitting physical measures with intermingled basins.

1. INTRODUCTION

Measures of maximal entropy (M.M.E) are global maxima for the Kolmogorov-Sinai entropy map $\mu \to h_{\mu}(f)$ where $f: M \to M$ is a continuous transformation defined on a compact metric space M. In this paper by m.m.e we mean **ergodic** measures of maximal entropy. These measures, when exist, are considered among natural invariant measures of f. Newhouse [17] studied an upper bound on the defect in uppersemicontinuity of topological and metric entropy and proved upper semi continuity of $\mu \to h_{\mu}(f)$ under C^{∞} regularity condition on f. As a consequence any C^{∞} diffeomorphism of a compact manifold admits at least one measure of maximal entropy.

However, there are examples of (with high regularity C^r , $0 < r < \infty$) diffeomorphisms without any measure of maximal entropy (See for instance [16], [6].)

It is natural to study the existence of maximal entropy measures among diffeomorphisms which preserve some additional structure. In the context of dynamical systems admitting some hyperbolicity, the question of the existence of maximal measures has been studied by several authors. After the classical results of existence and uniqueness in the uniformly hyperbolic case, a next step is to study the partially hyperbolic setting. See for instance [8], [23], [5], [20] among other results in the references.

For a diffeomorphism $f: M \to M$ of a compact manifold to itself recall the norm and conorm with respect to a subspace of $V \subset T_x M$ for some $x \in M$: $\|Df|V\| := \max\{\|Tf(v)\| : v \in V, \|v\| = 1\}$ and

$$\operatorname{conorm}(Df|V) := \min\{\|Tf(v)\| : v \in V, \|v\| = 1\}.$$

A splitting $E \oplus F$ is dominated¹ if it is nontrivial, invariant, and if there is some $N \ge 1$ such that, for all $x \in M$:

$$||Df^N|E_x|| < \frac{1}{2}\operatorname{conorm}(Df^N|F_x).$$

Definition 1.1. A diffeomorphism is (strongly) partially hyperbolic if there is an invariant splitting of the tangent bundle: $TM = E^s \oplus E^c \oplus E^u$ such that

A.T is supported by FAPESP 107/06463-3 and CNPq (PQ) 303025/2015-8.

¹This is sometimes called *pointwise* domination, see [1].

 $E^s \oplus (E^c \oplus E^u)$ and $(E^s \oplus E^c) \oplus E^u$ are dominated, E^s is uniformly contracted, and E^u is uniformly expanded.

It is a well-known fact that there are foliations \mathcal{F}^* tangent to the distributions E^* for * = s, u. The leaf of \mathcal{F}^* containing x will be denoted by $\mathcal{F}^*(x)$, for * = s, u.

In general it is not true that there is a foliation tangent to E^c . It can fail to be true even if dim $E^c = 1$ ([21]) for partially hyperbolic dynamics defined on \mathbb{T}^3 . We shall say that f is *dynamically coherent* if there exist invariant foliations tangent to $E^{c*} = E^c \oplus E^*$ for * = s, u (and then, to E^c). Along this paper all partially hyperbolic diffeomorphisms will be dynamically coherent.

We shall say that a set X is *-saturated if it is a union of leaves of the strong foliations \mathcal{F}^* for * = s or u. We also say that X is su-saturated if it is both s- and u-saturated. The accessibility class Acc(x) of the point $x \in M$ is the minimal susaturated set containing x. Note that the accessibility classes form a partition of M. In case there is some $x \in M$ whose accessibility class is M, then the diffeomorphism f is said to have the accessibility property. This is equivalent to say that any two points of M can be joined by a path which is piecewise tangent to E^s or to E^u .

If the center bundle is one dimensional or decomposes into one dimensional subbundles the results in [9], [15] show the existence of m.m.e's. However the quest for uniqueness or finiteness of ergodic measures of maximal entropy is a much more delicate problem in this setting.

In this paper we address the problem of determining the number of ergodic measures of maximal entropy of C^2 partially hyperbolic diffeomorphisms on \mathbb{T}^3 with compact center leaves. There is another important class of partially hyperbolic diffeomorphisms on \mathbb{T}^3 which are isotopic to Anosov diffeomorphisms. For this class R. Ures ([23]) proved uniqueness of measure of maximal entropy (See also a similar result in [5]).

In [20] the authors proved the following dichotomy for $C^{1+\alpha}$, $\alpha > 0$ dynamically coherent partially hyperbolic with accessibility property and compact center leaves: Either f is "rotation type" with a unique non hyperbolic measure of maximal entropy or there exist only finitely many hyperbolic ergodic m.m.e and there exists at least one measure with negative and another with positive center Lyapynov exponent. More precisely they proved the following theorem.

Theorem 1.2. [20] Let $f: M \to M$ be a $C^{1+\alpha}$ partially hyperbolic diffeomorphism of a 3-dimensional closed manifold M. Assume that f is dynamically coherent with compact one dimensional central leaves and has the accessibility property. Then fhas finitely many ergodic measures of maximal entropy. There are two possibilities:

- (1) (rotation type) f has a unique entropy maximizing measure μ . The center Lyapunov exponent $\lambda_c(\mu)$ vanishes and (f,μ) is isomorphic to a Bernoulli shift,
- (2) (generic case) f has more than one ergodic entropy maximizing measure, all of which with non vanishing center Lyapunov exponent. The center Lyapunov exponent $\lambda_c(\mu)$ is nonzero and (f,μ) is a finite extension of a Bernoulli shift for any such measure μ . Some of these measures have positive central exponent and some have negative central exponent.

Moreover, the diffeomorphisms fulfilling the conditions of the second item form a C^1 -open and C^{∞} -dense subset of the dynamically coherent partially hyperbolic diffeomorphisms with compact one dimensional central leaves.

In the generic case of the above theorem (second item), the number of ergodic m.m.e's is larger than one. However still one can hope uniquenss of m.m.e fixing the sign of center Lyapunov exponent. In this paper we analyse the number of

m.m.e with positive (negative) exponent assuming some condition on the dynamics of one periodic leaf.

Ures, Viana and J. Yang [25] proved an optimal quantitative result when $M \neq \mathbb{T}^3$ is a nil-manifold:

Theorem 1.3. [25] Let f be a C^2 partially hyperbolic diffeomoprhism on a 3-dimensional nilmanifold $M \neq \mathbb{T}^3$. Then

- either f has a unique maximal meaure, in which case f is conjugate to rotation extension of an Anosov diffeomorphism (rotation type) and the maximal measure is supported on the whole manifold and has vanishing center exponent;
- or f has exactly two ergodic maximal measures μ⁺, μ⁻, with positive and negative center exponents, respectively.

We emphasize that partially hyperbolic diffeomorphisms on 3-nilmanifolds other than \mathbb{T}^3 have some nice topological properties ([22], [12]) which are essential to the proof of the above theorem: Any such diffeomorphism is accessible, dynamically coherent and it has a unique compact, invariant, *u*-saturated (respectively, *s*-saturated) minimal subset.

In the setting of partially hyperbolic diffeomorphisms of \mathbb{T}^3 there is no "general result" on the number of measures of maximal entropy. Multiplying Anosov diffeomorphism on \mathbb{T}^2 to an appropriate one dimensional dynamics it is easy to get simples examples with any (countable)number of hyperbolic m.m.e and co-existence of hyperbolic and non-hyperbolic maximal measures. So, the interesting question is to verify the number of m.m.e's under some restriction hypothesis on the center dynamics or accessibility of the partially hyperbolic system.

In Theorem 2.1 (resp. 2.7) we assume that the dynamics of a center leaf is Morse-Smale (resp. with irrational rotation number). We also give examples (Theorem 2.6 and section 4.4) to show the optimality of results.

2. Statement of results

Let $f: \mathbb{T}^3 \to \mathbb{T}^3$ be a dynamically coherent partially hyperbolic diffeomorphism with compact center leaves. Let $\mathcal{F}^c(x)$ be a periodic leaf with period n. We say fhas k-Morse-Smale dynamics on this leaf if $f^n: \mathcal{F}^c(x) \to \mathcal{F}^c(x)$ is a Morse-Smale dynamics with k sinks and k sources. As an easy example of such dynamics consider direct product $f := A \times g$ where A is an Anosov diffeomorphism on \mathbb{T}^2 and gis an apropriate (to guarantee partial hyperbolicity) 1–Morse-Smale dynamics on the circle. It is clear that such an example admits exactly one ergodic measures of maximal entropy with positive center Lyapunov exponent and another ergodic m.m.e's with negative exponent. However, f_k is not accessible and it is not topologically transitive. For non trivial and transitive example we may consider Kan-Type diffeomorphisms (See Section 4.3). These examples have two center leaves with 1-Morse Smale dynamics.

Here we prove a dichotomy for all partially hyperbolic dynamics which admit a 1–Morse-Smale dynamics. In the following theorem by an su-torus we mean a 2–torus which is an accessibility class.

Theorem 2.1. Let $f : \mathbb{T}^3 \to \mathbb{T}^3$ be a C^2 dynamically coherent partially hyperbolic diffeomorphism with compact center leaves. Suppose that f has a periodic center leaf with 1–Morse-Smale dynamics then at least one of the following occurs:

- (1) Either the number of ergodic maximal measures is precisely two (one with positive center exponent and one with negative center exponent) or
- (2) there exists an invariant su-torus.

Corollary 2.2. If f satisfies the hypothesis of the above theorem and is accessible then f admits one m.m.e with positive center Lyapunov exponent and one m.m.e with negative center Lyapunov exponent.

Addendum 2.3. In the above theorem if f admits a periodic center leaf with k-Morse-Smale dynamics then either f admits at most $2(k+\lfloor\frac{k}{2}\rfloor)$ hyperbolic m.m.e or there exists an invariant su-torus.

Remark 2.4. Let us emphasize that even assuming accessibility, without our hypothesis on Morse-Smale periodic leaf, no general result on the number of m.m.e's is known. It is well worth to recall that for any $l \ge 1$ after a C^l -local perturbation one can change the dynamics of a periodic leaf to obtain a k-Morse-Smale dynamics for some $k \in \mathbb{N}$.

To prove the above theorem we consider two following disjoint cases:

- f admits an ergodic measure of maximal entropy with zero center exponent.
- f admits no ergodic measure of maximal entropy with vanishing center exponent.

In the first case we will show the existence of an su-torus. In the second case we prove that either there exists a unique m.m.e with negative center exponent (and a unique one with positive center exponent) or there exists an invariant su-torus.

Another result which is immediate consequence of Proposition 3.3 is the following:

Proposition 2.5. If f satisfies the hypothesis of Theorem 2.1 and all m.m.e's are hyperbolic then there are at most two m.m.e with negative center exponents (Similarly there are at most two m.m.e with positive center exponent.)

Proof. Suppose μ_i , i = 1, 2, 3 be three m.m.e with negative center Lyapunov exponent. We know that the support of each μ_i intersects any center leaf (See for instance inside the proof of Theorem 2.1). In particular supp (μ_i) intersects the periodic center leaf admitting 1–Morse-Smale dynamics. By definition of 1–Morse-Smale dynamics there exists one sink and one source on this periodic center leaf. As the support is closed invariant set and by Proposition 3.3 supports of μ_i are disjoint we get a contradiction.

In the sequel we give examples of such dynamics with four ergodic measures of maximal entropy (two with positive and two with negative center exponent). We also give a topologically transitive example with three ergodic measures of maximal entropy. The construction is based on examples of non-invertible maps appeared in the work of Nuñez-Madariaga, S. Ramirez and C. Vasquez [19]. The celebrated Kan example in the annulus is an endomorphism defined on $\mathbb{S}^1 \times [0, 1]$ with two physical measures with intermingled basins. Nuñez-Madariaga, S. Ramirez and C. Vasquez show that besides these two physical measures (Lebesgue measure on \mathcal{S}^1) which are measure of maximal entropy (besides being physical) there exists a third measure of maximal entropy. They describe this third measure as limit of periodic measures. We interpret this third measure as a twin measure in the diffeomorphism setting.

Theorem 2.6. There exist partially hyperbolic diffeomorphisms satisfying the hypothesis of the above theorem with 4 m.m.e's two of them with negative and two others with positive center exponent. There is also topologically transitive example with 2 m.m.e with negative center exponent and one with positive center exponent.

Observe that although the second example in the above theorem is topologically transitive, still we are not able to find an accessible example:

Question 1. Is there any accessible partially hyperbolic diffeomorphism satisfying hypothesis of Theorem 2.1 with more than one ergodic m.m.e with positive center exponent?

We still do not know a clear dichotomy on the hyperbolicity of m.m.e's under the hypothesis of Theorem 2.1.

Question 2. Is there any example of partially hyperbolic diffeomorphism satisfying hypothesis of Theorem 2.1 admitting both hyperbolic and non-hyperbolic measures of maximal entropy?

We emphasize that in the Kan tye examples stated in Theorem 2.6 the coexistence of hyperbolic and non-hyperbolic m.m.e's is ruled out using the intermingled property of hyperbolic m.m.e's and invariance principle argument.

In the next theorem we assume the existence of a periodic leaf with irrational rotation number dynamics and prove a dichotomy.

Theorem 2.7. Let $f : \mathbb{T}^3 \to \mathbb{T}^3$ be a C^2 -partially hyperbolic diffeomorphism, dynamically coherent with compact central leaves with one periodic leaf with irrational rotation number then:

- (1) Either there is unique measure of maximal entropy μ , moreover $\lambda^{c}(\mu) = 0$ and f is conjugate to rotation extension of Anosov homeomorphism or
- (2) f admits exactly two ergodic measures of maximal entropy. Both measures are hyperbolic and have opposite sign of center Lyapunov exponent.

Observe that in the above theorem we are not assuming accessibility of f. Instead we are assuming existence of a periodic leaf with irrational rotation dynamics. We also give examples (subsection 4.4) for each item of the dichotomy announced in the above theorem.

3. QUOTIENT DYNAMICS AND CONDITIONAL MEASURES

Let $f : \mathbb{T}^3 \to \mathbb{T}^3$ be a C^{1-} partially hyperbolic diffeomorphism with compact center leaves. We define the quotient space M_c and quotien dynamics f_c by means of the natural projection $\pi : \mathbb{T}^3 \to M_c := \mathbb{T}^3/\mathcal{F}^c$ which sends every point in $\mathcal{F}^c(x)$ to [x]. It is known that M_c is homeomorphim to \mathbb{T}^2 and f_c is an Anosov homeomorphism (See Theorem 3 in [20]). So it is topologically conjugate to an Anosov diffeomorphism on \mathbb{T}^2 .

From now on we fix some notations: $\mathcal{F}^s, \mathcal{F}^c$ and \mathcal{F}^u are respectively stable, center and the unstable foliation invariant by f. The points in M_c are denoted by [.]. For any point $[x] \in M_c$ by $\mathcal{W}^s([x])$ and $\mathcal{W}^u([x])$ we mean the stable and unstable set of [x]. As f is conjugate to an Anosov diffeomorphism on \mathbb{T}^2 it comes out that both (topological) foliations \mathcal{W}^u and \mathcal{W}^s are minimal. Observe that

- $\pi(\mathcal{F}^{cs})(x) = \mathcal{W}^s(\pi(x)),$
- $\pi(\mathcal{F}^{cu})(x) = \mathcal{W}^u(\pi(x)).$

Moreover, f_c has a unique measure of maximal entropy η which is fully supported on M_c . This measure has local product structure: For any $[x] \in M_c$ there exists an open set U around [x], a homeomorphism

$$\alpha: \mathcal{W}^s_{loc}([x]) \times \mathcal{W}^u_{loc}([x]) \to U$$

and measures $\eta_{[x]}^u, \eta_{[x]}^s$ supported on respectively $\mathcal{W}_{loc}^s([x])$ and $\mathcal{W}_{loc}^u([x])$ such that:

$$\eta|_U \sim \alpha_*(\eta^u_{[x]} \times \eta^s_{[x]}),$$

where \sim denotes equivalence of measures.

Take any invariant measure μ for f and let $\nu = \mu \circ \pi^{-1}$. By the Ledrappier-Walters variational principle [14]

$$\sup_{\hat{\mu}:\hat{\mu}\circ\pi^{-1}=\nu}h_{\hat{\mu}}(f)=h_{\nu}(f_c)+\int_{M_c}h(f,\pi^{-1}(y))d\nu(y).$$

Since $\pi^{-1}(y), y \in M_c$, is a circle and its iterates have bounded length we have that $h_{top}(f, \pi^{-1}(y)) = 0$, that is, fibers does not contribute to the entropy. Hence, by the above equality and the well-known fact that $h_{\mu}(f) \geq h_{\nu}(f_c)$ we conclude that $h_{\mu}(f) = h_{\nu}(f_c)$. Using the usual variational principle this implies that the topological entropies of f and f_c coincide. In particular, the set of entropy maximizing measures of f coincides with the subset of ergodic measures which projects down to η , the entropy maximizing measure of f_c .

3.1. Invariance principle and support of m.m.e. In the setting we are working, as center foliation is given by compact leaves and the quotient space is homeomorphic to \mathbb{T}^2 which is a separable metric space we can apply Rokhlin disintegration theorem and conclude that for any probability μ there is a unique family of conditional measures $\mu_{[x]}^c$ (probability supported on $\mathcal{F}^c(x)$) defined $\eta := \pi_* \mu$ -almost every where and

$$\mu = \int_{M_c} \mu_{[x]}^c d\eta([x]).$$

Sometimes, it is more conveniente to use the notation $\{\mu_x^c\}_{x\in M}$ for conditional measures along center foliation. Using this notation we have $\mu_x^c = \mu_y^c$ for $y \in \mathcal{F}^c(x)$ and μ_x^c 's are probability measures where

$$\mu = \int_M \mu_x^c d\mu(x).$$

As we mentioned before, if μ is any measure of maximal entropy then $\nu = \pi_* \mu$ is maximal entropy measure for the quotient dynamics which is Anosov. So ν has local product structure and one may apply the following invariance principle due to Avila-Viana. Recall that by center Lyapunov exponent $\lambda_c(x)$ we mean the following limit:

$$\lim_{n \to \infty} \frac{1}{n} \log \|Df^n\|_{E^c(x)}\|.$$

By Oseledets theorem, given any invariant probability measure the above limit exists for almost every point x.

Now we announce an Invariance principle due to Avila-Viana [2]. Let us concentrate on a class of partially hyperbolic dynamics which include the systems under consideration in this paper. Let $f: M \to M$ be a partially hyperbolic dynamics satisfying the following conditions:

- H1. f is dynamically coherent with all center leaves compact,
- H2. f admits global holonomies, that is, for any $y \in \mathcal{F}^u(x)$ the holonomy map $H^u_{x,y} : \mathcal{F}^c(x) \to \mathcal{F}^c(y)$ is a homeomorphism. For any $z \in \mathcal{F}^c(x)$, $H^u_{x,y}(z) = \mathcal{F}^u(z) \cap \mathcal{F}^c(y)$.
- H3. f_c is a transitive topological Anosov homeomorphism, where f_c is the induced dynamics satisfying $f_c \circ \pi = \pi \circ f$ and $\pi : M \to M/\mathcal{F}^c$ is the natural projection to the space of central leaves. In particular there are two foliations \mathcal{W}^s and \mathcal{W}^u which are stable and unstable sets for f_c .

Theorem 3.1 ([2]). (Invariance Principle) Let $f: M \to M$ be a partially hyperbolic diffeomorphism with one-dimensional compact center leaves satisfying H1, H2 and H₃. Let μ be an f-invariant probability measures whose projection $\nu = \pi_*\mu$ is probability measure that has local product structure. Assume that $\lambda_c(x) = 0$ for μ -almost every point. Then μ admits a disintegration $\{\mu_{[x]} : [x] \in M_c\}$ which is sinvariant and u-invariant and whose conditional probabilities $\mu_{[x]}$ vary continuously with [x] on the support of ν .

The following lemma is an immediate corollary of the uniqueness of Rokhlin disintegration theorem and above invariance principle result.

Lemma 3.2. Let $f : \mathbb{T}^3 \longrightarrow \mathbb{T}^3$ be C^2 partially hyperbolic diffeomorphism, dynamically coherent with compact central leaves and suppose that f has a maximal entropy ergodic measure μ with zero exponent. If $\{\mu_{[x]}\}_{[x]\in\mathbb{T}^3/\mathcal{F}^c}$ is a continuous disintegration of μ then, for all [x]:

$$f_*\mu_{[x]} = \mu_{f_c([x])}$$

Proof. Observe that by invariance of μ and uniqueness of disintegration for almost every x we have $f_*\mu_x = \mu_{f(x)}$. As $[x] \to \mu_{[x]}$ is continuous on $\operatorname{supp}(\pi_*\mu) = \mathbb{T}^3/\mathcal{F}^c$, we get the desired result.

3.2. Support of maximal entropy measures. Let us remind some basic properties of support of measures of maximal entropy.

Ures-Viana-Yang proved two crucial facts which hold for all systems satisfying hypothesis of Theorem 1.3.

Although the main theorem in [25] is formulated for $M \neq \mathbb{T}^3$, the following proposition holds in the case of partially hyperbolic $f : \mathbb{T}^3 \to \mathbb{T}^3$ with one dimensional compact center leaves which is the setting we are interested here. The proof is exactly the same as in [25].

Proposition 3.3. ([25] Lemma 4.4 and Proposition 3.12) The supports of two ergodic m.m.e with negative (positive) exponent are disjoint in the absence of m.m.e with zero central exponent. Moreover the support of any m.m.e with non-positive (resp. non-negative) center exponent is saturated by \mathcal{F}^u (resp. \mathcal{F}^s) leaves.

We just observe that in the case of $M \neq \mathbb{T}^3$, as any partially hyperbolic diffeomorphism is accessible, by Theorem 1.2 if there exists any hyperbolic m.m.e, then there does not exist any non-hyperbolic m.m.e. As a consequence, for $M \neq \mathbb{T}^3$, the disjointness of supports in the above proposition does not need the hypothesis of absence of non-hyperbolic m.m.e in [25].

Remark 3.4. Observe that in the case of $M \neq \mathbb{T}^3$ every partially hyperbolic diffeomorphism is accessible and the support of m.m.e with zero center exponent is the whole M (if it exists). In our setting $M = \mathbb{T}^3$ by above proposition one conclude that the support of any m.m.e with vanishing center exponent is saturated by accessibility classes.

The assumption M being nil-manifold different from \mathbb{T}^3 in [25] is crucial to conclude that f is dynamically coherent and there exists unique compact invariant, u-saturated (respectively *s*-saturated) minimal subset (Proposition 1.9 and 6.4 in [12]). So by the above proposition the authors can prove that there exist at most one m.m.e with negative (resp. positive) center exponent.

3.3. Twin measure construction. In this subsection we recall a method to obtain new measures of maximal entropy beginning from hyperbolic m.m.e's which initially appeared in [20] and denoted by twin measure construction.

Let f be as in Theorem 1.2 (without assuming accessibility property). Suppose μ is an hyperbolic measure of maximal entropy. Then using an argument as in Ruelle-Wilkinson one concludes that conditional measures of μ along center leaves

are atomic. As μ is hyperbolic, the center Lyapunov exponent is non-zero and we suppose that it is negative. Then for typical (for measure μ) x the intersection of Pesin stable manifold with center leaf is an open curve $\mathcal{W}^{ws}(x)$ (weak stable manifold) inside $\mathcal{F}^c(x)$. So there exists Z with full μ measure where points in Zadmit Pesin stable manifold. This enables us to define an injective measurable map $\beta: Z \to \mathbb{T}^3$ by $\beta(x) := \partial^+ \mathcal{W}^{ws}(x)$ where $\partial^+ \mathcal{W}^{ws}(x)$ stands for the boudary point of $\mathcal{W}^{ws}(x)$ going from x to positive orientation. As we are assuming that fpreserves the orientation of \mathcal{F}^c we have that $\beta \circ f = f \circ \beta$ and β is an isomorphism between (f, μ) and $(f, \beta_* \mu)$. We call $\beta_* \mu$ as a twin measure of μ . Clearly there may be constructed another twin measure just substituing $\partial^+ \mathcal{W}^{ws}(x)$ by $\partial^- \mathcal{W}^{ws}(x)$.

It is clear that $\beta_*\mu$ and μ has the same entropy and so if μ is m.m.e, its twin is too.

4. Proof of results

4.1. **Proof of Theorem 2.1.** Let $\mathcal{F}^{c}(a)$ be a periodic leaf through a where $f^{l}\mathcal{F}^{c}(a)$ is a 1-Morse-Smale dynamic with attractor a and repeller r. Without loss of generality we assume that a is a fixed point and so $\mathcal{F}^{c}(a)$ is a fixed leaf. We consider two following disjoint cases:

- f admits an ergodic measure of maximal entropy with zero center exponent.
- f admits no ergodic measure of maximal entropy with vanishing center exponent.

In the first case we will show the existence of an su-torus. In the second case we prove that either there exists a unique m.m.e with negative center exponent (and a unique one with positive center exponent) or there exists an invariant su-torus.

Let us deal with first case: let m be maximal measure with zero exponent and $\{m_{[x]}\}_{[x]\in\mathbb{T}^3/\mathcal{F}^c}$ be the disintegration along center leaves. By Lemma 3.2 we have $m_{\pi(a)} = c_1\delta_a + c_2\delta_r$. Suppose that $c_1 \neq 0$. As the conditional measures $m_{[x]}$ are invariant under stable and unstable holonomies we have:

$$\{h_{\gamma}(a) \cap \mathcal{F}^{c}(a) : \gamma \text{ is } su - \text{path}\} \subseteq \text{supp} \, m_{\pi(a)} \subseteq \{a, r\}.$$

Indeed, As h_{γ} is a homeomorphism for any su-path γ and the quotient dynamic is transitive Anosov homeomorphism and accessible, we conclude that:

$$1 \le |\{h_{\gamma}(a) \cap \mathcal{F}^{c}(p) : \gamma \quad su - \text{path}\}| \le 2,$$

for any $p \in \mathbb{T}^3$ and any su-path γ . In other words,

$$Card(Acc(a) \cap \mathcal{F}^{c}(p)) = Card(Acc(a) \cap \mathcal{F}^{c}(a)) \leq 2$$

for any $p \in \mathbb{T}^3$. Using this we can show that Acc(a) is compact. Then

$$\pi|_{Acc(a)} : Acc(a) \to \mathbb{T}^2$$

is a covering map with one or two sheets. As Acc(a) is compact and connected, it is a torus.

Now we deal with the second case: First of all, we know that f at least admits an m.m.e with negative center exponent. Suppose that there are two ergodic measures of maximal entropy μ_1, μ_2 with negative center exponent (the argument for two measures with positive exponent is similar). Observe that for any maximal measure η , $\operatorname{supp}(\eta) \cap \mathcal{F}^c(x) \neq \emptyset$ for all $x \in \mathbb{T}^3$. Indeed, as $\pi_*\eta$ is the measure of maximal entropy for a topologically transitive Anosov homeomorphism, it is fully supported on $\mathbb{T}^3/\mathcal{F}^c$ and by definition $\pi(\operatorname{supp}(\eta)) = \operatorname{supp}(\pi_*(\eta))$ and this implies $\operatorname{supp}(\eta) \cap \mathcal{F}^c(x) \neq \emptyset$.

Lemma 4.1. There exists $i \in \{1, 2\}$ such that $supp(\mu_i)$ intersects each center leaf in exactly one point.

Proof. There is a neighbourhood $U \subset \mathbb{T}^2$ around $\pi(r)$ such that for any $x \in U$ by local product structure $\mathcal{W}_{loc}^u(x) \cap \mathcal{W}_{loc}^s(\pi(r)) \neq \emptyset$. We prove the lemma by contradiction. As $\pi_*(\mu_1) = \pi_*(\mu_2)$ we may suppose that there exists $[p] \in U$ with $Card(\pi^{-1}([p]) \cap \operatorname{supp}(\eta) \geq 2$ for $\eta \in \{\mu_1, \mu_2\}$. Let $p_1, q_1 \in \pi^{-1}([p]) \cap \operatorname{supp}(\mu_1)$ and $p_2, q_2 \in \pi^{-1}([p]) \cap \operatorname{supp}(\mu_2)$.

Consider the local unstable holonomy:

$$h^{u}: \pi^{-1}([p]) \to \pi^{-1}(\mathcal{W}^{u}_{loc}([p]) \cap \mathcal{W}^{s}_{loc}(\pi(r))).$$

By *u*-saturation of support of m.m.e with negative center exponent (Proposition 3.3), we have $\hat{p}_i := h^u(p_i), \hat{q}_i := h^u(q_i) \in \text{supp}(\mu_i)$.

Now consider local stable holonomy:

$$h^s: \pi^{-1}(\mathcal{W}^u_{loc}([p]) \cap \mathcal{W}^s_{loc}(\pi(r))) \to \mathcal{F}^c(r).$$

As $\mathcal{F}^{c}(r)$ is one dimensional with 1–Morse-Smale dynamics for each $i \in \{1, 2\}$ we have

- $d(f^n(h^s(\hat{p_i}), a) \to 0 \text{ or}$
- $d(f^n(h^s(\hat{q}_i), a) \to 0.$

As $\hat{p}_i, \hat{q}_i \in \text{supp } \mu_i$ and support of a measure is closed invariant subset then $a \in \text{supp}(\mu_i)$ for both i = 1, 2 and this gives a contradiction as $\text{supp}(\mu_1) \cap \text{supp}(\mu_2) = \emptyset$ again using Proposition 3.3 (Observe that we are in the case where no m.m.e has zero center Lyapunov exponent).

By the above lemma we conclude that there exists $i \in \{1, 2\}$ such that

$$Card(\operatorname{supp}(\mu_i) \cap \{a, r\}) = 1.$$

and for simplicity we denote it by μ and assume that $r = \operatorname{supp}(\mu) \cap \mathcal{F}^c(r)$.

Lemma 4.2. $\operatorname{supp}(\mu)$ contains both stable and unstable leaf of r.

Proof. The inclusion of $\mathcal{F}^u(x) \subset \operatorname{supp}(\mu)$ is by u-saturtion of support. For the inclusion of the stable leaf, observe that for any point $z \in \mathcal{F}^{cs}(r)$ we have $\operatorname{supp}(\mu) \cap \mathcal{F}^c(z)$ is a unique point. We claim that this unique point should be $h^s(r) = \mathcal{F}^s(r) \cap \mathcal{F}^c(z)$. Indeed, if not $y \in \operatorname{supp}(\mu) \cap \mathcal{F}^c(z)$ with $\lim_{k \to \infty} d(f^k(y), a) = 0$. This implies that $a \in \operatorname{supp}(\mu)$ which is a contradiction.

To finish the proof of the theorem let $S:\mathbb{T}^3/\mathcal{F}^c\to\mathbb{T}^3$ be the section which associates

$$[p] \to S([p]) := \operatorname{supp}(\mu) \cap \pi^{-1}([p])$$

for any $[p] \in \mathbb{T}^3 / \mathcal{F}^c$.

As $\operatorname{supp}(\mu)$ is u-saturated we have that $S(\mathcal{W}^u([p])) = \mathcal{F}^u(S([p]))$. We have also proved that $\mathcal{F}^s(r) \subset \operatorname{supp}(\mu)$ which yields

(1)
$$S(\mathcal{W}^s(\pi(r))) = \mathcal{F}^s(r).$$

Finally we prove that $\sup(\mu)$ is an su-saturated set. Let $q \in \sup(\mu)$. We need to prove that $\mathcal{F}^s(q) \in \sup(\mu)$. We claim that there is a sequence p_n with $\mathcal{F}^s(p_n) \in \operatorname{supp}(\mu)$. As support is a closed set and stable foliation is continuous we conclude that $\mathcal{F}^s(q) \in \operatorname{supp}(\mu)$. Indeed, note that $\mathcal{W}^s(\pi(r)) \cap \mathcal{W}^u(\pi(q))$ accumulates on $\pi(q)$. Let $[z_n] \in \mathcal{W}^s(\pi(r)) \cap \mathcal{W}^u(\pi(q))$ converging to $\pi(q)$. Observe that as μ has negative center exponent

(2)
$$S(\mathcal{W}^u(\pi(q))) = \mathcal{F}^u(q).$$

Using (1), (2) we have:

(3)
$$p_n := S([z_n]) = \mathcal{F}^s(r) \cap \mathcal{F}^u(q) \cap \pi^{-1}([z_n]).$$

This concludes the proof of the claim and finishes the proof of Theorem 2.1 in the case where f admits 1–Morse-Smale dynamics on a periodic center leaf.

The proof of Addendum 2 is similar to the above theorem. In fact if there are $k + \lfloor \frac{k}{2} \rfloor + 1$ m.m.e with negative center exponent using similar argument as in Lemma 4.1 we get at least one μ_i such that $\operatorname{supp}(\mu_i)$ intersects each center leaf at just one point and repeating the above arguments we get an su-torus.

4.2. Proof of theorem 2.7.

Proof. We consider following disjoint cases:

- f admits an ergodic measure of maximal entropy with zero center exponent.
- *f* admits no ergodic measure of maximal entropy with vanishing center exponent.

First, Let us suppose that f has an ergodic maximal entropy measure m with zero center Lyapunov exponent. Then, by the invariance principle Theorem 3.1, there exists continuous disintegration $\{m_x^c\}$ along the central foliation which is invariant be stable and unstable holonomies.

The disintegration $\{m_x^c\}_x$ is invariant and $f^{\tau}|_{\mathcal{F}^c(a)} \circ h = h \circ R_{\alpha}$ for $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Since R_{α} is uniquely ergodic, then $h_*Leb_{\mathbb{S}}$ is the unique maximal entropy measure for $f^{\tau}|_{\mathcal{F}^c(a)}$. Moreover, this measure has no atoms and satisfies $\operatorname{supp}(m_a) =$ $\operatorname{supp} h_*Leb_{\mathbb{S}} = \mathcal{F}^c(a)$. Using accessibility of the quotient map $f_c : \mathbb{T}^3/\mathcal{F}^c \longrightarrow \mathbb{T}^3/\mathcal{F}^c$, we conclude that m_x^c has no atoms and is fully supported on center leaf for every $x \in \mathbb{T}^3$.

Then, we define an action of S on M that commutes with f. Take an orientation for \mathcal{F}^c . Now, identifying S with [0,1]/mod1, define $\Gamma : S \times M \longrightarrow M$ in such a way that $m_x([\Gamma(\theta, x), x]) = \theta$. This action is well defined and continuous because $\supp(m_x) = \mathcal{F}^c(x)$, the conditional measures has no atoms and the disintegration is continuous. Thus, by the invariance of the disintegration, we conclude that $f(\Gamma(\theta, x)) = \Gamma(\theta, f(x))$.

Similar to what have been done in [20] we conclude that f is conjugate to \bar{f} where \bar{f} is a rigid rotation extension of Anosov homeomorphism f_c . As f has a periodic leaf (with period τ with irrational rotation number dynamics, then there exists a periodic leaf $\mathcal{F}^c(a)$ where \bar{f}^{τ} is an irrational rotation.

For the uniqueness, it is enough to show that \overline{f} has a unique measure of maximal entropy m such that the conditional measures m_x along center leaves are Lebesgue and the quotient measure is the measure of maximal entropy for f_c . Let us suppose that η is a maximal entropy ergodic measure. Then, $\lambda^c(\eta) = 0$ because \overline{f} acts isometrically on center leaves. If $\{\eta_x\}_x$ is a disintegration of η , then η_a is the unique that satisfies $\eta_a = f_*^{\tau} \eta_a$ and by consequence, η_a is Lebesgue. Since the quotient map f_c is accessible and the disintegration of η is invariant by holonomies, we get $\eta_x = m_x$ for every $x \in \mathbb{T}^3$. This concludes the first part.

Now, let us suppose that there is no maximal entropy ergodic measure with zero central expoent and let μ be an m.m.e with $\lambda_c(\mu) < 0$. We will show that μ is the unique m.m.e with negative central exponent. If $\eta \neq \mu$, is another m.m.e with $\lambda_c(\eta) < 0$, we must have $\operatorname{supp}(\eta) \cap \operatorname{supp}(\mu) = \emptyset$. To get uniqueness is enough to show that $\operatorname{supp}(\mu) = \mathbb{T}^3$.

Let be $\mathcal{F}^c(p)$ the periodic leaf with irrational dynamics. As observed before, we have $\operatorname{supp}(\mu) \cap \mathcal{F}^c(x) \neq \emptyset$ for all $x \in \mathbb{T}^3$, in particular, $\operatorname{supp}(\mu) \cap \mathcal{F}^c(p) \neq \emptyset$. By consequence, $\mathcal{F}^c(p) \subset \operatorname{supp}(\mu)$ since $\mathcal{F}^c(p)$ has irrational dynamics and $\operatorname{supp}(\mu)$ is invariant. Once the $\operatorname{supp}(\mu)$ is saturated by unstable leaves, we have $\mathcal{F}^{cu}(p) \subset$ $\operatorname{supp}(\mu)$. Besides, $\overline{\mathcal{F}^{cu}(p)} = \mathbb{T}^3$ which gives $\operatorname{supp}(\mu) = \mathbb{T}^3$ and concludes the proof.

A twin argument shows that there exists also a unique measure of maximal entropy with positive exponent. 4.3. **Proof of Theorem 2.6, Kan type examples.** The Kan example [13] is a map of the cylinder $\mathbb{A} = \mathbb{S}^1 \times [0, 1] \to \mathbb{A}$, defined as:

$$f(x,t) = (3x \operatorname{mod-}\mathbb{Z}, t + \frac{t(1-t)}{32}\cos(2\pi x)), \quad \forall (x,t) \in \mathbb{S}^1 \times [0,1].$$

This map is well known because of its intricate property of admitting two physical measures with intermingled basins. Recall that for the basin $B(\mu)$ of an invariant measure μ is the set of points x such that

$$\lim_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n-1} \delta_{f^j(x)} = \mu.$$

An invariant measure is physical if $Leb(B(\mu)) > 0$. Two physical measures are called intermingled if for every open set U we have $Leb(B(\mu_1) \cap U) > 0$ and $Leb(B(\mu_2) \cap U) > 0$.

In the above Kan's example, the (one dimensional) Lebesgue measures μ_1, μ_2 on the boundary circles are invariant measures and Kan proved that μ_1 and μ_2 are physical measures with $Leb(B(\mu_1) \cup B(\mu_2)) = 1$ and more strikely, the basins are intermingled.

Similarly one may define an skew product on $\mathbb{T}^2 \times [0,1]$ as follows:

$$F(x,t) := (A(x), f_x(t)),$$

where $A: \mathbb{T}^2 \to \mathbb{T}^2$ is a linear Anosov diffeomorphism with eigenvalues $\mu < 1 < \lambda$. For each $x \in \mathbb{T}^2$ the function $f_x: [0,1] \to [0,1]$ is a diffeomorphism fixing the boundary of [0,1]. Take two fixed points of A called p,q. We require that f_p and f_q have exactly two fixed points each, a source at t = 1 (respectively t = 0) and a sink at t = 0 (respectively t = 1). Furthermore, $\mu < |f'_x(t)| < \lambda$ and

$$\int \log f_x'(0) dx < 0 \quad \text{and} \quad \int \log f_x'(1) dx < 0.$$

Under these conditions F has two interminged SRB measures which are normalized Lebesgue measure of each boundary torus. Under some more conditions F is also transitive (see [3].) Observe that as $f_x, x \in \mathbb{T}^2$ are orientation preserving if $(x, t) \in$ $B(\mu_1)$ then for all $0 \leq s \leq t$ we have $(x, s) \in B(\mu_1)$. Similarly if $(x, t) \in B(\mu_2)$ then $(x, s) \in B(\mu_2)$ for all $1 \geq s \geq t$. As $B(\mu_1) \cup B(\mu_2)$ has full Lebesgue measure we conclude that for Lebesgue almost every $x \in \mathbb{T}^2$ there exists $\sigma(x) \in (0, 1)$ where

(4)
$$(x,s) \in B(\mu_1)$$
 if $s < \sigma(x)$ and $(x,s) \in B(\mu_2)$ if $s > \sigma(x)$.

To get a diffeomorphism on a boundaryless manifold, we consider two such examples and glue them to find a partially hyperbolic diffeomorphism of \mathbb{T}^3 admitting two SRB measures (See also [7] and [24]).

Take $\tilde{G}: \mathbb{T}^2 \times [0,1] \to \mathbb{T}^2 \times [0,1]$ as follows:

(5)
$$\tilde{G}(x,t) = \begin{cases} (A(x), 1 - \frac{1}{2}f_x(2t)) & 0 \le t \le \frac{1}{2} \\ (A(x), \frac{1}{2}f_x(2(1-t))) & \frac{1}{2} \le t \le 1. \end{cases}$$

Clearly the above map defines a diffeomorphism on $G:\mathbb{T}^3\to\mathbb{T}^3$ just gluing the boundaries.

Although G is topologically transitive, as G^2 is not transitive we conclude that G is not topologically mixing. So it is still a question whether one can construct topologically mixing partially hyperbolic diffeomorphism with intermingled basin physical measures. However, see the result of Gan and Shi [11] where they prove C^2 robust topological mixing of Kan example on the annulus. We observe also that Bonatti-Potrie [4] had constructed examples of mixing diffeomorphisms with

intermingles basins physical measures, however their examples are not strongly partially hyperbolic. In fact in their examples $T\mathbb{T}^3 = E^{cs} \oplus E^u$ is the invariant tangent bundle splitting.

Observe that the two torus $\{t = 0\}, \{t = 1/2\}$ are invariant and support the SRB measures with intermingled basins on the 2-torus. We mention that Ures and Vasquez [24] proved the existence of su-torus for any partially hyperbolic $C^r, r > 1$ diffeomorphism in \mathbb{T}^3 with intermingled physical measures. In particular such diffeomorphisms are not accessible and so the set of these diffeomorphisms has empty interior in C^r topology.

The Lebesgue measures supported on these tori μ_1, μ_2 besides being physical, are also ergodic measures of maximal entropy for F. (and G) Both μ_1 and μ_2 have negative center Lyapunov exponent. Observe that μ_1 and μ_2 as measures defined on \mathbb{T}^3 have atomic disintegration along the center foliation.

Besides these two measures F (also G) admits another (unique) measure of maximal entropy ν with positive center exponent. In fact ν is twin measure of both μ_1 and μ_2 .

Indeed, $\nu_1 := \int_{\mathbb{T}^2} \delta_{\sigma(x)} dLeb(x)$ is an ergodic *F*-invariant measure such that $\pi_*\nu_1 = Leb$ which implies that ν_1 is a maximal entropy measure. Observe that $\sigma(x)$ is the boundary point of the weak (Pesin)stable set of (x, 0) which coincides with the boundary of weak (Pesin) stable set of (x, 1) for Lebesgue almost every x. So, the center Lyapunov exponent of ν_1 is non-negative.

We claim that $\lambda^c(\nu) \neq 0$. If it is not the case, then by Invariance principle the disintegration of ν is defined continuously and this implies that $x \to \sigma(x)$ can be extended continuously to whole \mathbb{T}^2 and this contradicts the fact that the basins of μ_1 and μ_2 are intermingled.

By construction of G we conclude that $\nu = \frac{1}{2}(\nu_1 + G_*\nu_1)$ is ergodic and maximal entropy for G.

Finally to construct an example with four m.m.e we just consider $H = G^2$ where μ_1, μ_2 (resp. ν_1, ν_2) are respectively two measures of negative (resp. positive) center Lyapunov exponent.

4.4. Examples for each item of theorem 2.7. For item 1, it is enough to consider a Anosov map $A : \mathbb{T}^2 \to \mathbb{T}^2$, a irrational rotation $R : \mathbb{S}^1 \to \mathbb{S}^1$ and then, take the product $A \times R : \mathbb{T}^3 \to \mathbb{T}^3$.

In order to give example for the second item of the dichotomy, note that if f has a periodic leaf with Morse Smale dynamics, then, it cannot be conjugate to rotation extension. Thus, it is enough to take a Anosov $A : \mathbb{T}^2 \to \mathbb{T}^2$ with two fixed points p and q and a family $f_x : \mathbb{S} \to \mathbb{S}$ of C^2 maps in a way that f_p is a irrational map and f_q is a Morse-Smale map.

In the theorem 2.7, we do not have the hypothesis of accessibility. One can ask whether there exists any example of accessible system that satisfies the hypothesis of 2.7 and still has two maximal entropy ergodic measures. Here we give a positive answer to this question.

The idea of the construction is to perturb a rotation extension in a neighborhood of a center leaf using a bump- function. If necessary, we rotate the system afterwards.

Consider an accessible rotation extension $F : \mathbb{T}^2 \times \mathbb{S}^1 \longrightarrow \mathbb{T}^2 \times \mathbb{S}^1$ (See [18] for such examples). The property of accessibility is open in C^1 topology (See [10]).

By definition $F = A \ltimes R_x$, where R_x is a rotation, i.e $F(x, \theta) = (A(x), R_x(\theta))$ We can assume that $R_0 = Id_{\mathbb{S}^1}$, if not, consider $G = (Id_{\mathbb{T}^2}, R_0^{-1}) \circ F$. Once the action of rotation preserves the unstable and stable foliation, the new system G is still accessible For $t \in \mathbb{R}$, we define $g_t : \mathbb{S}^1 \to \mathbb{S}^1$ by $g_t(\xi) := \xi + t \sin(4\pi\xi)$. For small t, g is a 2-Morse-Smale C^1 diffeomorphism close to the identity.

Consider R > 0 such that, there is a periodic point $p \in \mathbb{T}^2$ with orbit outside B(0, R). Now, consider a bump-function $\eta : \mathbb{T}^2 \longrightarrow [0, 1]$ with $\eta(0) = 1$ and $\eta(x) = 0$ if |x| > R.

Define $H = A \ltimes (g_{l(x)} \circ R_x)$, where $l(x) = t_0 \eta(x)$ for some fixed $t_0 > 0$.

As F is stably accessible in C^1 -topology and for t_0 small enough, then $d_{C^1}(H, F)$ is small H is accessible.

Let $\tau(p)$ be the period of p. If $F^{\tau(p)}|\mathcal{F}^c(p)$ is an irrational rotation, then H satisfies the properties that we are looking for. If not, perturb a little to obtain irrational rotation. An small perturbation does not destroy the Morse-Smale dynamics on $\mathcal{F}^c(0)$.

References

- [1] Flavio Abdenur and Marcelo Viana. Flavors of partial hyperbolicity. unpublished???, 2015.
- [2] Artur Avila and Marcelo Viana. Extremal Lyapunov exponents: an invariance principle and applications. *Invent. Math.*, 181(1):115–189, 2010.
- [3] Christian Bonatti, Lorenzo J. Díaz, and Marcelo Viana. Dynamics beyond uniform hyperbolicity, volume 102 of Encyclopaedia of Mathematical Sciences. Springer-Verlag, Berlin, 2005. A global geometric and probabilistic perspective, Mathematical Physics, III.
- [4] Christian Bonatti and Rafael Potrie. Many intermingled basins in dimension 3. Israel J. Math., 224(1):293–314, 2018.
- [5] J. Buzzi, T. Fisher, M. Sambarino, and C. Vásquez. Maximal entropy measures for certain partially hyperbolic, derived from Anosov systems. *Ergodic Theory Dynam. Systems*, 32(1):63–79, 2012.
- [6] Jérôme Buzzi. C^r surface diffeomorphisms with no maximal entropy measure. Ergodic Theory Dynam. Systems, 34(6):1770–1793, 2014.
- [7] Cheng Cheng, Shaobo Gan, and Yi Shi. A robustly transitive diffeomorphism of Kan's type. Discrete Contin. Dyn. Syst., 38(2):867–888, 2018.
- [8] Vaughn Climenhaga, Yakov Pesin, and Agnieszka Zelerowicz. Equilibrium measures for some partially hyperbolic systems. J. Mod. Dyn., 16:155–205, 2020.
- [9] Lorenzo J. Díaz, Todd Fisher, Maria José Pacifico, and José L. Vieitez. Entropy-expansiveness for partially hyperbolic diffeomorphisms. *Discrete Contin. Dyn. Syst.*, 32(12):4195–4207, 2012.
- [10] Ph. Didier. Stability of accessibility. Ergodic Theory Dynam. Systems, 23(6):1717–1731, 2003.
- [11] Shaobo Gan and Yi Shi. Robustly topological mixing of Kan's map. J. Differential Equations, 266(11):7173–7196, 2019.
- [12] A. Hammerlindl and R. Potrie. Pointwise partial hyperbolicity in three-dimensional nilmanifolds. J. Lond. Math. Soc. (2), 89(3):853–875, 2014.
- [13] Ittai Kan. Open sets of diffeomorphisms having two attractors, each with an everywhere dense basin. Bull. Amer. Math. Soc. (N.S.), 31(1):68–74, 1994.
- [14] François Ledrappier and Peter Walters. A relativised variational principle for continuous transformations. J. London Math. Soc. (2), 16(3):568–576, 1977.
- [15] Gang Liao, Marcelo Viana, and Jiagang Yang. The entropy conjecture for diffeomorphisms away from tangencies. J. Eur. Math. Soc. (JEMS), 15(6):2043–2060, 2013.
- [16] M. Misiurewicz. Diffeomorphism without any measure with maximal entropy. Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys., 21:903–910, 1973.
- [17] Sheldon Newhouse. On the continuity properties of entropy. Annals. Math. (2), 129(2):215– 235, 1989.
- [18] Viorel Niţică and Andrei Török. An open dense set of stably ergodic diffeomorphisms in a neighborhood of a non-ergodic one. *Topology*, 40(2):259–278, 2001.
- [19] Bárbara Núñez-Madariaga, Sebastián A. Ramírez, and Carlos H. Vásquez. Measures maximizing the entropy for kan endomorphisms. arXiv:2009.11956v1.
- [20] F. Rodriguez Hertz, M. A. Rodriguez Hertz, A. Tahzibi, and R. Ures. Maximizing measures for partially hyperbolic systems with compact center leaves. *Ergodic Theory Dynam. Systems*, 32(2):825–839, 2012.
- [21] F. Rodriguez Hertz, M. A. Rodriguez Hertz, and R. Ures. A non-dynamically coherent example on T³. Ann. Inst. H. Poincaré Anal. Non Linéaire, 33(4):1023–1032, 2016.
- [22] Federico Rodriguez Hertz, Maria Alejandra Rodriguez Hertz, and Raul Ures. Partial hyperbolicity and ergodicity in dimension three. J. Mod. Dyn., 2(2):187–208, 2008.

- [23] Raúl Ures. Intrinsic ergodicity of partially hyperbolic diffeomorphisms with a hyperbolic linear part. Proc. Amer. Math. Soc., 140(6):1973–1985, 2012.
- [24] Raúl Ures and Carlos H. Vásquez. On the non-robustness of intermingled basins. Ergodic Theory Dynam. Systems, 38(1):384–400, 2018.
- [25] Raul Ures, Marcelo Viana, and Jiagang Yang. Maximal entropy measures of diffeomorphisms of circle fiber bundles. *Journal of the London Mathematical Society*.

A. TAHZIBI, INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO (USP), *E-mail address:* tahzibi@icmc.usp.br

J. ROCHA, UNIDADE ACADÊMICA DE CABO DE SANTO AGOSTINHO, UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO, *E-mail address:* joas.rocha@ufrpe.br

14