COOLING DOWN STOCHASTIC DIFFERENTIAL EQUATIONS: ALMOST SURE CONVERGENCE

STEFFEN DEREICH AND SEBASTIAN KASSING

ABSTRACT. We consider almost sure convergence of the SDE $dX_t = \alpha_t dt + \beta_t dW_t$ under the existence of a C^2 -Lyapunov function $F: \mathbb{R}^d \to \mathbb{R}$. More explicitly, we show that on the event that the process stays local we have almost sure convergence in the Lyapunov function $(F(X_t))_{t\geq0}$ as well as $\nabla F(X_t) \to 0$, if $|\beta_t| = \mathcal{O}(t^{-\beta})$ for a $\beta > 1/2$. If, additionally, one assumes that F is a Lojasiewicz-function, we get almost sure convergence of the process itself, given that $|\beta_t| = \mathcal{O}(t^{-\beta})$ for a $\beta > 1$. The assumptions are shown to be optimal in the sense that there is a divergent counterexample where $|\beta_t|$ is of order t^{-1} .

1. INTRODUCTION

Let $(\Omega, (\mathcal{F}_t)_{t>0}, \mathcal{F}, \mathbb{P})$ be a filtered probability space satisfying the usual conditions and suppose that $(X_t)_{t\geq0}$ is a continuous semimartingale satisfying the equation

(1)
$$
X_t - X_0 = \int_0^t \alpha_s ds + \int_0^t \beta_s dW_s,
$$

for all $t \geq 0$, where

- $(\alpha_t)_{t\geq 0}$ is a progressive, \mathbb{R}^d -valued process,
- $(\beta_t)_{t\geq 0}$ is a progressive $\mathbb{R}^{d \times d'}$ -valued process and
- $(W_t)_{t\geq 0}$ is an $\mathbb{R}^{d'}$ -valued $(\mathcal{F}_t)_{t\geq 0}$ -Brownian motion with initial value $W_0 = 0$.

In this article, we analyse convergence of $(X_t)_{t\geq 0}$ under the assumption that the drift term satisfies a Lyapunov kind of condition and the diffusivity converges to zero sufficiently fast. More explicitly, we denote by $F: \mathbb{R}^d \to \mathbb{R}$ a C^2 -function, by $f := \nabla F$ its gradient and by $H := \text{Hess } F$ the Hessian of F and consider the following event

$$
\mathbb{C} := \left\{ \liminf_{t \to \infty} \frac{\langle f(X_t), -\alpha_t \rangle}{|f(X_t)|^2} > 0, \liminf_{t \to \infty} \frac{|f(X_t)|}{|\alpha_t|} > 0, \frac{\int_0^\infty |\beta_s|_F^2 ds < \infty \text{ and } \limsup_{t \to \infty} |\beta_t| < \infty \right\},\right\}
$$

where we interpret $\frac{0}{0}$ as ∞ and, for a matrix $A \in \mathbb{R}^{d \times d'}$, $|A|_F$ (resp. $|A|$) denotes the Frobenius norm (resp. spectral norm). Intuitively, on $\mathbb C$ the drift term $(\alpha_t)_{t>0}$ is comparable in size and direction to the negative gradient of the Lyapunov function, i.e., $(-\nabla F(X_t))_{t>0}$, and the size of the diffusivity $(\beta_t)_{t\geq 0}$ can be controlled at late times. The first and second condition in the definition of the event $\mathbb C$ are obviously satisfied if we consider the solution to the SDE

⁽²⁾ $dX_t = -\nabla F(X_t) dt + \beta_t dW_t,$

²⁰²⁰ Mathematics Subject Classification. Primary 60J60; Secondary 60H10, 65C35.

Key words and phrases. Stochastic gradient flow; Brownian particle; cooling down; Lojasiewicz-inequality; almost sure convergence.

which represents the canonical stochastic version of the gradient flow ODE

$$
\dot{x}_t = -\nabla F(x_t).
$$

However, the current framework allows more flexibility in the drift term. We will restrict attention to certain events: set

 λ

$$
\mathbb{B} := \left\{ \liminf_{t \to \infty} F(X_t) > -\infty, \limsup_{t \to \infty} |\alpha_t| < \infty \text{ and } \limsup_{\varepsilon \to 0} \limsup_{\substack{t \to \infty \\ y \in B(X_t, \varepsilon)}} |H(y)| < \infty \right\}
$$

and

$$
\mathbb{L} := \Big\{ \limsup_{t \to \infty} |X_t| < \infty \Big\}.
$$

We separately show almost sure convergence of $(F(X_t))_{t\geq 0}$, $(f(X_t))_{t\geq 0}$ and $(X_t)_{t\geq 0}$ under mild assumptions on the diffusivity and the underlying Lyapunov function which are shown to be optimal (c.f. Chapter [4\)](#page-15-0). The SDE [\(2\)](#page-0-0) is the continuous-time counterpart for the stochastic gradient descent scheme (when $(\beta_t)_{t>0}$ is chosen properly, see [\[LTE17\]](#page-20-0)) and, thus, a helpful tool for gaining intuition and insight on the behaviour of SGD. It also appears naturally in applications from Physics and Chemistry. For example, [\(2\)](#page-0-0) can be viewed as the pathwise solution to the Fokker–Planck equation (see [\[Gar85\]](#page-20-1)) in a physical system, where F models the energy and $(\beta_t)_{t\geq0}$ an external source of perturbation^{[1](#page-1-0)}, or as the chemical Langevin equation describing a chemical reaction evolving in time ([\[vK92\]](#page-20-2), [\[Gil00\]](#page-20-3), [\[SSG14\]](#page-20-4)). In our analysis, we allow general non-convex energy landscapes and focus on the case where the perturbation fades within time.

There are several asymptotic results for the solution $(X_t)_{t>0}$ of [\(2\)](#page-0-0) when choosing a deterministic diffusivity $(\beta_t)_{t>0}$. Inspired by simulated annealing, [\[GH86\]](#page-20-5) and [\[CHS87\]](#page-20-6) proved that for $(\beta_t)_{t\geq 0} = (\sqrt{2c/\log(t+1)})_{t\geq 0}$ with sufficiently large $c > 0$ and under additional assumptions on the target function F, the solution of [\(2\)](#page-0-0) converges weakly to a distribution π concentrated on the global minima of F independent of the initial value $X_0 = x \in \mathbb{R}^d$, even in the case of non-convex target functions F. More precisely, π is the weak limit of the Gibbs densities $\pi_t(x) \propto \exp(-2F(x)/\beta_t^2)$. In this regime, [\[HKS89\]](#page-20-7) showed that if the diffusion process is defined on a compact manifold, then $(F(X_t))_{t>0}$ converges to the optimal value in probability if and only if $c > c^*$, where c^* denotes the minimal potential energy necessary to connect every point with a global minimum using a continuous path. Recently, this result has been generalised by [\[FT21\]](#page-20-8) to diffusions on \mathbb{R}^d in the case where $F \in C^\infty$ is assumed to satisfy $\lim_{|x| \to \infty} F(x) = \infty$ and $\int_{\mathbb{R}^d} e^{-\alpha_0 F(x)} dx < \infty$ for some $\alpha_0 > 0$. In that case one has $\liminf_{t \to \infty} |X_t| < \infty$, almost surely.

However, in this particular scenario we cannot hope for the individual paths of the solution $(X_t)_{t\geq0}$ to converge. Conversely, the diffusivity $(\beta_t)_{t\geq0}$ considered here converges significantly faster to zero. Therefore, we cannot guarantee that the process $(X_t)_{t\geq0}$ started in the basin of attraction of a bad local minimum is able to overcome a barrier in the landscape in order to converge to a global minimum (see Section [3.3\)](#page-10-0) and the distribution of the limit points of $(X_t)_{t>0}$ depends heavily on the distribution of the initial value X_0 . As we will indicate below, the gradient flow converges if F is a Lojasiewicz-landscape, in the sense of Definition [1.4,](#page-2-0) and the gradient flow does not escape to infinity. The main contribution of this article is the analysis in how far this is still the case when additionally incorporating a stochastic perturbation. We stress that we analyse the asymptotic behaviour of the system as time tends to infinite which is significantly different from the analysis of families of SDEs where the diffusivity decays with a family index $\varepsilon > 0$, see e.g. [\[FW12\]](#page-20-9).

 $1(2/\beta_t^2)_{t\geq0}$ is sometimes referred to as the *inverse temperature*, e.g. [\[CHS87,](#page-20-6) [JKO98\]](#page-20-10)

First, we will provide a short proof for almost sure convergence of $(F(X_t))_{t\geq0}$ and $(f(X_t))_{t\geq0}$ with $\lim_{t\to\infty} f(X_t) = 0$ on the event $\mathbb{C} \cap \mathbb{B}$. We conclude that we have a.s. convergence of $(X_t)_{t\geq0}$ on the event $\mathbb{C} \cap \mathbb{L}$ in the case where where the set of critical points $\mathcal{C} := \{x \in \mathbb{R}^d : f(x) = 0\}$ is at most countable.

Theorem 1.1. On $\mathbb{C} \cap \mathbb{B}$, almost surely, $(F(X_t))_{t\geq 0}$ converges to a finite value and $\lim_{t\to\infty} f(X_t)$ 0 as well as $\limsup_{t_0 \to \infty} \int_{t_0}^{\infty} \langle f(X_s), -\alpha_s \rangle ds < \infty$. If the set of critical points of F is at most countable, then, on $\mathbb{C} \cap \mathbb{L}$, almost surely, $(X_t)_{t\geq 0}$ converges.

Theorem [1.1](#page-2-1) is an immediate consequence of the more general Proposition [2.1](#page-4-0) below.

Remark 1.2. Note that as F is C^2 we clearly have $\mathbb{C} \cap \mathbb{L} \subset \mathbb{C} \cap \mathbb{B}$ so that the first statement of Theorem [1.1](#page-2-1) also holds on $\mathbb{C} \cap \mathbb{L}$. Moreover, for a Lyapunov function F with $\liminf_{|x| \to \infty} |f(x)| >$ 0 the first statement implies that

$$
\mathbb{C} \cap \mathbb{B} = \mathbb{C} \cap \mathbb{L}.
$$

Remark 1.3. A standard approach for proving convergence statements similar to those given in Theorem [1.1](#page-2-1) is the theory of asymptotic pseudotrajectories (see [\[Ben99\]](#page-20-11), especially Corollary 6.7). While these methods may lead to different and typically weaker assumptions on the perturbation $(\beta_t)_{t>0}$, Theorem [1.1](#page-2-1) will mainly be used to prove Theorem [1.6](#page-3-0) below. In order to be self-contained we stated Theorem [1.1](#page-2-1) in such a way that it can easily be applied in Theorem [1.6](#page-3-0) and has a direct and brief proof.

In the case where the set of critical points of F contains a continuum of elements the situation is more subtle. Even in the case without random perturbations convergence of the solution $(x_t)_{t\geq 0}$ to the ODE

$$
\dot{x}_t = -\nabla F(x_t)
$$

is a non-trivial issue: one can find C^{∞} -functions F together with solutions $(x_t)_{t>0}$ that stay on compact sets but do not converge, see example 3 on page 14 of [\[PdM82\]](#page-20-12). Counterexamples of this structure have been known for a long time (see e.g. [\[Cur44\]](#page-20-13)) and include the famous Mexican hat function [\[AMA05\]](#page-20-14). To guarantee convergence (at least in the case where the solution stays on a compact set), one needs to impose additional assumptions. An appropriate assumption is the validity of a Lojasiewicz-inequality, see Definition [1.4](#page-2-0) below. This assumption has the appeal that it is satisfied by analytic functions, see [Loj63, [Loj65\]](#page-20-16). With the help of the inequality (4) Lojasiewicz showed in Loj84 that any integral curve of the gradient flow equation [\(3\)](#page-2-3) for an analytic function F that has an accumulation point has a unique limit. A proof of that assertion can also be found in [\[Har12\]](#page-20-18) or [\[CMI14\]](#page-20-19).

In this article, we device a probabilistic counterpart to the classical analytic approach. We will derive moment estimates of $F(X_t)$ for Lojasiewicz-functions $F: \mathbb{R}^d \to \mathbb{R}$ on $\mathbb{C} \cap \mathbb{L}$ under the additional assumption that we can asymptotically bound the diffusivity $(\beta_t)_{t\geq 0}$ by a locally-bounded deterministic function $\sigma : [0, \infty) \to [0, \infty)$ to deduce almost sure convergence of $(\int_0^t \alpha_s ds)_{t\geq 0}$ and thus, almost sure convergence of $(X_t)_{t\geq 0}$.

Definition 1.4. We call the function $F : \mathbb{R}^d \to \mathbb{R}$ Lojasiewicz-function, if for every $x \in \mathcal{C} :=$ $f^{-1}(\{0\})$, the Lojasiewicz-inequality is true on a neighbourhood U_x of x with parameters $L > 0$ and $\theta \in [\frac{1}{2}]$ $(\frac{1}{2}, 1)$, *i.e.*, for all $y \in U_x$

(4)
$$
|f(y)| \ge L |F(y) - F(x)|^{\theta}.
$$

Remark 1.5. Note that for all $x \notin \mathcal{C}$ and $\theta \in \left[\frac{1}{2}\right]$ $(\frac{1}{2}, 1)$ there trivially exist a neighbourhood U_x and constant $L > 0$ such that the Lojasiewicz-inequality with parameter L and θ hold on U_x .

The main contribution of this article concerns convergence of the SDE in the case where the Lyapunov function is a Lojasiewicz-function.

Theorem 1.6. Let F be a Lojasiewicz-function and $(\sigma_t)_{t\geq 0} = ((t+1)^{-\sigma})_{t\geq 0}$ with $\sigma > 1$. Then, on $\mathbb{C} \cap \mathbb{L} \cap \{ \limsup_{t \to \infty} \frac{|\beta_t|}{\sigma_t} \}$ $\frac{d\mathcal{F}_t}{dt} < \infty$, the process $(X_t)_{t\geq 0}$ satisfying [\(1\)](#page-0-1) converges, almost surely, to a critical point of F.

The proof of the theorem is accomplished in Section [3.](#page-6-0) The theorem is optimal in the sense that its conclusion is not true for the choice $(\sigma_t)_{t\geq 0} = ((t+1)^{-1})_{t\geq 0}$. More explicitly, we provide an SDE for which $\mathbb{C} \cap \mathbb{L} \cap \{\limsup_{t \to \infty} (t+1)|\overline{\beta}_t| < \infty\}$ is an almost sure set and for which $(X_t)_{t>0}$ diverges, almost surely, in Section [4.](#page-15-0)

Remark 1.7. Choose $(\alpha_t)_{t>0} = (-f(X_t))_{t>0}$ and consider the solution to the SDE [\(2\)](#page-0-0). In that case we have for $(\sigma_t)_{t\geq 0}$ as in Theorem [1.6](#page-3-0) that $\{\limsup_{t\to\infty} \frac{|\beta_t|}{\sigma_t}$ $\frac{\beta_t|}{\sigma_t} < \infty$ $\} \subset \mathbb{C}$. Now, assume that $F \in C^2$ satisfies $\lim_{|x| \to \infty} F(x) = \infty$. In [A](#page-19-0)ppendix A we derive sufficient conditions that guarantee locality of the process $(X_t)_{t>0}$. In particular, we show that if the Hessian of F is uniformly bounded (i.e. $\sup_{x \in \mathbb{R}^d} |H(x)| < \infty$) then $\{\int_0^\infty |\beta_s|^2_F ds < \infty\} \subset \mathbb{L}$. Also, we get for any compact set $K \subset \mathbb{R}^d$

$$
\left\{ \mathbb{1}_{\{X_t \notin K\}}(-|f(X_t)|^2 + \frac{1}{2} \text{tr}((\beta_t)^{\dagger} H(X_t)\beta_t)) \le 0 \text{ for all large } t \right\}
$$

and
$$
\int_0^{\infty} \mathbb{1}_{\{X_s \in K\}} |\beta_s|_F^2 ds < \infty \right\} \subset \mathbb{L}.
$$

The proof of Theorem [1.6](#page-3-0) is based on moment estimates for $F(X_t)$ restricted to particular events, see Proposition [3.1](#page-7-0) below. In the particular case where we approach a local minimum these estimates entail the following moment estimate.

Theorem 1.8. Let $K \subset \mathbb{R}^d$ be a compact set, $L > 0$, $\theta \in (1/2, 1)$, $\ell \in \mathbb{R}$, $C_\beta > 0$ and $\sigma \in [0, \infty)$. Suppose that for all $y \in K$

$$
F(y) \ge \ell \ \text{and} \ \ |f(y)| \ge L(F(y) - \ell)^{\theta}.
$$

Consider the stopping time

$$
T := \inf \Big\{ t \geq 0 : X_t \notin K, \frac{\langle f(X_t), -\alpha_t \rangle}{|f(X_t)|^2} \leq \rho , \ |\beta_t| \geq C_\beta (t+1)^{-\sigma} \Big\}.
$$

Then there exists $C \geq 0$ so that for all $t \geq 0$

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell)] \le C(t+1)^{-(\frac{\sigma}{\theta}\wedge \frac{1}{2\theta-1})}.
$$

Theorem [1.8](#page-3-1) will be a consequence of Proposition [3.1](#page-7-0) and Remark [3.2](#page-9-0) below. Note that if $\sigma \geq \theta/(2\theta-1)$ we essentially get the same order of convergence as in the case without random perturbations. If the drift terms vanishes at a slower rate, i.e. $\sigma < \theta/(2\theta - 1)$, this effects the order of decay of the expected function value $(F(X_t) - \ell)_{t\geq 0}$.

Notation. For a vector $v \in \mathbb{R}^d$ we denote by |v| the Euclidean norm induced by the scalar product, i.e. $|v|^2 = \langle v, v \rangle$. For a matrix $A \in \mathbb{R}^{d \times d'}$ we denote by |A| the spectral norm and by $|A|_F$ the Frobenius norm, *i.e.*

$$
|A| := \max_{|x| \neq 0} \frac{|Ax|}{|x|} \quad \text{and} \quad |A|_F := \sqrt{\text{tr}(A^\dagger A)}.
$$

For $t_0 \geq 0$ and a real-valued continuous semimartingale $(Y_s)_{s \geq t_0}$ we denote by $\langle Y \rangle = (\langle Y \rangle_s)_{s \geq t_0}$ its quadratic variation process.

2. Proof of Theorem [1.1](#page-2-1)

In this section we prove a slight generalisation of Theorem [1.1.](#page-2-1)

Proposition 2.1. Consider

$$
\mathbb{C}_1 := \Big\{ \liminf_{t \to \infty} F(X_t) > -\infty, \limsup_{t \to \infty} |\beta_t| < \infty, \liminf_{t \to \infty} \frac{\langle f(X_t), -\alpha_t \rangle}{|f(X_t)|^2} > 0 \text{ and}
$$

$$
\Big(\int_0^t \text{tr}(\beta_s^\dagger H(X_s)\beta_s) \, ds \Big)_{t \ge 0} \text{ converges to a finite value} \Big\}
$$

$$
\mathbb{C}_2 := \mathbb{C}_1 \cap \Big\{ \limsup_{t \to \infty} |\alpha_t| < \infty, \limsup_{\varepsilon \to 0} \limsup_{\substack{t \to \infty \\ y \in B(X_t, \varepsilon)}} |H(y)| < \infty, \int_0^\infty |\beta_s|_F^2 < \infty \Big\},\
$$

and

$$
\mathbb{C}_3 := \mathbb{C}_2 \cap \left\{ \liminf_{t \to \infty} |X_t| < \infty \right\},\
$$

where we interpret $\frac{0}{0}$ as ∞ . On \mathbb{C}_1 , almost surely, $(F(X_t))_{t\geq 0}$ converges to a finite value and $\limsup_{t_0 \to \infty} \int_{t_0}^{\infty} \langle f(X_s), -\alpha_s \rangle ds < \infty$. On \mathbb{C}_2 , almost surely, $\lim_{t \to \infty} f(X_t) = 0$. If the set of critical points of F is at most countable, then, on \mathbb{C}_3 , almost surely, $(X_t)_{t\geq0}$ converges.^{[2](#page-4-1)}

Note that $\mathbb{C} \cap \mathbb{B} \subset \mathbb{C}_2$ since $\int_0^\infty |\beta_s|_F^2 ds < \infty$ and $\limsup_{t \to \infty} |H(X_t)| < \infty$ imply convergence of

$$
\left(\int_0^t \text{tr}(\beta_s^{\dagger} H(X_s)\beta_s) \, ds\right)_{t\geq 0}
$$

.

Therefore, Theorem [1.1](#page-2-1) directly follows from Proposition [2.1.](#page-4-0)

Proof of Proposition [2.1.](#page-4-0) Let $t_0, C_1, C_2 \geq 0$ and set

$$
T := T^{(t_0, C_1, C_2)} := \inf \Big\{ t \ge t_0 : F(X_t) < -C_1 \vee \int_{t_0}^t \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) \, ds > C_1 \vee |\beta_t| > C_1,
$$
\n
$$
\text{or } \langle f(X_t), \alpha_t \rangle > -C_2 |f(X_t)|^2 \Big\}
$$

By Itô's formula one has

(5)
$$
dF(X_t) = \langle f(X_t), \alpha_t \rangle dt + \langle f(X_t) \rangle, \beta dW_t \rangle + \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) dt.
$$

We let

$$
(M_t)_{t \ge t_0} := \left(\int_{t_0}^{t \wedge T} \langle f(X_t), \beta \, dW_t \rangle \right)_{t \ge t_0}
$$

and note that its variation process satisfies

$$
\langle M \rangle_t = \int_{t_0}^{t \wedge T} |\beta_s^{\dagger} f(X_s)|^2 ds \le C_1 \int_{t_0}^{t \wedge T} |f(X_s)|^2 ds.
$$

On the other hand,

$$
\int_{t_0}^{t \wedge T} \langle f(X_s), \alpha_s \rangle ds \leq -C_2 \int_{t_0}^{t \wedge T} |f(X_s)|^2 ds.
$$

Consequently,

$$
\int_{t_0}^{t \wedge T} \langle f(X_s), \alpha_s \rangle \, ds + M_t \le M_t - \frac{C_2}{C_1} \langle M \rangle_t
$$

²Note that \mathbb{C}_3 entails \mathbb{C}_1 and \mathbb{C}_2 .

We conclude that for $t \geq t_0$

$$
\underbrace{F(X_{t\wedge T})}_{\geq -C_1} - F(X_{t_0}) = \underbrace{\int_{t_0}^{t\wedge T} \langle f(X_s), \alpha_s \rangle ds + M_t}_{\leq M_t - \frac{C_2}{C_1} \langle M \rangle_t} + \frac{1}{2} \underbrace{\int_{t_0}^{t\wedge T} \text{tr}(\beta_s^\dagger H(X_s)\beta_s) ds}_{\leq C_1}.
$$

Note that on $\{ \langle M \rangle_{\infty} = \infty \}$, almost surely, $M_t - \frac{C_2}{C_1}$ $\frac{C_2}{C_1} \langle M \rangle_t \to -\infty$. Since the left-hand side is bounded from below we get that $\langle M \rangle_{\infty} < \infty$, a.s., so that the martingale $(M_t)_{t \geq t_0}$ converges almost surely to a finite value. Consequently,

$$
-\int_{t_0}^{t \wedge T} \langle f(X_s), \alpha_s \rangle ds \le F(X_{t_0}) + M_t + \frac{3}{2}C_1
$$

and the left-hand side is increasing in t with a finite limit. Consequently, on

$$
\Big\{ \Big(\int_0^t \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) \, ds \Big)_{t \geq 0} \text{ converges} \Big\},
$$

we have that $(F(X_{t\wedge T}))_{t\geq 0}$ converges and $-\int_{t_0}^T \langle f(X_s), \alpha_s \rangle ds < \infty$, almost surely. We note that

$$
\mathbb{C}_1 = \left\{ \left(\int_0^t \text{tr}(\beta_s^\dagger H(X_s)\beta_s) \, ds \right)_{t \ge 0} \text{ converges} \right\} \cap \bigcup_{n \in \mathbb{N}} \{ T^{(n,n,1/n)} = \infty \}
$$

which implies the first statement.

To show the second statement we set

$$
\tilde{T} := \tilde{T}^{(t_0, C_1, C_2)} := T^{(t_0, C_1, C_2)} \wedge \inf \left\{ t \ge t_0 : |\alpha_t| \vee \sup_{y \in B(X_t, C_2)} |H(y)| \vee \int_{t_0}^t |\beta_s|_F^2 ds > C_1 \right\}
$$

and note that it satisfies to show that $f(X_t) \to 0$, almost surely, on $\{\tilde{T}^{(t_0,C_1,C_2)} = \infty\}$, since

$$
\mathbb{C}_2=\bigcup_{n\in\mathbb{N}}\{\tilde{T}^{(n,n,1/n)}=\infty\}.
$$

Consider

$$
(\tilde{M}_t)_{t \ge t_0} := \left(\int_{t_0}^{t \wedge \tilde{T}} \beta_s dW_s \right)_{t \ge t_0} \text{ and } (B_t)_{t \ge t_0} := \left(\int_{t_0}^{t \wedge \tilde{T}} \langle f(X_s), \alpha_s \rangle ds \right)_{t \ge t_0}
$$

Note that $(\tilde{M}_t)_{t \geq t_0}$ is a multivariate martingale with

$$
\langle \tilde{M} \rangle_t = \int_0^{t \wedge \tilde{T}} |\beta_s|_F^2 ds \le C_1.
$$

Hence, $(\tilde{M}_t)_{t\geq t_0}$ converges almost surely and together with the proof of the first statement we get that

$$
\Omega_0 := \{ (\tilde{M}_t)_{t \ge t_0} \text{ and } (B_t)_{t \ge t_0} \text{ converge} \}
$$

is an almost sure event. Suppose now that there exists $\omega \in \Omega_0 \cap {\tilde{T}} = \infty$ for which $(f(X_t(\omega)))_{t>0}$ does not converge to zero. Then there exist $\delta > 0$ and an increasing $[t_0, \infty)$ -valued sequence $(t_k)_{k\in\mathbb{N}}$ converging to infinity with

$$
|f(X_{t_k}(\omega))| > \delta
$$

for all $k \in \mathbb{N}$. Pick $\varepsilon \in (0, C_2)$ with $C_1 \varepsilon \leq \delta/2$. By thinning the original sequence $(t_k)_{k \in \mathbb{N}}$ we can ensure that $([t_k, t_k + \varepsilon/(2C_1)] : k \in \mathbb{N})$ are disjoint intervals and that for all $k \in \mathbb{N}$

$$
\sup_{t\geq t_k} |\tilde{M}_t(\omega) - \tilde{M}_{t_k}(\omega)| < \varepsilon/2.
$$

For $k \in \mathbb{N}$ and $t \in [t_k, t_k + \varepsilon/(2C_1)]$ one has

$$
|X_t(\omega) - X_{t_k}(\omega)| \leq \int_{t_k}^t |\alpha_s(\omega)| ds + |\tilde{M}_t(\omega) - \tilde{M}_{t_k}(\omega)|
$$

$$
\leq C_1(t - t_k) + \varepsilon/2 \leq \varepsilon
$$

Moreover, since $\varepsilon < C_2$ we conclude for $x = X_t(\omega)$ and $y = X_{t_k}(\omega)$ that

$$
|f(y) - f(x)| \le |H(\xi)| \, |y - x| \le C_1 |y - x|,
$$

where ξ lies on the segment joining x and y. Hence,

$$
|f(X_t(\omega))| \ge |f(X_{t_k}(\omega))| - \underbrace{|f(X_t) - f(X_{t_k}(\omega))|}_{\le \delta/2} \ge \delta/2,
$$

so that

$$
\int_{t_k}^{t_k+\varepsilon/(2C_1)} |f(X_t(\omega))|^2 dt \ge \left(\frac{\delta}{2}\right)^2 \frac{\varepsilon}{2C_1}.
$$

This entails that $\int_{t_0}^{\infty} |f(X_t)|^2 dt = \infty$ and, hence,

$$
-B_t(\omega) = \int_{t_0}^t \langle f(X_s(\omega)), -\alpha_s(\omega) \rangle ds \ge C_2 \int_{t_0}^t |f(X_s(\omega)|^2 ds \to \infty.
$$

This contradicts that $(B_t(\omega))_{t\geq t_0}$ converges to a finite value, by choice of ω .

It remains to prove almost sure convergence of $(X_t)_{t>0}$ on \mathbb{C}_3 in the case where the set $\mathcal{C} := \{x \in \mathbb{R}^d : f(x) = 0\}$ is at most countable. Take $\omega \in \overline{\Omega}$ for which $\lim_{t \to \infty} f(X_t(\omega)) = 0$ and set $r := \limsup_{t\to\infty} |X_t(\omega)| < \infty$. We show by contradiction that $(X_t(\omega))_{t\geq 0}$ converges. Suppose that $(X_t(\omega))_{t>0}$ would not converge. Then there would exist a coordinate $i \in \{1,\ldots,d\}$ and reals $a < b$ such that the set of accumulation points

$$
\mathcal{K} := \bigcap_{n \in \mathbb{N}} \{ X_t^{(i)}(\omega) : t \ge n, X_t(\omega) \in \overline{B(0, r+1)} \}
$$

is a compact set containing $[a, b]$. By applying the principle of nested intervals we can produce for every $u \in [a, b]$ an accumulation point $\eta(u)$ of $(X_t(\omega))_{t>0}$ with ith coordinate equal to u. Recall that $f(X_t) \to 0$ so that $\eta(u)$ is a critical point of F. We thus observe that there is an injective mapping taking $[a, b]$ to the set of critical points of F which contradicts the assumption. \Box

3. Proof of Theorem [1.6](#page-3-0)

The proof of Theorem [1.6](#page-3-0) is arranged as follows. In Section [3.1,](#page-7-1) we provide moment estimates for the Lyapunov function seen by the SDE in a neighbourhood U of a point $y \in \mathbb{R}^d$ so that the Lojasiewicz-inequality is satisfied on U with parameters $L > 0$ and $\theta \in (\frac{1}{2})$ $(\frac{1}{2}, 1)$. To that purpose we fix a particular critical level and "take out" realisations that undershoot the critical level by a certain amount (lower dropout). In the case where no perturbation is present ($\beta_t \equiv 0$), we recover the convergence rate for the respective ODE (3) derived in [Loj84], i.e.

$$
|F(x_t) - F(y)| \le C(t+1)^{-\frac{1}{2\theta-1}}.
$$

We show that the convergence rate is maintained in the SDE setting as long as $|\beta_t| \leq \tilde{C}(t +$ $(1)^{-\frac{\theta}{2\theta-1}}$ for sufficiently large t. In Section [3.2,](#page-9-1) we provide estimates for the drift term $(\int_0^t \alpha_s ds)_{t\geq 0}$ in the setting of Section [3.1](#page-7-1) on the event that no dropout occurs. In Section [3.3,](#page-10-0) we provide estimates that later allow us to show that when a dropout occurs for a particular critical level

it is very likely that the limit of the Lyapunov function lies strictly below that critical level. Finally, in Section [3.4](#page-12-0) we combine the results of the previous subsections to achieve the proof of Theorem [1.6.](#page-3-0)

3.1. Convergence rate for the target value in the case without lower dropout.

Proposition 3.1. Suppose that the following assumptions are satisfied:

Assumptions on the error term. Let $\theta \in (\frac{1}{2})$ $(\frac{1}{2}, 1)$, $(v_t)_{t\geq 0}$ be a continuously differentiable, decaying, positive function and $(\sigma_t)_{t\geq 0}$ be a right-continuous and locally integrable function, so that there exist constants $\kappa_1, \kappa_2 > 0$ such that for all $t \geq 0$

(6)
$$
-\frac{\dot{v}_t}{v_t} \le \kappa_1 v_t^{2\theta - 1} \quad \text{and} \quad \kappa_2 v_t^{2\theta} \ge \sigma_t^2.
$$

No lower dropout/Lojasiewicz-inequality. Let $\ell \in \mathbb{R}$, $L > 0$ and $(w_t)_{t>0} = (C_w v_t)_{t>0}$ for $a C_w \geq 0$. Let T be a stopping time, such that for all $t \geq 0$, on $\{T > t\}$,

$$
|f(X_t)| \ge L|F(X_t) - \ell|^{\theta} , F(X_t) - \ell \ge -w_t
$$

as well as

$$
-\langle f(X_t), \alpha_t \rangle \ge \rho |f(X_t)|^2, \quad \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) \le \sigma_t^2
$$

for a $\rho > 0$. Additionally, suppose that for every $t > 0$

(7)
$$
\mathbb{E}\Big[\int_0^{t\wedge T}|\beta^\dagger_s f(X_s)|^2 ds\Big] < \infty.
$$

Result. Suppose that the above assumptions hold and

$$
\mathbb{E}[\mathbb{1}_{\{T>0\}}(F(X_0)-\ell)] \leq R.
$$

Then there exist $\alpha, \eta > 0$ such that, for all $t \geq 0$, one has

(8)
$$
\mathbb{E}\big[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell+w_t)\big]\leq \Phi_t^{(R)}+\alpha v_t,
$$

where

$$
\Phi_t^{(R)} = R((2\theta - 1)\eta R^{2\theta - 1})t + 1)^{-1/(2\theta - 1)}.
$$

Proof. We assume without loss of generality that $\ell = 0$. By Itô's formula

(9)
$$
dF(X_t) = \langle f(X_t), \alpha_t \rangle dt + \langle f(X_t), \beta_t dW_t \rangle + \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) dt.
$$

Let $(Z_t)_{t\geq 0} := (\mathbb{1}_{\{T>t\}}(F(X_t)+w_t))_{t\geq 0}$. By choice of $T, (Z_t)_{t\geq 0}$ is a non-negative process and we have

(10)
$$
dZ_t = \mathbb{1}_{\{T>t\}} \big(\langle f(X_t), \alpha_t \rangle + \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) + \dot{w}_t \big) dt + dM_t - d\xi_t,
$$

where $(M_t)_{t\geq 0}$ denotes the L^2 -martingale

$$
(M_t)_{t\geq 0} := \Big(\int_0^{T\wedge t} \langle f(X_s)), \beta_s dW_s \rangle\Big)_{t\geq 0},
$$

see assumption [\(7\)](#page-7-2), and $(\xi_t)_{t\geq 0}$ is an increasing process given by

$$
\xi_t := \begin{cases} 0, & \text{if } t < T \text{ or } T = 0, \\ F(X_T) + w_T, & \text{else.} \end{cases}
$$

For all $t \geq 0$, we let $\zeta_t := \mathbb{E}[Z_t]$. In the first step we show that $(\zeta_t)_{t \geq 0}$ is right-continuous and has no upward jumps. By definition, $(Z_t)_{t>0}$ is right-continuous and it is straight-forward to

verify the continuity theorem for integration (Note that Z_t is non-negative and $\langle f(X_t), \alpha_t \rangle \leq 0$ on $\{t < T\}$). To verify that $(\zeta_t)_{t\geq 0}$ has no upward jump at $t > 0$ we note that for $\varepsilon \in (0, t)$

$$
\zeta_t - \zeta_{t-\varepsilon} = \mathbb{E}[Z_t - Z_{t-\varepsilon}] \le \mathbb{E}\Big[\int_{t-\varepsilon}^t \mathbb{1}_{\{s
$$

as $\varepsilon \downarrow 0$.

Take $\alpha \ge C_w$ and $\eta > 0$ such that for all $a, b > 0$

(11)
$$
\rho L^2 2^{-(2\theta-1)} (a+b)^{2\theta} > \eta a^{2\theta} + \frac{\rho L^2 C_w^{2\theta} + \kappa_2 + \kappa_1 \alpha}{\alpha^{2\theta}} b^{2\theta}.
$$

We will now show that $\zeta_t \leq \bar{\zeta}_t := \Phi_t^{(R)} + \alpha v_t$ for all $t \geq 0$. Note that $\zeta_0 \leq R + C_w v_0 \leq \Phi_0^{(R)} + \alpha v_0 =$ $\bar{\zeta}_0$ since $\alpha \geq C_w$ so that the inequality is at least true for $t = 0$. We prove the statement by contradiction. Suppose the time

$$
\tau := \inf\{s \ge 0 : \zeta_s > \bar{\zeta}_s\}
$$

is finite. Since $\zeta - \bar{\zeta}$ has no upwards jumps and starts in a non-positive value we have $\zeta_{\tau} = \bar{\zeta}_{\tau}$. We consider the slope of the right secant of ζ with supporting points τ and $\tau + \varepsilon$ for $\varepsilon > 0$. Using the Lojasiewicz-inequality we get that on $\{T > t\}$

$$
\langle f(X_t), \alpha_t \rangle \le -\rho |f(X_t)|^2 \le -\rho L^2 |F(X_t)|^{2\theta}
$$

$$
\le -\rho L^2 \left(2^{-(2\theta - 1)} |F(X_t) + w_t|^{2\theta} - w_t^{2\theta} \right)
$$

so that with [\(10\)](#page-7-3) and $\frac{1}{2}$ tr $(\beta_t^{\dagger} H(X_t) \beta_t) \leq \sigma^2(t)$

$$
\frac{\zeta_{\tau+\varepsilon}-\zeta_{\tau}}{\varepsilon} \leq \mathbb{E}\Big[\frac{1}{\varepsilon}\int_{\tau}^{\tau+\varepsilon}1\!\!1_{\{T>t\}}(-\rho L^2(2^{-(2\theta-1)}|Z_t|^{2\theta}-w_t^{2\theta})+\sigma_t^2)\,dt\Big].
$$

By non-negativity and right-continuity of $(Z_t)_{t>0}$ and convexity of $|\cdot|^{2\theta}$ one has

$$
\liminf_{\varepsilon \downarrow 0} \mathbb{E} \Big[\frac{1}{\varepsilon} \int_{\tau}^{\tau+\varepsilon} \mathbb{1}_{\{T > t\}} |Z_t|^{2\theta} dt \Big] \geq \mathbb{E} [|Z_{\tau}|^{2\theta}] \geq |\mathbb{E} [Z_{\tau}]|^{2\theta} = \overline{\zeta}_{\tau}^{2\theta}.
$$

Consequently,

(12)
$$
\limsup_{\varepsilon \downarrow 0} \frac{\zeta_{\tau+\varepsilon} - \zeta_{\tau}}{\varepsilon} \leq -\rho L^2 (2^{-(2\theta-1)} \bar{\zeta}_{\tau}^{2\theta} - w_{\tau}^{2\theta}) + \sigma_{\tau}^2
$$

Conversely, $(\Phi_t^{(R)})_{t \geq 0}$ is the solution of the differential equation

$$
\dot{\Phi}_t^{(R)} = -\eta (\Phi_t^{(R)})^{2\theta}, \Phi_0^{(R)} = R
$$

so that

$$
\dot{\bar{\zeta}}_{\tau} = -\eta (\Phi_{\tau}^{(R)})^{2\theta} + \alpha \dot{v}_{\tau}.
$$

Note that if the right-hand side of [\(12\)](#page-8-0) is strictly smaller than $\dot{\bar{\zeta}}_7$, i.e., if

(13)
$$
-\rho L^2 (2^{-(2\theta-1)} \bar{\zeta}_\tau^{2\theta} - w_\tau^{2\theta}) + \sigma_\tau^2 < -\eta (\Phi_\tau^{(R)})^{2\theta} + \alpha \dot{v}_\tau,
$$

then there exists an interval $[\tau, \tau + \varepsilon]$ on which $\zeta_t \leq \overline{\zeta}_t$ which entails that ζ cannot overtake $\overline{\zeta}$ at time τ which is a contradiction to the choice of τ . Using that $w_{\tau} = C_w v_{\tau}$, $\dot{v}_{\tau} \ge -\kappa_1 v_{\tau}^{2\theta}$ and $\sigma_{\tau}^2 \leq \kappa_2 v_{\tau}^{2\theta}$ we conclude that [\(13\)](#page-8-1) is satisfied, if

$$
\rho \mathrm{L}^2 2^{-(2\theta-1)} \big(\underbrace{\Phi_\tau^{(R)} + \alpha v_\tau}_{= \bar{\zeta}_\tau} \big)^{2\theta} > \eta(\Phi_\tau^{(R)})^{2\theta} + \frac{\rho \mathrm{L}^2 C_w^{2\theta} + \kappa_2 + \kappa_1 \alpha}{\alpha^{2\theta}} (\alpha v_\tau)^{2\theta}.
$$

By choice of α and η the latter inequality holds so that we produced a contradiction. This finishes the proof. \Box

Remark 3.2. We discuss Proposition [3.1](#page-7-0) in the case where the critical level ℓ is a local minimum and choose $C_w = 0$ in the proposition. For given $\theta \in (\frac{1}{2})$ $(\frac{1}{2}, 1)$, we note that for $v_t = (t+1)^{-1/(2\theta-1)}$ the condition on the left-hand side of [\(6\)](#page-7-4) is satisfied for an appropriate constant κ_1 and if $\sigma_t = \mathcal{O}((t+1)^{-\theta/(2\theta-1)})$, then also the second condition is satisfied for an appropriate κ_2 . In this case, $(v_t)_{t\geq 0}$ is of the same order as $(\Phi_t^{(R)})_{t\geq 0}$ so that

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell)]=\mathcal{O}((t+1)^{-1/(2\theta-1)}).
$$

This agrees with the corresponding estimates for the ODE convergence. So if $(\sigma_t)_{t\geq 0}$ is of order $\mathcal{O}((t+1)^{-\theta/(2\theta-1)})$ we essentially get the same order of convergence as in the deterministic setting.

Now suppose that $\sigma_t \approx (t+1)^{-\sigma}$ with $0 \leq \sigma < \theta/(2\theta-1)$. Then one can choose $v_t = (t+1)^{-\sigma/\theta}$ and constants κ_1 and κ_2 appropriately so that [\(6\)](#page-7-4) is satisfied. So in that case

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell)]=\mathcal{O}\big((t+1)^{-\sigma/\theta}\big).
$$

Altogether, we thus get for the choice $\sigma_t \approx (t+1)^{-\sigma}$ with $\sigma \in [0,\infty)$ that

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell)]=\mathcal{O}\big((t+1)^{-(\frac{\sigma}{\theta}\wedge \frac{1}{2\theta-1})}\big).
$$

Observing that on a compact set F , f and the Hessian are uniformly bounded it is straightforward to infer the statement of Theorem [1.8.](#page-3-1)

3.2. Bounding the drift term in the case where no dropout occurs.

Proposition 3.3. Suppose that T is such that on $\{T \ge t\}$, $\langle f(X_t), -\alpha_t \rangle \ge 0$ and $(F(X_t))_{t>0}$ converges on $\{T = \infty\}$, almost surely. Let $\Phi : [0, \infty) \to [0, \infty)$ be a decreasing, continuously differentiable function so that $t \mapsto -\dot{\Phi}_t$ is decreasing and

$$
\lim_{t \to \infty} \frac{\Phi_t}{\sqrt{-\dot{\Phi}_t}} = 0.
$$

If for every $t \geq 0$,

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t) - F(X_T))] + \frac{1}{2}\mathbb{E}\Big[\int_t^\infty \mathbb{1}_{\{s \le T\}} \text{tr}(\beta_s^\dagger H(X_s)\beta_s) ds\Big] \le \Phi_t
$$

and $(M_t)_{t>0}$ given by

$$
M_t := \int_0^{t \wedge T} \langle f(X_s), \beta_s dW_s \rangle
$$

is a regular martingale, then

$$
\mathbb{E}\Big[\int_0^\infty 1\!\!1_{\{s
$$

Proof. Since $(M_t)_{t>0}$ is a regular martingale we get with Itô's formula, see [\(9\)](#page-7-5), that

$$
\Psi(t) := \mathbb{E}\Big[\int_t^{\infty} \mathbb{1}_{\{s < T\}} \langle f(X_s), -\alpha_s \rangle ds\Big]
$$

=
$$
\mathbb{E}\big[\mathbb{1}_{\{T > t\}} (F(X_t) - F(X_T))\big] + \frac{1}{2} \mathbb{E}\Big[\int_t^{\infty} \mathbb{1}_{\{T > s\}} \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) ds\Big] \leq \Phi_t.
$$

The function Ψ is right-continuous and monotonically decreasing and, hence, càdlàg. We let $\varphi_s := (-\dot{\Phi}_s)^{-1/2}$ and note that by Cauchy-Schwarz

$$
\mathbb{E}\Bigl[\int_0^\infty 1\!\!1_{\{s
$$

Moreover, using partial integration we get that

$$
\int_0^\infty \varphi_s \mathbb{E} \big[\mathbb{1}_{\{s < T\}} \langle f(X_s), -\alpha_s \rangle \big] ds = -\int_0^\infty \varphi_s d\Psi_s
$$

=
$$
-[\varphi \Psi]_0^\infty + \int_0^\infty \Psi_s d\varphi_s.
$$

 $(\varphi_s)_{s\geq 0}$ is increasing by assumption and $\varphi_s\Phi_s\to 0$ as $s\to\infty$ so that

$$
-[\varphi\Psi]_0^{\infty} + \int_0^{\infty} \Psi_s d\varphi_s \le -\varphi_0(\Phi_0 - \Psi_0) - [\varphi\Phi]_0^{\infty} + \int_0^{\infty} \Phi_s d\varphi_s
$$

= $-\varphi_0(\Phi_0 - \Psi_0) + \int_0^{\infty} \underbrace{\varphi_s(-\dot{\Phi}_s)}_{=\sqrt{-\dot{\Phi}_s}} ds.$

Altogether we get that

$$
\mathbb{E}\Bigl[\int_0^\infty 1\!\!1_{\{s
$$

3.3. Technical analysis of lower dropouts. Roughly speaking, the following two lemmas will later be used to show that for a certain critical level of the target function, a lower dropout (in the sense of the previous two propositions) entails that the Lyapunov function converges to a value strictly below the respective critical level with high probability.

Lemma 3.4. Let $(M_t)_{t\geq 0}$ be a continuous local martingale started in zero. Then for every $\kappa > 0$

$$
\mathbb{P}\Big(\sup_{t\geq 0} (M_t - \langle M\rangle_t) \geq \kappa\Big) \leq \frac{1}{\kappa^2} + \sum_{n\in\mathbb{N}_0} \frac{2^{n+1}}{(2^n + \kappa)^2} =: \phi(\kappa).
$$

In particular, for every $\varepsilon > 0$ there exists a $\kappa > 0$ such that the above right-hand side is smaller than ε .

Proof. For $n \in \mathbb{N}_0$, let $T_n := \inf\{t \geq 0 : \langle M \rangle_t > 2^n\}$. Then we have

$$
\sup_{t \in [T_n, T_{n+1})} (M_t - \langle M \rangle_t) \le \sup_{t \in [0, T_{n+1})} M_t - 2^n
$$

and

$$
\sup_{t\in[0,T_0)}(M_t-\langle M\rangle_t)\leq \sup_{t\in[0,T_0)}M_t.
$$

We use Doob's L^2 -inequality to deduce that

$$
\mathbb{P}\Big(\sup_{t\in[0,T_{n+1})} M_t \ge 2^n + \kappa\Big) \le (2^n + \kappa)^{-2} \mathbb{E}[M_{T_{n+1}}^2] \le \frac{2^{n+1}}{(2^n + \kappa)^2}
$$

$$
\mathbb{P}\Big(\sup_{t\in[0,T_0)} M_t \ge \kappa\Big) \le \frac{1}{\kappa^2}.
$$

and

$$
f_{\rm{max}}
$$

Therefore,

$$
\mathbb{P}\Big(\sup_{t\geq 0} (M_t - \langle M \rangle_t) \geq \kappa\Big) \leq \frac{1}{\kappa^2} + \sum_{n \in \mathbb{N}_0} \frac{2^{n+1}}{(2^n + \kappa)^2}.
$$

Lemma 3.5. Let $t_0 \geq 0$, ρ , $\kappa : [t_0, \infty) \to (0, \infty)$ be functions and $t_0 \leq T' \leq T$ be two stopping times such that, for every $t \ge t_0$, on $\{T' \le t \le T\}$,

$$
|\beta_t^{\dagger} f(X_t)|^2 \leq \rho_{T'} \langle f(X_t), -\alpha_t \rangle \quad \text{and} \quad \frac{1}{2} \int_{T'}^t \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) \, ds \leq \kappa_{T'}.
$$

Then,

$$
\mathbb{P}\Big(\sup_{t\in[T',T]} F(X_t) - F(X_{T'}) \ge 2\kappa_{T'}\Big) \le \phi\Big(\inf_{t\ge t_0} \frac{\kappa_t}{\rho_t}\Big).
$$

Proof. We use again representation [\(5\)](#page-4-2):

$$
dF(X_t) = \langle f(X_t), \alpha_t \rangle dt + \langle f(X_t), \beta_t dW_t \rangle + \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) dt.
$$

Consider the local martingale

$$
(M_t)_{t \ge t_0} := \left(\int_0^t \mathbb{1}_{\{T' \le s \le T\}} \langle f(X_s), \beta_s dW_s \rangle \right)_{t \ge t_0}
$$

and note that

$$
\langle M \rangle_t = \int_0^t \mathbb{1}_{\{T' \le s \le T\}} |\beta_s^{\dagger} f(X_s)|^2 ds \le \rho_{T'} \int_0^t \mathbb{1}_{\{T' \le s \le T\}} \langle f(X_s), -\alpha_s \rangle ds.
$$

Hence,

$$
\int_0^t \mathbb{1}_{\{T' \le s \le T\}} \langle f(X_s), \alpha_s \rangle \, ds + \int_0^t \mathbb{1}_{\{T' \le s \le T\}} \langle f(X_s), \beta_s \, dW_s \rangle
$$

$$
\le M_t - \frac{1}{\rho_{T'}} \langle M \rangle_t =: \Xi_t.
$$

Note that $\Xi_t = \rho_{T}(\frac{1}{\rho_{T'}}M_t - \langle \frac{1}{\rho_{T'}}M \rangle_t)$ with $(\frac{1}{\rho_{T'}}M_t)_{t \geq t_0}$ being a continuous local martingale started in zero so that with Lemma [3.4](#page-10-1)

$$
\mathbb{P}\Big(\sup_{t\geq t_0} \Xi_t \geq \kappa_{T'}\Big) = \mathbb{P}\Big(\sup_{t\geq t_0} \frac{1}{\rho_{T'}} M_t - \langle \frac{1}{\rho_{T'}} M \rangle_t \geq \frac{\kappa_{T'}}{\rho_{T'}}\Big) \leq \mathbb{P}\Big(\sup_{t\geq t_0} \frac{1}{\rho_{T'}} M_t - \langle \frac{1}{\rho_{T'}} M \rangle_t \geq \inf_{t\geq t_0} \frac{\kappa_t}{\rho_t}\Big) \leq \phi\Big(\inf_{t\geq t_0} \frac{\kappa_t}{\rho_t}\Big).
$$

Altogether, using

$$
\sup_{T' \le t \le T} F(X_t) - F(X_{T'}) \le \sup_{T' \le t \le T} \Xi_t + \underbrace{\sup_{T' \le t \le T} \frac{1}{2} \int_{T'}^t \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) ds}_{\le \kappa_{T'}}
$$

we get that

$$
\mathbb{P}\Big(\sup_{T'\leq t\leq T} F(X_t)-F(X_{T'})\geq 2\kappa_{T'}\Big)\leq \mathbb{P}\Big(\sup_{t\geq t_0}\Xi_t\geq \kappa_{T'}\Big)\leq \phi\Big(\inf_{t\geq t_0}\frac{\kappa_t}{\rho_t}\Big).
$$

3.4. Proof of Theorem [1.6.](#page-3-0) We cite Lemma 3.7 from [\[DK21\]](#page-20-20).

Lemma 3.6. Let $F : \mathbb{R}^d \to \mathbb{R}$ be a Lojasiewicz-function with differential f and $K \subset \mathbb{R}^d$ be an arbitrary compact set.

(1) The set of critical levels

$$
\mathcal{L}_K := \{ F(x) : x \in \mathcal{C}_F \cap K \}
$$

is finite so that F has at most a countable number of critical levels.

(2) For every critical level $\ell \in \mathcal{L}_K$ there exists an open neighbourhood $U \supset F^{-1}(\{\ell\}) \cap K$, $L > 0, \ \theta \in \left[\frac{1}{2}\right]$ $(\frac{1}{2}, 1)$ such that for every $y \in U$

$$
|f(y)| \ge L|F(y) - \ell|^{\theta}.
$$

(3) For a neighbourhood as in (2), there exists $\varepsilon > 0$ such that

$$
F^{-1}((\ell-\varepsilon,\ell+\varepsilon))\cap K\subset U.
$$

Now we are able to prove the main result of this article.

Proof of Theorem [1.6.](#page-3-0) For $\rho, C, t_0 > 0$ and a set $K \subset \mathbb{R}^d$ let

$$
T_{\rho,C,t_0,K} := \inf \Big\{ t \ge t_0 : \langle f(X_t), -\alpha_t \rangle < \rho |f(X_t)|^2, \ |\alpha_t| > C |f(X_t)|, \ |f(X_t)| > C,
$$
\n
$$
|\beta_t|^2 > C\sigma_t^2, \ \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t)\beta_t) > C\sigma_t^2 \ \text{or} \ X_t \notin K \Big\}.
$$

Set $\theta_0 := \frac{\sigma}{2\sigma - 1} \in \left(\frac{1}{2}\right)$ $(\frac{1}{2}, 1).$

1) First let U be an open and bounded set such that for a $\theta \in (\theta_0, 1)$, an $\ell \in \mathbb{R}$ and $L > 0$, for all $y \in U$

$$
|f(y)| \ge \mathcal{L}|F(y) - \ell|^{\theta}.
$$

1.a) We show that on $\{T_{\rho,C,t_0,U}=\infty\}$, almost surely, the random set

$$
S := \{ s \ge 0 : F(X_s) < \ell - (s+1)^{-\frac{1}{2\theta - 1}} \}
$$

is bounded.

By Proposition [2.1](#page-4-0) we have on $\{T_{\rho,C,t_0,U}=\infty\}$ that, almost surely, $(F(X_t))_{t\geq0}$ converges and $f(X_t) \to 0$, which together with the Lojasiewicz-inequality implies that $F(X_t) \to \ell$.

By choice of θ one has $2\sigma > \frac{2\theta}{2\theta - 1}$. Thus, for $t_1 \ge t_0$ large enough we get for all $t \ge t_1$

$$
C\sigma_t^2 \le \frac{1}{4\theta - 2}(t+1)^{-\frac{2\theta}{2\theta - 1}}.
$$

Let $T'_{t_1} := \inf\{t \ge t_1 : F(X_t) < \ell - (t+1)^{-\frac{1}{2\theta-1}}\}$, then one has

$$
\sup_{T'_{t_1} \leq t \leq T_{\rho,C,t_0,U}} \int_{T'_{t_1}}^t \frac{1}{2} \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) ds \leq \frac{1}{4\theta - 2} \int_{T'_{t_1}}^{\infty} (s+1)^{-\frac{2\theta}{2\theta - 1}} ds = \frac{1}{2} (T'_{t_1} + 1)^{-\frac{1}{2\theta - 1}}.
$$

Furthermore, we have for t with $T'_{t_1} \leq t < T_{\rho,C,t_0,U}$

$$
|\beta_t^{\dagger} f(X_t)|^2 \leq |\beta_t^{\dagger}|^2 |f(X_t)|^2 \leq \frac{C}{\rho} (T'_{t_1} + 1)^{-2\sigma} \langle f(X_t), -\alpha_t \rangle
$$

so that, using Lemma [3.5,](#page-11-0) we get

$$
\mathbb{P}(S \cap [t_1, \infty) \neq \emptyset, T_{\rho, C, t_0, U} = \infty) = \mathbb{P}(T'_{t_1} < \infty, T_{\rho, C, t_0, U} = \infty, F(X_t) \to \ell)
$$

$$
\leq \mathbb{P}\Big(\sup_{t \in [T'_{t_1}, T_{\rho, C, t_0, U})} F(X_t) - F(X_{T'_{t_1}}) \geq (T'_{t_1} + 1)^{-\frac{1}{2\theta - 1}}\Big)
$$

$$
\leq \phi\Big(\frac{\rho}{2C}(t_1 + 1)^{2\sigma - \frac{1}{2\theta - 1}}\Big) \stackrel{t_1 \to \infty}{\longrightarrow} 0
$$

and for all $t_1 \geq t_0$

$$
\mathbb{P}(S \text{ is unbounded}, T_{\rho,C,t_0,U} = \infty) \leq \mathbb{P}(S \cap [t_1,\infty) \neq \emptyset, T_{\rho,C,t_0,U} = \infty) \to 0.
$$

1.b) Let

$$
\bar{T}_{\rho,C,t_0,U} := T_{\rho,C,t_0,U} \wedge \inf\{t \ge t_0 : X_t < \ell - (t+1)^{-\frac{1}{2\theta-1}}\}.
$$

We show that on $\{\bar{T}_{\rho,C,t_0,U}=\infty\}$ the process $(X_t)_{t\geq0}$ converges, almost surely.

We set $\sigma'_t := \sqrt{C}\sigma_t$ and $v_t := w_t := (t+1)^{-\frac{1}{2\theta-1}}$ (so that $C_w = 1$) and $T := \overline{T}_{\rho,C,t_0,U}$ and show that the functions satisfy the Assumptions of Proposition [3.1,](#page-7-0) Proposition [3.3](#page-9-2) and the right-hand side of the estimate in Proposition [3.3](#page-9-2) is finite.

First note that one can apply Proposition [3.1](#page-7-0) for the process $(X_t)_{t\geq t_0}$ started at time t_0 in X_{t_0} . In particular, we have

$$
-\dot{v}_t = \frac{1}{2\theta - 1}(t+1)^{-\frac{2\theta}{2\theta - 1}} = \frac{1}{2\theta - 1}v_t^{2\theta}
$$

and with $2\sigma > \frac{2\theta}{2\theta - 1}$

$$
v_t^{2\theta} = (t+1)^{-\frac{2\theta}{2\theta-1}} \ge \frac{1}{C} (\sigma'_t)^2
$$

so that [\(6\)](#page-7-4) is satisfied. Further, with the definition of $T_{\rho,C,t_0,K}$

$$
\mathbb{E}\Big[\int_{t_0}^T |\beta_s^{\dagger} f(X_s)|^2 ds\Big] \leq \int_{t_0}^{\infty} C^3 \sigma_t^2 dt < \infty,
$$

so that [\(7\)](#page-7-2) is satisfied. Now, we can choose $\eta > 0$ sufficiently small and $\alpha > 0$ sufficiently large so that (11) is satisfied and with the boundedness of U and Proposition [3.1](#page-7-0) we get for sufficiently large $R, C' > 0$ that for all $t \ge t_0$

$$
\mathbb{E}\big[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell+w_t)\big]\leq \Phi_t^{(R)}+\alpha v_t\leq C'(t-t_0+1)^{-\frac{1}{2\theta-1}}.
$$

Regarding Proposition [3.3](#page-9-2) we recall that, on $\{T = \infty\}$, we almost surely have $F(X_t) \to \ell$ so that $F(X_T)$ is well-defined. By the definition of T we have $-F(X_T) \le -\ell + w_T$, where $w_{\infty} = 0$, and the monotonicity of (w_t) gives

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-F(X_T))] \leq \mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t)-\ell+w_t)] \leq C'(t-t_0+1)^{-\frac{1}{2\theta-1}}.
$$

Thus, we have

$$
\mathbb{E}[\mathbb{1}_{\{T>t\}}(F(X_t) - F(X_T))] + \frac{1}{2} \mathbb{E}\Big[\int_t^\infty \mathbb{1}_{\{s \le T\}} \text{tr}(\beta_s^\dagger H(X_s)\beta_s) ds\Big]
$$

$$
\le C'(t - t_0 + 1)^{-\frac{1}{2\theta - 1}} + C \int_t^\infty \sigma_s^2 ds \le \Phi_t,
$$

where $\Phi_t := C''(t-t_0+1)^{-\frac{1}{2\theta-1}}$ for sufficiently large C'' . Moreover, $\dot{\Phi}_t$ is an increasing function so that $t \mapsto -\dot{\Phi}_t$ is decreasing and using that $\theta < 1$ we get that

$$
\Phi_t/\sqrt{-\dot{\Phi}_t} = \sqrt{2\theta - 1} \left(C'' \right)^{\theta - 1/2} (\Phi_t)^{1-\theta} \to 0.
$$

Moreover, $(M_t)_{t\geq t_0} := \left(\int_{t_0}^{t\wedge T} \langle f(X_s), \beta_s dW_s \rangle\right)_{t\geq t_0}$ is a regular martingale as

$$
\langle M_t \rangle_{\infty} = \int_{t_0}^T |\beta_s^{\dagger} f(X_s)|^2 ds \le \int_{t_0}^{\infty} C^3 \sigma_t^2 dt < \infty.
$$

Therefore, Proposition [3.3](#page-9-2) is applicable and we get

$$
\mathbb{E}\Big[\int_{t_0}^{\infty} \mathbb{1}_{\{s
\n
$$
\leq C\sqrt{\rho} \mathbb{E}\Big[\int_{t_0}^{\infty} \mathbb{1}_{\{s
\n
$$
\leq C\sqrt{\rho} \int_{t_0}^{\infty} \sqrt{-\dot{\Phi}_s} ds = C\sqrt{\frac{\rho C''}{2\theta - 1}} \int_{t_0}^{\infty} (t - t_0 + 1)^{-\frac{\theta}{2\theta - 1}} ds
$$

\n
$$
= C\sqrt{\rho C''} \frac{\sqrt{2\theta - 1}}{1 - \theta} < \infty.
$$
$$
$$

This implies almost sure convergence of $(\int_{t_0}^t \alpha_s ds)_{t\geq t_0}$, on $\{\bar{T}_{\rho,C,t_0,U} = \infty\}$. Note that also $(\int_{t_0}^{\infty} \beta_t dW_t)_{t \geq t_0}$ converges almost surely on $\{T_{\rho, C, t_0, U} = \infty\}$, since

$$
\int_{t_0}^{\bar{T}_{\rho,C,t_0,U}} |\beta_s|_F^2 ds \leq \int_{t_0}^{\infty} C \sigma_s^2 ds < \infty.
$$

Hence, we have almost sure convergence of

$$
X_t - X_{t_0} = \int_{t_0}^t \alpha_t dt + \int_{t_0}^t \beta_t dW_t,
$$

on $\{\bar{T}_{\rho,C,t_0,U}=\infty\}.$

1.c) Combining 1.a) and 1.b) we conclude that on $\{T_{\rho,C,t_0,U} = \infty\}$ we have, almost sure, convergence of $(X_t)_{t>0}$. Indeed, by 1.a) we have that up to nullsets $\{T_{\rho,C,t_0,U}=\infty\}$ is contained in

$$
\bigcup_{n\in\mathbb{N}}\{\bar{T}_{\rho,C,t_0+n,U}=\infty\}
$$

and by 1.b) on each of the latter sets $(X_t)_{t\geq0}$ converges almost surely.

2) Now let $K \subset \mathbb{R}^d$ be a compact subset. We show that we have almost sure convergence of $(X_t)_{t>0}$ on $\{T_{\rho,C,t_0,K}=\infty\}$. The statement then follows by observing that

$$
\mathbb{C} \cap \mathbb{L} \cap \{ \limsup_{t \to \infty} \frac{|\beta_t|}{\sigma_t} < \infty \} = \bigcup_{n \in \mathbb{N}} \{ T_{\frac{1}{n}, n, n, [-n, n]^d} = \infty \}.
$$

By Lemma [3.6,](#page-12-1) K contains finitely many critical levels, say $\ell_1, \ldots, \ell_m \in \mathbb{R}$, and there exists $\varepsilon > 0, L > 0, \theta \in \left[\frac{\sigma}{2\sigma-1}, 1\right)$ and bounded open neighbourhoods $U_k \supset F^{-1}((\ell_k - \varepsilon, \ell_k + \varepsilon)) \cap K$ such that for each $k = 1, ..., m$ and $y \in U_k$,

$$
|f(y)| \ge \mathcal{L}|F(y) - \ell_k|^{\theta}.
$$

By Theorem [1.1,](#page-2-1) we have that on $\{T_{\rho,C,t_0,K}=\infty\}$, almost surely, the limit $\lim_{t\to\infty} F(X_t)$ exists and $\lim_{t\to\infty} f(X_t) = 0$. Since $\inf_{y\in K\setminus (U_1\cup...\cup U_m)} |f(y)| > 0$, we get that on $\{T_{\rho,C,t_0,U} = \infty\}$, almost surely, $(X_t)_{t\geq0}$ attains from a random time onwards only values in $U_1\cup\ldots\cup U_m$ and using the Lojasiewicz-inequality we get that $(F(X_t))_{t\geq 0}$ converges to one of the levels ℓ_k and,

in particular, attains from a random time onwards only values in a U_k (which may be random). Hence, up to nullsets, $\{T_{\rho,C,t_0,K}=\infty\}$ is contained in

$$
\bigcup_{n\in\mathbb{N}}\bigcup_{k=1}^m\{T_{\rho,C,t_0+n,U_k}=\infty\}.
$$

By 1.c) we have almost sure convergence of $(X_t)_{t>0}$ on each of the countably many latter events. \Box

4. Optimality of Theorem [1.6](#page-3-0)

In this section, we provide an example which satisfies the assumptions of Theorem [1.6](#page-3-0) when choosing $\sigma_t = (t+1)^{-1}$, but for which $(X_t)_{t\geq 0}$ does not converge.

We consider a rotationally invariant Lyapunov function F on \mathbb{R}^2 . Let

$$
\psi : [0, \infty) \to [0, \infty), r \mapsto \begin{cases} (1-r)^2, & \text{if } r \ge \frac{1}{2}, \\ \frac{8}{3}r^3 - 3r^2 + \frac{2}{3}, & \text{else,} \end{cases}
$$

and set $F: \mathbb{R}^2 \to \mathbb{R}$, $F(x) := \psi(|x|)$. Note that F is a C^2 -function with critical points

$$
\mathcal{C} := f^{-1}(\{0\}) = \{x \in \mathbb{R}^2 : |x| \in \{0, 1\}\},\
$$

where we again denote by $f := \nabla F$ the gradient of F. For every $(x_1, x_2) \in \mathbb{R}^2$ we let

$$
ort(x_1, x_2) := (-x_2, x_1).
$$

and denote by $\varphi : \mathbb{R}^2 \to [0, 1]$ a rotationally invariant C^{∞} -function with

$$
\varphi|_{B(0,2)\setminus B(0,\frac{2}{3})}\equiv 1
$$
 and $\varphi|_{B(0,\frac{1}{2})\cup B(0,3)^c}\equiv 0.$

Consider the SDE

$$
dX_t = \left(-f(X_t) + \operatorname{ort}(X_t) ||X_t| - 1|\right) dt + \frac{1}{t+1} \varphi(X_t) dW_t
$$

started in a point $x_0 \in \mathbb{R}^2 \setminus \{0\}$, where $(W_t)_{t \geq 0}$ is a 2-dimensional Brownian motion with initial value $W_0 = 0$.

Theorem 4.1. (i) F is a Lojasiewicz-function.

(ii) $\mathbb{P}(\mathbb{C} \cap \mathbb{L}) = 1$.

(iii) Almost surely, $(X_t)_{t\geq0}$ does not converge.

The proof is based on the following proposition.

Proposition 4.2. For $t \geq 0$ consider the stopping time

$$
T_t := \inf\big\{s \ge t : X_s \not\in B(0,2) \backslash B(0,\frac{2}{3})\big\}.
$$

- (i) $\mathbb{P}(\mathbb{C} \cap \mathbb{L}) = 1$ and, almost surely, $\lim_{t \to \infty} |X_t| = 1$. In particular, almost surely, for all sufficiently large t, $T_t = \infty$.
- (ii) For every $t \geq 0$, $(\bar{Z}_s)_{s \geq t} := (|X_s| 1)_{s \geq t}$ solves on the random time interval $[t, T_t]$ the SDE

$$
d\bar{Z}_s = \left(-2\bar{Z}_s + \frac{1}{2}\frac{1}{\bar{Z}_s + 1}\frac{1}{(s+1)^2}\right)ds + \frac{1}{s+1}dB_s
$$

where $B_s := \int_t^s$ 1 $\frac{1}{|X_u|}\langle X_u, dW_u \rangle$ is a Brownian motion. (iii) For $t \geq 0$ let $(Z_s^{(t)})_{s \geq t}$ denote the solution to

(14)
$$
Z_t^{(t)} = \bar{Z}_t \text{ and } dZ_s^{(t)} = -2Z_s^{(t)} ds + \frac{1}{s+1} dB_s
$$

Then,

$$
\int_{t}^{T_t} |\bar{Z}_s - Z_s^{(t)}| ds \le \frac{1}{2} \frac{1}{t+1}.
$$

Proof. (i): For the choice

$$
\alpha_t := -f(X_t) + \operatorname{ort}(X_t) ||X_t| - 1| \text{ and } \beta_t := \frac{1}{t+1} \varphi(X_t) \mathbb{I}_2
$$

 $(X_t)_{t\geq0}$ solves the SDE [\(1\)](#page-0-1). We verify that the events B and C are almost sure events. Note that the diffusivity on $B(0, \frac{1}{2})$ $\frac{1}{2}$)∪ $B(0, 3)^c$ is zero and that the vector field pushes $(X_t)_{t\geq 0}$ towards values in $\overline{B(0,3)}\backslash B(0,\frac{1}{2})$ $\frac{1}{2}$). Hence, from a deterministic time, say t_0 , onwards the process $(X_t)_{t\geq 0}$ stays on the compact set $\overline{B(0,3)}\backslash B(0,\frac{1}{2})$ $\frac{1}{2}$). On the latter compact set F, ∇F and the Hessian H of F are uniformly bounded since F is C^2 . Therefore, B is an almost sure set.

For $t \geq t_0$, by orthogonality of f and ort one has

$$
\langle f(X_t), -\alpha_t \rangle = |f(X_t)|^2
$$

and, additionally,

$$
|\alpha_t| = |-f(X_t) + \text{ort}(X_t)||X_t| - 1|| \le \underbrace{|f(X_t)|}_{=2|1-|X_t|} + \underbrace{|X_t|}_{\le 3} ||X_t| - 1| \le 5|1-|X_t|| = \frac{5}{2}|f(X_t)|
$$

Moreover,

$$
\int_0^\infty |\beta_s|_F^2 ds \le \int_0^\infty \frac{1}{(s+1)^2} |\mathbb{I}_2|_F^2 ds = 2 \int_0^\infty \frac{1}{(s+1)^2} ds < \infty.
$$

Consequently, $\mathbb C$ is an almost sure event. By Theorem [1.1,](#page-2-1) $(f(X_t))_{t\geq0}$ converges, almost surely, to zero so that $\lim_{t\to\infty} |X_t| = 1$, almost surely.

(ii): Note that $|\cdot| : \mathbb{R}^2 \setminus \{0\} \to \mathbb{R}$ is C^{∞} and, on $[t, T_t], -f(X_t) = -2\overline{Z}_t \frac{X_t}{|X_t|}$ $\frac{X_t}{|X_t|}$ as well as $\langle X_t, \text{ort}(X_t) \rangle = 0$, so that with the Itô-formula we get

$$
d\bar{Z}_s = \frac{1}{|X_s|} \langle X_s, dX_s \rangle + \frac{1}{2} \frac{1}{(s+1)^2} \frac{1}{|X_s|} ds
$$

= $\left(-2\bar{Z}_s + \frac{1}{2} \frac{1}{\bar{Z}_s + 1} \frac{1}{(s+1)^2} \right) ds + \frac{1}{s+1} dB_s.$

(iii): Fix $t \geq 0$ and consider $(\Upsilon_s^{(t)})_{s \geq t} := (\bar{Z}_s - Z_s^{(t)})_{s \geq t}$. Then on $[t, T_t]$

$$
d\Upsilon_s^{(t)} = -2\Upsilon_s^{(t)} ds + \frac{1}{2} \frac{1}{\bar{Z}_s + 1} \frac{1}{(s+1)^2} ds.
$$

With $\Upsilon_t^{(t)} = 0$ we thus get that for $t \leq s \leq T_t$

$$
\Upsilon_s^{(t)} = \frac{1}{2} \int_t^s e^{-2(s-u)} \frac{1}{\bar{Z}_u + 1} \frac{1}{(u+1)^2} du \le \int_t^s e^{-2(s-u)} \frac{1}{(u+1)^2} du,
$$

where we used that for $t \le u \le T_t$, $\frac{1}{\bar{Z}_u+1} \le 2$. We thus get with Fubini that

$$
\int_{t}^{T_t} |\bar{Z}_s - Z_s^{(t)}| ds \le \int_{t}^{\infty} \int_{t}^{s} e^{-2(s-u)} \frac{1}{(u+1)^2} du ds
$$

=
$$
\int_{t}^{\infty} \underbrace{\int_{u}^{\infty} e^{-2(s-u)} ds}_{= \frac{1}{2}} \frac{1}{(u+1)^2} du = \frac{1}{2} \frac{1}{t+1}.
$$

Now, let $t \geq 0$ and $x_0 \in \mathbb{R}$ and let $(Z_s)_{s \geq t}$ be the solution of the SDE

(15)
$$
dZ_s = -2Z_s ds + (s+1)^{-1} dB_s, \text{ with } Z_t = x_0.
$$

Using Itô's lemma and the Dubins-Schwarz theorem it is straight-forward to see that one can represent the solution as

$$
(Z_s)_{s \geq t} = (e^{-2s}\tilde{B}_{g_s})_{s \geq t},
$$

where $(\tilde{B}_s)_{s\geq g_t}$ is a Brownian motion started at time g_t in $\tilde{B}_{g_t}=e^{2t}x_0$ and $(g_s)_{s\geq 0}$ is given by

$$
g_s := \int_0^s (u+1)^{-2} e^{4u} du \sim \frac{1}{4} (s+1)^{-2} e^{4s}.
$$

Note that $g_s \leq \frac{e^{4s}}{(s+1)^2}$.

Lemma 4.3. For every $t \geq 0$ and $x_0 \in \mathbb{R}$ the solution $(Z_s)_{s \geq t}$ of [\(15\)](#page-17-0) satisfies

$$
\int_t^\infty |Z_s| \, ds = \infty, \qquad \text{almost surely.}
$$

Proof. We let for $s \geq t$, $\overline{B}_s := \widetilde{B}_{g_s} - e^{2t}x_0$ and note that

$$
\left| \int_{t}^{u} |Z_{s}| ds - \int_{t}^{u} |e^{-2s} \bar{B}_{s}| ds \right| \leq |x_{0}| \int_{0}^{\infty} e^{-2s} ds = \frac{1}{2} |x_{0}|.
$$

Hence, it suffices to show that $\int_t^{\infty} |e^{-2s}\bar{B}_s| ds = \infty$, almost surely. Set

$$
\kappa := \int_{\mathbb{R}} \frac{1}{\sqrt{2\pi}} |x| \, e^{-x^2/2} \, dx = \sqrt{\frac{2}{\pi}}.
$$

One has

$$
\mathbb{E}\Big[\int_t^u e^{-2s}|\bar{B}_s|ds\Big] = \int_t^u e^{-2s}\mathbb{E}[|\bar{B}_s|]ds = \kappa \int_t^u e^{-2s}\sqrt{g_s - g_t}ds.
$$

Using that $g_s \sim \frac{1}{4}$ $\frac{1}{4}(s+1)^{-2}e^{4s}$ we get that

$$
\kappa \int_t^u e^{-2s} \sqrt{g_s - g_t} \, ds \sim \frac{1}{2} \kappa \int_t^u (s+1)^{-1} \, ds \sim \frac{1}{2} \kappa \log u.
$$

To estimate the variance we first show that for $v\geq u\geq t$ we have

(16)
$$
\text{cov}(|\bar{B}_u|, |\bar{B}_v|) \le g_v - g_t.
$$

 \Box

Indeed,

$$
\mathbb{E}[|\bar{B}_v \bar{B}_u|] \leq \mathbb{E}[\bar{B}_u^2] + \mathbb{E}[|\bar{B}_v - \bar{B}_u|] \mathbb{E}[|\bar{B}_u|]
$$

$$
\leq g_u - g_t + \kappa^2 \sqrt{g_v - g_u} \sqrt{g_u - g_t}
$$

and

$$
\mathbb{E}[|\bar{B}_v|] \mathbb{E}[|\bar{B}_u|] = \kappa^2 \sqrt{g_v - g_t} \sqrt{g_u - g_t}.
$$

Thus [\(16\)](#page-17-1) follows since $g_u \geq g_t$. We conclude that for all $\ell \geq t$

$$
\operatorname{Var}\Bigl(\int_t^{\ell} e^{-2s} |\bar{B}_s| ds\Bigr) = \int_t^{\ell} \int_t^{\ell} e^{-2(u+v)} \operatorname{cov}(|\bar{B}_u|, |\bar{B}_v|) du dv
$$

$$
\leq 2 \int_t^{\ell} \int_u^{\ell} e^{-2(u+v)} g_u dv du
$$

$$
\leq \int_0^{\ell} e^{-4u} g_u du \leq \int_0^{\ell} \frac{1}{(u+1)^2} du \leq 1.
$$

Hence, as a consequence of the Chebyshev inequality one gets that

$$
\int_{t}^{\infty} e^{-2s} |\bar{B}_s| ds = \lim_{u \to \infty} \int_{t}^{u} e^{-2s} |\bar{B}_s| ds = \infty, \text{ almost surely.}
$$

Proof of Theorem [4.1.](#page-0-2) (i): We first verify that F is a Lojasiewicz-function. We need to analyse the critical points only. One has

$$
\operatorname{Hess} F(0) = \begin{pmatrix} -6 & 0\\ 0 & -6 \end{pmatrix}
$$

so that the Hessian has full rank at 0 which implies validity of a Lojasiewicz-inequality with $\theta = \frac{1}{2}$ $\frac{1}{2}$ on an appropriate neighbourhood of 0. Next, let $x \in \mathbb{R}^2$ with $|x| = 1$. Since $\psi''(1) = 2$, ψ satisfies a Lojasiewicz-inequality

$$
|\psi'(r)| \ge \mathcal{L} |\psi(r) - \psi(1)|^{\frac{1}{2}}
$$

for all $r \in [1-\varepsilon, 1+\varepsilon)$ and appropriately fixed $\varepsilon \in (0, \frac{1}{2})$ $\frac{1}{2}$ and $L > 0$. Consequently, for $y \in B(0, 1+\varepsilon) \backslash B(0, 1-\varepsilon)$ one has

$$
|f(y)| = |\psi'(|y|)| \ge \mathcal{L} |\psi(|y|) - \psi(1)|^{1/2} = \mathcal{L} |F(y) - F(x)|^{1/2}.
$$

(ii): has been shown in Prop. [4.2.](#page-15-1)

(iii): We conceive $(X_t)_{t\geq 0}$ as complex-valued process by letting

$$
Y_t := X_t^{(1)} + iX_t^{(2)}
$$

and note that (since $(Y_t)_{t\geq 0}$ does not hit 0) there is a continuous adapted process $(\Phi_s)_{s\geq 0}$ satisfying

$$
Y_s = |X_s|e^{i\Phi_s}.
$$

Now on $[t, T_t]$

$$
d\Phi_s = ||X_s| - 1| ds + \frac{1}{|X_s|^2} \frac{1}{s+1} \langle \text{ort}(X_s), dW_s \rangle
$$

so that

$$
d\langle\Psi\rangle_s=\frac{1}{|X_s|^2}\frac{1}{(s+1)^2}\,ds.
$$

 \Box

We thus get that

$$
\int_{t}^{T_t} d\langle \Psi \rangle_s = \int_{t}^{T_t} \frac{1}{|X_s|^2} \frac{1}{(s+1)^2} ds \le \frac{9}{4} \int_{t}^{\infty} \frac{1}{(s+1)^2} ds < \infty
$$

so that $\int_t^{T_t\wedge s}$ 1 $\frac{1}{|X_u|^2} \frac{1}{u+1} \langle \text{ort}(X_u), dW_u \rangle$ converges almost surely as $s \to \infty$, say to $\bar{\Psi}^{(t)}$. Consequently, on $\{T_t = \infty\}$ one has as $s \to \infty$

$$
\Phi_s = \int_t^s ||X_u| - 1| \, du + \bar{\Phi}^{(t)} + o(1),
$$

where $o(1)$ stands for a term converging to zero as $s \to \infty$. By Proposition [4.2](#page-15-1) (iii) and Lemma [4.3,](#page-17-2) $\int_t^s ||X_u| - 1| du \to \infty$ as $s \to \infty$ which shows that $(X_t)_{t \geq 0}$ does not converge on $\{T_t = \infty\}$. The result follows by noticing that $\bigcup_{t \in \mathbb{N}} \{T_t = \infty\}$ is an almost sure set by Proposition [4.2](#page-15-1) (i). \Box

Appendix A. Criterion for staying local

Assume that F satisfies $\lim_{|x|\to\infty} F(x) = \infty$ and consider the set

$$
\mathbb{L}_1 := \Big\{ \sup_{t \geq 0} \int_0^t \big(\langle f(X_s), \alpha_s \rangle + \frac{1}{2} \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) \big) ds < \infty \Big\}.
$$

We will show that $\limsup_{t\to\infty} |X_t| < \infty$, almost surely, on \mathbb{L}_1 . By Itô's formula

$$
dF(X_t) = \langle f(X_t), \alpha_t \rangle dt + \langle f(X_t), \beta_t dW_t \rangle + \frac{1}{2} \text{tr}(\beta_t^{\dagger} H(X_t) \beta_t) dt.
$$

Now, let $n \in \mathbb{N}$ and consider the stopping time

$$
T_n := \inf \Big\{ t \geq 0 : \int_0^t \big(\langle f(X_s), \alpha_s \rangle + \frac{1}{2} \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) \big) ds \geq n \Big\}.
$$

and $(Y_t)_{t>0}$ given by

$$
Y_t := F(X_t) - \int_0^t \left(\langle f(X_s), \alpha_s \rangle + \frac{1}{2} \text{tr}(\beta_s^{\dagger} H(X_s) \beta_s) \right) ds.
$$

Then, $(Y_t^{T_n})_{t\geq 0}$ is a local martingale that is bounded from below and, thus, a supermartingale. With the martingale convergence theorem we get almost sure convergence of $(Y_t^{T_n})_{t\geq 0}$ and, thus, on $\{T_n = \infty\}$ we have that $\sup_{t>0} F(X_t)$ is finite, almost surely. Finally, note that

$$
\mathbb{L}_1 = \bigcup_{n \in \mathbb{N}} \{T_n = \infty\},\
$$

so that $\limsup_{t\to\infty} |X_t| < \infty$, almost surely, on \mathbb{L}_1 . Note that for the choice $(\alpha_t)_{t\geq 0} = (-f(X_t))_{t\geq 0}$ we have

$$
\left\{\sup_{t\geq 0}|H(X_t)| < \infty \text{ and } \int_0^\infty |\beta_s|_F^2 ds < \infty \right\} \subset \mathbb{L}_1
$$

as well as for a compact set $K \subset \mathbb{R}^d$

$$
\left\{ \mathbb{1}_{\{X_t \notin K\}}(-|f(X_t)|^2 + \frac{1}{2}\text{tr}((\beta_t)^{\dagger}H(X_t)\beta_t)) \le 0 \text{ for all large } t \right\}
$$

and
$$
\int_0^{\infty} \mathbb{1}_{\{X_s \in K\}} |\beta_s|_F^2 ds < \infty \right\} \subset \mathbb{L}_1.
$$

Acknowledgement. The authors would like to thank the two anonymous referees for their valuable comments. Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) under Germany's Excellence Strategy EXC 2044–390685587, Mathematics Münster: Dynamics–Geometry–Structure.

REFERENCES

- [AMA05] P.-A. Absil, R. Mahony, and B. Andrews. Convergence of the iterates of descent methods for analytic cost functions. SIAM J. Optim., 16(2):531–547, 2005.
- [Ben99] M. Benaïm. Dynamics of stochastic approximation algorithms. In Seminaire de probabilites XXXIII, pages 1–68. Springer, 1999.
- [CHS87] T.-S. Chiang, C.-R. Hwang, and S. J. Sheu. Diffusion for global optimization in \mathbb{R}^n . SIAM Journal on Control and Optimization, 25(3):737–753, 1987.
- [CMI14] T. Colding and W. Minicozzi II. Lojasiewicz inequalities and applications. In Surveys in Differential Geometry, XIX, pages 63–82. 2014.
- [Cur44] H. B. Curry. The method of steepest descent for non-linear minimization problems. Quarterly of Applied Mathematics, 2(3):258–261, 1944.
- [DK21] S. Dereich and S. Kassing. Convergence of stochastic gradient descent schemes for Lojasiewiczlandscapes. arXiv:2102.09385, 2021.
- [FT21] N. Fournier and C. Tardif. On the simulated annealing in \mathbb{R}^d . J. Funct. Anal., 281(5): Paper No. 109086, 30, 2021.
- [FW12] M. I. Freidlin and A. D. Wentzell. Random Perturbations of Dynamical Systems. Grundlehren der mathematischen Wissenschaften. Springer New York, 2012.
- [Gar85] C. W. Gardiner. Handbook of stochastic methods, volume 3. springer Berlin, 1985.
- [GH86] S. Geman and C.-R. Hwang. Diffusions for global optimization. SIAM Journal on Control and Optimization, 24(5):1031–1043, 1986.
- [Gil00] D. T. Gillespie. The chemical Langevin equation. The Journal of Chemical Physics, 113(1):297–306, 2000.
- [Har12] A. Haraux. Some applications of the Lojasiewicz gradient inequality. Communications on Pure \mathcal{C} Applied Analysis, 11(6):2417, 2012.
- [HKS89] R. A. Holley, S. Kusuoka, and D. W Stroock. Asymptotics of the spectral gap with applications to the theory of simulated annealing. Journal of functional analysis, 83(2):333–347, 1989.
- [JKO98] R. Jordan, D. Kinderlehrer, and F. Otto. The variational formulation of the fokker–planck equation. SIAM journal on mathematical analysis, 29(1):1–17, 1998.
- [Loj63] S. Lojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. Les équations aux dérivées partielles, 117:87–89, 1963.
- [Loj65] S. Lojasiewicz. Ensembles semi-analytiques. Lectures Notes IHES (Bures-sur-Yvette), 1965.
- [Loj84] S. Lojasiewicz. Sur les trajectoires du gradient d'une fonction analytique. (Trajectories of the gradient of an analytic function). Semin. Geom., Univ. Studi Bologna, 1982/1983:115–117, 1984.
- [LTE17] Q. Li, C. Tai, and W. E. Stochastic modified equations and adaptive stochastic gradient algorithms. In Proceedings of the 34th International Conference on Machine Learning, volume 70, pages 2101–2110. PMLR, 06–11 Aug 2017.
- [PdM82] J. Palis, Jr. and W. de Melo. Geometric theory of dynamical systems: an introduction. Springer Science & Business Media, 1982.
- [SSG14] D. Schnoerr, G. Sanguinetti, and R. Grima. The complex chemical Langevin equation. The Journal of Chemical Physics, 141(2):07B606 1, 2014.
- [vK92] N. G. van Kampen. Stochastic processes in physics and chemistry, volume 1. Elsevier, 1992.

STEFFEN DEREICH, INSTITUT FÜR MATHEMATISCHE STOCHASTIK, FACHBEREICH 10: MATHEMATIK UND INformatik, Westfälische Wilhelms-Universität Münster, Orléans-Ring 10, 48149 Münster, Germany Email address: steffen.dereich@wwu.de

SEBASTIAN KASSING, INSTITUT FÜR MATHEMATISCHE STOCHASTIK, FACHBEREICH 10: MATHEMATIK UND IN-FORMATIK, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, ORLÉANS-RING 10, 48149 MÜNSTER, GERMANY Email address: sebastian.kassing@wwu.de