

# Circular swimming motility and disordered hyperuniform state in an algae system

Mingji Huang<sup>a</sup>, Wensi Hu<sup>b</sup>, Siyuan Yang<sup>a</sup>, Quan-Xing Liu<sup>b,c</sup>, and H. P. Zhang<sup>a,d,1</sup>

<sup>a</sup>School of Physics and Astronomy and Institute of Natural Sciences, Shanghai Jiao Tong University, Shanghai 200240, China; <sup>b</sup>State Key Laboratory of Estuarine and Coastal Research, School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China; <sup>c</sup>Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration & Center for Global Change and Ecological Forecasting, School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China; <sup>d</sup>Collaborative Innovation Center of Advanced Microstructures, Nanjing 210093, China

This manuscript was compiled on June 15, 2021

**Active matter comprises individually driven units that convert locally stored energy into mechanical motion. Interactions between driven units lead to a variety of non-equilibrium collective phenomena in active matter. One of such phenomena is anomalously large density fluctuations, which have been observed in both experiments and theories. Here we show that, on the contrary, density fluctuations in active matter can also be greatly suppressed. Our experiments are carried out with marine algae (*Effrenium voratum*) which swim in circles at the air-liquid interfaces with two different eukaryotic flagella. Cell swimming generates fluid flow which leads to effective repulsions between cells in the far field. Long-range nature of such repulsive interactions suppresses density fluctuations and generates disordered hyperuniform states under a wide range of density conditions. Emergence of hyperuniformity and associated scaling exponent are quantitatively reproduced in a numerical model whose main ingredients are effective hydrodynamic interactions and uncorrelated random cell motion. Our results demonstrate a new form of collective state in active matter and suggest the possibility to use hydrodynamic flow for self-assembly in active matter.**

hyperuniformity | circular microswimmer | hydrodynamic interaction | transverse flagellum | algae

**A**ctive matter exists over a wide range of spatial and temporal scales (1–6) from animal groups (7, 8) to robot swarms (9–11), to cell colonies and tissues (12–16), to cytoskeletal extracts (17–20), and man-made microswimmers (21–25). Constituent particles in active matter systems are driven out of thermal equilibrium at the individual level; they interact to develop a wealth of intriguing collective phenomena, including clustering (13, 22, 24), flocking (11, 26), swarming (12, 13), spontaneous flow (14, 20), and giant density fluctuations (10, 11). Many of these observed phenomena have been successfully described by particle-based or continuum models (1–6), which highlight the important roles of both individual motility and inter-particle interactions in determining system dynamics.

Current active matter research focuses primarily on linearly swimming particles which have a symmetric body and self-propel along one of the symmetry axes. However, a perfect alignment between the propulsion direction and body axis is rarely found in reality. Deviation from such a perfect alignment leads to a persistent curvature in the microswimmer trajectories; examples of such circle microswimmers include anisotropic artificial micromotors (27, 28), self-propelled nematic droplets (29, 30), magnetotactic bacteria and Janus particles in rotating external fields (31, 32), Janus particle in viscoelastic medium (33), sperm and bacteria near interfaces (34, 35). Chiral motility of circle microswimmers, as predicted

by theoretical and numerical investigations, can lead to a range of interesting collective phenomena in circular microswimmers, including vortex structures (36, 37), localization in traps (38), enhanced flocking (39), and hyperuniform states (40). However, experimental verifications of these predictions are limited (32, 35), a situation mainly due to the scarcity of suitable experimental systems.

Here we address this challenge by investigating marine algae *Effrenium voratum* (*E. voratum*) (41, 42). At air-liquid interfaces, *E. voratum* cells swim in circles via two eukaryotic flagella: a transverse flagellum encircling the cellular anteroposterior axis and a longitudinal one running posteriorly. Over a wide range of densities, circling *E. voratum* cells self-organize into disordered hyperuniform states with suppressed density fluctuations at large lengthscales. Hyperuniformity (43, 44) has been considered as a new form of material order which leads to novel functionalities (45–49); it has been observed in many systems, including avian photoreceptor patterns (50), amorphous ices (51), amorphous silica (52), ultracold atoms (53), soft matter systems (54–61), and stochastic models (62–64). Our work demonstrates the existence of hyperuniformity in active matter and shows that hydrodynamic interactions can be used to construct hyperuniform states.


## Significance Statement

Disordered hyperuniform materials suppress large scale density fluctuations like crystals and remain locally isotropic as liquids. These materials possess unique properties, such as isotropic photonic gap, and have attracted the attention of scientists from many disciplines. Here we show that circularly swimming marine algae can robustly self-organize into disordered hyperuniform states through long-range hydrodynamic interactions at air-liquid interfaces. Important properties measured from hyperuniform states can be quantitatively reproduced by a numerical model whose main parameters are obtained experimentally. Our work clearly demonstrates the possibility to create disordered hyperuniform states via hydrodynamic interactions and highlights the importance of such interactions in active matter systems.

M.H. and H.P.Z. designed research; M.H., W.H. and S.Y. performed research; M.H., W.H. and H.P.Z. analyzed data; and M.H., W.H., S.Y., Q.X.L. and H.P.Z. wrote the paper.

The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: hepeng\_zhang@sjtu.edu.cn



**Fig. 1.** Cell motility and flagellar dynamics. (A) Scanning electron micrograph of *E. voratum* (courtesy of Sung Yeon Lee and Todd LaJeunesse (41)). (B) 3D trajectory of a cell approaching an air-liquid interface (at  $z = 0$ ) from the bulk. Time history of cell coordinates in the laboratory frame ( $x$ ,  $y$ , and  $z$ ) is shown in the inset. (C) Circular trajectory plotted on an optical image of a cell at the interface. Undulations in the trajectory reflect beating phases of the longitudinal flagellum. A cell body frame ( $XY$ ) is defined with  $X$  being the direction of body axis. (D) Waveform of the longitudinal flagellum over a period. (E) Kymograph to show transverse flagellum dynamics. Intensity profiles of cell image are extracted along the white line (fixed in the cell body frame) in (C). Scale bars in (A) and (C) are  $5 \mu\text{m}$ . See supporting videos S1 and S2 for cell motion and flagellar dynamics.

## Results


*E. voratum* belongs to family Symbiodiniaceae (41, 42). Dinoflagellates in this family are among the most abundant eukaryotic microbes found in coral reef ecosystems; they convert sunlight and carbon dioxide into organic carbon and oxygen to fuel coral growth and calcification (65). Cell motility of *E. voratum* has been shown to be important for algal-invertebrate partnerships (66), though quantitative understanding of cell motility is still lacking.

**Circular cell motion and associated flow field.** We observe the cells at the air-liquid interface on an up-right microscope. As shown in Fig. 1A and C, *E. voratum* cells have an approximately elliptic shape and are equipped with both longitudinal and transverse flagella for motility. Away from interfaces, cells swim in helical trajectories which are typical for motile algae, see Fig. 1B and Movie S1. However, when cells get close to an air-liquid interface, they adhere to the interface and start to move in circles; all cells move in the counter-clockwise direction when viewed from the air side of the air-liquid interface. In Fig. S3, we show that cells also adhere to liquid-solid interfaces (67) and estimate the gap between cell and interface to be  $0.3 \mu\text{m}$ . A typical counter-clockwise circular trajectory

at air-liquid interface is plotted on an optical image of a cell in Fig. 1C, where we define the long symmetric axis as the cell body direction,  $X$  coordinate. Typical cell circling radius, translation and angular velocities are  $\langle a \rangle = 11.6 \mu\text{m}$ ,  $\langle v_c \rangle = 180 \mu\text{m/s}$  and  $\langle \omega \rangle = 16.2 \text{ rad/s}$ , respectively. These motility characteristics depend weakly on cell density and their variations are quantified in Fig. S2B.

As shown in Movie S2, the longitudinal flagellum produces a planar wave in a plane parallel to the air-liquid interface. Waveforms of the longitudinal flagellum in a period (15 ms) are shown in Fig. 1D. The transverse flagellum sits in a groove (68–71), as shown in Fig. 1A and Movie S2; we cannot separate the flagellum’s image from that of the cell body to extract all information about the flagellum’s waveform. Instead, we extract intensity profiles from optical images along a line (fixed in the cell body frame) cutting through the transverse flagellum, cf. the white line in Fig. 1C, and construct a kymograph from the extracted line profiles. As shown Fig. 1E, the kymograph shows a wave propagation to the negative  $Y$  direction with a period about 23 ms and a wave speed of  $124 \mu\text{m/s}$ .


The swimming cell generates fluid flow (denoted by  $\mathbf{v}$ ) in space. We measure the two flow components in the plane of



**Fig. 2.** Mean in-plane flow field,  $\mathbf{v}_{\parallel}$ , measured in experiment (A) and regularized Stokeslet model (B). Flow speed ( $|\mathbf{v}_{\parallel}|$ ) is represented by color and arrows show local flow direction. Cell symmetry axis is oriented along the X axis and an angle from X direction is defined as  $\phi$  in (B). (C) An optimal slip flow pattern obtained from our numerical procedure (see Text for details). The air-liquid interface is shown by a green line; the gap between cell body and interface is set to be  $0.3 \mu\text{m}$ . (D) Angular dependence of the magnitude of the in-plane velocity at two radii,  $30 \mu\text{m}$  and  $40 \mu\text{m}$  dashed lines in (A) and (B), from experiments (symbols) and numerics (lines). Angular dependence of far-field flow speed (at the radius of  $2000 \mu\text{m}$ , computed from regularized Stokeslet calculation) is shown in the inset; the far-field flow is dominated by a pair of orthogonal pusher-puller dipoles (see SI for detail). Experimental data are measured from tracer motion around a cell with a swimming speed  $v_c = 201 \mu\text{m/s}$  and radius  $a = 11.5 \mu\text{m}$ , cf. Movie S3.

cell motion by tracking tracer particles, cf. Movie S3. The measured fields at different times are then averaged in the cell body (XY) frame. A typical averaged field,  $\mathbf{v}_{\parallel} = v_X \hat{X} + v_Y \hat{Y}$ , is plotted in Fig. 2A. Though bearing some similarities with that of a source dipole, the field does not show any obvious (left-right or fore-aft) symmetries, which are frequently found in the cases of straight swimmers (72). To reproduce such complex flow, we use a regularized Stokeslet model (73–75). In the model, the cell body is represented by a sphere with a radius of  $5 \mu\text{m}$ , which is driven by both longitudinal and transverse flagella. As shown in Fig. 1C–D, the longitudinal flagellum has a conventional structure; its planar waveform can be readily quantified and is faithfully represented in the model. However, the transverse flagellum is hidden in the groove and difficult to observe; its structure and driving mechanism are still being debated (68–71). We can measure the wave period and speed from Fig. 1E, but the exact waveform of the transverse flagellum is unknown. Due to this lack of

information for the transverse flagellum, we represent it in our model by a slip flow pattern on the cell surface,  $\mathbf{v}_b$ . For a given slip pattern, we use the regularized Stokeslet method to compute cell (translation and rotation) velocities and a flow field corresponding to the experimental result in Fig. 2A,  $\mathbf{v}_{\parallel}$ . We vary the slip flow pattern,  $\mathbf{v}_b$ , and search for a pattern that optimizes the match between numerical and experimental results of cell velocities and flow fields. A resultant flow pattern,  $\mathbf{v}_b$ , from such a procedure is shown in Fig. 2C and maximal slip ( $500 \mu\text{m/s}$ ) occurs in the bright yellow region, which approximately corresponds to the location of the transverse flagellum. With this slip pattern, we numerically generate the in-plane flow field around a cell, shown in Fig. 2B; two angular profiles of the in-plane flow speed ( $|\mathbf{v}_{\parallel}|$ ) in Fig. 2D show good agreement between experiment and Stokeslet results. See SI for detailed discussions on the above procedure and results obtained in another cell, cf. Fig. S6.



**Fig. 3.** (A) Streak image of cell motion. Raw images are obtained at cell density  $178 \text{ mm}^{-2}$  and averaged over 10 sec to produce the streak image. Scale bar is  $200 \mu\text{m}$ . (B) Instantaneous cell coordinates (gray line,  $\mathbf{r}(t)$ ) are averaged over a sliding window of 2 s to highlight the non-circular motion (green line,  $\bar{\mathbf{r}}(t)$ ). (C) Temporal history of instantaneous (gray) and window-averaged (green) coordinates. (D) Probability distribution functions of window-averaged cell displacements (squared and normalized by time separation,  $s \equiv \Delta\bar{r}^2/4\Delta t$ ) from experiment (circles) and model (squares) for four time separations. The experiment and model results are in agreement; for presentation clarity, we rescale model results in (D) by a factor of 100.

**Non-circular cell motion.** Though circular motion is most frequently observed, cells also exhibit rare non-circular motion. To demonstrate that, we show a streak image of cell motion in Fig. 3A. While majority of cells move in circles and appear as white "donuts", the image also contains rare long streaks, corresponding to rapid translational motion of cells. The transition to non-circular motion is likely related to changes in the longitudinal flagellar dynamics, as depicted in Fig. S4 (76, 77). We use a procedure with empirically chosen parameters to identify non-circular motion from instantaneous cell positions (see Fig. S5 and related discussions in SI for details). This procedure shows that durations of non-circular motion are usually less than a few seconds and that the occurring rate of non-circular motion is approximately  $10^{-3} \text{ s}^{-1} \text{ cell}^{-1}$ .


To quantify non-circular motion, we first average instantaneous cell coordinates,  $\mathbf{r}(t)$ , over a sliding window of 2 seconds ( $\sim 5.5$  circling period). As shown in Fig. 3B-C, circular motion is smoothed out in the window-averaged (green) trajectory, denoted as  $\bar{\mathbf{r}}(t)$ . From the window-averaged trajectories, we measure probability distributions of cell displacements for different time separations  $\Delta t$ . After squared displacements  $\Delta\bar{r}^2 \equiv (\bar{\mathbf{r}}(t + \Delta t) - \bar{\mathbf{r}}(t))^2$  are normalized by the time separation  $\Delta t$ , all distributions of  $s \equiv \Delta\bar{r}^2/4\Delta t$  collapse onto a single curve and exhibit a power-law scaling for large  $s$ , cf. Fig. 3D. Similar probability distributions have been found under different density conditions, cf. Fig. S9.

**Experimental observations of Hyperuniform states.** We next investigate collective states of interacting cells at the air-liquid interface. Cells in the bulk suspension swim to adhere at the interface in the first few minutes of experiments and this leads to a random initial distribution of cells at the interface, with a cell density  $\rho$ . Then, cells at the interface slowly self-organize into a steady state after a relaxation period about several thousand seconds, cf. Fig. S11. We measure static and dynamic properties of these steady states. Fig. 4A-B show typical instantaneous configurations from two experiments, see also Movie S4-5. Though no obvious order can be detected in these configurations, spatial distribution of cells appears to be quite uniform at large length scales. Quantitatively, from instantaneous cell positions  $\mathbf{r}^{(j)}(t)$ , we compute density fluctuations for square interrogation windows of different sizes

$L$ . For a given window size, we find that density fluctuations follow a Gaussian distribution, as shown in the inset of Fig. 4C. Variances of density fluctuations are plotted against the window size in Fig. 4C; data follow the scaling determined by the central limit theorem  $\langle \delta\rho^2 \rangle \sim L^{-2}$  at small scales and decay faster with the window size at large scales:  $\langle \delta\rho^2 \rangle \sim L^{-2.6}$ . Similar physics is also reflected by the static structure factors,  $S(k) = \left\langle \frac{1}{N} \left| \sum_{j=1}^N \exp(-i\mathbf{k} \cdot \mathbf{r}^{(j)}) \right|^2 \right\rangle$ , where  $N$  is the total number of observed cells. In Fig. 4D,  $S(k)$  shows a liquid-like peak in large  $k$  region and a scaling  $S(k) \sim k^{0.6}$  for small  $k$ . Lengthscale corresponding to liquid-like peaks matches approximately to the transition length between  $\langle \delta\rho^2 \rangle \sim L^{-2}$  and  $\langle \delta\rho^2 \rangle \sim L^{-2.6}$  scalings. The same scalings,  $\langle \delta\rho^2 \rangle \sim L^{-2.6}$  and  $S(k) \sim k^{0.6}$ , are found for different cell densities. Increasing the cell density leads to a decrease of  $S(k)$  for small  $k$  and shifts the liquid-like peak to larger  $k$ , as shown by Fig. 4C-D. To check the robustness of observed hyperuniformity, we also use window-averaged cell position,  $\bar{\mathbf{r}}(t)$  defined in Fig. 3B, to compute density fluctuations and obtain similar results, shown in Fig. S11.

Beyond static structures, we also investigate the system's dynamic properties. To explore the possibility of local synchronization, spatial correlation functions of circling phase and velocity are computed; results in Fig. S10C-D show that instantaneous cell motions are not spatially correlated, suggesting a weak interaction between cells. This is confirmed by Fig. 2 which shows that the flow velocity at the nearest cells (at a distance  $\sim 30 - 100 \mu\text{m}$ ) is much smaller than cell swimming speed. Therefore, the cell-cell interaction is not strong enough to significantly affect the instantaneous cell motion. However, as shown below, this weak hydrodynamic interaction can modulate cell positions over a long time and its long-range nature leads to the formation of hyperuniform states.

**Particle-based model for hyperuniformity.** We construct a numerical model to illustrate the origin of observed hyperuniformity. Dynamics in our experiments evolves over a time scale that is much longer than cell circling periods ( $\sim 0.4$  s). This separation of time scales allows us to build a temporally



**Fig. 4.** (A-B) Cell configurations at two densities (41 and 369 mm<sup>-2</sup>). Typical output of simulation is shown in the inset. Scale bar is 500  $\mu\text{m}$ . (C) Cell density fluctuations plotted against interrogation window size and (D) static structure factors for four densities. Experimental and numerical results are shown by light symbols and dark lines, respectively. The maximal measurement lengthscale is 800  $\mu\text{m}$  in experiments and 10000  $\mu\text{m}$  in simulations. The inset in (C) shows distributions of normalized density fluctuations measured at density 198 mm<sup>-2</sup>; color symbols are experimental data with different window sizes ( $L = 200$  to 800  $\mu\text{m}$ ) and the thick line represent a normal distribution. The inset in (D) shows a two-dimensional static structure factor measured at density 198 mm<sup>-2</sup>.


coarse-grained (over a few circling periods) model to capture the emergence of hyperuniformity without fully resolving fast circular cell motion (78). Therefore, particle coordinates in our model represent window-averaged cell positions in experiments ( $\bar{\mathbf{r}}(t)$  in Fig. 3B) and particles interact through period-averaged flow field. Beyond flow advection, our model also includes stochastic non-circular particle motion and uses the following equation to determine the displacement of the  $i$ th particle at time  $n\tau$  during a time step  $\tau$ :

$$\begin{aligned} \bar{\mathbf{r}}^{(i)}((n+1)\tau) - \bar{\mathbf{r}}^{(i)}(n\tau) = & \\ \sum_{j \neq i} \bar{\mathbf{V}}\left(\bar{\mathbf{r}}^{(i)}(n\tau) - \bar{\mathbf{r}}^{(j)}(n\tau); v_c^{(j)}\right) \tau & + \boldsymbol{\eta}\left(D^{(i)}\right) \delta(\text{mod}(n,p), s^{(i)}) \end{aligned} \quad [1]$$

The period-averaged flow field in Eq. [1],  $\bar{\mathbf{V}}(\mathbf{R}; v_c)$ , is calculated by the regularized Stokeslet method for a cell circling with a radius  $a = 10 \mu\text{m}$  and velocity  $v_c$ . As shown in Fig. 5A,  $\bar{\mathbf{V}}(\mathbf{R}; v_c)$  has an outgoing component in the far field; see

Materials and Methods for detailed discussions on the period-averaged flow field. In our model, parameter  $v_c^{(i)}$  is sampled from an experimentally determined distribution of cell velocity in Fig. S2.

The second term on the right-hand-side of Eq.[1] represents stochastic jumps. The Kronecker delta function  $\delta()$  dictates that adjacent random jumps for a given particle are temporally separated by  $p$  (an integer constant) time steps, defining a waiting time  $T = p\tau$ . Specifically, the  $i$ th particle jumps at time  $n\tau$  if  $\text{mod}(n,p) = s^{(i)}$ ;  $\text{mod}()$  represents modulo operation and  $s^{(i)}$  is an integer constant between 0 and  $p-1$ , randomly assigned to all particles. Components of jumping displacements  $\boldsymbol{\eta}(D^{(i)})$  are independently drawn from a normal distribution with a standard deviation  $\sqrt{2D^{(i)}T}$ , as shown in Fig. 5B. Parameter  $D^{(i)}$  is the diffusivity for the  $i$ th particle and drawn from a Pareto distribution with a cut-off value  $D_0$



**Fig. 5.** (A) Radial and tangential components of period-averaged flow field around a swimming cell. Flow components are defined in the inset, showing a cell circling with a radius  $a$  and a velocity  $v_c$ . Flow strength is capped at  $R = 30\mu\text{m}$ , which approximately corresponds to the minimal distance between circular centers of two cells in experiments. Black dash lines show the asymptotic behavior in the far-field (see text). (B) Normal distribution of jumping displacements  $\eta(D^{(i)})$  (cf. Eq.[1]) and Pareto distribution of particle diffusivity ( $D^{(i)}$ ) (cf. Eq.[2]). Three stochastic trajectories with different diffusivity are shown in the inset. (C) Effects of four model parameters on structure factors. A default set of parameters is used unless specified:  $\rho = 198 \text{ mm}^{-2}$ ,  $T = 400 \text{ s}$ ,  $D_0 = 0.95 \mu\text{m}^2/\text{s}$  and  $\beta = 2$ .

and a power index  $\beta$  :

$$f(D; D_0, \beta) = \begin{cases} (\beta - 1) \frac{D_0^{\beta-1}}{D^\beta} & D \geq D_0 \\ 0 & D < D_0 \end{cases}, \quad [2]$$

where  $\beta = 2$  unless stated otherwise.

In a 2D periodic domain of size  $L_{\text{max}} = 20 \text{ mm}$ , we simulate  $\rho L_{\text{max}}^2$  particles following Eq. [1]. For a given experimental condition, the cut-off diffusivity  $D_0$  and the waiting time  $T$  are varied to match simulation results to experiments. The obtained values for  $D_0$  ( $\sim 1 \mu\text{m}^2/\text{s}$ ) and  $T$  ( $\sim 500 \text{ s}$ ) are listed for different cell densities in Table SIII.

We measure density fluctuations after randomly initialized particles evolve to a steady state. As shown in Fig. 4C-D, our simulations can generate hyperuniform states and quantitatively reproduce measured density fluctuation  $\delta\rho^2$  and structure factor  $S(k)$ , highlighting scaling laws  $\delta\rho^2 \sim L^{-2.6}$  and  $S(k) \sim k^{0.6}$  for hyperuniformity. Distribution functions of cell displacements are also well reproduced in Fig. 3D. The Pareto distribution of  $D^{(i)}$  with a power index  $\beta = 2$  leads to the observed power-law distribution for large displacements.

We systematically study effects of model parameters; results are shown in Fig. 5C. An increase of particle density  $\rho$  shifts the liquid-like peak to higher  $k$  and leads to a decrease of density fluctuation for small  $k$ , which mirrors experimental results in Fig. 4D. Decreasing the waiting time  $T$  introduces more fluctuations into the system and lead to an increase in  $S(k)$  for small  $k$ , cf. the second panel in Fig. 5C. Smaller cut-off diffusivity  $D_0$  allows fewer particles with large  $D^{(i)}$ ; this leads to fewer stochastic jumps with large displacement, less fluctuations in large scales and smaller  $S(k)$  values. As shown in the fourth panel in Fig. 5C, the power index of the Pareto distribution,  $\beta$  in Eq. [2], can change power-law scaling of  $S(k)$  at small  $k$ . A larger  $\beta$  means fewer particles with large diffusivity; this leads the system to approach the strong hyperuniformity limit (44) and a larger exponent in  $S(k)$  is observed. Fig. 5C also demonstrates that variations in  $\rho$ ,  $T$  and  $D_0$  have relatively weak effects on the scaling exponent for small  $k$  in our observation window (79).

## Discussion

In summary, we have studied individual motility and collective dynamics in marine algae *E. voratum*. Cells swim in circles at the air-liquid interface with a longitudinal flagellum and a transverse one. Combining experimental measurements and the regularized Stokeslet method, we showed that period-averaged flow generated by cells has a long-ranged and outgoing radial component that disperses cells uniformly and leads to a disordered hyperuniform state. Stochastic cell motion with a power-law displacement distribution (79) also plays an important role in determining the properties of density fluctuations.

Regularized Stokeslet results in Fig. 2 can be used to clarify the current confusion on the contributions of two flagella to propulsion in *E. voratum* (68–70). For that, we measure the stalled force and torque with both flagella or only one of them functioning; as shown in SI, the longitudinal flagellum provides about 30% of the total torque and less than 10% of the force. The regularized Stokeslet method is also used to calculate 3D period-average flow and show that self-generated flow can lead to directed nutrient/particle transport around cells, cf. Fig. S8. These results provide novel insight into the ecological function and evolutionary traits of two flagella in this ecologically important dinoflagellate (41, 42, 65, 66).

Previous studies have shown that hydrodynamic interactions lead to interesting self-organization (58). For example, bacteria, when oriented perpendicularly to an interface, can generate inward flow that assembles bacterial cells into compact crystals at the interface (80, 81). In contrast, circling *E. voratum* cells in current work generate repulsive interactions. This can be understood from the far-field instantaneous flow in the plane cell reside. As shown in the inset of Fig. 2D and analysis in SI, out-going flow in the far-field is stronger than its in-coming counterpart. This leads to a period-average repulsive interaction between cells in Fig. 5A. The same hydrodynamic mechanism may underlie the formation of sperm vortex in (35). Our work suggests a novel mechanism of using average along circular trajectories to generate isotropic hydrodynamic interactions between force-free microswimmers. Such isotropic hydrodynamic interactions in chiral active matter are different from anisotropic dipolar interactions in conventional systems with linearly swimming particles and may produce new collective phenomena.

Fig. 4 shows that hyperuniformity is observed under different cell concentration conditions with a similar scaling exponent. Such a density independence has also been observed in other hyperuniform systems with long-range interactions. In a one-component plasma, particles with the same electrostatic charge interact with repulsive Coulomb potential which imposes energy penalty on density fluctuations and leads to hyperuniformity under all particle density conditions (44). In sedimentation system of irregular objects (58), falling objects interact via long-range hydrodynamic (force monopole) flow and objects' irregular shape leads to an anisotropic response to the local flow. A combination of the long-range interaction and anisotropic response in this system produces hyperuniformity with a density-independent exponent.

Hyperuniformity has been observed in many systems exhibiting absorbing-state transition (40, 57, 62–64, 82). For example, a recent numerical work simulated a system of active particles which self-propel in circles (alike *E. voratum* cells

here) but interact via short-range repulsive forces; hyperuniformity in this system was only observed in high-density active states (40). In contrast, *E. voratum* cells self-organize into hyperuniform states under all densities, thanks to the long-range nature of hydrodynamic interactions (44, 48). Phoretic interactions in synthetic active matter system are also known to be long-ranged; both attractive and repulsive phoretic interactions have been realized (24, 83). These long-range interactions may provide a promising avenue to generate novel hyperuniform materials with active matter.

## Materials and Methods

**Cell growth and imaging procedure.** Spices *E. voratum* cells are cultured in artificial seawater with F/2 medium in a 100 ml flask which is placed in an incubator (INFORS HT Multitron pro, Switzerland) at 20°C. We use a daily light cycle which consists of 12 hours of cool light with an intensity of 2,000 Lux and 12 hours in dark. The algal cells in the experiments are in an exponential phase after 14-day growth and observed a few hours after the light period starts, when cells showed excellent motility (see Fig. S1, (41, 42)). During experiments, cell culture is placed in a disk-shape chamber fabricated by cover glass and plastic gasket (8-mm in diameter). Cells gather and form a mono-layer at the air-liquid interface. Cell motion in the central region of the sample is recorded by a high-speed camera (Basler acA2040-180km, 4 M pixel resolution) mounted on an up-right microscope (Nikon Ni-U) with a 4× or 40× magnification objective; the acquisition rate varies from 50 to 850 frame/s. To measure fluid flow, milk (Deluxe Milk, Mengniu) is added to provide passive flow tracers (diameter 1~2 μm). Holographic imaging technique is used to measure 3D cell motion.

**Particle-based model.** To obtain period-averaged flow field in Fig. 5A, we use the regularized Stokeslet results from Fig. 2 to compute instantaneous flow fields around a cell and average computed fields in the lab frame over cell positions in a circling period. The flow field is capped at  $R = 30 \mu\text{m}$ , which approximately corresponds to the minimal distance between circular center of two cells in experiments. In simulation, we interpolate data in Fig. 5A to find two flow components at any separation; asymptotic expressions for flow components (Eq. [S21]) with a cap at  $R = 30 \mu\text{m}$  can also reproduce experimental data, cf. Fig. S12 for details. To reduce the computing load, interacting flow is assumed to be zero beyond a cutoff length of 20000 μm, which is twice the maximal length of computed density fluctuations ( $\delta\rho^2$ ) and structure factor  $S(k)$  in simulations.

In our model, time step  $\tau$  is set to be 10 sec, during which typical particle displacements ( $\sim 1 \mu\text{m}$ ) are much smaller than typical particle separations ( $\sim 50 \mu\text{m}$ ). Analysis of experiments show that durations of random non-circular motion are less than 10 sec, cf. Fig. S5D; such events occur within a single time-step in simulation.

Particle positions in simulations represent temporally averaged cell positions  $\bar{\mathbf{r}}^{(i)}(t)$ ; we add a random circling phase (cf. Fig. S10C) to each particle position to obtain instantaneous cell positions,  $\mathbf{r}^{(i)}(t)$ , which are used to compute density fluctuations.

**ACKNOWLEDGMENTS.** We acknowledge financial supports from National Natural Science Foundation of China Grants (12074243, 11774222 and 32071609) and from the Program for Professor of Special Appointment at Shanghai Institutions of Higher Learning (Grant GZ2016004). We thank the Student Innovation Center at Shanghai Jiao Tong University for support.

1. E. Lauga and T. R. Powers. The hydrodynamics of swimming microorganisms. *Rep Prog Phys*, 72(9):096601, September 2009. .
2. S. Ramaswamy. The mechanics and statistics of active matter. *Annual Review of Condensed Matter Physics*, 1:323–345, 2010. .
3. T. Vicsek and A. Zafeiris. Collective motion. *Phys Rep*, 517(3-4):71–140, August 2012. .
4. M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, Madan Rao, and R. Aditi Simha. Hydrodynamics of soft active matter. *Rev Mod Phys*, 85:1143–1189, Jul 2013. .

5. Clemens Bechinger, Roberto Di Leonardo, Hartmut Löwen, Charles Reichhardt, and Giovanni Volpe, Giorgio and. Active particles in complex and crowded environments. *Rev Mod Phys*, 88:045006, Nov 2016. .
6. G. Gompper, R. G. Winkler, T. Speck, A. Solon, C. Nardini, F. Peruani, H. Lowen, R. Golestanian, U. B. Kaupp, L. Alvarez, T. Kiorbø, E. Lauga, W. C. K. Poon, A. DeSimone, S. Muiños-Landin, A. Fischer, N. A. Soker, F. Cichos, R. Kapral, P. Gaspard, M. Ripoll, F. Sagues, A. Doostmohammadi, J. M. Yeomans, I. S. Aranson, C. Bechinger, H. Stark, C. K. Hemelrijk, F. J. Nedelec, T. Sarkar, T. Aryakama, M. Lacroix, G. Duclos, V. Yashunsky, P. Silberzan, M. Arroyo, and S. Kale. The 2020 motile active matter roadmap. *Journal of Physics-condensed Matter*, 32(19):193001, May 2020. .
7. A. Cavagna and I. Giardina. Bird flocks as condensed matter. *Annual Review of Condensed Matter Physics*, Vol 5, 5:183–207, 2014. .
8. A. Berdahl, C. J. Torney, C. C. Ioannou, J. J. Faria, and I. D. Couzin. Emergent sensing of complex environments by mobile animal groups. *Science*, 339(6119):574–576, February 2013. .
9. Michael Rubenstein, Alejandro Cornejo, and Radhika Nagpal. Programmable self-assembly in a thousand-robot swarm. *Science*, 345(6198):795–799, August 2014. .
10. V. Narayan, S. Ramaswamy, and N. Menon. Long-lived giant number fluctuations in a swarming granular nematic. *Science*, 317(5834):105–108, July 2007. .
11. J. Deseigne, O. Dauchot, and H. Chate. Collective motion of vibrated polar disks. *Phys Rev Lett*, 105(9):098001, August 2010. .
12. C. Dombrowski, L. Cisneros, S. Chatkaew, R. E. Goldstein, and J. O. Kessler. Self-concentration and large-scale coherence in bacterial dynamics. *Phys Rev Lett*, 93(9):098103, August 2004. .
13. H. P. Zhang, A. Be'er, E. L. Florin, and H. L. Swinney. Collective motion and density fluctuations in bacterial colonies. *Proc. Natl. Acad. Sci. U. S. A.*, 107(31):13626–13630, August 2010. .
14. E. Lushi, H. Wioland, and R. E. Goldstein. Fluid flows created by swimming bacteria drive self-organization in confined suspensions. *Proc Natl Acad Sci U S A*, 111(27):9733–9738, July 2014. .
15. Chong Chen, Song Liu, Xia-qing Shi, Hugues Chate, and Yilin Wu. Weak synchronization and large-scale collective oscillation in dense bacterial suspensions. *Nature*, 542(7640):210–214, February 2017. .
16. Xingbo Yang, Dapeng Bi, Michael Czajkowski, Matthias Merkel, M. Lisa Manning, and M. Cristina Marchetti. Correlating cell shape and cellular stress in motile confluent tissues. *Proc Natl Acad Sci U S A*, 114(48):12663–12668, nov 2017. .
17. V. Schaller, C. Weber, C. Semmrich, E. Frey, and A. R. Bausch. Polar patterns of driven filaments. *Nature*, 467(7311):73–77, September 2010. .
18. Y. Sumino, K. H. Nagai, Y. Shitaka, D. Tanaka, K. Yoshikawa, H. Chate, and K. Oiwa. Large-scale vortex lattice emerging from collectively moving microtubules. *Nature*, 483(7390):448–452, March 2012. .
19. P. Guillamat, J. Ignés-Mullol, and F. Sagues. Control of active liquid crystals with a magnetic field. *Proc Natl Acad Sci U S A*, 113(20):5498–5502, May 2016. .
20. Kun-Ta Wu, Jean Bernard Hishamunda, Daniel T. N. Chen, Stephen J. DeCamp, Ya-Wen Chang, Alberto Fernandez-Nieves, Seth Fraden, and Zvonimir Dogic. Active matter transition from turbulent to coherent flows in confined three-dimensional active fluids. *Science*, 355(6331):1284–+, March 2017. .
21. H. R. Jiang, N. Yoshinaga, and M. Sano. Active motion of a janus particle by self-thermophoresis in a defocused laser beam. *Phys Rev Lett*, 105(26):268302, December 2010. .
22. I. Theurkauff, C. Cottin-Bizonne, J. Palacci, C. Ybert, and L. Bocquet. Dynamic clustering in active colloidal suspensions with chemical signaling. *Phys Rev Lett*, 108(26):268303, June 2012. .
23. Antoine Bricard, Jean-Baptiste Caussin, Nicolas Desreumaux, Olivier Dauchot, and Denis Bartolo. Emergence of macroscopic directed motion in populations of motile colloids. *Nature*, 503(7474):95–98, November 2013. ISSN 0028-0836.
24. J. Palacci, S. Sacanna, A. P. Steinberg, D. J. Pine, and P. M. Chaikin. Living crystals of light-activated colloidal surfers. *Science*, 339(6122):936–940, February 2013. .
25. Jing Yan, Ming Han, Jie Zhang, Cong Xu, Erik Luijten, and Steve Granick. Reconfiguring active particles by electrostatic imbalance. *Nat Mater*, 15:1095, July 2016. .
26. A. Cavagna, A. Cimarrelli, I. Giardina, G. Parisi, R. Santagati, F. Stefanini, and M. Viale. Scale-free correlations in starling flocks. *Proc. Natl. Acad. Sci. U. S. A.*, 107(26):11865–11870, June 2010. .
27. Felix Kummel, Borge ten Hagen, Raphael Wittkowski, Ivo Buttinoni, Ralf Eichhorn, Giovanni Volpe, Hartmut Lowen, and Clemens Bechinger. Circular motion of asymmetric self-propelling particles. *Phys Rev Lett*, 110:198302, May 2013. .
28. Sinan Du, Huaqiang Wang, Chao Zhou, Wei Wang, and Zexin Zhang. Motor and rotor in one: Light-active zno/au twinned rods of tunable motion modes. *J Am Chem Soc*, 142(5):2213–2217, February 2020. .
29. C. Kruger, G. Klos, C. Bahr, and C. C. Maass. Curling liquid crystal microswimmers: A cascade of spontaneous symmetry breaking. *Phys Rev Lett*, 117(4):048003, July 2016. .
30. F. Lancia, T. Yamamoto, A. Ryabchun, T. Yamaguchi, M. Sano, and N. Katsonis. Reorientation behavior in the helical motility of light-responsive spiral droplets. *Nat Commun*, 10:5238, November 2019. .
31. K. Erglis, Q. Wen, V. Ose, A. Zeltins, A. Sharipo, P. A. Janmey, and A. Cebers. Dynamics of magnetotactic bacteria in a rotating magnetic field. *Biophys J*, 93(4):1402–1412, August 2007. .
32. M. Han, J. Yan, S. Granick, and E. Luijten. Effective temperature concept evaluated in an active colloid mixture. *Proc Natl Acad Sci U S A*, 114(29):7513–7518, July 2017. .
33. N. Narinder, Clemens Bechinger, and Juan Ruben Gomez-Solano. Memory-induced transition from a persistent random walk to circular motion for achiral microswimmers. *Phys Rev Lett*, 121(7):078003, August 2018. .
34. E. Lauga, W. R. DiLuzio, G. M. Whitesides, and H. A. Stone. Swimming in circles: Motion of bacteria near solid boundaries. *Biophys J*, 90(2):400–412, January 2006. .
35. I. H. Riedel, K. Kruse, and J. Howard. A self-organized vortex array of hydrodynamically entrained sperm cells. *Science*, 309(5732):300–303, July 2005. .
36. A. Kaiser and H. Loewen. Vortex arrays as emergent collective phenomena for circle swimmers. *Physical Review E*, 87(3):032712, March 2013. .
37. Jonas Denk, Lorenz Huber, Emanuel Reithmann, and Erwin Frey. Active curved polymers form vortex patterns on membranes. *Phys Rev Lett*, 116(17):178301, April 2016. .
38. C. Hoell, H. Lowen, and A. M. Menzel. Dynamical density functional theory for circle swimmers. *New J Phys*, 19(12):125004 (18 pp.)–125004, 2017. .
39. B. Liebchen and D. Levis. Collective behavior of chiral active matter: Pattern formation and enhanced flocking. *Phys Rev Lett*, 119(5):058002, August 2017. .
40. Qun-Li Lei, Massimo Pica Ciamarra, and Ran Ni. Nonequilibrium strongly hyperuniform fluids of circle active particles with large local density fluctuations. *Sci Adv*, 5(1):eaau7423, January 2019. .
41. H. J. Jeong, S. Y. Lee, N. S. Kang, Y. Du Yoo, A. S. Lim, M. J. Lee, H. S. Kim, W. Yih, H. Yamashita, and T. C. LaJeunesse. Genetics and morphology characterize the dinoflagellate symbiodinium voratum, n. sp., (dinophyceae) as the sole representative of symbiodinium clade e. *J Eukaryot Microbiol*, 61(1):75–94, January 2014. .
42. Todd C. LaJeunesse, John Everett Parkinson, Paul W. Gabrielson, Hae Jin Jeong, James Davis Reimer, Christian R. Voelstra, and Scott R. Santos. Systematic revision of symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr Biol*, 28(16):2570–+, August 2018. .
43. S. Torquato and F. H. Stillinger. Local density fluctuations, hyperuniformity, and order metrics. *Physical Review E*, 68(4):041113, October 2003. .
44. S. Torquato. Hyperuniform states of matter. *Physics Reports-review Section of Physics Letters*, 745:1–95, June 2018. .
45. M. Florescu, S. Torquato, and P. J. Steinhardt. Designer disordered materials with large, complete photonic band gaps. *Proc Natl Acad Sci U S A*, 106(49):20658–20663, December 2009. .
46. R. B. Xie, G. G. Long, S. J. Weigand, S. C. Moss, T. Carvalho, S. Roorda, M. Hejra, S. Torquato, and P. J. Steinhardt. Hyperuniformity in amorphous silicon based on the measurement of the infinite-wavelength limit of the structure factor. *Proc Natl Acad Sci U S A*, 110(33):13250–13254, August 2013. .
47. Weining Man, Marian Florescu, Kazue Matsuyama, Polin Yadak, Gev Nahal, Seyed Hashemizad, Eric Williamson, Paul Steinhardt, Salvatore Torquato, and Paul Chaikin. Photonic band gap in isotropic hyperuniform disordered solids with low dielectric contrast. *Opt Express*, 21(17):19972–19981, August 2013. .
48. S. Torquato, G. Zhang, and F. H. Stillinger. Ensemble theory for stealthy hyperuniform disordered ground states. *Phys Rev X*, 5(2):021020, May 2015. .
49. J. Ricouvier, P. Tabeling, and P. Yazhgor. Foam as a self-assembling amorphous photonic band gap material. *Proc Natl Acad Sci U S A*, 116(19):9202–9207, May 2019. .
50. Yang Jiao, Timothy Lau, Haralampos Hatzikirou, Michael Meyer-Hermann, Joseph C. Corbo, and Salvatore Torquato. Avian photoreceptor patterns represent a disordered hyperuniform solution to a multiscale packing problem. *Physical Review E*, 89(2):022721, February 2014. .
51. Fausto Martelli, Salvatore Torquato, Nicolas Giovambattista, and Roberto Car. Large-scale structure and hyperuniformity of amorphous ices. *Phys Rev Lett*, 119(13):136002, September 2017. .
52. Yu Zheng, Lei Liu, Hanqing Nan, Zhen-Xiong Shen, Ge Zhang, Duyu Chen, Lixin He, Wenxiang Xu, Mohan Chen, Yang Jiao, and Houlong Zhuang. Disordered hyperuniformity in two-dimensional amorphous silica. *Sci Adv*, 6(16):eaab0826, April 2020. .
53. Igor Lesanovsky and Juan P. Garrahan. Out-of-equilibrium structures in strongly interacting rydberg gases with dissipation. *Phys Rev A*, 90(1):011603, July 2014. .
54. A. Donev, F. H. Stillinger, and S. Torquato. Unexpected density fluctuations in jammed disordered sphere packings. *Phys Rev Lett*, 95(9):090604, August 2005. .
55. Alexandros Chremos and Jack F. Douglas. Hidden hyperuniformity in soft polymeric materials. *Phys Rev Lett*, 121(25):258002, December 2018. .
56. Elsen Tjhung and Ludovic Berthier. Hyperuniform density fluctuations and diverging dynamic correlations in periodically driven colloidal suspensions. *Phys Rev Lett*, 114(14):148301, April 2015. .
57. Joost H. Weijs, Raphael Jeanneret, Remi Dreyfus, and Denis Bartolo. Emergent hyperuniformity in periodically driven emulsions. *Phys Rev Lett*, 115(10):108301, September 2015. .
58. Tomer Goldfriend, Haim Diamant, and Thomas A. Witten. Screening, hyperuniformity, and instability in the sedimentation of irregular objects. *Phys Rev Lett*, 118(15):158005, April 2017. .
59. Joost H. Weijs and Denis Bartolo. Mixing by unstirring: Hyperuniform dispersion of interacting particles upon chaotic advection. *Phys Rev Lett*, 119(4):048002, July 2017. .
60. Joshua Ricouvier, Romain Pierrat, Remi Carminati, Patrick Tabeling, and Pavel Yazhgor. Optimizing hyperuniformity in self-assembled bidisperse emulsions. *Phys Rev Lett*, 119(20):208001, November 2017. .
61. Jikai Wang, J. M. Schwarz, and Joseph D. Paulsen. Hyperuniformity with no fine tuning in sheared sedimenting suspensions. *Nat Commun*, 9:2836, July 2018. .
62. Daniel Hexner and Dov Levine. Hyperuniformity of critical absorbing states. *Phys Rev Lett*, 114(11):110602, March 2015. .
63. Daniel Hexner and Dov Levine. Noise, diffusion, and hyperuniformity. *Phys Rev Lett*, 118(2):020601, January 2017. .
64. Daniel Hexner, Paul M. Chaikin, and Dov Levine. Enhanced hyperuniformity from random reorganization. *Proc Natl Acad Sci U S A*, 114(17):4294–4299, April 2017. .
65. Melissa S. Roth. The engine of the reef: photobiology of the coral-algal symbiosis. *Front Microbiol*, 5:422, August 2014. .
66. Yusuke Aihara, Shinichiro Maruyama, Andrew H. Baird, Akira Iguchi, Shunichi Takahashi, and Jun Minagawa. Green fluorescence from cnidarian hosts attracts symbiotic algae. *Proc Natl Acad Sci U S A*, 116(6):2118, February 2019. .
67. T. Ohmura, Y. Nishigami, A. Taniguchi, S. Nonaka, J. Manabe, T. Ishikawa, and M. Ichikawa. Simple mechanosense and response of cilia motion reveal the intrinsic habits of ciliates. *Proc*



- Natl Acad Sci U S A*, 115(13):3231–3236, March 2018. .
68. Gregory Gaines and F. J. R. Taylor. Form and function of the dinoflagellate transverse flagellum1. *The Journal of Protozoology*, 32(2):290–296, January 1985. ISSN 0022-3921. .
  69. T. Fenchel. How dinoflagellates swim. *Protist*, 152(4):329–338, December 2001. .
  70. I. Miyasaka, K. Nanba, K. Furuya, Y. Nimura, and A. Azuma. Functional roles of the transverse and longitudinal flagella in the swimming motility of prorocentrum minimum (dinophyceae). *J Exp Biol*, 207(17):3055–3066, August 2004. .
  71. Robert Edward Lee. *Phycology*. Cambridge University Press, Cambridge, 4 edition, 2008. .
  72. K. Drescher, J. Dunkel, L. H. Cisneros, S. Ganguly, and R. E. Goldstein. Fluid dynamics and noise in bacterial cell-cell and cell-surface scattering. *Proc. Natl. Acad. Sci. U. S. A.*, 108(27):10940–10945, July 2011. .
  73. J. Ainley, S. Durkin, R. Embid, P. Boindala, and R. Cortez. The method of images for regularized stokeslets. *J. Comput. Phys.*, 227(9):4600–4616, April 2008. .
  74. S. E. Spagnolie and E. Lauga. Hydrodynamics of self-propulsion near a boundary: predictions and accuracy of far-field approximations. *J. Fluid Mech.*, 700:105–147, June 2012. .
  75. G.S. Perkins and R.B. Jones. Hydrodynamic interaction of a spherical particle with a planar boundary i. free surface. *Physica A-statistical Mechanics and Its Applications*, 171(3):575–604, 1991. .
  76. R. Ma, G. S. Klindt, I. H. Riedel-Kruse, F. Julicher, and B. M. Friedrich. Active phase and amplitude fluctuations of flagellar beating. *Phys Rev Lett*, 113(4):048101, July 2014. .
  77. Kirsty Y. Wan and Raymond E. Goldstein. Rhythmicity, recurrence, and recovery of flagellar beating. *Phys Rev Lett*, 113(23):238103, December 2014. .
  78. Ken H. Nagai, Yutaka Sumino, Raul Montagne, Igor S. Aranson, and Hugues Chate. Collective motion of self-propelled particles with memory. *Phys Rev Lett*, 114(16):168001, April 2015. .
  79. Jaewuk Kim and Salvatore Torquato. Effect of imperfections on the hyperuniformity of many-body systems. *Phys Rev B*, 97(5):054105, February 2018. .
  80. Alexander P. Petroff, Xiao-Lun Wu, and Albert Libchaber. Fast-moving bacteria self-organize into active two-dimensional crystals of rotating cells. *Phys Rev Lett*, 114:158102, Apr 2015. .
  81. Xiao Chen, Xiang Yang, Mingcheng Yang, and H. P. Zhang. Dynamic clustering in suspension of motile bacteria. *Europhys Lett*, 111(5):54002 (6 pp.)–54002, 2015. .
  82. Qun-Li Lei and Ran Ni. Hydrodynamics of random-organizing hyperuniform fluids. *Proc Natl Acad Sci U S A*, 116(46):22983–22989, November 2019. .
  83. Jeffrey L. Moran and Jonathan D. Posner. Phoretic self-propulsion. *Annual Review of Fluid Mechanics*, Vol 49, 49:511–540, 2017. .