REGULARITY PROPERTIES OF k-BRJUNO AND WILTON FUNCTIONS

SEUL BEE LEE, STEFANO MARMI, IZABELA PETRYKIEWICZ, AND TANJA I. SCHINDLER

ABSTRACT. We study functions related to the classical Brjuno function, namely k-Brjuno functions and the Wilton function. Both appear in the study of boundary regularity properties of (quasi) modular forms and their integrals. We consider various possible versions of them, based on the α -continued fraction developments. We study their BMO regularity properties and their behaviour near rational numbers of their finite truncations. We then complexify the functional equations which they fulfill and we construct analytic extensions of the k-Brjuno and of the Wilton function to the upper half plane. We study their boundary behaviour using an extension of the continued fraction algorithm to the complex plane. We also prove that the harmonic conjugate of the real k-Brjuno function is continuous at all irrational numbers and has a decreasing jump of π/q^k at rational points p/q.

1. Introduction

Let G denote the Gauss map defined on the interval [0,1) by G(0)=0 and $G(x)=\left\{\frac{1}{x}\right\}$ otherwise, where $\{x\}=x-|x|$ and let

$$\beta_j(x) = \prod_{i=0}^j G^i(x)$$

for $j \ge 0$ with the convention $\beta_{-1}(x) = 1$.

1.1. **Brjuno function.** In 1988, Yoccoz introduced the following function - now called *Brjuno function* - defined for irrational numbers $x \in [0,1] \setminus \mathbb{Q}$ as

$$B_1(x) = \sum_{n=0}^{\infty} \beta_{n-1}(x) \log \left(\frac{1}{G^n(x)}\right),$$

see [Yoc88, MMY97, MMY06], see Figure 1 for its graph.

Date: September 16, 2021.

2020 Mathematics Subject Classification. 30B70, 37F50, 32A50, 46F15, 32A37, 11F03.

Key words and phrases. Brjuno function; Brjuno condition; Wilton function; bounded mean oscillation; complex boundary behaviour; hyperfunctions; modular group.

The first and the fourth author acknowledge the support of the Centro di Ricerca Matematica Ennio de Giorgi. The first, second and third author acknowledge the support of UniCredit Bank R&D group for financial support through the 'Dynamics and Information Theory Institute' at the Scuola Normale Superiore. The second and the fourth author acknowledge the support through the PRIN Grant "Regular and stochastic behaviour in dynamical systems" (PRIN 2017S35EHN).

Let $\frac{p_n(x)}{q_n(x)}$ denote the *n*th convergent of *x* with respect to its continued fraction expansion. The series $B_1(x)$ converges if and only if

$$\sum_{n=0}^{\infty} \frac{\log(q_{n+1}(x))}{q_n(x)} < \infty.$$

This condition is called Brjuno condition and was introduced by Brjuno in the study of certain problems in dynamical systems, see [Brj71, Brj72]. The points of convergence are called Brjuno numbers. The importance of Brjuno numbers comes from the study of analytic small divisors problems in dimension one. Indeed, extending previous fundamental work of C.L. Siegel [Sie42], Brjuno proved that all germs of holomorphic diffeomorphisms of one complex variable with an indifferent fixed point with linear part $e^{2\pi ix}$ are linearisable if x is a Brjuno number. Conversely, in 1988 J.-C. Yoccoz [Yoc88, Yoc95] proved that this condition is also necessary. Similar results hold for the local conjugacy of analytic diffeomorphisms of the circle [Yoc02] and for some complex area-preserving maps [Mar90, Day 94]. Moreover the Brjuno function provides a continuous interpolation of the radius of convergence of the quadratic polynomial [BC06] and the interpolation is conjectured to be Hölder continuous with exponent 1/2 [MMY97]. This has been recently proved for hightype irrational numbers [CC15]. The Brjuno condition has been of interest also in different contexts. For instance, it is conjectured that it is optimal for the existence of real analytic invariant circles in the standard family [Mac88, Mac89, MS92]. See also [BG01, Gen15] and references therein for related results.

Furthermore, the Brjuno function is Z-periodic and satisfies the functional equation

$$B_1(x) = -\log(x) + xB_1\left(\frac{1}{x}\right)$$

for $x \in (0,1)$. The second author together with Moussa and Yoccoz investigated the regularity properties of B_1 in [MMY97] and later constructed an analytic extension of B_1 to the upper-half plane $\mathbb{H} = \{z \in \mathbb{C} : \text{Im} z > 0\}$ [MMY01]. Let T denote the linear operator

$$Tf(x) = xf\left(\frac{1}{x}\right) \tag{1.1}$$

acting, for example, on measurable \mathbb{Z} -periodic functions on \mathbb{R} . Then in all L^p spaces the Brjuno function is the solution of the linear equation

$$[(1-T)B_1](x) = -\log x. (1.2)$$

By exploiting the fact that the operator T as in (1.1) acting on L^p spaces has spectral radius strictly smaller than 1, one can indeed obtain (1.1) by a Neumann series for $(1-T)^{-1}$, see [MMY97] for details.

Local properties of the Brjuno function have been recently investigated by M. Balazard and B. Martin [BM12] and its multifractal spectrum was determined by S. Jaffard and B. Martin in [JM18].

1.2. k-Brjuno functions. For the following, for $k \geq 2$ even, let E_k be the Eisenstein series of weight k defined in the upper-half plane \mathbb{H} . Then its Fourier expansion is given by

$$E_k(z) = 1 - \frac{2k}{b_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) e^{2\pi i n z},$$

where b_k is the kth Bernoulli number and $\sigma_{k-1}(n) = \sum_{d|n} d^{k-1}$. For all $k \geq 4$, E_k is modular of weight k under the action of $\mathrm{SL}_2(\mathbb{Z})$, and E_2 is quasi-modular of weight 2 under the action of $\mathrm{SL}_2(\mathbb{Z})$, see for example [Zag92]. The function E_2 can be viewed as a modular (or Eichler) integral on $\mathrm{SL}_2(\mathbb{Z})$ of weight 2 with the rational period function $-\frac{2\pi i}{z}$, see for example [Kno90].

For $k \geq 2$ even and $z \in \mathbb{H}$, denote $\varphi_k(z) = \sum_{n=1}^{\infty} \frac{\sigma_{k-1}(n)}{n^{k+1}} e^{2\pi i n z}$. We have that

$$\sum_{n=1}^{\infty} \sigma_{k-1}(n) e^{2\pi i n z} = \left(\frac{1}{2\pi i} \frac{\partial}{\partial z}\right)^{k+1} \varphi_k(z),$$

and

$$\varphi_k(z) = \frac{B_k(2\pi i)^{k+1}}{k!2k} \int_{i\infty}^z (z-t)^k (E_k(t) - 1) dt.$$

Consider the imaginary part of φ_k

$$F_k(x) = \sum_{n=1}^{\infty} \frac{\sigma_{k-1}(n)}{n^{k+1}} \sin(2\pi nx)$$
 for $x \in \mathbb{R}$.

Analytic properties, differentiability and Hölder regularity exponent, of the function F_k (and the real part of φ_k) were studied by the third author. It has been proved that for F_k the differentiability is related to a condition resembling the Brjuno condition. Considering the special case k=2, the third author proved that if $\sum_{n=0}^{\infty} \frac{\log(q_{n+1}(x))}{q_n^2(x)} < \infty$ and $\lim_{n\to\infty} \frac{\log(q_{n+4}(x))}{q_n^2(x)} = 0$, then F_2 is differentiable at $x \in \mathbb{R} \setminus \mathbb{Q}$, whereas if $\sum_{n=0}^{\infty} \frac{\log(q_{n+1}(x))}{q_n^2(x)}$ diverges, then F_2 is not differentiable at $x \in \mathbb{R} \setminus \mathbb{Q}$. It has been conjectured that for all $k \in \mathbb{N}$ even, F_k is differentiable at $x \in \mathbb{R} \setminus \mathbb{Q}$ if and only if it fulfills the k-Brjuno condition

$$\sum_{n=0}^{\infty} \frac{\log(q_{n+1}(x))}{q_n^k(x)} < \infty, \tag{1.3}$$

see [Pet14, Pet17]. The occurrence of a condition of this type motivates the following definition.

For $k \in \mathbb{N}$, let

$$B_k(x) = \sum_{n=0}^{\infty} (\beta_{n-1}(x))^k \log\left(\frac{1}{G^n(x)}\right)$$

$$\tag{1.4}$$

be called k-Brjuno function. From this equation we already get the implicit definition in terms of an analogue of the functional equation of the Brjuno function: if $x \in (0,1)$

$$B_k(x) = -\log(x) + x^k \cdot B_k(G(x)),$$

see [MMY97, MMY06]. The k-Brjuno function converges at an irrational x if and only if (1.3) holds, see also Proposition 2.5 for an even stronger statement about the relation between the k-Brjuno function and (1.3). Obviously, for k = 1, the function in (1.4) gives the Brjuno function introduced before.

1.3. α -continued fractions. Instead of considering the k-Brjuno function with respect to the Gauss map as in (1.4), it is also possible to use α -continued fractions instead. The classical Brjuno function associated to α -continued fractions was already investigated in [MMY97]. Let $\alpha \in \left[\frac{1}{2}, 1\right]$ and let $A_{\alpha} : (0, \alpha) \to [0, \alpha]$ the transformation of the α -continued fractions being given by

$$A_{\alpha}(x) = \left| \frac{1}{x} - \left| \frac{1}{x} - \alpha + 1 \right| \right|. \tag{1.5}$$

For $\alpha=1$, we obtain the Gauss map associated to the regular continued fraction transformation and for $\alpha=1/2$, we obtain the transformation associated to the nearest integer continued fractions. Nakada [Nak81] was the first to consider all these types of continued fractions as a one-parameter family, however he considered the 'unfolded' version of the α -continued fraction which is defined by the Gauss map $\widetilde{A}_{\alpha}(x) = \frac{1}{x} - \left\lfloor \frac{1}{x} - \alpha + 1 \right\rfloor$. The version as in (1.5) was little later considered in [TI81].

Further, let

$$B_{k,\alpha}(x) = \sum_{n=0}^{\infty} \left(\beta_{n-1}^{(\alpha)}(x) \right)^k \log \left(\frac{1}{A_{\alpha}^n(x)} \right),$$

where $\beta_j^{(\alpha)}(x) = \prod_{i=0}^j A_\alpha^i(x)$ for $j \geq 0$ and $\beta_{-1}^{(\alpha)}(x) = 1$, be the generalisation of (1.4) in the sense that we consider the k-Brjuno function not only for the Gauss transformation but also for other α -continued fraction transformations A_α with $\alpha \in [1/2, 1]$.

For given $\alpha \in [1/2, 1]$, any $x \in (0, 1]$ has the α -continued fraction expansion given by

$$x = \frac{1}{a_1 + \frac{\epsilon_1}{\cdots + \frac{\epsilon_{j-1}}{a_j + \cdots}}},$$

where $a_j := a_j^{(\alpha)}(x) := \left\lfloor \frac{1}{A_\alpha^{j-1}(x)} - \alpha + 1 \right\rfloor$ and $\epsilon_j := \epsilon_j^{(\alpha)}(x)$ is the sign of $\frac{1}{A_\alpha^{j-1}(x)} - a_j$.

Related to (1.1) we define an operator $T_{k,\alpha}$ acting on \mathbb{Z} -periodic measurable functions f such that f(-x) = f(x) for a.e. $x \in (0, 1 - \alpha)$ as

$$T_{k,\alpha}f(x) = x^k f\left(\frac{1}{x}\right), \quad x \in (0,\alpha).$$
 (1.6)

It is the operator $T_{\nu}^{(\alpha)}$ which is introduced in [MMY97], where ν corresponds to the exponent k in (1.6). It is understood that the function $T_{k,\alpha}f$ is completed outside the interval $(0,\alpha)$ by imposing to $T_{k,\alpha}f$ the same parity and periodicity conditions imposed to f.

Then we have

$$[(1 - T_{k,\alpha})B_{k,\alpha}](x) = -\log x, \quad x \in (0,\alpha)$$
(1.7)




FIGURE 1. Numerical computation of the Brjuno function B (left) and of the Wilton function W (right) when $\alpha = 1$. The asymmetric logarithmic singularities at rational points provide an intuitive justification for W not belonging to the BMO space, see Section 2.

which follows by a simple calculation.

1.4. **Wilton function.** Next we consider the related concept of the *Wilton function* which is given by

$$W(x) = \sum_{n=0}^{\infty} (-1)^n \beta_{n-1}(x) \log \left(\frac{1}{G^n(x)}\right),$$

namely by the alternate signs version of the Brjuno function series (1.1). (see Figure 1 for its graph). It converges if and only if it fulfills the *Wilton condition*

$$\left| \sum_{n=0}^{\infty} (-1)^n \frac{\log(q_{n+1}(x))}{q_n(x)} \right| < \infty,$$

see [BM13, Prop. 7] and Remark 2.7 for an even stronger connection between the Wilton function and the Wilton condition. The points of convergence are called *Wilton numbers* and appear in the work of Wilton, see [Wil33]. Clearly all Brjuno numbers are Wilton, but not vice versa (it is not difficult to build counterexamples by using the continued fraction).

The function W satisfies the functional equation for $x \in (0,1)$ being a Wilton number:

$$W(x) = \log\left(\frac{1}{x}\right) - xW(G(x)),$$

which by using the same linear operator T as in (1.1) can be written as

$$[(1+T)W](x) = -\log x. \tag{1.8}$$

We can extend the Wilton function in the same way we did for the k-Brjuno functions, i.e. replacing the Gauss map with the transformation of the α -continued fractions: for

6

 $\alpha \in \left[\frac{1}{2}, 1\right)$ and for all irrational x we define

$$W_{\alpha}(x) = \sum_{n=0}^{\infty} (-1)^n \beta_{n-1}^{(\alpha)}(x) \log \left(\frac{1}{A_{\alpha}^n(x)}\right).$$

We then have

$$[(1 - S_{\alpha})W_{\alpha}](x) = -\log x, \quad x \in (0, \alpha),$$

where the operator $S_{\alpha} = -T_{1,\alpha}$. Also in this case it is understood that S_{α} acts on \mathbb{Z} -periodic measurable functions f such that f(-x) = f(x) for a.e. $x \in (0, 1 - \alpha)$.

The Wilton function and its primitive have been studied recently by Balazard and Martin in terms of its convergence properties [BM19] and in the context of the Nyman and Beurling criterion, see [BM12, BM13] and [BBLS05].

1.5. **BMO** properties and complex k-Brjuno and Wilton functions. In [MMY97] the real regularity properties of the Brjuno function were systematically investigated by exploiting the associated functional equation, culminating with the proof that the Brjuno function belongs to the space BMO of functions with Bounded Mean Oscillation (we refer to [Gar07, GD11] for its definition and properties). Here (see Section 2) we extend those results to the k-Brjuno function, which turn out to belong to BMO, and to the Wilton function, which does not. Moreover we consider their generalizations obtained by replacing the Gauss map with α -continued fractions [Nak81], $\alpha \in \left[\frac{1}{2}, 1\right]$ and we prove that for all α the associated k-Brjuno functions belong to BMO and that the same holds for the associated Wilton functions when α is restricted to belong to the interval $\left[\frac{1}{2}, \frac{\sqrt{5}-1}{2}\right]$.

By Fefferman's duality theorem, see for example [Sar78, p. 39], one can add an L^{∞} function to a BMO function and obtain that the harmonic conjugate of the sum will be an essentially bounded function. One is then led to look for a periodic holomorphic function defined on the upper half plane whose imaginary parts, when looked upon \mathbb{R} , is the original function. As in [MMY01] we make this construction for the k-Brjuno functions and for the Wilton function. The associated complex k-Brjuno functions \mathcal{B}_k and complex Wilton function \mathcal{W} and their properties are summarized in the following Theorem:

Theorem 1.1. (1) The complex k-Brjuno function \mathcal{B}_k , $\mathbb{H} \to \mathbb{C}$ is given by the series

$$\mathcal{B}_{k}(z) = -\sum_{p/q \in \mathbb{Q}} \det \begin{pmatrix} p' & p \\ q' & q \end{pmatrix}^{k+1} \cdot \left\{ \frac{1}{\pi} \left[(p' - q'z)^{k} \operatorname{Li}_{2} \left(\frac{p' - q'z}{qz - p} \right) - (q''z - p'')^{k} \operatorname{Li}_{2} \left(\frac{p'' - q''z}{qz - p} \right) \right] + \sum_{n=0}^{k} \frac{\det \begin{pmatrix} p' & p \\ q' & q \end{pmatrix}^{n}}{n\pi q^{n}} \left[(p' - q'z)^{k-n} \left(\log \left(1 + \frac{q'}{q} \right) - \sum_{i=1}^{n-1} \frac{1}{i} \left(\frac{1}{(1 + q'/q)^{i}} - 1 \right) \right) - (q''z - p'')^{k-n} \left(\log \left(1 + \frac{q''}{q} \right) - \sum_{i=1}^{n-1} \frac{1}{i} \left(\frac{1}{(1 + q''/q)^{i}} - 1 \right) \right) \right] \right\}, \quad (1.9)$$

where $\left[\frac{p'}{q'},\frac{p''}{q''}\right]$ is the Farey interval such that $\frac{p}{q}=\frac{p'+p''}{q'+q''}$ (with the convention p'=p-1, q'=1, p''=1, q''=0 if q=1) and $\mathrm{Li}_2(z)$ is the dilogarithm. The complex Wilton function \mathcal{W} , $\mathbb{H} \to \mathbb{C}$ is given by the series

$$\mathcal{W}(z) = -\frac{1}{\pi} \sum_{p/q \in \mathbb{Q}} \left\{ \det \begin{pmatrix} p' & p \\ q' & q \end{pmatrix} (p' - q'z) \left[\operatorname{Li}_{2} \left(\frac{p' - q'z}{qz - p} \right) - \operatorname{Li}_{2} \left(-\frac{q'}{q} \right) \right] + \det \begin{pmatrix} p'' & p \\ q'' & q \end{pmatrix} (p'' - q''z) \left[\operatorname{Li}_{2} \left(\frac{p'' - q''z}{qz - p} \right) - \operatorname{Li}_{2} \left(-\frac{q''}{q} \right) \right] + \frac{1}{q} \log \left(\frac{q^{2}}{(q + q')(q + q'')} \right) \right\}.$$

$$(1.10)$$

- (2) The real part of \mathcal{B}_k is bounded on the upper half plane and its non-tangential limit on \mathbb{R} is continuous at all irrational points and has a decreasing jump of $\frac{\pi}{q^k}$ at each rational number $\frac{p}{q}$.
- (3) As one approaches the boundary the imaginary part of W behaves as follows:
 - (a) if $x \in \mathbb{R}$ is a Wilton number, then $\operatorname{Im} \mathcal{W}(x+w)$ converges to W(x) as $w \to 0$ in any domain with a finite order of tangency to the real axis;
 - (b) if $x \in \mathbb{R}$ is Diophantine, then in both cases one can allow domains with infinite order of tangency to the real line.

As the Wilton function is not BMO, it also becomes clear by Fefferman's theorem that its harmonic conjugate can not be an L^{∞} function.

1.6. Structure of the paper. The paper is organised as follows. In Section 2 we state and prove the BMO-properties of the k-Brjuno and the Wilton function and in Section 3 we give statements about the truncated real k-Brjuno and the truncated real Wilton function which corresponds to the finite part of the sum at rational numbers.

The subsequent sections deal with a complexification of the system. In Section 4 we introduce a complex continued fraction algorithm. In Section 5 we extend the operators $T_k := T_{k,1}$ and $S := S_1$ to the complex plane. The main findings concerning these operators are given in Section 6. Particularly, Proposition 6.9 indicates how T_k behaves. The analogous behaviour of S can then immediately be deduced by using the fact that $S = -T_1$.

Finally, with the help of these operators we define the complex k-Brjuno and the complex Wilton function in Section 7.

Those proofs which are very similar to the ones in [MMY01] are given in an appendix.

2. BMO properties of the real k-Brjuno and the real Wilton function

In this section, we study the bounded mean oscillation (BMO) properties of the real k-Brjuno and the Wilton function - both with respect to different transformations A_{α} with $\alpha \in [1/2, 1]$. Before stating the main results of this section, we will first recall the definition of a BMO function.

Let $L^1_{loc}(\mathbb{R})$ be the space of the locally integrable functions on \mathbb{R} . Recall that the mean value of a function $f \in L^1_{loc}(\mathbb{R})$ on an interval I is defined as

$$f_I = \frac{1}{|I|} \int_I f(x) dx.$$

For an interval U, we say that a function $f \in BMO(U)$ if

$$||f||_{*,U} := \sup_{I \subset U} \frac{1}{|I|} \int_{I} |f(x) - f_I| dx < \infty.$$
 (2.1)

For further properties of the BMO space, see for example [MMY97, Appendix] and the monographies [Gar07, GD11].

In the following, we will state the main properties of this section which show that the BMO properties fundamentally differ between the k-Brjuno functions and the Wilton function. We first give the statement for the k-Brjuno functions.

Proposition 2.1. For all $k \in \mathbb{N}$ and all $\alpha \in [1/2, 1]$, the k-Brjuno function $B_{k,\alpha}$ is a BMO function.

Contrarily, for the Wilton function we have the following statement:

Theorem 2.2. The Wilton function $W = W_1$ is not a BMO function.

On the other hand, we define for the following $g := \frac{\sqrt{5}-1}{2}$ and have:

Theorem 2.3. For all $\alpha \in \left[\frac{1}{2}, g\right]$, the function W_{α} is a BMO function.

Before we start with the proofs of the statements above, we first want to give some remarks about them: Proposition 2.1 is an extension of [MMY97, Thm. 3.2] from the classical Brjuno function to k-Brjuno functions.

As a comparison to Figure 1 showing $B_{1,1}$ and W_1 , in Figure 2, some numerical simulations of W_{α} with different values of α are shown: the numerical evidence supports the conjecture that also for $\alpha \in (g,1)$ the function W_{α} is a BMO function. Indeed, the graph for $\alpha = e - 2$ and $\alpha = 0.9$ (and nearby values, not displayed in Figure 1) suggest that the function W_{α} exhibits a mix of symmetric logarithmic singularities at rational points and of jumps, a singular behaviour which is compatible with belonging to BMO. However, unfortunately, the results from Proposition 2.3 can not immediately be transferred to $\alpha \in (g,1)$, see Remark 2.10 for an explanation which difficulties occur.

2.1. Proof that the real k-Brjuno functions are BMO functions. The main idea of the proof is to prove the statement for $B_{k,1/2}$, see Proposition 2.4, and to show then that $B_{k,\alpha}$ differs from $B_{k,1}$ only by an L^{∞} function which follows from Propositions 2.5 and 2.8. As the proof follows from very similar arguments as those in [MMY97, Thm. 3.2], we will only describe shortly the necessary changes in the proofs.

We start by introducing

$$X_* = \{ f \in \text{BMO}(\mathbb{R}) : f(x+1) = f(x) \text{ for all } x \in \mathbb{R}, \ f(-x) = f(x) \text{ for all } x \in [0, 1/2] \}.$$
(2.2)




FIGURE 2. Numerical computations of W_{α} for $\alpha = 1/2$ (upper left), $\alpha = \frac{\sqrt{5}-1}{2}$ (upper right), $\alpha = e-2$ (lower left) and $\alpha = 0.9$ (lower right).

endowed with a norm which is the sum of the BMO seminorm $\|\cdot\|_{*,[0,1/2]}$ as in (2.1) and of the L^2 norm on the interval (0,1/2) (w.r.t. the $A_{1/2}$ -invariant probability measure). Then one has:

Proposition 2.4 ([MMY97, Thm. 3.3]). For all $k \in \mathbb{N}$, the operator $T_{k,1/2}$ as in (1.6) is a bounded linear operator from X_* to X_* whose spectral radius is bounded above by $(\sqrt{2}-1)^k$.

To proceed, we prove an analog of [MMY97, Prop. 2.3, eq. (iv)].

Proposition 2.5. For all $k \in \mathbb{N}$, there exists a constant $C_{1,k} > 0$ such that for all $\alpha \in [1/2, 1]$ and $x \in \mathbb{R} \setminus \mathbb{Q}$, one has

$$\left| B_{k,\alpha}(x) - \sum_{j=0}^{\infty} \frac{\log q_{j+1}^{(\alpha)}(x)}{\left(q_j^{(\alpha)}(x) \right)^k} \right| < C_{1,k}.$$

Before we start with the proof, we recall the following property.

Remark 2.6. In [MMY97, Remark 1.7] it is showed that, for all $\alpha \in [1/2, 1]$,

$$\sum_{j=0}^{\infty} \frac{\log q_j^{(\alpha)}}{q_j^{(\alpha)}} \le 2/e =: c_1 \text{ and } \sum_{j=0}^{\infty} \frac{\log 2}{q_j^{(\alpha)}} \le 5 \log 2 =: c_2.$$

Proof of Proposition 2.5. For the following calculations we drop the dependence on α and x. We obtain by analogous calculations as in [MMY97, Prop. 2.3, eq. (iv)] that

$$\left| -B_{k,\alpha}(x) + \sum_{j=0}^{\infty} \frac{\log q_{j+1}}{q_j^k} \right|$$

$$= \left| \sum_{j=0}^{\infty} \beta_{j-1}^k \log \frac{\beta_j}{\beta_{j-1}} + \sum_{j=0}^{\infty} \left(\beta_{j-1} + \epsilon_j \frac{q_{j-1}}{q_j} \beta_j \right)^k \log q_{j+1} \right|$$

$$= \left| \sum_{j=0}^{\infty} \beta_{j-1}^k \log(\beta_j q_{j+1}) - \sum_{j=0}^{\infty} \beta_{j-1}^k \log \beta_{j-1} + \sum_{j=0}^{\infty} \left(\left(\beta_{j-1} + \epsilon_j \frac{q_{j-1}}{q_j} \beta_j \right)^k - \beta_{j-1}^k \right) \log q_{j+1} \right|$$

$$\leq \sum_{j=0}^{\infty} \left| \beta_{j-1}^k \log(\beta_j q_{j+1}) \right| + \sum_{j=0}^{\infty} \left| \beta_{j-1}^k \log \beta_{j-1} \right|$$

$$+ \sum_{j=0}^{\infty} \left| \left(\left(\beta_{j-1} + \epsilon_j \frac{q_{j-1}}{q_j} \beta_j \right)^k - \beta_{j-1}^k \right) \log q_{j+1} \right| .$$

We notice that $|\beta_i| \leq 1$ and thus

$$\sum_{j=0}^{\infty} \left| \beta_{j-1}^k \log(\beta_j q_{j+1}) \right| \le \sum_{j=0}^{\infty} \left| \beta_{j-1} \log(\beta_j q_{j+1}) \right| \le 2c_2,$$

where the last estimate follows as in the proof of [MMY97, Prop. 2.3, eq. (iv)] and c_2 is given in Remark (2.6). Furthermore,

$$\sum_{j=0}^{\infty} \left| \beta_{j-1}^k \log \beta_{j-1} \right| \le \sum_{j=0}^{\infty} \left| \beta_{j-1} \log \beta_{j-1} \right| \le 2(c_1 + c_2)$$

which also follows as in the proof of [MMY97, Prop. 2.3, eq. (iv)] and c_1 is given in Remark (2.6). Finally, we have

$$\sum_{j=0}^{\infty} \left| \left(\left(\beta_{j-1} + \epsilon_{j} \frac{q_{j-1}}{q_{j}} \beta_{j} \right)^{k} - \beta_{j-1}^{k} \right) \log q_{j+1} \right|$$

$$\leq \sum_{j=0}^{\infty} \left| \left(\sum_{\ell=1}^{k} {k \choose \ell} \beta_{j-1}^{k-\ell} \cdot \left(\epsilon_{j} \frac{q_{j-1}}{q_{j}} \beta_{j} \right)^{\ell} \right) \log q_{j+1} \right|$$

$$\leq \sum_{j=0}^{\infty} \left| (2^{k} - 1) \frac{q_{j-1}}{q_{j}} \beta_{j} \log q_{j+1} \right| \leq 2^{k+1} c_{1},$$

where the last estimate again follows analogously to [MMY97, Prop. 2.3, eq. (iv)] and the proof can be completed in the same way as in [MMY97, Prop. 2.3, eq. (iv)].

Remark 2.7. With analogous methods as above using the absolute values of the sum we also obtain the following statement: There exists a constant C > 0 such that for all $\alpha \in [1/2, 1]$ and $x \in \mathbb{R} \setminus \mathbb{Q}$, one has

$$\left| W_{\alpha}(x) - \sum_{j=0}^{\infty} (-1)^{j} \frac{\log q_{j+1}^{(\alpha)}(x)}{q_{j}^{(\alpha)}(x)} \right| < C.$$

The next proposition is an equivalent to [MMY97, Prop. 2.4].

Proposition 2.8. For all $k \in \mathbb{N}$, there exists a constant $C_{2,k} > 0$ such that for all $\alpha \in [1/2, 1]$ and for all $x \in \mathbb{R} \setminus \mathbb{Q}$ one has

$$\left| B_{k,\alpha}(x) - \sum_{j=0}^{\infty} \frac{\log q_{j+1}^{(1)}(x)}{(q_j^{(1)}(x))^k} \right| \le C_{2,k}.$$

Proof. The proof follows completely analogously to that of [MMY97, Prop. 2.4] with the only difference that we use Proposition 2.5 instead of [MMY97, Prop. 2.3] and each $q_j^{(1)}$ (which is denoted by Q_j in [MMY97]) in the denominator is replaced by $(q_j^{(1)})^k$.

Proof of Proposition 2.1. By Proposition 2.4, $1-T_{k,\alpha}$ is invertible on X_* . A \mathbb{Z} -periodic even function equal to $-\log x$ on (0,1/2] is in X_* . Thus, $B_{k,1/2}$ is BMO. Since by Proposition 2.8 for all $\alpha \in [1/2,1]$ the k-Brjuno function $B_{k,\alpha}$ differs from $B_{k,1/2}$ only by an L^{∞} function, $B_{k,\alpha}$ is a BMO function as well.

2.2. Proofs of the BMO properties of the Wilton function.

Proof of Theorem 2.2. For brevity, in the following we write $O_I(f) = \frac{1}{|I|} \int_I |f(x) - f_I| dx$. Furthermore, as we only consider $\alpha = 1$, we also always write W instead of W_1 . By [MMY95, Prop. A.7], if I_1 and I_2 are two consecutive intervals, then

$$O_{I_1 \cup I_2}(f) = \frac{|I_1|}{|I_1| + |I_2|} O_{I_1}(f) + \frac{|I_2|}{|I_1| + |I_2|} O_{I_2}(f) + \frac{2|I_1||I_2|}{(|I_1| + |I_2|)^2} |f_{I_1} - f_{I_2}|. \tag{2.3}$$

Let $I_n := \left[-\frac{1}{n}, \frac{1}{n}\right] = \left[-\frac{1}{n}, 0\right] \cup \left[0, \frac{1}{n}\right] =: I_n^- \cup I_n^+$. By (2.3), $O_{I_n}(f) \ge \frac{1}{2} |f_{I_n^-} - f_{I_n^+}|$ for any $f \in L^1_{loc}(\mathbb{R})$. By [BM13, Lem. 2], we have

$$\int_0^x W(t)dt = x \log(1/x) + x + O(x^2),$$

and

$$\int_{1-x}^{1} W(t)dt = -x \log(1/x) - x + O(x^{2} \log(2/x)),$$

thus we clearly have

$$W_{I_n^+} = \log(n) + 1 + O(1/n).$$

Since W is \mathbb{Z} -periodic, we also have

$$W_{I_n^-} = -\log(n) - 1 + O(n^{-1}\log(2n)).$$

Thus,

$$O_{I_n}(W) \ge \log(n) + 1 + O(n^{-1}\log(2n)),$$

which completes the proof of the proposition.

For the proof of Theorem 2.3, we can not use exactly the same strategy as for the proof of Proposition 2.1. The reason is that, as we have seen in Theorem 2.2, W_1 is not a BMO function. Hence, comparing W_{α} for $\alpha < 1$ with W_1 can not work. Instead, the underlying idea of the proof is to use that $W_{1/2}$, the Wilton function with respect to the nearest integer continued fraction, is a BMO function and compare $W_{1/2}$ with W_{α} for $\alpha \in [1/2, g]$, see Proposition 2.9.

Let X_* be defined as in (2.2). Since $S_{1/2} = -T_{1,1/2}$ clearly by Proposition 2.4 we have $W_{1/2} \in X_*$. On the other hand one has

Proposition 2.9. For $\alpha \in [\frac{1}{2}, g]$, we have $W_{1/2} - W_{\alpha} \in L^{\infty}$, where $g = \frac{\sqrt{5}-1}{2}$.


Proof. Let $x \in [0, \frac{1}{2}]$. Since $\alpha \in [\frac{1}{2}, g]$, we have $2 - \frac{1}{\alpha} \le 1 - g \le \frac{1}{2+\alpha} \le 1 - \alpha$. Recall that

$$A_{1/2}(x) = \begin{cases} \frac{1}{x} - k & \text{if } x \in (\frac{1}{k+1/2}, \frac{1}{k}], \\ (k+1) - \frac{1}{x} & \text{if } x \in (\frac{1}{k+1}, \frac{1}{k+1/2}], \end{cases} \text{ for } k \ge 2.$$

Since $\alpha \leq \frac{1}{1+\alpha}$, we have

$$A_{\alpha}(x) = \begin{cases} \frac{1}{x} - k & \text{if } x \in (\frac{1}{k+\alpha}, \frac{1}{k}], \\ (k+1) - \frac{1}{x} & \text{if } x \in (\frac{1}{k+1}, \frac{1}{k+\alpha}], & \text{for } k \ge 2. \\ 2 - \frac{1}{x} & \text{if } x \in (\frac{1}{2}, \alpha], \end{cases}$$

See Figure 3 for the graphs of $A_{1/2}$ and A_{α} for a typical $\alpha \in (1/2, g)$.






FIGURE 3. The graph of $A_{1/2}$ and A_{α} for $\alpha \in [\frac{1}{2}, g]$, where $g = \frac{\sqrt{5}-1}{2}$.

Let us denote by $(a_n^{(\alpha)}, \epsilon_n^{(\alpha)})$ the *n*th partial quotient of x given as a α -continued fraction, $x_n^{(\alpha)} = A_\alpha^n(x)$, where $x_0^{(\alpha)} = x$ and $p_n^{(\alpha)}/q_n^{(\alpha)}$ its *n*th principal convergent. We have $\epsilon_1^{(1/2)} \neq \epsilon_1^{(\alpha)}$ if and only if $x \in (\frac{1}{k+\alpha}, \frac{1}{k+1/2})$ for some $k \geq 2$. In this case,

$$(a_1^{(1/2)},\epsilon_1^{(1/2)}) = (k+1,-1), \ (a_1^{(\alpha)},\epsilon_1^{(\alpha)}) = (k,1) \ \text{ and } \ x_1^{(1/2)} = 1 - x_1^{(\alpha)}.$$

Let

$$t_i = \frac{1}{3 - \frac{1}{3 - \frac{1}{3 - \frac{1}{3 - \frac{1}{3}}}}}$$
 and $\frac{r_i}{s_i} = \frac{1}{3 - \frac{1}{3 - \frac{1}{3 - \frac{1}{3}}}}$,

where 3 in the continued fraction expansion appears i times. Note that $r_i = s_{i-1}$ and $2 - \frac{1}{1 - t_i} = \frac{r_i}{s_i}$. More precisely,

$$\{t_i\}_{i\geq 0} = \left\{\frac{1}{2}, \frac{2}{5}, \frac{5}{13}, \frac{13}{34}, \frac{34}{89}, \cdots\right\} \subset \left(1 - g, \frac{2}{5}\right) \cup \left\{\frac{1}{2}\right\},$$
$$\left\{\frac{r_i}{s_i}\right\}_{i\geq 0} = \left\{0, \frac{1}{3}, \frac{3}{8}, \frac{8}{21}, \frac{21}{55}, \cdots\right\} \subset \{0\} \cup \left[\frac{1}{3}, 1 - g\right).$$

Then, $A_{1/2}(t_i) = t_{i-1}$, $A_{\alpha}(\frac{r_i}{s_i}) = A_{1/2}(\frac{r_i}{s_i}) = \frac{r_{i-1}}{s_{i-1}}$ for $i \ge 1$ and $t_i \searrow 1 - g$ and $\frac{r_i}{s_i} \nearrow 1 - g$. Now, we suppose that n is the minimal index such that

$$x_1^{(1/2)} = x_1^{(\alpha)}, \ x_2^{(1/2)} = x_2^{(\alpha)}, \ \cdots, x_{n-1}^{(1/2)} = x_{n-1}^{(\alpha)}, \ \text{but } x_n^{(1/2)} \neq x_n^{(\alpha)}.$$

Then,

$$(a_i^{(1/2)}, \epsilon_i^{(1/2)}) = (a_i^{(\alpha)}, \epsilon_i^{(\alpha)}) \text{ and } q_i^{(1/2)} = q_i^{(\alpha)} \text{ for } 1 \le i \le n-1,$$

$$x_n^{(1/2)} = 1 - x_n^{(\alpha)}, \ a_n^{(1/2)} = a_n^{(\alpha)} + 1 \ge 3, \ \epsilon_n^{(1/2)} = -1, \ \epsilon_n^{(\alpha)} = 1 \text{ and } q_n^{(1/2)} = q_n^{(\alpha)} + q_{n-1}^{(1/2)}.$$

Since $x_n^{(1/2)} \neq x_n^{(\alpha)}$, $x_n^{(1/2)} = 1 - x_n^{(\alpha)}$ and the domain of $A_{1/2}$ is $[0, \frac{1}{2}]$, we have

$$x_n^{(1/2)} \in \left[1 - \alpha, \frac{1}{2}\right) \quad \text{and} \quad x_n^{(\alpha)} \in \left(\frac{1}{2}, \alpha\right].$$
 (2.4)

Since $[1-\alpha,\frac{1}{2})\subset \bigsqcup_{i=1}^{\infty}(t_i,t_{i-1}]$, there is $i\geq 1$ such that $x_n^{(1/2)}\in (t_i,t_{i-1}]$. This implies

$$x_{n+j}^{(1/2)} = 3 - \frac{1}{x_{n+j-1}^{(1/2)}} \in (t_{i-j}, t_{i-j-1}] \text{ and } (a_{n+j}^{(1/2)}, \epsilon_{n+j}^{(1/2)}) = (3, -1) \text{ for } 1 \le j \le i - 1,$$

$$x_{n+i}^{(1/2)} = \frac{1}{x_{n+i-1}^{(1/2)}} - 2 \text{ and } (a_{n+i}^{(1/2)}, \epsilon_{n+i}^{(1/2)}) = (2, 1).$$

On the other hand, $x_n^{(\alpha)} \in [1 - t_{i-1}, 1 - t_i)$, which implies

$$x_{n+1}^{(\alpha)} = 2 - \frac{1}{x_n^{(\alpha)}} \in \left(\frac{r_{i-1}}{s_{i-1}}, \frac{r_i}{s_i}\right] \text{ and } (a_{n+1}^{(\alpha)}, \epsilon_{n+1}^{(\alpha)}) = (2, -1),$$

$$x_{n+j}^{(\alpha)} = 3 - \frac{1}{x_{n+j-1}^{(\alpha)}} \in \left(\frac{r_{i-j}}{s_{i-j}}, \frac{r_{i-j+1}}{s_{i-j+1}}\right] \text{ and } (a_{n+j}^{(\alpha)}, \epsilon_{n+j}^{(\alpha)}) = (3, -1) \text{ for } 2 \le j \le i.$$

Then,

$$q_{n+j}^{(1/2)} - q_{n+j}^{(\alpha)} = q_{n+j-1}^{(1/2)}, \text{ for } 1 \le j \le i-1, \text{ and } q_{n+i}^{(1/2)} = q_{n+i}^{(\alpha)}.$$

In the following for $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{Z})$, we denote by $g.z = \frac{az+b}{cz+d}$ the Möbius transform applied on z. With this notation we have

$$x_{n+i}^{(1/2)} = \begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}^{i-1} . x_n^{(1/2)} \text{ and } x_{n+i}^{(\alpha)} = \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}^{i-1} \begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix} . x_n^{(\alpha)}.$$

Since
$$\begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} -1 & 1 \\ -1 & 2 \end{pmatrix}$$
,

$$x_{n+i}^{(1/2)} = \begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}^{i-1} \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}^{i-1} \begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix} \end{pmatrix}^{-1} .x_{n+i}^{(\alpha)}$$
$$= \begin{pmatrix} -2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -1 & 2 \end{pmatrix} .x_{n+i}^{(\alpha)} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} .x_{n+i}^{(\alpha)}.$$

It means that $\frac{1}{x_n^{(1/2)}} = \frac{1}{x_n^{(\alpha)}} - 1$, thus, for $0 \le c_1, c_2 \le 1$ and $k \in \mathbb{N}$ such that $k \ge 2$,

$$x_{n+i}^{(1/2)} \in \left(\frac{1}{k+c_1}, \frac{1}{k+c_2}\right] \text{ if and only if } x_{n+i}^{(\alpha)} \in \left(\frac{1}{k+1+c_1}, \frac{1}{k+1+c_2}\right].$$

If
$$x_{n+i}^{(1/2)} \in \left(\frac{1}{k+1}, \frac{1}{k+\alpha}\right] \cup \left(\frac{1}{k+1/2}, \frac{1}{k+1}\right]$$
, then

$$x_{n+i+1}^{(1/2)} = x_{n+i+1}^{(\alpha)}, \ a_{n+i+1}^{(1/2)} = a_{n+i+1}^{(\alpha)} - 1, \ \epsilon_{n+i+1}^{(1/2)} = \epsilon_{n+i+1}^{(\alpha)} \ \text{ and } q_{n+i+1}^{(1/2)} = q_{n+i+1}^{(\alpha)}.$$

If
$$x_{n+i}^{(1/2)} \in \left(\frac{1}{k+\alpha}, \frac{1}{k+1/2}\right]$$
, then

$$x_{n+i+1}^{(1/2)} = 1 - x_{n+i+1}^{(\alpha)}, \ a_{n+i+1}^{(1/2)} = a_{n+i+1}^{(\alpha)}, \ \epsilon_{n+i+1}^{(1/2)} = -1, \ \epsilon_{n+i+1}^{(\alpha)} = 1 \ \text{and}$$

$$q_{n+i+1}^{(1/2)} - q_{n+i+1}^{(\alpha)} = q_{n+i}^{(1/2)}.$$

Thus, by repeating the above process, we conclude that

$$q_i^{(1/2)} - q_i^{(\alpha)} = 0 \text{ or } q_{i-1}^{(1/2)}$$

and if
$$q_j^{(1/2)} - q_j^{(\alpha)} = q_{j-1}^{(1/2)}$$
, then $(a_{j+1}^{(1/2)}, \epsilon_{j+1}^{(1/2)}) = (3, -1)$ or $(2, 1)$.
Since $a_j^{(1/2)} \ge 2$ and $a_j^{(1/2)} = 2$ implies $\epsilon_j^{(1/2)} = 1$, we have

$$q_j^{(\alpha)} \ge q_j^{(1/2)} - q_{j-1}^{(1/2)} \ge q_{j-1}^{(1/2)}$$
 (2.5)

Hence,

$$\left| \frac{1}{q_j^{(1/2)}} - \frac{1}{q_j^{(\alpha)}} \right| = 0 \quad \text{or} \quad \left| \frac{1}{q_j^{(1/2)}} - \frac{1}{q_j^{(\alpha)}} \right| = \frac{q_{j-1}^{(1/2)}}{q_j^{(1/2)} q_j^{(\alpha)}} \le \frac{1}{q_j^{(1/2)}} \le \frac{4}{q_{j+1}^{(1/2)}}. \quad (2.6)$$

On the other hand, (2.5) also implies

$$\left|\log q_{j+1}^{(1/2)} - \log q_{j+1}^{(\alpha)}\right| = \log\left(1 + \frac{q_{j+1}^{(1/2)} - q_{j+1}^{(\alpha)}}{q_{j+1}^{(\alpha)}}\right) \le \log\left(1 + \frac{q_{j}^{(1/2)}}{q_{j+1}^{(\alpha)}}\right) \le \log 2.$$

Combining this with (2.6) and Remark 2.6 yields

$$\left| \sum_{j=0}^{\infty} (-1)^{j} \frac{\log q_{j+1}^{(1/2)}}{q_{j}^{(1/2)}} - \sum_{j=0}^{\infty} (-1)^{j} \frac{\log q_{j+1}^{(\alpha)}}{q_{j}^{(\alpha)}} \right|$$

$$\leq \sum_{j=0}^{\infty} \left| \frac{\log q_{j+1}^{(1/2)}}{q_{j}^{(1/2)}} - \frac{\log q_{j+1}^{(\alpha)}}{q_{j}^{(1/2)}} \right| + \sum_{j=0}^{\infty} \left| \frac{\log q_{j+1}^{(\alpha)}}{q_{j}^{(1/2)}} - \frac{\log q_{j+1}^{(\alpha)}}{q_{j}^{(\alpha)}} \right|$$

$$\leq \sum_{j=0}^{\infty} \frac{\log 2}{q_{j}^{(1/2)}} + \sum_{j=0}^{\infty} \frac{4 \log q_{j+1}^{(1/2)}}{q_{j+1}^{(1/2)}} \leq 4c_{1} + c_{2}.$$

For $x \in (\alpha, 1]$, the values of $W_{1/2}(x)$ and $W_{\alpha}(x)$ are defined symmetrically as

$$W_{1/2}(x) = W_{1/2}(1-x)$$
 and $W_{\alpha}(x) = W_{\alpha}(1-x)$.

By Remark 2.7, $W_{1/2} - W_{\alpha}$ is bounded on $[\alpha, 1]$.

For $x \in (1/2, \alpha]$, $1 - x \in [1 - \alpha, 1/2)$. Since $W_{1/2}(x) = W_{1/2}(1 - x)$, we can consider that n = 0 as in (2.4).

Proof of Theorem 2.3. As we already observed $1 - S^{(1/2)}$ is invertible in X_* , which together with the fact that the \mathbb{Z} -periodic even function equal to $-\log x$ on (0, 1/2] is in X_* implies that $W_{1/2}$ is BMO. By Proposition 2.9, W_{α} is BMO for $\alpha \in [1/2, g]$.

Finally, we give a remark which difficulties occur if one wants to extend the results of Proposition 2.3 to $\alpha \in (g, 1)$.

Remark 2.10. For the case $\alpha > g$, we cannot directly apply the same argument as in the proof of Proposition 2.9. If $\alpha > g$, then A_{α} has a branch which is defined by 1/x - 1 (see Figure 4 for the graph of A_{α}) contrary to the case of $\alpha \leq g$. It causes a different behaviour of the orbits of the points under A_{α} . In the proof, we showed a relation between $x_n^{(1/2)}$ and $x_n^{(\alpha)}$. By following the same argument for $\alpha > g$, we only obtain a relation between $x_n^{(1/2)}$ and $x_{n+N}^{(\alpha)}$, where N depends on the number of consecutive points in the orbit of x visiting $\left[\frac{1}{1+\alpha},\alpha\right]$.




FIGURE 4. Graph of A_{α} when $\alpha > g$.

3. Behaviour of the truncated real Brjuno function and the truncated real Wilton function

For $x \in \mathbb{R}$, recall that $\beta_{-1} = 1$ and

$$\beta_i(x) = \{x\} G(\{x\}) \cdots G^j(\{x\}) = |p_i(\{x\}) - q_i(\{x\}) \{x\}| \quad \text{for } j \ge 0,$$

where G is the Gauss map and $p_j(\lbrace x\rbrace)/q_j(\lbrace x\rbrace)$ is the jth principal convergent of $\lbrace x\rbrace$ with respect to the regular continued fraction algorithm. Here, contrary to the previous section, we omit the α in $\beta_i^{(\alpha)}$ as we will always assume α to be one.

In this section, we are interested to compare a finite k-Brjuno sum or finite Wilton sum with the k-Brjuno or Wilton of its principal convergent. For doing so we first have to define the finite k-Brjuno or finite Wilton function respectively for a rational number.

Within this section, let p_j/q_j be a rational number whose continued fraction algorithm terminates after r steps, i.e. it can be written as

$$p_j/q_j = m_0 + \frac{1}{m_1 + \frac{1}{m_2 + \cdots + \frac{1}{m_r}}}$$

with $m_r \geq 2$ when $q_j > 1$. (Of course this can correspond to the rth principal convergent of a number whose continued fraction expansion starts with $[m_0; m_1, \ldots, m_r, \ldots]$.) With this we can define the truncated real Brjuno function by

$$B_{k,\text{finite}}(p/q) = \sum_{j=0}^{r-1} (\beta_{j-1}(p/q - m_0))^k \log \left(\frac{1}{G^j(p/q - m_0)}\right)$$

and the truncated real Wilton function by

$$W_{\text{finite}}(p/q) = \sum_{j=0}^{r-1} (-1)^j \beta_{j-1}(p/q - m_0) \log \left(\frac{1}{G^j(p/q - m_0)} \right).$$

Before stating the results of this section, we also introduce the notation $x_j = G^j(x)$ for $x \in (0,1)$. This enables us to state the next lemma which is an analog to [MMY01, Lem. 5.20].

Lemma 3.1. For each $k \in \mathbb{N}$ there exists $C_k > 0$ such that for all $x \in (0,1)$ and $r \in \mathbb{N}$, we have

$$\left| B_{k,\text{finite}} \left(\frac{p_r(x)}{q_r(x)} \right) - \sum_{j=0}^{r-1} (\beta_{j-1}(x))^k \log \frac{1}{x_j} \right| \le C_k x_r (q_r(x))^{-1}$$

and

$$\left| W_{\text{finite}} \left(\frac{p_r(x)}{q_r(x)} \right) - \sum_{j=0}^{r-1} (-1)^j \beta_{j-1}(x) \log \frac{1}{x_j} \right| \le C_1 x_r (q_r(x))^{-1}.$$

Proof. To ease notation we write in the following p_r and q_r instead of $p_r(x)$ and $q_r(x)$ when the dependence on x is clear. If r = 1, then

$$|B_{k,\text{finite}}(p_1/q_1) - \log(1/x)| = |W_{\text{finite}}(p_1/q_1) - \log(1/x)| = |\log(q_1/p_1) - \log(1/x)|$$
$$= |\log(m_1) - \log(m_1 + x_1)| = \log(1 + x_1/m_1) \le x_1/m_1.$$

Suppose that r > 1, then we have

$$\left| \sum_{j=0}^{r-1} \left[(\beta_{j-1}(p_r/q_r))^k \log \frac{1}{G^j(p_r/q_r)} - (\beta_{j-1}(x))^k \log \frac{1}{x_j} \right] \right|$$

$$\leq \sum_{j=0}^{r-1} \left| (\beta_{j-1}(p_r/q_r))^k \left[\log \frac{1}{G^j(p_r/q_r)} - \log \frac{1}{x_j} \right] \right|$$

$$+ \sum_{j=0}^{r-1} \left| \left[(\beta_{j-1}(p_r/q_r))^k - (\beta_{j-1}(x))^k \right] \log \frac{1}{x_j} \right|$$
(3.1)

and similarly for the Wilton case if k=1.

Note that

$$\beta_{j-1}(p_r/q_r) = \left| q_{j-1} \frac{p_r}{q_r} - p_{j-1} \right| \tag{3.2}$$

for $r \leq j-1$. Since x_j , $G^j(p_r/q_r) \in \left[\frac{1}{m_{j+1}+1}, \frac{1}{m_{j+1}}\right]$, we have $2^{-1} \leq \frac{x_j}{G^j(p_r/q_r)} \leq 2$. By using (3.2) and the fact that $\log(y_2/y_1) \leq (y_2 - y_1)y_1^{-1}$ for $y_1 < y_2 \in \mathbb{R}_{>0}$, we have

$$\left|\log \frac{1}{G^j(p_r/q_r)} - \log \frac{1}{x_j}\right| \le \max\left\{G^j(p_r/q_r), x_j\right\} \left|\frac{1}{G^j(p_r/q_r)} - \frac{1}{x_j}\right|$$

$$\leq 2 G^{j}(p_r/q_r) \frac{|x - p_r/q_r|}{|q_j \frac{p_r}{q_r} - p_j| |q_j x - p_j|} \leq \frac{2|x - p_r/q_r|}{|q_{j-1} \frac{p_r}{q_r} - p_{j-1}| |q_j x - p_j|}.$$

Furthermore, we note

$$x = \frac{p_{i-1}x_i + p_i}{q_{i-1}x_i + q_i} \quad \text{and} \quad x_i = -\frac{q_i x - p_i}{q_{i-1}x - p_{i-1}}$$
(3.3)

which imply $\prod_{i=0}^{\ell} (-x_i) = q_{\ell}x - p_{\ell}$ and

$$|p_{\ell} - q_{\ell}x| = \left|p_{\ell} - q_{\ell} \frac{p_{\ell}x_{\ell+1} + p_{\ell+1}}{q_{\ell}x_{\ell+1} + q_{\ell+1}}\right| = \frac{|p_{\ell}q_{\ell+1} - q_{\ell}p_{\ell+1}|}{|q_{\ell}x_{\ell+1} + q_{\ell+1}|} < \frac{1}{q_{\ell+1}}.$$
 (3.4)

Hence, for the first summand of (3.1), from (3.2), (3.3) and (3.4), we have

$$\sum_{j=0}^{r-1} \left| \left(q_{j-1} \frac{p_r}{q_r} - p_{j-1} \right)^k \left[\log \frac{1}{G^j(p_r/q_r)} - \log \frac{1}{x_j} \right] \right| \\
\leq \frac{2}{q_r} \sum_{j=0}^{r-1} \left| q_{j-1} \frac{p_r}{q_r} - p_{j-1} \right|^{k-1} \left| \frac{p_r - xq_r}{q_j x - p_j} \right| \leq \frac{2x_r}{q_r} \sum_{j=0}^{r-1} x_{j+1} \cdots x_{r-1} \\
\leq \frac{2x_r}{q_r} \sum_{j=0}^{r-1} \left(\frac{\sqrt{5} - 1}{2} \right)^{r-j-1} < 2C' x_r q_r^{-1}, \quad \text{(by Proposition 1.4-(iv) in [MMY97])},$$

where $C' = \sum_{j=0}^{\infty} \left(\frac{\sqrt{5}-1}{2}\right)^j = \frac{\sqrt{5}+3}{2}$. On the other hand, by setting $X_{j,r} = q_{j-1} \frac{p_r}{q_r} - p_{j-1}$ and $Y_{j,r} = q_{j-1}x - p_{j-1}$, noting that $|X_{j,r}| \leq 1/q_{j-1}$ and $|Y_{j,r}| \leq 1/q_{j-1}$ we obtain for the second term of (3.1) that

$$\sum_{j=0}^{r-1} \left| \left(q_{j-1} \frac{p_r}{q_r} - p_{j-1} \right)^k - \left(q_{j-1} x - p_{j-1} \right)^k \right| \log \frac{1}{x_j}$$

$$= \sum_{j=0}^{r-1} \left| X_{j,r} - Y_{j,r} \right| \left| X_{j,r}^{k-1} + X_{j,r}^{k-2} Y + \dots + X_{j,r} Y_{j,r}^{k-2} + Y_{j,r}^{k-1} \right| \log \frac{1}{x_j}$$

$$\leq \sum_{j=0}^{r-1} \frac{k}{q_{j-1}^{k-2}} \left| p_r / q_r - x \right| \log \frac{1}{x_j}$$

$$\leq \sum_{j=0}^{r-1} k q_{j-1} \left| p_r / q_r - x \right| \log \frac{1}{x_j} .$$
(3.6)

Since $\log(y) < y$ and using (3.4) and (3.3), the value in (3.6) is bounded above by

$$\frac{k}{q_r} \sum_{j=0}^{r-1} \frac{|p_r - xq_r|}{|q_{j-1}x - p_{j-1}|} \frac{1}{x_j} = \frac{k}{q_r} \sum_{j=0}^{r-1} x_{j+1} \cdots x_{r-1} x_r \le kC' x_r q_r^{-1}.$$

By letting $C_k := 2kC'$, we complete the proof.

4. Complex continued fractions


We consider a continued fraction on a compact subset of \mathbb{C} which is a complex analog to the regular continued fractions. At the beginning of this section we will first define some domains which will be important to define our complex continued fraction algorithm which we introduce in the sequel.

Similarly as in [MMY01], we consider the following sets:

$$D_0 = \left\{ z \in \mathbb{C} \mid |z + 1| \le 1, \text{Re}(z) \ge \frac{\sqrt{3}}{2} - 1 \right\}$$
 (Figure 5a), (4.1)

$$D_1 = \left\{ z \in \mathbb{C} \mid |z| \ge 1, \left| z - \frac{\sqrt{3}}{3} \right| \le \frac{\sqrt{3}}{3} \right\}$$
 (Figure 5a), (4.2)

$$D = \{ z \in \mathbb{C} \mid |z| \le 1, |z - i| \ge 1, |z + i| \ge 1, \text{Re}(z) > 0 \}$$
 (Figure 5b and 8), (4.3)




$$H_0 = \left\{ z \in \mathbb{C} \mid |z - i| \le 1, |z + 1| \ge 1, |\text{Im}(z)| \le \frac{1}{2} \right\}$$
 (Figure 6a), (4.4)

$$H'_{0} = \left\{ z \in \mathbb{C} \mid |z+i| \le 1, |z+1| \ge 1, |\text{Im}(z)| \le \frac{1}{2} \right\}$$
 (Figure 6a), (4.5)

$$H = H_0 \cup H_0'$$
 (Figure 6a), (4.6)


$$\Delta = D \cup H_0 \cup H_0' = \left\{ z \in \mathbb{C} \mid |z| \le 1, |z+1| \ge 1, |\text{Im}(z)| \le \frac{1}{2} \right\} \quad \text{(Figure 6b)}, \quad (4.7)$$




$$D_{\infty} = \overline{\mathbb{C}} \setminus (D_0 \cup \Delta \cup D)$$

$$= \left\{ z \in \overline{\mathbb{C}} \mid |\operatorname{Im}(z)| > \frac{1}{2} \right\} \cup \left\{ z \in \overline{\mathbb{C}} \mid \operatorname{Re}(z) > \frac{\sqrt{3}}{2} - 1 \right\}$$

$$\cup \left\{ z \in \overline{\mathbb{C}} \mid \operatorname{Re}(z) > \frac{\sqrt{3}}{2}, \ \left| z - \frac{\sqrt{3}}{3} \right| > \frac{\sqrt{3}}{3} \right\}$$
 (Figure 7a). (4.8)





(7b) $\iota(D_{\infty})$ where $\iota(z) = 1/z$.

We define an extension of the continued fraction to the complex plane as follows. Let $z \in D$ as in Figure 5b. Then $1/z \in \bigcup_{n \in \mathbb{N}} (n + \Delta)$, see Figure 6a. If $1/z \in n + \Delta$, then we take $m_1 = n$ and we set $z_1 := 1/z - m_1$. If $z_1 \in \Delta - D$, then we finish the process. If $z_1 \in D$, then we define m_2 by an integer n such that $1/z_1 \in n + \Delta$. Repeating this process, we obtain a continued fraction expansion $\{m_i\}_{i=1}^r$ such that

$$\frac{1}{z_i} = m_{i+1} + z_{i+1} \quad \text{for all } 0 < i \le r, \tag{4.9}$$

such that $z_i \in D$ for i < r, and $z_r \in \Delta$. Let $D(m_1, \dots, m_r)$ be the set of $z_0 \in D$ whose first r complex continued fraction entries equal $\{m_i\}_{i=1}^r$.

Note that this continued fraction algorithm does not coincide with Hurwitz' continued fractions, which are often used on the complex plane, but which are expanded with Gaussian integers [Hur87], see also [Hen06]. The goal here is very different: our complex continued fraction expansion follows as closely as possible the standard continued fraction of neighboring real points. For non-real numbers the iteration of our algorithm leads to an increasing sequence of imaginary parts and it stops after finitely many steps, since "it is not possible anymore to associate a neighboring real point". We will see in the following section why this continued fraction definition makes sense.

For $z \in \Delta$ and $i \geq 1$, let

$$\varepsilon_i = \begin{cases} 0 & \text{if } z \in D(m_1, \dots, m_i) \text{ and } m_i = 1, \\ 1 & \text{otherwise.} \end{cases}$$
 (4.10)

We note here that the just defined ε_i is independent to the definition of ϵ_i defined for the α -continued fractions. However, since in the following sections we only consider the regular continued fraction algorithm with the Gauss map G we do not expect too much confusion for the reader.

We define p_{ℓ}/q_{ℓ} by

$$\frac{p_{\ell}}{q_{\ell}} = \frac{1}{m_1 + \frac{1}{m_2 + \frac{1}{\cdots + \frac{1}{m_{\ell}}}}}$$

and $q_0 = p_{-1} = q_{-2} = 1$ and $p_0 = p_{-2} = q_{-1} = 0$. Then,

$$z_0 = \frac{p_{i-1}z_i + p_i}{q_{i-1}z_i + q_i} \quad \text{and} \quad z_i = -\frac{q_i z_0 - p_i}{q_{i-1}z_0 - p_{i-1}}.$$
 (4.11)

By (4.11), we can easily see that

$$\prod_{i=0}^{\ell} (-z_i) = q_{\ell} z_0 - p_{\ell}. \tag{4.12}$$




FIGURE 8. The sets $D(m_1, \dots, m_r)$. The left figure is the partition of D by D(n). The right figure is of the sets D(1, n) in D(1).

We have

$$p_{\ell} - q_{\ell} z_0 = p_{\ell} - q_{\ell} \frac{p_{\ell} z_{\ell+1} + p_{\ell+1}}{q_{\ell} z_{\ell+1} + q_{\ell+1}} = \frac{p_{\ell} q_{\ell+1} - q_{\ell} p_{\ell+1}}{q_{\ell} z_{\ell+1} + q_{\ell+1}} = \frac{(-1)^{\ell+1}}{q_{\ell} z_{\ell+1} + q_{\ell+1}}.$$
 (4.13)

Thus, we have

$$|p_{\ell} - q_{\ell} z_0| < \frac{1}{q_{\ell+1}}. (4.14)$$

We set

$$H(m_1, \cdots, m_r) = D(m_1, \cdots, m_r) \setminus \operatorname{int} \left(\bigcup_{m_{r+1} \ge 1} D(m_1, \cdots, m_{r+1}) \right)$$
(4.15)

where it is defined by H when r=0, see (4.6) for the definition of H. The sets $H(m_1, \dots, m_r)$ give a partition of D as in Figure 9. Then we have

$$\{z: |\mathrm{Im}z| \leq 1/2\} = \bigcup_{n \in \mathbb{Z}} \bigcup_{r>0} \bigcup_{m_1, \dots, m_r > 1} [H(m_1, \dots, m_r) + n] \sqcup \mathbb{R} \setminus \mathbb{Q},$$

where the sets in the right-hand term have disjoint interiors. A set $H(m_1, \dots, m_r) + n$ meets \mathbb{R} in a unique point which is rational.




FIGURE 9. The partition of D by the sets $H(m_1, \dots, m_r)$.

5. Complexification of the operators T_k and S

Let J be a closed interval and $k \in \mathbb{N}$. Let $\mathcal{O}^k(\overline{\mathbb{C}} \setminus J)$ be the complex vector space of holomorphic functions in $\mathbb{C} \setminus J$, meromorphic in $\overline{\mathbb{C}} \setminus J$ with a zero at infinity of order at least k. There exists $C_{V,k} > 0$ such that for each $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus J)$ and each neighbourhood V of J we have

$$|\varphi(z)| \le C_{V,k}|z|^{-k} \sup_{\mathbb{C}\setminus \overline{V}} |\varphi| \quad \text{for } z \in C \setminus \overline{V}.$$
 (5.1)

This fact can be easily proven in the following way. We obtain that the function $\widetilde{\varphi}$ defined as $\widetilde{\varphi}(w) = \varphi(w) \cdot w^k$ is still analytic on $\mathbb{C} \setminus \overline{V}$. By the maximum principle, we obtain that both functions $\widetilde{\varphi}$ and φ attain their maximum on the boundary of $\mathbb{C} \setminus \overline{V}$. Thus, we obtain for all $w \in \mathbb{C} \setminus \overline{V}$

$$|\varphi(w)| = \frac{\left|\tilde{\varphi}(w)\right|}{|w|^k} \le \frac{\sup_{\mathbb{C}\setminus\overline{V}}|\tilde{\varphi}|}{|w|^k} = \frac{\sup_{\partial(\mathbb{C}\setminus\overline{V})}|\tilde{\varphi}|}{|w|^k} \le \frac{C_{V,k}\sup_{\mathbb{C}\setminus\overline{V}}|\varphi|}{|w|^k}$$
(5.2)

with $C_{V,k} = \sup_{x \in \partial(\mathbb{C}\setminus \overline{V})} |x|^k$.

Let $\mathcal{O}^{-k}(\overline{\mathbb{C}}\setminus J)$ be the complex vector space of holomorphic functions in $\mathbb{C}\setminus J$, meromorphic in $\overline{\mathbb{C}}\setminus J$ with a pole at infinity of order at most k.

5.1. Hyperfunctions and extensions to spaces of complex analytic functions of the operators T_k and S. Recall that $T_k := T_{k,\alpha}$ as in (1.6) for given $\alpha \in [1/2, 1]$ and $S := S_{\alpha}$. We want to extend T_k and S for $\alpha = 1$ to the space of complex analytic functions.

We proceed as in [MMY01], namely we extend T_k and S to A'([0,1]) the space of hyperfunctions with support contained in [0,1] for which we will first introduce some definitions. In Proposition 6.1, we will give the definition of these operators and prove that they are indeed well defined. In Proposition 6.6, we will see that this definition makes sense in terms of T_k and S being a complex extension of T_k and S being studied in the previous sections.

5.1.1. Hyperfunctions. Let K be a non-empty compact set $K \subset \mathbb{R}$. Let us denote by $\mathcal{O}(K)$ the space of functions analytic in a neighbourhood V of K. A hyperfunction with support in K is a linear functional u on $\mathcal{O}(K)$, such that for all neighbourhoods V of K, there is a constant $C_V > 0$ such that

$$|u(\varphi)| \le C_V \sup_V |\varphi|, \quad \forall \varphi \in \mathcal{O}(V).$$

We denote by A'(K) the space of hyperfunctions with support in K.

The space A'([0,1]) is canonically isomorphic to $\mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ the complex vector space of holomorphic functions on $\overline{\mathbb{C}} \setminus [0,1]$ vanishing at infinity. Let

$$c_z(x) = \frac{1}{\pi(x-z)}. (5.3)$$

Given $u \in A'([0,1])$, the corresponding $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ is obtained by

$$\varphi(z) = u(c_z), \quad \forall z \in \mathbb{C} \setminus [0, 1].$$
 (5.4)

On the other hand, for every $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, the corresponding $u \in A'([0,1])$ is given by

$$u(\psi) = \frac{i}{2} \int_{\gamma} \varphi(z)\psi(z)dz, \quad \forall \psi \in \mathcal{O}(V),$$
 (5.5)

where V is a complex neighbourhood of [0,1] and γ is any piecewise C^1 path winding around [0,1] in the positive direction. That u as in (5.5) and c_z as in (5.3) are inverse to each other can be easily seen by using the substitution $\xi = 1/\omega$ resulting in $d\omega = -\xi^{-2}d\xi$ which gives

$$u(c_z) = \frac{i}{2} \int_{\gamma} \varphi(\omega) c_z(\omega) d\omega = \frac{i}{2} \int_{\gamma} \varphi(\omega) \frac{1}{\pi} \frac{1}{\omega - z} d\omega = \frac{i}{2\pi} \int_{1/\gamma} \frac{\varphi(1/\xi)}{(1/\xi - z)(-\xi^2)} d\xi$$
$$= \frac{1}{z} \int_{1/\gamma} \frac{i}{2\pi} \frac{\varphi(\xi^{-1})\xi^{-1}}{\xi - 1/z} d\xi = \frac{1}{z} \varphi(z) \left(\frac{1}{z}\right)^{-1} = \varphi(z),$$

where we denote by $1/\gamma$ the transformed path and notice that $1/\gamma$ goes along the negative direction. We remark that in (5.5) we correct the formula for u given in [MMY01, Appendix 2] where it is given with an additional factor π . We write $u(x) = \frac{1}{2i}(\varphi(x+i0) - \varphi(x-i0))$ (see [MMY01, Appendix 2] for details).

5.1.2. Formulas of the extensions of T_k and S. We obviously have $S = -T_1$. Hence, for the following we will introduce the main formulas for T_k only and give explanations where differences occur for S. For $k \in \mathbb{N}$, $m \in \mathbb{N}^*$ and $f \in L^2([0,1])$, let us consider

$$T_{k,m}f(x) = \begin{cases} x^k f\left(\frac{1}{x} - m\right), & \text{if } x \in \left[\frac{1}{m+1}, \frac{1}{m}\right], \\ 0, & \text{otherwise.} \end{cases}$$
 (5.6)

We then have $T_k = \sum_{m=1}^{\infty} T_{k,m}$. For φ , $\psi \in L^2([0,1])$, we define the L^2 -adjoint $T_{k,m}^*$ by

$$\int_0^1 T_{k,m}\varphi(x)\psi(x)dx = \int_0^1 \varphi(x)T_{k,m}^*\psi(x)dx.$$
 (5.7)

Therefore, we have

$$T_{k,m}^* \psi(x) = \frac{1}{(m+x)^{k+2}} \psi\left(\frac{1}{m+x}\right).$$
 (5.8)

By a slight generalisation of [MMY01, (1.10)], we have that if ψ is holomorphic in a neighbourhood V of [0,1], then $T_{k,m}^*\psi$ is holomorphic in V and we have

$$\sup_{V} |T_{k,m}^* \psi| \le \frac{1}{2m^{k+2}} \sup_{V} |\psi|,$$

where we may take $V = D_{\infty}^c$ equipped with the Poincaré metric on the hyperbolic Riemann surface $\overline{\mathbb{C}} \setminus [0,1]$. It follows that the series $\sum_{m=1}^{\infty} T_{k,m}u$ converges to a hyperfunction T_ku in A'([0,1]).

We have

$$T_{k,m}^* c_z(x) = -z^k \left(c_{\frac{1}{z}-m}(x) - c_{-m}(x) \right) + \sum_{n=1}^k \frac{z^{k-n}}{n!} \cdot \frac{\partial^n}{\partial z^n} c_z(x) \Big|_{z=-m},$$

where c_z is as in (5.3). For $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, let u be the corresponding hyperfunction in A'([0,1]) as in (5.4). Then, $T_{k,m}\varphi$ is defined by $(T_{k,m}u)(c_z)$. If $z \notin [\frac{1}{m+1}, \frac{1}{m}]$, then we have

$$(T_{k,m}u)(c_z) = u(T_{k,m}^*c_z).$$

This follows from (5.7) and the fact that for $z \notin [\frac{1}{m+1}, \frac{1}{m}]$ we have that $T_{k,m}u$ and $T_{k,m}^*c_z$ are analytic. Hence, (5.7) already implies the equality.

By (5.4) and (5.5), we have

$$u(T_{k,m}^*c_z) = -z^k \left(u(c_{\frac{1}{z}-m}(x)) - u(c_{-m}(x)) \right) + \sum_{n=1}^k \frac{z^{k-n}}{n!} u\left(\frac{\partial^n}{\partial z^n} c_z(x)|_{z=-m} \right)$$
$$= -z^k \left(\varphi\left(\frac{1}{z}-m\right) - \varphi(-m) \right) + \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}(-m).$$

For $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, the formula for T_k is given by

$$T_k \varphi(z) = -\sum_{m=1}^{\infty} z^k \left(\varphi\left(\frac{1}{z} - m\right) - \varphi(-m) \right) + \sum_{m=1}^{\infty} \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}(-m), \tag{5.9}$$

which will be shown in Proposition 6.1. By using $S = -T_1$, we can also deduce that a natural extension of S to the space of complex analytic functions is

$$S\varphi(z) = \sum_{m=1}^{\infty} z \left(\varphi\left(\frac{1}{z} - m\right) - \varphi(-m) \right) - \sum_{m=1}^{\infty} \varphi'(-m).$$
 (5.10)

5.1.3. Algebraic properties of the inversion of $(1-T_k)$ and (1-S), the monoid \mathcal{M} and its actions. Let us consider the monoid

$$\mathcal{M} = \left\{ g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{GL}_2(\mathbb{Z}) \mid d \ge b \ge a \ge 0 \text{ and } d \ge c \ge a \ge 0 \right\} \cup \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

which is generated by $g(m) := \begin{pmatrix} 0 & 1 \\ 1 & m \end{pmatrix}$ for $m \in \mathbb{N}$. For $g \in \mathcal{M}$ and $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, we define

$$L_{k,g}\varphi(z) := \det(g)^{k+1} \left[(a - cz)^k \varphi\left(\frac{dz - b}{a - cz}\right) - \sum_{n=0}^k (a - cz)^{k-n} \frac{\det(g)^n}{c^n n!} \varphi^{(n)}\left(-\frac{d}{c}\right) \right]. \tag{5.11}$$

and

$$\overline{L}_{g}\varphi(z) := \det(g)(a - cz) \left(\varphi\left(\frac{dz - b}{a - cz}\right) - \varphi\left(-\frac{d}{c}\right) \right) - \frac{1}{c}\varphi'\left(-\frac{d}{c}\right) = \det(g)L_{1,g}\varphi(z). \tag{5.12}$$

As we will see in Proposition 6.6, we have

$$(1 - T_k)^{-1}\varphi(z) = \sum_{j=0}^{\infty} T_k^j \varphi(z) = \sum_{g \in \mathcal{M}} L_{k,g}\varphi(z), \qquad (5.13)$$

and from analogous arguments we will see that

$$(1-S)^{-1}\varphi(z) = \sum_{j=0}^{\infty} S^{j}\varphi(z) = \sum_{g \in \mathcal{M}} \overline{L}_{g}\varphi(z).$$
 (5.14)

For $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $n \in \mathbb{Z}$, let

$$L_g^{(n)}\varphi(z) := \frac{\det(g)^{n+1}}{(a-cz)^n}\varphi\left(\frac{dz-b}{a-cz}\right)$$
(5.15)

and

$$\overline{L}_g^{(n)}\varphi(z) := \frac{\det(g)^n}{(a-cz)^n}\varphi\left(\frac{dz-b}{a-cz}\right) = \det(g)L_g^{(n)}\varphi(z). \tag{5.16}$$

Analogously to [MMY01, (2.2)], we are interested in a connection between $L_{k,g}$ and $L_g^{(k+2)}$ and between \overline{L}_g and $\overline{L}_g^{(3)}$ respectively. In the first case, we obtain by induction that for $t \in \mathbb{N}^*$ and $n \in \mathbb{Z}$, we have

$$(L_g^{(n)}\varphi)^{(t)} = \det(g)^{n+1} \sum_{\ell=0}^{t-1} (n+\ell)(n+\ell+1) \cdots (n+t-1) \binom{t}{\ell} c^{t-\ell} \det(g)^{\ell} L_g^{(n+t+\ell)} \varphi^{(\ell)} + \det(g)^{t+n+1} L_g^{(n+2t)} \varphi^{(t)},$$

where $\sum_{\ell=0}^{-1}$ denotes an empty sum. In particular, for all $k \in \mathbb{N}^*$, we have

$$(L_q^{(-k)}\varphi)^{(k+1)} = L_q^{(k+2)}\varphi^{(k+1)}. (5.17)$$

5.1.4. Polynomial corrections to the actions of \mathcal{M} . If $\varphi \in \mathcal{O}^{-k}(\overline{\mathbb{C}} \setminus [0,1])$, then it can be uniquely written as

$$\varphi(z) = \xi_k z^k + \xi_{k-1} z^{k-1} + \dots + \xi_0 + p_k(\varphi)(z), \tag{5.18}$$

with $p_k(\varphi) \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$. Note that if $\varphi \in \mathcal{O}^{-k}(\overline{\mathbb{C}} \setminus [0,1])$, then $\varphi^{(k+1)} = p_k(\varphi)^{(k+1)}$. If $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{Z})$ and $\psi(z) = \xi_k z^k + \xi_{k-1} z^{k-1} + \cdots + \xi_0$, then

$$L_q^{(-k)}\psi(z) = \det(g)^{1-k} \left(\xi_k (dz - b)^k + \xi_{k-1} (dz - b)^{k-1} (a - cz) + \dots + \xi_0 (a - cz)^k \right), \quad (5.19)$$

which means that $p_k(L_g^{(-k)}\psi) = 0$. Thus, p_k is a projection from $\mathcal{O}^{-k}(\overline{\mathbb{C}} \setminus [0,1])$ to $\mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$. We have the formula

$$L_{k,g}\varphi = p_k(L_g^{(-k)}\varphi) \tag{5.20}$$

for $g \in \mathcal{M}$, $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ which defines an action of \mathcal{M} on $\mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, where p_k depends on k. To verify this formula we define $\psi(u) := (L_g^{(-k)}\varphi)(\frac{a}{c} - \frac{1}{cu})$. Since $\lim_{z\to\infty} p_k(L_g^{(-k)}\varphi)(z) = 0$ is equivalent to $\lim_{u\to 0} \psi(u) = 0$, it is enough to look at a projection of ψ . We have

$$\psi(u) = \det(g)^{1-k} u^{-k} \varphi\left(\frac{u \det(g) - d}{c}\right)$$

and if we denote by \widetilde{p}_k the projection to a function vanishing at 0, then we obtain

$$\widetilde{p}_{k}(\psi(u)) = \psi(u) - \det(g)^{1-k} u^{-k} \sum_{n=0}^{k} \frac{\psi^{(n)}(0) u^{n}}{n!}$$

$$= \psi(u) - \det(g)^{1-k} \sum_{n=0}^{k} \frac{\det(g)^{n} \varphi^{(n)}(-\frac{d}{c}) u^{n-k}}{c^{n} n!}.$$
(5.21)

Substituting back $u = (a - cz)^{-1}$ gives (5.20). From the definition of $L_{k,g}$ as in (5.11) and (5.19), the following diagram commutes:

$$\mathcal{O}^{-k}(\overline{\mathbb{C}}\setminus[0,1]) \xrightarrow{L_g^{(-k)}} \mathcal{O}^{-k}(\overline{\mathbb{C}}\setminus[0,1])$$

$$p_k \downarrow \qquad \qquad \downarrow p_k$$

$$\mathcal{O}^1(\overline{\mathbb{C}}\setminus[0,1]) \xrightarrow{L_{k,g}} \mathcal{O}^1(\overline{\mathbb{C}}\setminus[0,1])$$

By (5.11) and (5.12), the connection between \overline{L}_g and $\overline{L}_g^{(-1)}$ is similar as they only differ by a factor $\det(g)$ from $L_{1,g}$ and $L_g^{(-1)}$.

By the calculations in (5.21), the (k+1)th derivative of the difference between $L_{k,g}\varphi$ and $L_g^{(-k)}\varphi$ is zero for $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$. Thus, it follows from (5.17) that

$$(L_{k,g}\varphi)^{(k+1)}(\xi) = L_g^{(k+2)}\varphi^{(k+1)}(\xi) = \frac{\det(g)^{k+1}}{(a-c\xi)^{k+2}}\varphi^{(k+1)}(g^{-1}.\xi), \tag{5.22}$$

where $\varphi^{(k+1)} \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$. By Taylor's theorem, we have for every $\omega \in \overline{\mathbb{C}} \setminus [0,1]$ that

$$L_{k,g}\varphi(z) = L_{k,g}\varphi(\omega) + (L_{k,g}\varphi)'(\omega)(z-\omega) + \dots + \frac{(L_{k,g}\varphi)^{(k-1)}}{(k-1)!}(z-\omega)^{k-1} + \int_{\omega}^{z} \frac{(L_{k,g}\varphi)^{(k+1)}(\xi)}{k!}(z-\xi)^{k} d\xi.$$

When ω goes to ∞ , then only the remainder term will be left since $\frac{(L_{k,g}\varphi)^{(\ell)}(\omega)}{\ell!}(z-\omega)^{\ell} \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ for all $0 \leq \ell \leq k-1$. From (5.22), the remainder term can be written as

$$\frac{\det(g)^{k+1}}{k!} \int_{-\infty}^{z} \frac{(z-\xi)^{k}}{(a-c\xi)^{k+2}} \varphi^{(k+1)}(g^{-1}.\xi) d\xi.$$

By a change of variable with $\xi = \xi(t) = g \cdot \left(-\frac{d}{c} + \frac{\det(g)t}{c(a-cz)}\right)$, which means that $\frac{1}{(a-c\xi)^2}d\xi = \frac{1}{c(a-cz)}dt$ and $\xi = -\frac{(a-cz)}{ct} + \frac{a}{c}$, we then deduce that the remainder term equals

$$\frac{\det(g)^{k+1}}{k!} \int_0^1 \left(\frac{1-t}{c}\right)^k \frac{\varphi^{(k+1)}(-\frac{d}{c} + \frac{\det(g)t}{c(a-cz)})}{c(a-cz)} dt.$$

Then, for $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1]), g \in \mathcal{M}$ and $z \notin \left[\frac{b}{d}, \frac{a}{c}\right]$, we have

$$L_{k,g}\varphi(z) = \frac{c^{-(k+1)}\det(g)^{k+1}}{k!}(a-cz)^{-1}\int_0^1 (1-t)^k \varphi^{(k+1)}\left(-\frac{d}{c} + \frac{\det(g)t}{c(a-cz)}\right)dt.$$
 (5.23)

We then have for the special case g = g(m) that

$$L_{k,g(m)}\varphi(z) = -z^k \left(\varphi\left(\frac{1}{z} - m\right) - \varphi\left(-m\right)\right) + \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}\left(-m\right)$$
 (5.24)

$$= \frac{(-1)^k}{z \, k!} \int_0^1 \varphi^{(k+1)} \left(\frac{t}{z} - m\right) (1 - t)^k dt.$$
 (5.25)

Connected to that we note that by (5.22) $L_g^{(k+2)}$ can be represented as

$$L_g^{(k+2)}\psi(z) = \frac{\det(g)^{k+1}}{(a-cz)^{k+2}}\psi\left(\frac{dz-b}{a-cz}\right) = \left(L_{k,g}\psi^{(-(k+1))}\right)^{(k+1)}(z),\tag{5.26}$$

for $\psi \in \mathcal{O}^{-k}(\overline{\mathbb{C}} \setminus [0,1])$, where $\psi^{(-(k+1))}$ denotes the (k+1)th primitive (i.e. an integration between ∞ and z) of ψ . In the special case that g=g(m), we have that $\det(g(m))=-1$ and thus

$$L_{g(m)}^{(k+2)}\psi(z) = -z^{-(k+2)}\psi\left(\frac{1}{z} - m\right). \tag{5.27}$$

For $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ we can immediately conclude from (5.12), (5.24) and (5.25) that

$$\overline{L}_{g(m)}\varphi(z) = z\left(\varphi\left(\frac{1}{z} - m\right) - \varphi\left(-m\right)\right) - \varphi'\left(-m\right) = \frac{1}{z} \int_0^1 \varphi''\left(\frac{t}{z} - m\right) (1 - t) dt. \tag{5.28}$$

We also note that for simplicity we were restricting us to $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$. However, the same calculations of this section hold true if we consider instead $\gamma_1 > \gamma_0 > -1$, $I = [\gamma_0, \gamma_1]$, $z \notin \left[0, \frac{1}{\gamma_0 + m}\right]$, and $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus I)$.

6. Convergence of the sum over the monoid and boundary behaviour

In this Section we will show how to adapt the results of Sections 3 and 4 of [MMY01] to our monoid action. Also in our case the complex continued fraction introduced in Section 4 will play a fundamental role.

6.1. Convergence of the sum over the monoid. From the following proposition we will be able to deduce that T_k is given by (5.9). It is an analog to [MMY01, Prop. 3.1].

Let
$$-1 < \gamma_0 < \gamma_1$$
, $I = [\gamma_0, \gamma_1]$, $J = \left[0, \frac{1}{(1+\gamma_0)}\right]$, and
$$U_{\varepsilon} := \{z \in \mathbb{C} \mid \operatorname{Re}(z) \in (-\infty, \gamma_0 - \varepsilon] \cup [\gamma_1 + \varepsilon, +\infty) \text{ or } \operatorname{Im}(z) \notin (-\varepsilon, \varepsilon)\}. \tag{6.1}$$

Proposition 6.1. We have the following statements.

- (i) For all $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus I)$, the series $\sum_{m=1}^{\infty} L_{k,g(m)}\varphi$ converges uniformly on compact subsets $K \subseteq \overline{\mathbb{C}} \setminus J$ to a function $T_k\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus J)$ and there exist $\varepsilon > 0$ and $C_{K,k} > 0$ such that $\sup_K |T_k\varphi| \leq C_{K,k} \sup_{U_{\varepsilon}} |\varphi|$.
- (ii) For all $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus I)$, the series $\sum_{m=1}^{\infty} L_{g(m)}^{(k+2)} \psi$ converges uniformly on compact subsets $K \subseteq \overline{\mathbb{C}} \setminus J$ to a function in $\mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus J)$, which we denote by $T_k^{(k+2)} \psi$ and there exist $\varepsilon > 0$ and $\overline{C}_{K,k} > 0$ such that $\sup_K |T_k^{(k+2)} \psi| \leq \overline{C}_{K,k} \sup_{U_{\varepsilon}} |\psi|$.
- (iii) For all $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus I)$, we have $T_k^{(k+2)}\varphi^{(k+1)} = (T_k\varphi)^{(k+1)}$.

Remark 6.2. Analogous statements can immediately be deduced for $\overline{L}_{g(m)}$ and S by using the relations $\overline{L}_{g(m)} = -L_{1,g(m)}$ and S = -T.

Proof of Proposition 6.1. Let $\varepsilon > 0$, then, from (5.1), there exists $c_{1,\varepsilon,k} > 0$ such that for all $z \in U_{\varepsilon}$ and $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus I)$ we have $|\psi(z)| \leq c_{1,\varepsilon,k}|z|^{-(k+2)} \sup_{U_{\varepsilon}} |\psi|$.

If $K \subseteq \overline{\mathbb{C}} \setminus J$ is compact, then there exists $\varepsilon(K) \in (0, \max\{|\gamma_0|, |\gamma_1|\})$ such that $\frac{1}{z} - m \in U_{\varepsilon(K)}$, for all $z \in K$ and $m \in \mathbb{N}$, which implies that $\varepsilon(K) \leq |1/z - m|$. Also, there exist $c_{2,K} > 0$ and $M_{\gamma_0} \in \mathbb{N}$ such that for all $z \in K$ and $m \geq M_{\gamma_0}$ we have that $\left|\frac{1}{z} - m\right|^{-1} \leq c_{2,K}m^{-1}$. Therefore, for all $z \in K$ and $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus I)$ we have

$$\sum_{m=1}^{\infty} \left| \psi \left(\frac{1}{z} - m \right) \right| \leq \sum_{m=1}^{\infty} c_{1,\varepsilon(K)} \left| \frac{1}{z} - m \right|^{-(k+2)} \sup_{U_{\varepsilon(K)}} |\psi|$$

$$=c_{1,\varepsilon(K),k} \sup_{U_{\varepsilon(K)}} |\psi| \left(\sum_{m=1}^{M_{\gamma_0}-1} \left(\left| \frac{1}{z} - m \right|^{-1} \right)^{k+2} + \sum_{m=M_{\gamma_0}}^{\infty} \left(\left| \frac{1}{z} - m \right|^{-1} \right)^{k+2} \right)$$

$$\leq c_{1,\varepsilon(K),k} \sup_{U_{\varepsilon(K)}} |\psi| \left(\sum_{m=1}^{M_{\gamma_0}-1} \left(\varepsilon(K)^{-1} \right)^{k+2} + \sum_{m=M_{\gamma_0}}^{\infty} \left(c_{2,K} m^{-1} \right)^{k+2} \right)$$

$$=c_{1,\varepsilon(K),k} \sup_{U_{\varepsilon(K)}} |\psi| \left((M_{\gamma_0} - 1)\varepsilon(K)^{-(k+2)} + c_{2,K}^{k+2} \sum_{m=M_{\gamma_0}}^{\infty} m^{-(k+2)} \right)$$

$$\leq \overline{C}_{K,k} \sup_{U_{\varepsilon(K)}} |\psi|,$$

$$\leq \overline{C}_{K,k} \sup_{U_{\varepsilon(K)}} |\psi|,$$

$$(6.2)$$

with $\overline{C}_{K,k} = c_{2,\varepsilon(K),k} \left((M_{\gamma_0} - 1)\varepsilon(K)^{-(k+2)} + c_{2,K}^{k+2} \sum_{m=M_{\gamma_0}}^{\infty} m^{-(k+2)} \right)$ and we obtain the first part of (i). By integrating k+1 times between ∞ and z and substituting $\varphi = \psi^{(-(k+1))}$, we obtain the first part of (ii). The assertion (iii) then follows immediately.

Hence, by (5.22), (5.27) and (6.2) we have for all $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus I)$ and all $z \in K$ that

$$|(T_k \varphi)^{(k+1)}(z)| = |(T_k^{(k+2)} \varphi^{(k+1)})(z)| \le \sum_{m=1}^{\infty} \left| L_{g(m)}^{(k+2)} \varphi^{(k+1)}(z) \right|$$

$$\le |z|^{-(k+2)} \cdot \sum_{k=1}^{\infty} \left| \varphi^{(k+1)} \left(\frac{1}{z} - m \right) \right| \le \overline{C}_{K,k} |z|^{-(k+2)} \sup_{U_{\varepsilon(K)}} |\varphi^{(k+1)}|,$$

implying (ii). We conclude (i) by using Cauchy's formula.

In particular, we immediately obtain from (5.24) that

$$T_k \varphi(z) = -\sum_{m=1}^{\infty} z^k \left(\varphi\left(\frac{1}{z} - m\right) - \varphi(-m) \right) + \sum_{m=1}^{\infty} \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}(-m)$$

proving (5.9).

Let D_{∞} be given as in (4.8) and for $\rho > 0$ let

$$V_{\rho}(D_{\infty}) = \{ z \in \overline{\mathbb{C}} \setminus [0,1] \mid d_{\text{hyper}}(z, D_{\infty}) < \rho \},$$

where d_{hyper} denotes the Poincaré metric on $\overline{\mathbb{C}} \setminus [0,1]$.

To define the complex k-Brjuno function, we deal with $(1 - T_k)^{-1}$ as in the real case, see (1.7) and also (7.9). The following proposition is an analogous property to [MMY97, Thm. 2.6] for the real Brjuno function and it is a generalization of [MMY01, Prop. 3.3] for the case of k = 1, which guarantees that $(1 - T_k)^{-1} = \sum_{r=1}^{\infty} T_k^r$ converges.

However, before we state this proposition we first give an estimate of the derivatives of φ using Cauchy's integral formula. For $\varphi \in \mathcal{O}^n(\overline{\mathbb{C}} \setminus [0,1])$, let $\tilde{\varphi}(z) = \varphi(z) \cdot z^n$. Then we have

$$\varphi^{(j)}(z) = \sum_{i=0}^{j} {j \choose i} \frac{(-1)^{i}(n+i-1)!}{(n-1)!} \cdot \frac{\tilde{\varphi}^{(j-i)}(z)}{z^{n+i}}.$$

By Cauchy's integral formula, for a circle $\{\omega : |\omega - z| = R\}$ contained in $V_{\rho}(D_{\infty})$ whose center is given by $z \in V_{\rho}(D_{\infty})$, we have

$$\left|\widetilde{\widetilde{\varphi}}^{(\ell)}\left(z\right)\right| = \left|\frac{\ell!}{2\pi i} \int_{|\omega-z|=R} \frac{\widetilde{\widetilde{\varphi}}\left(\omega\right)}{(\omega-z)^{\ell+1}} d\omega\right| \le \frac{\ell! \sup_{V_{\rho}(D_{\infty})} \left|\widetilde{\widetilde{\varphi}}\right|}{R^{\ell}}.$$

Since we have

$$\sup_{V_{\rho}(D_{\infty})} |\widetilde{\varphi}| \le \sup_{V_{\rho}(D_{\infty})} |\varphi| \cdot \sup_{z \in \partial V_{\rho}(D_{\infty})} |z|^n \le \sup_{V_{\rho}(D_{\infty})} |\varphi| \cdot \sup_{z \in \partial D_{\infty}} |z|^n,$$

we obtain

$$|\varphi^{(j)}(z)| \le c'_{n,j} \frac{j! \sup_{V_{\rho}(D_{\infty})} |\varphi|}{R^{n+j}}, \quad \text{where } c'_{n,j} := \sup_{z \in \partial D_{\infty}} |z|^n \cdot \sum_{i=0}^j \binom{n+i-1}{i}. \tag{6.3}$$

Proposition 6.3. Let $\rho \geq 0$. The following statements hold.

(i) For all $k \in \mathbb{N}$, there exists $C_{\rho,k} > 0$ such that, for all $r \geq 0$ and $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$, we have

$$\sup_{z \in V_{\rho}(D_{\infty})} |((T_k^{(k+2)})^r \psi)(z)| \le C_{\rho,k} \left(\frac{\sqrt{5} - 1}{2}\right)^{rk} \sup_{z \in V_{\rho}(D_{\infty})} |\psi(z)|.$$

(ii) For all $k \in \mathbb{N}$, there exists $\overline{C}_{\rho,k}$ such that, for all $r \geq 0$ and $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, we have

$$\sup_{z \in V_{\rho}(D_{\infty})} |(T_k^r \varphi)(z)| \le \overline{C}_{\rho,k} \left(\frac{\sqrt{5} - 1}{2}\right)^{rk} \sup_{z \in V_{\rho}(D_{\infty})} |\varphi(z)|.$$

Similarly to the last proposition we remark also here that analogous statements can immediately be deduced for $\overline{L}_{g(m)}$ and S by using the relations $\overline{L}_{g(m)} = -L_{1,g(m)}$ and S = -T.

The proof of the above proposition is given in the Appendix.

In order to state the next proposition, we define

$$Z = \left\{ \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \in GL_2(\mathbb{Z}) \mid n \in \mathbb{Z} \right\}. \tag{6.4}$$

From now on, we consider $L_{k,g}$ and \overline{L}_g for $g \in \operatorname{GL}_2(\mathbb{Z})$. For $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, since $\varphi_0^{(n)} \in \mathcal{O}^{n+1}(\overline{\mathbb{C}} \setminus [0,1])$, we have $\lim_{c \to 0} \varphi_0^{(n)}(-d/c)/c^n = 0$ for $n \ge 0$. Thus, we define $L_{k,g}$ when c = 0 by $L_{k,g}\varphi(z) = \det(g)^{k+1}a^k\varphi\left(\frac{dz-b}{a}\right)$. Especially, for $g \in \mathbb{Z}$, we have

$$L_{k,\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}} \varphi(z) = \overline{L}_{\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}} \varphi(z) = \varphi(z - n).$$

For $H \subset GL_2(\mathbb{Z})$, let us denote

$$\sum_{H}^{(k+2)} \psi := \sum_{g \in H} L_g^{(k+2)} \psi \quad \text{and} \quad \sum_{H,k} \varphi := \sum_{g \in H} L_{k,g} \varphi,$$

which are uniformly summable on compact subsets of $\overline{\mathbb{C}} \setminus [0,1]$.

The following three propositions which are a generalization of [MMY01, Coro. 3.6] will show us a relation between $\sum_{\mathcal{M},k}$ and T_k . They state that $\sum_{\mathcal{M},k} \varphi$ and $\sum_{Z \cdot \mathcal{M},k} \varphi$ are welldefined on any compact set of $\overline{\mathbb{C}} \setminus [0,1]$ for $\varphi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$ and $\sum_{\mathcal{M},k}$ is the inverse of $(1-T_k).$

Proposition 6.4. Let $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$. We have

$$\sum_{\mathcal{M}}^{(k+2)} \psi = \sum_{r=0}^{\infty} (T_k^{(k+2)})^r \psi. \tag{6.5}$$

Let $K \subset \overline{\mathbb{C}} \setminus [0,1]$ be compact. Then, for all $k \in \mathbb{N}$, there exist $\varepsilon(K) > 0$ and $C_{K,k}$ such that

$$\sup_{K} \left| \sum_{\mathcal{M}}^{(k+2)} \psi \right| \le C_{K,k} \sup_{U_{\varepsilon(K)}} |\psi|, \tag{6.6}$$

where $U_{\varepsilon(K)}$ is as in (6.1) with $\gamma_0 = 0$ and $\gamma_1 = 1$.

Furthermore, $\sum_{Z \in \mathcal{M}}^{(k+2)} \psi$ is uniformly summable on all domains of the form $\{|\operatorname{Re}(z)| < 1\}$ $A, |\operatorname{Im}(z)| \geq \delta\}, \ \overline{for\ some}\ A, \delta > 0. \ \ It\ is\ holomorphic\ in\ \mathbb{C}\setminus\mathbb{R},\ \mathbb{Z}$ -periodic, bounded in the neighbourhood of $\pm i\infty$, and

$$\sum_{Z \cdot \mathcal{M}}^{(k+2)} \psi = \sum_{Z}^{(k+2)} \sum_{\mathcal{M}}^{(k+2)} \psi. \tag{6.7}$$

Proof. Equation (6.5) follows from (A.2), and equation (6.6) follows directly from Proposition 6.3 taking ε and ρ such that $K \subset V_{\rho}(D_{\infty}) \subset U_{\varepsilon(K)} \subset \overline{\mathbb{C}} \setminus [0,1]$.

Furthermore, making use of (5.26), we have for $g' = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \in Z$ that $L_{q'}^{(k+2)}\psi(z) =$ $\psi(z-n)$. Then, we have for all $g' \in Z$ and $g \in \mathcal{M}$ that $L_{g' \cdot g}^{(k+2)} \psi = L_{g'}^{(k+2)} \left(L_g^{(k+2)} \psi \right)$ giving (6.7) in case that at least one side is uniformly summable which we will show in the following.

We have by Proposition 6.1 that $(T_k^{(k+2)})^r \psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$ for any $r \geq 0$. Furthermore, by the above consideration, we have that $\sum_{Z}^{(k+2)} \sum_{\mathcal{M}}^{(k+2)} \psi(z) = \sum_{n=1}^{\infty} \left(\sum_{\mathcal{M}}^{(k+2)} \psi(z-n)\right)$. Since $\sum_{n=1}^{\infty} f(z-n) \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$ holds if $f \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$, it follows that $\sum_{Z}^{(k+2)} \sum_{\mathcal{M}}^{(k+2)} \psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$.

Proposition 6.5. Let $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$. We have

$$\sum_{\mathcal{M},k} \varphi = \sum_{r=0}^{\infty} T_k^r \varphi. \tag{6.8}$$

Furthermore, $\sum_{Z:\mathcal{M},k} \varphi$ is holomorphic in $\mathbb{C} \setminus \mathbb{R}$, \mathbb{Z} -periodic, vanishing at $\pm i\infty$ and we have

$$\left(\sum_{\mathcal{M},k}\varphi\right)^{(k+1)} = \sum_{\mathcal{M}}^{(k+2)}\varphi^{(k+1)} \quad and \quad \left(\sum_{Z\cdot\mathcal{M},k}\varphi\right)^{(k+1)} = \sum_{Z\cdot\mathcal{M}}^{(k+2)}\varphi^{(k+1)}. \tag{6.9}$$

Proof. By definition of $\sum_{H,k}$, $\sum_{H}^{(k+2)}$ and (5.22), the equations in (6.9) hold. Equation (6.8) follows from integrating the expression in Proposition 6.4 with $\psi = \varphi^{(k+1)}$, combining with 6.1-(iii) and the first equation of (6.9). The remaining statements follow immediately from the properties of φ and the uniform summability.

Proposition 6.6. We have

$$(1 - T_k) \sum_{\mathcal{M}, k} = \sum_{\mathcal{M}, k} (1 - T_k) = id, \quad and$$

$$(1 - T_k^{(k+2)}) \sum_{\mathcal{M}}^{(k+2)} = \sum_{\mathcal{M}}^{(k+2)} (1 - T_k^{(k+2)}) = id.$$

Proof. It follows from Propositions 6.4 and 6.5.

For $H \subset GL_2(\mathbb{Z})$, We denote by

$$\overline{\sum}_{H}^{(3)} \psi := \sum_{g \in H} \overline{L}_{g}^{(3)} \psi \quad \text{and} \quad \overline{\sum}_{H} \varphi := \sum_{g \in H} \overline{L}_{g} \varphi, \tag{6.10}$$

which are uniformly summable on compact subsets of $\mathbb{C} \setminus [0,1]$. As an analog to the statements above and to [MMY01, Coro. 3.6] we obtain the same statements as in Propositions 6.4, 6.5 and 6.6 with k=1 where we have to replace T_1 by S, \sum by $\overline{\sum}$ and $L_{1,g}$ and $L_{1,g}^{(3)}$ by \overline{L}_g and $\overline{L}_g^{(3)}$. In particular it shows that (6.10) is summable when $H = \mathcal{M}$ or $H = Z \cdot \mathcal{M}$.

6.2. Boundary behaviour of the sums over the monoid $\sum_{\mathcal{M},k} \varphi$ and $\overline{\sum}_{\mathcal{M}} \varphi$. We will consider the boundary behaviour of $\sum_{\mathcal{M},k} \varphi$ for $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$. The following proposition is a generalization of [MMY01, Prop. 4.1] and explains the behaviour of $\sum_{\mathcal{M},k} \varphi$ near 0.

Proposition 6.7. Let $-1 < \gamma_0 < \gamma_1$ and $I = [\gamma_0, \gamma_1]$. Also let

$$U = \{ z \in \mathbb{C} \mid \text{Re}(z) \in (-\infty, \gamma_0 - 1] \cup [\gamma_1 + 1, +\infty) \text{ or } \text{Im}(z) \notin (-1/2, 1/2) \}.$$
 (6.11)

There exists $C_{I,k} > 0$ such that, for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus I)$ and all $z \in D_0 \cup H_0 \cup H'_0$, we have

$$\left| T_k \varphi(z) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \varphi^{(n)}(-m) \right| \le C_{I,k} |z|^k \log(1 + |z|^{-1}) \sup_{U} |\varphi|. \tag{6.12}$$

Moreover, there exists $C_k > 0$ such that, for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and all $z \in D_0 \cup H_0 \cup H'_0$, we have

$$\left| \sum_{\mathcal{M},k} \varphi(z) - \varphi(z) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \left(\sum_{\mathcal{M},k} \varphi \right)^{(n)} (-m) \right| \le C_k |z|^k \log(1 + |z|^{-1}) \sup_{D_{\infty}} |\varphi|. \quad (6.13)$$

We will give the proof of this proposition in much detail as it differs significantly from the case k = 1.

Proof. As the proofs of (6.12) and (6.13) are similar, we will first prove (6.13) and later give the differences for the proof of (6.12). For the following, we set $\tilde{\varphi} = \sum_{\mathcal{M},k} \varphi$. Then, Proposition 6.6 and Proposition 6.1-(i) imply

$$\widetilde{\varphi} = \varphi + T_k \widetilde{\varphi} = \varphi + \sum_{m=1}^{\infty} L_{k,g(m)} \widetilde{\varphi}.$$

By the above equation and (5.9), the expression of the left hand-side of (6.13) can thus be summarized as

$$\left| \sum_{\mathcal{M},k} \varphi(z) - \varphi(z) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \left(\sum_{\mathcal{M}} \varphi \right)^{(n)} (-m) \right|$$

$$= \left| T_k \widetilde{\varphi}(z) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)} (-m) \right| = \left| \sum_{m=1}^{\infty} z^k \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) \right|. \quad (6.14)$$

We will split the sum in the following way:

$$\left| \sum_{\mathcal{M},k} \varphi(z) - \varphi(z) \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \left(\sum_{\mathcal{M},k} \varphi \right)^{(n)} (-m) \right|$$

$$\leq \left| \sum_{m=1}^{\lfloor 3|z|^{-1} \rfloor + 1} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) \right|$$

$$+ \left| \sum_{m=\lfloor 3|z|^{-1} \rfloor + 2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)} (-m) \right|$$

$$+ \left| \sum_{m=\lfloor 3|z|^{-1} \rfloor + 2}^{\infty} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) - \sum_{m=\lfloor 3|z|^{-1} \rfloor + 2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)} (-m) \right| .$$

$$(6.17)$$

We first estimate the first summand (6.15). We note that for $m \in \mathbb{N}_{\leq \lfloor 3|z|^{-1}\rfloor+1}$ and $z \in D_0 \cup H_0 \cup H_0'$ we have that $-m \in D_\infty$ and $1/z - m \in D_\infty$. Applying then (5.1) with $\mathbb{C}\backslash \overline{V} = D_\infty$ yields

$$\left| \sum_{m=1}^{\lfloor 3|z|^{-1}\rfloor + 1} z^k \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) \right|$$

$$\leq \sum_{m=1}^{\lfloor 3|z|^{-1}\rfloor + 1} c_{1,k} |z|^k \cdot \left(\frac{1}{m^k} + \left| -m + \frac{1}{z} \right|^{-k} \right) \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|$$

$$\leq c_{2,k} |z|^k \cdot \left(\sum_{m=1}^{\lfloor 3|z|^{-1}\rfloor + 1} \left(\frac{1}{m} + \left| -m + \frac{1}{z} \right|^{-1} \right) \right) \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|$$

with $c_{2,k} = c_{1,k} \sup_{z \in D_0 \cup H_0 \cup H'_0, m \in \mathbb{N}} \left| -m + \frac{1}{z} \right|^{-k+1} = c_{1,k} 2^{k-1}$. Noting that for all $z \in D_0 \cup H_0 \cup H'_0$ we have

$$\sum_{m=1}^{\lfloor 3|z|^{-1}\rfloor+1} \frac{1}{m} \le 1 + \int_{1}^{\lfloor 3|z|^{-1}\rfloor+1} \frac{1}{m} \, \mathrm{d}m \le 1 + \log(\lfloor 3|z|^{-1}\rfloor + 1),$$

and since $\left|\frac{1}{z} - \left(\left|\operatorname{Re}\left(\frac{1}{z}\right)\right| - n\right)\right| \ge \left|\frac{1}{z} - \left(\left|\operatorname{Re}\left(\frac{1}{z}\right)\right| + n\right)\right|$ for $1 \le n \le \left|\operatorname{Re}(1/z)\right|$ and $\operatorname{Re}(z) \ge 0$, we have

$$\sum_{m=1}^{\left\lfloor 3|z|^{-1}\right\rfloor+1} \left| m - \frac{1}{z} \right|^{-1} = \sum_{\substack{\{m: \ 1 \le m \le \lfloor 3|z|^{-1}\rfloor+1, \\ |m-\lfloor \operatorname{Re}(1/z)\rfloor| \le 2\}}} \left| m - \frac{1}{z} \right|^{-1} + \sum_{\substack{\{m: \ 1 \le m \le \lfloor 3|z|^{-1}\rfloor+1, \\ |m-\lfloor \operatorname{Re}(1/z)\rfloor| > 2\}}} \left| m - \frac{1}{z} \right|^{-1}$$

$$\leq 10 + 2 \sum_{\substack{m=\lfloor \operatorname{Re}(1/z)\rfloor+3 \\ m=1}}^{\left\lfloor 3|z|^{-1}\right\rfloor+1} \frac{1}{m - \lfloor \operatorname{Re}(1/z)\rfloor - 1}$$

$$\leq 10 + 2 \int_{m=1}^{\left\lfloor 3|z|^{-1}\right\rfloor-\lfloor \operatorname{Re}(1/z)\rfloor} \frac{1}{m} \, \mathrm{d}m \le 10 + 2 \log(\lfloor 3|z|^{-1}\rfloor - \lfloor \operatorname{Re}(1/z)\rfloor).$$

The last two inequalities even hold true if $\operatorname{Re}(1/z) < 0$. The fact that $\lfloor 3|z|^{-1} \rfloor + 1 \leq (|z|^{-1} + 1)^3$ implies that there exists another constant $c_{3,k} > 0$ such that, for all $\widetilde{\varphi} \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and $z \in D_0 \cup H_0 \cup H_0'$, we have

$$\left| \sum_{m=1}^{\left\lfloor 3|z|^{-1}\right\rfloor + 1} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) \right| \leq c_{3,k} |z|^{k} \cdot \log \left(1 + |z|^{-1} \right) \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|. \tag{6.18}$$

For the second summand (6.16), from (6.3), for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and $z \in D_0 \cup H_0 \cup H_0'$ we have

$$\left| \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)}(-m) \right| \leq \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \sum_{n=1}^{k} \frac{|z|^{k-n}}{n!} \cdot c'_{k,n} n! \cdot \frac{1}{m^{k+n}} \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|$$

$$\leq \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} |z|^{k-1} \cdot 2^{k-1} c'_{k,n} \cdot \frac{1}{m^{k+1}} \cdot \sup_{D_{\infty}} |\widetilde{\varphi}| .$$

Noting that

$$\sum_{m=|3|z|^{-1}|+2}^{\infty} m^{-k-1} < \int_{\lfloor 3|z|^{-1}\rfloor+1}^{\infty} m^{-k-1} dm = k^{-1} \left(\lfloor 3|z|^{-1}\rfloor + 1 \right)^{-k} < \left(\frac{|z|}{3} \right)^k$$
 (6.19)

yields that for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and $z \in D_0 \cup H_0 \cup H_0'$ we have

$$\left| \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)}(-m) \right| \le |z|^{2k-1} \cdot c'_{k,n} \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|. \tag{6.20}$$

Finally, we estimate the last summand (6.17) using (5.25)

$$\left| \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) - \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)}(-m) \right|$$

$$\leq \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \left| \frac{1}{z \, k!} \int_{0}^{1} \widetilde{\varphi}^{(k+1)} \left(\frac{t}{z} - m \right) (1-t)^{k} dt \right|.$$

$$(6.21)$$

Using Cauchy's integral formula yields for R sufficiently small (possibly depending on m and z) to be defined later that

$$\left| \int_{0}^{1} \widetilde{\varphi}^{(k+1)} \left(\frac{t}{z} - m \right) (1 - t)^{k} dt \right|$$

$$\leq \left| \int_{0}^{1} \frac{(k+1)!}{2\pi i} \int_{|\omega - \left(\frac{t}{z} - m\right)| = R} \frac{\widetilde{\varphi}(\omega)}{\left(\omega - \left(\frac{t}{z} - m\right)\right)^{k+2}} d\omega \cdot (1 - t)^{k} dt \right|$$

$$\leq \sup_{|\omega - \left(\frac{t}{z} - m\right)| = R} \left| \widetilde{\varphi}(\omega) \right| \cdot \frac{(k+1)!}{R^{k+2}} \int_{0}^{1} (1 - t)^{k} dt = \sup_{|\omega - \left(\frac{t}{z} - m\right)| = R} \left| \widetilde{\varphi}(\omega) \right| \cdot \frac{2\pi k!}{R^{k+1}}$$

Now, we specify the radius R. It is sufficient that $-m+t/|z|+R < \sqrt{3}/2-1$ is fulfilled, i.e. it is sufficient that $R < \sqrt{3}/2-1+m-1/|z|$ is fulfilled and thus we can choose R = m/3. Hence, using the above estimate and inserting it into (6.21) yields for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$

$$\left| \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) - \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)}(-m) \right|$$

$$\leq \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} 2\pi \cdot 3^{k+1} \cdot |z|^{-1} \cdot m^{-k-1} \cdot \sup_{\left|\omega - \left(\frac{t}{z} - m\right)\right| = m/3} |\widetilde{\varphi}|.$$

$$(6.22)$$

Next, we further estimate $\sup_{|\omega-\left(\frac{t}{z}-m\right)|=m/3} |\widetilde{\varphi}|$. Using (5.1) and noting that we have $D_{z,m,t}:=\{\omega: |\omega-\left(\frac{t}{z}-m\right)|=m/3\}\subset D_{\infty} \text{ and thus } \inf_{\omega\in D_{z,m,t}} |\omega|\geq \left|\frac{t}{z}-m\right|-\frac{m}{3}\geq m-|z|^{-1}-\frac{m}{3}\geq \frac{m}{3} \text{ yields that there exists } c_{4,k} \text{ such that}$

$$\sup_{D_{z,m,t}} |\widetilde{\varphi}| \le c_{4,k} \sup_{D_{\infty}} |\widetilde{\varphi}| \cdot \sup_{D_{z,m,t}} |\omega|^{-k} \le c_{4,k} \sup_{D_{\infty}} |\widetilde{\varphi}| \cdot \left(\frac{m}{3}\right)^{-k}$$
(6.23)

and for $m \ge |3|z|^{-1}| + 2$, we have

$$\sup_{D_{z,m,t}} |\widetilde{\varphi}| \le c_{4,k} \sup_{D_{\infty}} |\widetilde{\varphi}| \cdot |z|^{k}.$$

Combining this with (6.22) and (6.19) yields

$$\left| \sum_{m=\lfloor 3|z|^{-1}\rfloor+2}^{\infty} z^{k} \left(\widetilde{\varphi} \left(\frac{1}{z} - m \right) - \widetilde{\varphi}(-m) \right) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \widetilde{\varphi}^{(n)}(-m) \right|$$

$$\leq 2\pi c_{4,k} \cdot 3^{k+1} \cdot |z|^{-1} \cdot \left(\frac{|z|}{3} \right)^{2k} \cdot \sup_{D_{\infty}} |\widetilde{\varphi}| \leq 2\pi c_{4,k} \cdot |z|^{2k-1} \cdot \sup_{D_{\infty}} |\widetilde{\varphi}|.$$
 (6.24)

Combining the sum of (6.15), (6.16), and (6.17) estimated in (6.18), (6.20), and (6.24), we obtain that there exists a constant $C_k > 0$ such that for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and $z \in D_0 \cup H_0 \cup H'_0$ we have the statement of (6.13).

The left handside of (6.14) equals the left handside of (6.12) if we replace φ by $\widetilde{\varphi}$. All the remaining calculations remain the same with the difference that we have to take the supremum over U instead of D_{∞} . However, there are only two instances where this is important. The first is the choice of R which has to fulfill $R < \gamma_0 + m - 1/|z|$ instead of $R < \sqrt{3}/2 - 1 + m - 1/|z|$ and for both $R \le m/3$ is sufficient. The second is the occurrence of $\inf_{x \in D_{\infty}} |x|^{-k}$ in (6.23) which we were estimating by $c_{1,k}$ and which has to be changed to an estimate by a constant depending on I and k.

Our next concern will be the behaviour of $\sum_{\mathcal{M},k} \varphi$ for $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ near 1. The following proposition is a generalization of [MMY01, Prop. 4.2].

Proposition 6.8. There exists $C_k > 0$ such that for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$ and all $z \in D_1$ we have

$$\left| T_k \varphi(z) + z^k \varphi\left(\frac{1}{z} - 1\right) - \sum_{m=1}^{\infty} \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}(-m) \right| \le C_k |z - 1| \sup_{D_{\infty}} |\varphi|.$$
 (6.25)

Proof. We first notice that (5.9) implies

$$\left| T_{k}\varphi(z) + z^{k}\varphi\left(\frac{1}{z} - 1\right) - \sum_{m=1}^{\infty} \sum_{n=1}^{k} \frac{z^{k-n}}{n!} \varphi^{(n)}(-m) \right|
= \left| \sum_{m=2}^{\infty} z^{k}\varphi\left(\frac{1}{z} - m\right) - \sum_{m=1}^{\infty} z^{k}\varphi(-m) \right|
\leq |z|^{k} \cdot \left| \sum_{m=2}^{\infty} \left(\varphi\left(\frac{1}{z} - m\right) - \varphi(1 - m)\right) \right|.$$
(6.26)

For $z \in D_1$, we have $1/z - 1 \in D_0$. Equation (6.3) implies that there exists a constant $c_{1,k} > 0$ such that $|\varphi'(w)| \le c_{1,k} m^{-k-1} \sup_{D_{\infty}} |\varphi|$. This implies

$$\left| T_k \varphi(z) + z^k \varphi\left(\frac{1}{z} - 1\right) - \sum_{m=1}^{\infty} \sum_{n=1}^k \frac{z^{k-n}}{n!} \varphi^{(n)}(-m) \right|$$

$$\leq |z|^k \cdot \sum_{m=2}^{\infty} c_{1,k} m^{-k-1} \sup_{D_{\infty}} |\varphi| \cdot \left| \frac{1}{z} - 1 \right| = |z|^{k-1} \cdot |1 - z| \sup_{D_{\infty}} |\varphi| \sum_{m=2}^{\infty} c_{1,k} m^{-k-1}$$

$$= C_k \cdot |1 - z| \sup_{D_{\infty}} |\varphi|,$$

where
$$C_k := \sup_{z \in D_1} |z|^{k-1} c_{1,k} \sum_{m=2}^{\infty} m^{-k-1}$$
.

We can finally use the previous results and the complex continued fraction algorithm to study the behaviour of $T_k^n \varphi(z)$ when z is close to the boundary. We recall that $D(m_1, \dots, m_n)$ is the set of $z_0 \in D$ whose first n complex continued fraction entries equal $\{m_i\}_{i=1}^n$. Next, we will prove a proposition which is an analog to [MMY01, Prop. 4.4].

Proposition 6.9. For all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0,1])$, $n \in \mathbb{N}$, and $z_0 \in D(m_1, \dots, m_n)$ we have

$$T_k^n \varphi(z_0) = (-1)^{n(k+1)} (p_{n-1} - q_{n-1} z_0)^k (\varphi(z_n) + \varphi(z_n - 1) + \varepsilon_n \varphi(z_n + 1))$$
$$- (-1)^{(n-1)(k+1)} (p_{n-2} - q_{n-2} z_0)^k (1 + z_{n-1})^k \varepsilon_{n-1} \varphi\left(-\frac{z_{n-1}}{1 + z_{n-1}}\right) + R_k^{[n]} \varphi(z_0). \quad (6.27)$$

The remainder term $R_k^{[n]}$ is holomorphic in the interior of $D(m_1, \dots, m_n)$ and continuous in $D(m_1, \dots, m_n)$ and there exists $C_k > 0$ such that for all $\varphi \in \mathcal{O}^k(\overline{\mathbb{C}} \setminus [0, 1])$, $n \in \mathbb{N}$, and $z_0 \in D(m_1, \dots, m_n)$ we have

$$\left| R_k^{[n]} \varphi(z_0) \right| \le C_k \, n \left(\frac{\sqrt{5} - 1}{2} \right)^{nk} \sup_{D_\infty} |\varphi|. \tag{6.28}$$

Proof. We start by an estimate for n=1. For $z_0 \in D(m_1)$, by definition, we have

$$T_k \varphi(z_0) = -\sum_{m=1}^{\infty} z_0^k \left(\varphi\left(\frac{1}{z_0} - m\right) - \varphi(-m) \right) + \sum_{m=1}^{\infty} \sum_{j=1}^k \frac{z_0^{k-j}}{j!} \varphi^{(j)}(-m)$$
$$= -z_0^k (\varphi(z_1) + \varphi(z_1 - 1) + \varepsilon_1 \varphi(z_1 + 1)) + R_k^{(m_1)}(\varphi)(z_0),$$

with

$$R_k^{(m_1)}(\varphi)(z_0) = \sum_{\substack{m \ge 1 \\ |m-m_1| \le 1}} z_0^k \varphi(-m) + \sum_{m=1}^{\infty} \sum_{j=1}^k \frac{z_0^{k-j}}{j!} \varphi^{(j)}(-m)$$
$$- \sum_{\substack{m \ge 1 \\ |m-m_1| > 1}} z_0^k (\varphi(m_1 + z_1 - m) - \varphi(-m))$$
$$=: I_0 + I_1 - I_2.$$

Then, $R_k^{(m_1)}(\varphi)$ is holomorphic in a neighbourhood of $D(m_1)$.

We will first show that there is a constant $c_{1,k}$ such that

$$|R_k^{(m_1)}(\varphi)(z_0)| \le c_{1,k} \sup_{D_\infty} |\varphi| \quad \text{for all } z_0 \in D(m_1).$$
 (6.29)

Note that this however is not an estimate of (6.27) for n = 1 as the terms in the second line are all contained in the error term $R_k^{(m_1)}(\varphi)(z_0)$. We start by estimating I_0 . Since $|z_0| \le 1$ and $I_0 = z_0^k (\varphi(-m_1 - 1) + \varphi(-m_1) + \varepsilon_1 \varphi(-m_1 + 1))$, we have

$$|I_0| \le 3 \sup_{D_\infty} |\varphi|. \tag{6.30}$$

From (6.3) with $\{\omega : |\omega + m| = m - 0.8\} \subset D_{\infty}$, we have

$$|\varphi^{(j)}(-m)| \le c'_{k,j} \frac{j! \sup_{D_{\infty}} |\varphi|}{(m-0.8)^{k+j}}.$$

Let $c_{2,k} = \sum_{m=1}^{\infty} \sum_{j=1}^{k} \frac{c'_{k,j}}{(m-0.8)^{k+j}}$. Since $\sum_{m=1}^{\infty} \frac{1}{(m-0.8)^{k+j}} < \infty$ for $k, j \ge 1$,

$$c_{2,k} = \sum_{m=1}^{\infty} \sum_{j=1}^{k} \frac{c'_{k,j}}{(m-0.8)^{k+j}} = \sum_{j=1}^{k} \sum_{m=1}^{\infty} \frac{c'_{k,j}}{(m-0.8)^{k+j}} < \infty.$$

Then we have

$$|I_1| \le c_{2,k} \sup_{D_{\infty}} |\varphi|. \tag{6.31}$$

Since $z_0 \in D(m_1)$, we have $1/z_0 = z_1 + m_1 \in \Delta + m_1$ (see (4.7) for the definition of Δ). Furthermore, $\frac{1}{m_1+1} \leq |z_0| \leq \frac{1}{m_1-1}$ if $m_1 \neq 1$ and $1/2 \leq |z_0| \leq 1$ if $m_1 = 1$. Let us consider

$$I_2' := \sum_{m > m_1 + 2} z_0^k (\varphi(1/z_0 - m) - \varphi(-m)),$$

and $I_2'' := I_2 - I_2'$. Then, $I_2'' = 0$ if $m_1 \le 2$, and

$$I_2'' = \sum_{1 \le m \le m_1 - 2} z_0^k (\varphi(1/z_0 - m) - \varphi(-m))$$
 if $m_1 \ge 3$.

If $m \ge m_1 + 2$, then $\text{Re}(1/z_0 - m) \le -1$. For z in the segment $[-m, 1/z_0 - m] \subset D_{\infty}$, the circle $\{\omega : |\omega - z| = m - 0.8\}$ is contained in D_{∞} . Thus, by (6.3), we have $|\varphi'(z)| \le c'_{k,1} \frac{\sup_{D_{\infty}} |\varphi|}{(m-0.8)^{k+1}}$ and thus

$$|I_2'| \le \sum_{m \ge m_1 + 2} \frac{c_{k,1}'}{(m - 0.8)^{k+1}} \sup_{D_\infty} |\varphi|.$$

If $m_1 \geq 3$, then $1/z_0 - m \in \Delta + (m_1 - m) \subset D_{\infty}$ for $1 \leq m \leq m_1 - 2$. Thus,

$$|I_2''| \le \frac{2(m_1 - 2)\sup_{D_\infty} |\varphi|}{(m_1 - 1)^k} \le 2\sup_{D_\infty} |\varphi|.$$

Then, $|I_2| \leq c_{3,k} \sup_{D_{\infty}} |\varphi|$, where $c_{3,k} := \sum_{m=3}^{\infty} \frac{c'_{k,1}}{(m-0.8)^{k+1}} + 2$. Hence, by combining the last estimate with (6.30) and (6.31) and setting $c_{1,k} := c_{2,k} + c_{3,k} + 3$, we obtain (6.29).

In the following we will estimate $T_k^n \varphi(z_0)$ itself which we split into a number of summands to be estimated separately. Recall z_i as in (4.9) and (4.11). Iterating T_k n times, we get

$$T_k^n \varphi(z_0) = (\varphi(z_n) + \varphi(z_n - 1) + \varepsilon_n \varphi(z_n + 1)) \prod_{i=0}^{n-1} (-z_i^k)$$

$$+ \sum_{j=1}^{n-1} T_k^{n-j} \varphi(z_j - 1) \prod_{i=0}^{j-1} (-z_i^k)$$

$$+ \sum_{j=1}^{n-1} \varepsilon_j T_k^{n-j} \varphi(z_j + 1) \prod_{i=0}^{j-1} (-z_i^k)$$

$$+ \sum_{j=1}^{n} R_k^{(m_j)} (T_k^{n-j} \varphi)(z_{j-1}) \prod_{i=0}^{j-2} (-z_i^k)$$

$$=: J_1 + J_2 + J_3 + J_4,$$

where $z_{-1} = 1$. We recall (4.12) and (4.14), i.e., $\prod_{i=0}^{n-1} (-z_i^k) = (-1)^{n(k+1)} (p_{n-1} - q_{n-1}z_0)^k$ and $|p_{n-1} - q_{n-1}z_0| < q_n^{-1}$. Thus J_1 is the first term of (6.27). In the following steps (1)-(3) we estimate the summands J_2 , J_3 , and J_4 .

(1) We continue by estimating J_4 . By using (4.12), (4.14) and (6.29) and by applying Proposition 6.3-(ii) to each of the summands and noting that $q_j \geq \left(\frac{\sqrt{5}-1}{2}\right)^{-j}$, we obtain

$$\left| \sum_{j=1}^{n} R_{k}^{(m_{j})}(T_{k}^{n-j}\varphi)(z_{j-1}) \prod_{i=0}^{j-2} (-z_{i}^{k}) \right| \leq \sum_{j=1}^{n} \frac{c_{1,k} \sup_{D_{\infty}} |T_{k}^{n-j}\varphi|}{q_{j-1}^{k}}$$

$$\leq c_{1,k} \sum_{j=1}^{n} \overline{C}_{0,k} \left(\frac{\sqrt{5}-1}{2} \right)^{(n-1)k} \sup_{D_{\infty}} |\varphi|$$

$$\leq c_{4,k} \cdot n \left(\frac{\sqrt{5}-1}{2} \right)^{nk} \sup_{D_{\infty}} |\varphi|$$
(6.32)

with $c_{4,k} = c_{1,k} \overline{C}_{0,k} \left(\frac{\sqrt{5}+1}{2} \right)^k$.

(2) In the next steps, we will show that $|J_2| \leq c_{5,k} n(\frac{\sqrt{5}-1}{2})^{nk} \sup_{D_{\infty}} |\varphi|$ for some $c_{5,k}$. If $z_j - 1 \notin D_0$ for $1 \leq j < n$, then we have

$$|T_k^{n-j}\varphi(z_j-1)| \le C_{K,k} \sup_{D_\infty} |(T_k^{n-j-1}\varphi)|$$
 (6.33)

by Proposition 6.1-(i) with I = [0, 1] and $K = \overline{(D-1) \setminus D_0}$.

On the other hand, let us consider the case of $z_j - 1 \in D_0$. From (6.12), we deduce that

$$|T_k^{n-j}\varphi(z_j-1)| \le C_{I,k}|z_j-1|^k\log(1+|z_j-1|^{-1})\sup_{D_\infty}|(T_k^{n-j-1}\varphi)|$$
$$+\left|\sum_{m=1}^{\infty}\sum_{i=1}^k\frac{(z_j-1)^{k-i}}{i!}(T_k^{n-j-1}\varphi)^{(i)}(-m)\right|.$$

From (6.31) and $|z_j - 1|^k \log(1 + |z_j - 1|^{-1}) \le \log 2$, we see that

$$|T_k^{n-j}\varphi(z_j-1)| \le (C_{I,k}\log 2 + c_{2,k}) \sup_{D_\infty} |(T_k^{n-j-1}\varphi)|.$$
 (6.34)

Let $c_{6,k} = \max\{C_{K,k}, C_{I,k} \log 2 + c_{2,k}\}$. Then, by combining the calculations leading to (6.32), (6.33) and (6.34), we get

$$\left| \sum_{j=1}^{n-1} T_k^{n-j} \varphi(z_j - 1) \prod_{i=0}^{j-1} (-z_i^k) \right| \le c_{5,k} \, n \left(\frac{\sqrt{5} - 1}{2} \right)^{nk} \sup_{D_{\infty}} |\varphi|,$$

where $c_{5,k} = \overline{C}_{0,k} c_{6,k} \left(\frac{\sqrt{5}+1}{2} \right)^k$.

(3) To complete the proof, it is enough to show that there is a constant $c_{7,k}$ such that

$$\left| J_3 + \prod_{i=0}^{n-2} (-z_i^k) (1 + z_{n-1})^k \varepsilon_{n-1} \varphi \left(-\frac{z_{n-1}}{1 + z_{n-1}} \right) \right| \le c_{7,k} \, n \left(\frac{\sqrt{5} - 1}{2} \right)^{nk} \sup_{D_\infty} |\varphi|. \tag{6.35}$$

We have

$$J_{3} + \prod_{i=0}^{n-2} (-z_{i}^{k})(1+z_{n-1})^{k} \varepsilon_{n-1} \varphi \left(-\frac{z_{n-1}}{1+z_{n-1}}\right)$$

$$= \sum_{j=1}^{n-2} \varepsilon_{j} T_{k}^{n-j} \varphi(z_{j}+1) \prod_{i=0}^{j-1} (-z_{i}^{k})$$

$$+ \left(T_{k} \varphi(z_{n-1}+1) + (1+z_{n-1})^{k} \varphi\left(-\frac{z_{n-1}}{1+z_{n-1}}\right)\right) \prod_{i=0}^{n-2} (-z_{i}^{k})$$

$$=: I_{4} + I_{5}.$$

(3a) We will estimate I_4 . If m_{j+1} is small enough so that $z_j + 1 \notin D_1$, then by Proposition 6.1-(i) applied to I = [0,1] and $K' = \overline{(D+1) \setminus D_1}$ we have

$$|T_k^{n-j}\varphi(z_j+1)| \le C_{K',k} \sup_{D_\infty} |(T_k^{n-j-1}\varphi)|.$$

If m_{j+1} is large so that $z_j + 1 \in D_1$, then by Proposition 6.8 and (6.31), we deduce that

$$|T_k^{n-j}\varphi(z_j+1)|$$

$$\leq |1+z_{j}|^{k} \left| (T_{k}^{n-j-1}\varphi) \left(-\frac{z_{j}}{1+z_{j}} \right) \right| + C_{k}|z_{j}| \sup_{D_{\infty}} |T_{k}^{n-j-1}\varphi|$$

$$+ \left| \sum_{m=1}^{\infty} \sum_{i=1}^{k} \frac{(z_{j}+1)^{k-i}}{i!} (T_{k}^{n-j-1}\varphi)^{(i)} (-m) \right|$$

$$\leq |1+z_{j}|^{k} \left| (T_{k}^{n-j-1}\varphi) \left(-\frac{z_{j}}{1+z_{j}} \right) \right| + (C_{k} + c_{2,k}(2\sqrt{3}/3)^{k}) \sup_{D_{\infty}} |T_{k}^{n-j-1}\varphi|.$$

Since $-\frac{z_j}{1+z_j} \in D_0$ and $j \leq n-2$, by Proposition 6.7, we have

$$|1+z_{j}|^{k} \left| (T_{k}^{n-j-1}\varphi) \left(-\frac{z_{j}}{1+z_{j}} \right) \right|$$

$$\leq C_{I,k}|1+z_{j}|^{k} \left| -\frac{z_{j}}{1+z_{j}} \right|^{k} \log \left(1 + \left| -\frac{z_{j}}{1+z_{j}} \right|^{-1} \right) \sup_{D\infty} |T_{k}^{n-j-2}\varphi|$$

$$+ |1+z_{j}|^{k} \left| \sum_{m=1}^{\infty} \sum_{i=1}^{k} \frac{(-z_{j}/(1+z_{j}))^{k-i}}{i!} (T_{k}^{n-j-2}\varphi)^{(i)} (-m) \right|.$$

Note that

$$|1 + z_j|^k \left| -\frac{z_j}{1 + z_j} \right|^k \log \left(1 + \left| -\frac{z_j}{1 + z_j} \right|^{-1} \right)$$

$$= |z_j|^k \log \left(1 + \left| 1 + \frac{1}{z_j} \right| \right) \le |z_j|^k \log \left(2 + \frac{1}{|z_j|} \right).$$

Since $x^k \log(2 + 1/x) \to 0$ as $x \to 0$, it is increasing and $|z_j| < 1$, it follows that $|z_j|^k \log(2 + 1/|z_j|) \le \log 3$. Thus, by a similar way to the proof of (6.34), it follows that

$$|1+z_j|^k \left| (T_k^{n-j-1}\varphi) \left(-\frac{z_j}{1+z_j} \right) \right| \le \left(C_{I,k} \log 3 + c_{2,k} \left(2\sqrt{3}/3 \right)^k \right) \sup_{D_\infty} |(T_k^{n-j-2}\varphi)|.$$

By letting $c_{8,k} = \max \{C_{K',k}, C_k + c_{2,k} (2\sqrt{3}/3)^k, C_{I,k} \log 3 + c_{2,k} (2\sqrt{3}/3)^k \}$, we have

$$|I_4| \le c_{8,k} \sum_{i=1}^{n-2} \frac{\sup_{D_\infty} |T_k^{n-j-1}\varphi| + \sup_{D_\infty} |T_k^{n-j-2}\varphi|}{q_i^k}.$$
 (6.36)

(3b) We will estimate I_5 . If $z_{n-1} + 1 \notin D_1$, then $\operatorname{Re}\left(\frac{1}{(z_{n-1}+1)} - 1\right) < \frac{\sqrt{3}}{2} - 1$. Thus, $\frac{1}{(z_{n-1}+1)} - 1 \in D_{\infty}$. By Proposition 6.1-(i), we have

$$\left| T_k \varphi(z_{n-1} + 1) + (1 + z_{n-1})^k \varphi\left(-\frac{z_{n-1}}{1 + z_{n-1}} \right) \right| \le (C_{K',k} + 2^k) \sup_{D_{\infty}} |\varphi|.$$

By Proposition 6.8 in the same manner as showing (6.31), we obtain that $z_{n-1} + 1 \in D_1$ implies

$$\left| T_{k} \varphi(z_{n-1} + 1) + (1 + z_{n-1})^{k} \varphi\left(-\frac{z_{n-1}}{1 + z_{n-1}}\right) \right| \\
\leq C_{k} |z_{n-1}| \sup_{D_{\infty}} |\varphi| + \left| \sum_{m=1}^{\infty} \sum_{i=1}^{k} \frac{(z_{n-1} + 1)^{k-i}}{i!} \varphi^{(i)}(-m) \right| \\
\leq \left(C_{k} + c_{2,k} (2\sqrt{3}/3)^{k} \right) \sup_{D_{\infty}} |\varphi|.$$

Let $c_{9,k} = \max\{C_{K',k} + 2^k, C_k + c_{2,k}(2\sqrt{3}/3)^k\}$. Then, we have

$$|I_5| \le \frac{c_{9,k} \sup_{D_{\infty}} |\varphi|}{q_{n-1}^k}.$$
 (6.37)

Letting $c_{10,k} = \max\{c_{8,k}, c_{9,k}\}$, from (6.36), (6.37) and (6.32), we obtain

$$|I_4| + |I_5| \le 2c_{10,k} \sum_{j=0}^{n-1} \frac{\sup_{D_\infty} |T_k^{n-j-1}\varphi|}{q_{j-1}^k} \le c_{7,k} n \left(\frac{\sqrt{5}-1}{2}\right)^{nk} \sup_{D_\infty} |\varphi|,$$

where $c_{7,k} = 2c_{10,k}\overline{C}_{0,k}(\frac{\sqrt{5}+1}{2})^{2k}$. Note that $c_{7,k}$ depends only on k.

The next proposition is an analog to [MMY01, Prop. 4.1]. We can show the following proposition in the same manner of the proof in [MMY01] by using (5.28) and Proposition 6.3-(2) for k = 1 and the fact that S = -T.

Proposition 6.10. The following two statements hold true:

(i) Let $I = [\gamma_0, \gamma_1], \ \gamma_0 > -1$. There exists $C_I > 0$ such that for all $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus I)$ and for all $z \in D_0 \cup H_0 \cup H_0'$, one has

$$|S\varphi(z) + \sum_{m>1} \varphi'(-m)| \le C_I |z| \log(1+|z|^{-1}) \sup_{U} |\varphi|,$$
 (6.38)

where U is as in (6.11).

(ii) There exists C > 0 such that for all $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ and for all $z \in D_0 \cup H_0 \cup H'_0$, one has

$$|\overline{\sum_{\mathcal{M}}}\varphi(z) - \varphi(z) + \sum_{m \ge 1} (\overline{\sum_{\mathcal{M}}}\varphi)'(-m)| \le C|z|(1 + \log|z|^{-1}) \sup_{D_{\infty}} |\varphi|.$$
 (6.39)

Concerning the boundary behaviour of the sum leading to the complex Wilton function, we simply observe that since S = -T Propositions 4.1 and 4.4 of [MMY01] hold. The same is true for Proposition 4.11 of [MMY01], concerning the action of T = -S on the space of analytic functions on $\overline{\mathbb{C}} \setminus [0,1]$ with bounded real part. However the Wilton function will not have a bounded real part we thus omit the discussion of it.

7. Complex k-Brjuno function and complex Wilton function

In this section we finally define the complex k-Brjuno function and the complex Wilton function. Let

$$\varphi_0(z) := -\frac{1}{\pi} \operatorname{Li}_2\left(\frac{1}{z}\right) \quad \text{with} \quad \operatorname{Li}_2(z) := \sum_{n=1}^{\infty} \frac{z^n}{n^2}.$$
 (7.1)

Note that $\varphi_0 \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$, $\operatorname{Im}\varphi_0(x) = 0$ if $x \in \mathbb{R} \setminus [0,1]$ and

$$\operatorname{Im}\varphi_0(x \pm i0) = \pm \log\left(\frac{1}{x}\right) \quad \text{for } x \in (0, 1]. \tag{7.2}$$

We define $\tau := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$,

$$\varphi_{1,k} := (L_{k,g(1)} + L_{k,\tau}) \varphi_0$$
 and $\overline{\varphi}_1 := (\overline{L}_{g(1)} + \overline{L}_{\tau}) \varphi_0$.

Note that $\overline{\varphi}_1 = (-L_{1,g(1)} + L_{1,\tau})\varphi_0$. Further, we note that $\varphi_0 \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ implies $\varphi_{1,k}, \ \overline{\varphi}_1 \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [1/2,2])$.

Similarly as in [MMY01, Section 5] we note that from (6.8) we have that $\sum_{\mathcal{M},k} \varphi_{1,k} = \varphi_{1,k} + \sum_{\mathcal{M},k} T_k \varphi_{1,k}$. Since $Z\mathcal{M} = Z\mathcal{M}g(1) \sqcup Z\mathcal{M}\tau$, we have

$$\sum_{ZM,k} \varphi_0(z) = \sum_{Z,k} [\varphi_{1,k}(z) + \sum_{M,k} (T_k \varphi_{1,k}(z))] = \sum_{ZM,k} \varphi_{1,k}(z).$$
 (7.3)

Similarly, we have $\overline{\sum_{ZM}}\varphi_0(z) = \overline{\sum_{ZM}}\overline{\varphi}_1(z)$. We then deduce by simply applying the definitions of $L_{k,g}$ and \overline{L}_g

$$\varphi_{1,k}(z) = -z^k \varphi_0\left(\frac{1}{z} - 1\right) + \varphi_0(z - 1) + \sum_{n=0}^k \frac{z^{k-n}}{n!} \varphi_0^{(n)}(-1)$$

and

$$\overline{\varphi}_1(z) = z \left(\varphi_0 \left(\frac{1}{z} - 1 \right) - \varphi_0(-1) \right) + \varphi_0(z - 1) - \varphi_0'(-1). \tag{7.4}$$

We deduce that

$$\varphi_{1,k}(z) = \frac{1}{\pi} \left[z^k \text{Li}_2\left(\frac{z}{1-z}\right) - \text{Li}_2\left(\frac{1}{z-1}\right) \right] + \sum_{n=0}^k \frac{z^{k-n}}{n!} \varphi_0^{(n)}(-1), \tag{7.5}$$

and since we additionally have $\varphi_0(-1) = \frac{\pi}{12}$ and $\varphi_0'(-1) = \frac{\log 2}{\pi}$, we also have

$$\overline{\varphi}_1(z) = -\frac{1}{\pi} \left[z \operatorname{Li}_2\left(\frac{z}{1-z}\right) + \operatorname{Li}_2\left(\frac{1}{z-1}\right) \right] - z \frac{\pi}{12} - \frac{\log 2}{\pi}. \tag{7.6}$$

By (7.2), we have

$$\operatorname{Im} \varphi_{1,k}(x \pm i0) = \begin{cases} \pm x^k \log \frac{x}{1-x}, & \text{if } 1/2 \le x < 1, \\ \pm \log \frac{1}{x-1}, & \text{if } 1 < x \le 2 \end{cases}$$
 (7.7)

and

$$\operatorname{Im}\overline{\varphi}_{1}(x \pm i0) = \begin{cases} \mp x \log \frac{x}{1-x} & \text{if } 1/2 \le x < 1, \\ \pm \log \frac{1}{x-1} & \text{if } 1 < x \le 2. \end{cases}$$
 (7.8)

Since T_k is defined exactly in such a way that $T_k\varphi(x) = x^k \varphi(1/x)$ as an operator on the space of real functions and the k-Brjuno function fulfills the recursion $B_k(x) = -\log x + x^k B_k(1/x)$, we have $(1 - T_k) B_k(x) = -\log x$ and therefore

$$B_k(x) = ((1 - T_k)^{-1} f)(x)$$
 with $f(x) = \sum_{n \in \mathbb{Z}} \text{Im} \varphi_0(x + i0 - n).$ (7.9)

We note here that by the above considerations $f(x) = \sum_{n \in \mathbb{Z}} \operatorname{Im} \varphi_0(x + i0 - n) = \log\{x\}$. Then, the complex analytic extension of the real k-Brjuno function is $(1 - T_k)^{-1} (\sum_Z \varphi_0)$ and we have $(1 - T_k)^{-1} (\sum_Z \varphi_0) = \sum_{ZM,k} \varphi_0$ by applying (5.13), here T_k is the extended operator as in (5.9). Hence, it is natural to define the complex k-Brjuno function by

$$\mathcal{B}_k := \sum_{ZMk} \varphi_0.$$

By applying the above consideration to W instead of B_k and S instead of T_k , we may define the complex analytic extension of W by

$$W := \overline{\sum_{ZM}} \varphi_0, \tag{7.10}$$

which we call the complex Wilton function.

There is a two-to-one correspondence between $Z\mathcal{M}$ and \mathbb{Q} . From (5.11) and (5.12), we obtain analogous formulations for \mathcal{B}_k and \mathcal{W} respectively such that

$$\mathcal{B}_{k}(z) = -\sum_{p/q \in \mathbb{Q}} \det \begin{pmatrix} p' & p \\ q' & q \end{pmatrix}^{k+1}$$

$$\left\{ \frac{1}{\pi} \left[(p' - q'z)^{k} \operatorname{Li}_{2} \left(\frac{p' - q'z}{qz - p} \right) - (q''z - p'')^{k} \operatorname{Li}_{2} \left(\frac{p'' - q''z}{qz - p} \right) \right] + \sum_{n=0}^{k} \frac{1}{n!} \det \begin{pmatrix} p' & p \\ q' & q \end{pmatrix}^{n}$$

$$\left[\frac{(p' - q'z)^{k-n}}{(q')^{n}} \varphi_{0}^{(n)} \left(-\frac{q}{q'} \right) - \frac{(q''z - p'')^{k-n}}{(q'')^{n}} \varphi_{0}^{(n)} \left(-\frac{q}{q''} \right) \right] \right\}, \quad (7.11)$$

where $\left[\frac{p'}{q'}, \frac{p''}{q''}\right]$ is the Farey interval such that $\frac{p}{q} = \frac{p'+p''}{q'+q''}$ (with the convention p' = p-1, q' = 1, p'' = 1, q'' = 0 if q = 1). To obtain the representation as in (1.9) we remember the definition of φ_0 from (7.1) and note that

$$\varphi_0'(z) = -\frac{1}{\pi} \cdot \frac{1}{z} \log\left(1 - \frac{1}{z}\right).$$

If we let $f(z) = \frac{1}{z}$ and $g(z) = \log\left(1 - \frac{1}{z}\right)$, then $\varphi'_0(z) = -\frac{1}{\pi}f(z)g(z)$ and $f^{(j)}(z) = \frac{(-1)^j j!}{z^{j+1}} \text{ for } j \ge 0,$

and

$$g^{(j)}(z) = \frac{(-1)^{j-1}(j-1)!}{z^j} \left(\left(\frac{z}{z-1}\right)^j - 1 \right) \text{ for } j \ge 1$$

holds. Since

$$(fg)^{(n)} = \sum_{i=0}^{n} \binom{n}{i} f^{(n-i)} g^{(i)},$$

we have, for $n \geq 1$,

$$\varphi_0^{(n)}(z) = -\frac{1}{\pi} \sum_{i=0}^{n-1} \binom{n-1}{i} f^{(n-1-i)} g^{(i)}$$

$$= -\frac{1}{\pi} \left[\frac{(-1)^{n-1}(n-1)!}{z^n} \log \left(1 - \frac{1}{z}\right) + \sum_{i=1}^{n-1} \frac{(n-1)!}{(n-i-1)!i!} \frac{(-1)^{n-1-i}(n-1-i)!}{z^{n-i}} \frac{(-1)^{i-1}(i-1)!}{z^i} \right]$$

$$\cdot \left(\left(\frac{z}{z-1}\right)^i - 1 \right) \right]$$

$$= -\frac{1}{\pi} \left[\frac{(-1)^{n-1}(n-1)!}{z^n} \log \left(1 - \frac{1}{z}\right) + \frac{(-1)^n(n-1)!}{z^n} \sum_{i=1}^{n-1} \frac{1}{i} \left(\left(\frac{z}{z-1}\right)^i - 1 \right) \right]$$

$$= \frac{(-1)^n(n-1)!}{\pi z^n} \left[\log \left(1 - \frac{1}{z}\right) - \sum_{i=1}^{n-1} \frac{1}{i} \left(\left(\frac{z}{z-1}\right)^i - 1 \right) \right]$$

and by (7.11) we obtain (1.9). For the complex Wilton function, we obtain by the same considerations as above the representation as in (1.10).

7.1. Behaviour of \mathcal{B}_k and \mathcal{W} at rational points.

7.1.1. Behaviour of \mathcal{B}_k at rational points. Let

$$R_k(z) := \sum_{n=0}^k \frac{z^{k-n}}{n!} \varphi_0^{(n)}(-1),$$

which coincides with the last term in (7.5). Note that $R_k(1) \in \mathbb{R}$ is well defined for all $k \in \mathbb{N}$, as $\varphi_0 \in C^{\infty}$.

Lemma 7.1. The function $\varphi_{1,k}(z) + i \log(1-z)$ is continuous on $\overline{\mathbb{H}} = \mathbb{H} \cup \mathbb{R} \cup \{\infty\}$ and its value at 1 is $R_k(1) + \frac{\pi}{2}$.

Proof. Recall that for $t \in \mathbb{C} \setminus [0, +\infty)$ we have Euler's functional equation

$$\operatorname{Li}_{2}(t) + \operatorname{Li}_{2}\left(\frac{1}{t}\right) = -\frac{1}{2}(\log(-t))^{2} - \frac{\pi^{2}}{6}.$$
 (7.12)

If $z \notin [0, +\infty)$, then $\frac{z}{1-z}, \frac{1}{z-1} \notin [0, +\infty)$. Thus, applying (7.12) with $t = \frac{z}{1-z}$ and $t = \frac{1}{z-1}$, and substituting it into (7.5) gives

$$\varphi_{1,k}(z) = R_k(z) - z^k \frac{\pi}{6} + \frac{\pi}{6} - \frac{1}{\pi} \left(z^k \operatorname{Li}_2\left(\frac{1-z}{z}\right) - \operatorname{Li}_2(z-1) \right) - \frac{1}{2\pi} \left(z^k \log^2\left(\frac{-z}{1-z}\right) - \log^2\left(\frac{1}{1-z}\right) \right).$$

The function $z^k \operatorname{Li}_2\left(\frac{1-z}{z}\right) - \operatorname{Li}_2\left(z-1\right)$ is continuous on $\overline{\mathbb{H}}$ and it is in O(|z-1|) in the neighbourhood of 1. We then observe that

$$z^{k} \log^{2} \left(\frac{-z}{1-z}\right) - \log^{2} \left(\frac{1}{1-z}\right)$$

$$= (z^{k} - 1) \log^{2} \left(\frac{-z}{1-z}\right) + \log^{2} \left(\frac{-z}{1-z}\right) - \log^{2} \left(\frac{1}{1-z}\right),$$

and $(z^k - 1)\log^2\left(\frac{-z}{1-z}\right) = O\left(|1-z|\log\left(\frac{1}{|1-z|}\right)\right)$ in the neighbourhood of 1. Also, we have

$$\log^2\left(\frac{-z}{1-z}\right) - \log^2\left(\frac{1}{1-z}\right) = \left(\log(-z) + \log\left(\frac{1}{1-z}\right)\right)^2 - \log^2\left(\frac{1}{1-z}\right)$$
$$= \log^2(-z) + 2\log(-z)\log\left(\frac{1}{1-z}\right).$$

In the neighbourhood of 1 in $\overline{\mathbb{H}}$, we have $\log(-z) + i\pi = O(|z-1|)$. Thus,

$$\log^{2}\left(\frac{-z}{1-z}\right) - \log^{2}\left(\frac{1}{1-z}\right) = -\pi^{2} - 2i\pi\log\left(\frac{1}{1-z}\right) + O\left(|z-1|\log\left(|1-z|\right)\right).$$

Therefore, in the neighbourhood of 1, we have

$$\varphi_{1,k}(z) = R_k(z) + \frac{\pi}{2} + i\log\left(\frac{1}{1-z}\right) + O(|z-1|\log(|1-z|))$$
(7.13)

and the result follows.

Corollary 7.2. The real part of $\varphi_{1,k}$ is bounded in $\overline{\mathbb{C}} \setminus [1/2, 2]$. It has an extension to a continuous function on $\overline{\mathbb{C}} \setminus \{1\}$ and

$$\lim_{x \to 1^{\pm}} \operatorname{Re}(\varphi_{1,k}(x)) = R_k(1) \mp \frac{\pi}{2}.$$

Proof. By (7.13), we have

$$\lim_{x \to 1^{\pm}} \operatorname{Re}(\varphi_{1,k}(x)) = R_k(1) + \frac{\pi}{2} + \lim_{x \to 1^{\pm}} \operatorname{Re}\left(i \log\left(\frac{1}{1-x}\right)\right)$$

$$= R_k(1) + \frac{\pi}{2} - \lim_{x \to 1^{\pm}} \arg\left(\frac{1}{1-x}\right).$$

The result follows from the fact that

$$\lim_{x \to 1^+} \arg\left(\frac{1}{1-x}\right) = \pi \quad \text{ and } \quad \lim_{x \to 1^-} \arg\left(\frac{1}{1-x}\right) = 0.$$

Theorem 7.3. The real part of the complex k-Brjuno function has a decreasing jump of $\frac{\pi}{a^k}$ at each rational number $\frac{p}{a}$.

Proof. The proof follows similar arguments as [MMY01, Section 5.2]. The space $\widehat{\mathbb{H}/\mathbb{Z}}$ is defined by

$$\widehat{\mathbb{H}/\mathbb{Z}} = \mathbb{H}/\mathbb{Z} \sqcup (\mathbb{R}\backslash\mathbb{Q})/\mathbb{Z} \sqcup \left(\overline{\mathbb{Q}/\mathbb{Z}} \times \left[-\frac{\pi}{2}, +\frac{\pi}{2}\right]\right),$$

where $\overline{\mathbb{Q}/\mathbb{Z}} = \mathbb{Q}/\mathbb{Z} \cup \{\infty\}$. The space corresponds to a compactification of $(\overline{\mathbb{H}}\backslash\mathbb{Q})/\mathbb{Z}$ by attaching a semicircle on each $q \in \mathbb{Q} \cup \{\infty\}$. Then the value of a function on $\overline{\mathbb{Q}/\mathbb{Z}} \times [-\pi/2, \pi/2]$ is defined by

$$\varphi(\alpha, \theta) := \lim_{\substack{z \to \alpha \\ z \in \ell}} \varphi(z),$$

where ℓ is a ray emitting from α and the angle from $i\mathbb{R}$ to ℓ in the clockwise direction is θ . We note that the topology induced by $\widehat{\mathbb{H}/\mathbb{Z}}$ is the same as the topology induced by $\mathbb{H}/\mathbb{Z} \sqcup (\mathbb{R}\backslash\mathbb{Q})/\mathbb{Z}$. This implies the continuity of $\operatorname{Re}\sum_{Z\mathcal{M},k}\varphi_0$ on $\widehat{\mathbb{H}/\mathbb{Z}}$ and hence the real part $\operatorname{Re}\sum_{Z\mathcal{M},k}\varphi_0$ of the complex k-Brjuno function is continuous on $\mathbb{H}/\mathbb{Z} \sqcup (\mathbb{R}\backslash\mathbb{Q})/\mathbb{Z}$ in the usual sense. For $(\alpha_0,\pi/2) \in \overline{\mathbb{Q}/\mathbb{Z}} \times [-\pi/2,\pi/2]$, the value $\operatorname{Re}\sum_{Z\mathcal{M},k}\varphi_0(\alpha_0,\pi/2)$ (resp. $(\alpha_0,-\pi/2)$), with $\alpha_0 \in \mathbb{Q}/\mathbb{Z}$, is the right (resp. left) limit of $\operatorname{Re}\sum_{Z\mathcal{M},k}\varphi_0(\alpha)$, as $\alpha \in (\mathbb{R}\backslash\mathbb{Q})/\mathbb{Z}$ tends to α_0 . Recalling Corollary 7.2, one has

$$\operatorname{Re}\varphi_{1,k}(1,\pi/2) - \operatorname{Re}\varphi_{1,k}(1,-\pi/2) = -\pi$$

and more precisely, by a similar argument as in the proof of Corollary 7.2, from (7.13), we have

$$\operatorname{Re}\varphi_{1,k}(1,\theta) - \operatorname{Re}\varphi_{1,k}(1,0) \\
= -\lim_{t \to 0^+} \operatorname{arg}\left(\frac{1}{1 - (1 + te^{i(\pi/2 - \theta)})}\right) + \lim_{t \to 0^+} \operatorname{arg}\left(\frac{1}{1 - (1 + te^{i(\pi/2)})}\right) \\
= -\lim_{t \to 0^+} \operatorname{arg}\left(t^{-1}e^{i(\pi/2 + \theta)}\right) + \lim_{t \to 0^+} \operatorname{arg}\left(t^{-1}e^{i(\pi/2)}\right) = -\theta.$$

If $\alpha_0 \in \mathbb{Q}$, $\alpha_0 \neq 1$, then $\text{Re}\varphi_{1,k}(\alpha_0, \theta) = \text{Re}\varphi_{1,k}(\alpha_0, 0)$ for all $\theta \in [-\pi/2, \pi/2]$. Thus, by (7.3) one obtains that for all $p/q \in \mathbb{Q}$, $(p \land q = 1)$

$$\operatorname{Re} \sum_{Z\mathcal{M},k} \varphi_0(p/q,\theta) = \operatorname{Re} \sum_{Z\mathcal{M},k} \varphi_0(p/q,0) - \theta/q^k. \tag{7.14}$$

To show (7.14), we use the fact that

$$\operatorname{Re}(L_{k,g}\varphi(\alpha,\theta)) = \det(g)^{k+1}(a - c\alpha)^{k} \left[\operatorname{Re}\varphi\left(\frac{d\alpha - b}{a - c\alpha}, \det(g)\theta\right) - \varphi\left(-\frac{d}{c}\right) \right] - \sum_{n=1}^{k} \frac{\det(g)^{k-n+1}(a - c\alpha)^{k-n}}{c^{n}n!} \varphi^{(n)}\left(-\frac{d}{c}\right).$$
(7.15)

To verify this formula we note that for any $m \in \mathbb{N}$ we have

$$\begin{split} &\lim_{t\to 0+} \arg\left(\begin{pmatrix} 0 & 1 \\ 1 & m \end{pmatrix}.\left(\alpha + t\cdot e^{i(\pi/2-\theta)}\right) - \begin{pmatrix} 0 & 1 \\ 1 & m \end{pmatrix}.\alpha\right) \\ &= \lim_{t\to 0+} \arg\left(-\frac{t\cdot e^{i(\pi/2-\theta)}}{(\alpha+m)(\alpha+m+t\cdot e^{i(\pi/2-\theta))}}\right) = -\theta - \frac{\pi}{2}. \end{split}$$

(This is the ray we get after applying the Möbius transform.) Now, if φ is analytic on the upper half plane and real on the real line, we can apply Schwarz' reflection principle and obtain

$$\operatorname{Re}\varphi(\alpha,\theta) = \lim_{t \to 0+} \operatorname{Re}\varphi(\alpha + t \cdot e^{i(\pi/2 - \theta)}) = \lim_{t \to 0+} \operatorname{Re}\overline{\varphi}(\alpha + t \cdot e^{-i(\pi/2 - \theta)})$$
$$= \lim_{t \to 0+} \operatorname{Re}\varphi(\alpha + t \cdot e^{-i(\pi/2 - \theta)}).$$

Hence, if $\psi(z) = \varphi(g(m).z)$, then $\text{Re}\psi(\alpha,\theta) = \text{Re}\varphi(g(m).\alpha,-\theta)$ and generally if $\psi(z) = \varphi(g.z)$ with $g \in \mathcal{M}$, then $\text{Re}\psi(\alpha,\theta) = \text{Re}\varphi(g.\alpha,\det(g)\theta)$.

Since every element of \mathcal{M} can be written as a product of matrices g(m), (7.15) follows. By (7.3), we have

$$\operatorname{Re} \sum_{\mathcal{ZM},k} \varphi_0(\alpha,\theta) - \operatorname{Re} \sum_{\mathcal{ZM},k} \varphi_0(\alpha,0)$$
$$= \sum_{n \in \mathbb{Z}, g \in \mathcal{M}} \operatorname{Re} \left[L_{k,g} \varphi_{1,k}(\alpha - n, \theta) - L_{k,g} \varphi_{1,k}(\alpha - n, 0) \right].$$

By Corollary 7.2, the real part $\operatorname{Re}\left[L_{k,g}\varphi_{1,k}(\alpha-n,\theta)-L_{k,g}\varphi_{1,k}(\alpha-n,0)\right]$ is 0 if and only if $\frac{d(\alpha-n)-b}{a-c(\alpha-n)}\neq 1$ and by (7.15),

$$\operatorname{Re}\left[L_{k,g}\varphi_{1,k}(\alpha-n,\theta)-L_{k,g}\varphi_{1,k}(\alpha-n,0)\right]=-\left(\det(g)(a-c(\alpha-n))\right)^{k}\theta,$$

if $\frac{d(\alpha-n)-b}{a-c(\alpha-n)}=1$. There exist unique $n\in\mathbb{N}$ and $g=\left(\begin{smallmatrix} a&b\\c&d\end{smallmatrix}\right)\in\mathcal{M}$ such that $\frac{d(\alpha-n)-b}{a-c(\alpha-n)}=1$, i.e. $\alpha=n+\frac{a+b}{c+d}$. If $\alpha=p/q$, then c+d=q. Thus,

$$\operatorname{Re} \sum_{\mathcal{ZM}} \varphi_0(p/q, \theta) - \operatorname{Re} \sum_{\mathcal{ZM}} \varphi_0(p/q, 0) = -\frac{\det(g)^{2k}}{(c+d)^k} \theta = -\frac{\theta}{q^k}.$$

Thus, $\operatorname{Re}(\mathcal{B}_k)$ has at each rational $p/q \in \mathbb{Q}/\mathbb{Z}$ a decreasing jump of π/q^k .

7.1.2. A comment on the Wilton function. It would be interesting to get to know the behaviour of the Wilton function at rational points. However, the results can not easily be transferred to the Wilton function. By a similar argument as in the proof of Lemma 7.1, we will get from the next lemma that $\overline{\varphi}_1(z) + i \log(1-z) - \frac{1}{2\pi}(z+1) \log^2\left(1-\frac{1}{z}\right)$ is continuous on $\overline{\mathbb{H}}$ and in a neighbourhood of 1 it behaves like

$$\overline{\varphi}_{1}(z) = \frac{3\pi}{4} - \frac{\log 2}{\pi} - i\log\left(1 - z\right) + \frac{1}{2\pi}(z+1)\log^{2}\left(1 - \frac{1}{z}\right) + O\left(|1 - z|\log\left(|1 - z|\right)\right).$$

Thus, we do not have an analog of Corollary 7.2 since $\operatorname{Re}\left(\log^2\left(1-\frac{1}{z}\right)\right)$ does not go to a finite value for $z \to 1$.

7.2. Behaviour of the imaginary part of W. In this section, we will explain the behaviour of the imaginary part of the Wilton function near the real axis, distinguishing rational points, Wilton numbers and Diophantine numbers. Many arguments are slight adaptations of those given in [MMY01], thus they will only be sketched and many proofs will be given in the Appendix.

The same behaviour which we observe for the imaginary part of Wilton function will also characterize the approach to the real axis of the imaginary part of the k-Brjuno function, but we will limit ourselves to prove the results for the former, being the arguments very similar.

The following lemma is an analog to [MMY01, Lem. 5.2] and Lemma 7.1. However, here we are considering the function $\overline{\varphi}_1(z) + i \log(1-z) - \frac{1}{2\pi}(z+1) \log^2\left(1-\frac{1}{z}\right)$ instead of $\varphi_1(z) + i \log(1-z)$ in [MMY01]. We will later remark how this further influences the proof of Theorem 7.5.

Lemma 7.4. The function $\overline{\varphi}_1(z) + i \log(1-z) - \frac{1}{2\pi}(z+1) \log^2\left(1-\frac{1}{z}\right)$ is continuous on $\overline{\mathbb{H}}$ and its value at 1 is $\frac{3\pi}{4} - \frac{\log 2}{\pi}$.

Proof. By combining (7.6) and Euler's functional equation (7.12) it follows that

$$\overline{\varphi}_{1}(z) = \frac{1}{\pi} \left[z \operatorname{Li}_{2} \left(\frac{1-z}{z} \right) + \operatorname{Li}_{2} (z-1) \right] + \frac{1}{2\pi} \left[z \log^{2} \left(\frac{1-z}{-z} \right) + \log^{2} (-z+1) \right] + \frac{\pi}{12} z + \frac{\pi}{6} - \frac{\log 2}{\pi}.$$
(7.16)

The function $z \operatorname{Li}_2\left(\frac{1-z}{z}\right) + \operatorname{Li}_2\left(z-1\right)$ is regular and vanishing at z=1. We have

$$z\log^2\left(\frac{1-z}{-z}\right) + \log^2(-z+1) = (z+1)\log^2\left(\frac{1-z}{-z}\right) + \log^2(-z+1) - \log^2\left(\frac{1-z}{-z}\right).$$

As in the proof of [MMY01, Lem. 5.2], it follows that

$$\log^2(-z+1) - \log^2\left(\frac{1-z}{-z}\right) = \pi^2 - 2\pi i \log(1-z) + O\left(|z-1|\log\left(|z-1|\right)\right),$$

which completes the proof.

From now on, we write $f(z) \lesssim g(z)$ if there exists a constant C > 0 such that $f(z) \leq Cg(z)$ for all z. The following is an analog to [MMY01, Thm. 5.10].

Theorem 7.5. For $n \geq 0$, $m_1, \dots, m_n \geq 1$ and $z_0 \in H(m_1, \dots, m_n)$, we have

$$\operatorname{Im} \mathcal{W}(z_0) = -W_{\text{finite}}\left(\frac{p_n}{q_n}\right) + (-1)^n (p_{n-1} - q_{n-1}\operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_n + 1) + r_n(z_0),$$

with
$$|r_n(z_0)| \lesssim \frac{\log q_n}{q_n} |z_n| \log^2(1 + |z_n|^{-1}).$$

As the proof is very similar to the one in [MMY01, Thm. 5.11], it is given in the Appendix. However, the summand and the remainder term differ slightly from those in [MMY01] about which we will comment in the following.

Remark 7.6. Here we give a remark of the summand $(-1)^n(p_{n-1}-q_{n-1}\operatorname{Re}(z_0))\operatorname{Im}\overline{\varphi}_1(z_n+1)$ of Theorem 7.5. We have

$$(-1)^{n}(p_{n-1} - q_{n-1}\operatorname{Re}(z_{0}))\operatorname{Im}\overline{\varphi}_{1}(z_{n} + 1) = \left(q_{n}^{-1}\log\frac{1}{|z_{n}|}\right)\left(1 + \frac{2}{\pi}\operatorname{Arg}(1 + 1/z_{n}) + o(1)\right).$$

It can be derived as follows.

By (4.13), we have

$$q_n(p_{n-1} - q_{n-1}\operatorname{Re}(z_0)) = (-1)^n \operatorname{Re}\left(\frac{q_n}{q_{n-1}z_n + q_n}\right) \to (-1)^n \text{ as } z_n \to 0.$$

By $\operatorname{Re}(\log^2(w)) = \log^2|w| - (\operatorname{Arg}(w))^2$ and $\operatorname{Im}(\log^2(x)) = 2\log|w|\operatorname{Arg}(w)$, we have

$$\operatorname{Im}\left((z_n+2)\log^2\left(1+\frac{1}{z_n}\right)\right) = 2(\operatorname{Re}(z_n)+2)\operatorname{Arg}(1+1/z_n)(\log|z_n+1|+\log|z_n|^{-1}) + \operatorname{Im}(z_n)(\log^2|1+1/z_n|-(\operatorname{Arg}(1+1/z_n))^2).$$

In the above equation, $2(\operatorname{Re}(z_n) + 2)\operatorname{Arg}(1 + 1/z_n)\log|z_n + 1| - \operatorname{Im}(z_n)(\operatorname{Arg}(1 + 1/z_n))^2$ is uniformly bounded. Since $|\operatorname{Im}(z_n)\log^2|1 + 1/z_n| \leq |z_n|\log^2(1 + 1/|z_n|) \to 0$ as $z_n \to 0$, from Lemma 7.4, we have

$$\frac{\operatorname{Im}\overline{\varphi}_{1}(z_{n}+1)}{\left(\log\frac{1}{|z_{n}|}\right)} = \frac{-\log|z_{n}| + \frac{1}{2\pi}2(\operatorname{Re}(z_{n})+2)\operatorname{Arg}(1+1/z_{n})\log|z_{n}|^{-1}}{\log|z_{n}|^{-1}} + o(1)$$

$$= 1 + \frac{2}{\pi}\operatorname{Arg}(1+1/z_{n}) + o(1) = 1 + O(1),$$

but not in 1 + o(1).

Finally, we remark here on the estimate of the remainder term in Theorem 7.5 which differs from the remainder term in [MMY01, Thm. 5.10] as we have an additional factor of $\log q_n \cdot \log(1+|z_n|^{-1})$. It is caused by the signs of the terms $z\varphi_0(1/z-1)$ and $\varphi_0(z-1)$ in the definition of $\overline{\varphi}_1$ in (7.4). The functions $\varphi_0(1/z-1)$ and $\varphi_0(z-1)$ are not continuous at 1 because they can be written as a sum of a function continuous near 1 and a constant

multiple of $\log^2(1-z)$ (see the proof of Lemma 7.4). Contrary to that in the case of the Brjuno function, the non-continuous parts cancel out (see [MMY01, Lem. 5.2]).

Finally, we will show an analog of [MMY01, Thm. 5.19]. We will give a sufficient condition for W(x) to be approximated by the imaginary part of W(z) if z approaches x. For R > 0 and 0 < r < 1/2, let

$$U_R = \{u \in \mathbb{H} : \operatorname{Im}(u) \ge |\operatorname{Re}(u)|^R\} \text{ and } \widetilde{U}_r = \left\{u \in \mathbb{H} : \operatorname{Im}(u) \ge \exp\left(-\frac{1}{|\operatorname{Re}(u)|^r}\right)\right\}.$$

Theorem 7.7. We have the following statements:

(i) For any Wilton number x and any R > 0, we have

$$\lim_{u\to 0,\ u\in U_R} \operatorname{Im} \mathcal{W}(u+x) = -W(x).$$

(ii) Let x be an irrational Diophantine number and 0 < r < 1/2 such that

$$\liminf_{q \to \infty} \|qx\|_{\mathbb{Z}} \ q^{1/r-1} = \infty,$$

where $\|\cdot\|_{\mathbb{Z}}$ denotes the distance from the nearest integer. Then,

$$\lim_{u\to 0,\ u\in \widetilde{U}_r} \text{Im} \mathcal{W}(u+x) = -W(x).$$

Since the singular behaviour of the k-Brjuno function as the real line is approached is tamer than the singular behaviour of the Brjuno function, we expect that the same properties proven in [MMY01] and stated in the theorem above for the Wilton function hold by a modest adaptation of the proofs given there.

Proof of Theorem 7.7. Let $x \in (0,1)$ be an irrational number whose continued fraction expansion is

$$x = \frac{1}{m_1 + \frac{1}{m_2 + \frac{1}{\cdots + \frac{1}{m_n + \cdots}}}}.$$

Let $x_{\ell} = G^{\ell}(x)$. Let $\{p_n/q_n\}_{n\geq 0}$ be the sequence of the partial quotients of x. Let u be a point near 0 such that $|\operatorname{Im}(u)| \leq 1/2$ and $z = u + x \in \Delta$. Then there are $N_1 \cdots, N_L$ such that $z \in H(N_1, \cdots, N_L)$. Let p/q be a rational number such that

$$p/q = \frac{1}{N_1 + \frac{1}{N_2 + \frac{1}{N_L}}}.$$

We distinguish two cases.

(I) Assume that $p/q = p_L/q_L$, i.e., $m_i = N_i$ for $i = 1, \dots, L$. From Theorem 7.5, Remark 7.6 and Lemma 3.1, there exist uniform constants $c_1, c_2, c_3 > 0$ such that

$$|\operatorname{Im} \mathcal{W}(u+x) + W(x)|$$

$$= \left| -W_{\text{finite}} \left(\frac{p_L}{q_L} \right) + \sum_{\ell=0}^{L-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} + \sum_{\ell=L}^{\infty} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right.$$

$$+ (-1)^n (p_{L-1} - q_{L-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_L + 1) + r_n(z_0) \right|$$

$$\leq c_1 x_L q_n^{-1} + c_2 q_L^{-1} \log |z_L|^{-1} + c_3 q_L^{-1} \log (q_L) |z_L| \log^2(1 + |z_L|^{-1})$$

$$+ \left| \sum_{\ell=L}^{\infty} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right|.$$

$$(7.17)$$

This case follows in an analogous way to the proof of [MMY01, Thm. 5.19(I)] where we note that the considered terms also stay small if we consider a Wilton instead of a Brjuno number.

(II) Assume that p/q is not one of the partial quotients of x. We denote by $\{p'_{\ell}/q'_{\ell}\}_{0 \leq \ell \leq L}$ the partial quotient of p/q and by n the largest integer such that $p'_n/q'_n = p_n/q_n$. Clearly, one has n < L and $p'_{\ell}/q'_{\ell} = p/q$.

Clearly, one has n < L and $p'_L/q'_L = p/q$. Note that $q_n^{-1}\log(q_n) \ge (q'_L)^{-1}\log(q'_L)$ if $q_n \ge 3$. As in the proof of [MMY01, Thm. 5.19], from the fact $|x-p/q| \ge (2q^2)^{-1}$, if $u \in U_R$, then there exists a uniform constant c > 0 such that

$$q^{-1}\log|z_L|^{-1} \le q^{-1}(c + (2R - 2)\log q) \le q_n^{-1}(c + (2R - 2)\log q_n)$$

and if $u \in \widetilde{U_r}$, then we have

$$q^{-1}\log|z_L|^{-1} \le q^{-1}cq^{2r} \le q_n^{-1}cq_n^{2r}$$
.

From Theorem 7.5, Remark 7.6 and Lemma 3.1, there exist uniform constants $c_1, c_2, C_1, C_2, C_3 > 0$ such that

 $|\mathrm{Im}\mathcal{W}(u+x) + W(x)|$

$$= \left| -W_{\text{finite}} \left(\frac{p'_L}{q'_L} \right) + \sum_{\ell=0}^{n-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} + \sum_{\ell=n}^{\infty} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right.$$

$$+ (-1)^{L} (p'_{L-1} - q'_{L-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_{1}(z_{L} + 1) + r_{L}(z_{0}) \right|$$

$$\leq \left| -W_{\text{finite}} \left(\frac{p'_L}{q'_L} \right) + W_{\text{finite}} \left(\frac{p_n}{q_n} \right) \right| + \left| -W_{\text{finite}} \left(\frac{p_n}{q_n} \right) + \sum_{\ell=0}^{n-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right.$$

$$+ \left| \sum_{\ell=n}^{\infty} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right| + c_{1} \frac{\log |z_{L}|^{-1}}{q'_{L}} + c_{2} \frac{\log q'_{L}}{q'_{L}} |z_{L}| \log^{2}(1 + |z_{L}|^{-1})$$

$$\leq \left| -W_{\text{finite}} \left(\frac{p'_L}{q'_L} \right) + W_{\text{finite}} \left(\frac{p_n}{q_n} \right) \right| + \left| \sum_{\ell=n}^{\infty} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right|$$

$$+ C_{3} \frac{x_{n}}{q_{n}} + C_{1} \frac{1}{q_{n}^{\gamma}} + C_{2} \frac{\log q_{n}}{q_{n}}$$

since $|z_L| \log^2(1+|z_L|^{-1})$ are bounded, where $\gamma = 1$ for the case of $u \in U_R$ and $\gamma = 1 - 2r$ for the case of $u \in \widetilde{U}_r$. Thus, it is enough to show that $W_{\text{finite}}(p/q)$ is close to $W_{\text{finite}}(p_n/q_n)$.

Let

$$\rho = \max_{n \le \ell < L} (q'_{\ell})^{-1} \log \frac{q'_{\ell+1}}{q'_{\ell}} (\ell - n + 1)^2$$

By Lemma 3.1, we have

$$|W_{\text{finite}}(p_n/q_n) - W_{\text{finite}}(p/q)|$$

$$\leq \left| W_{\text{finite}}(p_n/q_n) - \sum_{\ell=0}^{n-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} + \sum_{\ell=0}^{L-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} - W_{\text{finite}}(p/q) \right| \\
\leq C x_n q_n^{-1} + C x_L(q_L')^{-1} + \left| \sum_{\ell=n}^{L-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right| \leq C' q_n^{-1} + \left| \sum_{\ell=n}^{L-1} (-1)^{\ell} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \right|.$$

We have

$$\sum_{\ell=n}^{L-1} \beta_{\ell-1}(x) \log \frac{1}{x_{\ell}} \lesssim \sum_{\ell=n}^{L-1} \frac{1}{q'_{\ell}} \log \frac{q'_{\ell+1}}{q'_{\ell}} \le \rho \sum_{\ell=r}^{L-1} \frac{1}{(\ell-n+1)^2} \lesssim \rho.$$

With the same argument as in the proof of [MMY01, Thm. 5.19] we have the conclusion.

APPENDIX A. PROOF OF PROPOSITION 6.3

Proof. From the fact that \mathcal{M} is the free monoid generated by the matrices g(m), we have that $z \in D_{\infty}$ and $g \in \mathcal{M}$ imply $g^{-1}.z \in D_{\infty}$. Also, by this fact combined with [MMY01, Lem. 3.2], we then have that

$$z \in V_{\rho}(D_{\infty}) \text{ implies } g^{-1}.z \in V_{\rho}(D_{\infty}).$$
 (A.1)

Let $\mathcal{M}^{(r)} = \{g \in \mathcal{M} \mid g = g(m_1) \cdots g(m_r), m_i \in \mathbb{N}\}$. Then we have for all $\psi \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus I)$ that

$$(T_k^{(k+2)})^r \psi = \sum_{g \in \mathcal{M}^{(r)}} L_g^{(k+2)} \psi = \sum_{g' \in \mathcal{M}^{(r-1)}} L_{g'}^{(k+2)} (T_k^{(k+2)} \psi).$$
 (A.2)

Let $g' = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$, $z \in V_{\rho}(D_{\infty})$ and $z' = (g')^{-1}.z$. Then for all $m \in \mathbb{N}$, we have

$$g'g(m) = \begin{pmatrix} b' & a' + mb' \\ d' & c' + md' \end{pmatrix}.$$

Thus, by using the definition of $T_k^{(k+2)}$ given in Proposition 6.1 together with (5.26) and (5.27) we obtain

$$L_{g'}^{(k+2)}\left(T_k^{(k+2)}\psi\right) = L_{g'}^{(k+2)}\left(\sum_{m=1}^{\infty} L_{g(m)}^{(k+2)}\psi\right) = L_{g'}^{(k+2)}\left(\sum_{m=1}^{\infty} -z^{-(k+2)}\psi\left(\frac{1}{z}-m\right)\right)$$

$$= \det(g)^{k+1}\left(a'-c'z\right)^{-(k+2)}\left(-\sum_{m=1}^{\infty} \left(\frac{a'-c'z}{d'z-b'}\right)^{k+2}\psi\left(\frac{1}{z'}-m\right)\right)$$

$$= -\det(g)^{k+1}(d'z-b')^{-(k+2)}\sum_{m=1}^{\infty} \psi\left(\frac{1}{z'}-m\right).$$

We remark that if r = 1, then we consider $g' \in \mathcal{M}^{(0)} = \{id\}$ and the following arguments work well.

It follows from (5.1) and (A.1) that

$$\sum_{m=1}^{\infty} \left| \psi \left(\frac{1}{z'} - m \right) \right| \le \sum_{m=1}^{\infty} c_{1,\rho,k} \left| \frac{1}{z'} - m \right|^{-(k+2)} \sup_{V_{\rho}(D_{\infty})} \left| \psi \right|$$

for some constant $c_{1,\rho,k}$. Since $\sup_{z\in D_{\infty}} \operatorname{Re}(1/z) = 1$ (see Figure 7b), there exists $c_2 > 0$ such that $|1/z - m|^{-1} \le c_2 m^{-1}$ for all $z \in D_{\infty}$ and $m \ge 2$. Thus, a similar argument as the one showing (6.2) in the proof of Proposition 6.1 yields that for all $z' \in V_{\rho}(D_{\infty})$

$$\sum_{m=1}^{\infty} \left| \psi \left(\frac{1}{z'} - m \right) \right| \le c_{3,\rho,k} \sup_{V_{\rho}(D_{\infty})} |\psi|$$

for some constant $c_{3,\rho,k}$.

Also, there exists $c_{4,\rho} > 0$ such that $|z - \frac{b'}{d'}|^{-1} \le c_{4,\rho}$ for all $z \in V_{\rho}(D_{\infty})$ and $\frac{b'}{d'} \in [0,1]$. Here, d' can be considered as the denominator of the rth principal convergent of some number. Finally,

$$\min_{g' \in \mathcal{M}^{(r-1)}} d' \ge \frac{1}{2} \left(\frac{\sqrt{5} + 1}{2} \right)^{r-1} \text{ and } \sum_{g' \in \mathcal{M}^{(r-1)}} d'^{-2} \le c_5, \tag{A.3}$$

for some $c_5 > 0$, see for example [MMY01, Prop. A1.1 in Appendix 1]. Furthermore, since $g' \in \mathcal{M}^{(r-1)}$, we have that $\left| \det(g')^{k+1} \right| = 1$. Combining these results yields the statement of (i).

In the following we will show (ii) for $r \geq 2$, and we will discuss the case r = 1 separately. Starting in the analogous way as in the proof of (i), using (5.23), we obtain for $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus [0,1])$ that

$$L_{k,g'}(T_k\varphi) = L_{k,g'}\left(\sum_{m=1}^{\infty} L_{k,g(m)}\varphi\right)$$

$$= \frac{(c')^{-(k+1)}\det(g')^{k+1}}{k!}(a'-c'z)^{-1}$$

$$\cdot \int_0^1 (1-t)^k \sum_{m=1}^{\infty} L_{k,g(m)}^{(k+2)}\varphi^{(k+1)}\left(-\frac{d'}{c'} + \frac{\det(g')t}{c'(a'-c'z)}\right) dt.$$

We have

$$\xi_{z,t} := -\frac{d'}{c'} + \frac{\det(g')t}{c'(a'-c'z)} = -\frac{d'}{c'}t + \frac{\det(g')t}{c'(a'-c'z)} + \frac{d'}{c'}t - \frac{d'}{c'} = tg'^{-1}.z - (1-t)\frac{d'}{c'}.$$

If we denote $\iota(z):=1/z$, see Figure 7b, then we can inductively show that $g'^{-1}.z$ is contained in a ρ -neighbourhood of $\bigcup_{m\geq 1}(\iota(D_\infty)-m)\subset D_\infty$ with respect to the Poincaré metric. Since $d'/c'\in(-\infty,-1]$, the line segment between -d'/c' and $g^{-1}.z$ is contained in $V_\rho(D_\infty\cap\bigcup_{m\geq 1}(\iota(D_\infty)-m))$.

Thus, there is a constant $c_{6,\rho} > 0$ such that $|\xi_{z,t}| \ge c_{6,\rho}$. The constant $c_{6,\rho}$ increases when $\rho \to 0$. Since $\xi_{z,t}$ is in a ρ -neighbourhood of i, -i, or $\text{Re}(\xi_{z,t}) \le 0$, we have $|1/\xi_{z,t} - m| \ge m$. Since $\varphi^{(k+1)} \in \mathcal{O}^{k+2}(\overline{\mathbb{C}} \setminus [0,1])$, by using (5.26) and (6.3), we have

$$\sup_{z \in V_{\rho}(D_{\infty})} |L_{k,g'}(T_{k}\varphi)| \leq \frac{(c')^{-(k+2)}}{k! \inf_{z \in V_{\rho}(D_{\infty})} |z - \frac{a'}{c'}|} \sum_{m \geq 1} \int_{0}^{1} \frac{|\varphi^{(k+1)}(1/\xi_{z,t} - m)|}{|\xi_{z,t}|^{k+2}} dt$$

$$\leq \frac{(c')^{-(k+2)}}{k! \inf_{z \in V_{\rho}(D_{\infty})} |z - \frac{a'}{c'}|} \sum_{m \geq 1} \frac{c'_{1,k+1}(k+1)!}{c_{6,\rho}^{k+2}(m-0.2)^{k+2}} \sup_{z \in V_{\rho}(D_{\infty})} |\varphi(z)|$$

$$\leq \frac{c_{7,k}c'_{1,k+1}(k+1)}{c_{6,\rho}^{k+2}} \cdot \frac{(c')^{-(k+2)}}{\inf_{z \in V_{\rho}(D_{\infty})} |z - \frac{a'}{c'}|} \cdot \sup_{z \in V_{\rho}(D_{\infty})} |\varphi(z)|,$$

where $c_{7,k} = \sum_{m \ge 1} (m - 0.2)^{-(k+2)}$.

Since $a'/c' \in [0,1]$, we have $|z - \frac{a'}{c'}|^{-1} \le c_{4,\rho}$ for $z \in V_{\rho}(D_{\infty})$. Using (A.3) and (6.3), we obtain

$$\sup_{z \in V_{\rho}(D_{\infty})} |(T_{k}^{r}\varphi)(z)| \leq \frac{2^{k} c_{7,k} c_{1,k+1}'(k+1)}{c_{6,\rho}^{k+2}} \left(\frac{\sqrt{5}-1}{2}\right)^{k(r-2)} \sum_{g' \in \mathcal{M}^{(r-1)}} (c')^{-2} \cdot \sup_{z \in V_{\rho}(D_{\infty})} |\varphi(z)| \\
\leq \overline{C}_{\rho,k,\geq 2} \left(\frac{\sqrt{5}-1}{2}\right)^{rk} \sup_{z \in V_{\rho}(D_{\infty})} |\varphi(z)|, \tag{A.4}$$

where $\overline{C}_{\rho,k,\geq 2} = 2^k (k+1) (\frac{\sqrt{5}+1}{2})^{2k} c'_{1,k+1} c_5 c_{4,\rho} c_{6,\rho}^{-(k+2)} c_{7,k}$, implying (ii) for $r \geq 2$. For the case r = 1, we will estimate

$$|T_k \varphi(z)| \le \sum_{m \ge 1} |L_{k,g(m)} \varphi(z)|$$

by using (5.24) and (5.25). We consider ℓ denoting the the right arc of ∂D_{∞} (which is the arc of the circle $\left|z-\frac{\sqrt{3}}{3}\right|=\frac{\sqrt{3}}{3}$ between $\frac{\sqrt{3}}{2}+\frac{1}{2}i$ and $\frac{\sqrt{3}}{2}-\frac{1}{2}i$ containing $\frac{2\sqrt{3}}{3}$), see Figure 7a. Since $1/z \in V_{\rho}(\iota(D_{\infty}))$, from the shape of $\iota(D_{\infty})$, we can conclude that $t/z \in V_{\rho}(\iota(D_{\infty}))$ for $0 \le t \le 1$. Thus, $t/z-m \in V_{\rho}(\iota(D_{\infty})-m)$. If $m \ge 2$ or z is not in the neighbourhood of ℓ , then there is c > 0 such that $|t/z-m| \ge cm$, thus by Cauchy's estimate as in (6.3), we have

$$\left| \varphi^{(k+1)} \left(\frac{t}{z} - m \right) \right| \le c'_{1,k+1} \frac{(k+1)! \sup_{V_{\rho}(D_{\infty})} |\varphi|}{cm^{k+2}}.$$

Thus, if z is not in the neighbourhood of ℓ , then by using (5.25), we have

$$|T_k \varphi(z)| \leq \frac{1}{c_{6,\rho} k!} \sum_{m \geq 1} c'_{1,k+1} \frac{(k+1)! \sup_{V_{\rho}(D_{\infty})} |\varphi|}{cm^{k+2}}$$

$$\leq \overline{C}'_{\rho,k} \left(\frac{\sqrt{5} - 1}{2}\right)^k \sup_{V_{\rho}(D_{\infty})} |\varphi|,$$
(A.5)

where $\overline{C}'_{\rho,k} = \left(\frac{\sqrt{5}-1}{2}\right)^{-k} (k+1) c'_{1,k+1} c_{6,\rho}^{-1} c^{-1} \sum_{m\geq 1} \frac{1}{m^{k+2}}$. We are left to show the case that z is in a neighbourhood of ℓ . For $z \in V_{\rho}(D_{\infty})$ in a neighbourhood of ℓ , we can assume that $|z| \leq 2$. Thus, by (5.24) and (6.3), we have

$$|L_{k,g(1)}\varphi(z)| \leq |z|^k \left| \varphi\left(\frac{1}{z} - 1\right) \right| + \sum_{n=0}^k \frac{|z|^{k-n}}{n!} |\varphi^{(n)}(-1)|$$

$$\leq |z|^k \sup_{V_{\rho}(D_{\infty})} |\varphi| + \sum_{n=0}^k \frac{|z|^{k-n}}{n!} c'_{1,n} \frac{n! \sup_{V_{\rho}(D_{\infty})} |\varphi|}{0.8^{n+1}}$$

$$\leq \left(|z|^k + \sum_{n=1}^k \frac{c'_{1,n} |z|^{k-n}}{0.8^{n+1}} \right) \sup_{V_{\rho}(D_{\infty})} |\varphi| \leq C_k \left(\frac{\sqrt{5} - 1}{2} \right)^k \sup_{V_{\rho}(D_{\infty})} |\varphi|, \quad (A.6)$$

where $C_k = \left(\frac{\sqrt{5}-1}{2}\right)^{-k} 2^k \left(1 + \sum_{n=1}^k \frac{c'_{1,n}}{2^{n_0}.8^{n+1}}\right)$. By replacing the summand of m=1 in (A.5) with (A.6) and combining this result with the estimate for $r \geq 2$ in (A.4), we have the conclusion when we take $\overline{C}_{\rho,k} = \max\{C_k + \overline{C}'_{\rho,k}, \overline{C}_{\rho,k,\geq 2}\}$.

APPENDIX B. PROOF OF THEOREM 7.5.

Analogously as in [MMY01] the strategy of the proof of the theorem is to start from the formula

$$\operatorname{Im} \mathcal{W}(z) = \sum_{n \in \mathbb{Z}} \sum_{\ell > 0} \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z+n). \tag{B.1}$$

We will only give the parts of the proof which are different.

We have the following lemmas and propositions which are analogs to [MMY01, Lem. 5.11, 5.14 and Prop. 5.15, 5.16] in the same way by using S = -T and substituting φ_1 to $\overline{\varphi}_1$. For the following we denote for a bounded closed interval $I \subset \mathbb{R}$, the space of functions $\varphi \in \mathcal{O}^1(\overline{\mathbb{C}} \setminus I)$ whose real part is bounded we denote by E(I), endowed with the norm $\|\varphi\|_{E(I)} = \sup_{\mathbb{C} \setminus I} |\text{Re}\varphi|$. If it is clear which interval I is meant, we also write $\|\varphi\|$ instead of $\|\varphi\|_{E(I)}$. Note that $\overline{\varphi}_1 \in E([1/2, 2])$. If $\psi \in E([1/2, 2])$ is real on the real axis outside [1/2, 2], then ψ holds the arguments about φ_1 in the proof of [MMY01, Lem. 5.11, 5.14 and Prop. 5.15, 5.16].

Lemma B.1. For $z \in \Delta \setminus [0,1]$, if we write

$$\operatorname{Im} \mathcal{W}(z) = \operatorname{Im} \overline{\varphi}_{1}(z) + \operatorname{Im} \overline{\varphi}_{1}(z+1) + \operatorname{Im} \overline{\varphi}_{1}(z+2) + \operatorname{Im} S \overline{\varphi}_{1}(z-1) + \operatorname{Im} S \overline{\varphi}_{1}(z) + \sum_{\ell > 1} \left[\operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z-1) + \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z) + \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z+1) \right] + r^{0}(z),$$

then we have

$$|r^{0}(z)| \le C|\text{Im}(z)| \|\overline{\varphi}_{1}\|_{E([1/2,2])}.$$

Lemma B.2. For $z \in \Delta \setminus [0, 1]$, we have

$$\operatorname{Im} \mathcal{W}(z) = (1 - \varepsilon_1) \operatorname{Im} \overline{\varphi}_1(z) + \operatorname{Im} \overline{\varphi}_1(z+1) + [1 - \varepsilon_2(1 - \varepsilon_1)] \operatorname{Im} S \overline{\varphi}_1(z) + \sum_{\ell > 1} \operatorname{Im} S^{\ell} \overline{\varphi}_1(z) + r^0(z) + r^1(z),$$
(B.2)

and

$$|r^1(z)| \le C|\operatorname{Im}(z)|\log(1+|\operatorname{Im}(z)|^{-1})\|\overline{\varphi}_1\|_{E([1/2,2])}.$$

Proposition B.3. For $n \geq 1$, $m_1, \ldots, m_n \geq 1$, $z_0 \in D(m_1, \ldots, m_n)$, $\ell > 0$ we have

$$\left| (-1)^n \operatorname{Im} \left(S^{n+\ell} \overline{\varphi}_1(z_0) \right) - \left[\operatorname{Im} \left(S^{\ell} \overline{\varphi}_1(z_n) \right) \right] \left(p_{n-1} - q_{n-1} \operatorname{Re}(z_0) \right) \right|$$

$$\leq C q_n^{-1} |\operatorname{Im}(z_n)| \log(1 + |\operatorname{Im}(z_n)|^{-1}) \times \begin{cases} \| S^{\ell-2} \overline{\varphi}_1 \| & \text{if } \ell > 1, \\ \| \overline{\varphi}_1 \| & \text{if } \ell = 1. \end{cases}$$

Proposition B.4. For $n \geq 1$, $m_1, \ldots, m_n \geq 1$, $z_0 \in D(m_1, \ldots, m_n)$ we have

$$\begin{aligned} &|(-1)^n \mathrm{Im}(S^n \overline{\varphi}_1(z_0)) - (p_{n-1} - q_{n-1} \mathrm{Re}(z_0))[(1 - \varepsilon_{n+1}) \mathrm{Im}(\overline{\varphi}_1(z_n)) + \varepsilon_n \mathrm{Im}(\overline{\varphi}_1(z_n + 1))]| \\ &\leq C q_n^{-1} |\mathrm{Im}(z_n)| \log(1 + |\mathrm{Im}(z_n)|^{-1}) ||\overline{\varphi}_1||. \end{aligned}$$

The following is an analog to [MMY01, Prop. 5.17]. We remember the definition of $H(m_1, \dots, m_n)$ given in (4.15).

Proposition B.5. For $n \geq 0$, $m_1, \ldots, m_n \geq 1$, $z_0 \in H(m_1, \ldots, m_n)$ we have

$$\operatorname{Im} \mathcal{W}(z_0) = \sum_{\ell=0}^{n} \varepsilon_{\ell} (-1)^{\ell} (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{\ell} + 1)$$

$$+ \sum_{\ell=0}^{n-1} (1 - \varepsilon_{\ell+1}) (-1)^{\ell} (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{\ell}) + r_n(z_0),$$

with $\varepsilon_0 = 1$, and $|r_n(z_0)| \le Cq_n^{-1}|z_n|\log(1+|z_n|^{-1})$.

Proof. Since $\overline{\varphi}_1$ is real on the real axis outside [0, 1], by Proposition 6.10, when $z \in H$,

$$|\operatorname{Im} S^{n} \overline{\varphi}_{1}(z)| \lesssim |z| \log(1+|z|^{-1}) \sup_{D_{\infty}} |S^{n-1} \overline{\varphi}_{1}|. \tag{B.3}$$

We obtain $\sum_{n=1}^{\infty}\sup_{D_{\infty}}|S^n\overline{\varphi}_1|<\infty$ by using Proposition 6.3-(ii). Thus, from (B.3) and Lemma B.2, we get

$$\operatorname{Im} \mathcal{W}(z) = (1 - \varepsilon_1) \operatorname{Im} \overline{\varphi}_1(z) + \operatorname{Im} \overline{\varphi}_1(z+1) + r(z), \tag{B.4}$$

with $|r(z)| \lesssim |z| \log(1 + |z|^{-1})$.

Let us consider the case of $z_0 \in H$ which is corresponding to the case of n = 0. Since we set $\varepsilon_1 = 1$ for $z_0 \in \Delta \setminus D$, from (B.4), we obtain

$$|\text{Im}\mathcal{W}(z_0) - \text{Im}\overline{\varphi}_1(z_0 + 1)| \lesssim |z_0|\log(1 + |z_0|^{-1}).$$

Next, we consider $z_0 \in H(m_1, \dots, m_n)$ with $n \ge 1$ and $m_1, \dots, m_n \ge 1$. By Lemma B.2, we need to show that the modulus of

$$\left(\sum_{\ell=n+1}^{\infty} \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z_{0})\right) + (-1)^{n} (p_{n-1} - q_{n-1} \operatorname{Re}(z_{0})) (1 - \varepsilon_{n+1}) \operatorname{Im} \overline{\varphi}_{1}(z_{n}) + r^{0}(z_{0}) + r^{1}(z_{0})
+ \left[(1 - \varepsilon_{2}(1 - \varepsilon_{1})) \operatorname{Im} S \overline{\varphi}_{1}(z_{0}) + (p_{0} - q_{0} \operatorname{Re}(z_{0})) (\varepsilon_{1} \operatorname{Im} \overline{\varphi}_{1}(z_{1} + 1) + (1 - \varepsilon_{2}) \operatorname{Im} \overline{\varphi}_{1}(z_{1})) \right]
+ \sum_{\ell=2}^{n} \left[\operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z_{0}) - (-1)^{\ell} (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_{0})) (\varepsilon_{\ell} \operatorname{Im} \overline{\varphi}_{1}(z_{\ell} + 1) + (1 - \varepsilon_{\ell+1}) \operatorname{Im} \overline{\varphi}_{1}(z_{\ell})) \right]$$
(B.5)

is in $O(q_n^{-1}|z_n|\log(1+|z_n|^{-1}))$. From [MMY01, Prop. 4.11] using S=-T, we have $\sum_{i=0}^{\infty} ||S^i\overline{\varphi}_1|| < \infty$. Then, from Proposition B.3, we have

$$\left| \left(\sum_{\ell=n+1}^{\infty} \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z_{0}) \right) - \left(p_{n-1} - q_{n-1} \operatorname{Re}(z_{0}) \right) \sum_{\ell=1}^{\infty} \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z_{n}) \right|$$

$$\lesssim q_{n}^{-1} |\operatorname{Im}(z_{n})| \log(1 + |\operatorname{Im}(z_{n})|^{-1}).$$

Since $z_n \in H$, by using (B.3), the first term of (B.5) is bounded by

$$\sum_{\ell=n+1}^{\infty} \left| \operatorname{Im} S^{\ell} \overline{\varphi}_{1}(z_{0}) \right| \lesssim q_{n}^{-1} |z_{n}| \log(1 + |z_{n}|^{-1}).$$

For $z \in H$, we have

$$|\operatorname{Im}\overline{\varphi}_1(z)| \lesssim |\operatorname{Im}(z)|.$$
 (B.6)

For the second term of (B.5), by using (B.6) and (4.14), we have

$$|(p_{n-1} - q_n - 1\operatorname{Re}(z_0))\operatorname{Im}\overline{\varphi}_1(z_n)(1 - \varepsilon_{n+1})| \lesssim q_n^{-1}|\operatorname{Im}(z_n)|.$$

We have $|r^0(z_0) + r^1(z_0)| \lesssim |\operatorname{Im}(z_0)| \log(1 + |\operatorname{Im}(z_0)|^{-1}) \|\overline{\varphi}_1\|_{E([1/2,2])}$ in Lemma B.1 and B.2. Since

$$[1 - \varepsilon_2(1 - \varepsilon_1)](1 - \varepsilon_2) = 1 - \varepsilon_2$$
 and $[1 - \varepsilon_2(1 - \varepsilon_1)]\varepsilon_1 = \varepsilon_1$,

the third term of (B.5) is 0 or

$$\left[\operatorname{Im} S\overline{\varphi}_{1}(z_{0})+(p_{0}-q_{0}\operatorname{Re}(z_{0}))(\varepsilon_{1}\operatorname{Im} \overline{\varphi}_{1}(z_{1}+1)+(1-\varepsilon_{2})\operatorname{Im} \overline{\varphi}_{1}(z_{1}))\right].$$

By using Proposition B.4, the last terms of (B.5) are in

$$O\left(\sum_{\ell=0}^{n} q_{\ell}^{-1} |\operatorname{Im}(z_{\ell})| \log(1 + |\operatorname{Im}(z_{\ell})|^{-1}) \|\overline{\varphi}_{1}\|_{E([1/2,2])}\right)$$

and we have

$$\sum_{\ell=0}^{n} q_{\ell}^{-1} |\operatorname{Im}(z_{\ell})| \log(1 + |\operatorname{Im}(z_{\ell})|^{-1}) \le q_{n}^{-1} |\operatorname{Im}(z_{n})| \log(1 + |\operatorname{Im}(z_{n})|^{-1})$$

as in the last part of the proof of [MMY01, Prop. 5.17]. Since $|\text{Re}(z_0)| \leq 1$, we have the conclusion.

Proof of Theorem 7.5. Let $n \geq 0$ and $m_1, \dots, m_n \geq 1$. The domain $H(m_1, \dots, m_n)$ meets \mathbb{R} in a unique point p_n/q_n . Let us denote $x_0 = p_n/q_n$ and consider its continued fraction

$$x_i^{-1} = m_{i+1} + x_{i+1}, \quad 0 \le i < n \quad \text{and} \quad x_n = 0,$$

here, x_i is the point of intersection of $H(m_{i+1}, \dots, m_n)$ with \mathbb{R} .

Let $z_0 \in H(m_1, \dots, m_n)$ and $\operatorname{Im}(z_0) > 0$. We will then have $(-1)^{\ell} \operatorname{Im}(z_{\ell}) > 0$ for $0 \le \ell \le n$. From (7.8), we deduce that

$$\operatorname{Im}\overline{\varphi}_{1}(x_{\ell}+1+(-1)^{\ell}i0) = (-1)^{\ell}\log\frac{1}{x_{\ell}}, \quad \text{for } 0 \leq \ell < n.$$
 (B.7)

Since $\frac{x_{\ell}}{1-x_{\ell}} = \frac{1}{x_{\ell+1}}$ if $\varepsilon_{\ell+1} = 0$, we obtain from (7.8)

$$(1 - \varepsilon_{\ell+1}) \operatorname{Im}_{\overline{\varphi}_1}(x_{\ell} + (-1)^{\ell} i 0) = (1 - \varepsilon_{\ell+1})(-1)^{\ell+1} x_{\ell} \log \frac{1}{x_{\ell+1}}, \quad \text{for } 0 \le \ell < n-1. \text{ (B.8)}$$

The above equations imply for W_{finite} extended to the complex plane that

$$W_{\text{finite}}(p_n/q_n) = \sum_{\ell=0}^{n-1} \varepsilon_{\ell}(q_{\ell-1}x_0 - p_{\ell-1})(-1)^{\ell} \text{Im}\overline{\varphi}_1(x_{\ell} + 1 + (-1)^{\ell}i0)$$

$$+ \sum_{\ell=0}^{n-2} (1 - \varepsilon_{\ell+1})(q_{\ell-1}x_0 - p_{\ell-1})(-1)^{\ell} \text{Im}\overline{\varphi}_1(x_{\ell} + (-1)^{\ell}i0).$$
(B.9)

Combining Proposition B.5 with (B.9), we get

$$\operatorname{Im} \mathcal{W}(z_0) + W_{\text{finite}}(p_n/q_n) - (-1)^n (p_{n-1} - q_{n-1}\operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_n + 1) - r_n(z_0)$$

$$= \sum_{\ell=0}^{n-1} \varepsilon_{\ell}(-1)^{\ell} \Big[(p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{\ell} + 1) - (p_{\ell-1} - q_{\ell-1} x_0) \operatorname{Im} \overline{\varphi}_1(x_{\ell} + 1 + (-1)^{\ell} i 0) \Big]$$

$$+ \sum_{\ell=0}^{n-2} (1 - \varepsilon_{\ell+1})(-1)^{\ell} \Big[(p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{\ell}) - (p_{\ell-1} - q_{\ell-1} x_0) \operatorname{Im} \overline{\varphi}_1(x_{\ell} + (-1)^{\ell} i 0) \Big]$$

$$-(1-\varepsilon_n)(-1)^n \Big[(p_{n-1} - q_{n-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_n + 1) + (p_{n-2} - q_{n-2} \operatorname{Re}(z_0)) \operatorname{Im} (\overline{\varphi}_1(z_{n-1})) \Big].$$
(B.10)

Thus, we have to deal with the following expressions. For $0 \le \ell \le n-1$ such that $\varepsilon_{\ell} = 1$,

$$A_{\ell} := (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{\ell} + 1) - (p_{\ell-1} - q_{\ell-1} x_0) \operatorname{Im} \overline{\varphi}_1(x_{\ell} + 1 + (-1)^{\ell} i 0).$$

For $0 \le \ell \le n-2$ such that $\varepsilon_{\ell+1} = 0$,

$$B_{\ell} := (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \operatorname{Im}_{\overline{\varphi}_1}(z_{\ell}) - (p_{\ell-1} - q_{\ell-1}x_0) \operatorname{Im}_{\overline{\varphi}_1}(x_{\ell} + (-1)^{\ell} i 0).$$

When $\varepsilon_n = 0$,

$$C_n := (p_{n-1} - q_{n-1} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_n + 1) + (p_{n-2} - q_{n-2} \operatorname{Re}(z_0)) \operatorname{Im} \overline{\varphi}_1(z_{n-1}).$$

The first n digits of the continued fraction of $Re(z_0)$ are also m_1, \dots, m_n . This gives

$$|\operatorname{Re}(z_0) - x_0| \le |z_0 - p_n/q_n| = q_n^{-1}|q_{n-1}z_0 - p_{n-1}||z_n| \le q_n^{-2}|z_n|.$$

Since (7.6) implies that

$$\overline{\varphi}'_1(z) = \frac{1}{\pi} \left(-\text{Li}_2\left(\frac{z}{1-z}\right) + \frac{1}{1-z} \log\left(\frac{1-2z}{1-z}\right) + \frac{1}{1-z} \log\left(\frac{z-2}{z-1}\right) \right) - \frac{\pi}{12},$$

we have

$$|\overline{\varphi}'_1(z)| \le C|z-1|^{-1}\log\frac{1}{|z-1|}$$
 near 1. (B.11)

For $0 \le \ell \le n$, the distance of x_{ℓ} and z_{ℓ} from 0 are comparable. Thus, for $0 \le \ell \le n$,

$$|\operatorname{Im}\overline{\varphi}_{1}(z_{\ell}+1) - \operatorname{Im}\overline{\varphi}_{1}(x_{\ell}+1+(-1)^{\ell}i0)| \leq Cx_{\ell}^{-1}\log\frac{1}{x_{\ell}}|z_{\ell}-x_{\ell}|,$$

and, for $0 \le \ell \le n - 1$,

$$|\operatorname{Im}\overline{\varphi}_1(z_{\ell}) - \operatorname{Im}\overline{\varphi}_1(x_{\ell} + (-1)^{\ell}i0)| \le C|x_{\ell} - 1|^{-1}\log\frac{1}{1 - x_{\ell}}|z_{\ell} - x_{\ell}|.$$

We have $|z_{\ell} - x_{\ell}| \lesssim |z_n| q_{\ell}^2 q_n^{-2}, x_{\ell}^{-1} \lesssim m_{\ell+1}$ and $|x_{\ell} - 1|^{-1} \lesssim x_{\ell+1}^{-1} \lesssim m_{\ell+2}$. Thus, according to (B.7) and (B.8), for $0 \leq \ell < n$, we have

$$|A_{\ell}| = \left| (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \left(\operatorname{Im} \overline{\varphi}_1(z_{\ell} + 1) - \operatorname{Im} \overline{\varphi}_1(x_{\ell} + 1 + (-1)^{\ell} i 0) \right) \right.$$

$$\left. + q_{\ell-1} (\operatorname{Re}(z_0) - x_0) \operatorname{Im} \overline{\varphi}_1(x_{\ell} + 1 + (-1)^{\ell} i 0) \right|$$

$$\lesssim x_{\ell}^{-1} \log \frac{1}{x_{\ell}} |z_{\ell} - x_{\ell}| q_{\ell}^{-1} + q_{\ell-1} q_n^{-2} |z_n| \log \frac{1}{x_{\ell}}$$

$$\lesssim |z_n| q_n^{-2} (q_{\ell} m_{\ell+1} + q_{\ell-1}) \log (m_{\ell+1}) \lesssim |z_n| q_n^{-2} q_{\ell+1} \log (m_{\ell+1}),$$

and, for $0 \le \ell < n-1$ such that $\varepsilon_{\ell+1} = 0$,

$$|B_{\ell}| = \left| (p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)) \left(\operatorname{Im} \overline{\varphi}_1(z_{\ell}) - \operatorname{Im} \overline{\varphi}_1(x_{\ell} + (-1)^{\ell} i 0) \right) \right.$$

$$\left. + q_{\ell-1} (\operatorname{Re}(z_0) - x_0) \operatorname{Im} \overline{\varphi}_1(x_{\ell} + (-1)^{\ell} i 0) \right|$$

$$\lesssim |p_{\ell-1} - q_{\ell-1} \operatorname{Re}(z_0)||x_{\ell} - 1|^{-1} \log \frac{1}{|x_{\ell} - 1|} |z_{\ell} - x_{\ell}| + q_{\ell-1} q_n^{-2} |z_n| \log \frac{1}{x_{\ell+1}}$$

$$\lesssim |x_{\ell} - 1|^{-1} \log \frac{1}{|x_{\ell} - 1|} |z_{\ell} - x_{\ell}| q_{\ell}^{-1} + q_{\ell-1} q_n^{-2} |z_n| \log \frac{1}{x_{\ell+1}}$$

$$\lesssim |z_n| q_n^{-2} (q_{\ell} m_{\ell+2} + q_{\ell-1}) \log(m_{\ell+2}) \lesssim |z_n| q_n^{-2} q_{\ell+2} \log(m_{\ell+2}).$$

From $q_n \geq 2^{\lfloor (n-\ell)/2 \rfloor} q_{\ell}$, it follows that

$$\sum_{\ell=0}^{n-1} |A_{\ell}| + \sum_{\ell=0}^{n-2} |B_{\ell}| \lesssim |z_n| \frac{\log q_n}{q_n}.$$

Furthermore, when $\varepsilon_n = 0$, we have

$$\overline{\varphi}_1(z_{n-1}) = z_{n-1} \left(\varphi_0(z_n) - \varphi_0(-1) \right) - \varphi_0'(-1) + \varphi_0(z_{n-1} - 1).$$

$$\overline{\varphi}_1(1+z_n) = z_{n-1}^{-1} \left(\varphi_0(z_{n-1} - 1) - \varphi_0(-1) \right) - \varphi_0'(-1) + \varphi_0(z_n).$$

Since

$$\overline{\varphi}_1(z_{n-1}) = (1 - z_{n-1}) \left(\frac{\pi}{12} - \frac{1}{\pi} \log 2 \right) + z_{n-1} \overline{\varphi}_1(1 + z_n),$$

we have

$$C_n = \operatorname{Im}\overline{\varphi}_1(z_n + 1)[(p_{n-1} - q_{n-1}\operatorname{Re}(z_0)) + (p_{n-2} - q_{n-2}\operatorname{Re}(z_0))\operatorname{Re}(z_{n-1})] + (p_{n-2} - q_{n-2}\operatorname{Re}(z_0))\operatorname{Im}(z_{n-1}) \left[-\left(\frac{\pi}{12} - \frac{1}{\pi}\log 2\right) + \operatorname{Re}\overline{\varphi}_1(1 + z_n)\right) \right].$$

By (7.16), we have $|\overline{\varphi}_1(1+z_n)| \lesssim \log^2(1+|z_n|^{-1})$. As in the proof of [MMY01, Theorem 5.10],

$$|(p_{n-2} - q_{n-2}\operatorname{Re}(z_0))\operatorname{Im}(z_{n-1})| \lesssim q_n^{-1}|\operatorname{Im}(z_n)|,$$

and

$$[(p_{n-1} - q_{n-1}\operatorname{Re}(z_0)) + (p_{n-2} - q_{n-2}\operatorname{Re}(z_0))\operatorname{Re}(z_{n-1})] \lesssim q_n^{-1}|\operatorname{Im}(z_n)|.$$

Thus, we have $|C_n| \lesssim q_n^{-1} |\text{Im}(z_n)| \log^2(1 + |z_n|^{-1})$.

References

- [BBLS05] L. Báez-Duarte, M. Balazard, B. Landreau, and E. Saias. Étude de l'autocorrélation multiplicative de la fonction 'partie fractionnaire'. *Ramanujan J.*, 9(1-2):215–240, 2005.
- [BC06] X. Buff and A. Chéritat. The Brjuno function continuously estimates the size of quadratic siegel disks. *Ann. Math.*, 164(1):265–312, 2006.
- [BG01] A. Berretti and G. Gentile. Bryuno function and the standard map. Comm. Math. Phys., 220(3):623–656, 2001.
- [BM12] M. Balazard and B. Martin. Comportement local moyen de la fonction de Brjuno. Fund. Math., 218(3):193–224, 2012.
- [BM13] M. Balazard and B. Martin. Sur l'autocorrélation multiplicative de la fonction "partie fractionnaire" et une fonction définie par J. R. Wilton. Preprint, arXiv:1305.4395, 2013.
- [BM19] M. Balazard and B. Martin. Sur certaines équations fonctionnelles approchées, liées à la transformation de Gauss. *Aequationes Math.*, 93(3):563–585, 2019.
- [Brj71] A.D. Brjuno. Analytic form of differential equations. I (Russian). Trudy Moskov. Mat. Obšč., 25:119–262, 1971.
- [Brj72] A.D. Brjuno. Analytic form of differential equations. II (Russian). Trudy Moskov. Mat. Obšč., 26:199–239, 1972.
- [CC15] D. Cheraghi and A. Chéritat. A proof of the Marmi–Moussa–Yoccoz conjecture for rotation numbers of high type. *Invent. Math.*, 202(2):677–742, 2015.
- [Dav94] A.M. Davie. The critical function for the semistandard map. Nonlinearity, 7(1):219, 1994.
- [Gar07] J. Garnett. Bounded analytic functions, volume 236. Springer Science & Business Media, 2007.
- [GD11] J. García-Cuerva and J.R. De Francia. Weighted norm inequalities and related topics. Elsevier, 2011.
- [Gen15] G. Gentile. Invariant curves for exact symplectic twist maps of the cylinder with Bryuno rotation numbers. *Nonlinearity*, 28(7):2555, 2015.
- [Hen06] D. Hensley. The Hurwitz complex continued fraction. preprint, January, 9, 2006.
- [Hur87] A. Hurwitz. Über die Entwicklung complexer Grössen in Kettenbrüche. Acta Math., 11(1-4):187–200
- [JM18] S. Jaffard and B. Martin. Multifractal analysis of the Brjuno function. *Invent. Math.*, 212(1):109–132, 2018.
- [Kno90] M. Knopp. Modular integrals and their Mellin transforms. In Analytic number theory: Proceedings of a Conference in Honor of Paul T. Bateman, volume 85 of Progr. Math., pages 327–342. Birkhäuser, Boston, 1990.
- [Mac88] R.S. MacKay. Exact results for an approximate renormalisation scheme and some predictions for the breakup of invariant tori. *Phys. D*, 33(1-3):240–265, 1988. Progress in chaotic dynamics.
- [Mac89] R.S. MacKay. Erratum: "Exact results for an approximate renormalisation scheme and some predictions for the breakup of invariant tori". *Phys. D*, 36(3):358, 1989.
- [Mar90] S. Marmi. Critical functions for complex analytic maps. J. Phys. A: Math. Gen., 23(15):3447–3474, 1990.

[MMY95] S. Marmi, P. Moussa, and J.-C. Yoccoz. Continued fractions transformations, Brjuno functions and BMO spaces. *preprint*, *doi=10.1.1.32.4860*, previous version of MMY95 containing some extended calculations, 1995.

[MMY97] S. Marmi, P. Moussa, and J.-C. Yoccoz. The Brjuno functions and their regularity properties. Commun. Math. Phys., 186(2):265–293, 1997.

[MMY01] S. Marmi, P. Moussa, and J.-C. Yoccoz. Complex Brjuno functions. J. Amer. Math. Soc., 14(4):783–841 (electronic), 2001.

[MMY06] S. Marmi, P. Moussa, and J.-C. Yoccoz. Some properties of real and complex Brjuno functions. In *Frontiers in number theory, physics, and geometry. I*, pages 601–623. Springer, Berlin, 2006.

[MS92] S. Marmi and J. Stark. On the standard map critical function. *Nonlinearity*, 5(3):743, 1992.

[Nak81] H. Nakada. Metrical theory for a class of continued fraction transformations and their natural extensions. *Tokyo J. Math.*, 4(2):399–426

[Pet14] I. Petrykiewicz. Note on the differentiability of Fourier series arising from Eisenstein series. C. R. Math. Acad. Sci. Paris, 352(4):273–276, 2014.

[Pet17] I. Petrykiewicz. Differentiability of Fourier series arising from Eisenstein series. Ramanujan J., 42(3):527–581, 2017.

[Sar78] D. Sarason. Function theory on the unit circle. Virginia Polytechnic Institute and State University, Department of Mathematics, Blacksburg, Va., 1978. Notes for lectures given at a Conference at Virginia Polytechnic Institute and State University, Blacksburg, Va., June 19–23, 1978.

[Sie42] C.L. Siegel. Iteration of analytic functions. Ann. Math., 43(4):607–612, 1942.

[TI81] S. Tanaka and S. Ito. On a family of continued-fraction transformations and their ergodic properties. *Tokyo J. Math.*, 4(1):153–175, 1981.

[Wil33] J.R. Wilton. An approximate functional equation with applications to a problem of Diophantine approximation. J. Reine Angew. Math., 169:219–237, 1933.

[Yoc88] J.-C. Yoccoz. Linéarisation des germes de difféomorphismes holomorphes de (C,0). C. R. Acad. Sci. Paris Sér. I Math., 306(1):55–58, 1988.

[Yoc95] J.-C. Yoccoz. Petits diviseurs en dimension 1. Astérisque, 231(3), 1995.

[Yoc02] J.-C. Yoccoz. Analytic linearization of circle diffeomorphisms. In *Dynamical systems and small divisors*, pages 125–173. Springer, 2002.

[Zag92] D. Zagier. Introduction to modular forms. In From number theory to physics (Les Houches, 1989), pages 238–291. Springer, Berlin, 1992.

Centro di Ricerca Matematica Ennio De Giorgi, Scuola Normale Superiore, Piazza dei Cavalieri 3, 56126 Pisa, Italy

Email address: seulbee.lee@sns.it

Scuola Normale Superiore, Palazzo della Carovana, Piazza dei Cavalieri 7, 56126 Pisa, Italy

Email address: stefano.marmi@sns.it

Email address: ipetrykiewicz@gmail.com

Centro di Ricerca Matematica Ennio De Giorgi, Scuola Normale Superiore, Piazza dei Cavalieri 3, 56126 Pisa, Italy

Email address: tanja.schindler@sns.it