

BROWNIAN REGULARITY FOR THE KPZ LINE ENSEMBLE

XUAN WU

ABSTRACT. This paper seeks a quantitative comparison between the curves in the KPZ line ensemble [CH16] and a standard Brownian bridge under the $t^{1/3}$ vertical and $t^{2/3}$ horizontal scaling. The estimate we obtained is parallel to the one established in [Ham1], where the Airy line ensemble was studied. Our main tool is the soft Brownian Gibbs property enjoyed by the KPZ line ensemble. In view of the Gibbs property, the KPZ line ensemble differs from the Airy line ensemble mainly due to the intersecting nature of its curves, which results in the main technical difficulty in this paper. We develop a resampling framework, the soft jump ensemble, to tackle this difficulty. Our method is highly inspired by the jump ensemble technique developed in [Ham1].

1. INTRODUCTION

The Kardar-Parisi-Zhang (KPZ) line ensemble, $\{\mathcal{H}_n^t\}$, is a countable collection of continuous random curves. It may be viewed as a multi-layer extension to the well-known KPZ equation, associated with a universality class which bears the same name. The KPZ line ensemble has attracted extensive attention during the past ten years due to its rich algebraic and probabilistic structures, which also served as powerful tools to study the KPZ equation.

The KPZ line ensemble $\{\mathcal{H}_n^t\}$ is a Gibbsian line ensemble, a special class of Gibbs measures which have received considerable attention in the past two decades owing, in part, to their occurrence in integrable probability. A Gibbsian line ensemble can be thought of as a countable collection of independent random curves reweighed by a local interaction energy. Local means that the energies only depend on the values of nearby curves. A simple example of a Gibbsian line ensemble is a collection of Brownian motions conditioned never to collide, known as the Dyson Brownian motion with $\beta = 2$. Another well-known example is the Airy line ensemble [CH14], a central object in the KPZ universality class.

There is a large class of stochastic integrable models from random matrix theory, interacting particle systems, last passage percolation, directed polymers that naturally carry the structure of random paths with some Gibbsian resampling invariance. The structure of a Gibbsian line ensemble can be utilized to great benefit when studying their asymptotic scaling limits or path regularity. Starting with [CH14], there has been a fruitful development of techniques which leverage the Gibbs property of Gibbsian line ensembles to prove their tightness under various scalings given only one-point tightness information about their top curve – see for instance [CH16, CD18, Wu19, BCD21]. Initiated in [Ham1], the Gibbs property was further employed to study the path regularity for Gibbsian line ensemble. Furthermore, the understanding of the path regularity for the Brownian last passage percolation (BLPP) then helped study the geometry of the geodesics for (BLPP) and the space-time Airy sheet, see for instance [Ham2, Ham3, Ham4, BGH].

The Gibbs property enjoyed by the KPZ line ensemble ensures that curves in the KPZ line ensemble locally behave like Brownian motions. This paper is devoted to provide a quantitative comparison between the curves in the KPZ line ensemble and a standard Brownian bridge through utilizing such a Gibbs property. We aim to address the following question.

Question: Suppose an arbitrary event A is of small probability $\varepsilon > 0$ under the law of a standard Brownian bridge. How comparable is $\mathbb{P}[A]$ with respect to ε under the law of the KPZ line ensemble?

Highly inspired by the jump ensemble method, introduced in [Ham1], we develop a resampling framework, the soft jump ensemble, to study the Brownian regularity for the KPZ line ensemble and to answer the above question.

1.1. The Kardar-Parisi-Zhang equation. The KPZ equation was introduced in 1986 by Kardar, Parisi and Zhang [KPZ] as a model for random interface growth. In one-spatial dimension (sometimes also called (1+1)-dimension to emphasize that time is one dimension too), it describes the evolution of a function $\mathcal{H}(t, x)$ recording the height of an interface at time $t > 0$ above position $x \in \mathbb{R}$. The KPZ equation is written formally as a stochastic partial differential equation (SPDE),

$$(1.1) \quad \partial_t \mathcal{H} = \frac{1}{2} \partial_x^2 \mathcal{H} + \frac{1}{2} (\partial_x \mathcal{H})^2 + \mathcal{W},$$

where \mathcal{W} is a space-time white noise (for mathematical background or literature review, see [Cor, QS15] for instance).

The KPZ equation is a canonical member of the associated KPZ universality class and a model belongs to the KPZ universality class if it bears the same long-time, large-scale behavior as the KPZ equation. All models in the KPZ universality class can be transformed to a kinetically growing interface reflecting the competition between growth in a direction normal to the surface, a surface tension smoothing force, and a stochastic term which tends to roughen the interface. These features may be illustrated by the Laplacian $\frac{1}{2} \partial_x^2 \mathcal{H}$, non-linear term $\frac{1}{2} (\partial_x \mathcal{H})^2$ and white noise \mathcal{W} in the KPZ equation (1.1). Numerical simulations along with some theoretical results have confirmed that in the long time t scaling limit, fluctuations in the height of such evolving interfaces scale like $t^{1/3}$ and display non-trivial spatial correlations in the scale $t^{2/3}$ (known as the 3 : 2 : 1 KPZ scaling).

The KPZ equation is related to the stochastic heat equation (SHE) with multiplicative noise through the Hopf–Cole transformation. Denote $\mathcal{Z}(t, x)$ as the solution to the following SHE,

$$(1.2) \quad \partial_t \mathcal{Z} = \frac{1}{2} \partial_x^2 \mathcal{Z} + \mathcal{Z} \mathcal{W}.$$

The Hopf–Cole solution to the KPZ equation (1.1) is defined through taking

$$\mathcal{H}(t, x) = \log \mathcal{Z}(t, x).$$

It was first proved in [Mue] that $\mathcal{Z}(t, x)$ is almost surely strictly positive, which justifies the validity of the transform. The fundamental solution $\mathcal{Z}^{nw}(t, x)$ to SHE (1.2) is of great importance. It solves (1.2) with a delta mass initial value problem at origin, i.e. $\mathcal{Z}(0, x) = \delta_{x=0}$. Meanwhile $\mathcal{H}^{nw}(t, x) = \log \mathcal{Z}^{nw}(t, x)$ is known as the narrow wedge solution to the KPZ equation. The initial condition of $\mathcal{H}^{nw}(0, x)$ is not well-defined; however $\mathcal{H}^{nw}(t, x)$ is stationary around a parabola $\frac{-x^2}{2t}$, which resembles a sharp wedge for small t , hence known as the narrow wedge initial condition.

Using the Feynman-Kac representation, $\mathcal{Z}^{nw}(t, x)$ formally take the following expression,

$$(1.3) \quad \mathcal{Z}^{nw}(t, x) = p(t, x) \mathbb{E} \left[: \exp : \left(\int_0^t \mathcal{W}(s, B_s) ds \right) \right],$$

where $p(t, x) = (2\pi t)^{-1/2} \exp(-x^2/2t)$ is the heat kernel, the expectation is taken with respect to a Brownian bridge $(B_s, s \leq t)$ which starts at origin at time 0 and arrives at x at time t . The $: \exp :$ is the *Wick exponential*, see [Jan] for instance. This bridge representation arises because of the $\delta_{x=0}$ initial condition, and hence the factor $p(t, x)$. This Feynman-Kac representation is mostly formal since the integral of white noise over a Brownian path is not well-defined pathwise or to exponentiate the integral.

We adopt this representation to emphasize on its interpretation as being the partition function of a continuum directed random polymer (CDRP) that is modeled by a continuous path interacting

with a space-time white noise. We will make sense of the expression (1.3) through a chaos expansion in Section 1.2.

1.2. The KPZ line ensemble. We emphasize that the above description is very useful for a generalization to the multi-layer scenario. The KPZ line ensemble is defined through taking the logarithm of ratios of partition functions of the continuum directed random polymers (CDRP). The partition functions of the CDRP $\mathcal{Z}_n(t, x)$ are formally written as

$$(1.4) \quad \mathcal{Z}_n(t, x) = p(t, x)^n \mathbb{E} \left[: \exp : \left\{ \sum_{i=1}^n \int_0^t \mathcal{W}(s, B_i(s)) ds \right\} \right],$$

where the expectation is taken with respect to the law on n independent Brownian bridges $\{B_i\}_{i=1}^n$ starting at 0 at time 0 and ending at x at time t . Intuitively these path integrals represent energies of non-intersecting paths, and thus the expectation of their exponential represents the partition functions for this polymer model. It is worth noting that the first layer, \mathcal{Z}_1 , is the same as the fundamental solution to the stochastic heat equation (SHE). These partition functions $\mathcal{Z}_n(t, x)$ also solve a multi-layer extension of SHE, see [OW].

For $n \in \mathbb{N}$, $t \geq 0$ and $x \in \mathbb{R}$, $\mathcal{Z}_n(t, x)$ is rigorously defined via the following chaos expansion,

$$(1.5) \quad \mathcal{Z}_n(t, x) = p(t, x)^n \sum_{k=0}^{\infty} \int_{\Delta_k(t)} \int_{\mathbb{R}^k} R_k^{(n)}((t_1, x_1), \dots, (t_k, x_k)) \mathcal{W}(dt_1 dx_1) \cdots \mathcal{W}(dt_k dx_k),$$

where $\Delta_k(t) = \{0 < t_1 < \dots < t_k < t\}$, and $R_k^{(n)}$ is the k -point correlation function for a collection of n non-intersecting Brownian bridges each of which starts at 0 at time 0 and ends at x at time t . For notational simplicity set $\mathcal{Z}_0(t, x) \equiv 1$. For details about integration with respect to a white noise, we refer to [Jan].

[LW] show that for any $t > 0$, with probability 1, for all $x \in \mathbb{R}$ and all $n \in \mathbb{N}$, $\mathcal{Z}_n(t, x) > 0$. The positivity result permits the following important definition of the KPZ line ensemble $\mathcal{H}_n^t(x)$, a process given by the logarithm of ratios of partition functions \mathcal{Z}_n .

Definition 1.1. For $t > 0$ fixed, the KPZ _{t} line ensemble is a continuous \mathbb{N} -indexed line ensemble $\mathcal{H}^t = \{\mathcal{H}_n^t\}_{n \in \mathbb{N}}$ on \mathbb{R} given by

$$(1.6) \quad \mathcal{H}_n^t(x) = \log \left(\frac{\mathcal{Z}_n(t, x)}{\mathcal{Z}_{n-1}(t, x)} \right).$$

Sometimes we will omit t and just write the KPZ line ensemble as \mathcal{H} .

This paper continues the investigation of the KPZ line ensemble under the KPZ scaling in [Wu21]. The scaled KPZ line ensemble \mathfrak{H}^t is defined as follows.

Definition 1.2. For $t > 0$ fixed, the scaled KPZ _{t} line ensemble is a continuous \mathbb{N} -indexed line ensemble $\mathfrak{H}^t = \{\mathfrak{H}_n^t\}_{n \in \mathbb{N}}$ on \mathbb{R} given by

$$(1.7) \quad \mathfrak{H}_n^t(x) := \frac{\mathcal{H}_n^t(t^{2/3}x) + \frac{t}{24}}{t^{1/3}}.$$

The result we prove is uniform in $t \geq 1$ and sometimes we will omit t and just write \mathfrak{H} .

1.3. Main result. The KPZ line ensemble is known to have a beautiful Gibbs property [CH16], which provides a precise Radon-Nikodym description of the paths in \mathfrak{H} with respect to Brownian bridges. The Gibbs property immediately implies the absolute continuity of the paths in \mathfrak{H} with respect to Brownian bridge measure [CH16]. [Wu21] studied the KPZ line ensemble and showed that the curves in the KPZ line ensemble fluctuate on the same scale compared to the local fluctuation of Brownian bridges. In this paper we aim to investigate how general one may push such a comparison between the curves of the KPZ line ensemble and Brownian bridges.

To be precise, let $s > 0$ and $C_{0,0}([0, s])$ be the space of continuous functions on $[0, s]$ which vanish both at 0 and s . Let B be a standard Brownian bridge on $[0, s]$ and denote its law by \mathbb{P}_{free} . Denote $\mathfrak{H}_k^{t,[0,s]}$ as the bridge part of \mathfrak{H}_k on $[0, s]$, i.e.

$$\mathfrak{H}_k^{t,[0,s]}(x) := \mathfrak{H}_k^t(x) - s^{-1}x\mathfrak{H}_k^t(s) - s^{-1}(s-x)\mathfrak{H}_k^t(0).$$

We will write $\mathfrak{H}_k^{[0,s]}$ for notational simplicity sometimes.

Theorem 1.3. *For any $k \in \mathbb{N}$ and $s \geq 1$, there exists $D := D(k, s)$ such that the following statement holds uniformly in $t \geq 1$. Given a Borel set $A \subset C_{0,0}([0, s])$, let $\varepsilon = \mathbb{P}_{\text{free}}(B \in A)$. For $\varepsilon \in (0, e^{-s^3}]$, we have*

$$\mathbb{P}(\mathfrak{H}_k^{t,[0,s]} \in A) \leq D\varepsilon \cdot e^{D(\log \varepsilon^{-1})^{5/6}}.$$

Remark 1.4. *For notational simplicity, we state and prove the above result for interval $[0, s]$. This can be easily adapted to general intervals as \mathfrak{H} is stationary under horizontal shifts after subtracting the parabola $\frac{x^2}{2}$. The dependence of $D(k, s)$ on k and s can be traced through our arguments, which we do not pursue here.*

Theorem 1.3 gives an informative comparison between the curves of the scaled KPZ line ensemble and a standard Brownian bridge. This is parallel to the comparison established in [Ham1] between curves in the Airy line ensemble, BLPP line ensemble and a standard Brownian bridge. Such a comparison serves as an important ingredient in investigating geodesic coalescence and geodesic energy profiles for Brownian last passage percolation, see for instance [Ham2, Ham3, Ham4, BGH]. Another quantitative approximation of the Airy line ensemble by connecting independent Brownian bridges was studied in [DV] and served as a major input in the construction of the direct landscape (space-time Airy sheet) [DOV]. We hope Theorem 1.3 may similarly shed light on the further study of the KPZ line ensemble and its related directed random polymer models.

It is worth mentioning that [CHH] further removes the affine shift and strengthen the comparison to a comparison between the curves of Airy line ensemble or BLPP line ensemble and a **Brownian motion**. The energy landscape of last passage percolation models (together with another 5 applications) were also studied in [CHH] based on their comparison results. It is interesting to investigate whether Brownian motion comparison holds true for the scaled KPZ line ensemble and explore further applications.

1.4. Gibbs property, local fluctuation estimates and the high jump difficulty. The KPZ line ensemble has been studied in [Wu21] through its Gibbs property. [Wu21] obtains local fluctuation estimates, of which a key application is the convergence of the KPZ line ensemble to the Airy line ensemble, together with the work in [QS20, Vir]. In this section, we briefly introduce the Gibbs property and explain framework of the argument in [Wu21]. We emphasize on the limitation of the resampling techniques from [Wu21] towards the Brownian comparison pursued in this paper.

The Gibbs property for the scaled KPZ line ensemble \mathfrak{H} is a spatial Markov property which further specifies the law of \mathfrak{H} in a domain, given the boundary information. We illustrate with the $k = 1$ case. More specifically, given the values of $\mathfrak{H}_1(\ell), \mathfrak{H}_1(r)$ and \mathfrak{H}_2 on $[\ell, r]$, called the boundary data, we may recover the law of \mathfrak{H}_1 on $[\ell, r]$. The law of \mathfrak{H}_1 on $[\ell, r]$ satisfies the following Radon-Nikodym derivative relation with respect to a Brownian bridge connecting $(\ell, \mathfrak{H}_1(\ell))$ and $(r, \mathfrak{H}_1(r))$.

$$(1.8) \quad \frac{d\mathbb{P}_{\mathfrak{H}_1}}{d\mathbb{P}_{\text{free}}}(B) = \frac{W(B)}{Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2}}.$$

W is known as the Boltzmann weight and it penalizes the curves for going out of order. Moreover, W is always less than 1, while the exact expression of W does not matter here for our purpose.

See (2.1) for the explicit form of W . The normalizing constant $Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2}$ is necessary to ensure a probability measure and it takes the form of

$$Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2} := \mathbb{E}_{\text{free}}[W(B)].$$

See Section 2.3 for a detailed description.

Let us emphasize that the the normalizing constant $Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2}$ inherits randomness from the boundary data $\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2$. This causes one of the main difficulties towards understanding the marginal law of \mathfrak{H}_1 on $[0, s]$. Intuitively, a quantitative control of $Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2}$ will be very helpful for a quantitative comparison between the law of \mathfrak{H}_1 with respect to a Brownian bridge.

We now illustrate the above idea in detail. Take $[\ell, r] = [-2, 2]$. Recall that $A \in C_{0,0}([0, s])$ is an arbitrary Borel subset with probability ε under a standard Brownian bridge. We abbreviate $Z^{\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2}$ as Z_{bdd} to emphasize that $\mathfrak{H}_1(\ell), \mathfrak{H}_1(r), \mathfrak{H}_2$ serve as the boundary data. Using (1.8), we have

$$\mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A) = \mathbb{E}_{\mathfrak{H}}[\mathbb{1}\{\mathfrak{H} \in A\}] = \mathbb{E}_{\text{bdd}} \left[\frac{\mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W]}{Z_{\text{bdd}}} \right].$$

Here \mathbb{E}_{bdd} means taking expectation over all possible boundary data. We may simply bound the numerator using $W \leq 1$,

$$\mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W] \leq \mathbb{P}_{\text{free}}(B \in A) =: \varepsilon.$$

As long as Z_{bdd} has a lower bound, we may find some upper bound for $\mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A)$.

The following quantitative tail bound on Z_{bdd} was established in [Wu21].

$$\mathbb{P}_{\mathfrak{H}} \left(Z_{\text{bdd}} < e^{-O(1)K^2} \right) < e^{-K^{3/2}}.$$

See Proposition 4.8 for a precise statement. Therefore,

$$\mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A) \leq \varepsilon e^{O(1)K^2} + e^{-K^{3/2}}$$

Choosing $K = C^{-1} (\log \varepsilon^{-1})^{1/2}$ with a large constant C , we obtain

$$(1.9) \quad \mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A) \leq e^{-O(1)(\log \varepsilon^{-1})^{3/4}}.$$

This bound is far from satisfactory. In particular, $e^{-O(1)(\log \varepsilon^{-1})^{3/4}} \gg \varepsilon^\alpha$ for all $\alpha > 0$.

An possible remedy which naturally occurs is to optimize our choice of the resampling domain $[\ell, r]$. By allowing $T = r - \ell$ to vary, we obtained the following estimate about Z_{bdd} ,

$$(1.10) \quad \mathbb{P}_{\mathfrak{H}} \left(Z_{\text{bdd}} < e^{-O(1)T^3} \right) < e^{-T^3}.$$

See Proposition 4.8 for a precise statement. Therefore,

$$\mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A) \leq \varepsilon e^{O(1)T^3} + e^{-T^3}$$

By setting $e^{O(1)T^3} = \varepsilon^{-1/2}$, we may deduce that for some large $C \in \mathbb{R}$,

$$(1.11) \quad \mathbb{P}_{\mathfrak{H}}(\mathfrak{H} \in A) \leq \varepsilon^{1/C}.$$

This estimate is an improvement of the one in (1.9) but still not satisfactory enough. In this paper, we seek for a bound of the order $\varepsilon^{1-\alpha(1)}$, which implies that the Radon-Nikodym in (1.8) is L^p integrable for any $p > 0$, known as the L^∞ -regularity Brownian regularity.

Let us re-examine the estimates we did in the above argument. We expect that the bound on the normalizing constant is sharp. Let us briefly explain why. The scaled KPZ line ensemble is

stationary around a parabola $-2^{-1}x^2$, i.e. $\mathfrak{H}_k(x) + 2^{-1}x^2$ is a stationary process in x for each $k \in \mathbb{N}$. When we resampling a curve over a large interval of size T , having in mind that the penalization (W) is extremely high for going out of order, the curve tries to rise up and jump over the parabola $-2^{-1}x^2$. For a free Brownian motion, such movement consumes a kinetic cost of order e^{-T^3} . Therefore, we expect that typically $Z_{\text{bdd}} \approx e^{-T^3}$, which matches with what we have in (1.10). We admit that there is possible room to improve the estimates in (1.10) due to the presence of the term $O(1)$. However, it seems quite difficult to pursue in this direction and we do not attempt to do so in this paper.

Expecting that the estimate on Z_{bdd} is sharp, the overestimate in above argument comes mainly from the step where we estimated the numerator, i.e.

$$(1.12) \quad \mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W] \leq \mathbb{P}_{\text{free}}(B \in A).$$

In the next subsection, we explain how to improve this step by introducing the "soft" jump ensemble method. Let conclude this section by comparing the resampling methods developed in [Wu21] and this paper, see Table 1.

Level	Method	Results
Basic	Simple resampling	Uniform tail bound
Intermediate	Inductive middle reconstruction	Local fluctuation, Tightness
Advanced	"Soft" jump ensemble	L^∞ -Brownian regularity

TABLE 1. A brief summary of the different stages of the resampling methods developed in [Wu21] and in this paper. The first two stages were developed in [Wu21] and the more delicate stage is developed in this paper. The inputs for the first stage are the one point tail estimates ([CG1, CG2]) of the top layer \mathfrak{H}_1^t and stationarity of \mathfrak{H}^t around parabola $\frac{x^2}{2}$. Then the results of previous layers will serve as inputs for the next one. In the first stage, the resampling is straightforward and uniform tail estimates are established for each layer $\mathfrak{H}_k^t, k \in \mathbb{N}$. In the second stage, an inductive middle reconstruction is used to separate curves to obtain quantitative control of the normalizing constant. A more advanced resampling technique, the soft jump ensemble, is an adaption of the jump ensemble method developed in [Ham1] for non-intersecting line ensembles.

1.5. The soft jump ensemble. In order to sharpen the estimate in (1.12), a new ensemble J , called the "soft" jump ensemble, is introduced. This method is heavily inspired by the jump ensemble introduced in [Ham1] in which the author studies non-intersecting Brownian Gibbsian ensembles. Since curves in the scaled KPZ line ensemble do not avoid each other, the soft jump ensemble is invented to adapt this intersecting nature. We will just call it the jump ensemble in the rest of the paper to save us some effort.

The jump ensemble J interpolates between the scaled KPZ line ensemble \mathfrak{H} and the Brownian bridge ensemble and seeks a good balance between those two. The idea is to decompose W into

$$W = W_{\text{jump}} \cdot W_{\text{rest}}.$$

We will choose both W_{jump} and W_{rest} to be less than or equal to 1. There are two extremal cases, $W_{\text{jump}} \equiv 1$ or $W_{\text{jump}} \equiv W$. The first case reduces to the resampling method discussed in the

previous section. The second case is informative as the jump ensemble measure is equivalent to the conditional measure of the line ensemble, which is the measure we need to study. W_{jump} will be designed carefully to provide a meaningful resampling strategy.

Let us first explain the argument framework. We may rewrite

$$\begin{aligned} \frac{\mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W]}{\mathbb{E}_{\text{free}}[W]} &= \frac{\mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W_{\text{rest}} \cdot W_{\text{jump}}]}{\mathbb{E}_{\text{free}}[W_{\text{rest}} \cdot W_{\text{jump}}]} \bigg/ \frac{\mathbb{E}_{\text{free}}[W_{\text{jump}}]}{\mathbb{E}_{\text{free}}[W_{\text{jump}}]} \\ &= \frac{\mathbb{E}_J[\mathbb{1}\{J \in A\} \cdot W_{\text{rest}}]}{\mathbb{E}_J[W_{\text{rest}}]}. \end{aligned}$$

Here the law of the jump ensemble J is specified through the Radon-Nikodym relation

$$\frac{d\mathbb{P}_J}{d\mathbb{P}_{\text{free}}}(B) = \frac{W_{\text{jump}}(B)}{\mathbb{E}_{\text{free}}[W_{\text{jump}}]}.$$

Because $W_{\text{rest}} \leq 1$, it holds that

$$(1.13) \quad \frac{\mathbb{E}_{\text{free}}[\mathbb{1}\{B \in A\} \cdot W]}{\mathbb{E}_{\text{free}}[W]} \leq \frac{\mathbb{P}_J(J \in A)}{\mathbb{E}_J[W_{\text{rest}}]}.$$

The above computation reduces the one in Subsection 1.4 when $W_{\text{jump}} = 1$. See Section 3 for a detailed account of this framework.

Since $W \leq W_{\text{rest}} \leq 1$, (1.13) is sharper than (1.12). Unfortunately in (1.13), bounding $\mathbb{P}_J(J \in A)$ from above and $\mathbb{E}_J[W_{\text{rest}}]$ from below oppose each other. To achieve a bound of order $\varepsilon^{1-o(1)}$ as we wish, we seek for the following balance when estimating $\mathbb{P}_J(J \in A)$ and $\mathbb{E}_J[W_{\text{rest}}]$ in (1.13).

$$(1.14) \quad \mathbb{P}_J(J \in A) = \varepsilon^{1-o(1)},$$

and

$$(1.15) \quad \mathbb{E}_J[W_{\text{rest}}] = \varepsilon^{o(1)}.$$

In light of $\varepsilon = \mathbb{P}_{\text{free}}(B \in A)$, (1.14) indicates that the jump ensemble J is comparable to the Brownian bridge ensemble. On the other hand, (1.15) shows that the jump ensemble J is a good approximation to the scaled KPZ line ensemble \mathfrak{H} . In particular, W_{rest} needs not to be small. The two requirements are opposing each other since (1.14) wants to make W_{jump} close to 1 but (1.15) wants to make W_{jump} close to W .

Our main contribution when designing the soft jump ensemble J is to carefully balance W_{jump} and W_{rest} and to find a common ground between these two Boltzmann weights. This framework is further explained in more detail in Section 3. We refer the readers to Section 6 for more explanation about the design concept of W_{jump} and how it is constructed precisely as it requires significantly heavier notation.

1.6. Outline. Section 2.1 contains various definitions necessary to describe Gibbsian line ensembles. Section 2.2 records lemmas about stochastic monotonicity and strong Markov property. Section 3 provides a framework of the main argument and splits the proof into three key propositions. The propositions are further proved in Section 4, Section 7 and Section 8 respectively. The main tool, the soft jump ensemble is designed in Section 6.

1.7. Notation. We would like to explain some notation here. The natural numbers are defined to be $\mathbb{N} = 1, 2, \dots$. Events are denoted in a special font \mathbf{E} , their indicator function is written either as $\mathbb{1}_{\mathbf{E}}$ and their complement is written as \mathbf{E}^c .

Universal constants will be generally denoted by C and constants depending only on k will be denoted as $D(k)$. We label the ones in statements of theorems, lemmas and propositions with

subscripts (e.g. $D_1(k), D_2(k), \dots$ based on their order of occurrence) but the constants in proofs may change value from line to line.

1.8. Acknowledgment. The author extends thanks to Alan Hammond for valuable discussions and the three Minerva lectures that Alan Hammond has given at Columbia in Spring 2019 about Gibbsian resampling techniques. The author also thanks Ivan Corwin, Alan Hammond and Milind Hegde for valuable suggestions on an earlier draft of this paper.

2. **H**-BROWNIAN GIBBS PROPERTY

In this section we introduce the notion of **H**-Brownian Gibbsian line ensembles. We first introduce the basic definitions necessary to define Gibbsian line ensembles in Subsection 2.1 and then provide properties of such Gibbsian line ensembles.

2.1. Line ensembles and the **H**-Brownian Gibbs property.

Definition 2.1 (Line ensembles). *Let Σ be an interval of \mathbb{Z} and let Λ be a subset of \mathbb{R} . Consider the set $C(\Sigma \times \Lambda, \mathbb{R})$ of continuous functions $f : \Sigma \times \Lambda \rightarrow \mathbb{R}$ endowed with the topology of uniform convergence on compact subsets of $\Sigma \times \Lambda$, and let $\mathcal{C}(\Sigma \times \Lambda, \mathbb{R})$ denote the sigma-field generated by Borel sets in $C(\Sigma \times \Lambda, \mathbb{R})$. A $\Sigma \times \Lambda$ -indexed line ensemble \mathcal{L} is a random variable on a probability space $(\Omega, \mathcal{B}, \mathbb{P})$, taking values in $C(\Sigma \times \Lambda, \mathbb{R})$ such that \mathcal{L} is a measurable function from \mathcal{B} to $\mathcal{C}(\Sigma \times \Lambda, \mathbb{R})$.*

We think of such line ensembles as multi-layer random curves. For integers $k_1 < k_2$, let $[k_1, k_2]_{\mathbb{Z}} := \{k_1, k_1 + 1, \dots, k_2\}$. We will generally write $\mathcal{L} : \Sigma \times \Lambda \rightarrow \mathbb{R}$ even though it is not \mathcal{L} , but rather $\mathcal{L}(\omega)$ for each $\omega \in \Omega$ which is such a function. We will also sometimes specify a line ensemble by only giving its law without reference to the underlying probability space. We write $\mathcal{L}_i(\cdot) := (\mathcal{L}(\omega))(i, \cdot)$ for the label $i \in \Sigma$ curve of the ensemble \mathcal{L} .

We now start to formulate the **H**-Brownian Gibbs property, a key property of the KPZ line ensemble, [CH16]. We adopt the convention that all Brownian motions and bridges have diffusion parameter one.

Definition 2.2 (**H**-Brownian bridge line ensemble). *Fix $k_1 \leq k_2$ with $k_1, k_2 \in \mathbb{Z}$, an interval $[a, b] \subset \mathbb{R}$ and two vectors $\vec{x}, \vec{y} \in \mathbb{R}^{k_2 - k_1 + 1}$. A $[k_1, k_2]_{\mathbb{Z}} \times [a, b]$ -indexed line ensemble $\mathcal{L} = (\mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2})$ is called a free Brownian bridge line ensemble with entrance data \vec{x} and exit data \vec{y} if its law $\mathbb{P}_{\text{free}}^{k_1, k_2, [a, b], \vec{x}, \vec{y}}$ is that of $k_2 - k_1 + 1$ independent standard Brownian bridges starting at time a at the points \vec{x} and ending at time b at the points \vec{y} .*

A Hamiltonian \mathbf{H} is defined to be a measurable function $\mathbf{H} : \mathbb{R} \rightarrow [0, \infty]$. Given a Hamiltonian \mathbf{H} and two measurable function $f, g : (a, b) \rightarrow \mathbb{R} \cup \{\pm\infty\}$, we define the **H**-Brownian bridge line ensemble with entrance data \vec{x} , exit data \vec{y} and boundary data (f, g) to be a $[k_1, k_2]_{\mathbb{Z}} \times (a, b)$ -indexed line ensemble $\mathcal{L} = (\mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2})$ with law $\mathbb{P}_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}$ given according to the following Radon-Nikodym derivative relation:

$$\frac{d\mathbb{P}_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}}{d\mathbb{P}_{\text{free}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}}}(\mathcal{L}) := \frac{W_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}(\mathcal{L})}{Z_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}}.$$

Here we adopt conventions that $\mathcal{L}_{k_1-1} = f$, $\mathcal{L}_{k_2+1} = g$ and define the Boltzmann weight

$$(2.1) \quad W_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}(\mathcal{L}) := \exp \left\{ - \sum_{i=k_1-1}^{k_2} \int_a^b \mathbf{H}(\mathcal{L}_{i+1}(u) - \mathcal{L}_i(u)) du \right\},$$

and the normalizing constant

$$(2.2) \quad Z_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g} := \mathbb{E}_{\text{free}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}} \left[W_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}(\mathcal{L}) \right],$$

where \mathcal{L} in the above expectation is distributed according to the measure $\mathbb{P}_{\text{free}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}}$.

H-Brownian Gibbs property could be viewed as a spatial Markov property, more specifically, it provides a description of the conditional law inside a compact set.

Definition 2.3 (**H**-Brownian Gibbs property). *A $\Sigma \times \Lambda$ -indexed line ensemble \mathcal{L} satisfies the **H**-Brownian Gibbs property if for all $K = \{k_1, \dots, k_2\} \subset \Sigma$ and $(a, b) \subset \Lambda$, its conditional law inside K takes the following form,*

$$\text{Law} \left(\mathcal{L} \Big|_{K \times (a, b)} \text{ conditional on } \mathcal{L} \Big|_{(\Sigma \times \Lambda) \setminus (K \times (a, b))} \right) = \mathbb{P}$$

Here $f := \mathcal{L}_{k_1-1}$ and $g := \mathcal{L}_{k_2+1}$ with the convention that if $k_1 - 1 \notin \Sigma$ then $f \equiv +\infty$ and likewise if $k_2 + 1 \notin \Sigma$ then $g \equiv -\infty$; we have also set $\vec{x} = (\mathcal{L}_{k_1}(a), \dots, \mathcal{L}_{k_2}(a))$ and $\vec{y} = (\mathcal{L}_{k_1}(b), \dots, \mathcal{L}_{k_2}(b))$.

This following description of Gibbs property using conditional expectation is equivalent and is convenient for computations sometimes. For $K \subset \mathbb{Z}$ and $S \subset \mathbb{R}$, let $C(K \times S, \mathbb{R})$ be the space of continuous functions from $K \times S \rightarrow \mathbb{R}$. Then a $\Sigma \times \Lambda$ -indexed line ensemble \mathcal{L} enjoys the **H**-Brownian Gibbs property if and only if for any $K = \{k_1, \dots, k_2\} \subset \Sigma$ and $(a, b) \subset \Lambda$, and any Borel function F from $C(K \times (a, b), \mathbb{R}) \rightarrow \mathbb{R}$, \mathbb{P} -almost surely

$$\mathbb{E} \left[F(\mathcal{L} \Big|_{K \times (a, b)}) \mid \mathcal{F}_{\text{ext}}(K \times (a, b)) \right] = \mathbb{E}_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g} \left[F(\mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2}) \right],$$

where \vec{x}, \vec{y}, f and g are defined in the previous paragraph and where

$$(2.3) \quad \mathcal{F}_{\text{ext}}(K \times (a, b)) := \sigma(\mathcal{L}_i(s) : (i, s) \in \Sigma \times \Lambda \setminus K \times (a, b))$$

is the exterior sigma-field generated by the line ensemble outside $K \times (a, b)$. On the left-hand side of the above equality $\mathcal{L} \Big|_{K \times (a, b)}$ is the restriction to $K \times (a, b)$ of curves distributed according to \mathbb{P} , while on the right-hand side $\mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2}$ are curves on (a, b) distributed according to $\mathbb{P}_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}, \vec{y}, f, g}$.

Remark 2.4. *The scaled KPZ line ensemble enjoys an \mathbf{H}_t -Brownian Gibbs property with*

$$(2.4) \quad \mathbf{H}_t(x) = e^{t^{1/3}x}.$$

Throughout this paper, we only focus on this specific Hamiltonian.

2.2. Strong Gibbs property and stochastic monotonicity. We record some important properties, developed in [CH16, Section 2], about **H**-Gibbsian line ensembles in this section. We begin with the strong Gibbs property, which enable us to resample the trajectory within a stopping domain as opposed to a deterministic interval.

Definition 2.5. *Let Σ be an interval of \mathbb{Z} , and Λ be an interval of \mathbb{R} . Consider a $\Sigma \times \Lambda$ -indexed line ensemble \mathcal{L} which has the **H**-Brownian Gibbs property for some Hamiltonian \mathbf{H} . For $K = \{k_1, \dots, k_2\} \subseteq \Sigma$ and $(\ell, r) \subseteq \Lambda$, $\mathcal{F}_{\text{ext}}(K \times (\ell, r))$ denotes the sigma field generated by the data outside $K \times (\ell, r)$. The random variable $(\mathfrak{l}, \mathfrak{r})$ is called a K -stopping domain if for all $\ell < r$,*

$$\{\mathfrak{l} \leq \ell, \mathfrak{r} \geq r\} \in \mathcal{F}_{\text{ext}}(K \times (\ell, r)).$$

For $K = \{k_1, \dots, k_2\} \subset \mathbb{Z}$, define

$$C_K := \{(\ell, r, f_{k_1}, \dots, f_{k_2}) : \ell < r, (f_{k_1}, \dots, f_{k_2}) \in C(K \times (\ell, r))\}.$$

Lemma 2.6. *Consider a $\Sigma \times \Lambda$ -indexed line ensemble \mathcal{L} which has the **H**-Brownian Gibbs property. Fix $K = \{k_1, \dots, k_2\} \subseteq \Sigma$. For all random variables $(\mathfrak{l}, \mathfrak{r})$ which are K -stopping domains for \mathcal{L} , the following strong **H**-Brownian Gibbs property holds: for all Borel functions $F : C_K \rightarrow \mathbb{R}$, \mathbb{P} -almost surely,*

$$(2.5) \quad \mathbb{E} \left[F(\mathfrak{l}, \mathfrak{r}, \mathcal{L} \Big|_{K \times (\mathfrak{l}, \mathfrak{r})}) \mid \mathcal{F}_{\text{ext}}(K \times (\mathfrak{l}, \mathfrak{r})) \right] = \mathbb{E}_{\mathbf{H}}^{k_1, k_2, (\mathfrak{l}, \mathfrak{r}), \vec{x}, \vec{y}, f, g} \left[F(a, b, \mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2}) \right],$$

where $a = \mathfrak{l}$, $b = \mathfrak{r}$, $\vec{x} = \{\mathcal{L}_i(\mathfrak{l})\}_{i=k_1}^{k_2}$, $\vec{y} = \{\mathcal{L}_i(\mathfrak{r})\}_{i=k_1}^{k_2}$, $f(\cdot) = \mathcal{L}_{k_1-1}(\cdot)$ (or ∞ if $k_1 - 1 \notin \Sigma$), $g(\cdot) = \mathcal{L}_{k_2+1}(\cdot)$ (or $-\infty$ if $k_2 + 1 \notin \Sigma$). On the left-hand side $\mathcal{L}|_{K \times (\mathfrak{l}, \mathfrak{r})}$ is the restriction of curves distributed according to \mathbb{P} and on the right-hand side $\mathcal{L}_{k_1}, \dots, \mathcal{L}_{k_2}$ is distributed according to $\mathbb{P}_{\mathbf{H}}^{k_1, k_2, (\mathfrak{l}, \mathfrak{r}), \vec{x}, \vec{y}, f, g}$.

For a convex Hamiltonian \mathbf{H} (such as \mathbf{H}_t), \mathbf{H} -Brownian bridge line ensembles \mathbb{P} .

Lemma 2.7. Fix $k_1 \leq k_2 \in \mathbb{Z}$, $a < b$. Consider two pairs of vectors $\vec{x}^{(i)}, \vec{y}^{(i)} \in \mathbb{R}^{k_2 - k_1 + 1}$ for $i \in \{1, 2\}$ such that $x_j^{(1)} \leq x_j^{(2)}$ and $y_j^{(1)} \leq y_j^{(2)}$ for all $j = k_1, \dots, k_2$. Consider two pairs of measurable functions $(f^{(i)}, g^{(i)})$ for $i \in \{1, 2\}$ such that $f^{(i)} : (a, b) \rightarrow \mathbb{R} \cup \{\infty\}$, $g^{(i)} : (a, b) \rightarrow \mathbb{R} \cup \{-\infty\}$ and for all $s \in (a, b)$, $f^{(1)}(s) \geq f^{(2)}(s)$ and $g^{(1)}(s) \geq g^{(2)}(s)$. For $i \in \{1, 2\}$, let $\mathcal{Q}^{(i)} = \{\mathcal{Q}_j^{(i)}\}_{j=k_1}^{k_2}$ be a $\{k_1, \dots, k_2\} \times (a, b)$ -indexed line ensemble on a probability space $(\Omega^{(i)}, \mathcal{B}^{(i)}, \mathbb{P}^{(i)})$ where $\mathbb{P}^{(i)}$ equals $\mathbb{P}_{\mathbf{H}}^{k_1, k_2, (a, b), \vec{x}^{(i)}, \vec{y}^{(i)}, f^{(i)}, g^{(i)}}$ (i.e. $\mathcal{Q}^{(i)}$ has the \mathbf{H} -Brownian Gibbs property with entrance data $\vec{x}^{(i)}$, exit data $\vec{y}^{(i)}$ and boundary data $(f^{(i)}, g^{(i)})$).

If the Hamiltonian $\mathbf{H} : \mathbb{R} \rightarrow [0, \infty)$ is convex then there exists a coupling of the probability measures $\mathbb{P}^{(1)}$ and $\mathbb{P}^{(2)}$ such that almost surely $\mathcal{Q}_j^{(1)}(s) \leq \mathcal{Q}_j^{(2)}(s)$ for all $j \in \{k_1, \dots, k_2\}$ and all $s \in (a, b)$.

3. PROOF FRAMEWORK FOR THEOREM 1.3

In this section we attempt to illustrate the structure our soft jump ensemble resampling argument towards proving the main Theorem 1.3. We provide the framework of the argument, containing a core inequality (3.14) and three key propositions, i.e. Propositions 3.4, 3.5 and 3.6. Combining these three Propositions and the core inequality (3.14), the main Theorem 1.3 readily follows. We prove these Propositions in sections 4, 7 and 8 respectively.

In this section we list essential properties that are necessary for deriving (3.14) for the reader's ease. We postpone explicit definitions of various objects to sections 4 and 6 and refer readers to those sections.

3.1. Introducing the resampling domain. We begin with introducing the domain, a subset in $\mathbb{N} \times \mathbb{R}$, where we run the resampling for \mathfrak{H} , i.e. apply the Gibbs property. We also introduce some relevant subsets, in which the boundary conditions are encoded. See Figure 1 for an illustration of these region.

Fix $k \in \mathbb{N}$ and $s \geq 1$. Take $\ell_0 < 0$ and $r_0 > s$, whose precise definitions and dependence on k can be found in (4.2). Let $U \subset \mathbb{R}^k \times \mathbb{R}^k$ be the collection of $(\bar{\ell}, \bar{r}) = (\ell_1, \ell_2, \dots, \ell_k, r_1, r_2, \dots, r_k)$ such that

$$(3.1) \quad \ell_0 < \ell_1 < \ell_2 < \dots < \ell_k < 0, \quad s < r_k < r_{k-1} < \dots < r_1 < r_0.$$

We assume $(\bar{\ell}, \bar{r}) \in U$ throughout this section. Define

$$(3.2) \quad \text{dom}(\bar{\ell}, \bar{r}) := \bigcup_{j=1}^k \{j\} \times [\ell_j, r_j],$$

$$(3.3) \quad \text{bdd}(\bar{\ell}, \bar{r}) := \{1\} \times \{\ell_1, r_1\} \cup \bigcup_{j=2}^k \{j\} \times ([\ell_{j-1}, \ell_j] \cup [r_j, r_{j-1}]) \cup \{k+1\} \times [\ell_k, r_k],$$

$$(3.4) \quad \text{ext}(\bar{\ell}, \bar{r}) := \text{cl}(\mathbb{N} \times \mathbb{R} \setminus \text{dom}(\bar{\ell}, \bar{r})),$$

$$(3.5) \quad \text{Box} := [1, k+1]_{\mathbb{Z}} \times [\ell_0, r_0],$$

$$(3.6) \quad \text{Jct}(\bar{\ell}, \bar{r}) := \text{cl}(\text{Box} \setminus \text{dom}(\bar{\ell}, \bar{r})),$$




FIGURE 1. An illustration when $k = 3$. We will run the resampling for the first three curve $\mathfrak{H}_1, \mathfrak{H}_3, \mathfrak{H}_3$ inside dom , which resembles the shape of an upside-down pyramid. More precisely, $\text{dom} := (\{1\} \times [\ell_1, r_1]) \cup (\{2\} \times [\ell_2, r_2]) \cup (\{3\} \times [\ell_3, r_3])$. We use bdd to denote the region where the values of \mathfrak{H} involves in the resampling as the boundary data. ext is the the closure of the complement of dom . Box and Jct are introduced for later notational convenience.

where cl means taking the closure. We write $C(\text{dom}(\bar{\ell}, \bar{r}))$, $C(\text{bdd}(\bar{\ell}, \bar{r}))$, $C(\text{Jct}(\bar{\ell}, \bar{r}))$ and $C(\text{Box}(\bar{\ell}, \bar{r}))$ to denote the spaces of continuous functions defined on $\text{dom}(\bar{\ell}, \bar{r})$, $\text{bdd}(\bar{\ell}, \bar{r})$, $\text{Jct}(\bar{\ell}, \bar{r})$ and $\text{Box}(\bar{\ell}, \bar{r})$ respectively. We proceed to formulate the Gibbs property of \mathfrak{H} restricted to $\text{dom}(\bar{\ell}, \bar{r})$. Let $f = (f_1, f_2, \dots, f_{k+1}) : \text{bdd}(\bar{\ell}, \bar{r}) \rightarrow \mathbb{R}$ be a continuous function. Let $\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ be the law of k independent Brownian bridges (B_1, B_2, \dots, B_k) defined in $\text{dom}(\bar{\ell}, \bar{r})$ with $B_j(\ell_j) = f_j(\ell_j)$ and $B_j(r_j) = f_j(r_j)$. We write $\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ for its expectation.

Recall that $\mathbf{H}_t(x) = e^{t^{1/3}x}$. Define the law $\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ through the following Radon-Nikodym derivative relation,

$$(3.7) \quad \frac{d\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}}{d\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}}(\mathcal{L}) := \frac{W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L})}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f} \left[W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f} \right]}.$$

The above Boltzmann weight $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L})$ represents the interaction between the curves $\mathcal{L}_1, \dots, \mathcal{L}_k$ in $\text{dom}(\bar{\ell}, \bar{r})$ and their interaction with the boundary data f , defined as

$$(3.8) \quad \begin{aligned} W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) &:= \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_j, \ell_{j+1}] \cup [r_{j+1}, r_j]} \mathbf{H}_t(f_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ &\times \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_{j+1}, r_{j+1}]} \mathbf{H}_t(\mathcal{L}_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ &\times \exp \left(- \int_{[\ell_k, r_k]} \mathbf{H}_t(f_{k+1}(x) - \mathcal{L}_k(x)) dx \right). \end{aligned}$$

Now we are ready to formulate the \mathbf{H}_t -Brownian Gibbs property of \mathfrak{H}^t on $\text{dom}(\bar{\ell}, \bar{r})$. Conditional on the boundary data, inside the upside-down pyramid $\text{dom}(\bar{\ell}, \bar{r})$, the law of \mathfrak{H} is equivalent to $\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$.

Lemma 3.1. *Fix $t > 0$, $k \in \mathbb{N}$ and $(\bar{\ell}, \bar{r}) \in U$. Let $\mathcal{F}_{\text{ext}}(\bar{\ell}, \bar{r})$ be the σ -algebra generated by $\mathfrak{H}^t|_{\text{ext}(\bar{\ell}, \bar{r})}$, i.e.*

$$(3.9) \quad \mathcal{F}_{\text{ext}}(\bar{\ell}, \bar{r}) := \sigma(\mathfrak{H}^t|_{\text{ext}(\bar{\ell}, \bar{r})}).$$

Let $f = \mathfrak{H}^t|_{\text{bdd}(\bar{\ell}, \bar{r})}$. For any Borel function $F : C(\text{dom}(\bar{\ell}, \bar{r})) \rightarrow \mathbb{R}$, almost surely it holds that

$$(3.10) \quad \mathbb{E}[F(\mathfrak{H}^t|_{\text{dom}(\bar{\ell}, \bar{r})}) | \mathcal{F}_{\text{ext}}(\bar{\ell}, \bar{r})] = \mathbb{E}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}[F(\mathcal{L})].$$

Actually, we need to apply the \mathbf{H}_t -Brownian Gibbs property over stopping domains, known as the strong \mathbf{H}_t -Brownian Gibbs property. Let us first recall the definition of stopping domains in the current context. Consider a random variable $(\bar{\mathbf{l}}, \bar{\mathbf{r}}) = (\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_k, \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_k)$ that takes value in U . The explicit definition of $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ can be found in (4.4).

Definition 3.2. *The random variable $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is called a stopping domain if the event*

$$\mathbf{l}_j \leq \ell_j, \quad \mathbf{r}_j \geq r_j \quad \text{for all } j \in [1, k]_{\mathbb{Z}}$$

is $\mathcal{F}_{\text{ext}}(\bar{\ell}, \bar{r})$ -measurable with $\bar{\ell} = (\ell_1, \ell_2, \dots, \ell_k)$ and $\bar{r} = (r_1, r_2, \dots, r_k)$.

From now on, we assume $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is a stopping domain. See Lemma 4.4 for a proof. We record the strong \mathbf{H}_t -Brownian Gibbs property in the following lemma. Define

$$C_{\text{dom}} := \Pi \{(\bar{\ell}, \bar{r}) \times C(\text{dom}(\bar{\ell}, \bar{r})) \mid (\bar{\ell}, \bar{r}) \in U\}.$$

We equip C_{dom} with the topology induced from $U \times C([1, k+1]_{\mathbb{Z}} \times \mathbb{R})$.

Lemma 3.3. *Let $t > 0$, $k \in \mathbb{N}$ and let $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ be a stopping domain that takes value in U . Let $f = \mathfrak{H}^t|_{\text{bdd}(\bar{\mathbf{l}}, \bar{\mathbf{r}})}$. For any function $F : C_{\text{dom}} \rightarrow \mathbb{R}$, almost surely it holds that*

$$(3.11) \quad \mathbb{E}[F(\bar{\mathbf{l}}, \bar{\mathbf{r}}, \mathfrak{H}^t|_{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}})}) | \mathcal{F}_{\text{ext}}(\bar{\mathbf{l}}, \bar{\mathbf{r}})] = \mathbb{E}_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[F(\mathcal{L})].$$

3.2. The core inequality. In this section we apply the strong \mathbf{H}_t -Brownian Gibbs property and derive the core inequality (3.14). We begin with formulating the favorable event Fav which confines the behavior of \mathfrak{H} in $\text{Jct}(\bar{\mathbf{l}}, \bar{\mathbf{r}})$.

Define

$$C_{\text{Jct}} := \Pi \{(\bar{\ell}, \bar{r}) \times C(\text{Jct}(\bar{\ell}, \bar{r})) \mid (\bar{\ell}, \bar{r}) \in U\}.$$

We equip C_{Jct} with the topology induced from $U \times C([1, k+1]_{\mathbb{Z}} \times \mathbb{R})$. Let $\mathcal{G} \subset C_{\text{Jct}}$ be a Borel set which contains the realization of all nice boundary data for our purpose. The explicit form of \mathcal{G} can be found in Definition 4.5. See Lemma 4.6 for the measurability of \mathcal{G} . Define the favorable event

$$(3.12) \quad \text{Fav} := \{(\bar{\mathbf{l}}, \bar{\mathbf{r}}, \mathfrak{H}^t|_{\text{Jct}(\bar{\mathbf{l}}, \bar{\mathbf{r}})}) \in \mathcal{G}\}.$$

Fav is measurable because $\text{Fav} = \{\pi \circ (\bar{\mathbf{l}}, \bar{\mathbf{r}}, \mathfrak{H}) \in \mathcal{G}\}$ with $\pi : U \times C([1, k+1]_{\mathbb{Z}} \times \mathbb{R}) \rightarrow C_{\text{Jct}}$ being the restriction map. Moreover, Fav is $\mathcal{F}_{\text{ext}}(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ -measurable since $\text{Jct}(\bar{\ell}, \bar{r}) \subset \text{ext}(\bar{\ell}, \bar{r})$.

Next, we discuss the jump ensemble J . The jump ensemble J is a proxy of \mathfrak{H}^t . The jump ensemble J is obtained by replacing the Boltzmann weight $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ by W_{jump} . The new weight W_{jump} is less restrictive in the sense that $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f} \leq W_{\text{jump}} \leq 1$. Therefore, compared to \mathfrak{H}^t , J is closer to Brownian bridges. The extra price to pay is to compare the proxy J to \mathfrak{H}^t .

We prescribe the law of J below. Given $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$, we consider the Boltzmann weight $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ with $f = f_J|_{\text{bdd}(\bar{\ell}, \bar{r})}$. We decompose $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ into two parts,

$$W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) = W_{\text{jump}}(\mathcal{L}; f_J) \cdot W_{\text{rest}}(\mathcal{L}; f_J).$$

The explicit form of $W_{\text{jump}}(\mathcal{L}; f_J)$ and $W_{\text{rest}}(\mathcal{L}; f_J)$ can be found in (6.15) and (6.16). The only property we use here is that $0 < W_{\text{rest}}(\mathcal{L}; f_J) \leq 1$, see Lemma 6.4. The jump ensemble J consists

of continuous random curves defined on $\text{dom}(\bar{\ell}, \bar{r})$. The law of J , denoted by \mathbb{P}_J , is defined through the following Radon-Nikodym derivative relation,

$$(3.13) \quad \frac{d\mathbb{P}_J}{d\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}}(\mathcal{L}) := \frac{W_{\text{jump}}(\mathcal{L})}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}[W_{\text{jump}}]}.$$

We note that even though $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ depends only on f , W_{jump} and \mathbb{P}_J do depend on f_J .

We are now ready to formulate and to prove the core inequality. Let $A \subset C_{0,0}([0, s])$ be an arbitrary Borel set. It holds that

$$(3.14) \quad \mathbb{P}(\mathfrak{H}_k^{t, [0, s]} \in A) \leq \sup \left\{ \frac{\mathbb{P}_J(J_k^{[0, s]} \in A)}{\mathbb{E}_J[W_{\text{rest}}]} \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G} \right\} + \mathbb{P}(\text{Fav}^c).$$

Proof of (3.14). Note that

$$\mathbb{P}(\mathfrak{H}_k^{t, [0, s]} \in A) \leq \mathbb{P}(\mathfrak{H}_k^{t, [0, s]} \in A, \text{Fav}) + \mathbb{P}(\text{Fav}^c) = \mathbb{E}[\mathbb{1}_{\text{Fav}} \cdot \mathbb{E}[\mathbb{1}\{\mathfrak{H}_k^{[0, s]} \in A\} \mid \mathcal{F}_{\text{ext}}(\bar{\mathbf{l}}, \bar{\mathbf{r}})]] + \mathbb{P}(\text{Fav}^c).$$

From (3.1) and the strong \mathbf{H}_t -Brownian Gibbs property, it holds that

$$\mathbb{E}[\mathbb{1}\{\mathfrak{H}_k^{t, [0, s]} \in A\} \mid \mathcal{F}_{\text{ext}}(\bar{\mathbf{l}}, \bar{\mathbf{r}})] = \frac{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[\mathbb{1}\{B_k^{[0, s]} \in A\} \cdot W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]}.$$

Here $f = \mathfrak{H}^t|_{\text{bdd}(\bar{\mathbf{l}}, \bar{\mathbf{r}})}$. The event Fav implies $(\bar{\mathbf{l}}, \bar{\mathbf{r}}, \mathfrak{H}^t|_{\text{Jct}(\bar{\mathbf{l}}, \bar{\mathbf{r}})}) \in \mathcal{G}$. Hence the term above is bounded by

$$\sup \left\{ \frac{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[\mathbb{1}\{B_k^{[0, s]} \in A\} \cdot W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]} \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}, f = f_J|_{\text{bdd}(\bar{\ell}, \bar{r})} \right\}.$$

In short, we obtain

$$\mathbb{P}(\mathfrak{H}_k^{t, [0, s]} \in A) \leq \sup \left\{ \frac{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[\mathbb{1}\{B_k^{[0, s]} \in A\} \cdot W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]} \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}, f = f_J|_{\text{bdd}(\bar{\ell}, \bar{r})} \right\} + \mathbb{P}(\text{Fav}^c).$$

Because $W_{\text{rest}}(\mathcal{L}; f_J) \leq 1$,

$$\frac{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[\mathbb{1}\{B_k^{[0, s]} \in A\} \cdot W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}[W_{\mathbf{H}_t}^{\text{dom}(\bar{\mathbf{l}}, \bar{\mathbf{r}}), f}]} = \frac{\mathbb{E}_J[\mathbb{1}\{J_k^{[0, s]} \in A\} \cdot W_{\text{rest}}]}{\mathbb{E}_J[W_{\text{rest}}]} \leq \frac{\mathbb{P}_J(J_k^{[0, s]} \in A)}{\mathbb{E}_J[W_{\text{rest}}]}.$$

It immediately follows that

$$\mathbb{P}(\mathfrak{H}_k^{t, [0, s]} \in A) \leq \sup \left\{ \frac{\mathbb{P}_J(J_k^{[0, s]} \in A)}{\mathbb{E}_J[W_{\text{rest}}]} \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G} \right\} + \mathbb{P}(\text{Fav}^c).$$

□

3.3. Three key propositions. We state the three key propositions, based on which we prove the main Theorem 1.3. We first briefly explain the definitions necessary to state these propositions.

Fav is the favorable event defined in (3.12), which confines the boundary data, i.e. triples $(\bar{\ell}, \bar{r}, f_J)$ for the resampling. \mathcal{G} is the Borel subset consisting of good triples $(\bar{\ell}, \bar{r}, f_J)$. Given $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$, W_{jump} and W_{rest} are Boltzmann weights which are carefully designed, see in (6.15) and (6.16).

Let J be the jump ensemble defined in (3.13). Furthermore, recall that $\varepsilon = \mathbb{P}_{\text{free}}(B_k^{[0,s]} \in A)$ and D is used to denote a large constant that depends only on k and s . The exact value of D may change from line to line.

The following proposition shows that the favorable event is typical.

Proposition 3.4.

$$(3.15) \quad \mathbb{P}(\text{Fav}^c) \leq D\varepsilon.$$

For any realization of the favorable event, i.e. a triple $(\bar{\ell}, \bar{r}, f_J)$, we are able to estimate the denominator and numerator in the core inequality, (3.14), respectively in Proposition 3.5 and 3.6. The estimate of the denominator, Proposition 3.5 gauges how the jump ensemble J differs from free Brownian bridges while the estimate of the numerator, Proposition 3.6 measures the cost to pay for free Brownian bridges to become the jump ensemble.

Proposition 3.5.

$$(3.16) \quad \inf\{\mathbb{E}_J[W_{\text{rest}}] \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}\} \geq D^{-1} \exp\left(-D(\log \varepsilon^{-1})^{5/6}\right).$$

Proposition 3.6.

$$\sup\{\mathbb{P}_J(J^{[0,s]} \in A) \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}\} \leq D\varepsilon \exp\left(D(\log \varepsilon^{-1})^{5/6}\right).$$

Assuming Propositions 3.4, 3.5 and 3.6 hold, we may complete the proof of the main Theorem 1.3. The rest of this paper is devoted to prove these three key propositions.

Proof of Theorem 1.3. Combining (3.14) and Propositions 3.4, 3.5 and 3.6, we conclude

$$\mathbb{P}(\mathfrak{H}_k^{t,[0,s]} \in A) \leq D\varepsilon \exp\left(D(\log \varepsilon^{-1})^{5/6}\right).$$

□

4. FAVORABLE EVENT

In this section we explain in detail the definition of the favorable event

$$\text{Fav} := \{(\bar{\ell}, \bar{r}, \mathfrak{H}^t|_{\text{Jct}(\bar{\ell}, \bar{r})}) \in \mathcal{G}\}.$$

\mathcal{G} consists of nice boundary data that meets the four assumptions **C1-C4** in Definition 4.5. We begin with introducing the parameters involved in the assumptions. The final goal of this section is to prove Proposition 3.4, which bounds the probability of the complement of the favorable event.

Fix $k \in \mathbb{N}$ and $s \geq 1$. Let $E = E(k, s) \geq 10$ be a large constant to be determined later. Take

$$(4.1) \quad T = E \max\{(\log \varepsilon^{-1})^{1/3}, s\},$$

and

$$(4.2) \quad \ell_0 = -(k+1)T, \quad r_0 = (k+1)T.$$

In other words, T determines the width of the Box region. For ε small, T is comparable to $(\log \varepsilon^{-1})^{1/3}$. This choice of T was pioneered by Hammand [Ham1] for non-intersecting Brownian Gibbsian ensembles. In [Wu21], the same author studies the uniform tail bound of the normalizing




FIGURE 2. Illustration of the choice of l_j . l_j is chosen to be an extreme point of the concave majorant such that \mathfrak{H}_{j+1} can be bounded by a linear function. The key observation is that $\mathfrak{H}_{j+1}(x)$ stays close to a parabola $-2^{-1}x^2$ and therefore the concave majorant of \mathfrak{H}_{j+1} also stays close to $-2^{-1}x^2$ with derivative close to $-x$.

constant for the scaled KPZ line ensemble \mathfrak{H}^t , recorded in this paper as Proposition 4.8 for the reader's convenience. Roughly speaking, with probability $1-\varepsilon$, the normalizing constant is bounded from below by $\varepsilon^{O(1)}$ if it is evaluated in an interval of length $(\log \varepsilon^{-1})^{1/3}$. This bound allows us to pick $T \approx (\log \varepsilon^{-1})^{1/3}$ as the width of Box.

From now on, we will use T instead of ε to define various quantities. Define

$$(4.3) \quad \Delta := 8 \log T, \quad d := 64T^{-4}(\log T)^2.$$

We will explain how these parameters are chosen when it becomes clear.

Next, we discuss how to pick the stopping domains (\bar{l}, \bar{r}) . \mathfrak{H}_j^t will be resampled in the interval $[l_j, r_j]$. In order to ensure \mathfrak{H}_j^t lies above \mathfrak{H}_{j+1}^t , we need to control the growth of \mathfrak{H}_{j+1}^t . A crucial observation of Hammond [Ham1] is the following. If $\mathfrak{H}_{j+1}^t(x)$ is close to a parabola $-2^{-1}x^2$, then the concave majorant of \mathfrak{H}_{j+1}^t is also close to $-2^{-1}x^2$ with derivative close to $-x$. See Lemma A.8. Therefore, \mathfrak{H}_{j+1}^t can be bounded by a linear function by choosing l_j to be an extreme point of the concave majorant. See Figure 2 for an illustration.

Now we start to construct (\bar{l}, \bar{r}) . We first define (\bar{l}, \bar{r}) for continuous functions $f_B \in C(\text{Box})$ and then define (\bar{l}, \bar{r}) by

$$(4.4) \quad (\bar{l}, \bar{r}) := (\bar{l}, \bar{r})(\mathfrak{H}^t|_{\text{Box}}).$$

For $j \in [1, k]_{\mathbb{Z}}$, let

$$R_{j+1, \pm} := [\pm(k+1-j)T - 2^{-1}T, \pm(k+1-j)T + 2^{-1}T].$$

Given $f_B = (f_1, f_2, \dots, f_{k+1})$ in $C(\text{Box})$, we write f_{j+1} for the concave majorant of f_{j+1} in $R_{j+1, \pm}$. Let

$$\bar{R}_{j+1, \pm} := [\pm(k+1-j)T - 5^{-1}T, \pm(k+1-j)T + 5^{-1}T].$$

Define

$$\begin{aligned} I_{j+1, -} &:= \{x \in \bar{R}_{j+1, -} \mid (f_{j+1})'_+(x) \leq (k+1-j)T\}, \\ I_{j+1, +} &:= \{x \in \bar{R}_{j+1, +} \mid (f_{j+1})'_-(x) \geq -(k+1-j)T\}. \end{aligned}$$

and

$$\begin{aligned} l_j &:= \begin{cases} \inf I_{j+1, -} & \text{if } I_{j+1, -} \neq \phi, \\ -(k+1-j) + 5^{-1}T & \text{if } I_{j+1, -} = \phi, \end{cases} \\ r_j &:= \begin{cases} \sup I_{j+1, +} & \text{if } I_{j+1, +} \neq \phi, \\ ((k+1-j) - 5^{-1})T & \text{if } I_{j+1, +} = \phi, \end{cases} \end{aligned}$$

Remark 4.1. Suppose $I_{j+1,-}$ is non-empty and $I_{j+1,-} \subset \text{int}\bar{R}_{j+1,-}$, where int means taking the interior. Then \mathbf{l}_j is an extreme point of \mathfrak{f}_{j+1} . Therefore, $\mathfrak{f}_{j+1}(\mathbf{l}_j) = f_{j+1}(\mathbf{l}_j)$ and $(\mathfrak{f}_{j+1})'_+(\mathbf{l}_j) \leq (k+1-j)T$. See Lemma A.7 and Lemma A.1. By the concavity of \mathfrak{f}_{j+1} , it holds that for $x \in R_{j+1,-}$ and $x \geq \mathbf{l}_j$,

$$f_{j+1}(x) \leq f_{j+1}(\mathbf{l}_j) + (k+1-j)T(x - \mathbf{l}_j).$$

See Figure 2 for an illustration.

Putting \mathbf{l}_j and \mathbf{r}_j together, we define

$$(\bar{\mathbf{l}}, \bar{\mathbf{r}}) := (\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_k, \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_k).$$

Remark 4.2. To make sure $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ defined in (4.4) is a random variable, we check $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is Borel measurable. The map from $f_{\mathbb{B}}$ to $(\mathfrak{f}_2, \dots, \mathfrak{f}_{k+1})$ is continuous under the uniform topology. From the view of Lemma A.6, $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is semi-continuous and hence Borel measurable.

Remark 4.3. For any continuous function $f_{\mathbb{B}}$, it holds that $(\bar{\mathbf{l}}, \bar{\mathbf{r}}) \in U$. Actually, it always holds that $|\mathbf{l}_j + (k+1-j)T| \leq 5^{-1}T$ and $|\mathbf{r}_j - (k+1-j)T| \leq 5^{-1}T$.

Lemma 4.4. $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is a stopping domain which take values in U .

Proof. From Remark 4.3, $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ takes value in U . From the construction, $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ depends only on f_{j+1} in $R_{j+1,\pm}$. From Remark (4.3), it holds that $\{j+1\} \times R_{j+1,\pm} \subset \text{Jct}(\bar{\mathbf{l}}, \bar{\mathbf{r}})$. Because $\text{Jct}(\bar{\ell}, \bar{r}) \subset \text{ext}(\bar{\ell}, \bar{r})$, we conclude that $(\bar{\mathbf{l}}, \bar{\mathbf{r}})$ is a stopping domain. \square

Now we are ready to define \mathcal{G} .

Definition 4.5. \mathcal{G} is defined to be the collections of $(\bar{\ell}, \bar{r}, f_{\mathbb{J}}) \in C_{\text{Jct}}$ such that the following conditions hold.

- C1:** There exists $f_{\mathbb{B}} \in C(\text{Box})$ with $f_{\mathbb{B}}|_{\text{Jct}(\bar{\ell}, \bar{r})} = f_{\mathbb{J}}$ such that $(\bar{\ell}, \bar{r}) = (\bar{\mathbf{l}}, \bar{\mathbf{r}})(f_{\mathbb{B}})$.
- C2:** $|f_j(x) + 2^{-1}x^2| \leq 2^{-2}10^{-2}T^2$ for all $(j, x) \in \text{Jct}(\bar{\ell}, \bar{r})$.
- C3:** $|f_j(x) - f_j(y)| \leq \Delta$ for all $|x - y| \leq d$ and $(j, x), (j, y) \in \text{Jct}(\bar{\ell}, \bar{r})$.
- C4:** $f_j(x) \geq f_{j+1}(x) - 3\Delta$ for all $(j, x), (j+1, x) \in \text{Jct}(\bar{\ell}, \bar{r})$.

Condition **C1** actually holds automatically for the triple $(\bar{\mathbf{l}}, \bar{\mathbf{r}}, \mathfrak{f}|_{\text{Jct}(\bar{\mathbf{l}}, \bar{\mathbf{r}})})$. We include it still as we need to make use of it later. Condition **C2** requires each layer in $f_{\mathbb{J}}$ to be close to the parabola $-2^{-1}x^2$, which further ensures that f_{j+1} is bounded above by a linear function as in Figure 2. See the discussion in Remark 4.1. Conditions **C3** controls the local fluctuation of $f_{\mathbb{J}}$ within distance d . Condition **C4** ensures that layers in $f_{\mathbb{J}}$ may go out of order by an amount of at most 3Δ .

Lemma 4.6. \mathcal{G} is a Borel subset of C_{Jct} .

Proof. Let $\pi^{-1}(\mathcal{G})$ be the pre-image of \mathcal{G} in $U \times C(\text{Box})$. $\pi^{-1}(\mathcal{G})$ is the collection of triples $(\bar{\ell}, \bar{r}, f_{\mathbb{B}})$ that satisfy Conditions **C2-C4** and $(\bar{\ell}, \bar{r}) = (\bar{\mathbf{l}}, \bar{\mathbf{r}})(f_{\mathbb{B}})$. Conditions **C2-C4** induce a closed subset of $U \times C(\text{Box})$. Together with Remark 4.2, $\pi^{-1}(\mathcal{G})$ is a Borel subset of $U \times C(\text{Box})$. Hence \mathcal{G} is a Borel subset of C_{Jct} . \square

Proposition 4.7. There exist $E_1(k, s)$ and $D = D(k, s)$ such that the following statement holds. For all $t \geq 1$ and $E \geq E_1(k, s)$, we have

$$\mathbb{P}(\text{Fav}^c) \leq De^{-D^{-1}T^3}.$$

We are ready to prove Proposition 3.4.

Proof of Proposition 3.4. If $\varepsilon \geq e^{-s^3}$, the assertion holds easily by taking D large enough. Assume $\varepsilon < e^{-s^3}$. In particular, $T = E(\log \varepsilon^{-1})^{1/3}$. By taking E large enough, we obtain from Proposition 4.7 that

$$\mathbb{P}(\text{Fav}^c) \leq D\varepsilon.$$

□

To prove Proposition 4.7, we record [Wu21, Proposition 3.1, Corollary 3.2 and Corollary B.3] here respectively.

Proposition 4.8. *There exists a constant $D = D(k)$ such that the following statement holds. For all $t \geq 1$, $L \geq 1$ and $K \geq L^2$, we have*

$$\mathbb{P} \left(\mathbb{E}_{\text{free}}^{1,k,(-L,L),\vec{x},\vec{y}} [W_{\mathbf{H}_t}^{1,k,(-L,L),\vec{x},\vec{y},+\infty,\mathfrak{H}_{k+1}^t}] < D^{-1} e^{-DL^{-1}K^2} \right) < e^{-K^{3/2}},$$

where $\vec{x} = (\mathfrak{H}_i^t(-L))_{i=1}^k$ and $\vec{y} = (\mathfrak{H}_i^t(L))_{i=1}^k$.

Proposition 4.9. *There exists a constant $D = D(k)$ such that the following statement holds. For all $t \geq 1$, $L \geq 1$, $d_0 \in (0, 1]$ and $K \geq 4Ld_0^{1/2}$, we have*

$$\mathbb{P} \left(\sup_{x,y \in [-L,L], |x-y| \leq d_0} |\mathfrak{H}_k^t(x) - \mathfrak{H}_k^t(y)| \geq Kd_0^{1/2} \right) \leq d_0^{-1} L D e^{-D^{-1}K^{3/2}}.$$

Proposition 4.10. *There exists a constant $D = D(k)$ such that the following statement holds. For all $t \geq 1$, intervals $I \subset \mathbb{R}$ and $K \geq 0$, we have*

$$\mathbb{P} \left(\sup_{x \in I} |\mathfrak{H}_k^t(x) + 2^{-1}x^2| \geq K \right) \leq (|I|/2 + 1) D e^{-D^{-1}K^{3/2}}.$$

Proposition 4.8 bounds the normalizing constant for \mathfrak{H} and is used to obtain Condition **C4**. Proposition 4.9 bounds the local fluctuation of \mathfrak{H} and is used to ensure Condition **C3**. Proposition 4.10 implies \mathfrak{H} are close to a parabola and guarantees Condition **C2**.

We would like to briefly explain how the parameters are chosen. The numbers Δ and d are chosen to ensure that

$$(4.5) \quad \exp(-d \mathbf{H}_t(\Delta)) \ll e^{-T^3}$$

and that

$$(4.6) \quad d^{-1/2} \Delta = T^2.$$

(4.5) implies that it is costly for \mathfrak{H}_j^t to always stay below $\mathfrak{H}_{j+1}^t - \Delta$ for an interval of length d . (4.6) controls the local fluctuation of \mathfrak{H}_j^t within distance d . More precisely, its modulus of continuity over an interval of length d is bounded above by Δ , with probability $1 - e^{-T^3}$, see Proposition 4.9. We will employ (4.5) and (4.6) to show that $\mathfrak{H}_j^t \geq \mathfrak{H}_{j+1}^t - 3\Delta$ with high probability.

Proof of Proposition 4.7. Let $\mathcal{G}_2, \mathcal{G}_3$ and \mathcal{G}_4 be the subsets of $C(\text{Box})$ such that **C2**, **C3** and **C4** hold in Box respectively. To be explicit,

$$\begin{aligned} \mathcal{G}_2 &:= \{f_{\text{B}} \in C(\text{Box}) \mid |f_j(x) + 2^{-1}x^2| \leq 2^{-2}10^{-2}T^2 \text{ for all } (j, x) \in \text{Box}\}, \\ \mathcal{G}_3 &:= \{f_{\text{B}} \in C(\text{Box}) \mid |f_j(x) - f_j(y)| \leq \Delta \text{ for all } (j, x), (j, y) \in \text{Box} \text{ with } |x - y| \leq d\}, \\ \mathcal{G}_4 &:= \{f_{\text{B}} \in C(\text{Box}) \mid f_j(x) \geq f_{j+1}(x) - 3\Delta \text{ for all } (j, x), (j+1, x) \in \text{Box}\}. \end{aligned}$$

For $i = 2, 3, 4$, denote by Fav_i the event $\{\mathfrak{H}|_{\text{Box}} \in \mathcal{G}_i\}$. From the definition of (\bar{l}, \bar{r}) , condition **C1** holds automatically. Therefore

$$\bigcap_{i=2}^4 \text{Fav}_i \subset \text{Fav}.$$

In the following, we bound $\mathbb{P}(\text{Fav}_i^c)$.

To bound $\mathbb{P}(\text{Fav}_2^c)$, we apply Proposition 4.10. Set $I = [\ell_0, r_0]$ and $K = 2^{-2}10^{-2}T^2$, we deduce that for all $j \in [1, k+1]_{\mathbb{Z}}$,

$$\mathbb{P}\left(\sup_{x \in [\ell_0, r_0]} |\mathfrak{H}_j^t + 2^{-1}x^2| > 2^{-2}10^{-2}T^2\right) \leq ((r_0 - \ell_0)/2 + 1)De^{-D^{-1}2^{-3}10^{-3}T^2}.$$

From $r_0 - \ell_0 = 2(k+1)T$ and $T \geq 10$, we deduce that

$$\mathbb{P}(\text{Fav}_2^c) \leq De^{-D^{-1}T^3}$$

To bound $\mathbb{P}(\text{Fav}_3^c)$, we apply Proposition 4.9. Set $L = (k+1)T$, $K = T^2$ and $d_0 = d$. We verify that $K \geq 4Ld_0^{1/2}$ as long as T is large enough. We deduce that for all $j \in [1, k+1]_{\mathbb{Z}}$,

$$\mathbb{P}\left(\sup_{x, y \in [\ell_0, r_0], |x-y| \leq d} |\mathfrak{H}_j^t(x) - \mathfrak{H}_j^t(y)| \geq \Delta\right) \leq d_0^{-1}(2k+2)TDe^{-D^{-1}T^3}$$

Together with $T \geq 1$,

$$\mathbb{P}(\text{Fav}_3^c) \leq De^{-D^{-1}T^3}.$$

Lastly, we bound $\mathbb{P}(\text{Fav}_4^c)$. Let $\bar{\ell}^* = (\ell_0, \ell_0, \dots, \ell_0)$ and $\bar{r}^* = (r_0, r_0, \dots, r_0)$. Apply Proposition 4.8. Setting $L = (k+1)T$ and $K = (k+1)^2T^2$, we have

$$\mathbb{P}\left(\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}^*, \bar{r}^*), f^*} \left[W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}^*, \bar{r}^*), f^*} \right] < D^{-1}e^{-D(k+1)^3T^3}\right) \leq e^{-(k+1)^3T^3}.$$

Here $f^* = \mathfrak{H}^t|_{\text{bdd}(\bar{\ell}^*, \bar{r}^*)}$. Denote this event by Z^c . Note that $\text{Fav}_3 \cap \text{Fav}_4^c$ implies there exists an interval with length d in which $\mathfrak{H}_j < \mathfrak{H}_{j+1} - \Delta$ for some $j \in [1, k]_{\mathbb{Z}}$. Because $d\mathbf{H}_t(\Delta) \geq 64T^4(\log T)^2$, we have

$$\text{Fav}_3 \cap \text{Fav}_4^c \subset \left\{ W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}^*, \bar{r}^*), f^*}(\mathfrak{H}^t|_{\text{dom}(\bar{\ell}^*, \bar{r}^*)}) \leq e^{-64T^4(\log T)^2} \right\}.$$

We then deduce from the \mathbf{H}_t -Brownian Gibbs property that

$$\begin{aligned} \mathbb{P}(\text{Fav}_3 \cap \text{Fav}_4^c) &\leq \mathbb{P}(\text{Fav}_3 \cap \text{Fav}_4^c \cap Z) + \mathbb{P}(Z^c) \\ &= \mathbb{E}[\mathbb{1}_Z \cdot \mathbb{E}[\mathbb{1}_{\text{Fav}_3} \cdot \mathbb{1}_{\text{Fav}_4^c} | \mathcal{F}_{\text{ext}}(\bar{\ell}^*, \bar{r}^*)]] + \mathbb{P}(Z^c) \\ &\leq De^{-64T^4(\log T)^2 + D(k+1)^3T^3} + e^{-(k+1)^3T^3} \\ &\leq De^{-D^{-1}T^3}. \end{aligned}$$

Therefore,

$$\mathbb{P}(\text{Fav}_4^c) \leq \mathbb{P}(\text{Fav}_3 \cap \text{Fav}_4^c) + \mathbb{P}(\text{Fav}_3^c) \leq De^{-D^{-1}T^3}.$$

In conclusion,

$$\mathbb{P}(\text{Fav}^c) \leq \sum_{i=2}^4 \mathbb{P}(\text{Fav}_i^c) \leq De^{-D^{-1}T^3}.$$

□

5. CURVE SEPARATION OVER A FINITE SET

The goal of this section is to prove Theorem 5.1. Theorem 5.1 shows that, in loose terms, conditioned on jumping over a Lipschitz function at finite many locations, the Brownian bridges may remain separated provided they are separated on the boundary. Theorem 5.1 will be used in Section 7 to separate the curves in the jump ensemble J over the resampling domain. The details in the proof will not be needed in the remaining part of this paper. Skipping those will not affect the reader's ability to follow the rest of the arguments.

Fix $k \in \mathbb{N}$ and $\ell < r$. Let $a^+ = (a_1^+, a_2^+, \dots, a_k^+)$ and $a^- = (a_1^-, a_2^-, \dots, a_k^-)$ be two vectors in \mathbb{R}^k . Recall that $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$ is the law of k independent Brownian bridges defined on $[\ell, r]$ with boundary values given by a^\pm . Let $g \in C([\ell, r])$ be a Lipschitz function with $g(\ell) = g(r) = 0$. $g(x)$ will serve as the lower boundary. Let $P \subset (\ell, r)$ be a finite subset. Let $\mathbb{P}_{\mathcal{L}}$ be the law of $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$ conditioned on jumping over $g(x)$ at pole points in P . In terms of the Radon-Nikodym derivative relation, we have

$$\frac{d\mathbb{P}_{\mathcal{L}}}{d\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}}(\mathcal{L}) := \frac{W(\mathcal{L})}{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[W]}.$$

The Boltzmann weight W is defined through

$$W(\mathcal{L}) := \begin{cases} 1 & \mathcal{L}_j(p) > g(p) \text{ for all } p \in P \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0 & \text{otherwise.} \end{cases}$$

Next, we introduce two assumptions (5.3) and (5.4). We assume (5.3) and (5.4) hold throughout this section. Fix $T \geq 10$,

$$(5.1) \quad b_0, b_1, \lambda_0, \lambda_1 > 0,$$

and

$$(5.2) \quad b_2 > 0.$$

We assume that

$$(5.3) \quad \begin{aligned} r - \ell &\leq b_0 T, \quad |P| \leq b_0 T, \quad \sup_{x, y \in [\ell, r], x \neq y} \frac{|g(x) - g(y)|}{|x - y|} \leq b_1 T, \\ a_j^\pm - a_{j+1}^\pm &\geq \lambda_0 T^{1/2} \text{ for all } j \in [1, k-1]_{\mathbb{Z}} \\ a_k^- - g(\ell) &\geq \lambda_1 T \text{ and } a_k^+ - g(r) \geq \lambda_1 T. \end{aligned}$$

We also assume that

$$(5.4) \quad a_1^- - g(\ell) \leq b_2 T^2 \text{ and } a_1^+ - g(r) \leq b_2 T^2.$$

The assumption (5.3) bounds the distant between ℓ and r , the number of elements in P , the Lipschitz constant of g and the gaps in a^\pm . The assumption (5.4) further gives an upper bound for a_1^\pm .

For $\mu \in (0, 1)$, Consider the event

$$\begin{aligned} \mathbf{H} := & \{ \mathcal{L}_j(p) - \mathcal{L}_{j+1}(p) \geq \mu \lambda_0 T^{1/2} \text{ for all } p \in P \text{ and } j \in [1, k-1]_{\mathbb{Z}} \} \\ & \cap \{ \mathcal{L}_k(p) - g(p) \geq \mu \lambda_1 T \text{ for all } p \in P \}. \end{aligned}$$

Theorem 5.1. *There exists a constant C depending on constants in (5.1), (5.2) and μ such that*

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) \geq C^{-1} e^{-CT^{5/2}}.$$

The rest of the section is devoted to prove Theorem 5.1. In subsection 5.1, we first prove Theorem 5.1 for a special case. The general case is proved in subsection 5.2 up to Lemma 5.7. Lemma 5.7 is then proved in subsection 5.3.

5.1. A special case. In this subsection, we prove Theorem 5.1 for the special case

$$P \subset (\ell, \ell + T^{1/2}) \cup (r - T^{1/2}, r).$$

We begin by considering the situation that $P \subset (\ell, \ell + T^{1/2})$.

Lemma 5.2. *Assume that $P \subset (\ell, \ell + T^{1/2})$. Then there exists a constant C depending on constants in (5.1) and μ such that*

$$\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

From Lemma 5.2, we immediately get the following two corollaries. Note that by the time reversal symmetry of the Brownian motion, the same results in Corollary 5.3 and Corollary 5.4 also hold if $P \subset (r - T^{1/2}, r)$.

Corollary 5.3. *Assume that $P \subset (\ell, \ell + T^{1/2})$. Then there exists a constant C depending on constants in (5.1) and μ such that*

$$\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[W] \geq C^{-1}e^{-CT^{5/2}}.$$

Proof. This follows easily from Lemma 5.2 because $\mathbf{H} \subset \{W = 1\}$. \square

Corollary 5.4. *Assume that $P \subset (\ell, \ell + T^{1/2})$. Then there exists a constant C depending on constants in (5.1) and μ such that*

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

Proof. By $W \leq 1$ and $\mathbf{H} \subset \{W = 1\}$, we have

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) = \frac{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[\mathbb{1}_{\mathbf{H}} \cdot W]}{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[W]} = \frac{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[\mathbb{1}_{\mathbf{H}}]}{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[W]} \geq \mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}(\mathbf{H}).$$

Then the assertion follows from Lemma 5.2. \square

Proof of Lemma 5.2. Note that without loss of generality, we may assume $g(\ell) = g(r) = 0$. The reason is that the law of Brownian bridges are invariant under affine transformations. The event \mathbf{H} is also unaffected by affine transformations. The only thing differs is the Lipschitz bound of $g(x)$. When we replace $g(x)$ by $g(x) - \frac{x-\ell}{r-\ell}g(r) - \frac{r-x}{r-\ell}g(\ell)$, the Lipschitz bound of $g(x)$ may be doubled. Therefore, it is sufficient to prove the assertion for the case $g(\ell) = g(r) = 0$. From now on we assume $g(\ell) = g(r) = 0$.

The proof involves direct computations of Brownian bridges. We separate the discussion into two cases depending on the length of the interval $[\ell, r]$.

Case 1: $r - \ell \geq 2T^{1/2}$. Let $q = \ell + T^{1/2}$. Under the law $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$, $B_j(q)$ are Gaussian random variables with mean and variance

$$m_j = \frac{(r-q)a_j^- + (q-\ell)a_j^+}{r-\ell}, \quad \sigma^2 = \frac{(q-\ell)(r-q)}{r-\ell}.$$

For $x \in [\ell, q]$, we have

$$B_j(x) = \frac{(q-x)a_j^- + (x-\ell)B_j(q)}{q-\ell} + B_j^{[\ell,q]}(x).$$

In particular,

$$\begin{aligned}
B_k(x) - g(x) &\geq \left(\frac{(q-x)a_k^- + (x-\ell)B_k(q)}{q-\ell} + B_k^{[\ell,q]}(x) \right) - (x-\ell)b_1T \\
&= \frac{(q-x)a_k^- + (x-\ell)m_k}{q-\ell} + \frac{(x-\ell)(B_k(q) - m_k - b_1T^{3/2})}{q-\ell} + B_k^{[\ell,q]}(x) \\
&\geq \lambda_1T - |B_k(q) - (m_k + b_1T^{3/2})| - \sup_{x \in [\ell,q]} |B_k^{[\ell,q]}(x)|.
\end{aligned}$$

We have used the assumption (5.3) in the above inequalities.

Similarly, for $x \in [\ell, q]$ and $j \in [1, k-1]_{\mathbb{Z}}$, we have

$$\begin{aligned}
B_j(x) - B_{j+1}(x) &\geq \lambda_0T^{1/2} - |B_j(q) - (m_j + b_1T^{3/2})| - \sup_{x \in [\ell,q]} |B_j^{[\ell,q]}(x)| \\
&\quad - |B_{j+1}(q) - (m_{j+1} + b_1T^{3/2})| - \sup_{x \in [\ell,q]} |B_{j+1}^{[\ell,q]}(x)|.
\end{aligned}$$

We have used the assumption (5.3) in the above inequality.

In order to ensure \mathbf{H} , it is sufficient to require that for all $j \in [1, k]_{\mathbb{Z}}$, it holds that

$$|B_j(q) - (m_j + b_1T^{3/2})| \leq 4^{-1}(1-\mu)\lambda T^{1/2}, \quad \sup_{x \in [\ell,q]} |B_j^{[\ell,q]}(x)| \leq 4^{-1}(1-\mu)\lambda T^{1/2},$$

where $\lambda = \min\{\lambda_0, \lambda_1\}$. Let N be a Gaussian random variable with mean 0 and variance 1. It holds that

$$\begin{aligned}
&\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} \left(|B_j(q) - (m_j + b_1T^{3/2})| \leq 4^{-1}(1-\mu)\lambda T^{1/2} \right) \\
&= \mathbb{P} \left(|N - \sigma^{-1}b_1T^{3/2}| \leq 4^{-1}\sigma^{-1}(1-\mu)\lambda T^{1/2} \right).
\end{aligned}$$

Since $r - \ell \geq 2T^{1/2}$ and $q = \ell + T^{1/2}$, $\sigma^2 \in [2^{-1}T^{1/2}, T^{1/2})$. It holds that

$$\sigma^{-1}b_1T^{3/2} \leq 2^{1/2}b_1T^{5/4},$$

and

$$4^{-1}\sigma^{-1}(1-\mu)\lambda T^{1/2} \geq 4^{-1}(1-\mu)\lambda T^{1/4}.$$

As a result,

$$\begin{aligned}
&\mathbb{P}(|N - \sigma^{-1}b_1T^{3/2}| \leq 4^{-1}\sigma^{-1}(1-\mu)\lambda T^{1/2}) \\
&\geq \mathbb{P}(|N - 2^{1/2}b_1T^{5/4}| \leq 4^{-1}(1-\mu)\lambda T^{1/4}) \\
&\geq C^{-1}e^{-CT^{5/2}}.
\end{aligned}$$

Next, we bound the event $\sup_{x \in [\ell,q]} |B_j^{[\ell,q]}(x)| \leq 4^{-1}(1-\mu)\lambda T^{1/2}$. Recall that $q - \ell = T^{1/2}$. Through a direct computation,

$$\begin{aligned}
&\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} \left(\sup_{x \in [\ell,q]} |B_j^{[\ell,q]}(x)| \leq 4^{-1}(1-\mu)\lambda T^{1/2} \right) \\
&= \mathbb{P} \left(\sup_{x \in [0,1]} |B^{[0,1]}(x)| \leq 4^{-1}(1-\mu)\lambda T^{1/4} \right) \geq C^{-1}.
\end{aligned}$$

Combining the above estimates, we conclude that

$$\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

Case 2: $r - \ell < 2T^{1/2}$. Set $q = 2^{-1}(\ell + r)$. Under the law $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$, $B_j(q)$ are Gaussian random variables with mean and variance

$$m_j = 2^{-1}(a_j^- + a_j^+), \quad \sigma^2 = 4^{-1}(r - \ell).$$

For $x \in [\ell, q]$ and $y \in [q, r]$, it holds that

$$\begin{aligned} B_k(x) - g(x) &\geq \lambda_1 T - |B_k(q) - (m_k + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_k^{[\ell, q]}(x)|, \\ B_k(y) - g(y) &\geq \lambda_1 T - |B_k(q) - (m_k + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_k^{[q, r]}(y)|. \end{aligned}$$

We have used the assumption (5.3) in the above inequality.

Similarly, for $x \in [\ell, q]$, $y \in [q, r]$ and $j \in [1, k - 1]_{\mathbb{Z}}$, it holds that

$$\begin{aligned} B_j(x) - B_{j+1}(x) &\geq \lambda_0 T^{1/2} - |B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \\ &\quad - |B_{j+1}(q) - (m_{j+1} + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_{j+1}^{[\ell, q]}(x)|, \\ B_j(y) - B_{j+1}(y) &\geq \lambda_0 T^{1/2} - |B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_j^{[q, r]}(y)| \\ &\quad - |B_{j+1}(q) - (m_{j+1} + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_{j+1}^{[q, r]}(y)|. \end{aligned}$$

We have used the assumption (5.3) in the above inequality.

In order to ensure \mathbf{H} , it is sufficient to require that for $j \in [1, k]_{\mathbb{Z}}$, it holds that

$$\begin{aligned} |B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| &\leq 4^{-1}(1 - \mu)\lambda T^{1/2}, \\ \sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| &\leq 4^{-1}(1 - \mu)\lambda T^{1/2}, \end{aligned}$$

and

$$\sup_{x \in [q, r]} |B_j^{[q, r]}(x)| \leq 4^{-1}(1 - \mu)\lambda T^{1/2},$$

where $\lambda = \min\{\lambda_0, \lambda_1\}$. Let N be a Gaussian random variable with mean 0 and variance 1. It holds that

$$\begin{aligned} \mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} (|B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| \leq 4^{-1}(1 - \mu)\lambda T^{1/2}) \\ = \mathbb{P}(|N - (r - \ell)^{1/2}b_1 T| \leq 2^{-1}(1 - \mu)\lambda(r - \ell)^{-1/2}T^{1/2}). \end{aligned}$$

Because $r - \ell < 2T^{1/2}$, it holds that

$$(r - \ell)^{1/2}b_1 T < 2^{1/2}b_1 T^{5/4}$$

and

$$2^{-1}(1 - \mu)\lambda(r - \ell)^{-1/2}T^{1/2} > 2^{-3/2}(1 - \mu)\lambda T^{1/4}.$$

Therefore,

$$\begin{aligned} \mathbb{P}(|N - (r - \ell)^{1/2}b_1 T| \leq 2^{-1}(1 - \mu)\lambda(r - \ell)^{-1/2}T^{1/2}) \\ \geq \mathbb{P}(|N - 2^{1/2}b_1 T^{5/4}| \leq 2^{-3/2}(1 - \mu)\lambda T^{1/4}) \\ \geq C^{-1}e^{-CT^{5/2}}. \end{aligned}$$

Next, we bound the event $\sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \leq 4^{-1}(1 - \mu)\lambda T^{1/2}$. By a direct computation,

$$\begin{aligned} & \mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} \left(\sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \leq 4^{-1}(1 - \mu)\lambda T^{1/2} \right) \\ &= \mathbb{P} \left(\sup_{x \in [0, 1]} |B^{[0, 1]}(x)| \leq 2^{-3/2}(r - \ell)^{-1/2}(1 - \mu)\lambda T^{1/2} \right) \\ &\geq \mathbb{P} \left(\sup_{x \in [0, 1]} |B^{[0, 1]}(x)| \leq 4^{-1}(1 - \mu)\lambda T^{1/4} \right) \geq C^{-1}. \end{aligned}$$

We have used $r - \ell < 2T^{1/2}$. Similarly,

$$\mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} \left(\sup_{x \in [q, r]} |B_j^{[q, r]}(x)| \leq 2^{-2}(1 - \mu)\lambda T^{1/2} \right) \geq C^{-1}.$$

Combining the above estimates, we conclude

$$\mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

□

We are ready to prove Theorem 5.1 for a special case $P \subset (\ell, \ell + T^{1/2}) \cup (r - T^{1/2}, r)$.

Lemma 5.5. *Assume that $P \subset (\ell, \ell + T^{1/2}) \cup (r - T^{1/2}, r)$. Then there exists a constant C depending on constants in (5.1) and μ such that*

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

Proof. The idea is to decompose the Boltzmann weight into $W = \widetilde{W} \cdot \widehat{W}$. Roughly speaking, \widetilde{W} and \widehat{W} activate the interaction at the pole points in $(r - T^{1/2}, r)$ and $(\ell, \ell + T^{1/2})$ respectively. Then we apply Corollary 5.4 twice to finish the argument.

Let $\tilde{P} = P \cap (r - T^{1/2}, r)$ and $\hat{P} = P \setminus \tilde{P}$. If \tilde{P} is empty, the assertion of Lemma 5.5 follows from Corollary 5.4. From now on we assume \tilde{P} is non-empty.

Let $\mathbb{P}_{\tilde{\mathcal{L}}}$ be the law of $\mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-}$ conditioned on jumping over $g(x)$ at pole points in \tilde{P} . In terms of the Radon-Nikodym derivative relation, it holds that

$$\frac{d\mathbb{P}_{\tilde{\mathcal{L}}}}{d\mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-}}(\mathcal{L}) = \frac{\widetilde{W}(\mathcal{L})}{\mathbb{E}_{\text{free}}^{1, k, (\ell, r), a^+, a^-}[\widetilde{W}]}$$

Here the weight \widetilde{W} is defined by

$$\widetilde{W}(\mathcal{L}) := \begin{cases} 1, & \mathcal{L}_j(p) > g(p) \text{ for all } p \in \tilde{P} \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0, & \text{otherwise.} \end{cases}$$

Correspondingly, we define the remaining weight

$$\widehat{W}(\mathcal{L}) := \begin{cases} 1, & \mathcal{L}_j(p) > g(p) \text{ for all } p \in \hat{P} \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0, & \text{otherwise.} \end{cases}$$

Since $W = \widetilde{W} \cdot \widehat{W}$, we have

$$\frac{d\mathbb{P}_{\mathcal{L}}}{d\mathbb{P}_{\tilde{\mathcal{L}}}}(\mathcal{L}) = \frac{\widehat{W}(\mathcal{L})}{\mathbb{E}_{\tilde{\mathcal{L}}}[\widehat{W}]}$$

Consider the following events $\tilde{\mathbf{G}}$ and $\hat{\mathbf{G}}$,

$$\begin{aligned} \tilde{\mathbf{G}} &:= \{\mathcal{L}_j(p) - \mathcal{L}_{j+1}(p) \geq 2^{-1}(1 + \mu)\lambda_0 T^{1/2} \text{ for all } p \in \tilde{P} \text{ and } j \in [1, k - 1]_{\mathbb{Z}}\} \\ &\cap \{\mathcal{L}_k(p) - g(p) \geq 2^{-1}(1 + \mu)\lambda_1 T \text{ for all } p \in \tilde{P}\}, \end{aligned}$$

and

$$\begin{aligned} \hat{\mathbf{G}} := & \{\mathcal{L}_j(p) - \mathcal{L}_{j+1}(p) \geq \mu\lambda_0 T^{1/2} \text{ for all } p \in \hat{P} \text{ and } j \in [1, k-1]_{\mathbb{Z}}\} \\ & \cap \{\mathcal{L}_k(p) - g(p) \geq \mu\lambda_1 T \text{ for all } p \in \hat{P}\}. \end{aligned}$$

Roughly speaking, $\tilde{\mathbf{G}}$ and $\hat{\mathbf{G}}$ require conditions in \mathbf{H} to hold in \tilde{P} and \hat{P} respectively. The requirements in $\tilde{\mathbf{G}}$ is stronger than the ones in \mathbf{H} . The reason is that we will apply Corollary 5.4 and Corollary 5.3 to \tilde{P} conditioned on the occurrence of $\tilde{\mathbf{G}}$.

Because $\hat{W} \leq 1$, we have

$$\begin{aligned} \mathbb{P}_{\mathcal{L}}(\mathbf{H}) & \geq \mathbb{P}_{\mathcal{L}}(\mathbf{H} \cap \tilde{\mathbf{G}}) \\ & = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}) \times \mathbb{P}_{\mathcal{L}}(\tilde{\mathbf{G}}) = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}) \times \frac{\mathbb{E}_{\tilde{\mathcal{L}}}[\mathbb{1}_{\tilde{\mathbf{G}}} \cdot \hat{W}]}{\mathbb{E}_{\tilde{\mathcal{L}}}[\hat{W}]} \\ & \geq \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}) \times \mathbb{E}_{\tilde{\mathcal{L}}}[\mathbb{1}_{\tilde{\mathbf{G}}} \cdot \hat{W}] \\ & = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}) \times \mathbb{P}_{\tilde{\mathcal{L}}}(\tilde{\mathbf{G}}) \times \mathbb{E}_{\tilde{\mathcal{L}}}[\hat{W}|\tilde{\mathbf{G}}]. \end{aligned}$$

Since $\tilde{P} \subset (r - T^{1/2}, r)$, we can apply Corollary 5.4 to obtain

$$\mathbb{P}_{\tilde{\mathcal{L}}}(\tilde{\mathbf{G}}) \geq C^{-1} e^{-CT^{5/2}}.$$

Next, we turn to $\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}})$. Conditioned on a realization of $\tilde{\mathbf{G}}$, the law of $\mathbb{P}_{\mathcal{L}}$ restricted to $[\ell, r - T^{1/2}]$ is given by the law of free Brownian bridges conditioned on jumping over g at those pole points in \tilde{P} . Therefore, we can apply Corollary 5.4 to bound $\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}})$.

We proceed to establish estimates that are uniform in any realizations of $\tilde{\mathbf{G}}$. To simplify the notation, we fix a realization of $\tilde{\mathbf{G}}$. That is, we fix $\mathcal{L}_j(p)$ for $j \in [1, k]_{\mathbb{Z}}$ and $p \in \tilde{P}$ which satisfy the requirements in $\tilde{\mathbf{G}}$. We denote the event of such realization by $\tilde{\mathbf{G}}_0$. Because $\hat{\mathbf{G}} \cap \tilde{\mathbf{G}}_0 \subset \mathbf{H}$, it holds that $\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}_0) \geq \mathbb{P}_{\mathcal{L}}(\hat{\mathbf{G}}|\tilde{\mathbf{G}}_0)$.

In order to apply Corollary 5.4, we define the following quantities. Let p_0 be the smallest element of \tilde{P} . Define

$$\hat{\ell} := \ell, \hat{r} := p_0, \hat{a}_j^- := a_j^-, \hat{a}_j^+ := \mathcal{L}_j(p_0).$$

Conditioned on $\tilde{\mathbf{G}}_0$, the law of $\mathcal{L}|_{[\hat{\ell}, \hat{r}]}$, denoted by $\hat{\mathbb{P}}_{\mathcal{L}}$, is the law of $\mathbb{P}_{\text{free}}^{1,k,(\hat{\ell}, \hat{r}), \hat{a}^+, \hat{a}^-}$ conditioned on jumping over $g(x)$ at pole points in \tilde{P} . In terms of the Radon-Nikodym derivative relation, it holds that

$$\frac{d\hat{\mathbb{P}}_{\mathcal{L}}}{d\mathbb{P}_{\text{free}}^{1,k,(\hat{\ell}, \hat{r}), \hat{a}^+, \hat{a}^-}}(\mathcal{L}) = \frac{\hat{W}(\mathcal{L})}{\mathbb{E}_{\text{free}}^{1,k,(\hat{\ell}, \hat{r}), \hat{a}^+, \hat{a}^-}[\hat{W}]}.$$

To apply Corollary 5.4, we check the assumption (5.3). $\tilde{\mathbf{G}}_0$ implies that

$$\begin{aligned} \hat{a}_j^\pm - \hat{a}_{j+1}^\pm & \geq 2^{-1}(1 + \mu)\lambda_0 T^{1/2} \text{ for all } j \in [1, k-1]_{\mathbb{Z}}, \\ \hat{a}_k^- - g(\hat{\ell}) & \geq 2^{-1}(1 + \mu)\lambda_1 T, \text{ and } \hat{a}_k^+ - g(\hat{r}) \geq 2^{-1}(1 + \mu)\lambda_1 T. \end{aligned}$$

Therefore, the assumption (5.3) holds for these data with slightly different bounds. By Corollary 5.4, it holds that $\mathbb{P}_{\mathcal{L}}(\hat{\mathbf{G}}|\tilde{\mathbf{G}}_0) \geq C^{-1} e^{-CT^{5/2}}$. This ensures

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}_0) \geq \mathbb{P}_{\mathcal{L}}(\hat{\mathbf{G}}|\tilde{\mathbf{G}}_0) \geq C^{-1} e^{-CT^{5/2}}.$$

Because this holds for any realization $\tilde{\mathbf{G}}_0$ of $\tilde{\mathbf{G}}$, we obtain

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\tilde{\mathbf{G}}) \geq C^{-1} e^{-CT^{5/2}}.$$

Next, we turn to $\mathbb{E}_{\tilde{\mathcal{L}}}[\hat{W}|\mathbf{G}_0]$. Conditioned on $\tilde{\mathbf{G}}_0$, The law of $\tilde{\mathcal{L}}|_{[\hat{\ell}, \hat{r}]}$ is identical to $\mathbb{P}_{\text{free}}^{1,k,(\hat{\ell}, \hat{r}), \hat{a}^+, \hat{a}^-}$. The assumption (5.3) is already checked above. Applying Corollary 5.3, it holds that

$$\mathbb{E}_{\tilde{\mathcal{L}}}[\hat{W}|\mathbf{G}_0] = \mathbb{E}_{\text{free}}^{1,k,(\hat{\ell}, \hat{r}), \hat{a}^+, \hat{a}^-}[\hat{W}] \geq C^{-1}e^{-CT^{5/2}}.$$

Because this holds for any realization $\tilde{\mathbf{G}}_0$ of $\tilde{\mathbf{G}}$, we obtain

$$\mathbb{E}_{\tilde{\mathcal{L}}}[\hat{W}|\tilde{\mathbf{G}}] \geq C^{-1}e^{-CT^{5/2}}.$$

Putting the above estimates together, we conclude that

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

□

5.2. The general case. In this subsection, we consider the general case of P and prove Theorem 5.1. The approach is similar to the one in proving Lemma 5.5. We decompose the Boltzmann weight into $W = W' \cdot \check{W}$. The weights W' and \check{W} activate the interaction at the pole points in $[\ell + T^{1/2}, r - T^{1/2}]$ and $(\ell, \ell + T^{1/2}) \cup (r - T^{1/2}, r)$ respectively. Lemma 5.5 is then applicable to \check{W} .

Let $P' = P \cap [\ell + T^{1/2}, r - T^{1/2}]$ and $\check{P} = P \setminus P'$. Suppose P' is empty, then Theorem 5.1 follows from Lemma 5.5. From now on we assume P' is non-empty.

Let $\mathbb{P}'_{\mathcal{L}}$ be the law of $\mathbb{P}_{\text{free}}^{1,k,(\ell, r), a^+, a^-}$ conditioned on jumping over $g(x)$ at pole points in P' . In terms of the Radon-Nikodym derivative relation, it holds that

$$\frac{d\mathbb{P}'_{\mathcal{L}}}{d\mathbb{P}_{\text{free}}^{1,k,(\ell, r), a^+, a^-}}(\mathcal{L}) = \frac{W'(\mathcal{L})}{\mathbb{E}_{\text{free}}^{1,k,(\ell, r), a^+, a^-}[W']}.$$

Here the weight W' is defined by

$$W'(\mathcal{L}) := \begin{cases} 1, & \mathcal{L}_j(p) > g(p) \text{ for all } p \in P' \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0, & \text{otherwise} \end{cases}$$

Correspondingly, we define

$$\check{W}(\mathcal{L}) := \begin{cases} 1, & \mathcal{L}_j(p) > g(p) \text{ for all } p \in \check{P} \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0, & \text{otherwise} \end{cases}$$

Since $W' \cdot \check{W} = W$, we have

$$\frac{d\mathbb{P}_{\mathcal{L}}}{d\mathbb{P}'_{\mathcal{L}}}(\mathcal{L}) = \frac{\check{W}(\mathcal{L})}{\mathbb{E}'_{\mathcal{L}}[\check{W}]}.$$

We further define \mathcal{L}'_+ to be ordered at pole points in P' . Let $\mathbb{P}'_{\mathcal{L}'_+}$ be the law of $\mathbb{P}'_{\mathcal{L}'_+}$ conditioned being ordered at pole points in P' . In terms of the Radon-Nikodym derivative relation, it holds that

$$\frac{d\mathbb{P}'_{\mathcal{L}'_+}}{d\mathbb{P}_{\text{free}}^{1,k,(\ell, r), a^+, a^-}}(\mathcal{L}) = \frac{W'_+(\mathcal{L})}{\mathbb{E}_{\text{free}}^{1,k,(\ell, r), a^+, a^-}[W'_+]}$$

Here the weight W'_+ is defined by

$$W'_+(\mathcal{L}) := \begin{cases} 1, & \mathcal{L}_k(p) > g(p), \mathcal{L}_{j+1}(p) > \mathcal{L}_j(p) \text{ for all } p \in P' \text{ and } j \in [1, k]_{\mathbb{Z}}, \\ 0, & \text{otherwise} \end{cases}$$

Since $W'_+ = W' \cdot W'_+$, we have

$$\frac{d\mathbb{P}'_{\mathcal{L}'_+}}{d\mathbb{P}'_{\mathcal{L}'}}(\mathcal{L}) = \frac{W'_+(\mathcal{L})}{\mathbb{E}'_{\mathcal{L}'_+}[W'_+]}$$

Lemma 5.6. *There exists a constant C depending on constants in (5.1) such that*

$$\mathbb{E}_{\mathcal{L}'}[W'_+] \geq C^{-1}e^{-CT}.$$

Proof. If $a_j^\pm = a_{j+1}^\pm$ for $j \in [1, k-1]_{\mathbb{Z}}$, $\{\mathcal{L}'_j(p), j \in [1, k]_{\mathbb{Z}}\}$ are identically distributed and it follows that

$$\mathbb{P}_{\mathcal{L}'}(\mathcal{L}'_j(p) > \mathcal{L}'_{j+1}(p) \text{ for } j \in [1, k-1]_{\mathbb{Z}}) = \frac{1}{k!}.$$

Since $a_j^\pm > a_{j+1}^\pm$ for $j \in [1, k-1]_{\mathbb{Z}}$, we have for any $p \in P'$,

$$\mathbb{P}_{\mathcal{L}'}(\mathcal{L}'_j(p) > \mathcal{L}'_{j+1}(p) \text{ for } j \in [1, k-1]_{\mathbb{Z}}) \geq \frac{1}{k!}.$$

Because $|P'| \leq b_0T$, we inductively get

$$\mathbb{E}_{\mathcal{L}'}[W'_+] \geq (k!)^{-b_0T}.$$

□

Consider the event \mathbf{G} ,

$$\begin{aligned} \mathbf{G} := & \{\mathcal{L}_j(p) - \mathcal{L}_{j+1}(p) \geq 2^{-1}(1+\mu)\lambda_0T^{1/2} \text{ for all } p \in P' \text{ and } j \in [1, k-1]_{\mathbb{Z}}\} \\ & \cap \{\mathcal{L}_k(p) - g(p) \geq 2^{-1}(1+\mu)\lambda_1T \text{ for all } p \in P'\}. \end{aligned}$$

Compared to \mathbf{H} , the event \mathbf{G} requires a stronger gap at pole points in P' .

Lemma 5.7. *There exists a constant C depending on constants in (5.1), (5.2) and μ such that*

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}) \geq C^{-1}e^{-CT^{5/2}}.$$

We postpone the proof of Lemma 5.7 and prove Theorem 5.1 below.

Proof of Theorem 5.1. Because $\check{W} \leq 1$, we have

$$\begin{aligned} \mathbb{P}_{\mathcal{L}}(\mathbf{H}) & \geq \mathbb{P}_{\mathcal{L}}(\mathbf{H} \cap \mathbf{G}) \\ & = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G}) \times \mathbb{P}_{\mathcal{L}}(\mathbf{G}) = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G}) \times \frac{\mathbb{E}_{\mathcal{L}'}[\mathbb{1}_{\mathbf{G}} \cdot \check{W}]}{\mathbb{E}_{\mathcal{L}'}[\check{W}]} \\ & \geq \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G}) \times \mathbb{E}_{\mathcal{L}'}[\mathbb{1}_{\mathbf{G}} \cdot \check{W}] \\ & = \mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G}) \times \mathbb{P}_{\mathcal{L}'}(\mathbf{G}) \times \mathbb{E}_{\mathcal{L}'}[\check{W}|\mathbf{G}]. \end{aligned}$$

$\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G})$ and $\mathbb{E}_{\mathcal{L}'}[\check{W}|\mathbf{G}]$ could be bounded similarly as in the proof of Lemma 5.5. Conditioned on a realization of \mathbf{G} , the law $\mathbb{P}_{\mathcal{L}'}$ behaves like the law of free Brownian bridges. Furthermore, the law $\mathbb{P}_{\mathcal{L}}$ is given by the law of those free Brownian bridges conditioned on jumping over g at pole points in \check{P} . Therefore, we can apply Corollary 5.3 and Corollary 5.4 to bound $\mathbb{E}_{\check{\mathcal{L}}}[\check{W}|\mathbf{G}]$ and $\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G})$ respectively. From Corollary 5.4, we have $\mathbb{P}_{\mathcal{L}}(\mathbf{H}|\mathbf{G}) \geq C^{-1}e^{-CT^{5/2}}$. From Corollary 5.3, we have $\mathbb{E}_{\mathcal{L}'}[\check{W}|\mathbf{G}] \geq C^{-1}e^{-CT^{5/2}}$. We omit the detail here and refer the readers to the proof of Lemma 5.5 for a similar argument.

It suffices to show $\mathbb{P}_{\mathcal{L}'}(\mathbf{G}) \geq C^{-1}e^{-CT^{5/2}}$. Note that $W'_+ = 1$ when \mathbf{G} holds. Therefore, we have

$$\mathbb{P}_{\mathcal{L}'}(\mathbf{G}) = \mathbb{E}_{\mathcal{L}'}[\mathbb{1}_{\mathbf{G}} \cdot W'_+] = \frac{\mathbb{E}_{\mathcal{L}'}[\mathbb{1}_{\mathbf{G}} \cdot W'_+]}{\mathbb{E}_{\mathcal{L}'}[W'_+]} \times \mathbb{E}_{\mathcal{L}'}[W'_+] = \mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}) \times \mathbb{E}_{\mathcal{L}'}[W'_+]$$

Lemmas 5.6 and 5.7 implies that there exists $C > 0$ such that

$$\mathbb{P}_{\mathcal{L}'}(\mathbf{G}) \geq C^{-1}e^{-CT^{5/2}}.$$

Putting the above estimates together, we conclude that

$$\mathbb{P}_{\mathcal{L}}(\mathbf{H}) \geq C^{-1}e^{-CT^{5/2}}.$$

□

5.3. Proof of Lemma 5.7. The proof of Lemma 5.7 is based on a modification of an approach of Hammond. The key idea is the following simple calculation of normal distribution. Let N be a Gaussian random variable with mean 0 and variance σ^2 and let $M_1 \gg M_2 \gg \sigma$. The probability that $N \geq M_1 + M_2$, conditioned on $N \geq M_1$ is given by

$$\mathbb{P}(N \geq M_1 + M_2 \mid N \geq M_1) \approx e^{-\sigma^{-2}M_1M_2}.$$

In the current context, the curve \mathcal{L}'_+ behaves like N conditioned on greater than T^2 . This is because the largest possible value of $g(x)$ is of order T^2 . The variance is of order $T^{1/2}$ because $P' \subset (\ell + T^{1/2}, r - T^{1/2})$. The event \mathbf{G} behaves like $N \geq T^2 + T$ because it requires $\mathcal{L}_k \geq g + T$. Therefore, we obtain

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}) \approx e^{-T^{5/2}}.$$

We carry out the above heuristic argument in the rest of this subsection. We may assume without loss of generality that $g(\ell) = g(r) = 0$. See the discussion at the beginning of the proof of Lemma 5.2.

Lemma 5.8. *There exists a constant C_1 depending on constants in (5.1) such that*

$$\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}[W'_+] \geq C_1^{-1}e^{-C_1T^3}.$$

Proof. The proof involves direct computations of Brownian bridges and is similar to the one for Lemma Lemma 5.2.

Set $q = 2^{-1}(\ell + r)$. Under the law $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$, $B_j(q)$ are normal distributions with mean and variance

$$m_j = 2^{-1}(a_j^- + a_j^+), \quad \sigma^2 = 4^{-1}(r - \ell).$$

For all $x \in [\ell, q]$ and $y \in [q, r]$, we have

$$\begin{aligned} B_k(x) - g(x) &\geq \lambda_1 T - |B_k(q) - (m_k + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_k^{[\ell, q]}(x)|, \\ B_k(y) - g(y) &\geq \lambda_1 T - |B_k(q) - (m_k + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_k^{[q, r]}(y)|. \end{aligned}$$

Similarly, for $1 \leq j \leq k - 1$, $x \in [\ell, q]$ and $y \in [q, r]$, we have

$$\begin{aligned} B_j(x) - B_{j+1}(x) &\geq \lambda_0 T^{1/2} - |B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \\ &\quad - |B_{j+1}(q) - (m_{j+1} + 2^{-1}(r - \ell)b_1 T)| - \sup_{x \in [\ell, q]} |B_{j+1}^{[\ell, q]}(x)|, \\ B_j(y) - B_{j+1}(y) &\geq \lambda_0 T^{1/2} - |B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_j^{[q, r]}(y)| \\ &\quad - |B_{j+1}(q) - (m_{j+1} + 2^{-1}(r - \ell)b_1 T)| - \sup_{y \in [q, r]} |B_{j+1}^{[q, r]}(y)|. \end{aligned}$$

Therefore, to ensure for all $x \in [\ell, r]$,

$$B_1(x) > B_2(x) > \cdots > B_k(x) > g(x),$$

it suffices to require that for all $j \in [1, k]_{\mathbb{Z}}$,

$$|B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| < 4^{-1}\lambda T^{1/2},$$

and

$$\sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \leq 4^{-1} \lambda T^{1/2}, \quad \sup_{x \in [q, r]} |B_j^{[q, r]}(x)| \leq 4^{-1} \lambda T^{1/2}.$$

Here $\lambda = \min\{\lambda_0, \lambda_1\}$. Let N be a normal distribution with mean 0 and variance 1.

$$\begin{aligned} & \mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} (|B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| < 4^{-1} \lambda T^{1/2}) \\ &= \mathbb{P}(|N - (r - \ell)^{1/2} b_1 T| < 2^{-1} \lambda (r - \ell)^{-1/2} T^{1/2}). \end{aligned}$$

Because $r - \ell \leq b_0 T$,

$$(r - \ell)^{1/2} b_1 T \leq b_0^{1/2} b_1 T^{3/2}$$

and

$$2^{-1} \lambda (r - \ell)^{-1/2} T^{1/2} \geq 2^{-1} \lambda b_0^{-1/2}.$$

Thus

$$\begin{aligned} & \mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} (|B_j(q) - (m_j + 2^{-1}(r - \ell)b_1 T)| < 4^{-1} (1 - \mu) \lambda T^{1/2}) \\ & \geq \mathbb{P}(|N - b_0^{1/2} b_1 T^{3/2}| < 2^{-1} \lambda b_0^{-1/2}) \geq C^{-1} e^{-CT^3}. \end{aligned}$$

Next, we bound the event $\sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \leq 4^{-1} \lambda T^{1/2}$. It holds that

$$\begin{aligned} & \mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} \left(\sup_{x \in [\ell, q]} |B_j^{[\ell, q]}(x)| \leq 4^{-1} \lambda T^{1/2} \right) \\ &= \mathbb{P} \left(\sup_{x \in [0, 1]} |B^{[0, 1]}(x)| \leq 2^{-3/2} (r - \ell)^{-1/2} \lambda T^{1/2} \right) \\ & \geq \mathbb{P} \left(\sup_{x \in [0, 1]} |B^{[0, 1]}(x)| \leq 2^{-3/2} \lambda b_0^{-1/2} \right) \geq C^{-1}. \end{aligned}$$

Similarly,

$$\mathbb{P}_{\text{free}} \left(\sup_{x \in [q, r]} |B_j^{[q, r]}(x)| \leq 4^{-1} (1 - \mu) \lambda T^{1/2} \right) \geq C^{-1}.$$

Combining the above estimates, we conclude that

$$\mathbb{E}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} [W'_+] \geq C^{-1} e^{-CT^3}.$$

The assertion follows by taking C_1 large enough. \square

Let $m + 1 = |P'|$ be the number of elements in P' . We label them as $\{p_0 < p_1 < \dots < p_m\}$. Let C_2 be a large number to be determined. Consider the event

$$\text{Gap} := \{|\mathcal{L}_j(p_i) - \mathcal{L}_j(p_0)| \leq C_2 T^2 \text{ for all } j \in [1, k]_{\mathbb{Z}} \text{ and } i \in [1, m]_{\mathbb{Z}}\}.$$

In the lemma below, we show that for C_2 large enough, it holds that $\mathbb{P}_{\mathcal{L}'_+}(\text{Gap}) \geq 2^{-1}$.

Lemma 5.9. *There exists a constant C_2 depending on constants in (5.1), (5.2) such that*

$$\mathbb{P}_{\mathcal{L}'_+}(\text{Gap}) \geq 2^{-1}.$$

Proof. From the view of Lemma 5.8, it suffices to show that for large enough C_2 , we have

$$\mathbb{P}_{\text{free}}^{1, k, (\ell, r), a^+, a^-} (\text{Gap}^c) \leq 2^{-1} C_1^{-1} e^{-C_1 T^3},$$

where C_1 is the constant in Lemma 5.8. From the assumptions (5.3) and (5.4),

$$\begin{aligned} |B_j(p_i) - B_j(p_0)| &\leq \left| B_j^{[\ell,r]}(p_i) - B_j^{[\ell,r]}(p_0) \right| + \left| \frac{(p_i - p_0)(a_j^+ - a_j^-)}{r - \ell} \right| \\ &\leq \left| B_j^{[\ell,r]}(p_i) - B_j^{[\ell,r]}(p_0) \right| + b_2 T^2. \end{aligned}$$

Hence for given $j \in [1, k]_{\mathbb{Z}}$ and $i \in [1, m]_{\mathbb{Z}}$,

$$\begin{aligned} &\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} (|B_j(p_i) - B_j(p_0)| > C_2 T^2) \\ &\leq \mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} \left(\left| B_j^{[\ell,r]}(p_i) - B_j^{[\ell,r]}(p_0) \right| > (C_2 - b_2) T^2 \right) \\ &\leq \mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} \left(\sup_{x \in [\ell,r]} \left| B_j^{[\ell,r]}(x) \right| > 2^{-1} (C_2 - b_2) T^2 \right) \\ &= \mathbb{P} \left(\sup_{x \in [0,1]} \left| B^{[0,1]}(x) \right| > 2^{-1} (C_2 - b_2) (r - \ell)^{-1/2} T^2 \right) \\ &\leq \mathbb{P} \left(\sup_{x \in [0,1]} \left| B^{[0,1]}(x) \right| > 2^{-1} b_0^{-1/2} (C_2 - b_2) T^{3/2} \right). \end{aligned}$$

By taking C_2 large enough, we can obtain

$$\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} (|B_j(p_i) - B_j(p_0)| > C_2 T^2) \leq 2^{-1} k^{-1} m^{-1} C_1^{-1} e^{-C_1 T^3}.$$

Taking union over $j \in [1, k]_{\mathbb{Z}}$ and $i \in [1, m]_{\mathbb{Z}}$, it holds that

$$\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} (\text{Gap}^c) \leq 2^{-1} C_1^{-1} e^{-C_1 T^3}.$$

As a result, we conclude that

$$\mathbb{P}_{\mathcal{L}'_+} (\text{Gap}^c) = \frac{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} [\mathbb{1}_{\text{Gap}^c} \cdot W'_+]}{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} [W'_+]} \leq \frac{\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} (\text{Gap}^c)}{\mathbb{E}_{\text{free}}^{1,k,(\ell,r),a^+,a^-} [W'_+]} \leq 2^{-1}.$$

□

From now on we fix the choice of C_2 .

We write \mathbb{P}_0 for the law of $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$ conditioned on Gap and write \mathbb{E}_0 for the expectation of \mathbb{P}_0 . To simplify the notation, we further take a realization of Gap . That is, for $j \in [1, k]_{\mathbb{Z}}$ and $i \in [1, m]_{\mathbb{Z}}$, we fix $c_{j,i} \in \mathbb{R}$ with $|c_{j,i}| \leq C_2 T^2$. We abuse the notation and still use \mathbb{P}_0 to denote the law of $\mathbb{P}_{\text{free}}^{1,k,(\ell,r),a^+,a^-}$ conditioned on

$$B_j(p_i) - B(p_0) = c_{j,i}.$$

The estimates below will be uniform in any such choice.

Under the law of \mathbb{P}_0 , $B_j(p_0)$ are independent Gaussian random variables with mean and variance

$$\begin{aligned} m_j &= \frac{a_j^-}{1 + (r - p_m)^{-1} (p_0 - \ell)} + \frac{a_j^+ - c_{j,m}}{1 + (p_0 - \ell)^{-1} (r - p_m)}, \\ \sigma_0^2 &= \frac{1}{(p_0 - \ell)^{-1} + (r - p_m)^{-1}}. \end{aligned}$$

To see this, by a direct computation, the p.d.f. of $B_j(p_0)$ is proportional to

$$\exp\left(-\frac{(x - a_j^-)^2}{2(p_0 - \ell)} - \frac{(a_j^+ - c_{j,m} - x)^2}{2(r - p_m)}\right) \propto \exp\left(-\frac{(x - m_j)^2}{2\sigma_0^2}\right).$$

Furthermore, $W'_+ = 1$ if and only if

$$\begin{aligned} B_k(p_0) &> \max_{0 \leq i \leq m} \{g(p_i) - c_{k,i}\} =: h_k, \\ B_j(p_0) - B_{j+1}(p_0) &> \max_{0 \leq i \leq m} \{c_{j+1,i} - c_{j,i}\} =: h_j. \end{aligned}$$

Here we adapt the convention that $c_{j,0} = 0$. Denote such event by \mathbf{Q} . Explicitly,

$$\mathbf{Q} := \{B_k(p_0) > h_k, \text{ and } B_j(p_0) - B_{j+1}(p_0) > h_j \text{ for all } j \in [1, k-1]_{\mathbb{Z}}\}.$$

We check that

$$(5.5) \quad \mathbb{P}_{\mathcal{L}'_+}(\mathbf{G} \mid \text{Gap}) = \mathbb{P}_0(\mathbf{G} \mid \mathbf{Q}).$$

Lemma 5.10. *There exists a constant C_3 depending on constants in (5.1) and (5.2) such that*

$$\mathbb{P}_0(\mathbf{Q}) \geq C_3^{-1} e^{-C_3 \sigma_0^{-2} T^4}.$$

Proof. It can be checked directly that the event

$$B_j(p_0) \in \sum_{i=j}^k |h_i| + 2(k-j)\sigma_0 + (0, \sigma_0] \text{ for all } j \in [1, k]_{\mathbb{Z}}$$

is contained in \mathbf{Q} . Let N be a Gaussian random variable with mean 0 and variance 1.

$$\mathbb{P}_0\left(B_j(p_0) \in \sum_{i=j}^k |h_i| + 2(k-j)\sigma_0 + (0, \sigma_0]\right) = \mathbb{P}\left(N \in \sigma_0^{-1} \left(\sum_{i=j}^k |h_i| - m_j\right) + 2(k-j) + (0, 1]\right).$$

Under the assumptions (5.3) and (5.4), for $j \in [1, k]_{\mathbb{Z}}$ it holds that

$$\begin{aligned} |h_j| &\leq (2C_2 + b_0 b_1) T^2, \\ |m_j| &\leq (b_2 + C_2) T^2. \end{aligned}$$

Because $T^{-2} \leq p_0 - \ell, r - p_m \leq b_0 T$, it holds that

$$2^{1/2} b_0^{-1/2} T^{-1/2} \leq \sigma_0^{-1} \leq 2^{1/2} T^{-1/4}.$$

Therefore,

$$\left| \sigma_0^{-1} \left(\sum_{i=j}^k |h_i| - m_j\right) + 2(k-j) \right| \leq C \sigma_0^{-1} T^2.$$

And

$$\mathbb{P}_0\left(B_j(p_0) \in \sum_{i=j}^k |h_i| + 2(k-j)\sigma_0 + (0, \sigma_0]\right) \geq \mathbb{P}(N \in C \sigma_0^{-1} T^2 + (0, 1]) \geq C^{-1} e^{-C \sigma_0^{-2} T^4}.$$

Because $B_j(p_0)$ are independent under the law \mathbb{P}_0 , we conclude that

$$\mathbb{P}_0(\mathbf{Q}) \geq C^{-1} e^{-C \sigma_0^{-2} T^4}.$$

The assertion follows by taking C_3 large enough. \square

Let C_4 be a large constant to be determined.

Lemma 5.11. *Assume that $C_4 \geq 2(b_2 + C_2)$. Then there exists a universal constant C_5 such that*

$$\mathbb{P}_0(B_1(p_0) > C_4 T^2) \leq C_5 e^{-C_5^{-1} C_4^2 \sigma_0^{-2} T^4}.$$

Proof. Let N be a Gaussian random variable with mean 0 and variance 1. It holds that

$$\mathbb{P}_0(B_1(p_0) > C_4 T^2) = \mathbb{P}(N > \sigma_0^{-1}(C_4 T^2 - m_1)).$$

Using $|m_1| \leq (b_2 + C_2)T^2$, we have

$$\begin{aligned} \mathbb{P}_0(B_1(p_0) > C_4 T^2) &\leq \mathbb{P}(N > \sigma_0^{-1}(C_4 - C_2 - b_2)T^2) \\ &\leq \mathbb{P}(N > 2^{-1} \sigma_0^{-1} C_4 T^2). \end{aligned}$$

Hence the assertion follows. \square

Lemma 5.12. *Assume that $C_4 \geq 2(b_2 + C_2)$. Then there exists a constant C_6 depending on constants in (5.1) such that*

$$\mathbb{P}_0(\mathbf{G}^c \cap \{B_1(p_0) \leq C_4 T^2\} | \mathbf{Q}) \leq 1 - C_6^{-1} e^{-C_6 C_4 T^{5/2}}.$$

Proof. Let $\mathbf{J} = \mathbf{G}^c \cap \mathbf{Q} \cap \{B_1(p_0) \leq C_4 T^2\}$ and $\Omega \subset \mathbb{R}^k$ be the set such that

$$\mathbf{J} = \{(B_1(p_0), B_2(p_0), \dots, B_k(p_0)) \in \Omega\}.$$

If Ω is of measure zero, the assertion is clearly true. From now on we assume Ω has positive measure.

Let $\alpha = \lambda_0 T^{1/2}$ and $\beta = \lambda_1 T$. Define the map $\Phi : \mathbb{R}^k \rightarrow \mathbb{R}^k$ by

$$\Phi_j(x_1, x_2, \dots, x_k) := x_j + \alpha(k - j) + \beta.$$

Define

$$\Phi(\mathbf{J}) := \{(B_1(p_0), B_2(p_0), \dots, B_k(p_0)) \in \Phi(\Omega)\}.$$

It holds that $\Phi(\mathbf{J}) \cup \mathbf{J} \subset \mathbf{Q}$ and $\Phi(\mathbf{J}) \cap \mathbf{G}^c = \emptyset$. Hence we have

$$\mathbb{P}_0(\mathbf{G}^c \cap \{B_1(p_0) \leq M\} | \mathbf{Q}) \leq \frac{\mathbb{P}_0(\mathbf{J})}{\mathbb{P}_0(\mathbf{J}) + \mathbb{P}_0(\Phi(\mathbf{J}))}.$$

Recall that m_j and σ_0^2 are the means and variance of $B_j(p_0)$ under the law \mathbb{P}_0 .

$$\mathbb{P}_0(\mathbf{J}) = \int_{\Omega} \prod_{j=1}^k \rho_{\sigma_0^2, m_j}(x_j) \prod_{j=1}^k dx_j.$$

Because the Jacobian of Φ equals 1,

$$\mathbb{P}_0(\Phi(\mathbf{J})) = \int_{\Phi(\Omega)} \prod_{j=1}^k \rho_{\sigma_0^2, m_j}(x_j) \prod_{j=1}^k dx_j = \int_{\Omega} \prod_{j=1}^k \rho_{\sigma_0^2, m_j}(x_j + \alpha(k - j) + \beta) \prod_{j=1}^k dx_j.$$

Thus

$$\begin{aligned} \frac{\mathbb{P}_0(\Phi(\mathbf{J}))}{\mathbb{P}_0(\mathbf{J})} &\geq \inf_{\Omega} \prod_{j=1}^k \frac{\rho_{\sigma_0^2, m_j}(x_j + \alpha(k - j) + \beta)}{\rho_{\sigma_0^2, m_j}(x_j)} \\ &= \inf_{\Omega} \prod_{j=1}^k \exp(-\sigma_0^{-2} [(x_j - m_j)(\alpha(k - j) + \beta) + 2^{-1}(\alpha(k - j) + \beta)^2]). \end{aligned}$$

For $(x_1, x_2, \dots, x_k) \in \Omega$ and $C_4 \geq 2(b_2 + C_2)$, we have $x_j - m_j \leq 2C_4 T^2$. Together with $\sigma_0^{-2} \leq 2T^{-1/2}$, the above is bounded from below by $C^{-1} e^{-CC_4 T^{5/2}}$ for some large constant C . As a result,

$$\mathbb{P}_0(\mathbf{G}^c \cap \{B_1(p_0) \leq M\} | \mathbf{Q}) \leq \frac{1}{1 + C^{-1} e^{-CC_4 T^{5/2}}} \leq 1 - C^{-1} e^{-CC_4 T^{5/2}}.$$

□

We are ready to prove Lemma 5.7.

Proof of Lemma 5.7. From Lemma 5.9,

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}) \geq \mathbb{P}_{\mathcal{L}'_+}(\mathbf{G} \cap \mathbf{Gap}) = \mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}|\mathbf{Gap}) \times \mathbb{P}_{\mathcal{L}'_+}(\mathbf{Gap}) \geq 2^{-1}\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}|\mathbf{Gap}).$$

From (5.5), Lemmas 5.10, 5.11 and 5.12, we have

$$\begin{aligned} \mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}|\mathbf{Gap}) &= \mathbb{P}_0(\mathbf{G}|\mathbf{Q}) \geq \mathbb{P}_0(\mathbf{G} \cap \{B_1(p_0) \leq C_4 T^2\}|\mathbf{Q}) \\ &\geq 1 - \mathbb{P}_0(\mathbf{G}^c \cap \{B_1(p_0) \leq C_4 T^2\}|\mathbf{Q}) - \mathbb{P}_0(B_1(p_0) > C_1 T^2|\mathbf{Q}) \\ &\geq 1 - (1 - C_6 e^{-C_6^{-1} C_4 T^{5/2}}) - C_3 C_5 e^{-\sigma_0^2 T^4 (C_5^{-1} C_4^2 - C_3)}. \end{aligned}$$

Requiring $C_4^2 \geq 2^{-1} C_3 C_5$ and using $\sigma_0^{-2} \geq 2b_0^{-1} T^{-1}$, we have

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}|\mathbf{Gap}) \geq C_6^{-1} e^{-C_6 C_4 T^{5/2}} - C_3 C_5 e^{-b_0^{-1} C_5^{-1} C_4^2 T^3}.$$

Further requiring C_4 to be large enough, we can obtain $2^{-1} C_6^{-1} e^{-C_6 C_4 T^{5/2}} \geq C_3 C_5 e^{-b_0^{-1} C_5^{-1} C_4^2 T^3}$ and thus

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}|\mathbf{Gap}) \geq 2^{-1} C_6^{-1} e^{-C_6 C_4 T^{5/2}}.$$

For concreteness, taking

$$C_4 := \max\{2(b_2 + C_2), 2^{-1/2} C_3^{1/2} C_5^{1/2}, b_0 C_5 (C_6 + 1), \log(2C_3 C_5 C_6)\},$$

We have

$$\mathbb{P}_{\mathcal{L}'_+}(\mathbf{G}) \geq 4^{-1} C_6^{-1} e^{-C_6 C_4 T^{5/2}}.$$

The proof is finished. □

6. THE "SOFT" JUMP ENSEMBLE

In this section, we introduce the jump ensemble J . Inspired by the jump ensemble [Ham1], J is the intermediary between the scaled KPZ line ensemble \mathfrak{H}^t and the Brownian bridge ensemble. We design J to be adapted to the intersecting nature of \mathfrak{H}^t ; moreover J is built to meet the following requirements:

- (1) The scaled KPZ line ensemble \mathfrak{H}^t is well represented by J .
- (2) J is comparable to Brownian bridge ensemble on $[0, s]$.

Requirement (1) is realized in Proposition 3.5 and requirement (2) is entailed in Proposition 3.6. These requirements are competing with each other. We find an appropriate balance between these two. We proceed to explain the design concept of J in detail.

Recall that $\text{dom}(\bar{\ell}, \bar{r})$, $\text{bdd}(\bar{\ell}, \bar{r})$ and $\text{ext}(\bar{\ell}, \bar{r})$ are regions in $\mathbb{N} \times \mathbb{R}$ defined in Section 3. We run Gibbs resampling for \mathfrak{H}^t in $\text{dom}(\bar{\ell}, \bar{r})$. The law of $\mathfrak{H}_t|_{\text{dom}(\bar{\ell}, \bar{r})}$ is then given by $\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ with $f = \mathfrak{H}^t|_{\text{bdd}(\bar{\ell}, \bar{r})}$. More precisely, $\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ is specified via the following Radon-Nikodym derivative relation,

$$\frac{d\mathbb{P}_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}}{d\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}}(\mathcal{L}) = \frac{W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L})}{\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f} \left[W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f} \right]}.$$




FIGURE 3. Illustration of the Tent as a good approximation to the concave majorant $\mathfrak{c}(x)$.

Here $\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ is the law of independent Brownian bridges in $\text{dom}(\bar{\ell}, \bar{r})$. The Boltzmann weight $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ is defined in (3.8) as

$$\begin{aligned} W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) &= \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_j, \ell_{j+1}] \cup [r_{j+1}, r_j]} \mathbf{H}_t(f_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ &\quad \times \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_{j+1}, r_{j+1}]} \mathbf{H}_t(\mathcal{L}_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ &\quad \times \exp \left(- \int_{[\ell_k, r_k]} \mathbf{H}_t(f_{k+1}(x) - \mathcal{L}_k(x)) dx \right). \end{aligned}$$

The jump ensemble J is obtained by replacing $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ with a new weight W_{jump} . In order to achieve the requirement (2), W_{jump} is designed in the way that

- (i) The interaction in W_{jump} is only activated near a finite subset of \mathbb{R} , (the Pole set).
- (ii) There is no self-interaction within J . The interaction is only present between curves in J and the boundary data f .

In order to achieve (1), the location where we activate the interaction in W_{jump} needs to be carefully chosen. Our choice is inspired by the following crucial observation of Hammond [Ham1].

If a function $g(x)$ is close to a parabola $-2^{-1}x^2$, its concave majorant $\mathfrak{c}(x)$ is also close to $-2^{-1}x^2$ with derivative close to $-x$, see Lemma A.8. At extreme points of $\mathfrak{c}(x)$, \mathfrak{c} agrees with g . Let P a discrete subset of extreme points of $\mathfrak{c}(x)$. A linear interpolation between points in P , denoted by $\text{Tent}(x)$, gives a good approximation to \mathfrak{c} , see Lemma A.4. As a result, a Brownian bridge conditioned on jumping over $g(x)$ in an interval is well-approximated by a Brownian bridge conditioned on jumping over $g(x)$ only at points in P . See Figure 3 for an illustration.

There is an extra difficulty. In the current context, different indexed jump curves J_j travel in different intervals, e.g. (ℓ_j, r_j) for J_i . Near $x = \ell_j$, we need to have $\text{Tent}(x)$, up to a controllable error, bounded from above by $f_j(\ell_j)$. This scenario allows us to raise J_j above $\text{Tent}(x)$ within a short amount of travel time. Therefore, we need to construct multiple Pole sets P_j and multiple tent maps $\text{Tent}_j(x)$, one for each J_j .

In subsection 6.1 we list essential properties of P_{j+1} needed in the rest of the paper. The construction of P_{j+1} and the verification of those properties are postponed to subsection 6.2. Skipping subsection 6.2 will not affect the reader's understanding for the rest of the paper. The construction of W_{jump} and the jump ensemble J are done in subsection 6.3.

6.1. Pole sets. Throughout this section, we fix $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$. Denote $\bar{\ell} = (\ell_1, \ell_2, \dots, \ell_k)$, $\bar{r} = (r_1, r_2, \dots, r_k)$ and $f_J = (f_1, f_2, \dots, f_{k+1})$.

Define $\underline{g}(x)$ by

$$(6.1) \quad \underline{g}(x) := \begin{cases} f_2(x) & x \in [\ell_0, \ell_2) \cup (r_2, r_0], \\ f_{i+1}(x) & x \in (\ell_i, \ell_{i+1}) \cup (r_{i+1}, r_i), \quad i \in [2, k-1]_{\mathbb{Z}}, \\ f_{k+1}(x) & x \in (\ell_k, r_k), \\ \max\{f_i(x), f_{i+1}(x)\} & x \in \{\ell_i, r_i\}, \quad i \in [2, k]_{\mathbb{Z}}. \end{cases}$$

The function \underline{g} consists of parts in f_J that interact \mathfrak{H} . At $x = \ell_j$ or $x = r_j$, the value of $\underline{g}(x)$ is chosen such that $\underline{g}(x)$ is upper semi-continuous. See Figure 4 for an illustration.




FIGURE 4. An illustration of $\underline{g}(x)$ when $k = 3$. Recall that the resampling domain resembles an upside-down pyramid and the resampled curves only interact with values on the boundary of the resampling domain, i.e. f_2 on $[\ell_1, \ell_2]$, f_3 on $[\ell_2, \ell_3]$, f_4 on $[\ell_3, \ell_4]$.

For all $j \in [1, k]_{\mathbb{Z}}$ and $x \in (\ell_j, r_j)$, we have

$$(6.2) \quad \begin{aligned} \underline{g}(x) &\leq f_{j+1}(\ell_j) + (k+1-j)T(x - \ell_j), \\ \underline{g}(x) &\leq f_{j+1}(r_j) + (k+1-j)T(r_j - x). \end{aligned}$$

See Figure 5 for an illustration. The proof can be found in the next subsection.




FIGURE 5. This picture illustrates (6.2) near ℓ_j . Note that for $x \in (\ell_j, \ell_{j+1})$, $\underline{g}(x) = f_{j+1}(x)$.

In the next subsection, we will construct Pole sets P_{j+1} for $j \in [1, k]_{\mathbb{Z}}$. P_{j+1} consists of some extreme points of a concave function which captures f_{j+1} well near $x = \ell_j$ and $x = r_j$. P_{j+1} satisfies the following requirements.

P_{j+1} is a subset of $[\ell_j, r_j]$ which includes ℓ_j and r_j . The number of pole points in P_{j+1} is at most of order T .

$$(6.3) \quad P_{j+1} \subset [\ell_j, r_j], \ell_j, r_j \in P_{j+1} \text{ and } |P_{j+1}| \leq |r_j - \ell_j| + 1.$$

We require different indexed P_{j+1} agree each other in $(-2^{-1}T, 2^{-1}T)$. For $i, j \in [1, k]_{\mathbb{Z}}$,

$$(6.4) \quad P_{i+1} \cap (-2^{-1}T, 2^{-1}T) = P_{j+1} \cap (-2^{-1}T, 2^{-1}T).$$

In order to make J close to Brownian bridges in the interval $[0, s]$, the number of pole points in $[0, s]$ is limited to 1. Suppose $p_0 \in P_{j+1} \cap [0, s]$. Let $p_- = \max\{p \in P_{j+1} \mid p < p_0\}$ and $p_+ = \min\{p \in P_{j+1} \mid p > p_0\}$ be the elements in P_{j+1} next to p_0 . We require that

$$(6.5) \quad s \leq |p_0 - p_{\pm}| < 4^{-1}T.$$

Near ℓ_j and r_j , we need to keep the full interaction $-\mathbf{H}_t(f_{j+1}(x) - \mathcal{L}_j(x))$ unchanged in W_{jump} . See subsection 6.3 for more explanation. We want to make sure this full interaction is not interrupted by the pole points. To do so, we require P_{j+1} to avoid ℓ_i and r_i . Define

$$(6.6) \quad d' := T^{-3/2}.$$

d' determines the width of intervals near ℓ_j and r_j in which $-\mathbf{H}_t(f_{j+1}(x) - \mathcal{L}_j(x))$ is preserved in W_{jump} . We will explain the choice of scaling in more detail in Section 7. For all $j \in [1, k]_{\mathbb{Z}}$ and $i \in [j, k]_{\mathbb{Z}}$, we ask

$$(6.7) \quad P_{j+1} \cap (\ell_i, \ell_i + 2d') = P_{j+1} \cap (r_i - 2d', r_i) = \phi.$$

Define the tent maps

$$(6.8) \quad \text{Tent}_{j+1}(x) := \begin{cases} f_{j+1}(x) & x = \ell_j \text{ or } x = r_j, \\ \underline{g}(x) & x \in P_{j+1} \setminus \{\ell_j, r_j\}, \\ \text{linear interpolation} & x \in [\ell_j, r_j] \setminus P_{j+1}. \end{cases}$$

We need $\text{Tent}_{j+1}(x)$ to be a good approximation of $\underline{g}(x)$ in the following sense. For all $x \in (\ell_j, r_j)$,

$$(6.9) \quad \text{Tent}_{j+1}(x) \geq \underline{g}(x) - 3(k+1-j)sT.$$

Moreover, we require that the slope of $\text{Tent}_{j+1}(x)$ is of order T .

$$(6.10) \quad \sup_{x \neq y \in [\ell_j, r_j]} \left| \frac{\text{Tent}_{j+1}(x) - \text{Tent}_{j+1}(y)}{x - y} \right| \leq 2(k+1-j)T.$$

The construction of P_{j+1} and the verification of above properties can be found in the next subsection.

6.2. Construction of pole sets. For $j \in [1, k]_{\mathbb{Z}}$, define

$$(6.11) \quad g_{j+1}(x) := \begin{cases} f_{j+1}(x) & x \in [\ell_0, \ell_j] \cup [r_j, r_0], \\ \underline{g}(x) & x \in (\ell_j, r_j). \end{cases}$$

Let \mathbf{c}_{j+1} be the concave majorant of g_{j+1} in $[\ell_0, r_0]$.

$$\mathbf{c}_{j+1}(x_0) := \inf\{ax_0 + b \mid ax + b \geq g_{j+1}(x) \text{ for all } x \in [\ell_0, r_0]\}.$$

Define

$$\mathbf{xExt}_{j+1} := \{x \in (\ell_0, r_0) \mid \mathbf{c}_{j+1} \text{ is not linear near } x\} \cup \{\ell_0, r_0\}.$$

The reason to use $g_{j+1}(x)$ instead of $\underline{g}(x)$ to construct the concave majorant is to obtain a better fit to f_{j+1} near $x = \ell_j$ and $x = r_j$. Especially, we have the following lemma.

Lemma 6.1. *For all $j \in [1, k]_{\mathbb{Z}}$, $\ell_j, r_j \in \mathbf{xExt}_{j+1}$ and (6.2) holds.*

Proof. From Condition **C1** and the construction of (\mathbf{l}, \mathbf{r}) in Section 4, we have

$$|\ell_j + (k - j + 1)T| \leq 5^{-1}T, |r_j - (k - j + 1)T| \leq 5^{-1}T.$$

From (6.11), g_{j+1} and f_{j+1} agree on $R_{j+1, \pm} = [\pm(k + 1 - j)T - 2^{-1}T, \pm(k + 1 - j)T + 2^{-1}T]$. Recall that \mathbf{f}_{j+1} , defined in section 4, is the concave majorant of f_{j+1} on $R_{j+1, \pm}$. By Condition **C2**, we can apply Corollary A.10 with $\delta = 10^{-1}T$ to show that \mathbf{c}_{j+1} and \mathbf{f}_{j+1} agree on $\bar{R}_{j+1, \pm} = [\pm(k + 1 - j)T - 5^{-1}T, \pm(k + 1 - j)T + 5^{-1}T]$. In particular, ℓ_j and r_j are extreme points of \mathbf{c}_j . See Lemmas A.5 and A.8.

Next, we prove (6.2). From Lemma A.7 and (6.11),

$$\mathbf{c}_{j+1}(\ell_j) = g_{j+1}(\ell_j) = f_{j+1}(\ell_j).$$

Moreover, Condition **C2** and Lemma A.8 imply

$$|\ell_j + (k + 1 - j)T| \leq 10^{-1}T, |r_j - (k + 1 - j)T| \leq 10^{-1}T.$$

In particular, $(\mathbf{c}'_{j+1})_+(\ell_j) \leq (k + 1 - j)T$. By the concavity of \mathbf{c}_{j+1} , we have for all $x \in (\ell_j, r_j)$

$$\begin{aligned} \underline{g}(x) &= g_{j+1}(x) \leq \mathbf{c}_{j+1}(x) \\ &\leq \mathbf{c}_{j+1}(\ell_j) + (k + 1 - j)T(x - \ell_j) \\ &= f_{j+1}(\ell_j) + (k + 1 - j)T(x - \ell_j). \end{aligned}$$

This proves the first part of (6.2). The other part can be proved similarly. \square

Remark 6.2. *From Condition **C2** and Corollary A.10, $\mathbf{c}_{j+1}(x)$ agrees each other on $[-2^{-1}T, 2^{-1}T]$ for all $j \in [1, k]_{\mathbb{Z}}$. Hence $\mathbf{xExt}_{j+1} \cap (-2^{-1}T, 2^{-1}T)$ are also the same.*

For $j \in [1, k]_{\mathbb{Z}}$, let $\mathring{P}_{j+1} \subset \mathbf{xExt}_{j+1} \cap [\ell_j, r_j]$ be a subset satisfying the following properties.

- $\mathring{P}_{j+1} \cap (-2^{-1}T, 2^{-1}T)$ are identical for all $j \in [1, k]_{\mathbb{Z}}$.
- $\ell_j, r_j \in \mathring{P}_{j+1}$
- For any $x_1 \neq x_2 \in \mathring{P}_{j+1}$, $|x_1 - x_2| \geq s$
- For any $y \in \mathbf{xExt}_{j+1} \cap [\ell_j, r_j]$, there exists $x \in \mathring{P}_{j+1}$ such that $|x - y| < s$.

The sets \mathring{P}_{j+1} satisfy the conditions (6.3), (6.4) and (6.5). To get (6.7), we need further modification. Let

$$\begin{aligned} A_{j+1}^- &:= \{i \in [j+1, k]_{\mathbb{Z}} \mid P_{j+1} \cap (\ell_i, \ell_i + 2d') \neq \emptyset\}, \\ A_{j+1}^+ &:= \{i \in [j+1, k]_{\mathbb{Z}} \mid P_{j+1} \cap (r_i - 2d', r_i) \neq \emptyset\}. \end{aligned}$$

The sets A_{j+1}^{\pm} record the indices of i for which P_{j+1} has a pole point near $x = \ell_i$ or $x = r_i$ respectively. If $i \in A_{j+1}^-$, we replace the pole points in $\mathring{P}_{j+1} \cap (\ell_i, \ell_i + 2d')$ by ℓ_i . The same replacement is also done for $i \in A_{j+1}^+$. Concretely, we define the Pole set by

$$(6.12) \quad P_{j+1} := \mathring{P}_{j+1} \cup \{\ell_i \mid i \in A_{j+1}^-\} \cup \{r_i \mid i \in A_{j+1}^+\} \setminus \bigcup_{i=j+1}^k \left((\ell_i, \ell_i + 2d') \cup (r_i - 2d', r_i) \right).$$

By the construction, (6.7) holds.

From the view of Lemma A.8 and condition **C2**, two consecutive elements in xExt_{j+1} have distance less than $10^{-1}T$. Together with $T \geq 10$, (6.5) holds.

Next, we aim to show (6.9) and (6.10). Define the tent map from \mathring{P}_{j+1} as

$$\mathring{\text{Tent}}_{j+1}(x) := \begin{cases} f_{j+1}(x) & x = \ell_j \text{ or } x = r_j, \\ \underline{g}(x) & x \in \mathring{P}_{j+1} \setminus \{\ell_j, r_j\}, \\ \text{linear interperlation} & x \in [\ell_j, r_j] \setminus \mathring{P}_{j+1}. \end{cases}$$

Because $|(\mathbf{c}_{j+1})'_{\pm}(x)| \leq (k+1-j)T$ for $x \in [\ell_j, r_j]$, we have from Lemma A.4,

$$\mathring{\text{Tent}}_{j+1}(x) \geq \mathbf{c}_{j+1}(x) - 2(k+1-j)sT \geq \underline{g}_{j+1}(x) - 2(k+1-j)sT.$$

In particular, for $x \in (\ell_j, r_j)$, we have

$$\mathring{\text{Tent}}_{j+1}(x) \geq \underline{g}(x) - 2(k+1-j)sT.$$

Moreover,

$$\sup_{x \neq y \in [\ell_j, r_j]} \left| \frac{\mathring{\text{Tent}}_{j+1}(x) - \mathring{\text{Tent}}_{j+1}(y)}{x - y} \right| \leq (k+1-j)T.$$

The strategy of proving (6.9) and (6.10) is to compare Tent_{j+1} with $\mathring{\text{Tent}}_{j+1}$ using Lemma B.3.

Lemma 6.3. *Suppose $p \in \mathring{P}_{j+1} \cap (\ell_i, \ell_i + 2d')$ for some $j \in [1, k]_{\mathbb{Z}}$ and $i \in [j+1, k]_{\mathbb{Z}}$. We have*

$$|\underline{g}(\ell_i) - \underline{g}(p)| \leq 2(k+1-j)T^{-1/2}.$$

Similarly, suppose $p \in \mathring{P}_j \cap [r_i - 2d', r_i)$ for some $j \in [1, k]_{\mathbb{Z}}$ and $i \in [j+1, k]_{\mathbb{Z}}$. Then

$$|\underline{g}(r_i) - \underline{g}(p)| \leq 2(k+1-j)T^{-1/2}.$$

We postpone the proof of Lemma 6.3 and show (6.9) and (6.10) first. For (6.9), it remains to prove

$$\text{Tent}_{j+1}(x) \geq \mathring{\text{Tent}}_{j+1}(x) - (k+1-j)sT.$$

Note that $\text{Tent}_{j+1}(x)$ agrees with $\mathring{\text{Tent}}_{j+1}(x)$ except for x near ℓ_i and r_i .

We give the proof for the case x close to ℓ_i . We may assume that $\mathring{P}_j \cap (\ell_i, \ell_i + 2d') \neq \emptyset$. Otherwise, $\text{Tent}_{j+1}(x) = \mathring{\text{Tent}}_{j+1}(x)$ near ℓ_i . Let $p = \mathring{P}_j \cap (\ell_i, \ell_i + 2d')$, $q = \ell_i$, $p_- < p < p_+$ be the elements

in \mathring{P}_{j+1} next to p . In the interval $[p_-, p_+]$, the tent maps $\mathring{\text{Tent}}_{j+1}(x)$ and $\text{Tent}_{j+1}(x)$ are piecewise linear functions given by

$$\begin{aligned}\mathring{\text{Tent}}_{j+1}(x) &= \begin{cases} \underline{g}(x) & x = p \text{ or } x = p_{\pm}, \\ \text{linear interperlation} & x \in (p_-, p) \cup (p, p_+). \end{cases} \\ \text{Tent}_{j+1}(x) &= \begin{cases} \underline{g}(x) & x = q \text{ or } x = p_{\pm}, \\ \text{linear interperlation} & x \in (p_-, q) \cup (q, p_+). \end{cases}\end{aligned}$$

To compare $\mathring{\text{Tent}}_{j+1}(x)$ and $\text{Tent}_{j+1}(x)$, we apply Lemma B.3. From Lemma 6.3,

$$|\underline{g}(\ell_i) - \underline{g}(p)| \leq 2(k+1-j)T^{-1/2}.$$

Note that $|p - q| \leq 2d' = 2T^{-3/2}$ and $|p_{\pm} - p| \geq s \geq 1$. Applying Lemma B.3 yields

$$\text{Tent}_{j+1}(x) - \mathring{\text{Tent}}_{j+1}(x) \geq -2(k+1-j)T^{-1/2} - 2T^{-3/2} \max\{|\mathring{m}_-|, |\mathring{m}_+|\}.$$

Here \mathring{m}_{\pm} is the slope of $\mathring{\text{Tent}}_{j+1}$ in $[p_-, p]$ and $[p, p_+]$. Because the slope of $\mathring{\text{Tent}}_{j+1}$ is bounded by $(k+1-j)T$, we get

$$\text{Tent}_{j+1}(x) \geq \mathring{\text{Tent}}_{j+1}(x) - 4(k+1-j)T^{-1/2}.$$

The proof of (6.9) near ℓ_i is finished.

Next, we turn to (6.10). We again apply Lemma B.3. The difference between the slopes of $\mathring{\text{Tent}}_{j+1}$ and Tent_{j+1} is bounded by $8(k+1-j)T^{-1/2}$. Therefore, (6.10) holds true. The argument for the case near r_i is similar.

Proof of Lemma 6.3. By symmetry, it suffices to prove the case $p \in \mathring{P}_{j+1} \cap (\ell_i, \ell_i + 2d')$. We start by showing $\underline{g}(\ell_i) \leq \underline{g}(p)$. By the concavity of \mathbf{c}_{j+1} ,

$$\underline{g}(\ell_i) = g_{j+1}(\ell_i) \leq \mathbf{c}_{j+1}(\ell_i) \leq \mathbf{c}_{j+1}(p) + (\mathbf{c}'_{j+1})_+(p)(\ell_i - p).$$

Because p is an extreme point of \mathbf{c}_{j+1} ,

$$\mathbf{c}_{j+1}(p) + (\mathbf{c}'_{j+1})_+(p)(\ell_i - p) = g_{j+1}(p) - (\mathbf{c}'_{j+1})_+(p)(p - \ell_i).$$

From the definitions of \underline{g} and g_{j+1} , see (6.1) and (6.11),

$$g_{j+1}(p) - (\mathbf{c}'_{j+1})_+(p)(p - \ell_i) = \underline{g}(p) - (\mathbf{c}'_{j+1})_+(p)(p - \ell_i).$$

By Lemma A.8, $(\mathbf{c}'_{j+1})_+(p) \geq -p - 10^{-1}T \geq 0$. We conclude $\underline{g}(\ell_i) \leq \underline{g}(p)$.

Next, we show that $\underline{g}(p) \leq \underline{g}(\ell_i) + 2(k+1-j)T^{-1/2}$. Because $p \in (\ell_i, \ell_i + 2d')$,

$$\underline{g}(p) = f_{i+1}(p) = g_{i+1}(p).$$

By the concavity of \mathbf{c}_{i+1} ,

$$g_{i+1}(p) \leq \mathbf{c}_{i+1}(p) \leq \mathbf{c}_{i+1}(\ell_i) + (\mathbf{c}'_{i+1})_+(\ell_i)(p - \ell_i).$$

Because ℓ_i is an extreme point of \mathbf{c}_{i+1} , it holds that

$$\mathbf{c}_{i+1}(\ell_i) = g_{i+1}(\ell_i) = f_{i+1}(\ell_i) \leq \underline{g}(\ell_i).$$

From Lemma A.8, we have

$$\mathbf{c}'_{i+1}(\ell_i) \leq -\ell_i + 10^{-1}T \leq (k+2-i)T \leq (k+1-j)T.$$

Together with $p - \ell_i \leq 2T^{-3/2}$, we conclude that

$$\underline{g}(p) \leq \underline{g}(\ell_i) + 2(k+1-j)T^{-1/2}.$$

□

6.3. Jump ensemble. In this subsection, we use the Pole set P_{j+1} to construct the jump ensemble. We put all P_{j+1} together to form the joint Pole set.

$$(6.13) \quad P^* := \bigcup_{j=1}^k P_{j+1}.$$

Recall that $d = 64T^{-4}(\log T)^2$. For $j \in [1, k-1]_{\mathbb{Z}}$, define

$$\text{Int}_j := \left(\bigcup_{p \in P^* \cap (\ell_j, \ell_{j+1})} [p-d, p] \right) \cup \left(\bigcup_{p \in P^* \cap [r_{j+1}, r_j]} [p, p+d] \right)$$

and

$$\text{Int}_k := \left(\bigcup_{p \in P^* \cap (\ell_k, 0)} [p-d, p] \right) \cup \left(\bigcup_{p \in P^* \cap [0, r_k]} [p, p+d] \right)$$

From (6.7), we have $\text{Int}_j \subset (\ell_j, \ell_{j+1}) \cup [r_{j+1}, r_j]$ for $j \in [1, k-1]_{\mathbb{Z}}$ and $\text{Int}_k \subset (\ell_k, r_k)$. Int_j and Int_k indicates the region where J interacts with the boundary. The index j in Int_j records the layer of the boundary data. For instance, in region Int_j , the lower boundary is given by f_{j+1} .




FIGURE 6. This picture illustrates W_{jump}^j near ℓ_j . In the thick pillar with width d' , we use the Hamiltonian \mathbf{H}_t as in the scaled KPZ line ensemble. In the thin pillars with width d , a weaker Hamiltonian $\mathbf{H}_t^{j,j+1}$ is used. There is no interaction on other places.

Now we are ready to define the jump weight W_{jump} . Recall that we fix the triple $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$. Fix $j \in [1, k]_{\mathbb{Z}}$. Let

$$W_{\text{jump}}^j(\mathcal{L}_j; f|_J) := \exp \left(- \int_{[\ell_j, \ell_j + d'] \cup [r_j - d', r_j]} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_j(x)) dx \right) \\ \times \exp \left(- \sum_{i=j}^k \int_{\text{Int}_i} \mathbf{H}_t^{i,i+1}(\underline{g}(x) - \mathcal{L}_j(x)) dx \right).$$

Here $\underline{g}(x)$ is the lower boundary defined in (6.1) and $\mathbf{H}_t^{i,j+1}(x)$ is the Hamiltonian defined in (B.1). $\mathbf{H}_t^{i,j+1}(x)$ is chosen to guarantee Lemma 6.4.

For any $\ell < r$, we define the truncated version

$$(6.14) \quad W_{\text{jump}}^{j, [\ell, r]}(\mathcal{L}_j; f|_J) := \exp \left(- \int_{([\ell_j, \ell_j + d'] \cup [r_j - d', r_j]) \cap [\ell, r]} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_j(x)) dx \right) \\ \times \exp \left(- \sum_{i=j}^k \int_{\text{Int}_i \cap [\ell, r]} \mathbf{H}_t^{j, i+1}(\underline{g}(x) - \mathcal{L}_j(x)) dx \right).$$

The jump weight W_{jump} is defined by

$$(6.15) \quad W_{\text{jump}}(\mathcal{L}; f_J) := \prod_{j=1}^k W_{\text{jump}}^j(\mathcal{L}_j; f_J).$$

In W_{jump} , the random curves don't interact with each other. In particular, the jump curves J_i and J_j , defined in (3.13), are independent of each other if $i \neq j$.

The j -th jump curve J_j interacts with \underline{g} near ℓ_j and r_j through \mathbf{H}_t . This is the same Hamiltonian in $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$. See (3.8). The reason to keep the Hamiltonian function comes from the possibility that f_J may go out of order. If $f_{j+1} > f_j$ near ℓ_j , the interaction in $-\mathbf{H}_t(f_{j+1}(x) - \mathcal{L}_j(x))$ near ℓ_j becomes so drastic such that any mollification would make a huge difference. Thus we preserve the original Hamiltonian.

The other location where J_j interacts with $\underline{g}(x)$ is determined by the Pole set P^* , which depends on f_J .

Having the jump weight W_{jump} , W_{rest} is defined by

$$(6.16) \quad W_{\text{rest}}(\mathcal{L}; f_J) := W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) / W_{\text{jump}}(\mathcal{L}; f_J).$$

Here $f = f_J|_{\text{bdd}(\bar{\ell}, \bar{r})}$.

We finish this section by the following lemma.

Lemma 6.4. *It holds that*

$$W_{\text{rest}}(\mathcal{L}; f_J) \leq 1.$$

Proof. This is equivalent to $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) \leq W_{\text{jump}}(\mathcal{L}; f_J)$. From (6.1), we can rewrite $W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ in (3.8) as

$$W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}(\mathcal{L}) = \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_j, \ell_{j+1}] \cup [r_{j+1}, r_j]} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_j(x)) dx \right) \\ \times \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_{j+1}, r_{j+1}]} \mathbf{H}_t(\mathcal{L}_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ \times \exp \left(- \int_{[\ell_k, r_k]} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_k(x)) dx \right).$$

Suppose $x_0 \in (\ell_j, \ell_{j+1})$ for some $j \in [1, k]_{\mathbb{Z}}$, the integrand of $\log W_{\mathbf{H}_t}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ at x_0 is

$$(6.17) \quad -\mathbf{H}_t(\underline{g}(x_0) - \mathcal{L}_j(x_0)) - \sum_{i=1}^{j-1} \mathbf{H}_t(\mathcal{L}_{i+1}(x_0) - \mathcal{L}_i(x_0)).$$

We claim that it is smaller than the integrand of $\log W_{\text{jump}}(\mathcal{L}; f_J)$. Suppose $x_0 \in (\ell_j, \ell_j + d']$. Because of (6.7), for any $i \in [1, k]_{\mathbb{Z}}$,

$$\text{Int}_i \cap (\ell_j, \ell_j + d'] = \phi.$$

Hence the integrand of $\log W_{\text{jump}}(\mathcal{L}; f_J)$ at x_0 is $-\mathbf{H}_t(\underline{g}(x_0) - \mathcal{L}_j(x_0))$, which is larger than (6.17). Suppose $x_0 \in \text{Int}_j$, the integrand of $\log W_{\text{jump}}(\mathcal{L}; f_J)$ at x_0 is

$$-\sum_{i=1}^j \mathbf{H}_t^{i,j+1}(\underline{g}(x_0) - \mathcal{L}_i(x_0)).$$

In light of Lemma B.2, it is larger than (6.17). Suppose $x_0 \in (\ell_j + d', \ell_{j+1}) \setminus \text{Int}_j$. Then the integrand of $\log W_{\text{jump}}(\mathcal{L}; f_J)$ at x_0 is zero, which is also larger than (6.17). This completes the discussion for the case $x_0 \in (\ell_j, \ell_{j+1})$. Other cases, $x_0 \in (r_{j+1}, r_j)$ and $x_0 \in (\ell_{k+1}, r_{k+1})$ can be treated similarly and we omit the detail. The remaining end points $x_0 = \ell_i$ or $x_0 = r_i$ have zero contribution. The proof is finished. \square

7. PROOF OF PROPOSITION 3.5

In this section, we aim to show that

$$\inf\{\mathbb{E}_J[W_{\text{rest}}] \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}\} \geq D^{-1}e^{-DT^{5/2}}.$$

Here \mathcal{G} is defined in Definition 4.5 and W_{rest} is defined in (6.16). Throughout this section, we fix $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$. In below we explain the strategy of the proof.

Consider the event **NoTouch** that requires the jump ensemble J to stay ordered and to lie above \underline{g} except in neighborhoods near ℓ_j or r_j . Concretely,

$$\begin{aligned} \text{NoTouch} := & \{J_j(x) \geq \underline{g}(x) \text{ for } j \in [1, k]_{\mathbb{Z}} \text{ and } x \in [\ell_j + d', r_j - d']\}, \\ & \{J_j(x) \geq J_{j+1}(x) \text{ for } j \in [1, k-1]_{\mathbb{Z}} \text{ and } x \in [\ell_{j+1}, r_{j+1}]\}. \end{aligned}$$

When **NoTouch** occurs, the weight W_{rest} is bounded from below by e^{-T} . See Lemma 7.1. Therefore Proposition 3.5 is reduced to showing $\mathbb{P}_J(\text{NoTouch}) \geq e^{-T^{5/2}}$.

To begin, we show that each curve J_j can jump over \underline{g} in the interval $[\ell_j + d', r_j - d']$ with probability $e^{-T^{5/2}}$. Moreover, a gap of size T between J_j and \underline{g} can be created in the interval $[\ell_j + 1, r_j - 1]$. This event is denoted by \mathbf{A}_j .

Next, we apply induction on the number of curves. Take $j \in [2, k]_{\mathbb{Z}}$. We write \mathbf{F}_{j+1} for the event that J_1, J_2, \dots, J_{j-1} stay ordered and that $J_{j-1} \geq \underline{g} + T$ in $[\ell_{j-1} + 1, r_{j-1} - 1]$. Suppose that $\mathbb{P}(\mathbf{F}_{j-1}) \geq e^{-T^{5/2}}$. Because the j -th curve J_j is independent of other curves, $\mathbb{P}(\mathbf{F}_{j-1} \cap \mathbf{A}_j) \geq e^{-T^{5/2}}$. To get bounds on $\mathbb{P}_J(\mathbf{F}_j)$, we need to make sure J_j does not raise too high and does not touch J_{j-1} .

To do so, we apply a stopping domain argument. Consider the leftest and the rightest locations where $J_{j-1} = J_j + 2^{-1}T$. Denote them by ι_j'' and τ_j'' respectively. At these two points, we know $J_1 \geq J_2 \geq \dots \geq J_{j-1} > J_j > \underline{g}$. Furthermore, $J_{j-1} - J_j = 2^{-1}T$ and $J_j - \underline{g} \geq 2^{-1}T$. Applying the strong \mathbf{H}_t -Brownian Gibbs property, we resample J_1, \dots, J_j in $[\iota_j'', \tau_j'']$. Because the boundary data is now well spaced, we can show that the J_1, \dots, J_j , with probability $e^{-T^{5/2}}$, remain ordered and lie beyond \underline{g} in $[\iota_j'', \tau_j'']$. This finishes the proof.

Lemma 7.1. *There exists $D = D(k, s)$ such that for all $t \geq 1$, it holds that*

$$\mathbb{1}_{\text{NoTouch}} \cdot W_{\text{rest}}(\cdot; f_J) \geq D^{-1}e^{-DT} \cdot \mathbb{1}_{\text{NoTouch}}.$$

Proof. From the definition of W_{rest} in (6.16) and $\mathbf{H}_t^{i,j} \geq 0$, we have

$$\begin{aligned} W_{\text{rest}}(\mathcal{L}; f_J) &\geq \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_j+d', \ell_{j+1}] \cup [r_{j+1}, r_j-d']} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_j(x)) dx \right) \\ &\quad \times \prod_{j=1}^{k-1} \exp \left(- \int_{[\ell_{j+1}, r_{j+1}]} \mathbf{H}_t(\mathcal{L}_{j+1}(x) - \mathcal{L}_j(x)) dx \right) \\ &\quad \times \exp \left(- \int_{[\ell_k+d', r_k-d']} \mathbf{H}_t(\underline{g}(x) - \mathcal{L}_k(x)) dx \right). \end{aligned}$$

When NoTouch occurs, the integrands above are all bounded from below by -1 . Hence

$$W_{\text{rest}}(\mathcal{L}; f_J) \geq e^{-k(r_1-\ell_1)} \geq e^{-2k(k+1)T}.$$

□

Proposition 7.2. *There exists a constant $D = D(k, s)$ such that for E large enough, it holds that*

$$\mathbb{P}_J(\text{NoTouch}) \geq D^{-1} e^{-DT^{5/2}}.$$

Proof of Proposition 3.5. From Lemma 7.1 and Proposition 7.2

$$\mathbb{E}_J[W_{\text{rest}}(J; f_J)] \geq \mathbb{P}_J(\text{NoTouch}) \times D^{-1} e^{-DT} \geq D^{-1} e^{-DT^{5/2}}.$$

□

The rest of the section is devoted to prove Proposition 7.2. In subsection 7.1, we deal with a single curve J_j and show that A_j occurs with probability at least $e^{-T^{5/2}}$. In subsection 7.2, we show that, provided the boundary data is well separated, J remains ordered and lying beyond \underline{g} under resampling. Lastly, we combine the two and finish the induction argument in subsection 7.3.

7.1. Single curve. In this subsection, we focus on a single curve J_j and prove Proposition 7.3. Roughly speaking, Proposition 7.3 says the probability that J_j is larger than \underline{g} in $[\ell_j + d', r_j - d']$ is at least $e^{-T^{5/2}}$.

We start by defining A_j . Fix $j \in [1, k]_{\mathbb{Z}}$ and let A_j be the event that

- $J_j(x) > \underline{g}(x)$ for all $x \in [\ell_j + d', r_j - d']$,
- $J_j(x) > \text{Tent}_{j+1}(x) + 16k(k+1-j)sT$ for all $x \in [\ell_j + 1, r_j - 1]$.

Proposition 7.3. *There exists $D = D(k)$ such that for E large enough, we have*

$$\mathbb{P}_J(A_j) \geq D^{-1} e^{-DT^{5/2}}.$$

We prove Proposition 7.3 by gradually lift the curve J_j . In below we define the events which record the lifting process.

Let $A_{j,1}$ be the event such that

- $J_j(\ell_j + d') > f_j(\ell_j) + kT^{1/2}$, $J_j(r_j - d') > f_j(r_j) + kT^{1/2}$,
- $J_j(\ell_j + 1) > f_j(\ell_j) + 20k(k+1-j)sT$,
- $J_j(r_j - 1) > f_j(r_j) + 20k(k+1-j)sT$.

Let $A_{j,2}$ be the event such that

- $J_j(x) > f_{j+1}(x)$ for all $x \in [\ell_j + d', \ell_{j+1}] \cup [r_j - 1, r_j - d']$,
- $J_j(\ell_j + 1) > \text{Tent}_{j+1}(\ell_j + 1) + 18k(k+1-j)sT$
- $J_j(r_j - 1) > \text{Tent}_{j+1}(r_j - 1) + 18k(k+1-j)sT$.

Let $A_{j,3}$ be the event such that

- $J_j(x) > \text{Tent}_{j+1}(x) + 16k(k+1-j)sT$ for all $x \in [\ell_j + 1, r_j - 1]$.

Recall that J_j is sampled in $[\ell_j, r_j]$ with boundary given by $J_j(\ell_j) = f(\ell_j)$ and $J_j(r_j) = f(r_j)$. In the event $A_{j,1}$, J_j goes up by $T^{1/2}$ within distance d' and then goes up by sT within distance 1. We then resample J_j in $[\ell_j + d', \ell_j + 1]$ and $[r_j - 1, r_j - d']$. Conditioned on $A_{j,1}$ the boundary data is high enough to ensure $A_{j,2}$ occurs with probability D^{-1} . After that having $A_{j,2}$, we resample J_j in $[\ell_j + 1, r_j - 1]$. Applying Theorem 5.1 ensures that conditioned on $A_{j,2}$, $A_{j,3}$ occurs with probability $e^{-T^{5/2}}$. This argument is entailed in the following three lemmas. To simplify the notation, we denote $\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ and $\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ by \mathbb{P}_{free} and \mathbb{E}_{free} respectively.

Lemma 7.4. *There exists $D = D(k, s)$ such that*

$$\mathbb{P}_{\text{free}}(A_{j,1}) \geq D^{-1}e^{-DT^{5/2}}.$$

Lemma 7.5. *There exists $D = D(k, s)$ such that*

$$\mathbb{P}_{\text{free}}(A_{j,2} | A_{j,1}) \geq D^{-1}.$$

Lemma 7.6. *There exists $D = D(k, s)$ such that for E large enough, it holds that*

$$\mathbb{P}_J(A_{j,3} | A_{j,2}) \geq D^{-1}e^{-DT^{5/2}}.$$

Proof of Proposition 7.3. From (6.9), we check $A_{j,2} \cap A_{j,3} \subset A_j$. By Lemmas 7.4 and 7.5,

$$\mathbb{P}_{\text{free}}(A_{j,2}) \geq \mathbb{P}_{\text{free}}(A_{j,2} | A_{j,1}) \times \mathbb{P}_{\text{free}}(A_{j,1}) \geq D^{-1}e^{-D^{5/2}}.$$

By the stochastic monotonicity Lemma 2.7,

$$\mathbb{P}_J(A_{j,2}) \geq \mathbb{P}_{\text{free}}(A_{j,2}) \geq D^{-1}e^{-D^{5/2}}.$$

With Lemma 7.6, we conclude that

$$\mathbb{P}_J(A_j) \geq \mathbb{P}_J(A_{j,2} \cap A_{j,3}) = \mathbb{P}_J(A_{j,3} | A_{j,2}) \times \mathbb{P}_J(A_{j,2}) \geq D^{-1}e^{-DT^{5/2}}.$$

□

In the rest of this subsection, we prove Lemmas 7.4, 7.5 and 7.6.

Proof of Lemma 7.4. The first condition in $A_{j,1}$ requires J_j to raise by an amount $T^{1/2}$ within a distance $d' = T^{-3/2}$. This event occurs with probability about $e^{-T^{5/2}}$. The second and the third condition requires J_j to raise by an amount T within a distance 1. This event occurs with probability about e^{-T^2} . In below we carry out this calculation.

Under the law \mathbb{P}_{free} , $B_j(\ell_j + d')$ is a Gaussian random variable with mean and variance

$$m = \frac{(r_j - \ell_j - d')f_j(\ell_j) + d'f_j(r_j)}{r_j - \ell_j}, \quad \sigma^2 = \frac{(r_j - \ell_j - d')d'}{r_j - \ell_j}.$$

Let N be a Gaussian random variable with mean 0 and variance 1. It holds that

$$\mathbb{P}_{\text{free}}(B_j(\ell_j + d') > f_j(\ell_j) + kT^{1/2}) = \mathbb{P}(N > \sigma^{-1}(f_j(\ell_j) - m) + \sigma^{-1}kT^{1/2}).$$

From Condition **C1** and Remark 4.3, it holds that $|\ell_j + (k+j-1)T|, |r_j - (k+j-1)T| \leq 10^{-1}T$. Recall that $d' = T^{-3/2}$. Hence $\sigma^{-1} \leq 2^{1/2}T^{3/4}$ and then

$$\sigma^{-1}kT \leq DT^{5/4}.$$

From Condition **C2**, we have $\left| \frac{f(\ell_j) - f(r_j)}{\ell_j - r_j} \right| \leq DT$. Thus

$$\sigma^{-1} |f_j(\ell_j) - m| \leq DT^{1/4}.$$

As a result,

$$\mathbb{P}_{\text{free}}(B_j(\ell_j + d') > f_j(\ell_j) + kT^{1/2}) \geq \mathbb{P}(N \geq DT^{5/4}) \geq D^{-1} e^{-DT^{5/2}}.$$

Conditioned on $B_j(\ell_j + d') = y$, $B_j(\ell_j + 1)$ is a normal distribution with mean and variance

$$m = \frac{(r_j - \ell_j - 1)y + (1 - d')f_j(r_j)}{r_j - \ell_j - d'}, \quad \sigma^2 = \frac{(r_j - \ell_j - 1)(1 - d')}{r_j - \ell_j - d'}.$$

Thus

$$\begin{aligned} & \mathbb{P}_{\text{free}}(B_j(\ell_j + 1') \geq f_j(\ell_j) + 20k(k+1-j)sT \mid B_j(\ell_j + d') = y) \\ &= \mathbb{P}(N > \sigma^{-1}(f_j(\ell_j) - m) + 20\sigma^{-1}k(k+1-j)sT) \end{aligned}$$

Because $\sigma \approx 1$,

$$20\sigma^{-1}k(k+1-j)sT \leq DT.$$

When $y \geq f_j(\ell_j) + kT^{1/2}$,

$$f_j(\ell_j) - m \leq \frac{(1 - d')(f_j(r_j) - f_j(\ell_j))}{r_j - \ell_j - d'} - \frac{(r_j - \ell_j - 1)kT^{1/2}}{r_j - \ell_j - d'} \leq DT.$$

As a result,

$$\begin{aligned} & \mathbb{1}\{y \geq f_j(\ell_j) + kT^{1/2}\} \times \mathbb{P}_{\text{free}}(B_j(\ell_j + 1') > f_j(\ell_j) + 20k(k+1-j)sT \mid B_j(\ell_j + d') = y) \\ & \leq \mathbb{1}\{y \geq f_j(\ell_j) + kT^{1/2}\} \times D^{-1} e^{-DT^2}. \end{aligned}$$

This finishes the proof for $\ell_j + d'$ and $\ell_j + 1$. The one for $r_j - d'$ and $r_j - 1$ is similar and we omit the detail. \square

Proof of Lemma 7.5. We first bound f_{j+1} by a linear function. By (6.1) and (6.2), it holds that for $x \in (\ell_j, \ell_j + 1]$,

$$f_{j+1}(x) = \underline{g}(x) \leq f_{j+1}(\ell_j) + (x - \ell_j)(k + 1 - j)T.$$

By Conditions **C4**, we obtain

$$f_{j+1}(x) = \underline{g}(x) \leq f_j(\ell_j) + 3\Delta + (x - \ell_j)(k + 1 - j)T =: L(x).$$

Next, we show that when $A_{j,1}$ occurs, it holds that

$$\begin{aligned} L(\ell_j + d') + 2^{-1}kT^{1/2} &< J_j(\ell_j + d'), \\ L(\ell_j + 1) + 18k(k+1-j)sT &< J_j(\ell_j + 1). \end{aligned}$$

They can be deduced by

$$\begin{aligned} L(\ell_j + d') = f_j(\ell_j) + 3\Delta + (k+1-j)T^{-1/2} &\leq f_j(\ell_j) + 2^{-1}kT^{1/2} \\ &< J_j(\ell_j + d') - 2^{-1}kT^{1/2}, \end{aligned}$$

and

$$\begin{aligned} L(\ell_j + 1) = f_j(\ell_j) + 3\Delta + (k+1-j)T &\leq f_j(\ell_j) + 2k(k+1-j)T \\ &< J_j(\ell_j + 1) - 18k(k+1-j)sT. \end{aligned}$$

In particular, the second condition in $A_{j,2}$ is ensured by $A_{j,1}$.

Applying a similar argument near r_j , the third condition in $A_{j,2}$ is also ensured by $A_{j,1}$. Furthermore,

$$A_{j,1} \cap \left\{ \left| J_j^{[\ell_j+d', \ell_j+1]} \right| \leq 2^{-1}kT^{1/2} \right\} \cap \left\{ \left| J_j^{[r_j-1, r_j-d']} \right| \leq 2^{-1}kT^{1/2} \right\} \subset A_{j,2}.$$

We conclude that

$$\mathbb{P}_{\text{free}}(A_{j,2} | A_{j,1}) \geq D^{-1}.$$

□

Proof of Lemma 7.6. We first replace $A_{j,2}$ by $A_{j,4}$, which fixes the value of $J_j(x)$ at $x = \ell_j + 1$ and $x = r_j - 1$. Let

$$A_{j,4} := \{J_j(x) = \text{Tent}_{j+1}(x) + 18k(k+1-j)sT \text{ for } x \in \{\ell_j + 1, r_j - 1\}\}.$$

From the stochastic monotonicity Lemma 2.7, it holds that

$$\mathbb{P}_J(A_{j,3} | A_{j,2}) \geq \mathbb{P}_J(A_{j,3} | A_{j,4}).$$

It remains to prove that

$$\mathbb{P}_J(A_{j,3} | A_{j,4}) \geq D^{-1}e^{-DT^{5/2}}.$$

Consider the event

$$L_j := \{\mathcal{L}_j(p) \leq \text{Tent}_{j+1}(p) - \Delta_k \text{ for some } p \in P_{j+1} \cap [\ell_j + 1, r_j - 1]\}.$$

This event represents the scenario that J_j goes below \underline{g} by an amount Δ_k at some point in $P_{j+1} \cap [\ell_j + 1, r_j - 1]$.

Step 1: We show in this step that conditioned on $A_{j,4}$, L_j is unlikely.

$$\mathbb{P}_J(L_j | A_{j,4}) \leq 2^{-1}.$$

We begin by controlling the local fluctuation of J_j . Recall that we write \mathbb{P}_{free} and \mathbb{E}_{free} for $\mathbb{P}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ and $\mathbb{E}_{\text{free}}^{\text{dom}(\bar{\ell}, \bar{r}), f}$ respectively. From (6.9) and (6.10), we can apply Lemma B.4 to get

$$\mathbb{P}_{\text{free}}(B_j(x) \geq \underline{g}(x) \text{ for } x \in [\ell_j + 1, r_j - 1] | A_{j,4}) \geq D^{-1}e^{-DT^3}.$$

When such event occurs, $W_{\text{Jump}}^{j, [\ell_j+1, r_j-1]}$ is bounded from below by some constant D^{-1} . Therefore,

$$\mathbb{E}_{\text{free}} \left[W_{\text{Jump}}^{j, [\ell_j+1, r_j-1]} | A_{j,4} \right] \geq D^{-1}e^{-DT^3}.$$

Let

$$\text{Flct}_j := \{|J_j(x) - J_j(y)| \leq \Delta \text{ for all } x, y \in [\ell_j + 1, r_j - 1] \text{ with } |x - y| \leq d\}.$$

When $A_{j,4}$ occurs, $J_j(x) = \text{Tent}_{j+1}(x) + 18k(k+1-j)sT$ for $x \in \{\ell_j + 1, r_j - 1\}$. From (6.10), we have

$$\left| \frac{J_j(\ell_j + 1) - J_j(r_j - 1)}{r_j - \ell_j - 2} \right| = \left| \frac{\text{Tent}_{j+1}(\ell_j + 1) - \text{Tent}_{j+1}(r_j - 1)}{r_j - \ell_j - 2} \right| \leq 2kT.$$

Hence

$$\begin{aligned} \mathbb{P}_{\text{free}}(\text{Flct}_j^c | A_{j,4}) &\leq \mathbb{P}_{\text{free}}(\omega_{d, [\ell, r]}(B^{[\ell, r]}) \geq \Delta - 2kdT) \\ &\leq \mathbb{P}_{\text{free}}(\omega_{d, [\ell, r]}(B^{[\ell, r]}) \geq 2^{-1}\Delta) \leq De^{-D^{-1}T^4}. \end{aligned}$$

Here

$$\omega_{d, [\ell, r]}(B^{[\ell, r]}) := \max_{x, y \in [\ell, r], |x-y| \leq d} |B^{[\ell, r]}(x) - B^{[\ell, r]}(y)|.$$

As a result, for E large enough, we have

$$\mathbb{P}_J(\text{Flct}_j^c | \mathbf{A}_{j,4}) \leq De^{DT^3 - D^{-1}T^4} \leq 4^{-1}.$$

Next, we show that $\mathbf{L}_j \cap \text{Flct}_j$ is unlikely. From Condition **C3**, \mathbf{L}_j and Flct_j imply there exists an interval in $\text{Int}_j \cap [\ell_j + 1, r_j - 1]$ with length d where

$$J_j(x) \leq \underline{g}(x) - k\Delta - k \log k.$$

Together with

$$d\mathbf{H}_{j,k+1}(k\Delta + k \log k) \geq de^\Delta = 64T^4(\log T)^2,$$

it holds that

$$\mathbb{1}_{\mathbf{L}_j \cap \text{Flct}_j} \times W_{\text{jump}}^{j, [\ell_j+1, r_j-1]} \leq e^{-64T^4(\log T)^2} \times \mathbb{1}_{\mathbf{L}_j \cap \text{Flct}_j}.$$

As a result, for E large enough, it holds that

$$\begin{aligned} \mathbb{P}_J(\mathbf{L}_j \cap \text{Flct}_j | \mathbf{A}_{j,4}) &\leq \mathbb{E}_{\text{free}} \left[W_{\text{jump}}^{j, [\ell_j+1, r_j-1]} | \mathbf{A}_{j,4} \right]^{-1} \times e^{-64T^4(\log T)^2} \\ &\leq De^{DT^3 - 64T^4(\log T)^2} \leq 4^{-1}. \end{aligned}$$

Putting the above estimates together, we conclude that

$$\mathbb{P}_J(\mathbf{L}_j | \mathbf{A}_{j,4}) \leq \mathbb{P}_J(\mathbf{L}_j \cap \text{Flct}_j | \mathbf{A}_{j,4}) + \mathbb{P}_J(\text{Flct}_j^c | \mathbf{A}_{j,4}) \leq 2^{-1}.$$

Step 2: In this step, we want to prove

$$\mathbb{P}_{\text{free}}(\mathbf{A}_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c) \geq D^{-1}e^{-DT^{5/2}}.$$

Let

$$\mathbf{A}'_{j,3} := \{J_j(p) > \text{Tent}_{j+1}(p) + 17k(k+1-j)sT \text{ for } p \in P_{j+1} \cap (\ell_j + 1, r_j - 1)\}.$$

To apply Theorem 5.1, we rewrite the following objects. Set $\check{\ell} = \ell_j + 1, \check{r} = r_j - 1,$

$$\check{f}(x) := \text{Tent}_{j+1}(x) - \Delta_k, \check{\mathcal{L}}(x) := J_j(x), \check{P} = P_{j+1} \cap (\check{\ell}, \check{r})$$

and

$$\check{a}^- = \text{Tent}_{j+1}(\check{\ell}) + 18k(k+1-j)sT, \check{a}^+ = \text{Tent}_{j+1}(\check{r}) + 18k(k+1-j)sT.$$

The law of $\check{\mathcal{L}}$ in $[\check{\ell}, \check{r}]$ under $\mathbb{P}_{\text{free}}(\cdot | \mathbf{A}_{j,4})$ is given by $\mathbb{P}_{\text{free}}^{1,1,(\check{\ell}, \check{r}), \check{a}^-, \check{a}^+}$. Furthermore, it can easily verified that

$$\begin{aligned} \mathbf{L}_j^c &= \{\check{\mathcal{L}}(p) > \check{f}(p) \text{ for } p \in \check{P}\}, \\ \mathbf{A}'_{j,3} &= \{\check{\mathcal{L}}(p) > \check{f}(p) + 17k(k+1-j)sT + \Delta_k \text{ for } p \in \check{P}\}. \end{aligned}$$

Because

$$\check{a}^- - \check{f}(\ell_j) = \check{a}^+ - \check{f}(r_j) = 18k(k+1-j)sT + \Delta_k,$$

we can applying Theorem 5.1 to conclude that

$$\mathbb{P}_{\text{free}}(\mathbf{A}'_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c) \geq D^{-1}e^{-DT^{5/2}}.$$

Resample J_j between pole points in $P_{j+1} \cap [\ell_j + 1, r_j - 1]$, we obtain

$$\mathbb{P}_{\text{free}}(\mathbf{A}_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c) \geq \mathbb{P}_{\text{free}}(\mathbf{A}_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c \cap \mathbf{A}'_{j,3}) \times \mathbb{P}_{\text{free}}(\mathbf{A}'_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c) \geq D^{-1}e^{-DT^{5/2}}.$$

Step 3: We finish the proof in this step. When $\mathbf{A}_{j,3}$ occurs, $W_{\text{Jump}}^{j, [\ell_j+1, r_j-1]} \geq D^{-1}$. Hence

$$\mathbb{P}_J(\mathbf{A}_{j,3} | \mathbf{A}_{j,4} \cap \mathbf{L}_j^c) \geq D^{-1}e^{-DT^{5/2}}.$$

In conclusion,

$$\mathbb{P}_J(\mathbf{A}_{j,3} | \mathbf{A}_{j,4}) \geq \mathbb{P}_J(\mathbf{A}_{j,3} | \mathbf{A}_{j,4} \cap \mathcal{L}_j^c) \times \mathbb{P}_J(\mathcal{L}_j^c | \mathbf{A}_{j,4}) \geq D^{-1} e^{-DT^{5/2}}.$$

□

7.2. Curve separation over the resampling interval. In this subsection, we prove Proposition 7.7. Roughly speaking, Proposition 7.7 says the following. Suppose at two points ℓ and r , the jump ensemble J is ordered and lie above \underline{g} . Then the probability that J remains ordered and lie above \underline{g} in $[\ell, r]$ is at least $e^{-T^{5/2}}$.

We begin with introducing the notation in order to resample J in the interval $[\ell, r]$. Fix $j \in [1, k]_{\mathbb{Z}}$. Given $\ell_j \leq \ell < r \leq r_j$ and $a^\pm = (a_1^-, \dots, a_k^-, a_1^+, \dots, a_k^+)$, define

$$\frac{d\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],a^+,a^-}}{d\mathbb{P}_{\text{free}}^{1,j,[\ell,r],a^+,a^-}}(\mathcal{L}) \propto \prod_{i=1}^j W_{\text{jump}}^{i,[\ell,r]}(\mathcal{L}_i; f_J).$$

Next, we prescribe the boundary condition which is well-spaced. Let $bS_j(\ell, r)$ be the collection of $(a^+, a^-) \in \mathbb{R}^j \times \mathbb{R}^j$ which satisfies

$$(7.1) \quad a_i^\pm - a_{i+1}^\pm \geq (k+2-j)T^{1/2} \quad \text{for all } i \in [1, j-1]_{\mathbb{Z}},$$

$$(7.2) \quad a_j^- \geq \text{Tent}_{j+1}(\ell) + 8k(2k+3-2j)sT \quad \text{and} \quad a_j^+ \geq \text{Tent}_{j+1}(r) + 8k(2k+3-2j)sT,$$

and

$$(7.3) \quad a_1^\pm \leq jT^2.$$

Let \underline{a}^\pm be the smallest member is $bS_j(\ell, r)$. That is,

$$\begin{aligned} \underline{a}_i^\pm - \underline{a}_{i+1}^\pm &= (k+2-j)T^{1/2}, \quad i \in [1, j-1]_{\mathbb{Z}}, \\ \underline{a}_j^- &= \text{Tent}_{j+1}(\ell) + 8k(2k+3-2j)sT, \\ \underline{a}_j^+ &= \text{Tent}_{j+1}(r) + 8k(2k+3-2j)sT. \end{aligned}$$

The following two events capture the scenario that the curves are well-separated in $[\ell, r]$.

$$(7.4) \quad \begin{aligned} S_j(\ell, r) &:= \{ \mathcal{L}_i(x) > \mathcal{L}_{i+1}(x) + (k+1-j)T^{1/2} \text{ for all } i \in [1, j-1]_{\mathbb{Z}} \text{ and } x \in [\ell, r] \} \\ &\quad \cap \{ \mathcal{L}_j(x) > \text{Tent}_{j+1}(x) + 16k(k+1-j)T \text{ for all } x \in [\ell, r] \}. \end{aligned}$$

$$(7.5) \quad \text{Up}_j(\ell, r) := \{ \mathcal{L}_1(x) < (j+1)T^2 \text{ for all } x \in [\ell, r] \}.$$

Proposition 7.7. *There exists a constant $D = D(k, s)$ such that for E large enough, the following statement holds. For all $j \in [1, k]_{\mathbb{Z}}$, ℓ, r in $[\ell_j, r_j]$ and $(a^+, a^-) \in bS_j(\ell, r)$, we have*

$$\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],a^-,a^+}(S_j(\ell, r) \cap \text{Up}_j(\ell, r)) \geq D^{-1} e^{-DT^{5/2}}.$$

Proof. The proof involves three steps.

Step 1: We first show that

$$\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],a^-,a^+}(\text{Up}_j^c(\ell, r)) \leq D e^{-D^{-1}T^3}.$$

This can be achieved by a simple stopping domain argument. Consider the intervals

$$\begin{aligned} I_- &:= \{ x \in [\ell, r] \mid \mathcal{L}_1(y) < (j+2^{-1})T^2 \text{ for all } y \in [\ell, x] \}, \\ I_+ &:= \{ x \in [\ell, r] \mid \mathcal{L}_1(y) < (j+2^{-1})T^2 \text{ for all } y \in [x, r] \}. \end{aligned}$$

Define

$$\begin{aligned} \ell' &:= \begin{cases} \sup I_- & I_- \neq \phi, \\ r & I_- = \phi. \end{cases} \\ \mathfrak{r}' &:= \begin{cases} \inf I_+ & I_+ \neq \phi, \\ \ell & I_+ = \phi. \end{cases} \end{aligned}$$

The last condition in $bS_j(\ell, r)$ ensures I_\pm are non-empty. Moreover,

$$\mathbf{Up}_j^c(\ell, r) \subset \{\ell < \ell' < \mathfrak{r}' < r\}.$$

Because $(j + 2^{-1})T^2$ is much larger than $\underline{g}(x)$, we have

$$\mathbb{P}_{\text{free}}^{1,1, [\ell', \mathfrak{r}'], (j+2^{-1})T^2, (j+2^{-1})T^2} (B_1(x) \geq \underline{g}(x) \text{ for all } x \in [\ell', \mathfrak{r}']) \geq D^{-1}.$$

And then

$$\mathbb{E}_{\text{free}}^{1,1, [\ell', \mathfrak{r}'], (j+2^{-1})T^2, (j+2^{-1})T^2} [W_{\text{jump}}^{1, [\ell', \mathfrak{r}']}] \geq D^{-1}.$$

Therefore,

$$\begin{aligned} & \mathbb{P}_{\text{jump}}^{1,j, [\ell', \mathfrak{r}'], (j+2^{-1})T^2, (j+2^{-1})T^2} (\mathbf{Up}_j^c(\ell', \mathfrak{r}')) \\ &= \mathbb{E}_{\text{free}}^{1,1, [\ell', \mathfrak{r}'], (j+2^{-1})T^2, (j+2^{-1})T^2} [W_{\text{jump}}^{1, [\ell', \mathfrak{r}']}]^{-1} \\ & \quad \times \mathbb{P}_{\text{free}}^{1,j, [\ell', \mathfrak{r}'], (j+2^{-1})T^2, (j+2^{-1})T^2} (\mathbf{Up}_j^c(\ell', \mathfrak{r}')) \leq D e^{-D^{-1}T^3}. \end{aligned}$$

Through the strong Markov property,

$$\begin{aligned} \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+} (\mathbf{Up}_j^c(\ell, r)) &\leq \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+} (\mathbf{Up}_j^c(\ell', \mathfrak{r}') \cap \{\ell < \ell' < \mathfrak{r}' < r\}) \\ &\leq D e^{-D^{-1}T^3}. \end{aligned}$$

This finishes Step 1.

Step 2: Define the event

$$\text{Low}_j(\ell, r) := \bigcup_{i=1}^j \{\mathcal{L}_i(p) \leq \text{Tent}_{j+1}(p) - \Delta_k \text{ for some } p \in P_{j+1} \cap [\ell, r]\}.$$

In Step 2, we show that

$$\mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+} (\text{Low}_j(\ell, r)) \leq 2^{-1}.$$

By the stochastic monotonicity Lemma 2.7, it suffices to prove

$$\mathbb{P}_{\text{jump}}^{1,j, [\ell, r], \underline{a}^-, \underline{a}^+} (\text{Low}_j(\ell, r)) \leq 2^{-1}.$$

Fix $i \in [1, j]_{\mathbb{Z}}$. Because $\underline{a}_i^- \geq \text{Tent}_{j+1}(\ell) + 8k(2k+3-2j)sT$, $\underline{a}_i^+ \geq \text{Tent}_{j+1}(r) + 8k(2k+3-2j)sT$ and (6.10), it's straightforward to show that

$$\mathbb{P}_{\text{free}}^{i,i, [\ell, r], \underline{a}_i^-, \underline{a}_i^+} (B_i(x) \geq \text{Tent}_{j+1}(x) + 7k(2k+3-2j)sT \text{ for all } x \in [\ell, r]) \geq D^{-1} e^{-DT^3}.$$

This can be done by applying Lemma B.4 with $x_1 = \ell$, $x_2 = r$ and $m = 2kT$. Together with (6.9),

$$\mathbb{P}_{\text{free}}^{1,j, [\ell, r], \underline{a}^-, \underline{a}^+} (B_i(x) \geq \underline{g}(x) \text{ for all } x \in [\ell, r] \text{ and } i \in [1, j]_{\mathbb{Z}}) \geq D^{-1} e^{-DT^3}.$$

Thus,

$$\mathbb{E}_{\text{free}}^{1,j, [\ell, r], \underline{a}^-, \underline{a}^+} \left[\prod_{i=1}^j W_{\text{jump}}^{i, [\ell, r]} \right] \geq D^{-1} e^{-DT^3}.$$

Recall that $\Delta = 8 \log T$ and $d = 64T^{-4}(\log T)^2$. Let

$$(7.6) \quad \text{Flct}_i(\ell, r) := \{|\mathcal{L}_i(x) - \mathcal{L}_i(y)| \leq \Delta \text{ for all } x, y \in [\ell, r] \text{ with } |x - y| \leq d\}.$$

Because $\mathcal{L}_i(\ell) = \underline{a}_i^-$, $\mathcal{L}_i(r) = \underline{a}_i^+$ and (6.10),

$$\left| \frac{\mathcal{L}_i(r) - \mathcal{L}_i(\ell)}{r - \ell} \right| = \left| \frac{\text{Tent}_{j+1}(r) - \text{Tent}_{j+1}(\ell)}{r - \ell} \right| \leq 2kT.$$

Hence

$$\begin{aligned} \mathbb{P}_{\text{free}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^-}(\text{Flct}_i^c(\ell, r)) &\leq \mathbb{P}_{\text{free}}(\omega_{d,[\ell,r]}(B^{[\ell,r]}) \geq \Delta - 2kdT) \\ &\leq \mathbb{P}_{\text{free}}(\omega_{d,[\ell,r]}(B^{[\ell,r]}) \geq 2^{-1}\Delta) \leq De^{-D^{-1}T^4}. \end{aligned}$$

As a result,

$$\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+} \left(\bigcup_{i=1}^j \text{Flct}_i^c(\ell, r) \right) \leq De^{DT^3 - D^{-1}T^4} \leq 4^{-1}.$$

From Condition **C3**, $\text{Low}_j(\ell, r)$ and $\bigcap_{i=1}^j \text{Flct}_i(\ell, r)$ imply there exists $i \in [1, j]_{\mathbb{Z}}$ and an interval in $\text{Int}_j \cap [\ell, r]$ with length d in which

$$\mathcal{L}_i(x) \leq \underline{g}(x) - k\Delta - k \log k.$$

Together with

$$d\mathbf{H}_{i,j+1}(k\Delta + k \log k) \geq de^\Delta = 64T^4(\log T)^2,$$

we deduce that when $\text{Low}_j(\ell, r) \cap \bigcap_{i=1}^j \text{Flct}_i(\ell, r)$ occurs,

$$\prod_{i=1}^j W_{\text{jump}}^{i,[\ell,r]} \leq e^{-64T^4(\log T)^2}.$$

As a result,

$$\begin{aligned} \mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+} \left(\text{Low}_j(\ell, r) \cap \bigcap_{i=1}^j \text{Flct}_i(\ell, r) \right) &\leq \mathbb{E}_{\text{free}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+} \left[\prod_{i=1}^j W_{\text{jump}}^{i,[\ell,r]} \right]^{-1} \times e^{-64T^4(\log T)^2} \\ &\leq De^{DT^3 - 64T^4(\log T)^2} \leq 4^{-1}. \end{aligned}$$

Putting the above estimates together, we conclude that

$$\begin{aligned} &\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+}(\text{Low}_j(\ell, r)) \\ &\leq \mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+} \left(\text{Low}_j(\ell, r) \cap \bigcap_{i=1}^j \text{Flct}_i(\ell, r) \right) + \mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+} \left(\bigcup_{i=1}^j \text{Flct}_i^c(\ell, r) \right) \\ &\leq 2^{-1}. \end{aligned}$$

Step 3: In this step, we show that

$$\mathbb{P}_{\text{jump}}^{1,j,[\ell,r],\underline{a}^-, \underline{a}^+}(\mathbf{S}_j(\ell, r) \mid \text{Low}_j^c(\ell, r)) \geq D^{-1}e^{-DT^{5/2}}.$$

Consider the event $\mathbf{S}'_j(\ell, r)$ being defined by

- $\mathcal{L}_i(x) \geq \mathcal{L}_{i+1}(x) + (k + 3/2 - j)T^{1/2}$ for all $i \in [1, j-1]_{\mathbb{Z}}$ and $x \in P_{j+1} \cap (\ell, r)$,
- $\mathcal{L}_j(x) \geq \text{Tent}_{j+1}(x) + 8k(2k + 5/2 - 2j)sT$ for all $x \in P_{j+1} \cap (\ell, r)$.

Compared to $S_j(\ell, r)$, $S'_j(\ell, r)$ requires a larger gap between \mathcal{L} at points in $P_{j+1} \cap (\ell, r)$.

In order to apply Theorem 5.1, we rewrite the following objects. Set

$$\check{f}(x) := \text{Tent}_{j+1}(x) - \Delta_k, \quad \check{\mathcal{L}}_i(x) := \mathcal{L}_i(x), \quad \check{P} = P_{j+1} \cap (\ell, r), \quad \check{a}_i^\pm = a_i^\pm.$$

It can easily verified that

$$\begin{aligned} \text{Low}_j^c &= \{\check{\mathcal{L}}_i(p) > \check{f}(p) \text{ for } i \in [1, \mathbb{Z}] \text{ and } p \in \check{P}\}, \\ S'_j(\ell, r) &= \{\check{\mathcal{L}}_i(p) \geq \check{\mathcal{L}}_{i+1}(p) + (k + 3/2 - j)T^{1/2} \text{ for all } i \in [1, j-1]_{\mathbb{Z}} \text{ and } p \in \check{P}\} \\ &\quad \cap \{\check{\mathcal{L}}_j(x) \geq \check{f}(x) + 8k(2k + 5/2 - 2j)sT + \Delta_k \text{ for all } x \in \check{P}_{j+1}\}. \end{aligned}$$

From the conditions in $bS_j(\ell, r)$

$$\begin{aligned} \check{a}_i^\pm - \check{a}_{i+1}^\pm &\geq (k + 2 - j)T^{1/2} \quad \text{for all } i \in [1, j-1]_{\mathbb{Z}}, \\ \check{a}_j^- - \check{f}(\check{\ell}), \check{a}_j^+ - \check{f}(\check{r}) &\geq 8k(2k + 3 - 2j)sT + \Delta_k \end{aligned}$$

and

$$\check{a}_1^\pm \leq jT^2.$$

Together with (6.3) and (6.10), the assumptions (5.3) and (5.4) hold. By Theorem 5.1, we conclude that

$$\mathbb{P}_{\text{free}}^{1,j, [\ell, r], a^-, a^+}(S'_j(\ell, r) \mid \text{Low}_j^c(\ell, r)) \geq D^{-1}e^{-DT^{5/2}}.$$

Resampling the curves between pole points in $P_{j+1} \cap (\ell, r)$, we deduce

$$\mathbb{P}_{\text{free}}^{1,j, [\ell, r], a^-, a^+}(S_j(\ell, r) \mid \text{Low}_j^c(\ell, r)) \geq D^{-1}e^{-DT^{5/2}}.$$

When $S_j(\ell, r)$ occurs, $\prod_{i=1}^j W_{\text{jump}}^{i, [\ell, r]} \geq D^{-1}$. Therefore

$$\mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(S_j(\ell, r) \mid \text{Low}_j^c(\ell, r)) \geq D^{-1}e^{-DT^{5/2}}.$$

In conclusion,

$$\begin{aligned} &\mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(S_j(\ell, r) \cap \text{Up}_j(\ell, r)) \geq \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(S_j(\ell, r)) - \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(\text{Up}_j^c(\ell, r)) \\ &\geq \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(S_j(\ell, r) \mid \text{Low}_j^c(\ell, r)) \times \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(\text{Low}_j^c(\ell, r)) - \mathbb{P}_{\text{jump}}^{1,j, [\ell, r], a^-, a^+}(\text{Up}_j^c(\ell, r)) \\ &\geq D^{-1}e^{-DT^{5/2}} - De^{-D^{-1}T^3} \geq D^{-1}e^{-DT^{5/2}}. \end{aligned}$$

The proof is finished. \square

7.3. Proof of Proposition 3.5. In this subsection, we combine Propositions 7.3 and 7.7 to prove Proposition 7.8. Proposition 7.2 follows easily from Proposition 7.8.

For $j \in [1, k]_{\mathbb{Z}}$, Let F_j be the event

- $J_i(x) > \underline{g}(x)$ for all $i \in [1, j]_{\mathbb{Z}}$ and $x \in [\ell_i + d', r_i - d']$,
- $J_j(x) > \text{Tent}_{j+1}(x) + 16k(k+1-j)sT$ for all $x \in [\ell_j + 1, r_j - 1]$,
- $J_i(x) > J_{i+1}(x) + (k+1-j)T^{1/2}$ for all $i \in [1, j-1]_{\mathbb{Z}}$ and $x \in [\ell_{i+1}, r_{i+1}]$,
- $J_1(x) < (j+1)T^2$ for all $x \in [\ell_1, r_1]$.

Roughly speaking, F_j is the event that $J_i > \underline{g}$ in $[\ell_i + d', r_i - d']$ and (J_1, J_2, \dots, J_j) stays ordered.

Proposition 7.8. *There exists a constant $D = D(k, s)$ such that for E large enough, the following statement holds. For all $j \in [1, k]_{\mathbb{Z}}$,*

$$(7.7) \quad \mathbb{P}_J(F_j) \geq D^{-1}e^{-DT^{5/2}}.$$

Proof of Proposition 7.2. It follows from $F_k \subset \text{NoTouch}$. \square

Proof of Proposition 7.8. We use induction on j and start with $j = 1$. Comparing A_1 and F_1 , the extra requirement in F_1 is that $J_1(x) < 2T^2$. By a stopping domain argument we can show that $\mathbb{P}_J(J_1(x) < 2T^2) \geq 1 - De^{-D^{-1}T^3}$. See Step 1 in the proof of Proposition 7.7 for a similar argument. We omit the detail here. For T large enough, it holds that

$$\mathbb{P}(F_1) \geq \mathbb{P}(A_1) - De^{-D^{-1}T^3} \geq D^{-1}e^{-DT^{5/2}}.$$

Let $j \in [2, k]_{\mathbb{Z}}$ and assume that

$$\mathbb{P}_J(F_{j-1}) \geq D^{-1}e^{-DT^{5/2}}.$$

Under the law of \mathbb{P}_J , F_{j-1} and A_j are independent. From Propositions 7.3, it holds that

$$\mathbb{P}_J(F_{j-1} \cap A_j) = \mathbb{P}_J(F_{j-1}) \times \mathbb{P}_J(A_j) \geq D^{-1}e^{-DT^{5/2}}.$$

In other words, with probability $e^{-T^{5/2}}$, J_1, \dots, J_{j+1} are well-separated and J_j jumps over \underline{g} .

Comparing $F_{j-1} \cap A_j$ with F_j , the only requirement in F_j which is not guaranteed by $F_{j-1} \cap A_j$ is

$$(7.8) \quad J_{j-1}(x) > J_j(x) + (k+1-j)T^{1/2} \text{ for all } x \in [\ell_j, r_j].$$

In other words, the only situation $F_{j-1} \cap A_j \cap F_j^c$ can occur is that J_j raises to high and intersects with J_{j-1} . We use a stopping domain argument to resolve this issue. In particular, we will consider the stopping domain given by the leftmost and rightmost locations where (7.8) fails.

Define a random set $I'_{j,-} \subset [\ell_1, r_1]$ which captures the requirements in $F_{j-1} \cap A_j$ from the left. Explicitly, $I'_{j,-} \subset [\ell_1, r_1]$ is the collection of $x_0 \in [\ell_1, r_1]$ such that

- $J_i(x) > \underline{g}(x)$ for all $i \in [1, j]_{\mathbb{Z}}$ and $x \in [\ell_i + d', r_i - d'] \cap (-\infty, x_0]$.
- $J_i(x) > J_{i+1}(x) + (k+2-j)T^{1/2}$ for all $i \in [1, j-2]_{\mathbb{Z}}$ and $x \in [\ell_{i+1}, r_{i+1}] \cap (-\infty, x_0]$.
- $J_{j-1}(x) > \text{Tent}_j(x) + 16k(k+2-j)sT$ for all $x \in [\ell_{j-1} + 1, r_{j-1} - 1] \cap (-\infty, x_0]$.
- $J_j(x) > \text{Tent}_{j+1}(x) + 16k(k+1-j)sT$ for all $x \in [\ell_j + 1, r_j - 1] \cap (-\infty, x_0]$.
- $J_1(x) < jT^2$ for all $x \in [\ell_1, r_1] \cap (-\infty, x_0]$.

Furthermore, we define $I''_{j,-}$ to be the subset of $I'_{j,-}$ with an extra requirement that

$$J_{j-1}(x) > J_j(x) + (k+2-j)T^{1/2} \text{ for all } x \in [\ell_j, r_j] \cap (-\infty, x_0].$$

Define $I'_{j,+}, I''_{j,+} \subset [\ell_1, r_1]$ by replacing $(-\infty, x_0]$ with $[x_0, \infty)$. Define

$$\ell''_j := \begin{cases} \sup I''_{j,-}, & \text{if } I''_{j,-} \neq \emptyset \\ \ell_1, & \text{if } I''_{j,-} = \emptyset. \end{cases} \quad \text{and} \quad r''_j := \begin{cases} \inf I''_{j,+}, & \text{if } I''_{j,+} \neq \emptyset \\ r_1, & \text{if } I''_{j,+} = \emptyset. \end{cases}$$

As $F_{j-1} \cap A_j$ occurs, $I'_{j,\pm} = [\ell_1, r_1]$. Moreover, $\ell_i \in I''_{j,-}$. It can be shown as the following. By the second requirement of F_{j-1} , it holds that

$$J_{j-1}(\ell_j) \geq \text{Tent}_j(\ell_j) + 16k(k+2-j)sT.$$

By (6.9), it holds that

$$\text{Tent}_j(\ell_j) + 16k(k+2-j)sT \geq \underline{g}(\ell_j) - 3ksT + 16k(k+2-j)sT.$$

Because $\underline{g}(\ell_j) = \max\{f_j(\ell_j), f_{j+1}(\ell_j)\} \geq f_j(\ell_j) = J_j(\ell_j)$, we have

$$\underline{g}(\ell_j) - 3ksT + 16k(k+2-j)sT \geq J_j(\ell_j) - 3ksT + 16k(k+2-j)sT > J_j(\ell_j) + (k+2-j)T^{1/2}.$$

Similarly, $F_{j-1} \cap A_j$ implies $r_i \in I''_{j,+}$.

Next, we discuss the implications of $F_{j-1} \cap A_j \cap F_j^c$. $F_{j-1} \cap A_j \cap F_j^c$ implies (7.8) fails at a point in (ℓ_j, r_j) . In particular, $\ell_j < \ell_j'' < \mathfrak{r}_j'' < r_j$,

$$J_j(\ell_j'') = J_{j-1}(\ell_j'') - (k+2-j)T^{1/2},$$

and

$$J_j(\mathfrak{r}_j'') = J_{j-1}(\mathfrak{r}_j'') - (k+2-j)T^{1/2}.$$

Moreover, all the conditions in $I_{j,\pm}'$ hold at ℓ_j'' and \mathfrak{r}_j'' .

Define the event K_j by

$$K_j := \{\ell_j < \ell_j'' < \mathfrak{r}_j'' < r_j\} \cap \{(J_1(\ell_j''), \dots, J_j(\ell_j''), J_1(\mathfrak{r}_j''), \dots, J_j(\mathfrak{r}_j'')) \in bS_j(\ell_j'', \mathfrak{r}_j'')\}.$$

Here $bS_j(\ell, r)$ defined by (7.1), (7.2) and (7.3). We claim that $F_{j-1} \cap A_j \cap F_j^c \subset K_j$. Comparing the condition in $bS_j(\ell_j'', \mathfrak{r}_j'')$ and in $I_{j,\pm}'$, we only need to check

$$\begin{aligned} F_{j-1} \cap A_j \cap F_j^c \subset \{ & J_j(\ell_j'') \geq \text{Tent}_{j+1}(\ell_j'') + 8k(k+3-2j)sT \\ & \cap \{ J_j(\mathfrak{r}_j'') \geq \text{Tent}_{j+1}(\mathfrak{r}_j'') + 8k(k+3-2j)sT \}. \end{aligned}$$

We give the proof for ℓ_j'' . The argument for \mathfrak{r}_j'' is similar. Suppose $\ell_j'' \in [\ell_j + 1, r_j - 1]$, then the assertion holds by the fourth requirement of $I_{j,-}'$. Assume $\ell_j'' \in [\ell_j, \ell_j + 1)$. By the third requirement of $I_{j,-}'$,

$$J_{j-1}(\ell_j'') \geq \text{Tent}_j(\ell_j'') + 16k(k+2-j)sT.$$

Because of (6.10),

$$\text{Tent}_j(\ell_j'') + 16k(k+2-j)sT \geq \text{Tent}_j(\ell_j) - 2ksT + 16k(k+2-j)sT.$$

From (6.9) and $\underline{g}(\ell_j) \geq f_{j+1}(\ell_j) = \text{Tent}_{j+1}(\ell_j)$,

$$\text{Tent}_j(\ell_j) - 2ksT + 16k(k+2-j)sT \geq \text{Tent}_{j+1}(\ell_j) - 5ksT + 16k(k+2-j)sT.$$

Using (6.10), again, we have

$$J_{j-1}(\ell_j'') \geq \text{Tent}_{j+1}(\ell_j'') - 7ksT + 16k(k+2-j)sT.$$

As a result, $F_{j-1} \cap A_j \cap F_j^c$ implies

$$\begin{aligned} J_j(\ell_j'') &= J_{j-1}(\ell_j'') - (k+2-j)T^{1/2} \\ &\geq \text{Tent}_{j+1}(\ell_j'') - 8ksT + 16k(k+2-j)sT \\ &= \text{Tent}_{j+1}(\ell_j'') + 8k(2k+3-2j)sT. \end{aligned}$$

The argument for $\ell_j'' \in (r_j - 1, r_j]$ is the similar.

Next, we check that

$$S_j(\ell_j'', \mathfrak{r}_j'') \cap \text{Up}_j(\ell_j'', \mathfrak{r}_j'') \cap K_j \subset F_j \cap K_j.$$

The events $S_j(\ell_j'', \mathfrak{r}_j'')$ and $\text{Up}_j(\ell_j'', \mathfrak{r}_j'')$ are defined in (7.4) and (7.5). The definition of K_j ensures the requirements in F_j holds outside the interval $(\ell_j'', \mathfrak{r}_j'')$. $S_j(\ell_j'', \mathfrak{r}_j'') \cap \text{Up}_j(\ell_j'', \mathfrak{r}_j'')$ then takes cares of the interval $[\ell_j'', \mathfrak{r}_j'']$.

We are ready to give a lower bound for $\mathbb{P}_J(F_j)$. Let $\delta := \mathbb{P}_J(F_{j-1} \cap A_j \cap F_j^c)$. From the above discussion,

$$\mathbb{P}_J(F_j) \geq \mathbb{P}_J(F_j \cap K_j) \geq \mathbb{P}_J(S_j(\ell_j'', \mathfrak{r}_j'') \cap \text{Up}_j(\ell_j'', \mathfrak{r}_j'') \cap K_j).$$

From the strong Markov property and Proposition 7.7, it holds that

$$\begin{aligned} \mathbb{P}_J(\mathbf{S}_j(\mathfrak{l}_j'', \mathfrak{r}_j'') \cap \mathbf{Up}_j(\mathfrak{l}_j'', \mathfrak{r}_j'') \cap \mathbf{K}_j) &= \mathbb{P}_J(\mathbf{S}_j(\mathfrak{l}_j'', \mathfrak{r}_j'') \cap \mathbf{Up}_j(\mathfrak{l}_j'', \mathfrak{r}_j'') \mid \mathbf{K}_j) \times \mathbb{P}_J(\mathbf{K}_j) \\ &\geq D^{-1} e^{-DT^{5/2}} \times \mathbb{P}_J(\mathbf{K}_j). \end{aligned}$$

Together with $\mathbf{F}_{j-1} \cap \mathbf{A}_j \cap \mathbf{F}_j^c \subset \mathbf{K}_j$, we have

$$\mathbb{P}_J(\mathbf{F}_j) \geq \delta \times D^{-1} e^{-DT^{5/2}}.$$

On the other hand,

$$\mathbb{P}_J(\mathbf{F}_j) \geq \mathbb{P}_J(\mathbf{F}_{j-1} \cap \mathbf{A}_j \cap \mathbf{F}_j) = \mathbb{P}_J(\mathbf{F}_{j-1} \cap \mathbf{A}_j) - \delta \geq D^{-1} e^{-DT^{5/2}} - \delta,$$

We conclude that

$$\mathbb{P}_J(\mathbf{F}_j) \geq \inf_{\delta \geq 0} \max\{\delta \times D^{-1} e^{-DT^{5/2}}, D^{-1} e^{-DT^{5/2}} - \delta\} \geq D^{-1} e^{-DT^{5/2}}.$$

The induction argument is finished. \square

8. PROOF OF PROPOSITION 3.6

In this section we prove Proposition 3.6. The jump ensemble J is defined through (3.13). Recall that $A \subset C_{0,0}([0, s])$ is a Borel set and $\varepsilon = \mathbb{P}_{\text{free}}(B^{[0,s]} \in A)$. The goal is to prove

$$\sup\{\mathbb{P}_J(J_k^{[0,s]} \in A) \mid (\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}\} \leq D\varepsilon \exp\left(D(\log \varepsilon^{-1})^{5/6}\right).$$

Starting from now, we fix a realization of the favorable event $(\bar{\ell}, \bar{r}, f_J) \in \mathcal{G}$ (see Definition 4.5). Suppose $P^* \cap (0, s) = \phi$, i.e. no pole in $(0, s)$. By the construction of the jump ensemble J , the law of $J_k^{[0,s]}$ is equivalent to that of a free Brownian bridge. Then $\mathbb{P}_J(J_k^{[0,s]} \in A) = \varepsilon$ and Proposition 3.6 follows easily.

From now on we assume $P^* \cap (0, s) \neq \phi$. From the construction of the pole set P^* , see (6.5) and (6.13), $P^* \cap (0, s)$ consists of only one element, denoted by p_0 . We enlarge the interval from $[0, s]$ to $[q_1, q_2]$ to prevent p_0 from being too close to the boundary. Let

$$\begin{aligned} q_1 &:= p_0 - \max\{p_0, 2^{-1}s\}, \\ q_2 &:= p_0 + \max\{s - p_0, 2^{-1}s\}. \end{aligned}$$

Let $\pi : C_{0,0}([q_1, q_2]) \rightarrow C_{0,0}([0, s])$ be a restriction map, defined as

$$\pi[h](x) := h(x) - s^{-1}xh(s) - s^{-1}(s-x)h(0).$$

Note that $\pi \circ B^{[q_1, q_2]} = B^{[0, s]}$ and $\pi \circ J_k^{[q_1, q_2]} = J_k^{[0, s]}$. For $A' = \pi^{-1}(A)$, it holds that

$$\begin{aligned} \mathbb{P}_{\text{free}}(B^{[q_1, q_2]} \in A') &= \mathbb{P}_{\text{free}}(B^{[0, s]} \in A) = \varepsilon, \\ \mathbb{P}_J(J_k^{[q_1, q_2]} \in A') &= \mathbb{P}_J(J_k^{[0, s]} \in A). \end{aligned}$$

In the remaining of this section, we consider the event $\{J_k^{[q_1, q_2]} \in A'\}$ instead of $\{J_k^{[0, s]} \in A\}$.

Let $p_- = \max\{p \in P^* \mid p < p_0\}$ and $p_+ = \min\{p \in P^* \mid p > p_0\}$ be the elements in P^* next to p_0 . From (6.5), it holds that

$$-2^{-1}T < p_- < q_1 < p_0 < q_2 < p_+ < 2^{-1}T.$$

Recall $\underline{g}(x)$ in (6.1). Define the tent map

$$\text{Tent}(x) := \begin{cases} \underline{g}(x) & x = p_0 \text{ or } x = p_{\pm}, \\ \text{linear interpolation} & x \in (p_-, p_0) \cup (p_0, p_+). \end{cases}$$

We remark that Tent agrees with Tent_{j+1} in $[p_-, p_+]$ for all $j \in [1, k]_{\mathbb{Z}}$.

Recall that $\Delta_k = (k+2)\Delta + k \log k$. Define the quantities (see Figure 7 for an illustration).

$$\begin{aligned} \underline{m}_0 &= \text{Tent}(p_0) - \Delta_k, \quad \underline{m}_{\pm} = \text{Tent}(p_{\pm}) - \Delta_k, \\ \underline{m}_1 &= \text{Tent}(q_1) - \Delta_k, \quad \underline{m}_2 = \text{Tent}(q_2) - \Delta_k, \\ \sigma_1^2 &= \frac{(p_0 - q_1)(q_1 - p_-)}{p_0 - p_-}, \quad \sigma_2^2 = \frac{(p_+ - q_2)(q_2 - p_0)}{p_+ - p_0}, \quad \sigma_3^2 = \frac{(q_2 - p_0)(p_0 - q_1)}{q_2 - q_1}. \end{aligned}$$




FIGURE 7. Tent is constructed using values of \underline{g} at poles p_-, p_0, p_+ with p_0 being the single pole in $[0, s]$. We expect the values of J_k stay around Tent , hence above $\text{Tent} - \Delta_k$ with high probability.

The following lemma shows that $J_k(p_0) \geq \underline{m}_0$ and $J_k(p_{\pm}) \geq \underline{m}_{\pm}$ occur with high probability. Define the event

$$\Upsilon := \{J_k(p_0) \geq \underline{m}_0, J_k(p_{\pm}) \geq \underline{m}_{\pm}\}.$$

Lemma 8.1. *There exists a constant $D = D(k)$ such that for E large enough, we have*

$$\mathbb{P}_J(\Upsilon^c) \leq D e^{-D^{-1}T^4}.$$

Proof. The idea is that if $J_k(p_0) < \underline{m}_0$ or $J_k(p_{\pm}) < \underline{m}_{\pm}$ occurs, then with high probability J_k will stay below $\underline{g} - k\Delta - k \log k$ in an interval with length d . Because $d \mathbf{H}_t^{k, k+1}(k\Delta + k \log k) \approx T^4$, the Boltzmann weight W_{jump}^k is less than e^{-T^4} which is unlikely. In below we fill in details of this argument.

From the definition of \mathbb{P}_J (see (3.13) and (6.15)), J_k is independent of other curves and is distributed according to

$$\frac{d\mathbb{P}_{J_k}}{d\mathbb{P}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k}}(\mathcal{L}) = \frac{W_{\text{jump}}^k(\mathcal{L})}{\mathbb{E}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k} [W_{\text{jump}}^k]}.$$

Here $x_k = f_k(\ell_k)$ and $y_k = f_k(r_k)$. We know that when Υ^c occurs, W_{jump}^k is, with high probability, less than e^{-T^4} . To conclude Υ^c is unlikely, we need to bound the normalizing constant. To do so, we turn off the interaction in W_{jump}^k except near p_0 and p_{\pm} to get a new curve J'_k . The law of J'_k satisfies the following Radon-Nikodym derivative relation,

$$\frac{d\mathbb{P}_{J'_k}}{d\mathbb{P}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k}}(\mathcal{L}) = \frac{W_{\text{jump}}^k(\mathcal{L})}{\mathbb{E}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k} [W_{\text{jump}}^k]}.$$

Here the weight $W_{\text{jump}}'^k$ is defined by

$$W_{\text{jump}}'^k(\mathcal{L}) := \exp \left(- \int_{[p_- - d, p_-] \cup [p_0, p_0 + d] \cup [p_+, p_+ + d]} \mathbf{H}_t^{k, k+1}(\underline{g}(x) - \mathcal{L}(x)) dx \right).$$

By the stochastic monotonicity Lemma 2.7, it holds that

$$\mathbb{P}_{J_k}(\mathbf{Y}^c) \leq \mathbb{P}_{J'_k}(\mathbf{Y}^c).$$

From condition **C2** and Remark 4.3, it is straightforward to check that

$$\mathbb{P}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k}(\{B(x) \geq \underline{g}(x) \text{ for } x \in [p_- - d, p_-] \cup [p_0, p_0 + d] \cup [p_+, p_+ + d]\}) \geq C^{-1} e^{-CT^3}.$$

Here $x_k = f_k(\ell_k)$ and $y_k = f_k(r_k)$. When this event occurs, $W_{\text{jump}}'^k$ is bounded from below. As a result,

$$\mathbb{E}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k} [W_{\text{jump}}'^k] \geq D^{-1} e^{-DT^3}.$$

Next, we bound the local fluctuation of J'_k . Recall the event $\text{Flct}_k(\ell_k, r_k)$ is defined in (7.6) as

$$\text{Flct}_k(\ell_k, r_k) := \{|\mathcal{L}_k(x) - \mathcal{L}_k(y)| \leq \Delta \text{ for all } x, y \in [\ell_k, r_k] \text{ with } |x - y| \leq d\}.$$

Because $\Delta^2/d = T^2$,

$$\mathbb{P}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k}(\text{Flct}_k^c(\ell_k, r_k)) \leq D e^{-D^{-1}T^4}.$$

With the bound on the normalizing constant, it holds that

$$\mathbb{P}_{J'_k}(\text{Flct}_k^c(\ell_k, r_k)) \leq D e^{DT^3 - D^{-1}T^4} \leq D e^{-D^{-1}T^4}.$$

When $\text{Flct}_k(\ell_k, r_k)$ and \mathbf{Y}^c both occur, there exists an interval with length d and on which $\underline{g}(x) - J_k(x) \geq k\Delta + k \log k$. From

$$d \mathbf{H}_t^{k, k+1}(k\Delta + k \log k) \geq d e^\Delta = 64T^4(\log T)^2,$$

we get

$$\mathbb{E}_{\text{free}}^{k, k, (\ell_k, r_k), x_k, y_k} [W_{\text{jump}}'^k \cdot \mathbb{1}_{\mathbf{Y}^c} \cdot \mathbb{1}_{\text{Flct}_k(\ell_k, r_k)}] \leq D e^{-D^{-1}T^4(\log T)^2}.$$

Using the bound on the normalizing constant again, we conclude that

$$\begin{aligned} \mathbb{P}_{J'_k}(\mathbf{Y}^c) &\leq \mathbb{P}_{J'_k}(\mathbf{Y}^c \cap \text{Flct}_k(\ell_k, r_k)) + \mathbb{P}_{J'_k}(\text{Flct}_k^c(\ell_k, r_k)) \\ &\leq D e^{-D^{-1}T^4(\log T)^2} + D e^{-D^{-1}T^4}. \end{aligned}$$

□

We now start to derive the expression (8.1) about $\mathbb{P}_J(J_k^{[q_1, q_2]} \in A')$, which will be the starting point for the estimates. For any $(x_1, x_2) \in \mathbb{R}^2$, define

$$F(x_1, x_2) := \mathbb{P}_J(J_k^{[q_1, q_2]} \in A' | J_k(q_1) = x_1, J_k(q_2) = x_2).$$

Let $Q(x_1, x_2)$ be the joint p.d.f. of $(J_k(q_1), J_k(q_2))$. Then

$$\mathbb{P}_J(J_k^{[q_1, q_2]} \in A') = \int_{\mathbb{R}^2} F(x_1, x_2) Q(x_1, x_2) dx_1 dx_2.$$

From the definition (6.14), $W_{\text{jump}}^{k, [p_-, p_0]}(\mathcal{L}_k; f_J) = 1$. Also,

$$W_{\text{jump}}^{k, [p_0, p_+]}(\mathcal{L}_k; f_J) = \exp \left(- \int_{[p_0, p_0 + d]} \mathbf{H}_t^{k, k+1}(\underline{g}(x) - \mathcal{L}_k) dx \right).$$

For any $\mathbf{b} = (b_0, b_\pm) \in \mathbb{R}^3$, let

$$m_1(\mathbf{b}) := \frac{b_-(p_0 - q_1) + b_0(q_1 - p_-)}{p_0 - p_-}, \quad m_2(\mathbf{b}) := \frac{b_0(p_+ - q_2) + b_+(q_2 - p_0)}{p_+ - p_0}.$$

Conditioned on $J_k(p_0) = b_0$ and $J_k(p_\pm) = b_\pm$, $J_k(q_1)$ and $J_k(q_2)$ are independent. Let $\rho_{\sigma,m}(x) = (2\pi\sigma^2)^{-1/2}e^{-x^2/(2\sigma^2)}$ be the p.d.f. of a normal distribution with mean m and variance σ^2 . The p.d.f. of $J_k(q_1)$ is given by

$$\rho_{\sigma_1,m_1(\mathbf{b})}(x_1) =: \rho_{\sigma_1,\underline{m}_1}(x_1)f_1(x_1; \mathbf{b}).$$

The function $f_1(x_1; \mathbf{b})$ is defined through the last equality. Note that as $m_1(\mathbf{b}) \geq \underline{m}_1$, it is straightforward to check that $f_1(x_1; \mathbf{b})$ is non-decreasing in x_1 . Similarly, the p.d.f. of $J_k(q_2)$ is given by

$$\begin{aligned} & \rho_{\sigma_2,m_2(\mathbf{b})}(x_2) \left(\mathbb{E}_{\text{free}}^{1,1,(p_0,q_2),b_0,x_2} [W_{\text{jump}}^{k,[p_0,p_0+d]}] \right) / \mathbb{E}_{\text{free}}^{1,1,(p_0,p_+),b_0,b_+} [W_{\text{jump}}^{k,[p_0,p_0+d]}] \\ & =: \rho_{\sigma_2,\underline{m}_2}(x_2)f_2(x_2; \mathbf{b}). \end{aligned}$$

The function $f_2(x_2; \mathbf{b})$ is defined through the last equality and is non-decreasing provided $m_2(\mathbf{b}) \geq \underline{m}_2$. Write $\bar{Q}(\mathbf{b})$ for the joint p.d.f of $(J(p_0), J(p_\pm))$. Then

$$Q(x_1, x_2) = \int_{\mathbb{R}^3} \rho_{\sigma_1,\underline{m}_1}(x_1)\rho_{\sigma_2,\underline{m}_2}(x_2)f_1(x_1; \mathbf{b})f_2(x_2; \mathbf{b})\bar{Q}(\mathbf{b})d\mathbf{b}.$$

The following expression serves as a starting point for proving Proposition 3.6.

$$(8.1) \quad \mathbb{P}_J(J_k^{[q_1,q_2]} \in A') = \int_{\mathbb{R}^3} \int_{\mathbb{R}^2} F(x_1, x_2)\rho_{\sigma_1,\underline{m}_1}(x_1)\rho_{\sigma_2,\underline{m}_2}(x_2)f_1(x_1; \mathbf{b})f_2(x_2; \mathbf{b})\bar{Q}(\mathbf{b}) dx_1 dx_2 d\mathbf{b}.$$

Let $\underline{\mathbf{m}} := (\underline{m}, \underline{m}_\pm)$ and denote $\{\mathbf{b} \geq \underline{\mathbf{m}}\} := \{b_0 \geq \underline{m}_0, b_\pm \geq \underline{m}_\pm\}$. Because $F(x_1, x_2) \leq 1$, together with Lemma 8.1, we could bound $\mathbb{P}_J(J_k^{[q_1,q_2]} \in A')$ by

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{R}^2} F(x_1, x_2)\rho_{\sigma_1,\underline{m}_1}(x_1)\rho_{\sigma_2,\underline{m}_2}(x_2)f_1(x_1; \mathbf{b})f_2(x_2; \mathbf{b})\mathbb{1}_{\{\mathbf{b} \geq \underline{\mathbf{m}}\}}\bar{Q}(\mathbf{b})dx_1 dx_2 d\mathbf{b} + \mathbb{P}_J(\mathcal{Y}^c) \\ & \leq \sup_{\mathbf{b} \geq \underline{\mathbf{m}}} \int_{\mathbb{R}^2} F(x_1, x_2)\rho_{\sigma_1,\underline{m}_1}(x_1)\rho_{\sigma_2,\underline{m}_2}(x_2)f_1(x_1; \mathbf{b})f_2(x_2; \mathbf{b})dx_1 dx_2 + \mathbb{P}_J(\mathcal{Y}^c). \end{aligned}$$

From now on we fix $\mathbf{b} \geq \underline{\mathbf{m}}$. The estimates we derive below are uniform in any such choice of \mathbf{b} . By the Gibbs property, $F(x_1, x_2)$ can be rewritten as

$$\begin{aligned} F(x_1, x_2) &= \mathbb{E}_{\text{free}}^{k,k,(q_1,q_2),x_1,x_2} \left[\mathbb{1}\{B^{[q_1,q_2]} \in A\} W_{\text{jump}}^{k,[p_0,p_0+d]} \right] / \mathbb{E}_{\text{free}}^{k,k,(q_1,q_2),x_1,x_2} \left[W_{\text{jump}}^{k,[p_0,p_0+d]} \right] \\ &\leq \mathbb{P}_{\text{free}}^{k,k,(q_1,q_2),0,0}(\{B^{[q_1,q_2]} \in A\}) \left(\mathbb{E}_{\text{free}}^{k,k,(q_1,q_2),x_1,x_2} [W_{\text{jump}}^{k,[p_0,p_0+d]}] \right)^{-1} \\ &= \varepsilon \left(\mathbb{E}_{\text{free}}^{1,1,(q_1,q_2),x_1,x_2} [W_{\text{jump}}^{k,[p_0,p_0+d]}] \right)^{-1}. \end{aligned}$$

Here we have used $W_{\text{jump}}^{k,[p_0,p_0+d]} \leq 1$.

Let

$$\bar{F}(x_1, x_2) := \min \left\{ 1, \varepsilon \left(\mathbb{E}_{\text{free}}^{k,k,(q_1,q_2),x_1,x_2} [W_{\text{jump}}^{k,[p_0,p_0+d]}] \right)^{-1} \right\}.$$

Hence $F(x_1, x_2) \leq \bar{F}(x_1, x_2)$. Since $\bar{F}(x_1, x_2)$ is non-increasing in x_1 for fixed x_2 and $f_1(x_1; \mathbf{b})$ is non-decreasing, for any $x_2 \in \mathbb{R}$,

$$\begin{aligned}
& \int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x_1) \bar{F}(x_1, x_2) f_1(x_1; \mathbf{b}) dx_1 - \int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x_1) \bar{F}(x_1, x_2) dx_1 \\
&= \left(\int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x_1) \bar{F}(x_1, x_2) f_1(x_1; \mathbf{b}) dx_1 \right) \left(\int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x'_1) dx'_1 \right) \\
&\quad - \int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x_1) \bar{F}(x_1, x_2) dx_1 \left(\int_{\mathbb{R}} \rho_{\sigma_1, \underline{m}_1}(x'_1) f_1(x'_1; \mathbf{b}) dx'_1 \right) \\
&= 2^{-1} \int_{\mathbb{R}^2} \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_1, \underline{m}_1}(x'_1) (\bar{F}(x_1, x_2) - \bar{F}(x'_1, x_2)) (f_1(x_1; \mathbf{b}) - f_1(x'_1; \mathbf{b})) dx_1 dx'_1 \\
&\leq 0.
\end{aligned}$$

Since $\bar{F}(x_1, x_2)$ is non-increasing in x_2 for fixed x_1 and $f_2(x_2; \mathbf{b})$ is non-decreasing, we can repeat the same deduction to get

$$\begin{aligned}
& \int_{\mathbb{R}^2} \bar{F}(x_1, x_2) \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) f_1(x_1; \mathbf{b}) f_2(x_2; \mathbf{b}) dx_1 dx_2 \\
&\leq \int_{\mathbb{R}^2} \bar{F}(x_1, x_2) \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) f_2(x_2; \mathbf{b}) dx_1 dx_2 \\
&\leq \int_{\mathbb{R}^2} \bar{F}(x_1, x_2) \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) dx_1 dx_2 \\
&\leq \int_{\underline{m}_2 - \sigma_2 T^{3/2}}^{\infty} \int_{\underline{m}_1 - \sigma_1 T^{3/2}}^{\infty} \bar{F}(x_1, x_2) \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) dx_1 dx_2 + C e^{-C^{-1} T^3}.
\end{aligned}$$

We have used $\bar{F}(x_1, x_2) \leq 1$ and $\int_{-\infty}^{\underline{m}_i - \sigma_i T^{3/2}} \rho_{\sigma_i, \underline{m}_i}(x) dx \leq C e^{-C^{-1} T^3}$ in the last inequality.

Now it suffices to estimate $\left(\mathbb{E}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} [W_{\text{jump}}^{k, [p_0, p_0 + d]}] \right)^{-1}$. Let B be a Brownian bridge with $B(q_1) = x_1$ and $B(q_2) = x_2$. We start by considering an event which ensures $B(x) \geq \underline{g}(x)$ for $x \in [p_0, p_0 + d]$.

Lemma 8.2. *For E large enough, the following statement holds. Let $B : [q_1, q_2] \rightarrow \mathbb{R}$ be a continuous curve with $B(q_2) = x_2$. Suppose that*

- $x_2 \geq \underline{m}_2 - \sigma_2 T^{3/2}$,
- $B(p_0) \geq \underline{g}(p_0) + 3\Delta$,
- $\inf_{x \in [p_0, q_2]} B^{[p_0, q_2]}(x) > -\Delta$.

Then $B(x) \geq \underline{g}(x)$ in $[p_0, p_0 + d]$.

Proof. It is sufficient to prove the assertion for the case $x_2 = \underline{m}_2 - \sigma_2 T^{3/2}$ and $B(p_0) = \underline{g}(p_0) + 3\Delta$.

From Condition **C3** and (6.1), for all $x \in [p_0, p_0 + d]$ we have

$$\underline{g}(x) \leq \underline{g}(p_0) + \Delta.$$

Under the assumptions of this lemma, we bound the slope of the secant of B in $[p_0, q_2]$ as

$$\begin{aligned} \left| \frac{x_2 - B(p_0)}{q_2 - p_0} \right| &= \left| \frac{(\underline{m}_2 - \sigma_2 T^{3/2}) - (g(p_0) + 3\Delta)}{q_2 - p_0} \right| \\ &= \left| \frac{\text{Tent}(q_2) - \text{Tent}(p_0) - \sigma_2 T^{3/2} - (k+5)\Delta - k \log k}{q_2 - p_0} \right|. \end{aligned}$$

Using (6.10), $\sigma_2 \leq s^{1/2}$ and $q_2 - p_0 \geq 2^{-1}s$, the above is bounded by

$$2T + 2T^{3/2} + 2(k+5)\Delta + 2(k \log k)T^2.$$

For large enough T , it holds that

$$\left| \frac{x_2 - B(p_0)}{q_2 - p_0} \right| \leq 4T^{3/2}.$$

As a result, for $x \in [p_0, p_0 + d]$, we have

$$\begin{aligned} B(x) &= B(p_0) + B^{[p_0, q_2]}(x) + \frac{x_2 - B(p_0)}{q_2 - p_0}(x - p_0) \\ &\geq \underline{g}(p_0) + 3\Delta + (-\Delta) - 4T^{3/2}d \\ &\geq \underline{g}(x) + (\Delta - 4T^{3/2}d) \\ &= \underline{g}(x) + (8 \log T - 256T^{-5/2}(\log T)^2). \end{aligned}$$

Here we have used $\Delta = 8 \log T$ and $d = 64T^{-4}(\log T)^2$. For T large enough, we conclude that $B(x) \geq \underline{g}(x)$ for all $x \in [p_0, p_0 + d]$. \square

When $B(x) \geq g(x)$ in $[p_0, p_0 + d]$, we have

$$W_{\text{jump}}^{k, [p_0, p_0 + d]}(B) \geq e^{-d\mathbf{H}_t^{k, k+1}(0)} \geq 2^{-1}.$$

Therefore, provided $x_2 \geq \underline{m}_2 - \sigma_2 T^{3/2}$,

$$\begin{aligned} &\mathbb{E}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} [W_{\text{jump}}^{k, [p_0, p_0 + d]}] \\ &\geq 2^{-1} \mathbb{P}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} (B(p_0) \geq \underline{g}(p_0) + 3\Delta, \inf_{x \in [p_0, q_2]} B^{[p_0, q_2]}(x) > -\Delta) \\ &= 2^{-1} \mathbb{P}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} (B(p_0) \geq \underline{g}(p_0) + 3\Delta) \mathbb{P}_{\text{free}}(\inf_{x \in [p_0, q_2]} B^{[p_0, q_2]}(x) > -\Delta) \\ &\geq 4^{-1} \mathbb{P}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} (B(p_0) \geq \underline{g}(p_0) + 3\Delta). \end{aligned}$$

Under the law $\mathbb{P}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2}$, $B(p_0)$ is a normal distribution with variance σ_3^2 and mean

$$m_3(x_1, x_2) := \frac{x_1(q_2 - p_0) + x_2(p_0 - q_1)}{q_2 - q_1}.$$

Therefore,

$$\begin{aligned} \mathbb{E}_{\text{free}}^{k, k, (q_1, q_2), x_1, x_2} [W_{\text{jump}}^{[p_0, p_0 + d]}] &\geq 4^{-1} \int_{\underline{g}(p_0) + 3\Delta - m_3(x_1, x_2)}^{\infty} \rho_{\sigma_3, 0}(z_3) dz_3 \\ &=: 4^{-1} \nu_{\sigma_3, 0} [\underline{g}(p_0) + 3\Delta - m_3(x_1, x_2)]. \end{aligned}$$

This implies

$$\bar{F}(x_1, x_2) \leq 4\varepsilon \nu_{\sigma_3, 0} [\underline{g}(p_0) + 3\Delta - m_3(x_1, x_2)]^{-1}.$$

Thus

$$\begin{aligned} & \int_{\underline{m}_2 - \sigma_2 T^{3/2}}^{\infty} \int_{\underline{m}_1 - \sigma_1 T^{3/2}}^{\infty} \bar{F}(x_1, x_2) \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) dx_1 dx_2 \\ & \leq 4\varepsilon \int_{\underline{m}_2 - \sigma_2 T^{3/2}}^{\infty} \int_{\underline{m}_1 - \sigma_1 T^{3/2}}^{\infty} \rho_{\sigma_1, \underline{m}_1}(x_1) \rho_{\sigma_2, \underline{m}_2}(x_2) \nu_{\sigma_3, 0} [g(p_0) + 3\Delta - m_3(x_1, x_2)]^{-1} dx_1 dx_2. \end{aligned}$$

To simplify the expression, we perform a change of variables $z_i = x_i - \underline{m}_i$.

$$m_3(x_1, x_2) = \frac{z_1(q_2 - p_0) + z_2(p_0 - q_1)}{q_2 - q_1} + \frac{\text{Tent}(q_1)(q_2 - p_0) + \text{Tent}(q_2)(p_0 - q_1)}{q_2 - q_1} - \Delta_k.$$

From (6.10), the slope of Tent is bounded by $\pm 2T$. Hence

$$\begin{aligned} & \frac{\text{Tent}(q_1)(q_2 - p_0) + \text{Tent}(q_2)(p_0 - q_1)}{q_2 - q_1} \\ & \geq \frac{(\text{Tent}(p_0) - (p_0 - q_1)2T)(q_2 - p_0) + (\text{Tent}(p_0) - (q_2 - p_0)2T)(p_0 - q_1)}{q_2 - q_1} \\ & = \underline{g}(p_0) - 4\sigma_3^2 T. \end{aligned}$$

It follows that

$$m_3(x_1, x_2) \geq \frac{z_1(q_2 - p_0) + z_2(p_0 - q_1)}{q_2 - q_1} + \underline{g}(p_0) - 4\sigma_3^2 T - \Delta_k.$$

Take $M = \sigma_3^{-1}(4\sigma_3^2 T + \Delta_k + 3\Delta)$. In particular, $4\sigma_3^2 T + \Delta_k + 3\Delta = \sigma_3 M$. Then the upper bound becomes

(8.2)

$$4\varepsilon \int_{-\sigma_2 T^{3/2}}^{\infty} \int_{-\sigma_1 T^{3/2}}^{\infty} \rho_{\sigma_1, 0}(z_1) \rho_{\sigma_2, 0}(z_2) \cdot \left[\nu_{\sigma_3, 0} \left(-\frac{z_1(q_2 - p_0) + z_2(p_0 - q_1)}{q_2 - q_1} + \sigma_3 M \right) \right]^{-1} dz_1 dz_2.$$

Let N_1 and N_2 be two independent Gaussian random variables with mean zero and variances σ_1^2 and σ_2^2 respectively. Then

$$N_4 := \frac{(q_2 - p_0)N_1 + (p_0 - q_1)N_2}{q_2 - q_1}$$

is a Gaussian random variable with mean zero and variance

$$\sigma_4^2 := \left(\frac{q_2 - p_0}{q_2 - q_1} \right)^2 \sigma_1^2 + \left(\frac{p_0 - q_1}{q_2 - q_1} \right)^2 \sigma_2^2.$$

Moreover, as $N_1 \geq -\sigma_1 T^{3/2}$ and $N_2 \geq -\sigma_2 T^{3/2}$,

$$N_4 \geq - \left(\left(\frac{q_2 - p_0}{q_2 - q_1} \right) \sigma_1 + \left(\frac{p_0 - q_1}{q_2 - q_1} \right) \sigma_2 \right) T^{3/2}.$$

Before continuing the estimate, we give a simple bound to the above.

Lemma 8.3.

- (1) $4^{-1}\sigma_3^2 \leq \sigma_4^2 \leq \sigma_3^2$,
- (2) $\left(\frac{q_2 - p_0}{q_2 - q_1} \right) \sigma_1 + \left(\frac{p_0 - q_1}{q_2 - q_1} \right) \sigma_2 \leq 2\sigma_3$

Proof. (1) From a direct computation,

$$\begin{aligned} \sigma_4^2 &= \frac{(q_2 - p_0)^2}{(q_2 - q_1)^2} \times \frac{(p_0 - q_1)(q_1 - p_-)}{p_0 - p_-} + \frac{(p_0 - q_2)^2}{(q_2 - q_1)^2} \times \frac{(p_+ - q_2)(q_2 - p_0)}{p_+ - p_0} \\ &= \sigma_3^2 \left(\frac{q_1 - p_-}{p_0 - p_-} \times \frac{q_2 - p_0}{q_2 - q_1} + \frac{p_+ - q_2}{p_+ - p_0} \times \frac{p_0 - q_1}{q_2 - q_1} \right). \end{aligned}$$

Because $p_- < q_1 < p_0 < q_2 < p_+$, it holds that

$$\frac{q_1 - p_-}{p_0 - p_-} \leq 1, \quad \frac{p_+ - q_2}{p_+ - p_0} \leq 1.$$

Therefore,

$$\sigma_4^2 \leq \sigma_3^2 \left(\frac{q_2 - p_0}{q_2 - q_1} + \frac{p_0 - q_1}{q_2 - q_1} \right) = \sigma_3^2.$$

Let $\alpha = \frac{q_2 - p_0}{q_2 - q_1}$ and $\beta = \frac{p_0 - q_1}{q_2 - q_1}$. Because $q_2 - q_1 \leq \frac{3}{2}s$ and $p_0 - p_- \geq s$, it holds that

$$\frac{q_1 - p_-}{p_0 - p_-} = 1 - \frac{p_0 - q_1}{p_0 - p_-} \geq 1 - \frac{3}{2} \frac{p_0 - q_1}{q_2 - q_1} = 1 - \frac{3}{2} \beta.$$

Similarly,

$$\frac{p_+ - q_2}{p_+ - p_0} \geq 1 - \frac{3}{2} \alpha.$$

Together with $\alpha + \beta = 1$, it holds that

$$\begin{aligned} & \frac{q_1 - p_-}{p_0 - p_-} \times \frac{q_2 - p_0}{q_2 - q_1} + \frac{p_+ - q_2}{p_+ - p_0} \times \frac{p_0 - q_1}{q_2 - q_1} \\ & \geq \left(1 - \frac{3}{2}\beta\right) \alpha + \left(1 - \frac{3}{2}\alpha\right) \beta \\ & = 3(\alpha - 2^{-1})^2 + 4^{-1} \geq 4^{-1}. \end{aligned}$$

We conclude that $\sigma_4^2 \geq 4^{-1} \sigma_3^2$.

(2) By a direct computation,

$$\begin{aligned} & \left(\frac{q_2 - p_0}{q_2 - q_1} \right) \sigma_1 + \left(\frac{p_0 - q_1}{q_2 - q_1} \right) \sigma_2 \\ & = \left(\frac{(q_2 - p_0)(p_0 - q_1)}{q_2 - q_1} \right)^{1/2} \left(\left(\frac{(q_2 - p_0)(q_1 - p_-)}{(q_2 - q_1)(p_0 - p_-)} \right)^{1/2} + \left(\frac{(p_0 - q_1)(p_+ - q_2)}{(q_2 - q_1)(p_+ - p_0)} \right)^{1/2} \right) \end{aligned}$$

Using $\sigma_3^2 = \frac{(q_2 - p_0)(p_0 - q_1)}{q_2 - q_1}$, $\alpha = \frac{q_2 - p_0}{q_2 - q_1}$ and $\beta = \frac{p_0 - q_1}{q_2 - q_1}$, it becomes

$$\sigma_3 \left(\alpha^{1/2} \left(\frac{q_1 - p_-}{p_0 - p_-} \right)^{1/2} + \beta^{1/2} \left(\frac{p_+ - q_2}{p_+ - p_0} \right)^{1/2} \right)$$

Because $p_- < q_1 < p_0 < q_2 < p_+$, it holds that

$$\frac{q_1 - p_-}{p_0 - p_-} \leq 1, \quad \frac{p_+ - q_2}{p_+ - p_0} \leq 1.$$

Therefore,

$$\left(\frac{q_2 - p_0}{q_2 - q_1} \right) \sigma_1 + \left(\frac{p_0 - q_1}{q_2 - q_1} \right) \sigma_2 \leq \sigma_3 (\alpha^{1/2} + \beta^{1/2}) \leq 2\sigma_3.$$

□

From Lemma 8.3, (8.2) is bounded by

$$\begin{aligned} & 4\varepsilon \int_{-2\sigma_3 T^{3/2}}^{\infty} \rho_{\sigma_4, 0}(z_4) \cdot \left[\nu_{\sigma_3, 0}(-z_4 + \sigma_3 M) \right]^{-1} dz_4 \\ & = 4\varepsilon \int_{-\infty}^{2\sigma_3 T^{3/2}} \rho_{\sigma_4, 0}(z_4) \cdot \left[\nu_{\sigma_3, 0}(z_4 + \sigma_3 M) \right]^{-1} dz_4. \end{aligned}$$

Because $\left[\nu_{\sigma_3,0}(z_4 + \sigma_3 M)\right]^{-1}$ is non-decreasing in z_4 ,

$$\begin{aligned} & \int_{-\infty}^0 \rho_{\sigma_4,0}(z_4) \cdot \left[\nu_{\sigma_3,0}(z_4 + \sigma_3 M)\right]^{-1} dz_4 \leq 2^{-1} \left[\nu_{\sigma_3,0}(\sigma_3 M)\right]^{-1} \\ & \leq 2^{-1} Z^{-1} \int_0^{2\sigma_3 T^{3/2}} \rho_{\sigma_4,0}(z_4) \cdot \left[\nu_{\sigma_3,0}(z_4 + \sigma_3 M)\right]^{-1} dz_4, \end{aligned}$$

Here

$$Z = \int_0^{2\sigma_3 T^{3/2}} \rho_{\sigma_4,0}(z_4) dz_4.$$

By Lemma 8.3, for large enough T , it holds that

$$Z \geq \int_0^{2\sigma_4 T^{3/2}} \rho_{\sigma_4,0}(z_4) dz_4 \geq 4^{-1}.$$

As a result, (8.2) is bounded by

$$12\varepsilon \int_0^{2\sigma_3 T^{3/2}} \rho_{\sigma_4,0}(z_4) \cdot \left[\nu_{\sigma_3,0}(z_4 + \sigma_3 M)\right]^{-1} dz_4.$$

Next, we replace $\nu_{\sigma_3,0}(z_4 + \sigma_3 M)$ with $\rho_{\sigma_3,0}(z_4 + \sigma_3 M)$. We first show that $M \geq 1$. Recall that $M = \sigma_3^{-1}(4\sigma_3^2 T + \Delta_k + 3\Delta)$. Because $T \geq 10$,

$$M \geq 4\sigma_3 T + \sigma_3^{-1}\Delta \geq 4(T\Delta)^{1/2} \geq 1.$$

For $z \geq 1$, [Wil91, Section 14.8] we have

$$\nu_{1,0}(z) \geq \frac{z}{z^2 + 1} (2\pi)^{-1/2} e^{-z^2/2} \geq (2z)^{-1} (2\pi)^{-1/2} e^{-z^2/2} = (2z)^{-1} \rho_{1,0}(z).$$

By the Brownian scaling, for $z \geq \sigma$ it holds that

$$\nu_{\sigma,0}(z) = \nu_{1,0}(\sigma^{-1}z) \geq \sigma(2z)^{-1} \rho_{1,0}(\sigma^{-1}z) = \sigma^2(2z)^{-1} \rho_{\sigma,0}(z).$$

Because $M \geq 1$ and $z_4 \geq 0$, we deduce

$$\nu_{\sigma_3,0}(z_4 + \sigma_3 M) \geq 2^{-1} \sigma_3^2 (z_4 + \sigma_3 M)^{-1} \rho_{\sigma_3,0}(z_4 + \sigma_3 M).$$

Hence (8.2) is bounded by

$$\begin{aligned} & 24\varepsilon \sigma_3^{-2} \int_0^{2\sigma_3 T^{3/2}} \rho_{\sigma_4,0}(z_4) \rho_{\sigma_3,0}(z_4 + \sigma_3 M)^{-1} (z_4 + \sigma_3 M) dz_4 \\ & = 24\varepsilon \int_0^{2\sigma_3 T^{3/2}} \rho_{\sigma_4,0}(z_4) \rho_{\sigma_3,0}(z_4)^{-1} e^{M\sigma_3^{-1}z_4 + M^2/2} (\sigma_3^{-1}z_4 + M) \sigma_3^{-1} dz_4. \end{aligned}$$

Because $4^{-1}\sigma_3^2 \leq \sigma_4^2 \leq \sigma_3^2$,

$$\rho_{\sigma_4,0}(z_4) \rho_{\sigma_3,0}(z_4)^{-1} = \sigma_4^{-1} \sigma_3 \exp(-2^{-1}(\sigma_4^{-2} - \sigma_3^{-2})z_4^2) \leq 2.$$

(8.2) is bounded by

$$\begin{aligned} & 48\varepsilon \int_0^{2\sigma_3 T^{3/2}} e^{M\sigma_3^{-1}z_4 + M^2/2} (\sigma_3^{-1}z_4 + M) \sigma_3^{-1} dz_4 \\ & = 48\varepsilon \times e^{M^2/2} \int_0^{2T^{3/2}} e^{Mw_4} (w_4 + M) dw_4. \\ & \leq 48\varepsilon \times 2T^{3/2} (2T^{3/2} + M) e^{2MT^{3/2} + M^2/2}. \end{aligned}$$

We proceed to bound M . Because $\frac{3}{2}s \geq q_2 - q_1$ and $q_2 - p_0, p_0 - q_1 \geq \frac{1}{2}s$, it holds that $\frac{3}{8}s \geq \sigma_3^2 \geq \frac{1}{6}s$. Therefore $M = \sigma_3^{-1}(4\sigma_3^2 T + \Delta_k + 3\Delta) \leq DT$. We arrive at

$$(8.2) \leq \varepsilon \times De^{DT^{5/2}}.$$

In conclusion,

$$\sup_{\mathbf{b} \geq \underline{\mathbf{m}}} \int_{\mathbb{R}^2} F(x_1, x_2) \rho_{\sigma_1, \underline{\mathbf{m}}_1}(x_1) \rho_{\sigma_2, \underline{\mathbf{m}}_2}(x_2) f_1(x_1; \mathbf{b}) f_2(x_2; \mathbf{b}) dx_1 dx_2 \leq \varepsilon \times De^{DT^{5/2}} + Ce^{-C^{-1}T^3}.$$

We are ready to complete the proof of Proposition 3.6.

Proof of Proposition 3.6. From the above discussion,

$$\begin{aligned} \mathbb{P}_J(J_k^{[q_1, q_2]} \in A') &\leq \sup_{\mathbf{b} \geq \underline{\mathbf{m}}} \int_{\mathbb{R}^2} F(x_1, x_2) \rho_{\sigma_1, \underline{\mathbf{m}}_1}(x_1) \rho_{\sigma_2, \underline{\mathbf{m}}_2}(x_2) f_1(x_1; \mathbf{b}) f_2(x_2; \mathbf{b}) dx_1 dx_2 + \mathbb{P}_J(Y^c) \\ &\leq \varepsilon \times De^{DT^{5/2}} + Ce^{-C^{-1}T^3} + De^{-D^{-1}T^4}. \end{aligned}$$

Recall that $T = E \max\{(\log \varepsilon^{-1})^{1/3}, s\}$. Therefore by taking E large enough,

$$\mathbb{P}_J(J_k^{[0, s]} \in A) = \mathbb{P}_J(J_k^{[q_1, q_2]} \in A') \leq \varepsilon \times D \exp\left(D (\log \varepsilon^{-1})^{5/6}\right).$$

□

APPENDIX A. CONCAVE FUNCTIONS

In this section we record basic properties of concave functions. Throughout this section, $h : [\ell, r] \rightarrow \mathbb{R}$ is a continuous concave function.

For $x \in (\ell, r)$, the left/right derivative of h at x is defined by

$$h'_\pm(x) := \lim_{y \rightarrow x^\pm} \frac{h(y) - h(x)}{y - x}.$$

We may extend $h'_+(x)$ to $x = \ell$ by allowing it to take value ∞ . We may also extend $h'_-(r)$ to $x = r$ by allowing it to take value $-\infty$. $h'_\pm(x)$ are monotone non-increasing where they are defined. Moreover, for all $x \in (\ell, r)$, $h'_-(x) \geq h'_+(x)$.

Lemma A.1. $h'_+(x) : [\ell, r) \rightarrow \mathbb{R} \cup \{\infty\}$ is right continuous and $h'_-(x) : (\ell, r] \rightarrow \mathbb{R} \cup \{-\infty\}$ is left continuous.

Proof. Fix $x_0 \in [\ell, r)$ and take $x_j \searrow x_0$. Given $y > x_0$, we have for $x_j < y$,

$$\frac{h(y) - h(x_j)}{y - x_j} \leq h'_+(x_j).$$

By the continuity of $h(x)$ at x_0 ,

$$\frac{h(y) - h(x_0)}{y - x_0} \leq \liminf_{j \rightarrow \infty} h'_+(x_j).$$

Hence $h'_+(x_0) \leq \liminf_{j \rightarrow \infty} h'_+(x_j)$. By the monotonicity of $h'_+(x)$, $h'_+(x_0) \geq \limsup_{j \rightarrow \infty} h'_+(x_j)$. This shows $h'_+(x)$ is right continuous. Left continuity of $h'_+(x)$ can be derived similarly. □

Define the extreme points of $h(x)$ as

$$\text{xExt}(h) := \{\ell, r\} \cup \left\{ x \in (\ell, r) \mid h|_{[x-\delta, x+\delta]} \text{ is not linear for any } \delta > 0 \right\}.$$

For any $x_0 \in [\ell, r] \setminus \text{xExt}(h)$, by considering the largest interval containing x_0 in which $h(x)$ is linear, we get the following lemma.

Lemma A.2. For all $x_0 \in [\ell, r] \setminus \text{xExt}(h)$, there exist $x_1, x_2 \in \text{xExt}(h)$ such that $x_0 \in (x_1, x_2)$ and $h|_{[x_1, x_2]}$ is linear.

As a direct corollary, we have

Corollary A.3. Let $\bar{h} : [\ell, r] \rightarrow \mathbb{R}$ be another concave function. Suppose $\bar{h}(x) \geq h(x)$ for all $x \in \text{xExt}(h)$. Then $\bar{h}(x) \geq h(x)$ for all $x \in [\ell, r]$.

Let $P \subset \text{xExt}(h)$ be a subset that satisfies the following properties

- $\ell, r \in P$
- For any $y \in \text{xExt}(h) \cap [\ell, r]$, there exists $x \in P$ such that $|x - y| < s$.

Lemma A.4. Suppose that $|h'_\pm| \leq m$ for some $m \geq 0$. Further assume that $\bar{h} : [\ell, r] \rightarrow \mathbb{R}$ is a concave function with $\bar{h}(x) \geq h(x)$ for all $x \in P \cap [\ell, r]$. Then $\bar{h}(x) + 2ms \geq h(x)$ for all $x \in [\ell, r]$.

Proof. It suffices to show that $\bar{h}(x) + 2m \geq h(x)$ for all $x \in \text{xExt}(h)$. Given $x \in \text{xExt}(h)$. If $x \in P$, by the assumption $\bar{h}(x) \geq h(x)$. Suppose $x \in \text{xExt}(h) \setminus P$. There exists $y \in P$ with $|x - y| < d$. We further assume that $x < y$. Because $\bar{h}(y) \geq h(y)$, $\bar{h}(\ell) \geq h(\ell)$ and \bar{h} is concave, we have

$$\begin{aligned} \bar{h}(x) &\geq \frac{(y-x)\bar{h}(\ell) + (x-\ell)\bar{h}(y)}{y-\ell} \geq \frac{(y-x)h(\ell) + (x-\ell)h(y)}{y-\ell} \\ &= h(y) + \frac{h(\ell) - h(y)}{y-\ell}(y-x) \geq h(y) - ms. \end{aligned}$$

Together with $h(x) \leq h(y) + ms$, we obtain that $\bar{h}(x) \geq h(x) - 2ms$. The argument for $y < x$ is similar. The proof is finished. \square

Lemma A.5. Let $m \in \mathbb{R}$. Assume that $\{x \in [\ell, r] \mid h'_+(x) \leq m\}$ is non-empty. Then

$$x_1 := \inf\{x \in [\ell, r] \mid h'_+(x) \leq m\} \in \text{xExt}(h)$$

and $h'_+(x_1) \leq m$. Similarly, assume $\{x \in [\ell, r] \mid h'_-(x) \geq m\}$ is non-empty. Then

$$x_2 := \sup\{x \in [\ell, r] \mid h'_-(x) \geq m\} \in \text{xExt}(h)$$

and $h'(x_2) \geq m$.

Proof. $h'_+(x_1) \leq m$ and $h'_-(x_2) \geq m$ follows from one-sided continuity of h'_\pm . If $x_1 \notin \text{xExt}(h)$, then h is linear around x_1 which contradicts the definition of x_1 . The proof for x_2 is similar. \square

Fix $m \in \mathbb{R}$ and $[\ell', r'] \subset (\ell, r)$. Define

$$\mathbf{l}(h) := \begin{cases} \inf_{r'}\{x \in [\ell', r'] \mid h'_+(x) \leq m\} & \{x \in [\ell', r'] \mid h'_+(x) \leq m\} \neq \emptyset, \\ & \{x \in [\ell', r'] \mid h'_+(x) \leq m\} = \emptyset. \end{cases}$$

Similarly,

$$\mathbf{r}(h) := \begin{cases} \sup_{\ell'}\{x \in [\ell', r'] \mid h'_-(x) \geq m\} & \{x \in [\ell', r'] \mid h'_-(x) \geq m\} \neq \emptyset, \\ & \{x \in [\ell', r'] \mid h'_-(x) \geq m\} = \emptyset. \end{cases}$$

The following lemma shows that \mathbf{l} and \mathbf{r} are semi-continuous with respect to the uniform topology.

Lemma A.6. Given a sequence of concave functions h_i converging to h_0 uniformly on $[\ell, r]$, we have

$$\liminf_{i \rightarrow \infty} \mathbf{l}(h_i) \geq \mathbf{l}(h_0), \quad \limsup_{i \rightarrow \infty} \mathbf{r}(h_i) \leq \mathbf{r}(h_0).$$

Proof. We give the proof for \mathbf{l} . The argument for \mathbf{r} is similar. We assume $\mathbf{l}(h_0) > \ell'$ otherwise the assertion is clearly true. For any $\ell' \leq x_0 < \mathbf{l}(h_0)$, we have $(h_0)'_+(x_0) > m$. Hence there exists $\varepsilon > 0$ such that

$$\varepsilon^{-1}(h_0(x_0 + \varepsilon) - h_0(x_0)) > m.$$

Because h_i converges to h_0 uniformly, for i large enough,

$$\varepsilon^{-1}(h_i(x_0 + \varepsilon) - h_i(x_0)) > m.$$

By the concavity of h_i , $(h_i)'_+(x_0) > m$. For such i , $\mathbf{l}(h_i) \geq x_0$. Hence

$$\liminf_{i \rightarrow \infty} \mathbf{l}(h_i) \geq x_0.$$

Because the above holds for all $\ell' \leq x_0 < \mathbf{l}(h_0)$, we have

$$\liminf_{i \rightarrow \infty} \mathbf{l}(h_i) \geq \mathbf{l}(h_0).$$

The proof is finished. \square

Let $g : [\ell, r] \rightarrow \mathbb{R}$ be an upper semi-continuous function. Let $h(x) : [\ell, r] \rightarrow \mathbb{R}$ be the concave majorant of g in $[\ell, r]$. In other words, for any $x_0 \in [\ell, r]$

$$h(x_0) := \inf \{ax_0 + b \mid ax + b \geq g(x) \text{ for all } x \in [\ell, r]\}.$$

We note that $h(x)$ is continuous in $[\ell, r]$.

Lemma A.7. *For all $x \in \text{xExt}(h) \setminus \{\ell, r\}$, we have $h(x) = g(x)$.*

Proof. Suppose that $x_0 \in \text{xExt}(h) \setminus \{\ell, r\}$ and that $h(x_0) - g(x_0) > 0$. Let $L_{\pm}(x) = h'_{\pm}(x_0)(x - x_0) + h(x_0)$. We have $L_{\pm}(x) \geq h(x) \geq g(x)$ for all $x \in [\ell, r]$. Suppose that $L_+(x) > g(x)$ for all $x \in (x_0, r]$. By the upper semi-continuity of $g(x)$,

$$\varepsilon = \inf_{[x_0, r]} (L_+(x) - g(x)) > 0.$$

Then for all $x \in [\ell, r]$,

$$L_+(x) - \frac{\varepsilon}{r - x_0}(x - x_0) \geq g(x).$$

Hence $L_+(x) - \frac{\varepsilon}{r - x_0}(x - x_0) \geq h(x)$. This implies $h'_+(x_0) - \frac{\varepsilon}{r - x_0} \geq h'_+(x_0)$ which is a contradiction. Therefore there exists $x_+ \in (x_0, r]$ such that $L_+(x_+) = g(x_+)$. Similarly, there exists $x_- \in [\ell, x_0)$ such that $L_-(x_-) = g(x_-)$. Because $x_0 \in \text{xExt}(h)$, we must have $h'_+(x_0) < h'_-(x_0)$. Take $L_0(x) = \frac{1}{2}(h'_+(x_0) + h'_-(x_0))(x - x_0) + h(x_0)$. At $x = x_0$, $L_0(x_0) = h(x_0) > g(x_0)$. For $x \in (x_0, r]$, $L_0(x) > L_+(x) \geq g(x)$. For $x \in [\ell, x_0)$, $L_0(x) > L_-(x) \geq g(x)$. In particular,

$$\varepsilon_0 = \inf_{x \in [\ell, r]} L_0(x) - g(x) > 0.$$

Then $h(x) \leq L_0(x) - \varepsilon_0$, which is a contradiction at $x = x_0$. \square

Next, we consider the situation that $h(x)$ or $g(x)$ is close to a parabola $-2^{-1}x^2$.

Lemma A.8. *Let $\delta > 0$ be a positive number. Let $h : [\ell, r] \rightarrow \mathbb{R}$ be a continuous concave function with $|h(x) + 2^{-1}x^2| \leq 4^{-1}\delta^2$. Then for all $x \in [\ell + \delta, r - \delta]$, we have*

$$(A.1) \quad -x - \delta \leq h'_+(x) \leq h'_-(x) \leq -x + \delta.$$

Proof. Let $x \in [\ell + \delta, r - \delta]$. By the concavity of h , we have

$$h'_+(x) \geq \frac{h(x) - h(x - \delta)}{\delta} \geq \delta^{-1} ((2^{-1}x^2 - 4^{-1}\delta^2) - (2^{-1}(x - \delta)^2 + 4^{-1}\delta^2)) = -x - \delta.$$

Similarly,

$$h'_-(x) \leq \frac{h(x+\delta) - h(x)}{\delta} \leq \delta^{-1} ((2^{-1}(x+\delta)^2 + 4^{-1}\delta^2) - (2^{-1}x^2 - 4^{-1}\delta^2)) = -x + \delta.$$

□

Lemma A.9. *Let $g : [\ell, r] \rightarrow \mathbb{R}$ be a function with $|g(x) + 2^{-1}x^2| \leq 4^{-1}\delta^2$. Take $x_0 \in [\ell + 3\delta, r - 3\delta]$. Let h_1 and h_2 be the concave majorant of $g(x)$ in $[\ell, r]$ and $[x_0 - 3\delta, x_0 + 3\delta]$ respectively. Then $h_1(x_0) = h_2(x_0)$.*

Proof. Directly from the definition, we have $h_1(x_0) \geq h_2(x_0)$. Consider the line

$$L(x) := h_2(x_0) + (h'_2)_+(x_0)(x - x_0).$$

It suffices to show that $L(x) \geq g(x)$ for all $x \in [\ell, r]$. Because $|h_2(x) + 2^{-1}x^2| \leq 4^{-1}\delta^2$ for all $x \in [x_0 - \delta, x_0 + \delta]$, from Lemma A.8 we have

$$|(h'_2)_+(x_0) + x_0| \leq \delta.$$

As a result, for all $x \geq x_0$, we have

$$\begin{aligned} L(x) - g(x) &\geq [-2^{-1}x_0^2 - 4^{-1}\delta^2 + (-x_0 + \delta)(x - x_0)] - [-2^{-1}x^2 + 4^{-1}\delta^2] \\ &= 2^{-1}(x - x_0)^2 - \delta(x - x_0) - 2^{-1}\delta^2. \end{aligned}$$

In particular, for $x \geq x_0 + 3\delta$, we have $L(x) \geq g(x)$. Similar argument can show that $L(x) \geq g(x)$ for all $x \leq x_0 - 3\delta$. Together with $L(x) \geq h_2(x) \geq f(x)$ for all $x \in [x_0 - 3\delta, x_0 + 3\delta]$, we conclude that $L(x) \geq f(x)$ for all $x \in [\ell, r]$. Thus

$$h_2(x_0) = L(x_0) \geq h_1(x_0).$$

The proof is finished. □

Corollary A.10. *Let $\ell < \ell' < r' < r$ be real numbers and $g : [\ell, r] \rightarrow \mathbb{R}$ be a function with $|g(x) + 2^{-1}x^2| \leq 4^{-1}\delta^2$. Let h_1 and h_2 be the concave majorant of $g(x)$ in $[\ell, r]$ and $[\ell', r']$ respectively. Then $h_1(x) = h_2(x)$ for all $x \in [\ell' + 3\delta, r' - 3\delta]$.*

APPENDIX B. MISCELLANEOUS

Lemma B.1. *For any $k \in \mathbb{N}$ and $(y_1, y_2, \dots, y_k, y_{k+1}) \in \mathbb{R}^{k+1}$,*

$$-\sum_{j=1}^k e^{y_{j+1} - y_j} \leq -k e^{(y_{k+1} - y_1)/k}.$$

Proof. The assertion holds clearly when $k = 1$. Assume $k \geq 2$ and the assertion holds for $k - 1$. Then

$$-\sum_{j=1}^k e^{y_{j+1} - y_j} = -\sum_{j=1}^{k-1} e^{y_{j+1} - y_j} - e^{y_{k+1} - y_k} \leq -(k-1)e^{(y_k - y_1)/(k-1)} - e^{y_{k+1} - y_k}.$$

It is straightforward to show that

$$-(k-1)e^{(y_k - y_1)/(k-1)} - e^{y_{k+1} - y_k} \leq -k e^{(y_{k+1} - y_1)/k}$$

and the inequality is achieved when $y_k = \frac{1}{k}y_1 + \frac{k-1}{k}y_{k+1}$. The proof is finished. □

Recall that $\mathbf{H}_t(x) = e^{t^{1/3}x}$. For integers $1 \leq j < k$, define

$$(B.1) \quad \mathbf{H}_t^{j,k}(x) := (k-1)^{-1} e^{t^{1/3}x/(k-j)}.$$

Lemma B.2. For any $k \in \mathbb{N}$, and $(y_1, y_2, \dots, y_{k+1}) \in \mathbb{R}^{k+1}$, we have

$$-\sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) \leq -\sum_{j=1}^k \mathbf{H}_t^{j,k+1}(y_{k+1} - y_j).$$

Proof. When $k = 1$, $\mathbf{H}_t^{1,2}(x) = \mathbf{H}_t(x)$ and the assertion clearly holds. Assume $k \geq 2$ and the assertion holds for $k - 1$.

$$\begin{aligned} -\sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) &= -\frac{1}{k} \sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) - \frac{k-1}{k} \sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) \\ &\leq -\frac{1}{k} \sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) - \frac{k-1}{k} \sum_{j=2}^k \mathbf{H}_t(y_{j+1} - y_j). \end{aligned}$$

By Lemma B.1,

$$-k^{-1} \sum_{j=1}^k \mathbf{H}_t(y_{j+1} - y_j) \leq -k \mathbf{H}_t^{1,k+1}(y_{k+1} - y_1) \leq -\mathbf{H}_t^{1,k+1}(y_{k+1} - y_1).$$

For $j \in [1, k-1]_{\mathbb{Z}}$, let $z_j = y_{j+1}$. Then by the induction hypothesis,

$$\begin{aligned} -\frac{k-1}{k} \sum_{j=2}^k \mathbf{H}_t(y_{j+1} - y_j) &= -\frac{k-1}{k} \sum_{j=1}^{k-1} \mathbf{H}_t(z_{j+1} - z_j) \leq -\frac{k-1}{k} \sum_{j=1}^{k-1} \mathbf{H}_t^{j,k}(z_k - z_j) \\ &= -\frac{k-1}{k} \sum_{j=2}^k \mathbf{H}_t^{j-1,k}(y_{k+1} - y_j) = -\sum_{j=2}^k \mathbf{H}_t^{j,k+1}(y_{k+1} - y_j). \end{aligned}$$

The proof is finished. \square

Lemma B.3. Fix $p_- < p < p_+$. Let $\mathring{L} : [p_-, p_+] \rightarrow \mathbb{R}$ be a continuous function which is linear in $[p_-, p]$ and $[p, p_+]$. Fix $q \in (p_-, p_+)$ and $b \in \mathbb{R}$. Let L be a continuous function which is linear in $[p_-, q]$ and $[q, p_+]$ with $L(p_{\pm}) = \mathring{L}(p_{\pm})$ and $L(q) = b$. Define

$$\mathring{m}_{\pm} := \frac{\mathring{L}(p) - \mathring{L}(p_{\pm})}{p - p_{\pm}}, \quad m_{\pm} := \frac{L(q) - \mathring{L}(p_{\pm})}{q - p_{\pm}}.$$

Then for all $x \in [p_-, p_+]$,

$$(B.2) \quad \mathring{L}(x) - L(x) \leq |L(p) - b| + |p - q| \max\{|\mathring{m}_-|, |\mathring{m}_+|\}$$

Furthermore, suppose $|p - q| \leq 2^{-1} \min\{p - p_-, p_+ - p\}$. Then

$$|m_- - \mathring{m}_-| \leq 2 \left| \frac{b - L(p)}{p - p_-} \right| + 2 \left| \frac{p - q}{p - p_-} \right| |\mathring{m}_-|.$$

$$|m_+ - \mathring{m}_+| \leq 2 \left| \frac{b - L(p)}{p_+ - p} \right| + 2 \left| \frac{p - q}{p_+ - p} \right| |\mathring{m}_+|.$$

Proof. Without loss of generality, we assume that $q < p$. It suffice to prove (B.2) at $x = p$ and $x = q$. For $x = q$,

$$\mathring{L}(q) - L(q) = [\mathring{L}(p) - \mathring{m}_-(p - q)] - b \leq |\mathring{L}(p) - b| + |p - q| |\mathring{m}_-|.$$

For $x = p$,

$$\begin{aligned}\dot{L}(p) - L(p) &= \dot{L}(p) - \left[\frac{(p_+ - p)b + (p - q)\dot{L}(p_+)}{p_+ - q} \right] \\ &= \frac{p_+ - p}{p_+ - q} (\dot{L}(p) - b) + (p - q) \frac{p_+ - p}{p_+ - q} \dot{m}_+ \\ &\leq |\dot{L}(p) - b| + |p - q| |\dot{m}_+|.\end{aligned}$$

This shows (B.2).

Under the assumption that $|p - q| \leq 2^{-1} \min\{p - p_-, p_+ - p\}$,

$$\begin{aligned}|m_- - \dot{m}_-| &= \left| \frac{b - \dot{L}(p_-)}{q - p_-} - \frac{\dot{L}(p) - \dot{L}(p_-)}{p - p_-} \right| \\ &= \left| \frac{b - \dot{L}(p)}{q - p_-} + \frac{(p - q)(\dot{L}(p) - \dot{L}(p_-))}{(q - p_-)(p - p_-)} \right| \\ &\leq 2 \left| \frac{b - L(p)}{p - p_-} \right| + 2 \left| \frac{p - q}{p - p_-} \times \frac{\dot{L}(p) - \dot{L}(p_-)}{p - p_-} \right| \\ &= 2 \left| \frac{b - L(p)}{p - p_-} \right| + 2 \left| \frac{p - q}{p - p_-} \right| |\dot{m}_-|.\end{aligned}$$

The bound for $|m_+ - \dot{m}_+|$ is similar. □

Lemma B.4. *Let $x_1 < x_2 \in \mathbb{R}$, $b_1, b_2 \in \mathbb{R}$ and $m > 0$. Assume that $|b_2 - b_1| \leq m(x_2 - x_1)$. Let*

$$\begin{aligned}x_0 &= 2^{-1}(x_1 + x_2) + (2m)^{-1}(b_2 - b_1) \in [x_1, x_2], \\ b_0 &= 2^{-1}(b_1 + b_2) + 2^{-1}m(x_2 - x_1).\end{aligned}$$

The point (x_0, b_0) is determined by

$$\frac{b_0 - b_1}{x_0 - x_1} = m, \quad \frac{b_2 - b_0}{x_2 - x_0} = -m.$$

Then

$$\mathbb{P}(B(x_0) \geq b_0 \mid B(x_1) = b_1, B(x_2) = b_2) \geq \mathbb{P}(N \geq (x_2 - x_1)^{1/2}m).$$

Proof. Then mean and variance of $B(x_0)$ is given by

$$\begin{aligned}M &= 2^{-1}(b_2 + b_1) + (2m(x_2 - x_1))^{-1}(b_2 - b_1)^2, \\ \sigma^2 &= (4m^2)^{-1}((x_2 - x_1)m^2 - (x_2 - x_1)^{-1}(b_2 - b_1)^2).\end{aligned}$$

Then

$$\sigma^{-1}(b_0 - M) = ((x_2 - x_1)m^2 - (x_2 - x_1)^{-1}(b_2 - b_1)^2)^{1/2} \leq (x_2 - x_1)^{1/2}m. \quad \square$$

REFERENCES

- [BCD21] G. Barraquand, I. Corwin and E. Dimitrov. Spatial tightness at the edge of Gibbsian line ensembles. *arxiv:2101.03045*.
- [BGH] R. Basu, S. Ganguly and A. Hammond. Fractal geometry of Airy₂ processes coupled via the Airy sheet. *Ann. Probab., to appear*.
- [CD18] I. Corwin and E. Dimitrov. Transversal fluctuations of the ASEP, stochastic six vertex model, and Hall-Littlewood Gibbsian line ensembles. *Comm. Math. Phys.*, 363(2) (2018).
- [CG1] I. Corwin and P. Ghosal. Lower tail of the KPZ equation. *Duke Math. J.*, 169(7) (2020).

- [CG2] I. Corwin and P. Ghosal. KPZ equation tails for general initial data. *Electron. J. Probab.* 25 (2020).
- [CH14] I. Corwin and A. Hammond. Brownian Gibbs property for Airy line ensembles. *Invent. Math.*, 195 (2014).
- [CH16] I. Corwin and A. Hammond. KPZ Line ensemble. *Probab. Theory Relat. Fields*, 166 (2016).
- [CHH] J. Calvert, A. Hammond and M. Hegde. Brownian structure in the KPZ fixed point. *arXiv:1912.00992*.
- [Cor] I. Corwin. The Kardar-Parisi-Zhang equation and universality class. *Random Matrices Theory Appl.*, 1(1), 76 (2012).
- [DM] E. Dimitrov and K. Matetski. Characterization of Brownian Gibbsian line ensembles. *arxiv:2002.00684*.
- [DOV] D. Dauvergne, J. Ortmann and B. Virág. The directed landscape. *arxiv:1812.00309*.
- [DV] D. Dauvergne and B. Virág. Bulk properties of the Airy line ensemble. *Ann. Probab.* 49(4) (2021).
- [Ham1] A. Hammond. Exponents governing the rarity of disjoint polymers in Brownian last passage percolations. *Proc. London Math. Soc.* 120(3) (2020).
- [Ham2] A. Hammond. Modulus of continuity of polymer weight profiles in Brownian last passage percolation. *Ann. Probab.* 47(6) (2019).
- [Ham3] A. Hammond. A patchwork quilt sewn from Brownian fabric: regularity of polymer weight profiles in Brownian last passage percolation. *Forum Math. Pi*, 7, e2 (2019).
- [Ham4] A. Hammond. Brownian regularity for the Airy line ensemble, and multi-polymer watermelons in Brownian last passage percolation. *Mem. Amer. Math. Soc.*, to appear.
- [Jan] S. Janson, S. Gaussian Hilbert spaces. *Cambridge Texts in Mathematics 129*. Cambridge Univ. Press, Cambridge.
- [KPZ] K. Kardar, G. Parisi and Y.Z. Zhang. Dynamic scaling of growing interfaces. *Phys. Rev. Lett.*, 56 (1986).
- [KS] I. Karatzas and S. Shreve. Brownian motion and stochastic calculus. *Volume 113 of Graduate Texts in Mathematics*. Springer (1988).
- [LW] C.H. Lun and J. Warren. Continuity and strict positivity of the multi-layer extension of the stochastic heat equation. *Electron. J. Probab.* Volume 25 (2020).
- [Mue] C. Mueller. On the support of solutions to the heat equation with noise. *Stochastics* 37(4) (1991).
- [OW] N. O’Connell and J. Warren. A Multi-Layer Extension of the Stochastic Heat Equation. *Commun. Math. Phys.* 341(1) (2016).
- [QS15] J. Quastel and H. Spohn. The one-dimensional KPZ equation and its universality class. *J. Statist. Phys.*, 160(4) (2015).
- [QS20] J. Quastel and S. Sarkar. Convergence of exclusion processes and KPZ equation to the KPZ fixed point. *arXiv:2008.06584*.
- [Vir] B. Virág. The heat and the landscape I. *arxiv.org/abs/2008.07241*.
- [Wu19] Tightness of discrete Gibbsian line ensembles with exponential interaction Hamiltonians. *arxiv:1909.00946*.
- [Wu21] Tightness and local Fluctuation of the KPZ line ensemble. *In progress*.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO IL, 60637
 Email address: xuanw@uchicago.edu