# Anisotropic percolation in high dimensions: the non-oriented case

Pablo A. Gomes\* Alan Pereira<sup>†</sup> Remy Sanchis<sup>‡</sup>

#### Abstract

We consider inhomogeneous non-oriented Bernoulli bond percolation on  $\mathbb{Z}^d$ , where each edge has a parameter, depending on its direction. We prove that, under certain conditions, if the sum of the parameters is strictly greater than 1/2, we have percolation for sufficiently high dimensions. The main tool is a dynamical coupling between the models in different dimensions with different set of parameters.

**Keywords:** anisotropic percolation; high dimensional systems; phase diagram; meanfield

**AMS-subject:** 60K35; 82B43

# 1 Introduction

The theory of bond percolation on  $\mathbb{Z}^d$  originated in [1]. In this model, each edge is declared open, independently of the others, with probability p, and closed otherwise. The primary question is to know if there is an infinite open connected component for a given value of p, in which case we say that *percolation* occurs. The existence of a non-trivial phase transition for  $d \geq 2$  was established in this seminal article with a *critical parameter*  $p_c(\mathbb{Z}^d) \in (0,1)$ , such that percolation occurs for  $p > p_c(\mathbb{Z}^d)$  and does not occur for  $p < p_c(\mathbb{Z}^d)$ .

It is well known that for d=2,  $p_c(\mathbb{Z}^d)=1/2$  (see [9]). Although the precise value of  $p_c(\mathbb{Z}^d)$  is unknown when  $d\geq 3$ , the asymptotic behavior  $2dp_c(\mathbb{Z}^d)=1+o(1/d)$  was obtained independently in [11] and [5]. Since  $\mathbb{Z}^d$  is locally a (2d)-ary tree, the model can be locally seen as a Galton-Watson process with i.i.d. branches, and one could say that the critical parameters of the two models are asymptotically equal, as  $d\to\infty$ .

In this paper we will consider inhomogeneous non-oriented Bernoulli percolation, also known as *anisotropic percolation*, where edges in each direction have distinct parameters. The

<sup>\*</sup>Universidade de São Paulo, Brasil. pagomes@usp.br

<sup>&</sup>lt;sup>†</sup>Universidade Federal de Alagoas, Brasil. alan.pereira@im.ufal.br

<sup>&</sup>lt;sup>‡</sup>Universidade Federal de Minas Gerais, Brasil. rsanchis@mat.ufmg.br

existence of a non-trivial critical hypersurface is well established and some of its features are known; it is continuous and strictly monotonic in each parameter (see [2]) and its behavior near the border (i.e. when some of the parameters are zero) is related to the so-called dimensional crossover phenomenon (see for instance [4]). For d=2, the phase-diagram was completely described by Kesten (see page 54 of [10]) who proved that percolation occurs if and only if the sum of the two parameters is greater than one, except when one of the parameters is trivial. For  $d \geq 3$ , such a sharp result is not expected and we will focus on how the asymptotic behavior of the critical hypersurface can resemble the inhomogeneous Galton-Watson process on the (2d)-ary tree. We will prove that if the mean number of open edges incident to the origin is greater than one, percolation occur under some regularity conditions on the parameters. The main tool we use is a monotonic coupling between anisotropic percolation in different dimensions, with distinct set of parameters.

The remainder of the text is organized as follows: in Section 2 we define more precisely the model and state the main result, in Section 3 we establish the dynamical coupling and in Section 4 we prove the main theorem.

## 2 The model and main results

We now briefly define the model. Let  $\{e_1, \ldots, e_d\}$  be the set of canonical vectors of  $\mathbb{Z}^d$ . For each  $i = 1, \ldots, d$ , let  $E_i = \{\langle x, x \pm e_i \rangle : x \in \mathbb{Z}^d\}$  be the set of edges parallel to  $e_i$ . We denote our edge set by  $E(\mathbb{Z}^d) := \bigcup_i E_i$ .

Given  $p_1, \ldots, p_d \in [0, 1]$ , consider a family of independent random variables  $\{X_e\}_{e \in E(\mathbb{Z}^d)}$ , where, for each  $e \in E_i$ ,  $X_e$  has  $Ber(p_i)$  distribution,  $i = 1, \ldots, n$ . Let  $\mu_e$  be the law of  $X_e$  and  $\mathbb{P} = \prod_{e \in E} \mu_e$  the resulting product measure. We declare an edge e to be open if  $X_e = 1$  and closed otherwise. The model is said to be homogeneous whenever all the parameters  $p_i$  are equal, and inhomogeneous otherwise.

We denote by  $\{x \leftrightarrow y\}$  the event where  $x, y \in \mathbb{Z}^d$  are connected by a open path, i.e., there exist  $x_0, \ldots, x_n$  such that  $x_0 = x$ ,  $x_n = y$  and each  $\langle x_{j-1}, x_j \rangle$  belongs to  $E(\mathbb{Z}^d)$  and is open for  $j = 1, \ldots, n$ . Let  $\mathcal{C}_0^d = \{x \in \mathbb{Z}^d : 0 \leftrightarrow x\}$  be the cluster of the origin, and  $|\mathcal{C}_0^d|$  its size. We define

$$\theta_d(p_1,\ldots,p_d) := \mathbb{P}(|\mathcal{C}_0^d| = \infty).$$

In what follows,  $p_c(\mathbb{Z}^d) = \sup\{p \geq 0 : \theta_d(p, \ldots, p) = 0\}$  denotes the critical point for the non-oriented homogeneous model.

Our main result is the following:

**Theorem 1.** Consider inhomogeneous non-oriented Bernoulli bond Percolation on  $\mathbb{Z}^d$  with parameters  $p_1, \ldots, p_d \in [0, 1]$ . There exists a constant C > 0 such that, for every  $d \geq 2$  and  $\varepsilon > 0$ , if the following conditions are satisfied

C1) 
$$p_1 + \cdots + p_d \ge \frac{1}{2} + \varepsilon$$
, and

C2) 
$$\max_{1 \le i \le d} p_i \le C\varepsilon^2$$
,

then  $\theta_d(p_1,\ldots,p_d)>0$ .

Remark 1. An analogous result was proved in [3] for inhomogeneous oriented Bernoulli bond Percolation. In that case, a martingale approach was used to weaken Condition C2 to  $\max_{1\leq i\leq d} p_i \leq C'\varepsilon$ , for some constant C'. In fact, if we apply the coupling approach presented in the proof of Theorem 1 to the oriented case, we obtain a bound  $C\varepsilon^{3/2}$  in Condition C2, which would be worse than the aforementioned result. We also observe that the methods in [3] are valid only for  $d\geq 4$ . However, since the constant C from our present Theorem 1 is not explicit, there is no gain in extending our result for the oriented case in dimensions d=2,3.

# 3 The Dynamical Couplings

The first step in proving Theorem 1 is to construct a monotonic coupling between inhomogeneous percolation on  $\mathbb{Z}^d$  with parameters  $(p_1, \ldots, p_{d-1}, \tilde{p_d})$ , where  $\tilde{p_d} = 1 - (1 - p_d)(1 - p_{d+1})$  and the inhomogeneous percolation in  $\mathbb{Z}^{d+1}$  with parameters  $(p_1, \ldots, p_{d-1}, p_d, p_{d+1})$ . Monotonic couplings between different percolation processes have been used before, see for instance [6].

**Proposition 1.** Consider inhomogeneous non-oriented Bernoulli bond Percolation on  $\mathbb{Z}^d$ . Let  $p_1, \ldots, p_{d+1} \in [0, 1]$  and let  $\tilde{p}_d \in [0, 1]$  be such that

$$(1 - \tilde{p}_d) = (1 - p_d)(1 - p_{d+1}).$$

Then 
$$\theta_{d+1}(p_1, \dots, p_d, p_{d+1}) \ge \theta_d(p_1, \dots, p_{d-1}, \tilde{p}_d)$$
.

Let us start with a description of the proof. We will construct a dynamic coupling between the percolation process on  $\mathbb{Z}^{d+1}$  with parameters  $p_1, \ldots, p_{d+1}$  and an infection process over  $\mathbb{Z}^d$ . We will do it in such a way that the law of infected sites in  $\mathbb{Z}^d$  is the same as the law of the open cluster of the origin for anisotropic percolation on  $\mathbb{Z}^d$  with parameters  $p_1, \ldots, p_{d-1}, \tilde{p}_d$  and also that, if the infection process survives, the open cluster of the origin of the process in  $\mathbb{Z}^{d+1}$  must be infinite.

The coupling will be built based on a susceptible-infected strategy described as follows. First, we declare the origin of  $\mathbb{Z}^d$  as the *initial* infected component. Next, at each time-step, we possibly grow the infected component and associate each new vertex v of the infected component in  $\mathbb{Z}^d$  to a vertex x(v) in the open cluster of the origin in  $\mathbb{Z}^{d+1}$ . More precisely, consider a vertex v in the infected component of  $\mathbb{Z}^d$  and a neighbor v + u out of the infected component, where  $u \in \{\pm e_1, \ldots, \pm e_d\}$ . The vertex v, in the infected component in  $\mathbb{Z}^d$ , will be associated with some vertex x(v) in the open cluster of the origin in  $\mathbb{Z}^{d+1}$ . If  $\langle x(v), x(v) + u \rangle$  is open, we infect v + u (and write x(v + u) = x(v) + u). If it happens that  $u \in \{\pm e_d\}$  and  $\langle x(v), x(v) + u \rangle$  is closed, we give a second chance to infect v + u. For  $u = \pm e_d$ , in this second

chance, we declare v + u infected in  $\mathbb{Z}^d$  if  $\langle x(v), x(v) \pm e_{d+1} \rangle$  is open in  $\mathbb{Z}^{d+1}$ . In the last case we set  $x(v+u) = x(v) + e_{d+1}$ , or  $x(v+u) = x(v) - e_{d+1}$ , depending on whether  $u = e_d$  or  $u = -e_d$ .

*Proof.* First, we give a precise description of the susceptible-infected type algorithm. Consider the sequence of sets  $(I_n, x(I_n), R_n, S_n)$ , where,  $I_n$  represents the *infected vertices* in  $\mathbb{Z}^d$  up to time n,  $x(I_n)$  represents the vertices in  $\mathbb{Z}^{d+1}$  associated with the infected vertices up to time n and  $R_n$  represents the *removed edges* in  $\mathbb{Z}^d$  up to time n. Finally,  $S_n$  represents the *susceptible edges*, defined as follows. Given  $I_n$ ,  $x(I_n)$  and  $R_n$ , let

$$S_n := \{\langle v, u \rangle : v \in I_n \text{ and } u \notin I_n\} \cap R_n^c.$$

In words,  $S_n$  is the set of edges not removed by time n and that are composed of an infected vertex and a non-infected vertex. We choose to consider susceptible edges instead of vertices, since it simplifies our description.

The dynamic of the process is as follows. We start with the following settings at time 0:

- $I_0 = \{0\},$
- $R_0 = \emptyset$ ,
- $x(0) = 0 \in \mathbb{Z}^{d+1}$ ,
- $S_0 := \{ \langle v, u \rangle : v \in I_0 \text{ and } u \notin I_0 \} \cap R_0^c = \{ \langle 0, \pm e_1 \rangle, \dots, \langle 0, \pm e_d \rangle \}.$

This means that, at time n = 0, only the vertex 0 is infected, and it can potentially infect any of its neighbours, so all edges with the origin as an end-vertex are susceptible. After that, in each step, an infected vertex tries to infect a non-infected vertex through a susceptible edge (if the latter exists). Given a fixed ordering of the edges of  $\mathbb{Z}^d$ , we proceed by the following rules.

Suppose that  $I_n$ ,  $R_n$  and  $S_n$  are already defined. In case there is no susceptible edge, i.e.  $S_n = \emptyset$ , the process stops and we set, for all  $k \ge 1$ ,

- $\bullet \ I_{n+k} = I_n,$
- $\bullet \ R_{n+k} = R_n.$
- $\bullet \ S_{n+k} = S_n.$

Otherwise, if there exists at least one susceptible edge, i.e.  $S_n \neq \emptyset$ , then the infected vertex incident to the smallest (in the previously fixed ordering) such edge tries to infect its non-infected neighbour. More precisely, let  $g_n$  be the smallest edge in  $S_n$ . Since  $g_n \in S_n$ , it has to correspond to some  $\langle v, v + u_n \rangle$ , where  $v \in I_n$ ,  $u_n \in \{\pm e_1, \ldots, \pm e_d\}$  and  $v + u_n \notin I_n$ . Thus we have two options: either  $u_n \in \{\pm e_1, \ldots, \pm e_{d-1}\}$ , or  $u_n \in \{\pm e_d\}$ .

Let us see the first case above. Suppose  $u_n \in \{\pm e_1, \ldots, \pm e_{d-1}\}$ . Then v infects  $v + u_n$  if  $\langle x(v), x(v) + u_n \rangle$  is open in  $\mathbb{Z}^{d+1}$ . More precisely, if  $\langle x(v), x(v) + u_n \rangle$  is open in  $\mathbb{Z}^{d+1}$  then we set

$$I_{n+1} := I_n \cup \{v + u_n\},$$

and define

$$x(v+u_n) := x(v) + u_n$$
.

Otherwise, if  $\langle x(v), x(v) + u_n \rangle$  is closed in  $\mathbb{Z}^{d+1}$ , we set  $I_{n+1} := I_n$ .

In case  $u_n = e_d$ , v has two chances of infecting  $v + u_n$ , that is, either  $\langle x(v), x(v) + e_d \rangle$  is open, and we set

$$x(v + u_n) = x(v + e_d) := x(v) + e_d,$$

or  $\langle x(v), x(v) + e_d \rangle$  is closed and  $\langle x(v), x(v) + e_{d+1} \rangle$  is open, and we write

$$x(v + u_n) = x(v + e_d) := x(v) + e_{d+1}.$$

In both cases, we set

$$I_{n+1} := I_n \cup \{v + u_n\}.$$

On the other hand, if  $\langle x(v), x(v) + e_d \rangle$  and  $\langle x(v), x(v) + e_{d+1} \rangle$  are closed, we write

$$I_{n+1} = I_n.$$

We proceed analogously when  $u_n = -e_d$ .

Now that we have explored  $g_n$ , we remove it and write

$$R_{n+1} := R_n \cup \{g_n\}.$$

To conclude our induction step, we set

$$S_{n+1} := \{ \langle v, u \rangle : v \in I_{n+1} \text{ and } u \notin I_{n+1} \} \cap R_{n+1}^c.$$

Observe that the function  $x: \cup_j I_j \to \mathbb{Z}^{d+1}$  is injective. In fact, let  $v = (v_1, \dots, v_d) \in I_n$ . By construction, we have that  $x(v) = (x_1, \dots, x_d, x_{d+1})$  satisfies

- $x_i = v_i$  for i = 1, ..., d 1,
- $\bullet \ x_d + x_{d+1} = v_d.$

Note that the image of x is contained in the open cluster of the origin of  $\mathbb{Z}^{d+1}$ . Since x is injective, if  $|\cup_n I_n|$  is infinite, then the open cluster of the origin of  $\mathbb{Z}^{d+1}$  must be infinite. Also, note that  $\cup_n I_n$  has the same law as  $\mathcal{C}_0^d$ , where  $\mathcal{C}_0^d$  is the open cluster of the percolation process on  $\mathbb{Z}^d$  with parameters  $p_1, \ldots, p_{d-1}, 1 - (1 - p_d)(1 - p_{d+1})$ . Therefore,

$$\theta_d(p_1, \dots, p_{d-1}, \tilde{p}_d) = \theta_d(p_1, \dots, p_{d-1}, 1 - (1 - p_d)(1 - p_{d+1}))$$

$$\leq \theta_{d+1}(p_1, \dots, p_{d+1}),$$

and the proof of Proposition 1 is complete.

## 4 Proof of Theorem 1

The proof of Theorem 1 will be based on successive applications of Proposition 1.

In this section, we will use the function  $q:[0,1)\to[0,\infty)$  defined by

$$q(p) = -\log(1-p).$$

For i = 1, ..., d, we denote  $q(p_i) = q_i$ ,  $\tilde{q}_d = q(\tilde{p}_d)$  and  $q(p_c(\mathbb{Z}^d)) = q_c(\mathbb{Z}^d)$ . With this notation, the condition of Proposition 1

$$(1 - p_d)(1 - p_{d+1}) = 1 - \tilde{p}_d,$$

simplifies to

$$q_d + q_{d+1} = \tilde{q}_d.$$

Before giving a proof of the Theorem 1, we claim that for sufficiently large values of  $\varepsilon$ , Condition C2 is not necessary, and we have the following non-asymptotic result.

**Proposition 2.** For any  $d \ge 1$ , if the parameters  $p_1, \ldots, p_d \in [0, 1]$  are such that

$$p_1 + \dots + p_d > 3\log 2,\tag{1}$$

then  $\theta_d(p_1,\ldots,p_d)>0$ .

**Remark 2.** In [10] (see page 54), Kesten proved that, for d = 2, the critical curve is  $p_1 + p_2 = 1$  and non-rigorous simulations in low dimensions suggest that the critical hypersurface is convex. If the latter were the case, Proposition 2 could be stated with constant 1 instead of  $3 \log 2$ . It would be nice to have such a result.

*Proof.* First of all, we recall that  $p_c(\mathbb{Z}^2) = 1/2$  and  $q(1/2) = \log 2$ . The idea is to apply Proposition 1 several times in order to compare our d-dimensional system with a supercritical 2-dimensional one. We will consider two cases:  $p_i < 1/2$  for all  $i = 1, \ldots, d$  and  $p_i \ge 1/2$  for some  $i = 1, \ldots, d$ .

If  $p_i < 1/2$ , for all  $i \ge 1$ , then  $q_i < \log 2$  for all  $i \ge 1$ . Also, since  $p_1 + \cdots + p_d \ge 3 \log 2$  and  $q_i \ge p_i$ , it follows that

$$q_1 + \dots + q_d \ge 3\log 2. \tag{2}$$

Let

$$m := \inf \left\{ j : \sum_{i=1}^{j} p_i > \log 2 \right\},$$
 (3)

$$\mathcal{D}_1 = \{1, \dots, m\} \text{ and } \mathcal{D}_2 = \{m+1, \dots, d\}.$$

Using (2) and the fact that  $q_i < \log 2$ , we have that

$$\sum_{i \in \mathcal{D}_1} q_i > \log 2 \text{ and } \sum_{i \in \mathcal{D}_2} q_i > \log 2,$$

and thus, by successive applications of Proposition 1 we obtain that  $\theta_d(p_1, \dots, p_d) > \theta_2(p, p)$ , for some p > 1/2. Since  $p_c(\mathbb{Z}^2) = 1/2$ , the last expression is strictly positive.

In the second case, we can assume, without loss of generality, that  $p_1 \ge 1/2$ . From (1), we have  $p_2 + \cdots + p_d > \log 2$ , hence  $q_2 + \cdots + q_d > \log 2$ .

Therefore, 
$$\theta_d(p_1,\ldots,p_d) > \theta_2(p_1,p) > 0$$
, for some  $p > 1/2$ .

By Proposition 2, it is enough to prove Theorem 1 with  $0 < \varepsilon < \varepsilon^*$ , where  $\varepsilon^* = 3 \log 2 - 1/2$ .

Proof of Theorem 1. Our strategy will be to partition  $\{1, \ldots, d\}$  into m < d subsets. We then apply Proposition 1 to each subset of directions so that inhomogeneous percolation on  $\mathbb{Z}^d$  will dominate a supercritical homogeneous percolation on  $\mathbb{Z}^m$ .

We say that  $(\mathcal{D}_1, \ldots, \mathcal{D}_m)$  is a partition of  $\{1, \ldots, d\}$ , whenever

$$\bigcup_{i=1}^{m} \mathcal{D}_i = \{1, \dots, d\} \text{ and } \mathcal{D}_i \cap \mathcal{D}_j = \emptyset, \ \forall i \neq j.$$

Consider the parameters  $p_1, \ldots, p_d \in [0, 1]$ . If for some  $m \in \mathbb{N}$ , there exists a partition  $(\mathcal{D}_1, \ldots, \mathcal{D}_m)$  of  $\{1, \ldots, d\}$ , such that

$$\sum_{i \in \mathcal{D}_i} q_i > q_c(\mathbb{Z}^m), \quad i = 1, \dots, m,$$
(4)

then, by successive applications of Proposition 1, we obtain  $\theta(p_1, \ldots, p_d) > 0$ . Thus, it is sufficient to prove the existence of a partition  $(\mathcal{D}_1, \ldots, \mathcal{D}_d)$  with the property given in (4). We start by showing a sufficient condition for the existence of such a partition and then, we show that the hypotheses of Theorem 1 imply the sufficient condition we are seeking.

Write  $q_{\text{max}} := \max_{1 \le i \le d} q_i$  and suppose that, for some m < d,

$$q_1 + \dots + q_d > (q_c(\mathbb{Z}^m) + q_{\max})(m-1) + q_c(\mathbb{Z}^m).$$
 (5)

We claim that there exists a partition with the property given by (4). Indeed, take  $i_0 = 0$  and, for each  $\ell = 1, \ldots, m-1$ , let

$$i_{\ell} := \inf \left\{ j : \sum_{i=i_{\ell-1}+1}^{j} q_i > q_c(\mathbb{Z}^m) \right\}.$$
 (6)

Also let  $i_m = d$  and define

$$\mathcal{D}_{\ell} := \{i_{\ell-1} + 1, \dots, i_{\ell}\}, \ \ell = 1, \dots, m.$$

By construction, the partition  $(\mathcal{D}_1, \ldots, \mathcal{D}_d)$  has the desired property and, therefore, we just need to show that it is well defined. Indeed, it follows from (6) that, for each  $\ell = 1, \ldots, m-1$ ,

$$\sum_{i=1}^{i_{\ell}} q_i \le (q_c(\mathbb{Z}^m) + q_{\max})\ell,$$

so, by (5), we have

$$\sum_{i=i_{\ell}+1}^{d} q_{i} > (q_{c}(\mathbb{Z}^{m}) + q_{\max})(m-1-\ell) + q_{c}(\mathbb{Z}^{m}) \ge q_{c}(\mathbb{Z}^{m}), \tag{7}$$

which guarantees the existence of j as in (6). Therefore,  $i_0, \ldots, i_m$  and  $(\mathcal{D}_1, \ldots, \mathcal{D}_m)$  are well defined.

To finish the proof, we will show that the conditions of Theorem 1 imply the existence of some m as in (5). Since, by hypothesis,

$$q_1 + \dots + q_d > p_1 + \dots + p_d \ge \frac{1}{2} + \varepsilon,$$

it is sufficient to find m such that

$$\frac{1}{2} + \varepsilon \ge m(q_c(\mathbb{Z}^m) + q_{\max}),$$

that is,

$$\frac{1}{2m} + \frac{\varepsilon}{m} \ge q_c(\mathbb{Z}^m) + q_{\text{max}}.$$
 (8)

In [8] it is shown that

$$p_c(\mathbb{Z}^d) = \frac{1}{2d} + \frac{1}{4d^2} + \frac{7}{16d^3} + O\left(\frac{1}{d^4}\right).$$

In particular, there exists a constant  $C_1$  such that

$$q_c(\mathbb{Z}^d) \le \frac{1}{2d} + \frac{C_1}{d^2}, \forall d \ge 2.$$

Now, given  $\varepsilon > 0$ , let

$$m(\varepsilon) := \left\lceil \frac{2C_1}{\varepsilon} \right\rceil.$$

Observe that, by the choice of  $m(\varepsilon)$ , we have  $\varepsilon \geq 2C_1/m(\varepsilon)$ , which gives

$$\frac{1+\varepsilon}{2m(\varepsilon)} \ge \frac{1}{2m(\varepsilon)} + \frac{C_1}{m(\varepsilon)^2} \ge q_c(\mathbb{Z}^{m(\varepsilon)}). \tag{9}$$

Moreover, there exists  $C_2 > 0$  such that

$$C_2 \varepsilon^2 \le \frac{\varepsilon}{2m(\varepsilon)}, \ \forall \varepsilon > 0.$$
 (10)

The ratio q(p)/p is increasing in p, so there exists a constant C>0 such that

$$\max_{1 \le i \le d} p_i \le C\varepsilon^2 \implies q_{\max} \le C_2\varepsilon^2, \quad \forall \varepsilon \le \varepsilon^*.$$
 (11)

Combining (9), (10) and (11), we obtain (8), which concludes the proof.

**Remark 3.** Since  $C_1$  could be taken close to 1/4 as long as we take a sufficiently high dimension, we conclude that the constant C could be taken as close to 1 as one wishes. Unfortunately, for a given value of C, we do not have an estimate of the least dimension for which the theorem holds.

Remark 4. Condition C2 may not be optimal, but certainly some condition on the maximal value of the parameters is necessary. For instance, let  $p = 1/2d + 1/8d^2$ . We know that there exists  $d_0$  large enough, such that  $p < p_c(d)$  for every  $d \ge d_0$ . In this case  $\varepsilon = 1/4d$ , and  $p = 2\varepsilon(1+\varepsilon)$ . Thus, Condition C2 could not be replaced by  $\max_i p_i < C\varepsilon^{1-\delta}$  for any constant C > 0 and any  $\delta > 0$ .

### Acknowledgements

The authors thank Roger Silva for valuable comments on the manuscript. P.A. Gomes has been supported by São Paulo Research Foundation (FAPESP), grant 2020/02636-3 and grant 2017/10555-0. R. Sanchis has been partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), CAPES and by FAPEMIG (PPM 00600/16).

# References

- [1] Broadbent S.R., Hammersley, J.M. *Percolation processes: I. Crystals and mazes.* In Mathematical Proceedings of the Cambridge Philosophical Society. Vol. 53, No. 3, pp. 629–641, (1957).
- [2] Couto, R.G., de Lima, B.N.B. and Sanchis, R. Anisotropic percolation on slabs. Markov Process. Related Fields, 20 (1), 145–154, (2014).
- [3] Gomes, P.A., Pereira, A., and Sanchis, R. Anisotropic oriented percolation in high dimensions. ALEA, Lat. Am. J. Probab. Math. Stat. 17, 531–543 (2020).
- [4] Gomes P.A., Sanchis, R. and Silva, R.W.C. A note on the dimensional crossover critical exponent. Lett. Math. Phys. 110, 3427–3434 (2020).
- [5] Gordon, D.M. *Percolation in high dimensions*. Journal of the London Mathematical Society V.2-44 (2), 373-384, (1991).
- [6] Grimmett, G.R., Stacey, A.M.: Critical probabilities for site and bond percolation models. Ann. Probab. 26, 1788–1812 (1998)

- [7] Grimmett, G.R. Percolation. Springer-Verlag (1999).
- [8] Hara, T., and Slade, G. The self-avoiding-walk and percolation critical points in high dimensions. Combinatorics, Probability and Computing 4.3, 197-215 (1995).
- [9] Kesten, H. The Critical Probability of Bond Percolation on the Square Lattice Equals 1/2, Commun. Math. Phys. 74, 41-59 (1980).
- [10] Kesten, H. Percolation Theory for Mathematicians. Bikhäuser (1982).
- [11] Kesten, H. Asymptotics in high dimensions for percolation. Disorder in physical systems, 219-240 (1990).