

---

# DISCOVERY OF INTERSTELLAR GLYCINE IN THE HOT MOLECULAR CORE G10.47+0.03

---

RESEARCH ARTICLE

**Arijit Manna**

Midnapore City College  
Kuturia, Bhadutala, Paschim Medinipur,  
West Bengal, India 721129  
mannaarijit@hotmail.com

**Sabyasachi Pal**

Indian Centre for Space Physics  
43 Chalantika, Garia Station Road,  
Kolkata, India 700084

Midnapore City College  
Kuturia, Bhadutala, Paschim Medinipur,  
West Bengal, India 721129  
sabya.pal@gmail.com

**Soumyadip Banerjee**

Midnapore City College  
Kuturia, Bhadutala, Paschim Medinipur,  
West Bengal, India 721129  
soumyadipbanerjee@hotmail.com

April 14, 2022

**ABSTRACT**

Amino acids are the essential keys in chemistry that contribute to the study of the formation of life. The complex organic molecule glycine ( $\text{NH}_2\text{CH}_2\text{COOH}$ ) is the simplest amino acid that has been investigated in the interstellar medium for a long period to search for a potential connection between the Universe and the origin of life. Several attempts have failed to search for glycine in the last forty years, which made the researcher look for some glycine precursor in the interstellar medium as an alternative approach[1]. We report the successful detection of the rotational emission lines of interstellar glycine with confirmer I and II in the hot molecular core G10.47+0.03 between the frequency range of  $\nu = 158.6\text{--}160.4$  GHz with Atacama Large Millimeter/Submillimeter Array (ALMA) observation. In hot molecular core G10.47+0.03, the fractional abundance of glycine is found between the range of  $(4.01\text{--}4.61)\times 10^{-10}$  which refers to the "medium warm-up" case[2]. The detection of glycine in the interstellar medium is very complicated but many theoretical and laboratory studies indicated the possibilities of the presence of glycine and its precursors in hot molecular cores[2]. We also detected the emission lines of complex organic molecules  $\text{CHOCHOHCH}_2\text{OH}$ ,  $^{13}\text{CH}_2\text{OHCHO}$ ,  $\text{CHD}(\text{OH})\text{CHO}$ ,  $\text{CH}_2\text{OH}^{13}\text{CHO}$ ,  $\text{cis-CH}_2\text{OHCHO}$ ,  $\text{G}'\text{Gg}'\text{-CH}_2(\text{OH})\text{CH}(\text{OH})\text{CH}_2\text{OH}$ , and  $\text{CH}_2\text{DOH}$  in the hot molecular core G10.47+0.03.

**Keywords** astrochemistry – ISM: individual objects: G10.47+0.03 – ISM: molecules – ISM: abundances

**Introduction**

Life most likely originated about 700 million years after the formation of Earth[3]. The three biopolymers RNA, DNA, and nucleic acids are responsible to store the genetic information and proteins. The evolution of prebiotic chemicals from small to massive and complex molecules is believed to have lead to the beginning of life. The very famous Miller-Urey's experiment[4] proved that the organic materials could be synthesized from inorganic materials. During the experiment, many organic molecules, including the amino acids, like alanine and glycine were synthesized,

by discharging a gas mixture of H<sub>2</sub>, NH<sub>3</sub>, H<sub>2</sub>O, and CH<sub>4</sub> which was believed to be the early Earth's atmospheric composition. Many surveys of amino acid glycine (NH<sub>2</sub>CH<sub>2</sub>COOH) in Sgr-B2 and other interstellar medium sources were unsuccessful[7, 8, 9]. The first detection of glycine conformer I line was claimed in hot molecular cores like Sgr B2(N-LMH), Orion KL, and W51 e1/e2[5] but several follow-up observations denied the detection[6]. Earlier, the investigations of the emission lines of glycine have been hampered by the weakness and instability of the glycine lines as well as due to the low sensitivity of the telescopes. Glycine detectability towards NGC6334 IRS1 using ALMA with one hour of on-source integration time was predicted[2]. A three-phase chemical model that completely simulates bulk ice chemistry, coupled gas phase, and grain surface predicted the chances of detecting glycine from the interstellar medium sources using ALMA[2].

## Result and discussions

We detected total twenty-one rotational emission lines of glycine (NH<sub>2</sub>CH<sub>2</sub>COOH) with conformer I and II between the frequency range of  $\nu = 158.6\text{--}160.4$  GHz in the hot molecular core G10.47+0.03. The statistical column density of the emission lines of glycine is found between the range of  $(1.05\text{--}1.2)\times 10^{15}$  cm<sup>-2</sup> (see Methods) and calculated fractional abundance of glycine is found between the range of  $(4.01\text{--}4.61)\times 10^{-10}$  which satisfied the 'medium warm-up' case[2]. The rotational emission spectrum of glycine with conformer I and II are shown in Fig.1. The molecular properties of detected glycine with conf. I and conf. II are shown in Tab.1.

Table 1: Molecular properties of detected glycine with conformer I and II in hot molecular core G10.47+0.03.

Species	Frequency (GHz)	Transition	Lower energy (E <sub>l</sub> ) (K)	log(A <sub>ij</sub> ) (s <sup>-1</sup> )
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.051	24(3,21)–23(3,20), F=24–24	66.338	-7.494
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.211	14(12,3)–14(11,4), F=14–14	51.869	-5.474
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.236	24(5,19)–23(6,18), F=24–23	774.434	-6.718
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.273	13(12,2)–13(11,3)	48.679	-5.891
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.296	25(3,23)–24(2,22), F=25–24	140.342	-4.873
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.327	12(12,1)–12(11,2), F=12–12	45.718	-6.394
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.412	33(29,5)–34(28,6), F=33–34	387.304	-7.262
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.437	29(27,3)–30(26,4), F=29–30	332.966	-7.492
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.547	18(7,11)–18(5,14), F=18–18	117.167	-6.798
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.628	26(0,26)–25(1,25), F=26–25	1554.680	-7.304
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.742	21(5,16)–20(5,15), F=21–21	757.119	-7.333
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.777	14(6,8)–14(3,11), F=14–14	99.016	-7.119
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.895	48(14,34)–48(13,35)	310.934	-5.277
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.909	27(0,27)–26(1,26), F=27–27	70.482	-7.906
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.014	21(5,16)–20(5,15), F=21–20	1544.184	-5.850
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.069	41(8,34)–41(7,35)	215.850	-5.050
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.107	41(8,34)–41(6,35), F=40–40	915.851	-6.038
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.142	21(7,14)–21(5,17)	60.987	-5.351
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	160.157	40(13,28)–40(12,29), F=39–39	222.082	-5.417
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.170	20(6,14)–20(4,17)	53.427	-5.667
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.305	39(6,33)–39(5,34), F=39–39	891.832	-6.604

The complex organic molecules are produced on the grain surface of dust which is evaporated by ultraviolet photons from a central star. The Clausius-Clapeyron equation with a theoretically established boiling point of  $487 \pm 23$  K can be used to compute the evaporation temperature[1]. The evaporation temperature is determined between the temperature range of 82 and 90 K, which is calculated from the fractional abundance of glycine, i.e.  $4.6 \times 10^{-10}$  and a normal number density of H<sub>2</sub> is  $10^7$  cm<sup>-3</sup> in case of a hot core[17]. The measured evaporation temperature is very close to the water ice evaporation temperature on the dust ( $\sim 100$  K) which suggests that the glycine solid phase can evaporate with the help of water ice. As a result, we conclude that glycine's excitation temperature ( $T_{ex}$ ) is  $\sim 100$  K. The source size would come as a significant factor when we consider the detectability of molecular lines. If glycine and water both evaporate at 100 K, the size of the source could be equated to an area with a 100 K or higher temperature. The hot molecular core G10.47+0.03 will have an angular size of roughly 2''. At the distance of hot molecular core G10.47+0.03, the angular dimension equates to a linear size of 0.1 pc, which is equivalent to the size of a typical hot molecular core[1]. Since the beam filling factor for the IRAM single-dish radio telescope is 0.01 between 150–160 GHz, so the probability




Figure 1: The unblended rotational molecular emission spectrums of NH<sub>2</sub>CH<sub>2</sub>COOH with confimer I and II between the frequency range of 158.6–160.4 GHz respectively in the hot molecular core G10.47+0.03 using ALMA band 4 observation. The yellow thick lines indicate the best fit LTE model over the original transition of glycine conformer I and II. The 3-D diagram of the glycine is shown in the center of the figure in which white, red, blue, and black lines are indicative of the hydrogen, oxygen, nitrogen, and carbon atoms respectively.




Figure 2: Suggested possible chemical pathways to the formation of interstellar glycine in hot molecular core G10.47+0.03. In reaction, the “lq” represents the hydrolysis process to formation of NH<sub>2</sub>CH<sub>2</sub>COOH from NH<sub>2</sub>CH<sub>2</sub>CN.

of the detection of glycine is very poor with IRAM[1]. As a result, the beam-averaged brightness temperatures will decrease from 1.2 to 1.6 mK, meaning that single dish telescopes will be unable to detect glycine[1]. So, we will require an interferometer like ALMA to detect the interstellar glycine and we successfully detected the emission lines of glycine with confirmer I and II in G10.47+0.03.

There have been several attempts to find the most essential amino acid glycine (NH<sub>2</sub>CH<sub>2</sub>COOH) in Sgr B2 and other high mass star-forming regions, but none have succeeded. In this condition, searching for glycine precursors is one of the best solutions to explore the formation mechanism of glycine. Although the chemical origin of interstellar N-bearing Complex organic Molecules (COMs) is unknown, so researchers proposed to observe methylamine (CH<sub>3</sub>NH<sub>2</sub>) which is known as a precursor of glycine. The precursor of glycine methylamine (CH<sub>3</sub>NH<sub>2</sub>) was successfully found in the hot molecular core G10.47+0.03 with the fractional abundance of  $1.5 \pm 1.1 \times 10^{-8}$  using Nobeyama 45 m radio telescope[1]. Under cosmic ray irradiation, CH<sub>3</sub>NH<sub>2</sub> can be produced from two well-known abundant compounds, CH<sub>4</sub> and NH<sub>3</sub> on icy dust surfaces[10]. The hydrogenation (adding hydrogen) of HCN generates of CH<sub>2</sub>NH and again hydrogenation of CH<sub>2</sub>NH leads to the generation of CH<sub>3</sub>NH<sub>2</sub> in hot molecular core (HCN → CH<sub>2</sub>NH → CH<sub>3</sub>NH<sub>2</sub>)[11, 10, 12]. Additionally, under the vacuum UV irradiation, CH<sub>3</sub>NH<sub>2</sub> can thermally react with CO<sub>2</sub> in the solid phase to create a carbamate, which could then be converted to a glycine salt.

It is also possible that CH<sub>2</sub>NH is formed during radical reactions in the gas phase[13, 14]. The excitation temperatures of CH<sub>2</sub>NH and CH<sub>3</sub>NH<sub>2</sub> are different which suggests different formation paths for CH<sub>2</sub>NH and CH<sub>3</sub>NH<sub>2</sub>[14]. In another way, if the H atom and HCN would react with CH<sub>2</sub>NH on the grain surface, then as a result the aminoacetonitrile (NH<sub>2</sub>CH<sub>2</sub>CN) would be created[15]. In this way, the glycine would be formed via hydrolysis of NH<sub>2</sub>CH<sub>2</sub>CN[16]. The probable chemical pathways to the formation of detected glycine in the hot molecular core are shown in Fig.2.

We successfully discovered the emission lines of glycine with confirmer I and II towards the hot molecular core G10.47+0.03 with fractional abundance  $(4.01-4.61) \times 10^{-10}$ . The other transition lines of glycine are not observed due to weak signals. The interstellar glycine is detected by ALMA band 4 whereas it would have been very difficult to detect it by the single-dish radio telescope[1]. In the future, we suggest for conducting surveys to the observation of glycine and other prebiotic organic molecules in the comets, protoplanetary disks, solar planets, and extrasolar planets to find the connection between the Universe and the origin of life.

## 1 Supplementary materials

### 1.1 ALMA observations and data reductions:

The interstellar glycine in hot molecular core G10.47+0.03 was observed with the high-resolution Atacama Large Millimeter/submillimeter Array (ALMA) as a part of cycle 3. The observation of interstellar glycine was carried out on 2017 March 7, with phase center of  $(\alpha, \delta)_{J2000} = (18:08:38.232, -19:51:50.400)$  and using thirty-nine antennas. The on-source integration time was 1723.680 sec with band 4 observation. During the observation, Saturn moon Titan was used as a flux calibrator, J1924–2914 acted as a bandpass calibrator, and J1832–2039 acted as a phase calibrator. The correlator (XX–YY) was set up to observe the emission lines of glycine with four spectral windows (SPWs) between the frequency range of 158.49–160.43 GHz. The spectral windows 0 and 1 were set up with 1920 channels (corresponding bandwidth 0.46875 GHz) but other spectral windows like 2 and 3 were set up with 960 channels (corresponding bandwidth 0.9375 GHz).




Figure 3: Millimeter continuum images of hot molecular core G10.47+0.03 at frequency  $\nu = 158.84$  and  $160.02$  GHz in ALMA band 4 observation. The synthesized beam of the continuum image is shown by the red color elliptical circles. The synthesized beam size of displayed images are shown in upper-right corner of each images. The contour levels begin at  $3\sigma$  (1.5 mJy/beam), with each map's contour rising by a factor of  $\sqrt{2}$ .

The uncalibrated raw data of G10.47+0.03 were calibrated by the standard procedure with the python script pipeline in Common Astronomy Software Application (CASA 5.4.1) which was delivered by ALMA observatory. After the initial data reduction, we further performed the self-calibration procedure on the calibrated data in amplitude and phase. The continuum subtraction is applied using the line-free channels with task `uvcontsub`. After the initial data reduction, we generated the synthesized images of G10.47+0.03 using task `tclean` with Briggs weighting robust parameter of 0.5. The resultant millimeter continuum image of G10.47+0.03 at the frequency  $\nu = 158.84$  and  $160.02$  GHz are shown in Fig. 3. After the creation of the emission map, the disk average spectrum is extracted. The corresponding disk average spectrum is shown in Fig. 1. The summary of molecular properties of glycine are shown in Tab. 1. The resultant velocity maps of glycine at frequency  $\nu = 160.30$  GHz with cell size  $0.29''$  are shown in Fig. 4. We also detected the emission lines of complex organic molecules CHOCHOHCH<sub>2</sub>OH, <sup>13</sup>CH<sub>2</sub>OHCHO, CHD(OH)CHO, CH<sub>2</sub>OH<sup>13</sup>CHO, *cis*-CH<sub>2</sub>OHCHO, G'Gg'-CH<sub>2</sub>(OH)CH(OH)CH<sub>2</sub>OH, and CH<sub>2</sub>DOH which are shown in Fig. 5. The molecular properties of other detected complex organic molecules are shown in Tab. 2. The spectral resolution of the spectral windows 0 (159.64–160.11 GHz) and 1 (159.18–159.65 GHz) is 488.28 kHz but in the spectral window 3 (158.49–159.43 GHz) and 4 (159.49–160.43 GHz) are 1128.91 kHz.

## 1.2 Calculation of column density and excitation temperature of glycine:

We use the MADrid Data CUBe Analysis (MADCUBA)[20] to classify the complex organic molecular lines in the hot molecular core G10.47+0.03. The radiative transfer equation can be solved using Spectral Line Identification and Modelling (SLIM) after the identification of NH<sub>2</sub>CH<sub>2</sub>COOH with Conf I and II molecular lines in the hot molecular core. It generates the synthetic spectra of defined molecular species in a Local Thermodynamic Equilibrium (LTE) environment. The SLIM program in MADCUBA employs the Cologne Database for Molecular Spectroscopy (CDMS)[18] and Jet Propulsion Laboratory (JPL)[19] databases to identify the correct spectral line. We used the AUTOFIT tool in SLIM to calculate the physical quantities like column density ( $N$ ), excitation temperature ( $T_{ex}$ ) of the identified molecular emission spectra, which performs a nonlinear least-squares fitting of simulated LTE spectra. To decrease the  $\chi^2$  function, the Levenberg–Marquardt technique is utilized, which combines gradient descent with the Gauss-Newton approach. We first established the spectral feature of the NH<sub>2</sub>CH<sub>2</sub>COOH line by fixing the FWHM (Full-width Half Maximum) manually for each line, which accurately reproduces the known spectrum profiles of the NH<sub>2</sub>CH<sub>2</sub>COOH transitions and it is consistent with other molecules in the region. After fitting the synthetic LTE spectra over original transition of NH<sub>2</sub>CH<sub>2</sub>COOH, we found the column density of glycine  $N(\text{NH}_2\text{CH}_2\text{COOH}) = (1.05\text{--}1.2) \times 10^{15} \text{ cm}^{-2}$  with excitation temperature  $T_{ex} = 100$  K. The dipole moment of the confirmer II line is  $\sim 5$  times higher than that of the confirmer I line at higher energy. The relative abundance of glycine in the hot molecular core concerning H<sub>2</sub> is  $(4.01\text{--}4.61) \times 10^{-10}$ . The resultant LTE fitting parameters are shown in Tab. 3.




Figure 4: Channel map of glycine in the hot molecular core G10.47+0.03 in  $\text{km s}^{-1}$ , which was obtained at the frequency  $\nu = 160.03$  GHz. The synthesized beam of the emission map is  $2.409'' \times 1.460''$ . Synthesized beams are shown by the yellow colour elliptical circles. The contour levels begin at  $3\sigma$  (1.5 mJy/beam), which increased by a factor of  $\sqrt{2}$ .




Figure 5: Disk average rotational molecular emission spectrum of complex organic molecules CHOCHOHCH<sub>2</sub>OH, <sup>13</sup>CH<sub>2</sub>OHCHO, CHD(OH)CHO, CH<sub>2</sub>OH<sup>13</sup>CHO, cis-CH<sub>2</sub>OHCHO, G'Gg'-CH<sub>2</sub>(OH)CH(OH)CH<sub>2</sub>OH, CH<sub>2</sub>DOH, and NH<sub>2</sub>CH<sub>2</sub>COOH with conf I and II in the hot molecular core G10.47+0.03 between the frequency range 158.6–160.4 GHz.

## References

- [1] Ohishi, M., Suzuki, T., Hirota, T., Saito, M. & Kaifu, N. Detection of a new methylamine (CH<sub>3</sub>NH<sub>2</sub>) source: Candidate for future glycine surveys, *Publications of the Astronomical Society of Japan*, **71** (2019).
- [2] Garrod, R. T. A three-phase chemical model of hot cores: the formation of glycine, *ApJ*, **765**, 60 (2013).
- [3] Mojzsis, S. J., Arrhenius, G., McKeegan, K. D. et al. Evidence for life on Earth before 3,800 million years ago, *Nature*, **384**, 5559 (1996).
- [4] Miller, S. L. A Production of Amino Acids Under Possible Primitive Earth Conditions, *Science*, **117**, 528 (1953).
- [5] Kuan Y., Charnley S. B., Huang H., Tseng W. & Kisiel Z., Interstellar Glycine, *ApJ*, **593**, 848 (2003).
- [6] Jones. F. A., Cunningham. M. R., Godfrey P. D. & Cragg D. M., A search for biomolecules in Sagittarius B2 (LMH) with the Australia telescope compact array, *MNRAS*, **347**, 579 (2007).
- [7] Brown, R. D., Godfrey, P. D., Storey J., W., V. & Bassez Marie-Paule. A search for interstellar glycine, *MNRAS*, **186**, 5P (1979).
- [8] Snyder, L. E., Hollis, J. M., Suenram, R. D. et al. An extensive galactic search for conformer II glycine, *ApJ*, **268**, 123 (1983).

Table 2: Properties of detected other complex organic molecules in hot molecular core G10.47+0.03.

Species	Frequency (GHz)	Transition	Lower energy ( $E_l$ ) (K)	$\log(A_{ij})$ ( $s^{-1}$ )
CHOCHOHCH <sub>2</sub> OH	158.622	50(18,33)–49(19,30)	247.605	–6.520
<sup>13</sup> CH <sub>2</sub> OHCHO	158.656	4(3,2)–3(0,3)	2.259	–5.577
CHOCHOHCH <sub>2</sub> OH	158.667	49(4,45)–48(5,43)	205.766	–5.194
CHOCHOHCH <sub>2</sub> OH	158.898	77(42,36)–76(43,34)	683.638	–5.322
CHD(OH)CHO	158.935	27(8,20)–27(7,21)	162.757	–4.129
<sup>13</sup> CH <sub>2</sub> OHCHO	159.201	21(8,14)–22(5,17)	109.979	–6.753
<sup>13</sup> CH <sub>2</sub> OHCHO	159.254	32(9,23)–31(10,22)	232.528	–5.901
CHD(OH)CHO	159.286	16(0,16)–15(1,15)	40.891	–4.756
CHOCHOHCH <sub>2</sub> OH	159.370	44(13,32)–43(14,30)	183.647	–4.904
G'Gg'–CH <sub>2</sub> (OH)CH(OH)CH <sub>2</sub> OH	159.380	34(16,19)–33(17,16)	122.319	–6.779
CH <sub>2</sub> OH <sup>13</sup> CHO	159.489	75(28,47)–74(29,46)	1411.695	–4.906
cis–CH <sub>2</sub> OHCHO	159.556	50(11,39)–49(13,36)	548.621	–8.695
cis–CH <sub>2</sub> OHCHO	159.571	37(11,26)–36(12,25)	319.025	–6.428
G'Gg'–CH <sub>2</sub> (OH)CH(OH)CH <sub>2</sub> OH	159.678	30(4,27)–29(4,26)	80.412	–5.773
CHOCHOHCH <sub>2</sub> OH	159.749	67(16,52)–67(13,54)	417.669	–5.261
G'Gg'–CH <sub>2</sub> (OH)CH(OH)CH <sub>2</sub> OH	159.802	31(20,11)–31(19,12)	120.619	–6.741
CHD(OH)CHO	159.851	26(7,19)–25(8,18)	147.673	–4.814
CH <sub>2</sub> DOH	159.990	16(4,12)–17(2,15)	247.155	–7.627

Table 3: Molecular line properties after fitting LTE model over original transition of glycine with confirmer I and II in hot molecular core G10.47+0.03.

Species	Frequency (GHz)	FWHM (km s <sup>–1</sup> )	$\chi^2$	$N$ (cm <sup>–2</sup> )	$T_{ex}$ (K)	Relative <sup>a</sup> abundance
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.051	6.514±0.611	0.041	1.051×10 <sup>15</sup>	100	4.042×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.211	7.631±0.284	0.034	1.068×10 <sup>15</sup>	100	4.107×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.236	5.394±0.875	0.030	1.182×10 <sup>15</sup>	100	4.546×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.273	5.904±0.500	0.430	1.125×10 <sup>15</sup>	100	4.326×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.296	6.003±0.094	0.032	1.112×10 <sup>15</sup>	100	4.276×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.327	7.719±0.088	0.098	1.135×10 <sup>15</sup>	100	4.365×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.412	7.624±0.140	0.021	1.159×10 <sup>15</sup>	100	4.457×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.437	7.003±0.864	0.131	1.127×10 <sup>15</sup>	100	4.334×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.547	6.432±0.672	0.059	1.176×10 <sup>15</sup>	100	4.523×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.628	7.116±0.184	0.021	1.195×10 <sup>15</sup>	100	4.596×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	159.742	6.003±0.502	0.018	1.192×10 <sup>15</sup>	100	4.584×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.777	7.152±0.451	0.035	1.185×10 <sup>15</sup>	100	4.557×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.895	8.746±0.563	0.058	1.152×10 <sup>15</sup>	100	4.430×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	159.909	7.579±0.530	0.013	1.136×10 <sup>15</sup>	100	4.369×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.014	8.028±0.532	0.033	1.196×10 <sup>15</sup>	100	4.601×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.069	7.523±0.411	0.059	1.186×10 <sup>15</sup>	100	4.561×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.107	8.067±0.463	0.130	1.179×10 <sup>15</sup>	100	4.536×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.142	7.534±0.316	0.573	1.193×10 <sup>15</sup>	100	4.588×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. I	160.157	8.713±0.819	0.138	1.145×10 <sup>15</sup>	100	4.403×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.170	7.498±0.732	0.893	1.199×10 <sup>15</sup>	100	4.611×10 <sup>–10</sup>
NH <sub>2</sub> CH <sub>2</sub> COOH Conf. II	160.305	7.107±0.478	0.637	1.187×10 <sup>15</sup>	100	4.565×10 <sup>–10</sup>

<sup>a</sup>The relative abundance of glycine is calculated with respect to H<sub>2</sub>, where  $N(\text{H}_2) = 2.6 \times 10^{24} \text{ cm}^{-2}$ .[9] Combes, F., Nguyen Q. R., Wlodarczak, G., Search for interstellar glycine, *A&A*, **308**, 618 (1996)[10] Kim, Y. S. & Kaiser, R. I. On the formation of amines (RNH<sub>2</sub>) and the cyanide anion (CN<sup>–</sup>) in electron-irradiated



- ammonia-hydrocarbon interstellar model ices, *ApJ*, **729**, 68 (2011).
- [11] Dickens, J. E., Irvine, W. M., DeVries, C. H. & Ohishi, M. Hydrogenation of interstellar molecules: A survey for methylenimine (CH<sub>2</sub>NH), *ApJ*, **479**, 307 (1997).
- [12] Theule, P., Borget, F., Mispelaer, F., Danger, G., Duvernay, F., Guillemin, J. C. & Chiavassa, T. Hydrogenation of solid hydrogen cyanide HCN and methanimine CH<sub>2</sub>NH at low temperature, *A&A*, **534**, A64 (2011).
- [13] Turner, B. E., Terzieva, R. & Herbst, E. The physics and chemistry of small translucent molecular clouds. XII. More complex species explainable by gas-phase processes, *ApJ*, **518**, 699 (1999).
- [14] Halfen, D. T., Iyushin, V. V., & Ziurys, L. M. Insights into surface hydrogenation in the interstellar medium: Observations of methanimine and methyl amine in Sgr-B2 (N), *ApJ*, **767**, 66 (2013).
- [15] Danger, G., Borget, F., Chomat, M. et al. Experimental investigation of aminoacetonitrile formation through the Strecker synthesis in astrophysical-like conditions: reactivity of methanimine (CH<sub>2</sub>NH), ammonia (NH<sub>3</sub>), and hydrogen cyanide (HCN), *A&A*, **535**, A47 (2011).
- [16] Peltzer, E. T., Bada, J. L., Schlesinger, G. & Miller, S. L. The chemical conditions on the parent body of the Murchison meteorite: Some conclusions based on amino, hydroxy and dicarboxylic acids, *Advances in Space Research*, **4**, 69 (1984).
- [17] Cesaroni, R., Olmi, L., Walmsley, C. M., Churchwell, E. & Hofner, P. A. Massive young embedded object associated with the UC H II region G31.41+0.31, *ApJL*, **435**, L137–L140 (1994).
- [18] Müller, H. S. P., Schlemöder, F., Stutzki, J. & Winnewisser, G. The Cologne Database for Molecular Spectroscopy, CDMS: a useful tool for astronomers and spectroscopists, *Journal of Molecular Structure*, **742**, 215–227 (2005).
- [19] Pickett, H. M., Poynter, R. L., Cohen, E. A. et al. Submillimeter, millimeter, and microwave spectral line catalog, *Journal of Quantitative Spectroscopy and Radiative Transfer*, **60**, 883 (1998).
- [20] Martín, S., Martín, P. J., Blanco, S. C., Rivilla, V., Rodríguez, F. A. & Rico, V. F. Spectral Line Identification and Modelling (SLIM) in the MAdrid Data CUBe Analysis (MADCUBA) package: An interactive software for data cube analysis, *A&A*, **A159**, 17 (2019).

## Acknowledgement

This paper makes use of the following ALMA data: ADS/JAO.ALMA#2016.1.00929.S. ALMA is a partnership of ESO (representing its member states), NSF (USA), and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO, and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.