Large Deviations For Randomly Weighted Sums of Random Measures^{*}

Shui Feng

Department of Mathematics and Statistics McMaster University Hamilton, Ontario Canada L8S 4K1 shuifeng@mcmaster.ca

Abstract

Let $\{Z_n : n \ge 1\}$ be a sequence of i.i.d. random probability measures. Independently, for each $n \ge 1$, let (X_{n1}, \ldots, X_{nn}) be a random vector of positive random variables that add up to one. This paper studies the large deviation principles for the randomly weighted sum $\sum_{i=1}^{n} X_{ni}Z_i$. In the case of finite Dirichlet weighted sum of Dirac measures, we obtain an explicit form for the rate function. It provides a new measurement of divergence between probabilities. As applications, we obtain the large deviation principles for a class of randomly weighted means including the Dirichlet mean and the corresponding posterior mean. We also identify the minima of relative entropy with mean constraint in both forward and reverse directions.

Key words: Bayesian non-parametrics, Dirichlet process, Dirichlet means, gamma process, large deviation, relative entropy.

MSC2020-Mathematics Subject Classifications: Primary: 60G57; Secondary: 62F15.

1 Introduction

Let (Ω, \mathcal{F}, P) be a probability space, S be a compact subset of real line \mathbb{R} , and \mathcal{S} be the Borel σ -algebra. Let C(S) and B(S) denote the spaces of continuous functions and bounded

^{*}Research supported by the Natural Science and Engineering Research Council of Canada

measurable functions on S, respectively. Let $M_1(S)$ denote the space of probability measures on (S, \mathcal{S}) equipped with the weak topology.

For any $n \ge 1$, set

$$\triangle_n = \{(x_1, \dots, x_n) : x_i \ge 0, \sum_{k=1}^n x_k = 1\}.$$

Let $\{Z_i : i \geq 1\}$ be i.i.d. $M_1(S)$ -valued random probability measures with common distribution Π , and, independently, $\mathbf{W} := \{(W_{n1}, W_{n2}, \dots, W_{nn}) : n \geq 1\}$ be a triangular array of positive random variables. Set

$$X_{ni} = \frac{W_{ni}}{\sum_{k=1}^{n} W_{nk}}, \ i = 1, \dots, n.$$

and

$$\mathcal{W}_n(\Pi; \mathbf{W}) = \sum_{i=1}^n X_{ni} \mathcal{Z}_i.$$
 (1.1)

The random distributions \mathcal{Z}_i usually carry the information of location distributions and the coefficients X_{ni} are the corresponding weights or proportions. The collection of random measures $\mathcal{W}_n(\Pi, \mathbf{W})$ is a subclass in the family of randomly weighted averages. The latter has drawn a great deal of interests from researchers due to a wide range of applications including Bayesian nonparametrics, genetics, and the path behaviour of diffusion processes. See the survey papers [12],[14], and the references therein for a comprehensive coverage on the historical and recent development of the subjects.

The main concern of this paper is the large deviations of $\mathcal{W}_n(\Pi; \mathbf{W})$ as *n* tends to infinity. To motivate the study, consider the following subclass of models.

For each $\theta > 0$, let **W** be the triangular array where for each $n \ge 1$ W_1^n, \ldots, W_n^n are i.i.d. exponential with parameter θ/n . Independently, let ξ_1, ξ_2, \ldots be i.i.d. with common distribution ν_0 in $M_1(S)$, δ_x be the Dirac measure concentrated at x, and $\mathcal{Z}_i = \delta_{\xi_i}$.

Then, as θ tends to infinity, $\mathcal{W}_n(\Pi, \mathbf{W})$ converges in distribution to the empirical distribution of ξ_1, \ldots, ξ_n

$$\mathcal{L}_{n,\nu_0} = \frac{1}{n} \sum_{i=1}^n \delta_{\xi_i}.$$

Now letting n go to infinity, we obtain the limit ν_0 .

On the other hand, let U_1, U_2, \ldots be i.i.d. with $\text{Beta}(1, \theta)$ distribution. The Dirichlet process ([7]) with base distribution ν_0 and concentration parameter θ is the random measure

$$\mathcal{V}_{\theta,\nu_0} = \sum_{i=1}^{\infty} V_i \delta_{\xi_i},\tag{1.2}$$


where $V_1 = U_1, V_i = (1 - U_1) \cdots (1 - U_{i-1})U_i, i \ge 2.$

It is known ([10]) that $\mathcal{W}_n(\Pi, \mathbf{W})$ converges in distribution to the Dirichlet process $\mathcal{V}_{\theta,\nu_0}$ as *n* tends to infinity. Taking the limit of θ going to infinity, we obtain the same limit ν_0 .

Instead of taking limits of θ and n separately, we can consider the special case $\theta = n$. Denote $\mathcal{W}_n(\Pi, \mathbf{W})$ in this case by

$$\mathcal{W}_{n,\nu_0} = \sum_{i=1}^n X_{ni} \delta_{\xi_i}.$$
(1.3)

One can show that \mathcal{W}_{n,ν_0} also converges in distribution to ν_0 as *n* tends to infinity. The following diagram is a summery of these relations.



It is natural to investigate and to compare the asymptotic behaviours of the three different estimation of ν_0 . The large deviations of \mathcal{L}_{n,ν_0} from ν_0 when *n* tends to infinity is given by Sanov's theorem (cf. [18]) with the following good rate function ([5])

$$H(\mu|\nu_0) = \sup\{\langle \mu, f \rangle - \log\langle \nu_0, e^f \rangle : f \in C(S)\},$$
(1.5)

where $\langle \mu, f \rangle = \int f(x)\mu(dx)$. The function $H(\mu|\nu_0)$ is the relative entropy of μ with respect to ν_0 .

When θ converges to infinity, the large deviations of $\mathcal{V}_{\theta,\nu_0}$ from ν_0 ([3],[13]) has a good rate function given by the reversed relative entropy $H(\nu_0|\mu)$ when the support of μ is contained in the support of ν_0 .

Our new result of large deviations for \mathcal{W}_{n,ν_0} corresponds to the diagonal limit in (1.4). The good rate function will be identified as the minimum of the sum of relative entropies with respect to μ and ν_0 . Like the relative entropy, it provides an alternate measurement of divergence between probability measures. Comparisons among the three good rate functions reveal refined differences of the three sequences of random measures.

The random probability measure \mathcal{W}_{n,ν_0} also appears in the Dirichlet posterior. More specifically, let η_1, η_2, \ldots , be i.i.d. given the Dirichlet process $\mathcal{V}_{\theta,\nu_0}$. Then the posterior distribution of $\mathcal{V}_{\theta,\nu_0}$ given η_1, \ldots, η_n is

$$U\mathcal{V}_{\theta,\nu_0} + (1-U)\sum_{i=1}^n X_i\delta_{\eta_i}$$

where U is Beta (θ, n) distributed, (X_{n1}, \ldots, X_{nn}) has Dirichlet $(1, \ldots, 1)$ distribution, and all random variables appearing on the right hand side are independent given η_1, \ldots, η_n . Clearly the term $\sum_{i=1}^n X_{ni}\delta_{\eta_i}$ is the same as \mathcal{W}_{n,ν_0} if η_1, \ldots, η_n are i.i.d. with common distribution ν_0 . If θ is fixed, then the large deviation principle for \mathcal{W}_{n,ν_0} is associated with the annealed large deviation principle for the Dirichlet posterior.

The development of the paper is as follows. We starts with the large deviation principles for $\mathcal{W}_n(\Pi; \mathbf{W})$ in Section 2. Our focus will then shift in Section 3 to the study of the large deviations of \mathcal{W}_{n,ν_0} , and its connections to Sanov's theorem and the large deviations of $\mathcal{V}_{\theta,\nu_0}$. Several applications will be discussed in Section 4 including large deviations for Dirichlet and Dirichlet posterior means. We also obtain some minimization properties of the relative entropy. The logarithm log in this article is the natural logarithm, with base *e*. All terms and definitions regarding large deviations are found in the reference [4].

2 Large Deviations for $\mathcal{W}_n(\Pi; \mathbf{W})$

In this section, we discuss the large deviations for $\mathcal{W}_n(\Pi; \mathbf{W})$ defined in (1.1) with S = [0, 1]. We also assume in the sequel that

$$\lim_{n \to \infty} \mathbb{E}\left[\sum_{i=1}^{n} X_{ni}^{2}\right] = 0.$$
(2.1)

Let $\{f_i \in C(S) : i = 1, 2, ...\}$ be a countable dense subset of C(S). The metric on $M_1(S)$ is given by

$$\rho(\upsilon,\mu) = \sum_{i=1}^{\infty} \frac{|\langle \upsilon - \mu, f_i \rangle| \wedge 1}{2^i}.$$

Let ν_0 denote the mean measure of \mathcal{Z}_1 . Then for any f in C(S)

$$\mathbb{E}\left[\langle \mathcal{W}_n(\Pi; \mathbf{W}), f \rangle\right] = \langle \nu_0, f \rangle, \quad \operatorname{Var}\left[\langle \mathcal{W}_n(\Pi; \mathbf{W}), f \rangle\right] = \mathbb{E}\left[\sum_{i=1}^n X_{ni}^2\right] \operatorname{Var}\left(\langle \mathcal{Z}_1, f \rangle\right).$$

It follows from (2.1) that for any $\varepsilon > 0$

$$\lim_{n \to \infty} P\{\rho(\mathcal{W}_n(\Pi; \mathbf{W}), \nu_0) \ge \varepsilon\} = 0.$$

Thus $\mathcal{W}_n(\Pi; \mathbf{W})$ converges in probability to ν_0 in $M_1(S)$. For any $\delta > 0$ and any μ in $M_1(S)$, let

$$B(\mu, \delta) = \{ v \in M_1(S) : \rho(v, \mu) < \delta \}, \quad \overline{B}(\mu, \delta) = \{ v \in M_1(S) : \rho(v, \mu) \le \delta \},$$

and

$$S_{\mu} = \{t \in S : t \text{ is a continuity point of } \mu\}.$$

For any $m \ge 1$, and any $0 = t_0 < t_1 < \ldots < t_m < 1$ in S_{μ} , set

$$B_{t_1,...,t_m}(\mu,\delta) = \{ v \in M_1(S) : \rho_{t_1,...,t_m}(v,\mu) < \delta \}$$

$$\overline{B}_{t_1,...,t_m}(\mu,\delta) = \{ v \in M_1(S) : \rho_{t_1,...,t_m}(v,\mu) \le \delta \},$$

where

$$\rho_{t_1,\dots,t_m}(\upsilon,\mu) = \upsilon([t_m,1]) - \mu([t_m,1])| + \sum_{i=1}^m |\upsilon([t_{i-1},t_i)) - \mu([t_{i-1},t_i))|.$$

Then the following holds.

Theorem 2.1 For each $n \ge 1$, let Π_n denote the law of $\mathcal{W}_n(\Pi, \mathbf{W})$. Assume that for any μ in $M_1(S)$, $m \ge 1$, and any $0 < t_1 < \ldots < t_m < 1$ in S_{μ} , there exists $I_{t_1,\ldots,t_m}(\mu)$ such that

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log \prod_{n} \{ B_{t_1, \dots, t_m}(\mu, \delta) \}$$

=
$$\lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log \prod_{n} \{ \overline{B}_{t_1, \dots, t_m}(\mu, \delta) \}$$

=
$$-I_{t_1, \dots, t_m}(\mu).$$
 (2.2)

Then the sequence $\{\Pi_n : n \ge 1\}$ satisfies a large deviation principle with good rate function

$$I(\mu) = \sup_{t_1, \dots, t_m \in S_{\mu}} \{ I_{t_1, \dots, t_m}(\mu) \}.$$

Proof: Since the space $M_1(S)$ is compact, the sequence $\{\Pi_n : n \ge 1\}$ is exponentially tight. It follows from Theorem (P) in [16] that every subsequence of $\{\Pi_n : n \ge 1\}$ has a further subsequence that satisfies a large deviation principle with a good rate function. To prove the theorem it suffices to prove that all these good rate functions are the same as $I(\cdot)$. This is clearly true if we can show that

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log \prod_n \{ B(\mu, \delta) \}$$
$$= \lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log \prod_n \{ \overline{B}(\mu, \delta) \}$$
$$= -I(\mu).$$

For any μ in $M_1(S)$ there exist f_1, \dots, f_r in C(S) and $\varepsilon > 0$ such that

$$\{v \in M_1(S) : |\langle v, f_j \rangle - \langle \mu, f_j \rangle| < \varepsilon : j = 1, \cdots, r\} \subset B(\mu, \delta).$$

Let

$$c = \sup\{|f_j(x)| : x \in S, j = 1, \cdots, r\}$$

and choose the continuity points $0 < t_1 < \cdots < t_m < 1$ in S_{μ} such that

$$\sup\{|f_j(x) - f_j(y)| : x, y \in [t_{i-1}, t_i], i = 1, \dots, m+1; j = 1, \dots, r\} < \varepsilon/4$$

where $t_0 = 0, t_{m+1} = 1$. Let $0 < \delta_1 < \frac{\varepsilon}{4c}$. Then for any υ in $B_{t_1,\dots,t_m}(\mu, \delta_1)$ and any f_j , we have

$$\begin{aligned} |\langle v, f_j \rangle - \langle \mu, f_j \rangle| &= |\int_{[t_m, 1]} [f_j(x) - f_j(t_m) + f_j(t_m)](v(dx) - \mu(dx)) \\ &+ \sum_{i=1}^m \int_{[t_{i-1}, t_i)} [f_j(x) - f_j(t_{i-1}) + f_j(t_{i-1})](v(dx) - \mu(dx))| \\ &< \frac{\varepsilon}{2} + c\delta_1 < \varepsilon, \end{aligned}$$

which implies that

$$B_{t_1,\cdots,t_m}(\mu,\delta_1) \subset \{\mu \in M_1(S) : |\langle v, f_j \rangle - \langle \mu, f_j \rangle| < \varepsilon : j = 1, \cdots, r\} \subset B(\mu,\delta).$$

Noting that the continuity points t_1, \ldots, t_m does not depend on δ_1 , it follows from (2.2) that

$$-I(\mu) \leq -I_{t_1,\dots,t_m}(\mu)$$

=
$$\lim_{\delta_1 \to 0} \liminf_{n \to \infty} \frac{1}{n} \log \prod_n \{B_{t_1,\dots,t_m}(\mu,\delta_1)\}$$

$$\leq \liminf_{n \to \infty} \frac{1}{n} \log \prod_n \{B(\mu,\delta)\}.$$

It now follows by letting δ go to zero that

$$\liminf_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log \prod_n \{ B(\mu, \delta) \} \ge -I(\mu).$$
(2.3)

On the other hand, for any μ in $M_1(S)$, $\delta_2 > 0$, $m \ge 1$, and any continuity points $0 < t_1 < \cdots < t_m < 1$ of μ , the set $B_{t_1,\ldots,t_m}(\mu, \delta_2)$ is open in $M_1(S)$. One can choose δ small enough so that

$$B(\mu,\delta) \subset \overline{B}(\mu,\delta) \subset B_{t_1,\dots,t_m}(\mu,\delta_2) \subset \overline{B}_{t_1,\dots,t_m}(\mu,\delta_2).$$

Letting δ go to zero we obtain that

$$\limsup_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log \prod_n \{\overline{B}(\mu, \delta)\} \le \limsup_{n \to \infty} \frac{1}{n} \log \prod_n \{\overline{B}_{t_1, \dots, t_m}(\mu, \delta_2)\}$$

Letting δ_2 go to zero followed by taking the infimum of $-I_{t_1,\dots,t_m}(\mu)$ over the continuity points of μ we obtain

$$\limsup_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log \prod_{n} \{ \overline{B}(\mu, \delta) \} \le -I(\mu).$$
(2.4)

Putting together (2.3) and (2.4) we obtain the result.

Remark 2.1 The space S may be generalized to a compact set in a Polish space. But the compactness is crucial in general.

Remark 2.2 The limit $I_{t_1,...,t_m}(\mu)$ appearing in (2.2) is just a non-negative number. There is no other conditions imposed. This makes the verification a bit easier.

Theorem 2.1 works well for random probabilities that have explicit forms when projected to finite partitions of S. We demonstrate this through two examples.

Example 2.1 Let $\xi_1, \ldots, \xi_n, \ldots$ be *i.i.d.* with common law $\nu_0, W_{n1} = \cdots = W_{nn} = 1$, and $\mathcal{Z}_i = \delta_{\xi_i}$. Then $X_{n1} = \cdots = X_{nn} = \frac{1}{n}$, Π is the law of δ_{ξ_1} , and $\mathcal{W}_n(\Pi, \mathbf{W})$ is the usual empirical distribution

$$\mathcal{L}_{n,\nu_0} = \frac{1}{n} \sum_{i=1}^n \delta_{\xi_i}$$

Assume that ν_0 is positive on all open intervals. For any $m \geq 2$ and any

$$0 = t_0 < t_1 < \dots < t_{m-1} < 1,$$

let

$$p_m = \nu_0([t_{m-1}, 1]), \ p_i = \nu_0([t_{i-1}, t_i)), \ i = 1, \dots, m-1.$$

We have

$$\left(\mathcal{W}_n(\Pi, \mathbf{W})([0, t_1)), \dots, \mathcal{W}_n(\Pi, \mathbf{W})([t_{m-1}, 1])\right) \stackrel{d}{=} \frac{1}{n}(Y_1^n, \dots, Y_m^n)$$

where $\stackrel{d}{=}$ denotes equality in distribution, and (Y_1^n, \ldots, Y_m^n) is multinomial with parameters n, p_1, \ldots, p_m . Then the condition (2.2) holds for

$$I_{t_1,\dots,t_m}(\mu) = \sum_{i=1}^m q_i \ln \frac{q_i}{p_i}$$

where

$$q_m = \mu([t_{m-1}, 1]), \ q_i = \mu([t_{i-1}, t_i), i = 1, \dots, m-1.$$

Example 2.2 Let $\gamma(t)$ denote the gamma subordinator on $[0, +\infty)$ and set

$$W_i = \gamma(i) - \gamma(i-1), i = 1, 2, \dots$$

Let $\{J_{ik} : k \ge 1\}$ denote all the jump sizes of $\gamma(t)$ over the interval [i-1,i]. Then it follows from the Gamma-Dirichlet algebra (Theorem 1.1 in [6]) that the Dirichlet process, defined in (1.2), with base distribution ν_0 on [0,1] and concentration parameter n has the form

$$\mathcal{V}_{n,\nu_0} = \frac{1}{\gamma(n)} \sum_{i=1}^n \sum_{k=1}^\infty J_{ik} \delta_{\xi_{ik}}$$

where $\{\xi_{ik} : i = 1, ..., n; k = 1, ...\}$ are *i.i.d.* with common distribution ν_0 . Reorganizing the terms and applying the Gamma-Dirichlet algebra again we obtain

$$\mathcal{V}_{n,\nu_0} = \frac{1}{\sum_{l=1}^{n} W_l} \sum_{i=1}^{n} W_i \sum_{k=1}^{\infty} \frac{J_{ik}}{W_i} \delta_{\xi_{ik}}$$

$$= \frac{1}{\sum_{l=1}^{n} W_l} \sum_{i=1}^{n} W_i \mathcal{Z}_i$$
(2.5)

where $\mathcal{Z}_1, \ldots, \mathcal{Z}_n$ are *i.i.d.* Dirchlet processes with base distribution ν_0 and concentration parameter one. Thus the Dirichlet process \mathcal{V}_{n,ν_0} corresponds to $\mathcal{W}_n(\Pi, \mathbf{W})$ where W_1, W_2, \ldots are *i.i.d.* exponential with parameter one and each \mathcal{Z}_i is a copy of \mathcal{V}_{1,ν_0} . Assume that the support of ν_0 is [0,1]. Then the condition (2.2) holds ([3]) for

$$I_{t_1,\dots,t_m}(\mu) = \sum_{i=1}^m p_i \ln \frac{p_i}{q_i}$$

where p_i 's and q_i 's are defined as in Example 2.1. Note that the order between p_i 's and q_i 's is switched here.

3 Large Deviations for W_{n,ν_0}

In this section we establish the large deviations for \mathcal{W}_{n,ν_0} defined in (1.3). The key in the proof is to verify the condition (2.2) in Theorem 2.1. Noting that the weights of \mathcal{W}_{n,ν_0} are the same as the weights of the Dirichlet process \mathcal{V}_{n,ν_0} in (2.5). But the location distributions are different. This innocent-looking difference leads to very different large deviation behaviours as shown below.

For any $m \geq 1$ and any partition A_1, \ldots, A_m of S, define the map

$$\pi: M_1(S) \to \triangle_m, \upsilon \to (\upsilon(A_1), \dots, \upsilon(A_m))$$

Let $p_i = \nu_0(A_i), i = 1, ..., m$, and

$$n_i = \#\{1 \le k \le n : \xi_k \in A_i\}, \ i = 1, \dots, m.$$

Then $\pi(\mathcal{W}_{n,\nu_0})$ has Dirichlet (n_1,\ldots,n_m) distribution, and $\pi(\mathcal{L}_{n,\nu_0})$ has multinomial distribution with parameters n, p_1, \ldots, p_m . Without loss of generality, we assume $n_i \ge 1, p_i > 0$ for $i = 1, \ldots, m$. For μ, ν in $M_1(S)$, set

$$\pi(\mu) = (q_1, \ldots, q_n), \pi(\upsilon) = (o_1, \ldots, o_n)$$

Define

$$H(\pi(\upsilon)|\pi(\mu)) = \sum_{i=1}^{m} o_i \log \frac{o_i}{q_i}, \ H(\pi(\upsilon)|\pi(\nu_0)) = \sum_{i=1}^{m} o_i \log \frac{o_i}{p_i}$$

where $0 \log 0 = 0 \log \frac{0}{0} = 0$.

Lemma 3.1 For any μ, v in $M_1(S)$, we have

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| < \delta \right\} \\
= \lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| \le \delta \right\} \quad (3.1)$$

$$= -\left[H(\pi(\upsilon)|\pi(\mu)) + H(\pi(\upsilon)|\pi(\nu_0)) \right]$$

where

$$\left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| = \sum_{i=1}^m \left[\left| \mathcal{W}_{n,\nu_0}(A_i) - q_i \right| + \left| \mathcal{L}_{n,\nu_0}(A_i) - o_i \right| \right].$$

Proof: We only give the proof of equality

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| < \delta \right\}$$

$$= -\left[H(\pi(\upsilon)|\pi(\mu)) + H(\pi(\upsilon)|\pi(\nu_0)) \right].$$
(3.2)

The proof for $P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| \le \delta \right\}$ is similar. For any $\delta > 0, 1 \le l \le m$, and $r_i \ge 1, \sum_{i=1}^l r_i = n$, let

$$F(r_1,\ldots,r_l;\delta) = \frac{\Gamma(n)}{\Gamma(r_1)\cdots\Gamma(r_l)} \int_{\mathbf{D}_{\delta,l}} \prod_{i=1}^l x_i^{r_i-1} dx_1 \ldots dx_{l-1}$$

where

$$\mathbf{D}_{\delta,l} = \{(x_1,\ldots,x_l) \in \triangle_l : \sum_{i=1}^l |x_i - q_i| < \delta\}.$$

We can extend the domain of F to zero integers by defining

$$F(n_1,\ldots,n_m;\delta) = F(n_{k_1},\ldots,n_{k_r};\delta)$$

where $\{n_{k_j} : j = 1, ..., r\}$ consists of all non-zero elements of $\{n_i : i = 1, ..., m\}$.

For any μ in $M_1(S)$, we have

$$P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| < \delta \right\}$$

$$= \sum_{\sum_{k=1}^m |\frac{n_i}{n} - o_i| < \delta} A(n_1, \dots, n_m; \delta)$$
(3.3)

where

$$A(n_1,\ldots,n_m;\delta) = \binom{n}{n_1\cdots n_m} F(n_1,\ldots,n_m;\delta) \prod_{i=1}^m p_i^{n_i}$$

Noting that $A(n_1, \ldots, n_m; \delta)$ depends on positive n_i 's only. We thus assume, without loss of generality, that $n_i \ge 1$ for $i = 1, \ldots, m$. The arguments for general cases are similar except the replacement of (n_1, \ldots, n_m) with its subset consisting of non-zero elements.

By Stirling's formula for the gamma function, there exist positive constants $c_1 < c_2$ such that for all $n \ge 1$ and $n_i \ge 1, \sum_{i=1}^m n_i = n$

$$c_1 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-n_i} \sqrt{\frac{n_1 \dots n_m}{n}} \le \frac{\Gamma(n)}{\Gamma(n_1) \cdots \Gamma(n_m)} \le c_2 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-n_i} \sqrt{\frac{n_1 \dots n_m}{n}}$$

Noting that for any integer $n\geq 1$

$$\sqrt{2\pi}n^{n+\frac{1}{2}}e^{-n} \le n! \le en^{n+\frac{1}{2}}e^{-n},$$

it follows that there exist constants $c_3 < c_4$ such that

$$c_3 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-n_i} \sqrt{\frac{n}{n_1 \dots n_m}} \le \binom{n}{n_1 \dots n_m} \le c_4 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-n_i} \sqrt{\frac{n}{n_1 \dots n_m}}.$$

Putting these together it follows that

$$c_5 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-2n_i} \le \binom{n}{n_1 \dots n_m} \frac{\Gamma(n)}{\Gamma(n_1) \cdots \Gamma(n_m)} \le c_6 \prod_{i=1}^m \left(\frac{n_i}{n}\right)^{-2n_i}$$

for some positive constants $c_5 < c_6$.

For any $\mathbf{x} = (x_1, \ldots, x_m), \mathbf{y} = (y_1, \ldots, y_m)$ in Δ_m , define the function

$$\Psi(\mathbf{x}, \mathbf{y}) = H(\mathbf{y}|\mathbf{x}) + H(\mathbf{y}|\pi(\nu_0)).$$

Then we have for $\mathbf{y} = \left(\frac{n_1}{n}, \dots, \frac{n_m}{n}\right)$

$$c_{5} \int_{\mathbf{D}_{\delta,m}} \exp\{-n(\Psi(\mathbf{x},\mathbf{y}) + n^{-1}\log\prod_{i=1}^{m} x_{i})\}dx_{1}, \cdots dx_{m-1}$$
$$\leq A(n_{1}, \dots, n_{m}; \delta)$$
(3.4)

$$\leq c_6 \int_{\mathbf{D}_{\delta,m}} \cdots \int \exp\{-n(\Psi(\mathbf{x},\mathbf{y}) + n^{-1}\log\prod_{i=1}^m x_i\}dx_1, \cdots dx_{m-1}\}$$

where

$$\mathbf{D}_{\delta,m} = \{(x_1,\ldots,x_m) \in \triangle_m : \sum_{i=1}^m |x_i - q_i| < \delta\}.$$

We now divide the upper estimations into three cases.

Case 1. $q_i > 0$ for all *i*. For any $\varepsilon > 0$ one can choose *n* large and δ small such that

$$\Psi(\mathbf{x}, \mathbf{y}) + n^{-1} \log \prod_{i=1}^{m} x_i - \Psi(\pi(\mu), \pi(\upsilon)) - n^{-1} \log \prod_{i=1}^{m} q_i \bigg| \le \varepsilon.$$

It follows that

$$A(n_1, \dots, n_m; \delta) \leq c_6 [2\delta]^m \exp\left\{-n\left(\Psi(\pi(\mu), \pi(\upsilon)) + n^{-1}\log\prod_{i=1}^m q_i - \varepsilon\right)\right\}.$$
(3.5)

Since the total number of terms in (3.3) is at most $(n+1)^m$ and ε is arbitrary, we obtain that

$$\lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log P \left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| < \delta \right\} \\
\leq -\sum_{i=1}^{l} o_i \left(\log \frac{o_i}{q_i} + \log \frac{o_i}{p_i} \right) \\
= - \left[H(\pi(\upsilon) | \pi(\mu)) + H(\pi(\upsilon) | \pi(\nu_0)) \right].$$
(3.6)

Next we turn to the situation where $q_i = 0$ for some *i*. Without loss of generality, we assume that there exists an 1 < l < m such that $q_i > 0$ for $1 \le i \le l$ and $q_i = 0$ for i > l. **Case 2.** $o_i > 0$ for some i > l. In this case the term $\frac{n_i - 1}{n} \log \frac{n_i / n}{x_i}$ in

$$\Psi(\mathbf{x}, \mathbf{y}) + n^{-1} \log \prod_{i=1}^{m} x_i$$

converges to infinity as δ tends to zero. Thus

$$\lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| \le \delta \right\} \le -\infty.$$

The result (3.2) follows from the fact that

$$H(\pi(\upsilon)|\pi(\mu)) + H(\pi(\upsilon)|\pi(\nu_0)) = \infty,$$

Case 3. $o_i = 0$ for i > l. It follows from direct calculation that

$$\begin{split} \exp\left\{-n\left(\Psi(\mathbf{x},\mathbf{y})+n^{-1}\log\prod_{i=1}^{m}x_{i}\right)\right\} \\ &=\left(\prod_{i=l+1}^{m}x_{i}^{n_{i}-1}\right)\exp\left\{-n\left(\sum_{i=1}^{l}y_{i}\left(\log\frac{y_{i}}{x_{i}}+\log\frac{y_{i}}{p_{i}}\right)+n^{-1}\log\prod_{i=1}^{l}x_{i}\right)\right\} \\ &\quad \times\exp\left\{-n\left(\sum_{i=l+1}^{m}y_{i}\left(\log y_{i}+\log\frac{y_{i}}{p_{i}}\right)\right)\right\} \\ &\leq \exp\left\{-n\left(\sum_{i=1}^{l}y_{i}\left(\log\frac{y_{i}}{x_{i}}+\log\frac{y_{i}}{p_{i}}\right)+n^{-1}\log\prod_{i=1}^{l}x_{i}\right)\right\} \\ &\quad \times\exp\left\{-n\left(\sum_{i=l+1}^{m}y_{i}\left(\log y_{i}+\log\frac{y_{i}}{p_{i}}\right)\right)\right\} \end{split}$$

on the domain $\mathbf{D}_{\delta,m}$. The exponential term

$$\exp\left\{-n\left(\sum_{i=1}^{l} y_i\left(\log\frac{y_i}{x_i} + \log\frac{y_i}{p_i}\right) + n^{-1}\log\prod_{i=1}^{l} x_i\right)\right\}$$

can be estimated by an argument similar that used in deriving (3.5).

For the second exponential term we have

$$\lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log \exp \left\{ -n \left(\sum_{i=l+1}^{m} y_i \left(\log y_i + \log \frac{y_i}{p_i} \right) \right) \right\}$$
$$\leq -\lim_{\delta \to 0} \inf_{\sum_{i=l+1}^{m} y_i \leq \delta} \left\{ \sum_{i=l+1}^{m} y_i \left(\log y_i + \log \frac{y_i}{p_i} \right) \right\}$$
$$= 0.$$

Thus (3.6) also holds in this case. It remains to check the lower bound in **Cases 1** and **3**.

In **Case 1**, the function $\Psi(\mathbf{x}, \mathbf{y}) + n^{-1} \log \prod_{i=1}^{m} x_i$ is continuous at (\mathbf{q}, \mathbf{o}) . Thus for any $\varepsilon > 0$ one can choose δ small so that for

$$|(\mathbf{x}, \mathbf{y}) - (\mathbf{q}, \mathbf{o})| < \delta$$

one has

$$|\Psi(\mathbf{x}, \mathbf{y}) - \Psi(\mathbf{q}, \mathbf{o}) + n^{-1} (\log \prod_{i=1}^{m} x_i - \log \prod_{i=1}^{m} q_i)| < \varepsilon$$

By (3.4) we have

$$A(n_1, \dots, n_m; \delta)$$

$$\geq c_5 \int_{\mathbf{D}_{\delta,m}} \cdots \int exp\{-n[\Psi(\mathbf{x}, \mathbf{y}) + n^{-1}\log\prod_{i=1}^m x_i]\}dx_1, \cdots dx_m$$

$$\geq c_5 e^{-n\varepsilon} \exp\{-n[\Psi(\mathbf{q}, \mathbf{o}) + n^{-1}\prod_{i=1}^m q_i]\} \int_{\mathbf{D}_{\delta,m}} \cdots \int dx_1 \cdots dx_m$$

which implies

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log P\left\{ \left| \left(\pi(\mathcal{W}_{n,\nu_0}), \pi(\mathcal{L}_{n,\nu_0}) \right) - \left(\pi(\mu), \pi(\upsilon) \right) \right| < \delta \right\} \qquad (3.7)$$

$$\geq - \left[H(\pi(\upsilon)|\pi(\mu)) + H(\pi(\upsilon)|\pi(\nu_0)) \right].$$

In Case 3, let $T = \{1 \le i \le m : q_i = 0\} (o_i = 0 \text{ for } i \in T) \text{ and define}$

$$\Psi_1(\mathbf{x}, \mathbf{y}) = \sum_{i \notin T} y_i \log \frac{y_i}{x_i}, \ \Psi_2(\mathbf{x}, \mathbf{y}) = \sum_{i \in T} y_i \log \frac{y_i}{x_i}.$$

The function $\Psi_1(\mathbf{x}, \mathbf{y}) + n^{-1} \log \prod_{i \notin T} x_i$ is clearly continuous at (\mathbf{q}, \mathbf{o}) . On the other hand, set

$$\mathbf{C}_{\delta,m} = \{ \mathbf{x} = (x, \dots, x_m) \in \Delta_m : \frac{1}{2}(q_i + \frac{\delta}{m}) < x_i < q_i + \frac{\delta}{m}, \text{ for all } i \}.$$

Then the following holds on $\mathbf{C}_{\delta,m}$.

$$\Psi_2(\mathbf{x}, \mathbf{y}) + n^{-1} \log \prod_{i \in T} x_i \le -m\delta \log \frac{\delta}{2m} + \frac{|T|}{n} \log \frac{\delta}{m}$$

where |T| is the cardinality of T.

Since $C_{\delta,m}$ is a subset of $D_{\delta,m}$, it follows from (3.4) and an argument similar to **Case 1** that (3.7) holds in **Case 3**.

Putting together (3.6) and (3.7) we obtain (3.2) and the lemma.

For any μ in $M_1(S)$, and any $0 < t_1 < \cdots < t_m < 1$ in S_{μ} , define

$$\pi_{t_1,\dots,t_m}(\mu) = (\mu([0,t_1)),\dots,\mu([t_k,t_{k+1})),\dots,\mu([t_m,1])).$$

Lemma 3.2 For any v, μ in $M_1(S)$, we have

$$H(\upsilon|\mu) + H(\upsilon|\nu_0) = \sup_{0 < t_1 < \dots < t_m < 1 \in S_{\upsilon} \cap S_{\mu}} \{ H(\pi_{t_1,\dots,t_m}(\upsilon)|\pi_{t_1,\dots,t_m}(\mu)) + H(\pi_{t_1,\dots,t_m}(\upsilon)|\pi_{t_1,\dots,t_m}(\nu_0)\} \}.$$
(3.8)

Proof: It is known ([9], [3]) that for any v, μ in $M_1(S)$

$$H(v|\mu) = \sup_{0 < t_1 < \dots < t_m < 1 \in S_{\mu}} \{ H(\pi_{t_1,\dots,t_m}(v) | \pi_{t_1,\dots,t_m}(\mu) \}.$$

Since the supremum of sums is less than or equal to the sum of supremums, it follows that

$$H(v|\mu) + H(v|\nu_0) \geq \sup_{\substack{0 < t_1 < \cdots < t_m < 1 \text{ in } S_v \cap S_\mu \\ + H(\pi_{t_1, \dots, t_m}(v) | \pi_{t_1, \dots, t_m}(\nu_0) \}.} \{H(\pi_{t_1, \dots, t_m}(v) | \pi_{t_1, \dots, t_m}(v) \}.$$

To prove the other direction, we first recall the variational form (1.5) of the relative entropy

$$H(\upsilon|\mu) = \sup_{g \in C(S)} \{ \langle \upsilon, g \rangle - \log \langle \mu, e^g \rangle \}.$$

For any $\varepsilon > 0$, there are g, h in C(S) such that

$$H(\upsilon|\mu) \le \langle \upsilon, g \rangle - \log \langle \mu, e^g \rangle + \varepsilon$$

and

$$H(v|\nu_0) \le \langle v, h \rangle - \log \langle \mu, e^h \rangle + \varepsilon.$$

Since $S_v \cap S_\mu$ is dense in S, there exist $0 < t_{n1} < \cdots < t_{nn} < 1$ in $S_v \cap S_\mu$ such that

$$\lim_{n \to \infty} \max_{1 \le i \le n+1} \left\{ \max\{|g(x) - g(y)| : x, y \in [t_{n(i-1)}, t_{ni}] \right\} = 0,$$
$$\lim_{n \to \infty} \max_{1 \le i \le n+1} \left\{ \max\{|h(x) - h(y)| : x, y \in [t_{n(i-1)}, t_{ni}] \} \right\} = 0.$$

where $t_{n0} = 0, t_{n(n+1)} = 1$. Set

$$\alpha_{ni} = g(t_{ni}), \beta_{ni} = h(t_{ni}), i = 1, \dots, n+1$$

and

$$A_{n(n+1)} = [t_{nn}, 1], A_{ni} = [t_{n(i-1)}, t_{ni}), i = 1, \dots, n.$$

Then there exists $c_n(g, h)$ such that

$$\lim_{n \to \infty} c_n(g, h) = 0$$

$$H(\upsilon | \mu) \leq \sum_{i=1}^{n+1} \alpha_{ni} \upsilon(A_{ni}) - \log \sum_{i=1}^{n+1} e^{\alpha_{ni}} \mu(A_{ni}) + \varepsilon + c_n(g, h)$$

$$\leq H(\pi_{t_{n1},\dots,t_{nn}}(\upsilon) | \pi_{t_{n1},\dots,t_{nn}}(\mu)) + \varepsilon + c_n(g, h)$$

and

$$H(\upsilon|\nu_0) \le \sum_{i=1}^{n+1} \beta_{ni} \upsilon(A_{ni}) - \log \sum_{i=1}^{n+1} e^{\beta_{ni}} \nu_0(A_{ni}) + \varepsilon + c_n(g,h) \le H(\pi_{t_{n1},\dots,t_{nn}}(\upsilon)|\pi_{t_{n1},\dots,t_{nn}}(\nu_0)) + \varepsilon + c_n(g,h).$$

Putting all these together we obtain (3.8).

Theorem 3.3 Assume that the topological support of ν_0 is S. Then the family of the laws of $\{\mathcal{W}_{n,\nu_0} : n \geq 1\}$ satisfies a large deviation principle on $M_1(S)$ with good rate function

$$J(\mu,\nu_0) = \inf_{\nu \in M_1(S)} \{ H(\nu|\mu) + H(\nu|\nu_0) \}.$$
(3.9)

Proof: For any v, μ in $M_1(S)$, and any $0 < t_1 < \cdots < t_m < 1$ in $S_v \cap S_{\mu}$, let

$$A_{m+1} = [t_m, 1], A_i = [t_{i-1}, t_i), i = 1, \dots, m$$

Then it follows from (3.1) that

$$\lim_{\delta \to 0} \liminf_{n \to \infty} \frac{1}{n} \log P \bigg\{ \bigg| \bigg(\pi_{t_1, \dots, t_m}(\mathcal{W}_{n, \nu_0}), \pi_{t_1, \dots, t_m}(\mathcal{L}_{n, \nu_0}) \bigg) - \bigg(\pi_{t_1, \dots, t_m}(\mu), \pi_{t_1, \dots, t_m}(\upsilon) \bigg) \bigg| < \delta \bigg\} = \lim_{\delta \to 0} \limsup_{n \to \infty} \frac{1}{n} \log P \bigg\{ \bigg| \bigg(\pi_{t_1, \dots, t_m}(\mathcal{W}_{n, \nu_0}), \pi_{t_1, \dots, t_m}(\mathcal{L}_{n, \nu_0}) \bigg) - \bigg(\pi_{t_1, \dots, t_m}(\mu), \pi_{t_1, \dots, t_m}(\upsilon) \bigg) \bigg| \le \delta \bigg\} = - \bigg[H \bigg(\pi_{t_1, \dots, t_m}(\upsilon) | \pi_{t_1, \dots, t_m}(\mu) \bigg) + H \bigg(\pi_{t_1, \dots, t_m}(\upsilon) | \pi_{t_1, \dots, t_m}(\nu_0) \bigg) \bigg].$$

Since $M_1(S) \times M_1(S)$ is compact, it follows, by Lemma 3.2 and an arguments similar to that used in the proof of Theorem 2.1, that the family of the laws of $(\mathcal{W}_{n,\nu_0}, \mathcal{L}_{n,\nu_0})$ satisfies a large deviation principle with rate function $H(v|\mu) + H(v|\nu_0)$.

Noting that \mathcal{W}_{n,ν_0} is the continuous image of $(\mathcal{W}_{n,\nu_0}, \mathcal{L}_{n,\nu_0})$ through projection, we obtain the result by the contraction principle.

Remark 3.1 The Dirichlet posterior given η_1, \ldots, η_n is a Beta mixture of the Dirichlet process and the conditional distribution of $\{W_{n,\nu_0} : n \ge 1\}$ given $\xi_i = \eta_i, i = 1, \ldots, n$. Since the coefficient the Dirichlet process part converges to zero as n tends to infinity, the large deviation behaviour of the Dirichlet posterior is the same as the large deviation for the conditional distribution of $\{W_{n,\nu_0} : n \ge 1\}$. Integrating the posterior with respect to independent product of a probability measure ν_0 leads to an annealed Dirichlet posterior. The large deviation principle in [8] with rate function $H(\nu_0|\mu)$ can be viewed as a quenched large deviation principle for the annealed Dirichlet posterior. The annealed rate function is smaller than the quenched rate function.

Recall that the large deviation rate functions for the empirical distributions and the Dirichlet processes are $H(\mu|\nu_0)$ and $H(\nu_0|\mu)$, respectively. Since

$$H(\mu|\nu_0) = H(\mu|\mu) + H(\mu|\nu_0), \quad H(\nu_0|\mu) = H(\nu_0|\mu) + H(\nu_o|\nu_0),$$

it follows that $J(\mu, \nu_0)$ is less than both $H(\mu|\nu_0)$ and $H(\nu_0|\mu)$.

Let

$$\mathcal{D}_1 = \{ \mu \in M_1(S) : H(\mu|\nu_0) < H(\nu_0|\mu) \}$$

$$\mathcal{D}_2 = \{ \mu \in M_1(S) : H(\mu|\nu_0) > H(\nu_0|\mu) \}$$

$$\mathcal{D}_3 = \{ \mu \in M_1(S) : H(\mu|\nu_0) = H(\nu_0|\mu) \}.$$

Consider the value of rate function at μ as the energy required to move from ν_0 to μ . Then it requires less energy to move into region \mathcal{D}_1 for the empirical distributions than the Dirichlet processes. The deviation into \mathcal{D}_2 is easier for the Dirichlet processes than the empirical distributions. There will be no difference between the two for region \mathcal{D}_3 . The energy required for a deviation of $\{\mathcal{W}_{n,\nu_0} : n \geq 1\}$ from ν_0 is always the lowest among the three.

For any μ, ν in $M_1(S)$, the relative entropy $H(\mu|\nu)$ originated in information theory as the Kullback-Leibler divergence. It describes the information gain when μ is used in comparison with ν . It is non-negative, convex, and equals to zero only when $\mu = \nu$. But it is not a metric since it is asymmetric, and does not obey the triangle inequality. The asymmetry can be rectified by considering the symmetric divergence $H(\mu|\nu) + H(\nu|\mu)$. But the triangle inequality still does not hold. However, one does have the following Pinsker's inequality:

$$\rho_{tv}^2(\mu,\nu) \le 2\min\{H(\mu|\nu), H(\nu|\mu)\}$$

where $\rho_{tv}(\mu, \nu)$ denote the total variation distance between μ and ν .

The function $J(\mu, \nu)$ defined as in (3.9) provides a new measurement of divergence between μ and ν , which we call the *J*-divergence. It is non-negative, symmetric, convex, and equals to zero only when $\mu = \nu$. The finiteness of $J(\mu, \nu)$ does not require the absolute continuity between μ and ν . This can be seen from the following example:

$$\nu(d\,x) = d\,x, \quad \mu(d\,x) = \frac{1}{2}[d\,x + \delta_{\{0\}}(d\,x)], \quad \nu(d\,x) = \frac{1}{2}[d\,x + \delta_{\{1\}}(d\,x)].$$

It follows from direct calculation that

$$H(v|\mu) = \log 2 = H(v|\nu), \quad J(\mu, \nu) \le 2\log 2.$$

Clearly $H(\mu|\nu) = H(\nu|\mu) = \infty$. Thus $J(\mu, \nu)$ can be strictly less than the minimum of $H(\mu|\nu)$ and $H(\nu|\mu)$. This helps in quantifying the relative information between probabilities that have no absolute continuity relation.

The *J*-divergence does not obey the triangle inequality either. But it satisfies an inequality similar to Pinsker's inequality.

Theorem 3.4 For any μ, ν in $M_1(S)$,

$$\rho_{tv}^2(\mu,\nu) \le 4J(\mu,\nu). \tag{3.10}$$

Proof: One can prove the result using Pinsker's inequality. We choose a direct proof below. Fix μ, ν in $M_1(S)$. For any ν in $M_1(S)$, and any measurable subset A of S, set

$$p = \upsilon(A), \ q = \mu(A), \ \nu(A) = r,$$

$$\pi(\upsilon) = (p, 1 - p), \ \pi(\mu) = (q, 1 - q), \ \pi(\nu) = (r, 1 - r),$$

and

$$L(p) = H(\pi(\upsilon)|\pi(\mu)) + H(\pi(\upsilon)|\pi(\nu)).$$

If q = r, then $|\mu(A) - \nu(A)| = 0$. If q = 0, 0 < r < 1 (the case 0 < q < 1, r = 0 is similar), then L(p) achieves minimum at p = 0, and

$$L(0) = -\log(1-r) \ge r = |\mu(A) - \nu(A)| \ge |\mu(A) - \nu(A)|^2.$$

Next we assume that 0 < q, r < 1.

Solving the equation L'(p) = 0 we obtain

$$\hat{p} = \frac{\sqrt{\frac{qr}{(1-q)(1-r)}}}{1 + \sqrt{\frac{qr}{(1-q)(1-r)}}}.$$

Since $L''(p) \ge 0$, it follows that \hat{p} is the minima for $L(\cdot)$, and

$$L(\hat{p}) = -2\log[\sqrt{(1-q)(1-r)} + \sqrt{qr}].$$

Without loss of generality we assume q > r and set $\delta = q - r$. It is easily checked by direct calculation that the function

$$\sqrt{(1-q)(1-r)} + \sqrt{qr} = \sqrt{(1-r)^2 - \delta(1-r)} + \sqrt{r^2 + \delta r}$$

reaches its maximum $\sqrt{1-\delta^2}$ at $r=\frac{1-\delta}{2}$. Thus

$$L(\hat{p}) \ge -\log[1-\delta^2] \ge \delta^2.$$

Putting all these together we obtain

$$J(\mu, \nu) \geq \inf\{H(\pi(\nu)|\pi(\mu)) + H(\pi(\nu)|\pi(\nu)) : \nu \in M_1(S)\} \\ \geq \sup\{|\mu(A) - \nu(A)|^2 : A \subset S\} = \frac{\rho_{tv}^2(\mu, \nu)}{4}$$

and thus (3.10)

4 Applications

For any f in C(S), consider the random weighted averages $\langle \mathcal{L}_{n,\nu_0}, f \rangle$, $\langle \mathcal{V}_{n,\nu_0}, f \rangle$, and $\langle \mathcal{W}_{n,\nu_0}, f \rangle$. Let ξ have distribution ν_0 . It follows from direct calculation that

$$\mathbb{E}[\langle \mathcal{L}_{n,\nu_0}, f \rangle] = \mathbb{E}[\langle \mathcal{V}_{n,\nu_0}, f \rangle] = \mathbb{E}[\langle \mathcal{W}_{n,\nu_0}, f \rangle] = \langle \nu_0, f \rangle,$$

$$\operatorname{Var}[\langle \mathcal{L}_{n,\nu_0}, f \rangle] = \frac{1}{n} \operatorname{Var}[f(\xi)],$$

$$\operatorname{Var}[\langle \mathcal{V}_{n,\nu_0}, f \rangle] = \frac{1}{n} \operatorname{Var}[f(\xi)] + \frac{n-1}{n(n+1)} \langle \nu_0, f^2 \rangle$$

$$\operatorname{Var}[\langle \mathcal{W}_{n,\nu_0}, f \rangle] = \frac{1}{n+1} \operatorname{Var}[f(\xi)].$$

Thus all three sequences converge to $\langle \nu_0, f \rangle$ in probability as n tends to infinity. The sequence $\{\langle \mathcal{V}_{n,\nu_0}, f \rangle : n \geq 1\}$ has a bigger variance among the three.

For f(x) = x, $\langle \mathcal{L}_{n,\nu_0}, f \rangle$, $\langle \mathcal{V}_{n,\nu_0}, f \rangle$, and $\langle \mathcal{W}_{n,\nu_0}, f \rangle$ are, respectively, the sample mean, the well studied random Dirichlet mean ([12]), and the finite Dirichlet weighted mean. The objective of this section is to compare the large deviation behaviours of the three means when *n* tends to infinity.

Since the map

$$\Phi_f: M_1(S) \to \mathbb{R}, \ \mu \to \langle \mu, f \rangle$$

is continuous, the next result follows by a direct application of Theorem 3.3, the large deviations for the Dirichlet process, and the contraction principle.

Theorem 4.1 Both the family of laws of $\{\langle \mathcal{V}_{n,\nu_0}, f \rangle : n \geq 1\}$ and the family of laws of $\{\langle \mathcal{W}_{n,\nu_0}, f \rangle : n \geq 1\}$ satisfy large deviation principles with respective good rate functions

$$I_{1,f}(u;\nu_0) = \inf_{\mu \in M_1(S)} \{ H(\nu_0|\mu) : \langle \mu, f \rangle = u \}$$

and

$$I_{2,f}(u;\nu_0) = \inf_{\mu \in M_1(S)} \{ J(\mu,\nu_0) : \langle \mu, f \rangle = u \}.$$

Let ν_1 denote the law of $M_{1,\nu_0} = \langle \mathcal{V}_{1,\nu_0}, x \rangle$, $I_i(\cdot, \nu) = I_{i,f}(\cdot, \nu)$ for f(x) = x, i = 1, 2. Then we have

$$I_1(u;\nu_0) = I_2(u;\nu_1).$$
(4.1)

Proof: It suffices to prove (4.1). Let $\{\xi_i : i \ge 1\}$ be i.i.d copies of M_{1,ν_0} with common distribution ν_1 . It follows from (2.5) that

$$M_{n,\nu_0} = \langle \mathcal{V}_{n,\nu_0}, x \rangle$$

= $\frac{1}{\sum_{k=1}^n W_k} \sum_{i=1}^n W_i \xi_i$
= $\langle \mathcal{W}_{n,\nu_1}, x \rangle.$

Since both $I_{1,f}(u;\nu_0)$ and $I_{2,f}(u;\nu_1)$ are good rate functions, the result follows from the uniqueness of good rate function.

The rate function $I_1(u)$ involves the minimization of the relative entropy $H(\nu_0|\mu)$ over the set of probability measures $\{\mu \in M_1(S) : \langle \mu, x \rangle = u\}$. This is called the reverse information projection in comparison with the forward information projection of minimizing $H(\mu|\nu_0)$ for μ over a a subset of $M_1(S)$. The forward information projection over convex set has an explicit solution ([1]), and the reverse information projection has a solution over any log-convex domain([2]). Since the set $\{\mu \in M_1(S) : \langle \mu, x \rangle = u\}$ is not log-convex, it is not clear whether one can identify the minimizer, and thus $I_1(u)$, explicitly in general. Our final result will give an explicit form for $I_1(u)$ in the case $\nu_0(dx) = dx$.

Theorem 4.2 Let ν_0 be the uniform distribution over S. Then

$$I_1(u,\nu_0) = \int_0^1 \log(\alpha + \beta x)\nu_0(dx)$$

= $F^{-1}(u)(u-1) + \log(1 + F^{-1}(u)u),$ (4.2)

where

$$\alpha + \beta = \alpha e^{\beta}, \quad \alpha + \beta u = 1.$$

and

$$F(\lambda) = \begin{cases} \frac{1}{2} & \lambda = 0, \\ \frac{e^{\lambda}}{e^{\lambda} - 1} - \frac{1}{\lambda} & else. \end{cases}$$

Proof: For any μ in $M_1(S)$, let $\mu = \mu_1 + \mu_2$ be the Lebesgue decomposition of μ with respect to ν_0 , with $\mu_1 \ll \nu_0$ and $\mu_2 \perp \nu_0$. Set $f(x) = \frac{d\mu_1}{d\nu_0}$, the Radon-Nikodym derivative of μ_1 with respect to ν_0 . Then for any u in S we have

$$I_{1}(u;\nu_{0}) = \inf\{\langle\nu_{0}, -\log f(x)\rangle : \langle\mu_{1} + \mu_{2}, x\rangle = u, \mu_{1} \equiv \nu_{0}\}$$

= $\inf\{\langle\nu_{0}, -\log f(x)\rangle : f > 0, \nu_{0} - a.s., \langle\nu_{0}, f(x)\rangle \leq 1, \langle\nu_{0}, xf(x)\rangle \leq u\}$ (4.3)
= $\inf_{a \in [0,1], b \in [0, a \land u]} \inf_{f \in \Gamma_{a,b}} \{\langle\nu_{0}, -\log f(x)\rangle\}$

where

$$\Gamma_{a,b} = \{ f \in B(S) : f > 0, \ \nu_0 - a.s., \langle \nu_0, f(x) \rangle = a, \langle \nu_0, xf(x) \rangle = b \}$$

If b = 0, then f(x) = 0 and $\langle \nu_0, -\log f(x) \rangle = \infty$.

For any $0 < a \leq 1$, and $0 < b \leq u \wedge a$, let $\lambda_1(a, b)$ and $\lambda_2(a, b)$ be the solution to the following equations

$$\lambda_1 + \lambda_2 = \lambda_1 e^{a\lambda_2}, \quad a\lambda_1 + b\lambda_2 = 1.$$

Since $a \ge b$, it follows that

$$\lambda_1(a,b) > 0, \ \lambda_1(a,b) + \lambda_2(a,b) > 0.$$

Thus the nonnegative function

$$g_{a,b}(x) = \frac{1}{\lambda_1(a,b) + \lambda_2(a,b)x}$$

is well-defined. It follows from direct calculation that

$$\langle \nu_0, g_{a,b}(x) \rangle = a, \ \langle \nu_0, xg_{a,b}(x) \rangle = b.$$

Thus $g_{a,b}$ is in $\Gamma_{a,b}$. For any f in $\Gamma_{a,b}$, we obtain

$$\int_0^1 \frac{f(x)}{g_{a,b}(x)} \nu_0(dx) = \int_0^1 (\lambda_1(a,b) + \lambda_2(a,b)x) f(x) \nu_0(dx) = 1$$

and

$$\int_0^1 \log \frac{f(x)}{g_{a,b}(x)} \nu_0(d\,x) \le \log \left(\int_0^1 \frac{f(x)}{g_{a,b}(x)} \nu_0(d\,x)\right) = 0.$$

Hence

$$\langle \nu_0, -\log f(x) \rangle \ge \langle \nu_0, -\log g_{a,b}(x) \rangle,$$

and the infimum of $\langle \nu_0, -\log f(x) \rangle$ is achieved at $g_{a,b}$. It is not difficult to see that the map from (a, b) to (λ_1, λ_2) is one-to-one, and

$$\frac{\partial a}{\partial \lambda_1} < 0, \quad \frac{\partial a}{\partial \lambda_2} < 0, \quad \frac{\partial b}{\partial \lambda_1} < 0, \quad \frac{\partial b}{\partial \lambda_2} < 0.$$

It follows that $\langle \nu_0, -\log g_{a,b}(x) \rangle$ is decreasing in both a and b, and the infimum is achieved for a = 1, b = u. Thus by (4.3) we obtain

$$I_1(u;\nu_0) = -\int_0^1 \log g_{1,u}(x)\nu_0(d\,x)$$

which implies (4.2) and the theorem by taking $\alpha = \lambda_1(1, u), \beta = \lambda_2(1, u).$

Remark 4.1 The function $F(\cdot)$ is the cumulative distribution function of a random variable.

Remark 4.2 Let $\nu_0(dx) = dx$, and $\{\xi_i : i \ge 1\}$ be *i.i.d.* with common distribution ν_0 . Then the large deviation rate function for the sample mean

$$\frac{1}{n}\sum_{i=1}^{n}\xi_{i} = \langle \mathcal{L}_{n,\nu_{0}}, x \rangle$$

is

$$I_3(u,\nu_0) = \inf\{H(\mu|\nu_0) : \langle \mu, x \rangle = u\}.$$

By the minimum discrimination information theorem (pages 36–39 in [11]), the infimum in $I_3(u)$ is achieved at measure μ_0 satisfying

$$\mu_0(d\,x) = ce^{rx}\nu_0(d\,x).$$

The constraints

$$\langle \nu_0, ce^{rx} \rangle = 1, \quad \langle \nu_0, cxe^{rx} \rangle = u$$

imply that

$$c = \frac{r}{e^r - 1}, \ ce^r - 1 = ur,$$

and

$$u = \frac{e^r}{e^r - 1} - \frac{1}{r} = F(r).$$

By direct calculation,

$$I_{3}(u) = H(\mu_{0}|\nu_{0}) = \int_{0}^{1} ce^{rx} (\log c + rx) dx$$

= $\log c + ce^{r} - 1 = r(u - 1) + \log(1 + ru)$
= $F^{-1}(u)(u - 1) + \log(1 + F^{-1}(u)u)$

which is the same as $I_1(u, \nu_0)$. Thus the sample means and the Dirichlet means have the same large deviation behaviour.

Remark 4.3 It is known ([17]) that the law of the Dirichlet mean M_{n,ν_0} for $n \geq 2$ is absolutely continuous with respect to ν_0 with the Radon-Nikodym derivative

$$q_n(x) = \frac{n-1}{\pi} \int_{-\infty}^x (x-y)^{n-2} e^{-n \int_0^1 \log|y-z|dz} \sin(n\pi y) dy$$

But it is not clear how one can use this to obtain the explicit form of $I_1(u, \nu_0)$.

References

- I. Csiszár (1975). I-divergence geometry of probability distributions and minimization problems. Ann. Probab. 3, 146–158.
- [2] I. Csiszár and F. Matúš (2003). Information projections revisited. *IEEE Trans. Inf.* Theory 49, No.6, 1474–1490.
- [3] D.A. Dawson and S. Feng (2001). Large deviations for the Fleming–Viot process with neutral mutation and selection, II. *Stoch. Proc. Appl.* **92**, 131–162.
- [4] A. Dembo and O. Zeitouni. Large Deviations Techniques and Applications, 2nd Edition. Springer, New York, 1998.
- [5] M.D. Donsker and S.R.S. Varadhan (1975). Asymptotic evaluation of certain Markov process expectations for large time, I. Comm. Pure Appl. Math., 28, pp. 1-47.
- [6] S. Feng. *The Poisson-Dirichlet Distribution and Related Topics*. Springer, Heidelberg, 2010.
- [7] T.S. Ferguson (1973). A Baysian analysis of some nonparametric problems. Ann. Statist., 1:209–230.
- [8] A.J. Ganesh and N. O'Connell (2000). A large-deviation principle for Dirichlet posteriors. *Bernoulli* 6, No. 6, 1021–1034.
- [9] H.-O. Georgii. *Gibbs Measures and Phase Transitions*. Berlin, De Gruyter, 1988.
- [10] H. Ishwaran and M. Zarapour (2002). Exact and approximate sum-representations for the Dirichlet process. *Canad. J. Statist.* 36, 269–283.
- [11] S. Kullback. Information Theory and Statistics. Wiley, New York, 1959.
- [12] A. Lijoi and I. Prünster (2009). Distributional properties of means of random probability measures. Stat. Surv., 3, 47–95.
- [13] J. Lynch and J. Sethuraman (1987). Large deviations for processes with independent increments. Ann. Probab. 15, 610–627.
- [14] J. Pitman (2018). Random weighted averages, partition structures and generalized arcsine laws. arXiv:1804.07896v1.
- [15] J. Pitman and M. Yor (1997). The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. Ann. Probab., Vol. 25, No. 2, 855-900.

- [16] A.A. Pukhalskii. On functional principle of large deviations. In New Trends in Probability and Statistics, ed. V. Sazonov and T. Shervashidze, pages 198–218, VSP Moks'las, Moskva, 1991.
- [17] E. Regazzini, A. Guglielmi, and G. Di Nunno (2002). Theory and numerical analysis for exact distribution of functionals of a Dirichlet process. *Ann. Statist.*, **30**, 1376–1411.
- [18] I.N. Sanov (1957). On the probability of large deviations of random variables. (Russian)Mat. Sb. N. S. 42 (84), 11–44. (English translation in Select. Transl. Math. Statist. and Probability, Vol. 1 pp. 213–244, 1961.)