

AN EXPONENTIAL INEQUALITY FOR ORTHOMARTINGALE DIFFERENCES RANDOM FIELDS AND SOME APPLICATIONS

DAVIDE GIRAUDO

ABSTRACT. In this paper, we establish an exponential inequality for random fields, which is applied in the context of convergence rates in the law of large numbers and Hölderian weak invariance principle.

1. STATEMENT OF THE EXPONENTIAL INEQUALITY

1.1. Goal and motivations. Understanding the asymptotic behavior of sums of random variables is an important topic in probability. A useful tool for establishing limit theorems is to control the probability that the partial sums or the maximum of the absolute values of the partial sums exceeds a fixed number. A direct application can give convergence rates in the law of large numbers, and such an inequality can be used to check tightness criteria in some functional spaces.

In this paper, we will be concerned by establishing and applying an exponential inequality for the so-called orthomartingale differences random fields introduced in [2]. We make an assumption of invariance in law of the partial sums, which is weaker than stationarity. We control the tails of maximum of partials sums on rectangles of \mathbb{N}^d by two quantities: an exponential term and an other one involving the common distribution function of the increments. We will then formulate two applications of the result. One will concern the convergence rates in the strong law of large numbers. The second one will be about the invariance principle in Hölder spaces, which will be restricted to the case of stationary processes for technical reasons.

Orthomartingales are well-suited for the summation over rectangles of \mathbb{Z}^d because we can use a one dimensional martingale property when we sum according to each fixed coordinate. Moreover, the approximation of stationary random fields by this class of martingales can be done, like in [3, 12, 23, 34]. However, for rates in the law of large numbers and the functional central limit theorem in Hölder spaces, only a few results are available in the literature. Indeed, rates on the law of large number for orthomartingales with polynomial moment have been given in [14, 18, 19], but it seems that the question of exponential moments was only addressed for martingale difference sequences. For the functional central limit theorem in Hölder spaces for random fields, the i.i.d. case was addressed for moduli of regularity of the form t^α , $0 < \alpha < 1/2$ in the case of random fields, and for more general moduli in the case of sequences. It turns

Date: June 25, 2021.

Key words and phrases. orthomartingales; exponential inequality; random fields; law of large numbers; invariance principles in function spaces.

Acknowledgement: “The work is supported by Ministry of Science and Higher Education of the Russian Federation, agreement N 075–15–2019–1619”.

out that the inequality we will present in this paper is a well-suited tool for dealing with these problems.

A key ingredient for the aforementioned limit theorems is an exponential inequality for orthomartingale random fields, that is, an inequality putting into play an exponential term and the tail of the common distribution of the increments. At a first glance, it seems that multi-indexed martingales could be treated like sequences, up to some technical and notational obstacles. However, it turns out that the standard tools for proving deviation inequalities for martingales, like martingale transforms or uses of exponential supermartingale, do not extend easily. Nevertheless, it is possible to apply induction arguments when the random field satisfies good properties.

The paper is organized as follows: in Subsection 1.2, we state the definition of orthomartingale difference random fields and an exponential inequality for such random fields, with an assumption of invariance of the law of the sum on rectangles by translation. Subsections 2.1 (respectively 2.2) provide an application to the rates in the strong law of large numbers (respectively, to the invariance principle in Hölder spaces). Section 3 is devoted to the proof of the previously stated results.

1.2. Exponential inequality for orthomartingales. We start by defining the concept of orthomartingale. Before that, we need to introduce the notion of filtration with respect to the coordinatewise order \preceq : we say that for $\mathbf{i}, \mathbf{j} \in \mathbb{Z}^d$, $\mathbf{i} \preceq \mathbf{j}$ if $i_q \leq j_q$ for each $q \in \{1, \dots, d\} =: [d]$.

Definition 1.1. A filtration indexed by \mathbb{Z}^d on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a collection of σ -algebras $(\mathcal{F}_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ such that for each $\mathbf{i} \preceq \mathbf{j}$, the inclusion $\mathcal{F}_{\mathbf{i}} \subset \mathcal{F}_{\mathbf{j}}$ take place.

Definition 1.2. We say that the filtration $(\mathcal{F}_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is commuting if for all integrable random variable Y and all $\mathbf{i}, \mathbf{j} \in \mathbb{Z}^d$, the equality

$$\mathbb{E}[\mathbb{E}[Y | \mathcal{F}_{\mathbf{i}}] | \mathcal{F}_{\mathbf{j}}] = \mathbb{E}[Y | \mathcal{F}_{\min\{\mathbf{i}, \mathbf{j}\}}], \quad (1.1)$$

takes place, where the minimum is taken coordinatewise.

Example 1.3. Let $(\varepsilon_{\mathbf{j}})_{\mathbf{j} \in \mathbb{Z}^d}$ be an i.i.d. random field and let $\mathcal{F}_{\mathbf{j}} := \sigma(\varepsilon_{\mathbf{i}}, \mathbf{i} \preceq \mathbf{j})$. Then the filtration $(\mathcal{F}_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is commuting.

Example 1.4. Let $(\varepsilon_{\mathbf{k}}^{(q)})_{\mathbf{k} \in \mathbb{Z}^d}$, $q \in [d]$ be independent copies of an i.i.d. sequences $(\varepsilon_k)_{k \in \mathbb{Z}}$ and let $\mathcal{F}_{\mathbf{j}} := \sigma(\varepsilon_{k_q}^{(q)}, k_q \leq j_q, 1 \leq q \leq d)$. Then the filtration $(\mathcal{F}_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is commuting, and connected to decoupled U -statistics.

Definition 1.5. We say that the random field $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is an orthomartingale differences random field with respect to the commuting filtration $(\mathcal{F}_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ if for all $\mathbf{i} \in \mathbb{Z}^d$, $X_{\mathbf{i}}$ is integrable, $\mathcal{F}_{\mathbf{i}}$ -measurable and

$$\mathbb{E}[X_{\mathbf{i}} | \mathcal{F}_{\mathbf{i} - \mathbf{e}_q}] = 0, \quad (1.2)$$

where \mathbf{e}_q is the q -th vector of the canonical basis of \mathbb{R}^d .

Theorem 1.6. Let $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ be an orthomartingale differences random field such that for all $\mathbf{n} \in \mathbb{N}^d$ and all $\mathbf{k} \in \mathbb{Z}^d$,

$$S_{\mathbf{n}} := \sum_{\mathbf{1} \preceq \mathbf{i} \preceq \mathbf{n}} X_{\mathbf{i}} \text{ and } \sum_{\mathbf{1} \preceq \mathbf{i} \preceq \mathbf{n}} X_{\mathbf{i} + \mathbf{k}} \text{ have the same distribution.} \quad (1.3)$$

Then the following inequality holds for all $x > 0$ and all y such that $x^2/y^2 > 3^d$:

$$\mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \leq A_d \exp \left(- \left(\frac{x}{y} \right)^{2/d} \right) + B_d \int_1^{+\infty} \mathbb{P} \{ |X_{\mathbf{1}}| > yu C_d \} u (\log(1+u))^{p_d} du, \quad (1.4)$$

where A_d , B_d and C_d depend only on d , $p_d = 3d + d(d-1)/4$ and $|\mathbf{n}| = \prod_{q=1}^d n_q$.

Remark 1.7. Assume that $(X_i)_{i \in \mathbb{Z}^d}$ is bounded, that is, there exists a constant K such that $|X_i| \leq K$ almost surely for all $i \in \mathbb{Z}^d$. If $x > 3^{d/2} K/C_d$ then we can choose $y = K/C_d$ and inequality (1.4) simplifies as

$$\mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \leq A_d \exp \left(- \left(\frac{C_d x}{K} \right)^{2/d} \right). \quad (1.5)$$

Remark 1.8. The exponent $2/d$ in the exponential term of (1.4) is not improvable, even in the bounded case. To see this, consider d i.i.d. sequences $(\varepsilon_i^{(q)})_{i \in \mathbb{Z}}$ which are independent of each others, in the sense that the collection $\{\varepsilon_{i_q}^{(q)}, i_q \in \mathbb{Z}\}$ is independent. Assume that $\varepsilon_0^{(q)}$ takes the values 1 and -1 with probability $1/2$ for all $q \in [d]$. Let $X_i = \prod_{q=1}^d \varepsilon_{i_q}^{(q)}$. Then

$$\mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \geq \mathbb{P} \left\{ |S_{\mathbf{n}}| > x |\mathbf{n}|^{1/2} \right\} = \mathbb{P} \left\{ \left| \prod_{q=1}^d \frac{1}{\sqrt{n_q}} \sum_{i_q=1}^{n_q} \varepsilon_{i_q}^{(q)} \right| > x \right\}. \quad (1.6)$$

The vector $\left(\frac{1}{\sqrt{n_q}} \sum_{i_q=1}^{n_q} \varepsilon_{i_q}^{(q)} \right)_{q=1}^d$ converges in distribution to $(N_q)_{q=1}^d$, where $(N_q)_{q=1}^d$ is independent and each N_q has a standard normal distribution. Therefore, if $f(x)$ is a function such that for each bounded by 1 orthomartingale differences random field satisfying (1.3) and each $x > 0$

$$\mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \leq f(x), \quad (1.7)$$

then letting $Y := \prod_{q=1}^d |N_q|$, the following inequality should hold for all x : $\mathbb{P}\{Y > x\} \leq f(x)$. Since the \mathbb{L}^p -norm of Y behaves like $p^{d/2}$, the function f cannot decay quicker than $\exp(-Kx^{2/d})$ for some K .

2. APPLICATION TO LIMIT THEOREMS

2.1. Convergence rates in the law of large numbers. A sequence $(X_i)_{i \geq 1}$ satisfies the law of large numbers if the sequence $(n^{-1} \sum_{i=1}^n X_i)_{i \geq 1}$ converges almost surely to zero. This is for example the case of a strictly stationary ergodic sequence where X_1 is integrable. Then arises the question of evaluating the speed of convergence and finding bounds for the large deviation probabilities, namely,

$$\mathbb{P} \left\{ \frac{1}{n} \left| \sum_{i=1}^n X_i \right| > x \right\}. \quad (2.1)$$

This question has been treated in the independent case in [6, 16, 30] under conditions on the \mathbb{L}^p -norm of X_i . The case of martingale differences has also been adressed: under boundedness

of moments of order p and a Cramer-type condition $\sup_{i \geq 1} \mathbb{E}[\exp(|X_i|)] < +\infty$ in [20], a conditional Cramer condition in [21] and under finite exponential moments in [7].

For random fields, we can consider consider large deviation probabilities defined by

$$\mathbb{P} \left\{ \frac{1}{|\mathbf{N}|} \left| \sum_{\mathbf{1} \leq \mathbf{n} \leq \mathbf{N}} X_{\mathbf{i}} \right| > x \right\}, \mathbf{N} \succcurlyeq \mathbf{1}, x > 0. \quad (2.2)$$

Result for orthomartingales with polynomial moment have been given in [14, 18, 19].

Theorem 2.1. *Let $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ be an orthomartingale difference random field satisfying (1.3). Suppose that for some $\gamma > 0$,*

$$\sup_{s > 0} \exp(s^\gamma) \mathbb{P}\{|X_{\mathbf{1}}| > s\} \leq 1. \quad (2.3)$$

Then for each positive x , the following inequality takes place:

$$\mathbb{P} \left\{ \frac{1}{|\mathbf{N}|} \left| \sum_{\mathbf{1} \leq \mathbf{n} \leq \mathbf{N}} X_{\mathbf{i}} \right| > x \right\} \leq C_{1,d,\gamma} \exp \left(-C_{2,d,\gamma} |\mathbf{N}|^{\frac{\gamma}{2+d\gamma}} x^{\frac{2\gamma}{2+d\gamma}} \right), \quad (2.4)$$

where $C_{1,d,\gamma}$ and $C_{2,d,\gamma}$ depend only on d and γ .

2.2. Application to Hölderian invariance principle. Given a sequence of random variables $(X_i)_{i \geq 1}$, a way to understand the asymptotic behavior of the partial sums given by $S_k = \sum_{i=1}^k X_i$ is to define a sequence of random functions $(W_n)_{n \geq 1}$ on the unit interval $[0, 1]$ in the following way: $W_n(k/n) = n^{-1/2} S_k$ for $k \in \{1, \dots, n\}$, $W_n(0) = 0$ and on $((k-1)/n, k/n)$, W_n is linearly interpolated. When the sequence $(X_i)_{i \geq 1}$ is i.i.d., centered and has unit variance, the sequence $(W_n)_{n \geq 1}$ converges in law in $C[0, 1]$. The result has been extended to strictly stationary martingale difference sequences in [1, 17]. Then numerous papers treated the case of weakly dependent strictly stationary sequence, see [22] and the references therein for an overview.

There are also two other possibilities of extension of such result. The first one is to consider other functional spaces, in order to establish the convergence of $(F(W_n))_{n \geq 1}$ for a larger class of functional than the continuous functionals $F: C[0, 1] \rightarrow \mathbb{R}$. In other words, we view W_n as an element of an element of a Hölder space \mathcal{H}_ρ with modulus of regularity ρ and investigate the convergence of $(W_n)_{n \geq 1}$ in this function space. Applications to epidemic changes can be given, for example in [27]. The question of the invariance principle in Hölder spaces has been treated for i.i.d. sequences for modulus of regularity of the form t^α in [25] and also for more general ones, of the form $t^{1/2} \log(ct)^\beta$ in [26]. Some results are also available for stationary weakly dependent sequences, for example mixing sequences [10, 15] or by the use of a martingale approximation [9, 11].

A second development of these invariance principles is the consideration of partial sum processes built on random fields. Given a random field $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$, one can define the following random function on $[0, 1]^d$:

$$W_{\mathbf{n}}(\mathbf{t}) = \frac{1}{\sqrt{|\mathbf{n}|}} \sum_{\mathbf{i} \in \mathbb{Z}^d} \lambda \left(R_{\mathbf{i}} \cap \prod_{q=1}^d [0, n_q t_q] \right) X_{\mathbf{i}}, \mathbf{n} \succcurlyeq \mathbf{1}, \mathbf{t} \in [0, 1]^d, \quad (2.5)$$

where λ denotes the Lebesgue measure on \mathbb{R}^d and $R_i = \prod_{q=1}^d (i_q - 1, i_q)$. Notice that the map $\mathbf{t} \mapsto W_{\mathbf{n}}(\mathbf{t})$ is 1-Lipschitz continuous and that $W_{\mathbf{n}}\left(\left(\frac{k_q}{n_q}\right)_{q=1}^d\right) = S_{k_1, \dots, k_d}$ hence $W_{\mathbf{n}}$ takes into account the values of all the partial sums $S_{\mathbf{k}}$ for $\mathbf{1} \preceq \mathbf{k} \preceq \mathbf{n}$.

Convergence of $W_{\mathbf{n}}$ in the space of continuous functions has been investigated in [33] for an i.i.d. random field, [5] for martingales for the lexicographic order and in [31, 32] for orthomartingales.

In this section, we will study the convergence of $W_{\mathbf{n}}$ in some Hölder spaces for orthomartingales random fields. First, an observation is that for $\mathbf{t}, \mathbf{t}' \in [0, 1]^d$,

$$\left| \lambda \left(R_i \cap \prod_{q=1}^d [0, n_q t_q] \right) - \lambda \left(R_i \cap \prod_{q=1}^d [0, n_q t'_q] \right) \right| \leq \max_{1 \leq q \leq d} n_q |t_q - t'_q| \leq |\mathbf{n}| \|\mathbf{t} - \mathbf{t}'\|_{\infty} \quad (2.6)$$

hence

$$|W_{\mathbf{n}}(\mathbf{t}) - W_{\mathbf{n}}(\mathbf{t}')| \leq \sqrt{|\mathbf{n}|} \|\mathbf{t} - \mathbf{t}'\|_{\infty} \sum_{\mathbf{1} \preceq \mathbf{i} \preceq \mathbf{n}} |X_{\mathbf{i}}| \quad (2.7)$$

hence for almost every ω , the map $\mathbf{t} \mapsto W_{\mathbf{n}}(\mathbf{t})(\omega)$ is Lipschitz-continuous. As pointed out in the i.i.d. case in [33], the finite dimensional distributions of $W_{\mathbf{n}}$ converge to those of a standard Brownian sheet, that is, a centered Gaussian process $(W(\mathbf{t}))_{\mathbf{t} \in [0, 1]^d}$ whose covariance function is given by

$$\text{Cov}(W(\mathbf{t}), W(\mathbf{t}')) = \prod_{q=1}^d \min\{t_q, t'_q\}. \quad (2.8)$$

Given an increasing continuous function $\rho: [0, 1] \rightarrow \mathbb{R}$ such that $\rho(0) = 0$, let \mathcal{H}_{ρ} be the space of all the functions x from $[0, 1]^d$ to \mathbb{R} such that $\sup_{\mathbf{t}, \mathbf{t}' \in [0, 1]^d, \mathbf{t} \neq \mathbf{t}'} |x(\mathbf{t}) - x(\mathbf{t}')| / \rho(\|\mathbf{t} - \mathbf{t}'\|_{\infty})$ is finite.

The Brownian sheet has trajectories in \mathcal{H}_{ρ} with $\rho(h) = h^{\alpha}$ for all $\alpha \in (0, 1/2)$ but not for $\alpha \geq 1/2$. Therefore, it is not possible to expect to show the convergence of $(W_{\mathbf{n}})_{\mathbf{n} \geq \mathbf{1}}$ in all the possible Hölder spaces and some restriction have to be made. In order to treat a class of modulus of regularity larger than the power function, we need to introduce the slowly varying functions. We say that a function $L: (0, +\infty) \rightarrow (0, +\infty)$ is slowly varying if for all $c > 0$, the quantity $L(ct)/L(t)$ goes to 1 as t goes to infinity. For example, functions which behave asymptotically as a power of the logarithm are slowly varying. We now define the moduli of regularity and the associated Hölder spaces of interest from the point of view of the convergence of the partial sum process $W_{\mathbf{n}}$.

Definition 2.2. *Let $d \geq 1$ be an integer. We say that ρ belongs to the class $\mathcal{R}_{1/2, d}$ if there exists a slowly varying function L such that $L(t) \rightarrow \infty$ as t goes to infinity and a constant c such that*

$$\rho(h) = h^{1/2} \left(\ln \left(\frac{c}{h} \right) \right)^{d/2} L \left(\frac{1}{h} \right), \quad h \in [0, 1], \quad (2.9)$$

and ρ is increasing on $[0, 1]$.

We now give a sufficient condition for tightness of the partial sum process associated to a strictly stationary random field, that is, a random field $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ such that for each integer N and each $\mathbf{i}_1, \dots, \mathbf{i}_N, \mathbf{j} \in \mathbb{Z}^d$, the vectors $(X_{\mathbf{i}_k + \mathbf{j}})_{k=1}^N$ and $(X_{\mathbf{i}_k})_{k=1}^N$ have the same distribution. The criterion puts into play tails of the maximum of partial sums of the rectangles.

Proposition 2.3. *Let $d \geq 1$ and ρ be an element of $\mathcal{R}_{1/2,d}$. If $(X_i)_{i \in \mathbb{Z}^d}$ is a strictly stationary random field such that for each $i \in \{1, \dots, d\}$ and each positive ε ,*

$$\lim_{J \rightarrow \infty} \limsup_{\min \mathbf{n} \rightarrow \infty} \sum_{j=J}^{n_i} 2^j \mathbb{P} \left\{ \max_{1 \leq l_i \leq 2^{n_i-j}} \max_{\substack{1 \leq \ell_k \leq 2^{n_k} \\ k \neq i}} \left| \sum_{\mathbf{1} \leq \mathbf{s} \leq \ell} X_{\mathbf{s}} \right| > \varepsilon \rho (2^{-j}) \prod_{q=1}^d 2^{n_q/2} \right\} = 0, \quad (2.10)$$

then the net $(W_{\mathbf{n}})_{\mathbf{n} \geq \mathbf{1}}$ is asymptotically tight in $\mathcal{H}_\rho([0, 1]^d)$.

No other assumption than stationarity is done but of course, some dependence will be required for this condition to be satisfied since one need a good control of the tails of the partial sums on rectangles normalized by the square root of the number of elements in the rectangle.

Now, we state a result for the weak convergence of $(W_{\mathbf{n}})_{\mathbf{n} \geq \mathbf{1}}$ in the space H_ρ , where $\rho \in \mathcal{R}_{1/2,d}$.

Theorem 2.4. *Let $d \geq 1$ and let $(X_i)_{i \in \mathbb{Z}^d}$ be a strictly stationary orthomartingale differences random field. Let ρ be an element of $\mathcal{R}_{1/2,d}$ be given by (2.9), where L is slowly varying.*

Suppose that

$$\forall A > 0, \quad \sum_{j \geq 1} 2^j \mathbb{P} \{ |X_{\mathbf{1}}| > L(2^j) A \} < \infty. \quad (2.11)$$

Then $(W_{\mathbf{n}})_{\mathbf{n} \geq \mathbf{1}}$ is asymptotically tight in $\mathcal{H}_\rho([0, 1]^d)$.

If moreover $(X_i)_{i \in \mathbb{Z}^d}$ is such that any $A \in \mathcal{B}(\mathbb{R}^{\mathbb{Z}^d})$ such that $\{(X_i)_{i \in \mathbb{Z}^d} \in A\} = \{(X_{i+e_d})_{i \in \mathbb{Z}^d} \in A\}$, $\mathbb{P} \{(X_i)_{i \in \mathbb{Z}^d} \in A\} \in \{0, 1\}$, then $(W_{\mathbf{n}})_{\mathbf{n} \geq \mathbf{1}}$ converges to $\|X_{\mathbf{0}}\|_2 W$ in $\mathcal{H}_\rho([0, 1]^d)$, where W is a standard Brownian sheet.

3. PROOFS

3.1. Proof of Theorem 1.6. The proof will be done by induction on the dimension d . For $d = 1$, we need the following deviation inequality for one dimensional martingales. This is a combination of Theorem 2.1 in [8] and Theorem 6 in [29].

Proposition 3.1 (Proposition 2.3 in [13]). *Let $(D_j)_{j \geq 1}$ be a martingale differences sequence with respect to the filtration $(\mathcal{F}_i)_{i \geq 0}$. Suppose that $\mathbb{E}[D_i^2]$ is finite for all $i \geq 1$. Suppose that there exists a random variable Y such that for all $1 \leq i \leq n$, $\mathbb{E}[\varphi(D_i^2)] \leq \mathbb{E}[\varphi(Y^2)]$ for all convex increasing function $\varphi: [0, \infty) \rightarrow [0, \infty)$. Then for all $x, y > 0$ and each $n \geq 1$, the following inequality holds:*

$$\mathbb{P} \left\{ \max_{1 \leq k \leq n} \left| \sum_{i=1}^k D_i \right| > xn^{1/2} \right\} \leq 2 \exp \left(-\frac{1}{2} \left(\frac{x}{y} \right)^2 \right) + 2 \int_1^{+\infty} \mathbb{P} \{ Y^2 > y^2 u/4 \} du. \quad (3.1)$$

Assume that Theorem 1.6 holds in dimension $d - 1$ with $d \geq 2$. We have to prove (1.4) for all d -dimensional orthomartingale differences fields satisfying (1.3). We first get rid of the maximum over the coordinate d , apply the inequality of the $d - 1$ dimension to the obtained orthomartingale and we are then reduced to control the tails of a one dimensional martingale. More concretely, the induction step is done as follows.

(1) Step 1: let $M := \max_{\mathbf{1} \leq i \leq \mathbf{n}} |S_i|$ and

$$M' := \max_{1 \leq i_q \leq n_q, 1 \leq q \leq d-1} |S_{i_1, \dots, i_{d-1}, n_d}|. \quad (3.2)$$

Then we will show that

$$\mathbb{P}\{M > x\} \leq \int_1^\infty \mathbb{P}\{M' > xu/2\} du. \quad (3.3)$$

(2) Step 2: the tails of M' are controlled by applying the result for $(d-1)$ -dimensional random fields.

(3) Step 3: it remains to control the tails of $\sum_{i_d=1}^{n_d} X_{1, \dots, 1, i_d}$, which can be done by using the one dimensional result.

The proof will be quite similar to that of Theorem 1.1 in [13]. The latter gave an exponential inequality in the spirit of those of the paper for U -statistics, that is, sum of terms of the form $h(X_1, \dots, X_r)$ where $(X_i)_{i \geq 1}$ is i.i.d. The connection with orthomartingales is the following. First, using decoupling (see [4]), the following inequality takes place:

$$\mathbb{P}\left\{\left|\sum_{1 \leq i_1 < \dots < i_r \leq n} h(X_{i_1}, \dots, X_{i_r})\right| > x\right\} \leq C \mathbb{P}\left\{\left|\sum_{1 \leq i_1 < \dots < i_r \leq n} h(X_{i_1}^{(1)}, \dots, X_{i_r}^{(r)})\right| > Cx\right\}, \quad (3.4)$$

where C is a universal constant and the sequences $(X_i^{(k)})_{i \in \mathbb{Z}}$, $1 \leq k \leq r$ are mutually independent copies of $(X_i)_{i \in \mathbb{Z}}$. If we assume that $\mathbb{E}[h(X_1, \dots, X_r) | \sigma(X_i, i \in \{1, \dots, r\} \setminus \{k\})] = 0$ for a all $k \in \{1, \dots, r\}$, then the random field $(h(X_{i_1}^{(1)}, \dots, X_{i_r}^{(r)}))_{i_1, \dots, i_r \in \mathbb{Z}}$ is an orthomartingale differences random field for the filtration $(\mathcal{F}_{i_1, \dots, i_r})_{i_1, \dots, i_r \in \mathbb{Z}}$ given by $\mathcal{F}_{i_1, \dots, i_r} = \sigma(\varepsilon_{j_q}^{(q)}, j_q \leq i_q)$.

Let us now go into the details of the proof. Let x, y be positive numbers such that $x/y > 3^d$.

(1) Step 1. Let M and M_N be defined as $M := \max_{\mathbf{1} \leq i \leq \mathbf{n}} |S_i|$, $\mathbf{n} = (n_1, \dots, n_d)$ and

$$M_N := \max_{1 \leq i_q \leq n_q, 1 \leq q \leq d-1} |S_{i_1, \dots, i_{d-1}, N}|. \quad (3.5)$$

Define the filtration \mathcal{G}_N as $\mathcal{F}_{n_1, \dots, n_{d-1}, N}$. We check that $(M_N)_{N \geq 1}$ is a submartingale with respect the filtration $(\mathcal{G}_N)_{N \geq 1}$. Indeed,

$$\mathbb{E}[M_N | \mathcal{G}_{N-1}] \geq \max_{1 \leq i_q \leq n_q, 1 \leq q \leq d-1} |\mathbb{E}[S_{i_1, \dots, i_{d-1}, N} | \mathcal{G}_{N-1}]| \quad (3.6)$$

and by the orthomartingale property,

$$\mathbb{E}[S_{i_1, \dots, i_{d-1}, N} | \mathcal{G}_{N-1}] = S_{i_1, \dots, i_{d-1}, N-1}. \quad (3.7)$$

By Doob's inequality, we derive that

$$x \mathbb{P}\{M > x\} \leq \mathbb{E}[M_{n_d} \mathbf{1}\{M > x\}]. \quad (3.8)$$

Expressing the latter expectation as an integral of the tail and cutting this integral at $x/2$ gives the bound

$$\mathbb{P}\left\{\max_{\mathbf{1} \leq i \leq \mathbf{n}} |S_i| > x |\mathbf{n}|^{1/2}\right\} \leq \int_1^{+\infty} \mathbb{P}\left\{\max_{1 \leq i_q \leq n_q, 1 \leq q \leq d-1} |S_{i_1, \dots, i_{d-1}, n_d}| > x |\mathbf{n}|^{1/2} u/2\right\} du. \quad (3.9)$$

(2) Step 2. Define for i_1, \dots, i_{d-1} the random variable

$$X'_{i_1, \dots, i_{d-1}} := \sum_{i_d=1}^{n_d} X_{i_1, \dots, i_{d-1}, i_d}. \quad (3.10)$$

Then $(X'_{i_1, \dots, i_{d-1}})_{i_1, \dots, i_{d-1} \in \mathbb{Z}}$ is an orthomartingale differences random field with respect to the commuting filtration $(\mathcal{F}_{i_1, \dots, i_{d-1}, n_d})_{i_1, \dots, i_{d-1} \in \mathbb{Z}}$ satisfying (1.3). We thus apply the induction hypothesis with

$$\tilde{x} := xn_d^{1/2}u/2, \quad \tilde{y} := n_d^{1/2}x^{1/d}y^{\frac{d-1}{d}}u(1+2\ln(u))^{-(d-1)\frac{1}{2}}. \quad (3.11)$$

We get in view of (3.9) and

$$\left(\frac{\tilde{x}}{\tilde{y}}\right)^{\frac{2}{d-1}} = \left(\frac{x}{y}\right)^{2/d} (1+2\ln u) > 3 \quad (3.12)$$

that

$$\begin{aligned} & \mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \leq A_{d-1} \int_1^{+\infty} \exp \left(- \left(\frac{x}{y} \right)^{2/d} (1+2\ln u) \right) du \\ & + B_{d-1} \int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ |X'_{1, \dots, 1}| > C_{d-1} v n_d^{1/2} x^{1/d} y^{\frac{d-1}{d}} u (1+2\ln(u))^{-(d-1)\frac{1}{2}} \right\} v (\log(1+v))^{p_{d-1}} dudv. \end{aligned} \quad (3.13)$$

For the first term, using that $\left(\frac{x}{y}\right)^{2/d} > 3$, we get

$$\int_1^{+\infty} \exp \left(- \left(\frac{x}{y} \right)^{2/d} (1+2\ln u) \right) du \leq \exp \left(- \left(\frac{x}{y} \right)^{2/d} \right) \int_1^{+\infty} \exp(-6\ln u) du \quad (3.14)$$

hence

$$\begin{aligned} & \mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} \leq \frac{A_{d-1}}{5} \exp \left(- \left(\frac{x}{y} \right)^{2/d} \right) \\ & + B_{d-1} \int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ \left| \sum_{i_d=1}^{n_d} X_{1, \dots, 1, i_d} \right| > C_{d-1} v n_d^{1/2} x^{1/d} y^{\frac{d-1}{d}} u (1+2\ln(u))^{-(d-1)\frac{1}{2}} \right\} v (\log(1+v))^{p_{d-1}} dudv. \end{aligned} \quad (3.15)$$

(3) Step 3. It remains to find a bound for the double integral. For fixed u and $v > 1$, we apply Proposition 3.1 in the following setting: x is replaced by \tilde{x} defined by

$$\tilde{x} := C_{d-1} v n_d^{1/2} x^{1/d} y^{\frac{d-1}{d}} u (1+2\ln(u))^{-(d-1)\frac{1}{2}} \quad (3.16)$$

and y by \tilde{y} defined by

$$\tilde{y} := C_{d-1} y u v (1+2\ln u)^{-\frac{d-1}{2}} 2^{-1/2} (1+2\ln(uv))^{-1/2}. \quad (3.17)$$

In this way,

$$\frac{1}{2} \left(\frac{\tilde{x}}{\tilde{y}} \right)^2 = \left(\frac{x}{y} \right)^{2/d} (1+2\ln(uv)) \quad (3.18)$$

and (3.1) gives

$$\begin{aligned} \mathbb{P} \left\{ \max_{1 \leq i \leq n} |S_i| > x |\mathbf{n}|^{1/2} \right\} &\leq \frac{A_{d-1}}{5} \exp \left(- \left(\frac{x}{y} \right)^{2/d} \right) \\ &+ B_{d-1} \exp \left(- \left(\frac{x}{y} \right)^{2/d} \right) \int_1^{+\infty} \int_1^{+\infty} \exp(-2 \ln(uv)) \, dudv \\ &+ B_{d-1} \int_1^{+\infty} \int_1^{+\infty} \int_1^{+\infty} h(u, v, w) \, dudvdw, \quad (3.19) \end{aligned}$$

where

$$h(u, v, w) := \mathbb{P} \left\{ |X_1| > C_{d-1} y u v w (1 + 2 \ln u)^{-\frac{d-1}{2}} 2^{-1/2} (1 + 2 \ln(uv))^{-1/2} \right\} v w (\log(1+v))^{p_{d-1}}. \quad (3.20)$$

It remains to rearrange the last triple integral. To this aim, we first use the substitution $u' = u^2$ and $v' = v^2$ in order to get

$$\begin{aligned} &\int_1^{+\infty} \int_1^{+\infty} \int_1^{+\infty} h(u, v, w) \, dudvdw \\ &\leq \int_1^{+\infty} \int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ |X_1| > C_{d-1} y \sqrt{u'v'} w (1 + \ln u')^{-\frac{d}{2}} 2^{1/2} (1 + \ln(v'))^{-1/2} \right\} w (\log(1+v^2))^{p_{d-1}} \, dudvdw. \end{aligned}$$

Then we will use the following lemma.

Lemma 3.2 (Lemma 2.4 in [13]). *Let X be a non-negative random variable and let q, q' be non-negative numbers. Define*

$$\kappa_q = \begin{cases} 1 & \text{if } q \leq 1 \\ e^{q-1}/q^q & \text{if } q > 1. \end{cases} \quad (3.21)$$

Then

$$\begin{aligned} &\int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ X > s (1 + \ln(s))^{-q_1} t \right\} (1 + \ln(t))^{q_2} \, dt ds \\ &\leq \frac{2^{q_1}}{\kappa_{q_1}} \left(1 + \ln \frac{\kappa_{2q_1}}{\kappa_{q_1}} \right)^{q_1} \int_1^{+\infty} \mathbb{P} \left\{ X > t/\kappa_{q_1} \right\} (1 + \ln t)^{q_1+q_2+1} \, dt. \quad (3.22) \end{aligned}$$

To conclude, we apply Lemma 3.2: for a fixed v , with

$$X = X_1^2 / \left(2C_{d-1} y \sqrt{v} (1 + \ln(v))^{-1/2} \right)^2, \quad (3.23)$$

$q_1 = d/2$ and $q_2 = 0$ to get

$$\begin{aligned} &\int_1^{+\infty} \int_1^{+\infty} \int_1^{+\infty} h(u, v, w) \, dudvdw \\ &\leq c_d \int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ X_1^2 / \left(2C_{d-1} y \sqrt{v} (1 + \ln(v))^{-1/2} \right)^2 > t/c_d \right\} (1 + \ln t)^{1+d/2} (\log(1+v^2))^{p_{d-1}} \, dt dv. \quad (3.24) \end{aligned}$$

Using the same lines of the proof of Lemma 3.2, we can show that

$$\begin{aligned} \int_1^{+\infty} \int_1^{+\infty} \mathbb{P} \left\{ X > s (1 + \ln(s))^{-1/2} t \right\} (1 + \ln(s))^{q_2} (1 + \ln(t))^{q_2} dt ds \\ \leq C_{q_1, q_2} \int_1^{+\infty} \mathbb{P} \{ X > t/C_{q_1, q_2} \} (1 + \ln t)^{q_1 + q_2 + 2} dt. \end{aligned} \quad (3.25)$$

Using this with $q_1 = 1 + d/2$ and $q_2 = p_{d-1}$ and noticing that $p_d = 3 + d/2 + p_{d-1}$ ends the proof of Theorem 1.6.

3.2. Proof of Theorem 2.1. We apply Theorem 1.6 to $y = |\mathbf{N}|^{1/(2+d\gamma)} x^{2/(2d+\gamma)}$. Bounding the resulting integral term by a constant times $\exp(-y^\gamma)$, one sees that in this case, the two terms of the right hand side of (1.4) have a similar contribution.

3.3. Proof of Proposition 2.3. The proof of the tightness criterion rests on the Schauder decomposition of the spaces $\mathcal{H}_\rho^o([0, 1]^d)$. In order to state it, we need to introduce the following notations.

Set for $j \geq 0$,

$$W_j := \{k2^{-j}, 0 \leq k \leq 2^j\}^d \quad (3.26)$$

and

$$V_0 := W_0, \quad V_j := W_j \setminus W_{j-1}, j \geq 1. \quad (3.27)$$

We define for $\mathbf{v} \in V_j$ the pyramidal function $\Lambda_{j, \mathbf{v}}$ by

$$\Lambda_{j, \mathbf{v}}(\mathbf{t}) := \Lambda(2^j(\mathbf{t} - \mathbf{v})), \quad \mathbf{t} \in [0, 1]^d, \quad (3.28)$$

where

$$\Lambda(\mathbf{t}) := \max \left\{ 0, 1 - \max_{t_i < 0} |t_i| - \max_{t_i > 0} |t_i| \right\}, \quad \mathbf{t} = (t_i)_{i=1}^d \in [-1, 1]^d. \quad (3.29)$$

For $x \in \mathcal{H}_\rho^o([0, 1]^d)$, we define the coefficients $\lambda_{j, \mathbf{v}}(x)$ by $\lambda_{0, \mathbf{v}}(x) = x(\mathbf{v})$, $\mathbf{v} \in V_0$ and for $j \geq 1$ and $\mathbf{v} \in V_j$,

$$\lambda_{j, \mathbf{v}}(x) := x(\mathbf{v}) - \frac{1}{2} (x(\mathbf{v}^-) + x(\mathbf{v}^+)), \quad (3.30)$$

where \mathbf{v}^+ and \mathbf{v}^- are define in the following way. Each $\mathbf{v} \in V_j$ is represented in a unique way by $\mathbf{v} = (k_i 2^{-j})_{i=1}^d$. Then $\mathbf{v}^+ := (v_i^+)_{i=1}^d$ and $\mathbf{v}^- := (v_i^-)_{i=1}^d$ are defined by

$$\mathbf{v}_i^- := \begin{cases} v_i - 2^{-j}, & \text{if } k_i \text{ is odd;} \\ v_i, & \text{if } k_i \text{ is even} \end{cases} \quad \mathbf{v}_i^+ := \begin{cases} v_i + 2^{-j}, & \text{if } k_i \text{ is odd;} \\ v_i, & \text{if } k_i \text{ is even.} \end{cases} \quad (3.31)$$

The sequential norm is defined by

$$\|x\|_\rho^{\text{seq}} := \sup_{j \geq 0} \rho (2^{-j})^{-1} \max_{\mathbf{v} \in V_j} |\lambda_{j, \mathbf{v}}(x)|, \quad x \in \mathcal{H}_\rho^o([0, 1]^d). \quad (3.32)$$

By [24], the norm $\|\cdot\|_\rho^{\text{seq}}$ is equivalent to $\|\cdot\|_\rho$ on $\mathcal{H}_\rho^o([0, 1]^d)$.

A general tightness criterion is available for moduli of the form $\rho: h \mapsto h^\alpha$. The criterion rests on a Schauder decomposition of \mathcal{H}_ρ^o as $\bigoplus_{j \geq 1} E_j$, where E_j is the vector space generated by the functions $\Lambda_{j, \mathbf{v}}$, $\mathbf{v} \in V_j$.

Theorem 3.3 (Theorem 6, [28]). *Let $\{\zeta_n, \mathbf{n} \in \mathbb{N}^d\}$ and ζ be random elements with values in the space $\mathcal{H}_\alpha([0, 1]^d)$. Assume that the following conditions are satisfied.*

- (1) For each dyadic $\mathbf{t} \in [0, 1]^d$, the net $\{\zeta_{\mathbf{n}}(\mathbf{t}), \mathbf{n} \in \mathbb{N}^d\}$ is asymptotically tight on \mathbb{R} .
(2) For each positive ε ,

$$\lim_{J \rightarrow \infty} \limsup_{\min \mathbf{n} \rightarrow \infty} \mathbb{P} \left\{ \sup_{j \geq J} 2^{\alpha j} \max_{\mathbf{v} \in V_j} |\lambda_{j, \mathbf{v}}(\zeta_{\mathbf{n}})| > \varepsilon \right\} = 0. \quad (3.33)$$

Then the net $\{\zeta_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$ is asymptotically tight in the space $\mathcal{H}_\alpha([0, 1]^d)$.

First observe that by bounding from below the sum over j by the term at index $j = J$ and taking $i = d$, (2.10) implies that

$$\lim_{J \rightarrow \infty} \limsup_{\min \mathbf{n} \rightarrow \infty} 2^J \mathbb{P} \left\{ \max_{1 \leq \ell_d \leq 2^{n_d - J}} \max_{\substack{1 \leq \ell_k \leq 2^{n_k} \\ 1 \leq k \leq d-1}} \left| \sum_{1 \leq s \leq \ell} X_s \right| > \varepsilon \rho (2^{-J}) \prod_{q=1}^{d-1} 2^{n_q/2} 2^{(n_d - J)/2} \right\} = 0, \quad (3.34)$$

and doing the replacement of index $n'_d = n_d - J$ for a fixed J gives asymptotic tightness of $(W_{\mathbf{n}}(\mathbf{t}))_{\mathbf{n} \geq 1}$ for each \mathbf{t} .

It remains to check the second condition of Theorem 3.3. Since the Schauder decomposition $\mathcal{H}_\rho = \bigoplus_{j \geq 1} E_j$ is also valid for $\rho \in \mathcal{R}_{1/2, d}$, this theorem also holds with the map $h \mapsto h^\alpha$ replaced by ρ . Therefore, it suffices to prove that

$$\lim_{J \rightarrow \infty} \limsup_{\min \mathbf{n} \rightarrow \infty} \mathbb{P} \left\{ \sup_{j \geq J} \rho (2^{-j})^{-1} \max_{\mathbf{v} \in V_j} |\lambda_{j, \mathbf{v}}(W_{\mathbf{n}})| > \varepsilon \right\} = 0. \quad (3.35)$$

Consider for $\mathbf{s} := (s_2, \dots, s_d) \in [0, 1]^{d-1}$ and $t, t' \in [0, 1]$ the quantity

$$\Delta_n(t, t', \mathbf{s}) := \sqrt{|\mathbf{n}|} |W_{\mathbf{n}}(t', \mathbf{s}) - W_{\mathbf{n}}(t, \mathbf{s})|. \quad (3.36)$$

We recall the following lemma:

Lemma 3.4 (Lemma 11, [28]). *For any $t', t \in [0, 1]$, $t' > t$, the following inequality holds:*

$$\begin{aligned} \sup_{\mathbf{s} \in [0, 1]^{d-1}} \Delta_n(t, t', \mathbf{s}) &\leq 3^d \mathbf{1} \left\{ t' - t \geq \frac{1}{n_1} \right\} \max_{\substack{1 \leq k_l \leq n_l \\ 2 \leq l \leq d}} \left| \sum_{i_1 = [n_1 t] + 1}^{[n_1 t']} \sum_{\substack{1 \leq i_l \leq k_l \\ 2 \leq l \leq d}} X_{i_l} \right| + \\ &\quad + 3^d \min \{1, n_1(t' - t)\} \max_{1 \leq i_1 \leq n_1} \max_{\substack{1 \leq k_l \leq n_l \\ 2 \leq l \leq d}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ 2 \leq l \leq d}} X_{i_l} \right|. \end{aligned} \quad (3.37)$$

Now, we define for $q \in [d]$ and $\mathbf{s} = (s_\ell)_{\ell \in [d] \setminus \{q\}} \in [0, 1]^{d-1}$,

$$\Delta_n^{(q)}(t, t', \mathbf{s}) := |W_{\mathbf{n}}(s_1, \dots, s_{q-1}, t', s_{q+1}, \dots, s_d) - W_{\mathbf{n}}(s_1, \dots, s_{q-1}, t, s_{q+1}, \dots, s_d)|.$$

By definition of $\lambda_{j, \mathbf{v}}$ and V_j , the inequality

$$\begin{aligned} \mathbb{P} \left\{ \sup_{j \geq J} \rho (2^{-j})^{-1} \max_{\mathbf{v} \in V_j} |\lambda_{j, \mathbf{v}}(W_{\mathbf{n}})| > 2^{d+1} 3^d x \right\} &\leq \\ &\sum_{q=1}^d \mathbb{P} \left\{ \sup_{j \geq J} \rho (2^{-j})^{-1} \max_{\substack{0 \leq k < 2^j \\ \mathbf{0} \leq \mathbf{u} \leq 2^j \mathbf{1}}} \Delta_n^{(q)}(t_{k+1}, t_k; \mathbf{s}_{\mathbf{u}}) > 2 \cdot 3^d x \right\} \end{aligned} \quad (3.38)$$

takes place, where $t_k = k2^{-j}$ and $\mathbf{s}_u = (u_i 2^{-j})_{i \in [d] \setminus \{q\}}$.

We have in view of Lemma 3.4 that for each $q \in [d]$,

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{j \geq J} \rho(2^{-j})^{-1} \max_{\substack{0 \leq k < 2^j \\ \mathbf{0} \preceq \mathbf{u} \preceq 2^j \mathbf{1}}} \Delta_n^{(q)}(t_{k+1}, t_k; \mathbf{s}_u) > 2x |\mathbf{n}|^{1/2} \right\} \\
& \leq \mathbb{P} \left\{ \sup_{j \geq J} \max_{0 \leq a < 2^j} \rho(2^{-j})^{-1} \mathbf{1} \left\{ 2^{-j} \geq \frac{1}{n_q} \right\} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q(a+1)2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| + \right. \\
& \quad \left. + \min\{1, n_q 2^{-j}\} \max_{1 \leq i_q \leq n_q} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > 2x |\mathbf{n}|^{1/2} \right\} \\
& \leq \mathbb{P} \left\{ \sup_{j \geq J} \max_{0 \leq a < 2^j} \rho(2^{-j})^{-1} \mathbf{1} \left\{ 2^{-j} \geq \frac{1}{n_q} \right\} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q(a+1)2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} + \\
& \quad + \mathbb{P} \left\{ \sup_{j \geq J} \rho(2^{-j})^{-1} \min\{1, n_q 2^{-j}\} \max_{1 \leq i_q \leq n_q} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\}. \quad (3.39)
\end{aligned}$$

Since the indicator in the first term of the right hand side of (3.39) vanishes if $j > \log n_q$, we have

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{j \geq J} \max_{0 \leq a < 2^j} \rho(2^{-j})^{-1} \mathbf{1} \left\{ 2^{-j} \geq \frac{1}{n_q} \right\} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q(a+1)2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} \\
& \leq \mathbb{P} \left\{ \sup_{J \leq j \leq \log n_q} \max_{0 \leq a < 2^j} \rho(2^{-j})^{-1} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q(a+1)2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} \\
& \leq \sum_{j=J}^{\log n_q} 2^j \max_{0 \leq a < 2^j} \mathbb{P} \left\{ \rho(2^{-j})^{-1} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q(a+1)2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\}, \quad (3.40)
\end{aligned}$$

and by stationarity, it follows that

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{j \geq J} \max_{0 \leq a < 2^j} \rho (2^{-j})^{-1} \mathbf{1} \left\{ 2^{-j} \geq \frac{1}{n_q} \right\} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{i_q = [n_q a 2^{-j} + 1]}^{[n_q (a+1) 2^{-j}]} \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} \\ & \leq \sum_{j=J}^{\log n_q} 2^j \mathbb{P} \left\{ \rho (2^{-j})^{-1} \max_{0 \leq i_q \leq 2n_q 2^{-j}} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\}. \end{aligned} \quad (3.41)$$

For the second term of the right hand side of (3.39), notice that

$$\sup_{j \geq J} \rho (2^{-j})^{-1} \min \{1, n_q 2^{-j}\} \leq c_\rho \rho (1/n_q)^{-1}. \quad (3.42)$$

Indeed, if $j \leq \log n_q$, then $2^j \leq n_q$ hence $\rho (2^{-j})^{-1} \min \{1, n_q 2^{-j}\} = \rho (2^{-j})^{-1}$ and the sequence $(\rho (2^{-j})^{-1})_j$ is increasing, and if $j > \log n_q$, then $2^j > n_q$, hence $\min \{1, n_q 2^{-j}\} = n_q 2^{-j}$ and for such j 's, we have $\rho (2^{-j})^{-1} n_q 2^{-j}$ and we use decreasingness of the sequence $(\rho (2^{-j})^{-1} 2^{-j})_j$. As a consequence, after having bounded the probability of the max over i_q by the sum of probabilities and used stationarity, we obtain

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{j \geq J} 2^{\alpha j} \min \{1, n_q 2^{-j}\} \max_{1 \leq i_q \leq n_q} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} \\ & \leq n_q \mathbb{P} \left\{ \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \rho (1/n) / c \right\} \end{aligned} \quad (3.43)$$

hence

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{j \geq J} 2^{\alpha j} \min \{1, n_q 2^{-j}\} \max_{1 \leq i_q \leq n_q} \max_{\substack{1 \leq k_l \leq n_l \\ l \neq q}} \left| \sum_{\substack{1 \leq i_l \leq k_l \\ l \neq q}} X_i \right| > x |\mathbf{n}|^{1/2} \right\} \\ & \leq C_\rho \sum_{j=J}^{n_i} 2^j \mathbb{P} \left\{ \max_{1 \leq l_i \leq 2^{n_i-j}} \max_{\substack{1 \leq \ell_k \leq 2^{n_k} \\ k \neq i}} \left| \sum_{1 \leq s \leq \ell} X_s \right| > \varepsilon \rho (2^{-j}) \prod_{q=1}^d 2^{n_q/2} \right\}. \end{aligned} \quad (3.44)$$

Combining inequalities (3.39) with (3.41) and (3.44), we obtain (3.35). This ends the proof of Proposition 2.3.

3.4. Proof of Theorem 2.4. We have to check that (2.10) is satisfied. For simplicity, we will do this for $i = d$; the general case can be done similarly. To this aim, we apply inequality (1.4) for fixed $\mathbf{n} \succcurlyeq \mathbf{1}$, $J \geq 1$ and $j \in \{J, \dots, n_d\}$ in the following setting: $\tilde{\mathbf{n}} = (2^{n_1}, \dots, 2^{n_{d-1}}, 2^{n_d-j})$, $x = \varepsilon \rho (2^{-j}) 2^{j/2}$ and $y = \varepsilon / 2 \rho (2^{-j}) 2^{j/2} j^{d/2}$. The sum of the exponential terms in (1.4) can be

bounded by the remainder of a convergent series. With the assumption on ρ , the sum of the obtained integral terms does not exceed $\sum_{j \geq J} 2^j \int_1^{+\infty} \mathbb{P}\{|X_{\mathbf{1}}| > L(2^j) u C_d\} u (\log(1+u))^{p_d} du$ where C depends only on ρ and ε .

Now, we will show that (2.11) guarantees the convergence to zero of the previous term as J goes to infinity. As there exists constants κ and q such that $u (\log(1+u))^{p_d} \leq \kappa u^q$ for each $u \geq 1$, it suffices to prove that for each C ,

$$\sum_{j \geq 1} 2^j \int_1^{+\infty} \mathbb{P}\{|X_{\mathbf{1}}| > L(2^j) u C_d\} u^q du < \infty \quad (3.45)$$

This will be a consequence of the following lemmas.

Lemma 3.5. *Let $L: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a slowly varying increasing function. There exists a constant C_L such that for each $k \geq 1$,*

$$\sum_{j=1}^k \frac{2^j}{L(2^j)} \leq C_L \frac{2^k}{L(2^k)}. \quad (3.46)$$

Proof. Since L is slowly varying, there exists k_0 be such that for each $k \geq k_0$, $L(2^k) > 2L(2^{k+1})/3$. We show by induction over $k \geq k_0$ that for each $k \geq k_0$,

$$\sum_{j=k_0}^k \frac{2^j}{L(2^j)} \leq 4 \frac{2^k}{L(2^k)}. \quad (3.47)$$

The initial case $k = k_0$ is obvious. If (3.47) is true for some $k \geq k_0$, let us show it for $k+1$. First, splitting the sum and using the induction assumption gives

$$\sum_{j=k_0}^{k+1} \frac{2^j}{L(2^j)} \leq 4 \frac{2^k}{L(2^k)} + \frac{2^{k+1}}{L(2^{k+1})} \quad (3.48)$$

and since $k \geq k_0$,

$$\sum_{j=k_0}^{k+1} \frac{2^j}{L(2^j)} \leq 4 \frac{2^k}{\frac{2}{3}L(2^{k+1})} + \frac{2^{k+1}}{L(2^{k+1})} = 4 \frac{2^{k+1}}{L(2^{k+1})}. \quad (3.49)$$

Let $a_j = 2^j/L(2^j)$, $s_k = \sum_{j=1}^k a_j$ and $c := s_{k_0}/\inf_{k \geq k_0} a_k$. Then for $k \geq k_0$, $s_k \leq (4+c)a_k$ and for $k < k_0$, $s_k \leq \max_{1 \leq \ell \leq k_0-1} s_\ell/a_\ell a_k$ hence we can take

$$c_L := \max \left\{ 4 + c, \max_{1 \leq \ell \leq k_0-1} s_\ell/a_\ell \right\}.$$

This ends the proof of Lemma 3.5. \square

Lemma 3.6. *Let X be a non-negative random variable and let $L: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a slowly varying increasing function such that $L(x) \rightarrow \infty$ as $x \rightarrow \infty$. Suppose that*

$$\forall A > 0, \sum_{j \geq 1} \mathbb{P}\{X > L(2^j) A\} < \infty. \quad (3.50)$$

Then for each $p \geq 1$ and each $C > 0$,

$$\sum_{j \geq 1} 2^j \int_1^{+\infty} \mathbb{P}\{X > L(2^j) u C\} u^p du < \infty \quad (3.51)$$

Proof. Without loss of generality, we can assume that $C = 1$. First, observe that

$$\int_1^{+\infty} \mathbb{P}\{X > L(2^j)u\} u^p du \leq \mathbb{E} \left[\left(\frac{X}{L(2^j)} \right)^{p+1} \mathbf{1}\{X > L(2^j)\} \right]. \quad (3.52)$$

Therefore,

$$\begin{aligned} & \sum_{j \geq 1} 2^j \int_1^{+\infty} \mathbb{P}\{X > L(2^j)u\} u^p du \\ & \leq \sum_{j \geq 1} 2^j \mathbb{E} \left[\left(\frac{X}{L(2^j)} \right)^{p+1} \sum_{k \geq j} \mathbf{1}\{L(2^k) < X \leq L(2^{k+1})\} \right] \\ & \leq \sum_{j \geq 1} 2^j \mathbb{E} \left[\left(\frac{L(2^{k+1})}{L(2^j)} \right)^{p+1} \sum_{k \geq j} \mathbf{1}\{L(2^k) < X \leq L(2^{k+1})\} \right]. \end{aligned} \quad (3.53)$$

Switching the sums over j and k and using Lemma 3.6 reduces use to show that

$$\sum_{k \geq 1} 2^k \mathbb{P}\{L(2^k) < X \leq L(2^{k+1})\} < \infty, \quad (3.54)$$

which follows from the assumption and the obvious bound $\mathbb{P}\{L(2^k) < X \leq L(2^{k+1})\} \leq \mathbb{P}\{L(2^k) < X\}$. □

The last part of the statement of Theorem 2.4 follow from [32]. This ends the proof of Theorem 2.4.

REFERENCES

- [1] Patrick Billingsley, *The Lindeberg-Lévy theorem for martingales*, Proc. Amer. Math. Soc. **12** (1961), 788–792. MR 0126871 (23 #A4165) 4
- [2] Renzo Cairoli, *Un théorème de convergence pour martingales à indices multiples*, C. R. Acad. Sci. Paris Sér. A-B **269** (1969), A587–A589. MR 0254912 (40 #8119) 1
- [3] Ch. Cuny, J. Dedecker, and D. Volný, *A functional CLT for fields of commuting transformations via martingale approximation*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **441** (2015), no. Veroyatnost’i Statistika. 22, 239–262. MR 3504508 1
- [4] Victor H. de la Peña and S. J. Montgomery-Smith, *Decoupling inequalities for the tail probabilities of multivariate U -statistics*, Ann. Probab. **23** (1995), no. 2, 806–816. MR 1334173 7
- [5] Jérôme Dedecker, *Exponential inequalities and functional central limit theorems for a random fields*, ESAIM Probab. Statist. **5** (2001), 77–104. MR 1875665 (2003a:60054) 5
- [6] P. Erdős, *On a theorem of Hsu and Robbins*, Ann. Math. Statistics **20** (1949), 286–291. MR 30714 3
- [7] Xiequan Fan, Ion Grama, and Quansheng Liu, *Large deviation exponential inequalities for supermartingales*, Electron. Commun. Probab. **17** (2012), no. 59, 8. MR 3005732 4
- [8] ———, *Exponential inequalities for martingales with applications*, Electron. J. Probab. **20** (2015), no. 1, 22. MR 3311214 6
- [9] Davide Giraud, *Holderian weak invariance principle under a Hannan type condition*, Stochastic Process. Appl. **126** (2016), no. 1, 290–311. MR 3426520 4
- [10] ———, *Holderian weak invariance principle for stationary mixing sequences*, J. Theoret. Probab. **30** (2017), no. 1, 196–211. MR 3615086 4
- [11] ———, *Hölderian weak invariance principle under the Maxwell and Woodroffe condition*, Braz. J. Probab. Stat. **32** (2018), no. 1, 172–187. MR 3770868 4

- [12] ———, *Invariance principle via orthomartingale approximation*, Stoch. Dyn. **18** (2018), no. 6, 1850043, 29. MR 3869881 [1](#)
- [13] Davide Giraud, *An exponential inequality for U -statistics of i.i.d. data*, arXiv e-prints (2019), arXiv:1911.05502. [6](#), [7](#), [9](#)
- [14] Davide Giraud, *Deviation inequalities for Banach space valued martingales differences sequences and random fields*, ESAIM Probab. Stat. **23** (2019), 922–946. MR 4046858 [1](#), [4](#)
- [15] D. Hamadouche, *Invariance principles in Hölder spaces*, Portugal. Math. **57** (2000), no. 2, 127–151. MR 1759810 (2001g:60075) [4](#)
- [16] P. L. Hsu and Herbert Robbins, *Complete convergence and the law of large numbers*, Proc. Nat. Acad. Sci. U.S.A. **33** (1947), 25–31. MR 19852 [3](#)
- [17] I. A. Ibragimov, *A central limit theorem for a class of dependent random variables*, Teor. Veroyatnost. i Primenen. **8** (1963), 89–94. MR 0151997 (27 #1978) [4](#)
- [18] Anna Kuczmaszewska and Zbigniew A. Lagodowski, *Convergence rates in the SLLN for some classes of dependent random fields*, J. Math. Anal. Appl. **380** (2011), no. 2, 571–584. MR 2794415 [1](#), [4](#)
- [19] Zbigniew A. Lagodowski, *An approach to complete convergence theorems for dependent random fields via application of Fuk-Nagaev inequality*, J. Math. Anal. Appl. **437** (2016), no. 1, 380–395. MR 3451971 [1](#), [4](#)
- [20] Emmanuel Lesigne and Dalibor Volný, *Large deviations for martingales*, Stochastic Process. Appl. **96** (2001), no. 1, 143–159. MR 1856684 (2002k:60080) [4](#)
- [21] Quansheng Liu and Frédérique Watbled, *Exponential inequalities for martingales and asymptotic properties of the free energy of directed polymers in a random environment*, Stochastic Process. Appl. **119** (2009), no. 10, 3101–3132. MR 2568267 [4](#)
- [22] Florence Merlevède, Magda Peligrad, and Sergey Utev, *Recent advances in invariance principles for stationary sequences*, Probab. Surv. **3** (2006), 1–36. MR 2206313 (2007a:60025) [4](#)
- [23] Magda Peligrad and Na Zhang, *Martingale approximations for random fields*, Electron. Commun. Probab. **23** (2018), Paper No. 28, 9. MR 3798239 [1](#)
- [24] A. Račkauskas and C. Suquet, *Central limit theorems in Hölder topologies for Banach space valued random fields*, Teor. Veroyatn. Primen. **49** (2004), no. 1, 109–125. MR 2141332 (2006b:60011) [10](#)
- [25] Alfredas Račkauskas and Charles Suquet, *Necessary and sufficient condition for the Lamperti invariance principle*, Teor. Īmovīr. Mat. Stat. (2003), no. 68, 115–124. MR 2000642 (2004g:60050) [4](#)
- [26] Alfredas Račkauskas and Charles Suquet, *Necessary and sufficient condition for the functional central limit theorem in Hölder spaces*, J. Theoret. Probab. **17** (2004), no. 1, 221–243. MR 2054586 (2005b:60085) [4](#)
- [27] ———, *Hölderian invariance principles and some applications for testing epidemic changes*, Long memory in economics, Springer, Berlin, 2007, pp. 109–128. MR 2265057 [4](#)
- [28] Alfredas Račkauskas, Charles Suquet, and Vaidotas Zemlys, *A Hölderian functional central limit theorem for a multi-indexed summation process*, Stochastic Process. Appl. **117** (2007), no. 8, 1137–1164. MR 2340883 (2008e:60086) [10](#), [11](#)
- [29] Ludger Rüschendorf, *Ordering of distributions and rearrangement of functions*, Ann. Probab. **9** (1981), no. 2, 276–283. MR 606989 [6](#)
- [30] Frank Spitzer, *A combinatorial lemma and its application to probability theory*, Trans. Amer. Math. Soc. **82** (1956), 323–339. MR 79851 [3](#)
- [31] Dalibor Volný, *A central limit theorem for fields of martingale differences*, C. R. Math. Acad. Sci. Paris **353** (2015), no. 12, 1159–1163. MR 3427925 [5](#)
- [32] ———, *On limit theorems for fields of martingale differences*, Stochastic Process. Appl. **129** (2019), no. 3, 841–859. MR 3913270 [5](#), [15](#)
- [33] Michael J. Wichura, *Inequalities with applications to the weak convergence of random processes with multi-dimensional time parameters*, Ann. Math. Statist. **40** (1969), 681–687. MR 0246359 (39 #7663) [5](#)
- [34] Na Zhang, Lucas Reding, and Magda Peligrad, *On the quenched central limit theorem for stationary random fields under projective criteria*, J. Theoret. Probab. **33** (2020), no. 4, 2351–2379. MR 4166203 [1](#)