# ELLIPTIC RUIJSENAARS DIFFERENCE OPERATORS, SYMMETRIC POLYNOMIALS, AND WESS-ZUMINO-WITTEN FUSION RINGS

JAN FELIPE VAN DIEJEN AND TAMÁS GÖRBE

ABSTRACT. The fusion ring for  $\hat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten conformal field theories is known to be isomorphic to a factor ring of the ring of symmetric polynomials presented by Schur polynomials. We introduce a deformation of this factor ring associated with eigenpolynomials for the elliptic Ruijsenaars difference operators. The corresponding Littlewood-Richardson coefficients are governed by a Pieri rule stemming from the eigenvalue equation. The orthogonality of the eigenbasis gives rise to an analog of the Verlinde formula. In the trigonometric limit, our construction recovers the refined  $\hat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion ring associated with the Macdonald polynomials.

### 1. INTRODUCTION

It is well-known that the structure constants for the ring of symmetric polynomials in n variables in the basis of Schur polynomials  $s_{\lambda}(x_1, \ldots, x_n)$ , the Littlewood-Richardson coefficients, count the tensor multiplicities in the decomposition of tensor products of irreducible representations for the Lie algebra  $\mathfrak{su}(n;\mathbb{C})$  (cf. e.g. [M95, P07]). Here the partitions  $\lambda = (\lambda_1, \ldots, \lambda_n)$  label the dominant weights and one divides out by an ideal generated by the relation  $x_1 x_2 \cdots x_n = 1$ . If additionally—upon fixing a level  $m \in \mathbb{N}$ —the ideal generated by the Schur polynomials corresponding to partitions such that  $\lambda_1 - \lambda_n = m + 1$  is divided out, then one arrives at a finite-dimensional factor ring that is isomorphic to the fusion ring for  $\widehat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten conformal field theories [DMS97, F95, G91, GW90, K90, KS10]. The factor ring in question is also often referred to as the Verlinde algebra in the literature and it has a natural basis of Schur classes labeled by partitions  $\lambda$  with  $\lambda_1 - \lambda_n \leq m$  (which encode the dominant weights of the underlying affine Lie algebra  $\widehat{\mathfrak{su}}(n)_m$ ). The corresponding structure constants, i.e. the affine counterparts of the Littlewood-Richard coefficients at level m, describe the fusion rules for primary fields of the associated conformal field theories. From a mathematical point of view these structure constants compute, cf. e.g. [GN91], the Littlewood-Richardson coefficients of Hecke algebras at roots of unity [GW90] and the dimensions of spaces of conformal blocks of three-point functions in Wess-Zumino-Witten conformal field theories [TUY89].

Generalizations of the  $\widehat{\mathfrak{su}}(n)_m$  fusion ring have been constructed by means of parameter deformations of the Schur polynomials; important examples arise this

Date: June 2021.

<sup>2010</sup> Mathematics Subject Classification. Primary: 05E05; Secondary: 05E10, 33D52, 81T40, 81T45, 81Q80.

Key words and phrases. symmetric functions, elliptic Ruijsenaars system, Macdonald polynomials, Wess-Zumino-Witten fusion ring, Verlinde algebra.

way from the Hall-Littlewood polynomials [K13] and from the Macdonald polynomials [AS15, C05, K96, N14]. While the deformed fusion ring stemming from the Hall-Littlewood polynomials has been related to a deformation of the Verlinde algebra appearing in connection with an index formula of Teleman and Woodward [AGP16, OY14, T04, TW09], in the case of the Macdonald polynomials one deals rather with a Verlinde algebra that is intimately connected to the computation of refined Chern-Simons invariants for torus knots [AS15, C16, H16, GN15, N14].

From the point of view of quantum integrable particle dynamics, the Verlinde algebra for  $\hat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten conformal field theories can be interpreted as a Hilbert space for the phase model of impenetrable bosons on the one-dimensional periodic lattice [KS10]. From this perspective, the deformations of the Verlinde algebra associated with the Hall-Littlewood polynomials and with the Macdonald polynomials correspond in turn to Hilbert spaces for the periodic *q*-boson model [K13] and for the quantized trigonometric Ruijsenaars-Schneider system on  $\mathbb{CP}^{n-1}$ [DV98, GH18] (cf. also [BDM15]), respectively.

The idea of the present work is to introduce a further generalization of the Verlinde algebra originating from the *elliptic* Ruijsenaars operators; these are commuting difference operators with coefficients built from products of Weierstrass' sigma functions [R87, R99] that reduce to the Macdonald difference operators [M95] in the trigonometric limit. The study of their eigenfunctions points towards an elliptic counterpart of Macdonald's theory of symmetric polynomials [EK95, LNS20, MMZ21, RSV18]. Following [DG21], we will discretize the Ruijsenaars operators on a lattice of points labeled by partitions. Specifically, the lattice points are given by the  $\mathfrak{su}(n)$  dominant weights shifted by a Weyl vector that is rescaled linearly (as usual) with the multiplicity (or coupling) parameter. The eigenfunctions of these discrete Ruijsenaars operators are subsequently constructed in terms of polynomials determined by a recurrence stemming from the eigenvalue equation. The polynomials at issue turn out to provide a basis for the ring of symmetric polynomials. The associated Littlewood-Richardson coefficients constitute an elliptic deformation of Macdonald's (q, t)-Littlewood-Richardson coefficients [M95]. In this context, the eigenvalue equation for the elliptic Ruijsenaars operator entails an explicit formula for the elliptic deformation of the Pieri rule. Next, we construct the corresponding elliptic analog of the fusion ring and compute its structure constants in terms of elliptic Littlewood-Richardson coefficients. Now this deformed Verlinde algebra constitutes the Hilbert space for the compact elliptic Ruijsenaars model on  $\mathbb{CP}^{n-1}$ , whose classical and quantum dynamics was studied in [FG16] and [DG21], respectively. This particle interpretation gives rise to an orthogonality relation for the eigenpolynomials of the elliptic Ruijsenaars operators [DG21], from which we derive a Verlinde formula for the structure constants upon identifying the pertinent elliptic deformation of the  $\widehat{\mathfrak{su}}(n)_m$  Kac-Peterson modular S-matrix [K90].

The material is organized as follows. We start by discretizing the elliptic Ruijsenaars operators onto partitions in Section 2. Next, in Section 3, we construct an eigenbasis for these discrete Ruijsenaars operators; this gives rise to an elliptic deformation of Littlewood-Richardson coefficients generated by explicit Pieri rules. By dividing out the ideal generated by basis polynomials labeled by partitions with  $\lambda_1 - \lambda_n = m + 1$ , we arrive in Section 4 at an elliptic deformation of the  $\hat{\mathfrak{su}}(n)_m$ Wess-Zumino-Witten fusion ring; we compute its structure constants in terms of elliptic Littlewood-Richardson coefficients. In Section 5, a Verlinde formula for these structure constants is presented in terms of the relevant elliptic deformation of the Kac-Peterson modular S-matrix. We wrap up by detailing how the structure constants for the  $\widehat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion ring and its refined deformation associated with the Macdonald polynomials are recovered from their elliptic counterparts via parameter degenerations.

## 2. DISCRETE RUIJSENAARS OPERATORS ON PARTITIONS

2.1. Ruijsenaars' commuting difference operators. The Ruijsenaars difference operators  $D_1, \ldots, D_n$  are of the form

$$D_r = \sum_{\substack{J \subset \{1,\dots,n\} \\ |J|=r}} V_J(x) T_J, \quad V_J(x) = \prod_{\substack{j \in J \\ k \notin J}} \frac{[x_j - x_k + g]}{[x_j - x_k]} \qquad (r = 1,\dots,n).$$
(2.1)

Here  $|\cdot|$  denotes the cardinality of the set in question and  $T_J$  acts by translation on complex functions  $f(x) = f(x_1, \ldots, x_n)$ :

$$(T_J f)(x) = f(x + \varepsilon_J)$$
 with  $\varepsilon_J = \sum_{j \in J} \varepsilon_j$ 

(where  $\varepsilon_1, \ldots, \varepsilon_n$  refer to the standard unit basis of  $\mathbb{C}^n$ ). For  $g \in \mathbb{C}$ , these difference operators commute if the coefficients are built from a function [z] that factorizes into the product of a Weierstrass sigma function  $\sigma(z)$  and a Gaussian of the form  $\exp(az^2 + bz)$  (for any  $a, b \in \mathbb{C}$ ) [R87, R99]. For our purposes it is convenient to pick

$$[z] = [z; p] = \frac{\vartheta_1(\frac{\alpha}{2}z; p)}{\frac{\alpha}{2}\vartheta'_1(0; p)} \qquad (z \in \mathbb{C}, \, \alpha > 0, \, 0$$

where  $\vartheta_1$  denotes the Jacobi theta function

$$\vartheta_1(z;p) = 2\sum_{l\geq 0} (-1)^l p^{(l+\frac{1}{2})^2} \sin(2l+1)z, \qquad (2.2b)$$
$$= 2p^{1/4} \sin(z) \prod_{l\geq 1} (1-p^{2l})(1-2p^{2l}\cos(2z)+p^{4l}).$$

The conversion to the Weierstrass sigma function associated with the period lattice  $\Omega = 2\omega_1\mathbb{Z} + 2\omega_2\mathbb{Z}$  is governed by the relation (cf. e.g. [OLBC10, §23.6(i)]):

$$[z;p] = \sigma(z)e^{-\frac{\zeta(\omega_1)}{2\omega_1}z^2}$$

with  $\alpha = \frac{\pi}{\omega_1}$ ,  $p = e^{i\pi\tau}$ ,  $\tau = \frac{\omega_2}{\omega_1}$ , and  $\zeta(z) = \frac{\sigma'(z)}{\sigma(z)}$ . Below we will often use that [z; p] extends analytically in p to the interval  $-1 with <math>[z; 0] = \frac{2}{\alpha} \sin(\frac{\alpha z}{2})$ .

2.2. Discretization on partitions. Let  $\lambda = (\lambda_1, \ldots, \lambda_n)$  denote a partition of length  $\ell(\lambda) \leq n$ , i.e.  $\lambda$  belongs to

$$\Lambda^{(n)} = \{ \lambda \in \mathbb{Z}^n \mid \lambda_1 \ge \lambda_2 \ge \dots \ge \lambda_n \ge 0 \}.$$
(2.3)

Unless explicitly stated otherwise, it will be assumed that the value of g is chosen generically in  $\mathbb R$  such that

$$jg \notin \mathbb{Z}_{\leq 0} + \frac{2\pi}{\alpha} \mathbb{Z} \quad \text{for } j = 1, \dots, n$$
 (2.4a)

(where  $\mathbb{Z}_{\leq 0} = \mathbb{Z} \setminus \mathbb{N} = \{0, -1, -2, -3, \ldots\}$ ), which ensures in particular that

$$\prod_{1 \le j < k \le n} [\lambda_j - \lambda_k + (k - j)g] \ne 0 \qquad (\forall \lambda \in \Lambda^{(n)}).$$
(2.4b)

The following lemma now allows us to restrict  $D_r$  (2.1) to a discrete difference operator acting on functions supported on partitions shifted by

$$\rho_{\rm g} = ((n-1)g, (n-2)g, \dots, g, 0).$$
(2.5)

**Lemma 2.1** (Boundary Condition). For any  $\lambda \in \Lambda^{(n)}$  and  $J \subset \{1, \ldots, n\}$  (with g generic as detailed above), one has that

$$V_J(\rho_g + \lambda) = 0 \quad \text{if } \lambda + \varepsilon_J \notin \Lambda^{(n)}. \tag{2.6}$$

Proof. If  $\lambda \in \Lambda^{(n)}$  and  $\mu = \lambda + \varepsilon_J \notin \Lambda^{(n)}$  then  $\mu_j - \mu_{j+1} < 0$  for some  $1 \leq j < n$ , which implies that  $j \notin J$ ,  $j+1 \in J$  and  $\lambda_j - \lambda_{j+1} = 0$ . Since the denominators do not vanish because of Eq. (2.4b), one then picks up a zero of  $V_J(x)$  at  $x = \rho_g + \lambda$  from the factor  $[x_{j+1} - x_j + g]$ .

Specifically, by means of Lemma 2.1 we cast the corresponding action of  $D_r$  (2.1) in terms of a discrete difference operator in the space  $\mathcal{C}(\Lambda^{(n)})$  of complex lattice functions  $\lambda \xrightarrow{f} f_{\lambda}$ :

$$(D_r f)_{\lambda} = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + r}} B_{\nu/\lambda}(\alpha, \mathbf{g}; p) f_{\nu} \qquad (f \in \mathcal{C}(\Lambda^{(n)}), \ \lambda \in \Lambda^{(n)}), \tag{2.7a}$$

with

$$B_{\nu/\lambda}(\alpha, \mathbf{g}; p) = \prod_{1 \le j < k \le n} \frac{[\lambda_j - \lambda_k + \mathbf{g}(k - j + \theta_j - \theta_k)]}{[\lambda_j - \lambda_k + \mathbf{g}(k - j)]} \quad \text{and } \theta = \nu - \lambda, \quad (2.7b)$$

through the dictionary  $f(\rho_{\rm g} + \lambda) = f_{\lambda}$ ,  $\lambda + \varepsilon_J = \nu$  (which implies that  $V_J(\rho_{\rm g} + \lambda) = B_{\nu/\lambda}(\alpha, {\rm g}; p)$ ). Here and below we have employed the following standard notational conventions regarding partitions:

$$|\lambda| = \lambda_1 + \dots + \lambda_n, \qquad m^r = (\underbrace{m, \dots, m}_r, \underbrace{0, \dots, 0}_{n-r}),$$

and  $\forall \lambda, \mu \in \Lambda^{(n)}$ :  $\lambda \subset \mu$  iff  $\lambda_j \leq \mu_j$  for j = 1, ..., n. Since  $D_r$  (2.7a), (2.7b) amounts to a discretization of the elliptic Ruijsenaars operator  $D_r$  (2.1), the commutativity is inherited automatically.

**Corollary 2.2** (Commutativity). The discrete Ruijsenaars operators  $D_1, \ldots, D_n$  (2.7a), (2.7b) commute in  $\mathcal{C}(\Lambda^{(n)})$ .

## 3. Eigenpolynomials

3.1. Joint eigenfunctions. We now define polynomials  $P_{\mu}(\mathbf{e}) = P_{\mu}(\mathbf{e}; \alpha, \mathbf{g}; p), \mu \in \Lambda^{(n)}$  in the variables  $\mathbf{e} = (\mathbf{e}_1, \dots, \mathbf{e}_n)$  by means of the recurrence relation

$$P_{\mu}(\mathbf{e}) = e_r P_{\lambda}(\mathbf{e}) - \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n, \, |\nu| = |\mu| \\ \text{s.t. } \nu \in \Lambda^{(n)} \setminus \{\mu\}}} \psi'_{\nu/\lambda}(\alpha, \mathbf{g}; p) P_{\nu}(\mathbf{e}) \quad \text{if } \mu \neq 0,$$
(3.1a)

and  $P_{\mu}(\mathbf{e}) = 1$  if  $\mu = 0$ . Here  $\lambda = \mu - 1^r$ , where

$$r = r_{\mu} = \min\{1 \le j \le n \mid \mu_j - \mu_{j+1} > 0\}$$
(3.1b)

(with the convention  $\mu_{n+1} \equiv 0$ ) and

$$\psi_{\nu/\lambda}'(\alpha, \mathbf{g}; p) = \prod_{\substack{1 \le j < k \le n \\ \theta_j - \theta_k = -1}} \frac{[\nu_j - \nu_k + \mathbf{g}(k-j+1)]}{[\nu_j - \nu_k + \mathbf{g}(k-j)]} \frac{[\lambda_j - \lambda_k + \mathbf{g}(k-j-1)]}{[\lambda_j - \lambda_k + \mathbf{g}(k-j)]} \quad \text{with } \theta = \nu - \lambda.$$
(3.1c)

Let

$$d_{\mu} = \mu_1 - \mu_n \tag{3.2}$$

and let  $\leq$  denote the dominance partial order on  $\Lambda^{(n)}$ , i.e.

$$\forall \lambda, \mu \in \Lambda^{(n)} : \lambda \preceq \mu \Leftrightarrow |\lambda| = |\mu| \text{ and } \sum_{1 \leq j \leq r} \lambda_j \leq \sum_{1 \leq j \leq r} \mu_j \text{ for } r = 1, \dots, n-1$$

(and  $\lambda \prec \mu$  if  $\lambda \preceq \mu$  and  $\lambda \neq \mu$ ).

**Proposition 3.1** (Triangularity). The polynomials  $P_{\mu}(\mathbf{e})$ ,  $\mu \in \Lambda^{(n)}$  are uniquely determined by the recurrence (3.1a)–(3.1c) (from the initial condition  $P_0(\mathbf{e}) = 1$ ), and their expansion in the monomial basis is unitriangular with respect to the dominance order:

$$P_{\mu}(\mathbf{e}) = \mathbf{e}_{\mu} + \sum_{\nu \in \Lambda^{(n)}, \nu \prec \mu} u_{\mu,\nu} \, \mathbf{e}_{\nu} \quad with \ u_{\mu,\nu} = u_{\mu,\nu}(\alpha, \mathbf{g}; p) \in \mathbb{R}$$
(3.3a)

and

$$\mathbf{e}_{\mu} = \prod_{1 \le j \le n} \mathbf{e}_{j}^{\mu_{j} - \mu_{j+1}}.$$
(3.3b)

*Proof.* The proof is by lexicographical induction in  $(d_{\mu}, r_{\mu})$ , with  $r_{\mu}$  and  $d_{\mu}$  as in Eqs. (3.1b) and (3.2), respectively.

If  $d_{\mu} = 0$ , then either  $\mu = 0$  or r = n, i.e.  $\mu = m^n$  with  $m = \mu_1$ . When m = 0 we have that  $\mu = 0$ , so  $P_{\mu}(\mathbf{e}) = \mathbf{e}_{\mu} = 1$  by the initial condition, while for m > 0 the recurrence entails that  $P_{\mu}(\mathbf{e}) = P_{m^n}(\mathbf{e}) = \mathbf{e}_n P_{(m-1)^n}(\mathbf{e}) = \mathbf{e}_n^m = \mathbf{e}_{\mu}$ .

If  $d_{\mu} > 0$ , then we have that  $\mu = \lambda + 1^r$  with  $r = r_{\mu} < n$  and  $\lambda \in \Lambda^{(n)}$ . Since  $d_{\lambda} = d_{\mu} - 1$ , the induction hypothesis now ensures that on the RHS of the recurrence (3.1a) the term  $e_r P_{\lambda}(\mathbf{e})$  expands as  $e_r e_{\lambda} = e_{\mu}$  plus a  $\mathbb{R}$ -linear combination of monomials of the form  $e_r e_{\nu} = e_{\nu+1^r}$  with  $\nu \prec \lambda$ , i.e.  $\nu + 1^r \prec \mu$ ; the coefficients in this expansion stem from  $P_{\lambda}(\mathbf{e})$  and are thus uniquely determined from the recurrence relation (by the induction hypothesis). The remaining terms on the RHS of the recurrence (3.1a) consist in turn of a  $\mathbb{R}$ -linear combination of  $P_{\nu}(\mathbf{e})$  with  $d_{\nu} \leq d_{\mu}$  and  $\nu \prec \mu$ . Moreover, one *either* has that  $d_{\nu} < d_{\mu}$  or that  $d_{\nu} = d_{\mu}$  with  $r_{\nu} < r_{\mu}$ . Hence, in either case the induction hypothesis guarantees that  $P_{\nu}(\mathbf{e})$  is uniquely determined by the recurrence relations through a monomial expansion consisting of  $e_{\nu}$  perturbed by a linear combination of  $e_{\tilde{\nu}}$  with  $\tilde{\nu} \prec \nu$ .

Upon combining all these terms appearing on the RHS of the recurrence (3.1a), one confirms that monomial expansion of  $P_{\mu}(\mathbf{e})$  is of the form asserted in Eq. (3.3a) with expansion coefficients  $u_{\mu,\nu}(\alpha, \mathbf{g}; p) \in \mathbb{R}$  that are determined uniquely by the recurrence relation.

For  $\mathbf{e} \in \mathbb{C}^n$ , we define  $p(\mathbf{e}) = p(\mathbf{e}; \alpha, \mathbf{g}; p) \in \mathcal{C}(\Lambda^{(n)})$  in terms of the normalized polynomials

$$p_{\mu}(\mathbf{e}) = c_{\mu} P_{\mu}(\mathbf{e}) \quad (\mu \in \Lambda^{(n)}), \tag{3.4a}$$

with

$$c_{\mu} = c_{\mu}(\alpha, \mathbf{g}; p) = \prod_{1 \le j < k \le n} \frac{[(k-j)\mathbf{g}]_{\mu_j - \mu_k}}{[(k-j+1)\mathbf{g}]_{\mu_j - \mu_k}},$$
(3.4b)

where  $[z]_k, k = 0, 1, 2, ...$  denotes the *elliptic factorial* 

$$[z]_k = \prod_{0 \le l < k} [z+l]$$
 with  $[z]_0 = 1$ .

Notice that the regularity assumption on the parameter g in Eq. (2.4a) ensures that both the numerator and the denominator of  $c_{\mu}$  do not vanish.

**Theorem 3.2** (Joint Eigenfunctions). (i) The functions  $p(\mathbf{e})$ ,  $\mathbf{e} \in \mathbb{C}^n$  constitute a family of joint eigenfunctions for the elliptic Ruijsenaars operators  $D_1, \ldots, D_n$ (2.7a), (2.7b) in  $\mathcal{C}(\Lambda^{(n)})$ :

$$D_r p(\mathbf{e}) = \mathbf{e}_r p(\mathbf{e}) \quad for \ r = 1, \dots, n.$$
(3.5)

(ii) The vector of joint eigenvalues  $\mathbf{e} = (\mathbf{e}_1, \dots, \mathbf{e}_n) \in \mathbb{C}^n$  for the simultaneous eigenvalue problem in Eq. (3.5) is multiplicity free in  $\mathcal{C}(\Lambda^{(n)})$ .

*Proof.* When evaluating at  $\mu \in \Lambda^{(n)}$ , the sth eigenvalue equation in Eq. (3.5) reads

$$\sum_{\substack{\mu \subset \nu \subset \mu + 1^n \\ |\nu| = |\mu| + s}} B_{\nu/\mu}(\alpha, \mathbf{g}; p) \, p_{\nu}(\mathbf{e}) = \mathbf{e}_s p_{\mu}(\mathbf{e}). \tag{3.6a}$$

The explicit product formulas for  $B_{\mu/\lambda}(\alpha, \mathbf{g}; p)$  (2.7b),  $\psi'_{\mu/\lambda}(\alpha, \mathbf{g}; p)$  (3.1c) and  $c_{\mu}$  (3.4b) reveal that for all  $\nu \in \Lambda^{(n)}$  such that  $\mu \subset \nu \subset \mu + 1^{n+1}$ :

$$\psi'_{\nu/\mu}(\alpha, \mathbf{g}; p)c_{\mu} = B_{\nu/\mu}(\alpha, \mathbf{g}; p)c_{\nu}.$$
 (3.6b)

Hence, Eq. (3.6a) can be rewritten in terms of  $P_{\mu}(\mathbf{e})$  as follows:

$$\sum_{\substack{\mu \subset \nu \subset \mu + 1^n \\ |\nu| = |\mu| + s}} \psi'_{\nu/\mu}(\alpha, \mathbf{g}; p) P_{\nu}(\mathbf{e}) = \mathbf{e}_s P_{\mu}(\mathbf{e}).$$
(3.6c)

Since  $\psi'_{\mu+1^s/\mu}(\alpha, \mathbf{g}; p) = 1$ , the equality in Eq. (3.6c) is immediate from the recurrence for  $P_{\mu+1^s}(\mathbf{e})$  if  $s \leq r_{\mu}$  with the convention that  $r_0 = n$  (cf. Eqs. (3.1a)–(3.1c)). This settles the proof of Eq. (3.6a) for  $s \leq r_{\mu}$ , but it remains to check that the identity in question also holds if  $r_{\mu} < s \leq n$ . To this end we perform induction with respect to the lexicographical order on  $(d_{\mu}, r_{\mu})$  as in the proof of Proposition 3.1:

$$\mathbf{e}_{s}p_{\mu}(\mathbf{e}) = c_{\mu}\mathbf{e}_{s}P_{\mu}(\mathbf{e})$$

$$\stackrel{\text{Eq. (3.1a)}}{=} c_{\mu}\mathbf{e}_{s}\left(\mathbf{e}_{r}P_{\lambda}(\mathbf{e}) - \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^{n}, |\nu| = |\mu| \\ \text{s.t. } \nu \in \Lambda^{(n)} \setminus \{\mu\}}} \psi_{\nu/\lambda}'(\alpha, \mathbf{g}; p)P_{\nu}(\mathbf{e})\right)$$

(where  $r = r_{\mu}$  and  $\lambda = \mu - 1^r$ )

$$\stackrel{\text{induction}}{=} c_{\mu} \Big( e_{s} e_{r} P_{\lambda}(\mathbf{e}) - \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^{n}, |\nu| = |\mu| \\ \text{s.t. } \nu \in \Lambda^{(n)} \setminus \{\mu\}}} \psi_{\nu/\lambda}'(\alpha, \mathbf{g}; p) c_{\nu}^{-1} \big( D_{s} p(\mathbf{e}) \big)_{\nu} \Big)$$

$$\stackrel{\text{Eq. (3.6b)}}{=} \Big( D_{s} p(\mathbf{e}) \Big)_{\mu} + \frac{c_{\mu}}{c_{\lambda}} \Big( \underbrace{e_{s} e_{r} p_{\lambda}(\mathbf{e}) - \big( D_{r} D_{s} p(\mathbf{e}) \big)_{\lambda}}_{=0} \Big) = \Big( D_{s} p(\mathbf{e}) \Big)_{\mu}$$

as desired. To verify the cancellation of the underbraced terms it is essential to exploit the commutativity of the elliptic Ruijsenaars operators:

$$(D_r D_s p(\mathbf{e}))_{\lambda} \stackrel{\text{Cor. 2.2}}{=} (D_s D_r p(\mathbf{e}))_{\lambda} = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + s}} B_{\nu/\lambda}(\alpha, \mathbf{g}; p) (D_r p(\mathbf{e}))_{\nu}$$

$$\stackrel{*}{=} \mathbf{e}_r \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + s}} B_{\nu/\lambda}(\alpha, \mathbf{g}; p) p(\mathbf{e})_{\nu} = \mathbf{e}_r (D_s p(\mathbf{e}))_{\lambda} \stackrel{\text{induction}}{=} \mathbf{e}_r \mathbf{e}_s p_{\lambda}(\mathbf{e}).$$

In step \* we used that either  $(d_{\nu}, r_{\nu}) < (d_{\mu}, r_{\mu})$  in the lexicographical order (in which case the equality  $(D_r p(\mathbf{e}))_{\nu} = e_r p(\mathbf{e})_{\nu}$  stems from the induction hypothesis), or else  $r = r_{\mu} \leq r_{\nu}$  (in which case the equality in question is plain from the recurrence relation for  $P_{\nu+1r}(\mathbf{e})$  in combination with Eq. (3.6b)).

This completes the proof of part (i) of the Theorem. Part (ii) follows in turn from the observation that, upon normalizing such that  $p_0(\mathbf{e}) = 1$ , any joint eigenfunction  $p(\mathbf{e}) \in \mathcal{C}(\Lambda^{(n)})$  solving the eigenvalue equations in Eq. (3.5) automatically gives rise to polynomials  $P_{\mu}(\mathbf{e})$  (3.4a), (3.4b) obeying the recurrence relation (3.1a)–(3.1c). Proposition 3.1 thus implies that the joint eigenfunction is unique, i.e. the vector of joint eigenvalues  $\mathbf{e} = (e_1, \ldots, e_n)$  is multiplicity free in  $\mathcal{C}(\Lambda^{(n)})$ .

3.2. Elliptic Littlewood-Richardson coefficients. It is clear from Proposition 3.1 that the polynomials  $P_{\mu}(\mathbf{e}), \mu \in \Lambda^{(n)}$  form a basis for the polynomial ring  $\mathbb{R}[\mathbf{e}_1, \ldots, \mathbf{e}_n]$ . The corresponding structure constants give rise to an elliptic generalization of the Littlewood-Richardson coefficients:

$$P_{\lambda}P_{\mu} = \sum_{\nu \in \Lambda^{(n)}} c_{\lambda,\mu}^{\nu}(\alpha, \mathbf{g}; p) P_{\nu}$$
(3.7)

(where the arguments **e** are suppressed). For  $\mu = 1^r$ , an explicit product formula for these elliptic Littlewood-Richardson coefficients is immediate from Theorem 3.2.

**Corollary 3.3** (Pieri Rule). For  $\mu \in \Lambda^{(n)}$  and  $1 \leq r \leq n$ , one has that

$$P_{\lambda}P_{1^{r}} = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^{n} \\ |\nu| = |\lambda| + r}} \psi_{\nu/\lambda}'(\alpha, \mathbf{g}; p) P_{\nu}.$$
(3.8)

*Proof.* Since  $P_{1^r}(\mathbf{e}) = \mathbf{e}_{1^r} = \mathbf{e}_r$ , the asserted Pieri rules encode the eigenvalue equations of Theorem 3.2 in the reformulation of Eq. (3.6c) (obtained via Eq. (3.6b)).

With the aid of the Pieri rule, it is readily seen that the classical Littlewood-Richardson coefficients  $c_{\lambda,\mu}^{\nu}$  for the Schur polynomials [M95, Chapter I.9] and their two-parameter deformation  $f_{\lambda,\mu}^{\nu}(q,t)$  associated with the Macdonald polynomials [M95, Chapter VI.7] arise as suitable parameter specializations of the threeparameter elliptic Littlewood-Richardson coefficients  $c_{\lambda,\mu}^{\nu}(\alpha, \mathbf{g}; p)$  (3.7).

**Proposition 3.4** (Degenerations). The classical Littlewood-Richardson coefficients  $c^{\nu}_{\lambda,\mu}$  and Macdonald's (q,t)-Littlewood-Richardson coefficients  $f^{\nu}_{\lambda,\mu}(q,t)$  are recovered from  $c^{\nu}_{\lambda,\mu}(\alpha, \mathbf{g}; p)$  (3.7) in the following way:

$$\lim_{\mathbf{g}\to\mathbf{1}} c_{\lambda,\mu}^{\nu}(\alpha,\mathbf{g};p) = c_{\lambda,\mu}^{\nu} \tag{3.9a}$$

(provided  $\frac{2\pi}{\alpha} > 0$  is irrational), and

$$\lim_{p \to 0} c^{\nu}_{\lambda,\mu}(\alpha, \mathbf{g}; p) = f^{\nu}_{\lambda,\mu}(q, q^{\mathbf{g}}) \quad with \quad q = e^{i\alpha}$$
(3.9b)

(provided  $jg \notin \mathbb{Z}_{\leq 0} + \frac{2\pi}{\alpha}\mathbb{Z}$  for j = 1, ..., n). In particular, for  $\lambda, \nu \in \Lambda^{(n)}$  such that  $\lambda \subset \nu \subset \lambda + 1^n$  one has with these genericity assumptions in place that

$$\lim_{\mathbf{g}\to 1}\psi'_{\nu/\lambda}(\alpha,\mathbf{g};p) = 1 \tag{3.10a}$$

and

$$\lim_{p \to 0} \psi'_{\nu/\lambda}(\alpha, \mathbf{g}; p) = \prod_{\substack{1 \le j < k \le n \\ \theta_j - \theta_k = -1}} \frac{[\nu_j - \nu_k + \mathbf{g}(k-j+1)]_q}{[\nu_j - \nu_k + \mathbf{g}(k-j)]_q} \frac{[\lambda_j - \lambda_k + \mathbf{g}(k-j-1)]_q}{[\lambda_j - \lambda_k + \mathbf{g}(k-j)]_q},$$
(3.10b)

where  $\theta = \nu - \lambda$  and

$$[z]_q = \frac{\sin(\frac{\alpha z}{2})}{\sin(\frac{\alpha}{2})} = \frac{q^{\frac{z}{2}} - q^{-\frac{z}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}.$$

*Proof.* By definition, the polynomial ring  $\mathbb{R}[e_1, \ldots, e_n]$  is generated by the monomials  $e_r = P_{1r}(\mathbf{e}), r = 1, \dots, n$ . It is therefore sufficient to verify the limits in Eqs. (3.9a), (3.9b) for  $\lambda = 1^r$  (r = 1, ..., n), which—by the Pieri rule of Corollary 3.3—amounts to checking the limits in Eqs. (3.10a), (3.10b). The genericity assumptions on the parameters ensure that none of the denominators vanish.

Specifically, in the product formula for  $\psi'_{\nu/\lambda}$  (3.1c) one has that  $\lambda_j - \lambda_k =$  $\nu_j - \nu_k + 1$  if  $\theta_j - \theta_k = -1$ , so the limit in Eq. (3.10a) is evident. We thus recover in this manner the (dual) Pieri rules for the Schur polynomials  $s_{\mu}(x)$  [M95, Ch. I, Eq. (5.17) from Corollary 3.3, which in turn implies the limit in Eq. (3.9a).

Similarly, since [z; p] (2.2a), (2.2b) extends analytically to -1 with $[z;0] = \frac{\alpha}{2}\sin(\frac{\alpha z}{2})$ , the limit in Eq. (3.10b) is also manifest from Eq. (3.1c). Upon comparing with [M95, Ch. VI, Eqs. (6.7'), (6.13)], we see that at p = 0 the Pieri rules for the Macdonald polynomials  $P_{\mu}(x;q,q^{\rm g})$  are recovered from Corollary 3.3, therewith settling the limit in Eq. (3.9b). 

The Pieri rule also confirms that some well-known vanishing properties enjoyed by the Littlewood-Richardson coefficients persist at the elliptic level.

**Proposition 3.5** (Vanishing Terms). For  $\lambda, \mu, \nu \in \Lambda^{(n)}$ , the elliptic Littlewood-Richardson coefficient  $c^{\nu}_{\lambda,\mu}(\alpha, \mathbf{g}; p)$  vanishes unless  $\lambda \subset \nu$  and  $\mu \subset \nu$  with  $|\lambda| + |\mu| =$  $|\nu|$ .

*Proof.* It suffices to mimic the proof for the corresponding statement at p = 0 from [M95, Ch. VI, Eq. (7.4)]. Since the monomial expansion of  $P_{\mu}(\mathbf{e})$  in Proposition 3.1 involves only monomials  $e_{\kappa}$  with  $|\kappa| = |\mu|$  and  $e_{\kappa}e_{\tilde{\kappa}} = e_{\kappa+\tilde{\kappa}}$ , it is clear that  $c_{\lambda,\mu}^{\nu}(\alpha, \mathbf{g}; p)$  (3.7) can only be nonzero provided  $|\nu| = |\lambda| + |\mu|$ . Moreover, let  $\mathcal{I}_{\lambda}$ denote the subspace of  $\mathbb{R}[e_1, \ldots, e_n]$  spanned by the  $P_{\kappa}(\mathbf{e})$  with  $\lambda \subset \kappa$ . It is manifest from the Pieri rule that  $e_r \mathcal{I}_{\lambda} \subset \mathcal{I}_{\lambda}$  for  $r = 1, \ldots, n$ , so  $\mathcal{I}_{\lambda}$  is an ideal in  $\mathbb{R}[e_1, \ldots, e_n]$ . It thus follows that  $P_{\lambda}(\mathbf{e})P_{\mu}(\mathbf{e}) \in \mathcal{I}_{\lambda} \cap \mathcal{I}_{\mu}$ . 

3.3. Symmetric polynomials. Let us recall (cf. the proof of Proposition 3.4) that  $s_{\mu}(x) = s_{\mu}(x_1, \ldots, x_n)$  and  $P_{\mu}(x;q,t) = P_{\mu}(x_1, \ldots, x_n;q,t)$  refer to the Schur polynomials [M95, Chapter I] and the Macdonald polynomials [M95, Chapter VI],

respectively. The symmetric polynomials in question are monic in the sense that their leading monomial is given by

$$m_{\mu}(x) = m_{\mu}(x_1, \dots, x_n) = \sum_{\nu \in S_n(\mu)} x_1^{\nu_1} \cdots x_n^{\nu_n} \qquad (\mu \in \Lambda^{(n)}),$$
(3.11)

where the sum is over all compositions reordering the parts of  $\mu$  (i.e. over the orbit of  $\mu$  with respect to the action of the permutation-group  $S_n$  of permutations  $\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ \sigma_1 & \sigma_2 & \cdots & \sigma_n \end{pmatrix}$  on  $\mu_1, \mu_2, \ldots, \mu_n$ ). It is instructive to describe the precise relation between our elliptic eigenpolynomials and these two standard bases for the ring  $\mathcal{A}^{(n)} = \mathbb{R}[x_1, \ldots, x_n]^{S_n}$  of symmetric polynomials in the variables  $x_1, \ldots, x_n$ .

To this end, let us observe that the polynomials  $P_{\mu}(\mathbf{e}), \mu \in \Lambda^{(n)}$  give rise to a (monic) basis for  $\mathcal{A}^{(n)}$  through the ring isomorphism  $\mathbb{R}[\mathbf{e}_1, \ldots, \mathbf{e}_n] \cong \mathcal{A}^{(n)}$  determined by the injection  $\mathbf{e}_r \to m_{1^r}(x), r = 1, \ldots, n$ :

$$R_{\mu}(x;\alpha,\mathbf{g};p) = P_{\mu}(\mathbf{e}) \quad \text{with } \mathbf{e} = (m_{1^{1}}(x), m_{1^{2}}(x), \dots, m_{1^{n}}(x)).$$
(3.12)

From the limits in Proposition 3.4, it is then clear that the corresponding degenerations of the recurrence relations (3.1a)-(3.1c) reproduce the Schur polynomials and the Macdonald polynomials respectively (cf. [M95, Ch. I, Eq. (5.17)] and [M95, Ch. VI, Eqs. (6.7'), (6.13)]).

**Corollary 3.6** (The Schur and Macdonald Limits). For any  $\mu \in \Lambda^{(n)}$ , the Schur polynomial  $s_{\mu}(x)$  and the Macdonald polynomial  $P_{\mu}(x;q,q^{\rm g})$  are recovered from  $R_{\mu}(x;\alpha,{\rm g};p)$  (3.12) in the following way:

$$\lim_{\mathbf{a}\to\mathbf{1}} R_{\mu}(x;\alpha,\mathbf{g};p) = s_{\mu}(x) \tag{3.13a}$$

(provided  $\frac{2\pi}{\alpha} > 0$  is irrational), and

$$\lim_{p \to 0} R_{\mu}(x; \alpha, \mathbf{g}; p) = P_{\mu}(x; q, q^{\mathbf{g}}) \quad with \quad q = e^{i\alpha}$$
(3.13b)

(provided  $jg \notin \mathbb{Z}_{\leq 0} + \frac{2\pi}{\alpha}\mathbb{Z}$  for  $j = 1, \ldots, n$ ).

4. GENERIC ELLIPTIC DEFORMATION OF THE FUSION RING FOR 
$$\widehat{\mathfrak{su}}(n)_m$$

4.1. Character ring for  $\mathfrak{su}(n)$ . When r = n the Pieri rule (3.8) simply states that  $P_{\mu+1^n}(\mathbf{e}) = P_{\mu}(\mathbf{e})P_{1^n}(\mathbf{e}) = P_{\mu}(\mathbf{e})\mathbf{e}_n$ . Hence, one has more generally that

$$P_{\mu}(\mathbf{e}) = P_{\underline{\mu}}(\mathbf{e}) \mathbf{e}_{n}^{\mu_{n}} \quad \text{with } \underline{\mu} = (\mu_{1} - \mu_{n}, \mu_{2} - \mu_{n}, \dots, \mu_{n-1} - \mu_{n}, 0).$$
(4.1)

To divide out this translational symmetry one substitutes  $e_n = 1$ , therewith reducing to the ring

$$\mathcal{R}_0^{(n)} = \mathbb{R}[\mathbf{e}_1, \dots, \mathbf{e}_n] / \langle \mathbf{e}_n - 1 \rangle$$
(4.2a)

for which (the corresponding specialization of) the polynomials

$$P_{\mu}(\mathbf{e}) \quad \text{with } \mu \in \Lambda_0^{(n)} = \{\lambda \in \Lambda^{(n)} \mid \lambda_n = 0\}$$
(4.2b)

provide a basis. In view of Proposition 3.5, it is immediate from Eq. (4.1) that the corresponding structure constants for  $\mathcal{R}_0^{(n)}$  can be expressed in terms of elliptic Littlewood-Richardson coefficients  $c_{\lambda,\mu}^{\nu}(\alpha, \mathbf{g}; p)$  (3.7) as follows:

$$P_{\lambda}P_{\mu} = \sum_{\substack{\nu \supset \lambda, \ \nu \supset \mu \\ |\nu| = |\lambda| + |\mu|}} c_{\lambda,\mu}^{\nu}(\alpha, \mathbf{g}; p) P_{\underline{\nu}} \qquad (\lambda, \mu \in \Lambda_0^{(n)}, \nu \in \Lambda^{(n)}).$$
(4.3a)

In particular, for  $1 \leq r < n$  and  $\lambda \in \Lambda_0^{(n)}$  one retrieves from Corollary 3.3 that

$$P_{\lambda}P_{1^{r}} = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^{n} \\ |\nu| = |\lambda| + r}} \psi_{\nu/\lambda}'(\alpha, \mathbf{g}; p) P_{\underline{\nu}}.$$
(4.3b)

Proposition 3.4 and Corollary 3.6 entail that at p = 0 Eq. (4.3a) recovers the structure constants for the multiplication in the basis of Macdonald polynomials associated with (the root system of) the complex simple Lie algebra  $\mathfrak{su}(n)$ [M00, M03]. Indeed, Eq. (4.3b) degenerates to the Pieri rule for the  $\mathfrak{su}(n)$  Macdonald polynomials in this situation (which is obtained from the Pieri formula for  $P_{\mu}(x_1, \ldots, x_n; q, q^{\mathrm{g}})$  in [M95, Ch. VI, Eqs. (6.7'), (6.13)] by dividing out the ideal generated by  $m_{1^n}(x_1, \ldots, x_n) - 1$ ). Here partitions  $\lambda \in \Lambda_0^{(n)}$  are identified with dominant weight vectors for  $\mathfrak{su}(n)$  in the standard way:

$$\lambda \leftrightarrow \sum_{1 \le r < n} (\lambda_r - \lambda_{r+1}) \overline{\omega}_r, \tag{4.4}$$

where  $\varpi_r \leftrightarrow 1^r$ , r = 1, ..., n-1 refers to the corresponding basis of fundamental weight vectors (labeled in accordance with the plates of [B68]).

Similarly, for  $g \to 1$  Eq. (4.3a) encodes the structure constants for the character ring of  $\mathfrak{su}(n)$  in the basis of the irreducible characters. For instance, in this limit the Pieri rule (4.3b) counts the tensor multiplicities for tensoring with a fundamental representation, cf. e.g. [P07, Chapter 9.10].

4.2. **Fusion ideal.** We now scale  $\alpha$  in terms of  $g \in \mathbb{R} \setminus \mathbb{Q}$  as follows:

$$\alpha = \frac{2\pi}{m + ng} \quad \text{with } m \in \mathbb{N}.$$
(4.5)

The irrationality of g then guarantees that the regularity requirement in Eq. (2.4b) is satisfied, so the polynomials  $P_{\mu}(\mathbf{e})$  (3.1a)–(3.1c) are well-defined for this parameter specialization. Let us consider the following ideal in  $\mathcal{R}_0^{(n)}$  (4.2a):

$$\mathcal{I}^{(n,m)} = \langle P_{\mu}(\mathbf{e}) \mid \mu \in \Lambda_0^{(n)} \text{ with } d_{\mu} = m + 1 \rangle.$$
(4.6)

This ideal should be viewed as an elliptic (g, p)-deformation associated with the elliptic Ruijsenaars model of the fusion ideal for  $\widehat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten conformal field theories (cf. e.g. [DMS97, F95, G91, GW90, KS10] and references therein).

With the aid of the Pieri rule (4.3b) and following lemma, one arrives at a convenient a basis for  $\mathcal{I}^{(n,m)}$  in terms of the eigenpolynomials associated with the elliptic Ruijsenaars lattice model.

**Lemma 4.1** (Level *m* Boundary Condition). Let  $\lambda \in \Lambda_0^{(n)}$  with  $d_{\lambda} = m + 1$  and let  $\lambda \subset \nu \subset \lambda + 1^n$ . Then one has that

$$\psi_{\nu/\lambda}'(\frac{2\pi}{m+ng}, g; p) = 0 \quad if \ d_{\nu} \le m \tag{4.7}$$

(assuming  $g \in \mathbb{R} \setminus \mathbb{Q}$ ).

*Proof.* The conditions imply that  $\nu = \lambda + \theta$  with  $\theta$  a vertical *r*-strip  $(1 \le r < n)$ , so  $d_{\nu} = d_{\lambda} + \theta_1 - \theta_n$  with  $\theta_1, \theta_n \in \{0, 1\}$ . If  $d_{\nu} \le m$ , we must in fact have that  $\theta_1 = 0$ ,  $\theta_n = 1$  and  $d_{\nu} = m$  as  $d_{\lambda} = m + 1$ . In this situation,  $\psi'_{\nu/\lambda}(\frac{2\pi}{m+ng}, g; p)$  (3.1c) picks up a zero from the factor  $[\nu_j - \nu_k + (k - j + 1)g]$  in the numerator for j = 1 and k = n:  $[\nu_1 - \nu_n + ng] = [m + ng] = [\frac{2\pi}{\alpha}] = 0$ .

**Proposition 4.2** (Basis for  $\mathcal{I}^{(n,m)}$ ). For  $\alpha = \frac{2\pi}{m+ng}$  and  $g \in \mathbb{R} \setminus \mathbb{Q}$ , the polynomials  $P_{\mu}(\mathbf{e})$  with  $\mu \in \Lambda_0^{(n)}$  such that  $d_{\mu} > m$  constitute a basis for  $\mathcal{I}^{(n,m)}$ .

*Proof.* By definition, the ideal  $\mathcal{I}^{(n,m)}$  (4.6) consists of polynomials of the form

$$\sum_{\lambda \in \Lambda_0^{(n)}, \, d_{\lambda} = m+1} a_{\lambda}(\mathbf{e}) P_{\lambda}(\mathbf{e}) \quad \text{with } a_{\lambda}(\mathbf{e}) \in \mathcal{R}_0^{(n)}.$$
(4.8)

Since the monomials  $\mathbf{e}_r = P_{1r}(\mathbf{e})$   $(1 \leq r < n)$  generate  $\mathcal{R}_0^{(n)}$ , it is immediate from the Pieri rule (4.3b) and Lemma 4.1 that all products  $a_\lambda(\mathbf{e})P_\lambda(\mathbf{e})$  in the sum of Eq. (4.8) expand as  $\mathbb{R}$ -linear combinations of basis polynomials  $P_\mu(\mathbf{e})$  with  $\mu \in \Lambda_0^{(n)}$ and  $d_\mu > m$ . It remains to check that the pertinent basis polynomials  $P_\mu(\mathbf{e})$  indeed belong to  $\mathcal{I}^{(n,m)}$ . If  $d_\mu = m+1$  this is the case by definition, while for  $d_\mu > m+1$  it follows by lexicographical induction in  $(d_\mu, r_\mu)$  from the recurrence in Eqs. (3.1a)– (3.1c) with the aid of Lemma 4.1 (using also that  $P_\nu(\mathbf{e}) = P_{\underline{\nu}}(\mathbf{e})$  in  $\mathcal{R}_0^{(n)}$ ). Indeed, on the RHS of (3.1a) one has that  $d_\lambda = d_\mu - 1 > m$ , so  $\mathbf{e}_r P_\lambda(\mathbf{e}) \in \mathcal{I}^{(n,m)}$  since  $P_\lambda(\mathbf{e}) \in \mathcal{I}^{(n,m)}$  by virtue of the induction hypothesis. The remaining terms on the RHS of Eq. (3.1a) involve polynomials  $P_\nu(\mathbf{e}) = P_{\underline{\nu}}(\mathbf{e})$  with *either*  $d_\nu = d_\mu > m$  and  $r_\nu < r_\mu$  or (using Lemma 4.1) with  $d_\mu > d_\nu > m$ ; the induction hypothesis therefore guarantees again that all of these terms belong to  $\mathcal{I}^{(n,m)}$ . We may thus conclude that  $P_\mu(\mathbf{e})$  (3.1a) lies in  $\mathcal{I}^{(n,m)}$ , therewith completing the induction step.  $\Box$ 

4.3. Fusion ring. Upon dividing out  $\mathcal{I}^{(n,m)}$ , one arrives in turn at a corresponding elliptic deformation of the fusion ring for  $\mathfrak{su}(n)_m$  Wess-Zumino-Witten conformal field theories:

$$\mathcal{R}_{0}^{(n,m)} = \mathcal{R}_{0}^{(n)} / \mathcal{I}^{(n,m)}$$
(4.9)

(cf. again [DMS97, F95, G91, GW90, KS10] and references therein). For  $P \in \mathcal{R}_0^{(n)}$ , we will denote its coset  $P + \mathcal{I}^{(n,m)}$  in  $\mathcal{R}_0^{(n,m)}$  by [P]. The cosets of the elliptic eigenpolynomials labeled by bounded partitions in

$$\Lambda_0^{(n,m)} = \{\lambda \in \Lambda_0^{(n)} \mid d_\lambda \le m\}$$
(4.10)

provide a basis for our elliptic fusion ring. Notice in this connection that the bijection (4.4) maps the bounded partitions in question to (the nonaffine parts of) the dominant weights of the affine Lie algebra  $\hat{\mathfrak{su}}(n)_m$  (cf. e.g. [KS10, Section 2.1]).

**Proposition 4.3** (Basis for  $\mathcal{R}_0^{(n,m)}$ ). For  $\alpha = \frac{2\pi}{m+ng}$  and  $g \in \mathbb{R} \setminus \mathbb{Q}$ , the cosets  $[P_{\mu}(\mathbf{e})], \ \mu \in \Lambda_0^{(n,m)}$  constitute a basis for  $\mathcal{R}_0^{(n,m)}$ .

Proof. Since the polynomials  $P_{\mu}(\mathbf{e}), \ \mu \in \Lambda_0^{(n)}$  form a basis for  $\mathcal{R}_0^{(n)}$ , the assertion in the proposition is immediate from Proposition 4.2. Indeed, the kernel of the ring homomorphism  $P \to [P]$  from  $\mathcal{R}_0^{(n)}$  onto  $\mathcal{R}_0^{(n,m)}$  is equal to the  $\mathbb{R}$ -span of the polynomials  $P_{\mu}(\mathbf{e}), \ \mu \in \Lambda_0^{(n)} \setminus \Lambda_0^{(n,m)}$  (by Proposition 4.2), so the cosets  $[P_{\mu}(\mathbf{e})], \ \mu \in \Lambda_0^{(n,m)}$  constitute a basis for  $\mathcal{R}_0^{(n,m)}$ .

It is now straightforward to express the structure constants of  $\mathcal{R}_0^{(n,m)}$  in the basis  $[P_{\mu}(\mathbf{e})], \ \mu \in \Lambda_0^{(n,m)}$  in terms of elliptic Littlewood-Richardson coefficients.

**Theorem 4.4** (Structure Constants of  $\mathcal{R}_0^{(n,m)}$ ). For  $g \in \mathbb{R} \setminus \mathbb{Q}$ , the structure constants of  $\mathcal{R}_0^{(n,m)}$  in the basis  $[P_{\mu}(\mathbf{e})], \ \mu \in \Lambda_0^{(n,m)}$  can be expressed in terms of elliptic Littlewood-Richardson coefficients as follows:

$$[P_{\lambda}][P_{\mu}] = \sum_{\substack{\nu \supset \lambda, \ \nu \supset \mu \\ |\nu| = |\lambda| + |\mu|, \ d_{\nu} \le m}} c_{\lambda,\mu}^{\nu} \left(\frac{2\pi}{m + ng}, g; p\right)[P_{\underline{\nu}}] \qquad (\lambda, \mu \in \Lambda_{0}^{(n,m)}, \ \nu \in \Lambda^{(n)}).$$

$$(4.11a)$$

In particular, for  $1 \leq r < n$  and  $\lambda \in \Lambda_0^{(n,m)}$  one has explicitly that

$$[P_{\lambda}][P_{1^r}] = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + r, \ d_{\nu} \le m}} \psi'_{\nu/\lambda} \left(\frac{2\pi}{m + ng}, g; p\right) [P_{\underline{\nu}}].$$
(4.11b)

*Proof.* In view of Propositions 4.2 and 4.3, the asserted formulas in Eqs. (4.11a) and (4.11b) follow from Eqs. (4.3a) and (4.3b) upon applying the ring homomorphism  $P \to [P]$  from  $\mathcal{R}_0^{(n)}$  onto  $\mathcal{R}_0^{(n,m)}$ .

# 5. $\widehat{\mathfrak{su}}(n)_m$ Verlinde algebras from elliptic Ruijsenaars systems

5.1. Analyticity. For g > 0 and  $\mu \in \Lambda_0^{(n)}$  with  $d_{\mu} \leq m+1$ , we now employ analytic continuation in the parameters of  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) \in \mathcal{R}_0^{(n)}$  so as to remove the restriction that g be irrational.

**Lemma 5.1** (Positivity of the Recurrence Coefficients). Let  $\alpha = \frac{2\pi}{m+ng}$ , g > 0,  $-1 , and <math>\nu \in \Lambda^{(n)}$ .

(i) For  $1 \le j < k \le n$ , one has that  $[\nu_j - \nu_k + (k - j)g; p] > 0$  if  $\nu_j - \nu_k \le m$ .

(ii) For  $\lambda \in \Lambda_0^{(n,m)}$  and  $\lambda \subset \nu \subset \lambda + 1^n$ , the recurrence coefficient  $\psi'_{\nu/\lambda}(\frac{2\pi}{m+ng}, g; p)$ (3.1c) is analytic in  $g \in (0, \infty)$  and positive.

*Proof.* (i) The asserted positivity of the theta factor is plain from the product expansion (2.2b) via the estimate

$$0 < g \le \nu_j - \nu_k + (k - j)g < m + ng = \frac{2\pi}{\alpha}.$$

(*ii*) The analyticity of  $\psi'_{\nu/\lambda}(\frac{2\pi}{m+ng}, g; p)$  in g > 0 follows in turn from the fact that the zeros in the denominators are avoided as a consequence of part (*i*). Indeed, in all factors of  $\psi'_{\nu/\lambda}(\frac{2\pi}{m+ng}, g; p)$  (3.1c) the denominators stay positive because  $\lambda_j - \lambda_k \leq d_\lambda \leq m$  and  $\nu_j - \nu_k = \lambda_j - \lambda_k + \theta_j - \theta_k \leq m - 1$ . The positivity of the corresponding numerators is subsequently seen from the estimates

$$0 \le \nu_j - \nu_k + (k - j - 1)g < \lambda_j - \lambda_k + (k - j - 1)g < m + ng$$
  
and  $0 < \nu_j - \nu_k + (k - j + 1)g < m + ng.$ 

**Proposition 5.2** (Analyticity of Basis Polynomials). For  $\mu \in \Lambda_0^{(n)}$  the polynomial  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) \in \mathcal{R}_0^{(n)}$  is analytic, both in  $g \in (0, \infty)$  and in  $p \in (-1, 1)$ , provided  $d_{\mu} \leq m+1$ .

*Proof.* By Lemma 5.1, the conditions make sure that all coefficients  $\psi'_{\nu/\lambda}(\frac{2\pi}{m+ng}, g; p)$  in the recurrence (3.1a)–(3.1c) for  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p)$  enjoy the asserted analyticity, which is therefore inherited by the polynomials in question.

For later reference, let us also explicitly check that the pertinent normalization coefficients  $c_{\mu}(\alpha, \mathbf{g}; p)$  from Eq. (3.4b) permit analytic continuation to  $\mathbf{g} > 0$ .

**Lemma 5.3** (Positivity of the Normalization Constants). For  $\mu \in \Lambda_0^{(n,m)}$ , g > 0and  $-1 , the normalization coefficient <math>c_{\mu}(\frac{2\pi}{m+ng}, g; p)$  (3.4b) is positive.

*Proof.* The arguments of the theta functions in the numerator and the denominator of  $c_{\mu}(\frac{2\pi}{m+ng}, g; p)$  are of the form l + kg, with  $0 \le l < m$  and  $1 \le k \le n$ , so  $0 < l + kg < m + ng = \frac{2\pi}{\alpha}$ . The corresponding values of the theta function are thus positive by the product formula (2.2b).

5.2. Spectral variety. In [DG21] it was shown that for  $\alpha$  (4.5) with g > 0, the elliptic Ruijsenaars operators truncate to commuting discrete difference operators of the form (cf. Eqs. (2.7a), (2.7b)):

$$(D_r f)_{\lambda} = \sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + r, \, d_{\nu} \le m}} B_{\nu/\lambda} \left(\frac{2\pi}{m + ng}, g; p\right) f_{\underline{\nu}}, \quad 1 \le r < n.$$
(5.1)

These operators turn out to be normal in the Hilbert space  $\ell^2(\Lambda_0^{(n,m)}, \Delta)$  with the inner product

$$\langle f,g\rangle_{\Delta} = \sum_{\lambda \in \Lambda_0^{(n,m)}} f_{\lambda} \overline{g_{\lambda}} \Delta_{\lambda} \qquad \left(f,g \in \ell^2(\Lambda_0^{(n,m)},\Delta)\right),\tag{5.2a}$$

where

$$\Delta_{\lambda} = \Delta_{\lambda}(\alpha, \mathbf{g}; p) = \prod_{1 \le j < k \le n} \frac{[\lambda_j - \lambda_k + (k-j)\mathbf{g}]}{[(k-j)\mathbf{g}]} \frac{[(k-j+1)\mathbf{g}]_{\lambda_j - \lambda_k}}{[1 + (k-j-1)\mathbf{g}]_{\lambda_j - \lambda_k}} \tag{5.2b}$$

(cf. [DG21, Proposition 6]). Their joint spectrum is moreover given by  $\binom{n-1+m}{m}$  multiplicity-free vectors (cf. [DG21, Corollary 10])

$$\mathbf{e}_{\nu} = \mathbf{e}_{\nu} \left( \frac{2\pi}{m+ng}, g; p \right) = \left( e_{1,\nu}, \dots, e_{n-1,\nu}, 1 \right) \in \mathbb{C}^n \quad (\nu \in \Lambda_0^{(n,m)})$$
(5.3a)

such that

$$D_r p(\mathbf{e}_{\nu}) = \mathbf{e}_{r,\nu} p(\mathbf{e}_{\nu}) \quad (1 \le r < n, \, \nu \in \Lambda_0^{(n,m)}), \tag{5.3b}$$

which are analytic in  $p \in (-1, 1)$  with

$$\lim_{p \to 0} e_{r,\nu} \left( \frac{2\pi}{m+ng}, g; p \right) =$$

$$q^{-r \left( \frac{|\nu|}{n} + \frac{(n-1)g}{2} \right)} m_{1r} \left( q^{\nu_1 + (n-1)g}, q^{\nu_2 + (n-2)g}, \dots, q^{\nu_{n-1} + g}, 1 \right).$$
(5.3c)

Here  $p(\mathbf{e})$  is given by Eqs. (3.4a), (3.4b) and  $q = e^{\frac{2\pi i}{m+ng}}$ .

Let us now define the *spectral variety* as the zero locus  $V(\mathcal{I}^{(n,m)})$  of the fusion ideal  $\mathcal{I}^{(n,m)}$  (4.6):

$$V(\mathcal{I}^{(n,m)}) = \{ \mathbf{e} = (\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, 1) \in \mathbb{C}^n \mid P(\mathbf{e}) = 0, \, \forall P \in \mathcal{I}^{(n,m)} \}.$$
(5.4)

**Proposition 5.4** (Spectral Variety). For  $\alpha = \frac{2\pi}{m+ng}$  with  $g \in (0, \infty)$ , the spectral variety  $V(\mathcal{I}^{(n,m)})$  (5.4) is given by the joint spectrum  $\mathbf{e}_{\nu}\left(\frac{2\pi}{m+ng}, g; p\right)$  (5.3a)–(5.3c) of the truncated elliptic Ruijsenaars operators  $D_r$  (5.1):

$$V(\mathcal{I}^{(n,m)}) = \mathbb{E}_0^{(n,m)} = \{ \mathbf{e}_{\nu} \left( \frac{2\pi}{m+ng}, g; p \right) \mid \nu \in \Lambda_0^{(n,m)} \}.$$
(5.5)

*Proof.* Upon specializing the parameters in the Pieri rule (4.3b) with the aid of Proposition 5.2, one sees that for  $\lambda \in \Lambda_0^{(n,m)}$ ,  $g \in (0,\infty)$  and  $\mathbf{e} \in V(\mathcal{I}^{(n,m)})$  (5.4):

$$\mathbf{e}_{r} P_{\lambda}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) =$$

$$\sum_{\substack{\lambda \subset \nu \subset \lambda+1^{n} \\ |\nu|=|\lambda|+r, d_{\nu} \leq m}} \psi_{\nu/\lambda}'(\frac{2\pi}{m+ng}, \mathbf{g}; p) P_{\underline{\nu}}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) \quad \text{for } 1 \leq r < n.$$
(5.6)

As in the proof of Theorem 3.2, we can now employ Eq. (3.6b) and Lemma 5.3 to rewrite Eq. (5.6) in the form  $D_r p(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) = \mathbf{e}_r p(\mathbf{e}; \frac{2\pi}{m+ng}, g; p)$  with  $D_r$  and  $p_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p)$  taken from Eq. (5.1) and Eqs. (3.4a), (3.4b), respectively. Since  $p_0(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) = 1 \neq 0$ , this implies that  $\mathbf{e}$  must belong to the joint spectrum  $\mathbb{E}_0^{(n,m)}$  (5.5) of the operators  $D_1, \ldots, D_{n-1}$  in  $\ell^2(\Lambda_0^{(n,m)}, \Delta)$ . Reversely, if we assume that  $\mathbf{e} \in \mathbb{E}_0^{(n,m)}$  (5.5) then the eigenvalue equation entails that Eq. (5.6) holds for  $1 \leq r < n$ . The recurrence relations (3.1a)–(3.1c) thus yield in this situation that  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) = 0$  if  $d_{\mu} = m+1$  (where we use that  $\psi'_{\lambda+1r/\lambda}(\frac{2\pi}{m+ng}, g; p) = 1$  and that the solution of the recurrence is unique). In other words, the vector of joint eigenvalues  $\mathbf{e}$  necessarily lies on the zero locus  $V(\mathcal{I}^{(n,m)})$  (5.4) of  $\mathcal{I}^{(n,m)}$  (4.6).  $\Box$ 

5.3. Verlinde algebra. Let us now define the Verlinde algebra associated with the elliptic Ruijsenaars system as the  $\binom{n-1+m}{m}$ -dimensional algebra of complex functions on the joint spectrum  $\mathbb{E}_{0}^{(n,m)}$  (5.5):

$$\mathcal{F}_{0}^{(n,m)} = \{ f : \mathbb{E}_{0}^{(n,m)} \to \mathbb{C} \},$$
(5.7)

and let us write  $\mathcal{R}_{0,\mathbb{C}}^{(n)}$  and  $\mathcal{R}_{0,\mathbb{C}}^{(n,m)}$  for the algebras obtained by complexifying  $\mathcal{R}_{0}^{(n)}$ (4.2a) and  $\mathcal{R}_{0}^{(n,m)}$  (4.9).

**Proposition 5.5** (Basis for  $\mathcal{F}_0^{(n,m)}$ ). For  $\alpha = \frac{2\pi}{m+ng}$  with g > 0, the restrictions of the polynomials  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p), \ \mu \in \Lambda_0^{(n,m)}$  on the joint spectrum  $\mathbb{E}_0^{(n,m)}$  (5.5) constitute a basis for the Verlinde algebra  $\mathcal{F}_0^{(n,m)}$  (5.7).

Proof. It is immediate from the fact that the eigenfunctions  $p(\mathbf{e}_{\nu}), \nu \in \Lambda_0^{(n,m)}$  of the normal operators  $D_1, \ldots, D_{n-1}$  (5.1) provide an orthogonal basis for the Hilbert space  $\ell^2(\Lambda_0^{(n,m)}, \Delta)$  (cf. [DG21, Theorem 8]) that the square matrix  $[P_{\mu}(\mathbf{e}_{\nu})]_{\mu,\nu\in\Lambda_0^{(n,m)}}$  is of full rank.

It is plain from Propositions 5.4 and 5.5 that for  $\mathbf{g} \in (0, \infty)$  the vanishing ideal  $\mathcal{I}(\mathbb{E}_{0}^{(n,m)}) = \{P \in \mathcal{R}_{0}^{(n)} \mid P(\mathbf{e}) = 0, \forall \mathbf{e} \in \mathbb{E}_{0}^{(n,m)}\}$  is equal to  $\mathcal{I}^{(n,m)}$  (4.6). It means that in this situation the Verlinde algebra  $\mathcal{F}_{0}^{(n,m)}$  and the  $\widehat{\mathfrak{su}}(n)_{m}$  elliptic fusion algebra  $\mathcal{R}_{0,\mathbb{C}}^{(n,m)}$  are isomorphic, as the kernel of the evaluation homomorphism  $P(\mathbf{e}) \to P(\mathbf{e}_{\nu})$  from  $\mathcal{R}_{0,\mathbb{C}}^{(n)}$  onto  $\mathcal{F}_{0}^{(n,m)}$  coincides with the complexification of  $\mathcal{I}^{(n,m)}$  (4.6).

**Corollary 5.6**  $(\mathcal{R}_{0,\mathbb{C}}^{(n,m)} \cong \mathcal{F}_0^{(n,m)})$ . For  $\alpha = \frac{2\pi}{m+ng}$  with  $g \in (0,\infty)$ , the evaluation homomorphism  $P(\mathbf{e}) \to P(\mathbf{e}_{\nu})$  from  $\mathcal{R}_{0,\mathbb{C}}^{(n)}$  onto  $\mathcal{F}_0^{(n,m)}$  induces an algebra isomorphism  $[P] \to P(\mathbf{e}_{\nu})$  from  $\mathcal{R}_{0,\mathbb{C}}^{(n,m)}$  onto  $\mathcal{F}_0^{(n,m)}$ . Corollary 5.6 makes it trivial to determine further algebraic properties of  $\mathcal{R}_0^{(n,m)}$ and  $\mathcal{I}^{(n,m)}$ . For instance, it is manifest from the isomorphism that the factor ring  $\mathcal{R}_0^{(n,m)}$  does not have nilpotents, i.e. it is a reduced ring and  $\mathcal{I}^{(n,m)}$  is a radical ideal. However, clearly the factor ring  $\mathcal{R}_0^{(n,m)}$  has zero divisors (stemming from functions in  $\mathcal{F}_0^{(n,m)}$  with disjoint support), i.e. it is not an integral domain and the ideal  $\mathcal{I}^{(n,m)}$  is therefore neither maximal nor prime.

With the aid of the Verlinde algebra  $\mathcal{F}_0^{(n,m)}$  we are now in the position to extend the elliptic fusion rules in Theorem 4.4 so as to include the case of positive rational values for g.

**Theorem 5.7** (Structure Constants of  $\mathcal{F}_0^{(n,m)}$ ). For  $\alpha = \frac{2\pi}{m+ng}$  with g > 0, the structure constants of the Verlinde algebra  $\mathcal{F}_0^{(n,m)}$  in the basis  $P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p)$ ,  $\mu \in \Lambda_0^{(n,m)}$  can be expressed in terms of elliptic Littlewood-Richardson coefficients as follows:

$$P_{\lambda}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p) =$$

$$\sum_{\mathbf{c} \in \mathbb{R} \setminus \mathbb{Q}} \left( \lim_{\mathbf{c} \in \mathbb{R} \setminus \mathbb{Q}} c_{\lambda,\mu}^{\nu} \left( \frac{2\pi}{m+nc}, \mathbf{c}; p \right) \right) P_{\underline{\nu}}(\mathbf{e}; \frac{2\pi}{m+ng}, g; p)$$
(5.8a)

 $(with \ \lambda, \mu \in \Lambda_0^{(n,m)}, \ \nu \in \Lambda^{(n)} \ and \ \mathbf{e} \in \mathbb{E}_0^{(n,m)}).$ 

particular, for 
$$1 \leq r < n$$
,  $\lambda \in \Lambda_0^{(n;m)}$  and  $\mathbf{e} \in \mathbb{E}_0^{(n;m)}$  one has explicitly that  
 $P_{\lambda}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) P_{1^r}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) =$ 

$$\sum_{\substack{\lambda \subset \nu \subset \lambda + 1^n \\ |\nu| = |\lambda| + r, d_{\nu} \leq m}} \psi'_{\nu/\lambda} \left(\frac{2\pi}{m+ng}, \mathbf{g}; p\right) P_{\underline{\nu}}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p).$$
(5.8b)

*Proof.* If we pick g > 0 irrational, then it is clear from Theorem 4.4 and Corollary 5.6 that on  $\mathbb{E}_0^{(n,m)}$ :

$$P_{\lambda}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) P_{\mu}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) = \sum_{\substack{\nu \supset \lambda, \ \nu \supset \mu \\ |\nu| = |\lambda| + |\mu|, \ d_{\nu} \le m}} c_{\lambda,\mu}^{\nu} \left(\frac{2\pi}{m+ng}, \mathbf{g}; p\right) P_{\underline{\nu}}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p),$$

which establishes in particular the validity of the Pieri rule (5.8b) in this situation. Rational values of g > 0 can now be included by analytic continuation, which gives rise to Eq. (5.8a). Indeed, in the case of the Pieri rule the analytic continuation is achieved through the explicit formulas with the aid of Lemma 5.1 and Proposition 5.2 (where one also uses that the spectral points  $\mathbf{e}_{\nu} = \mathbf{e}_{\nu}(\mathbf{g}, p)$  are analytic in  $\mathbf{g} > 0$ by the normality of  $D_r$  (5.1) in  $\ell^2(\Lambda_0^{(n,m)}, \Delta)$ , cf. e.g. [K95, Chapter 2, Theorem 1.10]). Since the monomials  $P_{1r}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p) = \mathbf{e}_r$ ,  $1 \leq r < n$  generate  $\mathcal{F}_0^{(n,m)}$ , the analyticity of the Pieri rule is inherited in turn by the structure constants of  $\mathcal{F}_0^{(n,m)}$  in general. To see this, it suffices to expand one of the two factors in the product on the LHS of Eq. (5.8a) in monomials (cf. Proposition 3.1). Since the corresponding expansion coefficients are analytic by virtue of Proposition 5.2, iterated application of the Pieri rules then leads to the analyticity of the structure constants as claimed. 5.4. Verlinde formula. Let us assume that g > 0 and denote the squared norms of the eigenbasis  $p(\mathbf{e}_{\nu}), \nu \in \Lambda_0^{(n,m)}$  for the elliptic Ruijsenaars operators  $D_r$  (5.1) in the Hilbert space by  $\ell^2(\Lambda_0^{(n,m)}, \Delta)$  by (cf. Eqs. (5.2a), (5.2b)):

$$\hat{\Delta}_{\nu} = 1/\langle p(\mathbf{e}_{\nu}), p(\mathbf{e}_{\nu}) \rangle_{\Delta} \qquad (\nu \in \Lambda_0^{(n,m)}).$$
(5.9)

We now endow the Verlinde algebra  $\mathcal{F}_0^{(n,m)}$  (5.7) with the following inner product:

$$\langle f, g \rangle_{\hat{\Delta}} = \sum_{\nu \in \Lambda_0^{(n,m)}} f(\mathbf{e}_{\nu}) \overline{g(\mathbf{e}_{\nu})} \hat{\Delta}_{\nu} \qquad (f, g \in \mathcal{F}_0^{(n,m)}).$$
 (5.10)

The basis of the Verlinde algebra in Proposition 5.5 is then orthogonal with respect to this inner product [DG21, Corollary 11], viz. one has that

$$\forall \lambda, \mu \in \Lambda_0^{(n,m)}: \qquad \langle P_\lambda, P_\mu \rangle_{\hat{\Delta}} = \begin{cases} \frac{1}{c_\lambda^2 \Delta_\lambda} & \text{if } \lambda = \mu, \\ 0 & \text{if } \lambda \neq \mu, \end{cases}$$
(5.11)

with  $c_{\lambda}$  and  $\Delta_{\lambda}$  given by Eqs. (3.4b) and (5.2b). With the aid of this dual orthogonality relation one arrives at a Verlinde formula expressing the structure constants

$$P_{\lambda}\left(\mathbf{e}; \frac{2\pi}{m+ng}, g; p\right) P_{\mu}\left(\mathbf{e}; \frac{2\pi}{m+ng}, g; p\right) =$$

$$\sum_{\kappa \in \Lambda_{0}^{(n,m)}} N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+ng}, g; p\right) P_{\kappa}\left(\mathbf{e}; \frac{2\pi}{m+ng}, g; p\right) \qquad (\mathbf{e} \in \mathbb{E}_{0}^{(n,m)})$$

$$(\mathbf{e} \in \mathbb{E}_{0}^{(n,m)})$$

for  $\mathcal{F}_0^{(n,m)}$  in terms of a corresponding S-matrix stemming from the elliptic Ruijsenaars model.

**Theorem 5.8** (Verlinde Formula). For g > 0 the structure constants  $N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+ng}, g; p\right)$  (5.12) for the Verlinde algebra  $\mathcal{F}_{0}^{(n,m)}$  (5.7) are given by

$$N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+ng}, g; p\right) = \sum_{\nu \in \Lambda_0^{(n,m)}} \frac{S_{\lambda,\nu} S_{\mu,\nu} S_{\nu,\kappa}^{-1}}{S_{0,\nu}}$$
(5.13a)

with

$$S_{\lambda,\nu} = S_{\lambda,\nu} \left(\frac{2\pi}{m+ng}, g; p\right) = \frac{P_{\lambda} \left(\mathbf{e}_{\nu}; \frac{2\pi}{m+ng}, g; p\right)}{c_{\nu} \left(\frac{2\pi}{m+ng}, g; p\right)}$$
(5.13b)

and

$$S_{\lambda,\nu}^{-1} = c_{\lambda}^2 \hat{\Delta}_{\lambda} \overline{S_{\nu,\lambda}} c_{\nu}^2 \Delta_{\nu}.$$
 (5.13c)

*Proof.* Let us first observe that it is clear from the orthogonality relation in Eq. (5.11) that the inverse of the S-matrix  $S_{\lambda,\nu}$  (5.13b) is given by the matrix  $S_{\lambda,\nu}^{-1}$  (5.13c). Hence, upon pairing both sides of Eq. (5.12) against  $P_{\kappa}(\mathbf{e}; \frac{2\pi}{m+ng}, \mathbf{g}; p), \kappa \in \Lambda_0^{(n,m)}$  with  $\langle \cdot, \cdot \rangle_{\hat{\Delta}}$  (5.10), the Verlinde formula readily follows via the orthogonality relation (5.11):

$$N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+ng}, g; p\right) = \frac{\langle P_{\lambda}P_{\mu}, P_{\kappa}\rangle_{\hat{\Delta}}}{\langle P_{\kappa}, P_{\kappa}\rangle_{\hat{\Delta}}} = c_{\kappa}^{2}\Delta_{\kappa}\sum_{\nu\in\Lambda_{0}^{(n,m)}} P_{\lambda}(\mathbf{e}_{\nu})P_{\mu}(\mathbf{e}_{\nu})\overline{P_{\kappa}(\mathbf{e}_{\nu})}\hat{\Delta}_{\nu}$$
$$= \sum_{\nu\in\Lambda_{0}^{(n,m)}} \frac{S_{\lambda,\nu}S_{\mu,\nu}S_{\nu,\kappa}^{-1}}{S_{0,\nu}}.$$

It is useful to recall though that we already know from Theorem 5.7 that many of the structure constants in Eq. (5.12) in fact vanish:

$$\begin{split} \mathbf{N}_{\lambda,\mu}^{\kappa} \Big( \frac{2\pi}{m+n\mathbf{g}},\mathbf{g};p \Big) &= \\ \begin{cases} \lim_{\substack{\mathbf{c} \in \mathbb{R} \setminus \mathbb{Q} \\ \mathbf{c} \to \mathbf{g}}} c_{\lambda,\mu}^{\nu} \Big( \frac{2\pi}{m+n\mathbf{c}},\mathbf{c};p \Big) & \text{if } \kappa = \underline{\nu} \text{ with } \nu \supset \lambda, \, \nu \supset \mu, \, \text{and } |\nu| = |\lambda| + |\mu|, \\ 0 & \text{otherwise.} \end{split}$$

Notice also that it follows from Eq. (5.13c) that the absolute value of the determinant of the S-matrix is given by

$$|\det S| = \left(\prod_{\lambda \in \Lambda_0^{(n,m)}} c_\lambda^2 \sqrt{\Delta_\lambda \hat{\Delta}_\lambda}\right)^{-1}.$$
(5.14)

5.5. **Degenerations.** To see how the structure constants of the  $\widehat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion ring and its refined deformation are recovered from  $\mathcal{F}_0^{(n,m)}$ , let us put

$$q = e^{\frac{2\pi i}{m+ng}}$$
 and  $e_r = (x_1 \cdots x_n)^{-r/n} m_{1^r}(x)$   $1 \le r < n.$  (5.15)

Then for g > 0,  $-1 and <math>\lambda, \mu \in \Lambda_0^{(n,m)}$  we have that:

$$\psi_{\nu/\lambda}'\left(\frac{2\pi}{m+ng}, g; p\right) =$$

$$\begin{cases} \prod_{\substack{1 \le j < k \le n \\ \theta_j - \theta_k = -1}} \frac{[\nu_j - \nu_k + g(k-j+1)]_q}{[\nu_j - \nu_k + g(k-j)]_q} \frac{[\lambda_j - \lambda_k + g(k-j-1)]_q}{[\lambda_j - \lambda_k + g(k-j)]_q} & \text{if } p = 0, \\ 1 & \text{if } g = 1 \end{cases}$$
(5.16a)

(for  $\lambda \subset \nu \subset \lambda + 1^n$  with  $\theta = \nu - \lambda$  and  $\underline{\nu} \in \Lambda_0^{(n,m)}$ ),

$$P_{\mu}\left(\mathbf{e}; \frac{2\pi}{m+ng}, g; p\right) = \begin{cases} (x_1 \cdots x_n)^{-|\mu|/n} P_{\mu}(x; q, q^g) & \text{if } p = 0, \\ (x_1 \cdots x_n)^{-|\mu|/n} s_{\mu}(x) & \text{if } g = 1, \end{cases}$$
(5.16b)

and

$$e_{r,\nu}\left(\frac{2\pi}{m+ng}, g; p\right) =$$

$$\begin{cases} q^{-r\left(\frac{|\nu|}{n} + \frac{(n-1)g}{2}\right)} m_{1r}(q^{\nu_1 + (n-1)g}, q^{\nu_2 + (n-2)g}, \dots, q^{\nu_{n-1} + g}, 1) & \text{if } p = 0, \\ q^{-r\left(\frac{|\nu|}{n} + \frac{n-1}{2}\right)} m_{1r}(q^{\nu_1 + n-1}, q^{\nu_2 + n-2}, \dots, q^{\nu_{n-1} + 1}, 1) & \text{if } g = 1 \end{cases}$$
(5.16c)

(for  $\nu \in \Lambda_0^{(n,m)}$ ). Indeed, Eqs. (5.16a) and (5.16b) follow from Proposition 3.4 and Corollary 3.6 by Lemma 5.1, while Eq. (5.16c) is immediate from Eq. (5.3c) when p = 0, and hence also when g = 1 (since the *p*-dependence drops out at g = 1).

At p = 0, Theorem 5.7 therefore computes the structure constants of the refined Verlinde algebra [AS15, C05, K96, N14] in terms of Macdonald's (q, t)-Littlewood-Richardson coefficients (cf. Proposition 3.4):

$$N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+ng}, g; 0\right) =$$

$$\begin{cases} \lim_{\substack{c \in \mathbb{R} \setminus \mathbb{Q} \\ c \to g}} f_{\lambda,\mu}^{\nu}\left(e^{\frac{2\pi i}{m+nc}}, e^{\frac{2\pi i c}{m+nc}}\right) & \text{if } \kappa = \underline{\nu} \text{ with } \nu \supset \lambda, \nu \supset \mu, \text{ and } |\nu| = |\lambda| + |\mu|, \\ 0 & \text{otherwise.} \end{cases}$$
(5.17a)

In the case of the Pieri rule this becomes explicitly

$$N_{\lambda,1^{r}}^{\kappa}\left(\frac{2\pi}{m+ng}, g; 0\right) =$$

$$\begin{cases} \psi_{\nu/\lambda}^{\prime}\left(\frac{2\pi}{m+ng}, g; 0\right) & \text{if } \kappa = \underline{\nu} \text{ with } \lambda \subset \nu \subset \lambda + 1^{n}, \text{ and } |\nu| = |\lambda| + r, \\ 0 & \text{otherwise.} \end{cases}$$
(5.17b)

At g = 1, on the other hand, we see from the Pieri rule in Theorem 5.7 that

$$N_{\lambda,1^{r}}^{\kappa}\left(\frac{2\pi}{m+n},1;p\right) = N_{\lambda,1^{r}}^{\kappa}\left(\frac{2\pi}{m+n},1;0\right)$$

$$\begin{cases} 1 & \text{if } \kappa = \underline{\nu} \text{ with } \lambda \subset \nu \subset \lambda + 1^{n}, \text{ and } |\nu| = |\lambda| + r, \\ 0 & \text{otherwise.} \end{cases}$$

$$(5.18a)$$

More generally, Theorem 5.7 thus states that

$$N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+n},1;p\right) = N_{\lambda,\mu}^{\kappa}\left(\frac{2\pi}{m+n},1;0\right) =$$

$$\begin{cases} \lim_{\substack{c \in \mathbb{R} \setminus \mathbb{Q} \\ c \to 1 \\ 0 \end{cases}} f_{\lambda,\mu}^{\nu}\left(e^{\frac{2\pi i}{m+nc}}, e^{\frac{2\pi i c}{m+nc}}\right) & \text{if } \kappa = \underline{\nu} \text{ with } \nu \supset \lambda, \nu \supset \mu, \text{ and } |\nu| = |\lambda| + |\mu|, \\ 0 & \text{otherwise,} \end{cases}$$

$$(5.18b)$$

which retrieves the structure constants of the  $\widehat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion ring [DMS97, F95] through Macdonald's (q, t)-Littlewood-Richardson coefficients. In other words, (q, t)-deformation can be used as a vehicle for computing structure constants in the fusion ring (i.e. modulo the fusion ideal) by degeneration from (deformed) Littlewood-Richardson coefficients in the ring of symmetric polynomials itself, as was previously pointed out in [D20].

Finally, we see from Eqs. (5.16b), (5.16c) that the corresponding degenerations of the elliptic S-matrix (5.13b) are given by

$$S_{\lambda,\nu}\left(\frac{2\pi}{m+ng}, g; 0\right) = q^{-\frac{1}{n}|\lambda||\nu| - \frac{1}{2}(n-1)(|\lambda|+|\nu|)g}$$

$$\times P_{\lambda}(q^{\nu_{1}+(n-1)g}, q^{\nu_{2}+(n-2)g}, \dots, q^{\nu_{n-1}+g}, 1; q, q^{g})$$

$$\times P_{\nu}(q^{(n-1)g}, q^{(n-2)g}, \dots, q^{g}, 1; q, q^{g}),$$
(5.19a)

and

$$S_{\lambda,\nu}\left(\frac{2\pi}{m+n}, 1; p\right) = S_{\lambda,\nu}\left(\frac{2\pi}{m+n}, 1; 0\right) \frac{c_{\nu}\left(\frac{2\pi}{m+n}, 1; 0\right)}{c_{\nu}\left(\frac{2\pi}{m+n}, 1; p\right)}$$
(5.19b)

with

$$S_{\lambda,\nu}\left(\frac{2\pi}{m+n},1;0\right) = q^{-\frac{1}{n}|\lambda||\nu|-\frac{1}{2}(n-1)(|\lambda|+|\nu|)} \\ \times s_{\lambda}(q^{\nu_{1}+n-1},q^{\nu_{2}+n-2},\ldots,q^{\nu_{n-1}+1},1) \\ \times s_{\nu}(q^{n-1},q^{n-2},\ldots,q,1).$$

Notice that the gauge factor multiplying the S-matrix in Eq. (5.19b) from the right cancels in the Verlinde formula (5.13a), and observe also that we have rewritten the final expression with the aid of the principal specialization formulas for the Macdonald and Schur polynomials [M95, Chapter VI, §6]:

$$q^{-\frac{1}{2}|\nu|(n-1)g} P_{\nu}(q^{(n-1)g}, q^{(n-2)g}, \dots, q^{g}, 1; q, q^{g}) = \prod_{1 \le j < k \le n} \frac{[(k-j+1)g]_{q,\nu_{j}-\nu_{k}}}{[(k-j)g]_{q,\nu_{j}-\nu_{k}}}$$

$$(=1/c_{\nu}(\frac{2\pi}{m+ng}, \mathbf{g}; \mathbf{0})) \text{ and}$$
$$q^{-\frac{1}{2}|\nu|(n-1)}s_{\nu}(q^{n-1}, q^{n-2}, \dots, q, \mathbf{1}) = \prod_{1 \le j < k \le n} \frac{[k-j+\nu_j-\nu_k]_q}{[k-j]_q}$$

 $(=1/c_{\nu}(\frac{2\pi}{m+n},1;0))$ , where  $[z]_{q,k} = \prod_{0 \le l < k} [z+l]_q$  with  $[z]_{q,0} = 1$ . We thus observe that  $S_{\lambda,\nu}(\frac{2\pi}{m+n},1;0)$  coincides up to normalization with (the

We thus observe that  $S_{\lambda,\nu}(\frac{2\pi}{m+n}, 1; 0)$  coincides up to normalization with (the adjoint of) the Kac-Peterson modular S-matrix for  $\hat{\mathfrak{su}}(n)_m$  (cf. e.g. [K90, Theorem 13.8], [DMS97, Equation (14.217)] and [KS10, Proposition 6.15]), while the S-matrix  $S_{\lambda,\nu}(\frac{2\pi}{m+ng}, g; 0)$  recovers in turn the trigonometric S-matrix from the refined Chern-Simons theory of knot invariants (cf. e.g. [AS15, Section 5.2], [GN15, Section 2.6], [K96, Section 5] and [N14, Section 3]). In other words, Theorem 5.8 reproduces in these two situations, respectively, the classical Verlinde formula for the structure constants of the  $\hat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion ring (cf. e.g. [K90, Exercise 13.34], [DMS97, Equation (16.3)], [KS10, Equation (6.40]) as well as its refinement stemming from Macdonald theory [AS15, D20]. The conventional normalization of the S-matrix corresponds to the following rescaling:

$$S_{\lambda,\nu}\left(\frac{2\pi}{m+ng}, g; p\right) \to n^{-\frac{1}{2}}\left(\frac{2\pi}{m+ng}, g; p\right) S_{\lambda,\nu}\left(\frac{2\pi}{m+ng}, g; p\right)$$
(5.20a)

with

$$n\left(\frac{2\pi}{m+ng}, g; p\right) = \sum_{\lambda \in \Lambda_0^{(n,m)}} \Delta_\lambda\left(\frac{2\pi}{m+ng}, g; p\right).$$
(5.20b)

Notice in this connection that at the classical parameter specialization (g, p) = (1, 0)this normalization recovers (the adjoint of) the unitary Kac-Peterson S-matrix for  $\widehat{\mathfrak{su}}(n)_m$ :

$$n\left(\frac{2\pi}{m+n}, 1; 0\right) = \sum_{\lambda \in \Lambda_0^{(n,m)}} \prod_{1 \le j < k \le n} \frac{[\lambda_j - \lambda_k + k - j]_q^2}{[k-j]_q^2} = \sum_{\lambda \in \Lambda_0^{(n,m)}} \prod_{1 \le j < k \le n} S_{0,\lambda}^2\left(\frac{2\pi}{m+n}, 1; 0\right)$$
  
(so  $n\left(\frac{2\pi}{m+n}, 1; 0\right) = \frac{(2\sin\frac{\pi}{m+n})^{-n(n-1)}n(n+m)^{n-1}}{\prod_{1 \le j < k \le n} [k-j]_q^2}$ , cf. e.g. [KS10, Proposition 6.15]).

#### Acknowledgements

Helpful feedback from Stephen Griffeth is gratefully acknowledged.

The work of JFvD was supported in part by the *Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT)* Grant # 1210015. TG was supported in part by the NKFIH Grant K134946.



This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 795471.

## References

- AS15. M. Aganagic and S. Shakirov, Knot homology and refined Chern-Simons index, Comm. Math. Phys. 333 (2015), 187–228.
- AGP16. J.E. Andersen, S. Gukov, and D. Pei, The Verlinde formula for Higgs bundles, arXiv:1608.01761.
- BDM15. O. Blondeau-Fournier, P. Desrosiers, and P. Mathieu, Supersymmetric Ruijsenaars-Schneider model, Phys. Rev. Lett. 114 (2015), 121602.
- B68. N. Bourbaki, Groupes et algèbres de Lie, Chapitres 4-6, Hermann, Paris, 1968.
- C05. I. Cherednik, Double Affine Hecke Algebras, London Mathematical Society Lecture Note Series 319, Cambridge University Press, Cambridge, 2005.

- C16. I. Cherednik, DAHA-Jones polynomials of torus knots, Selecta Math. (N.S.) **22** (2016), 1013–1053.
- D20. J.F. van Diejen,  $\hat{\mathfrak{su}}(n)_m$  Wess-Zumino-Witten fusion rules via Macdonald polynomials, Preprint April 2020.
- DG21. J.F. van Diejen and T. Görbe, Elliptic Ruijsenaars difference operators on bounded partitions, arXiv:2106.06512
- DV98. J.F. van Diejen and L. Vinet, The quantum dynamics of the compactified trigonometric Ruijsenaars-Schneider model, Comm. Math. Phys. **197** (1998), 33–74.
- DMS97. P. Di Francesco, P. Mathieu, and D. Sénéchal, Conformal Field Theory, Graduate Texts in Contemporary Physics, Springer-Verlag, 1997.
- EK95. P.I. Etingof and A. Kirillov Jr., On the affine analogue of Jack and Macdonald polynomials, Duke Math. J. 78 (1995), 229–256.
- FG16. L. Fehér and T.F. Görbe, Trigonometric and elliptic Ruijsenaars-Schneider systems on the complex projective space, Lett. Math. Phys. **106** (2016), 1429–1449.
- F95. J. Fuchs, Affine Lie Algebras and Quantum Groups. An Introduction, with Applications in Conformal Field Theory. Corrected reprint, Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1995.
- G91. D. Gepner, Fusion rings and geometry, Comm. Math. Phys. 141 (1991), 381–411.
- GN91. F.M. Goodman and T. Nakanishi, Fusion algebras in integrable systems in two dimensions, Phys. Lett. B 262 (1991), 259–264.
- GW90. F.M. Goodman and H. Wenzl, Littlewood-Richardson coefficients for Hecke algebras at roots of unity, Adv. Math. 82 (1990), 244–265.
- GH18. T.F. Görbe and M. Hallnäs, Quantization and explicit diagonalization of new compactified trigonometric Ruijsenaars-Schneider systems, J. Integrable Syst. 3 (2018), no. 1, xyy015, 29 pp.
- GN15. E. Gorsky and A. Neguţ, Refined knot invariants and Hilbert schemes, J. Math. Pures Appl. 104 (2015), 403–435.
- H16. J. Haglund, The combinatorics of knot invariants arising from the study of Macdonald polynomials. In: *Recent Trends in Combinatorics*, A. Beveridge, J.R. Griggs, L. Hogben, G. Musiker and P. Tetali (eds.), The IMA Volumes in Mathematics and its Applications **159**, Springer, Cham, 2016, 579–600.
- K90. V.G. Kac, Infinite-dimensional Lie Algebras, Third Edition, Cambridge University Press, Cambridge, 1990.
- K95. T. Kato, Perturbation Theory for Linear Operators, Reprint of the 1980 edition, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
- K96. A.A. Kirillov Jr., On an inner product in modular tensor categories, J. Amer. Math. Soc. 9 (1996), 1135–1169.
- K13. C. Korff, Cylindric versions of specialised Macdonald functions and a deformed Verlinde algebra, Comm. Math. Phys, **318** (2013), 173–246.
- KS10. C. Korff and C. Stroppel, The  $\hat{\mathfrak{sl}}(n)_k$ -WZNW fusion ring: a combinatorial construction and a realisation as quotient of quantum cohomology, Adv. Math. **225** (2010), 200–268.
- LNS20. E. Langmann, M. Noumi, and J. Shiraishi, Construction of eigenfunctions for the elliptic Ruijsenaars difference operators, arXiv:2012.05664
- M95. I.G. Macdonald, Symmetric Functions and Hall Polynomials, Second Edition, Clarendon Press, Oxford, 1995.
- M00. I.G. Macdonald, Orthogonal polynomials associated with root systems, Sém. Lothar. Combin. 45 (2000/01), Art. B45a.
- M03. I.G. Macdonald, Affine Hecke Algebras and Orthogonal Polynomials, Cambridge University Press, Cambridge, 2003.
- MMZ21. A. Mironov, A. Morozov, and Y. Zenkevich, Duality in elliptic Ruijsenaars system and elliptic symmetric functions, Eur. Phys. J. C 81 (2021), 461.
- N14. H. Nakajima, Refined Chern-Simons theory and Hilbert schemes of points on the plane.
  In: Perspectives in Representation Theory, P. Etingof, M. Khovanov, and A. Savage (eds.), Contemp. Math. 610, Amer. Math. Soc., Providence, RI, 2014, 305–331.
- OY14. S. Okuda and Y. Yoshida, G/G gauged WZW-matter model, Bethe Ansatz for qboson model and commutative Frobenius algebra, J. High Energ. Phys. 2014:3 (2014), https://doi.org/10.1007/JHEP03(2014)003

ELLIPTIC RUIJSENAARS OPERATORS AND WESS-ZUMINO-WITTEN FUSION RINGS 21

- OLBC10. F.W.J. Olver, D.W. Lozier, R.F. Boisvert and C.W. Clark. (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press, Cambridge, 2010.
- P07. C. Procesi, Lie Groups. An Approach through Invariants and Representations, Springer, New York, 2007.
- RSV18. E.M. Rains, Y. Sun, and A. Varchenko, Affine Macdonald conjectures and special values of Felder-Varchenko functions, Selecta Math. (N.S.) **24** (2018), 1549–1591.
- R87. S.N.M. Ruijsenaars, Complete integrability of relativistic Calogero-Moser systems and elliptic function identities, Comm. Math. Phys. 110 (1987), 191–213.
- R99. S.N.M. Ruijsenaars, Systems of Calogero-Moser type. In: Particles and Fields (Banff, AB, 1994), G.W. Semenoff and L. Vinet (eds.), CRM Ser. Math. Phys., Springer, New York, 1999, 251–352.
- T04. C. Teleman, K-theory and the moduli space of bundles on a surface and deformations of the Verlinde algebra. In: *Topology, Geometry and Quantum Field Theory*, U. Tillmann (ed.), London Math. Soc. Lecture Note Ser. **308**, Cambridge Univ. Press, Cambridge, 2004, 358–378.
- TW09. C. Teleman and C.T. Woodward, The index formula for the moduli of *G*-bundles on a curve, Ann. of Math. (2) **170** (2009), 495–527.
- TUY89. A. Tsuchiya, K. Ueno, and Y. Yamada, Conformal field theory on universal family of stable curves with gauge symmetries. In: *Integrable Systems in Quantum Field Theory* and Statistical Mechanics, M. Jimbo, T. Miwa, and A. Tsuchiya (eds.), Adv. Stud. Pure Math. 19, Academic Press, Boston, MA, 1989, 459–566.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD DE TALCA, CASILLA 747, TALCA, CHILE *Email address*: diejen@inst-mat.utalca.cl

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UK *Email address*: T.Gorbe@leeds.ac.uk