Spectral concepts in genome informational analysis

V. Bonnici¹, G. Franco¹, V. Manca^{1,*}

^aUniversity of Verona, Department of Computer Science, Strada Le Grazie, 15 - 37134 Verona, Italy

Abstract

The concept of k-spectrum for genomes is here investigated as a basic tool to analyze genomes. Related spectral notions based on k-mers are introduced with some related mathematical properties which are relevant for informational analysis of genomes. Procedures to generate spectral segmentations of genomes are provided and are tested (under several values of length k for k-mers) on cases of real genomes, such as some human chromosomes and Saccharomyces cervisiae.

Keywords: Alignment-free method, genome, k-mer, k-spectrum, informational genomics

1. Introduction

The notion of spectrum comes from physics, usually related to a function of frequency (namely frequency spectrum, or power spectrum of a signal). The term applies to any signal decomposed along a variable, such as energy in electron spectroscopy or mass-to-charge ratio in mass spectrometry. In a wider sense, a spectrum is a distribution associated to a phenomenon, where a measure (intensity, density, frequency) is associated to each value taken from a variable. In fact, spectroscopy originated as the study of light intensity at the different wavelength of its components.

A linearized genome (that is, observed by neglecting the secondary and tertiary structure of the double helix) is a long sequence of characters from the DNA alphabet $\{A, T, C, G\}$, which may be seen as an information source, composed by a finite set of signals (e.g., words) emitted (or occurring) at a given "frequency". In particular, if we restrict ourself to consider all genomic substrings of a fixed length k (factors of length k, here called k-mers, in other contexts also called k-grams), we may call k-spectrum of a genome the distribution which associates to each k-mer the (normalized) number of times that it occurs in the genome (that is, its multiplicity, which is called frequency when normalized). If we denote by $D_k(\mathbb{G})$ the set of k-mers occurring in a genome \mathbb{G} , and by $mult_{\mathbb{G}}(\alpha)$ the multiplicity of a k-mer α in \mathbb{G} , the k-spectrum of a genome \mathbb{G} , is the multiset

$$spec_k(\mathbb{G}) = \{(\alpha, mult_{\mathbb{G}}(\alpha)) \mid \alpha \in D_k(\mathbb{G})\}.$$

In sequence analysis, the term "spectrum" is used in many contexts and with several meanings (see for example, [30, 32, 21, 19, 31]), in particular to tackle the problem of reconstructing a string from the compositions of its substrings. Other terms are used with a similar meaning, as bags of words, where a k-spectrum is represented by a multiplicity vector, where the i-component gives the multiplicity of the k-mer of position i in some prefixed (for example lexicographic) order [23]. The genomic profile is instead a related concept with a different meaning, as it accounts for the distribution of quantity of k-mers over their multiplicity, by ignoring the information on the specific k-mer sequences.

In general, string reconstruction refers to the long standing problem of recovering a string based on some information about its substrings (i.e, sequences composed of consecutive elements of the string) or

^{*}Corresponding author

 $Email\ addresses:$ vincenzo.bonnici@univr.it (V. Bonnici), giuditta.franco@univr.it (G. Franco), vincenzo.manca@univr.it (V. Manca)

subsequences (i.e, sequences composed of possibly non-consecutive elements of the string) [33]. Above problems may be formulated under specific hypotheses, for example whether or not we are given the order of the substrings in the string, or the order of the bits in each substring [3], also, predefined constraints may be imposed on the properties of the strings one seeks to reconstruct (see coded strings in [19] or [16] for a model of DNA self-assembly). The first instance of a coded sequence reconstruction problem was studied by Levenshtein [22], who posed the sequence reconstruction problem for strings drawn from an error-correcting codebook (setup in which not all substrings in the k-spectrum are received). Reconstruction of a string from a few substrings was considered in [14, 30] as well as the case where, for each substring, only the composition is given, neither the order of the symbols within it nor the substring's location in the original string. For example, in [31] reconstruction of strings based upon their error-prone substrings spectrum is developed, while the noisy setup of this problem was first studied by Gabrys and Milenkovic [19].

In several works the initial information is given in terms of the sequence spectrum, over a binary alphabet, comprising all distinct substrings (of variable length) appearing in the string: the multiset spectrum [21], comprising the multiset of the substrings of the string; the n-deck [22], comprising all subsequences of the string of length n; sequence traces [20], corresponding to randomly selected subsequences of the string; or multiset compositions [3], providing information about the composition of substrings only. These approaches are often motivated by molecular biology problems: mass-spectrometry protein sequencing [2, 3], DNA-based data storage, where the strings to be sequenced are user-defined and synthetically generated, and hence allowed to have arbitrary content [22], and account for types of errors that are encountered in both DNA synthesis and sequencing [2]. For example, the nanopore sequencing technology (developed by Oxford university) motivated the assumption of "traces", that are genomic subsequences long k which have passed through k deletion channels, deleting each character of the original string with a given probability.

In our (above defined) notion of genomic k-spectrum we are given all the error-free k-mers of the genome, with their multiplicity value, whereas we do not know their order on the genome nor their exact position. We focus on the relevance of the k-spectrum for genomes analysis, by exploiting the information contained in a k-spectrum to define new spectral notions which allow compact representations of genomes.

Many concepts based on k-mers, or analogous notions of scattered factors and de Bruijn sequences, are shared by many fields in Computer Science, e.g. combinatorics on words, frequently with different names and aims [4, 29, 35]. However, in this paper, k-mers and related spectra are mainly considered in the perspective of their possible elongation within the genomes, a feature which turns out very significant for their enormous lengths (from millions to billions of characters). Apparently related to the aim of this paper there are also the articles [3] and [2], motivated by mass-spectrometry protein sequencing. They report the simply-stated problem of reconstructing a string from the multiset of its substring compositions. However, the assumed substring composition multiset contains substrings of variable lengths, and the interesting results in [3] are found only for short strings (e.g., strings of length 7 can be reconstructed uniquely up to reversal), while the general problem formulation being a combinatorial simplification of the well-known turnpike problem. Interestingly, general classes of strings that cannot be distinguished from their substring compositions are provided in [2].

Works dealing with (real) genome recombinations and reconstructions often assume to start with a set of substrings for which multiplicity is not known, as it is the case for ciliates rearrangements [9, 10]. On the other hand, current methods for DNA assembling which belong to "next-generation" (of second, third, and fourth generation) sequencing (NGS) allow us to have several duplicates of given k-mers.

In this paper we aim at providing correspondences between informational profiles and biological interpretations, as in [37, 36], where genomic sequence entropy and complexity are investigated to discover evolution patterns, also discussed in [28]. Some concepts related to genomic k-spectra have been introduced in our previous work [11, 15, 25, 26, 27, 6, 5, 7, 8, 26, 15, 18] and are here recalled in Section 2. In particular, the distinction between hapax and repeat substring, as defined later on, and all the notions related to them [12, 17]. Their importance in genome analyses emerged in the course of investigations developed by means of information theory concepts. After a brief discussion in Section 2 on genomic k-spectra, in Section 3 we introduce related notions of spectral segments, spectral segmentation, k-univocal genomes and genome spectrality, by showing their relevance to recognize long portions of genomes univocally identified by spectra. Section 4 concludes our contribution with a few final comments and observations.

2. Properties related to k-spectra

In this section we present some informational properties of a genome \mathbb{G} , which are given within its k-spectrum:

$$spec_k(\mathbb{G}) = \{(\alpha, mult_{\mathbb{G}}(\alpha)) \mid \alpha \in D_k(\mathbb{G})\}.$$

The usual operations of multiset-sum and multiset-difference are assumed for k-spectra. In particular, the elements of a multiset-sum A+B are those of A or B with the multiplicities given by the sum of their multiplicities in A and B, while those of A-B have as multiplicities the differences of multiplicities, by setting to zero the negative values. Moreover, a multiset does not change if a new element with zero multiplicity is added or removed, therefore a multiset having only zero multiplicities can be considered equivalent to the empty set.

The support of a k-spectrum of a genome \mathbb{G} is the k-dictionary $D_k(\mathbb{G})$, that is, the set of all k-mers occurring at least once in \mathbb{G} . Notice that we are interested to study this specific dictionary with a prefixed k, rather than the dictionary of genomic words

$$D(\mathbb{G}) = \bigcup_{1 \le k \le |\mathbb{G}|} D_k(\mathbb{G}).$$

However, all the notions introduced here can be extended to this dictionary D, by replacing $spec_k(\mathbb{G})$ by $spec_D(\mathbb{G})$:

$$spec_D(\mathbb{G}) = \{(\alpha, mult_{\mathbb{G}}(\alpha)) \mid \alpha \in D\}.$$

Here we point out that reversal strings have k-spectra with reversal words and identical multiplicities, and two strings with the same k-spectra have the same length. We indicate by $|\mathbb{G}|$ the length of \mathbb{G} . The count of possible positions for k-mers on the genome \mathbb{G} is $|\mathbb{G}| - k + 1$, which corresponds to the sum of the multiplicities (over $D_k(\mathbb{G})$) present in the k-spectrum.

A k-mer α such that $mult_{\mathbb{G}}(\alpha)$ = 1 in the literature is called a hapax (plural hapaxes) or unique string, while α is called a repeat if it occurs many times, that is, $mult_{\mathbb{G}}(\alpha) > 1$. Then, the k-dictionary of a genome may be decomposed in two disjoint subsets, the set $H_k(\mathbb{G})$ of hapaxes and its complementary set $R_k(\mathbb{G})$ of repeats, by omitting (\mathbb{G}) when the genome is clear from the context.

For example, in the genome abcbbabcaa substrings ab, bc are repeats occurring two times; all the substrings shorter than 2 are repeats; the substring abc is a maximal repeat, because it occurs two times and no longer repeat occurs in the genome; the substrings cb, bb, ba, ca, aa are hapaxes because each of them occurs only once in the genome; all the substrings longer than 3 are hapaxes, moreover cb, bb, ba, ca, aa are also minimal hapaxes, because no hapax shorter than them occurs in the genome.

If a genomic string includes a k-hapax as a substring, for some k, it is a hapax too, while if it is a substring of a k-repeat, for some k, it is a repeat too.

Given a genomic k-spectum, some informational indexes over the genome may be defined [11, 24, 25, 26, 7, 27], which have nice mathematical properties, as in the following. The average multiplicity of k-mers $LX_k(\mathbb{G})$ is called k-lexical index, the maximal repeat length is denoted by $mrl(\mathbb{G})$, the minimal hapax length is denoted by $mrl(\mathbb{G})$. For example, in the genome considered above, we have that mrl(abcbbabcaa) = 3 and mhl(abcbbabcaa) = 2. By definition, any genomic string longer than its mrl is a hapax, and shorter than its mrl is a repeat.

Proposition 1. Maximal repeats over genomes have multiplicity equal or less than 5.

Proof If α is a maximal repeat, none of its extensions αx with $x \in \{A, C, G, T\}$ can occur twice, otherwise α would not be maximal. This means that symbols x after the occurrences of α in the genome have to be different, that is α may occur at most four times in the middle of the genome, and additionally α may occur as a suffix of the genome. In conclusion, a maximal repeat can have at most five possibilities of occurring. \square

An interesting speculation to study genomic strings may be developed around the inclusion of all k long words in the k-dictionary, or some possible absence of such words. To this purpose, over a genome \mathbb{G} , we

define the maximal complete length $mcl(\mathbb{G})$ as the maximal k such that all possible k long strings occur in \mathbb{G}), and the minimal forbidden length $mfl(\mathbb{G})$ as the minimal length k such that at least one k-mer does not belong to the genomic k-dictionary (in the literature this is called k-forbidden word or k-nullomer).

By definition, thus, in any genome, mcl = mfl - 1. Moreover, it holds the following

Proposition 2. In all genomes $mfl \leq mhl + 1$.

Proof Any hapax α is followed over the genome by only one symbol of the alphabet. Therefore, if $|\alpha| = mhl$ then we may find three forbidden words long mhl + 1.

Since $mcl \leq mhl$ (as a consequence of the observations above), any genomic string shorter than the mcl is a repeat. If we call $LG = \lg_4(|\mathbb{G}|)$, the logarithmic length of \mathbb{G} , we have also the following property for the mcl index:

Proposition 3. In any genome \mathbb{G} the inequality mcl < LG holds.

Proof Let n be the length of genome \mathbb{G} . Since, by definition of mcl, we have $4^{mcl} \leq n - k + 1$, then $mcl \leq \lg_4(n - k + 1) < LG$.

A genome \mathbb{G} is a k-hapax genome when its dictionary $D_k(\mathbb{G})$ consists only of hapaxes, it is also a k-complete genome if $D_k(\mathbb{G})$ consists of all 4^k k-mers (in this case the length of \mathbb{G} is $4^k + k - 1$).

The toy genome aaccggttagatctg is a 2-hapax genome, while aaccggttagatctgca is a 2-hapax complete genome. It is complete because all pairs occur and it is hapax because it has the minimum length $17 = 2^4 + 2 - 1$ capable of containing all the possible pairs. However, if we permute the last two symbols the result is not anymore a hapax genome: aaccggttagatctgac.

Given the k-spectrum of a genome \mathbb{G} , the k-entropy $E_k(\mathbb{G})$ is defined as the Shannon entropy of the probability distribution assigning to any k-mer its frequency:

$$E_k(\mathbb{G}) = \sum_{\alpha \in D_k(\mathbb{G})} p(\alpha) \log_2 p(\alpha) \text{ where } p(\alpha) = \frac{mult_{\mathbb{G}}(\alpha)}{|\mathbb{G}| - k + 1}$$

According to the equipartition property of Entropy [34, 7] we may deduce the following result:

Proposition 4. Any k-hapax genome \mathbb{G} of length n has the maximum value of $E_k(\mathbb{G})$ among all genomes of the same length.

We conclude this section with the observation that the notion of k-spectrum can be extended to any genomic distribution, that is, to any variable X associated to a genome and assuming a (finite) set A of values, by considering the number of times any value is assumed by X. In this sense X can be considered an information source, as defined within Shannon theory [34], therefore the informational nature of genomic spectra clearly appears as a powerful feature to analyse and compare genomes.

3. Spectral segmentation of genomes

In this section some notions based on genomic spectra are introduced that shed a new light on the structure of genomes.

Intuitively, we aim at finding long genomic segments constructed by overlap concatenation of k-mers (out of the k-spectrum, then coming with their multiplicity), with an overlapping long k-1. Namely, we look for unary paths (paths of nodes with degree 1) in de Brujin graph (essential tools in genome sequencing [13]) under the constraint given by the multiplicity: each k-mer node may be passed through only the number of times given by the k-mer multiplicity in the k-spectrum.

Two k-mers (α, β) may be k-concatenated if the (k-1)-long suffix of α equals the (k-1)-long prefix of β , that is, there exist γ such that $\alpha = x\gamma$ and $\beta = \gamma y$, with x and y being symbols from the alphabet. The

concatenation results in $\alpha \cdot_k \beta = x \gamma y$, which is the string αy obtained by right elongation of the string α with the symbol y (suffix of β).

In a k-spectrum one may find couples of strings which may be k-concatenated, either according to the way they are k-concatenated over the genome or differently, and this process may be iterated, to assemble either the genome or other different strings. Even under the constraint given by the multiplicity, to use any k-mer only the number of times allowed by its multiplicity, strings that do not occur in the genome may be obtained.

For example, the two different strings: G = aaggccgaagggcacccaa and G' = aagggcacccaaggccgaa (the second being obtained by the first one by swapping the prefix aaggccgaa of G with the suffix aa of G') share the following 2-spectrum:

$$\{(aa,3),(ag,2),(gg,3),(gc,2),(cc,3),(cg,1),(ga,1),(ca,2),(ac,1)\}.$$

These two different strings are obtained as in the following, as a result on the right of \rightarrow , by iterating the 2-concatenation in the sequences reported on the left of \rightarrow , where the same 2-mers occur, with the multiplicities of their common 2-spectrum, but with different orders:

 $aa, ag, gg, gc, cc, cg, ga, aa, ag, gg, gg, gc, ca, ac, cc, cc, ca, aa \rightarrow aaggccgaagggcacccaa$

 $aa, ag, gg, gg, gc, ca, ac, cc, cc, ca, aa, ag, gg, gc, cc, cg, ga, aa \rightarrow aagggcacccaaggccgaa.$

Since the k-spectrum of a genome in general does not identify it in a univocal way, we say k-univocal (in spectrum) a genome \mathbb{G} if no genome different from \mathbb{G} exists that has the same k-spectrum of \mathbb{G} . Of course, a k-univocal genome may always be obtained for suitably high values of k (two trivial cases being the length of the genome itself and $\operatorname{mrl} + 2$, since k-mers and their possible overlapping are all hapaxes). Given a genome \mathbb{G} , it would be interesting to know which is the minimal k for which it is possible a unique reconstruction from the k-spectrum by k-concatenation. The following discussion and toy examples of k-spectra may help to familiarize with these concepts.

Strings $a^3b^2a^5$ and $a^5b^2a^3$ have the same 2-spectrum: $\{(aa,6),(ab,1),(bb,1),(ba,1)\}$, then they are not 2-univocal, but they are 5-univocal. In particular, they are both 5-hapax, with

$$spec_5(a^3b^2a^5) \neq spec_5(a^5b^2a^3)$$

since aaaab occurs only in $a^5b^2a^3$ while baaaa occurs only in $a^3b^2a^5$. However, a k-hapax genome is not necessarily k-univocal, as it may easily seen from the sequences aabcac and acaabc having both the 2-spectrum $\{(aa,1),(ab,1),(bc,1),(ca,1),(ac,1)\}$.

If \mathbb{G} is not k-univocal, by definition there is at least one different genome \mathbb{G}' having the same spectrum of \mathbb{G} . This means that at least some of the k-mers of the spectrum occur in the two genomes \mathbb{G} and \mathbb{G}' with different reciprocal orders. Also, from the examples above, we may deduce that two (different) genomes with the same k-spectrum may be obtained by k-concatenation of the k-mers in a different order.

An interesting question here is: which are the longest arrangements of k-mers having, in both genomes \mathbb{G} and \mathbb{G}' , the same k-mers occurrences with the same order? In an extreme case these arrangements coincide with the k-mers, however in general these portions can be longer than k, as we will see in the following subsection, where spectral maximal segments are defined. It could be interesting to find such common components occurring in genomes having the same k-spectrum.

3.1. Spectral segments

Given a k-spectrum H, we call k-spectral segment an iterated k-concatenation of k-mer copies in H. Being each k-mer present in H with a multiplicity (number of copies), all the k-spectral segments that can be formed have to involve k-mers not more times than the number of their occurrences in H. This process is like having bricks in a limited number of copies, and compose them until they may be used up.

A k-spectral segment may result in a string which is not a substring of any genome having H as a k-spectrum. For example, the string canegattogallogattone has the 4-spectral segment canegattone which is

not a substring occurring in the string. This feature is due to the possibility to have more than one way to elongate a k-mer out of the k-spectrum.

Let us here point out that this phenomenon is present also for k-hapax genomes. As an instance, see genome abcabbaba, with 3-spectrum:

$$\{(abc, 1), (bca, 1), (cab, 1), (abb, 1)(bba, 1), (bab, 1)(aba, 1)\}.$$

In this spectrum where the 3-spectral segment *abbabca* can be obtained by iterating 3-concatenation over the following sequence on the left of \rightarrow :

$$(abb, bba, bab, abc, bca) \rightarrow abbabca$$

however the obtained string (on the right of \rightarrow) is not a substring of the genome *abcabbaba*, which in turn can be obtained by concatenating in a different way the hapax 3-mers of the spectrum:

$$(abc, bca, cab, abb, bba, bab, aba) \rightarrow abcabbaba.$$

These examples (the last one and of genome canegattogallogattone) show an important not so obvious aspect of k-spectral segments.

Of course, if a k-mer is a hapax by definition it may be univocally elongated *over the genome*, but the converse does not hold, because a k-mer may always be elongated by the same symbol (in all its occurrences) without any necessity to be a hapax.

Furthermore, we notice that in the example above we may "maximally" elongate the 3-spectral segment abcabbaba (by 3-concatenation) and obtain another genome abbabcaba (of the same length). This segment is maximal because "we have consumed all the bricks", that is, the sum of multiplicities in the k-spectrum is an upper bound to the number of times we may iterate the k-concatenation.

The capability to obtain k-spectral segments which are not genomic substrings depends on the k value as well. Indeed, the above genome abcabbaba (which is of course also 5-hapax) has 5-spectral segments (such as abbaba) which are all genomic substrings.

Due to the observations and discussion so far, we need to introduce some helpful more specific definitions. In particular, if we include k-mers of the k-spectrum in the definition of k-spectral segments, we say that a k-spectral segment α of H is univocally elongated (on the right) in H if the k-mer suffix of α can be k-concatenated with only one k-mer β in M can be k-concatenated with the k-mer prefix of α . Moreover, α is maximally univocally elongated when, after its last unique elongation, on the right or on the left, α cannot be further univocally elongated in H (on the right or on the left).

We shortly call k-spectral maximal segments the maximally univocally elongated k-spectral segments ("spectral maximal segments" or only "maximal segments" when k-spectrum is clear from the context). They represent genomic portions that are blocks common to all the genomes having the same k-spectrum, a sort of signature associated to a genomic k-spectrum. In the following, experimental results on some human chromosomes are reported along with seven tables, where the term "Maximals" stands for "Maximal segments", which have been specifically computed and counted in number.

The k-spectral maximal segments are the longest arrangements of k-mers having the same k-mers occurrences, with the same order, in all genomes with the same k-spectrum. We say k-segmentation of H the multiset of the k-spectral maximal segments of H (a maximal segment may occur in H with a multiplicity greater than 1). The following two procedures provide the k-segmentation of a k-spectrum H according to different strategies.

Procedure 1 to compute k-segmentations

The procedure starts by assigning to the current string w one k-mer chosen out of H and, at each step, searches in H for the k-mer that uniquely elongates w, on the right, or on the left, by updating H after removing the k-mer occurrence used in the elongation. If more than one possible k-mers of H can be concatenated to both the extremities of the current string, then the concatenation process halts, and the string

obtained so far is produced in output, as a k-spectral maximal segment. Then, the procedure restarts with a similar process with another k-mer of H. The procedure halts when H becomes empty.

In the procedure above it is not essential to establish the choice criterium for the initial string w (the first k-mer in the lexicographic order could be one possibility). Some of the k-spectral maximal segments produced in output during the whole process above can be generated with a multiplicity greater than 1. The multiset of these segments with their respective multiplicities is the k-segmentation of H. Clearly, in the genomes with k-spectrum H, the k-spectral maximal segments of the obtained k-segmentation of H occur with different possible orders that are specific of each genome.

If we change perspective, and assume to start from the genomic sequence itself, we may compute its spectral segmentation in a more efficient way. This is indeed the strategy we have followed by procedure 2, in order to compute the k-segmentations of real genomes: given a genome \mathbb{G} , a length k, and its k-dictionary $D_k(G)$, the k-spectral maximal segments of \mathbb{G} are retrieved by mapping univocally elongated words of $D_k(G)$ at their starting positions in \mathbb{G} .

Procedure 2 to compute k-segmentations

A k-mer $\alpha \in D_k(\mathbb{G})$ is univocally elongated in G if $|\{\beta \in D_k(G) : \alpha[2...k] = \beta[1...k-1]\}| = 1$. A Boolean array A, such that $|A| = |\mathbb{G}|$, is initialized to be false in every position. Let $pos(\alpha, \mathbb{G})$ to be the set of starting positions in G where α occurs, namely $pos(\alpha, \mathbb{G}) = \{i : \mathbb{G}[i, ..., i+k-1] = \alpha\}$. For each $\alpha \in D_k(G)$ such that α is univocally elongated in \mathbb{G} , the algorithm sets as true all the positions where an occurrence of α starts in \mathbb{G} , namely A[i] = true for each $i \in pos(\alpha, G)$. Then, the k-segmentation is retrieved by scanning for consecutive runs of true values in A, and a k-spectral maximal segment is a substring $\mathbb{G}[i...j]$ of \mathbb{G} such that $A[l] = true \ \forall l : i \leq l \leq j$.

4. Computational results

Table 1 reports the data for human chromosome 1 up to k=40. It appears remarkable that already in the range [28-40] for the k values, the genomic coverage of the k-segmentation is almost total (over positions where there is no N), and almost all the k-mers present in the chromosome are involved by k-spectral maximal segments (see the normalized cardinality of set U_k), that is, they have the property to be univocally elongated in the k-spectrum. Essentially, we see that for k=27 human chromosome 1 may be covered by 4 millions of relatively short (80 bp long) spectral maximal segments, which have of course a high value of multiplicity in the segmentation, whereas for k=44 about half of the spectral maximal segments (1.985.225) of average length 157 cover the chromosome. We observe that he total coverage is reached for k=216 with 47.237 maximal segments of average length 5.093 (and maximal length 1.978.600). In this context it is worth to notice that the unique reconstruction of the genome is proved to be possible for k=mrl+2, which corresponds to the value 29.265 for human chromosome 1.

A k-segmentation was computed also for Saccharomyces cerevisiae genome (around 12 Millions bp with 16 chromosomes) where no N characters are present. In all chromosomes, we have verified that for k = mrl + 2, only one spectral segment was found covering the whole chromosome, while for k close to 60, in almost all the cases, spectral segments are less than 100. Table 2 reports the data for chromosome IV (which is the longest one) of Saccharomyces cerevisiae (with mrl equal to 3.573) up to k = 40. We remarkably notice that k = 21 is sufficient to entirely cover the genomic sequence (1.531.933 long) by a segmentation of 1.584 maximal segments (in average shorter than one thousand bp), and that k = 16 corresponds to a 16-spectral segmentation with a very good coverage, which involves most of the 16-mers present in the genome, into only 8.958 maximal segments having an average length equal to 185,8.

The k-segmentations were also computed, by the previous algorithm, for chromosomes 2, 10, 22, X, Y of Human genome (Hg 38), for k-values ranging from 10 to more than 500, where segments are measured (in number and lengths), Tables 3, 4, 5, 6, 7 report the obtained values up to k = 40. The results are uniform for all considered chromosomes.

The algorithm of Procedure 2 was used to compute the data reported in all tables, where the minimal values of k indicate the length of the minimal k-mer which could be univocally elongated in the k-spectrum.

\overline{k}	D_k	U_k	U_k/D_k	Coverage	Maximals	AvgL	MaxL
10	1.045.927	38	0,0000	2,25615E-06	52	11,00	11
11	4.046.584	13.683	0,0034	0,000724155	15.955	12,08	17
12	13.963.535	313.027	0,0224	0,014264785	318.364	13,21	24
13	40.048.761	3.259.722	0,0814	0,099454848	2.645.335	14,44	39
14	88.918.637	15.853.072	$0,\!1783$	$0,\!417332951$	10.984.163	15,66	54
15	137.433.303	59.962.772	$0,\!4363$	0,843980167	33.214.551	17,01	103
16	166.201.314	119.093.792	0,7166	0,921733136	35.952.622	19,59	165
17	179.785.859	157.738.137	$0,\!8774$	0,939876978	22.706.197	$24,\!43$	375
18	186.272.565	176.141.059	0,9456	0,949814751	12.897.425	$32,\!53$	883
19	189.945.022	184.536.009	0,9715	0,957262414	8.346.098	$42,\!45$	1.677
20	192.510.989	188.930.042	0,9814	0,963687855	6.456.253	50,99	3.903
21	194.595.977	191.777.370	0,9855	0,969077973	5.624.104	57,07	6.407
22	196.426.475	193.995.985	0,9876	0,973513497	5.165.406	$61,\!69$	14.233
23	198.093.250	195.897.366	0,9889	0,977337135	4.864.712	$65,\!52$	14.887
24	199.637.221	197.618.157	0,9899	0,980668464	4.623.956	69,09	16.450
25	201.077.240	199.212.471	0,9907	0,98370779	4.424.312	$72,\!46$	16.511
26	202.425.001	200.690.961	0,9914	0,986368122	4.248.537	75,75	21.393
27	203.689.655	202.070.404	0,9921	0,98849874	4.072.258	79,21	23.140
28	204.878.861	203.365.399	0,9926	0,990262556	3.907.943	82,71	23.289
29	206.000.589	204.579.712	0,9931	0,99181704	3.746.919	$86,\!35$	25.316
30	207.060.369	205.720.228	0,9935	0,993210426	3.601.305	89,95	25.317
31	208.061.649	206.793.925	0,9939	0,994265107	3.457.401	93,71	26.342
32	209.008.844	207.810.544	0,9943	0,995187959	3.315.966	$97,\!65$	32.156
33	209.904.984	208.781.444	0,9946	0,995923721	3.169.314	101,95	32.159
34	210.752.552	209.694.818	0,9950	0,996654705	3.046.401	105,98	32.162
35	211.553.055	210.557.940	0,9953	0,997153583	2.918.996	$110,\!36$	37.175
36	212.308.870	211.371.979	0,9956	0,997652279	2.795.085	114,94	37.176
37	213.022.579	212.138.601	0,9959	0,997991136	2.680.028	$119,\!56$	37.179
38	213.696.481	212.864.835	0,9961	0,998288267	2.565.984	$124,\!45$	37.180
39	214.332.908	213.548.776	0,9963	0,998509309	2.457.660	$129,\!48$	37.359
40	214.934.436	214.196.288	0,9966	0,99873295	2.350.398	134,82	37.372

Table 1: Values of spectral segmentation for Hg38 Chr 1, computed by Procedure 2 and using the IGTools platform [5]. From the left: k-column gives the length of k-mers; D_k -column provides the cardinality of D_k (number of different k-mers in the chromosome), U_k -column reports the number of k-mers occurring in the k-spectral maximal segments; Coverage(%)-column the percentages of positions of Hg38 Chr 1 covered by the k-spectral segmentation; Maximals-column shows the number of k-spectral maximal segments, AvgL-column and MaxL-column the average and maximum lengths of k-spectral maximal segments. The length of the chromosome 1 (including the 18.475.410 N characters which were ignored from the collection of k-mers) is 248.956.422 bp.

\overline{k}	D_k	U_k	U_k/D_k	Coverage	Maximals	AvgL	MaxL
8	65337	617	$0,\!45915$	0,3133	6	9	9
9	239715	33262	0,006695	0,010416	1837	10,037	12
10	630466	279271	0,0920684	0,2700	53070	11,2373	22
11	1066662	776349	$0,\!352781$	0,8457	231242	12,7622	38
12	1328523	1177684	$0,\!672753$	0,9934	288803	$15,\!2555$	129
13	1434547	1363931	0,875117	0,9994	166821	20,8055	148
14	1470635	1430564	0,958397	0,9998	66411	35,8881	439
15	1482371	1452182	0,986831	0,9999	23344	79,4844	876
16	1486575	1459635	0,99548	0,9999	8958	185,843	2440
17	1488399	1462631	0,998181	0,9999	4356	$367,\!469$	6536
18	1489415	1464150	0,999085	0,9999	2718	580,377	11110
19	1490140	1465161	0,999397	0,9999	2061	761,024	25903
20	1490735	1465965	0,999531	0,9999	1785	876,975	65133
21	1491246	1466635	0,999597	1	1584	$986,\!89$	69154
22	1491694	1467215	0,999663	1	1421	$1098,\!87$	69155
23	1492103	1467741	0,999694	1	1307	$1193,\!91$	85413
24	1492492	1468250	0,999705	1	1213	1285,75	85414
25	1492849	1468714	0,99974	1	1118	$1394,\!09$	85417
26	1493188	1469161	0,999752	1	1059	$1471,\!44$	85420
27	1493510	1469594	0,999763	1	1014	$1536,\!66$	85455
28	1493813	1470004	0,999783	1	967	1611,12	95170
29	1494104	1470401	0,999793	1	925	$1684,\!07$	118445
30	1494380	1470780	0,999803	1	877	1775,72	118446
31	1494639	1471130	0,999813	1	829	$1877,\!86$	118447
32	1494885	1471461	0,999829	1	790	1970,09	118448
33	1495122	1471781	0,999834	1	767	$2029,\!24$	118449
34	1495349	1472088	0,999843	1	729	$2134,\!35$	146444
35	1495571	1472389	0,999846	1	711	$2188,\!55$	146499
36	1495786	1472682	0,999849	1	689	$2258,\!35$	154915
37	1495992	1472961	0,99986	1	653	2381,92	154916
38	1496194	1473234	0,999866	1	639	$2434,\!32$	154917
39	1496396	1473507	0,999859	1	639	$2435,\!32$	154918
40	1496593	1473775	0,999862	1	617	2521,8	154919

Table 2: Values of spectral segmentation for Saccharomyces cerevisiae Chr IV, computed by Procedure 2 and using the IGTools platform [5]. Columns are as in Table 1. The length of the chromosome (where no N characters appear) is 1531933 bp.

k	D_k	U_k	U_k/D_k	Coverage	Maximals	AvgL	MaxL
10	1.045.356	62	0,0001	0,000003613	79	11	11
11	4.037.486	15.091	0,0037	0,000815795	17.279	12	16
12	13.970.306	308.947	0,0221	$0,\!013846816$	301.339	13	25
13	40.326.969	3.210.596	0,0796	0,097668722	2.559.598	14	38
14	91.475.583	15.421.454	$0,\!1686$	0,399774431	10.398.399	16	54
15	145.002.519	59.930.671	0,4133	0,846513054	33.438.889	17	109
16	178.168.268	123.940.839	0,6956	0,934553290	38.598.600	19	188
17	194.126.835	168.126.622	0,8661	0,951892994	25.112.141	24	338
18	201.601.752	189.796.905	0,9414	0,960309510	14.035.702	32	829
19	205.618.151	199.614.013	0,9708	0,966795324	8.619.987	43	1.778
20	208.273.542	204.520.815	0,9820	0,971930785	6.328.018	54	3.767
21	210.348.652	207.521.297	0,9866	0,976231049	5.322.938	62	6.466
22	212.126.808	209.757.547	0,9888	0,980012274	4.799.273	67	11.899
23	213.720.398	211.619.396	0,9902	0,983189018	4.464.885	72	14.695
24	215.179.675	213.269.242	0,9911	0,985755501	4.202.799	77	17.310
25	216.525.819	214.780.861	0,9919	0,987908009	3.980.457	81	21.586
26	217.774.255	216.168.103	0,9926	0,989823405	3.790.270	85	31.139
27	218.936.692	217.448.199	0,9932	0,991326612	3.606.846	89	31.144
28	220.022.185	218.640.791	0,9937	0,992583745	3.432.766	94	31.158
29	221.039.081	219.751.927	0,9942	0,993689170	3.272.001	98	31.159
30	221.993.683	220.786.995	0,9946	0,994613704	3.126.091	103	35.072
31	222.890.490	221.756.576	0,9949	0,995382290	2.984.030	108	35.078
32	223.733.892	222.668.355	0,9952	0,996121593	2.844.942	113	35.079
33	224.527.803	223.532.397	0,9956	0,996735362	2.713.682	118	35.519
34	225.274.155	224.343.871	0,9959	0,997212601	2.594.990	123	35.520
35	225.976.292	225.103.137	0,9961	0,997588970	2.477.302	128	35.521
36	226.635.933	225.818.025	0,9964	0,997944325	2.369.341	134	35.848
37	227.256.084	226.488.028	0,9966	0,998234288	2.267.537	140	35.860
38	227.839.791	227.119.333	0,9968	0,998574569	2.168.482	145	35.861
39	228.389.481	227.711.681	0,9970	0,998894409	2.071.522	152	35.862
40	228.906.606	228.271.367	0,9972	0,999011699	1.976.489	158	38.236

Table 3: Values of spectral segmentation for Hg38 Chr $\bf 2$

\overline{k}	D_k	U_k	U_k/D_k	Coverage	Maximals	AvgL	MaxL
10	1.041.093	286	0,0000	0,000030241	367	11	12
11	3.936.809	25.449	0,0023	0,002252922	27.322	12	18
12	12.817.030	480.917	0,0368	0,036765759	462.859	13	31
13	34.145.871	3.580.802	0,1749	0,174926481	2.628.021	15	36
14	65.646.514	17.099.963	0,6659	0,665857037	11.413.628	16	55
15	90.151.885	50.398.126	0,9116	0,911648152	22.403.605	17	121
16	102.828.231	82.036.963	0,9472	0,947221269	18.212.617	21	198
17	108.607.958	99.323.101	0,9590	0,959014141	10.462.638	27	449
18	111.483.717	107.084.861	0,9669	0,966874359	6.111.992	36	1.546
19	113.244.161	110.709.583	0,9728	0,972821263	4.250.781	46	2.204
20	114.553.598	112.755.264	0,9784	0,978430121	3.489.199	54	4.035
21	115.654.605	114.180.258	0,9824	0,982428689	3.125.739	59	10.768
22	116.633.603	115.341.284	0,9858	0,985826249	2.918.936	63	14.196
23	117.528.956	116.353.857	0,9885	0,988479620	2.762.778	67	18.967
24	118.357.515	117.277.400	0,9908	0,990840418	2.634.631	70	19.354
25	119.126.857	118.133.609	0,9928	0,992771187	2.519.577	74	20.823
26	119.843.613	118.923.757	0,9943	0,994315060	2.420.266	77	20.825
27	120.512.866	119.658.553	0,9955	0,995469199	2.318.595	81	25.320
28	121.139.473	120.344.058	0,9965	0,996514958	2.223.277	84	25.322
29	121.726.829	120.985.037	0,9973	0,997292661	2.131.640	88	33.768
30	122.278.403	121.582.101	0,9980	0,997975957	2.049.112	92	33.771
31	122.796.035	122.142.297	0,9985	0,998514103	1.962.984	96	33.773
32	123.282.548	122.667.660	0,9989	0,998881107	1.886.814	99	33.775
33	123.739.101	123.167.264	0,9992	0,999177739	1.806.606	104	33.777
34	124.166.975	123.634.195	0,9994	0,999430442	1.730.945	108	33.779
35	124.568.421	124.070.103	0,9996	0,999605749	1.660.329	112	33.781
36	124.944.815	124.478.890	0,9997	0,999722136	1.590.536	117	33.783
37	125.297.872	124.860.960	0,9998	0,999798684	1.524.071	122	36.729
38	125.629.183	125.221.245	0,9999	0,999872643	1.456.539	127	38.370
39	125.940.266	125.557.286	0,9999	0,999924443	1.398.753	132	60.547
40	126.231.729	125.873.920	1,0000	0,999959411	1.341.799	137	60.549

Table 4: Values of spectral segmentation for Hg38 Chr 10

1		T 7	11 /D	(01)	11 . 1	A T	M T
k	D_k	U_k	U_k/D_k	Coverage $(\%)$	Maximals	AvgL	MaxL
9	261.825	5	0,0000	0,000001788	7	10	10
10	1.019.673	1.999	0,0020	0,000756797	2.756	11	14
11	3.506.660	79.689	0,0227	0,023291501	86.240	12	20
12	9.548.425	866.147	0,0907	0,172177385	735.492	13	29
13	18.068.493	4.891.914	$0,\!2707$	0,650453372	3.467.102	15	46
14	24.506.912	14.053.286	$0,\!5734$	0,885121409	6.352.781	16	82
15	27.801.432	22.415.019	0,8063	0,922426014	5.120.141	20	141
16	29.318.528	26.906.274	0,9177	0,938427867	3.034.727	26	314
17	30.110.958	28.924.250	0,9606	0,949208214	1.883.877	34	668
18	30.636.017	29.900.266	0,9760	0,958403696	1.419.137	41	1.685
19	31.056.553	30.492.368	0,9818	0,965737956	1.236.907	46	3.585
20	31.426.643	30.940.639	0,9845	0,971911536	1.149.500	49	6.485
21	31.766.989	31.324.505	0,9861	0,976692104	1.092.487	52	9.854
22	32.085.192	31.673.457	0,9872	0,980757296	1.050.098	55	14.755
23	32.385.495	31.997.039	0,9880	0,984112984	1.015.327	57	17.495
24	32.669.603	32.304.683	0,9888	0,987362364	980.900	60	19.464
25	32.938.199	32.596.243	0,9896	0,990175685	946.474	62	21.031
26	33.192.010	32.871.397	0,9903	0,992043162	912.734	65	22.724
27	33.431.987	33.130.689	0,9910	0,993624555	878.002	68	29.616
28	33.659.147	33.375.097	0,9916	0,994783065	845.236	71	29.617
29	33.874.638	33.605.890	0,9921	0,995783148	813.740	73	29.618
30	34.078.824	33.824.614	0,9925	0,996856749	785.507	76	33.620
31	34.272.239	34.030.192	0,9929	0,997512652	758.686	79	33.621
32	34.454.766	34.226.804	0,9934	0,997979738	728.923	82	33.626
33	34.626.863	34.413.820	0,9938	0,998461074	697.974	86	33.784
34	34.789.125	34.589.584	0,9943	0,998695498	670.865	89	33.830
35	34.942.372	34.754.380	0,9946	0,998919529	645.406	92	33.831
36	35.086.703	34.910.137	0,9950	0,999043534	618.723	96	33.836
37	35.222.644	35.056.110	0,9953	0,999186563	595.130	100	35.249
38	35.350.607	35.194.803	0,9956	0,999260210	571.318	103	36.970
39	35.471.346	35.324.799	0,9959	0,999324000	549.538	107	36.976
40	35.585.086	35.447.664	0,9961	0,999460058	528.884	111	36.985

Table 5: Values of spectral segmentation for Hg38 Chr 22

\overline{k}	D_k	U_k	U_k/D_k	Coverage (%)	Maximals	AvgL	MaxL
10	1.037.774	491	0,0005	0,000043888	623	11	13
11	3.889.504	31.867	0,0082	0,002465811	35.538	12	18
12	12.504.932	519.962	0,0416	0,035696493	530.923	13	26
13	33.746.525	3.471.711	0,1029	$0,\!145729870$	2.581.493	15	39
14	67.004.374	16.083.058	0,2400	0,603250867	11.410.193	16	70
15	94.629.309	50.176.817	0,5302	0,887882178	24.597.625	17	121
16	110.045.379	85.018.232	0,7726	0,937308793	22.040.843	20	451
17	117.654.131	105.593.254	0,8975	0,951868860	14.160.125	25	896
18	121.730.602	115.579.606	0,9495	0,960595541	9.168.860	32	1.641
19	124.355.508	120.620.193	0,9700	0,967176250	6.814.620	38	2.347
20	126.354.205	123.628.203	0,9784	0,972524051	5.744.645	43	4.280
21	128.036.423	125.792.642	0,9825	0,976775856	5.193.767	47	7.205
22	129.518.262	127.571.616	0,9850	0,980340251	4.803.598	50	9.871
23	130.854.734	129.111.191	0,9867	0,983110260	4.536.017	53	14.486
24	132.074.054	130.491.374	0,9880	0,985695050	4.298.488	56	19.626
25	133.189.081	131.756.797	0,9892	0,987901502	4.072.902	59	20.102
26	134.213.903	132.905.141	0,9902	0,989864328	3.880.410	62	21.965
27	135.159.044	133.956.732	0,9911	0,991405462	3.702.602	65	21.966
28	136.032.999	134.928.613	0,9919	0,992785602	3.529.928	68	21.968
29	136.844.664	135.822.987	0,9925	0,993861409	3.376.285	71	21.970
30	137.598.873	136.651.222	0,9931	0,994778932	3.233.787	74	22.045
31	138.300.893	137.420.171	0,9936	0,995770642	3.101.559	78	22.046
32	138.955.736	138.134.801	0,9941	0,996451125	2.971.323	81	30.578
33	139.566.748	138.806.672	0,9946	0,996995920	2.850.829	84	30.579
34	140.137.310	139.430.458	0,9950	0,997497434	2.746.491	87	30.584
35	140.670.461	140.011.539	0,9953	0,997900325	2.643.366	90	39.481
36	141.169.081	140.554.194	0,9956	0,998246067	2.553.300	94	39.513
37	141.635.366	141.061.342	0,9959	0,998532329	2.462.499	97	39.514
38	142.072.342	141.536.092	0,9962	0,998691768	2.375.435	100	39.515
39	142.482.391	141.980.427	0,9965	0,998991672	2.294.023	104	39.516
40	142.867.125	142.397.483	0,9967	0,999122853	2.222.933	107	39.561

Table 6: Values of spectral segmentation for Hg38 Chr $\bf X$

\overline{k}	D	U_k	U_k/D_k	Canamaga (%)	Maximals	AnaI	MaxL
$\frac{\kappa}{9}$	$D_k = 258.851$	0_k 184	0.0007	Coverage $(\%)$ 0,000094794	252	AvgL 10	MaxL 12
10	952.028	9.178	0,0007	0.00094794 0.004327837	11.235	10	12 17
11	2.950.639	161.600	0,0090 $0,0548$	0.059458998	171.130	12	23
$\frac{11}{12}$	7.349.099	925.005	0,0348 0,1259	0,059458998 $0,260281537$	772.716	14	23 38
13	12.604.575		0,1259 $0,3452$,		15	69
$\frac{13}{14}$	16.150.413	4.350.557		0,797859424	3.258.408	15 17	
$\frac{14}{15}$	10.130.413 $17.999.177$	$10.425.123 \\ 15.058.625$	0,6455 $0,8366$	0,942926688 0,967826477	4.617.874 $3.437.514$	20	$\frac{115}{217}$
				,			
16	18.975.234	17.499.482	0,9222	0,977689796	2.223.413	25	713
17	19.582.524	18.720.914	0,9560	0,983727378	1.584.051	31	1.591
18	20.032.378	19.425.139	0,9697	0,987848742	1.278.129	36	2.341
19	20.404.478	19.918.053	0,9762	0,990960416	1.116.113	40	4.899
20	20.728.314	20.311.145	0,9799	0,993522138	1.010.214	43	8.835
21	21.017.072	20.649.604	0,9825	0,995278940	924.063	47	8.836
22	21.276.934	20.947.952	0,9845	0,996511003	853.382	50	14.256
23	21.512.015	21.215.272	0,9862	0,997469662	788.915	54	14.290
24	21.725.180	21.457.857	0,9877	0,998096577	728.442	58	14.294
25	21.918.376	21.677.034	0,9890	0,998614237	675.225	62	14.295
26	22.094.064	21.876.610	0,9902	0,999058832	626.810	66	14.296
27	22.254.431	22.056.528	0,9911	0,999357298	584.288	70	14.323
28	22.400.981	22.221.033	0,9920	0,999647057	544.951	74	22.075
29	22.535.244	22.371.147	0,9927	0,999741132	507.782	79	24.908
30	22.658.352	22.507.948	0,9934	0,999864471	475.866	83	25.084
31	22.771.307	22.634.213	0,9940	0,999906985	444.811	88	25.094
32	22.875.207	22.749.604	0,9945	0,999936665	416.799	93	25.096
33	22.971.055	22.855.636	0,9950	0,999952716	390.904	99	25.101
34	23.059.696	22.953.556	0,9954	0,999964679	366.515	104	25.102
35	23.141.808	23.043.565	0,9958	0,999973765	344.966	110	28.304
36	23.217.796	23.127.687	0,9961	0,999980163	324.812	115	28.305
37	23.288.437	23.204.700	0,9964	0,999986788	306.251	121	28.306
38	23.354.165	23.276.355	0,9967	0,999990574	287.691	128	29.584
39	23.415.300	23.343.464	0,9969	0,999993261	271.426	135	29.585
40	23.472.435	23.405.683	0,9972	0,999995722	256.379	141	29.586

Table 7: Values of spectral segmentation for Hg38 Chr ${\bf Y}$

These computational experiments were implemented within the platform IGTools [5], specifically designed to develop informational analyses of genomes. Data were globally obtained in less than 10 hours running computations. The algorithm was implemented using a suffix array representation of genomes [1] with a complexity linear with respect the size of genomes.

5. Conclusions

The notion of distribution and of information source has an immense scientific value and appears everywhere, in an enormous number of different contexts. This ubiquity and variety is a sort of evidence of its fundamental role. In the paper we studied a special case of genomic distribution, based on substring occurrences in genomes. The notions of k-spectrum, spectral segments and segmentations, are showed to be strictly related to the internal organisation of genomes, through experimental results on human chromosomes. Only some computational experiments on real genomes are here reported, however many other related ones may be conceived (for example to study the correlation among the informational indexes discussed in this paper, including those of k-spectral segmentations), which could assess the real biological importance of the concepts analysed. Specific experiments could suggest further clues on the deep nature of genomic sequences, and on the internal principles that rule their coherence, their plasticity, their equilibria and their instability, by pushing them toward continuous transformations.

The topics developed in the paper are mainly of theoretical nature, and the genomic terminology is used for the motivation underlying the investigated arguments. As an example we mention the algorithm of Procedure 1, which may become more interesting in the case that some efficient technology providing directly k-spectra, independently from sequencing procedures, should be available in the future.

References

- [1] Abouelhoda, M. I., Kurtz, S. & Ohlebusch, E. Replacing suffix trees with enhanced suffix arrays. *J Discrete Algorithms*, 2, 53–86 (2004).
- [2] Acharya J., Das H., Milenkovic O., Orlitsky A., Pan S., On Reconstructing a String from its Substring Compositions, *IEEE International Symposium on Information Theory*, pp. 1238-1242, (2010).
- [3] Acharya J., Das H., Milenkovic O., Orlitsky A., Pan S., String reconstruction from substring compositions, SIAM Journal on Discrete Mathematics 29(3): 1340-1371, (2015).
- [4] Berstel, J., Karhumaki, J.: Combinatorics on Words A Tutorial. http://www-igm.univ-mlv.fr/~berstel/Articles/2003TutorialCoWdec03.pdf (2003).
- [5] Bonnici, V., Manca, V.: Infogenomics tools: A computational suite for informational analysis of genomes. J. Bioinformatics & Proteomics Review 1, 8–14 (2015).
- [6] Bonnici, V., Manca, V.: Recurrence distance distributions in computational genomics. American J. of Bioinformatics & Computational Biology, 3, 5–23 (2015).
- [7] Bonnici, V., Manca, V.: Informational laws of genome structures. *Scientific Reports*, 6, 28840, http://www.nature.com/articles/srep28840 (2016).
- [8] Bonnici, V., Manca, V.: An informational test for random finite strings. Entropy, 20(12), 934 (2018).
- [9] Brijder, R., Daley, M., Harju, T., Jonoska, N., Petre, I., and Rozenberg, G.: The computational nature of gene assembly in ciliates, in *Handbook of natural computing*, Vol. 3, G. Rozenberg, T. Bäck, J.N. Kok (eds), Springer, 1233–1281 (2012).
- [10] J. Burns, D. Kukushkin, K. Lindblad, X. Chen, N. Jonoska, L. F. Landweber, A database of ciliate genome rearrangements, *Nucleic Acids Res.* 44: 703–709, (2016).
- [11] Castellini, A., Franco, G. & Manca, V.: A dictionary based informational genome analysis. BMC genomics 13, 485 (2012).
- [12] Castellini, A., Franco, G. & Milanese, A.: A genome analysis based on repeat sharing gene networks, *Natural Computing* 14(3), pp 403–420, (2015).
- [13] Compeau, P. E. C.; Pevzner, P. A., Glenn Tesler, G.: Why are de Bruijn graphs useful for genome assembly? Nat Biotechnol., 29(11): 987–991 (2011).
- [14] Dudik M., Schulman L.: Reconstruction from Subsequences, Journal of Combinatorial Theory, 103(2): 337-348, (2003).
- [15] Franco G. (2014) Perspectives in Computational Genome Analysis. In: Jonoska N., Saito M. (eds) Discrete and Topological Models in Molecular Biology. Natural Computing Series. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-40193-0
- [16] Franco, G. & Jonoska, N.: Forbidding enforcing conditions in DNA self-assembly of graphs, book chapter in J. Chen, N. Jonoska, G. Rozenberg Eds: Nanotechnology, Science and Computation, Springer, pp 105 118, (2006).
- [17] Franco, G. & Milanese, A.: An investigation on genomic repeats, Proceedings of 9th Conference on Computability in Europe, CiE 2013, Milano, July 1-5, 2013, in The Nature of Computation. Logic, Algorithms, Applications, LNCS 7921 Springer Verlag Germany, pp 149–160, (2013).

- [18] Franco G., Manca V.: Decoding Genomic Information. In: Stepney S., Rasmussen S., Amos M. (eds) Computational Matter. Natural Computing Series. Springer, Cham. (2018) https://doi.org/10.1007/978-3-319-65826-1
- [19] Gabrys R, Milenkovic O.: Unique Reconstruction of Coded Strings from Multiset Substring Spectra, *IEEE Trans. Inf. Theory* 65(12): 7682-7696, IEEE press (2019).
- [20] Holden N., Pemantle R., and Peres Y.: Subpolynomial trace reconstruction for random strings and arbitrary deletion probability, Proceedings of the 31st Conference On Learning Theory, PMLR 75:1799-1840, (2018).
- [21] Kiah H. M., Puleo G. J., Milenkovic O.: Codes for DNA Sequence Profiles, *IEEE Transactions on Information Theory*, 62(6): 3125-3146, (2016).
- [22] Levenshtein V.: Efficient Reconstruction of Sequences from their Subsequences or Supersequences, Journal of Combinatorial Theory, Series A 93(2): 310-332, (2001).
- [23] M. Lan, C. L. Tan, J. Su, and Y. Lu. Supervised and traditional term weighting methods for automatic text categorization. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 31(4):721–735, (2009).
- [24] Manca, V.: Infobiotics. Information in biotic systems. Springer (2013).
- [25] Manca, V.: Infogenomics, genomes as information sources. Chapter 21, 317–324. Elsevier, Morgan Kauffman (2016).
- [26] Manca, V.: Information theory in genome analysis. In Membrane Computing, LNCS 9504, 3–18. Springer (2016).
- [27] Manca, V.: The Principles of Informational Genomics. Theoretical Computer Science (2017)
- [28] Manca V.: A Marvelous Accident: The Birth of Life. J Proteomics Bioinform 11: 135-137 (2018).
- [29] Manuch, J.: Characterization of a word by its subwords, Proc. of DLT, 210-219, World Scientific, (2000).
- [30] Margaritis D., Skiena S.: Reconstructing Strings from Substrings in Rounds, Foundations of Computer Science, Proceedings of IEEE 36th Annual Foundations of Computer Science, Milwaukee, WI, USA, 1995, pp. 613-620, doi: 10.1109/SFCS.1995.492591.
- [31] Marcovich S., Yaakobi E., Reconstruction of Strings from their Substrings Spectrum, 2020 IEEE International Symposium on Information Theory (ISIT), Los Angeles, CA, USA, 2020, pp. 658-663, doi: 10.1109/ISIT44484.2020.9174113.
- [32] A. S. Motahari, G. Bresler and D. Tse, Information theory of DNA shotgun sequencing, IEEE Transactions on Information Theory, 59(10: pp. 6273-6289, (2013).
- [33] Scott A.: Reconstructing sequences, Discrete Mathematics, 175(1-3): 231–238, (1997)
- [34] Shannon, C. E. A mathematical theory of communication. Bell Sys Tech J 27, 623–656 (1948).
- [35] Sievers, A., Wenz, F., Hausmann, M., Hildenbrand, G.: Conservation of k-mer Composition and Correlation Contribution between Introns and Intergenic Regions of Animalia Genomes, *Genes*, 9(10): 482, (2018).
- [36] Simões R., Wolf I., Correa B., Valente G.: Uncovering patterns of the evolution of genomic sequence entropy and complexity, Molecular Genetics and Genomics, October 2020, https://doi.org/10.1007/s00438-020-01729-y
- [37] Yang, Z., Li, H., Jia, Y. et al. Intrinsic laws of k-mer spectra of genome sequences and evolution mechanism of genomes. BMC Evol Biol 20, 157 (2020). https://doi.org/10.1186/s12862-020-01723-3