The Kinetic and Hydrodynamic Bohm Criterions for Plasma Sheath Formation

Masahiro Suzuki¹ and Masahiro Takayama²

¹Department of Computer Science and Engineering, Nagoya Institute of Technology, Gokiso-cho, Showa-ku, Nagoya, 466-8555, Japan

> ²Department of Mathematics, Keio University, Hiyoshi, Kohoku-ku, Yokohama, 223-8522, Japan

> > March 16, 2022

Abstract

The purpose of this paper is to mathematically investigate the formation of a plasma sheath, and to analyze the Bohm criterions which are required for the formation. Bohm derived originally the (hydrodynamic) Bohm criterion from the Euler–Poisson system. Boyd and Thompson proposed the (kinetic) Bohm criterion from a kinetic point of view, and then Riemann derived it from the Vlasov–Poisson system. In this paper, we prove the solvability of boundary value problems of the Vlasov–Poisson system. On the process, we see that the kinetic Bohm criterion is a necessary condition for the solvability. The argument gives a simpler derivation of the criterion. Furthermore, the hydrodynamic criterion can be derived from the kinetic criterion. It is of great interest to find the relation between the solutions of the Vlasov–Poisson systems. To clarify the relation, we also study the delta mass limit of solutions of the Vlasov–Poisson system.

Keywords: Vlasov–Poisson system, Euler–Poisson system, Boundary value problem, Delta mass limit.

2020 Mathematics Subject Classification: 35A01, 35M32, 35Q35, 76X05

1 Introduction

The purpose of this paper is to mathematically investigate the formation of a plasma boundary layer, called as a *sheath*, near the surface of materials immersed in a plasma, and to analyze the Bohm criterions which are required for the formation. The sheath appears when a material is surrounded by a plasma and the plasma contacts with its surface. Because the thermal velocities of electrons are much higher than those of ions, more electrons tend to hit the material compared with ions. This makes the material negatively charged with respect to the surrounding plasma. Then the material with a negative potential attracts and accelerates ions toward the surface, while repelling electrons away from it. Eventually, there appears a non-neutral potential region near the surface, where a nontrivial equilibrium of the densities is achieved. This non-neutral region is referred as to the sheath. For more details of physicality of the sheath development, we refer the reader to [5, 7, 25, 26, 32, 33].

For the formation of sheath, Langmuir [25] observed that positive ions must enter the sheath region with a sufficiently large flow velocity. Using the Euler–Poisson system (see (1.5) below), Bohm [5] proposed the original *Bohm criterion* which states that the flow velocity of positive ions at the plasma edge must exceed the ion acoustic speed. In this paper, we call the criterion *a hydrodynamic Bohm criterion*. Nowadays there are many mathematical results which investigated the sheath formation by using the Euler–Poisson system. The studies [1, 2, 28, 40–42] established the existence and stability of stationary solutions assuming the hydrodynamic Bohm criterion. These results validated mathematically the criterion. From different perspectives to the those results', the sheath formation was discussed by considering the quasi-neutral limit as letting the Debye length in the Euler–Poisson system tend to zero [10, 11, 21–23]. Furthermore, Feldman–Ha–Slemrod [9] studied the sheath formation adopting a certain hydrodynamic model which describes the dynamics of an interface between the plasma and sheath (see also [34]).

From a kinetic point of view, Boyd–Thompson [6] proposed a Bohm criterion. After that Riemann [32] derived it from the Vlasov–Poisson system (see (1.1) below). We call it *a kinetic Bohm criterion* in this paper. There is no mathematical result which investigates the sheath formations by using the Vlasov–Poisson system with rigorous proofs. One of our goals is to prove the solvability of boundary value problems of the Vlasov–Poisson system under the kinetic Bohm criterion. On the process, we see that the criterion is a necessary condition for the solvability. The argument gives a simpler derivation of the criterion.

It is worth pointing out that the hydrodynamic Bohm criterion can be derived from the kinetic Bohm criterion. We are also interested in finding the relation between the solutions of the Vlasov–Poisson and Euler–Poisson systems. To clarify the relation, we study some limit of solutions of the Vlasov–Poisson system.

After a suitable nondimensionalization, the stationary Vlasov-Poisson system is written by

$$\xi_1 \partial_x f + \partial_x \phi \partial_{\xi_1} f = 0, \quad x > 0, \ \xi \in \mathbb{R}^3,$$
(1.1a)

$$\partial_{xx}\phi = \int_{\mathbb{R}^3} f d\xi - n_e(\phi), \quad x > 0, \tag{1.1b}$$

where x > 0 and $\xi = (\xi_1, \xi_2, \xi_3) = (\xi_1, \xi') \in \mathbb{R}^3$ are the space variable and velocity, respectively. The unknown functions $f = f(x, \xi)$ and $-\phi = -\phi(x)$ stand for the velocity distribution of positive ions and the electrostatic potential, respectively. The given function $n_e(\cdot) \in C^2(\mathbb{R})$ denotes the number density of electrons. We assume that

$$n_e(0) = 1, \quad n'_e(0) = -1.$$

One of typical functions is the Boltzmann relation $n_e(\phi) = e^{-\phi}$.

We study the boundary value problem of (1.1a)–(1.1b) with the boundary conditions

$$f(0,\xi) = f_b(\xi) + \alpha f(0, -\xi_1, \xi'), \quad \xi_1 > 0,$$
(1.1c)

$$\phi(0) = \phi_b, \tag{1.1d}$$

$$\lim_{x \to \infty} f(x,\xi) = f_{\infty}(\xi), \quad \xi \in \mathbb{R}^3,$$
(1.1e)

$$\lim_{x \to \infty} \phi(x) = 0. \tag{1.1f}$$

The constants $\alpha \in [0,1]$ and $-\phi_b \in \mathbb{R}$ denote the rate of refraction and the voltage on the boundary, respectively. Furthermore, $f_b = f_b(\xi)$ and $f_{\infty} = f_{\infty}(\xi)$ are given nonnegative functions. Physically speaking, the case $f_b = 0$ and $\alpha = 0$ corresponds to *a completely absorbing wall*. In addition, $\phi_b > 0$ and $\phi_b < 0$ mean that the wall is negatively and positively charged, respectively. Let us say *attractive* and *repulsive* boundaries for positive ions if $\phi_b > 0$ and $\phi_b < 0$, respectively.

Riemann [32] studied essentially the same boundary value problem as (1.1) to derive the kinetic Bohm criterion

$$\int_{\mathbb{R}^3} \xi_1^{-2} f_\infty(\xi) d\xi \le 1.$$
(1.2)

He also assumed that the number density of electrons is given by a function of the electrostatic potential as in (1.1b). On the other hand, he did not impose any boundary condition for the potential ϕ at x = 0, and supposed implicitly a situation that ϕ is monotone. The condition (1.1d) is one of simplest boundary conditions that can create the situation. In stead of (1.1d), we can also impose the following boundary condition:

$$n_e(\phi(0))v_e = (1-\alpha) \int_{\mathbb{R}^3} \xi_1 f(0,\xi) d\xi, \qquad (1.3)$$

where v_e is a constant. This condition means physically that the outgoing fluxes of electrons and ions coincide at the boundary. Even if we consider a boundary value problem of (1.1a)–(1.1b) with conditions (1.1c), (1.1e), (1.1f), and (1.3), it is seen that a value $\phi(0)$ is determined a priori, and thus we can reduce the problem to the boundary value problem (1.1) (for more details, see Appendix A). Therefore, we focus ourself on the study of the problem (1.1) in this paper.

The derivation of (1.2) by Riemann [32] is clear to understand, but it seems to be simplified. Indeed he divided the two cases $\xi_1 > 0$ and $\xi_1 < 0$, and then changed the coordinates according to $\xi_1 \ge 0$. Some expansion of the unknown function ϕ was also used. One of our purposes is to find a simpler derivation avoiding the use of such coordinate transformations and expansions.

The Vlasov–Poisson system in (1.1) is given by a system of partial and ordinary differential equations, while the stationary Euler–Poisson system is given by a system of just ordinary differential equations. Besides (1.1b) has a non-local term. The idea to resolve these difficulties is to reduce the problem (1.1) to a boundary value problem of a first-order ordinary differential

equation only for ϕ by combining the characteristics method and the technique used in [40]. The reduction also enables us to derive more simply the kinetic Bohm criterion (1.2), which is a necessary condition of the solvability of (1.1). We will construct the solution under (1.2) (see Theorems 2.2, 2.5, and 2.6 below).

It is worth pointing out that letting $f_{\infty}(\xi)$ be a delta function $\delta(\xi - (-u_{\infty}, 0, 0))$ in the kinetic Bohm criterion (1.2), one can obtain the hydrodynamic Bohm criterion:

$$u_{\infty}^2 \ge 1, \tag{1.4}$$

where u_{∞} is a positive constant and $-u_{\infty}$ means the flow velocity of positive ions at infinite distance. Bohm derived originally the criterion (1.4) by using the stationary Euler-Poisson system of cold plasma:

$$(\rho u)' = 0, \quad uu' = \phi', \quad \phi'' = \rho - e^{-\phi}, \quad x > 0,$$
 (1.5a)

where $\rho = \rho(x)$, u = u(x), and $-\phi = -\phi(x)$ represent the number density and flow velocity of positive ions and the electrostatic potential, respectively. Suzuki [40] showed the unique existence of solutions of the system (1.5a) with the conditions

$$\inf_{x \in \mathbb{R}_+} \rho(x) > 0, \quad \phi(0) = \phi_b, \quad \lim_{x \to \infty} (\rho, u, \phi)(x) = (1, -u_{\infty}, 0), \tag{1.5b}$$

where $u_{\infty} > 0$ and $\phi_b \in \mathbb{R}$ are constants. For more details, see Proposition 2.3 below.

It is of great interest to investigate the relation between the solutions of the Vlasov–Poisson system (1.1) and the Euler–Poisson system (1.5). To clarify the relation, we choose some approximate functions of the delta function $\delta(\xi - (-u_{\infty}, 0, 0))$ for f_{∞} , and then show in Theorem 2.4 that the solution of (1.1) converges to that of (1.5) by taking the limit of approximate functions to the delta function. We call the limit *a delta mass limit*.

We review mathematical results on the Vlasov–Poisson system describing the motion of plasma. For the Cauchy problem, early references [3, 27, 29, 35] investigated the time-global solvability and dispersive analysis (see also textbooks [12, 31]). Guo–Strauss [15] established spatially periodic stationary solutions and also investigated its instability (see also [16] studying the relativistic Vlasov–Maxwell system). For the the initial–boundary value problem, Guo [14] and Hwang–Velázquez [20] showed the time-global solvability adopting the specular reflection boundary condition. Han-Kwan–Rousset [19] and Han-Kwan–Iacobelli [17, 18] analyzed the quasi-neutral limit of solutions satisfying a periodic boundary condition. Furthermore, Skubachevskii [36, 37] and Skubachevskii–Tsuzuki [39] focused on the analysis of toroidal magnetic plasma containment devices (tokamak), and established the existence of solutions of the initial–boundary value problem with an external magnetic field, where supports supp f do not contact with boundaries.

The stationary problem has also been extensively studied. In an infinite cylinder and a halfspace, Belyaeva [4] and Skubachevskii [38] constructed stationary solutions whose supports supp f do not contact with boundaries by applying an external magnetic field. Similarly, Knopf [24] established the stationary solution in the whole space, where the support supp f is contained in an infinite cylinder. Let us introduce results which treated more similar settings to this paper. Greengard–Raviart [13] and Rein [30] constructed the stationary solutions in bounded domains adopting the Dirichlet and the specular refection boundary conditions, respectively. Recently, Esentürk–Hwang–Strauss [8] showed the solvability of the stationary problem with diffusive boundary conditions for various domains including a half-space. For the half-space case, they considered a situation that there are only particles whose energies $|\xi|^2/2$ are bounded by some finite number, and the boundary is not electrically charged. This point is one of the differences between settings in [8] and this paper. We also emphasize that there is no research which studies the delta mass limit mentioned above.

This paper is organized as follows. Section 2 provides our main theorems on the solvability and delta mass limit. In Section 3, we treat the completely absorbing and attractive boundary, i.e. $f_b = \alpha = 0$ and $\phi_b > 0$. Subsection 3.1 is devoted to discussion on a simpler derivation of the kinetic Bohm criterion (1.2). Subsection 3.2 establishes the solvability of the problem (1.1). We also justify the delta mass limit in subsection 3.3. In Section 4, we study general boundaries, i.e. $(f_b, \alpha) \neq (0,0)$ and $\phi_b \neq 0$.

Before closing this section, we give our notation used throughout this paper.

Notation. For $1 \le p \le \infty$, $L^p(\Omega)$ is the Lebesgue space equipped with the norm $\|\cdot\|_{L^p(\Omega)}$. Let us denote by $(\cdot, \cdot)_{L^2(\Omega)}$ the inner product of $L^2(\Omega)$. For $1 < r < \infty$ and $-\infty \le a < b \le \infty$, the function spaces $L^r(a, b; L^1(\mathbb{R}^2))$ and $L^r_{loc}(\mathbb{R}; L^1(\mathbb{R}^2))$ are defined by

$$\begin{split} L^{r}(a,b;L^{1}(\mathbb{R}^{2})) &:= \left\{ f \in L^{1}((a,b) \times \mathbb{R}^{2}) \left| \|f\|_{L^{r}(a,b;L^{1}(\mathbb{R}^{2}))} < \infty \right\}, \\ L^{r}_{loc}(\mathbb{R};L^{1}(\mathbb{R}^{2})) &:= \left\{ f \in L^{1}_{loc}(\mathbb{R}^{3}) \left| \|f\|_{L^{r}(a,b;L^{1}(\mathbb{R}^{2}))} < \infty \text{ for } -\infty < \forall a < \forall b < \infty \right\}, \\ \|f\|_{L^{r}(a,b;L^{1}(\mathbb{R}^{2}))} &:= \left\{ \int_{a}^{b} \left(\int_{\mathbb{R}^{2}} |f(\xi_{1},\xi')| d\xi' \right)^{r} d\xi_{1} \right\}^{1/r}. \end{split}$$

Furthermore, $\mathbb{R}_+ := \{x > 0\}$ stands for a one-dimensional half space; $\mathbb{R}^3_+ := \{\xi \in \mathbb{R}^3; \xi_1 > 0\}$ stands for a three-dimensional upper half space; $\mathbb{R}^3_- := \{\xi \in \mathbb{R}^3; \xi_1 < 0\}$ stands for a three-dimensional lower half space. We also use the one-dimensional indicator function $\chi(s)$ of the set $\{s > 0\}$.

2 Main Results

We focus ourself on the analysis of solutions whose potential ϕ is monotone, since the potential is observed as a monotone function when the plasma sheath is formed. Let us give a definition of solutions of the boundary value problem (1.1).

Definition 2.1. We say that (f, ϕ) is a solution of the boundary value problem (1.1) if it satisfies *the following:*

- (i) $f \in L^1_{loc}(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \cap C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$ and $\phi \in C^1(\overline{\mathbb{R}_+}) \cap C^2(\mathbb{R}_+)$.
- (*ii*) $f(x) \ge 0$, and either $\partial_x \phi(x) > 0$ or $\partial_x \phi(x) < 0$.
- (iii) f solves

$$(f,\xi_1\partial_x\psi+\partial_x\phi\partial_{\xi_1}\psi)_{L^2(\mathbb{R}_+\times\mathbb{R}^3)} + (f_b,\xi_1\psi(0,\cdot))_{L^2(\mathbb{R}^3_+)} = 0 \quad for \ \forall \psi \in \mathscr{X},$$
(2.1a)
$$\lim_{x \to \infty} \|f(x,\cdot) - f_\infty\|_{L^1(\mathbb{R}^3)} = 0,$$
(2.1b)

where
$$\mathscr{X} := \{ f \in C_0^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \mid \alpha f(0, \xi_1, \xi') = f(0, -\xi_1, \xi') \text{ for } (\xi_1, \xi') \in \mathbb{R}^3_+ \}.$$

(iv) ϕ solves (1.1b) with (1.1d) and (1.1f) in the classical sense.

The equation (2.1a) is a standard weak form of the equation (1.1a) and boundary condition (1.1c). We also remark that it is possible to replace *the classical sense* in the condition (iv) by *the weak sense*. Indeed a weak solution ϕ of the problem of (1.1b) with (1.1d) and (1.1f) is a classical solution if $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$.

Next we discuss some necessary conditions for the solvability of the problem (1.1), which are used to state our main results. To solve the Poisson equation (1.1b) with (1.1f), we must require the quasi-neutral condition

$$\int_{\mathbb{R}^3} f_{\infty}(\xi) d\xi = 1.$$
(2.2)

We remark that the boundary value problem (1.1) is overdetermined. Let us explain briefly for the case $\phi_b = 0$. First ϕ must be zero, and then the equation (1.1a) implies that f is independent of x. This fact together with the boundary condition (1.1e) means that $f = f_{\infty}$. On the other hand, due to the boundary condition (1.1c), we have a necessary condition for f_{∞} and f_b :

$$f_{\infty}(\xi)=f_b(\xi)+lpha f_{\infty}(-\xi_1,\xi'), \quad \xi=(\xi_1,\xi')\in\mathbb{R}^3_+.$$

Consequently, we cannot choose independently f_{∞} and f_b . For the case $\phi_b \neq 0$, the ¹following are necessary conditions:

$$f_{\infty}(\xi) = f_{b}(\sqrt{\xi_{1}^{2} + 2\phi_{b}}, \xi') + \alpha f_{\infty}(-\xi_{1}, \xi'), \quad \xi \in \mathbb{R}^{3}_{+} \quad \text{if } \phi_{b} > 0, \quad (2.3)$$

$$f_{\infty}(\xi) = f_{b}(\sqrt{\xi_{1}^{2} + 2\phi_{b}}, \xi') + \alpha f_{\infty}(-\xi_{1}, \xi'), \quad \xi \in (\sqrt{2|\phi_{b}|}, \infty) \times \mathbb{R}^{2},$$

$$f_{\infty}(\xi_{1}, \xi') = f_{\infty}(-\xi_{1}, \xi'), \quad \xi \in (-\sqrt{2|\phi_{b}|}, \sqrt{2|\phi_{b}|}) \times \mathbb{R}^{2} \quad \text{if } \phi_{b} < 0. \quad (2.4)$$

In particular, for the completely absorbing and attractive boundary, i.e. $f_b = \alpha = 0$ and $\phi_b > 0$, it is written by

$$f_{\infty}(\xi) = 0, \quad \xi_1 > 0.$$
 (2.5)

We will show that (2.3), (2.4), and (2.5) are necessary conditions in Lemmas 4.1, 4.3, and 3.1, respectively. From the above observation, we also see that $(f, \phi) = (f_{\infty}, 0)$ is a unique solution of (1.1) with $\phi_b = 0$, and hence suppose $\phi_b \neq 0$ hereafter.

We state our main results for the completely absorbing and attractive boundary, i.e. $f_b = \alpha = 0$ and $\phi_b > 0$ in subsection 2.1. It is one of the most discussed situation in plasma physics. Subsection 2.2 provides the main results for general boundaries, i.e. $(f_b, \alpha) \neq (0, 0)$ and $\phi_b \neq 0$.

¹We may not find any physical meaning of (2.3)–(2.5), but mathematically speaking there is no solution without them.

2.1 The Completely Absorbing and Attractive Boundary

We first discuss the results of the solvability of the problem (1.1) with the completely absorbing and attractive boundary. After that we also study the delta mass limit of the solution. It validates rigorously the relation of the kinetic and hydrodynamic Bohm criterions.

The solvability is summarized in Theorem 2.2 below. Here the set B is defined for f_{∞} as

$$B := \{ \varphi > 0; V(\phi) > 0 \text{ for } \phi \in (0, \varphi] \},\$$

where

$$V(\phi) := \int_0^{\phi} \rho_i(\phi) - n_e(\phi) d\phi, \quad \rho_i(\phi) := \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{-\xi_1}{\sqrt{\xi_1^2 + 2\phi}} d\xi.$$
(2.6)

The function ρ_i is well-defined for $f_{\infty} \in L^1(\mathbb{R})$. Indeed,

$$|\rho_i(\phi)| \le ||f_{\infty}||_{L^1(\mathbb{R}^3)} \quad \text{for } \phi \ge 0.$$
 (2.7)

Theorem 2.2. Let $f_b = \alpha = 0$ and $\phi_b > 0$. Suppose that $f_{\infty} \in L^1(\mathbb{R}^3)$ satisfies $f_{\infty} \ge 0$ and the necessary conditions (2.2) and (2.5).

(i) Assume that

$$\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi < 1.$$
 (2.8)

Then the set *B* is not empty. Furthermore, if and only if $\phi_b < \sup B$ holds, the problem (1.1) has a unique solution (f, ϕ) . There also hold that

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x),\xi'})\chi(\xi_1^2 - 2\phi(x))\chi(-\xi_1), \qquad (2.9)$$

$$|\partial_x^l \phi(x)| \le C e^{-cx} \quad for \ l = 0, 1, 2,$$
 (2.10)

where $\chi(s)$ is the one-dimensional indicator function of the set $\{s > 0\}$, and c and C are positive constants independent of x. In addition, if $f_{\infty} \in C^2(\mathbb{R}^3)$, then the solution $(f, \phi) \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \times C^2(\overline{\mathbb{R}_+})$ is a classical solution.

(ii) Assume that

$$\int_{\mathbb{R}^3} \xi_1^{-2} f_\infty(\xi) d\xi = 1, \quad B \neq \emptyset.$$

If and only if $\phi_b < \sup B$ holds, the problem (1.1) has a unique solution (f, ϕ) . Further, (2.9) holds. If $f_{\infty} \in C^2(\mathbb{R}^3)$, then the solution $(f, \phi) \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \times C^2(\overline{\mathbb{R}_+})$ is a classical solution.

(iii) Assume that

$$\int_{\mathbb{R}^3} \xi_1^{-2} f_\infty(\xi) d\xi = 1, \quad B = \emptyset.$$

Then the problem (1.1) *admits no solution.*

(iv) Assume ²that

$$\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi > 1.$$
 (2.11)

Then the problem (1.1) *admits no solution.*

This theorem covers all possible cases of $\phi_b > 0$ and $f_{\infty} \in L^1(\mathbb{R}^3)$ with $f_{\infty} \ge 0$, and clarifies completely when there is a solution or not. Assertion (iv) means that the kinetic Bohm criterion (1.2) is a necessary condition for the solvability of the problem (1.1), since (2.11) is the negation of (1.2). Assertion (iv) will be shown in subsection 3.1. The proof gives simultaneously a simpler derivation of the kinetic Bohm criterion than that of [32]. Subsection 3.1 also provides the proof of Assertion (iii). In subsection 3.2, we will prove Assertions (i) and (ii).

Next we discuss the relation between the solutions of the Vlasov–Poisson system (1.1) and the Euler–Poisson system (1.5). As mentioned in Section 1, the hydrodynamic Bohm criterion (1.4) can be derived from the kinetic Bohm criterion (1.2) by plugging a delta function $\delta(\xi - (-u_{\infty}, 0, 0))$ into f_{∞} . To investigate the relation of solutions, we use

$$f_{\infty,\varepsilon}(\xi) := \frac{1}{\varepsilon^3} \varphi\left(\frac{\xi - (-u_{\infty}, 0, 0)}{\varepsilon}\right), \qquad (2.12)$$

where u_{∞} is a positive constant and

$$0 < \varepsilon < 1, \quad \varphi \in C_0^{\infty}(\mathbb{R}^3), \quad \varphi(\xi) \ge 0, \quad \operatorname{supp} \varphi \subset \{|\xi| < 1\}, \quad \int_{\mathbb{R}^3} \varphi(\xi) d\xi = 1.$$
 (2.13)

Note that

$$\lim_{\varepsilon \to 0} \int_{\mathbb{R}^3} f_{\infty,\varepsilon}(\xi) \psi(\xi) d\xi = \psi(-u_{\infty},0,0), \quad \forall \psi \in C(\mathbb{R}^3).$$

This means that $f_{\infty,\varepsilon}(\xi)$ converges to the delta function $\delta(\xi - (-u_{\infty}, 0, 0))$ as $\varepsilon \to 0$. Let us denote by $(f_{\varepsilon}, \phi_{\varepsilon})$ the solution of the problem (1.1) with $f_{\infty} = f_{\infty,\varepsilon}$, where $f_{\infty,\varepsilon} \in C_0^{\infty}(\mathbb{R}^3)$ satisfies the conditions (2.2), (2.5), and (2.8) being in Theorem 2.2 for $u_{\infty} > 1$ and $\varepsilon \ll 1$. We also introduce the moments

$$\rho_{\varepsilon}(x) := \int_{\mathbb{R}^3} f_{\varepsilon}(x,\xi) d\xi, \quad u_{\varepsilon}(x) := \frac{1}{\rho_{\varepsilon}(x)} \int_{\mathbb{R}^3} \xi_1 f_{\varepsilon}(x,\xi) d\xi.$$

It is expected that the moments converge to (ρ, u) of the Euler–Poisson system (1.5) by taking the delta mass limit as $\varepsilon \to 0$. The solvability of the problem (1.5) is summarized in the following proposition.

Proposition 2.3 ([40]). Let (1.4) and $\phi_b \in (-u_{\infty}^2/2, \infty)$ hold. Then the problem (1.5) has a unique monotone solution $(\rho, u, \phi) \in C^{\infty}(\overline{\mathbb{R}_+})$. Furthermore, there hold that

$$\rho = \frac{u_{\infty}}{\sqrt{u_{\infty}^2 + 2\phi}}, \quad u = -\sqrt{u_{\infty}^2 + 2\phi}, \quad \partial_x \phi(x) \neq 0, \quad (2.14)$$
$$|\partial_x^l(\rho - 1, u + u_{\infty}, \phi)(x)| \le Ce^{-cx}, \quad l = 0, 1, 2,$$

where c and C are positive constants independent of x.

²This includes the case $\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi = \infty$.

We are now in a position to state our results on the delta mass limit.

Theorem 2.4. Let $f_b = \alpha = 0$, $n_e(\phi) = e^{-\phi}$, and $u_{\infty} > 1$. There exist positive constants δ_0 and C_0 such that if $0 < \phi_b < \delta_0$, then the following holds:

$$\sup_{x \in \mathbb{R}_+} |\rho_{\varepsilon}(x) - \rho(x)| + \sup_{x \in \mathbb{R}_+} |\rho_{\varepsilon}u_{\varepsilon}(x) - \rho_{\varepsilon}u(x)| + \sup_{x \in \mathbb{R}_+} |\phi_{\varepsilon}(x) - \phi(x)| \le C_0 \varepsilon$$
(2.15)

for any $\varepsilon \in (0, \varepsilon_0)$, where $\varepsilon_0 := (u_{\infty} - 1)/2$.

Now we mention some remarks. We cannot choose directly a Maxwellian for f_{∞} due to the necessary condition (2.5) for the solvability of the problem (1.1), but it allows products of the Maxwellian and cut-off functions. For such products, the delta mass limit as $\varepsilon \to 0$ corresponds to that the temperature of the Maxwellian tends to zero. In this sense, it is reasonable that the limit is a solution of the Euler–Poisson system of cold plasma.

2.2 General Boundaries

This section provides the main results for general boundary conditions, i.e. $(f_b, \alpha) \neq (0, 0)$ and $\phi_b \neq 0$. Similarly as Theorem 2.2, we use notation

$$B^{+} := \{ \varphi > 0; V^{+}(\phi) > 0 \text{ for } \phi \in (0, \varphi] \}, \quad V^{+}(\phi) := \int_{0}^{\phi} \left(\rho_{i}^{+}(\varphi) - n_{e}(\varphi) \right) d\varphi, \quad (2.16)$$

$$B^{-} := \{ \varphi < 0; V^{-}(\phi) > 0 \text{ for } \phi \in [\varphi, 0) \}, \quad V^{-}(\phi) := \int_{0}^{\phi} \left(\rho_{i}^{-}(\varphi) - n_{e}(\varphi) \right) d\varphi, \quad (2.17)$$

where

$$\rho_{i}^{+}(\phi) := \int_{\mathbb{R}^{3}} f_{\infty}(\xi) \frac{|\xi_{1}|}{\sqrt{\xi_{1}^{2} + 2\phi}} d\xi + \frac{2}{1 - \alpha} \int_{\sqrt{2(\phi_{b} - \phi)\chi(\phi_{b} - \phi)}}^{\sqrt{2\phi_{b}}} \int_{\mathbb{R}^{2}} f_{b}(\xi) \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2(\phi - \phi_{b})}} d\xi_{1} d\xi' \quad \text{for } \phi \geq 0, \quad (2.18)$$

$$\rho_i^-(\phi) := \int_{\mathbb{R}^3} f_\infty(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi}} \chi(\xi_1^2 + 2\phi) d\xi \qquad \text{for } \phi \le 0.$$
(2.19)

The functions ρ_i^{\pm} are well-defined for $f_b \in L^1(\mathbb{R}^3_+)$ and $f_{\infty} \in L^1(\mathbb{R}^3)$ with $f_b, f_{\infty} \ge 0$, since all the integrants are nonnegative. Furthermore, for r > 2 and M > 0, it is seen ³that

$$|\rho_i^+(\phi)| \le \|f_\infty\|_{L^1(\mathbb{R}^3)} + C\|f_b\|_{L^r(0,\sqrt{2\phi_b};L^1(\mathbb{R}^2))} \qquad \text{for } \phi \ge 0,$$
(2.20)

$$|\rho_i^-(\phi)| \le \sqrt{2} \|f_\infty\|_{L^1(\mathbb{R}^3)} + C_M \|f_\infty\|_{L^r(-2\sqrt{M}, 2\sqrt{M}; L^1(\mathbb{R}^2))} \quad \text{for } \phi \in [-M, 0], \quad (2.21)$$

where *C* is a positive constant, and C_M is a positive constant depending on *M*. The proofs of the estimates are postponed until Appendix B. In the definition (2.18), the indicator function $\chi(\phi_b - \phi)$ is used to extend continuously ρ_i^+ beyond $\phi = \phi_b$.

Now we state main theorems on the solvability by employing the function spaces $L^r(a,b;L^1(\mathbb{R}^2))$ and $L^r_{loc}(\mathbb{R};L^1(\mathbb{R}^2))$. Here the set B^+ depends on ϕ_b whereas the set B^- is independent of ϕ_b .

³If the problem (1.1) has a solution (f, ϕ) , then $\rho_i^+ \in C([0, \phi_b])$ and $\rho_i^- \in C([\phi_b, 0])$ must hold, whether or not $f_b \in L^r(0, \sqrt{2\phi_b}; L^1(\mathbb{R}^2))$ and $f_\infty \in L^r_{loc}(\mathbb{R}; L^1(\mathbb{R}^2))$ hold. For more details, see Lemmas 4.1 and 4.3.

Theorem 2.5. (Attractive boundary) Let $\phi_b > 0$, $f_b \in L^1(\mathbb{R}^3_+) \cap L^r(0, \sqrt{2\phi_b}; L^1(\mathbb{R}^2))$, ${}^4\alpha \neq 1$, $f_{\infty} \in L^1(\mathbb{R}^3)$, and $f_b, f_{\infty} \ge 0$ for some r > 2. Suppose ⁵that (2.2), (2.3), and $V^+ \in C^2([0, \phi_b])$ hold.

(i) Assume that $d^2V^+/d\phi^2(0) > 0$. Then the set B^+ is not empty. Furthermore, ⁶if and only if $\phi_b < \sup B^+$ holds, the problem (1.1) has a unique solution (f, ϕ) . There also hold that

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_{1}^{2} - 2\phi(x),\xi'})\chi(\xi_{1}^{2} - 2\phi(x))\chi(-\xi_{1}) + \frac{1}{1-\alpha}f_{b}(\sqrt{\xi_{1}^{2} - 2\phi(x) + 2\phi_{b}},\xi')\chi(-\xi_{1}^{2} + 2\phi(x))$$
(2.22)
$$+ f_{\infty}(\sqrt{\xi_{1}^{2} - 2\phi(x)},\xi')\chi(\xi_{1}^{2} - 2\phi(x))\chi(\xi_{1}), |\partial_{x}^{l}\phi(x)| \leq Ce^{-cx}, \quad l = 0, 1, 2,$$
(2.23)

where *c* and *C* are positive constants independent of *x*.

- (ii) Assume that $d^2V^+/d\phi^2(0) = 0$ and $B^+ \neq \emptyset$. If and only if $\phi_b < \sup B^+$ holds, the problem (1.1) has a unique solution (f, ϕ) . Furthermore, (2.22) holds.
- (iii) Assume that $d^2V^+/d\phi^2(0) = 0$ and $B^+ = \emptyset$. Then the problem (1.1) admits no solution.
- (iv) Assume that $d^2V^+/d\phi^2(0) < 0$. Then the problem (1.1) admits no solution.

Theorem 2.6. (Repulsive boundary) Let $\phi_b < 0$, $f_b \in L^1(\mathbb{R}^3_+)$, $f_\infty \in L^1(\mathbb{R}^3) \cap L^r_{loc}(\mathbb{R}; L^1(\mathbb{R}^2))$, and $f_b, f_\infty \ge 0$ for some r > 2. Suppose ⁷that (2.2), (2.4), and $V^- \in C^2([\phi_b, 0])$ hold.

(i) Assume that $d^2V^-/d\phi^2(0) > 0$. Then the set B^- is not empty. Furthermore, if and only if $\phi_b > \inf B^-$ holds, the problem (1.1) has a unique solution (f, ϕ) . There also hold that (2.23) and

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x)}, \xi')\chi(-\xi_1) + f_{\infty}(\sqrt{\xi_1^2 - 2\phi(x)}, \xi')\chi(\xi_1).$$
(2.24)

- (ii) Assume that $d^2V^-/d\phi^2(0) = 0$ and $B^- \neq \emptyset$. If and only if $\phi_b > \inf B^-$ holds, the problem (1.1) has a unique solution (f, ϕ) . Furthermore, (2.24) holds.
- (iii) Assume that $d^2V^-/d\phi^2(0) = 0$ and $B^- = \emptyset$. Then the problem (1.1) admits no solution.
- (iv) Assume that $d^2V^-/d\phi^2(0) < 0$. Then the problem (1.1) admits no solution.

From these theorems, we conclude that the following conditions are general representations of the kinetic Bohm criterion:

$$\frac{d^2 V^{\pm}}{d\phi^2}(0) \ge 0. \tag{2.25}$$

⁴We can construct multiple solutions for the case $\alpha = 1$, and therefore we exclude it.

⁵If a solution exists, then (2.2), (2.3), and $V^+ \in C^1([0, \phi_b])$ must hold. For more details, see Lemma 4.1.

⁶If supp $f_b \subset (c, \infty) \times \mathbb{R}^2$ for some c > 0, the second term in the definition of ρ_i^+ vanishes for $\phi_b < c^2/2$. This means that B^+ is independent of ϕ_b . In this case, $\phi_b < \sup B^+$ holds for $\phi_b \ll 1$.

⁷If a solution exists, then (2.2), (2.4), and $V^- \in C^1([\phi_b, 0])$ must hold. For more details, see Lemma 4.3.

Indeed, it can be rewritten by the kinetic Bohm criterion (1.2) for the case $f_b = 0$, $\alpha = 0$, and $\phi_b > 0$ (for details, see (3.6)). We can find some functions (f_b, f_{∞}) so that $V^{\pm} \in C^2$ and (2.25) hold, for example, the functions $(g_{b,\varepsilon}, g_{\infty,\varepsilon})$ defined in (2.26) and (2.27) below. We introduce some more general functions in Appendix C. Theorems 2.5 and 2.6 will be shown in subsections 4.1 and 4.2, respectively.

We also state the results on the delta mass limit. Let us define the functions ${}^8g_{b,\varepsilon}$ and $g_{\infty,\varepsilon}$ as

$$g_{b,\varepsilon}(\xi) := \frac{m_b}{\varepsilon^3} \varphi \left(\frac{(\sqrt{\xi_1^2 - 2\phi_b, \xi') - (v_b, 0, 0)}}{\varepsilon} \right) \chi(\xi_1^2 - 2\phi_b), \quad \xi \in \mathbb{R}^3_+, \quad (2.26)$$

$$g_{\infty,\varepsilon}(\xi) := \frac{m_\infty}{\varepsilon^3} \varphi \left(\frac{\xi - (-v_\infty, 0, 0)}{\varepsilon} \right) + \frac{m_b}{\varepsilon^3} \varphi \left(\frac{\xi - (v_b, 0, 0)}{\varepsilon} \right) + \alpha \frac{m_\infty}{\varepsilon^3} \varphi \left(\frac{(-\xi_1, \xi') + (v_\infty, 0, 0)}{\varepsilon} \right), \quad \xi \in \mathbb{R}^3, \quad (2.27)$$

where ε and φ are defined in (2.13), and $m_b \ge 0$, $m_{\infty} > 0$, $v_b > 0$, and $v_{\infty} > 0$ are constants. It is supposed that

$$m_b + (1+\alpha)m_{\infty} = 1, \quad m_b v_b^{-2} + (1+\alpha)m_{\infty}v_{\infty}^{-2} < 1,$$
 (2.28)

which ensures (2.2) and $\int_{\mathbb{R}^3} \xi_1^{-2} g_{\infty,\varepsilon}(\xi) d\xi < 1$ for $\varepsilon \ll 1$. Let us denote by $(f_{\varepsilon}, \phi_{\varepsilon})$ the solution of the problem (1.1) with $(f_b, f_{\infty}) = (g_{b,\varepsilon}, g_{\infty,\varepsilon})$. The important thing to note here is that $g_{b,\varepsilon}$ and $g_{\infty,\varepsilon}$ satisfy the conditions (2.2)–(2.4) and $d^2V^{\pm}/d\phi^2(0) = -\int_{\mathbb{R}^3} \xi_1^{-2} g_{\infty,\varepsilon}(\xi) d\xi + 1 > 0$ being in Theorems 2.5 and 2.6 for the case $|\phi_b| \ll 1$ and $\varepsilon \ll 1$. In this case, $\sup B^+$ is independent of ϕ_b , and also $\phi_b < \sup B^+$ holds, since the second term in the definition (2.18) of ρ_i^+ vanishes. Note that $\inf B^-$ is always independent of ϕ_b . We also introduce the moments

$$\rho_{\varepsilon}(x) := \int_{\mathbb{R}^3} f_{\varepsilon}(x,\xi) d\xi, \quad u_{\varepsilon}(x) := \frac{1}{\rho_{\varepsilon}(x)} \int_{\mathbb{R}^3} \xi_1 f_{\varepsilon}(x,\xi) d\xi.$$

Theorem 2.7. Let $\phi_b \neq 0$, $\alpha \neq 1$, and (2.28) hold. There exist positive constants ε_0 , δ_0 , and C_0 such that if $|\phi_b| < \delta_0$, then the following holds:

$$\sup_{x \in \mathbb{R}_+} |\rho_{\varepsilon}(x) - \rho(x)| + \sup_{x \in \mathbb{R}_+} |\rho_{\varepsilon}u_{\varepsilon}(x) - \rho_{\varepsilon}u(x)| + \sup_{x \in \mathbb{R}_+} |\phi_{\varepsilon}(x) - \phi(x)| \le C_0 \varepsilon$$
(2.29)

for any $\varepsilon \in (0, \varepsilon_0)$, where $\phi = \phi(x)$ solves a boundary value problem

$$\partial_{xx}\phi = \frac{m_b v_b}{\sqrt{v_b^2 + 2\phi}} + \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^2 + 2\phi}} - n_e(\phi), \quad x > 0,$$
(2.30a)

$$\phi(0) = \phi_b, \quad \lim_{x \to 0} \phi(x) = 0,$$
 (2.30b)

and $\rho = \rho(x)$ and u = u(x) are defined as

$$\rho(x) := \frac{m_b v_b}{\sqrt{v_b^2 + 2\phi(x)}} + \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^2 + 2\phi(x)}}, \quad u(x) := \frac{m_b v_b + (\alpha - 1)m_{\infty}v_{\infty}}{\rho(x)}.$$
 (2.31)

⁸We first determine $g_{\infty,\varepsilon}$ as (2.27), and then find a suitable $g_{b,\varepsilon}$ as (2.26) so that the necessary conditions (2.3) and (2.4) hold. This choice of $g_{\infty,\varepsilon}$ is one of the simplest extensions of $f_{\infty,\varepsilon}$ in (2.12).

For the case $m_b = 0$, i.e. $f_b = g_{b,\varepsilon} = 0$, the functions (ρ, u, ϕ) in Theorem 2.7 solve a problem

$$(\rho u)' = 0, \quad \frac{(\alpha + 1)^2}{(\alpha - 1)^2} u u' = \phi', \quad \phi'' = \rho - n_e(\phi), \quad x > 0,$$
 (2.32a)

$$\inf_{x \in \mathbb{R}_+} \rho(x) > 0, \quad \lim_{x \to \infty} (\rho, u, \phi)(x) = \left(1, \frac{\alpha - 1}{\alpha + 1} v_{\infty}, 0\right), \quad \phi(0) = \phi_b, \tag{2.32b}$$

which is similar to (1.5). We conclude that (2.32) is a hydrodynamic model taking the reflection of positive ions on the boundary into account.

3 The Completely Absorbing and Attractive Boundary

In this section, we study the solvability and delta mass limit in Theorems 2.2 and 2.4 for completely absorbing and attractive boundary, i.e. $f_b = \alpha = 0$ and $\phi_b > 0$. We first show Assertion (iv) in Theorem 2.2 in subsection 3.1. The proof gives simultaneously a simpler derivation of the kinetic Bohm criterion (1.2) than that of [32], and also shows that (2.5) is a necessary condition. In subsection 3.1, we also prove Assertion (iii). Subsections 3.2 deals with the proofs of Assertions (i) and (ii) of Theorem 2.2. We show Theorem 2.4 in subsection 3.3.

3.1 A Derivation of the Kinetic Bohm Criterion

We discuss necessary conditions for the solvability by assuming that the solution (f, ϕ) of the problem (1.1) with $f_b = \alpha = 0$ and $\phi_b > 0$ exists. First we show the following lemma which ensures immediately Assertion (iv) in Theorem 2.2. Note that (1.2) is the negation of (2.11).

Lemma 3.1. Let $f_b = \alpha = 0$, $\phi_b > 0$, $f_{\infty} \in L^1(\mathbb{R}^3)$, $f_{\infty} \ge 0$, and (2.2) hold. Suppose that the problem (1.1) has a solution (f, ϕ) . Then the function f_{∞} satisfies the condition (2.5) and the kinetic Bohm criterion (1.2). Furthermore, ϕ solves (3.5) below with (1.1d) and (1.1f), and f is written as (2.9) by ϕ .

Proof. First $\partial_x \phi(x) < 0$ holds thanks to $\phi_b > 0$ and the condition (ii) in Definition 2.1. Owing to (1.1f), (2.1b), $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$, and $\phi \in C(\overline{\mathbb{R}_+})$, it follows from (1.1b) that $\partial_{xx}\phi$ is bounded and therefore $\partial_x \phi$ is uniformly continuous on $\overline{\mathbb{R}_+}$. Then the fact together with (1.1f) and $\partial_x \phi(x) < 0$ leads to a necessary condition

$$\lim_{x \to \infty} \partial_x \phi(x) = 0. \tag{3.1}$$

Let us first show that f_{∞} satisfies (2.5) and f is written as (2.9). Regarding ϕ as a given function and then applying the characteristics method to (2.1a), we see that the value of f must remain a constant along the following characteristic curves:

$$\frac{1}{2}\xi_1^2 - \phi(x) = c,$$

where c is some constant. We draw the illustration of characteristics in Figure 1 below. It tells

Figure 1: characteristic curves for the case $\partial_x \phi < 0$

us that

$$f(y, -\sqrt{\eta_1^2 + 2\phi(y)}, \eta') = f_{\infty}(\eta), \quad (y, \eta_1, \eta') \in X_1,$$
 (3.2a)

$$f(y, \pm \sqrt{2\phi(y) - \eta_1^2, \eta'}) = 0, \quad (y, \eta_1, \eta') \in X_2,$$
 (3.2b)

$$f(y, \sqrt{\eta_1^2 + 2\phi(y), \eta'}) = f_{\infty}(\eta) = 0, \quad (y, \eta_1, \eta') \in X_3,$$
(3.2c)

where $\boldsymbol{\eta} = (\eta_1, \eta_2, \eta_3) = (\eta_1, \eta')$ and

$$X_1 := \overline{\mathbb{R}_+} \times \mathbb{R}^3_-, \quad X_2 := \{ (y, \eta_1, \eta') \in \overline{\mathbb{R}_+} \times \mathbb{R}^3 \mid \eta_1^2 < 2\phi(y) \le 2\phi_b \}, \quad X_3 := \overline{\mathbb{R}_+} \times \mathbb{R}^3_+.$$

The last inequality (3.2c) means that (2.5) must hold. Furthermore, we conclude from (3.2) that f must be written by (2.9), i.e.

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x)},\xi')\chi(\xi_1^2 - 2\phi(x))\chi(-\xi_1),$$

where $\chi(s)$ is the one-dimensional indicator function of the set $\{s > 0\}$. Next we show that ϕ satisfies (3.5) below. Integrating (2.9) over \mathbb{R}^3 , and using (2.5) and the change of variable $\sqrt{\xi_1^2 - 2\phi} = -\zeta_1$, we see that

$$\int_{\mathbb{R}^3} f(x,\xi) d\xi = \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{-\xi_1}{\sqrt{\xi_1^2 + 2\phi(x)}} d\xi = \rho_i(\phi(x)),$$
(3.3)

where ρ_i is the same function defined in (2.6). Substituting (3.3) into (1.1b) yields an ordinary differential equation for ϕ :

$$\partial_{xx}\phi = \rho_i(\phi) - n_e(\phi), \quad x > 0.$$
(3.4)

Multiply (3.4) by $\partial_x \phi$, integrate it over (x, ∞) , and use (1.1f) and (3.1) to obtain

$$(\partial_x \phi)^2 = 2V(\phi), \quad V(\phi) = \int_0^\phi \left(\rho_i(\phi) - n_e(\phi)\right) d\phi, \tag{3.5}$$

where V is the same function defined in (2.6). Thus ϕ must satisfy (3.5).

To obtain the kinetic Bohm criterion (1.2), we divide the proof into two cases $\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi < \infty$ and $\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi = \infty$. For the former case, $V \in C^2([0, \phi_b])$ follows from the monotone convergence theorem and the fact $f_{\infty}, \xi_1^{-2} f_{\infty} \in L^1(\mathbb{R}^3)$. We also see from (2.2) and (2.5) that

$$V(0) = \frac{dV}{d\phi}(0) = 0, \quad \frac{d^2V}{d\phi^2}(0) = -\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi + 1.$$
(3.6)

Hence, we arrive at the criterion (1.2) from (3.5) and (3.6) with the aid of the Taylor theorem. Furthermore, we show that the other case does not occur. Suppose that $\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi = \infty$ holds. Then there hold that $V \in C^2((0, \phi_b])$, $V(0) = \frac{dV}{d\phi}(0) = 0$, and

$$rac{d^2 V}{d\phi^2}(\phi) < -C_0 \quad ext{for } \phi \in (0,c_0],$$

where c_0 and C_0 are some positive constants. These facts imply that $V(\phi) < 0$ holds for $\phi \in (0, c_0]$. It contradicts to (3.5), and thus $\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi < \infty$ holds. Consequently, (1.2) must hold.

This proof provides a simpler derivation of the kinetic Bohm criterion (1.2). Next we show Assertion (iii).

Proof of Assertion (iii) in Theorem 2.2. In the case $B \neq \emptyset$, either one of the following holds:

(a) $V(\phi) < 0$ around $\phi = 0$.

(b) There exists a sequence $\{\phi_n\}_{n=1}^{\infty}$ such that $V(\phi_n) = 0$ and $\lim_{n \to \infty} \phi_n = 0$.

On the other hand, it is clear from (3.5) that the case (a) does not occur. It suffices to consider the case (b). Suppose that a solution (f, ϕ) of the problem (1.1) exists. For sufficiently large *n*, there exists a sequence $\{x_n\}_{n=1}^{\infty}$ such that $\phi(x_n) = \phi_n$ and $\lim_{n\to\infty} x_n = \infty$. Then we see from (3.5) that $(\partial_x \phi)^2(x_n) = 2V(\phi(x_n)) = 2V(\phi_n) = 0$. This fact contradicts to $\partial_x \phi(x) < 0$ in Definition 2.1. Therefore, the problem (1.1) admits no solution.

3.2 Solvability

We prove Assertions (i) and (ii) in Theorem 2.2 on the solvability of the problem (1.1).

Proof of Assertions (i) and (ii) in Theorem 2.2. We first note that (3.6) holds owing to $\xi_1^{-2} f_{\infty} \in$ $L^1(\mathbb{R}^3)$. It is seen that $B \neq \emptyset$ if (2.8) holds, since the Taylor theorem with (2.8) and (3.6) ensures that V is positive around $\phi = 0$. It is easy to see from (3.5) and $\partial_x \phi < 0$ in Definition 2.1 that the condition $\phi_b < \sup B$ is necessary for the solvability stated in Assertions (i) and (ii).

Hereafter we discuss simultaneously Assertions (i) and (ii). Suppose that $\phi_b < \sup B$ holds, which implies that $V(\phi) > 0$ for $\phi \in (0, \phi_b]$. Let us construct ϕ with $\partial_x \phi < 0$ by solving (3.5) with (1.1d) and (1.1f). To this end, we rewrite (3.5) into the equivalent equation

$$\partial_x \phi = -\sqrt{2V(\phi)}.\tag{3.7}$$

The Taylor theorem together with (3.6) and the assumption $\int_{\mathbb{R}^3} \xi_1^{-2} f_\infty(\xi) d\xi \leq 1$ ensures that \sqrt{V} is Lipschitz continuous around $\phi = 0$. Combining this and the fact that $V(\phi) > 0$ for $\phi \in (0, \phi_b]$, we deduce that \sqrt{V} is Lipschitz continuous on $[0, \phi_b]$. Therefore, a standard theory of ordinary differential equations gives the unique solvability of (3.7) with (1.1d) and (1.1f). Then it is straightforward to see $\partial_x \phi < 0$ and $\phi \in C(\overline{\mathbb{R}_+}) \cap C^2(\mathbb{R}_+)$ from the equivalent equation. Thus we have the desired ϕ satisfying (3.5) with (1.1d) and (1.1f). Furthermore, it is easy to show by differentiating (3.5) and using $\partial_x \phi < 0$ that ϕ satisfies (3.4).

Now we define f as (2.9) by using ϕ , and prove that f satisfies the conditions (i)–(iii) in Definition 2.1. Owing to (2.5) and $f \in L^1(\mathbb{R}^3)$, there exists a sequence $\{f_{\infty}^k\} \subset C_0^{\infty}(\mathbb{R}^3)$ such

that $f_{\infty}^{k}(\xi) = 0$ for $\xi_{1} > -1/k$, and $f_{\infty}^{k} \to f_{\infty}$ in $L^{1}(\mathbb{R}^{3})$ as $k \to \infty$. Let us show the condition (i), i.e. $f \in C(\overline{\mathbb{R}_{+}}; L^{1}(\mathbb{R}^{3})) \cap L_{loc}^{1}(\mathbb{R}_{+} \times \mathbb{R}^{3})$. First $f(x, \cdot) \in L^{1}(\mathbb{R}^{3})$ follows from (2.7) and (3.3). To investigate the continuity of f, we set $G(x, \xi) = \chi(\xi_{1}^{2} - \chi)))$ $2\phi(x))\chi(-\xi_1)$ and observe that for $x, x_0 \in \overline{\mathbb{R}_+}$,

$$\begin{split} \|f(x) - f(x_0)\|_{L^1(\mathbb{R}^3)} &= \int_{\mathbb{R}^3} \left| f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x)}, \xi') G(x, \xi) - f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x_0)}, \xi') G(x_0, \xi) \right| d\xi \\ &\leq \int_{\mathbb{R}^3} \left| f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x)}, \xi') G(x, \xi) - f_{\infty}^k(-\sqrt{\xi_1^2 - 2\phi(x)}, \xi') G(x, \xi) \right| d\xi \\ &+ \int_{\mathbb{R}^3} \left| f_{\infty}^k(-\sqrt{\xi_1^2 - 2\phi(x)}, \xi') G(x, \xi) - f_{\infty}^k(-\sqrt{\xi_1^2 - 2\phi(x_0)}, \xi') G(x_0, \xi) \right| d\xi \\ &+ \int_{\mathbb{R}^3} \left| f_{\infty}^k(-\sqrt{\xi_1^2 - 2\phi(x_0)}, \xi') G(x_0, \xi) - f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x_0)}, \xi') G(x_0, \xi) \right| d\xi \\ &=: K_1 + K_2 + K_3. \end{split}$$

Using the change of variable $\zeta_1 = -\sqrt{\xi_1^2 - 2\phi(x)}$ and the fact that $f_{\infty}(\xi) = f_{\infty}^k(\xi) = 0$ holds for $\xi_1 > 0$, we can estimate K_1 as

$$K_{1} = \int_{\mathbb{R}^{3}} |f_{\infty}(\xi) - f_{\infty}^{k}(\xi)| \frac{|\xi_{1}|}{\sqrt{\xi_{1}^{2} + 2\phi(x)}} d\xi \le \|f_{\infty} - f_{\infty}^{k}\|_{L^{1}(\mathbb{R}^{3})} \to 0 \quad \text{as } k \to \infty,$$

where we have also used the fact $\phi(x) > 0$ in deriving the inequality. Similarly, $K_3 \rightarrow 0$ as $k \to \infty$. Thus K_1 and K_3 can be arbitrarily small for suitably large k. For the fixed k, the dominated convergence theorem ensures that K_2 converges to zero as $x \to x_0$. Hence, we deduce $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$. Now it is straightforward to show $f \in L^1_{loc}(\mathbb{R}_+ \times \mathbb{R}^3)$.

It is clear that the condition (ii), i.e. $f \ge 0$, holds. Let us prove (2.1a) and (2.1b) in the condition (iii). Obviously, (2.1b) follows from the same manner as above. It is also evident that the function f^k defined by replacing f_{∞} by f_{∞}^k in (2.9) belongs to $C_0^{\infty}(\mathbb{R}^3)$, and satisfies the weak form (2.1a) for each k. Using the same change of variable as above and letting $k \to \infty$, we see that f also satisfies (2.1a). Consequently, all the conditions (i)–(iii) hold.

The condition (iv) is validated by (2.9), (3.3), and (3.4). The uniqueness of a solution (f, ϕ) follows from the uniqueness of the solution of (3.7) with (1.1d) and (1.1f). Using (2.5), (2.9), and the fact $\phi \in C(\overline{\mathbb{R}_+}) \cap C^2(\mathbb{R}_+)$, we can show that $f \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3)$ if $f_{\infty} \in C^2(\mathbb{R}^3)$.

The decay estimate (2.10) follows from (2.8), (3.6), and (3.7). The proof is complete. \Box

For the proof of the delta mass limit as $\varepsilon \to 0$, we need to show that sup *B* in Assertion (i) of Theorem 2.2 is bounded from below by some positive constant δ_0 independent of ε for the problem (1.1) with $f_{\infty} = f_{\infty,\varepsilon}$. Here $f_{\infty,\varepsilon}$ is defined in (2.12).

Corollary 3.2. Let $f_b = \alpha = 0$ and $u_{\infty} > 1$. There exists a positive constant δ_0 such that if $0 < \phi_b < \delta_0$, then the problem (1.1) with $f_{\infty} = f_{\infty,\varepsilon}$ has a unique solution $(f_{\varepsilon}, \phi_{\varepsilon}) \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \times C^2(\overline{\mathbb{R}_+})$ for any $\varepsilon \in (0, \varepsilon_0)$, where $\varepsilon_0 := (u_{\infty} - 1)/2$.

Proof. We first note that $f_{\infty,\varepsilon} \in C_0^{\infty}(\mathbb{R}^3)$ satisfies the conditions (2.2), (2.5), and (2.8) being in Theorem 2.2 for any $\varepsilon \in (0, \varepsilon_0)$. It suffices to show that there exists $\delta_0 > 0$ independent of ε such that

$$V_{\varepsilon}(\phi) > 0, \quad \phi \in (0, \delta_0], \tag{3.8}$$

where V_{ε} denotes the function replaced f_{∞} by $f_{\infty,\varepsilon}$ in (2.6). Indeed, repeating the proof of Assertions (i) and (ii) of Theorem 2.2 with the aid of (3.8), we can conclude that the problem (1.1) with $f_{\infty} = f_{\infty,\varepsilon}$ has a unique solution $(f_{\varepsilon}, \phi_{\varepsilon}) \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \times C^2(\overline{\mathbb{R}_+})$ for $\phi_b \in (0, \delta_0)$ and $\varepsilon \in (0, \varepsilon_0)$.

Let us complete the proof by showing (3.8). Using $\int_{\mathbb{R}^3} f_{\infty,\varepsilon}(\xi) d\xi = 1$, we observe that

$$\begin{aligned} \left| \frac{d^{3}V_{\varepsilon}}{d\phi^{3}}(\phi) \right| &= \left| 3 \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) \frac{-\xi_{1}}{(\xi_{1}^{2}+2\phi)^{5/2}} d\xi - n_{e}''(\phi) \right| \leq 3 \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) |\xi_{1}|^{-4} d\xi + C_{0} \\ &\leq 3 (\sup_{|\xi_{1}+u_{\infty}|\leq\varepsilon_{0}} |\xi_{1}|^{-4}) \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) d\xi + C_{0} \leq C_{0} \end{aligned}$$

for any $\phi \in (0,1]$, where C_0 is independent of ε . Recalling (3.6), we also see that

$$\frac{d^2 V_{\varepsilon}}{d\phi^2}(0) = -\int_{\mathbb{R}^3} f_{\infty,\varepsilon}(\xi) |\xi_1|^{-2} d\xi + 1 \ge -(\sup_{|\xi_1+u_{\infty}| \le \varepsilon_0} |\xi_1|^{-2}) + 1 > 0,$$

where we have used $\varepsilon_0 = (u_{\infty} - 1)/2$ and $u_{\infty}^2 > 1$. By the Taylor theorem and the above two inequalities, we conclude that there exists δ_0 independent of ε such that

$$\frac{d^2 V_{\varepsilon}}{d\phi^2}(\phi) \ge -(\sup_{|\xi_1+u_{\infty}| \le \varepsilon_0} |\xi_1|^{-2}) + 1 - C_0 \delta_0 > 0, \quad \phi \in [0, \delta_0],$$

which together with (3.6) leads to (3.8). The proof is complete.

3.3 The Delta Mass Limit

This section deals with the proof of Theorem 2.4 on the delta mass limit.

Proof of Theorem 2.4. Let $(f_{\varepsilon}, \phi_{\varepsilon})$ be a solution in Corollary 3.2, and also $(\rho_{\varepsilon}, u_{\varepsilon})$ be the moments of f_{ε} . We first show the estimate of $\phi_{\varepsilon} - \phi$ in (2.15). From (1.5a), (2.14), (3.3), and (3.4), we observe that

$$\partial_{xx}(\phi_{\varepsilon} - \phi) = W(\phi_{\varepsilon}) - W(\phi) + R_{\varepsilon}(x), \quad x > 0, \tag{3.9}$$

where

$$W(\phi) := \frac{u_{\infty}}{\sqrt{u_{\infty}^2 + 2\phi}} - e^{-\phi}, \quad R_{\varepsilon}(x) := \int_{\mathbb{R}^3} f_{\infty,\varepsilon}(\xi) \frac{-\xi_1}{\sqrt{\xi_1^2 + 2\phi_{\varepsilon}(x)}} d\xi - \frac{u_{\infty}}{\sqrt{u_{\infty}^2 + 2\phi_{\varepsilon}(x)}}.$$

Set $M_{\varepsilon} := \sup_{x \in \mathbb{R}_+} |R_{\varepsilon}(x)|$. Owing to the assumption $u_{\infty} > 1$, we have $dW/d\phi(0) > 0$. Then it is seen by taking $\delta_0 > 0$ and $c_0 > 0$ small enough that

$$c_0 \le \frac{dW}{d\phi}(\phi) \le c_0^{-1}, \quad \phi \in [0, \delta_0].$$
 (3.10)

Now we claim that

$$\sup_{x \in \mathbb{R}_+} |(\phi_{\varepsilon} - \phi)(x)| \le M_{\varepsilon}/c_0.$$
(3.11)

To show this, we observe from (3.9) that

$$-\partial_{xx}(\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0) + (W(\phi_{\varepsilon}) - W(\phi) - M_{\varepsilon}) \le 0.$$
(3.12)

Furthermore, $(\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+ := \max\{0, \phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0\} \in H^1(\mathbb{R}_+)$ follows from (1.1f) and (1.5b). Multiply (3.12) by $(\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+$, integrate it over $(0,\infty)$, and use $(\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+(0) = 0$ to obtain

$$\|\partial_x(\phi_{\varepsilon}-\phi-M_{\varepsilon}/c_0)_+\|_{L^2(\mathbb{R}_+)}^2 + \int_0^\infty (W(\phi_{\varepsilon})-W(\phi)-M_{\varepsilon})(\phi_{\varepsilon}-\phi-M_{\varepsilon}/c_0)_+dx \le 0.$$
(3.13)

The second term on the left hand side is positive. Indeed, using the mean value theorem, (3.10), and $0 \le \phi, \phi_{\varepsilon} \le \delta_0$, we arrive at

$$\int_0^\infty (W(\phi_{\varepsilon}) - W(\phi) - M_{\varepsilon}) (\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+ dx$$

=
$$\int_0^\infty \left(\int_0^1 \frac{dW}{d\phi} (\theta \phi_{\varepsilon} + (1 - \theta)\phi) d\theta (\phi_{\varepsilon} - \phi) - M_{\varepsilon} \right) (\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+ dx \ge 0.$$

Therefore, (3.13) gives $(\phi_{\varepsilon} - \phi - M_{\varepsilon}/c_0)_+ = 0$ which means $\phi_{\varepsilon} - \phi \leq M_{\varepsilon}/c_0$. Similarly, $\phi_{\varepsilon} - \phi \geq -M_{\varepsilon}/c_0$ holds. Thus the claim (3.11) is valid.

Next let us estimate M_{ε} by $C_0 \varepsilon$ as

$$M_{\varepsilon} = \sup_{x \in \mathbb{R}_{+}} \left| \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) \left(\frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{u_{\infty}}{\sqrt{u_{\infty}^{2} + 2\phi_{\varepsilon}(x)}} \right) d\xi \right|$$

$$\leq \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (-u_{\infty}, 0, 0)| \le \varepsilon} \left| \frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{u_{\infty}}{\sqrt{u_{\infty}^{2} + 2\phi_{\varepsilon}(x)}} \right| \right) \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) d\xi$$

$$\leq C_{0}\varepsilon,$$

where we have used the fact $\int_{\mathbb{R}^3} f_{\infty,\varepsilon}(\xi) d\xi = 1$ in deriving the equality, and C_0 is a positive constant independent of ε . This together with (3.11) leads to the estimate of $\phi_{\varepsilon} - \phi$ in (2.15).

We complete the proof by showing the other estimates in (2.15). Let us first handle $\rho_{\varepsilon} - \rho$. It is clear that $\rho_{\varepsilon}(x) = \rho_i(\phi_{\varepsilon}(x))$ holds owing to (3.3). With the aid of this and (2.14), we estimate $\rho_{\varepsilon} - \rho$ as

$$\begin{split} \sup_{x\in\mathbb{R}_{+}} |\rho_{\varepsilon}(x) - \rho(x)| &= \sup_{x\in\mathbb{R}_{+}} \left| \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) \left(\frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{u_{\infty}}{\sqrt{u_{\infty}^{2} + 2\phi(x)}} \right) d\xi \right| \\ &\leq \sup_{x\in\mathbb{R}_{+}} \left(\sup_{|\xi - (-u_{\infty},0,0)| \le \varepsilon} \left| \frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{u_{\infty}}{\sqrt{u_{\infty}^{2} + 2\phi(x)}} \right| \right) \int_{\mathbb{R}^{3}} f_{\infty,\varepsilon}(\xi) d\xi \\ &\leq C_{0}(\varepsilon + \sup_{x\in\mathbb{R}_{+}} |\phi(x) - \phi_{\varepsilon}(x)|) \le C_{0}\varepsilon. \end{split}$$

Let us treat $\rho_{\varepsilon}u_{\varepsilon} - \rho u$. By the change of variable $\sqrt{\xi_1^2 - 2\phi(x)} = -\zeta_1$, the term $\rho_{\varepsilon}u_{\varepsilon}$ is rewritten as

$$(\rho_{\varepsilon}u_{\varepsilon})(x) = \int_{\mathbb{R}^3} \xi_1 f_{\infty,\varepsilon}(\xi) d\xi$$

It is easy to see $\rho u = -u_{\infty}$ from (2.14). Then we arrive at

$$\sup_{x\in\mathbb{R}_+}|(\rho_{\varepsilon}u_{\varepsilon})(x)-(\rho u)(x)|=\sup_{x\in\mathbb{R}_+}\left|\int_{\mathbb{R}^3}f_{\infty,\varepsilon}(\xi)(\xi_1+u_{\infty})d\xi\right|\leq\varepsilon.$$

Consequently, (2.15) holds.

4 General Boundaries

In this section, we prove Theorems 2.5–2.7 on the solvability and delta mass limit for general boundaries. Subsections 4.1 and 4.2 deal with the solvability for the attractive and repulsive boundaries, respectively. In subsection 4.3, we study the delta mass limit.

4.1 Solvability for the Attractive Boundary

We start from studying the necessary conditions for the solvability of the problem (1.1) similarly as in Section 3. Specifically, we show the following lemma which ensures immediately Assertion (iv) in Theorem 2.5.

Lemma 4.1. Let $\phi_b > 0$, $f_b \in L^1(\mathbb{R}^3_+)$, $\alpha \neq 1$, $f_\infty \in L^1(\mathbb{R}^3)$, $f_b \ge 0$, $f_\infty \ge 0$, and (2.2) hold. Suppose that the problem (1.1) has a solution (f, ϕ) . Then $\rho^+ \in C([0, \phi_b])$, $V^+ \in C^1([0, \phi_b])$, and the function f_∞ satisfies the condition (2.3). Furthermore, ϕ solves (4.4) below with (1.1d) and (1.1f), and f is written as (2.22) by ϕ . If $V^+ \in C^2([0, \phi_b])$, the generalized Bohm criterion holds:

$$\frac{d^2 V^+}{d\phi^2}(0) \ge 0. \tag{4.1}$$

Proof. In much the same way as in the proof of Lemma 3.1, we see $\partial_x \phi(x) < 0$ and $\lim_{x\to\infty} \partial_x \phi(x) = 0$. Let us first show that f_{∞} satisfies (2.3) and f is written by (2.22). Applying the characteristics method to (2.1a), we see that the value of f must remain a constant along the characteristic curves $\xi_1^2/2 - \phi(x) = c$, where c is some constant. The characteristics are drawn as in Figure 1. Therefore, there hold that

$$f(y, -\sqrt{\eta_1^2 + 2\phi(y)}, \eta') = f_{\infty}(\eta), \quad (y, \eta_1, \eta') \in X_1,$$

$$f(y, \pm\sqrt{2\phi(y) - \eta_1^2}, \eta') = \frac{1}{1 - \alpha} f_b(\sqrt{2\phi_b - \eta_1^2}, \eta'), \quad (y, \eta_1, \eta') \in X_2,$$

$$f(y, \sqrt{\eta_1^2 + 2\phi(y)}, \eta') = f_{\infty}(\eta) = f_b(\sqrt{\eta_1^2 + 2\phi_b}, \eta') + \alpha f_{\infty}(-\eta_1, \eta'), \quad (y, \eta_1, \eta') \in X_3,$$

where $\boldsymbol{\eta} = (\eta_1, \eta_2, \eta_3) = (\eta_1, \eta')$ and

$$X_1 := \overline{\mathbb{R}_+} \times \mathbb{R}_-^3, \quad X_2 := \{ (y, \eta_1, \eta') \in \overline{\mathbb{R}_+} \times \mathbb{R}^3 \mid \eta_1^2 < 2\phi(y) \le 2\phi_b \}, \quad X_3 := \overline{\mathbb{R}_+} \times \mathbb{R}_+^3.$$

The last equality means that (2.3) must hold. Furthermore, we conclude from these three equalities 9 that *f* must be written by (2.22), i.e.

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_{1}^{2} - 2\phi(x)},\xi')\chi(\xi_{1}^{2} - 2\phi(x))\chi(-\xi_{1}) + \frac{1}{1 - \alpha}f_{b}(\sqrt{\xi_{1}^{2} - 2\phi(x) + 2\phi_{b}},\xi')\chi(-\xi_{1}^{2} + 2\phi(x)) + f_{\infty}(\sqrt{\xi_{1}^{2} - 2\phi(x)},\xi')\chi(\xi_{1}^{2} - 2\phi(x))\chi(\xi_{1}).$$

Next we show that ϕ satisfies (4.4) below. Integrating (2.22) over \mathbb{R}^3 and using the change of variables $\sqrt{\xi_1^2 - 2\phi} = -\zeta_1$, $\sqrt{\xi_1^2 - 2\phi + 2\phi_b} = \zeta_1$, and $\sqrt{\xi_1^2 - 2\phi} = \zeta_1$ for the first, second,

⁹If $\alpha = 1$ and $f_b = 0$, we have multiple choices for the second term on the right hand side of (2.22).

and third terms on the right hand side of (2.22), respectively, we see that

$$\begin{aligned} \int_{\mathbb{R}^3} f(x,\xi) d\xi &= \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi(x)}} d\xi \\ &+ \frac{2}{1-\alpha} \int_{\sqrt{2(\phi_b - \phi(x))}}^{\sqrt{2\phi_b}} \int_{\mathbb{R}^2} f_b(\xi) \frac{\xi_1}{\sqrt{\xi_1^2 + 2\phi(x) - 2\phi_b}} d\xi_1 d\xi' \\ &= \rho_i^+(\phi(x)), \end{aligned}$$
(4.2)

where $\phi(x) \in [0, \phi_b]$ and ρ_i^+ is the same function defined in (2.18). Substituting (4.2) into (1.1b) yields an ordinary differential equation for ϕ :

$$\partial_{xx}\phi = \rho_i^+(\phi) - n_e(\phi), \quad x > 0.$$
(4.3)

Multiply (4.3) by $\partial_x \phi$, integrate it over (x, ∞) , and use (1.1f) and $\lim_{x\to\infty} \partial_x \phi(x) = 0$ to obtain

$$(\partial_x \phi)^2 = 2V^+(\phi), \quad V^+(\phi) = \int_0^\phi \left(\rho_i^+(\phi) - n_e(\phi)\right) d\phi, \tag{4.4}$$

where the function V^+ is the same function defined in (2.16). Thus ϕ must satisfy (4.4).

Now we show $\rho_i^+ \in C([0, \phi_b])$ which immediately gives $V^+ \in C^1([0, \phi_b])$. Owing to $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$ in Definition 2.1 and (2.1b), there hold that

$$\rho_i^+(\phi(x)) = \|f(x)\|_{L^1} \in C(\overline{\mathbb{R}_+}), \quad \lim_{x \to \infty} \|f(x)\|_{L^1} = \|f_\infty\|_{L^1} = 1.$$

These imply $\rho_i^+ \in C([0, \phi_b])$ with the aid of (1.1f), $\phi \in C(\overline{\mathbb{R}_+})$, and $\partial_x \phi < 0$ in Definition 2.1.

What is left is to obtain the generalized Bohm criterion (4.1). We see from $n_e(0) = 1$ and (2.2) that

$$V^{+}(0) = \frac{dV^{+}}{d\phi}(0) = 0.$$
(4.5)

If $V^+ \in C^2([0, \phi_b])$, we arrive at (4.1) from (4.4) and (4.5) with the aid of the Taylor theorem. The proof is complete.

Assertion (iii) in Theorem 2.5 can be shown by the same method as in the proof of Assertion (iii) in Theorem 2.2 with the aid of (4.4). We omit the proof. Let us show Assertions (i) and (ii) in Theorem 2.5.

Proof of Assertions (i) and (ii) in Theorem 2.5. We first note that (4.5) holds. It is seen that $B^+ \neq \emptyset$ if $d^2V^+/d\phi^2(0) > 0$, since the Taylor theorem with (4.5) ensures that V^+ is positive around $\phi = 0$. Let us show that $\phi_b < \sup B^+$ is a necessary condition for the solvability stated in Assertions (i) and (ii). Suppose that a solution (f, ϕ) exists for $\phi_b \ge \sup B^+$. Then $\sup B^+ \in (0, \phi_b]$ and $V^+(\sup B^+) = 0$. Furthermore, there exists a unique $x_* \in \mathbb{R}_+$ so that $\phi(x_*) = \sup B^+$. On the other hand, due to (4.4), it is seen that $(\partial_x \phi(x_*))^2 = 2V^+(\sup B^+) = 0$. This contradicts to $\partial_x \phi(x) < 0$ in Definition 2.1. Hence, $\phi_b < \sup B^+$ is necessary.

Hereafter we discuss simultaneously Assertions (i) and (ii). Suppose that $\phi_b < \sup B^+$ holds, which implies that $V^+(\phi) > 0$ for $\phi \in (0, \phi_b]$. The Taylor theorem with (4.5) and the assumption

 $d^2V^+/d\phi^2(0) \ge 0$ ensures that $\sqrt{V^+}$ is Lipschitz continuous around $\phi = 0$. Combining this and the fact that $V^+(\phi) > 0$ for $\phi \in (0, \phi_b]$, we deduce that $\sqrt{V^+}$ is Lipschitz continuous on $[0, \phi_b]$. Hence, we can define ϕ by solving (4.4) with (1.1d) and (1.1f). Then it is seen that ϕ satisfies $\partial_x \phi < 0$, $\phi \in C(\overline{\mathbb{R}_+}) \cap C^2(\mathbb{R}_+)$, and (4.3) by following the proof of Assertions (i) and (ii) of Theorem 2.2 in subsection 3.2.

Now by using ϕ , we define f as (2.22), i.e.

$$\begin{split} f(x,\xi) &= f_{\infty}(-\sqrt{\xi_{1}^{2} - 2\phi(x),\xi'})\chi(\xi_{1}^{2} - 2\phi(x))\chi(-\xi_{1}) \\ &+ \frac{1}{1 - \alpha}f_{b}(\sqrt{\xi_{1}^{2} - 2\phi(x) + 2\phi_{b}},\xi')\chi(-\xi_{1}^{2} + 2\phi(x)) \\ &+ f_{\infty}(\sqrt{\xi_{1}^{2} - 2\phi(x)},\xi')\chi(\xi_{1}^{2} - 2\phi(x))\chi(\xi_{1}) \\ &=:F_{1}(x,\xi) + F_{2}(x,\xi) + F_{3}(x,\xi), \end{split}$$

and prove that f satisfies the conditions (i)–(iii) in Definition 2.1. Owing to (2.3), $f_b \in L^1(\mathbb{R}^3_+) \cap L^r(0, \sqrt{2\phi_b}; L^1(\mathbb{R}^2))$, and $f_{\infty} \in L^1(\mathbb{R}^3)$, there exist sequences $\{f_{\infty}^k\}, \{f_b^k\} \subset C_0^{\infty}(\mathbb{R}^3)$ such that (2.3) with $(f_{\infty}, f_b) = (f_{\infty}^k, f_b^k)$ holds; $f_{\infty}^k(\xi) = 0$ holds for $|\xi_1| < 1/k$; $f_b^k(\xi) = 0$ holds for $|\xi_1 - \sqrt{2\phi_b}| < 1/k$; $f_{\infty}^k \to f_{\infty}$ in $L^1(\mathbb{R}^3)$ as $k \to \infty$; $f_b^k \to f_b$ in $L^1(\mathbb{R}^3_+) \cap L^r(0, \sqrt{2\phi_b}; L^1(\mathbb{R}^2))$ as $k \to \infty$. Let us show the condition (i), i.e. $f \in C(\mathbb{R}_+; L^1(\mathbb{R}^3)) \cap L_{loc}^1(\mathbb{R}_+ \times \mathbb{R}^3)$. It is seen that $F_1, F_3 \in C(\mathbb{R}_+; L^1(\mathbb{R}^3))$ in much the same way as the proof of Assertions (i) and (ii) of Theorem 2.2. The fact $F_2(x, \cdot) \in L^1(\mathbb{R}^3)$ follows from (2.20) and (4.2). To investigate the continuity of F_2 , we set $G(x, \xi) = \chi(-\xi_1^2 + 2\phi(x))$ and observe that for $x, x_0 \in \mathbb{R}_+$,

$$\begin{aligned} &(1-\alpha)\|F_{2}(x)-F_{2}(x_{0})\|_{L^{1}(\mathbb{R}^{3})} \\ &= \int_{\mathbb{R}^{3}} \left| f_{b}(\sqrt{\xi_{1}^{2}-2\phi(x)+2\phi_{b}},\xi')G(x,\xi) - f_{b}(\sqrt{\xi_{1}^{2}-2\phi(x_{0})+2\phi_{b}},\xi')G(x_{0},\xi) \right| d\xi \\ &\leq \int_{\mathbb{R}^{3}} \left| f_{b}(\sqrt{\xi_{1}^{2}-2\phi(x)+2\phi_{b}},\xi')G(x,\xi) - f_{b}^{k}(\sqrt{\xi_{1}^{2}-2\phi(x)+2\phi_{b}},\xi')G(x,\xi) \right| d\xi \\ &+ \int_{\mathbb{R}^{3}} \left| f_{b}^{k}(\sqrt{\xi_{1}^{2}-2\phi(x)+2\phi_{b}},\xi')G(x,\xi) - f_{b}^{k}(\sqrt{\xi_{1}^{2}-2\phi(x_{0})+2\phi_{b}},\xi')G(x_{0},\xi) \right| d\xi \\ &+ \int_{\mathbb{R}^{3}} \left| f_{b}(\sqrt{\xi_{1}^{2}-2\phi(x_{0})+2\phi_{b}},\xi')G(x_{0},\xi) - f_{b}(\sqrt{\xi_{1}^{2}-2\phi(x_{0})+2\phi_{b}},\xi')G(x_{0},\xi) \right| d\xi \\ &=: K_{1}+K_{2}+K_{3}. \end{aligned}$$

Using the change of variable $\zeta_1 = \sqrt{\xi_1^2 - 2\phi(x) + 2\phi_b}$, we can estimate K_1 as

$$\begin{split} K_{1} &= 2 \int_{\sqrt{2(\phi_{b} - \phi(x))}}^{\sqrt{2\phi_{b}}} \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi(x) - 2\phi_{b}}} \left(\int_{\mathbb{R}^{2}} |f_{b}(\xi) - f_{b}^{k}(\xi)| d\xi' \right) d\xi_{1} \\ &\leq 2 \sup_{x \in \mathbb{R}_{+}} \left(\int_{\sqrt{2(\phi_{b} - \phi(x))}}^{\sqrt{2\phi_{b}}} \frac{\xi_{1}^{r'}}{(\xi_{1}^{2} + 2\phi(x) - 2\phi_{b})^{r'/2}} d\xi_{1} \right)^{1/r'} \|f_{b} - f_{b}^{k}\|_{L^{r}(0,\sqrt{2\phi_{b}};L^{1}(\mathbb{R}^{2}))} \\ &\leq C \|f_{b} - f_{b}^{k}\|_{L^{r}(0,\sqrt{2\phi_{b}};L^{1}(\mathbb{R}^{2}))} \to 0 \quad \text{as } k \to \infty, \end{split}$$

where r' < 2 is the Hölder conjugate of r > 2, and *C* is some positive constant. Similarly, $K_3 \to 0$ as $k \to \infty$. Thus K_1 and K_3 can be arbitrarily small for suitably large *k*. For the fixed *k*, the dominated convergence theorem ensures that K_2 converges to zero as $x \to x_0$. Hence, we deduce that $F_2 \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$ holds and so does $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$. Now it is straightforward to show $f \in L^1_{loc}(\mathbb{R}_+ \times \mathbb{R}^3)$.

It is clear that the condition (ii), i.e. $f \ge 0$, holds. Let us prove (2.1a) and (2.1b) in the condition (iii). Obviously, (2.1b) follows from the same manner as above. It is also evident that the function f^k defined by replacing (f_{∞}, f_b) by (f_{∞}^k, f_b^k) in (2.22) belongs to $C_0^{\infty}(\mathbb{R}^3)$, and satisfies the weak form (2.1a) for each k. Letting $k \to \infty$, we see that f also satisfies (2.1a). Consequently, all the conditions (i)–(iii) hold.

The condition (iv) is validated by (2.22), (4.2), and (4.3). The uniqueness of a solution (f, ϕ) follows from the uniqueness of the solution of (4.4) with (1.1d) and (1.1f). We have the decay estimate (2.23) from (4.4), (4.5), and $d^2V^+/d\phi^2(0) > 0$. The proof is complete.

We also have Corollary 4.2 stating that $\sup B^+$ in Assertion (i) of Theorem 2.5 is bounded from below by some positive constant δ_0 independent of ε and ϕ_b for the problem (1.1) with $(f_b, f_\infty) = (g_{b,\varepsilon}, g_{\infty,\varepsilon})$, where $g_{b,\varepsilon}$ and $g_{\infty,\varepsilon}$ are defined in (2.26) and (2.27). As mentioned just after (2.28), the functions $g_{b,\varepsilon}$ and $g_{\infty,\varepsilon}$ satisfy the conditions (2.2), (2.3), and $d^2V^+/d\phi^2(0) > 0$ being in Theorem 2.5 for $|\phi_b| \ll 1$ and $\varepsilon \ll 1$ provided that (2.28) holds. Furthermore, B^+ is independent of ϕ_b , and also $\phi_b < \sup B^+$ holds for $\phi_b \ll 1$. We omit the proof of Corollary 4.2, since it is the same as the proof of Corollary 3.2.

Corollary 4.2. Let $\alpha \neq 1$ and (2.28) hold. There exist positive constants ε_0 and δ_0 such that if $0 < \phi_b < \delta_0$, then the problem (1.1) with $(f_b, f_\infty) = (g_{b,\varepsilon}, g_{\infty,\varepsilon})$ has a unique solution $(f_{\varepsilon}, \phi_{\varepsilon}) \in C^1(\overline{\mathbb{R}_+} \times \mathbb{R}^3) \times C^2(\overline{\mathbb{R}_+})$ for any $\varepsilon \in (0, \varepsilon_0)$.

4.2 Solvability for the Repulsive Boundary

We first study the necessary conditions for the solvability of the problem (1.1) similarly as in subsection 4.1. Specifically, we show the following lemma which ensures immediately Assertion (iv) in Theorem 2.6.

Lemma 4.3. Let $\phi_b < 0$, $f_b \in L^1(\mathbb{R}^3_+)$, $f_\infty \in L^1(\mathbb{R}^3)$, $f_b \ge 0$, $f_\infty \ge 0$, and (2.2) hold. Suppose that the problem (1.1) has a solution (f, ϕ) . Then $\rho^- \in C([\phi_b, 0])$, $V^- \in C^1([\phi_b, 0])$, and the function f_∞ satisfies the condition (2.4). Furthermore, ϕ solves (4.9) below with (1.1d) and (1.1f), and f is written as (2.24) by ϕ . If $V^- \in C^2([\phi_b, 0])$, the generalized Bohm criterion also holds:

$$\frac{d^2 V^-}{d\phi^2}(0) \ge 0. \tag{4.6}$$

Proof. In the same manner as in subsection 3.1, we see $\partial_x \phi(x) > 0$ and $\lim_{x\to\infty} \partial_x \phi(x) = 0$. Let us first show that f_{∞} satisfies (2.4) and f is written by (2.24). Applying the characteristics method to (2.1a), we see that the value of f must remain a constant along the characteristic curves $\xi_1^2/2 - \phi(x) = c$, where c is some constant. We draw the illustration of characteristics in Figure 2 below. It tells us that Figure 2: characteristics curves for the case $\partial_x \phi > 0$

$$\begin{aligned} f(y, -\sqrt{\eta_1^2 + 2\phi(y)}, \eta') &= f_{\infty}(\eta), \quad (y, \eta_1, \eta') \in Y_1, \\ f(y, \pm\sqrt{\eta_1^2 + 2\phi(y)}, \eta') &= f_{\infty}(\eta), \quad (y, \eta_1, \eta') \in Y_2, \\ f(y, \sqrt{\eta_1^2 + 2\phi(y)}, \eta') &= f_{\infty}(\eta) = f_b(\sqrt{\eta_1^2 + 2\phi_b}, \eta') + \alpha f_{\infty}(-\eta_1, \eta'), \quad (y, \eta_1, \eta') \in Y_3, \end{aligned}$$

where

$$\begin{split} Y_1 &:= \overline{\mathbb{R}_+} \times (-\infty, -\sqrt{2|\phi_b|}) \times \mathbb{R}^2, \\ Y_2 &:= \{ (y, \eta_1, \eta') \in \overline{\mathbb{R}_+} \times \mathbb{R}^3 \mid 2|\phi(y)| < \eta_1^2 < 2|\phi_b| \}, \\ Y_3 &:= \overline{\mathbb{R}_+} \times (\sqrt{2|\phi_b|}, \infty) \times \mathbb{R}^2. \end{split}$$

The second and third equalities mean that (2.4) must hold. Furthermore, we conclude from these three equalities that f must be written by (2.24), i.e,

$$f(x,\xi) = f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(x)},\xi')\chi(-\xi_1) + f_{\infty}(\sqrt{\xi_1^2 - 2\phi(x)},\xi')\chi(\xi_1).$$

Next we show that ϕ satisfies (4.9) below. Integrating (2.24) over \mathbb{R}^3 and using the change of variables $\sqrt{\xi_1^2 - 2\phi} = -\zeta_1$ and $\sqrt{\xi_1^2 - 2\phi} = \zeta_1$ for the first and second terms of the right hand side of (2.24), respectively, we see that

$$\int_{\mathbb{R}^3} f(x,\xi) d\xi = \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi(x)}} \chi(\xi_1^2 + 2\phi(x)) d\xi = \rho_i^-(\phi(x)), \quad (4.7)$$

where ρ_i^- is the same function defined in (2.19). Substituting (4.7) into (1.1b) yields an ordinary differential equation for ϕ :

$$\partial_{xx}\phi = \rho_i^-(\phi) - n_e(\phi), \quad x > 0.$$
(4.8)

Multiply (4.8) by $\partial_x \phi$, integrate it over (x, ∞) , and use (1.1f) and $\lim_{x\to\infty} \partial_x \phi(x) = 0$ to obtain

$$(\partial_x \phi)^2 = 2V^-(\phi), \quad V^-(\phi) = \int_0^\phi \left(\rho_i^-(\phi) - n_e(\phi) \right) d\phi, \tag{4.9}$$

where the function V^- is the same function defined in (2.17). Thus ϕ must satisfy (4.9).

Now we show $\rho_i^- \in C([\phi_b, 0])$ which immediately gives $V^- \in C^1([\phi_b, 0])$. Owing to $f \in C(\overline{\mathbb{R}_+}; L^1(\mathbb{R}^3))$ and (2.1b) in Definition 2.1, there hold that

$$\rho_i^-(\phi(x)) = \|f(x)\|_{L^1} \in C(\overline{\mathbb{R}_+}), \quad \lim_{x \to \infty} \|f(x)\|_{L^1} = \|f_\infty\|_{L^1} = 1.$$

These imply $\rho_i^- \in C([\phi_b, 0])$ with the aid of (1.1f), $\phi \in C(\overline{\mathbb{R}_+})$, and $\partial_x \phi > 0$ in Definition 2.1.

What is left is to obtain the generalized Bohm criterion (4.6). We see from $n_e(0) = 1$ and (2.2) that

$$V^{-}(0) = \frac{dV^{-}}{d\phi}(0) = 0.$$
(4.10)

If $V^- \in C^2([\phi_b, 0])$, we arrive at (4.6) from (4.9) and (4.10) with the aid of the Taylor theorem. The proof is complete.

Assertion (iii) in Theorem 2.6 can be shown by the same method as in the proof of Assertion (iii) in Theorem 2.2 with the aid of (4.9). We omit the proof. Let us show Assertions (i) and (ii) in Theorem 2.6.

Proof of Assertions (i) and (ii) in Theorem 2.6. In the same way as in the proof of Assertions (i) and (ii) in Theorem 2.5, we first see that $B^- \neq \emptyset$ if $d^2V^-/d\phi^2(0) > 0$; $\phi_b > \inf B^-$ is necessary; $\sqrt{V^-}$ is Lipschitz continuous on $[\phi_b, 0]$. To find a solution ϕ with $\partial_x \phi > 0$ of the equation (4.9) with (1.1d) and (1.1f), we rewrite (4.9) into the equivalent equation

$$\partial_x \phi = \sqrt{2V^-(\phi)}.$$

Then we complete the proof in the same manner as that of Assertions (i) and (ii) of Theorems 2.2 and 2.5. \Box

Similarly as the proof of Corollary 3.2, we also see that $\inf B^-$ in Assertion (i) of Theorem 2.6 is bounded from above by some negative constant $-\delta_0$ independent of ε for the problem (1.1) with $(f_b, f_\infty) = (g_{b,\varepsilon}, g_{\infty,\varepsilon})$. Here $g_{b,\varepsilon}$ and $g_{\infty,\varepsilon}$ defined in (2.26) and (2.27) satisfy the conditions (2.2), (2.4), and $d^2V^-/d\phi^2(0) > 0$ being in Theorem 2.6 for $|\phi_b| \ll 1$ and $\varepsilon \ll 1$ provided that (2.28) holds. The result is summarized in the following corollary.

Corollary 4.4. Let (2.28) hold. There exist positive constants ε_0 and δ_0 such that if $-\delta_0 < \phi_b < 0$, then the problem (1.1) with $(f_b, f_\infty) = (g_{b,\varepsilon}, g_{\infty,\varepsilon})$ has a unique solution $(f_{\varepsilon}, \phi_{\varepsilon}) \in C^1(\mathbb{R}_+ \times \mathbb{R}^3) \times C^2(\mathbb{R}_+)$ for any $\varepsilon \in (0, \varepsilon_0)$.

4.3 The Delta Mass Limit

This section provides the proof of Theorem 2.7 on the delta mass limit. We start from showing the next lemma on the solvability of the problem (2.30).

Lemma 4.5. Let (2.28) hold. There exists a positive constant δ_0 such that if $|\phi_b| < \delta_0$, then the problem (2.30) has a unique monotone solution $\phi \in C^2(\overline{\mathbb{R}_+})$.

Proof. We first note that $\lim_{x\to\infty} \partial_x \phi(x) = 0$ must hold if such a solution ϕ exists. Now multiply (2.30a) by $\partial_x \phi$ and integrate it over (x, ∞) to obtain

$$(\partial_x \phi)^2 = 2\tilde{V}(\phi), \quad \tilde{V}(\phi) := \int_0^\phi \frac{m_b v_b}{\sqrt{v_b^2 + 2\varphi}} + \frac{(1+\alpha)m_\infty v_\infty}{\sqrt{v_\infty^2 + 2\varphi}} - n_e(\varphi) d\varphi$$

It is straightforward to see that $\tilde{V} \in C^2([-\delta_0, \delta_0])$ holds for some $\delta_0 > 0$. From $n_e(0) = 1$, $n'_e(0) = -1$, and (2.28), one can know that

$$ilde{V}(0)=rac{d ilde{V}}{d\phi}(0)=0,\quad rac{d^2 ilde{V}}{d\phi^2}(0)>0.$$

Then following the proofs of Assertions (i) and (ii) of Theorems 2.2, 2.5, and 2.6, we can complete the proof. \Box

We are now in a position to prove Theorem 2.7.

Proof of Theorem 2.7. Let $(f_{\varepsilon}, \phi_{\varepsilon})$ be solutions in Corollaries 4.2 and 4.4, and also $(\rho_{\varepsilon}, u_{\varepsilon})$ be the moments of f_{ε} . We first show the estimate of $\phi_{\varepsilon} - \phi$ in (2.29). From (2.30a), (4.3), and (4.8), we observe that

$$\partial_{xx}(\phi_{\varepsilon}-\phi)=W(\phi_{\varepsilon})-W(\phi)+R_{\varepsilon}(x), \quad x>0,$$

where we have used the fact that the second term vanishes in (2.20) for $f_b = g_{b,\varepsilon}$ and $0 < \phi_b \ll 1$, and $W(\phi)$ and $R_{\varepsilon}(x)$ are defined as

$$\begin{split} W(\phi) &:= \frac{m_b v_b}{\sqrt{v_b^2 + 2\phi}} + \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^2 + 2\phi}} - n_e(\phi), \\ R_{\varepsilon}(x) &:= \int_{\mathbb{R}^3} g_{\infty,\varepsilon}(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi_{\varepsilon}(x)}} d\xi - \frac{m_b v_b}{\sqrt{v_b^2 + 2\phi_{\varepsilon}(x)}} - \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^2 + 2\phi_{\varepsilon}(x)}}. \end{split}$$

Set $M_{\varepsilon} := \sup_{x \in \mathbb{R}_+} |R_{\varepsilon}(x)|$. Owing to $n'_{\varepsilon}(0) = -1$ and (2.28), we have $dW/d\phi(0) > 0$. Then it is seen by taking $\delta_0 > 0$ and $c_0 > 0$ small enough that

$$c_0 \leq rac{dW}{d\phi}(\phi) \leq c_0^{-1}, \quad \phi \in [-\delta_0, \delta_0].$$

Now following the proof of (3.11), we arrive at

$$\sup_{x \in \mathbb{R}_+} |(\phi_{\varepsilon} - \phi)(x)| \le M_{\varepsilon}/c_0.$$
(4.11)

For sufficiently small $\varepsilon > 0$, let us estimate M_{ε} by $C_0 \varepsilon$ as

$$\begin{split} M_{\varepsilon} &= \sup_{x \in \mathbb{R}_{+}} \left| \int_{\mathbb{R}^{3}} g_{\infty,\varepsilon}(\xi) \left(\frac{|\xi_{1}|}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{m_{b}v_{b}}{\sqrt{v_{b}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi_{\varepsilon}(x)}} \right) d\xi \right| \\ &\leq \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (-v_{\infty}, 0, 0)| \leq \varepsilon} \left| \frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi_{\varepsilon}(x)}} \right| \right) m_{\infty} \right. \\ &+ \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (v_{b}, 0, 0)| \leq \varepsilon} \left| \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{b}}{\sqrt{v_{b}^{2} + 2\phi_{\varepsilon}(x)}} \right| \right) m_{b} \right. \\ &+ \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (v_{\infty}, 0, 0)| \leq \varepsilon} \left| \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi_{\varepsilon}(x)}} \right| \right) \alpha m_{\infty} \right. \\ &\leq C_{0}\varepsilon, \end{split}$$

where we have used the fact $\int_{\mathbb{R}^3} g_{\infty,\varepsilon}(\xi) d\xi = m_b + (1+\alpha)m_{\infty} = 1$ in deriving the equality, and C_0 is a positive constant independent of ε . This together with (4.11) leads to the estimate of $\phi_{\varepsilon} - \phi$ in (2.29).

We complete the proof by showing the other estimates in (2.29). It is clear that $\rho_{\varepsilon}(x) = \rho_i^{\pm}(\phi_{\varepsilon}(x))$ holds owing to (4.2) and (4.7). We estimate $\rho_{\varepsilon} - \rho$ as

$$\begin{split} \sup_{x \in \mathbb{R}_{+}} |\rho_{\varepsilon}(x) - \rho(x)| \\ &= \sup_{x \in \mathbb{R}_{+}} \left| \int_{\mathbb{R}^{3}} g_{\infty,\varepsilon}(\xi) \left(\frac{|\xi_{1}|}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{m_{b}v_{b}}{\sqrt{v_{b}^{2} + 2\phi(x)}} - \frac{(1+\alpha)m_{\infty}v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi(x)}} \right) d\xi \\ &\leq \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (-v_{\infty}, 0, 0)| \le \varepsilon} \left| \frac{-\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi(x)}} \right| \right) m_{\infty} \\ &+ \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (v_{b}, 0, 0)| \le \varepsilon} \left| \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{b}}{\sqrt{v_{b}^{2} + 2\phi(x)}} \right| \right) m_{b} \\ &+ \sup_{x \in \mathbb{R}_{+}} \left(\sup_{|\xi - (v_{\infty}, 0, 0)| \le \varepsilon} \left| \frac{\xi_{1}}{\sqrt{\xi_{1}^{2} + 2\phi_{\varepsilon}(x)}} - \frac{v_{\infty}}{\sqrt{v_{\infty}^{2} + 2\phi(x)}} \right| \right) \alpha m_{\infty} \\ &\leq C_{0}(\varepsilon + \sup_{x \in \mathbb{R}_{+}} |\phi(x) - \phi_{\varepsilon}(x)|) \le C_{0}\varepsilon. \end{split}$$

Let us treat $\rho_{\varepsilon}u_{\varepsilon} - \rho u$. By the change of variables $\sqrt{\xi_1^2 - 2\phi(x)} = \pm \zeta_1$, the term $\rho_{\varepsilon}u_{\varepsilon}$ is rewritten as

$$(\boldsymbol{\rho}_{\varepsilon}\boldsymbol{u}_{\varepsilon})(\boldsymbol{x}) = \int_{\mathbb{R}^3} \boldsymbol{\xi}_1 g_{\infty,\varepsilon}(\boldsymbol{\xi}) d\boldsymbol{\xi}.$$

It is easy to see $\rho u = m_b v_b + (\alpha - 1) m_{\infty} v_{\infty}$ from (2.31). Then we arrive at

$$\sup_{x\in\mathbb{R}_+}|(\rho_{\varepsilon}u_{\varepsilon})(x)-(\rho u)(x)|=\sup_{x\in\mathbb{R}_+}\left|\int_{\mathbb{R}^3}g_{\infty,\varepsilon}(\xi)(\xi_1-m_bv_b-(\alpha-1)m_{\infty}v_{\infty})d\xi\right|\leq\varepsilon.$$

Consequently, (2.29) holds.

Acknowledge. This work was supported by JSPS KAKENHI Grant Numbers 18K03364 and 21K03308.

A Reduction

In this section, we reduce the boundary value problem of (1.1a)–(1.1b) with conditions (1.1c), (1.1e), (1.1f), and (1.3) to the boundary value problem (1.1). For simplicity, we treat the reduction only for the completely absorbing boundary, i.e. $f_b = \alpha = 0$.

Suppose that the former boundary value problem has a solution (f, ϕ) with $\partial_x \phi(x) < 0$. It is sufficient to show that a value $\phi(0)$ is determined a priori by f_{∞} . Indeed we can construct the solution of the former problem by solving the problem (1.1) with $\phi_b = \phi(0)$. Let us find an a priori value $\phi(0)$. Following the proof of Lemma 3.1, we see that f must be written by the form (2.9) even for the former problem. Substituting (2.9) into (1.3) with $\alpha = 0$ yields the ¹⁰ following condition:

$$n_e(\phi(0))v_e = \int_{\mathbb{R}^3} \xi_1 f_{\infty}(-\sqrt{\xi_1^2 - 2\phi(0)}, \xi') \chi(\xi_1^2 - 2\phi(0)) \chi(-\xi_1) d\xi$$

By solving this with respect to $\phi(0)$, we can know the a priori value $\phi(0)$. Consequently, the former problem can be reduced to the problem (1.1) with $\phi_b = \phi(0)$.

B Estimates of ρ^{\pm}

This section provides the proofs of estimates (2.20) and (2.21). First we can obtain (2.20) by using the Hölder inequality as follows:

$$\begin{split} |\rho_{i}^{+}(\phi)| &\leq \|f_{\infty}\|_{L^{1}(\mathbb{R}^{3})} + C \int_{\sqrt{2(\phi_{b}-\phi)\chi(\phi_{b}-\phi)}}^{\sqrt{2\phi_{b}}} \frac{\xi_{1}}{\sqrt{\xi_{1}^{2}+2\phi-2\phi_{b}}} \left(\int_{\mathbb{R}^{2}} f_{b}(\xi) \ d\xi'\right) d\xi_{1} \\ &\leq \|f_{\infty}\|_{L^{1}(\mathbb{R}^{3})} + C \left(\int_{\sqrt{2(\phi_{b}-\phi)\chi(\phi_{b}-\phi)}}^{\sqrt{2\phi_{b}}} \frac{|\xi_{1}|^{r'}}{(\xi_{1}^{2}+2\phi-2\phi_{b})^{r'/2}} d\xi_{1}\right)^{1/r'} \|f_{b}\|_{L^{r}(0,\sqrt{2\phi_{b}};L^{1}(\mathbb{R}^{2}))} \\ &\leq \|f_{\infty}\|_{L^{1}(\mathbb{R}^{3})} + C \|f_{b}\|_{L^{r}(0,\sqrt{2\phi_{b}};L^{1}(\mathbb{R}^{2}))} \end{split}$$

¹⁰If there is no value $\phi(0)$ so that this condition holds, the boundary value problem of (1.1a)–(1.1b) with conditions (1.1c), (1.1e), (1.1f), and (1.3) admits no solution.

for $\phi \ge 0$, where r' < 2 is the Hölder conjugate of r > 2. Similarly, we observe that for $\phi \in [-M, 0]$,

$$\begin{aligned} |\rho_i^-(\phi)| &= \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi}} \chi(\xi_1^2 - 4M) d\xi \\ &+ \int_{\mathbb{R}^3} f_{\infty}(\xi) \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi}} \{ \chi(\xi_1^2 + 2\phi) - \chi(\xi_1^2 - 4M) \} d\xi \\ &\leq \sqrt{2} \|f_{\infty}\|_{L^1(\mathbb{R}^3)} + \int_{-2\sqrt{M}}^{2\sqrt{M}} \frac{|\xi_1|}{\sqrt{\xi_1^2 + 2\phi}} \chi(\xi_1^2 + 2\phi) \left(\int_{\mathbb{R}^2} f_{\infty}(\xi) d\xi' \right) d\xi_1 \\ &\leq \sqrt{2} \|f_{\infty}\|_{L^1(\mathbb{R}^3)} + C_M \|f_{\infty}\|_{L^r(-2\sqrt{M}, 2\sqrt{M}; L^1(\mathbb{R}^2))}. \end{aligned}$$

Thus (2.21) holds. The proofs are complete.

C Properties of V^{\pm}

We investigate properties of the functions V^{\pm} defined in (2.16) and (2.17).

Lemma C.1. (Attractive boundary) Let $\phi_b > 0$, $\alpha \neq 1$, and $f_b \in L^1(\mathbb{R}^3_+)$ satisfy $f_b \geq 0$ and

$$f_b(\xi) = 0, \quad 0 < \xi_1 < c_0$$
 (C.1)

for some $c_0 > 0$. Suppose that $f_{\infty} \in L^1(\mathbb{R}^3)$ satisfies $f_{\infty} \ge 0$, (2.2), (2.3), and (2.8). Then there exists a positive constant δ such that if $0 < \phi_b < \delta$, the function V^+ belongs to $C^2([0, \phi_b])$ and satisfies $d^2V^+/d\phi^2(0) > 0$.

(Repulsive boundary) Let $\phi_b < 0$, $f_b \in L^1(\mathbb{R}^3_+)$, and $f_b \ge 0$. Suppose that $f_\infty \in L^1(\mathbb{R}^3)$ satisfies $f_\infty \ge 0$, (2.2), (2.4), (2.8), and

$$f_\infty(\xi) = 0, \quad -c_0 < \xi_1 < c_0$$

for some $c_0 > 0$. Then there exists a constant δ such that if $-\delta < \phi_b < 0$, the function V^- belongs to $C^2([\phi_b, 0])$ and satisfies $d^2V^-/d\phi^2(0) > 0$.

Proof. We show only the assertion for the attractive boundary, since the other one can be shown similarly. Recalling the definition of ρ_i^+ in (2.18), we see from (C.1) that the second term vanishes for sufficiently small ϕ_b . Therefore, we can complete the proof by following the same argument as in the last paragraph of the proof of Lemma 3.1, and also noting that

$$\frac{d^2V^+}{d\phi^2}(0) = -\int_{\mathbb{R}^3} \xi_1^{-2} f_{\infty}(\xi) d\xi + 1 > 0,$$

where we have used (2.8) in deriving the last inequality.

28

References

- A. Ambroso, Stability for solutions of a stationary Euler–Poisson problem, Math. Models Methods Appl. Sci. 16 (2006), 1817–1837.
- [2] A. Ambroso, F. Méhats and P.-A. Raviart, *On singular perturbation problems for the nonlinear Poisson equation*, Asympt. Anal. **25** (2001), 39–91.
- [3] C. Bardos and P. Degond, Global existence for the Vlasov–Poisson equation in 3 space variables with small initial data, Ann. Inst. H. Poincaré Anal. Non Linéaire 2 (1985), 101–118.
- [4] Y. O. Belyaeva, Stationary solutions of the Vlasov–Poisson system for two-component plasma under an external magnetic field in a half-space, Math. Model. Nat. Phenom. 12 (2017), 37–50.
- [5] D. Bohm, *Minimum ionic kinetic energy for a stable sheath*, in The characteristics of electrical discharges in magnetic fields, A. Guthrie and R.K.Wakerling eds., McGraw-Hill, New York, (1949), 77–86.
- [6] R. L. F. Boyd and J. B. Thompson, *The Operation of Langmuir Probes in Electro-Negative Plasmas*, Proc. R. Soc. Lond. A **252** (1959), 102–119.
- [7] F. F. Chen, Introduction to Plasma Physics and Controlled Fusion, 2nd edition edition, Springer, 1984.
- [8] E. Esentürk, H. J. Hwang, and W. A. Strauss, *Stationary solutions of the Vlasov–Poisson system with diffusive boundary conditions*, J. Nonlinear Sci. **25** (2015), 315–342.
- [9] M. Feldman, S.-Y. Ha, and M. Slemrod, *A geometric level-set formulation of a plasma-sheath interface*, Arch. Ration. Mech. Anal. **178** (2005), 81–123.
- [10] D. Gérard-Varet, D. Han-Kwan, and F. Rousset, *Quasineutral limit of the Euler–Poisson* system for ions in a domain with boundaries, Indiana Univ. Math. J. **62** (2013), 359–402.
- [11] D. Gérard-Varet, D. Han-Kwan, and F. Rousset, *Quasineutral limit of the Euler–Poisson* system for ions in a domain with boundaries II, J. Éc. polytech. Math. 1 (2014), 343–386.
- [12] R. T. Glassey, *The Cauchy problem in kinetic theory*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1996.
- [13] C. Greengard and P.-A. Raviart, *A boundary value problem for the stationary Vlasov– Poisson equations: The plane diode*, Comm. Pure Appl. Math. **43** (1990), 473–507.
- [14] Y. Guo, Regularity for the Vlasov equations in a half space, Indiana Univ. Math. J. 43 (1994), 255–320.
- [15] Y. Guo and W. Strauss, *Instability of periodic BGK equilibria*, Comm. Pure Appl. Math. 48 (1995), 861–894.

- [16] Y. Guo and W. Strauss, Unstable BGK solitary waves and collisionless shocks, Comm. Math. Phys. 195 (1998), 267–293.
- [17] D. Han-Kwan and M. Iacobelli, Quasineutral limit for Vlasov–Poisson via Wasserstein stability estimates in higher dimension, J. Differential Equations **263** (2017), 1–25.
- [18] D. Han-Kwan and M. Iacobelli, The quasineutral limit of the Vlasov–Poisson equation in Wasserstein metric, Commun. Math. Sci. 15 (2017), 481–509.
- [19] D. Han-Kwan and F. Rousset, Quasineutral limit for Vlasov–Poisson with Penrose stable data, Ann. Sci. Éc. Norm. Supér. 49 (2016), 1445–1495.
- [20] H. J. Hwang and J. J. L. Velázquez, On global existence for the Vlasov–Poisson system in a half space, J. Differential Equations 247 (2009), 1915–1948.
- [21] C.-Y. Jung, B. Kwon, and M. Suzuki, *Quasi-neutral limit for the Euler–Poisson system* in the presence of plasma sheaths with spherical symmetry, Math. Models Methods Appl. Sci. 26 (2016), 2369–2392.
- [22] C.-Y. Jung, B. Kwon, and M. Suzuki, *Quasi-neutral limit for Euler–Poisson system in the presence of boundary layers in an annular domain*, J. Differential Equations 269 (2020), 8007–8054.
- [23] C.-Y. Jung, B. Kwon, and M. Suzuki, On approximate solutions to the Euler–Poisson system with boundary layers, Commun. Nonlinear Sci. Numer. Simul. 96 (2021), 105717.
- [24] P. Knopf, Confined steady states of a Vlasov–Poisson plasma in an infinitely long cylinder, Math. Methods Appl. Sci. 42 (2019), 6369–6384.
- [25] I. Langmuir, *The interaction of electron and positive ion space charges in cathode sheaths*, Phys. Rev. **33** (1929), 954–989.
- [26] M. A. Lieberman and A. J. Lichtenberg, Principles of Plasma Discharges and Materials Processing, 2nd edition, Wiley-Interscience, 2005.
- [27] P.-L. Lions and B. Perthame, *Propagation of moments and regularity for the 3dimensional Vlasov–Poisson system*, Invent. Math. **105** (1991), 415–430.
- [28] S. Nishibata, M. Ohnawa, and M. Suzuki, Asymptotic stability of boundary layers to the Euler–Poisson equations arising in plasma physics, SIAM J. Math. Anal. 44 (2012), 761–790.
- [29] K. Pfaffelmoser, *Global classical solutions of the Vlasov–Poisson system in three dimensions for general initial data*, Journal of Differential Equations **95** (1992), 281–303.
- [30] G. Rein, *Existence of stationary, collisionless plasmas in bounded domains*, Math. Methods Appl. Sci. **15** (1992), 365–374.
- [31] G. Rein, *Collisionless kinetic equations from astrophysics—the Vlasov–Poisson system*, Handbook of differential equations: evolutionary equations, Vol. III, 383–476, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2007.

- [32] K.-U. Riemann, *The Bohm criterion and sheath formation*, J. Phys. D: Appl. Phys. **24** (1991), 493–518.
- [33] K.-U. Riemann, *The Bohm Criterion and Boundary Conditions for a Multicomponent System*, IEEE Trans. Plasma Sci. **23** (1995), 709–716.
- [34] K.-U. Riemann and T. Daube, *Analytical model of the relaxation of a collisionless ion matrix sheath*, J. Appl. Phys. **86** (1999), 1201–1207.
- [35] J. Schaeffer, *Global existence of smooth solutions to the Vlasov–Poisson system in three dimensions*, Communications in Partial Differential Equations **16** (1991), 1313–1335.
- [36] A. L. Skubachevskii, On the unique solvability of mixed problems for the system of Vlasov–Poisson equations in a half-space, Dokl. Math. **85** (2012), 255–258.
- [37] A. L. Skubachevskii, Initial–Boundary Value Problems for the Vlasov–Poisson equations in a half-space, Proc. Steklov Inst. Math. 283 (2013), 197–225.
- [38] A. L. Skubachevskii, *Vlasov–Poisson equations for a two-component plasma in a homo*geneous magnetic field, Russ. Math. Surv. **69** (2014), 291–330.
- [39] A. L. Skubachevskii and Y. Tsuzuki, Classical solutions of the Vlasov–Poisson equations with external magnetic field in a half-space, Comput. Math. Math. Phys., 57 (2017), 541– 557.
- [40] M. Suzuki, Asymptotic stability of stationary solutions to the Euler–Poisson equations arising in plasma physics, Kinet. Relat. Models 4 (2011), 569–588.
- [41] M. Suzuki, Asymptotic stability of a boundary layer to the Euler–Poisson equations for a multicomponent plasma, Kinet. Relat. Models 9 (2016), 587–603.
- [42] M. Suzuki and M. Takayama, Stability and existence of stationary solutions to the Euler– Poisson equations in a domain with a curved boundary, Arch. Ration. Mech. Anal. 239 (2021), 357–387.