A SEMI-FINITE PROOF OF JACOBI'S TRIPLE PRODUCT IDENTITY

JUN-MING ZHU

Abstract. Jacobi's triple product identity is proved from one of Euler's *q*-exponential functions in an elementary way.

1. Introduction

We suppose $|q| < 1$ and, as usual, the *q*-Pochhammer symbols are defined respectively by

$$
(a;q)_{\infty} = \prod_{n=0}^{\infty} (1 - aq^n) \qquad \text{and} \qquad (a;q)_n = \frac{(a;q)_{\infty}}{(aq^n;q)_{\infty}}
$$

for any integer *n*. Jacobi's triple product identity is one of the most important series-product identities. We state it in the following theorem.

Theorem 1.1. *For any complex number* $z \neq 0$ *, we have*

(1.1)
$$
(q;q)_{\infty}(-q/z;q)_{\infty}(-z;q)_{\infty} = \sum_{n=-\infty}^{\infty} q^{n(n-1)/2} z^n.
$$

Bellman said in [\[3,](#page-1-0) p. 42] that there are no simple proofs known of the complete result. After that, Andrews [\[1\]](#page-1-1) proved Theorem [1.1](#page-0-0) using both of Euler's *q*-exponential functions, which are, respectively,

(1.2)
$$
(-z;q)_{\infty} = \sum_{n=0}^{\infty} \frac{q^{n(n-1)/2} z^n}{(q;q)_n}, \ z \in \mathbb{C},
$$

and

$$
\frac{1}{(-z;q)_{\infty}} = \sum_{n=0}^{\infty} \frac{(-1)^n z^n}{(q;q)_n}, \ |z| < 1.
$$

In this short note, we will prove Jacobi's triple product identity only using [\(1.2\)](#page-0-1). This seems also a proof of the complete triple product identity required by Bellman. We use a method which is often called semi-finite method (see [\[3\]](#page-1-0)).

Keywords: Jacobi's triple product identity; Euler's *q*-exponential function; semi-finite. MSC (2010): 33D15, 11F27.

2. The proof

Proof. For a nonnegative integer *m* and $z \neq 0$,

$$
(2.1) \quad \sum_{n=-m}^{\infty} \frac{q^{n(n-1)/2} z^n}{(q^{m+1};q)_n} = \sum_{n=0}^{\infty} \frac{q^{(n-m)(n-m-1)/2} z^{n-m}}{(q^{m+1};q)_{n-m}}
$$

$$
= \frac{q^{m(m+1)/2} z^{-m}}{(q^{m+1};q)_{-m}} \sum_{n=0}^{\infty} \frac{q^{n(n-1)/2} (zq^{-m})^n}{(q;q)_n}
$$

$$
= q^{m(m+1)/2} z^{-m} (q;q)_m (-zq^{-m};q)_{\infty} \qquad \text{(by (1.2))}
$$

$$
= q^{m(m+1)/2} z^{-m} (q;q)_m (-zq^{-m};q)_m (-z;q)_{\infty}
$$

$$
= (q;q)_m (-q/z;q)_m (-z;q)_{\infty}.
$$

Note that $\frac{1}{(q^{m+1};q)_{-n}} = (q^{m+1-n};q)_n = 0$ when $n > m$. The left hand side of [\(2.1\)](#page-1-2) can be rewritten as

$$
\sum_{n=0}^{\infty} \frac{q^{n(n-1)/2} z^n}{(q^{m+1};q)_n} + \sum_{n=1}^{\infty} (q^{m+1-n};q)_n q^{n(n+1)/2} z^{-n}.
$$

Restricting *z* in any compact subset of $0 < |z| < \infty$ and letting $m \to \infty$ in [\(2.1\)](#page-1-2), we get [\(1.1\)](#page-0-2). By analytic continuation, the restriction on *z* may be relaxed. \Box

Acknowledgment. The author wishes to thank the editor and the referee for their valuable comments and advices. The author is partially supported by the National Science Foundation of China (Grant No. 11371184).

REFERENCES

- [1] G. E. Andrews, A simple proof of Jacobi's triple product identity, *Proc. Amer. Math. Soc.* 16 no. 2 (1965) 333–334.
- [2] R. Bellman, *A brief introduction to theta functions*, Holt, Rinehart and Winston, New York, 1961.
- [3] W. Y. C. Chen, A. M. Fu, Semi-finite forms of bilateral basic hypergeometric series, *Proc. Amer. Math. Soc.* 134 (2006) 1719–1725.

Department of Mathematics, Luoyang Normal University, Luoyang City, Henan Province 471022, **CHINA**

Email address: junming zhu@163.com