

# Generalized Gibbs ensemble of the Ablowitz-Ladik lattice, Circular $\beta$ -ensemble and double confluent Heun equation

T. Grava <sup>\*</sup>, G. Mazzuca <sup>†</sup>

October 14, 2021

## Abstract

We consider the discrete defocusing nonlinear Schrödinger equation in its integrable version, which is called defocusing Ablowitz-Ladik lattice. We consider periodic boundary conditions with period  $N$  and initial data sample according to the Generalized Gibbs ensemble. In this setting, the Lax matrix of the Ablowitz-Ladik lattice is a random CMV-periodic matrix and it is related to the Killip-Nenciu Circular  $\beta$ -ensemble at high-temperature. We obtain the generalized free energy of the Ablowitz-Ladik lattice and the density of states of the random Lax matrix by establishing a mapping to the one-dimensional log-gas. For the Gibbs measure related to the Hamiltonian of the Ablowitz-Ladik flow, we obtain the density of states via a particular solution of the double-confluent Heun equation.

## 1 Introduction

The defocusing Ablowitz-Ladik (AL) lattice for the complex functions  $\alpha_j(t)$ ,  $j \in \mathbb{Z}$  and  $t \in \mathbb{R}$ , is the system of nonlinear equations

$$i\dot{\alpha}_j = -(\alpha_{j+1} + \alpha_{j-1} - 2\alpha_j) + |\alpha_j|^2(\alpha_{j-1} + \alpha_{j+1}), \quad (1.1)$$

where  $\dot{\alpha}_j = \frac{d\alpha_j}{dt}$ . We assume  $N$ -periodic boundary conditions  $\alpha_{j+N} = \alpha_j$ , for all  $j \in \mathbb{Z}$ . The AL lattice was introduced by Ablowitz and Ladik [2,3] as the spatial integrable discretization of the cubic nonlinear Schrödinger Equation (NLS) for the complex function  $\psi(x, t)$ ,  $x \in S^1$  and  $t \in \mathbb{R}$ :

$$i\partial_t \psi(x, t) = -\partial_x^2 \psi(x, t) + 2|\psi(x, t)|^2 \psi(x, t).$$

The cubic NLS equation was proved to be integrable by Zakharov and Shabat [66].

It is straightforward to verify that the quantity

$$K^{(0)} := \prod_{j=1}^N (1 - |\alpha_j|^2), \quad (1.2)$$

is a constant of motion for the AL lattice, namely  $\frac{d}{dt} K^{(0)} = 0$ . This implies that if  $|\alpha_j(0)| < 1$  for all  $j \in \mathbb{Z}$ , then  $|\alpha_j(t)| < 1$  for all  $t > 0$ . We chose the  $N$ -dimensional disc  $\mathbb{D}^N$  as the phase space of the AL system, here  $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ . On  $\mathbb{D}^N$  we introduce the symplectic form [21,27]

$$\omega = i \sum_{j=1}^N \frac{1}{\rho_j^2} d\alpha_j \wedge d\bar{\alpha}_j, \quad \rho_j = \sqrt{1 - |\alpha_j|^2}. \quad (1.3)$$

The corresponding Poisson bracket is defined for functions  $f, g \in C^\infty(\mathbb{D}^N)$  as

$$\{f, g\} = i \sum_{j=1}^N \rho_j^2 \left( \frac{\partial f}{\partial \bar{\alpha}_j} \frac{\partial g}{\partial \alpha_j} - \frac{\partial f}{\partial \alpha_j} \frac{\partial g}{\partial \bar{\alpha}_j} \right).$$

<sup>\*</sup>International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy and School of Mathematics, University of Bristol, Fry Building, BS8 1UG, UK

Email: [grava@sissa.it](mailto:grava@sissa.it)

<sup>†</sup>International School for Advanced Studies (SISSA), Via Bonomea 265, 34136 Trieste, Italy

Email: [guido.mazzuca@sissa.it](mailto:guido.mazzuca@sissa.it)

The phase shift  $\alpha_j(t) \rightarrow e^{-2it}\alpha_j(t)$  transforms the AL lattice into the equation

$$i\dot{\alpha}_j = -\rho_j^2(\alpha_{j+1} + \alpha_{j-1}), \quad \rho_j = \sqrt{1 - |\alpha_j|^2}, \quad (1.4)$$

which we call the *reduced* AL equation. We remark that the quantity  $-2\ln(K^{(0)})$  is the generator of the shift  $\alpha_j(t) \rightarrow e^{-2it}\alpha_j(t)$ , while  $H_1 = K^{(1)} + \overline{K^{(1)}}$  with

$$K^{(1)} := -\sum_{j=1}^N \alpha_j \bar{\alpha}_{j+1},$$

generates the flow (1.4). The AL equations (1.1) have the Hamiltonian structure

$$\dot{\alpha}_j = \{\alpha_j, H_{AL}\}, \quad H_{AL}(\alpha_j, \bar{\alpha}_j) = -2\ln(K^{(0)}) + K^{(1)} + \overline{K^{(1)}}. \quad (1.5)$$

**Integrability.** The AL lattice was discovered by Ablowitz and Ladik by discretizing the  $2 \times 2$  Zakharov-Shabat Lax pair [1, 2] of the cubic nonlinear Schrödinger equation. For a comprehensive review see [4]. The integrability of the Ablowitz-Ladik system has also been proved by constructing a bi-Hamiltonian structure [9, 21].

Using the connection between orthogonal polynomials on the unit circle and the AL lattice, Nenciu and Simon [49, 57] constructed a new Lax pair for the AL lattice. The link between orthogonal and biorthogonal polynomials on the unit circle and the Ablowitz-Ladik hierarchy (see also [5], [38]) is the analogue of the celebrated link between the Toda hierarchy and orthogonal polynomials on the real line (see e.g. [17]). This link was also generalized to the non-commutative case [14]. Generalization of this construction to other integrable equations has been considered in [50, 53].

Following [49, 57] we double the size of the chain according to the periodic boundary condition, thus we consider a chain of  $2N$  particles  $\alpha_1, \dots, \alpha_{2N}$  such that  $\alpha_j = \alpha_{j+N}$  for  $j = 1, \dots, N$ . Define the  $2 \times 2$  unitary matrix  $\Xi_j$

$$\Xi_j = \begin{pmatrix} \bar{\alpha}_j & \rho_j \\ \rho_j & -\alpha_j \end{pmatrix}, \quad j = 1, \dots, 2N,$$

and the  $2N \times 2N$  matrices

$$\mathcal{M} = \begin{pmatrix} -\alpha_{2N} & & & & & & & \rho_{2N} \\ & \Xi_2 & & & & & & \\ & & \Xi_4 & & & & & \\ & & & \ddots & & & & \\ & & & & \Xi_{2N-2} & & & \\ \rho_{2N} & & & & & & & \bar{\alpha}_{2N} \end{pmatrix}, \quad \mathcal{L} = \begin{pmatrix} \Xi_1 & & & & & \\ & \Xi_3 & & & & \\ & & \ddots & & & \\ & & & \Xi_{2N-1} & & \end{pmatrix}.$$

Now let us define the unitary Lax matrix

$$\mathcal{E} = \mathcal{L}\mathcal{M}, \quad (1.6)$$

that has the structure of a 5-band diagonal matrix

$$\begin{pmatrix} * & * & * & & & & & * \\ * & * & * & & & & & * \\ & * & * & * & * & & & \\ & * & * & * & * & & & \\ & & & & \ddots & \ddots & & \\ & & & & & * & * & * & * \\ & & & & & * & * & * & * \\ * & & & & & * & * & * & * \\ * & & & & & * & * & * & * \end{pmatrix}.$$

The matrix  $\mathcal{E}$  is a periodic CMV matrix (after Cantero Morales and Velsquez [15]). The  $N$ -periodic equation (1.4) is equivalent to the following Lax equation for the matrix  $\mathcal{E}$ :

$$\dot{\mathcal{E}} = i[\mathcal{E}, \mathcal{E}^+ + (\mathcal{E}^+)^{\dagger}], \quad (1.7)$$

where  $\dagger$  stands for hermitian conjugate and

$$\mathcal{E}_{j,k}^+ = \begin{cases} \frac{1}{2}\mathcal{E}_{j,j} & j = k \\ \mathcal{E}_{j,k} & k = j + 1 \bmod 2N \text{ or } k = j + 2 \bmod 2N \\ 0 & \text{otherwise.} \end{cases}$$

We observe that  $(\mathcal{E}^+)^\dagger + (\mathcal{E}^\dagger)^+ = \mathcal{E}^\dagger$  and  $[\mathcal{E}, \mathcal{E}^\dagger] = 0$  because  $\mathcal{E}$  is unitary. Therefore, one can write the equations (1.7) in the equivalent form

$$\dot{\mathcal{E}} = i [\mathcal{E}, \mathcal{E}^+ - (\mathcal{E}^\dagger)^+].$$

The formulation (1.7) implies that the quantities

$$K^{(\ell)} = \frac{\text{Tr}(\mathcal{E}^\ell)}{2}, \quad \ell = 1, \dots, N-1, \quad (1.8)$$

are constants of motion for the defocusing AL system. For example

$$K^{(1)} = -\sum_{j=1}^N \alpha_j \bar{\alpha}_{j+1}, \quad K^{(2)} = \sum_{j=1}^N [(\alpha_j \bar{\alpha}_{j+1})^2 - 2\alpha_j \bar{\alpha}_{j+2} \rho_{j+1}^2].$$

Furthermore,  $K^{(0)}, K^{(1)}, \dots, K^{(N-1)}$  are functionally independent and in involution, showing that the  $N$ -periodic AL system is integrable [1–4, 49].

**Remark 1.1.** We remark that the quantity  $2\Re(K^{(1)})$  generates the reduces AL equation (1.4), while the quantity  $-2\Im(K^{(1)})$  generates the flow

$$\dot{\alpha}_j = (1 - |\alpha_j|^2)(\alpha_{j+1} - \alpha_j),$$

which is called Schur flow. The Schur flow emerges in [3] as a spatial discretization of the defocusing modified Korteweg de Vries equation

$$\partial_t f - 6f \partial_x f + \partial_x^3 f = 0.$$

For the integration of the Schur flow and its relation to orthogonal polynomial on the unit circle see [28].

**Generalized Gibbs Ensemble for the Ablowitz-Ladik Lattice.** The symplectic form  $\omega$  in (1.3) induces on  $\mathbb{D}^N$  the volume form  $\text{dvol} = \frac{1}{K^{(0)}} \text{d}^2\alpha$ , with  $\text{d}^2\alpha = \prod_{j=1}^N (i \text{d}\alpha_j \wedge \text{d}\bar{\alpha}_j)$ . We observe that  $\int_{\mathbb{D}^N} \text{dvol} = \infty$ , however we can define the Gibbs measure with respect to the Hamiltonian  $H_{AL}$  in (1.5):

$$\frac{1}{Z_\beta} e^{-\frac{\beta}{2} H_{AL}} \text{dvol} = \frac{1}{Z_\beta} e^{\beta \Re(K^{(1)})} \prod_{j=1}^N (1 - |\alpha_j|^2)^{\beta-1} \text{d}^2\alpha, \quad \beta > 0, \quad (1.9)$$

where  $Z_\beta = \int_{\mathbb{D}^N} e^{\beta \Re(K^{(1)})} \prod_{j=1}^N (1 - |\alpha_j|^2)^{\beta-1} \text{d}^2\alpha < \infty$  is the normalizing constant. The above probability measure is clearly invariant under the Hamiltonian flow  $\alpha_j(0) \rightarrow \alpha_j(t)$  associated to the Ablowitz-Ladik equation (1.1).

Since the Ablowitz-Ladik lattice posses several conserved quantities (1.2)-(1.8), one can introduce a Generalized Gibbs Ensemble on the phase space  $\mathbb{D}^N$  in the following way. Fix  $\mathbb{N} \ni \kappa < N-1$  and let us define

$$V(z) = \sum_{m=1}^{\kappa} \eta_m \Re(z^m), \quad (1.10)$$

where  $\eta_m \in \mathbb{R}$  are called chemical potentials. Then

$$\text{Tr}(V(\mathcal{E})) = \sum_{m=1}^{\kappa} \eta_m (K^{(m)} + \overline{K^{(m)}}),$$

where  $K^{(m)}$  are the AL conserved quantities (1.8). The finite volume Generalized Gibbs measure can be written as:

$$\text{d}\mathbb{P}_{AL}(\alpha_1, \dots, \alpha_N) = \frac{1}{Z_N^{AL}(V, \beta)} \prod_{j=1}^N (1 - |\alpha_j|^2)^{\beta-1} \exp(-\text{Tr}(V(\mathcal{E}))) \text{d}^2\alpha, \quad (1.11)$$

where  $Z_N^{AL}(V, \beta)$  is the partition function of the system:

$$Z_N^{AL}(V, \beta) = \int_{\mathbb{D}^N} \prod_{j=1}^N (1 - |\alpha_j|^2)^{\beta-1} \exp(-\text{Tr}(V(\mathcal{E}))) d^2\alpha. \quad (1.12)$$

Choosing the initial data of the Ablowitz-Ladik lattice according to the Generalized Gibbs measure, the Lax matrix  $\mathcal{E}$  turns into a random matrix. In [46] Mendl and Sphon study the dynamic of the Ablowitz-Ladik lattice at non-zero temperature. They study numerically correlation functions and in particular, introducing the density  $\delta_j = \Re(\alpha_{j+1}\bar{\alpha}_j)$ , they study the density-density correlation function

$$\mathbf{E}[\delta_j(t)\delta_1(0)] - \mathbf{E}[\delta_j(t)]\mathbf{E}[\delta_1(0)]$$

where  $\mathbf{E}[\cdot]$  is the expectation with respect to Gibbs measure (1.9). They showed numerically that density-density time correlations in thermal equilibrium have symmetrically located peaks, which travel in opposite directions at constant speed, broaden ballistically and decay as  $1/t^\gamma$  when  $t \rightarrow \infty$ , where the scaling exponent  $\gamma$  is approximately equal to one. This behaviour is believed to be typical of integrable nonlinear systems.

More quantitative results have been obtained for linear (integrable) systems and for the Toda lattice, which is a nonlinear integrable system. It was shown in [29] that the fastest peaks of the correlation functions of harmonic oscillators with short range interactions have a Airy type scaling. Regarding nonlinear integrable systems in [58] Spohn was able to connect the Gibbs ensemble of the Toda lattice to the Dumitriu-Edelman  $\beta$ -ensemble [19]. In this way, the generalized Gibbs free energy of the Toda chain turns out to be related to the  $\beta$ -ensembles of random matrix theory in the mean-field regime [7,20]. The behaviour of the correlation functions of the Toda chains has been derived by applying the theory of generalized hydrodynamic [18,60]. We mention also the recent work [31], where the authors derive a large deviation principle for the mean density of states for the Generalized Gibbs measure of the Toda lattice.

## 2 Statement of the results

In this manuscript we derive the mean density of states  $\mu_{AL}^\beta$  of the random Lax matrix  $\mathcal{E}$  sampled according to generalized Gibbs measure (1.11) and we determine the free energy of the AL generalized Gibbs ensemble

$$F_{AL}(V, \beta) = \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_N^{AL}(V, \beta).$$

This is achieved by connecting the generalized Gibbs ensemble of the Ablowitz-Ladik lattice to the Killip-Nenciu [37] matrix Circular  $\beta$ -ensemble at high-temperature investigated by Hardy and Lambert [32]. Let  $\mathcal{M}(\mathbb{T})$  be the space of probability measures on the torus  $\mathbb{T} = [-\pi, \pi]$  and for  $\mu \in \mathcal{M}(\mathbb{T})$  let us consider the functional

$$\mathcal{F}^{(V, \beta)}(\mu) = 2 \int_{\mathbb{T}} V(\theta) \mu(\theta) d\theta - \beta \int \int_{\mathbb{T} \times \mathbb{T}} \ln \sin \left( \frac{|\theta - \phi|}{2} \right) \mu(\theta) \mu(\phi) d\theta d\phi + \int_{\mathbb{T}} \ln(\mu(\theta)) \mu(\theta) d\theta + \ln(2\pi). \quad (2.1)$$

Such functional has a unique minimizer  $\mu_{HT}^\beta(d\theta) = \mu_{HT}^\beta(\theta)d\theta$ , [56], that describes the density of states of the  $\beta$ -ensembles at high temperature [32]. For finite  $\beta$  and smooth potentials  $V(\theta)$ , it has been shown by Hardy and Lambert in [32] that the minimizer  $\mu_{HT}^\beta(d\theta)$  has a smooth density and its support is the whole torus  $\mathbb{T}$ .

**Theorem 2.1** (First Main theorem). *Consider  $\beta > 0$  and a smooth potential  $V(e^{i\theta})$  as in (1.10) on the unit circle  $\mathbb{T}$ . The mean density of states  $\mu_{AL}^\beta(d\theta) := \mu_{AL}^\beta(\theta)d\theta$  of the Ablowitz Ladik Lax matrix  $\mathcal{E}$  in (1.6) endowed with the probability (1.11) is absolutely continuous with respect to the Lebesgue measure and takes the form*

$$\mu_{AL}^\beta(\theta) = \partial_\beta \left( \beta \mu_{HT}^\beta(\theta) \right) \text{ a.e.}, \quad (2.2)$$

where  $\mu_{HT}^\beta$  is the unique minimizer of the functional (3.12).

To prove the above theorem we derive a relation (see Proposition 4.2) between the free energy of the  $\beta$ -ensembles at high temperature, namely the minimum value of the minimizer (2.1)

$$F_{HT}(V, \beta) := \mathcal{F}^{(V, \beta)}(\mu_{HT}^\beta),$$

and the free energy  $F_{AL}(V, \beta)$  of the AL lattice:

$$F_{AL}(V, \beta) = \partial_\beta (\beta F_{HT}(V, \beta)) + \ln(2).$$

Such relation is obtained via transfer operator techniques.

The particular case  $V(e^{i\theta}) = 2\eta \cos \theta$  corresponds to the free energy associated to the AL equation (1.1), and we show that the minimizer of the functional (2.1) is obtained via a particular solution of the Double Confluent Heun (DCH) equation.

**Theorem 2.2** (Second main theorem). *Fix  $\beta > 0$  and let  $V(e^{i\theta}) = \eta \cos \theta$ , where  $\eta$  is a real parameter. There exists  $\varepsilon > 0$  such that for all  $\eta \in (-\varepsilon, \varepsilon)$ , the minimizer  $\mu_{HT}^\beta(d\theta) = \mu_{HT}^\beta(\theta)d\theta$  of the functional (2.1) takes the form*

$$\mu_{HT}^\beta(\theta) = \frac{1}{2\pi} + \frac{1}{\pi\beta} \Re \left( \frac{zv'(z)}{v(z)} \right) \Big|_{z=e^{i\theta}}, \quad (2.3)$$

where  $v(z)$  is the unique solution (up to a multiplicative non-zero constant) of the Double Confluent Heun (DCH) equation

$$z^2 v''(z) + (-\eta + z(\beta + 1) + \eta z^2) v'(z) + \eta\beta(z + \lambda)v(z) = 0 \quad (2.4)$$

analytic for  $|z| \leq r$  with  $r \geq 1$ . Such solution is differentiable in the parameter  $\eta$  and  $\beta$ . The parameter  $\lambda = \lambda(\eta, \beta)$  in (2.4) is determined for  $\eta \in (-\varepsilon, \varepsilon)$  by the solution of the equation

$$\lambda(R_1)_{11} + \frac{\eta}{\beta + 1}(R_1)_{21} = 0,$$

with the condition  $\lambda(\eta = 0, \beta) = 0$ . In the above expression  $(R_1)_{jk}$  is the  $jk$  entry of the matrix  $R_1$  which is defined by the infinite product

$$R_1 = M_1 M_2 \dots M_k \dots, \quad M_k = \begin{pmatrix} 1 + \frac{\lambda\beta\eta}{k(k+\beta)} & \frac{\eta^2}{k(k+\beta+1)} \\ 1 & 0 \end{pmatrix}.$$

We remark that the solution of the double confluent Heun equation has generically an essential singularity at  $z = 0$  and  $z = \infty$ , and one needs to tune the *accessory parameter*  $\lambda$  to obtain an analytic solution, for a review see [54]. In our derivation of (2.4) the parameter  $\lambda$  coincides with the first moment of the measure  $\mu(\theta)$ , namely  $\lambda = \int_{\mathbb{T}} \mu(\theta) e^{i\theta} d\theta$ . It is a transcendental function of  $\beta$  and  $\eta$  and it is related to the Painlevé III equation [22, 42].

**Remark 2.3.** *Under the change of variable*

$$v(z) = \exp\left(-\frac{\eta}{2}\left(z + \frac{1}{z}\right)\right) z^{-\frac{\beta+1}{2}} f(z),$$

the DCH equation (2.4) takes the form of a Schrödinger equation

$$f''(z) + q(z)f(z) = 0,$$

with potential  $q(z)$  singular at the origin

$$q(z) = \frac{1}{z^2} \left( \eta \frac{\beta-1}{2} \left( z + \frac{1}{z} \right) - \frac{\eta^2 (z^2 - 1)^2}{4z^2} - \frac{1}{4}(\beta^2 + 4\beta + 3) + \eta\beta\lambda \right).$$

**Remark 2.4.** *For the case  $V = 0$  it was shown in [32] that the minimizer of the functional (3.12) is the uniform measure on the circle, while for the case  $V(e^{i\theta}) \rightarrow \beta V(e^{i\theta})$  and  $\beta \rightarrow \infty$  the minimizer of (2.1) was considered in [44]. The particular case  $V(e^{i\theta}) \rightarrow \beta\eta \cos \theta$  and  $\beta \rightarrow \infty$  has first been considered by Gross Witten [30] and Baik-Deift-Johansson [11]. Despite Theorem 2.2 is proved for small values of  $\eta$ , the measure (2.3) generalizes the result of Gross and Witten [30] and Baik-Deift-Johansson [11] to the high-temperature regime (see Remark 5.6).*

This manuscript is organized as follows. In section 3 we introduce the Circular  $\beta$  ensemble and its high-temperature limit. We review old results and we derive some technical results needed to prove our main theorems. In section 4 we prove our first main theorem, namely Theorem 2.1 and in section 5 we prove Theorem 2.2. Finally, the most technical parts of our arguments are deferred to the appendix.



With the above changes, we arrive to the probability distribution of the Circular ensemble at high-temperature, and with an external potential:

$$d\mathbb{P}_\beta^V(\theta_1, \dots, \theta_N) = \frac{1}{Z_N^{HT}(V, \beta)} \prod_{j < \ell} |e^{i\theta_\ell} - e^{i\theta_j}|^{\frac{2\beta}{N}} e^{-2\sum_{j=1}^N V(\theta_j)} d\boldsymbol{\theta}, \quad (3.3)$$

where  $Z_N^{HT}(V, \beta)$  is the normalizing constant or the partition function of the system. Also in this case, we can associate to the above probability distribution a random CMV matrix. The lemma below has probably already appeared in the literature, but for completeness we provide the proof.

**Lemma 3.3.** *Let  $E$  be the CMV matrix (3.2). Consider the block  $2N \times 2N$  matrix*

$$\tilde{E} = \text{diag}(E, E), \quad (3.4)$$

whose entries are distributed according to

$$d\mathbb{P}(\alpha_1, \dots, \alpha_N) = \frac{1}{Z_N^{HT}(V, \beta)} \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1-\frac{j}{N})-1} e^{-\text{Tr}(V(\tilde{E}))} \prod_{j=1}^{N-1} d^2\alpha_j \frac{d\alpha_N}{i\alpha_N}. \quad (3.5)$$

Then the eigenvalues of  $\tilde{E}$  are all double, they lie on the unit circle and are distributed according to (3.3). Moreover

$$Z_N^{HT}(V, \beta) = 2^{1-N} \frac{\Gamma\left(\frac{\beta}{N}\right)^N}{\Gamma(\beta)} Z_N^{HT}(V, \beta), \quad (3.6)$$

where  $Z_N^{HT}(V, \beta)$  is the norming constant of the probability distribution (3.3) and  $Z_N^{HT}(V, \beta)$  is the norming constant of the probability distribution (3.5).

*Proof.* First, we notice that the eigenvalues of  $\tilde{E}$  are all double, since it is a block diagonal matrix with two identical blocks.

We consider the change of variables  $\alpha_N \rightarrow e^{i\varphi}$ , thus (3.5) becomes:

$$d\mathbb{P}(\alpha_1, \dots, \alpha_{N-1}, \varphi) = \frac{1}{Z_N^{HT}(V, \beta)} \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1-\frac{j}{N})-1} e^{-\text{Tr}(V(\tilde{E}))} \prod_{j=1}^{N-1} d^2\alpha_j d\varphi. \quad (3.7)$$

Now, let  $e^{i\theta_1}, \dots, e^{i\theta_N}$  be the eigenvalues of the CMV matrix  $E$  endowed with probability (3.5), and let  $q_1, \dots, q_N$  be the entries of the first row of the unitary matrix  $Q$  such that  $Q^\dagger \Theta Q = E$  where  $\Theta = \text{Diag}(e^{i\theta_1}, \dots, e^{i\theta_N})$  and  $\sum_{k=1}^N |q_k|^2 = 1$ . Introduce the variable  $\gamma_j = |q_j|^2$  for  $j = 1, \dots, N$ , then From [37, Lemma 4.1, and Proposition 4.2 relation (4.14)] we have

$$|\Delta(e^{i\theta})|^2 \prod_{j=1}^N \gamma_j = \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{(N-j)}, \quad (3.8)$$

$$\left| \frac{\partial(\alpha_1, \dots, \alpha_{N-1}, \varphi)}{\partial(\boldsymbol{\theta}, \boldsymbol{\gamma})} \right| = 2^{1-N} \frac{\prod_{j=1}^{N-1} (1 - |\alpha_j|^2)}{\prod_{j=1}^N \gamma_j}, \quad (3.9)$$

here  $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_{N-1})$ , and  $\Delta(e^{i\theta}) = \prod_{j < \ell} (e^{i\theta_j} - e^{i\theta_\ell})$ . Applying the previous equalities to (3.7) we derive

$$\begin{aligned} d\mathbb{P}(\alpha_1, \dots, \alpha_{N-1}, \varphi) &= \frac{e^{-\text{Tr}(V(\tilde{E}))}}{Z_N^{HT}(V, \beta)} d\varphi \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1-\frac{j}{N})-1} d\alpha_j d\bar{\alpha}_j \\ &\stackrel{(3.9)}{=} \frac{1}{Z_N^{HT}(V, \beta)} \frac{2^{1-N}}{\prod_{j=1}^N \gamma_j} \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1-\frac{j}{N})} e^{-2\sum_{j=1}^N V(e^{i\theta_j})} d\boldsymbol{\theta} d\boldsymbol{\gamma} \\ &\stackrel{(3.8)}{=} \frac{1}{Z_N^{HT}(V, \beta)} 2^{1-N} |\Delta(e^{i\theta})|^{\frac{2\beta}{N}} \prod_{j=1}^N \gamma_j^{\frac{\beta}{N}-1} e^{-2\sum_j V(e^{i\theta_j})} d\boldsymbol{\theta} d\boldsymbol{\gamma}. \end{aligned}$$

Thus, we deduce the relation

$$Z_N^{HT}(V, \beta) = 2^{1-N} \mathcal{Z}_N^{HT}(V, \beta) \int_{\Delta} \prod_{j=1}^N \gamma_j^{\frac{\beta}{N}-1} d\gamma_1 \dots d\gamma_{N-1},$$

here  $\Delta$  is the simplex  $\sum_{j=1}^N \gamma_j = 1$ . The above integral is a well-known Dirichlet integral that can be computed explicitly (see [37, Lemma 4.4])

$$\int_{\Delta} \prod_{j=1}^N \gamma_j^{\frac{\beta}{N}-1} d\gamma_1 \dots d\gamma_{N-1} = \frac{\Gamma\left(\frac{\beta}{N}\right)^N}{\Gamma(\beta)},$$

proving (3.6). □

Let  $e^{i\theta_1}, \dots, e^{i\theta_N}$  be the double eigenvalues of the CMV Matrix  $\tilde{E}$  in (3.4), whose entries are distributed according to (3.7). The *empirical measure* is the random probability measure

$$\mu_N = \frac{1}{N} \sum_{j=1}^N \delta_{e^{i\theta_j}}. \quad (3.10)$$

The mean density of state  $\mu_{HT}^{\beta}$  is defined as the non-random probability measure such that

$$\int_{\mathbb{T}} f(\theta) \mu_{HT}^{\beta}(d\theta) = \lim_{N \rightarrow \infty} \mathbf{E} \left[ \int_{\mathbb{T}} f(\theta) \mu_N(d\theta) \right], \quad (3.11)$$

for all continuous function  $f$  on the torus  $\mathbb{T}$ , and the expected value is taken with respect to (3.5). In order to discuss the large  $N$  limit of  $\mu_N$  we have to introduce several quantities. Let  $\mathcal{M}(\mathbb{T})$  be the set of probability measures on the one-dimensional torus  $\mathbb{T}$  and for  $\mu \in \mathcal{M}(\mathbb{T})$  we consider the logarithmic energy [56]

$$E(\mu) := \int_{\mathbb{T} \times \mathbb{T}} \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right|^{-1} \mu(d\theta) \mu(d\phi).$$

We define the relative entropy with respect to  $\mu_0(d\theta) = \frac{d\theta}{2\pi}$  as

$$K(\mu|\mu_0) := \int_{\mathbb{T}} \log \left( \frac{\mu}{\mu_0} \right) \mu(d\theta) \in [0, +\infty],$$

when  $\mu$  is absolutely continuous with respect to  $\mu_0$  and otherwise  $K(\mu|\mu_0) := +\infty$ . The relevant functional is

$$\mathcal{F}^{(V, \beta)}(\mu) := \beta E(\mu) + K(\mu|\mu_0) + 2 \int_{\mathbb{T}} V(\theta) \mu(d\theta).$$

When  $\mathcal{F}^{(V, \beta)}(\mu)$  is finite, it follows that  $\mu$  is absolutely continuous with respect to the Lebesgue measure  $\mu_0$  and we can write  $\mu(d\theta) = \mu(\theta)d\theta$ . We denote by  $C^{n,1}(\mathbb{T})$  with  $n = 0, 1, 2, \dots$  the space of  $n$ -times differentiable functions whose  $n$ -derivative is also Lipschitz continuous.

The following result describes the limiting measure  $\mu_{HT}^{\beta}$  in (3.11).

**Theorem 3.4.** (cf. [32, Proposition 2.1 and 2.5]) *Let  $\mathcal{M}(\mathbb{T})$  be the set of probability measures on the one-dimensional torus and  $V : \mathbb{T} \rightarrow \mathbb{R}$  be a measurable and bounded function. For any  $\beta > 0$  consider the functional  $\mathcal{F}^{(V, \beta)} : \mathcal{M}(\mathbb{T}) \rightarrow [0, \infty]$*

$$\mathcal{F}^{(V, \beta)}(\mu) = 2 \int_{\mathbb{T}} V(\theta) \mu(\theta) d\theta + \beta E(\mu) + \int_{\mathbb{T}} \ln(\mu(\theta)) \mu(\theta) d\theta + \ln(2\pi). \quad (3.12)$$

Then

(a) *the functional  $\mathcal{F}^{(V, \beta)}(\mu)$  has a unique minimizer  $\mu_{HT}^{\beta}(d\theta) = \mu_{HT}^{\beta}(\theta)d\theta$  in  $\mathcal{M}(\mathbb{T})$ ;*

(b)  $\mu_{HT}^\beta$  is absolutely continuous with respect to the Lebesgue measure and there is  $0 < \delta < 1$  such that

$$\delta \leq \frac{\mu_{HT}^\beta(\theta)}{2\pi} \leq \delta^{-1}, \quad \text{a.e.};$$

(c) if  $V = 0$ , then  $\mu_{HT}^\beta(d\theta) = \frac{1}{2\pi}d\theta$ ;

(d) if  $V \in C^{m,1}(\mathbb{T})$ , then  $\mu_{HT}^\beta \in C^{m,1}(\mathbb{T})$ ;

(d) the empirical measure  $\mu_N$  in (3.10) converges weakly and almost surely to the measure  $\mu_{HT}^\beta$  as  $N \rightarrow \infty$ .

From the above theorem when the potential  $V$  is at least  $C^{2,1}(\mathbb{T})$  the minimizer of the functional  $\mathcal{F}^{(V,\beta)}$  is characterized by the Euler-Lagrange equations

$$\frac{\delta \mathcal{F}^{(V,\beta)}}{\delta \mu} = 2V(\theta) - 2\beta \int_{\mathbb{T}} \ln \sin \left( \frac{|\theta - \phi|}{2} \right) \mu(\phi) d\phi + \ln \mu(\theta) + 1 = C(V, \beta) \quad (3.13)$$

where  $C(V, \beta)$  is a constant in  $\theta$ . Below we derive further properties of the minimizer  $\mu_{HT}^\beta$ .

**Lemma 3.5.** *For any  $V(z)$  as in (1.10), any  $\beta > 0$  the following holds*

- a) *The map  $\beta \rightarrow \inf(\mathcal{F}^{(V,\beta)}(\mu))$  is Lipschitz;*
- b) *The map  $t \rightarrow \inf(\mathcal{F}^{(V+tz^m,\beta)}(\mu))$  is Lipschitz;*
- c) *Let  $D$  be the distance on  $\mathcal{M}(\mathbb{T})$  given by*

$$\begin{aligned} D(\mu, \mu') &= \left( - \int \int \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| (\mu - \mu')(d\theta)(\mu - \mu')(d\phi) \right)^{1/2} \\ &= \sqrt{\sum_{k \geq 1} \frac{1}{k} |\hat{\mu}_k - \hat{\mu}'_k|^2}, \end{aligned}$$

where  $\hat{\mu}_k = \int_{\mathbb{T}} e^{ik\theta} \mu(d\theta)$ . Then for any  $\varepsilon > 0$  there exists a finite constant  $C_\varepsilon$  such that for all  $\beta, \beta' > \varepsilon$

$$D(\mu_{HT}^\beta, \mu_{HT}^{\beta'}) \leq C_\varepsilon |\beta - \beta'|. \quad (3.14)$$

**Remark 3.6.** *We notice that if  $f \in L^2(\mathbb{T})$  with derivative in  $L^2(\mathbb{T})$ , we can set  $\|f\|_{\frac{1}{2}} = \sqrt{\sum_{k \geq 1} k |\hat{f}_k|^2}$ . So, for any measure  $\nu$  with zero mass we deduce that*

$$\int_{\mathbb{T}} f(\theta) \nu(d\theta) = \sum_{k \neq 0} \hat{f}_k \widehat{\nu}_k = \sum_{k \neq 0} k \hat{f}_k \frac{\widehat{\nu}_k}{k}.$$

Then, by Cauchy-Schwartz inequality, we obtain

$$\left| \int_{\mathbb{T}} f(\theta) \nu(d\theta) \right|^2 \leq \left| \sum_{k \neq 0} k \hat{f}_k \sum_{k \neq 0} \frac{\widehat{\nu}_k}{k} \right|^2 \leq 4 \|f\|_{\frac{1}{2}}^2 D(\nu, 0)^2. \quad (3.15)$$

Combining (3.14) and (3.15) we conclude that for any function  $f$  with finite  $\|f\|_{\frac{1}{2}}$  norm, the map  $\beta \rightarrow \int_{\mathbb{T}} f d\mu_{HT}^\beta$  is Lipschitz for  $\beta > 0$ . As a consequence, the moments of  $\mu_{HT}^\beta$  are almost surely differentiable with respect to  $\beta$ .

The proof of Lemma 3.5 follows the lines of the corresponding one in [31].

*Proof of Lemma 3.5.* The proofs of points a) and b) exploit the same ideas, thus we prove just point a).

For all  $\beta > \beta' > 0$  we have

$$\mathcal{F}^{(V,\beta)}(\mu_{HT}^\beta) \leq \mathcal{F}^{(V,\beta)}(\mu_{HT}^{\beta'}) = \mathcal{F}^{(V,\beta')}(\mu_{HT}^{\beta'}) + (\beta - \beta') E(\mu_{HT}^{\beta'})$$

and

$$\mathcal{F}^{(V,\beta')}(\mu_{HT}^{\beta'}) \leq \mathcal{F}^{(V,\beta')}(\mu_{HT}^{\beta}) = \mathcal{F}^{(V,\beta)}(\mu_{HT}^{\beta}) + (\beta' - \beta)E(\mu_{HT}^{\beta}),$$

so that

$$(\beta - \beta')E(\mu_{HT}^{\beta}) \leq \mathcal{F}^{(V,\beta)}(\mu_{HT}^{\beta}) - \mathcal{F}^{(V,\beta')}(\mu_{HT}^{\beta'}) < (\beta - \beta')E(\mu_{HT}^{\beta'}).$$

Since  $E(\mu_{HT}^{\beta})$ , and  $E(\mu_{HT}^{\beta'})$  are finite we obtain the claim.

We now move to the proof of point c). Setting  $\Delta\mu = \mu_{HT}^{\beta'} - \mu_{HT}^{\beta}$  we deduce that

$$\begin{aligned} 0 &\geq \mathcal{F}^{(\beta,V)}(\mu_{HT}^{\beta}) - \mathcal{F}^{(\beta,V)}(\mu_{HT}^{\beta'}) \\ &= 2 \int_{\mathbb{T}} V(\theta) \Delta\mu(d\theta) - 2\beta \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \mu_{HT}^{\beta'}(d\theta) \Delta\mu(d\phi) - \beta \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \Delta\mu(d\theta) \Delta\mu(d\phi) \\ &\quad + \int_{\mathbb{T}} \ln(\mu_{HT}^{\beta}(\theta)) \mu_{HT}^{\beta}(d\theta) - \int_{\mathbb{T}} \ln(\mu_{HT}^{\beta'}(\theta)) \mu_{HT}^{\beta'}(d\theta) \\ &= \int_{\mathbb{T}} \ln \left( \frac{\mu_{HT}^{\beta}(\theta)}{\mu_{HT}^{\beta'}(\theta)} \right) \mu_{HT}^{\beta}(d\theta) + 2(\beta' - \beta) \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \mu_{HT}^{\beta'}(d\theta) \Delta\mu(d\phi) \\ &\quad - \beta \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \Delta\mu(d\theta) \Delta\mu(d\phi), \end{aligned}$$

where in the third line we used (3.13). Since  $\int_{\mathbb{T}} \ln \left( \frac{\mu_{HT}^{\beta}(\theta)}{\mu_{HT}^{\beta'}(\theta)} \right) \mu_{HT}^{\beta}(d\theta) \leq 0$  by Jensen's inequality, we deduce that

$$\beta D(\mu_{HT}^{\beta}, \mu_{HT}^{\beta'})^2 \leq 2(\beta - \beta') \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \mu_{HT}^{\beta'}(d\theta) \Delta\mu(d\phi). \quad (3.16)$$

For convenience, we introduce a new probability measure  $\nu \in \mathcal{M}(\mathbb{T})$  in the previous expression, in this way we deduce

$$\begin{aligned} \beta D(\mu_{HT}^{\beta'}, \mu_{HT}^{\beta'})^2 &\leq 2(\beta - \beta') \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| (\mu_{HT}^{\beta'} - \nu)(d\theta) \Delta\mu(d\phi) \\ &\quad + 2(\beta - \beta') \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \nu(d\theta) \Delta\mu(d\phi). \end{aligned}$$

We chose  $\nu$  in such a way that the function  $g_{\nu}(\phi) = \int_{\mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \nu(d\theta)$  is in  $L^2(\mathbb{T})$  with derivative in  $L^2(\mathbb{T})$ . With this choice of  $\nu$  and applying (3.15) we conclude that there exists a constant  $C$  such that

$$\left| \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| \nu(d\theta) \Delta\mu(d\phi) \right| \leq CD(\mu_{HT}^{\beta}, \mu_{HT}^{\beta'}).$$

Moreover, taking the Fourier transform and apply again the Cauchy-Schwartz inequality (3.15) we obtain that

$$\left| \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right| (\mu_{HT}^{\beta'} - \nu)(d\theta) \Delta\mu(d\phi) \right| = \left| \sum_{k \geq 1} \frac{1}{k} \left( \widehat{\mu_{HT}^{\beta'} - \nu} \right)_k \overline{\widehat{\Delta\mu}_k} \right| \leq D(\mu_{HT}^{\beta'}, \nu) D(\mu_{HT}^{\beta'}, \mu_{HT}^{\beta}),$$

since  $D(\mu_{HT}^{\beta'}, \nu)$  is bounded, from (3.16) we conclude that there exists a constant  $R$  such that

$$D(\mu_{HT}^{\beta}, \mu_{HT}^{\beta'}) \leq \frac{R}{\beta} (\beta - \beta'),$$

from which (3.14) follows. □

For convenience, we define  $F_{HT}(V, \beta)$  as the value of the functional at the minimizer, namely

$$F_{HT}(V, \beta) := \mathcal{F}^{(V, \beta)}(\mu_{HT}^\beta).$$

The quantity  $F_{HT}(V, \beta)$  is referred to as free energy of the Circular beta ensemble at high temperature. It is a standard result that (see e.g. [26])

$$F_{HT}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_N^{HT}(V, \beta) \quad (3.17)$$

where the partition function  $Z_N^{HT}(V, \beta)$  of the Circular beta ensemble at high-temperature is defined in (3.3).

**Remark 3.7.** We notice that from (3.6) and (3.17) we can also obtain the free energy  $F_{HT}(V, \beta)$  from the partition function  $Z_N^{HT}(V, \beta)$  of the CMV matrix ensemble (3.3), namely:

$$F_{HT}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{\ln(Z_N^{HT}(V, \beta))}{N} - \ln(2).$$

The literature related to the high-temperature regime of the classical beta ensemble is quite broad. For completeness, we mention some of the results in the field. In [7, 8, 20, 25, 32, 62, 63] the authors explicitly computed the mean density of states for the classical Gaussian, Laguerre, Jacobi, and Circular beta ensemble at high-temperature. In [7, 8, 25, 32] the densities of states are computed as a solution of some particular ordinary differential equations. On the other hand, in [20, 62, 63] the authors reconstruct the densities from the moment generating functions. Several authors [12, 40, 47, 48, 51, 64] investigated the local fluctuations of the eigenvalues, they observed that in this regime they are described by a Poisson process. In particular, in [40] Lambert studied the local fluctuations for general Gibbs ensembles on  $N$ -dimensional manifolds, moreover he also studied the asymptotic behaviour of the maximum eigenvalue for the classical beta ensemble at high-temperature. In [24, 25] the loop equations for the classical beta ensemble at high-temperature are studied, in particular in [24] a special kind of duality between high and low temperature is underlined. There are also some results for higher dimensional Coulomb gases [6, 55]. It is worth to mention also the work [43], where some new tridiagonal random matrix ensembles related to the classical  $\beta$  one at high temperature are defined.

## 4 Proof of Theorem 2.1

The generalized Gibbs ensemble of the Ablowitz-Ladik lattice in (1.11) is very close to the probability distribution (3.5) of the Circular beta ensemble at high-temperature with an external source. Indeed, the only difference between the two ensembles is the exponent of the terms  $(1 - |\alpha_j|)$ . Our first main result contained in Theorem 2.1 relates the mean density of states of the Ablowitz-Ladik lattice to the mean density of states of the Circular beta ensemble at high-temperature.

To prove the result, we use the *moment matching technique* and the following lemma to prove that the two measures in (2.2) coincide.

**Lemma 4.1.** ([10, Lemma B.1 - B.2]) *Let  $d\sigma, d\sigma'$  be two measures with the same moment sequence  $\{u^{(\ell)}\}_{\ell \geq 0}$ . If*

$$\liminf_{\ell \rightarrow \infty} \frac{(u^{(2\ell)})^{\frac{1}{2\ell}}}{\ell} < \infty,$$

*then  $d\sigma = d\sigma'$ .*

Next, we define the *free energy* for the Ablowitz-Ladik lattice as:

$$F_{AL}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z_N^{AL}(V, \beta), \quad (4.1)$$

where the partition function  $Z_N^{AL}(V, \beta)$  is defined in (1.12). The next proposition shows that the free energy  $F_{AL}(V, \beta)$  of the Generalized Gibbs ensemble of the Ablowitz-Ladik lattice and the free energy  $F_{HT}(V, \beta)$  in (3.17) of the Circular beta ensemble at high-temperature are related. This fact allows us to calculate the moments of the mean density of states of the CMV matrix  $E$  in (3.2) and of the Lax matrix  $\mathcal{E}$  in (1.6).

**Proposition 4.2.** *The free energy  $F_{AL}(V, \beta)$  in (4.1) of the AL lattice and the free energy  $F_{HT}(V, \beta)$  in (3.17) of the Circular beta ensemble at high-temperature are analytic with respect to  $\beta > 0$ , and are related by*

$$\partial_\beta (\beta F_{HT}(V, \beta)) + \ln(2) = F_{AL}(V, \beta). \quad (4.2)$$

*The moments of the density of states  $\mu_{AL}^\beta$  of the Lax matrix  $\mathcal{E}$  in (1.6) endowed with the probability measure (1.11) and the moments of the density of states  $\mu_{HT}$  of the Circular beta ensemble in the high-temperature regime (3.5) are related to the free energies  $F_{AL}(V, \beta)$  and  $F_{HT}(V, \beta)$  by*

$$\int_{\mathbb{T}} e^{i\theta m} \mu_{AL}^\beta(d\theta) = \partial_t F_{AL} \left( V + \frac{t}{2} z^m, \beta \right) \Big|_{t=0}, \quad \int_{\mathbb{T}} e^{i\theta m} \mu_{HT}^\beta(d\theta) = \partial_t F_{HT} \left( V + \frac{t}{2} z^m, \beta \right) \Big|_{t=0}. \quad (4.3)$$

Since the proof of this proposition is rather technical, we postpone it to the Appendix A. We are now ready to prove the first main Theorem 2.1.

*Proof of Theorem 2.1.* First, we define  $c_n(\beta) := \int e^{i\theta n} \mu_{AL}^\beta(d\theta)$ ,  $d_n(\beta) := \int e^{i\theta n} \mu_{HT}^\beta(d\theta)$ . Since the eigenvalues of  $\mathcal{E}$  lie on the unit circle, we deduce the following chain of inequalities:

$$|c_n(\beta)| = \frac{|\mathbf{E}[\mathrm{Tr}(\mathcal{E}^n)]|}{2N} \leq \frac{\mathbf{E}[|\mathrm{Tr}(\mathcal{E}^n)|]}{2N} \leq 1.$$

Thus, from Lemma 4.1, we obtain that the measure  $\mu_{AL}^\beta(d\theta)$  is uniquely characterized by its moments.

Next we observe that from Proposition 4.2 we obtain the relation

$$c_n(\beta) = \partial_\beta (\beta d_n(\beta)) \quad (4.4)$$

between the moments of the measures  $\mu_{AL}^\beta(\theta)$  and  $\mu_{HT}^\beta(\theta)$  respectively.

This, together with (4.4) and Remark 3.6 implies

$$\mu_{AL}^\beta(\theta) = \partial_\beta \left( \beta \mu_{HT}^\beta(\theta) \right) \quad a.e..$$

□

Our next main result provides an explicit expression of the mean density of states  $\mu_{HT}(\theta)$  for the potential  $V(z) = \eta \Re(z)$ . This generalizes the result by Gross and Witten [30] and Baik-Deift-Johansson [11] obtained for finite temperature to the high-temperature regime.

## 5 Proof of Theorem 2.2

The proof of Theorem 2.2 consists of mainly two parts: we first derive from the variational equations with respect to the functional  $\mathcal{F}^{(V, \beta)}$ , the double confluent Heun equation (2.4). Then we show that such equation admits an analytic solution in any compact sets of the complex plane containing the origin. From Theorem 3.4 we know that the density  $\mu_{HT}^\beta$  is characterized as the unique minimizer of the functional (3.12). We follow the ideas developed in [7, 8, 16, 25] to find this minimizer explicitly. We consider the Euler-Lagrange equation of the functional (3.12), namely

$$\frac{\delta \mathcal{F}^{(V, \beta)}}{\delta \mu} = 2V(\theta) - 2\beta \int_{\mathbb{T}} \ln \sin \left( \frac{|\theta - \phi|}{2} \right) \mu(\phi) d\phi + \ln \mu(\theta) = C(V, \beta), \quad a.e. \quad (5.1)$$

where the equation holds almost everywhere and where  $C(V, \beta)$  is a constant depending on the potential and  $\beta$ , but not on the variable  $\theta$ . Differentiating the Euler-Lagrange equation (5.1) at the minimizer  $\mu_{HT}^\beta(\theta)$  with respect to  $\theta$  we obtain the following integral equation (see [32, Proposition 2.5]):

$$\partial_\theta \mu_{HT}^\beta(\theta) + \mu_{HT}^\beta(\theta) [2\partial_\theta (V(\theta)) + \beta H \mu_{HT}^\beta(\theta)] = 0, \quad (5.2)$$

where  $H$  is the Hilbert transform defined on  $L^2(\mathbb{T})$  as

$$H \mu_{HT}^\beta(\theta) = -\text{p.v.} \int_{\mathbb{T}} \cot \left( \frac{\theta - \phi}{2} \right) \mu_{HT}^\beta(\phi) d\phi$$

and p.v. is the Cauchy principal value, that is the limit as  $\varepsilon \rightarrow 0$  of the integral on the torus  $\mathbb{T}$  restricted to the domain  $|e^{i\theta} - e^{i\phi}| > \varepsilon$ . We notice that the Hilbert transform  $H$  is diagonal on the bases of exponential  $\{e^{in\theta}\}_{n \in \mathbb{Z}}$ , meaning that

$$He^{in\theta} = 2\pi i \operatorname{sgn}(n)e^{in\theta},$$

where  $\operatorname{sgn}(\cdot)$  is the sign function with the convention that  $\operatorname{sgn}(0) = 0$ . Setting  $e^{i\theta} = z$  and  $e^{i\phi} = w$ , we recognize the Riesz - Herglotz kernel  $\frac{z+w}{z-w}$  expressed as

$$\frac{z+w}{z-w} = -i \cot\left(\frac{\theta-\phi}{2}\right).$$

Therefore

$$\int_{\mathbb{T}} \cot\left(\frac{\theta-\phi}{2}\right) \mu(\phi) d\phi = i + 2 \int_{S^1} \frac{\mu(\phi)|_{e^{i\phi}=w} dw}{z-w},$$

where  $S^1$  is the anticlockwise oriented circle, and we used the normalization condition  $\int_{\mathbb{T}} \mu(\phi) d\phi = 1$ . In the following, in order to simplify the notation, we indicate  $\mu(\phi)|_{e^{i\phi}=w}$  just as  $\mu(w)$ . We can recast (5.2) in the form

$$z\partial_z\mu(z) + \mu(z) \left[ 2z\partial_z V(z) - \beta + 2i\beta \text{p.v.} \int_{S^1} \mu(w) \frac{dw}{z-w} \right] = 0.$$

For  $z \in \mathbb{C} \setminus S^1$  let us define

$$G(z) := \int_{S^1} \mu(w) \frac{dw}{w-z} = \frac{i}{2} - \frac{1}{2} \int_{\mathbb{T}} \cot\left(\frac{\theta-\phi}{2}\right) \mu(\phi) d\phi,$$

and for  $z \in S^1$  let  $G_{\pm}(z) = \lim_{\tilde{z} \rightarrow z} G(\tilde{z})$  for  $\tilde{z}$  inside and outside the unit circle respectively. Then

$$\begin{aligned} G_{\pm}(z) &= \pm\pi i\mu(z) + \text{p.v.} \int_{S^1} \mu(w) \frac{dw}{w-z} \\ &= \pm\pi i\mu(z) + \frac{i}{2} - \frac{2iz\partial_z V(z)}{2\beta} - \frac{iz\partial_z\mu(z)}{2\beta\mu(z)}. \end{aligned}$$

This implies that for  $z \in S^1$  one has

$$\begin{aligned} G_+(z) + G_-(z) &= i - \frac{2iz\partial_z V(z)}{\beta} - \frac{iz\partial_z\mu(z)}{\beta\mu(z)}, \\ G_+(z) - G_-(z) &= 2\pi i\mu(z). \end{aligned}$$

Multiplying the two previous expressions one obtains:

$$G_+(z)^2 - G_-(z)^2 = 2\pi i\mu(z) \left( i - \frac{2iz\partial_z V(z)}{\beta} - \frac{zi\partial_z\mu(z)}{\beta\mu(z)} \right).$$

In order to proceed we have to specify the potential  $V(z)$ , in our case we will consider

$$V(z) = \frac{\eta}{2} \left( z + \frac{1}{z} \right).$$

Applying the Sokhotski-Plemelj formula to the above boundary value problem one obtains

$$G^2(z) = i \int_{S^1} \frac{\mu(w)}{w-z} dw - \frac{i\eta}{\beta} \int_{S^1} \frac{(w-\bar{w})\mu(w)}{w-z} dw - \frac{i}{\beta} \int_{S^1} \frac{w\partial_w\mu(w)}{w-z} dw. \quad (5.3)$$

The second term in the r.h.s. of the above expression gives

$$\begin{aligned} \int_{S^1} \frac{(w-\bar{w})\mu(w)}{w-z} dw &= \int_{S^1} \frac{w \pm z}{w-z} \mu(w) dw + \frac{1}{z} \int_{S^1} \mu(w) \left( -\frac{1}{w-z} + \frac{1}{w} \right) dw \\ &= \left( zG(z) + i\lambda - \frac{G(z)}{z} + \frac{i}{z} \right), \end{aligned}$$

where we have defined

$$\lambda := -i \int_{S^1} \mu(w) dw, \quad \lambda \in \mathbb{R}. \quad (5.4)$$

The third term in the r.h.s. of (5.3) gives

$$\begin{aligned} \int_{S^1} \frac{w \partial_w \mu(w)}{w-z} dw &= \int_{S^1} \partial_w \mu(w) dw + z \int_{S^1} \frac{\partial_w \mu(w)}{w-z} dw \\ &= z \int_{S^1} \frac{\mu(w)}{(w-z)^2} dw = z \partial_z G(z), \end{aligned}$$

where in these last relations we use the results of Theorem 3.4 about the regularity of  $\mu$ . Now we can rewrite (5.3) as

$$G^2(z) = iG(z) - \frac{i\eta}{\beta} \left( zG(z) + i\lambda - \frac{G(z)}{z} + \frac{i}{z} \right) - \frac{iz \partial_z G(z)}{\beta}. \quad (5.5)$$

**Remark 5.1.** In the above ODE, the parameter  $\lambda = \lambda(\eta, \beta)$  depends via (5.4) implicitly on the function  $G(z)$ . Our strategy to solve the above equation is to consider  $\lambda$  as a free parameter that is uniquely fixed by the analytic properties of the function  $G(z)$ .

We can now turn the non-linear first order ODE (5.5) into a linear second order ODE through the substitution

$$G(z) = i + \frac{izv'(z)}{\beta v(z)}, \quad (5.6)$$

getting:

$$z^2 v''(z) + (-\eta + z(\beta + 1) + \eta z^2) v'(z) + \eta \beta (z + \lambda) v(z) = 0, \quad (5.7)$$

which is the DCH equation in (2.4). The solutions to this equation have generically essential singularities at  $z = 0$  and  $z = \infty$  and the local description near the singularities depends on the parameter  $\eta$  and  $\beta$ . Indeed we have that the two fundamental solutions near  $z = 0$  have the following asymptotic behaviour

$$v_1^{(0)}(z) = e^{\eta(z+\frac{1}{z})} z^{1-\beta} \kappa_1(\eta, \beta, \lambda; z), \quad -\frac{3\pi}{2} < \arg(\eta z) < \frac{3\pi}{2}, \quad (5.8)$$

$$v_2^{(0)}(z) = \kappa_2(\eta, \beta, \lambda, z), \quad -\frac{\pi}{2} < \arg(\eta z) < \frac{5\pi}{2}, \quad (5.9)$$

where  $\kappa_j(\eta, \beta, \lambda; z)$ ,  $j = 1, 2$ , are asymptotic series in a neighbourhood of  $z = 0$ .

The quantity  $\lambda$  is usually referred to as *accessory parameter*. Since  $G(z)$  is analytic in the unit disk, continuous up to the boundary, and  $G(0) = i$ , we deduce that

$$v(z) = v_0 \exp \left[ -i \int_0^z \beta \frac{G(s) - i}{s} ds \right], \quad v_0 \neq 0,$$

has to be analytic in the unit disk. For this reason we seek for a solution  $v(z)$  of the DCH equation that is analytic in the unit disk and such that  $v(z) \xrightarrow{z \rightarrow 0} v_0$ , where  $v_0$  is a nonzero constant.

**Construction of the analytic solution.** Of the fundamental solutions (5.8) and (5.9), only the solution (5.9) has a chance of being analytic near  $z = 0$ . This occurs if we are able to make the asymptotic series defined by  $\kappa_2(\eta, \beta, \lambda, z)$ , into a convergent series. To simplify notation instead of writing  $\kappa_2(\eta, \beta, \lambda, z)$ , we simply write  $v(z) = v(\eta, \beta, \lambda; z)$  and we look for a solution of (5.7) in the form

$$v(z) = \sum_{k=0}^{\infty} a_k z^k. \quad (5.10)$$

This implies the following recurrence relations for the coefficients  $\{a_k\}_{k \in \mathbb{N}}$

$$\eta(a_0 \lambda \beta - a_1) = 0, \quad (5.11)$$

$$a_k(k^2 + k\beta + \lambda\beta\eta) + \eta(k-1+\beta)a_{k-1} - \eta(k+1)a_{k+1} = 0, \quad k > 0, \quad (5.12)$$

where we have the freedom to chose  $\lambda$  and  $a_0$ . Generically, the above recurrence relation for the coefficients  $\{a_k\}_{k \in \mathbb{N}}$  gives a divergent series in (5.10). To obtain a convergent series, we follow the ideas in [13, 61].

We start by considering the  $2 \times 2$  matrices  $R_k^{(s)}$  defined as

$$R_k^{(s)} = M_k M_{k+1} \dots M_s, \quad s \geq k, \quad M_k = \begin{pmatrix} 1 + \frac{\lambda\beta\eta}{k(k+\beta)} & \frac{\eta^2}{k(k+\beta+1)} \\ 1 & 0 \end{pmatrix}, \quad (5.13)$$

which satisfy the recurrence relation  $R_k^{(s)} = R_k^{(s-1)} M_s$ . The next lemma shows that the limit of  $R_k^{(s)}$  as  $s \rightarrow \infty$  exists.

**Lemma 5.2.** *Let  $R_k^{(s)}$  be the matrix defined in (5.13). Then the limit of  $R_k^{(s)}$  as  $s \rightarrow \infty$  exists and*

$$R_k := \lim_{s \rightarrow \infty} R_k^{(s)}. \quad (5.14)$$

The matrices  $R_k$ ,  $k \geq 1$  satisfy the descending recurrence relation:

$$R_k = M_k R_{k+1} \quad k \geq 1. \quad (5.15)$$

Furthermore each entry of the matrix  $R_k = R_k(\beta, \eta, \lambda)$  is differentiable with respect to the parameters  $\beta$ ,  $\eta$ , and  $\lambda$ .

Since the proof of this lemma is rather technical, we defer it to the appendix B.

Further, let us define the following function:

$$\xi(\eta, \beta, \lambda) := \left( \lambda \quad \frac{\eta}{\beta+1} \right) R_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (5.16)$$

We are now ready to prove the following result that will give us a necessary condition to fix the value of  $\lambda$ .

**Proposition 5.3.** *For the values of  $\lambda$  such that*

$$\xi(\eta, \beta, \lambda) = 0, \quad (5.17)$$

where  $\xi(\eta, \beta, \lambda)$  is defined in (5.16), the Double Confluent Heun equation (5.7) admits a non-zero solution  $v = v(z, \eta, \beta)$  defined by the series (5.10) that is uniformly convergent in  $|z| \leq r$  with  $r \geq 1$ . The corresponding coefficients  $\{a_k\}_{k \in \mathbb{N}}$  of the Taylor expansion (5.10) are given by the relation

$$a_0 = \frac{1}{\beta} \begin{pmatrix} 1 & 0 \end{pmatrix} R_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (5.18)$$

$$a_k = (-1)^k \frac{\eta^k}{k!(k+\beta)} \begin{pmatrix} 0 & 1 \end{pmatrix} R_k \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad k \geq 1, \quad (5.19)$$

where the matrices  $R_k$  are defined in (5.14). For each  $\lambda$  satisfying (5.17), the solution  $v(z)$  of the DCH equation (5.7), analytic at zero is unique up to a multiplicative factor.

*Proof.* First, we show that choosing  $a_k$  according to (5.18)-(5.19) we obtain a solution of the recurrence (5.12). We notice that due to the recurrence relation for the matrices  $R_k$  (5.15), we have that:

$$\begin{pmatrix} 0 & 1 \end{pmatrix} R_k \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \end{pmatrix} R_{k+1} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Thus, applying the previous equation and (5.18)-(5.19), we can recast (5.12) as:

$$\begin{aligned} & \left[ (-1)^{k-1} \frac{\eta^k}{(k-1)!} \begin{pmatrix} 1 & 0 \end{pmatrix} R_k + (-\eta)^k \frac{k(k+\beta) + \eta\lambda\beta}{k!(k+\beta)} \begin{pmatrix} 1 & 0 \end{pmatrix} R_{k+1} \right. \\ & \left. + (-1)^k \frac{\eta^{k+2}}{k!(k+1+\beta)} \begin{pmatrix} 0 & 1 \end{pmatrix} R_{k+1} \right] \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ & = \frac{(-\eta)^k}{(k-1)!} \left[ - \begin{pmatrix} 1 & 0 \end{pmatrix} R_k + \left( 1 + \frac{\lambda\beta\eta}{k(k+\beta)} \quad \frac{\eta^2}{k(k+1+\beta)} \right) R_{k+1} \right] \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 0, \end{aligned}$$

where in the last equality we have enforced (5.15). Next we can rewrite (5.11) in terms of the matrix  $R_1$  exploiting (5.18)-(5.19), namely

$$0 = \left( \lambda \quad \frac{\eta}{\beta+1} \right) R_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \xi(\eta, \beta, \lambda),$$

which is exactly (5.16). Since the entries of the matrices  $R_k$  are uniformly bounded, the solution  $v(z) = \sum_{k \geq 0} a_k z^k$  with  $a_k$  as in (5.19), defines a uniformly convergent Taylor series in  $|z| < r$  for any  $r \geq 0$  and in particular for any  $r > 1$ .

To show that the solution analytic at  $z = 0$  is unique up to a constant, we consider the Wronkian  $W(v, \tilde{v})(z)$  of two independent solution  $v$  and  $\tilde{v}$  of the Double Confluent Heun equation (5.7), namely

$$W(z) = e^{-\eta(z+\frac{1}{z})} z^{-(\beta+1)} (v'(z)\tilde{v}(z) - v(z)\tilde{v}'(z)).$$

Since  $W'(z) = 0$ , it follows that  $W(z) = C$  a constant. If by contradiction we suppose that there are two analytic solutions at  $z = 0$ , then from the above relation we obtain

$$e^{-\eta z} (v'(z)\tilde{v}(z) - v(z)\tilde{v}'(z)) = C e^{\frac{\eta}{z}} z^{\beta+1}.$$

If  $\eta \neq 0$  the left-hand side of the above equation is analytic and the right-hand side is not, that is clearly a contradiction. If  $\eta = 0$  then (5.7) becomes:

$$z^2 v''(z) + z(\beta + 1)v'(z) = 0.$$

The above equation has two independent solutions, one is the constant solution, which is analytic, the other one is  $v(z) = C z^{-\beta}$  which is not analytic since  $\beta > 0$ . □

**Remark 5.4.** We observe that the equation (5.17) does not uniquely determine  $\lambda$ . Indeed, as it is shown in Figure 1 the function  $\xi(\eta, \beta, \lambda)$  may have several zeros for given  $\eta$  and  $\beta$ .




Figure 1: Plots of  $\xi(\eta, \beta, \lambda)$  for various values of  $\eta, \beta$

**Choice of the parameter  $\lambda$ .** We will now prove that the parameter  $\lambda$  is uniquely determined in a neighbourhood of  $\eta = 0$  by requiring that the solution  $v = v(z, \eta, \beta)$  depends continuously on the parameter  $\eta$ .

**Lemma 5.5.** *There exists an  $\varepsilon > 0$  such that for all  $\eta \in (-\varepsilon, \varepsilon)$  and  $\beta > 0$  there is a unique  $\lambda = \lambda(\eta, \beta)$  such that  $\xi(\eta, \beta, \lambda(\eta, \beta)) = 0$ .*

*Proof.* When  $\eta = 0$  the matrix  $R_j = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$  so that the only solution of the equation (5.17)  $\xi(\eta = 0, \beta, \lambda) = 0$  is  $\lambda = 0$ . To show the existence of the solution (5.17) for  $\lambda = \lambda(\eta, \beta)$  near  $\eta = 0$ , we use

the implicit function theorem. We have to show that  $\partial_\lambda \xi(\eta, \beta, \lambda)|_{(0, \beta, 0)} \neq 0$ . For the purpose, we need to evaluate

$$\partial_\lambda (M_k)_{(\eta=0, \lambda=0)} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix},$$

where  $M_k$  is defined in (5.14). This equation implies that

$$\partial_\lambda (\xi(\eta, \beta, \lambda))|_{(0, \beta, 0)} = (1 \ 0) \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 1.$$

Thus we can apply the implicit function theorem, and we obtain the claim.  $\square$

We conclude the proof of Theorem 2.2. When  $\eta = 0$  the only analytic solution of DCH equation is  $v(z) = c$ ,  $c \in \mathbb{C}$ . In this case, in principle  $\lambda$  is undetermined. However, from Theorem 3.4 the minimizer  $\mu_{HT}^\beta$  of (3.12) is the uniform measure on the circle and therefore from equation 5.4 one has  $\lambda = 0$ . From Lemma 5.5 when  $\eta \in (-\varepsilon, \varepsilon)$ , there exists a unique  $\lambda(\eta, \beta)$  that satisfies equation (5.17) and such that  $\lambda(\eta = 0, \beta) = 0$  and therefore by Proposition 5.3 we obtain for  $\eta \in (-\varepsilon, \varepsilon)$ , the unique solution  $v(z, \eta, \beta)$  of the DCH equation analytic in any compact set  $|z| \leq r$ , with  $r > 0$  and in particular when  $r = 1$ . Because of lemma 5.2 the solution  $v(z, \eta, \beta)$  is differentiable with respect to the parameters  $\eta$  and  $\beta$ .

We remark that  $v(z) \neq 0$  on the unit disc  $\mathbb{D}$  because of the relation (5.6) between the analytic function  $G(z)$  and  $v(z)$ .

To complete our proof we recover the explicit expression of  $\mu_{HT}^\beta(\theta)$  from  $G(z)$  and  $v(z)$  using the *Poisson representation formula* (see for example [57, Chapter 1]):

$$\mu_{HT}^\beta(\theta) = -\frac{1}{2\pi} - \frac{\Re(iG(e^{i\theta}))}{\pi\beta} = \frac{1}{2\pi} + \frac{1}{\pi\beta} \Re \left( \frac{zv'(z)}{v(z)} \Big|_{z=e^{i\theta}} \right). \quad (5.20)$$

$\square$




Figure 2: The mean density of states  $\mu_{HT}^\beta$  for different parameters.

In Figure 2 we plotted the density of states of the Circular beta ensemble in the high-temperature regime with potential  $V(z) = \eta \Re(z)$ . To produce this picture and Figure 1, we used extensively the NumPy [33], and matplotlib [34] libraries.

**Remark 5.6.** *The Gross-Witten [30] and Baik-Deift-Johannson [11] solution is obtained by making the substitution  $\eta \rightarrow \beta\eta$  and  $\beta \rightarrow \infty$  in equation (3.12) which gives the functional*

$$\mathcal{F}^{(\eta)}(\mu) := \int_{\mathbb{T} \times \mathbb{T}} \int_{\mathbb{T} \times \mathbb{T}} \ln \left| \sin \left( \frac{\theta - \phi}{2} \right) \right|^{-1} \mu(d\theta) \mu(d\phi) + 2\eta \int_{\mathbb{T}} \cos(\theta) \mu(d\theta).$$

The minimizer is  $\mu(\theta) = \frac{1}{2\pi} (1 - 2\eta \cos \theta)$  with  $0 \leq 2\eta \leq 1$ . To recover such solution from the DCH equation (5.7) and the expression (5.20) it is sufficient to make the ansatz  $v(z) = e^{\beta\lambda z}$  that solves the DCH equation in the limit  $\beta \rightarrow \infty$  and when  $\lambda = -\eta$ .

## A Proof of Proposition 4.2

First, we prove the relation between the free energies (4.2) namely:

$$\partial_\beta(\beta F_{HT}(V, \beta)) + \ln(2) = F_{AL}(V, \beta), \quad (\text{A.1})$$

and we show that  $F_{HT}(V, \beta)$  is analytic with respect to  $\beta > 0$ .

The above expression is equivalent to:

$$\partial_\beta \left( \beta \lim_{N \rightarrow \infty} \frac{\ln(Z_N^{HT}(V, \beta))}{N} \right) = \lim_{N \rightarrow \infty} \frac{\ln(Z_N^{AL}(V, \beta))}{N}.$$

To prove this relation, we will use the so-called transfer operator technique [36, 39, 52]. We are considering a potential of the form  $\text{Tr}(V(\mathcal{E}))$  as in (1.10) which is of finite range  $K$ , meaning that it can be expressed as a sum of local quantities, i.e. depending on a finite number  $2K$  of variables, with  $K$  independent of  $N$  [49]. For example, if  $V(z) = \Re(z)$ , then  $\text{Tr}(\mathcal{E}) = -2 \sum_{j=1}^N \Re(\alpha_j \bar{\alpha}_{j+1})$  and in this case the range is  $K = 1$ . Let  $N = KM + L$  with  $M, L \in \mathbb{N}$  and  $L < K$ . We split the coordinates  $(\alpha_1, \dots, \alpha_N)$  into  $M$  blocks of length  $K$  and a reminder of length  $L$ , and we define  $\tilde{\alpha}_j = (\alpha_{K(j-1)+1}, \alpha_{K(j-1)+2}, \dots, \alpha_{Kj})$ . In this notation,

$$(\alpha_1, \dots, \alpha_N) = (\overbrace{\tilde{\alpha}_1, \dots, \tilde{\alpha}_M}^{KM}, \overbrace{\alpha_{KM+1}, \dots, \alpha_N}^L),$$

and we can rewrite the potential as

$$\text{Tr}(V(\mathcal{E})) = \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) + W(\tilde{\alpha}_M, \overbrace{\alpha_{KM+1}, \dots, \alpha_N, \alpha_1, \dots, \alpha_{K-L}}^K),$$

where  $W$  is a continuous function  $W : \mathbb{D}^K \times \mathbb{D}^K \rightarrow \mathbb{R}$ , and  $W(\tilde{\alpha}_1, \tilde{\alpha}_2) = W(\tilde{\alpha}_2, \tilde{\alpha}_1)$ . The last term in the above expression is different from the others since we may have an off-set of length  $L$ , due to periodicity. In the case  $V(z) = \Re(z)$ , then  $W(\alpha_1, \alpha_2) = -2\Re(\alpha_1 \bar{\alpha}_2)$  and there is no off-set.

For convenience, we define

$$\tilde{\alpha}_{M+1} = (\alpha_{KM+1}, \dots, \alpha_N, \alpha_1, \dots, \alpha_{K-L}).$$

We can now rewrite  $Z_N^{AL}(V, \beta)$  in (1.12) as

$$Z_N^{AL}(V, \beta) = \int_{\mathbb{D}^N} \prod_{j=1}^N (1 - |\alpha_j|^2)^{\beta-1} \exp \left( - \sum_{\ell=1}^M W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) \right) d^2 \alpha.$$

We are now in position to apply the transfer operator technique to compute this partition function. On  $L^2(\mathbb{D}^K)$  we introduce the scalar product

$$(f, g) = \int_{\mathbb{D}^K} f(\mathbf{z}) \overline{g(\mathbf{z})} d\mathbf{z}, \quad (\text{A.2})$$

where  $\mathbf{z} = (z_1, \dots, z_K)$ . This scalar product induces a norm on  $L^2(\mathbb{D})$  and also a norm on bounded operators  $T : L^2(\mathbb{D}^K) \rightarrow L^2(\mathbb{D}^K)$  as

$$\|T\|_2 := \sup_{f: \|f\|_2=1} \|Tf\|_2,$$

where  $\|f\|_2$  is the standard  $L^2$  norm.

Let  $\zeta = (\zeta_1, \dots, \zeta_{2K})$  with  $\zeta_{K+j} = \zeta_j > 0$  for  $j = 1, \dots, K$ , we define the continuous family of transfer operators  $\mathcal{T}_\zeta : L^2(\mathbb{D}^K) \rightarrow L^2(\mathbb{D}^K)$  as

$$(\mathcal{T}_\zeta f)(\tilde{\alpha}_2) = \int_{\mathbb{D}^K} f(\tilde{\alpha}_1) \prod_{j=1}^{2K} (1 - |\alpha_j|^2)^{\frac{\zeta_j-1}{2}} \exp(-W(\tilde{\alpha}_1, \tilde{\alpha}_2)) d^2 \tilde{\alpha}_1. \quad (\text{A.3})$$

We notice that  $\mathcal{T}_\zeta$  is symmetric with respect to the scalar product (A.2), indeed  $(f, \mathcal{T}g) = (\mathcal{T}f, g)$ . Furthermore,  $\mathcal{T}_\zeta$  is an integral operator whose kernel  $\prod_{j=1}^{2K} (1 - |\alpha_j|^2)^{\frac{\zeta_j-1}{2}} \exp(-W(\tilde{\alpha}_1, \tilde{\alpha}_2))$  belongs to

$L^2(\mathbb{D}^K \times \mathbb{D}^K)$ , and therefore  $\mathcal{T}_\zeta$  is an Hilbert-Schmidt operator. We conclude that there exists a complete set of normalized eigenfunctions  $\{\psi_j\}_{j \geq 1}$  with real eigenvalues  $\{\lambda_j\}_{j \geq 1}$  in non-increasing order, such that:

$$(\mathcal{T}_\zeta \psi_j)(\mathbf{z}, V, \zeta) = \lambda_j(V, \zeta) \psi_j(\mathbf{z}', V, \zeta), \quad (\text{A.4})$$

$$\sum_{n=1}^{\infty} \bar{\psi}_n(\mathbf{z}, V, \zeta) \psi_n(\mathbf{z}', V, \zeta) = \delta_{\mathbf{z}}(\mathbf{z}'), \quad (\text{A.5})$$

where  $\delta_{\mathbf{z}}(\cdot)$  is the Dirac delta function at  $\mathbf{z} \in \mathbb{D}^K$ . For clearness, we collect a series of properties that the operator  $\mathcal{T}_\zeta$  fulfils:

- $\sum_{j=1}^{\infty} |\lambda_j(V, \zeta)|^2 < \infty$ , since  $\mathcal{T}_\zeta$  is Hilbert-Schmidt, therefore it is in particular trace-class (see [35, Chapter V.2.4]);
- The biggest eigenvalue  $\lambda_1(\zeta, V)$  is simple, positive and  $\lambda_1(\zeta, V) > |\lambda_n(\zeta, V)|$  for all  $n \geq 2$  (see [65, Theorem 137.4]);
- The biggest eigenvalue  $\lambda_1(\zeta, V)$  and its eigenfunction  $\psi_1(\mathbf{z}, \zeta, V)$  are analytic functions of the parameters  $\zeta$  (see [35, Chapter VII, Theorem 1.8]).

We artificially rewrite  $Z_N^{AL}$  as

$$\begin{aligned} Z_N^{AL}(V, \beta) &= \int_{\mathbb{D}^{N+K}} \delta_{\tilde{\alpha}_1}(\gamma) \prod_{\ell=1}^K (1 - |\gamma_\ell|^2)^{\frac{\beta-1}{2}} \prod_{\ell=1}^K (1 - |\alpha_j|^2)^{\frac{\beta-1}{2}} \prod_{\ell=K+1}^N (1 - |\alpha_j|^2)^{\beta-1} \\ &\quad \times \exp \left( - \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \gamma_1, \dots, \gamma_{K-L}) \right) d^2 \alpha d^2 \gamma, \end{aligned}$$

where  $\gamma = (\gamma_1, \dots, \gamma_K)$  and  $\gamma \in \mathbb{D}^K$ . We can use (A.5) with  $\zeta = \beta = \overbrace{(\beta, \dots, \beta)}^{2K}$  to rewrite the previous equation as:

$$\begin{aligned} Z_N^{AL}(V, \beta) &= \sum_{n=1}^{\infty} \int_{\mathbb{D}^{N+K}} \bar{\psi}_n(\gamma, V, \beta) \psi_n(\tilde{\alpha}_1, V, \beta) \prod_{\ell=1}^K (1 - |\gamma_\ell|^2)^{\frac{\beta-1}{2}} \prod_{\ell=1}^K (1 - |\alpha_j|^2)^{\frac{\beta-1}{2}} \prod_{\ell=K+1}^N (1 - |\alpha_j|^2)^{\beta-1} \\ &\quad \times \exp \left( - \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \gamma_1, \dots, \gamma_{K-L}) \right) d^2 \alpha d^2 \gamma \\ &= \sum_{n=1}^{\infty} \int_{\mathbb{D}^N} \bar{\psi}_n(\gamma, V, \beta) (\mathcal{T}_\beta \psi_n)(\tilde{\alpha}_2) \prod_{\ell=K+1}^{2K} (1 - |\alpha_j|^2)^{\frac{\beta-1}{2}} \prod_{\ell=2K+1}^N (1 - |\alpha_j|^2)^{\beta-1} \\ &\quad \times \exp \left( - \sum_{\ell=2}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \gamma_1, \dots, \gamma_{K-L}) \right) d^2 \alpha d^2 \gamma \\ &= \sum_{n=1}^{\infty} \lambda_n(V, \beta) \int_{\mathbb{D}^N} \bar{\psi}_n(\gamma, V, \beta) \psi_n(\tilde{\alpha}_2) \prod_{\ell=K+1}^{2K} (1 - |\alpha_j|^2)^{\frac{\beta-1}{2}} \prod_{\ell=2K+1}^N (1 - |\alpha_j|^2)^{\beta-1} \\ &\quad \times \exp \left( - \sum_{\ell=2}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \gamma_1, \dots, \gamma_{K-L}) \right) d^2 \alpha d^2 \gamma. \end{aligned}$$

In the above integral, from the first to the second relation we identify the integral operator  $\mathcal{T}_\beta$  where

$\beta = \overbrace{(\beta, \dots, \beta)}^{2K}$ . We repeatedly apply  $\mathcal{T}_\beta$  and (A.4) another  $M-2$  times to the above integral, to obtain:

$$Z_N^{AL}(V, \beta) = \sum_{n=1}^{\infty} (\lambda_n(V, \beta))^{M-1} R_n, \quad (\text{A.6})$$

$$\begin{aligned} R_n &= \int_{\mathbb{D}^{2K+L}} \bar{\psi}_n(\gamma, V, \beta) \psi_n(\tilde{\alpha}_M, V, \beta) \prod_{\ell=1}^K (1 - |\gamma_\ell|^2)^{\frac{\beta-1}{2}} \prod_{\ell=(M-1)K+1}^{MK} (1 - |\alpha_j|^2)^{\frac{\beta-1}{2}} \\ &\quad \times \prod_{\ell=MK+1}^N (1 - |\alpha_j|^2)^{\beta-1} \exp(-W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \gamma_1, \dots, \gamma_{K-L})) \prod_{\ell=(M-1)K+1}^N d^2 \alpha_j \prod_{\ell=1}^K d^2 \gamma_\ell. \end{aligned} \quad (\text{A.7})$$

The modulus of the reminder  $|R_n|$  in (A.7) can be bounded from above and below by two constants  $C_1, C_2 > 0$  independent of  $N$ , therefore we conclude from (A.6) that

$$F_{AL}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{1}{N} \ln (Z_N^{AL}(V, \beta)) = - \frac{1}{K} \ln (\lambda_1(V, \beta)).$$

Since  $\lambda_1(V, \beta)$  is analytic for  $\beta > 0$ , see [35, Chapter VII, Theorem 1.8], and strictly positive, see [65, Theorem 137.4], we conclude that  $F_{AL}(V, \beta)$  is analytic with respect to  $\beta$ .

We can apply the same procedure to the partition function  $Z_N^{HT}(V, \beta)$  in (3.5). Also in this case the potential  $\text{Tr}(V(\tilde{E}))$  with  $V$  as in (1.10) and the matrix  $\tilde{E}$  as in (3.4) is of finite range  $K$ , meaning that it can be expressed as a sum of local quantities [49]. More precisely, assuming  $N = KM + L$  with  $L < K$  and  $M, N, L \in \mathbb{N}$  we have

$$\text{Tr}(V(\tilde{E})) = \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) + W(\overbrace{0, \dots, 0}^{K-1}, -1, \tilde{\alpha}_1) + W(\tilde{\alpha}_M, \underbrace{\alpha_{KM+1}, \dots, \alpha_N}_{K}, \overbrace{0, \dots, 0}^{K-L}).$$

For example for  $V(z) = z^2 + \bar{z}^2$  one has  $K = 2$  and  $N = 2M + L$  where  $L = 0, 1$ , depending on the parity of  $N$ . The vector  $\tilde{\alpha}_\ell$  takes the form  $\tilde{\alpha}_\ell = (\alpha_{2\ell-1}, \alpha_{2\ell})$  for  $\ell = 1, \dots, M$ . In this notation, we can rewrite the potential as

$$\text{Tr}(V(\tilde{E})) = \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) + \delta_{L,0} W(\tilde{\alpha}_M, \alpha_N, 0) + \underbrace{2\Re(\alpha_1^2 + 2\bar{\alpha}_2 \rho_1^2)}_{=W(0, -1, \alpha_1, \alpha_2)},$$

where in this case  $W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) = 2\Re \left[ \sum_{s=0}^1 (\alpha_{2\ell-1+s} \bar{\alpha}_{2\ell+s})^2 - 2\alpha_{2\ell-1+s} \bar{\alpha}_{2\ell+1+s} \rho_{2\ell+s}^2 \right]$ . In this way, the partition function can be written in the form

$$\begin{aligned} Z_N^{HT}(V, \beta) &= \int_{\mathbb{D}^{N-1} \times S^1} \frac{d\alpha_N}{i\alpha_N} \prod_{j=1}^{N-1} d^2 \alpha_j (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N})-1} \\ &\times \exp \left( - \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \overbrace{0, \dots, 0}^{K-L}) - W(\overbrace{0, \dots, 0}^{K-1}, -1, \tilde{\alpha}_1) \right). \end{aligned} \quad (\text{A.8})$$

We want to apply the same technique as in the previous case, but we have to pay attention to one important detail: in this situation the eigenvalues and the eigenfunctions of the transfer operators will be dependent on the block number. Indeed, in this case the exponents of  $(1 - |\alpha_j|^2)$  are not identical, but they depend on the index  $j$  as in (A.8).

For this reason, we define

$$\zeta^{(1)} = \overbrace{\left( \beta \left( 1 - \frac{1}{N} \right), \beta \left( 1 - \frac{2}{N} \right), \dots, \beta \left( 1 - \frac{K}{N} \right), \beta \left( 1 - \frac{1}{N} \right), \beta \left( 1 - \frac{2}{N} \right), \dots, \beta \left( 1 - \frac{K}{N} \right) \right)}^{2K},$$

and

$$\zeta^{(j)} = \zeta^{(1)} - \beta^{\frac{j-1}{N}} \mathbf{K}, \quad j = 1, \dots, M-1,$$

where  $\mathbf{K}_j = K$  for  $j = 1, \dots, 2K$ . For  $K$  integer and  $K < N$  we introduce the multiplication operator  $\mathcal{M}_K : L^2(\mathbb{D}^K) \rightarrow L^2(\mathbb{D}^K)$  defined as

$$(\mathcal{M}_K f)(\alpha) = \prod_{j=1}^K (1 - |\alpha_j|^2)^{-\frac{K}{2N}} f(\alpha).$$

We observe that  $\mathcal{M}_{-K} = (\mathcal{M}_K)^{-1}$  and the operators  $\mathcal{T}_{\zeta^{(j)}} : L^2(\mathbb{D}^K) \rightarrow L^2(\mathbb{D}^K)$  defined in (A.3) satisfy the relation

$$\mathcal{T}_{\zeta^{(j+1)}} = \mathcal{M}_K \mathcal{T}_{\zeta^{(j)}} \mathcal{M}_K, \quad j = 1, \dots, M-1.$$

We recall that the operators  $\mathcal{T}_{\zeta^{(j)}}$  are compact and symmetric.

Let us define the operator

$$\tilde{\mathcal{T}}_M = \mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \mathcal{T}_{\zeta^{(M-2)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(1)}}, \quad (\text{A.9})$$

we notice that it is a compact operator, since all  $\mathcal{T}_{\zeta^{(j)}}$  are Hilbert-Schmidt and the multiplication operator  $\mathcal{M}_K$  is bounded. We will now prove the following technical result:

**Proposition A.1.** *Let  $\tilde{\mathcal{T}}_M$  as in (A.9) and  $Z_N^{HT}$  as in (A.8) then:*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left( \frac{Z_N^{HT}}{\text{Tr}(\tilde{\mathcal{T}}_M)} \right) = 0, \quad (\text{A.10})$$

here by  $\text{Tr}(\tilde{\mathcal{T}}_M)$  we indicate the standard trace on  $L^2$ .

*Proof.* We will estimate both  $Z_N^{HT}$ , and  $\text{Tr}(\tilde{\mathcal{T}}_M)$  from above and below, then combining these estimates we will obtain (A.10). We start with  $Z_N^{HT}$ .

$$\begin{aligned} Z_N^{HT} &= \int_{\mathbb{D}^{N-1} \times S^1} \prod_{j=1}^{N-1} d^2 \alpha_j \frac{d\alpha_N}{i\alpha_N} \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \\ &\times \exp \left( - \sum_{\ell=1}^{M-1} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \overbrace{0, \dots, 0}^{K-L}) - W(\overbrace{0, \dots, 0}^{K-1}, -1, \tilde{\alpha}_1) \right). \end{aligned}$$

We notice that there exists two constant  $C(V, \beta)$  and  $c(V, \beta)$ , independent of  $N$ , such that

$$c(V, \beta) \leq \exp \left( -W(\tilde{\alpha}_1, \tilde{\alpha}_2) - W(\tilde{\alpha}_M, \alpha_{KM+1}, \dots, \alpha_N, \overbrace{0, \dots, 0}^{K-L}) - W(\overbrace{0, \dots, 0}^{K-1}, -1, \tilde{\alpha}_1) \right) \leq C(V, \beta).$$

From the previous inequalities, we deduce that

$$\begin{aligned} c(V, \beta) &\int_{\mathbb{D}^{N-1} \times S^1} \frac{d\alpha_N}{i\alpha_N} \prod_{j=1}^{N-1} d^2 \alpha_j (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp \left( - \sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) \right) \leq Z_N^{HT} \\ &\leq C(V, \beta) \int_{\mathbb{D}^{N-1} \times S^1} \frac{d\alpha_N}{i\alpha_N} \prod_{j=1}^{N-1} d^2 \alpha_j \prod_{j=1}^{N-1} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp \left( - \sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) \right) \end{aligned}$$

We can explicitly integrate in  $\alpha_j$  for  $j = 1, \dots, K$  and  $j = (M-2)K+1, \dots, N$  using the formula

$$\int_{\mathbb{D}} (1 - |z|^2)^{t-1} d^2 z = \pi t^{-1}, \quad (\text{A.11})$$

obtaining that there exists a constant  $C_1(V, \beta)$  and  $c_1(V, \beta)$  depending on  $V, \beta$ , but not on  $N$ , such that

$$Z_N^{HT} \leq C_1(V, \beta) N^{2K+L-1} \int_{\mathbb{D}^{(M-3)K}} \prod_{j=K+1}^{(M-2)K} d^2 \alpha_j \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp \left( - \sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) \right), \quad (\text{A.12})$$

and

$$Z_N^{HT} \geq c_1(V, \beta) N^{2K+L-1} \int_{\mathbb{D}^{(M-3)K}} \prod_{j=K+1}^{(M-2)K} d^2 \alpha_j \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp \left( - \sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1}) \right). \quad (\text{A.13})$$

We can proceed analogously to estimate the trace of  $\tilde{\mathcal{T}}_M$ :

$$\begin{aligned} \text{Tr}(\tilde{\mathcal{T}}_M) &= \int_{\mathbb{D}^{(M-2)K}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{j}{N}) - \frac{1}{2}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{(M-2)K+j}{N}) - \frac{1}{2}} \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \\ &\quad \times \exp\left(-\sum_{j=1}^{M-3} W(\tilde{\alpha}_j, \tilde{\alpha}_{j+1}) - W(\tilde{\alpha}_{M-2}, \tilde{\alpha}_1)\right) \prod_{j=1}^{M-2} d^2 \tilde{\alpha}_j. \end{aligned}$$

As before, we notice that there exist two constants  $\tilde{C}(V, \beta)$ , and  $\tilde{c}(V, \beta)$ , independent of  $N$ , such that

$$\tilde{c}(V, \beta) < \exp(-W(\tilde{\alpha}_1, \tilde{\alpha}_2) - W(\tilde{\alpha}_{M-2}, \tilde{\alpha}_1)) < \tilde{C}(V, \beta).$$

From these inequalities, we deduce that

$$\begin{aligned} \tilde{c}(V, \beta) &\int_{\mathbb{D}^{(M-2)K}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{j}{N}) - \frac{1}{2}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{(M-2)K+j}{N}) - \frac{1}{2}} \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \\ &\quad \times \exp\left(-\sum_{j=2}^{M-3} W(\tilde{\alpha}_j, \tilde{\alpha}_{j+1})\right) \prod_{j=1}^{M-2} d^2 \tilde{\alpha}_j < \text{Tr}(\tilde{\mathcal{T}}_M) \\ &< \tilde{C}(V, \beta) \int_{\mathbb{D}^{(M-2)K}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{j}{N}) - \frac{1}{2}} \prod_{j=1}^K (1 - |\alpha_j|^2)^{\frac{\beta}{2}(1 - \frac{(M-2)K+j}{N}) - \frac{1}{2}} \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \\ &\quad \times \exp\left(-\sum_{j=2}^{M-3} W(\tilde{\alpha}_j, \tilde{\alpha}_{j+1})\right) \prod_{j=1}^{M-2} d^2 \tilde{\alpha}_j. \end{aligned}$$

We can now explicitly integrate in  $\alpha_j$  for  $j = 1, \dots, K$ , obtaining the following inequalities

$$\text{Tr}(\tilde{\mathcal{T}}_M) \leq \tilde{C}_1(V, \beta) \int_{\mathbb{D}^{(M-3)K}} \prod_{j=1}^{(M-2)K} d^2 \alpha_j \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp\left(-\sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1})\right), \quad (\text{A.14})$$

$$\text{Tr}(\tilde{\mathcal{T}}_M) \geq \tilde{c}_1(V, \beta) \int_{\mathbb{D}^{(M-3)K}} \prod_{j=1}^{(M-2)K} d^2 \alpha_j \prod_{j=K+1}^{(M-2)K} (1 - |\alpha_j|^2)^{\beta(1 - \frac{j}{N}) - 1} \exp\left(-\sum_{\ell=2}^{M-3} W(\tilde{\alpha}_\ell, \tilde{\alpha}_{\ell+1})\right), \quad (\text{A.15})$$

here  $\tilde{C}_1(V, \beta)$ , and  $\tilde{c}_1(V, \beta)$  are constants depending on  $V, \beta$ , but not on  $N$ . Combining (A.12)-(A.13)-(A.14)-(A.15) we deduce (A.10).  $\square$

Applying the previous proposition, we can express the Free energy of the Circular beta ensemble in the high-temperature regime in terms of  $\text{Tr}(\tilde{\mathcal{T}}_M)$ :

$$\begin{aligned} F_{HT}(V, \beta) &= -\lim_{N \rightarrow \infty} \frac{1}{N} \ln(Z_N^{HT}) = -\lim_{N \rightarrow \infty} \frac{1}{N} \left( \ln\left(\frac{Z_N^{HT}}{\text{Tr}(\tilde{\mathcal{T}}_M)}\right) + \ln(\text{Tr}(\tilde{\mathcal{T}}_M)) \right) \\ &= -\lim_{N \rightarrow \infty} \frac{\ln(\text{Tr}(\tilde{\mathcal{T}}_M))}{N}, \end{aligned} \quad (\text{A.16})$$

where in the last equality we used Proposition A.1.

Thus, we have to understand the behaviour of  $\text{Tr}(\tilde{\mathcal{T}}_M)$ , and for the purpose we need to carefully analyse the compact operators  $\mathcal{T}_{\zeta^{(j)}}$ .

Let  $\psi_n(\mathbf{z}, V, \zeta^{(j)})$  be the eigenfunctions of  $\mathcal{T}_{\zeta^{(j)}}$  with corresponding eigenvalues  $\lambda_n(V, \zeta^{(j)})$  in not increasing order. From a generalized version of Jentzsch's Theorem (see [65, Theorem 137.4]), we deduce

that  $|\lambda_n(V, \zeta^{(j)})| < \lambda_1(V, \zeta^{(j)})$  for all  $n \geq 2$ . Moreover, the largest eigenvalues of each  $\mathcal{T}_{\zeta^{(j)}}$  is an analytic function of the parameters  $\zeta^{(j)}$  [35, Chapter VII, Theorem 1.8], so we conclude that

$$\lambda_1(V, \zeta^{(j+1)}) = \lambda_1(V, \zeta^{(j)}) \left( 1 + O\left(\frac{1}{N}\right) \right).$$

Furthermore, we claim that

$$\left( \psi_1(\mathbf{z}, V, \zeta^{(j+1)}), \mathcal{M}_K \psi_n(\mathbf{z}, V, \zeta^{(j)}) \right) = \delta_{1,n} + O\left(\frac{1}{N}\right).$$

Indeed, let us consider the following chain of equality

$$\begin{aligned} \left( \psi_1(\mathbf{z}, V, \zeta^{(j+1)}), \mathcal{M}_K \psi_n(\mathbf{z}, V, \zeta^{(j)}) \right) &= \left( \psi_1(\mathbf{z}, V, \zeta^{(j)}) \left( 1 + O\left(\frac{1}{N}\right) \right), \psi_n(\mathbf{z}, V, \zeta^{(j)}) \left( 1 + O\left(\frac{1}{N}\right) \right) \right) \\ &= \delta_{1,n} + O\left(\frac{1}{N}\right) \end{aligned}$$

where we used the analyticity of  $\psi_1(\mathbf{z}, \zeta^{(j+1)}, V)$  in the parameter  $\zeta^{(j+1)}$ , we expanded the operator  $\mathcal{M}_K$  in power series of  $1/N$ , and the orthogonality of  $\{\psi_n(\mathbf{z}, \zeta^{(j)}, V)\}_{n=1}^{\infty}$ .

We are now in position to prove the following proposition

**Proposition A.2.** *Let  $\{\psi_n(\mathbf{z}, V, \zeta^{(j)})\}_{n=1}^{\infty}$  be the eigenfunctions of  $\mathcal{T}_{\zeta^{(j)}}$  (A.3) with corresponding eigenvalues  $\{\lambda_n(V, \zeta^{(j)})\}_{n=1}^{\infty}$  in not increasing order. Consider the operator  $\tilde{\mathcal{T}}_M$  in (A.9), then there exists some constants  $c, a_j, b_j, j = 1, \dots, M-1$  independent of  $N$ , and so on  $M$ , such that :*

$$\left( \psi_1(\mathbf{z}, V, \zeta^{(1)}), \tilde{\mathcal{T}}_M \psi_1(\mathbf{z}, V, \zeta^{(1)}) \right) = \prod_{j=1}^{M-1} \lambda_1(V, \zeta^{(j)}) \left( 1 + \frac{a_j}{N} + O\left(\frac{1}{N^2}\right) \right), \quad (\text{A.17})$$

$$\left| \sum_{\ell \geq 2} \left( \psi_\ell(\mathbf{z}, V, \zeta^{(1)}), \tilde{\mathcal{T}}_M \psi_\ell(\mathbf{z}, V, \zeta^{(1)}) \right) \right| \leq c \prod_{j=2}^{M-1} \lambda_1(V, \zeta^{(j)}) \left( 1 + \frac{b_j}{N} + O\left(\frac{1}{N}\right) \right) \left( |\text{Tr}(\mathcal{T}_{\zeta^{(1)}})| + \lambda_1(V, \zeta^{(1)}) \right) \quad (\text{A.18})$$

*Proof.* To simplify the notation, we will drop the  $V$  dependence of the eigenvalues  $\lambda_n(V, \zeta^{(j)})$ , and of the eigenfunctions  $\psi_n(\mathbf{z}, V, \zeta^{(j)})$ . First, we will focus on (A.17). We will prove it by induction on  $M$ . For  $M = 2$ , we have that  $\tilde{\mathcal{T}}_M = \mathcal{M}_K \mathcal{T}_{\zeta^{(1)}}$ , so we have to compute:

$$\left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(1)}} \psi_1(\mathbf{z}, \zeta^{(1)}) \right) = \lambda_1(\zeta^{(1)}) \left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \psi_1(\mathbf{z}, \zeta^{(1)}) \right).$$

We can expand the operator  $\mathcal{M}_K$  in series of  $1/N$ , so, for  $N$  big enough, we deduce that there exists a constant  $a_1$  independent of  $N$  such that:

$$\left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \psi_1(\mathbf{z}, \zeta^{(1)}) \right) = 1 + \frac{a_1}{N} + O\left(\frac{1}{N^2}\right),$$

so the first inductive step is proved.

For general  $M$ , we have to compute:

$$\begin{aligned} &\left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(1)}} \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(1)}} \psi_1(\mathbf{z}, \zeta^{(1)}) \right) \\ &= \lambda_1(\zeta^{(1)}) \left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(2)}} \mathcal{M}_K \psi_1(\mathbf{z}, \zeta^{(1)}) \right) \\ &= \lambda_1(\zeta^{(1)}) \left( \psi_1(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(2)}} \psi_1(\mathbf{z}, \zeta^{(1)}) \right) \left( 1 + \frac{\tilde{a}_1}{N} + O\left(\frac{1}{N}\right) \right). \end{aligned}$$

In the second equality, we have expanded the operator  $\mathcal{M}_K$  in  $1/N$  powers as we did in the previous proof. Thanks to [35, Chapter VII, Theorem 1.8], we know that the eigenfunctions  $\psi_1(\mathbf{z}, \zeta^{(j)})$  are analytic functions of the parameter  $\zeta^{(j)}$ , so, for  $N$  big enough, there exists a constant  $\xi_1$  independent of  $N$  such that:

$$\psi_1(\mathbf{z}, \zeta^{(1)}) = \psi_1(\mathbf{z}, \zeta^{(2)}) \left( 1 + \frac{\xi_1}{N} + O\left(\frac{1}{N^2}\right) \right).$$

So we conclude that there exists a constant  $a_1$  independent of  $N$  such that

$$\begin{aligned} & \left( \psi_1(\mathbf{z}, \zeta^{(1)}), \tilde{\mathcal{T}}_M \psi_1(\mathbf{z}, \zeta^{(1)}) \right) \\ &= \lambda_1(\zeta^{(1)}) \left( 1 + \frac{a_1}{N} + O\left(\frac{1}{N^2}\right) \right) \left( \psi_1(\mathbf{z}, \zeta^{(2)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(2)}} \psi_1(\mathbf{z}, \zeta^{(2)}) \right). \end{aligned}$$

Applying the inductive hypothesis, we deduce (A.17).

We move to the proof of (A.18). To prove this inequality, we need some preparation. First, we notice that there exist constants  $b_j$ ,  $\mathbf{d}$ , and  $\mathbf{c}$  independent of  $N$ , such that the following holds

$$\|\mathcal{M}_K \mathcal{T}_{\zeta^{(j)}}\|_2 \leq \lambda_1(\zeta^{(j)}) \left( 1 + \frac{b_j}{N} + O\left(\frac{1}{N^2}\right) \right), \quad (\text{A.19})$$

$$\|\mathcal{M}_K\|_2 \leq \mathbf{d} \left( \frac{\pi^K}{1 - \frac{2\beta K}{N}} \right)^{\frac{K}{2}} \leq \mathbf{c}. \quad (\text{A.20})$$

The first inequality is just a consequence of the expansion of  $\mathcal{M}_K$  in series of  $1/N$ . For the second one, one has just to compute the  $L^2$  norm of  $\mathcal{M}_K$  exploiting (A.11). Applying the previous inequalities, we obtain that

$$\begin{aligned} & \left| \sum_{\ell \geq 2} \left( \psi_\ell(\mathbf{z}, V, \zeta^{(1)}), \tilde{\mathcal{T}}_M \psi_\ell(\mathbf{z}, V, \zeta^{(1)}) \right) \right| \\ & \leq \left| \sum_{\ell \geq 2} \lambda_\ell(\zeta^{(1)}) \left( \psi_\ell(\mathbf{z}, \zeta^{(1)}), \mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(2)}} \mathcal{M}_K \psi_\ell(\zeta^{(1)}) \right) \right| \\ & \leq \|\mathcal{M}_K \mathcal{T}_{\zeta^{(M-1)}} \mathcal{M}_K \cdots \mathcal{M}_K \mathcal{T}_{\zeta^{(2)}} \mathcal{M}_K\|_2 \left| \sum_{\ell \geq 2} \lambda_\ell(\zeta^{(1)}) \right| \\ & \leq \|\mathcal{M}_K\|_2 \prod_{j=2}^{M-1} \|\mathcal{T}_{\zeta^{(j)}} \mathcal{M}_K\|_2 \left( |\text{Tr}(\mathcal{T}_{\zeta^{(1)}})| + \lambda_1(\zeta^{(1)}) \right) \\ & \leq \mathbf{c} \prod_{j=2}^{M-1} \lambda_1(V, \zeta^{(j)}) \left( 1 + \frac{b_j}{N} + O\left(\frac{1}{N}\right) \right) \left( |\text{Tr}(\mathcal{T}_{\zeta^{(1)}})| + \lambda_1(V, \zeta^{(1)}) \right). \end{aligned}$$

Here we have used the inequalities (A.19)-(A.20) in the last passage. Thus, we proved our claim.  $\square$

We remark that both  $\text{Tr}(\mathcal{T}_{\zeta^{(1)}})$ , and  $\lambda_1(V, \zeta^{(1)})$  are uniformly bounded, with a bound independent of  $N$ .

Applying Proposition (A.2) to (A.16) we obtain that:

$$\begin{aligned} F_{HT}(V, \beta) &= - \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left( \text{Tr}(\tilde{\mathcal{T}}) \right) = - \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left( \sum_{n \geq 1} \left( \psi_n(\mathbf{z}, V, \zeta^{(1)}), \tilde{\mathcal{T}} \psi_n(\mathbf{z}, V, \zeta^{(1)}) \right) \right) \\ &= - \lim_{N \rightarrow \infty} \left[ \frac{1}{N} \ln \left( \prod_{j=1}^{M-1} \lambda_1(V, \zeta^{(j)}) \left( 1 + \frac{a_j}{N} + O\left(\frac{1}{N^2}\right) \right) \right) \right. \\ & \quad \left. + \frac{1}{N} \ln \left( 1 + \frac{\sum_{\ell \geq 2} \left( \psi_\ell(\mathbf{z}, V, \zeta^{(1)}), \tilde{\mathcal{T}}_M \psi_\ell(\mathbf{z}, V, \zeta^{(1)}) \right)}{\prod_{j=1}^{M-1} \lambda_1(V, \zeta^{(j)}) \left( 1 + \frac{a_j}{N} + O\left(\frac{1}{N^2}\right) \right)} \right) \right], \end{aligned} \quad (\text{A.21})$$

applying (A.20), we deduce that the last term goes to zero, and since  $a_j$  are uniformly bounded we obtain that

$$F_{HT}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left( \prod_{j=1}^{M-1} \lambda_1(V, \zeta^{(j)}) \right).$$

Since the maximum eigenvalue of each  $\mathcal{T}_{\zeta^{(j)}}$  is positive and an analytic function of the parameters, we deduce that  $\lambda_1(V, \zeta^{(j)}) = \lambda_1 \left( V, \beta \left( 1 - \frac{jK}{N} \right) \right) + O(N^{-1})$ . Therefore, we can rewrite (A.21) as

$$F_{HT}(V, \beta) = - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^{M-1} \ln \left( \lambda_1 \left( V, \beta \left( 1 - \frac{jK}{N} \right) \right) \right) = - \frac{1}{K} \int_0^1 \ln(\lambda_1(V, \beta x)) dx.$$

This leads to (A.1), moreover, as a consequence of the previous relation, we deduce that  $F_{HT}(V, \beta)$  is analytic in  $\beta$  for  $\beta > 0$ .

We notice that the proof is heavily based on the assumption that the potential that we are considering is of finite range, otherwise our approach would not work.

We now prove the moments relations (4.3). Thus, we want to prove that

$$\int_{\mathbb{T}} e^{i\theta m} \mu_{HT}^\beta(d\theta) = \partial_t F_{HT} \left( V + \frac{t}{2} z^m, \beta \right) \Big|_{t=0}, \quad (\text{A.22})$$

$$\int_{\mathbb{T}} e^{i\theta m} \mu_{AL}^\beta(d\theta) = \partial_t F_{AL} \left( V + \frac{t}{2} z^m, \beta \right) \Big|_{t=0}. \quad (\text{A.23})$$

We focus on (A.22). From Remark 3.7, we know that  $F_{HT}(V, \beta) = \mathcal{F}^{(V; \beta)}(\mu_{HT}^\beta(\theta))$ , we recall that  $\mu_{HT}^\beta(\theta)$  is the density of the mean density of states of Circular beta ensemble at high-temperature and the functional  $\mathcal{F}^{(V; \beta)}$  is defined in (3.12). We write the Euler-Lagrange equation for this functional, getting that  $\mu_{HT}^\beta(\theta)$  satisfies:

$$2V(\theta) - 2\beta \int_{\mathbb{T}} \ln \left( \sin \left( \frac{|\theta - \gamma|}{2} \right) \right) \mu_{HT}^\beta(\gamma) d\gamma + \ln(\mu_{HT}^\beta(\theta)) + C(V, \beta) = 0, \quad (\text{A.24})$$

where  $C(V, \beta)$  is a constant not depending on  $\theta$ .

Consider the same functional as before, but with potential  $\tilde{V}(\theta) = V(\theta) + \frac{t}{2} e^{im\theta}$ :

$$\begin{aligned} \mathcal{F}^{(V(\theta) + \frac{t}{2} e^{im\theta}, \beta)}(\mu) &= 2 \int_{\mathbb{T}} V(\theta) \mu(\theta) d\theta + t \int_{\mathbb{T}} e^{im\theta} \mu(\theta) d\theta - \beta \int \int_{\mathbb{T} \times \mathbb{T}} \ln \sin \left( \frac{|\theta - \gamma|}{2} \right) \mu(\theta) \mu(\gamma) d\theta d\gamma \\ &\quad + \int_{\mathbb{T}} \ln(\mu(\theta)) \mu(\theta) d\theta + \ln(2\pi). \end{aligned}$$

Also this functional has a unique minimizer  $\mu^{(t)}(\theta)$ , and we notice that  $\mu^{(0)}(\theta) = \mu_{HT}^\beta(\theta)$ . Evaluating the previous functional at  $\mu^{(t)}(\theta)$ , and computing its derivative at  $t = 0$ , we deduce the following relation:

$$\begin{aligned} \partial_t \mathcal{F}^{(V(\theta) + \frac{t}{2} e^{im\theta}, \beta)}(\mu) \Big|_{t=0} &= 2 \int_{\mathbb{T}} V(\theta) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta + \int_{\mathbb{T}} e^{im\theta} \mu_{HT}^\beta(\theta) d\theta \\ &\quad - \beta \int \int_{\mathbb{T} \times \mathbb{T}} \ln \sin \left( \frac{|\theta - \gamma|}{2} \right) \mu_{HT}^\beta(\gamma) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta d\gamma + \int_{\mathbb{T}} \ln(\mu_{HT}^\beta(\theta)) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta. \end{aligned} \quad (\text{A.25})$$

Testing (A.24) against  $\partial_t \mu^{(t)}(\theta) \Big|_{t=0}$  we obtain

$$2 \int_{\mathbb{T}} V(\theta) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta - \beta \int \int_{\mathbb{T} \times \mathbb{T}} \ln \sin \left( \frac{|\theta - \gamma|}{2} \right) \mu_{HT}^\beta(\gamma) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta d\gamma + \int_{\mathbb{T}} \ln(\mu_{HT}^\beta(\theta)) \partial_t \mu^{(t)}(\theta) \Big|_{t=0} d\theta = 0,$$

where we used that  $\int_{\mathbb{T}} \partial_t \mu^{(t)}(\theta) d\theta = 0$ . Thus, we can simplify (A.25) as :

$$\partial_t \mathcal{F}^{(V(\theta) + \frac{t}{2} e^{im\theta}, \beta)}(\mu) \Big|_{t=0} = \int_{\mathbb{T}} e^{im\theta} \mu_{HT}^\beta(\theta) d\theta,$$

which is equivalent to (A.22).

To complete the proof of Proposition 4.2 we have to show that (A.23) holds. From the definition of mean density of states (3.11) we obtain that:

$$\int_{\mathbb{T}} e^{im\theta} \mu_{AL}^\beta(d\theta) = \lim_{N \rightarrow \infty} \frac{\mathbf{E}[\text{Tr}(\mathcal{E}^m)]}{2N} = - \lim_{N \rightarrow \infty} \frac{\partial_t \left( Z_N^{(AL)} \left( V + \frac{t}{2} z^m, \beta \right) \right) \Big|_{t=0}}{N Z_N^{(AL)}(V, \beta)},$$

where the expected value is taken with respect to the generalized Gibbs ensemble of the Ablowitz-Ladik lattice.

Exploiting (A.6) we can rewrite the previous expression as:

$$\begin{aligned} - \lim_{N \rightarrow \infty} \frac{\partial_t \left( Z_N^{(AL)} \left( V + \frac{t}{2} z^m, \beta \right) \right) \Big|_{t=0}}{N Z_N^{(AL)}(V, \beta)} &= - \lim_{N \rightarrow \infty} \frac{\partial_t \left( \sum_{\ell=1}^{\infty} \lambda_{\ell}^{M-1} \left( V + \frac{t}{2} z^m, \beta \right) \right) \Big|_{t=0}}{N \sum_{\ell=1}^{\infty} \lambda_{\ell}^{M-1}(V, \beta)} \\ &= - \frac{\partial_t \lambda_{\max} \left( \left( V + \frac{t}{2} z^m, \beta \right) \right) \Big|_{t=0}}{K \lambda_{\max}(V, \beta)} = \partial_t \left( F_{AL} \left( V + \frac{t}{2} z^m, \beta \right) \right) \Big|_{t=0}. \end{aligned}$$

Thus, we have completed the proof of Proposition 4.2.  $\square$

## B Proof of lemma 5.5

We prove (5.14) for  $k = 1$  and the cases  $k > 1$  easily follow. For convenience we consider the more general case of  $(\lambda, \eta, \beta) \in \mathbb{C} \times \mathbb{C} \times \{z \in \mathbb{C} | \Re z > 0\}$ .

Let us define  $R_1^{(s)} = \begin{pmatrix} f_s & h_s \\ p_s & q_s \end{pmatrix}$  where  $s \geq 1$ . It follows from (5.13) that

$$\begin{pmatrix} f_s & h_s \\ p_s & q_s \end{pmatrix} = \begin{pmatrix} f_{s-1} & h_{s-1} \\ p_{s-1} & q_{s-1} \end{pmatrix} \begin{pmatrix} 1 + \frac{\lambda\beta\eta}{s(s+\beta)} & \frac{\eta^2}{s(s+\beta+1)} \\ 1 & 0 \end{pmatrix}, \quad s > 1.$$

Note that in the case  $\eta = 0$  the lemma is trivially satisfied. We will show that all the sequences  $\{f_s, h_s, p_s, q_s\}_{s \geq 1}$  converge as  $s \rightarrow \infty$ , moreover  $h_s, q_s \xrightarrow{s \rightarrow \infty} 0$ . First of all, we notice that  $h_s = \frac{\eta^2 f_{s-1}}{s(s+\beta+1)}$  and  $q_s = \frac{\eta^2 p_{s-1}}{s(s+\beta+1)}$ , thus the convergence to zero of these two sequences follows from the convergence of  $p_s$  and  $f_s$  as  $s \rightarrow \infty$ . Moreover, the terms of these last two sequences obey to the 3-terms recurrence:

$$f_s = \left( 1 + \frac{\lambda\beta\eta}{s(s+\beta)} \right) f_{s-1} + \frac{\eta^2}{(s-1)(s+\beta)} f_{s-2}, \quad (\text{B.1})$$

and the same holds for  $p_s$  in place of  $f_s$ . Thus, we have just to prove that the sequence  $\{f_s\}_{s \geq 1}$  converges. We assume that  $(\lambda, \eta, \beta) \in \Omega$  where  $\Omega \subset \mathbb{C} \times \mathbb{C} \times \{z \in \mathbb{C} | \Re z > 0\}$  is a compact set. With this assumption we can give a bound to  $|f_s|$  from above as:

$$|f_s| \leq \left( 1 + \frac{2\eta^2 + |\lambda\beta\eta|}{s(s+\beta)} \right) \max(|f_{s-1}|, |f_{s-2}|).$$

Inductively, we deduce that there exists a constant  $C = C(\Omega)$  such that:

$$|f_s| \leq C \prod_{\ell=1}^s \left( 1 + \frac{2\eta^2 + |\lambda\beta\eta|}{\ell(\ell+\beta)} \right) \leq C \prod_{\ell=1}^{\infty} \left( 1 + \frac{2\eta^2 + |\lambda\beta\eta|}{\ell(\ell+\beta)} \right). \quad (\text{B.2})$$

Furthermore the infinite product on the right-hand side of (B.2) is convergent by a classical result, see for example [41, Chapter XIII, Lemma 1], this implies that the sequence  $\{f_s\}_{s \geq 1}$  is uniformly bounded. Moreover, we have that:

$$|f_{s+1} - f_s| \leq \frac{|f_s \lambda \beta \eta|}{(s+1)(s+1+\beta)} + \frac{\eta^2 |f_{s-1}|}{s(s-1+\beta)} \leq \tilde{C} \frac{\eta^2 + |\lambda\beta\eta|}{s(s-1+\beta)},$$

for some constant  $\tilde{C} > 0$  that depends on the compact set  $\Omega$ . This last equation implies that the sequence  $\{f_s\}_{s \geq 1}$  is a Cauchy sequence, thus it is convergent. So we get the claim (5.13). The claim (5.15) easily follows from (5.13).

Regarding the differentiability in the parameters  $\lambda, \eta$  and  $\beta$ , it follows from (B.1) that  $f_s = f_s(\lambda, \eta, \beta)$  is analytic in  $\Omega$ . Since  $f_s(\lambda, \eta, \beta) \rightarrow f(\lambda, \eta, \beta)$  as  $s \rightarrow \infty$  uniformly, then by Weierstrasse convergence theorem,  $f(\lambda, \eta, \beta)$  is analytic in  $\Omega$ .  $\square$

### Acknowledgements

We thank Herbert Spohn and Gaultier Lambert for the many discussions and suggestions during our time at MSRI. We thank Rostyslav Kozhan for useful comments on the manuscript. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930 while the author participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2-21 semester "Universality and Integrability in Random Matrix Theory and Interacting Particle Systems". This project has received funding from the European Union's H2020 research and innovation programme under the Marie Skłodowska–Curie grant No. 778010 *IPaDEGAN*. TG acknowledge the support of GNFM-INDAM group.

**Added note.** Independently, H. Spohn [59] discovered the connection between the Ablowitz-Ladik lattice and the circulant  $\beta$  ensemble at high temperature. He also calculated Generalized Gibbs Ensemble averaged field and currents and the associated hydrodynamic equations.

## References

- [1] M. J. ABLOWITZ, D. J. KAUP, AND A. C. NEWELL, *Coherent pulse propagation, a dispersive, irreversible phenomenon*, J. Math. Phys., 15 (1973), pp. 1852–1858.
- [2] M. J. ABLOWITZ AND J. F. LADIK, *Nonlinear differential-difference equations*, J. Math. Phys., 16 (1974), pp. 598–603.
- [3] ———, *Nonlinear differential-difference equations and Fourier analysis*, J. Math. Phys., 17 (1975), pp. 1011–1018.
- [4] M. J. ABLOWITZ, B. PRINARI, AND A. D. TRUBATCH, *Discrete and Continuous Nonlinear Schrödinger Systems*, Cambridge University Press, dec 2003.
- [5] M. ADLER AND P. VAN MOERBEKE, *Integrals over classical groups, random permutations, Toda and Toeplitz lattices*, Comm. Pure Appl. Math., 54 (2001), pp. 153–205.
- [6] G. AKEMANN AND S.-S. BYUN, *The high temperature crossover for general 2D Coulomb gases*, J. Stat. Phys., 175 (2019), pp. 1043–1065.
- [7] R. ALLEZ, J. P. BOUCHAUD, AND A. GUIONNET, *Invariant beta ensembles and the gauss-Wigner crossover*, Phys. Rev. Lett., 109 (2012), pp. 1–5.
- [8] R. ALLEZ, J. P. BOUCHAUD, S. N. MAJUMDAR, AND P. VIVO, *Invariant  $\beta$ -Wishart ensembles, crossover densities and asymptotic corrections to the Marčenko-Pastur law*, J. Phys. A Math. Theor., 46 (2013), pp. 1–26.
- [9] Y. ANGELOPOULOS, R. KILLIP, AND M. VISAN, *Invariant measures for integrable spin chains and an integrable discrete nonlinear Schrödinger equation*, SIAM J. Math. Anal., 52 (2020), pp. 135–163.
- [10] Z. BAI AND J. W. SILVERSTEIN, *Spectral Analysis of Large Dimensional Random Matrices*, Springer Series in Statistics, Springer New York, New York, NY, 2010.
- [11] J. BAIK, P. DEIFT, AND K. JOHANSSON, *On the distribution of the length of the longest increasing subsequence of random permutations*, J. Amer. Math. Soc., 12 (1999), pp. 1119–1178.
- [12] F. BENAYCH-GEORGES AND S. PÉCHÉ, *Poisson statistics for matrix ensembles at large temperature*, J. Stat. Phys., 161 (2015), pp. 633–656.
- [13] V. M. BUCHSTABER AND S. I. TERTYCHNYI, *Holomorphic solutions of the double confluent Heun equation associated with the RSJ model of the Josephson junction*, Theor. Math. Physics (Russian Fed.), 182 (2015), pp. 329–355.
- [14] M. CAFASSO, *Matrix biorthogonal polynomials on the unit circle and non-abelian Ablowitz-Ladik hierarchy*, J. Phys. A, 42 (2009), pp. 365211, 20.
- [15] M. J. CANTERO, L. MORAL, AND L. VELÁZQUEZ, *Minimal representations of unitary operators and orthogonal polynomials on the unit circle*, Linear Algebra Appl., 408 (2005), pp. 40–65.

- [16] P. DEIFT, T. KRIECHERBAUER, AND K. T.-R. MCLAUGHLIN, *New results on the equilibrium measure for logarithmic potentials in the presence of an external field.*, J. Approx. Theory, 95 (1998), pp. 388–475.
- [17] P. A. DEIFT, *Orthogonal polynomials and random matrices: a Riemann–Hilbert approach*, vol. 3 of Courant Lecture Notes in Mathematics, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [18] B. DOYON, *Lecture Notes On Generalised Hydrodynamics*, SciPost Phys. Lect. Notes, (2020), p. 18.
- [19] I. DUMITRIU AND A. EDELMAN, *Matrix models for beta ensembles*, J. Math. Phys., 43 (2002), pp. 5830–5847.
- [20] K. T. DUJ AND T. SHIRAI, *The mean spectral measures of random Jacobi matrices related to Gaussian beta ensembles*, Electron. Commun. Probab., 20 (2015), pp. 1–15.
- [21] N. M. ERCOLANI AND G. LOZANO, *A bi-Hamiltonian structure for the integrable, discrete non-linear Schrödinger system*, Physica D, 218 (2006), pp. 105–121.
- [22] A. S. FOKAS, A. R. ITS, A. A. KAPAEV, AND V. Y. NOVOKSHENOV, *Painlevé transcendents. The Riemann–Hilbert approach.*, American Mathematical Society, Providence, RI, 2006.
- [23] P. J. FORRESTER, *Log-gases and random matrices*, vol. 34 of London Mathematical Society Monographs Series, Princeton University Press, Princeton, NJ, 2010.
- [24] P. J. FORRESTER, *High-low temperature dualities for the classical  $\beta$ -ensembles*, arXiv preprint: 2103.11250, (2021).
- [25] P. J. FORRESTER AND G. MAZZUCA, *The classical  $\beta$ -ensembles with  $\beta$  proportional to  $1/n$ : From loop equations to dyson’s disordered chain*, Journal of Mathematical Physics, 62 (2021), p. 073505.
- [26] D. GARCÍA-ZELADA, *A large deviation principle for empirical measures on Polish spaces: application to singular Gibbs measures on manifolds*, Ann. Inst. Henri Poincaré Probab. Stat., 55 (2019), pp. 1377–1401.
- [27] M. GEKHTMAN AND I. NENCIU, *Multi-hamiltonian structure for the finite defocusing Ablowitz–Ladik equation*, Comm. Pure Appl. Math., 62 (2009), pp. 147–182.
- [28] L. GOLINSKII, *Schur flows and orthogonal polynomials on the unit circle*, Sb. Math., 197 (2006), pp. 1145 – 1165.
- [29] T. GRAVA, T. KRIECHERBAUER, G. MAZZUCA, AND K. D. T.-R. MCLAUGHLIN, *Correlation functions for a chain of short range oscillators.*, J. Stat. Phys., 183 (2021).
- [30] D. GROSS AND E. WITTEN, *Possible third-order phase transition in the large  $N$ -lattice gauge theory*, Physical Review D, 21 (1980), pp. 446 – 453.
- [31] A. GUIONNET AND R. MEMIN, *Large deviations for generalized gibbs ensembles of the classical Toda chain*, arXiv preprint: 2103.04858, (2021), pp. 1–30.
- [32] A. HARDY AND G. LAMBERT, *CLT for circular beta-ensembles at high temperature*, J. Funct. Anal., 280 (2021), pp. 108869, 40.
- [33] C. R. HARRIS, K. J. MILLMAN, S. J. VAN DER WALT, R. GOMMERS, P. VIRTANEN, D. COURNAPEAU, E. WIESER, J. TAYLOR, S. BERG, N. J. SMITH, R. KERN, M. PICUS, S. HOYER, M. H. VAN KERKWIJK, M. BRETT, A. HALDANE, J. F. DEL RÍO, M. WIEBE, P. PETERSON, P. GÉRARD-MARCHANT, K. SHEPPARD, T. REDDY, W. WECKESSER, H. ABBASI, C. GOHLKE, AND T. E. OLIPHANT, *Array programming with NumPy*, Nature, 585 (2020), pp. 357–362.
- [34] J. D. HUNTER, *Matplotlib: A 2d graphics environment*, Computing in Science & Engineering, 9 (2007), pp. 90–95.
- [35] T. KATO, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.

- [36] P. G. KEVREKIDIS, K. Ø. RASMUSSEN, AND A. R. BISHOP, *The Discrete Nonlinear Schrödinger Equation: A Survey of Recent Results*, Int. J. Mod. Phys. B, 15 (2001), pp. 2833–2900.
- [37] R. KILLIP AND I. NENCIU, *Matrix models for circular ensembles*, Int. Math. Res. Not., 2004 (2004), p. 2665.
- [38] ———, *CMV: the unitary analogue of Jacobi matrices*, Comm. Pure Appl. Math., 60 (2007), pp. 1148–1188.
- [39] J. A. KRUMHANSL AND J. R. SCHRIEFFER, *Dynamics and statistical mechanics of a one-dimensional model Hamiltonian for structural phase transitions*, Phys. Rev. B, 11 (1975), pp. 3535–3545.
- [40] G. LAMBERT, *Poisson statistics for Gibbs measures at high temperature*, Annales de l’Institut Henri Poincaré, Probabilités et Statistiques, 57 (2021), pp. 326 – 350.
- [41] S. LANG, *Complex Analysis*, vol. 103 of Graduate Texts in Mathematics, Springer New York, New York, NY, 1999.
- [42] O. LISOVYY AND A. NAIDIUK, *Accessory parameters in confluent heun equations and classical irregular conformal blocks*, ArXiv preprint: 2101.05715, (2021).
- [43] G. MAZZUCA, *On the mean density of states of some matrices related to the beta ensembles and an application to the Toda lattice*, arXiv preprint: 2008.04604, (2021), pp. 1–26.
- [44] K. T.-R. MCLAUGHLIN AND P. D. MILLER, *The  $\bar{\partial}$  steepest descent method and the asymptotic behavior of polynomials orthogonal on the unit circle with fixed and exponentially varying nonanalytic weights*, IMRP Int. Math. Res. Pap., (2006), pp. Art. ID 48673, 1–77.
- [45] M. L. MEHTA, *Random matrices*, vol. 142 of Pure and Applied Mathematics (Amsterdam), Elsevier/Academic Press, Amsterdam, third ed., 2004.
- [46] C. MENDEL AND H. SPHON, *Low temperature dynamics of the one-dimensional discrete nonlinear Schrödinger equation*, J. Stat. Mech. Theory Exp., 8 (2015), pp. P08028, 35 pp.
- [47] F. NAKANO AND K. D. TRINH, *Gaussian beta ensembles at high temperature: eigenvalue fluctuations and bulk statistics*, J. Stat. Phys., 173 (2018), pp. 295–321.
- [48] ———, *Poisson statistics for beta ensembles on the real line at high temperature*, J. Stat. Phys., 179 (2020), pp. 632–649.
- [49] I. NENCIU, *Lax pairs for the Ablowitz-Ladik system via orthogonal polynomials on the unit circle*, Int. Math. Res. Not., 2005 (2005), pp. 647–686.
- [50] F. W. NIJHOFF, V. G. PAPAGEORGIOU, H. W. CAPEL, AND G. R. W. QUISPEL, *The lattice Gelfand-Dikii hierarchy*, Inverse Problems, 8 (1992), p. 5970621.
- [51] C. PAKZAD, *Poisson statistics at the edge of Gaussian  $\beta$ -ensemble at high temperature*, ALEA Lat. Am. J. Probab. Math. Stat., 16 (2019), pp. 871–897.
- [52] M. PEYRARD AND A. R. BISHOP, *Statistical mechanics of a nonlinear model for DNA denaturation*, Phys. Rev. Lett., 62 (1989), pp. 2755–2758.
- [53] G. QUISPEL, F. NIJHOFF, H. CAPEL, AND J. VAN DER LINDEN, *Linear integral equations and nonlinear difference-difference equations*, Physica A: Statistical Mechanics and its Applications, 125 (1984), pp. 344–380.
- [54] A. RONVEAUX, *Heun’s differential equations*, Oxford University Press, Oxford, 1995.
- [55] N. ROUGERIE AND S. SERFATY, *Higher-dimensional Coulomb gases and renormalized energy functionals*, Comm. Pure Appl. Math., 69 (2016), pp. 519–605.
- [56] E. B. SAFF AND V. TOTIK, *Logarithmic potentials with external fields*, vol. 316 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 1997. Appendix B by Thomas Bloom.

- [57] B. SIMON, *Orthogonal Polynomials on the Unit Circle*, vol. 54.1 of Colloquium Publications, American Mathematical Society, Providence, Rhode Island, 2005.
- [58] H. SPOHN, *Generalized Gibbs ensembles of the classical Toda chain*, J. Stat. Phys., 180 (2020), pp. 4–22.
- [59] H. SPOHN, *Hydrodynamic equations for the Ablowitz-Ladik*, arXiv preprint:2107.04866, (2021).
- [60] H. SPOHN, *Hydrodynamic equations for the Toda lattice*, arXiv preprint: 2101.06528, (2021).
- [61] S. I. TERTYCHNIY, *General solution of overdamped Josephson junction equation in the case of phase-lock*, Electron. J. Differ. Equations, 2007 (2007), pp. 1–20.
- [62] H. D. TRINH AND K. D. TRINH, *Beta Laguerre ensembles in global regime*, arXiv preprint: 1907.12267, (2019).
- [63] ———, *Beta Jacobi ensembles and associated Jacobi polynomials*, arXiv preprint:2005.01100, (2020).
- [64] K. D. TRINH, *Global spectrum fluctuations for Gaussian beta ensembles: a Martingale approach*, J. Theoret. Probab., 32 (2019), pp. 1420–1437.
- [65] A. C. ZAAANEN, *Riesz spaces. II*, vol. 30 of North-Holland Mathematical Library, North-Holland Publishing Co., Amsterdam, 1983.
- [66] V. ZKHAROV AND A. SHABAT, *Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media*, Soviet Physics JETP, 34 (1972), pp. 62–69.