

Low-regularity Seiberg-Witten moduli spaces on manifolds with boundary

Piotr Suwara *

December 7, 2021

Abstract

For a compact spin^c manifold X with boundary $b_1(\partial X) = 0$, we consider moduli spaces of solutions to the Seiberg-Witten equations in a generalized double Coulomb slice in $W^{1,2}$ Sobolev regularity. We prove they are Hilbert manifolds, prove denseness and "semi-infinite-dimensionality" properties of the restriction to ∂X , and establish a gluing theorem.

To achieve these, we prove a general regularity theorem and a strong unique continuation principle for Dirac operators, and smoothness of a restriction map to configurations of higher regularity on the interior, all of which are of independent interest.

Contents

1	Introduction	2
2	Analytical preparation	6
2.1	A regularity theorem for Dirac operators	6
2.2	A strong UCP for Dirac operators	9
3	Seiberg-Witten moduli spaces in split Coulomb slice	10
3.1	Coulomb slice on 3-manifolds	10
3.2	Split Coulomb slice on 4-manifolds	12
3.3	Restriction to the boundary and twisting	21
3.4	Seiberg-Witten moduli in split Coulomb slice	25

*Institute of Mathematics, Polish Academy of Sciences, ul. Śniadeckich 8, 00-656 Warszawa, email: psuwara@impan.pl

4	Properties of moduli spaces	26
4.1	Extended linearized SW operator	26
4.2	Regularity and semi-infinite-dimensionality of moduli spaces	30
5	Gluing along a boundary component	34
5.1	Smoothness of restrictions	34
5.2	Proof of the gluing theorem	37
A	Appendix	42
	References	44

1 Introduction

This article is motivated by an effort to provide a new framework for constructing so-called Floer theories, which are used to construct invariants of low-dimensional manifolds, knots, as well as Lagrangians in symplectic manifolds. Originally, Floer implanted ideas from finite-dimensional Morse theory (cf. [Sch93, Wit82]) to the study of functions on infinite-dimensional spaces (often called *functionals* on *configuration spaces*), leading to the resolution of the Arnold conjecture in symplectic topology [Flo87, Flo88]. Inspired by Donaldson’s proof of the diagonalization theorem for 4-manifolds [Don83], Floer used these ideas to construct homological invariants of 3-manifolds, aiming to establish gluing formulas for Donaldson’s invariants. An analogous construction for knots in 3-manifolds has been carried out by Kronheimer and Mrowka and plays the key role in the proof that Khovanov homology detects the unknot [KM11]. An analogue for the Seiberg-Witten equations (in place of the anti-self-duality (ASD) equations of Donaldson’s theory) is the central ingredient of Manolescu’s disproof of the triangulation conjecture in dimensions ≥ 5 [Man13].

While the idea of using Morse theory in infinite-dimensional settings dates back to Atiyah and Bott [AB83], Floer’s novelty was dealing with functionals having critical points of infinite index. The key observation was that the unstable manifolds of critical points were of finite index with respect to a certain subbundle of the tangent space, allowing one to define finite relative indices of critical points and eventually leading to a computation of “middle-dimensional” homology groups of the space. As noted by Atiyah [Ati88] (cf. [CJS95]), Floer theory was, from the very beginning, understood as describing the behavior of so-called semi-infinite-dimensional cycles.

As in Morse homology (cf. [Sch93]), one needs to overcome analytical difficulties to even define Floer theories and prove they are well-defined and independent of choices –

one needs to establish regularity and compactness of moduli spaces of trajectories between critical points. Moreover, Morse theory is generally not well-suited for equivariant constructions since one in general cannot guarantee regularity of the moduli spaces without breaking the symmetry coming from a group action. To address these problems and to allow general constructions of equivariant (cf. [DS19, KM07, Lin18, Mil19]) and generalized Floer homologies (cf. [Man14, Lin15, AB21]), Lipyanskiy [Lip08] introduced a framework for using such semi-infinite-dimensional cycles as a tool for defining Floer theories and the author has further developed these methods in [Suw20]. This article contains results required to rigorously define the relative invariants of 4-manifolds with boundary and maps induced by cobordisms in this construction, using the Seiberg-Witten equations. Moreover, the key result is a gluing theorem which is fundamental for establishing functoriality of the cobordism maps.

Results. Consider a 4-manifold X with boundary a nonempty collection of rational homology spheres (i.e., $b_1(\partial X) = 0$) and a spin^c structure \hat{s} on X . We study the moduli space of solutions to the Seiberg-Witten equations for pairs (A, Φ) of a spin^c connection A on X and a section Φ of the spinor bundle S^+ over X associated to the spin^c structure \hat{s} :

$$\begin{cases} \frac{1}{2}F_{A^t}^+ - \rho^{-1}((\Phi\Phi^*)_0) = 0, \\ \not{D}_A^+ \Phi = 0. \end{cases}$$

(Definition 3.4). The split Coulomb slice with respect to a reference connection A_0 is given by the equations

$$\begin{cases} d^*(A - A_0) = 0, \\ d^*(\iota_{\partial X}^*(A - A_0)) = 0, \end{cases}$$

(where $\iota_{\partial X} : \partial X \hookrightarrow X$ denotes the inclusion) together with a condition restricting $A - A_0$ to a subset of codimension $b_0(\partial X) - 1$ (Definition 3.15). This additional condition depends on a choice of a gauge splitting s (Definition 3.11). The moduli space $\mathcal{M}_s^\circ(X, \hat{s})$ (Definition 3.31) is the quotient of the space of L_1^2 -solutions to the Seiberg-Witten equations in this split Coulomb slice by the (discrete) action of the split gauge group (Definition 3.18); this gauge group preserves the split Coulomb slice as well as the set of solutions to the Seiberg-Witten equations. There is also a residual action of S^1 on $\mathcal{M}_s^\circ(X, \hat{s})$ given by multiplication of the spinor component, $\Phi \mapsto z\Phi$, by complex numbers in the unit circle $z \in S^1 \subset \mathbb{C}$.

Firstly, we choose a gauge twisting τ (Definition 3.24) and define the (S^1 -equivariant) twisted restriction map $R_\tau : \mathcal{M}_s^\circ(X, \hat{s}) \rightarrow \mathcal{C}_{cC}(\partial X, \mathfrak{s})$, taking values in the configurations on ∂X in the Coulomb slice (Definition 3.2). The gauge splittings and gauge twistings we introduce generalize the double Coulomb slice used in [Lip08, Kha]

and twistings utilized in [KLS18]. We prove regularity, denseness and “semi-infinite-dimensionality” of the Seiberg-Witten moduli spaces:

Semi-infinite-dimensionality Theorem. *The moduli spaces $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$ are Hilbert manifolds. The differential of the twisted restriction map $R_\tau : \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{cC}(\partial X, \mathfrak{s})$ decomposes into $\Pi^- DR_\tau : T\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow H^-(\partial X, \mathfrak{s})$ which is Fredholm and $\Pi^+ DR_\tau : T\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow H^+(\partial X, \mathfrak{s})$ which is compact.*

Moreover, if $b_0(\partial X) > 1$, then for any connected component $Y_0 \subset \partial X$ the restriction R_{τ, Y_0} to Y_0 has dense differential.

The maps Π^\pm come from a decomposition $\mathcal{C}_{cC}(\partial X, \mathfrak{s}) = H^+(\partial X, \mathfrak{s}) \oplus H^-(\partial X, \mathfrak{s})$ according to the eigenvalues of the operator $(\star d, \not{D}_{B_0})$, also called a polarization (Definition 3.3).

The proof utilizes the Atiyah-Patodi-Singer boundary value problem for an extended linearized Seiberg-Witten operator (Definition 4.1). Then we prove its properties transfer to the restriction to the Coulomb slice.

Our low regularity setting requires us to prove a regularity theorem (Regularity Theorem) for an operator of the form $D = D_0 + K : L_1^2 \rightarrow L^2$, where D_0 is a Dirac operator and K is any compact operator, making it a result of independent interest. We also prove a strong version of the unique continuation principle for D (Strong UCP Theorem).

Secondly, we prove a gluing theorem for a composite cobordism. Assume X splits as $X = X_1 \cup_Y X_2$ along a rational homology sphere Y . If the auxiliary data of gauge splittings and gauge twistings are compatible in a suitable sense (see Proposition 3.29 and Proposition 5.5), then the moduli space $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$ can be recovered from the fiber product of $\mathcal{M}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1)$ and $\mathcal{M}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2)$ over the configuration space of Y , in a way compatible with the twisted restriction maps:

Gluing Theorem. *Assume $s_{\mathbb{Z}}$ and $(s_{1, \mathbb{Z}}, s_{2, \mathbb{Z}})$ are compatible. Then there is an S^1 -equivariant diffeomorphism $F : \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{M}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times_Y \mathcal{M}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2)$ such that R_τ is S^1 -equivariantly homotopic to $(R_{\tau_1} \times_Y R_{\tau_2}) \circ F$.*

The proof uses the following fact of independent interest. We show that the restriction map of solutions in $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$ to a submanifold in the interior of X is smooth as a map into a configuration space of *higher* regularity (Theorem 5.2). While it is easy to prove that its image lies in the space of smooth configurations (Lemma 5.1), the proof of *smoothness* of this map is non-trivial.

Applications. The [Semi-infinite-dimensionality Theorem](#) together with compactness of moduli spaces proved in [KM07] (with minor modifications to account for the double Coulomb slice instead of the Coulomb-Neumann slice used in [KM07]) show that the maps $R_\tau : \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{cC}(Y, \mathfrak{s})$ are *semi-infinite-dimensional cycles* in $\mathcal{C}_{cC}(\partial X, \mathfrak{s})$

as defined in [Lip08, Suw20], establishing the existence of relative Seiberg-Witten invariants of X . If the boundary ∂X is connected, these do not depend on the choice of an integral splitting (Lemma 3.30). This also implies that cobordisms $\partial W = -Y_1 \cup Y_2$ induce *correspondences* (defined in [Lip08, Suw20]) between the configuration spaces over Y_1 and Y_2 .

Our methods apply to perturbed equations as well, which we did not include for the sake of simplicity. Varying the metrics and perturbations gives cobordisms between the relevant moduli spaces on X , and changing the reference connections on X gives isomorphic moduli spaces with homotopic restriction maps. This means that the relative invariant of X , up to a suitable cobordism relation, does not depend on the choices of perturbations and metric.

Crucially, the [Gluing Theorem](#) says that the correspondence induced by a composite cobordism is (homotopic to) the composition of the respective correspondences, proving the theory comes with a TQFT-like structure. We hope to prove that this theory recovers a non-equivariant version of monopole Floer homology \widehat{HM}_* and describe a construction supposed to recover all of the flavors of monopole Floer homology (for rational homology spheres) in an upcoming work.

Applications of Seiberg-Witten-Floer theory suggest including the case $b_1(\partial X) > 0$ or allowing X to have cylindrical or conical ends (like in [KM97]) with certain asymptotic conditions on the configurations. The methods presented here should suffice to prove regularity and semi-infinite-dimensionality of the corresponding moduli spaces. However, to establish the semi-infinite-dimensional theory in full one needs to carefully select a slice for the gauge action and deal with compactness issues which we hope to address in further work.

Finally, the methods presented here should be applicable (with appropriate adjustments) for defining semi-infinite-dimensional Floer theories in other contexts, e.g., in the Yang-Mills-Floer theory.

Organization. In [section 2](#) we prove the regularity theorem for a Dirac operator with compact perturbation, the [Regularity Theorem](#), and a strong unique continuation principle for a Dirac operator with a low-regularity potential, the [Strong UCP Theorem](#).

In [section 3](#) we introduce the basic notions of Seiberg-Witten theory on 3- and 4-manifolds. For a chosen gauge splitting the split Coulomb slice is defined, and for a gauge twisting a twisted restriction map to the boundary is introduced. The choices of splittings and twistings are shown to be equivalent to a choice of an integral splitting, one which does not require any twisting.

[Section 4](#) provides proofs of some of the key properties of the moduli spaces: regularity, semi-infinite-dimensionality and denseness of the restriction map ([Semi-infinite-dimensionality Theorem](#)).

Finally, [section 5](#) contains the proof of the [Gluing Theorem](#), showing that moduli on a composite cobordism are a fiber product of the moduli on its components.

Notations. In this article we use the notation L_k^p for Sobolev spaces of regularity k , in accordance with the literature in gauge theory and Floer theory in low dimensional topology. These are often denoted by $W^{k,p}$ or $H^{k,p}$ or, when $p = 2$, simply by H^k .

All manifolds are assumed to be smooth, submanifolds to be smoothly embedded, and manifolds with boundary to have smooth boundary.

Acknowledgments. I would like to express gratitude to my graduate advisor Tom Mrowka for his encouragement, support and sharing many insights on geometry, topology and analysis. I would like to thank Maciej Borodzik and Michał Miśkiewicz for helpful comments on a draft of this paper, and an anonymous referee for pointing out an inaccuracy in the statement of [Regularity Theorem](#).

Most results of this paper have been part of the author's Ph.D. thesis [\[Suw20\]](#), although the proofs have been revised and some results have been generalized. In particular, [\[Suw20\]](#) does not consider 4-manifolds with more than two boundary components, or the general split Coulomb slice on them.

2 Analytical preparation

We prove two results which are fundamental for the surjectivity proofs in [section 4](#).

The first one, the [Regularity Theorem](#), is a regularity theorem for operators of the form $D = D_0 + K : L_1^2 \rightarrow L^2$, where D_0 is a first-order elliptic operator and K is a compact operator. In our applications K is a multiplication by an L_1^2 -configuration on a 4-manifold and thus it does not factor as a map $L_1^2 \rightarrow L_1^2$. If it did, elliptic regularity would immediately imply the regularity result. The novelty here is the general form of the perturbation K which is only assumed to be compact as a map $L_1^2 \rightarrow L^2$.

The second result, the [Strong UCP Theorem](#), is a strong unique continuation principle in a similar low regularity setting. This can be understood as a strengthening of the unique continuation results of [\[KM07, Section 7\]](#).

2.1 A regularity theorem for Dirac operators

Let X be a smooth Riemannian manifold with asymptotically cylindrical ends (without boundary). Let D_0 be an elliptic operator $D_0 : C^\infty(X; E) \rightarrow C^\infty(X; F)$ of order 1 which is asymptotically cylindrical on the ends of X . Let $K : L_1^2(X; E) \rightarrow L^2(X; F)$ be a compact operator which has a compact formal adjoint $K^* : L_1^2(X; E) \rightarrow L^2(X; F)$

and which extends to a continuous operator $K : L^2(X; E) \rightarrow L^2_{-l}(X; F)$ for some l . Consider the operator

$$D = D_0 + K : L^2_1(X; E) \rightarrow L^2(X; F).$$

The aim of this subsection is to prove the following regularity theorem:

Regularity Theorem. *Assume that $v \in L^2(X; E)$ satisfies $Dv = h$ for some $h \in L^2(X; F)$. Then $v \in L^2_1(X; E)$.*

Moreover, if $v, h \in L^2_{loc}(X; E)$ instead, then $v \in L^2_{1,loc}(X; E)$.

In the course of the proof we will make use of the following simple lemma (cf. Lipyanskiy [Lip08, Lemma 44]):

Lemma 2.1. *Suppose the sequence of Hilbert space operators $\{A_i : V \rightarrow W\}$ is uniformly bounded and weakly convergent to A in the sense that for any $v \in V$ we have $A_i(v) \rightarrow A(v)$.*

If $K : W \rightarrow U$ is compact, then $A_i \circ K \rightarrow A \circ K$.

We also need the following version of the Gårding inequality, proven in [Shu92, Appendix 1, Lemma 1.4]:

Proposition 2.2 (cylindrical Gårding inequality). *For every $s, t \in \mathbb{R}$ there exists $C > 0$ such that for any $u \in C_0^\infty(X; E)$*

$$\|u\|_{L^2_{s+1}} \leq C(\|D_0 u\|_{L^2_s} + \|u\|_{L^2_t}).$$

With these in hand, we are ready to prove the [Regularity Theorem](#).

Proof (Proof of Regularity Theorem). By considering

$$\begin{pmatrix} 0 & D_0^* \\ D_0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & K^* \\ K & 0 \end{pmatrix} : L^2_1(X; E \oplus F) \rightarrow L^2(X; E \oplus F)$$

where D_0^* is the formal adjoint of D_0 we may assume, without loss of generality, that $E = F$ and D_0 is formally self-adjoint. Since X has bounded geometry and D_0 is uniformly elliptic, Proposition 4.1 in [Shu92, Section 1] implies that the minimal and maximal operators D_0^{min} and D_0^{max} of $D_0 : L^2(X; E) \rightarrow L^2(X; E)$ coincide, and their domains are equal to $L^2_1(X; E)$. Thus D_0 is essentially self-adjoint.

Consider the equation

$$(D_0 + ic)\psi + K\psi = h + icv \tag{1}$$

for large $c \in \mathbb{R}$. Certainly, $\psi = v \in L^2(X; E)$ solves the equation. We will prove that for large c equation (1) has a unique solution ψ in $L^2_1(X; E)$, which is also unique in $L^2(X; E)$, so that $v = \psi \in L^2_1(X; E)$, as wished.

Firstly, we want to prove $D_0 + ic : L_1^2(X; E) \rightarrow L^2(X; E)$ is invertible. Since the spectrum of D_0 is real, $(D_0 + ic)^{-1} : L^2(X; E) \rightarrow L^2(X; E)$ exists and is bounded by $\frac{1}{|c|}$ (cf. [Lan93, Chapter XIX, Theorem 2.4]). Therefore by Proposition 2.2 we have the first inequality:

$$\begin{aligned} \|(D_0 + ic)^{-1}u\|_{L_1^2} &\leq \|D_0(D_0 + ic)^{-1}u\|_{L^2} + \|(D_0 + ic)^{-1}u\|_{L^2} \\ &\leq \|u - ic(D_0 + ic)^{-1}u\|_{L^2} + (1/|c|)\|u\|_{L^2} \\ &\leq (2 + 1/|c|)\|u\|_{L^2} \end{aligned}$$

and therefore $(D_0 + ic)^{-1}$ is a bounded operator $L^2(X; E) \rightarrow L_1^2(X; E)$.

Secondly, we want to prove weak convergence of $(D_0 + ic)^{-1}$ to 0 as $|c| \rightarrow \infty$, i.e., that for each $v \in L^2(X; E)$ we have $(D_0 + ic)^{-1}(v) \rightarrow 0$ in $L_1^2(X; E)$ as $|c| \rightarrow \infty$. The spectral theorem for unbounded self-adjoint operators [Lan93, Chapter XIX, Theorem 2.7] provides us with an orthogonal decomposition $L^2(X; E) = \hat{\bigoplus}_n H_n$ such that the restriction $D_n = D_0|_{H_n} : H_n \rightarrow H_n$ is a bounded operator (in L^2 -norm on H_n) and $D_0 = \hat{\bigoplus}_n D_n$. In particular, $H_n \subset \text{Dom}(D_0) = L_1^2(X; E)$. For each $v_n \in H_n$ we thus have, by Proposition 2.2,

$$\|v_n\|_{L_1^2} \leq C(\|D_0 v_n\|_{L^2} + \|v_n\|_{L^2}) \leq C(C_n + 1)\|v_n\|_{L^2} = C'_n\|v_n\|_{L^2},$$

where $C_n = \|D_n\|_{L^2}$. Therefore

$$\|(D_0 + ic)^{-1}v_n\|_{L_1^2} \leq C'_n\|(D_0 + ic)^{-1}v_n\|_{L^2} \leq \frac{C'_n}{|c|}\|v_n\|_{L^2} \xrightarrow{|c| \rightarrow \infty} 0$$

which proves that for any $v \in L^2(X; E)$, $(D_0 + ic)^{-1}v \xrightarrow{|c| \rightarrow \infty} 0$ in $L_1^2(X; E)$, i.e., $(D_0 + ic)^{-1}$ converges weakly to 0, as wished.

We proceed to proving existence and uniqueness of solutions to (1) in $L_1^2(X; E)$ for large $|c|$. Using Lemma 2.1 we conclude that $T = (D_0 + ic)^{-1}K : L_1^2(X; E) \rightarrow L_1^2(X; E)$ converges strongly to 0. Thus, for large $|c|$, the operator $\text{Id} + (D_0 + ic)^{-1}K : L_1^2(X; E) \rightarrow L_1^2(X; E)$ is invertible. Composing with $(D_0 + ic)$ we conclude that $D_0 + ic + K = D + ic : L_1^2(X; E) \rightarrow L^2(X; E)$ is invertible for large $|c|$. Existence and uniqueness of $\psi \in L_1^2(X; E)$ solving (1) follows.

To conclude the first part of the proof we need to establish uniqueness of solutions to (1) in $L^2(X; E)$. We have that $(D_0 + ic + K)(\psi - v) = 0$. This implies that $\psi - v$ is perpendicular in $L^2(X; E)$ to the image of $D_0^* - ic + K^* = D_0 - ic + K : L_1^2(X; E) \rightarrow L^2(X; E)$. However, we already established that for large $|c|$ the latter operator is invertible and it follows that $\psi - v = 0$; thus $v = \psi \in L_1^2(X; E)$.

Finally, it remains to consider the more general case $v, h \in L^2_{loc}$. For any compact $A \subset X$ we can take a compactly supported bump function $\rho : X \rightarrow [0, 1]$ with $\rho|_A = 1$ and define $v' = \rho v \in L^2(X; E)$. Then $Dv' = h'$ for some $h' \in L^2(X; E)$ and therefore $v' \in L^2_1(X; E)$ as proven above, so $v|_A \in L^2_1(A; E|_A)$. Repeating the argument for every compact $A \subset X$ shows that $v \in L^2_{1,loc}(X; E)$. \square

2.2 A strong UCP for Dirac operators

Let M be a connected Riemannian manifold and S a real (resp. complex) Dirac bundle over it (e.g., the real (resp. complex) spinor bundle associated to a spin^c -structure on M) with connection A . Denote by $\not{D}_A : \Gamma(S) \rightarrow \Gamma(S)$ the corresponding Dirac operator. Let $V \in L^n(M; \mathbb{R})$ be a potential. Here we prove the following unique continuation theorem for spinors and potentials of low regularity.

Strong UCP Theorem. *The differential inequality*

$$|\not{D}_A \Phi| \leq V|\Phi|$$

has unique continuation property in $L^{\frac{2n}{n+2}}_{1,loc}(M; S)$.

The version of this theorem for $d + d^*$ instead of \not{D}_A has been proven in [Wol92, Theorem 2]:

Theorem 2.3 (Wolff, [Wol92, Theorem 2]). *Suppose M is an n -dimensional manifold, $n \geq 3$, $p = \frac{2n}{n+2}$, and $\omega \in L^p_{1,loc}(\Omega^*(M))$ such that $|d\omega| + |d^*\omega| \leq V|\omega|$ with $V \in L^n_{loc}(M)$. Then if ω vanishes on an open set, it vanishes identically.*

It thus suffices to reduce our problem to Wolff's result, following the idea of [Man94]. Since the proof in [Man94] does not explain the reduction rigorously, we describe the procedure below.

Proof. The problem is local, thus we need only to consider the case of M being \mathbb{R}^n with some metric g . If A_0 is the flat connection on S , $\not{D}_A - \not{D}_{A_0}$ is a smooth operator of order 0, so (again using locality) we can assume that A is flat. Moreover, by contractibility of \mathbb{R}^n , we can decompose S into irreducible components, and irreducible components must be isomorphic to the real (resp. complex) spinor bundle \tilde{S} (resp. $\tilde{S}_{\mathbb{C}}$) associated to the unique spin structure. Thus we have reduced the problem to the case where $S = \tilde{S} \otimes \mathbb{R}^k$ (resp. $S = \tilde{S} \otimes \mathbb{C}^k$) for some k .

The real (resp. complex) spinor bundle \tilde{S} (resp. $\tilde{S}_{\mathbb{C}}$) embeds into $\text{Cl}_n(\mathbb{R}^n)$ (resp. $\text{Cl}_n(\mathbb{R}^n)$). Furthermore, [LM89, Theorem 5.12] implies that the Dirac operator on the Clifford bundle $\text{Cl}_n(\mathbb{R}^n)$ (resp. $\text{Cl}_n(\mathbb{R}^n) = \text{Cl}(\mathbb{R}^n) \otimes \mathbb{C}$) is equivalent to $d + d^*$ on $\Lambda^*(\mathbb{R}^n)$ (resp. $\Lambda^*(\mathbb{R}^n) \otimes \mathbb{C}$) via the canonical isomorphism $\text{Cl}(\mathbb{R}^n) \simeq \Lambda^*(\mathbb{R}^n)$. Thus, \not{D}_{A_0} is equivalent to $(d + d^*) \otimes 1_{\mathbb{R}^k} \Lambda^*(\mathbb{R}^n) \otimes \mathbb{R}^k \rightarrow \Lambda^*(\mathbb{R}^n) \otimes \mathbb{R}^k$ (resp. $(d + d^*) \otimes 1_{\mathbb{C}^k} \Lambda^*(\mathbb{R}^n) \otimes \mathbb{C}$).

$\mathbb{C}^k \rightarrow \Lambda^*(\mathbb{R}^n) \otimes \mathbb{C}^k$). The proof of [Theorem 2.3](#) goes through for differential forms with coefficients in \mathbb{R}^k (resp. \mathbb{C}^k), establishing the unique continuation property for $(d + d^*) \otimes \mathbb{R}^k$ (resp. $(d + d^*) \otimes 1_{\mathbb{C}^k}$). This finishes the proof. \square

In the article we will use the following special case:

Corollary 2.4 (*UCP for Dirac operators in 4d*). *In the setting of the [Strong UCP Theorem](#), assume $n = 4$ and $V \in L^2_1(M; \text{Aut}(S))$. Then any solution $\Phi \in L^2_{1,loc}(M; S)$ to*

$$\not{D}_A \Phi + V\Phi = 0$$

which is zero on some open set is identically zero.

3 Seiberg-Witten moduli spaces in split Coulomb slice

In this section we introduce the moduli spaces of the Seiberg-Witten equations on a Riemannian 4-manifold with boundary a collection of rational homology spheres, together with the restriction maps to the boundary. For 3-manifolds, we introduce the Coulomb slice and its polarization, a decomposition of the tangent space into a sum of two infinite-dimensional subspaces. For 4-manifolds, we introduce the double Coulomb slice and what we call the split Coulomb slice together with the split gauge group. Since the restriction maps are generally not invariant with respect to the split gauge group, we need to introduce appropriate twisted restriction maps as well.

The split gauge fixing is a key novel element that generalizes the gauge slice introduced by Khandhawit [[Kha](#)] and Lipyanskiy [[Lip08](#)]. It simplifies the proof of the [Gluing Theorem](#) letting us to reduce it to the case of untwisted restriction maps.

3.1 Coulomb slice on 3-manifolds

We begin by introducing polarizations on the Seiberg-Witten configuration space in Coulomb gauge on an oriented rational homology sphere Y and collections of such.

Let g be a Riemannian metric and \mathfrak{s} be a spin^c structure on Y . Denote by S_Y the associated spinor bundle and choose a smooth spin^c connection B_0 . The Seiberg-Witten configuration space on Y is the space

$$\mathcal{C}(Y, \mathfrak{s}) = (B_0, 0) + L^2_{1/2}(i\Omega^1(Y) \oplus \Gamma(S_Y))$$

consisting of pairs (B, Ψ) of a spin^c connection and a spinor on Y .

In Seiberg-Witten theory one investigates the Chern-Simons-Dirac functional \mathcal{L}

$$\mathcal{L}(B, \Psi) = -\frac{1}{8} \int_Y (B^t - B_0^t) \wedge (F_{B^t} + F_{B_0^t}) + \frac{1}{2} \int_Y \langle \not{D}_B \Psi, \Psi \rangle.$$

on the configuration space. The gauge group $\mathcal{G}(Y) = L^2_{3/2}(Y; S^1)$ acts on $\mathcal{C}(Y, \mathfrak{s})$ via $u(A, \Phi) = (A - u^{-1}du, u\Phi)$ where $u \in \mathcal{G}(Y)$, leaving \mathcal{L} invariant. If one used spaces of higher regularity, one could work with the quotient of the configuration space by the action of the gauge group. However, in the low regularity setting the action of $\mathcal{G}(Y)$ on the spinors is not continuous. Because of that (and in applications concerning the Seiberg-Witten stable homotopy type, cf. [KLS18]), it is preferable to take the Coulomb slice as the model for the quotient by the identity component of the gauge group. Indeed, for Y a rational homology sphere the Hodge decomposition gives the L^2 -orthogonal decomposition

$$\Omega^1(Y) = \Omega^1_C(Y) \oplus \Omega^1_{cC}(Y)$$

where $\Omega^1_{cC}(Y) = \{b \in \Omega^1(Y) | d^*b = 0\} = d^*(\Omega^2(Y))$ is the space of (smooth) co-closed forms and $\Omega^1_C(Y) = \{b \in \Omega^1(Y) | db = 0\} = d(\Omega^0(Y))$ is the space of (smooth) closed forms. Denote by Π_d the projection $\Pi_d : L^2_{1/2}(i\Omega^1(Y)) \rightarrow L^2_{1/2}(i\Omega^1_C(Y)) = d\left(L^2_{3/2}(i\Omega^0(Y))\right)$ along $L^2_{1/2}(i\Omega^1_{cC}(Y))$.

Lemma 3.1 (gauge fixing in 3d). *On Y , there is a continuous choice of based and contractible gauge transformations putting forms in the Coulomb slice, i.e., a homomorphism*

$$\begin{aligned} L^2_{1/2}(i\Omega^1(Y)) &\rightarrow \mathcal{G}^{e,\circ}(Y) \\ a &\mapsto u_a \end{aligned}$$

such that $a - u_a^{-1}du_a = (1 - \Pi_d)a \in L^2_{1/2}(i\Omega^1_{cC}(Y))$, where $\mathcal{G}^{e,\circ}(Y) = \left\{ e^f \mid f \in L^2_{3/2}(i\Omega^0(Y)), \int_Y f = 0 \right\} \subset \mathcal{G}(Y)$. For each a , there is exactly one such u_a .

Proof. Denote $\Omega^0_0(Y) = \{f \in \Omega^0(Y) | \int_Y f = 0\}$. The exterior derivative $d : L^2_{3/2}(\Omega^0_0(Y)) \rightarrow L^2_{1/2}(i\Omega^1_C(Y))$ has inverse G_d . Denote by Π_d the orthogonal projection $\Omega^1(Y) \rightarrow d(\Omega^0(Y))$. We take $u_a = e^{G_d \Pi_d a}$ which has the desired properties. Uniqueness follows from the fact that $df = 0$ and $\int_Y f = 0$ imply $f = 0$. \square

Moreover, for Y a rational homology sphere we have $\mathcal{G}(Y) = \mathcal{G}^e(Y) = \left\{ e^f \mid f \in L^2_{3/2}(Y; i\mathbb{R}) \right\}$. Indeed, taking $\tilde{u} = u_{-u^{-1}du}u$ gives us \tilde{u} with $d(\tilde{u}^{-1}d\tilde{u}) = 0$, which implies $d\tilde{u} = 0$ and thus \tilde{u} is constant. It follows that there are bijections

$$\begin{aligned} \mathcal{C}_{cC}(Y, \mathfrak{s}) &\leftrightarrow (\mathcal{C}(Y, \mathfrak{s})) / \mathcal{G}^{e,\circ}(Y), \\ (\mathcal{C}_{cC}(Y, \mathfrak{s})) / S^1 &\leftrightarrow (\mathcal{C}(Y, \mathfrak{s})) / \mathcal{G}(Y), \end{aligned}$$

justifying the restriction to the Coulomb slice:

Definition 3.2 (Coulomb slice). The Coulomb slice on Y with respect to the reference connection B_0 is the space of configurations

$$\mathcal{C}_{cC}(Y, \mathfrak{s}) = (B_0, 0) + L^2_{1/2}(i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y)) \subset \mathcal{C}(Y, \mathfrak{s}).$$

The following subspaces are crucial to the analysis of Atiyah-Patodi-Singer boundary value problem for the Seiberg-Witten equations on 4-manifolds with boundary Y .

Definition 3.3 (polarization on the Coulomb slice). We define $H^+(Y, \mathfrak{s})$ (resp. $H^-(Y, \mathfrak{s})$) $\subset L^2_{1/2}(i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y))$ to be the closure of the span of positive (resp. nonpositive) eigenvalues of

$$(\star d) \oplus \not{D}_{B_0} : L^2_{1/2}(i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y)) \rightarrow L^2_{1/2}(i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y)).$$

We denote by $\Pi^\pm : L^2_{1/2}(i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y)) \rightarrow H^\pm(Y, \mathfrak{s})$ the projection onto $H^\pm(Y, \mathfrak{s})$ along $H^\mp(Y, \mathfrak{s})$.

One of our goals is to prove that the moduli spaces of solutions to the Seiberg-Witten equations on X are, in a precise sense, comparable to the negative subspace $H^-(Y, \mathfrak{s})$ via the restriction map (cf. [Semi-infinite-dimensionality Theorem](#)).

Finally, if $Y = \bigsqcup_i Y_i$ is a disjoint sum of oriented rational homology spheres Y_i then we define

$$\begin{aligned} \mathcal{C}(Y, \mathfrak{s}) &= \prod_i \mathcal{C}(Y_i, \mathfrak{s}_i), & \mathcal{C}_{cC}(Y, \mathfrak{s}) &= \prod_i \mathcal{C}_{cC}(Y_i, \mathfrak{s}_i), \\ \mathcal{L}(\prod_i b_i, \prod_i \Psi_i) &= \sum_i \mathcal{L}(b_i, \Psi_i) & H^\pm(Y, \mathfrak{s}) &= \prod_i H^\pm(Y_i, \mathfrak{s}_i). \end{aligned}$$

Moreover, note that there are a natural identifications between configuration spaces for Y and oppositely oriented $-Y$. For a spin^c structure \mathfrak{s} with its spinor bundle S_Y there is the conjugate spin^c structure $\bar{\mathfrak{s}}$ determined by the conjugate bundle \bar{S}_Y , and the anti-linear isomorphism $S_Y \simeq \bar{S}_Y$ induces natural affine isomorphisms

$$\mathcal{C}(Y, \mathfrak{s}) \simeq \mathcal{C}(-Y, \bar{\mathfrak{s}}) \quad \mathcal{C}_{cC}(Y, \mathfrak{s}) \simeq \mathcal{C}_{cC}(-Y, \bar{\mathfrak{s}}) \quad H^\pm(Y, \mathfrak{s}) \simeq H^\mp(-Y, \bar{\mathfrak{s}}).$$

3.2 Split Coulomb slice on 4-manifolds

We turn our attention to the Seiberg-Witten equations and gauge fixings for configurations on a connected oriented 4-manifold X with nonempty boundary $\partial X \neq \emptyset$ satisfying $b_1(\partial X) = 0$, oriented using the outward normal. As explained by Khandawit [\[Kha\]](#), the most convenient slice for these is a kind of a *double* Coulomb slice (which was already used by Lipyanskiy [\[Lip08\]](#)), which imposes both coclosedness of the connection 1-form on both X and ∂X , as well as an auxiliary condition near ∂X . We drop

this auxiliary condition from the definition of the double Coulomb slice and instead introduce the split Coulomb slice which generalizes the constructions of Khandhawit and Lipyanskiy. This allows one to choose a gauge fixing which do not require twisting or ones that are more geometric in nature, depending on one's needs. Indeed, twisting is necessary in Khandhawit's and Lipyanskiy's gauge fixing, in which the restriction map may not commute with the residual gauge group action.

We begin by introducing the Seiberg-Witten equations on 4-manifolds.

Definition 3.4 (Seiberg-Witten equations). The Seiberg-Witten map is defined by

$$\begin{aligned} \text{SW} : \mathcal{C}(X, \hat{\mathfrak{s}}) &\rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)), \\ \text{SW}(A, \Phi) &= \left(\frac{1}{2}F_{A^t}^+ - \rho^{-1}((\Phi\Phi^*)_0), \mathcal{D}_A^+ \Phi \right), \end{aligned}$$

where A^t denotes the connection induced by A on $\det(S_X^+)$ and $F_{A^t}^+$ denotes the self-dual part of its curvature, according to the splitting $\Lambda^2(X) = \Lambda^+(X) \oplus \Lambda^-(X)$ by the eigenspaces of the Hodge star \star .

The Seiberg-Witten equations are the equations given by $\text{SW}(A, \Phi) = 0$, that is,

$$\begin{cases} \frac{1}{2}F_{A^t}^+ - \rho^{-1}((\Phi\Phi^*)_0) = 0, \\ \mathcal{D}_A^+ \Phi = 0. \end{cases}$$

Note that continuity and smoothness of the map SW follow from [Theorem A.1](#) and the fact that continuous multilinear maps on Banach spaces are smooth.

These equations are equivariant with respect to the action of the gauge group $\mathcal{G}(X) = L_2^2(X; S^1)$. As is easily seen, the solution set is invariant under this action.

Lemma 3.5 (gauge group action on a 4-manifold). *The gauge group $\mathcal{G}(X)$ acts on $\mathcal{C}(X, \hat{\mathfrak{s}})$ via $u(A, \Phi) = (A - u^{-1}du, u\Phi)$ where $u \in \mathcal{G}(X)$. Moreover, $\text{SW}(A, \Phi) = 0$ if and only if $\text{SW}(u(A, \Phi)) = 0$.*

Note that this action is not continuous since the multiplication $L_2^2(X) \times L_1^2(X) \rightarrow L_1^2(X)$ is not continuous. It is well-defined since $\mathcal{G}(X) \subset L^\infty(X) \cap L_2^2(X)$ and the multiplication $(L_2^2(X) \cap L^\infty(X)) \times L_1^2(X) \rightarrow L_1^2(X)$ is continuous.

In order to prove that the moduli spaces of solutions are manifolds we need to investigate the differential of SW.

$$\begin{aligned} D_{(A, \Phi)} \text{SW} : L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)) &\rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)), \\ D_{(A, \Phi)} \text{SW}(a, \phi) &= (d^+a, \mathcal{D}_{A_0}^+ \phi) + (-\rho^{-1}(\phi\Phi^* + \Phi\phi^*)_0, \rho(a)\Phi + \rho(A^\delta)\phi). \end{aligned}$$

Similarly, at (e, A, Φ) the differential of the gauge group action is:

$$\begin{aligned} T\mathcal{G}(X) &= L_2^2(X; \mathbb{R}) \rightarrow L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)), \\ f &\mapsto (-df, f\Phi). \end{aligned}$$

As in dimension 3, we can fix gauge using the Coulomb condition, i.e., require that the 1-form is coclosed. Adding the same condition on the boundary ∂X ensures that the restriction to the boundary lies in the previously defined Coulomb slice (cf. [Definition 3.2](#)):

Definition 3.6 (double Coulomb slice). We define the double Coulomb slice:

$$\Omega_{CC}^1(X) = \{a \in \Omega^1(X) \mid d^*a = 0, d^*(\iota_{\partial X}^*a) = 0\}.$$

The gauge group preserving it is called the harmonic gauge group:

$$\mathcal{G}^h(X) = \{u : X \rightarrow S^1 \mid u^{-1}du \in L_1^2(i\Omega_{CC}^1(X))\}.$$

While the action of the full gauge group $\mathcal{G}(X)$ is not continuous, the action of $\mathcal{G}^h(X)$ is, which will be proven in [Lemma 3.9](#).

To define the split Coulomb slice we need to first understand the harmonic gauge group and its relation to harmonic functions and forms on X . Notice that we have a well-defined homomorphism

$$\begin{aligned} \delta : \mathcal{G}(X) &\longrightarrow L_1^2(i\Omega^1(X)) \\ u &\longmapsto \delta(u) = u^{-1}du \end{aligned} \tag{2}$$

which, restricted to harmonic gauge transformations, induces

$$\delta : \mathcal{G}^h(X) \rightarrow i\mathcal{H}_D^1(X)$$

where

$$\mathcal{H}_D^1(X) = \{a \in \Omega^1(X) \mid da = 0, d^*a = 0, \iota_{\partial X}^*a = 0\}$$

is the space of harmonic 1-forms with Dirichlet boundary conditions. Note that δ (both on $\mathcal{G}(X)$ and on $\mathcal{G}^h(X)$) is an inclusion modulo S^1 , i.e., $\ker \delta = S^1$, the group of constant gauge transformations.

On the other hand, the exponential map

$$\begin{aligned} \exp : L_2^2(i\Omega^0(X)) &\longrightarrow \mathcal{G}(X) \\ f &\longmapsto e^f \end{aligned}$$

restricted to the space of doubly harmonic functions

$$\mathcal{H}(X) = \{f \in \Omega^0(X) \mid \Delta f = 0, \Delta(f|_{\partial X}) = 0\}$$

yields a homomorphism

$$\exp : i\mathcal{H}(X) \rightarrow \mathcal{G}^h(X)$$

since the conditions $\Delta f = 0$ and $\Delta(f|_{\partial X}) = 0$ for an imaginary-valued function f are equivalent to $df \in L_1^2(i\Omega_{CC}^1(X))$. Importantly, the composition $\delta \circ \exp : L_2^2(i\Omega^0(X)) \rightarrow L_1^2(i\Omega^1(X))$ is exactly the differential $f \mapsto df$.

Denote the image of this exponential map by $\mathcal{G}^{h,e}(X) = \exp(i\mathcal{H}(X))$. As the next proposition explains, $\mathcal{G}^{h,e}(X)$ is the identity component of $\mathcal{G}^h(X)$. Thus, our goal will be to find a gauge fixing that dispenses with the action of this identity component, saving only the action by S^1 , the constant elements.

Proposition 3.7 (sequence of harmonic gauge groups). *The following sequence is exact:*

$$0 \longrightarrow \mathcal{G}^{h,e}(X) \longrightarrow \mathcal{G}^h(X) \longrightarrow \pi_0\mathcal{G}^h(X) \longrightarrow 0. \quad (3)$$

The identity component $\mathcal{G}^{h,e}(X)$ is isomorphic to $S^1 \times \mathbb{R}^{b_0(\partial X)-1}$ and the group of components $\pi_0\mathcal{G}^h(X)$ is naturally isomorphic to $H^1(X; \mathbb{Z})$.

Remark 3.8. Recall that $H^1(X; \mathbb{Z}) \simeq \text{Hom}(\pi_1(X), \mathbb{Z})$ has no torsion.

Proof. Crucial to the understanding of the gauge group is the homomorphism (2). Hodge theory provides an identification $\mathcal{H}_D^1(X) \simeq H^1(X, \partial X; \mathbb{R})$. Thus, we will consider δ as a map $\delta : \mathcal{G}^h(X) \rightarrow H^1(X, \partial X; i\mathbb{R})$ with kernel S^1 . This will be used to establish a map of horizontal short exact sequences

$$\begin{array}{ccccccc} & & S^1 & \xlongequal{\quad} & S^1 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{G}^{h,e}(X) & \hookrightarrow & \mathcal{G}^h(X) & \longrightarrow & \pi_0\mathcal{G}^h(X) \rightarrow 0 \\ & & \downarrow & & \downarrow \frac{\delta}{2\pi i} & & \downarrow \frac{[\delta]}{2\pi i} \\ 0 & \rightarrow & H^0(\partial X; \mathbb{R})/H^0(X; \mathbb{R}) & \rightarrow & H^1(X, \partial X; \mathbb{R}) & \longrightarrow & H^1(X; \mathbb{R}) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & H^1(X; \mathbb{R})/H^1(X; \mathbb{Z}) = H^1(X; \mathbb{R})/H^1(X; \mathbb{Z}) & & \downarrow \end{array} \quad (4)$$

where the vertical sequences are also exact, as will be shown in the course of the proof.

Firstly, we prove that $\pi_0\mathcal{G}^h(X) \simeq H^1(X; \mathbb{Z})$. Notice that for any closed loop $\gamma \subset X$ the period of $\delta(u)$, i.e., the integral $\int_\gamma \delta(u)$, is an integer multiple of $2\pi i$; and it is zero whenever γ is contractible. (In fact it is the obstruction to lifting $u|_\gamma : \gamma \rightarrow S^1$ to a map $u|_\gamma : \gamma \rightarrow \mathbb{R}$.) This way any $u \in \mathcal{G}^h(X)$ determines an element $[u] \in H^1(X; 2\pi i\mathbb{Z})$ and we get a homomorphism $\mathcal{G}^h(X) \rightarrow H^1(X; 2\pi i\mathbb{Z})$. Since the periods (having values in

$2\pi i\mathbb{Z}$) do not change under homotopy, this descends to a map $\pi_0\mathcal{G}^h(X) \rightarrow H^1(X; 2\pi i\mathbb{Z})$ and from its construction it follows that it coincides with the composition

$$\mathcal{G}^h(X) \xrightarrow{\delta} H^1(X, \partial X; i\mathbb{R}) \rightarrow H^1(X; i\mathbb{R})$$

which has image in $H^1(X; 2\pi i\mathbb{Z})$. It remains to notice that any element in $x \in H^1(X; 2\pi i\mathbb{Z})$ can be lifted to an element $\tilde{x} = H^1(X, \partial X; i\mathbb{R}) \simeq i\mathcal{H}_D^1(X)$ and then integrated along curves to obtain an element $u_{\tilde{x}} \in \mathcal{G}^h(X)$ mapping to x (see (6)). After dividing 1-forms by $2\pi i$ we obtain a natural isomorphism $\pi_0\mathcal{G}^h(X) \simeq H^1(X; \mathbb{Z})$, as wished.

Moreover, this shows that the kernel K of the map $\mathcal{G}^h(X) \rightarrow H^1(X; 2\pi i\mathbb{Z})$ maps via δ to the kernel of $H^1(X, \partial X; i\mathbb{R}) \rightarrow H^1(X; i\mathbb{R})$. We thus get the map

$$\delta|_K : K \rightarrow \text{im}(H^0(\partial X; i\mathbb{R}))$$

to the image of $H^0(\partial X; i\mathbb{R})$ in $H^1(X, \partial X; i\mathbb{R})$. The map $\delta|_K$ itself has kernel S^1 .

With this in mind, we turn our focus to $\mathcal{G}^{h,e}(X) = \exp(\mathcal{H}(X))$. Since any harmonic function f on X is determined by its restriction $f|_{\partial X}$ to ∂X , and harmonic functions on ∂X are locally constant, we have an isomorphism

$$H^0(\partial X; \mathbb{R}) \simeq \mathcal{H}(X)$$

and we will denote any element g in $H^0(\partial X; \mathbb{R})$ by $f|_{\partial X}$ for the unique $f \in \mathcal{H}(X)$ such that $f|_{\partial X} = g$. We obtain a canonically defined surjection

$$\begin{aligned} \exp : H^0(\partial X; i\mathbb{R}) &\longrightarrow \mathcal{G}^{h,e}(X) \\ f|_{\partial X} &\longmapsto e^f \end{aligned}$$

with kernel generated by restrictions of $H^0(X; 2\pi i\mathbb{Z})$. After dividing by $2\pi i$ we get an isomorphism $\mathcal{G}^{h,e}(X) \simeq H^0(\partial X; \mathbb{R})/H^0(X; \mathbb{Z}) \simeq S^1 \times (H^0(\partial X; \mathbb{R})/H^0(X; \mathbb{R}))$, as wished.

Finally, recall that the composition

$$H^0(\partial X; i\mathbb{R}) \simeq \mathcal{H}(X) \xrightarrow{\exp} \mathcal{G}^{h,e}(X) \xrightarrow{\delta} H^1(X, \partial X; i\mathbb{R})$$

is given by $f|_{\partial X} \mapsto df$ and therefore, by Hodge theory, represents the boundary map in the exact sequence

$$0 \rightarrow H^0(X; \mathbb{R}) \rightarrow H^0(\partial X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R}) \rightarrow H^1(X; \mathbb{R}) = 0$$

Thus $\delta(\mathcal{G}^{h,e}(X)) = \text{im}(H^0(\partial X; i\mathbb{R})) = \delta(K)$. Since $\ker \delta|_{\mathcal{G}^{h,e}(X)} = S^1 = \ker \delta|_K$ and $\mathcal{G}^{h,e}(X) \subseteq K$, we obtain that $\mathcal{G}^{h,e}(X) = K$, i.e., the sequence (3) is exact, as wished. \square

Lemma 3.9 (*\mathcal{G}^h acts continuously*). *The action of $\mathcal{G}^h(X)$ on $\mathcal{C}(X, \hat{s})$ is smooth.*

Proof. By Proposition 3.7 it suffices prove that the action of $\mathcal{G}^{h,e}(X)$ is smooth. Further, by Theorem A.1 it suffices to prove that $\mathcal{G}^{h,e}(X) \subset L^2_3(X; S^1)$ and that this injection is continuous. Since $f \in i\mathcal{H}(X)$ and $\mathcal{H}(X)$ is finite-dimensional, there are constants C, C' such that $\|\exp(f)\|_{L^2_3} \leq C\|f\|_{L^2_3} \leq CC'\|f\|_{L^2_2}$, finishing the proof. \square

Let us compare different splittings. If s, s' are two different splittings, then for any $[u] \in \pi_0\mathcal{G}^h(X)$ we have $s(s')^{-1} \in \mathcal{G}^{h,e}(X)$. Therefore any two gauge splittings differ by a homomorphism $\pi_0\mathcal{G}^h(X) \rightarrow \mathcal{G}^{h,e}(X)$.

Our goal is to reduce the gauge group action to the action of S^1 and the action of a chosen lift of $\pi_0\mathcal{G}^h(X)$ to $\mathcal{G}^h(X)$. Precisely, we will consider splittings $s : \pi_0\mathcal{G}^h(X) \rightarrow \mathcal{G}^h(X)$ of (3). In order to choose the gauge fixing we need to understand that a gauge splitting induces another splitting on the level of homology.

Proposition 3.10 (*homological splitting*). *Any splitting $s : \pi_0\mathcal{G}^h(X) \rightarrow \mathcal{G}^h(X)$ of (3) induces a splitting $s^H : H^1(X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R})$ of the exact sequence*

$$0 \rightarrow H^0(\partial X; \mathbb{R})/H^0(X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R}) \rightarrow H^1(X; \mathbb{R}) \rightarrow 0. \quad (5)$$

Proof. Composing $H^1(X; \mathbb{Z}) \xrightarrow{2\pi i} \pi_0\mathcal{G}^h(X) \xrightarrow{s} \mathcal{G}^h(X) \xrightarrow{\frac{\delta}{2\pi i}} H^1(X, \partial X; \mathbb{R})$ we get a linear map which by linearity uniquely extends to a section $s_H : H^1(X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R})$ of the aforementioned exact sequence. \square

Definition 3.11 (*gauge splitting*). By a gauge splitting we call a splitting $s : \pi_0\mathcal{G}^h(X) \rightarrow \mathcal{G}^h(X)$ of the exact sequence (3). We denote by $s^H : H^1(X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R})$ the associated homological splitting.

We clarify the relationship between gauge splittings and homological splittings.

Proposition 3.12 (*gauge splittings from homological splittings*).

Let $\sigma : H^1(X; \mathbb{R}) \rightarrow H^1(X, \partial X; \mathbb{R})$ be a homological splitting, i.e., a splitting of (5). Then up to action of S^1 there exists a unique gauge splitting s such that $\sigma = s^H$.

Proof. For existence, choose $x_0 \in X$ and consider the map

$$I_{x_0} : \{\eta \in \mathcal{H}_D^1(X) \mid [\eta] \in H^1(X; 2\pi i\mathbb{Z})\} \longrightarrow \mathcal{G}^h(X), \quad (6)$$

$$\eta \longmapsto \left(I_{x_0}(\eta)(x) = \exp\left(\int_{x_0}^x \eta\right) \right).$$

Notice that $\delta(I_{x_0}(\eta)) = \eta$. Therefore taking $s([u]) = I_{x_0}(\sigma([\delta(u)]))$ we get a gauge

splitting $s : \pi_0 \mathcal{G}^h(X) \rightarrow \mathcal{G}^h(X)$ with $\sigma = s^H$.

The uniqueness up to action of S^1 follows from the exactness of the middle vertical sequence in (4). \square

In order to find the appropriate gauge fixing we need the following analogue of [Proposition 3.7](#) for 1-forms.

Lemma 3.13 (decomposing 1-forms).

$$\Omega^1(X) = \Omega_{CC}^1(X) + d(\Omega^0(X)) \text{ and } \Omega_{CC}^1(X) \cap d(\Omega^0(X)) = d(\mathcal{H}(X)).$$

Proof. This follows from the proof of [[Kha](#), Proposition 2.2]. (Note that our definition of $\Omega_{CC}^1(X)$ differs from Khandhawit's, which we denote by $\Omega_{s^\perp}^1(X)$ (cf. [Definition 3.21](#)).) \square

In particular, we can decompose 1-forms as

$$\Omega^1(X) = \Omega_{CC}^1(X) \oplus d(\Omega_\partial^0(X))$$

where

$$\Omega_\partial^0(X) = \left\{ f \in \Omega^0(X) \left| \int_{Y_i} f = 0 \text{ for each component } Y_i \subset \partial X \right. \right\}.$$

Denoting by Π_{CC} the projection onto $\Omega_{CC}^1(X)$ along $d(\Omega_\partial^0(X))$ we obtain the following analog of [Lemma 3.1](#).

Lemma 3.14 (Coulomb gauge fixing in 4d).

There is a unique homomorphism

$$\begin{aligned} L_1^2(i\Omega^1(X)) &\rightarrow \mathcal{G}^{e,\partial}(X) = \exp(L_2^2(i\Omega_\partial^0(X))) \\ a &\mapsto u_a^{CC} \end{aligned}$$

such that

$$a - (u_a^{CC})^{-1} du_a^{CC} = \Pi_{CC} a \in L_1^2(i\Omega_{CC}^1(X)).$$

Proof. The projection $(1 - \Pi_{CC})$ on $\Omega^1(X)$ has image in $d(\Omega_\partial^0(X))$ and that d is injective on $\Omega_\partial^0(X)$. Therefore there is a unique homomorphism $L_1^2(i\Omega^1(X)) \rightarrow L_2^2(i\Omega_\partial^0(X))$ sending a to the unique $f_a^{CC} \in \Omega_\partial^0(X)$ such that $a - df_a^{CC} \in \Omega^1(X)$. Then we take $u_a^{CC} = \exp(f_a^{CC})$. \square

We can further decompose

$$\Omega_{CC}^1(X) = \left(\Omega_{CC}^1(X) \cap (\mathcal{H}_D^1(X))^\perp \right) \oplus \mathcal{H}_D^1(X). \quad (7)$$

A homological splitting s^H provides a decomposition

$$\mathcal{H}_D^1(X) = s^H(H^1(X; \mathbb{R})) \oplus d(\mathcal{H}(X)) \quad (8)$$

which is an analogue of (3) for $\mathcal{H}_D^1(X)$. With these in hand, we are ready to define the split Coulomb slice.

Definition 3.15 (split Coulomb slice). Let s be a gauge splitting and s^H its associated homological splitting. The split Coulomb slice is

$$\Omega_s^1(X) = \{a \in \Omega_{CC}^1(X) \mid a \in (\Omega_{CC}^1(X) \cap (\mathcal{H}_D^1(X))^\perp) \oplus s^H(H^1(X; i\mathbb{R}))\}.$$

In particular, we have that

$$\Omega^1(X) = \Omega_s^1(X) \oplus d(\Omega^0(X)) \quad (9)$$

and parallel to Lemma 3.14 and Lemma 3.1 we can use the projection Π_s onto the first factor along the second one to obtain:

Lemma 3.16 (split gauge fixing in 4d). *There is a unique homomorphism*

$$\begin{aligned} L_1^2(i\Omega^1(X)) &\rightarrow \mathcal{G}^{e,\circ}(X) \\ a &\mapsto u_a^s \end{aligned}$$

such that

$$a - (u_a^s)^{-1} du_a^s = \Pi_s a \in L_1^2(i\Omega_s^1(X)).$$

Proof. The projection $(1 - \Pi_s)$ on $\Omega^1(X)$ has image in $d(\Omega^0(X))$ and d is injective on $\Omega_0^0(X)$. Therefore there is a unique homomorphism $L_1^2(i\Omega^1(X)) \rightarrow L_2^2(i\Omega_0^0(X))$ sending a to the unique $f_a^s \in \Omega_0^0(X)$ such that $a - df_a^s \in \Omega^1(X)$. Then we take $u_a^s = \exp(f_a^s)$. \square

Remark 3.17 (continuous gauge fixing within double Coulomb slice). If we only consider $a \in \Omega_{CC}^1(X)$, then the above map has image in $\mathcal{G}^h(X)$, which is finite-dimensional. Since the latter gauge group acts continuously on the configuration space by Lemma 3.9, we conclude that putting $(A, \Phi) \in \mathcal{C}_{CC}(X, \hat{\mathfrak{s}})$ into split Coulomb slice $\mathcal{C}_s(X, \hat{\mathfrak{s}})$ can be done continuously with respect to (A, Φ) .

The gauge group acting on this split Coulomb slice is the product of S^1 and the split gauge group:

Definition 3.18 (split harmonic gauge group). Let s be a gauge splitting. The split gauge group is defined to be

$$\mathcal{G}_s^{h,\circ}(X) = s(\pi_0 \mathcal{G}^h(X)).$$

Lemma 3.19 (split gauge group preserves the split Coulomb slice). *For $u \in \mathcal{G}_s^{h,\circ}(X)$ we have $u^{-1} du \in L_1^2(i\Omega_s^1(X))$.*

Conversely, if $u^{-1} du \in L_1^2(i\Omega_s^1(X))$, then for some $z \in S^1$ we have $zu \in \mathcal{G}_s^{h,\circ}(X)$.

Proof. One direction follows directly from the definition of s^H : if $u = s([u])$, then $s^H([\delta(u)]) = \delta(u) = u^{-1}du \in \mathcal{H}_D^1(X) \simeq H^1(X, \partial X; i\mathbb{R})$, so $u^{-1}du \in \text{im } s^H \subset L_1^2(i\Omega_s^1(X))$.

The other direction follows by chasing arrows in the diagram (4). \square

The circle \circ in the superscript indicates that the only constant gauge transformation contained in $\mathcal{G}_s^{h,\circ}(X)$ is the identity. This way we do not forget the S^1 -action when taking the quotient by the split gauge group.

We want to compare different split slices together with the split gauge group actions. Choose two splittings s, s' . These determine a map $s' \cdot s^{-1} : \pi_0 \mathcal{G}^h(X) \rightarrow \mathcal{G}^{h,e}(X)$. Viewing $\pi_0 \mathcal{G}^h(X)$ as a sublattice $\pi_0 \mathcal{G}^h(X) \simeq H^1(X; 2\pi i\mathbb{Z}) \subset H^1(X; i\mathbb{R})$, let $\nu : H^1(X; i\mathbb{R}) \rightarrow \mathcal{G}^{h,e}(X)$ be any homomorphism extending $s' \cdot s^{-1}$. Define

$$\begin{aligned} F_\nu : \mathcal{C}_{s'}(X, \hat{\mathfrak{s}}) &\longrightarrow \mathcal{C}_s(X, \hat{\mathfrak{s}}) \\ (A, \Phi) &\longmapsto \nu(\Pi_{\text{im } s^H}(A - A_0)) \cdot (A, \Phi). \end{aligned}$$

where $\Pi_{\text{im } s^H} : i\Omega_{CC}^1(X) \rightarrow s^H(H^1(X; i\mathbb{R}))$ is the projection along $i(\Omega_{CC}^1(X) \cap \mathcal{H}_D^1(X)) \oplus \text{id}(\mathcal{H}(X))$ (cf. (7), (8)).

Proposition 3.20 (equivalence of slices). *The map F_ν is well-defined, a diffeomorphism, equivariant with respect to the action of $\pi_0 \mathcal{G}^h(X) \simeq \mathcal{G}_{s'}^{h,\circ}(X) \simeq \mathcal{G}_s^{h,\circ}(X)$.*

Proof. Firstly, we need to show that the image of F_ν actually lies in $\mathcal{C}_s(X, \hat{\mathfrak{s}})$. Equivalently, we want to show that

$$\begin{aligned} A_\nu : \Omega_{CC}^1(X) &\longrightarrow \Omega_{CC}^1(X) \\ a &\longmapsto \nu(\Pi_{\text{im } s^H} a) \cdot a = a + \delta(\nu(\Pi_{\text{im } s^H} a)) \end{aligned}$$

maps $\Omega_s^1(X)$ to $\Omega_{s'}^1(X)$. We have $\delta(\mathcal{G}^{h,e}(X)) = d(\mathcal{H}(X))$ and, moreover, $\delta \circ \nu$ is a homomorphism, thus a linear map $H^1(X; i\mathbb{R}) \rightarrow d(\mathcal{H}(X))$. What follows is that A_ν is a linear map. Now A_ν restricted to $\Omega_{CC}^1(X) \cap (\mathcal{H}_D^1(X))^\perp$ is identity by definition, so it suffices to show $A_\nu(\text{im } s^H) \subset \text{im}((s')^H)$. Furthermore, $H^1(X; i\mathbb{R})$ is spanned by $[\delta(\pi_0 \mathcal{G}^h(X))]$ and therefore $s^H(H^1(X; i\mathbb{R}))$ is spanned by $\delta(\mathcal{G}_s^{h,\circ}(X))$, so it suffices to show $A_\nu(\delta(\mathcal{G}_s^{h,\circ}(X))) \subset \text{im}((s')^H)$. But we defined ν so that for any $u \in \mathcal{G}_s^{h,\circ}(X)$ $\nu(\delta(u)) \cdot u \in \mathcal{G}_{s'}^{h,\circ}(X)$, which implies $A_\nu(\delta(u)) = \delta(\nu(\delta(u)) \cdot u) \in \text{im}((s')^H)$, as wished.

The map F_ν is smooth because the map ν is smooth and the action of the finite-dimensional $\mathcal{G}^h(X)$ on $\mathcal{C}(X, \hat{\mathfrak{s}})$ is smooth.

It is invertible because $F_{1/\nu}$ is its inverse. Indeed, since $\text{im } \nu \subset \mathcal{G}^{h,e}(X)$, we have that $\delta\nu \in d(\mathcal{H}(X))$, so $\Pi_{\text{im } s^H}(\delta\nu) \equiv 0$. Therefore

$$\Pi_{\text{im } s^H}(\nu(\Pi_{\text{im } s^H}(A - A_0))A - A_0) = \Pi_{\text{im } s^H}(A - A_0),$$

so

$$(\nu(\Pi_{\text{im } s^H}(\nu(\Pi_{\text{im } s^H}(A - A_0))A - A_0)))^{-1} = (\nu(\Pi_{\text{im } s^H}(A - A_0)))^{-1}$$

and $F_{1/\nu} \circ F_\nu = \text{id}$ follows. \square

Finally, we discuss the gauge slice used by Lipyanskiy [Lip08] and Khandhawit [Kha, KLS18]. They require that $a \in \Omega_{CC}^1(X)$ and that for each component $Y_i \subset \partial X$ we have $\int_{Y_i} \iota^*(\star a) = 0$. Using Stokes' theorem one can show that for $a \in \Omega_{CC}^1(X)$ this integral condition is equivalent to the condition that $\int_X df \wedge \star a = 0$ for any $f \in \mathcal{H}(X)$. This fits into our setup perfectly, since there is exactly one homological splitting s_\perp^H such that $\text{im } s_\perp^H = \mathcal{H}_D^1(X) \cap (d(\mathcal{H}(X)))^\perp$.

Definition 3.21 (orthogonal splitting). We call s_\perp^H the orthogonal homological splitting. We say that a splitting s is a orthogonal splitting if $s^H = s_\perp^H$.

3.3 Restriction to the boundary and twisting

Unless ∂X is connected, we are not guaranteed that the restriction to the boundary is invariant under the action of the split gauge group $\mathcal{G}_s^{h,\circ}(X)$. If it happens to be invariant for some s , we call such s an integral splitting. For a general s , we introduce and prove the existence of twistings of the restriction map, making it invariant under the action of $\mathcal{G}_s^{h,\circ}(X)$ action even for non-integral s . As mentioned before, integral splittings are utilized in the proof of the [Gluing Theorem](#), while non-integral splittings may be more convenient in other contexts (e.g., in constructions of [Lip08, Kha, KLS18]).

We start by defining the restriction maps for an embedding $\iota_Y : Y \hookrightarrow X$ of an oriented 3-manifold Y . Denote by \mathfrak{s} the restriction to Y of the spin^c structure $\hat{\mathfrak{s}}$ on X . We get canonical identifications $S_X^\pm|_Y \simeq S_Y$. Assuming Y is a geodesic codimension-1 submanifold of X , the spin^c connection A_0 induces a spin^c connection B_0 on Y by simple restriction: $\nabla^{B_0} = \iota_Y^* \nabla^{A_0}$. Let $a \in L_1^2(i\Omega^1(X))$, $A \in A_0 + L_1^2(i\Omega^1(X))$, $\Phi \in L_1^2(\Gamma(S_X^\pm))$ and $u \in \mathcal{G}(X)$. We define the restrictions:

$$\begin{aligned} R(a) &= \iota_Y^*(a) \in L_{1/2}^2(i\Omega^1(Y)), \\ R(A) &= B_0 + \iota_Y^*(A - A_0) \in B_0 + L_{1/2}^2(i\Omega^1(Y)), \\ R(\Phi) &= \Phi|_Y \in L_{1/2}^2(\Gamma(S_Y)), \\ R(u) &= u|_Y \in \mathcal{G}(Y). \end{aligned}$$

Integral splittings are the ones for which restriction maps are invariant under the split gauge group.

Definition 3.22 (integral splitting). We call a gauge splitting s integral if for each $u \in \mathcal{G}_s^{h,\circ}(X)$ we have $u|_{\partial X} \equiv 1$.

Equivalently, s is integral if the composition $\pi_0 \mathcal{G}^h(X) \xrightarrow{s} \mathcal{G}^h(X) \xrightarrow{R} \mathcal{G}^h(\partial X) \simeq (S^1)^{\pi_0(\partial X)}$ is trivial.

The integrality of s is closely connected to the integrality of s^H .

Proposition 3.23 (homological classification of integral splittings). *If s is integral, then $s^H(H^1(X; \mathbb{Z})) \subset H^1(X, \partial X; \mathbb{Z})$, i.e., s^H is integral as well.*

Given any integral homological splitting σ there exists a unique integral splitting s such that $\sigma = s^H$.

Proof. Assume s is integral. Choose $y_0 \in \partial X$ and consider the map I_{y_0} defined in (6). We know s and $I_{y_0} \circ s^H \circ [\delta]$ differ by action of S^1 , but since both are equal to 1 at y_0 , thus $s = I_{y_0} \circ s^H \circ [\delta]$. This implies that for any $y \in \partial X$ and any embedded curve $\gamma : [0, 1] \rightarrow X$ with $\gamma(0) = y_0$ and $\gamma(1) = y$ we have that $\exp\left(\int_{y_0}^y s^H([\delta(u)])\right) = 1$ and thus $\int_{y_0}^y s^H([\delta(u)]) \in 2\pi i\mathbb{Z}$. This proves that s^H is integral.

Similarly, if σ is integral, then $s = I_{y_0} \circ \sigma \circ [\delta]$ satisfies that $\sigma = s^H$ and $s(y) = \exp\left(\int_{y_0}^y s^H([\delta(u)])\right) = 1$ for any $y \in \partial X$. \square

To find a consistent way of twisting the boundary values of 1-forms we consider ways to “undo” the action of $\mathcal{G}_s^{h,\circ}(X)$ on the boundary “in a linear fashion”.

Definition 3.24 (gauge twisting). We call a homomorphism $\tau : H^1(X; i\mathbb{R}) \rightarrow \mathcal{G}^h(\partial X) \simeq (S^1)^{\pi_0(\partial X)}$ a gauge twisting for s if the composition

$$\pi_0 \mathcal{G}^h(X) \simeq H^1(X; 2\pi i\mathbb{Z}) \hookrightarrow H^1(X; i\mathbb{R}) \xrightarrow{\tau} \mathcal{G}^h(\partial X)$$

agrees with the action of the split gauge group on the boundary, $R \circ s : \pi_0 \mathcal{G}^h(X) \rightarrow \mathcal{G}^h(\partial X)$.

Continuous homomorphisms from a vector space to S^1 correspond to linear functionals on the vector space. Thus, every such twisting τ comes from a linear map $d\tau : H^1(X; i\mathbb{R}) \rightarrow H^0(\partial X; i\mathbb{R})$ and $\tau = \exp \circ (d\tau)$. We utilize it to prove the existence of gauge twistings for s , and one could use it to classify all possible gauge twistings for s . Actually, every such homomorphism τ is a gauge twisting for some s , but we do not use this fact in this article.

Lemma 3.25 (existence of gauge twistings). *For a given gauge splitting s there exists a gauge twisting τ .*

Proof. Since $\pi_0\mathcal{G}^h(X)$ is free, we can lift the map $R \circ s : H^1(X; 2\pi i\mathbb{Z}) \simeq \pi_0\mathcal{G}^h(X) \rightarrow \mathcal{G}^h(\partial X) \simeq (S^1)^{\pi_0(\partial X)}$ to a homomorphism $\tilde{\tau} : H^1(X; 2\pi i\mathbb{Z}) \rightarrow H^0(\partial X; i\mathbb{R})$:

$$\begin{array}{ccc} & & H^0(\partial X; i\mathbb{R}) \\ & \nearrow \tilde{\tau} & \downarrow \text{exp} \\ \pi_0\mathcal{G}^h(X) & \xrightarrow{R \circ s} & (S^1)^{\pi_0(\partial X)}, \end{array}$$

and this extends to a map $\tilde{\tau} : H^1(X; i\mathbb{R}) \rightarrow H^0(\partial X; i\mathbb{R})$ by linearity. Taking $\tau = \text{exp} \circ \tilde{\tau} : H^1(X; i\mathbb{R}) \rightarrow (S^1)^{\pi_0(\partial X)}$ gives a gauge twisting for s . \square

With τ in hand, there is a way of defining a twisting on the whole Coulomb slice, enabling us to finally define the twisted restriction maps.

Definition 3.26 (twisted restriction map). We define the Coulomb slice twisting $\tau_{CC} : L_1^2(i\Omega_{CC}^1(X)) \rightarrow \mathcal{G}^h(\partial X)$ associated to τ to be the composition

$$\begin{aligned} L_1^2(i\Omega_{CC}^1(X)) &\xrightarrow{\Pi_{L^2, \perp}} \mathcal{H}_D^1(X) \simeq H^1(X, \partial X; \mathbb{R}) \\ &\xrightarrow{i_X^*} H^1(X; \mathbb{R}) \xrightarrow{\tau} (S^1)^{\pi_0(\partial X)} \simeq \mathcal{G}^h(\partial X). \end{aligned}$$

We define the twisted restriction map

$$R_\tau : \mathcal{C}_{CC}(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{cC}(\partial X, \mathfrak{s})$$

by the formula $R_\tau(A, \Phi) = (R(A), \tau_{CC}(A - A_0)R(\Phi))$.

Remark 3.27. What is of importance for defining the twisted restriction maps is the map $\tau_{CC} : i\Omega_s^1(X) \rightarrow (S^1)^{\pi_0(\partial X)}$. The extension of τ_{CC} to the whole of $i\Omega_{CC}^1(X)$ is artificial: it does not undo the action of $\mathcal{G}^{h,e}(X)$ on the boundary as one might expect.

With more work, including a choice of a based gauge group $\mathcal{G}_o^h(X) \subset \mathcal{G}^h(X)$ (such that $\mathcal{G}^h(X)/\mathcal{G}_o^h(X) \simeq S^1$) and a more general twisting, one could work with the full $i\Omega_{CC}^1(X)$ and then quotient by the action of $\mathcal{G}_o^h(X)$. However, this would introduce unnecessary complications.

These twisted restriction maps are indeed invariant under $\mathcal{G}_s^{h,\circ}(X)$.

Lemma 3.28 (twisted restriction map is invariant under split gauge group).

Let τ be a gauge twisting for s . For any $(A, \Phi) \in \mathcal{C}_{CC}(X, \hat{\mathfrak{s}})$ (resp. $(a, \phi) \in L_1^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X^+))$) and $u \in \mathcal{G}_s^{h,\circ}(X)$ we have

$$R_\tau(u(A, \Phi)) = R_\tau(A, \Phi)$$

(resp. $R_\tau(u(a, \phi)) = R_\tau(a, \phi)$).

Proof. Since $u \in \mathcal{G}^h(X)$, we have $\iota_{\partial X}^*(u^{-1}du) = 0$, so $R(A - u^{-1}du) = R(A)$.

It remains to prove

$$\tau_{CC}(A - A_0 - u^{-1}du)R(u\Phi) = \tau_{CC}(A - A_0)R(\Phi)$$

but that is equivalent to

$$(\tau_{CC}(-u^{-1}du)R(u))\tau_{CC}(A - A_0)R(\Phi) = \tau_{CC}(A - A_0)R(\Phi)$$

so it suffices to prove $\tau_{CC}(u^{-1}du) = R(u)$. Since s splits (3), we have $u = s([u])$, where $[u] \in \pi_0\mathcal{G}^h(X)$ is the homotopy class of u . So we have to prove $\tau_{CC}(u^{-1}du) = R \circ s([u])$. This follows directly from Definition 3.24 of the twistings and Definition 3.26 of the twisted restriction map, since $u^{-1}du \in \mathcal{H}_D^1(X)$ and the isomorphism $\pi_0\mathcal{G}^h(X) \simeq H^1(X; 2\pi i\mathbb{Z})$ is given by $[u] \mapsto [u^{-1}du]$. \square

We conclude these sections by showing that choosing τ is essentially equivalent to choosing an integral splitting. In general, one can restrict themselves to considering integral splittings without any twisting at all.

Proposition 3.29 (twistings are integral splittings). *Let τ be a twisting for s . Then there is an integral splitting $s_{\mathbb{Z}}$ and an equivariant diffeomorphism*

$$F_{s,\tau} : \mathcal{C}_s(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{s_{\mathbb{Z}}}(X, \hat{\mathfrak{s}})$$

such that $R \circ F_{s,\tau} = R_{\tau}$.

Proof. Every function $f \in \mathcal{H}(X)$ is determined by its restriction to the boundary $f|_{\partial X}$, which is locally constant. We thus have the exact sequence

$$0 \rightarrow H^0(\partial X; 2\pi i\mathbb{Z}) \rightarrow H^0(\partial X; i\mathbb{R}) \simeq i\mathcal{H}(X) \xrightarrow{f \mapsto \exp(f|_{\partial X})} \mathcal{G}^h(\partial X) \rightarrow 0$$

as well as

$$0 \rightarrow H^0(X; 2\pi i\mathbb{Z}) \rightarrow i\mathcal{H}(X) \xrightarrow{\exp} \mathcal{G}^{h,e}(X) \rightarrow 0$$

and from these two it follows that

$$0 \rightarrow H^0(\partial X; 2\pi i\mathbb{Z})/H^0(X; 2\pi i\mathbb{Z}) \rightarrow \mathcal{G}^{h,e}(X) \xrightarrow{\cdot|_{\partial X}} \mathcal{G}^h(\partial X) \rightarrow 0 \quad (10)$$

is exact. Since the group to the left is discrete it follows that there exists a unique lift

$\tilde{\tau}$ of τ to $\mathcal{G}^{h,e}(X)$:

$$\begin{array}{ccc} & & \mathcal{G}^{h,e}(X) \\ & \nearrow \tilde{\tau} & \downarrow \cdot|_{\partial X} \\ H^1(X; i\mathbb{R}) & \xrightarrow{\tau} & \mathcal{G}^h(\partial X) \end{array}$$

We define

$$s_{\mathbb{Z}}([u]) = (\tilde{\tau}([u]))^{-1} \cdot s([u])$$

for any $[u] \in \pi_0\mathcal{G}(X) \simeq H^1(X; 2\pi i\mathbb{Z})$, and

$$F_{s,\tau} = F_{(\tilde{\tau})^{-1}}$$

using the construction of F_ν of [Proposition 3.20](#). This gives an equivariant diffeomorphism from $\mathcal{C}_s(X, \hat{\mathfrak{s}})$ to $\mathcal{C}_{s_{\mathbb{Z}}}(X, \hat{\mathfrak{s}})$.

The equality $R \circ F_{s,\tau} = R_\tau$ follows from the construction. \square

Even though the spaces $\mathcal{C}_{s_{\mathbb{Z}}}(X, \hat{\mathfrak{s}})$ and $\mathcal{C}_{s'}(X, \hat{\mathfrak{s}})$ are equivariantly diffeomorphic by [Proposition 3.20](#), the corresponding restriction maps differ by a twist. Thus, *a priori* we cannot get rid of the choice of a splitting. However, this is not relevant to most of the applications because for connected boundary there is no choice to make.

Lemma 3.30 (uniqueness of integral splittings). *If $\partial X \neq \emptyset$ is connected, there exists exactly one integral splitting $s_{\mathbb{Z}}$.*

Proof. In this case, the restriction map $\mathcal{G}^{h,e}(X) \rightarrow \mathcal{G}^h(\partial X)$ is an isomorphism (cf. [\(10\)](#)). Therefore for each element $\pi_0\mathcal{G}(X)$ there exists exactly one representative $u \in \mathcal{G}^h(X)$ such that $u|_{\partial X} = 1$. \square

3.4 Seiberg-Witten moduli in split Coulomb slice

We conclude this section by defining the Seiberg-Witten moduli spaces, the main object of study of this article. We also prove they only depend on the choice of $s_{\mathbb{Z}}$ associated to s and τ .

Thanks to [Lemma 3.19](#), we can define the following.

Definition 3.31 (moduli spaces on 4-manifolds with boundary). We define the moduli spaces in split slice:

$$\begin{aligned} \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}}) &= \{(A, \Phi) \in \mathcal{C}_s(X, \hat{\mathfrak{s}}) \mid \text{SW}(A, \Phi) = 0\}, \\ \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) &= \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}}) / \mathcal{G}_s^{h,\circ}(X). \end{aligned}$$

We also define a version of the moduli space using the full double Coulomb slice,

$$\widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}}) = \{(A, \Phi) \in \mathcal{C}_{CC}(X, \hat{\mathfrak{s}}) \mid \text{SW}(A, \Phi) = 0\},$$

which will be utilized in some of the proofs.

From [Lemma 3.28](#) it follows that

Corollary 3.32. *There is a well-defined restriction map*

$$R_{\tau} : \mathcal{M}_s^{\circ}(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{cC}(\partial X, \mathfrak{s}).$$

A direct consequence of [Proposition 3.29](#) is

Corollary 3.33 (dependence on twistings). *Given s and τ , there is an integral splitting $s_{\mathbb{Z}}$ and a diffeomorphism*

$$F_{s,\tau} : \mathcal{M}_s^{\circ}(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{M}_{s_{\mathbb{Z}}}^{\circ}(X, \hat{\mathfrak{s}})$$

such that $R \circ F_{s,\tau} = R_{\tau}$.

4 Properties of moduli spaces

In this section we prove that ([Semi-infinite-dimensionality Theorem](#)):

- the moduli spaces of solutions to the Seiberg-Witten equations on X are Hilbert manifolds,
- the restriction map to the boundary is “semi-infinite”, i.e., Fredholm in the negative direction and compact in the positive direction,
- if ∂X is disconnected, restriction to a single boundary component has dense differential.

This is done by analyzing the properties of the linearized Seiberg-Witten operator DSW . We start by investigating an extended version of this operator, \widetilde{DSW} . The reason is that the standard Atiyah-Patodi-Singer boundary value problem as well as the elliptic theory developed in [section 2](#) can be directly applied to the study of \widetilde{DSW} . Our understanding of the gauge action ([subsection 3.2](#)) will allow us to transfer these properties to DSW .

4.1 Extended linearized SW operator

Here we apply the Atiyah-Patodi-Singer boundary value problem to an extended version of the linearized Seiberg-Witten operator, \widetilde{DSW} . The properties we prove are the direct analogues of the properties of DSW which are proved in the next section.

Definition 4.1 (extended linearized SW operator). We define the extended linearized Seiberg-Witten operator

$$\begin{aligned} \widetilde{D}SW_{(A,\Phi)} : L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)) \\ \rightarrow L^2(X; i\mathbb{R}) \oplus L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \end{aligned}$$

by adding a component related to the linearization of gauge action:

$$\widetilde{D}SW_{(A,\Phi)}(a, \phi) = (d^*a, DSW_{(A,\Phi)}(a, \phi)).$$

In order to study the Atiyah-Patodi-Singer boundary value problem we need to introduce the appropriate operator on the boundary and consider its Calderón projector. Denote $Y = \partial X$ and define

$$\begin{aligned} \tilde{L} : i\Omega^1(Y) \oplus \Gamma(S_Y) \oplus i\Omega^0(Y) &\rightarrow i\Omega^1(Y) \oplus \Gamma(S_Y) \oplus i\Omega^0(Y), \\ \tilde{L}(b, \psi, c) &= (\star db - dc, \not{D}_{B_0} \psi, -d^*b). \end{aligned}$$

This is a first-order self-adjoint elliptic operator. Denote by $\tilde{H}^+(Y, \mathfrak{s})$ (resp. $\tilde{H}^-(Y, \mathfrak{s})$) the closure of the span of positive (resp. nonpositive) eigenspaces of \tilde{L} in $L_{1/2}^2(i\Omega^1(Y) \oplus \Gamma(S_Y) \oplus i\Omega^0(Y))$, and by $\tilde{\Pi}^\pm$ the projection onto $\tilde{H}^\pm(Y, \mathfrak{s})$ along $\tilde{H}^\mp(Y, \mathfrak{s})$. The proof of the following proposition follows a standard argument; we briefly recall it to set up the stage for the proofs in the rest of this section.

Proposition 4.2 (semi-infinite-dimensionality of $\widetilde{D}SW$).

The operator

$$\begin{aligned} \widetilde{D}SW_{(A,\Phi)} \oplus \tilde{\Pi}^- R : L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)) \\ \rightarrow L^2(X; i\mathbb{R}) \oplus L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \oplus \tilde{H}^-(Y, \mathfrak{s}) \end{aligned} \quad (11)$$

is Fredholm of index

$$2 \operatorname{ind}_{\mathbb{C}} \not{D}_{A_0} + b_1(X) - b^+(X) - b_1(Y) - 1. \quad (12)$$

Moreover, the positive part of the restriction map from the kernel of $\widetilde{D}SW$, $\tilde{\Pi}^+ R : \ker(\widetilde{D}SW_{(A,\Phi)}) \rightarrow \tilde{H}^+(Y, \mathfrak{s})$, is compact.

Proof. We can write $\widetilde{D}SW_{(A,\Phi)} = \tilde{D} + \tilde{K}$ where

$$\tilde{D}(a, \phi) = (d^+a, \not{D}_{A_0} \phi, d^*a)$$

and

$$\tilde{K}(a, \phi) = (0, -\rho^{-1}(\phi\Phi^* + \Phi\phi^*)_0, \rho(a)\Phi + \rho(A^\delta)\phi).$$

As explained in [Kha, Proposition 3.1], applying the Atiyah-Patodi-Singer boundary value problem [KM07, Theorem 17.1.3] to \tilde{D} proves that $\tilde{D} \oplus \tilde{\Pi}R$ is Fredholm with index equal to (12). Furthermore, [KM07, Theorem 17.1.3] implies that for any bounded sequence $(u_i) \subset L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+))$ such that $(\tilde{D}(u_i))$ is Cauchy, the sequence $(\tilde{\Pi}^+Ru_i)$ is precompact.

The operator \tilde{K} is compact by Theorem A.1. Since $\widetilde{DSW}_{(A,\Phi)} = \tilde{D} + \tilde{K}$, thus $\widetilde{DSW}_{(A,\Phi)}$ is Fredholm with the same index as \tilde{D} . Moreover, since \tilde{K} is compact, for any sequence $(u_i) \subset \ker(\widetilde{DSW}_{(A,\Phi)})$ we can choose a subsequence such that the sequence of $\tilde{D}(u_i) = -\tilde{K}(u_i)$ is convergent, thus Cauchy. By what we proved in the previous paragraph, the sequence $(\tilde{\Pi}^+Ru_i)$ is precompact. This shows that $\tilde{\Pi}^+R : \ker(\widetilde{DSW}_{(A,\Phi)}) \rightarrow \tilde{H}^+(Y, \mathfrak{s})$ is compact. \square

The proof of surjectivity utilizes both of the analytical results of section 2 (cf. [Lip08, Theorem 2]).

Proposition 4.3 (surjectivity of \widetilde{DSW}). *The operator $\widetilde{DSW}_{(A,\Phi)}$ is surjective.*

Proof. Assume $\widetilde{DSW}_{(A,\Phi)}$ is not surjective. Proposition 4.2 implies its image is closed, so there is $0 \neq \tilde{v} \in \mathcal{V}(X, \hat{\mathfrak{s}}) \oplus L^2(i\Omega^0(X))$ orthogonal to $\text{im } \widetilde{DSW}$. Recall that \tilde{K} is a certain multiplication by $p = (A - A_0, \Phi) \in L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+))$. Let $X^* = X \cup ([0, \infty) \times Y)$ with cylindrical metric on the end, and extend the spinor bundle S_X to S_{X^*} which is cylindrical on ends. Extend p to $p^* \in L_1^2(i\Omega^1(X^*) \oplus \Gamma(S_{X^*}^+))$ in an arbitrary way and \tilde{v} to $\tilde{v}^* \in \mathcal{V}(X^*, \hat{\mathfrak{s}}) \oplus L^2(i\Omega^0(X^*))$ by zero on $[0, \infty) \times Y$. We have

$$\langle \tilde{v}^*, (\tilde{D} + \tilde{K})(w) \rangle_{L^2(X^*)} = \langle \tilde{v}, \widetilde{DSW}_{(A,\Phi)}(w|_X) \rangle_{L^2(X)} = 0$$

for any $w \in \mathcal{TC}(X^*, \hat{\mathfrak{s}})$. Therefore \tilde{v}^* is a weak solution to $\tilde{D}^*\tilde{v}^* + \tilde{K}^*\tilde{v}^* = 0$ where \tilde{D}^*, \tilde{K}^* are formal adjoints of \tilde{D}, \tilde{K} , respectively. The map $\tilde{K}^* : L_1^2 \rightarrow L^2$ is compact by Theorem A.1. Thus from the Regularity Theorem it follows that $\tilde{v}^* \in L_1^2(i\Omega^+(X^*) \oplus \Gamma(S_{X^*}^-) \oplus i\Omega^0(X^*))$ and it is a solution to $(\tilde{D}^* + \tilde{K}^*)\tilde{v}^* = 0$. Furthermore, Corollary 2.4 implies that $\tilde{v}^* = 0$ and therefore $v = 0$. Thus, by contradiction, we have proved that DSW is surjective. \square

Finally, we focus on the density of the restriction map from the kernel of \widetilde{DSW} to one boundary component. The proof of this proposition utilizes some of the ideas we have just seen (cf. [Lip08, Lemma 5]).

Proposition 4.4 (density of moduli on one boundary component). *Assume the boundary Y has at least two connected components and let $Y_0 \subset Y$ be any one of these components. Then the restriction*

$$R : \ker \widetilde{D}\text{SW}_{(A,\Phi)} \rightarrow L^2_{1/2}(i\Omega^1(Y_0) \oplus \Gamma(S_{Y_0}))$$

is dense.

Proof. Assume, by contradiction, that it is not dense and choose a nonzero element $v \in L^2_{1/2}(i\Omega^1(Y_0) \oplus \Gamma(S_{Y_0}))$ which is $L^2_{1/2}$ -perpendicular to its image. Since $\widetilde{D}\text{SW}_{(A,\Phi)}$ is surjective by [Proposition 4.3](#), the map

$$\begin{aligned} \widetilde{D}\text{SW}_{(A,\Phi)} \oplus \Pi_v R : L^2_1(i\Omega^1(X) \oplus \Gamma(S_X^+)) \\ \rightarrow L^2(X; i\mathbb{R}) \oplus L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \oplus \mathbb{C}v \end{aligned}$$

has finite-dimensional cokernel, where Π_v is the $L^2_{1/2}$ -projection onto $v \in L^2_{1/2}(i\Omega^1(Y_0) \oplus \Gamma(S_{Y_0}))$.

From the definition of v it follows that $\widetilde{D}\text{SW}_{(A,\Phi)} \oplus \Pi_v R$ is not surjective, since otherwise there would be an element $w \in L^2_1(i\Omega^1(X) \oplus \Gamma(S_X^+))$ which solves $\widetilde{D}\text{SW}_{(A,\Phi)}(w) = 0$ such that $\Pi_v(R(w)) \neq 0$. Therefore, we can pick an element (a, v) which is orthogonal to the image of $\widetilde{D}\text{SW}_{(A,\Phi)} \oplus \Pi_v R$, where $a \in L^2(X; i\mathbb{R}) \oplus L^2(i\Omega^+(X) \oplus \Gamma(S_X^-))$.

As in the proof of [Proposition 4.3](#), attach a cylindrical end along Y to get $X^* = X \cup [0, \infty) \times Y$. Extend a to a^* by 0 on $[0, \infty) \times Y$. Since \tilde{K} is a certain multiplication by $p = (A - A_0, \Phi) \in L^2_1(i\Omega^1(X) \oplus \Gamma(S_X^+))$, extend p to $p^* \in L^2_1(i\Omega^1(X^*) \oplus \Gamma(S_{X^*}^+))$ in an arbitrary way to get \tilde{K} defined on X^* . By [Theorem A.1](#), \tilde{K} is a compact operator $L^2_1 \rightarrow L^2$.

Since $(a, v) \perp \text{im}(\widetilde{D}\text{SW}_{(A,\Phi)} \oplus \Pi_v R)$, we get that a^* is a weak solution to $(\tilde{D}^* + \tilde{K}^*)a^* = 0$ on the interior of $X^*_1 = X^* \setminus ([0, \infty) \times Y_0)$. Take any compact set $C \subset X^*_1$ and a smooth bump function $\eta : X^* \rightarrow [0, 1]$ such that $\eta|_C = 1$ and $\text{supp } \eta \subset X^*_1$. It follows that $(D^* + K^*)(\eta a) = \rho^*(d\eta)a \in L^2(X^*)$. Therefore by the [Regularity Theorem](#) we get that $\eta a \in L^2_1(X^*)$, and in particular $a \in L^2_1(\overset{\circ}{C})$. Varying C we obtain $a^* \in L^2_{1,loc}(X^*_1)$. Since $a^* \equiv 0$ on $[0, \infty) \times (Y \setminus Y_0) \subset X^*_1$, [Corollary 2.4](#) implies that $a^* = 0$ on X^*_1 , so $a = 0$ on X .

This is a contradiction since we can extend $v \in L^2_{1/2}(i\Omega^1(Y_0) \oplus \Gamma(S_{Y_0}))$ to $\tilde{v} \in L^2_1(i\Omega^1(X) \oplus \Gamma(S_X^+))$ such that $R(\tilde{v}) = v$. Then

$$\begin{aligned} 0 &= \langle (\widetilde{D}\text{SW}_{(A,\Phi)}(\tilde{v}), \Pi_v R\tilde{v}), (a, v) \rangle_{L^2_{1/2}} \\ &= \langle (\widetilde{D}\text{SW}_{(A,\Phi)}(\tilde{v}), v), (0, v) \rangle_{L^2_{1/2}} \\ &= 0 + \langle v, v \rangle_{L^2_{1/2}} \end{aligned}$$

implying $\|v\|_{L^2_{1/2}} = 0$, which contradicts the assumption that $v \neq 0$. \square

4.2 Regularity and semi-infinite-dimensionality of moduli spaces

We turn our focus to the operator DSW . The main difficulty in transferring the results of [subsection 4.1](#) from \widetilde{DSW} to DSW is the presence of the Coulomb condition on both X and ∂X . The split gauge condition introduces an additional twist to the story.

A key fact is that the differential of the gauge group action at e (cf. [Lemma 3.5](#)) preserves the kernel of DSW at a solution (A, Φ) .

Lemma 4.5. *Assume $\not{D}_A \Phi = 0$. Then for any $f \in L^2_2(i\Omega^0(X))$ we have $DSW_{(A,\Phi)}(df, -f\Phi) = 0$.*

Proof. We compute:

$$\begin{aligned} & (\hat{D} + \hat{K})(df, -f\Phi) \\ &= (\rho^{-1}(f\Phi\Phi^* + \Phi(f\Phi)^*)_0, \rho(df)\Phi - \not{D}_{A_0}(f\Phi) - \rho(A^\delta)f\Phi) \\ &= (\rho^{-1}(f\Phi\Phi^* - \Phi f\Phi^*)_0, \rho(df)\Phi - \rho(df)\Phi - f(\not{D}_{A_0}\Phi + \rho(A^\delta)\Phi)) \\ &= (0, -f\not{D}_A\Phi) \\ &= (0, 0), \end{aligned}$$

which finishes the proof. \square

Following the idea of [[Kha](#), Proposition 3.1], we deduce the semi-infinite-dimensionality and compute the index of DSW from [Proposition 4.2](#). These methods will be utilized to prove further results in this section, too.

Proposition 4.6 (semi-infinite-dimensionality of DSW). *The operator*

$$\begin{aligned} D_{(A,\Phi)}SW \oplus \Pi^- R : L^2_1(i\Omega^1_s(X) \oplus \Gamma(S^+_X)) & \quad (13) \\ & \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S^-_X)) \oplus H^-(Y, \mathfrak{s}) \end{aligned}$$

is Fredholm of index

$$2 \operatorname{ind}_{\mathbb{C}} + b_1(X) - b^+(X) - b_1(Y). \quad (14)$$

Moreover, the restriction $\Pi^+ R : \ker(D_{(A,\Phi)}SW) \rightarrow H^+(Y, \mathfrak{s})$ is compact, where $\ker(D_{(A,\Phi)}SW) \subset L^2_1(i\Omega^1_s(X) \oplus \Gamma(S^+_X))$.

Proof. Firstly, we compare the respective polarizations. Recall that the decomposition of $L^2_{1/2}(i\Omega^1(Y) \oplus \Gamma(S_Y))$ into $\tilde{H}^+(Y, \mathfrak{s}) \oplus \tilde{H}^-(Y, \mathfrak{s})$ is given by the eigenspaces of \tilde{L} . Decomposing

$$i\Omega^1(Y) \oplus \Gamma(S_Y) \oplus i\Omega^0(Y) = (i\Omega^1_{cC}(Y) \oplus \Gamma(S_Y)) \oplus (\Omega^1_C(Y) \oplus i\Omega^0(Y))$$

we see that \tilde{L} decomposes as $\begin{pmatrix} \star d & 0 \\ 0 & \not{D}_{B_0} \end{pmatrix} \oplus \begin{pmatrix} 0 & -d \\ -d^* & 0 \end{pmatrix}$. Denote by Π_1^\pm the spectral projections of $\begin{pmatrix} 0 & -d \\ -d^* & 0 \end{pmatrix}$ in $L^2_{1/2}(i\Omega^1_C(Y) \oplus i\Omega^0(Y))$. It follows that $\tilde{\Pi}^\pm = \Pi^\pm \oplus \Pi_1^\pm$.

This is enough to prove the statement about compactness. Indeed, the map $\Pi^+ R : \ker(D_{(A,\Phi)} \text{SW}) \rightarrow H^+(Y, \mathfrak{s})$ is just the composition

$$\ker(D_{(A,\Phi)} \text{SW}) \hookrightarrow \ker(\widetilde{D \text{SW}}_{(A,\Phi)}) \xrightarrow{\tilde{\Pi}^+ R} \tilde{H}^+(Y, \mathfrak{s}) \xrightarrow{\Pi^+} H^+(Y, \mathfrak{s})$$

where the map in the middle is compact by [Proposition 4.2](#).

For Fredholmness, denote by

$$\Pi_C : L^2_{1/2}(\Omega^1_C(Y) \oplus \Omega^0(Y)) \rightarrow L^2_{1/2}(\Omega^1_C(Y) \oplus \Omega^0(Y))$$

the projection onto $L^2_{1/2}(\Omega^1_C(Y) \oplus \mathbb{R}^{\pi_0(Y)})$ (where $\mathbb{R}^{\pi_0(Y)} \subset \Omega^0(Y)$ is the space of locally constant functions) along $\{0\} \oplus L^2_{1/2}(\Omega^0_0(Y))$ (where $\Omega^0_0(Y) = \{f \in \Omega^0(Y) \mid \forall_i \int_{Y_i} f = 0\}$). This projection can be used to define the split Coulomb slice for the orthogonal splitting s_\perp ([Definition 3.21](#)). Precisely, we have

$$L^2_1(i\Omega^1_{s_\perp}(X) \oplus \Gamma(S^+_X)) = \ker(d^*, \Pi_C).$$

Khandhawit [[Kha](#), Proposition 3.1] proves that $\text{im } \Pi_1^-$ and $\ker \Pi_C$ are complementary, and then [[KM07](#), Proposition 17.2.6] implies that

$$\begin{aligned} \widetilde{D \text{SW}}_{(A,\Phi)} \oplus (\Pi^- R) \oplus (\Pi_C R) : L^2_1(i\Omega^1(X) \oplus \Gamma(S^+_X)) \\ \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S^-_X)) \oplus L^2_{1/2}(i\Omega^1_C(Y) \oplus \mathbb{R}^{\pi_0(Y)} \oplus H^-(Y, \mathfrak{s})) \end{aligned} \quad (15)$$

is Fredholm since (11) is Fredholm. Since $\Pi_C|_{\text{im } \Pi_1^-}$ is an isomorphism onto $\text{im } \Pi_C$, therefore the proof of [[KM07](#), Proposition 17.2.6] implies that the index of (15) is the same as the index of (11).

The following lemma is a simple exercise in Fredholm theory.

Lemma 4.7. *Let $(F, G) : H \rightarrow A \oplus B$ be Fredholm. Then $\tilde{F} = F|_{\ker G} : \tilde{H} = \ker G \rightarrow A$ is Fredholm and has index equal to $\text{ind}((F, G)) + \dim \text{coker } G$.*

It implies that the operator $D_{(A,\Phi)} \text{SW} \oplus (\Pi^- R)$ is Fredholm as a map $\ker(\star d \oplus (\Pi_C R)) = L_1^2(i\Omega_{s_\perp}^1(X) \oplus \Gamma(S_X^+)) \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \oplus H^-(Y, \mathfrak{s})$ and has index

$$\begin{aligned} & \text{ind}\left(\widetilde{D\text{SW}}_{(A,\Phi)} \oplus (\Pi^- R) \oplus (\Pi_C R)\right) + \dim \text{coker}(\star d \oplus (\Pi_C R)) \\ &= [2 \text{ind}_{\mathbb{C}} \mathcal{D}_{A_0}^+ - b_0(X) + b_1(X) - b^+(X) - b_1(Y)] + b_0(X) \\ &= 2 \text{ind}_{\mathbb{C}} \mathcal{D}_{A_0}^+ + b_1(X) - b^+(X) - b_1(Y). \end{aligned}$$

Thus, we have proven the Proposition for a particular splitting, $s = s_\perp$.

For any splitting s , the inclusion $\Omega_s^1(X) \hookrightarrow \Omega_{CC}^1(X)$ is of codimension $\dim(H^1(X, Y; \mathbb{R})) - \dim(H^1(X; \mathbb{R})) = b_0(Y) - 1$. Therefore, the split double Coulomb slice $L_1^2(i\Omega_s^1(X) \oplus \Gamma(S_X^+))$ is a finite-dimensional subspace of the “full” double Coulomb slice $L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+))$. Therefore the Fredholmness of (13) for $s = s_\perp$ implies the Fredholmness of

$$\begin{aligned} D_{(A,\Phi)} \text{SW} \oplus \Pi^- R : L_1^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X^+)) & \quad (16) \\ \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \oplus H^-(Y, \mathfrak{s}) & \end{aligned}$$

and this, in turn, implies the Fredholmness of (13) for any splitting s . Moreover, the index of (16) is equal to $b_0(Y) - 1$ plus (14) for any splitting s , finishing the proof. \square

We turn to deducing the surjectivity of $D\text{SW}$ from the surjectivity of $\widetilde{D\text{SW}}$.

Proposition 4.8 (surjectivity of $D\text{SW}$). *For any gauge splitting s , the differential*

$$D_{(A,\Phi)} \text{SW} : L_1^2(i\Omega_s^1(X) \oplus \Gamma(S_X^+)) \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-))$$

is surjective.

Proof. We will prove a stronger statement, that this extended differential together with the exact part of the restriction to the boundary

$$\begin{aligned} (\widetilde{D\text{SW}}_{(A,\Phi)}, \Pi_d R, \Pi_{V_s}) &= (D\text{SW}_{(A,\Phi)}, d^*, \Pi_d R, \Pi_{V_s}) : & (17) \\ L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)) &\rightarrow \\ L^2(i\Omega^+(X) \oplus \Gamma(S_X^-)) \oplus L^2(i\Omega^0(X)) \oplus L_{1/2}^2(i\Omega_C^1(Y)) \oplus V_s & \end{aligned}$$

is surjective, where $\Pi_d : L_{1/2}^2(i\Omega^1(Y)) \rightarrow L_{1/2}^2(i\Omega_C^1(Y))$ is the projection along $L_{1/2}^2(i\Omega_{CC}^1(Y))$ and $\Pi_{V_s} : \Omega^1(X) \rightarrow V_s = (\text{im } s)^\perp$ is the orthogonal projection. The Proposition will follow since $L_1^2(i\Omega_s^1(X) \oplus \Gamma(S_X^+)) = \ker(d^*, \Pi_d R, \Pi_{V_s})$.

Proposition 4.3 implies that $\widetilde{D\text{SW}}_{(A,\Phi)} : L_1^2(i\Omega^1(X) \oplus \Gamma(S_X^+)) \rightarrow L^2(i\Omega^+(X) \oplus \Gamma(S_X^-))$ is surjective. To prove surjectivity of (17) it thus remains to prove that $\Pi_d R : \ker(\widetilde{D\text{SW}}_{(A,\Phi)}) \rightarrow L_{1/2}^2(i\Omega_C^1(Y))$ is surjective and that $\Pi_{V_s} : \ker(\widetilde{D\text{SW}}_{(A,\Phi)}, \Pi_d R) \rightarrow V_s$ is surjective.

We prove that $\Pi_d R|_{\ker \widetilde{D}SW_{(A,\Phi)}}$ is surjective. Take any $g \in L^2_{3/2}(i\Omega^0(Y))$ representing a given element $dg \in L^2_{1/2}(i\Omega^1_C(Y))$. Take the unique $f \in L^2_2(i\Omega^0(X))$ such that $\Delta f = 0$ and $f|_Y = g$. Then $\Pi_d R(df, -f\Phi) = df|_Y = dg$. The required surjectivity follows since $(df, -f\Phi) \in \ker \widetilde{D}SW_{(A,\Phi)}$, which follows from $d^*df = \Delta f = 0$ and [Lemma 4.5](#).

Similarly we prove $\Pi_{V_s}|_{\ker(\widetilde{D}SW_{(A,\Phi),\Pi_d R})}$ is surjective. By (8) and definition of V_s , the orthogonal projection $d(\mathcal{H}(X)) \rightarrow V_s$ is an isomorphism and therefore for any $v \in V_s$ there is $f \in \mathcal{H}(X)$ such that $\Pi_{V_s}(df) = v_s$. Moreover $\widetilde{D}SW(df, -f\Phi) = 0$ and $\Pi_d R(df, -f\Phi) = 0$, as wished. \square

Finally, we prove the density of the restriction map to a connected component of Y .

Proposition 4.9 (density of moduli on one boundary component). *The restriction*

$$R : \ker(D_{(A,\Phi)}SW) \rightarrow L^2_{1/2}(i\Omega^1_{cC}(Y_0) \oplus \Gamma(S_{Y_0}))$$

to a connected component $Y_0 \subset Y$ is dense, where we consider $\ker(D_{(A,\Phi)}SW) \subseteq L^2_1(i\Omega^1_s(X) \oplus \Gamma(S^+_X))$.

Proof. Take any $(b, \psi) \in L^2_{1/2}(i\Omega^1_{cC}(Y_0) \oplus \Gamma(S_{Y_0})) \subset L^2_{1/2}(i\Omega^1(Y_0) \oplus \Gamma(S_{Y_0}))$. It follows from [Proposition 4.4](#) that we can take a sequence $(\tilde{a}_k, \phi_k) \in \ker(\widetilde{D}SW_{(A,\Phi)})$ such that $R_{Y_0}(\tilde{a}_k, \phi_k) \rightarrow (b, \psi)$ in $L^2_{1/2}(i\Omega^1_{cC}(Y_0) \oplus \Gamma(S_{Y_0}))$. Using the decomposition (9) we can write $\tilde{a}_k = a_k + df_k$ for $f_k \in i\Omega^0(X)$ and $a_k \in \Omega^1_s(X)$. Since Y_0 is connected, we can change f_k by a constant to obtain $\int_{Y_0} f_k = 0$. Decomposing $R_{Y_0}(\tilde{a}_k) = R_{Y_0}(a_k) + R_{Y_0}(df_k)$ we get that $R_{Y_0}(a_k) \rightarrow b$ and $R_{Y_0}(df_k) \rightarrow 0$. This together with $\int_{Y_0} f_k = 0$ implies that $f_k|_{Y_0} \rightarrow 0$ and therefore

$$(b, \psi) = \lim_{k \rightarrow \infty} R_{Y_0}(\tilde{a}_k, \phi_k) = \lim_{k \rightarrow \infty} R_{Y_0}(a_k, \phi_k + if_k\Phi)$$

which finishes the proof because $a_k \in L^2_1(i\Omega^1_s(X))$ and $D SW(a_k, \phi_k + if_k\Phi) = D SW(a_k + idf_k, \phi_k) = 0$ by [Lemma 4.5](#). \square

The results of [Proposition 4.6](#), [Proposition 4.8](#) and [Proposition 4.9](#) can be summarized as follows.

Semi-infinite-dimensionality Theorem. *The moduli spaces $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$ are Hilbert manifolds. The differential of the twisted restriction map $R_\tau : \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_{cC}(\partial X, \mathfrak{s})$ decomposes into $\Pi^- DR_\tau : T\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow H^-(\partial X, \mathfrak{s})$ which is Fredholm and $\Pi^+ DR_\tau : T\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow H^+(\partial X, \mathfrak{s})$ which is compact.*

Moreover, if $b_0(\partial X) > 1$, then for any connected component $Y_0 \subset \partial X$ the restriction R_{τ, Y_0} to Y_0 has dense differential.

5 Gluing along a boundary component

This section is devoted to the proof of the main result of this article, the [Gluing Theorem](#), which relates the moduli spaces of solutions on X_1 , X_2 and $X = X_1 \cup_Y X_2$, where Y is a rational homology sphere, oriented as a component of the boundary of X_1 . Under the identification $\mathcal{C}_{cC}(Y, \mathfrak{s}) \simeq \mathcal{C}_{cC}(-Y, \bar{\mathfrak{s}})$ there are S^1 -equivariant twisted restriction maps $R_{\tau_i, Y} : \mathcal{M}_{s_i}^\circ(X_i, \hat{\mathfrak{s}}_i) \rightarrow \mathcal{C}_{cC}(Y, \mathfrak{s})$, where $\mathfrak{s} = \hat{\mathfrak{s}}|_Y$. One can expect the fiber product

$$\begin{aligned} \mathcal{M}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times_Y \mathcal{M}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2) &= \\ &= \{(A_1, \Phi_1, A_2, \Phi_2) \in \mathcal{M}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times \mathcal{M}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2) \\ &\quad | R_{\tau_1, Y}(A_1, \Phi_1) = R_{\tau_2, Y}(A_2, \Phi_2)\} \end{aligned}$$

to be diffeomorphic to $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$, and this turns out to be true. One would also like to have this map intertwine the twisted restriction maps to ∂X , but this is a bit too much: the splittings and twistings (s, τ) , (s_1, τ_1) , (s_2, τ_2) , need to enjoy certain compatibility, and even then the restriction maps may not match on the nose but need to be homotoped to each other. This reflects the fact that we did not quotient by the action of S^1 on the configuration spaces.

The proof utilizes the following fact which is of independent interest. Let $X' \subset X$ be a submanifold, the closure of which is contained in the interior of X . Then the restriction map from $\widetilde{\mathcal{M}}_{cC}^\circ(X, \hat{\mathfrak{s}})$ to L_k^2 -configurations on X' is well-defined and *smooth* for any $k \geq 0$. Well-definedness follows from a standard argument, but proving the smoothness of this map turns out to be a surprisingly delicate task which we tackle in [subsection 5.1](#).

The same strategy should work to prove smoothness of restriction maps to interior submanifolds for other types of moduli spaces appearing in gauge theory, e.g., for the space of anti-self-dual connections on $G \hookrightarrow P \rightarrow X$. The key is the ellipticity of the equations together with the gauge fixing.

5.1 Smoothness of restrictions

Assume $X' \subset \mathring{X}$ is a submanifold with closure contained in \mathring{X} . The goal is to show (cf. [Theorem 5.2](#)) that the restriction map $R : \widetilde{\mathcal{M}}_{cC}^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_k(X', \hat{\mathfrak{s}}|_{X'})$ is smooth for any k , where $\mathcal{C}_k(X', \hat{\mathfrak{s}}|_{X'}) = (A_0|_{X'}, 0) + L_k^2(i\Omega^1(X') \oplus \Gamma(S_{X'}^+))$. We restrict ourselves to the case $k = 2$, but the same strategy may be used iteratively, bootstrapping the result to any k , if needed.

Due to [Theorem A.2](#) we may assume, without loss of generality, that X' is of codimension 0. Since we require the closure of X' to be contained in the interior \mathring{X} , we may as well assume that X' is a closed submanifold.

The following fundamental fact shows that any element in the image of the restriction map is itself a smooth configuration.

Lemma 5.1 (interior smoothness of solutions). [KM07, Lemma 5.1.5] *Every $\gamma \in \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}})$ is smooth on \mathring{X} .*

We are ready to prove the main theorem of this subsection. Note that the surjectivity assumption is satisfied whenever $\partial X \neq \emptyset$ due to [Proposition 4.8](#).

Theorem 5.2 (restriction is smooth on solution sets). *Assume that for any $(A, \Phi) \in \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}})$ the operator $DSW_{(A, \Phi)}$ is surjective. Then the restriction map $R : \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{C}_k(X', \hat{\mathfrak{s}}|_{X'})$ is smooth.*

Proof. Choose a compact codimension-0 submanifold $X'' \subset \mathring{X}$ such that $X' \subset \mathring{X}''$. Let us introduce an intermediate space $L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))$ defined as the completion of $i\Omega_{CC}^1(X) \oplus \Gamma(S_X^+)$ with respect to the norm $\|v\|_{2, X''} = \sqrt{\|\tilde{D}v\|_{L_1^2(X'')}^2 + \|v\|_{L_1^2(X)}^2}$, so that $L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))$ is a Hilbert space. Define

$$\widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{\mathfrak{s}}) = \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}}) \cap ((A_0, 0) + L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))),$$

the set of solutions to the Seiberg-Witten equations in the corresponding configuration space. Since \tilde{D} is elliptic and $\tilde{D}v = (\hat{D}v, 0)$, by [Theorem A.3](#) the restriction map

$$L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X)) \rightarrow L_2^2(i\Omega^1(X') \oplus \Gamma(S_{X'}^+))$$

is continuous linear (thus smooth). It thus suffices to prove that the moduli space $\widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{\mathfrak{s}})$ is a smooth submanifold of the configuration space $(A_0, 0) + L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))$ and that the identity map $\text{Id}_2 : \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{\mathfrak{s}}) \rightarrow \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{\mathfrak{s}})$ is well-defined, continuous and smooth.

Firstly, it is well-defined by [Lemma 5.1](#).

Secondly, we prove that $\widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{\mathfrak{s}})$ is a smooth submanifold of $(A_0, 0) + L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))$. Define $L_{1, X''}^2(i\Omega^+(X) \oplus \Gamma(S_X^+))$ to be the Hilbert space obtained as the completion of $i\Omega^+(X) \oplus \Gamma(S_X^+)$ with respect to the norm $\|v\|_{L^2(X) \cap L_1^2(X'')} = \sqrt{\|v\|_{L^2(X)}^2 + \|v\|_{L_1^2(X'')}^2}$. By the [Implicit Function Theorem](#), it suffices to prove the following Lemma.

Lemma 5.3. The differential

$$DSW_{(A, \Phi)} : L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X)) \rightarrow L_{1, X''}^2(i\Omega^+(X) \oplus \Gamma(S_X^+))$$

is surjective at each $(A, \Phi) \in \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{\mathfrak{s}})$.

Proof. We assumed that the operator $DSW_{(A,\Phi)}$ is surjective. Choose any $w \in L^2_{1,X''}(i\Omega^+(X) \oplus \Gamma(S_X^+))$ and pick a solution $v \in L^2_1(i\Omega^1_{CC}(X) \oplus \Gamma(S_X^+))$ to $DSW_{(A,\Phi)}(v) = w$. Then $\hat{D}v|_{X''} = -\hat{K}v|_{X''} + w|_{X''}$ is in $L^2_1(X'')$ since $w|_{X''} \in L^2_1(X'')$, $v|_{X''} \in L^2_1(X'')$ and $\hat{K}|_{X''}$ is a certain multiplication by $(A - A_0, \Phi)|_{X''}$, which is smooth by [Lemma 5.1](#). \blacksquare

Finally, it remains to prove that the identity map

$$\widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) \rightarrow \widetilde{\mathcal{M}}_{CC,X''}^\circ(X, \hat{s})$$

is smooth. We start by identifying the tangent spaces at $p = (A, \Phi)$; we have

$$\begin{aligned} T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) &= \ker DSW_p, \\ T_p \widetilde{\mathcal{M}}_{CC,X''}^\circ(X, \hat{s}) &= (\ker DSW_p) \cap L^2_{2,X''}(i\Omega^1_{CC}(X) \oplus \Gamma(S_X)). \end{aligned}$$

Lemma 5.4 (isometry of the tangent spaces). *The identity map $T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) \rightarrow T_p \widetilde{\mathcal{M}}_{CC,X''}^\circ(X, \hat{s})$ is well-defined, and an isometry.*

Proof. Take any $v \in T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s})$; in particular, $DSW_{(A,\Phi)}(v) = 0$. Well-definedness follows since $\hat{D}v|_{X''} = -\hat{K}v|_{X''}$ and as before, $\hat{K}|_{X''}$ is a multiplication by a smooth configuration on X'' . This also implies

$$\begin{aligned} \|v\|_{L^2_1(X)}^2 &\leq \|v\|_{2,X''}^2 = \|v\|_{L^2_1(X)}^2 + \|\hat{D}v\|_{L^2_1(X'')}^2 \\ &= \|v\|_{L^2_1(X)}^2 + \|\hat{K}v\|_{L^2_1(X'')}^2 \\ &\leq \|v\|_{L^2_1(X)}^2 + C_p \|v\|_{L^2_1(X'')}^2 \\ &\leq (1 + C_p) \|v\|_{L^2_1(X)}^2 \end{aligned}$$

which proves this map is an isometry. \blacksquare

While the $L^2(X)$ norm is not complete on $L^2_1(i\Omega^1_{CC}(X) \oplus \Gamma(S_X^+))$, the $L^2(X)$ -orthogonal complement H to $T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s})$ is a closed subspace of $L^2_1(i\Omega^1_{CC}(X) \oplus \Gamma(S_X^+))$ such that $H + T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) = L^2_1(i\Omega^1_{CC}(X) \oplus \Gamma(S_X^+))$ and $H \cap T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) = \{0\}$, thus by the open mapping theorem

$$L^2_1(i\Omega^1_{CC}(X) \oplus \Gamma(S_X^+)) = T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) \oplus H.$$

By the [Implicit Function Theorem](#) there is a neighborhood U of p such that the affine projection

$$U \cap \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) \rightarrow U \cap \left((A_0, 0) + T_p \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{s}) \right)$$

along H is a diffeomorphism. Similarly, the [Implicit Function Theorem](#) implies that there is a neighborhood V of p such that the affine projection

$$U \cap \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s}) \rightarrow U \cap \left((A_0, 0) + T_p \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s}) \right)$$

along $H' = H \cap L_{2, X''}^2(i\Omega_{CC}^1(X) \oplus \Gamma(S_X))$ is a diffeomorphism since H' is the $L^2(X)$ -orthogonal complement to $T_p \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s})$. Thus the identity map $\widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{s}) \rightarrow \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s})$ near p factors as

$$\begin{aligned} U \cap \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{s}) &\rightarrow U \cap \left((A_0, 0) + T_p \widetilde{\mathcal{M}}_{CC}^{\circ}(X, \hat{s}) \right) \\ &\xrightarrow{\text{id}} V \cap \left((A_0, 0) + T_p \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s}) \right) \\ &\rightarrow V \cap \widetilde{\mathcal{M}}_{CC, X''}^{\circ}(X, \hat{s}) \end{aligned}$$

where the middle identity map is smooth by [Lemma 5.4](#) and the two other maps are smooth since they are parametrizations coming from the [Implicit Function Theorem](#), as we just showed. \square

5.2 Proof of the gluing theorem

We are ready to prove the gluing theorem.

Let $X = X_1 \cup_Y X_2$ with Y connected and $b_1(\partial X_i) = 0$. Let \hat{s} be a spin^c structure and A_0 be a reference spin^c connection on X . Let restrictions of \hat{s} be the spin^c structures used to define configuration spaces and the restrictions of A_0 to be the reference connections. Denote $Y_1 = \partial X_1 \setminus Y$, $Y_2 = \partial X_2 \setminus Y$, $\hat{s}_i = \hat{s}|_{X_i}$. Fix gauge splittings s, s_1, s_2 and twistings τ, τ_1, τ_2 on X, X_1, X_2 .

Denote by $s_{\mathbb{Z}}, s_{1, \mathbb{Z}}, s_{2, \mathbb{Z}}$ the associated integral splittings given by [Proposition 3.29](#). We say they are compatible if $s_{\mathbb{Z}}$ corresponds to $(s_{1, \mathbb{Z}}, s_{2, \mathbb{Z}})$ under the following identification.

Proposition 5.5 (*integral splittings on a composite cobordism*). *There is a canonical identification between the set of integral splittings on X and the set of pairs of integral splittings on X_1 and X_2 .*

Proof. Choose an integral splitting s . Take any $a \in \mathcal{H}_D^1(X)$. Denote by \tilde{a}_i its restriction to X_i . For each i there is a unique $f_i \in L_2^2(\Omega^0(X_i))$ such that $\Delta f_i = 0$, $f_i|_Y = G_d t_Y^*(a)$, and $f_i|_{Y_i} = 0$. The resulting $a_i = \tilde{a}_i - df_i$ is in $\mathcal{H}_D^1(X_i)$, thus we get a map

$$R_{\mathcal{H}} : \mathcal{H}_D^1(X) \rightarrow \mathcal{H}_D^1(X_1) \times \mathcal{H}_D^1(X_2)$$

sending a to (a_1, a_2) . Note that $\mathcal{H}_D^1(X_i) \subset \Omega_{CC}^1(X_i)$ and therefore $R_{\mathcal{H}}$ coincides with doing the gauge fixing of [Lemma 3.16](#) on both components, i.e., $a \mapsto (\Pi_{CC}(a|_{X_1}), \Pi_{CC}(a|_{X_2}))$.

The cohomology class $[a] \in H^1(X; \mathbb{R})$ restricts to $([\tilde{a}_1], [\tilde{a}_2]) = ([a_1], [a_2]) \in H^1(X_1; \mathbb{R}) \times H^1(X_2; \mathbb{R})$. It follows that the composition

$$\begin{aligned} H^1(X_1; \mathbb{R}) \times H^1(X_2; \mathbb{R}) &\xrightarrow{\iota_*} H^1(X; \mathbb{R}) \xrightarrow{s^H} \mathcal{H}_D^1(X) \rightarrow \\ &\xrightarrow{R_{\mathcal{H}}} \mathcal{H}_D^1(X_1) \times \mathcal{H}_D^1(X_2) \rightarrow H^1(X_1; \mathbb{R}) \times H^1(X_2; \mathbb{R}) \end{aligned}$$

is the identity, where ι_* is the inverse of the restriction map $H^1(X; \mathbb{R}) \rightarrow H^1(X_1; \mathbb{R}) \times H^1(X_2; \mathbb{R})$ (invertible by Mayer-Vietoris). We can thus choose (s_1^H, s_2^H) to be the composition $R_{\mathcal{H}} \circ s^H \circ \iota_*$. Since s^H was integral, thus s_i^H are integral and by [Proposition 3.23](#) there exist unique integral splittings s_1, s_2 inducing s_1^H, s_2^H .

On the other hand, given integral splittings s_1, s_2 on X_1, X_2 , we can choose s^H to be $(R_{\mathcal{H}})^{-1} \circ (s_1^H, s_2^H) \circ (\iota_*)^{-1}$ and by [Proposition 3.23](#) there is a unique integral splitting s inducing s^H . \square

[Proposition 4.8](#) guarantees that whenever $\partial X \neq \emptyset$, the moduli $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$ is a smooth Hilbert manifold, and the same follows for $\mathcal{M}_{s_i}^\circ(X_i, \hat{\mathfrak{s}})$. We want to include the case when X is a closed manifold, when it is well-known that to achieve surjectivity one, in general, needs to perturb the metric on X or perturb the Seiberg-Witten equations. Therefore, we assume that for any $(A, \Phi) \in \text{SW}^{-1}(0)$ the operator $D\text{SW}_{(A, \Phi)}$ is onto.

Remark 5.6. The careful reader may notice that we do not assume any transversality of the maps $R_{\tau_i, Y}$ and thus the fiber product may not *a priori* be a manifold. That it is a manifold follows from the proof of the theorem. What is not proven here, but may be useful in other contexts, is that the transversality is indeed equivalent to $D\text{SW}_{(A, \Phi)}$ being surjective for all $(A, \Phi) \in \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$.

Gluing Theorem. *Assume $s_{\mathbb{Z}}$ and $(s_{1, \mathbb{Z}}, s_{2, \mathbb{Z}})$ are compatible. Then there is an S^1 -equivariant diffeomorphism $F : \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \mathcal{M}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times_Y \mathcal{M}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2)$ such that R_τ is S^1 -equivariantly homotopic to $(R_{\tau_1} \times_Y R_{\tau_2}) \circ F$.*

Proof. By [Corollary 3.33](#) we can assume, without loss of generality, that $s = s_{\mathbb{Z}}$ and $s_i = s_{i, \mathbb{Z}}$, as well as $\tau \equiv 1$ and $\tau_i \equiv 1$. We thus drop τ 's from the notation entirely.

The plan is as follows. We will construct a map

$$F : \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow \widetilde{\mathcal{M}}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times \widetilde{\mathcal{M}}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2),$$

and a homotopy

$$H : \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}}) \rightarrow (\mathcal{C}_{cC}(Y_1, \mathfrak{s})) \times (\mathcal{C}_{cC}(Y_2, \mathfrak{s}))$$

between R and $(R_{Y_1} \times R_{Y_2}) \circ F$. Then we will prove that F intertwines the actions of $\mathcal{G}_s^{h, \circ}(X)$ and $\mathcal{G}_{s_1}^{h, \circ}(X_1) \times \mathcal{G}_{s_2}^{h, \circ}(X_2)$, and that H is $\mathcal{G}_s^{h, \circ}(X)$ -invariant; thus, both F and H

descend to $\mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$. Furthermore, we will prove F and H are continuous and smooth, and that F has image in the fiber product. Finally, we will show that F is a smooth immersion onto $\widetilde{\mathcal{M}}_{s_1}^\circ(X_1, \hat{\mathfrak{s}}_1) \times_Y \widetilde{\mathcal{M}}_{s_2}^\circ(X_2, \hat{\mathfrak{s}}_2)$. The S^1 -equivariance of F and H will be apparent from the construction.

Step 1. We start by constructing F , proving its smoothness and that its image lies in the fiber product. Take $(A, \Phi) \in \mathcal{M}_s^\circ(X, \hat{\mathfrak{s}})$. We would like to simply restrict it to the components X_i and then put into split Coulomb slice. By [Lemma 3.16](#) modulo S^1 there a unique way of doing that using a contractible gauge transformation. Here we make a different choice than in [Lemma 3.16](#), requiring $\int_Y f = 0$ instead of $\int_X f = 0$. Precisely, for $a \in L_1^2(i\Omega^1(X_i))$ choose $\tilde{u}_a^s = e^{f_a}$ such that $f_a \in L_2^2(i\Omega^0(X_i))$, $\int_Y f_a = 0$ and $a - (\tilde{u}_a^s)^{-1} d\tilde{u}_a^s \in L_1^2(i\Omega_s^1(X_i))$. Define

$$F(A, \Phi) = \left(\tilde{u}_{(A-A_0)|_{X_1}}^s (A, \Phi)|_{X_1}, \tilde{u}_{(A-A_0)|_{X_2}}^s (A, \Phi)|_{X_2} \right).$$

Since $A - A_0$ was in double Coulomb slice, thus $(A - A_0)|_{X_1}$ (resp. $(A - A_0)|_{X_2}$) is already coclosed on X_1 (resp. X_2) and on Y_1 (resp. Y_2), moreover $(A - A_0)|_{X_1}$ and $(A - A_0)|_{X_2}$ agree on Y . Denoting the restriction to Y by $b_{(A, \Phi)} = \iota_Y^*(A - A_0)$, [Theorem 5.2](#) together with [Theorem A.2](#) imply that $b_{(A, \Phi)}$ is smooth and depends smoothly on $(A, \Phi) \in \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}})$ as an element of $L_{3/2}^2$. Notice that $f_i = f_{(A-A_0)|_{X_i}}$ can be decomposed as $f_i = f_i^\partial + f_i^s$, where f_i^∂ are the unique solutions to

$$f_i^\partial|_{Y_i} = 0, \quad f_i|_Y = g, \quad \Delta f_i = 0, \tag{18}$$

where $g = G_d \Pi_d b_{(A, \Phi)}$ depends smoothly on (A, Φ) as an element of $L_{5/2}^2$. Moreover, the map $g \mapsto f_i^\partial$ is linear and continuous as a map $L_{s+1/2}^2 \rightarrow L_{s+1}^2$ for $s \geq 0$, so f_i^∂ depend smoothly on (A, Φ) as elements of L_3^2 . Furthermore, since $a|_{X_i} - df_i^\partial \in L_1^2(i\Omega_{CC}^1(X_i))$, f_i^s are the unique elements of $\mathcal{H}_D^1(X_i)$ such that $a|_{X_i} - df_i^\partial - df_i^s \in L_1^2(i\Omega_s^1(X_i))$ and $f_i^s|_Y = 0$. By [Remark 3.17](#), f_i^s depend continuously on $a|_{X_i} - df_i^\partial \in L_1^2$. Which proves that $f_i \in L_3^2$ depend continuously and linearly on $g \in L_{5/2}^2$, thus depend smoothly on g , so they depend smoothly on (A, Φ) .

This establishes the well-definedness and smoothness of F . That its image lies in the fiber product follows directly from the construction.

Step 2. We proceed to constructing H . By (18), the functions f_i are locally constant on Y_i . We also have

$$R_{Y_i}(F(A, \Phi)) = e^{f_i|_{Y_i}} \cdot R_{Y_i}(A, \Phi)$$

by the construction of F . Thus we can define

$$H(A, \Phi, t) = \left(e^{t\tilde{f}_1|_{Y_1}}, e^{t\tilde{f}_2|_{Y_2}} \right) R(A, \Phi)$$

which at $t = 0$ coincides with $R(A, \Phi)$ and at $t = 1$ coincides with $(R_{Y_1}, R_{Y_2}) \circ F(A, \Phi)$.

Step 3. We now investigate the equivariance of F under the actions of gauge groups. Let $u \in \mathcal{G}_s^{h,\circ}(X)$. From [Lemma 3.16](#) it follows that there is exactly one contractible $\tilde{u}_i = e^{\tilde{f}_i} \in \mathcal{G}^e(X_i)$ which puts $-u^{-1}du$ into $L_1^2(i\Omega_s^1(X_i))$ with $\int_Y \tilde{f}_i = 0$. Equivalently, this is the unique $\tilde{u}_i = e^{\tilde{f}_i} \in \mathcal{G}^e(X_i)$ with $\int_Y \tilde{f}_i = 0$ such that $\tilde{u}_i u|_{X_i} \in \mathcal{G}_{s_i}^{h,\circ}(X_i)$. Define $u_i = \tilde{u}_i u|_{X_i}$. Since \tilde{u}_i is contractible, thus $[u_{X_i}^*(u^{-1}du)] = [u_i^{-1}du_i]$ in $H^1(X_i; 2\pi i\mathbb{Z})$. Therefore the map $u \mapsto (u_1, u_2)$ provides the canonical isomorphism

$$\mathcal{G}_s^{h,\circ}(X) \simeq \mathcal{G}_{s_1}^{h,\circ}(X_1) \times \mathcal{G}_{s_2}^{h,\circ}(X_2) \quad (19)$$

which agrees with the isomorphism coming from the Mayer-Vietoris sequence $H^1(X; \mathbb{Z}) \simeq H^1(X_1; \mathbb{Z}) \times H^1(X_2; \mathbb{Z})$. From the construction of $F(A, \Phi)$ it follows that $(u|_{X_1}^{-1}, u|_{X_2}^{-1})F(u(A, \Phi))$ differs from $F(A, \Phi)$ exactly by the factor of $(e^{\tilde{f}_1}, e^{\tilde{f}_2})$. Thus $(u_1^{-1}, u_2^{-1})F(u(A, \Phi)) = F(A, \Phi)$. This proves that F commutes with the gauge group action as identified in [\(19\)](#).

Step 4. We prove the invariance of H under $\mathcal{G}_s^{h,\circ}(X)$. By the construction of H , and since $u|_{\partial X} = 1$ for $u \in \mathcal{G}_s^{h,\circ}(X)$, we have $H(u(A, \Phi), t) = (e^{t\tilde{f}_1|_{Y_1}}, e^{t\tilde{f}_2|_{Y_2}})H(A, \Phi, t)$ where \tilde{f}_i are as in the previous paragraph. Since $R_{\mathcal{H}}(\mathcal{H}_D^1(X)) \in (\text{im } s_1^H) \times (\text{im } s_2^H)$ we get that $R_{\mathcal{H}}(u^{-1}du)$ is already in the split Coulomb slice on X_1 and X_2 . Moreover, $a_i = \iota_{X_i}^*(u^{-1}du)$ is coclosed on X_i and Y_i . Since there are functions \hat{f}_i satisfying

$$\begin{aligned} \Delta \hat{f}_1 &= 0, & \hat{f}_1|_{Y_1} &= 0, & \hat{f}_1|_Y &= G_d \iota_Y^* a, \\ \Delta \hat{f}_2 &= 0, & \hat{f}_2|_{Y_2} &= 0, & \hat{f}_2|_Y &= G_d \iota_Y^* a, \end{aligned}$$

the uniqueness of \tilde{f}_i implies $\tilde{f}_i = \hat{f}_i$ and therefore $\tilde{f}_i|_{Y_i} = 0$. Thus $H(u(A, \Phi), t) = H(A, \Phi, t)$, as wished.

Step 5. We show that F is bijective onto the fiber product, following the argument in [\[Lip08\]](#). Let $(A_i, \Phi_i) \in \widetilde{\mathcal{M}}_{s_i}^\circ(X_i, \hat{s})$ such that $R_Y(A_1, \Phi_1) = R_Y(A_2, \Phi_2)$. These would give a configuration on X if the normal components of connections A_1 and A_2 agreed on Y . Let $h_1 dt$ and $h_2 dt$ be the dt -components of $(A_1 - A_0)|_Y$ and $(A_2 - A_0)|_Y$. We want to find harmonic functions $f_i \in L_2^2(X_i; i\mathbb{R})$ such that

$$\begin{aligned} f_1|_Y &= f_2|_Y, \\ \partial_t f_1|_Y + h_1 &= \partial_t f_2|_Y + h_2, \end{aligned}$$

$$f_1|_{Y_1} = 0, f_2|_{Y_2} = 0.$$

Take a tubular neighborhood $[-\varepsilon, \varepsilon] \times Y \subset X$ of Y . Let $\{\phi_\lambda\}_\lambda$ be an eigenbasis for Δ_Y and write $h_2 - h_1 = \sum_\lambda c_\lambda \phi_\lambda$. Since $h_2 - h_1 \in L^2_{1/2}$, therefore $\sum_\lambda \lambda^{1/2} |c_\lambda|^2 < \infty$ and thus the following are well-defined as elements of $L^2_2([-\varepsilon, \varepsilon] \times Y; i\mathbb{R})$:

$$\begin{aligned} g_1 &= \frac{1}{2} \sum c_\lambda \lambda^{-1/2} e^{\lambda^{1/2} t} \varphi_\lambda, \\ g_2 &= \frac{1}{2} \sum c_\lambda \lambda^{-1/2} e^{-\lambda^{1/2} t} \varphi_\lambda, \end{aligned}$$

which satisfies $\partial_t g_1|_Y - \partial_t g_2|_Y = h_1 - h_0$. Let $\rho \in C^\infty(X; \mathbb{R})$ be a bump function supported in $[-\varepsilon, \varepsilon] \times Y$ which is identically 1 in a neighborhood of Y . The configurations $e^{\rho g_1}(A_1, \Phi_1)$ and $e^{\rho g_2}(A_2, \Phi_2)$ patch to give a L^2_1 configuration (A', Φ') , but this is not necessarily in the Coulomb slice because $\rho f'_i$ are not necessarily harmonic. Take $f \in L^2_2(X; i\mathbb{R})$ such that

$$f|_{Y_1} = 0, \quad f|_{Y_2} = 0, \quad \Delta f = -\Delta(\rho g_1) - \Delta(\rho g_2).$$

Denote $(A'', \Phi'') = e^f(A', \Phi') \in \widetilde{\mathcal{M}}_{CC}^\circ(X, \hat{\mathfrak{s}})$. Finally, by [Remark 3.17](#) we can continuously deform (A'', Φ'') to a configuration (A, Φ) such that $F(A, \Phi) = ((A_1, \Phi_1), (A_2, \Phi_2))$.

Step 6. We need to prove that F^{-1} constructed in the previous step is continuous. Notice (g_1, g_2) as elements of L^2_2 depend continuously on $h_1 - h_0 \in L^2_{1/2}$ which in turn depends continuously on $A_1 - A_0$ and $A_2 - A_0 \in L^2_{1/2}$. Moreover, $f \in L^2_3$ depends continuously on $(g_1, g_2) \in L^2_2$. If the multiplication $L^2_2 \times L^2_1 \rightarrow L^2_1$ was continuous on 4-manifolds then we would have shown that the map $((A_1, \Phi_1), (A_2, \Phi_2)) \rightarrow (A, \Phi)$ which we constructed is continuous. Since it is not true, we need to show that $e^{g_i} \Phi_i \in L^2_1$ depends continuously on the initial configurations.

We will prove it depends continuously on $h_2 - h_1 \in L^2_{1/2}$ and $\Phi_i \in L^2_1$. Let (A'_1, Φ'_1) and (A'_2, Φ'_2) be another choice of configurations, and denote the corresponding harmonic functions on $[-\varepsilon, 0] \times Y$ and $[0, \varepsilon] \times Y$ by g'_1, g'_2 . Then

$$\begin{aligned} \|e^{g_1} \Phi_1 - e^{g'_1} \Phi'_1\|_{L^2_1} &\leq \|e^{g'_1}(\Phi_1 - \Phi'_1)\|_{L^2_1} + \|(e^{g_1} - e^{g'_1})\Phi_1\|_{L^2_1} \\ &\leq C(\|e^{g'_1}\|_{L^\infty} + \|e^{g'_1}\|_{L^2_2}) \|\Phi_1 - \Phi'_1\|_{L^2_1} \\ &\quad + C\|e^{g_1} - e^{g'_1}\|_{L^2_2} \|\Phi_1\|_{L^2_1} \\ &\quad + C\|e^{g_1} - e^{g'_1}\|_{L^\infty([-\varepsilon, -\delta] \times Y)} \|\Phi_1\|_{L^2_1([-\varepsilon, -\delta] \times Y)} \\ &\quad + C\|e^{g_1} - e^{g'_1}\|_{L^\infty([-\delta, 0] \times Y)} \|\Phi_1\|_{L^2_1([-\delta, 0] \times Y)} \end{aligned}$$

for any δ . One can choose δ to have $\|\Phi_1\|_{L^2_1([-\delta, 0] \times Y)}$ as small as one wants while

$\|e^{g_1} - e^{g'_1}\|_{L^\infty([- \delta, 0] \times Y)} \leq 2$. Moreover, we have

$$\|e^{g_1} - e^{g'_1}\|_{L^\infty([- \varepsilon, -\delta] \times Y)} \leq \|g_1 - g'_1\|_{L^\infty([- \varepsilon, -\delta] \times Y)} \leq C\|(h_1 - h_2) - (h'_1 - h'_2)\|_{L^2_{1/2}}$$

via a direct computation (or by interior regularity estimates following from [Theorem A.3](#)).

This finishes the proof that the inverse map is continuous.

Step 7. Finally, we prove that the differential of F is invertible. Assume this is not the case, so that there exists $(A, \Phi) \in \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}})$ such that for any $\varepsilon > 0$ there is $(A', \Phi') \in \widetilde{\mathcal{M}}_s^\circ(X, \hat{\mathfrak{s}})$ such that $0 < D = \|F(A, \Phi) - F(A', \Phi')\|_{L^2_1} < 1$ and $\|(A - A', \Phi - \Phi')\|_{L^2_1} \leq D\varepsilon$. We get that $\|g - g'\|_{L^2_{5/2}} \leq C_{5/2}D\varepsilon$ and thus

$$\begin{aligned} \|f_i - f'_i\|_{L^2_3} &\leq C\|g - g'\|_{L^2_{5/2}} \\ &\leq CC_{5/2}D\varepsilon. \end{aligned}$$

From this and [Theorem A.1](#) it follows that $D = \|F(A, \Phi) - F(A', \Phi')\|_{L^2_1} \leq C''\|(A - A', \Phi - \Phi')\|_{L^2_1} \leq C''D\varepsilon$ for some C'' depending on $\|\Phi\|_{L^2_1}$. Choosing $\varepsilon = \frac{1}{1+C''}$ gives the desired contradiction. \square

A Appendix

This section presents some standard analytical results which are used repeatedly in the article. We recall the Gårding inequality, Sobolev multiplication and trace theorems and the Implicit Function Theorem.

Theorem A.1 (Sobolev multiplication theorem). *Let M be a manifold with compact boundary and cylindrical ends.*

Assume $k, l \geq m$ and $1/p + 1/q \geq 1/r$ for $p, q, r \in (1, \infty)$. Then the multiplication

$$L^p_k(M) \times L^q_l(M) \rightarrow L^r_m(M)$$

is continuous if any of these hold:

- $(k - n/p) + (l - n/q) \geq m - n/r$ and both $k - n/p < 0, l - n/q < 0$,
- $\min(k - n/p, l - n/q) \geq m - n/r$ and either $k - n/p > 0$ or $l - n/q > 0$,
- $\min(k - n/p, l - n/q) > m - n/r$ and either $k - n/p = 0$ or $l - n/q = 0$.

What is more, whenever it is continuous, it restricts to a compact map on $\{f\} \times L^q_l(M) \rightarrow L^r_m(M)$ provided that $l > m$ and $l - n/q > m - n/r$.

Proof. We can reduce to the case of a manifold without boundary by considering a double of M , or by attaching cylindrical ends along ∂M .

To obtain continuity, combine [Pal68, Theorem 9.6] for compact manifolds together with [KM07, Theorem 13.2.2] for an infinite cylinder.

For the compactness statement on a compact manifold, for each sequence $g_i \xrightarrow{L_i^q(M)} g$ take any sequence $f_j \in C^\infty(M)$ such that $\|f - f_j\|_{L_k^p(M)} \leq \frac{1}{2^j}$. By continuity of multiplication there is a constant C such that

$$\|(f - f_j)g_i\|_{L_m^r(M)} \leq C\|f - f_j\|_{L_k^p(M)} \leq \frac{C}{2^j}.$$

Set $a_{0,i} = i$ and define inductively $a_{j,i}$ in the following manner. Take $a_{j,i} = a_{j-1,i}$ whenever $i < j$. Since $f_j \in C^\infty(M)$, the sequence $f_j g_i$ is bounded in $L_l^q(M)$ and by Sobolev embedding is precompact in $L_m^r(M)$. Thus, we can choose $(a_{j,i})_i$ to be a subsequence of $(a_{j-1,i})_i$ satisfying

$$\|f_j g_{a_{j,j}} - f_j g_{a_{j,i}}\|_{L_m^r(M)} \leq \frac{1}{2^j}$$

for any $i \geq j$. We finally get, for $i \geq 0$,

$$\begin{aligned} \|f g_{a_{j,j}} - f g_{a_{j+i,j+i}}\|_{L_m^r(M)} &\leq \|(f - f_j)(g_{a_{j,j}} - g_{a_{j+i,j+i}})\|_{L_m^r(M)} \\ &\quad + \|f_j(g_{a_{j,j}} - g_{a_{j+i,j+i}})\|_{L_m^r(M)} \\ &\leq \frac{2C}{2^j} + \frac{1}{2^j} \end{aligned}$$

making $f g_{a_{j,j}}$ a Cauchy sequence in $L_m^r(M)$, as wished.

Compactness for a cylinder is proved in [KM07, Theorem 13.2.2]. \square

Theorem A.2 (Sobolev trace theorem). *Suppose $N \subset M$ is a closed compact $n-j$ -dimensional smooth submanifold, $1 \leq p < \infty$, and $k, l \geq 0$ with $k - j/p \geq l > 0$. Then the restriction map extends to continuous*

$$L_k^p(M) \rightarrow L_l^p(N).$$

Proof. See [Pal68, Theorem 9.3]. \square

We will make use of the Implicit Function Theorem in the following form (cf. [Lan93]).

Implicit Function Theorem. Suppose A, B are Hilbert spaces and $\mathcal{F} : A \rightarrow B$ is a smooth map such that the derivative $D_p\mathcal{F}$ at p is surjective and that its kernel splits with C as a complementary subspace, $A = \ker(D_p\mathcal{F}) \oplus C$. Let $\Pi : A \rightarrow \ker(D_p\mathcal{F})$ denote the projection onto the kernel along C . Then there are open neighborhoods $U \subset A$ of p , $V \subset B$ of $\mathcal{F}(p)$ and $W \subset \ker(D_p\mathcal{F})$ of 0 and a smooth diffeomorphism $\mathcal{G} : V \times W \rightarrow U$ such that

$$V \times W \xrightarrow{\mathcal{G}} U \xrightarrow{(\mathcal{F}, \Pi)} B \oplus \ker(D_p\mathcal{F})$$

is the identity map (i.e., $\mathcal{G} = \mathcal{F}^{-1}$).

In particular, the projection

$$\Pi : \mathcal{F}^{-1}(0) \cap U \rightarrow \ker(D_p\mathcal{F})$$

along C is a local diffeomorphism.

We will also utilize the Gårding inequality.

Theorem A.3 (Gårding inequality). Let D be a first-order elliptic operator with smooth coefficients on a compact manifold M (possibly with boundary) and $M' \subset M$ be open with compact closure. Then there is a constant C such that for any $\gamma \in L_{k+1}^p(M)$ we have

$$\|\gamma\|_{L_{k+1}^p(M')} \leq C(\|D\gamma\|_{L_k^p(M)} + \|\gamma\|_{L_k^p(M)}).$$

Proof. This follows from [Shu92, Appendix 1, Lemma 1.4] by extending D to the cylindrical-end manifold $M^* = M \cup (\partial M) \times [0, \infty)$. Indeed, taking a smooth bump function ρ such that $\rho|_{M'} = 1$ and $\rho|_{M^* \setminus M} = 0$ we get

$$\begin{aligned} \|\gamma\|_{L_{k+1}^p(M')} &\leq \|\rho\gamma\|_{L_{k+1}^p(M^*)} \\ &\leq C(\|D(\rho\gamma)\|_{L_k^p(M^*)} + \|\rho\gamma\|_{L_k^p(M^*)}) \\ &\leq CC'_\gamma(\|D(\gamma)\|_{L_k^p(M)} + \|\gamma\|_{L_k^p(M)}) \end{aligned}$$

where the middle inequality follows from [Shu92]. \square

Remark A.4. The author's understanding is that without the use of twistings there is (in general) no choice of s, s_1, s_2 making F commute with the restriction maps on the nose. This problem does not show up in the construction of monopole Floer homology since there one quotients the moduli and configuration spaces by S^1 after blowing up.

References

- [Ati88] Michael Atiyah. New invariants of 3- and 4-dimensional manifolds. The mathematical heritage of Hermann Weyl (Durham, NC, 1987). Proc. Sympos. Pure Math., vol. 48. Amer. Math. Soc., Providence, RI., pages 285–299. 1988.

- [AB83] M. F. Atiyah and R. Bott. The Yang-Mills equations over Riemann surfaces. Philos. Trans. Roy. Soc. London Ser. A 308 (1505):523–615, 1983.
- [AB21] Mohammed Abouzaid and Andrew J. Blumberg. Arnold Conjecture and Morava K-theory, 2021, preprint.
- [CJS95] Ralph L. Cohen, John D. S. Jones, and Graeme B. Segal. Floer’s infinite-dimensional Morse theory and homotopy theory. The Floer memorial volume. *Progr. Math.*, vol. 133. Birkhäuser, Basel., pages 297–325. 1995.
- [Don83] Simon K. Donaldson. An application of gauge theory to four-dimensional topology. J. Differential Geom. 18 (2):279–315, 1983.
- [DS19] Aliakbar Daemi and Christopher Scaduto. Equivariant aspects of singular instanton Floer homology, 2019.
- [Lin18] Francesco Lin. A Morse-Bott approach to monopole Floer homology and the triangulation conjecture. Mem. Amer. Math. Soc. 255 (1221):v+162, 2018.
- [Flo87] Andreas Floer. Morse theory for fixed points of symplectic diffeomorphisms. Bull. Amer. Math. Soc. (N.S.) 16 (2):279–281, 1987.
- [Flo88] ———. Morse theory for Lagrangian intersections. J. Differential Geom. 28 (3):513–547, 1988.
- [Kha] Tirasan Khandhawit. A new gauge slice for the relative Bauer-Furuta invariants.
- [KLS18] Tirasan Khandhawit, Jianfeng Lin, and Hirofumi Sasahira. Unfolded Seiberg-Witten Floer spectra, I: Definition and invariance. Geom. Topol. 22 (4):2027–2114, 2018.
- [KM07] Peter Kronheimer and Tomasz Mrowka. Monopoles and three-manifolds. New Mathematical Monographs, vol. 10. Cambridge University Press, Cambridge, 2007.
- [KM97] P. B. Kronheimer and T. S. Mrowka. Monopoles and contact structures. Invent. Math. 130 (2):209–255, 1997.
- [KM11] Peter Kronheimer and Tomasz Mrowka. Khovanov homology is an unknot-detector. Publ. Math. Inst. Hautes Études Sci. 113:97–208, 2011.
- [Lan93] Serge Lang. Real and functional analysis. Graduate Texts in Mathematics, vol. 142. Springer-Verlag, New York, 3rd ed., 1993.
- [Lin15] Jianfeng Lin. Pin(2)-equivariant KO-theory and intersection forms of spin 4-manifolds. Algebr. Geom. Topol. 15 (2):863–902, 2015.
- [Lip08] Maksim Lipyanskiy, A semi-infinite cycle construction of Floer homology, 2008. Thesis (Ph.D.)–Massachusetts Institute of Technology.
- [LM89] H. Blaine Lawson Jr. and Marie-Louise Michelsohn. Spin geometry. Princeton Mathematical Series, vol. 38. Princeton University Press, Princeton, NJ, 1989.
- [Man94] Niculae Mandache. Some remarks concerning unique continuation for the Dirac operator. Lett. Math. Phys. 31 (2):85–92, 1994.
- [Man13] Ciprian Manolescu. The Conley index, gauge theory, and triangulations. J. Fixed Point Theory Appl. 13 (2):431–457, 2013.

- [Man14] _____ . On the intersection forms of spin four-manifolds with boundary. Math. Ann. 359 (3-4):695–728, 2014.
- [Pal68] Richard S. Palais. Foundations of global non-linear analysis. W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [Sch93] Matthias Schwarz. Morse homology. Progress in Mathematics, vol. 111. Birkhäuser Verlag, Basel, 1993.
- [Shu92] Mikhail A. Shubin. Spectral theory of elliptic operators on noncompact manifolds. Astérisque 207:5, 35–108, 1992. Méthodes semi-classiques, Vol. 1 (Nantes, 1991).
- [Mil19] Stephen Michael Miller. Equivariant Instanton Homology, 2019. Thesis (Ph.D.)–University of California, Los Angeles.
- [Suw20] Piotr Suwara, Semi-infinite Homology of Floer spaces, 2020. Thesis (Ph.D.)–Massachusetts Institute of Technology.
- [Wit82] Edward Witten. Supersymmetry and Morse theory. J. Differential Geometry 17 (4):661–692 (1983), 1982.
- [Wol92] T. H. Wolff. A property of measures in \mathbf{R}^N and an application to unique continuation. Geom. Funct. Anal. 2 (2):225–284, 1992.