Fock-space geometry and strong correlations in many-body localized systems

Christian P. Chen and Henning Schomerus Department of Physics, Lancaster University, Lancaster, LA1 4YB, United Kingdom (Dated: March 1, 2022)

We adopt a geometric perspective on Fock space to provide two complementary insights into the eigenstates in many-body-localized fermionic systems. On the one hand, individual many-body-localized eigenstates are well approximated by a Slater determinant of single-particle orbitals. On the other hand, the orbitals of different eigenstates in a given system display a varying, and generally imperfect, degree of compatibility, as we quantify by a measure based on the projectors onto the corresponding single-particle subspaces. We study this incompatibility between states of fixed and differing particle number, as well as inside and outside the many-body-localized regime. This gives detailed insights into the emergence and strongly correlated nature of quasiparticle-like excitations in many-body localized systems, revealing intricate correlations between states of different particle number down to the level of individual realizations.

Many-body localized (MBL) systems constitute a broad class of closed quantum systems that fail to equilibrate under their own dynamics [1–6]. This behaviour manifests itself as a violation of the eigenstate thermalization hypothesis [7, 8], a hypothesis that more conventional, ergodic systems obey. A pressing question [9] is whether the MBL phase can be explored equally via features of the wavefunctions, such as in the area law of entanglement [4, 5], and via energy-level statistics [10], as well as how firmly these approaches tie into key theoretical concepts. In particular, wavefunction-based characterizations so far approach each eigenstate individually, while energy-level statistics are generally based on energy-level spacings, and thereby involve the interplay of several states [5, 11, 12]. This leaves a conceptual gap between the most reliable characterizations of MBL and the most influential theoretical framework, the concept of robust emergent integrability [13–16], which ties the MBL phase transition to the emergence of an extensive set of local conserved quantities that are commonly referred to as l-bits [13–19]. These l-bits can be seen as highly structured constraints on the complete set of eigenstates of any given system, both energetically as well as in terms of their wavefunctions. In principle, an allencompassing description of MBL should therefore also allow to develop expectations for the joint characteristics of the wavefunctions of a given system.

In this work, we set out a framework to formulate and verify such expectations for the paradigmatic case of fermionic systems, and use this to uncover their strongly correlated nature from a transparent geometric perspective. Our framework rests on the observation that for these systems, the MBL wavefunctions are closely approximated by Slater determinants, a realization that previously proved useful to lift Fock-space localization into real space by means of the one-particle density matrix [20–30]. Here, we propose to capture this structure by considering the span of single-particle orbitals whose Slater determinant provides the largest overlap to a given eigenstate—known in quantum chemistry as Brueckner

orbitals [31–33]—and show that this indeed provides a natural avenue to study the unexplored geometric interrelations between the wavefunctions of a given system.

Our key insights are the following: While individual MBL wavefunctions are indeed well approximated by a single Slater determinant, when comparing different eigenstates of the same system the underlying orbitals are generally mutually incompatible with each other. We define and quantify this mutual incompatibility precisely using a natural geometric measure, based on the projection operators of the occupied Brueckner orbitals from different eigenstates. We then establish the finite extent of this incompatibility in a model system, use randommatrix theory as a benchmark to show that it scales systematically with system size, and develop this picture further by varying the disorder and interaction strength. Finally, we relate the observed incompatibility to features of the aforementioned l-bit operators, whose systematical dressing is revealed by geometric correlations of states with different particle numbers. The proposed framework therefore provides a natural perspective on the emergence of MBL that complements traditional insights, revealing strongly correlated features that can be precisely quantified.

Brueckner orbitals and orbital incompatibility. In a sector of N fermions, Brueckner orbitals are the name given to a set $S_m = \{\chi_n^{(m)}\}_{n=1}^N$ of single-particle orbitals which maximize the overlap

$$\mathcal{I}_m \equiv |\langle \psi_m | S_m \rangle|^2 \tag{1}$$

between the Slater determinant $|S_m\rangle$ constructed from these orbitals and a given many-body eigenstate $|\psi_m\rangle$ of the system. These orbitals enjoy a wide range of applications in quantum-chemical approaches to interacting quantum systems [31–34], and their use can also be extended to a systematic approximation by a series of Slater determinants [35]. For this work, an important feature of these orbitals is the fact that the Slater determinant $|S_m\rangle$ only depends on the subspace they span, so all their relevant features are uniquely captured by the projector

 P_m onto this subspace. In general, the Brueckner orbitals are distinct from the well-known natural orbitals $\varphi_{\alpha}^{(m)}$, which in a discrete basis of single-particle states with fermionic annihilation operators c_i are obtained by diagonalizing the one-particle density matrix (OPDM) $\rho_{ij}^{(m)} \equiv \langle \psi_m | c_i^{\dagger} c_j | \psi_m \rangle$, where the eigenvalues $n_{\alpha}^{(m)}$ represent the occupations of these natural orbitals. When the overlap \mathcal{I}_m is moderate or small, both types of orbitals can be very different to each other, to the extent that constructing the Brueckner orbitals from the natural orbitals can be a numerically unfavourable task [34, 35]. However, the situation changes when the overlap \mathcal{I}_m is large, and for $\mathcal{I}_m \to 1$, $\rho^{(m)} \to P_m$. In turn, from the perspective of the OPDM this limit is reached when the occupations $n_{\alpha}^{(m)} \to 0, 1$. This indeed applies well to individual eigenstates inside the many-body-localized regime [20–22], and we will expand on this observation further below.

Turning our attention to the relation between different eigenstates, our key premise is the following observation: Even when two (or perhaps all) eigenstates of an interacting system are exact Slater determinants, this does not quarantee that there is a common single-particle basis from which one can form the Brueckner orbitals of these eigenstates. In particular, such a single-particle basis is not guaranteed by the mere requirement that the corresponding many-body Slater determinants are orthogonal to each other. This is in contrast to noninteracting systems, where the single-particle eigenstates provide a joint set from which one can choose the Brueckner orbitals. Indeed, it is easy to quantify and detect this orbital incompatibility, as it can be established already on the level of pairs of many-body eigenstates $|\psi_l\rangle$ and $|\psi_m\rangle$. When the orbitals of both states are compatible with each other, their projectors P_l and P_m commute. We therefore introduce the following incompatibility measure,

$$C_{ml} \equiv \sqrt{\operatorname{tr}\left[P_l, P_m\right]\left[P_m, P_l\right]} = \sqrt{2\operatorname{tr}\left(P_l P_m - (P_l P_m)^2\right)},$$
(2)

which mathematically corresponds to the Frobenius norm of the commutator of both projectors. As the projectors P_m characterize subspaces in the single-particle Hilbert space, this provides us with a simple geometric picture of the interrelation between different wavefunctions of a given MBL system.

Application. To illustrate the geometric framework described above in a specific setting, we place it into the context of a paradigmatic fermionic model exhibiting many-body localization. This is a system consisting of N locally interacting spinless fermions confined to a one-dimensional lattice with L sites and subjected to a disordered onsite potential, where the Hamiltonian is




FIG. 1. Geometric features of eigenstates in the many-body localized system (3) (half-filled system of size L=14, with V=1.5,~W=8; total number of states denoted as \mathcal{N}). In the left column, states are ordered by descending overlap \mathcal{I}_m with a single Slater determinant, while in the right column they are ordered by ascending energy E_m ; see panels (b,c,f,g) for the averaged behaviour of these quantities. The observed large overlaps \mathcal{I}_m show that a large fraction of states are well approximated by a single Slater determinant. Panels (a,d) show the averaged incompatibility \mathcal{C}_{ml} defined in Eq. (2), which quantifies the geometric interrelation of the states. This incompatibility is appreciable even for states with $\mathcal{I}_m \approx 1$, which are very precisely approximated by a single Slater determinant.

written as

$$H = -\frac{1}{2} \sum_{i=1}^{L} \left(c_i^{\dagger} c_{i+1} + \text{h.c.} \right) + \sum_{i=1}^{L} \varepsilon_i \left(n_i - \frac{1}{2} \right) + V \sum_{i=1}^{L} \left(n_i - \frac{1}{2} \right) \left(n_{i+1} - \frac{1}{2} \right).$$
 (3)

Here c_i destroys a fermion on site i, and $n_i = c_i^{\dagger} c_i$ is the associated number operator, while we set $c_{i+L} \equiv c_i$ to enforce periodic boundary conditions. Disorder is introduced via a uniformly distributed uncorrelated onsite potential $\varepsilon_i \in [-W, W]$, where W denotes the disorder strength, while the interactions of strength V depend on the occupations of neighbouring sites. Earlier investigations of this system and its equivalent formulations in terms of interacting spins have established that for moderate interactions ($V \sim 1$) the critical disorder strength W_c of the MBL phase transition takes values in the range between 3 and 4 [4–6, 20, 36–38].




FIG. 2. Panel (a) shows the orbital incompatibility C_{ml} for half-filled systems as in Fig. 1, but for different sizes L=8,10,12,14 and additionally averaged over pairs with fixed l+m (states are ordered by descending overlap \mathcal{I}_m). The collapse of data in units of the random-matrix benchmark $C_0 = \sqrt{L/8}$ reveals that this incompatibility scales systematically as in a random system, and takes appreciable values. Panel (b) shows that the overlap \mathcal{I}_m itself only weakly depends on the system size.

To establish the incompatibility of MBL states numerically, we first calculate the Brueckner orbitals for each many-body eigenstate in a given realization [39], and determine from these the overlaps \mathcal{I}_m . We then order the states either by overlap \mathcal{I}_m or energy E_m , and next perform the disorder averages of these quantities, as well as of the incompatibility measures \mathcal{C}_{ml} for all pairs of eigenstates. All averages are carried out over 10^3 realizations.

The results of this procedure in a many-body localized system (V = 1.5, W = 8) of length L = 14 at half filling (N = 7) are shown in Fig. 1. In panels (a-c) the states are ordered by overlap. The averaged overlaps in panel (b) confirm that, as anticipated, a large fraction of states are exceedingly well approximated by a Slater determinant. Panel (a) then verifies that nonetheless, these states generically have noticeably incompatible orbitals, with the incompatibility roughly constant along the antidiagonal. The incompatibility is less pronounced for the states with the very largest overlaps, which as seen in panel (c) arise from the edge of the energy spectrum. Panels (d) and (e) show that these features are otherwise roughly independent of energy, provided again that we avoid the edges of the energy spectrum, according to which the states are ordered there [see panel (f)].

As shown in Fig. 2, the observed incompatibility is indeed sizable and scales systematically with the system size. As a benchmark, we consider the extreme case of a pair of states with randomly incompatible orbitals, and hence mutually uncorrelated projectors. Applying random-matrix theory (RMT; see the Appendix), we find that for such a pair of states, the squared incompatibility averages to $\overline{C_{ml}^2}|_{\rm RMT}=2N^2(L-N)^2/(L-1)L(L+\eta)$, where $\eta=1$ for complex orbitals and $\eta=2$ for real orbitals. Hence, as self-averaging suppresses statistical




FIG. 3. Dependence of (a) the averaged incompatibility \mathcal{C}_{ml} and (b) the overlap \mathcal{I}_m on the disorder strength W (half-filled systems of size L=8,12 at fixed V=1.5, obtained from eigenstates from the middle 10% of the energy spectrum). Panels (c,d) display the color-coded probability density function of \mathcal{I}_m for both system sizes.

fluctuations, for a half-filled large system $\overline{C_{ml}}|_{\rm RMT} \sim \sqrt{L/8} \equiv \mathcal{C}_0$. To compare the MBL system with this benchmark, we take the result from Fig. 1(a) averaged over the antidiagonals, and supplement this with the corresponding results for half-filled systems of size L=8,10,12. Figure 2(a) then confirms that the incompatibility in the MBL system is indeed sizeable, and in particular scales reliably as $\overline{C_{ml}} = O(L^{1/2})$, with a slight tendency to flatten out for larger systems.

Figure 3 illuminates how these characteristic features develop as a function of the disorder strength, where we focus on states from the central 10% of the energy spectrum. At values of $W \gtrsim 4$, where many-body localization sets in, a noticeable fraction of these states become very well approximated by a single Slater determinant. This occurs roughly independently of the system size, as can be seen both from the averages shown in panel (b) as well as, in particular, in the colour-coded probabilitydensity functions (pdfs) $P(\mathcal{I}_m)$ in panels (c,d). For smaller values of W, hence, in the ergodic phase, such well-approximated states are rare, and the pdf is strongly size dependent. On the other hand, the incompatibility measure \mathcal{C}_{ml} retains its systematic scaling for all disorder strengths, approaching its RMT limiting value \mathcal{C}_0 for very weak disorder. Most importantly, this measure is again still sizeable deep in the MBL phase, dropping only gradually to $\approx C_0/2$ at W=8.

Figure 4 further explores how the incompatibility develops as interactions V are switched on when one moves from Anderson localization to many-body localization




FIG. 4. Dependence of (a) the averaged incompatibility \mathcal{C}_{ml} and (b) the overlap \mathcal{I}_m on the interaction strength V (half-filled systems of size L=12, solid lines, and L=8, dashed lines, obtained from eigenstates from the middle 10% of the energy spectrum). The thick lines show data for W=8, where the system crosses over from Anderson localization to many-body localization. The thin lines contrast this to cases where the system transitions towards ergodic behaviour.

(W=8, thick curve), and contrasts this with systems developing ergodic many-body dynamics (W=2,3,4, thin curves). In the ergodic systems, the overlaps \mathcal{I}_m drop significantly with increasing interactions, and depend noticeably on system size. In contrast, in the cross-over to the MBL regime for W=8, the overlaps \mathcal{I}_m remain high while the incompatibility \mathcal{C}_{ml} increases, with both quantities being only weakly dependent on the system size.

Connection to l-bits. So far, we have used the Brueckner orbitals to provide a direct geometric view of the relation between different many-body localized eigenstates. In the remainder, we highlight the utility of this framework to further illuminate the general phenomenology of these systems, and formulate new questions. This applies particularly to the notion of emergent integrability involving an extensive set of local conserved quantities, the aforementioned l-bits. In the present work, these l-bits provide useful context when one interprets them as quasiparticle densities, $Q_i = q_i^{\dagger}q_i$, and seeks to construct the quasiparticle operators [21]

$$q_i^{\dagger} = p_i^{\dagger} + \sum_{l,m,n} K_{lmn} p_l^{\dagger} p_m^{\dagger} p_n + \dots \tag{4}$$

in terms of single-particle operators p_i . In the noninteracting case, where $q_i^{\dagger} = p_i^{\dagger}$, all eigenstates are Slater determinants constructed from a shared set of single-particle orbitals. In a many-body localized system, higher-order terms must be taken into account, and a key guiding question is then whether this expansion is perturbative in nature. Focusing on the first term in the series and assuming that it dominates, this would require the eigenstates of the system to be well approximated by Slater determinants. But as we have seen, this is not a sufficient condition: even though the eigenstates are all well approximated by Slater determinants, these Slater determinants cannot be constructed from a common set




FIG. 5. (a) Overlap $\mathcal{I}_{m,l}$ in individual realizations and (b) averaged orbital incompatibility in the system of Fig. 1, but with states taken from half-filling (solid curves, N=7, index m) and from the sector with a particle removed (dashed curves, N=6, index l; total number of states denoted as \mathcal{N}'). See the text for an interpretation of the systematic correlations in (a) and reduced incompatibility along the diagonal of (b) in terms of emergent integrability.

of single-particle orbitals, revealing that the construction of the l-bits is of a nonperturbative nature.

To further quantify this, we consider the relation of eigenstates $|\psi_m\rangle$ with N=L/2 particles to eigenstates $|\psi_l\rangle$ with N=L/2-1 particles. Some of these eigenstates should be related by the application of a single quasiparticle operator, $|\psi_m\rangle=q_i^\dagger|\psi_l\rangle$. In terms of the projectors P_m and P_l , the compatibility of these states will then depend on how much the higher-order dressing terms in q_i^\dagger fall into the space of orbitals that are already occupied in $|\psi_l\rangle$. If they fully do, the two states will have the same overlap with a single Slater determinant, $\mathcal{I}_m=\mathcal{I}_l$, and be highly compatible.

As shown in Fig. 5, this feature is indeed present in the numerical data, and manifests itself both on the level of individual realizations as well as in the statistical averages. Firstly, as shown in panel (a), in individual realizations, hence, for fixed given systems, the overlaps \mathcal{I}_m of states with different particle number follow each other closely. Secondly, as shown in panel (b), for pairs of states between which the generic compatibility is not already high, this interrelation shows up as a reduced averaged incompatibility of states along the diagonal. The threshold value $\overline{\mathcal{C}_{ml}} \approx 0.5$ at the boundary of the region where this effect is visible reveals the residual incompatibility of the described pairs of states. These geometric observations give direct quantitative insights into emergent integrability down to the level of individual systems, circumventing the challenging task of directly constructing the underlying quasi-particle operators.

Conclusions. In summary, we established a geometric framework to describe the interrelation of eigenstates in fermionic many-body localized systems, unraveling the intriguing structure and correlations exhibited by these paradigms of constrained complex quantum dynamics. Applied to a paradigmatic model system, this approach reveals that while individual eigenstates are well approximated by single Slater determinants, they collectively de-

part from a uniquely-defined single-particle picture. This supports the notion of strongly dressed quasiparticle excitations residing behind the local conserved quantities that characterize the emergent integrability of these systems. The results complement the existing phenomenology based on individual eigenstates, such as the emergence of an area law of entanglement, both on the level of individual realisations as well as in statistical averages. These considerations can also be usefully extended in a number of directions. For instance, based on our approach, one could inquire how well a single basis of orbitals can approximate a larger number of eigenstates, if not all of them [40]. Furthermore, to obtain additional insights into the ergodic phase and transition region, one could make use of the fact that the notion of Brueckner orbital could be extended to systematically approximate states by multiple Slater determinants. Finally, the described connection of our approach to the ubiquitous notion of l-bits, as well as generalizations of the one-particle density matrix [26, 28], could be explored to extend these considerations to non-fermionic systems, such as bosonic systems and spin chains.

This research was funded by UK Engineering and Physical Sciences Research Council (EPSRC) via Grant Nos. EP/P010180/1 and EP/L01548X/1. Computer time was provided by Lancaster University's High-End Computing facility.

Appendix: Random-matrix benchmark of orbital incompatibility

To determine how sizeable the numerically observed orbital incompatibility of MBL states is, we compare it to the extreme case of two random Slater determinants $|S_m\rangle$ and $|S_l\rangle$, with mutually uncorrelated projectors P_m and P_l , admitting for additional generality that they may represent states of possibly different particle number N and N'. As before, we denote the dimensionality of the single-particle Hilbert space as L.

We set out to obtain a compact result for the averaged squared incompatibility,

$$\overline{C_{ml}^2} = 2 \operatorname{tr} \overline{P_m P_l} - 2 \operatorname{tr} \overline{P_m P_l P_m P_l}$$
 (5)

$$= 2\operatorname{tr}\overline{(P_l P_m P_l)} - 2\operatorname{tr}\overline{(P_l P_m P_l)^2},\tag{6}$$

where the expression in the second line exploits that any projector fulfills $P_l^2 = P_l$. This is useful, as for any pair of states, we can adopt a single-particle basis in which P_l is diagonal and projects onto the first N states of the basis. In this basis, the combination

$$P_l P_m P_l = \begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix} \tag{7}$$

appearing in Eq. (6) has a finite subblock X of size $N \times N$,

and in terms of this

$$\overline{\mathcal{C}_{ml}^2} = 2\operatorname{tr}\overline{X} - 2\operatorname{tr}\overline{X^2},\tag{8}$$

where the trace is now of matrices of size N.

Diagonalizing P_m in this basis, we can further write $(P_m)_{rs} = \sum_{n=1}^{N'} U_{rn} U_{sn}^*$, where the columns of the unitary matrix U are the orthonormalized eigenstates of P_m , and the sum runs over the N' eigenstates with eigenvalue 1 (the remaining eigenvalues vanish). Therefore, $X = uu^{\dagger}$ be further written in terms of a rectangular $N \times N'$ -dimensional subblock u of the $L \times L$ -dimensional unitary matrix U. We then have to calculate

$$\overline{C_{ml}^2} = 2 \operatorname{tr} \overline{uu^{\dagger}} - 2 \operatorname{tr} \overline{(uu^{\dagger})^2}. \tag{9}$$

In random-matrix theory, we can evaluate these averages using a standard ensemble. Here we consider two cases, systems with real orbitals and systems with complex orbitals. For the case of real orbitals, we take U to be uniformly distributed over the orthogonal group O(L), which is also known as the circular real ensemble (CRE; distinct from the circular orthogonal ensemble in which $U = U^T$ is still complex) [41]. For the case of complex orbitals, we take U to be uniformly distributed over the unitary group U(L), which corresponds to the standard circular unitary ensemble (CUE). In both ensembles, all matrix elements of U are equivalent, which allows us to decompose the averaged traces

$$\operatorname{tr} \overline{uu^{\dagger}} = NN'A,\tag{10}$$

$$\operatorname{tr} \overline{(uu^{\dagger})^{2}} = NN'[B + (N+N'-2)C + (N-1)(N'-1)D]$$
(11)

of the truncated matrices into a small number of fundamental terms,

$$A \equiv \overline{|u_{rs}|^2}, \quad B \equiv \overline{|u_{rs}|^4}, \tag{12}$$

$$C \equiv \overline{|u_{rs}|^2 |u_{rt}|^2} = \overline{|u_{sr}|^2 |u_{tr}|^2} \quad (s \neq t), \tag{13}$$

$$D \equiv \overline{u_{rs}u_{ps}^*u_{pt}u_{rt}^*} \quad (r \neq p, s \neq t), \tag{14}$$

where unspecified indices are unconstrained. The same equivalence of matrix elements also implies the following sum rules,

$$\operatorname{tr} \overline{UU^{\dagger}} = L^2 A = L, \tag{15}$$

$$\operatorname{tr} \overline{(UU^{\dagger})^2} = L^2 B + 2L^2 (L-1)C + L^2 (L-1)^2 D = L,$$
(16)

$$\overline{[(UU^{\dagger})_{11}]^2} = LB + L(L-1)C = 1. \tag{17}$$

These expressions can all be combined to express the desired average as

$$\overline{\mathcal{C}_{ml}^2}|_{\text{RMT}} = -2DNN'(L-N)(L-N'). \tag{18}$$

Averages such as the ones presented above also appear in quantum transport, where u would be interpreted as a

block of a scattering matrix and the given combinations determine, for instance, universal conductance fluctuations and shot noise [42]. Furthermore, such averages also appear in the dynamics of quantum-chaotic systems, where U represents a time-evolution operator [43], while truncated versions appear in the description of leaky systems [44] and in Floquet-descriptions of quantum transport [45]. Here, A and B can be worked out by interpreting them as moments of a coordinate from a random L-dimensional unit vector, parameterized in hyperspherical coordinates, thereby relating it to integrals of the form

$$I_{n,m} = \int_0^{\pi} \cos^n(\varphi) \sin^m(\varphi) d\varphi, \tag{19}$$

while C and D then follow from the given sum rules. In the CRE, we then have

$$A = \frac{I_{2,L-2}}{I_{0,L-2}} = \frac{1}{L},\tag{20}$$

$$B = \frac{I_{4,L-2}}{I_{0,L-2}} = \frac{3}{L(L+2)},\tag{21}$$

$$C = \frac{1}{L(L+2)},\tag{22}$$

$$D = -\frac{1}{(L-1)L(L+2)},\tag{23}$$

while in the CUE we have

$$A = 2\frac{I_{2,2L-2}}{I_{0,2L-2}} = \frac{1}{L},\tag{24}$$

$$B = 2\frac{I_{4,2L-2}}{I_{0,2L-2}} + 2\frac{I_{2,2L}I_{2,2L-3}}{I_{0,2L-2}I_{0,2L-3}} = \frac{2}{L(L+1)}, \quad (25)$$

$$C = \frac{1}{L(L+1)},\tag{26}$$

$$D = -\frac{1}{L(L^2 - 1)}. (27)$$

(In both cases, A also follows from the given sum rule, and they can be related to each other using $B|_{\text{CUE}(L)} = 2[B|_{\text{CRE}(2L)} + C|_{\text{CRE}(2L)}]$.)

Therefore, as our final results, we obtain

$$\overline{C_{ml}^2}|_{\text{RMT}} = \frac{2NN'(L-N)(L-N')}{(L-1)L(L+2)}$$
 (CRE) (28)

for the averaged squared incompatibility of random Slater determinants with real orbitals, and

$$\overline{C_{ml}^2}|_{\text{RMT}} = \frac{2NN'(L-N)(L-N')}{L(L^2-1)}$$
 (CUE) (29)

for the case of complex orbitals. For large systems near half filling, both expressions approach $\overline{\mathcal{C}_{ml}^2}|_{\mathrm{RMT}} \sim L/8 \equiv C_0^2$. Level repulsion between the eigenvalues of X then guarantees that fluctuations about this value are suppressed [42]. Therefore, C_0 not only characterizes the

squared incompatibility, but can be used as a benchmark for the incompatibility C_{ml} itself.

- R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [2] E. Altman and R. Vosk, Universal dynamics and renormalization in many-body-localized systems, Annu. Rev. Condens. Matter Phys. 6, 383 (2015).
- [3] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. 91, 021001 (2019).
- [4] A. Pal and D. A. Huse, Many-body localization phase transition, Phys. Rev. B 82, 174411 (2010).
- [5] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, Phys. Rev. B 75, 155111 (2007).
- [6] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, Phys. Rev. B 77, 064426 (2008).
- [7] J. M. Deutsch, Quantum statistical mechanics in a closed system, Phys. Rev. A 43, 2046 (1991).
- [8] M. Srednicki, Chaos and quantum thermalization, Phys. Rev. E 50, 888 (1994).
- [9] D. Abanin, J. Bardarson, G. De Tomasi, S. Gopalakrishnan, V. Khemani, S. Parameswaran, F. Pollmann, A. Potter, M. Serbyn, and R. Vasseur, Distinguishing localization from chaos: Challenges in finite-size systems, Ann. Phys. (N.Y.) 427, 168415 (2021).
- [10] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Quantum chaos challenges many-body localization, Phys. Rev. E 102, 062144 (2020).
- [11] M. Serbyn and J. E. Moore, Spectral statistics across the many-body localization transition, Phys. Rev. B 93, 041424 (2016).
- [12] C. L. Bertrand and A. M. García-García, Anomalous Thouless energy and critical statistics on the metallic side of the many-body localization transition, Phys. Rev. B 94, 144201 (2016).
- [13] M. Serbyn, Z. Papić, and D. A. Abanin, Local conservation laws and the structure of the many-body localized states, Phys. Rev. Lett. 111, 127201 (2013).
- [14] D. A. Huse, R. Nandkishore, and V. Oganesyan, Phenomenology of fully many-body-localized systems, Phys. Rev. B 90, 174202 (2014).
- [15] V. Ros, M. Müller, and A. Scardicchio, Integrals of motion in the many-body localized phase, Nucl. Phys. B 891, 420 (2015).
- [16] J. Z. Imbrie, On many-body localization for quantum spin chains, J. Stat. Phys. 163, 998 (2016).
- [17] J. Z. Imbrie, Diagonalization and many-body localization for a disordered quantum spin chain, Phys. Rev. Lett. 117, 027201 (2016).
- [18] J. Z. Imbrie, V. Ros, and A. Scardicchio, Local integrals of motion in many-body localized systems, Ann. Phys. (Berl.) 529, 1600278 (2017).
- [19] L. Rademaker, M. Ortuño, and A. M. Somoza, Manybody localization from the perspective of integrals of motion, Ann. Phys. (Berl.) 529, 1600322 (2017).
- [20] S. Bera, H. Schomerus, F. Heidrich-Meisner, and J. H.

- Bardarson, Many-body localization characterized from a one-particle perspective, Phys. Rev. Lett. **115**, 046603 (2015).
- [21] S. Bera, T. Martynec, H. Schomerus, F. Heidrich-Meisner, and J. H. Bardarson, One-particle density matrix characterization of many-body localization, Ann. Phys. (Berl.) 529, 1600356 (2017).
- [22] B. Villalonga, X. Yu, D. J. Luitz, and B. K. Clark, Exploring one-particle orbitals in large many-body localized systems, Phys. Rev. B 97, 104406 (2018).
- [23] S.-H. Lin, B. Sbierski, F. Dorfner, C. Karrasch, and F. Heidrich-Meisner, Many-body localization of spinless fermions with attractive interactions in one dimension, SciPost Phys. 4, 002 (2018).
- [24] W. Buijsman, V. Gritsev, and V. Cheianov, Many-body localization in the Fock space of natural orbitals, SciPost Phys. 4, 38 (2018).
- [25] N. Macé, N. Laflorencie, and F. Alet, Many-body localization in a quasiperiodic Fibonacci chain, SciPost Phys. 6, 50 (2019).
- [26] C. P. Chen, M. Szyniszewski, and H. Schomerus, Manybody localization of zero modes, Phys. Rev. Research 2, 023118 (2020).
- [27] T. Orito, Y. Kuno, and I. Ichinose, Effects of power-law correlated disorders in XXZ spin chain: Many-body localized to thermal phase transition and its critical regime (2020), arXiv:2002.12575 [cond-mat.stat-mech].
- [28] M. Hopjan and F. Heidrich-Meisner, Many-body localization from a one-particle perspective in the disordered one-dimensional Bose-Hubbard model, Phys. Rev. A 101, 063617 (2020).
- [29] T. L. M. Lezama, S. Bera, H. Schomerus, F. Heidrich-Meisner, and J. H. Bardarson, One-particle density matrix occupation spectrum of many-body localized states after a global quench, Phys. Rev. B 96, 060202 (2017).
- [30] M. Hopjan, F. Heidrich-Meisner, and V. Alba, Scaling properties of a spatial one-particle densitymatrix entropy in many-body localized systems (2020), arXiv:2011.02200 [cond-mat.str-el].
- [31] K. A. Brueckner and C. A. Levinson, Approximate reduction of the many-body problem for strongly interacting particles to a problem of self-consistent fields, Phys. Rev. 97, 1344 (1955).
- [32] K. A. Brueckner and W. Wada, Nuclear saturation and two-body forces: Self-consistent solutions and the effects of the exclusion principle, Phys. Rev. 103, 1008 (1956).
- [33] R. K. Nesbet, Brueckner's theory and the method of superposition of configurations, Phys. Rev. **109**, 1632

- (1958).
- [34] S. Larsson and V. H. Smith, Analysis of the 2S ground state of Lithium in terms of natural and best overlap (Brueckner) spin orbitals with implications for the fermi contact term, Phys. Rev. 178, 137 (1969).
- [35] J. M. Zhang and M. Kollar, Optimal multiconfiguration approximation of an N-fermion wave function, Phys. Rev. A 89, 012504 (2014).
- [36] A. D. Luca and A. Scardicchio, Ergodicity breaking in a model showing many-body localization, EPL 101, 37003 (2013).
- [37] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field heisenberg chain, Phys. Rev. B 91, 081103 (2015).
- [38] Y. Bar Lev, G. Cohen, and D. R. Reichman, Absence of diffusion in an interacting system of spinless fermions on a one-dimensional disordered lattice, Phys. Rev. Lett. 114, 100601 (2015).
- [39] Brueckner orbitals can be constructed systematically in several ways. Here, we adopt the method of Ref. [35], where one starts from a set of trial orbitals and updates these iteratively one at a time, by which \mathcal{I}_m increases monotonously. This method can also be extended to approximate the state using M > N orbitals entering several Slater determinants.
- [40] For this, the geometric perspective could be deepened by exploiting the general mathematical properties of the projectors P_m. Mathematically, the space of these projectors is isomorphic to the Grassmannian G(H₁, N) of N dimensional subspaces in the single-particle Hilbert space H₁.
- [41] C. W. J. Beenakker, Random-matrix theory of majorana fermions and topological superconductors, Rev. Mod. Phys. 87, 1037 (2015).
- [42] C. W. J. Beenakker, Random-matrix theory of quantum transport, Rev. Mod. Phys. 69, 731 (1997).
- [43] F. Haake, M. Kus, H.-J. Sommers, H. Schomerus, and K. Zyczkowski, Secular determinants of random unitary matrices, J. Phys. A 29, 3641 (1996).
- [44] H. Schomerus and J. Tworzydło, Quantum-to-classical crossover of quasibound states in open quantum systems, Phys. Rev. Lett. 93, 154102 (2004).
- [45] J. Tworzydło, A. Tajic, H. Schomerus, and C. W. J. Beenakker, Dynamical model for the quantum-toclassical crossover of shot noise, Phys. Rev. B 68, 115313 (2003).