

Lifting the Convex Conjugate in Lagrangian Relaxations: A Tractable Approach for Continuous Markov Random Fields

Hartmut Bauermeister*[§] Emanuel Laude*[†] Thomas Möllenhoff[‡]
Michael Moeller[§] Daniel Cremers[†]

July 14, 2021

Abstract

Dual decomposition approaches in nonconvex optimization may suffer from a duality gap. This poses a challenge when applying them directly to nonconvex problems such as MAP-inference in a Markov random field (MRF) with continuous state spaces. To eliminate such gaps, this paper considers a reformulation of the original nonconvex task in the space of measures. This infinite-dimensional reformulation is then approximated by a semi-infinite one, which is obtained via a piecewise polynomial discretization in the dual. We provide a geometric intuition behind the primal problem induced by the dual discretization and draw connections to optimization over moment spaces. In contrast to existing discretizations which suffer from a grid bias, we show that a piecewise polynomial discretization better preserves the continuous nature of our problem. Invoking results from optimal transport theory and convex algebraic geometry we reduce the semi-infinite program to a finite one and provide a practical implementation based on semidefinite programming. We show, experimentally and in theory, that the approach successfully reduces the duality gap. To showcase the scalability of our approach, we apply it to the stereo matching problem between two images.

Keywords: Markov random fields · moment relaxation · sum of squares · polynomial optimization · generalized conjugacy · optimal transport

AMS Subject Classification: 49N15 · 49M29 · 65K10 · 90C23 · 90C26 · 90C35

1 Introduction and Motivation

1.1 Lagrangian Relaxations

In this paper, our goal is to develop a convex optimization framework à la dual decomposition for the MAP-inference problem in a continuous *Markov random field* (MRF). Continuous MRFs

*equal contribution

[†]Technical University of Munich, Department of Informatics, Munich, Germany

[‡]RIKEN Center for Advanced Intelligence Project, Tokyo, Japan.

[§]University of Siegen, Department of Electrical Engineering and Computer Science, Siegen, Germany

are versatile and therefore widely used as a model in image processing, computer vision and machine learning [42, 53, 55, 10, 13, 50, 2, 51, 12, 3, 26, 36, 49]. Due to the continuous nature of the state space, inference amounts to a continuous nonlinear optimization problem. For an undirected graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ it takes the following form:

$$\min_{x \in \Omega^{\mathcal{V}}} F(x) := \sum_{u \in \mathcal{V}} f_u(x_u) + \sum_{uv \in \mathcal{E}} f_{uv}(x_u, x_v). \quad (\text{P})$$

The objective function F is an additive composition of a separable part, $\sum_{u \in \mathcal{V}} f_u(x_u)$, with unary functions $f_u : (\Omega \subset \mathbb{R}^m) \rightarrow \mathbb{R}$ and a coupling part $\sum_{uv \in \mathcal{E}} f_{uv}(x_u, x_v)$ with pairwise functions $f_{uv} : \Omega \times \Omega \rightarrow \mathbb{R}$ and $\emptyset \neq \Omega \subset \mathbb{R}^m$ compact.

The coupling term $\sum_{uv \in \mathcal{E}} f_{uv}(x_u, x_v)$ introduces a major challenge for efficient optimization in particular when \mathcal{V} is large and therefore the problem is high-dimensional. For tractability, one therefore seeks to find a decomposable reformulation of the problem. In continuous optimization a viable approach is to derive a *Lagrangian relaxation* of the problem: The key idea is to introduce auxiliary variables $x_{uv} \in \mathbb{R}^m \times \mathbb{R}^m$ for each edge $uv \in \mathcal{E}$ and linear constraints $x_{uv} = (x_u, x_v)$. Dualizing the linear constraints with Lagrange multipliers $\lambda_{uv} \in \mathbb{R}^m \times \mathbb{R}^m$ one arrives at the Lagrangian dual problem which falls within the regime of convex optimization algorithms that can exploit its separable dual structure. Examples include subgradient ascent or the *primal-dual hybrid gradient* (PDHG) method [9].

However, the approach suffers from potentially large duality gaps since problem (P) is non-convex in general. Indeed, it can be shown that a direct Lagrangian relaxation of (P) leads to the following “naive” convexification of the original problem:

$$\min_{x \in (\mathbb{R}^m)^{\mathcal{V}}} \sum_{u \in \mathcal{V}} f_u^{**}(x_u) + \sum_{uv \in \mathcal{E}} f_{uv}^{**}(x_u, x_v). \quad (1)$$

With some abuse of notation, here, f_u and f_{uv} attain the value $+\infty$ whenever $x_u \notin \Omega$ or $(x_u, x_v) \notin \Omega^2$. Then f_u^{**} are the convex biconjugates which correspond to the largest convex lower semicontinuous under-approximations to f_u . Such component-wise convex envelopes can produce inaccurate or even trivial convex under-approximations to the global objective F .

1.2 The Local Marginal Polytope Relaxation

To overcome such duality gaps, we propose to reformulate the original nonlinear problem in terms of an infinite-dimensional linear program over the space of Radon measures and apply the Lagrangian relaxation afterwards.

The following key lemma reveals, that every minimization problem can be equivalently formulated in terms of an infinite-dimensional linear program. In the following, we denote $\mathcal{P}(X)$ as the space of Borel probability measures on $\emptyset \neq X \subset \mathbb{R}^m$ compact, and we denote by $\langle \mu, f \rangle = \int_X f(x) d\mu(x)$.

Lemma 1.1. *Let $f : X \rightarrow \mathbb{R}$ be lower semicontinuous (lsc) with $\emptyset \neq X \subset \mathbb{R}^m$ compact. Then we have*

$$\min_{x \in X} f(x) = \min_{\mu \in \mathcal{P}(X)} \langle \mu, f \rangle, \quad (2)$$

and $x^* \in \arg \min_{x \in X} f(x)$ is a solution to $\min_{x \in X} f(x)$ if and only if δ_{x^*} is a minimizer of $\min_{\mu \in \mathcal{P}(X)} \langle \mu, f \rangle$.

Proof. The result follows immediately via compactness of X and the properties of probability measures: By compactness of X and since f is lsc relative to X we know x^* exists. Let $f_{\min} = f(x^*)$. By the properties of the Lebesgue integral we have:

$$\int_X f(x) d\mu(x) \geq \int_X f_{\min} d\mu(x) = f_{\min} \int_X 1 d\mu(x) = \min_{x \in X} f(x),$$

for all $\mu \in \mathcal{P}(X)$ and $\min_{x \in \mathbb{R}^n} f(x) = \langle \delta_{x^*}, f \rangle = f(x^*)$. \square

Let $\pi_u : \Omega^{\mathcal{V}} \rightarrow \Omega$, $\pi_{uv} : \Omega^{\mathcal{E}} \rightarrow \Omega^2$ denote the canonical projections onto the u^{th} resp. u^{th} and v^{th} components and $\pi_u \# \mu$ is the pushforward of μ w.r.t. π_u defined by: $(\pi_u \# \mu)(A) = \mu((\pi_u)^{-1}(A))$ for all $A \subset \Omega$. Then, in our case, applying the reformulation from Lem. 1.1 directly to the cost function F with $X = \Omega^{\mathcal{V}}$ we obtain by linearity:

$$\begin{aligned} \min_{\mu \in \mathcal{P}(\Omega^{\mathcal{V}})} \langle \mu, F \rangle &= \min_{\mu \in \mathcal{P}(\Omega^{\mathcal{V}})} \left\langle \mu, \sum_{u \in \mathcal{V}} f_u \circ \pi_u + \sum_{uv \in \mathcal{E}} f_{uv} \circ \pi_{uv} \right\rangle \\ &= \min_{\mu \in \mathcal{P}(\Omega^{\mathcal{V}})} \sum_{u \in \mathcal{V}} \langle \pi_u \# \mu, f_u \rangle + \sum_{uv \in \mathcal{E}} \langle \pi_{uv} \# \mu, f_{uv} \rangle. \end{aligned}$$

This relaxation is known as the *full marginal polytope relaxation* which is, however, intractable if \mathcal{V} is large as one minimizes over probability measures on the product space $\mathcal{P}(\Omega^{\mathcal{V}})$. Instead, we consider the following linear programming relaxation of (P) which is also referred to as the *local marginal polytope relaxation* [26, 14, 48, 35] which is more tractable as the optimization variable lies in the product space of probability measures $\mathcal{P}(\Omega)^{\mathcal{V}}$:

$$\inf_{\mu \in \mathcal{P}(\Omega)^{\mathcal{V}}} \mathcal{F}(\mu) := \sum_{u \in \mathcal{V}} \langle \mu_u, f_u \rangle + \sum_{uv \in \mathcal{E}} \text{OT}_{f_{uv}}(\mu_u, \mu_v). \quad (\text{R-P})$$

Here, $\text{OT}_{f_{uv}}$ denotes the optimal transportation [15] with marginals (μ_u, μ_v) and cost f_{uv} defined by

$$\text{OT}_{f_{uv}}(\mu_u, \mu_v) = \inf_{\mu_{uv} \in \Pi(\mu_u, \mu_v)} \langle f_{uv}, \mu_{uv} \rangle. \quad (3)$$

The constraint set $\Pi(\mu_u, \mu_v)$ consists of all Borel probability measures on Ω^2 with specified marginals μ_u and μ_v :

$$\Pi(\mu_u, \mu_v) = \{ \mu_{uv} \in \mathcal{P}(\Omega^2) : \pi_u \# \mu_{uv} = \mu_u, \pi_v \# \mu_{uv} = \mu_v \}. \quad (4)$$

Note that for finite state-spaces (i.e., $|\Omega| < \infty$), the set $\mathcal{P}(\Omega)$ can be identified with the standard $(|\Omega| - 1)$ -dimensional probability simplex and $\Pi(\mu_u, \mu_v)$ with the set of nonnegative $|\Omega| \times |\Omega|$ matrices whose rows and columns sum up to μ_v and μ_u . In that case, the linear program given in (R-P) is equivalent to the well-known finite-dimensional local marginal polytope relaxation for MRFs, which is for example studied in [52]. We further remark that the case of finite Ω has been extensively studied in the literature, see, e.g., [16, 47] for recent overviews.

For the more challenging setting of continuous state-spaces, a major difficulty stems from the fact that the linear programming relaxation (R-P) is an infinite-dimensional optimization problem posed in the space of Borel probability measures. Perhaps due to this difficulty, discrete

MRF approaches are still routinely applied despite the continuous nature of Ω . This is typically done by considering a finite sample approximation of Ω , so that the infinite-dimensional linear program reduces to a finite-dimensional one. This, however, may lead to discretization errors and comes with an exponential complexity in the dimension of Ω .

Due to the fact that $\Pi(\delta_x, \delta_{x'}) = \{\delta_{(x,x')}\}$ one sees that restricting μ_u (and therefore μ_{uv}) to be Dirac probability measures, the formulation (R-P) reduces to the original problem (P). As one instead considers the larger convex set of all probability measures, it is a *relaxation* which lower bounds (P), i.e., we have the following important relation:

$$(R-P) \leq (P). \tag{5}$$

When considering a relaxation, the immediate question arises whether this lower bound is attained, i.e., the relaxation is tight and therefore the above inequality holds with equality. For finite and ordered Ω the situation is well-understood, see [52]. For infinite $\Omega \subset \mathbb{R}^m$, to our knowledge, the situation is less clear, unless f_u, f_{uv} are convex. An exception is, when f_u, f_{uv} are continuous piecewise linear: Then the continuous formulation is equivalent to a discrete MRF, see [14, Thm. 6], and existing results from the discrete case carry over. Despite the lack of theoretical tightness guarantees, in experiments, the infinite-dimensional local marginal polytope relaxation produces accurate lower relaxations, in particular when Ω is univariate.

A Lagrangian relaxation to the infinite-dimensional problem is obtained by dualizing the marginalization constraints (4) for each $e \in \mathcal{E}$ with Lagrange multipliers $\lambda_e \in \mathcal{C}(\Omega)^2$. This is equivalent to a substitution of the optimal transportation with its dual formulation. Adopting the approach of [14] a finite-dimensional problem is obtained by approximating the Lagrange multipliers in terms of finite linear combinations of certain basis functions. These approximations are chosen in such a way that the classical Lagrangian relaxation and the discrete approach, described above, are special cases of the considered framework: Let D be the Lagrange dual function. Then for any hierarchy of increasingly expressive dual subspaces $\Lambda_1 \subset \Lambda_2 \subset \dots \subset \mathcal{C}(\Omega)$, where Λ_1 is the space of affine functions, the induced hierarchy of dual problems

$$(1) = \max_{\lambda \in ((\Lambda_1)^2)^\mathcal{E}} D(\lambda) \leq \max_{\lambda \in ((\Lambda_2)^2)^\mathcal{E}} D(\lambda) \leq \dots \leq \max_{\lambda \in (\mathcal{C}(\Omega)^2)^\mathcal{E}} D(\lambda) = (R-P) \leq (P),$$

leads to a reduction of the duality gap. For a piecewise polynomial hierarchy with increasing degrees and/or number of pieces we will show that the gap eventually vanishes as the number of pieces and/or the degree goes to ∞ .

1.3 Contributions and Overview

Expanding upon existing related work our contributions are threefold:

Theory: We derive and study the primal optimization problem corresponding to the dual discretization and draw connections to optimization over moment spaces and aspects from variational analysis. In the same way a convex duality follows from a linear separation argument, a nonconvex duality can be formulated in terms of nonlinear separation boundaries. After a “lifting” to a higher-dimensional space these nonlinear separation boundaries become linear. This is analogous to the idea of feature maps for linear classifiers and gives rise to our notion of a “lifted” convex conjugate. More formally, we show that under a certain extremality condition, satisfied by a polynomial discretization with degree at least 2, the generalized biconjugate of a

potentially discontinuous, nonconvex function equals (up to closure) the original function. As a consequence, under extremality, the discretized energy agrees with the original energy, when restricted to a certain continuum of extreme points associated to Diracs. And, for every point in the continuous domain of the original energy such a correspondence exists. In contrast, in a piecewise linear discretization there only exist finitely many extreme points which makes such a one-to-one correspondence impossible. In addition, for a spline-based dual discretization we show that the nonconvex primal-dual gap vanishes as the degree and/or number of pieces goes to infinity.

First tractable formulation and implementation: We exploit optimal transport duality theory, and the aforementioned duality relations to optimization over moment spaces to derive, to our knowledge, the first tractable polynomial discretization of the infinite-dimensional local marginal polytope relaxation. More precisely, after discretization the problem amounts to a semi-infinite program which can be transformed into a separable finite-dimensional semidefinite program applying concepts from algebraic geometry, such as nonnegativity certificates of polynomials. This allows us to derive an efficient first-order primal-dual algorithm which, due to the separable problem structure, can be parallelized on a GPU to handle large problems.

Experimental evaluation and large scale capability: Experimentally, we provide numerical evidence for the strict reduction of the duality-gap and demonstrate the advantages of the nonlinear approximation over existing liftings in the literature in Sec. 5. We implement our algorithm on a GPU and apply it to the nonconvex problem of large-scale stereo matching. The results show that increasing the dual subspace improves both, the dual energy, and the quality of a recovered primal solution from a discretized measure.

Notation For a compact nonempty set $\Omega \subset \mathbb{R}^m$, denote by $\mathcal{P}(\Omega)$ the space of Borel probability measures on Ω and by $\mathcal{M}(\Omega)$ the space of Radon measures on Ω . The convex cone of nonnegative Radon measures is denoted by $\mathcal{M}_+(\Omega)$. Furthermore let $\mathcal{C}(\Omega)$ be the space of continuous functions on Ω . We will write $\langle f, \mu \rangle = \int_{\Omega} f(x) d\mu(x)$, for $f \in \mathcal{C}(\Omega)$ and $\mu \in \mathcal{M}(\Omega)$. For a set $C \subset \mathbb{R}^m$ we denote by $\iota_C : \mathbb{R}^m \rightarrow \overline{\mathbb{R}} = \mathbb{R} \cup \{\pm\infty\}$ the indicator function with $\iota_C(x) = 0$ if $x \in C$ and $\iota_C(x) = \infty$ if $x \notin C$ and by $\sigma_C(x) = \sup_{y \in C} \langle x, y \rangle$ the support function of C at $x \in \mathbb{R}^m$. We denote by $C^* = \{y \in \mathbb{R}^m : \langle y, x \rangle \geq 0, \forall x \in C\}$ the dual cone of C . These notions are defined analogously for the topologically paired spaces $\mathcal{M}(\Omega)$ and $\mathcal{C}(\Omega)$. The convex hull $\text{con } C$ of a set $C \subset \mathbb{R}^m$ is the smallest convex set that contains C . Equivalently, $\text{con } C$ is the set of all finite convex combinations of points in C . With some abuse of notation we write $\text{con } f$ for a function f to denote the largest convex function below f . Let δ_x denote the Dirac measure centered at $x \in \mathbb{R}^m$. We write lsc for lower semicontinuous and denote the epigraphical closure of some function f by $\text{cl } f$. Let $\llbracket \cdot \rrbracket$ denote the Iverson bracket, where $\llbracket P \rrbracket = 1$ if P is true and $\llbracket P \rrbracket = 0$, otherwise. For some extended real-valued function $f : \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$ let $f^*(y) = \sup_{x \in \mathbb{R}^m} \langle x, y \rangle - f(x)$ denote the Fenchel conjugate of f at y and $f^{**} = (f^*)^*$ the Fenchel biconjugate.

1.4 Related Work

MRFs Fix and Agarwal [14] are the first to propose a dual subspace approximation of the infinite-dimensional local marginal polytope relaxation. This allows for a generalization and unified treatment of convex piecewise linear formulations for MRFs with continuous state spaces proposed in [55, 54]. However, for the polynomial case, due to the semi-infinite problem struc-

ture, the existence of an implementable algorithm beyond subgradient ascent is posed as an open question by [14]. As a consequence, no numerical results are presented. Dual subspace approximations for related models in a spatially continuous setting have been considered by [23] and further employed for the global optimization of vector- or manifold-valued optimization problems in imaging and vision [24, 45, 46]. In these papers, piecewise linear dual approximations are used and the techniques developed in the present work allow one to go beyond the piecewise linear case in a tractable way. Besides applications in imaging and vision, MRFs with continuous state spaces have for example been used in protein folding [26].

Lifting Lifting to measures for global constrained polynomial optimization is used in [18, 19] to reformulate the infinite-dimensional *linear program* (LP) in terms of a linear objective over the *semidefinite programming* (SDP) characterization of the finite-dimensional space of moments. Its SDP-dual is connected to nonnegativity certificates of polynomials [17, 41, 39]. As a key difference to [18, 19] we consider possibly nonpolynomial objective functions which results in a generally nonlinear formulation over the space of moments. Similar techniques have been considered recently in [20, 11] to estimate Lipschitz constants in neural networks.

Lagrangian relaxation and decoupling by lower relaxations A component-wise lifting in problems with a partially separable structure results in a generalized dual decomposition approach. Dual decomposition and Lagrangian relaxation are general principles in optimization and appear as a useful tool across many disciplines, see [4, Ch. 5–6], [22] and the references therein for an overview. Traditionally, decomposition methods are applied directly in the non-convex setting without lifting, unlike the generalized scheme we present in this paper which is based on a lifted reformulation. Without lifting this typically results in a component-wise convex lower envelope. More recently, [40] consider a homotopy method based on component-wise Lasry–Lions envelopes which specialises to a component-wise convex envelope in a certain limit case.

Generalized conjugacy and duality Our notion of a lifted convex conjugate is closely related to generalized conjugate functions originally due to [25]. It was utilized by [1] to study nonconvex dualities, expanding upon the work of [32] on augmented Lagrangians for nonconvex optimization. The specific case of quadratic conjugate functions was developed by [27] as an analytic tool in a seminal proof that the proximal subgradient map of a lower semicontinuous extended real-valued function is monotone if and only if the function is convex. Our motivation, shared with [32], is to eliminate duality gaps in nonconvex optimization. We offer a somewhat complementary view through a primal-type lifting approach which allows us to establish the connection to optimization in spaces of measures.

2 Dual Discretization for the Continuous MRF

2.1 A Reduced Dual Formulation

In contrast to previous approaches [14], we restrict ourselves to metric pairwise terms $f_{uv}(x, y) = d(x, y)$ which eventually leads to a different dual formulation and turns out more tractable: Then, the optimal transportation $\text{OT}_{f_{uv}}(\mu_u, \mu_v)$ in Problem (R-P) is the Wasserstein-1 distance

$W_1^d(\mu_u, \mu_v)$ induced by the metric d between μ_u and μ_v . Furthermore we assume that $f_u : \Omega \rightarrow \mathbb{R}$ is lsc and $\Omega \subset \mathbb{R}^m$ is a compact nonempty set. In particular this implies that f_u is proper and bounded from below.

Thanks to optimal transport duality theory [44] we are therefore able to obtain a more compact dual formulation, which is instrumental to derive a tractable implementation for a piecewise polynomial discretization later on: More precisely, we substitute $\text{OT}_{f_{uv}}(\mu_u, \mu_v) = W_1^d(\mu_u, \mu_v)$ in Problem (R-P) with its dual formulation

$$W_1^d(\mu_u, \mu_v) = \sup_{\lambda \in \text{Lip}_d(\Omega)} \int \lambda(x) d(\mu_u - \mu_v)(x), \quad (6)$$

where λ is 1-Lipschitz with respect to the metric d , i.e.,

$$\lambda \in \text{Lip}_d(\Omega) = \{\lambda : \Omega \rightarrow \mathbb{R} : |\lambda(x) - \lambda(y)| \leq d(x, y)\}. \quad (7)$$

In contrast to the general dual formulation of optimal transport, which involves two Lagrange multipliers per edge, interacting via the constraint set, the Wasserstein-1 dual involves only a single dual variable for each edge, that satisfies a Lipschitz constraint. We show in Sec. 4 that even though this formulation still involves infinitely many constraints, thanks to convex algebraic geometry [5], there exists a tractable finite representation of this constraint set in terms of SDP.

For notational convenience, we assign an arbitrary orientation to the edges in \mathcal{G} . After introducing a graph divergence operator $\text{Div} : \mathcal{C}(\Omega)^\mathcal{E} \rightarrow \mathcal{C}(\Omega)^\mathcal{V}$ defined by:

$$-(\text{Div } \lambda)_u = \sum_{v:(u,v) \in \mathcal{E}} \lambda_{(u,v)} - \sum_{v:(v,u) \in \mathcal{E}} \lambda_{(v,u)}, \quad (8)$$

an interchange of min and sup yields the following reduced dual problem, which is the starting point for further discussion and a tractable implementation:

$$\sup_{\lambda \in \mathcal{C}(\Omega)^\mathcal{E}} D(\lambda) := - \sum_{u \in \mathcal{V}} \sigma_{\mathcal{P}(\Omega)}(-f_u + (\text{Div } \lambda)_u) - \sum_{e \in \mathcal{E}} \iota_{\mathcal{K}}(\lambda_e), \quad (\text{R-D})$$

where $\mathcal{K} = \text{Lip}_d(\Omega)$ and $\sigma_{\mathcal{P}(\Omega)}(-f_u + (\text{Div } \lambda)_u) = \sup_{\mu \in \mathcal{P}(\Omega)} \langle \mu, -f_u + (\text{Div } \lambda)_u \rangle$ is the support function of $\mathcal{P}(\Omega)$ at $-f_u + (\text{Div } \lambda)_u$, which, thanks to Lem. 1.1, can be rewritten:

$$-\sigma_{\mathcal{P}(\Omega)}(-f_u + (\text{Div } \lambda)_u) = \min_{x \in \Omega} (f_u - (\text{Div } \lambda)_u)(x). \quad (9)$$

Under mild assumptions we have strong duality:

Proposition 2.1. *Let $\emptyset \neq \Omega \subset \mathbb{R}^m$ be compact, let $f_u : \Omega \rightarrow \mathbb{R}$ be lsc, and $f_{uv} = d$, where $d : \Omega^2 \rightarrow \mathbb{R}$ is lsc and a metric. Then, the following strong duality holds:*

$$(\text{R-P}) = (\text{R-D}), \quad (10)$$

and a maximizer of (R-D) exists.

Proof. Define $f : \mathcal{M}(\Omega)^\mathcal{V} \rightarrow \overline{\mathbb{R}}$

$$f(\mu) := \sum_{u \in \mathcal{V}} \langle f_u, \mu_u \rangle + \iota_{\mathcal{P}(\Omega)}(\mu_u),$$

which is convex, proper and lsc due to [37, Lem. 1.1.3]. Likewise, define the functional $g : \mathcal{M}(\Omega)^\mathcal{E} \rightarrow \overline{\mathbb{R}}$

$$g(\nu) := \sum_{e \in \mathcal{E}} \sup_{\lambda_e \in \mathcal{K}} \langle \lambda_e, \nu_e \rangle = \sum_{e \in \mathcal{E}} \sigma_{\mathcal{K}}(\nu_e),$$

which is proper convex lsc, as it is a pointwise supremum over linear functionals. Define $\nabla : \mathcal{M}(\Omega)^\mathcal{V} \rightarrow \mathcal{M}(\Omega)^\mathcal{E}$

$$(\nabla \mu)_{(u,v)} = \mu_u - \mu_v,$$

which is bounded and linear. Then we compute the convex conjugates $f^* : \mathcal{C}(\Omega)^\mathcal{V} \rightarrow \overline{\mathbb{R}}$ to

$$f^*(\theta) = \sum_{u \in \mathcal{V}} \sigma_{\mathcal{P}(\Omega)}(\theta_u - f_u),$$

and the functional $g^* : \mathcal{C}(\Omega)^\mathcal{E} \rightarrow \overline{\mathbb{R}}$

$$g^*(\lambda) = \sum_{e \in \mathcal{E}} \iota_{\mathcal{K}}(\lambda_e),$$

and $\nabla^* : \mathcal{C}(\Omega)^\mathcal{E} \rightarrow \mathcal{C}(\Omega)^\mathcal{V} = -\text{Div}$.

Choose $x \in \Omega$ and define $\mu := (\delta_x)^\mathcal{V}$. Then $f(\mu) = \sum_{u \in \mathcal{V}} f_u(x) < \infty$. In addition we have $\langle \lambda_e, \delta_x - \delta_x \rangle = 0$ for all $\lambda_e \in \mathcal{C}(\Omega)$ and therefore $g(\nabla \mu) = 0$. Now consider a weakly*-convergent sequence $\mathcal{M}(\Omega)^\mathcal{E} \ni \nu^t \xrightarrow{*} \nabla \mu$. This means for all $e \in \mathcal{E}$ and $\lambda_e \in \mathcal{C}(\Omega)$ we have $\langle \lambda_e, \nu_e^t \rangle \rightarrow \langle \lambda_e, (\nabla \mu)_e \rangle = 0$. In particular this implies that $g(\nu^t) \rightarrow 0$, and therefore g is continuous at $\nabla \mu$. Then we can invoke the Fenchel–Rockafellar duality Theorem [30, Thm. 1] and obtain that (R-P) = (R-D) and a maximizer of (R-D) exists. \square

As discussed in Sec. 1 the next step in our strategy to obtain a tractable formulation is to restrict $\lambda_{(u,v)} \in \mathcal{C}(\Omega)$ in Problem (R-D) to a subspace $\Lambda = \langle \varphi_0, \dots, \varphi_n \rangle$ spanned by basis functions $\varphi_k \in \mathcal{C}(\Omega)$ and instead consider $\sup_{\lambda \in \Lambda^\mathcal{E}} D(\lambda)$.

Before we derive the primal problem corresponding to the discretized dual problem we discuss the duality between finite-dimensional subspaces of $\mathcal{C}(\Omega)$ and certain equivalence classes of measures and in particular moment spaces in the next section.

2.2 Duality of $\Lambda \subset \mathcal{C}(\Omega)$

Due to Riesz' theorem the dual space of continuous functions on a compact set $\mathcal{C}(\Omega)$ is given by the space of Radon measures $\mathcal{M}(\Omega)$. Note that conversely for any $\lambda \in \Lambda$ we obtain a linear functional on $\mathcal{M}(\Omega)$ by mapping $\mu \in \mathcal{M}(\Omega)$ to $\mu \mapsto \langle \mu, \lambda \rangle = \int \lambda(x) d\mu(x)$. Although any $\mu \in \mathcal{M}(\Omega)$ maps to a linear functional on Λ , this mapping is not injective. In particular there exist distinct measures $\mu \neq \nu$ such that $\langle \mu, \lambda \rangle = \langle \nu, \lambda \rangle$ for every $\lambda \in \Lambda$, i.e., μ and ν induce the same linear functional on Λ . Instead, invoking [34, Thm. 4.9] the dual space Λ^* can be related to a quotient space of $\mathcal{M}(\Omega)$ as follows: Λ^* is isometrically isomorphic to a quotient space on $\mathcal{M}(\Omega)$ via

$$\Lambda^* \cong \mathcal{M}(\Omega)/\Lambda^0 := \{ \mu + \Lambda^0 : \mu \in \mathcal{M}(\Omega) \}, \quad (11)$$

where $\Lambda^0 := \{\mu \in \mathcal{M} : \langle \mu, \lambda \rangle = 0 \text{ for all } \lambda \in \Lambda\}$ is the annihilator of Λ . Any $\mu \in \mathcal{M}(\Omega)$ generates an equivalence class in \mathcal{M}/Λ^0 denoted by $[\mu]$ and the corresponding element in Λ^* is denoted by $[\mu]_\Lambda$.

Further restricting to the cone of functions in Λ that are nonnegative on Ω , the corresponding dual cone has a nice interpretation: It can be identified with a certain space of moments: Given a measure $\mu \in \mathcal{M}_+(\Omega)$, for a particular choice of basis functions $\{\varphi_0, \varphi_1, \dots, \varphi_n\}$, we refer to $\langle \mu, \varphi_k \rangle = \int \varphi_k(x) d\mu(x)$ as the k^{th} moment of μ and likewise the moment space is

$$(\mathcal{M}_\Lambda)_+ = \{y \in \mathbb{R}^{n+1} : \exists \mu \in \mathcal{M}_+(\Omega) : y_k = \langle \mu, \varphi_k \rangle\} \subset \mathbb{R}^{n+1}. \quad (12)$$

This is discussed in Sec. 4 in the context of implementation. The above result then shows that the moment space $(\mathcal{M}_\Lambda)_+$ identifies the space of equivalence classes of nonnegative measures, where two nonnegative measures are considered equivalent if they have the same moments.

For the remainder of this paper, for simplicity, we fix the basis $\{\varphi_0, \varphi_1, \dots, \varphi_n\}$ and identify the dual variable $\lambda_{(u,v)}$ with its vector of coefficients $p_{(u,v)} \in \mathbb{R}^{n+1}$.

2.3 Dual Discretization for the Metric Pairwise MRF

In view of the Wasserstein-1 dual in Eq. (6) constant components in $\lambda_{(u,v)}$ cancel out and therefore, assuming φ_0 is the constant function, we can choose $p_0 = 0$. As a consequence we can represent $(\text{Div } \lambda)_u$ in terms of its coefficients $(\text{Div}_\Lambda p)_u$ w.r.t. the basis functions $\{\varphi_1, \dots, \varphi_n\}$, for an appropriately chosen linear mapping $\text{Div}_\Lambda : (\mathbb{R}^n)^\mathcal{E} \rightarrow (\mathbb{R}^n)^\mathcal{V}$.

We identify the basis functions φ_k as the component functions of the mapping $\varphi : \Omega \rightarrow \mathbb{R}^n$. Then, substituting the representation $\lambda_{(u,v)} = \langle p_{(u,v)}, \varphi(\cdot) \rangle$ in the support function we obtain thanks to Eq. (9) the following discretized dual problem:

$$\sup_{p \in (\mathbb{R}^n)^\mathcal{E}} D_\Lambda(p) := \sum_{u \in \mathcal{V}} \min_{x \in \Omega} -\langle (\text{Div}_\Lambda p)_u, \varphi(x) \rangle + f_u(x) - \sum_{e \in \mathcal{E}} \iota_{\mathcal{K}_\Lambda}(p_e), \quad (\text{dR-D})$$

where the constraint set \mathcal{K}_Λ is the set of coefficients of Lipschitz functions in Λ :

$$\mathcal{K}_\Lambda := \{p \in \mathbb{R}^n : \langle p, \varphi(\cdot) \rangle \in \text{Lip}_d(\Omega)\}. \quad (13)$$

In order to derive the primal problem note that for any $f_u : \Omega \rightarrow \mathbb{R}$ the expression $\sup_{x \in \Omega} \langle (\text{Div}_\Lambda p)_u, \varphi(x) \rangle - f_u(x)$, (up to the presence of $\varphi(x)$) resembles the form of a convex conjugate. Indeed, exploiting the notion of an extended real-valued function this is the convex conjugate of a lifted version $f_\Lambda : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ of $f : \Omega \rightarrow \mathbb{R}$ which is defined by

$$f_\Lambda(y) = \begin{cases} f(x), & \text{if } y = \varphi(x) \text{ for some } x \in \Omega, \\ \infty, & \text{otherwise,} \end{cases} \quad (14)$$

and gives rise to our notion of a *lifted convex conjugate* which is studied in depth in Sec. 3.

Since the “lifting” to a higher dimensional space happens through the application of the mapping φ , whose component functions are the elements of the basis, it is intuitive to refer to φ as the lifting aka feature map in this context. Likewise we refer to the curve described by $x \mapsto \varphi(x)$ as the moment curve.

With this construction at hand and via an application of Fenchel–Rockafellar duality we are ready to state the primal problem of the discretized dual problem in terms of the lifted biconjugates $(f_u)_\Lambda^{**}$:

$$\min_{y \in (\mathbb{R}^n)^\mathcal{V}} \mathcal{F}_\Lambda(y) := \sum_{u \in \mathcal{V}} (f_u)_\Lambda^{**}(y_u) + \sum_{e \in \mathcal{E}} \sigma_{\mathcal{K}_\Lambda}((\nabla_\Lambda y)_e), \quad (\text{dR-P})$$

where $\nabla_\Lambda := -\text{Div}_\Lambda^*$ is the negative adjoint of the graph divergence, the graph gradient operator. A comparison of (dR-P) with the convex relaxation (1) shows the effect of the dual discretization, where the classical convex biconjugates are replaced with biconjugates of the “lifted” functions $(f_u)_\Lambda$. Indeed, without lifting, i.e., $\varphi(x) = x$, we recover the classical biconjugates and therefore the convex relaxation (1).

In experiments, we will discretize the dual problem with a piecewise polynomial family of functions. For intervals, the following proposition shows that either by increasing the number of pieces or the degree of the polynomial the primal-dual gap can be reduced. In our case we approximate the Lipschitz dual variable in terms of a Lipschitz spline. As a consequence existing results such as [14, Thm. 2] do not apply. Instead, we use a construction based on Bernstein-polynomials. Then the result follows from [6, Thm. 1].

Proposition 2.2. *Assume that $\Omega = [a, b] \subset \mathbb{R}$, $a < b$, and let the metric d be given by $d(x, y) = |x - y|$. Furthermore, let $\Lambda \subset \mathcal{C}(\Omega)$ be the space spanned by continuous piecewise polynomials on intervals $[t_i, t_{i+1}]$ defined by a regularly spaced grid with nodes given by $t_i = a + (b - a) \cdot (i - 1) / K$, $i = 1, \dots, K + 1$. Then the optimality gap satisfies:*

$$(\text{R-P}) - (\text{dR-D}) = \mathcal{O}(1/(K \cdot \sqrt{\text{deg}})),$$

where deg is the degree of the polynomial on each piece.

Proof. We consider the discretized dual problem (dR-D) where K is the set of coefficients corresponding to 1-Lipschitz piecewise polynomials on $[a, b]$ of degree deg with k pieces. Also recall that we have the following relations between the dual and primal problems: $(\text{dR-P}) = (\text{dR-D}) \leq (\text{R-P}) = (\text{R-D})$.

Now, let us denote a maximizer of (R-D) as $\lambda^* \in \text{Lip}_d(\Omega)^\mathcal{E}$. Existence of such a dual maximizer follows by Prop. 2.1. Then, one has for any $\lambda \in \Lambda^\mathcal{E}$:

$$\begin{aligned} \min_{x \in \Omega} f_u(x) - (\text{Div } \lambda^*)_u(x) &= \min_{x \in \Omega} f_u(x) - (\text{Div } \lambda)_u(x) - (\text{Div } \lambda^*)_u(x) + (\text{Div } \lambda)_u(x) \\ &\leq \min_{x \in \Omega} f_u(x) - (\text{Div } \lambda)_u(x) + \| -(\text{Div}(\lambda^* - \lambda))_u \|_\infty. \end{aligned} \quad (15)$$

This allows us to bound the optimality gap by:

$$\begin{aligned} (\text{R-D}) - (\text{dR-D}) &\leq \sum_{u \in \mathcal{V}} \| -(\text{Div}(\lambda - \lambda^*))_u \|_\infty \\ &\leq \sum_{u \in \mathcal{V}} |d(u)| \cdot \sup_{e \in \mathcal{E}} \|\lambda_e - \lambda_e^*\|_\infty \\ &\leq 2|\mathcal{E}| \cdot \sup_{e \in \mathcal{E}} \|\lambda_e - \lambda_e^*\|_\infty, \end{aligned} \quad (16)$$




Figure 1: Different finite-dimensional approximations \mathcal{P}_Λ of the infinite-dimensional space of probability measures $\mathcal{P}([-1, 1])$ with $\Omega = [-1, 1]$. Left and middle: 2-dimensional probability simplex and right: Monomial moment space $\text{con}\{(x, x^2, x^3) : x \in [-1, 1]\}$ of degree 3. The approximations are obtained as the convex hulls of the black curves $x \mapsto \varphi(x)$ for 3 different choices of φ . From left to right cf. Ex. 3.3, Ex. 3.4 and Ex. 3.5. In all cases, the black curves themselves correspond to Dirac measures and the convex hulls of the curves correspond to the space of probability measures. In contrast to the simplex that only has a finite number of extreme points, the monomial moment curve comprises a continuum of extreme points so that no Dirac measure on the monomial moment curve can be expressed as a convex combination of other Diracs.

where $d(u)$ denotes the degree of the vertex u .

For a L -Lipschitz function $f : [0, 1] \rightarrow \mathbb{R}$ there exists a Bernstein polynomial $p : [0, 1] \rightarrow \mathbb{R}$ with $p(0) = f(0)$ and $p(1) = f(1)$ such that $\|p - f\|_\infty \leq \frac{3L}{2} \text{deg}^{-1/2}$ [8, Thm. 2.6]. By [6, Thm. 1], this polynomial is L -Lipschitz as well. For each $e \in \mathcal{E}$ we pick the coefficients of the function λ_e such that it approximates the optimal dual variable λ_e^* with such a polynomial individually on each interval $[t_i, t_{i+1}]$. Then one obtains an overall 1-Lipschitz polynomial with the following bound:

$$\|\lambda_e - \lambda_e^*\|_\infty \leq \frac{3(b-a)}{2k\sqrt{\text{deg}}}. \quad (17)$$

Inserting this into (16) yields via (R-P) = (R-D):

$$(\text{R-P}) - (\text{dR-D}) \leq 3|\mathcal{E}| \frac{(b-a)}{k\sqrt{\text{deg}}}, \quad (18)$$

which gives the stated $\mathcal{O}(1/(k \cdot \sqrt{\text{deg}}))$ rate. \square

3 Lifted Convex Conjugates

3.1 Lifting and Moments

A fundamental question in the study of the lifted biconjugate $(f_u)_\Lambda^{**}$ is the characterization of its domain $\text{dom}(f_u)_\Lambda^{**} = \text{con } \varphi(\Omega)$, which, assuming f_u is bounded from below, equals the convex

hull of the graph of φ . For that purpose we define the probability moment space, as the set of all vectors $y \in \mathbb{R}^n$ for which there exists a probability measure $\mu \in \mathcal{P}(\Omega)$ such that the k^{th} component of y is the k^{th} moment of μ :

$$\mathcal{P}_\Lambda = \{y \in \mathbb{R}^n : \exists \mu \in \mathcal{P}(\Omega), y_k = \langle \mu, \varphi_k \rangle\}, \quad (19)$$

In other words \mathcal{P}_Λ is the set of all “infinite” convex combinations of points $y \in \varphi(\Omega)$ that belong to the graph of φ . The following relation to $(\mathcal{M}_\Lambda)_+$ holds: If $\varphi_0 \equiv 1$, a vector $(y_0, y_1, \dots, y_n) \in (\mathcal{M}_\Lambda)_+$ if and only if $(y_1, \dots, y_n) \in \mathcal{P}_\Lambda$. Also see Lem. 4.1.

Furthermore, note that for some $x \in \Omega$ the moment vector $\varphi(x) = \int \varphi(x') d\delta_x(x')$ of a Dirac measure δ_x sets up a correspondence between lifted points $\varphi(x)$ and the Dirac measure δ_x itself. Thus the moment space can be written equivalently as the set of all finite convex combinations of moment vectors of Dirac measures, in the same way the unit simplex is the convex hull of the unit vectors:

Proposition 3.1. *Let $\emptyset \neq \Omega$ be compact and $\Lambda = \langle \varphi_0, \dots, \varphi_n \rangle \subset \mathcal{C}(\Omega)$ with $\varphi_0 \equiv 1$. Then every moment vector of a probability measure is a finite convex combination of moment vectors of Dirac measures, i.e.*

$$\text{con}(\varphi(\Omega)) = \mathcal{P}_\Lambda, \quad (20)$$

and \mathcal{P}_Λ is nonempty and compact.

Proof. Note that \mathcal{P}_Λ is convex and bounded. It is also closed: To this end consider a sequence $y^t \rightarrow y$ with $y^t \in \mathcal{P}_\Lambda$. This means for any t there exists $\mu^t \in \mathcal{P}(\Omega)$ with $y_k^t = \int_\Omega \varphi_k(x) d\mu^t(x)$ for $k = 1, \dots, n$. Since Ω is compact, due to Prokhorov’s theorem, see [37, Sec. 1.1], there exists a weakly* convergent subsequence $\mu^{t_j} \xrightarrow{*} \mu$ and, hence, $\int_\Omega \varphi_k(x) d\mu^{t_j}(x) = y_k^{t_j} \rightarrow \int_\Omega \varphi_k(x) d\mu(x) = y_k$. Therefore $y \in \mathcal{P}_\Lambda$. Next we show identity of the support functions of the convex sets $\text{con}(\varphi(\Omega))$ and \mathcal{P}_Λ as this implies the equality of the sets. To this end let $f \in \Lambda = \langle \varphi_0, \dots, \varphi_n \rangle$ with $\varphi_0 = 1$. We write $f(x) = \langle \varphi(x), f \rangle$. Assume that $f_0 = 0$. We have the identities:

$$\sigma_{\varphi(\Omega)}(-f) = -\min_{x \in \Omega} \langle \varphi(x), f \rangle = -\min_{\mu \in \mathcal{P}(\Omega)} \int_\Omega f(x) d\mu(x) = -\min_{y \in \mathcal{P}_\Lambda} \langle y, f \rangle = \sigma_{\mathcal{P}_\Lambda}(-f),$$

where the first equality follows by the definition of the support function, the second equality by Lem. 1.1 and the third equality by the definition of \mathcal{P}_Λ , the identity

$$\min_{\mu \in \mathcal{P}(\Omega)} \int_\Omega f(x) d\mu(x) = \min_{\mu \in \mathcal{P}(\Omega)} \sum_{k=1}^n f_k \int_\Omega \varphi_k(x) d\mu(x),$$

and the substitution $y_k = \int_\Omega \varphi_k(x) d\mu(x)$. Since for each such f the support functions are equal we have equality of the support functions of $\varphi(\Omega)$ and \mathcal{P}_Λ . Since Ω is compact and φ continuous and the convex hull of a compact set stays compact, cf. [33, Cor. 2.30], we can replace $\varphi(\Omega)$ with its convex hull $\text{con}(\varphi(\Omega))$ and the conclusion follows. \square

The proof of the above proposition also reveals that for $f \in \Lambda$, the lifted biconjugate is a linear function over the moment space,

$$f_\Lambda^{**}(y) = \langle y, f \rangle + \iota_{\mathcal{P}_\Lambda}(y), \quad (21)$$

whose minimization is actually equivalent to minimizing the original function.

Specializing φ_k to the monomial basis and Ω to a set defined via polynomial inequalities, this yields the formulation proposed by [18, 19] for constrained polynomial optimization.

More generally, a meaningful notion of moments y_k is induced by a lifting map φ which is a homeomorphism between Ω and $\varphi(\Omega)$.

Definition 3.2 (lifting map). *Let Ω be compact and be nonempty. Then we say the mapping $\varphi : \Omega \rightarrow \mathbb{R}^n$ is a lifting map if φ is continuous on Ω and injective with continuous inverse $\varphi^{-1} : \varphi(\Omega) \rightarrow \mathbb{R}^n$.*

It is instructive to discuss possible choices for φ including the ones that correspond to existing discretizations for the continuous MRF such as the discrete approach and the piecewise linear approach. In the latter two cases, the moment space is merely the unit simplex. In that sense, the monomial moment space can be interpreted as a “nonlinear probability simplex”, which, in contrast to the unit simplex, has infinitely many extreme points, see Fig. 1. For the same reason, as we will see in the course of this section, it better suits the continuous nature of our optimization problem.

The discrete sampling-based approach is recovered by the following choice of φ :

Example 3.3. We discretize the interval $\Omega = [a, b]$, $a < b$ and re-define $\Omega := \{t_1, \dots, t_n\}$ with $t_k \in [a, b]$, $t_k < t_{k+1}$. For any $t_k \in \Omega$ let $\varphi(t_k) = e_k$ with $e_k \in \mathbb{R}^n$ being the k^{th} unit vector. As a result φ is the canonical basis and spans the space of discrete functions $f : \{t_1, \dots, t_n\} \rightarrow \mathbb{R}$ and the moment space $\text{con}(\varphi(\Omega)) = \mathcal{P}_\Lambda$ is given by the unit simplex.

The above example can be extended by assigning any points $x \in (t_k, t_{k+1})$ to points on the connecting line between two corresponding Diracs which results in a more continuous formulation:

Example 3.4. Let $\Omega = [a, b]$, $a < b$: Let $t_k < t_{k+1}$ and $t_1 = a$, $t_n = b$ be a sequence of knots that subdivide the interval Ω into $n - 1$ subintervals $[t_k, t_{k+1}] =: \Omega_k$. We define $\varphi(x) = \alpha e_k + (1 - \alpha)e_{k+1}$, with $e_k \in \mathbb{R}^n$ being the k^{th} unit vector, $\alpha \in [0, 1]$ such that $\alpha t_k + (1 - \alpha)t_{k+1} = x$. This yields a so-called 2-sparse lifting map that has been used in the related work of [21]. It spans the space Λ of univariate piecewise linear functions on Ω and the moment space $\text{con}(\varphi(\Omega)) = \mathcal{P}_\Lambda$ is given by the unit simplex.

For φ_k being chosen as the monomials we obtain the classical notion of moments:

Example 3.5. For $\varphi_0 = 1$, the space of univariate polynomials $\Lambda = \mathbb{R}[x]$ with maximum degree n is spanned by the monomials:

$$\varphi(x) = (x, x^2, \dots, x^n), \quad (22)$$

and $\text{con}(\varphi(\Omega)) = \mathcal{P}_\Lambda$ is the monomial moment space.

Example 3.6. Identifying $\Omega = \{x \in \mathbb{R}^2 : \|x\|_2 = 1\}$ with the complex unit circle $\Omega \cong \{z \in \mathbb{C} : |z| = 1\}$ and again assuming $\varphi_0 = 1$, the mapping

$$\varphi(z) = (\text{Re}(z), \text{Im}(z), \dots, \text{Re}(z^n), \text{Im}(z^n)) \in \mathbb{R}^{2n} \quad (23)$$

spans the space Λ of real trigonometric polynomials of maximum degree n . Parametrizing elements $z \in \Omega$ via the bijection between $z = e^{i\omega}$ and its angle $\omega \in [0, 2\pi)$ the components of φ are the Fourier basis functions, which define the Carathéodory curve [7]. The convex hull $\text{con}(\varphi(\Omega)) = \mathcal{P}_\Lambda$ is the trigonometric moment space.

3.2 Extremal Subspaces

In contrast to the piecewise linear lifting, the polynomial lifting is extremal in the sense that no Dirac measure on the monomial moment curve can be expressed as a convex combination of other Diracs, which sets up an one-to-one correspondence between Diracs δ_x and lifted points $\varphi(x)$. This is illustrated in Fig. 1.

More formally, we call a lifting map φ an extremal curve if each $y \in \varphi(\Omega)$ is an extreme point of $\text{con}(\varphi(\Omega))$ defined according to [31, Sec. 18].

Definition 3.7 (extreme points). *Let C be a convex set and $y \in C$. Then y is called an extreme point of C if there is no way to express y as a convex combination $y = (1 - \alpha)x + \alpha z$ of $x, z \in C$ and $0 < \alpha < 1$, except by taking $y = x = z$.*

Definition 3.8 (extremal moment curve). *Let Ω be compact and be nonempty. Then we say the mapping $\varphi : \Omega \rightarrow \mathbb{R}^n$ is an extremal moment curve if φ is a lifting map and any point $y \in \varphi(\Omega) \subset \mathbb{R}^n$ is an extreme point of $\text{con} \varphi(\Omega)$.*

Via a change of basis it becomes clear that the definition of extremality is independent of a specific choice of a basis for Λ . Therefore extremality is rather a property of the subspace Λ . This also motivates the following lemma which shows that extremality is inherited along a hierarchy $\Theta \subset \Lambda \subset \mathcal{C}(\Omega)$.

Lemma 3.9 (extremal subspaces). *Let $\emptyset \neq \Omega \subset \mathbb{R}^m$ be compact. Let $\Theta \subset \Lambda \subset \mathcal{C}(\Omega)$ be a hierarchy of finite-dimensional subspaces of the space of continuous functions $\mathcal{C}(\Omega)$. Let $\Theta = \langle \theta_1, \dots, \theta_n \rangle$ such that $\theta : \Omega \rightarrow \mathbb{R}^n$ is an extremal curve. Then Λ is spanned by an extremal curve as well.*

Proof. $\{\theta_1, \dots, \theta_n\}$ is a basis of $\Theta \subset \Lambda$ and therefore linearly independent. Since Λ is finite-dimensional in view of the basis extension theorem $\{\theta_1, \dots, \theta_n\}$ can be extended to a basis $\varphi = (\theta_1, \dots, \theta_n, \psi_1, \dots, \psi_k)$ of Λ with vectors $\psi_i \in G$, where $\text{span } G = \Lambda$ and $|G| < \infty$ such that $\text{span } \varphi = \Lambda$.

Now choose $y \in \Omega$ and consider $\varphi(y)$. Let $\alpha \in (0, 1)$ and $\varphi(y) = \alpha x + (1 - \alpha)z$ for $x, z \in \text{con } \varphi(\Omega)$. Then there exist coefficients $\alpha_i, \beta_k > 0$ such that $x = \sum_{i=1}^m \alpha_i \varphi(x^{(i)})$ and $z = \sum_{k=1}^l \beta_k \varphi(z^{(k)})$, $z^{(k)}, x^{(i)} \in \Omega$ with $\sum_{i=1}^m \alpha_i = 1$, $\sum_{k=1}^l \beta_k = 1$. This implies that $\theta(y) = \alpha \sum_{i=1}^m \alpha_i \theta(x^{(i)}) + (1 - \alpha) \sum_{k=1}^l \beta_k \theta(z^{(k)})$. Extremality of θ implies that $\theta(y) = \sum_{i=1}^l \alpha_i \theta(x^{(i)}) = \sum_{k=1}^l \beta_k \theta(z^{(k)})$ and therefore $\theta(y) = \theta(x^{(i)}) = \theta(z^{(k)})$. Since θ is an extremal curve it is injective and therefore $y = z^{(k)} = x^{(i)}$. This implies that $\varphi(y) = x = z$. \square

Lemma 3.10 (extremality of quadratic subspace). *Let $\emptyset \neq \Omega \subset \mathbb{R}^m$ be compact. Let $\varphi(x) = (x_1, x_2, \dots, x_m, \|x\|^2)$. Assume that $\{x \mapsto \langle a, \varphi(x) \rangle : a \in \mathbb{R}^{m+1}\} \subset \Lambda \subset \mathcal{C}(\Omega)$. Then Λ is spanned by an extremal curve.*

Proof. Now choose $y \in \Omega$. Let $\alpha \in (0, 1)$ and $\varphi(y) = \alpha x + (1 - \alpha)z$ for $x, z \in \text{con } \varphi(\Omega)$. Then there exist coefficients $\alpha_i, \beta_k > 0$ such that $x = \sum_{i=1}^m \alpha_i \varphi(x^{(i)})$ and $z = \sum_{k=1}^l \beta_k \varphi(z^{(k)})$, $z^{(k)}, x^{(i)} \in \Omega$ with $\sum_{i=1}^m \alpha_i = 1$, $\sum_{k=1}^l \beta_k = 1$.

Now choose $f(x) = \|x - y\|^2 = \|x\|^2 - 2\langle y, x \rangle + \|y\|^2$. Then we have for $a = (-2y, 1)$ and $a_0 = \|y\|^2$:

$$\begin{aligned} 0 = f(y) &= \langle \varphi(y), a \rangle + a_0 = \left\langle \alpha \sum_{i=1}^m \alpha_i \varphi(x^{(i)}) + (1 - \alpha) \sum_{k=1}^l \beta_k \varphi(z^{(k)}), a \right\rangle + a_0 \\ &= \sum_{i=1}^m \alpha \cdot \alpha_i \cdot \langle \varphi(x^{(i)}), a \rangle + \sum_{k=1}^l (1 - \alpha) \cdot \beta_k \cdot \langle \varphi(z^{(k)}), a \rangle + a_0 \\ &= \sum_{i=1}^m \alpha \cdot \alpha_i \cdot f(x^{(i)}) + \sum_{k=1}^l (1 - \alpha) \cdot \beta_k \cdot f(z^{(k)}) \end{aligned}$$

As $f(x^{(i)}) > 0$ for $x^{(i)} \neq y$ and $\alpha > 0$ it holds that $x^{(i)} \neq y$ implies $\alpha_i = 0$. The same is true for $z^{(k)}$ and β_k . Hence $x = \varphi(y) = z$, and therefore $\varphi : \Omega \rightarrow \mathbb{R}^{m+1}$ is an extremal curve. In view of Lem. 3.9 Λ is spanned by an extremal curve. \square

This shows that in a piecewise polynomial discretization with degree at least 2 the corresponding basis inherits the extreme point property from the extremality of the subspace of quadratic functions. Extremal curves are key to preserve the cost function when restricted to the set of discretized Diracs $\varphi(x)$:

Theorem 3.11. *Let $\Omega \subset \mathbb{R}^m$ be nonempty and compact and let $f : \Omega \rightarrow \mathbb{R}$ be lsc. Furthermore, let $\varphi : \Omega \rightarrow \mathbb{R}^n$ be an extremal curve. Then we have*

$$f_{\Lambda}^{**} \circ \varphi = f \tag{24}$$

on Ω . In addition, we have that $\text{con } f_{\Lambda} = f_{\Lambda}^{**}$.

Proof. Since f is lsc and Ω compact f is bounded from below, i.e., there is $\gamma > -\infty$ so that $f(x) \geq \gamma$ for all $x \in \Omega$. We have $\text{dom } f_{\Lambda} = \varphi(\Omega) \subset \mathbb{R}^n$ and in view of [33, Prop. 2.31] it holds for any $y \in \mathbb{R}^n$,

$$\begin{aligned} (\text{con } f_{\Lambda})(y) &= \inf \left\{ \sum_{i=1}^{n+1} \lambda_i f_{\Lambda}(y^{(i)}) : \lambda_i \geq 0, \sum_{i=1}^{n+1} \lambda_i = 1, y = \sum_{i=1}^{n+1} \lambda_i y^{(i)}, y^{(i)} \in \mathbb{R}^n \right\} \\ &= \inf \left\{ \sum_{i=1}^{n+1} \lambda_i f(x^{(i)}) : \lambda_i \geq 0, \sum_{i=1}^{n+1} \lambda_i = 1, y = \sum_{i=1}^{n+1} \lambda_i \varphi(x^{(i)}), x^{(i)} \in \Omega \right\} \geq \gamma. \end{aligned}$$

Let $x \in \Omega$. Since the only possible convex combination of the extreme point $\varphi(x)$ from points $y^{(i)} \in \varphi(\Omega)$ is $\varphi(x)$ itself, we have

$$\begin{aligned} (\text{con } f_{\Lambda})(\varphi(x)) &= \inf \left\{ \sum_{i=1}^{n+1} \lambda_i f_{\Lambda}(y^{(i)}) : \lambda_i \geq 0, \sum_{i=1}^{n+1} \lambda_i = 1, \varphi(x) = \sum_{i=1}^{n+1} \lambda_i y^{(i)}, y^{(i)} \in \varphi(\Omega) \right\} \\ &= f_{\Lambda}(\varphi(x)) = f(x). \end{aligned}$$

This shows that $\text{con } f_{\Lambda} \circ \varphi = f$. Since $\text{con } f_{\Lambda}$ is bounded from below $\text{con } f_{\Lambda}$ is proper.




Figure 2: Lifted version f_Λ of the function f for 3 different choices of φ . In the top row the same nonconvex function f (blue curves) is depicted. The colored surfaces in the bottom row correspond to the lifted biconjugates f_Λ^{**} , where the gray shadow areas correspond to their domains $\text{dom } f_\Lambda^{**} = \mathcal{P}_\Lambda$ and the blue curves correspond to the lifted cost f_Λ , see Eq. (14). The black curves in the bottom row correspond to the moment curve described by Diracs $\varphi(\Omega)$, i.e., the domain of f_Λ . From left to right, cf. Ex. 3.3, Ex. 3.4 and Ex. 3.5. The nonlinear supporting dual functions $q_{\lambda,\beta}(x) = \langle \varphi(x), \lambda \rangle - \beta$ (red curves) to f in the top row (middle and right), are transformed into linear supporting hyperplanes $l_{\lambda,\beta}(y) = \langle y, \lambda \rangle - \beta$ (red surfaces) to f_Λ^{**} in the bottom row through the feature map φ . In the language of kernel methods, these functions can be interpreted as the nonlinear decision boundaries that separate individual points on the graph of f from the epigraph of f . Only in the most right case such a separation is possible. As a result the polynomial lifting preserves the nonconvex cost function $f_\Lambda^{**} \circ \varphi = f$ on $\varphi(\Omega)$ whereas the 2-sparse lifting (middle) only leads to a piecewise convex under-approximation $f_\Lambda^{**} \circ \varphi \leq f$.

Let f be lsc. Then f_Λ inherits its lower semicontinuity from f : Assume that $y \in \varphi(\Omega)$. Then there exists $x \in \Omega$ with $y = \varphi(x)$ and we have due to the continuity of φ and φ^{-1} :

$$\begin{aligned}
 \liminf_{y' \rightarrow \varphi(x)} f_\Lambda(y') &:= \lim_{\delta \rightarrow 0^+} \inf \{f_\Lambda(y') : y' \in B_\delta(\varphi(x))\} \\
 &= \lim_{\delta \rightarrow 0^+} \inf \{f_\Lambda(\varphi(x')) : \varphi(x') \in B_\delta(\varphi(x))\} = \lim_{\epsilon \rightarrow 0^+} \inf \{f(x') : x' \in B_\epsilon(x)\} \\
 &= \liminf_{x' \rightarrow x} f(x') \geq f(x) = f_\Lambda(y).
 \end{aligned}$$

We invoke [33, Cor. 3.47] and deduce that $\text{con } f_\Lambda$ is proper, lsc and convex. In view of [33, Thm. 11.1] we have $\text{con } f_\Lambda = \text{cl con } f_\Lambda = f_\Lambda^{**}$. \square

For a geometric intuition of this theorem we refer to Fig. 2.

For a piecewise polynomial discretization with degree at least 2, due to extremality, the primal discretized energy \mathcal{F}_Λ restricted to $\varphi(\Omega)^\mathcal{V}$ agrees with the original energy F . In particular, this implies that an obtained Dirac solution $(\varphi(x_u^*))_{u \in \mathcal{V}}$ of the discretization corresponds to a solution of the original problem in the same way integer solutions of LP relaxations are certificates of optimality for the corresponding ILP.

Proposition 3.12. *Let $\emptyset \neq \Omega \subset \mathbb{R}^m$. Let the metric d be induced by a norm and assume the space of linear functions on Ω is contained in Λ . Furthermore, assume Λ is spanned by an extremal curve $\varphi : \Omega \rightarrow \mathbb{R}^n$ and f_u lsc. Then for any $y \in (\mathbb{R}^n)^\mathcal{V}$, with $y_u = \varphi(x_u)$, $x_u \in \Omega$ the following identity holds true:*

$$\mathcal{F}_\Lambda(y) = F(x).$$

In particular, whenever y^ is a solution of problem (dR-P) such that $y^* \in (\mathbb{R}^n)^\mathcal{V}$, with $y_u^* = \varphi(x_u^*)$, for some $x_u^* \in \Omega$, x^* is a solution of the original problem (P).*

Proof. Let d be induced by some norm $\|\cdot\|$ and denote its dual norm by $\|\cdot\|_*$. As shown in Thm. 3.11, the unaries preserve the original cost functions at $\varphi(\Omega)$. Hence it remains to show that for the pairwise costs it holds $\sigma_{\mathcal{K}}((\nabla_\Lambda y)_{(u,v)}) = \|x_u - x_v\|$. By assumption $(\nabla_\Lambda y)_{(u,v)} = \varphi(x_u) - \varphi(x_v)$. We rewrite

$$\sigma_{\mathcal{K}_\Lambda}(\varphi(x_u) - \varphi(x_v)) = \sup_{\lambda \in \Lambda \cap \text{Lip}_d(\Omega)} \lambda(x_u) - \lambda(x_v) \leq \|x_u - x_v\|.$$

For any $x_u, x_v \in \Omega$ we have

$$\|x_u - x_v\| = \sup_{p \in \mathbb{R}^m: \|p\|_* \leq 1} |\langle x_u - x_v, p \rangle| = \langle x_u - x_v, p^* \rangle,$$

where p^* denotes the maximizer in the supremum, which exists due to the compactness of the unit ball in a finite-dimensional space. Define the linear function $\lambda^* := \langle \cdot, p^* \rangle$ and note that by assumption $\lambda^* \in \Lambda$. In addition we have shown that $\lambda^* \in \text{Lip}_d(\Omega)$ is 1-Lipschitz. This implies $\|x_u - x_v\| = \lambda^*(x_u) - \lambda^*(x_v) \leq \sup_{\lambda \in \Lambda \cap \text{Lip}_d(\Omega)} \lambda(x_u) - \lambda(x_v) \leq \|x_u - x_v\|$ and hence equality holds. \square

3.3 A Generalized Conjugacy Perspective

The previous results can be obtained from a generalized conjugacy point of view. In particular, the convex conjugate of the lifted function f_Λ^* is comprised by the notion of Φ -conjugacy, see [33, Ch. 11L*]:

Definition 3.13 (generalized conjugate functions). *Let X and Y be nonempty sets. Let $\Phi : X \times Y \rightarrow \overline{\mathbb{R}}$ be any function. Let $f : X \rightarrow \overline{\mathbb{R}}$. Then the Φ -conjugate of f on Y at $y \in Y$ is defined by*

$$f^\Phi(y) := \sup_{x \in X} \Phi(x, y) - f(x), \tag{25}$$

and the Φ -biconjugate of f back on X at $x \in X$ is given by

$$f^{\Phi\Phi}(x) := \sup_{y \in Y} \Phi(x, y) - f^\Phi(y). \tag{26}$$

We say that f is a Φ -envelope on X if f can be written in terms of a pointwise supremum of a collection of elementary functions $x \mapsto \Phi(x, y) - \alpha$, where $(\alpha, y) \in \overline{\mathbb{R}} \times Y$ is the parameter element.

Let $g : Y \rightarrow \overline{\mathbb{R}}$. Then, the Φ -conjugate of g on X at $x \in X$ is defined by

$$g^\Phi(x) := \sup_{y \in Y} \Phi(x, y) - g(y), \quad (27)$$

and the Φ -biconjugate of g back on Y at $y \in Y$ is given by

$$g^{\Phi\Phi}(y) := \sup_{x \in X} \Phi(x, y) - g^\Phi(x). \quad (28)$$

We say that g is a Φ -envelope on Y if g can be written in terms of a pointwise supremum of a collection of elementary functions $y \mapsto \Phi(x, y) - \beta$, where $(\beta, x) \in \overline{\mathbb{R}} \times X$ is the parameter element.

For $X = \Omega$, $Y = \mathbb{R}^n$, and $\Phi(x, y) = \langle y, \varphi(x) \rangle$, the convex conjugate f_Λ^* is identical to the Φ -conjugate f^Φ of f , while its biconjugate f_Λ^{**} is the tightest lsc convex extension of f_Λ to $\text{con } \varphi(\Omega)$. The Φ -biconjugate $f^{\Phi\Phi}$ of f at a point $x \in \Omega$ is equal to the classical biconjugate of f_Λ , evaluated at $\varphi(x)$, i.e., $f^{\Phi\Phi} = f_\Lambda^{**} \circ \varphi$ on Ω , showing that the Φ -biconjugate is a convexly composite function and, therefore, it is nonconvex in general. Actually, Φ -conjugacy also comprises lifting to measures via $\varphi(x) = \delta_x$, Φ the corresponding dual pairing and $Y = \mathcal{C}(\Omega)$.

As a consequence of [33, Ex. 11.63], the considered Φ -conjugacy can be interpreted in terms of under-approximation by functions in Λ . In analogy to the biconjugate f^{**} , which is the pointwise supremum of affine-linear functions majorized by f , the Φ -biconjugate $f^{\Phi\Phi}$ is the pointwise supremum of functions in Λ up to constant translation majorized by f . This point of view also relates $(f^\Phi)^*$ and $f^{\Phi\Phi}$ by each other.

Remark 3.14. The function $(f^\Phi)^* = f_\Lambda^{**}$ is the pointwise supremum of all affine-linear functions $l_{\lambda, \beta} := \langle \cdot, \lambda \rangle - \beta$ for which $q_{\lambda, \beta} := \Phi(\cdot, \lambda) - \beta$ is majorized by f . This can be seen as follows: The Legendre–Fenchel conjugate can be characterized via the identity $(f^\Phi)^*(y) = \sup_{(\lambda, \beta) \in \text{epi}(f^\Phi)} \langle y, \lambda \rangle - \beta$. The observation now follows from the fact that $(\lambda, \beta) \in \text{epi}(f^\Phi)$ if and only if $q_{\lambda, \beta}$ is majorized by f .

The correspondence between f_Λ^{**} and $(f^\Phi)^*$ and between the minorizers $l_{\lambda, \beta}$ and $q_{\lambda, \beta}$ is illustrated in Fig. 2. Note that this is closely related to the idea of feature maps φ in linear classifiers.

Thm. 3.11 identifies all lsc functions f as Φ -envelopes whenever φ is extremal:

Corollary 3.15. *Let $\Omega \subset \mathbb{R}^m$ be nonempty and compact and let $f : \Omega \rightarrow \mathbb{R}$ be bounded from below. Furthermore, let $\varphi : \Omega \rightarrow \mathbb{R}^n$ be an extremal curve. Then we have*

$$f^{\Phi\Phi} = f_\Lambda^{**} \circ \varphi = \text{cl } f, \quad (29)$$

on Ω .

Proof. Since f is finite-valued and bounded from below on Ω we have $(\text{cl } f)(x) > -\infty$ for $x \in \Omega$ and therefore $\text{cl } f$ is finite-valued. By [33, Ex. 11.63] $f^{\Phi\Phi}$ is the largest Φ -envelope below f . Since φ is continuous relative to Ω , $f^{\Phi\Phi} = f_\Lambda^{**} \circ \varphi$ is lsc relative to Ω . Since $\text{cl } f$ is the largest

lsc function below f we have $f^{\Phi\Phi} \leq \text{cl } f$. Since $\text{cl } f \leq f$ we also have $(\text{cl } f)^{\Phi\Phi} \leq f^{\Phi\Phi}$. Invoking Thm. 3.11 we have

$$\text{cl } f = (\text{cl } f)^{\Phi\Phi} \leq f^{\Phi\Phi} \leq \text{cl } f,$$

on Ω . Therefore $f^{\Phi\Phi} = \text{cl } f$ on Ω . \square

Up to the presence of the compact set Ω , this result generalizes the basic quadratic transform [33, Ex. 11.66] originally due to [27, Prop. 3.4] (for lsc functions only), which is obtained by choosing $\varphi(x) = (x_1, x_2, \dots, x_m, \|x\|^2)$.

4 A Tractable Conic Program for MRFs

4.1 Nonnegativity and Moments

After discretization a next step to obtain a practical implementation is to derive finite characterizations of the lifted biconjugates f_Λ^{**} and the constraint set \mathcal{K}_Λ . We will show that the formulations can be rewritten in terms of a semi-infinite conic program which can be implemented using semidefinite programming in the piecewise polynomial case.

For now let $f_u \in \Lambda$. Eq. (21) then shows, that the challenging part is to characterize the moment space \mathcal{P}_Λ : The following result shows that up to normalization, \mathcal{P}_Λ can be written in terms of the dual cone of the cone of functions in Λ that are nonnegative on Ω .

Lemma 4.1. *Let $(\mathcal{M}_\Lambda)_+$ be the cone of moments of nonnegative measures as defined in (12) and let \mathcal{N}_Λ be the cone of the coefficients of the functions in $\Lambda = \langle \varphi_0, \dots, \varphi_n \rangle$ that are nonnegative on Ω defined as:*

$$\mathcal{N}_\Lambda := \{p \in \mathbb{R}^{n+1} : \langle p, \varphi(x) \rangle \geq 0, \forall x \in \Omega\}. \quad (30)$$

Then $(\mathcal{M}_\Lambda)_+$ is equal to \mathcal{N}_Λ^ , where \mathcal{N}_Λ^* denotes the dual cone of \mathcal{N}_Λ .*

If, in addition, $\varphi_0 \equiv 1$ we also have $\mathcal{P}_\Lambda = \{y \in (\mathcal{M}_\Lambda)_+ : y_0 = 1\}$.

Proof. Let $y \in (\mathcal{M}_\Lambda)_+$. This means there exists $\mu \in \mathcal{M}_+(\Omega)$ such that $y_k := \int_\Omega \varphi_k(x) d\mu(x)$. Let $p \in \mathcal{N}_\Lambda$. Because of $\langle p, \varphi(\cdot) \rangle \in \Lambda \subset \mathcal{C}(\Omega)$ and $\langle p, \varphi(x) \rangle \geq 0$ for all $x \in \Omega$ and $\mu \in \mathcal{M}_+(\Omega)$ is a nonnegative measure it holds $\langle p, y \rangle = \sum_{k=0}^n p_k \int_\Omega \varphi_k(x) d\mu(x) = \int_\Omega \langle p, \varphi(x) \rangle d\mu(x) \geq 0$ for all $x \in \Omega$. Since $p \in \mathcal{N}_\Lambda$ was an arbitrary choice from \mathcal{N}_Λ we have $y \in \mathcal{N}_\Lambda^*$.

Next we show $(\mathcal{M}_\Lambda)_+^* \subseteq \mathcal{N}_\Lambda$ as this implies $\mathcal{N}_\Lambda^* \subseteq (\mathcal{M}_\Lambda)_+^{**} = (\mathcal{M}_\Lambda)_+$, where the last equality holds since $(\mathcal{M}_\Lambda)_+$ is closed and convex. Take $p \in (\mathcal{M}_\Lambda)_+^*$. Let $x \in \Omega$. Now we choose $y \in (\mathcal{M}_\Lambda)_+$ such that $y_k = \langle \delta_x, \varphi_k \rangle = \varphi_k(x)$. Then $p \in (\mathcal{M}_\Lambda)_+^*$ implies that $\langle p, y \rangle \geq 0$. Since the choice $x \in \Omega$ was arbitrary we have $\langle p, \varphi(x) \rangle \geq 0$ for all $x \in \Omega$ and therefore $p \in \mathcal{N}_\Lambda$.

Finally, $\mathcal{P}_\Lambda = \{y \in (\mathcal{M}_\Lambda)_+ : y_0 = 1\}$ follows from the fact that $\mu \in \mathcal{M}(\Omega)$ is an element of $\mathcal{P}(\Omega)$ if and only if $\mu \in \mathcal{M}_+(\Omega)$ and $\langle \mu, \varphi_0 \rangle = 1$. \square

Before we specialize Λ to the space of polynomials, we derive a cone programming formulation of the Lipschitz constraints $\lambda_{(u,v)} \in \text{Lip}_d(\Omega)$. Here we restrict $\Omega = [a, b]$ to be a compact interval. In the following, we provide an implementation for two specific metrics. As before, this boils down to nonnegativity of functions: Firstly, we consider total variation regularization, i.e. $d(x, y) = |x - y|$: Assume that Λ is closed under differentiation, i.e., φ is differentiable and

$\varphi'_k \in \Lambda$. Then, the condition $\lambda_{(u,v)} \in \text{Lip}_d([a, b])$ can be phrased in terms of the constraints $-1 \leq \lambda'_{(u,v)}(x) \leq 1$ for all $x \in [a, b]$, where $\lambda'_{(u,v)}$ is the derivative of $\lambda_{(u,v)}$. Equivalently, this means that the coefficients of the functions $1 + \lambda'_{(u,v)}$ and $1 - \lambda'_{(u,v)}$ are in \mathcal{N}_Λ . Secondly, we consider Potts regularization, i.e., $d(x, y) = \llbracket x = y \rrbracket$. Since $\lambda_{(u,v)}$ is a univariate function, and constant terms in the dual variable do not matter, the condition $\lambda_{(u,v)} \in \text{Lip}_d([a, b])$ can be equivalently phrased as $0 \leq \lambda_{(u,v)} \leq 1$, see, [43, Ex. 1.17]. Equivalently, this means that the coefficients of the functions $\lambda_{(u,v)}$ and $1 - \lambda_{(u,v)}$ are in \mathcal{N}_Λ .

4.2 Semidefinite Programming and Nonnegative Polynomials

As we have seen in the previous section, an important ingredient for a tractable formulation is the efficient characterization of nonnegativity of functions in a finite-dimensional subspace $\Lambda \subset \mathcal{C}(\Omega)$. A promising choice of Λ in that regards is the space of polynomials. Indeed, the characterization of nonnegativity of polynomials is a fundamental problem in *convex algebraic geometry* surveyed in [5]: Let $\mathbb{R}[x_1, \dots, x_m]$ denote the ring of possibly multivariate polynomials with $p \in \mathbb{R}[x_1, \dots, x_m]$. Then $p = \sum_{\alpha \in I} p_\alpha x^\alpha$ for monomials x^α . Let $\deg p$ denote its degree. A key result from real algebraic geometry is the Positivstellensatz due to [17] and [41] refined in [39] and [29]. It characterizes polynomials $p \in \mathbb{R}[x_1, \dots, x_m]$ that are positive on semi-algebraic sets Ω , i.e. $p(x) > 0$ for all $x \in \Omega$, where Ω is defined in terms of polynomial inequalities.

Key to such results is a *certificate* of nonnegativity of the polynomial p that involves *sum-of-squares* (SOS) multipliers q , where q is SOS if $q(x) = \sum_{i=1}^N q_i^2(x)$ for polynomials $q_i \in \mathbb{R}[x_1, \dots, x_m]$. For intervals $X = [a, b]$, thanks to [5, Thm. 3.72] originally due to [28, Cor. 2.3] we have following result:

Lemma 4.2. *Let $a < b$. Then the univariate polynomial $p \in \mathbb{R}[x]$ is nonnegative on $[a, b]$ if and only if it can be written as*

$$p(x) = \begin{cases} s(x) + (x - a) \cdot (b - x) \cdot t(x) & \text{if } \deg p \text{ is even,} \\ (x - a) \cdot s(x) + (b - x) \cdot t(x) & \text{if } \deg p \text{ is odd,} \end{cases} \quad (31)$$

where $s, t \in \mathbb{R}[x]$ are sum of squares. If $\deg p = 2n$, then we have $\deg s \leq 2n$, $\deg t \leq 2n - 2$, while if $\deg p = 2n + 1$, then $\deg s \leq 2n$, $\deg t \leq 2n$.

Remarkably, the above result provides us with explicit upper bounds of the degrees of the SOS multipliers s and t that are important to derive a practical implementation: Then the SOS constraints can be formulated in terms of semidefinite and affine inequalities: We adopt [5, Lem. 3.33] and [5, Lem. 3.34]:

Lemma 4.3. *A univariate polynomial $p \in \mathbb{R}[x]$ with $\deg p = 2n$, $n \geq 0$ is SOS if and only if there exists a positive semidefinite matrix $Q \in \mathbb{R}^{n+1 \times n+1}$ such that*

$$p_k = \sum_{\substack{0 \leq i, j \leq n, \\ i+j=k}} Q_{ij}, \quad \forall 0 \leq k \leq 2n. \quad (32)$$

Invoking the results above SDP-duality yields the following compact representation of \mathcal{M}_Λ :

Lemma 4.4. *Let $n \geq 0$. For odd degree $2n + 1$, $y \in \mathcal{M}_\Lambda$ if and only if*

$$bM_{0,n}(y) \succeq M_{1,n}(y) \succeq aM_{0,n}(y), \quad (33)$$

for Hankel matrices

$$M_{i,n}(y) := \begin{bmatrix} y_i & y_{i+1} & \cdots & y_{i+n} \\ y_{i+1} & y_{i+2} & \cdots & y_{i+n+1} \\ \vdots & & \cdots & \vdots \\ y_{i+n} & & \cdots & y_{i+2n} \end{bmatrix}. \quad (34)$$

For even degree $2n$, $y \in \mathcal{M}_\Lambda$ if and only if

$$M_{0,n}(y) \succeq 0, \quad (35)$$

$$(a + b) M_{1,n-1}(y) - abM_{0,n-1}(y) \succeq M_{2,n-1}(y). \quad (36)$$

Proof. Follows by Lem. 4.2 and Lem. 4.3 invoking elementary SDP-duality. \square

4.3 A First-Order Primal-Dual Algorithm

We are now ready to describe the algorithm for solving the resulting semidefinite program. We first consider the case $f_u \in \Lambda$ and Λ is the space of univariate polynomials. We propose to use the PDHG [9] algorithm, as it can exploit the partially separable structure of our SDP. The primal-dual algorithm optimizes the problem (dR-P) via alternating projected gradient descent/ascent steps applied to the saddle-point formulation of (dR-P):

$$\min_{y \in (\mathcal{P}_\Lambda)^\mathcal{V}} \max_{p \in (\mathcal{K}_\Lambda)^\mathcal{E}} \langle y, f - \text{Div}_\Lambda p \rangle, \quad (37)$$

which is obtained by expanding the support function in Problem (dR-P) and substituting the expression (21) for the lifted biconjugates. In each iteration the algorithm performs a projected gradient ascent step in the dual p followed by a projected gradient descent step in the primal variable y . Subsequently it performs an extrapolation step in the primal. The projections onto the sets $(\mathcal{P}_\Lambda)^\mathcal{V}$ and $(\mathcal{K}_\Lambda)^\mathcal{E}$ are separable and can therefore be carried out in parallel on a GPU using the SDP characterizations derived above. For practicality, we introduce additional auxiliary variables and linear constraints to decouple the affine constraints (32) and the SDP constraints. The projection operator of the semidefinite cone can then be solved using an eigenvalue decomposition.

4.4 Piecewise Polynomial Duals and Nonlinear Lifted Biconjugates

The polynomial discretization can be extended by means of a continuous piecewise polynomial representation of the dual variables resulting in a possibly more accurate approximation of the dual subspace Λ . Then, both, nonnegativity and Lipschitz continuity can be enforced on each piece Ω_k individually. Continuity of the piecewise polynomial dual variables can be enforced via linear constraints. The corresponding primal variable y belongs to $y \in (\mathcal{M}_\Lambda)_+ \times (\mathcal{M}_\Lambda)_+ \times \cdots \times (\mathcal{M}_\Lambda)_+$. Then the restriction that y is a moment vector of a probability measure supported on the whole space Ω yields an additional sum-to-one constraint on the 0th moments $1 = \sum_{k=1}^K y_{k,0}$.




Figure 3: Primal and dual energies for MAP-inference in a continuous MRF with TV regularization using a piecewise polynomial hierarchy of dual variables. (a) shows the dual energy for TV. In (b) the dashed lines correspond to the primal energy at the rounded solution and the solid lines correspond to the dual energy. (c) shows the gap between the nonconvex primal energy at the rounded solution and the dual energy for TV regularization. (d) shows the dual energies for Potts regularization.

Another issue to address is when $f_u \notin \Lambda$ which results in a nonlinear lifted biconjugate over the moment-space as in Fig. 2.

The formulation which is derived next addresses both: In particular it allows one to choose Λ independently from f_u which can even be discontinuous, as long as f_u has a piecewise polynomial structure. Key to the formulation is to rewrite the inner minimum in the dual formulation dR-D exploiting a duality between nonnegativity and minimization of functions: Let $\Omega = [a, b]$, $a < b$ be a compact interval. Let $a = t_1 < t_2 < t_3, \dots, < t_{K+1} = b$ be a sequence of knots, where $\Omega_k := [t_k, t_{k+1}]$. Let Θ be the space of univariate polynomials with some maximum degree n . Let $f : \Omega \rightarrow \mathbb{R}$ be a possibly discontinuous lsc piecewise polynomial function defined by $f(x) = \min_{1 \leq k \leq K} f_k(x) + \iota_{\Omega_k}(x)$ with $f_k \in \Theta$, i.e., $f_k(x) = \langle \varphi(x), w_k \rangle$ for coefficients $w_k \in \mathbb{R}^{n+1}$, where $\varphi_0 \equiv 1$. First observe the following duality between nonnegativity and minimization of a lsc function:

$$\min_{x \in \Omega} f(x) = \max_{q \in \mathbb{R}} q - \iota_{\mathcal{N}}(f - q),$$

where we denote by $\mathcal{N}(\Omega) = \{\lambda : \Omega \rightarrow \mathbb{R} : \lambda(x) \geq 0, \forall x \in \Omega\}$ the cone of nonnegative lsc functions on Ω . Then we obtain for $Aq = (e_0q, \dots, e_0q)$, where $e_0 = (1, 0, \dots, 0) \in \mathbb{R}^{n+1}$ is the 0th unit vector:

$$\min_{x \in \Omega} f(x) = \max_{q \in \mathbb{R}} q - \iota_{\mathcal{N}(\Omega)}(f - q) \quad (38)$$

$$= \max_{q \in \mathbb{R}} q - \sum_{k=1}^K \iota_{\mathcal{N}_{\Theta}}(w_k - A_k q). \quad (39)$$

Fenchel–Rockafellar duality then yields:

$$\begin{aligned} \min_{x \in \Omega} f(x) &= \min_{y \in (\mathbb{R}^{n+1})^K} \iota_{\{1\}}(A^*y) + \sum_{k=1}^K \langle y_k, w_k \rangle + \iota_{(\mathcal{M}_\Theta)_+}(y_k) \\ &= \min_{\substack{y \in ((\mathcal{M}_\Theta)_+)^K \\ \sum_{k=1}^K y_{k,0} = 1}} \sum_{k=1}^K \langle y_k, w_k \rangle. \end{aligned}$$

This formulation can be substituted in the dual problem (dR-D) and we obtain

$$\sup_{p \in ((\mathbb{R}^n)^K)^\mathcal{E}} \sum_{u \in \mathcal{V}} \min_{\substack{y \in ((\mathcal{M}_\Theta)_+)^K \\ \sum_{k=1}^K y_{k,0} = 1}} \sum_{k=1}^K \langle y_k, (w - \text{Div}_\Lambda p)_{u,k} \rangle - \sum_{e \in \mathcal{E}} \iota_{\mathcal{K}}(p_e). \quad (40)$$

Here the dual variables λ are chosen such that $(\text{Div}_\Lambda p)_u$ represents a piecewise polynomial with knots $a = t_1 < t_2 < t_3 < \dots < t_{K+1} = b$ such that for each piece we have $\langle (\text{Div}_\Lambda p)_{u,k}, \varphi(\cdot) \rangle \in \Theta$. Note that this does not require Λ to be equal to the whole space of continuous piecewise polynomials of degree n . Indeed, Λ can be a subspace thereof which covers the case where $f_u \notin \Lambda$.

5 Numerical Experiments

5.1 Empirical Convergence Study

In this first experiment we evaluate the local marginal polytope relaxation of the MRF formulation (P) using a piecewise polynomial hierarchy of dual variables. We choose a vertex grid of size 16×16 . We fix a random polynomial data term of degree 4 at each vertex by fitting a random sample of data points. To obtain a high-accuracy solution we solve the primal SDP formulation corresponding to the saddle-point formulation (40) with MOSEK¹. For recovering a primal solution at each vertex u we compute the mode w.r.t. the 0th moments to select the best interval denoted by $k^* = \arg \max_{1 \leq k \leq K} (y_u)_{k,0}$. Then we compute the mean of the discretized measure corresponding to the $(k^*)^{\text{th}}$ interval as $x_u = (y_u)_{k^*,1}$.

Fig. 3 visualizes the primal and dual energies for varying degrees and/or number of pieces of the dual variable. While the dual energy strictly increases with higher degrees and/or number of pieces the primal energy is evaluated at the rounded solution and therefore does not strictly decrease in general. While for TV increasing the degree vs. increasing the number of pieces (for $K \cdot \text{deg}$ constant) leads to similar performance, for Potts, in many situations, increasing the degree leads to larger dual energies, e.g., consider $\text{deg} = 4, K = 1$ vs. $\text{deg} = 1, K = 4$, red curve vs. blue curve in Fig. 3(d). In further experiments, we observed, that this holds in particular when the structure of the dual variables and the unaries match, i.e., $f_u, \lambda_e \in \Lambda$. Note that for Potts, since the dual variables are uniformly bounded on Ω and the derivative can be unbounded we drop the continuity constraint which leads to a more compact formulation and larger dual energies.

¹<https://www.mosek.com/products/academic-licenses>




Figure 4: Stereo disparity estimation from a stereo image pair: Left: Standard MRF/OT discretization implemented using a continuous piecewise linear under-approximation for the unaries and piecewise linear duals. middle: piecewise linear duals. right: piecewise polynomial duals. The visual appearance of the solution to the standard MRF/OT discretization shows a strong grid bias. The dual energy gap increases for increasing the degree and/or the number of pieces. Likewise the energy at the rounded solution decreases.

5.2 Stereo Matching

In this experiment we consider stereo matching using the anisotropic relaxation (R-P). We consider the Motorcycle and the Piano image pairs from the Middlebury stereo benchmark [38]. We downsample the images by factor 4. The disparity cost term is first calculated using 135 discrete disparities obtained by shifting the images by the corresponding amount of pixels and comparing the image gradients. Then, the cost data term is approximated from below in terms of a continuous piecewise cubic polynomial f_u using 30 pieces at each $u \in \mathcal{V}$. In Fig. 4 we compare the standard MRF/OT discretization as described in Ex. 3.3 with our framework using piecewise linear and piecewise quadratic dual variables. The standard MRF/OT discretization is equivalent to a piecewise linear approximation of the data term with piecewise linear duals in our framework. For a fair comparison we use a continuous piecewise linear under-approximation of the continuous piecewise cubic polynomial under-approximation f_u . As the resulting optimization problem is large-scale we solve the saddle-point formulation (37) with PDHG [9] as described in Sec. 4.3 using the GPU-based PDHG framework `prost`². In contrast to the previous experiment which uses a combined mode and mean rounding procedure we found the plain mean of the discretized measure to produce better results on real data: More explicitly we recover a solution according to $x_u = \sum_{k=1}^K t_k(y_u)_{k,0}$ at each vertex $u \in \mathcal{V}$. In Tab. 1 we compare both,

²<https://github.com/tum-vision/prost>

Dual energies				Energies rounded			
deg	$K = 1$	$K = 3$	$K = 5$	deg	$K = 1$	$K = 3$	$K = 5$
1	14180.08	15733.71	16227.80	1	28982.45	25005.05	22283.26
2	15052.18	16430.97	16773.75	2	31038.14	22525.12	21049.32
3	15601.89	16778.87	17055.25	3	28505.41	21500.65	20428.34
4	15938.92	16998.63	17235.21	4	27255.48	20841.62	20049.04
5	16191.40	17147.92	17346.49	5	25795.56	20344.17	19764.23
6	16369.95	17243.47	17422.26	6	24081.51	20032.77	19559.00
7	16480.22	17308.91	17472.13	7	23142.67	19869.71	19428.50

Table 1: Energies for stereo matching Motorcycle. Left: Dual energies. Right: Primal energies at the rounded solution.

dual and nonconvex primal energies, for a larger hierarchy of dual subspaces.

6 Discussion

We presented a method to reduce duality gaps in the Lagrangian relaxation of the MAP-inference problem in a continuous MRF taking a nonlinear optimization-driven approach. Our theoretical contribution identifies *extremality* of the lifting as a key component, as it leads to convexifications which do not discard any information for a wide range of nonconvex cost functions. Using results from convex algebraic geometry we provide, to our knowledge, the first tractable formulation of the polynomial discretization in terms of semidefinite programming. We have provided a parallel implementation of a first-order primal dual algorithm on a GPU which can handle large problems. Indeed, the approach of [18, 19] applied directly to the original problem (P) (with f_u, f_{uv} polynomial) attempts to solve the *full* marginal polytope relaxation which is tight but intractable for large \mathcal{V} as the number of coupling moments grows exponentially with $|\mathcal{V}|$. In contrast, our framework applies to the *local* marginal polytope relaxation which is not tight in general but leads to a tractable formulation as it exploits the sparse structure of the optimization problem.

As revealed by our experiments, both increasing the number of pieces and the degree of the dual variables successfully reduces the nonconvex duality gap. In particular, the algorithm is applied to the stereo matching problem between two images, showing significant improvements over piecewise constant or piecewise linear discretizations in the dual which demand a higher number of samples.

For total variation regularized problems, our theory suggests that the duality gap vanishes like $\mathcal{O}(1/(K \cdot \sqrt{\text{deg}}))$. The faster convergence in the number of pieces K is also confirmed by our experiments. However, setting the degree $\text{deg} \geq 2$ has the attractive theoretical property that all information of the original nonconvex cost is preserved. For Potts regularization, our experiments indicate that increasing the degree leads to better results than increasing the pieces.

Going beyond the piecewise-linear setting might be particularly promising for vector- or manifold-valued label-spaces with $\dim(\Omega) > 1$. Labeling problems in such higher-dimensional settings were recently considered in [21, 24, 46]. There, due to the piecewise-linear dual variables,

the complexity grows exponentially with $\dim(\Omega)$. Moreover, as discussed in [46], the piecewise-approach inherently requires an approximation of Ω by a triangulated manifold. Building upon the techniques developed in this paper, an interesting direction for future work is to investigate whether these two drawbacks can be overcome by considering suitably defined approximation spaces for the dual variables in vector- or manifold-valued settings.

Acknowledgments

We would like to thank Johannes Milz, Peter Ochs and Jan-Hendrik Lange for their valuable feedback on an early version of this manuscript.

References

- [1] E. J. Balder. An extension of duality-stability relations to nonconvex optimization problems. *SIAM J. Control Optim.*, 15(2):329–343, 1977.
- [2] R. Bergmann, R. H. Chan, R. Hielscher, J. Persch, and G. Steidl. Restoration of manifold-valued images by half-quadratic minimization. *arXiv preprint arXiv:1505.07029*, 2015.
- [3] F. Bernard, F. R. Schmidt, J. Thunberg, and D. Cremers. A combinatorial solution to non-rigid 3d shape-to-image matching. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2017.
- [4] D. Bertsekas. *Nonlinear Programming*. Athena Scientific, 1999.
- [5] G. Blekherman, P. A. Parrilo, and R. R. Thomas. *Semidefinite Optimization and Convex Algebraic Geometry*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2012.
- [6] B. Brown, D. Elliott, and D. Paget. Lipschitz constants for the Bernstein polynomials of a Lipschitz continuous function. *Journal of approximation theory*, 49(2):196–199, 1987.
- [7] C. Carathéodory. Über den Variabilitätsbereich der Fourier’schen Konstanten von positiven harmonischen Funktionen. *Rendiconti Del Circolo Matematico di Palermo (1884-1940)*, 32(1):193–217, 1911.
- [8] N. L. Carothers. A short course on approximation theory. 1998.
- [9] A. Chambolle and T. Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of mathematical imaging and vision*, 40(1):120–145, 2011.
- [10] Q. Chen and V. Koltun. Full flow: Optical flow estimation by global optimization over regular grids. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2016.
- [11] T. Chen, J.-B. Lasserre, V. Magron, and E. Pauwels. Semialgebraic optimization for Lipschitz constants of ReLU networks. In *Conference on Neural Information Processing Systems*, 2020.

- [12] D. J. Crandall, A. Owens, N. Snavely, and D. P. Huttenlocher. SfM with MRFs: Discrete-continuous optimization for large-scale structure from motion. *IEEE Trans. Pattern Anal. Mach. Intell. (PAMI)*, 35(12):2841–2853, 2012.
- [13] C. Domokos, F. R. Schmidt, and D. Cremers. MRF optimization with separable convex prior on partially ordered labels. In *European Conference on Computer Vision (ECCV)*, 2018.
- [14] A. Fix and S. Agarwal. Duality and the continuous graphical model. In *European Conference on Computer Vision (ECCV)*, 2014.
- [15] L. V. Kantorovich. Mathematical methods of organizing and planning production. *Management Science*, 6(4):366–422, 1960.
- [16] J. Kappes, B. Andres, F. Hamprecht, C. Schnorr, S. Nowozin, D. Batra, S. Kim, B. Kausler, J. Lellmann, N. Komodakis, et al. A comparative study of modern inference techniques for discrete energy minimization problems. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2013.
- [17] J.-L. Krivine. Anneaux préordonnés. *Journal d’analyse mathématique*, 12(1):307–326, 1964.
- [18] J. B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM Journal on Optimization*, 11(3):796–817, 2001.
- [19] J. B. Lasserre. Semidefinite programming vs. LP relaxations for polynomial programming. *Mathematics of Operations Research*, 27(2):347–360, 2002.
- [20] F. Latorre, P. T. Y. Rolland, and V. Cevher. Lipschitz constant estimation for neural networks via sparse polynomial optimization. In *8th International Conference on Learning Representations*, 2020.
- [21] E. Laude, T. Möllenhoff, M. Moeller, J. Lellmann, and D. Cremers. Sublabel-accurate convex relaxation of vectorial multilabel energies. In *European Conference on Computer Vision (ECCV)*, 2016.
- [22] C. Lemaréchal. Lagrangian relaxation. In *Computational Combinatorial Optimization*, pages 112–156. Springer, 2001.
- [23] T. Möllenhoff and D. Cremers. Sublabel-accurate discretization of nonconvex free-discontinuity problems. In *International Conference on Computer Vision (ICCV)*, 2017.
- [24] T. Möllenhoff and D. Cremers. Lifting vectorial variational problems: a natural formulation based on geometric measure theory and discrete exterior calculus. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2019.
- [25] J. J. Moreau. Inf-convolution, sous-additivité, convexité des fonctions numériques. *Journal de Mathématiques Pures et Appliquées*, (47):33–41, 1970.
- [26] J. Peng, T. Hazan, D. McAllester, and R. Urtasun. Convex max-product algorithms for continuous MRFs with applications to protein folding. In *International Conference on Machine Learning (ICML)*, 2011.

- [27] R. Poliquin. Subgradient monotonicity and convex functions. *Nonlinear Analysis: Theory, Methods & Applications*, 14(4):305–317, 1990.
- [28] V. Powers and B. Reznick. Polynomials that are positive on an interval. *Transactions of the American Mathematical Society*, 352(10):4677–4692, 2000.
- [29] M. Putinar. Positive polynomials on compact semi-algebraic sets. *Indiana University Mathematics Journal*, 42(3):969–984, 1993.
- [30] R. T. Rockafellar. Duality and stability in extremum problems involving convex functions. *Pacific J. Math.*, 21(1):167–187, 1967.
- [31] R. T. Rockafellar. *Convex Analysis*. Princeton University Press, New Jersey, 1970.
- [32] R. T. Rockafellar. Augmented Lagrange multiplier functions and duality in nonconvex programming. *SIAM J. Control Optim.*, 12(2):268–285, 1974.
- [33] R. T. Rockafellar and R. J.-B. Wets. *Variational Analysis*. Springer, New York, 1998.
- [34] W. Rudin. Functional analysis. *Internat. Ser. Pure Appl. Math*, 1991.
- [35] N. Ruozi. Exactness of approximate MAP inference in continuous MRFs. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2015.
- [36] M. Salzmann. Continuous inference in graphical models with polynomial energies. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2013.
- [37] F. Santambrogio. *Optimal Transport for Applied Mathematicians*. Birkhäuser, New York, 2015.
- [38] D. Scharstein, H. Hirschmüller, Y. Kitajima, G. Krathwohl, N. Nešić, X. Wang, and P. Westling. High-resolution stereo datasets with subpixel-accurate ground truth. In *German conference on pattern recognition*, pages 31–42. Springer, 2014.
- [39] K. Schmüdgen. The k -moment problem for compact semi-algebraic sets. *Mathematische Annalen*, 289(1):203–206, 1991.
- [40] M. Simões, A. Themelis, and P. Patrinos. Lasry-lions envelopes and nonconvex optimization: A homotopy approach. *arXiv preprint arXiv:2103.08533*, 2021.
- [41] G. Stengle. A Nullstellensatz and a Positivstellensatz in semialgebraic geometry. *Mathematische Annalen*, 207(2):87–97, 1974.
- [42] H. Trinh and D. McAllester. Particle-based belief propagation for structure from motion and dense stereo vision with unknown camera constraints. In *International Workshop on Robot Vision*, pages 16–28. Springer, 2008.
- [43] C. Villani. *Topics in optimal transportation*. Number 58. American Mathematical Soc., 2003.
- [44] C. Villani. *Optimal Transport: Old and New*. Springer, 2008.

- [45] T. Vogt, R. Haase, D. Bednarski, and J. Lellmann. On the connection between dynamical optimal transport and functional lifting. *arXiv:2007.02587*, 2020.
- [46] T. Vogt, E. Strekalovskiy, D. Cremers, and J. Lellmann. Lifting methods for manifold-valued variational problems. In *Handbook of Variational Methods for Nonlinear Geometric Data*, pages 95–119. Springer, 2020.
- [47] M. J. Wainwright, M. I. Jordan, et al. Graphical models, exponential families, and variational inference. *Foundations and Trends[®] in Machine Learning*, 1(1–2):1–305, 2008.
- [48] Y. Wald and A. Globerson. Tightness results for local consistency relaxations in continuous MRFs. In *Conference on Uncertainty in Artificial Intelligence (UAI)*, 2014.
- [49] S. Wang, A. Schwing, and R. Urtasun. Efficient inference of continuous Markov random fields with polynomial potentials. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2014.
- [50] A. Weinmann, L. Demaret, and M. Storath. Total variation regularization for manifold-valued data. *SIAM J. Imaging Sci.*, 7(4):2226–2257, 2014.
- [51] A. Weinmann, L. Demaret, and M. Storath. Mumford–Shah and Potts regularization for manifold-valued data. *Journal of Mathematical Imaging and Vision*, 55(3):428–445, 2016.
- [52] T. Werner. A linear programming approach to max-sum problem: A review. *IEEE Trans. Pattern Anal. Mach. Intell. (PAMI)*, 29(7):1165–1179, 2007.
- [53] K. Yamaguchi, T. Hazan, D. McAllester, and R. Urtasun. Continuous Markov random fields for robust stereo estimation. In *European Conference on Computer Vision (ECCV)*, 2012.
- [54] C. Zach. Dual decomposition for joint discrete-continuous optimization. In *International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2013.
- [55] C. Zach and P. Kohli. A convex discrete-continuous approach for Markov random fields. In *European Conference on Computer Vision (ECCV)*, 2012.