MinP Score Tests with an Inequality Constrained Parameter Space

Giuseppe Cavaliere^a, Zeng-Hua Lu^b, Anders Rahbek^c, Yuhong Yang^d

^aDepartment of Economics, University of Exeter Business School, Exeter EX4 4PU, UK And Department of Economics, University of Bologna, 40126 Bologna, Italy

^bUniversity of South Australia Business School, Adelaide, Australia

^cDepartment of Economics, University of Copenhagen, 1353, Copenhagen K, Denmark

^dSchool of Statistics, University of Minnesota, Minneapolis, USA

Abstract

Score tests have the advantage of requiring estimation alone of the model restricted by the null hypothesis, which often is much simpler than models defined under the alternative hypothesis. This is typically so when the alternative hypothesis involves inequality constraints. However, existing score tests address only jointly testing all parameters of interest; a leading example is testing all ARCH parameters or variances of random coefficients being zero or not. In such testing problems rejection of the null hypothesis does not provide evidence on rejection of specific elements of parameter of interest. This paper proposes a class of one-sided score tests for testing a model parameter that is subject to inequality constraints. Proposed tests are constructed based on the minimum of a set of *p*-values. The minimand includes the *p*-values for testing individual elements of parameter of interest using individual scores. It may be extended to include a *p*-value of existing score tests. We show that our tests perform better than/or perform as good as existing score tests in terms of joint testing, and has furthermore the added benefit of allowing for simultaneously testing individual elements of parameter of interest. The added benefit is appealing in the sense that it can identify a model without estimating it. We illustrate our tests in linear regression models, ARCH and random coefficient models. A detailed simulation study is provided to examine the finite sample performance of the proposed tests and we find that our tests perform well as expected. JEL classification: C12.

Keywords: Combined tests, Model identification, Multiple testing, One-sided tests

1. Introduction

Score tests were originally proposed by Rao (1948). They are also known as Lagrange multiplier (LM) tests due to the work of Aitchison and Silvey (1958) and Silvey (1959). Score tests have the advantage of requiring only estimate of the model restricted by the null hypothesis, which often is much simpler than models defined under the alternative hypothesis. This advantage becomes more attractive when models defined under the alternative hypothesis are subject to inequality constraints that complicate model estimation and statistical inference. One-sided tests are more appropriate than two-sided tests when dealing with inequality constraint (see, e.g., Andrews (2001), Francq and Zakoian (2009), Ketz (2018) and Cavaliere et al. (2020) and the early literature therein). One-sided score tests have been studied by many authors, including Gouriéroux et al. (1982), Wolak (1989a,b), Lee and King (1993), Silvapulle and Silvapulle (1995), King and Wu (1997), Demos and Sentana (1998) and Andrews (2001). However, existing one-sided score tests are designed for jointly testing on the parameter of interest. For example, they can be used for testing the null hypothesis of no autoregressive conditional heteroskedasticity (ARCH) effects by assessing whether *all* the ARCH parameters are *jointly* zero.

This paper proposes what we term 'MinP score' tests, where the test statistic is the minimum of a set of *p*-values. The minimand includes the *p*-values for testing individual elements of parameter of interest using individual scores. Importantly, the set may be extended to include a *p*-value of existing score tests. We shall show that, compared with existing tests, our tests have a competitive power. In addition, our test can have the further benefit of allowing, under some regularity conditions, for simultaneously testing individual components of the parameter of interest. For instance, when testing for ARCH the practitioner is able to detect which individual ARCH parameters are non-zero. In contrast, rejection of non-ARCH effects by existing tests does not reveal the evidence of which of the individual ARCH parameters are non-zero. This added benefit is appealing in particular as it allows for model identification within nested models through testing, without estimating candidate models except the model defined under the null hypothesis. To the best of our knowledge, no existing tests allow for model identification using only the restricted estimate under the null hypothesis. The competitiveness of our proposed tests with regard to joint testing on the parameter of interest is supported by the admissibility property we establish. Our admissibility result is applicable to a sort of combined tests for one-sided testing that includes the extended MaxT (EMaxT) tests recently proposed by Lu (2016). With regard to testing individual components of the parameter of interest, it is important to take into account multiplicity of Type I error (cf. Romano et al., 2010). We adopt the family wise error rate (FWER), i.e. the probability of wrongly rejecting at least one true individual hypotheses, for the Type I error control and show that the FWER control can be achieved under certain condition.

We illustrate our tests using three examples, all relevant in applications: linear regression models, ARCH models and random coefficient models. In the linear regression model example, we show that our tests can be used for selecting variables that only exert non-negative marginal effects on the dependent variable, as well as jointly testing if any such effects exist. In this application, the FWER is asymptotically controlled under conditions required for usual Type I error control in joint testing. Variable selection through testing in linear regression models has gained much interest in the recent literature (see, e.g., McKeague and Qian (2015) and related discussion papers). Our tests add contributions to the literature. In the context of ARCH (and random coefficient) models, we show our tests may be used for testing ARCH (and random coefficient) effects and selecting ARCH (and random coefficient) models if non-ARCH (non-random coefficient) effect is rejected. We provide sufficient conditions that lead to the asymptotic control of FWER in ARCH (and random coefficient) models. Simulation studies are carried out to compare finite sample performance of our tests with existing one-sided score tests in terms of joint testing, as well as the performance in model identification.

Structure of the paper

The remainder of the paper is organized as follows. In the next section we set up a general framework under maximisation of an objective function and introduce some assumptions. We provide a result of the limiting normal distribution for function of scores that serves as a basis for constructing our tests. Section 3 presents our MinP score tests and focus on illustrating them for joint testing. Section 4 studies multiple testing by our tests. Section 5 provides three applications

and conducts the simulation study. Concluding remarks are made in Section 6. Technical proofs are provided in the appendix.

Notation

The following notation is used throughout the paper. Denote by \xrightarrow{p} convergence in probability and by \xrightarrow{d} convergence in distribution. Let $C_k = [0, \infty)^k$ and $C_k^+ = (0, \infty)^k$. Denote by I_k the *k*-dimension identity matrix and 0_k the $k \times 1$ vector of 0 (we may simply use I and 0 without indicating the dimension if no confusion is deemed to arise). For a matrix B, B_{ij} indicates its (i, j)-th element. Similarly, for a vector s, s_i indicates its *i*-th element. A parameter symbol used in a subscript indicates the partitioned component(s) of a vector or a matrix corresponding to parameter involved. For example, $s = (s_{\gamma} : s_{\psi})$ indicates the partition of s according to the parameter $(\gamma : \psi)$. $\mathcal{J}_{\gamma\psi}$ is the left top corner block of \mathcal{J} corresponding to γ and ψ . When an inequality operation is applied to vector(s) it means an elementwise relationship. Finally, $\mathbb{1}(\cdot)$ denotes the usual indicator function and $\|\cdot\|$ denotes the Euclidean norm.

2. Set up

Let Y_T be the data at sample size T and $l_T(\theta)$ be the estimator objective function (e.g., the likelihood function) that depends on Y_T and the q dimensional parameter θ . With θ partitioned as $\theta = (\gamma', \psi')'$, interest is in inference on the $k \times 1$ vector γ , with ψ being a vector of nuisance parameters. We assume that a subset of γ may lie on the boundary of the parameter space, with all the remaining elements of γ being interior points. Specifically, with the partition $\gamma = (\gamma'_1, \gamma'_2)'$, we assume that the first subset $\gamma_1 \in \mathcal{C}_{k_1}, \gamma_2 \in \mathcal{C}^+_{k-k_1}$ and $\psi \in \Psi \subset \mathbb{R}^{q-k}$ for some $1 \leq k_1 \leq k \leq q < \infty$. That is, γ_1 may be on the boundary of the cone \mathcal{C}_{k_1} while γ_2 is assumed to be an interior point of the cone \mathcal{C}_{k-k_1} . The parameter space is denoted by $\Theta(k_1) = \mathcal{C}_{k_1} \times \mathcal{C}^+_{k-k_1} \times \Psi$, and $\Theta(k)$ implies $\mathcal{C}^+_{k-k_1} = \emptyset$ (with $k_1 = k$).

We consider two inference problems. The first, which we label 'joint testing' or 'global testing' interchangeably, concerns testing the null hypothesis $H_0: \gamma = 0$. This testing problem is complicated by the fact that under the null γ is on the boundary of the parameter space. The second is

model selection, i.e. identification of the non-zero elements of γ once the global null hypothesis is rejected. This inference problem requires to deal with multiple testing on the elements of γ .

Let $\theta_0 = (\gamma'_{01}, \gamma'_{02}, \psi'_0)'$ denote the true value of $\theta \in \Theta(k_1)$. Consider the parameter space restricted by

$$\bar{\Theta}(k_1) = \{0\}^{k_1} \times \mathcal{C}^+_{k-k_1} \times \Psi \subset \Theta(k_1).$$

Let $\tilde{\theta}(k_1)$ be the restricted parameter estimator, which satisfies

$$l_T(\tilde{\theta}(k_1)) = \sup_{\theta \in \bar{\Theta}(k_1)} l_T(\theta) + o_p(1)$$
(2.1)

Remark 1. Note that (2.1) is a weaker requirement than $l_T(\tilde{\theta}(k_1)) = \sup_{\theta \in \bar{\Theta}(k_1)} l_T(\theta)$. This allows for $\tilde{\theta}(k_1)$ not being the exact solution that maximises $l_T(\theta)$ in the constrained space $\bar{\Theta}(k_1)$.

We define $\bar{\theta}_0 = (0'_{k_1}, \bar{\gamma}'_{02}, \bar{\psi}'_0)'$ as $plim_{T \to \infty} \tilde{\theta}(k_1) \in \bar{\Theta}(k_1)$ and denote this by the "pseudo-true" value. When $\gamma_{01} = 0$, clearly the pseudo-true value $\bar{\theta}_0 = \theta_0$, the true value. However, when $\gamma_{01} \neq 0$, these differ as will be exploited below.

Denote by $\mathcal{N}(\theta_0, \epsilon) = \{\theta : \|\theta - \theta_0\| = \epsilon < aT^{-1/2}\}$ for some a > 0, such that $\epsilon = O(T^{-1/2})$. That is, $\mathcal{N}(\theta_0, \epsilon)$ is an open sphere neighbourhood centered at θ_0 with the radius $\epsilon = O(T^{-1/2})$. Let $\Theta_{\epsilon}(\theta_0, k_1) = \mathcal{N}(\theta_0, \epsilon) \cap \Theta(k_1)$ and $\overline{\Theta}_{\epsilon}(\theta_0, k_1) = \mathcal{N}(\theta_0, \epsilon) \cap \overline{\Theta}(k_1)$. Denote furthermore by $s_T(\theta) = \partial l_T(\theta) / \partial \theta$ the score function and by $\mathcal{J}_T(\theta) = -\partial^2 l_T(\theta) / \partial \theta \partial \theta'$ the negative Hessian matrix. We make the following assumption throughout.

Assumption 1. With $\theta \in \Theta(k_1)$, assume that $l_T(\theta)$ has continuous derivatives with respect to θ of order two, but $l_T(\theta)$ has continuous right partial derivatives with respect to γ_1 of order two when γ_1 is on the boundary of \mathcal{C}_{k_1} .

Assumption 2. Assume that

$$\sup_{\theta \in \Theta_{\epsilon}(\theta_{0},k_{1})} \left\| T^{-1} \mathcal{J}_{T}(\theta) - \mathcal{J}(\theta) \right\| \to_{p} 0,$$

where $\mathcal{J}(\theta)$ is symmetric, positive definite and uniformly continuous.

Assumption 3. $T^{-1/2}s_T(\theta_0) \xrightarrow{d} N(0, \mathcal{V}(\theta_0))$, where $T^{-1}s_T(\theta_0)s'_T(\theta_0) \xrightarrow{p} \mathcal{V}(\theta_0)$ and $\mathcal{V}(\theta_0)$ is positive definite.

Assumption 4. With $\theta_0 \in \Theta_{\epsilon}(\bar{\theta}_0, k_1)$, assume $\tilde{\theta}(k_1) - \theta_0 = O_p(T^{-1/2})$.

Remark 2. Without loss of generality we assume $T^{1/2}$ -rate constrained estimator $\tilde{\theta}(k_1)$ in this paper.

Remark 3. When $\gamma_{01} = 0$; that is $\theta_0 \in \overline{\Theta}_{\epsilon}(\overline{\theta}_0, k_1)$, Assumption 4 holds if $\tilde{\theta}(k_1)$ is a $T^{1/2}$ -rate constrained estimator. When $\gamma_{01} \neq 0$, we consider the local drifting sequences $\theta_0 = \overline{\theta}_0 + aT^{-1/2}$ for some a > 0. This implies that θ_0 depends on T. To simplify notation, we do not make such dependence explicit. Under the local drifting sequence Assumption 4 is met because

$$\left\|\tilde{\theta}(k_1) - \theta_0\right\| \le \left\|\tilde{\theta}(k_1) - \bar{\theta}_0\right\| + \left\|\bar{\theta}_0 - \theta_0\right\|.$$

We now introduce a result which will be used throughout. Let

$$U_T(\tilde{\theta}(k_1)) = \mathcal{J}_T^{-1}(\tilde{\theta}(k_1)) s_T(\tilde{\theta}(k_1)),$$

and

$$\mathcal{G}_T(\tilde{\theta}(k_1)) = \mathcal{J}_T^{-1}(\tilde{\theta}(k_1)) s_T(\tilde{\theta}(k_1)) s_T'(\tilde{\theta}(k_1)) \mathcal{J}_T^{-1}(\tilde{\theta}(k_1)).$$

Moreover, let $U_{T,\gamma_1}(\theta)$ be the sub-vector of $U_T(\theta)$ and $\mathcal{G}_{T,\gamma_1\gamma_1}(\theta)$ be the sub-block matrix of $\mathcal{G}_T(\theta)$ corresponding to γ_1 . Because ψ is not subject to constraint in obtaining $\tilde{\theta}(k_1)$ it implies $s_{T,\psi}(\tilde{\theta}(k_1)) = 0$. Also, $s_{T,\gamma_2}(\tilde{\theta}(k_1)) = 0$ as $\gamma_2 \in \mathcal{C}_{k-k_1}^+$ is assumed to be interior point of \mathcal{C}_{k-k_1} . Therefore,

$$U_{T,\gamma_1}(\tilde{\theta}(k_1)) = \mathcal{J}_{T,\gamma_1\gamma_1}^{-1}(\tilde{\theta}(k_1))s_{T,\gamma_1}(\tilde{\theta}(k_1)),$$

where $\mathcal{J}_{T,\gamma_1\gamma_1}^{-1}(\theta)$ is the sub-block matrix of $\mathcal{J}_T^{-1}(\theta)$ corresponding to γ_1 . The following lemma holds for θ_0 in a neighbourhood of $\bar{\theta}_0$.

Lemma 2.1. If Assumptions 1-4 hold, then for $\theta_0 \in \overline{\Theta}_{\epsilon}(\overline{\theta}_0, k_1)$ it follows,

$$T^{1/2}\{U_{T,\gamma_1}(\tilde{\theta}(k_1)) - \gamma_{01})\} \xrightarrow{d} N(0, \mathcal{G}_{\gamma_1\gamma_1}(\theta_0)), \qquad (2.2)$$

where $\mathfrak{G}(\theta) = \mathfrak{J}^{-1}(\theta) \mathcal{V}(\theta) \mathfrak{J}^{-1}(\theta).$

Our score tests for both global and multiple testings are constructed based on $U_{T,\gamma}(\tilde{\theta}(k))$, where $\tilde{\theta}(k)$ is the constrained estimator in $\bar{\Theta}(k) = \{0\}^k \times \Psi$. We do not estimate $\tilde{\theta}(k_1)$ for $k_1 < k$. Nevertheless, Lemma 2.1 provides the theoretical basis to study the FWER in multiple testing where the true null hypotheses may be only some of the elements of γ being zero. To illustrate let us consider the ARCH(3) model:

$$\epsilon_n = \sigma_n u_n,$$

$$\sigma_n^2 = \omega + \alpha_1 \epsilon_{n-1}^2 + \alpha_2 \epsilon_{n-1}^2 + \alpha_3 \epsilon_{n-3}^2,$$

where u_n is assumed to be independently drawn from a random distribution with the mean 0 and variance 1 and $\theta = (\gamma', \omega)' = (\alpha_1, \alpha_2, \alpha_3, \omega)'$. Our score tests are constructed based on $\tilde{\theta}(k) = (0, 0, 0, \tilde{\omega})'$ with $\tilde{\gamma} = \tilde{\gamma}_1 = (0, 0, 0)'$, $\tilde{\gamma}_2 = \emptyset$ and $\tilde{\psi} = \tilde{\omega}$. In the global testing $H_0: \gamma = 0$, Lemma 2.1 provides the null distribution of $T^{1/2}U_{T,\gamma}(\tilde{\theta}(k))$ for asymptotically controlling the Type I error. In the multiple testing $H_{0i}: \alpha_i = 0, i \in \{1, 2, 3\}$, suppose the true null is $\alpha_{01} = 0$, $\alpha_{02} = 0$ and $\alpha_{03} > 0$. Lemma 2.1 provides the asymptotic distribution of $T^{1/2}U_{T,\gamma_1}(\tilde{\theta}(k_1))$, where $\gamma_1 = (\alpha_1, \alpha_2)'$ and $\gamma_2 = \alpha_3$. Clearly, this distribution depends on not only $\alpha_{01} = 0$ and $\alpha_{02} = 0$, but $\alpha_{03} > 0$ and ω_0 . The FWER control requires the probability of rejecting $\alpha_1 = 0$ and/or $\alpha_2 = 0$ is bounded by a designated level. Because the researcher does not know the true θ_0 , the FWER should be controlled at any possible θ_0 . We shall study our tests in ARCH models in more details in Section 5.2.

In what follows, to simplify notation, $\tilde{\theta}$ should be understood as $\tilde{\theta}(k)$; if needed, $\tilde{\theta}(k_1)$ is used to stress result holds for $k_1 \leq k$.

3. Joint testing

The joint testing concerns all elements of γ being 0 or not. The global null and alternative hypotheses of interest are

$$\mathsf{H}_0: \theta \in \bar{\Theta}(k) \quad vs \quad \mathsf{H}_1: \theta \in \Theta(k) \setminus \{\gamma = 0\}.$$

$$(3.1)$$

Lemma 2.1 above implies $U_{T,\gamma}(\tilde{\theta}) = \gamma_0 + o_p(1)$. As $\gamma_0 \in \mathbb{C}_k \subset \Theta(k)$, it is desired for the test statistic constructed based on $U_{T,\gamma}(\tilde{\theta})$ to reflect the cone restriction $\Theta(k)$. One may construct one-sided tests based on the test statistic (see, e.g., Silvapulle and Sen, 2005, Sec. 3.4)

$$\tilde{t}_c = \bar{U}'_{T,\gamma}(\tilde{\theta}) \mathcal{G}_{T,\gamma\gamma}^{-1}(\tilde{\theta}) \bar{U}_{T,\gamma}(\tilde{\theta})$$

where

$$\bar{U}_{T,\gamma}(\tilde{\theta}) = \arg \inf_{u \in \mathcal{C}_k} (U_{T,\gamma}(\tilde{\theta}) - u)' \mathcal{G}_{T,\gamma\gamma}^{-1}(\tilde{\theta}) (U_{T,\gamma}(\tilde{\theta}) - u),$$

It then follows from Lemma 2.1 that under H_0

$$\tilde{t}_c \xrightarrow{d} \inf_{u \in \mathcal{C}_k} (Z_{\theta_0} - u)' \mathcal{G}_{\gamma_1 \gamma_1}^{-1}(\theta_0) (Z_{\theta_0} - u),$$

where Z_{θ_0} is the k-dimension multivariate normal random variable with the mean 0 and variance $\mathcal{G}_{\gamma\gamma}(\theta_0)$.

Alternatively, one may construct joint t type tests based on the test statistic

$$\tilde{t}_t = d'_T \mathcal{G}_{T,\gamma\gamma}^{-1}(\tilde{\theta}) U_{T,\gamma}(\tilde{\theta}),$$

where

$$d_T \in \{h \in \mathcal{C}_k^+ : h' \mathcal{G}_{T,\gamma\gamma}^{-1}(\tilde{\theta})h = 1\},\$$

(see e.g., Lu, 2013, for one-sided t tests). Under H_0 , $\tilde{t}_t \xrightarrow{d} N(0, 1)$. We suggest that d_T be computed by Algorithm 1 in Lu (2016) using the covariance $\mathcal{G}_{T,\gamma\gamma}(\tilde{\theta})$, so that the resulting t test has the optimality of asymptotically maximizing the minimum power within the class of tests based on \tilde{t}_t .

3.1. MinP score tests

We now discuss our proposed three MinP score tests. Let $U_{T,\gamma} = (U_{T,\gamma,1}, ..., U_{T,\gamma,k})'$ and $\tilde{t} = (\tilde{t}_1, ..., \tilde{t}_k)'$, where $\tilde{t}_i = U_{T,\gamma,i}(\tilde{\theta})/\sqrt{\mathcal{G}_{T,\gamma\gamma,ii}(\tilde{\theta})}$ and $\mathcal{G}_{T,\gamma\gamma,ii}(\tilde{\theta})$ is the (i, i)th-element of $\mathcal{G}_{T,\gamma\gamma}(\tilde{\theta})$. Let $\tilde{p}_i = 1 - F_{T,i}(\tilde{t}_i), i \in \{1, ..., k\}$, where $F_T(\cdot)$ is the cumulative distribution function (CDF) of \tilde{t} under H_0 and $F_{T,i}(\cdot)$ indicates the marginal distribution corresponding to \tilde{t}_i . Let $F_{T,c}(\cdot)$ and $F_{T,t}(\cdot)$

be the CDFs under H_0 corresponding to \tilde{t}_c and \tilde{t}_t , respectively. Finally, let $\tilde{p}_c = 1 - F_{T,c}{\{\tilde{t}_c\}}$ and $\tilde{p}_t = 1 - F_{T,t}{\{\tilde{t}_t\}}$.

The first proposed MinP score test is based on the test statistic

$$\tilde{p}_{m1} = \min(\tilde{p}_1, \dots, \tilde{p}_k). \tag{3.2}$$

The test statistic for the other two proposed MinP score tests is given by

$$\tilde{p}_{m2} = \min(\tilde{p}_g, \tilde{p}_1, ..., \tilde{p}_k),$$
(3.3)

where \tilde{p}_g takes the value of either \tilde{p}_c or \tilde{p}_t . We refer to the score test based on \tilde{p}_{m1} as the 'MinP-s' test, and the score tests based on \tilde{p}_{m2} using \tilde{p}_c and \tilde{p}_t as 'MinP-sc' and 'MinP-st' tests, respectively.

Let $\tilde{c}_{mj}(\alpha)$ be the critical value at the α significance level such that for $\theta \in \bar{\Theta}(k)$

$$\limsup_{T \to \infty} \Pr(\tilde{p}_{mj} < \tilde{c}_{mj}(\alpha)) \le \alpha, \quad j = 1, 2.$$

We then reject H_0 if $\tilde{p}_{mj} \leq \tilde{c}_{mj}(\alpha)$; otherwise H_0 is accepted. Algorithm 1 below provides an algorithm for numerically computing \tilde{p}_c , \tilde{p}_t , \tilde{p}_i , i = 1, ..., k, and $\tilde{c}_{mj}(\alpha)$ via a valid bootstrap method. Examples of bootstrap implementations are given in Section 5 below.

Algorithm 1.

- 1. For each bootstrapped sample indexed by $b \in \{1, ..., B\}$ compute $\tilde{t}_c^b, \tilde{t}_t^b$ and $\tilde{t}_i^b, i \in \{1, ..., k\}$.
- 2. Compute the empirical p-values \tilde{p}_c , \tilde{p}_t and \tilde{p}_i , i = 1, ..., k. For example, $\tilde{p}_c = B^{-1} \sum_{b=1}^{B} \mathbb{1}(\tilde{t}_c^b \leq \tilde{t}_c)$, where $\mathbb{1}(\cdot)$ is the usual indicator function. Similarly, compute the empirical p-values \tilde{p}_c^b , \tilde{p}_t^b and \tilde{p}_i^b for each $b \in \{1, ..., B\}$.
- 3. Compute $\tilde{p}_{m1}^b = \min(\tilde{p}_1^b, ..., \tilde{p}_k^b)$ and $\tilde{p}_{m2}^b = \min(\tilde{p}_g^b, \tilde{p}_1^b, ..., \tilde{p}_k^b), b = 1, ..., B.$
- 4. Compute the α quantile of the ordered sequence $\{\tilde{p}_{mj}^b, b = 1, ..., B\}, j = 1, 2, as \tilde{c}_{mj}(\alpha)$.

Remark 4. Because it always holds true that $\tilde{p}_{m2}^b \leq \tilde{p}_{m1}^b$ it follows $\tilde{c}_{m2}(\alpha) \leq \tilde{c}_{m1}(\alpha)$.

3.2. Admissibility

We establish the asymptotically locally admissibility property of our tests. Andrews (1996) studied the admissibility of one-sided LR tests, Wald tests and LM tests by showing the optimal power of the tests for a given weighting function of the parameter space under the alternative hypothesis. A similar technique was adopted by Andrews and Ploberger (1995), Andrews et al. (2006) and Chernozhukov et al. (2009) for the study of admissibility of rotation invariant tests. Marden (1982, 1985) studied the admissibility of tests that combine tests of the same kind but applied to independent samples. His admissibility results are based on the principle of Bayes tests. Our proposed tests are a combination of different kinds of tests applied to the same sample. We derive our admissibility result by extending the work of Birnbaum (1955) and Stein (1956) (see, e.g., Theorem 6.7.1 of Lehmann and Romano (2005)) to the case of one-sided testing.

Let $\varphi(Y_T)$ be a $\{0, 1\}$ -valued test for the global hypothesis H_0 . If $\lim_{T\to\infty} \sup_{\theta\in\bar{\Theta}_\epsilon(\bar{\theta}_0,k)} E_\theta(\varphi(Y_T)) \leq \alpha$, then the test $\varphi(Y_T)$ is said to be a locally asymptotically level α test. Denote by $\varphi_{mj}(Y_T) = \mathbb{1}(\tilde{p}_{mj} \leq \tilde{c}_{mj}(\alpha))$ the test function for the proposed MinP score tests. Let $\varphi'_{mj}(Y_T)$ be a locally asymptotically distinct test from $\varphi_{mj}(Y_T)$ such that for each $\theta \in \Theta_\epsilon(\bar{\theta}_0,k)$

$$\liminf_{T \to \infty} E_{\theta}[\{1 - \varphi'_{mj}(Y_T)\}\varphi_{mj}(Y_T) + \varphi'_{mj}(Y_T)\{1 - \varphi_{mj}(Y_T)\}] > 0.$$
(3.4)

Note that a distinct test $\varphi'_{mj}(Y_T)$ defined in (3.4) implies that $\varphi'_{mj}(Y_T)$ cannot take the same values as $\varphi_{mj}(Y_T)$ asymptotically, and the limiting acceptance region of the test $\varphi_{mj}(Y_T)$ is different from that of $\varphi'_{mj}(Y_T)$.

Suppose the limiting acceptance region of $\varphi_{mj}(Y_T)$ is a closed, convex and lower set E such that if $u \leq v$ and $v \in E$, then $u \in E$. Define a halfspace by $W_{e,\delta} = \{v : e' \mathcal{G}_{\gamma\gamma}^{-1}(\theta_0) v < \delta\}$, where $v \in \mathbb{R}^k$, $e \in \mathbb{C} \setminus \{\gamma = 0\}$ and $\delta \in \mathbb{R}$. Let $W = \{v : \bigcap_{e \in \mathbb{C} \setminus \{\gamma = 0\}} W_{e,\delta}\}$. Denote by $W^*(E) \subseteq W$ such that $E \subseteq W^*(E) \subseteq W'(E)$ for all W'(E) that satisfies $E \subseteq W'(E) \subseteq W$. Therefore, $W^*(E)$ can be viewed as the intersection of the halfspaces $W_{e,\delta} \subseteq W$ for all $e \in \mathbb{C} \setminus \{\gamma = 0\}$ that are tangent with E.

The limiting acceptance region of the MinP-s test is $E_s(c_s) = \{v : v < c_s\}, c_s > 0$. Consider the k halfspaces $W_{e,\delta}$ with e and v being the column vector of the identity matrix. Then $W^*(E_s) = E_s$. For joint t tests based on \tilde{t}_t the limiting acceptance region is the halfspace defined by $E_t(c_t) = \{v : d' \mathcal{G}_{\gamma\gamma}^{-1}(\theta_0)v < c_t, d \in \mathbb{C}^+ \setminus \{\gamma = 0\}, c_t > 0$. Because that d is a column vector with positive elements that E_t cuts through the top corner of E_s . Thus, for the MinP-st test the limiting acceptance region is $E_{st}(c) = \{v : E_s(c) \cap E_t(c)\}, c > 0$, and $W^*(E_{st}) = E_{st}$. For tests based on \tilde{t}_c the limiting acceptance region is $E_c(c_c) = \{v : v' \mathcal{G}_{\gamma\gamma}^{-1}(\theta_0)v < c_c\}, c_c > 0$. The limiting acceptance region of the MinP-sc test is then $E_{sc} = \{v : E_s(c_s) \cap E_c(c_c)\}, where c_s$ and c_c are positive constants such that the overall Type I error control for testing H_0 by MinP-sc tests is achieved. In this case it may be that $W^*(E_{sc}) = E_{sc}$ or $W^*(E_{sc}) = E_s$ (see Figure 1 for illustration).

[Figure 1 about here.]

Theorem 3.1. Under Assumptions 1, 2, 3 and 4 $\varphi_{mj}(Y_T)$ is asymptotically admissible in the sense that there does not exist a distinct test $\varphi'_{mj}(Y_T)$ whose limiting acceptance region E' is not contained in $W^*(E_{mj})$ such that for all $\theta \in \Theta_{\epsilon}(\bar{\theta}_0, k) \setminus \{\gamma = 0\}$

$$\liminf_{T \to \infty} E_{\theta} \varphi'_{mj}(Y_T) \ge \limsup_{T \to \infty} E_{\theta} \varphi_{mj}(Y_T).$$
(3.5)

Remark 5. For MinP-s and MinP-st tests because $W^*(E) = E$ Theorem 3.1 suggests that there does not exists any other test that is uniformly more powerful than MinP-s or MinP-st tests. This is also true for MinP-sc tests if $W^*(E_{sc}) = E_{sc}$. However, when $W^*(E_{sc}) = E_s$ Theorem 3.1 suggests that there does not exists another test whose limiting acceptance region is not contained in E_s , is uniformly more powerful than MinP-sc tests.

4. Multiple testing and model selection

The MinP score tests described previously may be used for simultaneously testing on individual components of $\gamma = (\gamma_1, ..., \gamma_k)'$. That is, consider testing multiple hypotheses

$$\mathsf{H}_{0i}: \gamma_i = 0, \psi \in \Psi \quad vs \quad \mathsf{H}_{1i}: \gamma_i > 0, \psi \in \Psi \quad i \in K = \{1, ..., k\}.$$
(4.1)

When a set of hypotheses is tested simultaneously it is important to control multiplicity of Type I errors as otherwise the probability of wrongly rejecting one hypothesis increases as the number of true hypotheses increases. The family-wise error rate (FWER) is a widely used Type I error probability control in multiple testing. Let

$$K_0 = \{i \in K : \gamma_{0i} = 0\}$$

be the set containing the indices of true H_{0i} . The FWER is the probability of rejecting at least one H_{0i} , $i \in K_0$ under $\bigcap_{i \in K_0} H_{0i}$. We shall show that the FWER control is bounded by α for MinP score tests under some additional assumptions.

To illustrate, consider the following simple example. Suppose the sample $\{y_n, n = 1, ..., T\}$ is independently randomly drawn from a multivariate normal distribution with mean γ and known variance Ω . Then, the objective (log-likelihood) function is

$$l_T(\theta) \propto -\frac{T}{2} \log |\Omega| - \frac{1}{2} \sum_{n=1}^T (y_n - \gamma)' \Omega^{-1}(y_n - \gamma).$$

As Ω is known, $s_T(\gamma) = \sum_{n=1}^T \Omega^{-1}(y_n - \gamma)$, $\mathcal{J}_T(\gamma) = T\Omega^{-1}$ and $U_T(\gamma) = T^{-1}\sum_{n=1}^T(y_n - \gamma)$. Therefore, under $\mathsf{H}_0, U_T(\tilde{\gamma}) = T^{-1}\sum_{n=1}^T y_n$ and $T^{1/2}U_T(\tilde{\gamma}) \sim N(\gamma, \Omega)$. The MinP score test based on \tilde{p}_{m1} is the same as MinP tests based on the sample average. In this case the joint distribution of $U_{T,i}(\tilde{\gamma}), i \in J \subset K$, for a subset J under $\bigcap_{i \in J} \mathsf{H}_{0i}$ is $N(0, \Omega_J)$, where Ω_J contains the blocks of Ω corresponding to J. Obviously, the joint distribution of $U_{T,i}(\tilde{\gamma})$ is not affected by the truth or falsehood of the remaining $\mathsf{H}_{0i}, i \in K \setminus J$. This is an example of what is known as the subset pivotality condition (see, e.g., Westfall and Young (1993)). As a result, the stepdown procedure of MinP-s score tests based on \tilde{p}_{m1} allows for the control of the FWER. However, this may not be the case in our set up as the covariance $\mathcal{G}_{\gamma\gamma}(\theta_0)$ in (2.2) may depend on γ_0 . By setting $\gamma = 0$ as the researcher would usually do in joint testing of the global null H_0 is normally referred to as the weak control of FWER in the literature of multiple testing, which does not guarantee the control of the FWER in multiple testing (see, e.g., Romano and Wolf (2005)).

Let $\tilde{c}_J(\alpha)$ be the critical value at the α significance level such that under $\bigcap_{i \in J} \mathsf{H}_{0i}, J \subset K$,

$$\limsup_{T \to \infty} \Pr[\min\{\tilde{p}_i(\tilde{\theta}(k)), i \in J\} < \tilde{c}_J(\alpha)] \le \alpha$$

where the notation $\tilde{p}_i(\tilde{\theta}(k))$ is used to stress $\tilde{p}_i(\tilde{\theta}(k))$ is based on the estimate $\tilde{\theta}(k)$, not the estimate by restricting $\tilde{\gamma}_i = 0, i \in J$. Note that if J = K then $\tilde{c}_J(\alpha)$ is equal to $\tilde{c}_{m1}(\alpha)$ in the global testing.

MinP score tests for multiple testing are as follows. If the global hypothesis H_0 is rejected then order \tilde{p}_i , i = 1, ..., k, as follows:

$$\tilde{p}_{(1)} \le \tilde{p}_{(2)} \le \dots \le \tilde{p}_{(k)}$$

and let $\mathsf{H}_{(01)}$, ..., $\mathsf{H}_{(0k)}$ be the corresponding hypotheses. Reject $\mathsf{H}_{(01)}$ if and only if $\tilde{p}_{(1)} \leq \tilde{c}_{mj}(\alpha)$. If $\mathsf{H}_{(01)}$ is rejected one would then continue to the stepdown procedure for the remaining $\mathsf{H}_{(02)}$, ..., $\mathsf{H}_{(0k)}$, as shown in Algorithm 2 below.

Algorithm 2.

- 1. In the first step if $\tilde{p}_{mj} > \tilde{c}_{mj}(\alpha)$ then accept H_{0i} for all $i \in K$, and stop; otherwise continue.
- 2. If $\tilde{p}_{(1)} > \tilde{c}_{mj}(\alpha)$ then accept $\mathsf{H}_{(0i)}$ for all $i \in K$; otherwise, reject $\mathsf{H}_{(01)}$ and continue.
- 3. For the remaining $\mathsf{H}_{(0i)}$, $i \in \{2, ..., k\}$, let K_i be the set of indices of the individual hypotheses subject to testing at the *i*th step for i = 2, ..., k. If $\tilde{p}_{(i)} > \tilde{c}_{K_i}(\alpha)$, then accept $\mathsf{H}_{(0i)}$ for all $i \in K_i$, and stop; otherwise reject $\mathsf{H}_{(0i)}$ if $\tilde{p}_{(i)} \leq \tilde{c}_{mi}(\alpha)$ and continue until K_i is the empty set.

Without loss of generality let the first k_0 elements of γ_0 containing the true $\mathsf{H}_{0i} : \gamma_{0i} = 0$, i.e., $K_0 = \{1, ..., k_0\}$. When $K_0 = \emptyset$, the FWER is set to 0 by default. When $K_0 \neq \emptyset$, let $c_{K_0}(\alpha)$ be the critical value such that

$$c_{K_0}(\alpha) = \sup\{c : \limsup_{T \to \infty} \sup_{\theta \in \bar{\Theta}_{\epsilon}(\bar{\theta}_0, k_0)} \Pr[\min\{\tilde{p}_i(\tilde{\theta}(k_0)), i \in K_0\} \le c] \le \alpha\}.$$

We show in Theorem 4.1 that, under the assumptions of Theorem 3.1, if also Assumption 5 holds, the FWER control for MinP score tests is bounded by α .

Assumption 5. $\tilde{c}_{K_i}(\alpha) \leq c_{K_0}(\alpha) + \varepsilon$ with probability approaching to 1 for $K_0 \subseteq K_i \subseteq K$ and any $\varepsilon > 0$.

Theorem 4.1. Under Assumptions 1, 2, 3, 4 and 5 $\limsup_{T\to\infty} \sup_{\theta\in\bar{\Theta}_{\epsilon}(\bar{\theta}_{0},k_{0})} \mathsf{FWER} \leq \alpha$ holds true for all three proposed MinP score tests, namely, MinP-s, MinP-sc and MinP-st tests.

Remark 6. In some cases $\tilde{c}_{K_i}(\alpha)$ may be computed once \tilde{p}_i^b , $i \in \{1, ..., k\}$, $b \in \{1, ..., B\}$, are generated for computing $\tilde{c}_{m1}(\alpha)$ in Algorithms 1; one simply takes \tilde{p}_i^b , $i \in K_i$, for computing $\tilde{c}_{K_i}(\alpha)$ without a need for further resampling. However, one needs to ensure that Assumption 5 holds under such resampling procedure.

Remark 7. For MinP-s tests based on \tilde{p}_{m1} the rejection of the global null hypothesis H_0 means the rejection of $\mathsf{H}_{(01)}$. But for MinP-sc and MinP-st tests based on \tilde{p}_{m2} the rejection of H_0 does not necessarily mean the rejection of $\mathsf{H}_{(01)}$; there may be the case $\tilde{p}_g \leq \tilde{c}_{m2}(\alpha)$ but $\tilde{p}_{(1)} > \tilde{c}_{m2}(\alpha)$. Therefore, MinP-s tests have an advantage over MinP-sc and MinP-st tests in the sense of what is known as the consonance property in the literature of multiple testing which states that the rejection of $\bigcap_{i \in J} \mathsf{H}_{0i}$ implies the rejection of at least one component H_{0i} (see, e.g., Finner and Strassburger (2002) and Romano et al. (2011)). However, MinP-sc and MinP-st tests may have global power advantages in some cases as shown in our applications.

Remark 8. If $H_{(01)}$ is rejected by MinP-sc or MinP-st tests, then it is rejected by MinP-s tests if $\tilde{c}_{mj}(\alpha)$, j = 1, 2, is computed by Algorithm 1 as $\tilde{c}_{m2}(\alpha) \leq \tilde{c}_{m1}(\alpha)$ discussed in Remark 4. Furthermore, once $H_{(01)}$ is rejected by MinP-s, MinP-sc and MinP-st tests, all the three tests would have the same rejection outcome in multiple testing as they proceed to the same stepdown procedure.

Multiple tests of the hypotheses (4.1) allow for model selection through testing. Let

$$\tilde{K} = \{i \in \{1, ..., k\} : \tilde{p}_{(i)} \le \tilde{c}_{K_i}(\alpha)\},\$$

where $\tilde{c}_{K_1}(\alpha) = \tilde{c}_{mj}(\alpha)$, be the set of the indices of H_{0i} that is rejected in the above stepdown procedure.

Theorem 4.2. Under Assumptions 1, 3, 4 and 5 we have (i) $\limsup_{T\to\infty} \sup_{\theta\in\bar{\Theta}_{\epsilon}(\bar{\theta}_{0},k_{0})} \Pr[(\hat{K}\cap K_{0})\neq \emptyset] \leq \alpha$, and (ii) $\lim_{T\to\infty} \Pr(\hat{K}\supseteq\bar{K}_{0})=1$, where $\bar{K}_{0}=K\setminus K_{0}\neq \emptyset$, for $\theta\in\Theta_{\epsilon}(\bar{\theta}_{0},k_{0})\setminus\{\cap_{i\in K_{0}}\gamma_{i}=0\}$.

Remark 9. The result in Theorem 4.2(i) says that the supremum of the probability of at least one true $\gamma_{0i} = 0, i \in K_0$, being selected in \hat{K} is asymptotically bounded above by the designated level Remark 10. The result in Theorem 4.2(ii) implies that the infimum of the probability of $i \in \bar{K}_0$ being selected in \hat{K} converges to 1 as the sample size T goes to infinity as long as \bar{K}_0 is not empty. This relates to the consistency property of multiple testing.

5. Applications and Monte Carlo studies

We implement our tests for applications in linear regression models, ARCH models and random coefficient models. Simulation studies are also conducted in order to examine the finite sample performance of our tests. The number of bootstrap repetitions B in implementing Algorithm 1 is set to 999. The number of Monte Carlo replications in our simulation studies is set to 2000. All the reported results are based on the nominal 0.05 size. In all three models considered below $x_n = (x_n^d, 1)'$, where x_n^d is a scalar, $\beta = (1, 1)'$ and we observe the sample $\{y_n, z_n, x_n^d,$ $n = 1, ...T\}$. Let $\tilde{\epsilon}_n$ be the regression residuals under H_0 . When implemented, the bootstrap is based on generating the bootstrap shocks ϵ_n^* by randomly drawn with replacement from $\tilde{\epsilon}_n$ and y_n^* from the regression model under H_0 using $\tilde{\theta}(k)$, ϵ_n^* and fixed covariates. Note that, in some applications, one may need to center the residuals in order to have bootstrap shocks with (conditionally on the original data) exact zero mean.

In all applications we report size and power with respect to global tests of H_0 . In terms of multiple testing of H_{0i} , $i \in K$, we report the FWER and the probability of rejecting individual H_{0i} by the stepdown multiple testing procedure described in Algorithm 2.

5.1. Linear models

Consider the model

$$y_n = z'_n \gamma + x'_n \beta + \epsilon_n$$

where ϵ_n is assumed to be independently drawn from a random distribution with the mean 0 and variance $\sigma^2 < \infty$, z_n and x_n are the $k \times 1$ and $(q - k) \times 1$ column vectors, respectively, and $\gamma \in \mathbb{C} \subset \mathbb{R}^k$ and $\beta \in \mathbb{R}^{q-k-1}$ with the last element being the intercept parameter. The objective function $l_T(\theta) = -\sum_{n=1}^T (y_n - z'_n \gamma - x'_n \beta)^2$, where $\theta = (\gamma', \beta')'$. Let $\tilde{\epsilon}_n$ be the estimated residuals based on the constrained Ordinary Least Squares (OLS) estimate $\tilde{\theta} = (0', \tilde{\beta}')'$. To check whether Assumption 5 holds in this example, we can make use of the following proposition.

Proposition 1. If Assumptions 1, 3 and 4 hold then Assumption 5 is satisfied. This is true even if Algorithm 2 is implemented by following Remark 6.

Remark 11. Let $\tilde{\sigma}^2(k_1)$ and $\tilde{\sigma}^2(k)$ be the restricted estimate of variance of ϵ_n in the models $y_n = z'_{n,2}\gamma_2 + x'_n\beta + \epsilon_n$ and $y_n = x'_n\beta + \epsilon_n$, respectively. The reason Assumption 5 is satisfied is that the limiting joint distribution $\{U_{T,i}(\tilde{\theta}(k_1)), i \in \{1, ..., k_1\} \subset K\}$ under $\bigcap_{i \in \{1,...,k_1\}} \mathsf{H}_{0i}$ is affected by the truth or falsehood of the remaining H_{0i} only through $\tilde{\sigma}^2(k_1)$, which is always no larger than $\tilde{\sigma}^2(k)$.

In our simulation study we generate $X^d = (z_1, ..., z_T, x_1^d, ..., x_T^d)'$ as i.i.d. from $N(0, \Sigma)$, where

$$\Sigma = (1 - \rho)^{-1} [I_{k+1} - \rho \{1 + k\rho\}^{-1} \mathbb{1}_{k+1} \mathbb{1}'_{k+1}],$$

 ρ is constant and 1_k is a $k \times 1$ vector of 1. Thus $E(T^{-1}X^{d'}X^d)^{-1} = \Sigma^{-1}$, which has the structure of a correlation matrix with all the off-diagonal elements being ρ .

Tables 1–2 report the estimated probabilities of rejecting H_0 , FWER and H_{0i} with ϵ_n being i.i.d. (and independent of X^d) and following either the N(0, 1) distribution or the Student's tdistribution with 5 degrees of freedom; k is set to 2. Notice that for this data generating process, Assumptions 1, 3 and 4 hold. Assumption 5 also holds by Proposition 1. Results show that MinPsc and MinP-st score tests have competitive finite-sample performance in both size and power when compared with score tests based on \tilde{t}_c and \tilde{t}_t in terms of global testing. However, MinP-sc and MinP-st score tests have an added benefit of testing individual components of γ with the FWER control. While MinP-s score tests may perform slightly better in multiple testing, they can have inferior power in global testing of H_0 , for example, in the case of $\rho = -0.45$.

[Table 1 about here.]

[Table 2 about here.]

5.2. ARCH models

Consider the regression model with ARCH disturbances

$$y_n = x'_n \beta + \epsilon_n,$$

$$\epsilon_n = \sigma_n u_n,$$

$$\sigma_n^2 = \omega + \sum_{j=1}^k \alpha_j \epsilon_{n-j}^2,$$
(5.1a)

where u_n is assumed to be independently drawn from a random distribution with the mean 0 and variance 1 and $\theta = (\lambda', \beta')'$, where $\lambda = (\gamma', \omega)' = (\alpha_1, ..., \alpha_k, \omega)'$. The quasi-log-likelihood function of the model under the Gaussian sequence $\{u_n\}$ is $l_T(\theta) \propto -\frac{1}{2} \sum_{n=1}^T (\log \sigma_n^2 + \frac{\epsilon_n^2}{\sigma_n^2})$. We assume that $E |x_n x'_n| < \infty$ and the ARCH process $\{\epsilon_n\}$ is stationary and ergodic with $E\epsilon_n^6 < \infty$.

Let $w_n = (\epsilon_{n-1}^2, ..., \epsilon_{n-k}^2, 1)'$. Since $\mathcal{J}_{T,\lambda\beta}(\theta) = T^{-1} \sum_{n=1}^T (2\sigma_n^4)^{-1} w_n x'_n \epsilon_n = o_p(1)$, $\mathcal{G}_{\lambda\lambda}(\theta)$ only involves $\mathcal{J}_{\lambda\lambda}(\theta)$ and $\mathcal{V}_{\lambda\lambda}(\theta)$. So let us consider $U_{T,\lambda}(\theta) = \mathcal{J}_{T,\lambda\lambda}^{-1}(\theta) s_{T,\lambda}(\theta)$ for contructing our tests with $s_{T,\lambda}(\theta) = \sum_{n=1}^T \frac{\epsilon_n^2 - \sigma_n^2}{2\sigma_n^4} w_n$ and $\mathcal{J}_{T,\lambda\lambda}(\theta) = \sum_{n=1}^T \frac{2\epsilon_n^2 - \sigma_n^2}{2\sigma_n^6} w_n w'_n$. As $Eu_n^2 = 1$, it follows that $T^{-1}\mathcal{J}_{T,\lambda\lambda}(\theta) = T^{-1} \sum_{n=1}^T (2\sigma_n^4)^{-1} w_n w'_n + o_p(1)$. Let $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$ be the block of $\mathcal{G}(\theta_0)$ corresponding to the true $\alpha_{0i} = 0$ for $k_0 = k_1 \leq k$.

Because $\tilde{\gamma} = 0$, $\tilde{\sigma}_n^2 = \tilde{\omega}$ and $E\epsilon_n^2 = \sigma_n^2$ under the global H_0 , it follows $T^{-1}\mathcal{J}_{T,\lambda\lambda}(\tilde{\theta}) = (2\tilde{\omega}^2)^{-1}T^{-1}\sum_{n=1}^T \tilde{w}_n \tilde{w}'_n + o_p(1)$ and $s_{T,\lambda}(\tilde{\theta}) = (2\tilde{\omega}^2)^{-1}\sum_{n=1}^T (\tilde{\epsilon}_n^2 - \tilde{\omega})\tilde{w}_n$, where \tilde{w}_n is computed using $\tilde{\epsilon}_n$. Therefore, $U_{T,\lambda}(\tilde{\theta})$ can be viewed as the OLS estimate of the regression coefficients of the regression of $\tilde{\epsilon}_n^2 - \tilde{\omega}$ on an intercept and the q lagged values of $\tilde{\epsilon}_n^2$, and $U_{T,\gamma}(\tilde{\theta})$ is the slope estimate. Since adding a constant to the dependent variable only affects the estimate of the intercept, $U_{T,\gamma}(\tilde{\theta})$ can be equivalently viewed as the estimate of the slope $(\hat{\gamma})$ of the regression of $\tilde{\epsilon}_n^2$ on an intercept and q lagged values of $\tilde{\epsilon}_n^2$.

For $i \neq j, i, j \in \{1, ..., T\}$ it follows $E\tilde{\epsilon}_n^2 = \omega_0, E\tilde{\epsilon}_i^2\tilde{\epsilon}_j^2 = E\tilde{\epsilon}_i^2E\tilde{\epsilon}_j^2$ and $E\tilde{\epsilon}_i^4 = E\tilde{\epsilon}_j^4$. It can then be shown that

$$T^{-1}s_{T,\lambda\lambda}(\tilde{\theta})s'_{T,\lambda\lambda}(\tilde{\theta}) = (E\eta_n^4 - 1)(2\tilde{\omega})^{-2}T^{-1}\sum_{n=1}^T \tilde{w}_n\tilde{w}'_n + o_p(1)$$

and $T\mathcal{G}_{T,\gamma\gamma}(\tilde{\theta}) = I_k + o_p(1)$. Assumptions 1, 3 and 4 are straightforward to verify. Assumption 5 follows under a sufficient condition as stated in the following proposition.

Proposition 2. Under Assumptions 1, 3 and 4 if $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$ is a non-positive matrix meaning all off diagonal elements are non-positive, then Assumption 5 is satisfied.

We compare the performance of MinP score tests with that of score tests proposed in Demos and Sentana (1998) and Lee and King (1993). The score test proposed in Demos and Sentana (1998) is asymptotically equivalent to the score test based on \tilde{t}_c that has the limiting null distribution $\sum_{i=0}^{k} 2^{-q} {k \choose i} \chi_i^2$. The score test proposed in Lee and King (1993) is asymptotically equivalent to the score test based on \tilde{t}_t . We generate $x_n = (x_n^d, 1)'$, $x_n^d = 0.8x_{n-1}^d + e_n$, $e_n \sim i.i.d.N(0, 4)$, and u_n from either the standard normal distribution or the Student's t with 5 degrees of freedom. We add a burn-in period of 1000 observations for ϵ_n , which are discarded in estimation. We adopt the constrained OLS estimate $\tilde{\theta} = (0', \tilde{\beta}', \tilde{\sigma}^2)'$.

Tables 3 and 7 report the estimated probabilities of rejecting H_0 , FWER and H_{0i} . The results show that the global powers of MinP tests are competitive with those of Demos and Sentana (1998)'s and Lee and King (1993)'s tests. Although the FWER control by our MinP tests may not always be guaranteed, the probability of identifying the false H_{0i} increases towards 1 as sample size increases.

[Table 3 about here.][Table 4 about here.][Table 5 about here.][Table 6 about here.][Table 7 about here.]

5.3. Random coefficient models

Consider the model

$$y_n = z'_n \xi_n + x'_n \beta + \epsilon_n,$$

where the random coefficient ξ_n has the form

 $\xi_n = \xi + \eta_n,$

and ξ and β are a column vector of a fixed coefficient with the dimension of k and (q - k - 1), respectively. The random variables $\epsilon_n \in \mathbb{R}$ and $\eta_n \in \mathbb{R}^k$ are unobserved independent errors across individual elements of η_n , between ϵ_n and η_n , as well as across n = 1, ..., T that satisfy $E\epsilon_n = 0$, $E\epsilon_n^2 = \sigma_{\epsilon}^2$, $E(\eta_n | x_n) = 0$ and $E(\eta_n \eta'_n | x_n) = diag(\gamma)$, where $\gamma = (\sigma_{\eta,1}^2, ..., \sigma_{\eta,k}^2)'$.

The model may be rewritten as

$$y_n = z'_n \xi + x'_n \beta + \varpi_n, \tag{5.2}$$

where

$$\varpi_n = z'_n \eta_n + \epsilon_n.$$

Consider the quasi-log-likelihood function under normality as $l_T(\theta) \propto -\frac{1}{2} \sum_{n=1}^T \{\log \sigma_{\varpi_n}^2 + \frac{\varpi_n^2}{\sigma_{\varpi_n}^2}\},$ where $\sigma_{\varpi_n}^2 = \gamma' z_n^2 + \sigma_{\epsilon}^2$ and $z_n^2 = (z_{n,1}^2, ..., z_{n,k}^2)'$. Let $\lambda = (\gamma', \sigma_{\epsilon}^2)', \ \psi = (\xi', \beta')', \ \theta = (\lambda', \psi')'$ and $d_n = ((z_n^2)', 1)'$. Since $E(z'_n, x'_n) \varpi_n = 0$ we have $T^{-1} \mathcal{J}_{T,\lambda\psi}(\theta) = \sum_{n=1}^T (2\sigma_{\varpi_n}^4)^{-1} d_n(z'_n, x'_n) \varpi_n =$ $o_p(1)$ and $\mathcal{G}_{\lambda\lambda}(\theta)$ only involves $\mathcal{J}_{\lambda\lambda}(\theta)$ and $\mathcal{V}_{\lambda\lambda}(\theta)$. So let's consider $U_{T,\lambda}(\theta) = \mathcal{J}_{T,\lambda\lambda}^{-1}(\theta) s_{T,\lambda}(\theta)$ for constructing our tests with $s_{T,\lambda}(\theta) = \sum_{n=1}^T \frac{\varpi_n^2 - \sigma_{\varpi_n}^2}{2\sigma_{\varpi_n}^4} d_n$ and $\mathcal{J}_{T,\lambda\lambda}(\theta) = \sum_{n=1}^T \frac{2\varpi_n^2 - \sigma_{\varpi_n}^2}{2\sigma_{\varpi_n}^6} d_n d'_n$. Hence, $U_{T,\gamma_1}(\tilde{\theta}(k_1)) = \mathcal{J}_{T,\gamma_1\gamma_1}^{-1}(\tilde{\theta}(k_1)) s_{T,\gamma_1}(\tilde{\theta}(k_1))$ and $\mathcal{G}_{\gamma_1\gamma_1}(\theta)$ is the upper left block corresponding to $\mathcal{G}_{\lambda\lambda}(\theta)$. Without loss of generality let $k_0 = k_1$, so $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$ corresponds to the true $\gamma_{0i} = 0$.

Under the global H_0 we have $\tilde{\gamma} = 0$, $\tilde{\sigma}_{\varpi_n}^2 = \tilde{\sigma}_{\epsilon}^2$ and $E\varpi_n^2 = \sigma_{\varpi_n}^2$, hence $T^{-1}\mathcal{J}_{T,\lambda\lambda}(\tilde{\theta}) = (2\tilde{\sigma}_{\epsilon}^4)^{-1}T^{-1}\sum_{n=1}^T d_n d'_n + o_p(1)$ and $s_{T,\lambda}(\tilde{\theta}) = (2\tilde{\sigma}_{\epsilon}^4)^{-1}\sum_{n=1}^T (\tilde{\varpi}_n^2 - \tilde{\sigma}_{\epsilon}^2)d_n$, where $\tilde{\varpi}_n$ is the residual computed based on $\tilde{\theta}$. For $i \neq j$, $i, j \in \{1, ..., T\}$, it follows $E\tilde{\epsilon}_n^2 = \omega_0$, $E\tilde{\epsilon}_i^2\tilde{\epsilon}_j^2 = E\tilde{\epsilon}_i^2E\tilde{\epsilon}_j^2$ and $E\tilde{\epsilon}_i^4 = E\tilde{\epsilon}_j^4$. Let $\tilde{\varpi}_n^2 = T^{-1}\sum_{n=1}^T (y_n - z'_n\tilde{\xi} + x'_n\tilde{\beta})^2$ and $T^{-1}\sum_{n=1}^T d_n d'_n = \Sigma_d + o_p(1)$. It can then be shown that $T\mathcal{G}_{T,\lambda\lambda}(\tilde{\theta}) = \tilde{\Sigma}_d + o_p(1)$, where $\tilde{\Sigma}_d = \sum_{n=1}^T (y_n - z'_n\tilde{\xi} + x'_n\tilde{\beta})^2(\sum_{n=1}^T d_n d'_n)^{-1}$. Assumption 5 can be checked through the following proposition (where Assumptions 1, 3 and 4 can be checked as in Andrews, 1999, Example 1).

Proposition 3. Under Assumptions 1, 3 and 4 if $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$ is asymptotically elementwise no greater than the upper left $k_0 \times k_0$ block of $\tilde{\Sigma}_d$, then Assumption 5 is satisfied.

Tables 8–9 report the estimated probabilities of rejecting H_0 , FWER and H_{0i} with $\epsilon_n \sim N(0, 1)$ or t_5 . We generate i.i.d x_n^d from N(0, 4) and i.i.d. z_n from the multivariate normal distribution with mean 0 and covariance matrix with diagonal entry 1 and all off-diagonal elements being 0.2. The results show that MinP tests have a competitive finite sample performance compared with score tests based on \tilde{t}_c in terms of global testing. Score tests based on \tilde{t}_t can have considerable worse global power in some cases. With regard to multiple testing MinP score tests identify the false H_{0i} with probability increases towards 1 as sample size increases although the FWER control may not always guaranteed. While MinP-s score tests may perform slightly better in multiple testing, such an advantage may be compromised by their global power in testing H_0 .

[Table 8 about here.]

[Table 9 about here.]

6. Conclusion

This paper proposes MinP score tests that allow for global and multiple testing of one-sided hypotheses. The simulation results suggest that the proposed tests are competitive with existing one-sided score tests with respect to global power. We further demonstrate that the proposed tests may be used for multiple testing and model identification among nested models. The feature of model identification is appealing when the model parameter is subject to inequality constraints. These tests allow for model identification without estimating candidate models except the one defined under the global null hypothesis, which is usually easy to obtain. We illustrated applications of the tests in linear regression, ARCH models, and random coefficient models. Admissibility and model identification properties of these tests provide a theoretical support of our testing approach to the practically important problem of one-side testing.

APPENDIX: PROOFS

Proof of Lemma 2.1. Consider the Taylor expansion at θ_0

$$T^{-1}s_T(\tilde{\theta}(k_1)) = T^{-1}s_T(\theta_0) - T^{-1}\mathcal{J}_T(\theta_0)(\tilde{\theta}(k_1) - \theta_0) + R_T$$
(A.1)

where

$$R_T = T^{-1} \{ \mathcal{J}_T(\theta_0) - \mathcal{J}_T(\theta_+) \} (\tilde{\theta}(k_1) - \theta_0).$$

and θ_+ is a point on the line segment joining θ_0 and $\tilde{\theta}(k_1)$.

We shall first show

$$||R_T|| = o_p(1). \tag{A.2}$$

By Assumption 4

$$\tilde{\theta}(k_1) - \theta_0 = o_p(1) \tag{A.3}$$

Hence,

$$\theta_+ - \theta_0 = o_p(1) \tag{A.4}$$

Consider

$$\begin{aligned} \left\| T^{-1} \mathcal{J}_T(\theta_+) - T^{-1} \mathcal{J}_T(\theta_0) \right\| \\ &\leq \left\| T^{-1} \mathcal{J}_T(\theta_+) - \mathcal{J}(\theta_+) \right\| + \left\| T^{-1} \mathcal{J}_T(\theta_0) - \mathcal{J}(\theta_0) \right\| + \left\| \mathcal{J}(\theta_+) - \mathcal{J}(\theta_0) \right\|. \end{aligned}$$

It follows from Assumption 2 that

$$\left\| T^{-1} \mathcal{J}_T(\theta_+) - \mathcal{J}(\theta_+) \right\| = o_p(1),$$
$$\left\| T^{-1} \mathcal{J}_T(\theta_0) - \mathcal{J}(\theta_0) \right\| = o_p(1).$$

It follows from (A.4) and the continuous mapping theorem that

$$\|\mathcal{J}(\theta_+) - \mathcal{J}(\theta_0)\| = o_p(1),$$

thus,

$$T^{-1}\mathcal{J}_T(\theta_+) - T^{-1}\mathcal{J}_T(\theta_0) = o_p(1).$$
(A.5)

and (A.2) follows from (A.3) and (A.5).

Next, it follows from (A.1) that

$$\mathcal{J}_T^{-1}(\theta_0)s_T(\tilde{\theta}(k_1)) - (\theta_0 - \tilde{\theta}(k_1)) = \mathcal{J}_T^{-1}(\theta_0)s_T(\theta_0) + o_p(1).$$

Because $\tilde{\gamma}_1 = 0$, $\theta_0 - \tilde{\theta}(k_1) = (\gamma'_{01}, (\gamma_{02} - \tilde{\gamma}_2)', (\psi_0 - \tilde{\psi})')'$. Furthermore, $T^{-1}\mathcal{J}_T(\tilde{\theta}(k_1)) = T^{-1}\mathcal{J}_T(\theta_0) + o_p(1)$ by following similar steps to show (A.5), and $T^{-1}\mathcal{J}_T(\theta_0) = \mathcal{J}(\theta_0) + o_p(1)$, the result in Lemma 2.1 follows by applying Assumption 3.

Lemma A.1. Let v be the k-dimensional normal random variable with mean γ and covariance Ω , whose probability density function is

$$f(v;\gamma,\Omega) = (2\pi)^{-\frac{k}{2}} |\Omega|^{-\frac{1}{2}} \exp\{-\frac{1}{2}(v-\gamma)'\Omega^{-1}(v-\gamma)\},\$$

where Ω is assumed known and positive definite. Suppose $\varphi(v)$ is a test of

$$\mathsf{H}_0: \gamma = 0 \quad vs \quad \mathsf{H}_1: \gamma \in \mathfrak{C}_1 = \mathfrak{C} \setminus \{\gamma = 0\}.$$

If $\varphi(v)$ has the acceptance region E that is closed, convex and of lower set, then $\varphi(v)$ is admissible among the class of tests whose limiting acceptance region E' is not contained in $W^*(E)$ in the sense that there cannot exist a test belonging to such the class of tests that is asymptotically uniformly more powerful than our tests.

Proof. Let $\varphi'(v)$ be any other test that has the acceptance region that is a closed and convex lower set and is not contained in $W^*(E)$. We shall prove that if

$$\Pr(v \in E') \le \Pr(v \in E)$$

for all $\gamma \in \mathfrak{C}_1$, then E' must be contained in $W^*(E)$. This would imply that if $\varphi'(v)$ is such a test that satisfies

$$E_{\gamma}\varphi(v) \leq E_{\gamma}\varphi'(v), \quad \forall \gamma \in \mathfrak{C}_1,$$

then $E_{\gamma}\varphi'(v) = E_{\gamma}\varphi(v)$ for all $\gamma \in \mathcal{C}_1$.

The proof of $E' \subseteq W^*(E)$ is by contradiction. Suppose this is not the case, i.e., $E' \subsetneq W^*(E)$. We shall then show that

$$\Pr(v \in E') > \Pr(v \in E), \tag{A.6}$$

holds for some $\gamma \in \mathcal{C}_1$, which provides the required contradiction.

Let $\gamma = \gamma^0 + \lambda e$. If $\gamma \in \mathbb{C}_1$ for any $\gamma^0 \in \mathbb{C}$ as $\lambda \to \infty$, it implies $e \in \mathbb{C}_1$. If $E' \subsetneq W^*(E)$ it follows from the definition of $W^*(E)$ that there exists a halfspace $W_{e,\delta}$ such that $E \subset W_{e,\delta}$ and $E' \cap \overline{W}_{e,\delta} \neq \emptyset$, where $\overline{W}_{e,\delta}$ is the complement set of $W_{e,\delta}$. That is

$$E \cap \overline{W}_{e,\delta} = \emptyset, \tag{A.7}$$

$$\Pr(v \in E' \cap \overline{W}_{e,\delta}) > 0. \tag{A.8}$$

Because

$$= \frac{\int \{\varphi(v) - \varphi'(v)\} f(v;\gamma,\Omega) dv}{C(\gamma^0)} \exp(\delta\lambda) \cdot \int \{\varphi(v) - \varphi'(v)\} \exp\{\lambda(e'\Omega^{-1}v - \delta)\} f(v;\gamma^0,\Omega) dv,$$

where $C(\gamma) = \exp(-0.5\gamma'\Omega^{-1}\gamma)$, which is bounded with a positive definite Ω . Let the integral be $I^- + I^+$, where I^- and I^+ denote the contributions over the integration regions $W_{e,\delta}$ and $\overline{W}_{e,\delta}$, respectively. Since $e'\Omega^{-1}v < \delta$ in the region $W_{e,\delta}$, $\exp\{\lambda(e'\Omega^{-1}v - \delta)\} \to -\infty$ as $\lambda \to \infty$, hence I^- is bounded as $\lambda \to \infty$. Therefore, we may only need to show $I^+ \to \infty$ as $\lambda \to \infty$. When $e'\Omega^{-1}v > \delta$ it follows from (A.7) that $\varphi(v) = 1$. Because $\varphi'(v)$ is bounded by 1 it follows that $\varphi(v) - \varphi'(v) \ge 0$. Therefore, by (A.8)

$$\Pr\{\varphi(v) - \varphi'(v) > 0 \text{ and } e'\Omega^{-1}v > \delta\} > 0.$$

Finally, when $e'\Omega^{-1}v > \delta$ it follows that $\exp\{\lambda(e'\Omega^{-1}v - \delta)\} \to \infty$ as $\lambda \to \infty$. This shows $I^+ \to \infty$ as $\lambda \to \infty$, hence (A.6).

Lemma A.2. Let $p_{mi}(v)$ be the test statistic of MinP score tests constructed analogous to (3.3),

but based on the random variable v. Let c_{α} be the critical value corresponding $p_{mj}(v)$. Let

$$\varphi_{mj}(v) = \mathbb{1}(p_{mj}(v) < c_{\alpha}),$$

where $0 < c_{\alpha} < 0.5$. Then, the acceptance region $\{v : p_{mj} \ge c_{\alpha}\}$ is a closed and convex lower set.

Proof. Let $t_g(v)$, g = c, t, be the test statistics constructed analogous to \tilde{t}_g based on v. Similarly, define by $p_g(v)$ the p-values analogous to \tilde{p}_g . We shall first show that the set $\{v : p_c(v) \ge c_c\}$, where c_c is a constant, and $\{v : p_t(v) \ge \alpha\}$ for $\alpha \in (0, 1)$ is a closed and convex lower set.

For g = c we have

$$t_c(v) = \bar{v}' \Omega^{-1} \bar{v},$$

where

$$\bar{v} = \arg \inf_{u \in \mathcal{C}} (v - u)' \Omega^{-1} (v - u)$$

Because Ω is of positive definite it follows $t_c(v) \ge 0$. Let

$$p_c(t_c(v)) = \sum_{m=1}^k \pi_m \{ 1 - F_{\chi_m^2}(t_c(v)) \}.$$

Clearly, $p_c(v)$ is a decreasing function of $t_c(v)$. It follows that the set $E_c = \{v : p_c(v) \ge \alpha\}$, where $0 < \alpha < 0.5$, is equivalent to the set $E_c = \{v : t_c(v) \le c_1\}$, where c_1 satisfies $p_c(c_1) = \alpha$.

Let A be a matrix with the row vector $a_i \in \mathbb{R}^k$, $i \in K$, such that $A'A = \Omega^{-1}$. Thus

$$E_c = \{v : (A\bar{v})'A\bar{v} \le c_1\}$$

is the intersection of the halfspaces $a_i \bar{v} \leq d_i$, $\forall i \in K$, where $d = (d_1, ..., d_k)'$ and $(d'd)^{1/2} = c_1$. Therefore, E_c is of closed, covex and lower set.

For g = t we have

$$t_t(v) = b' \Omega^{-1} v,$$

where

$$b \in \{b \in R^k : b \in \mathcal{C}_1, b' \Omega^{-1} b = 1\},\$$

which is computed by Algorithm 1 in Lu (2016) using Ω that ensures $\Omega^{-1}b \in C_1$ (see Remark 7 of Lu (2016) for a discussion). Therefore, $t_t(v)$ is an increasing function of elements of v. For $p_t(v) = 1 - \Phi(t_t(v))$ it follows that $E_t = \{v : p_t(v) \ge \alpha\}$ is equivalent to the halfspace $E_t = \{v : t_t(v) \le c_2\}$, where c_2 satisfies $1 - \Phi(c_2) = \alpha$. Therefore, E_t is of closed, convex and lower set.

Obviously, the region

$$E_{mk} = \{ v : p_i(v) = 1 - \Phi(v_i / \Omega_{ii}) \ge c_3, \forall i \in K \}$$

is also of closed, convex and lower set. Because at a given $\alpha \in (0, 0.5)$

$$\{v: p_g(v) \ge \alpha\} \cap \{v: p_i(v) \ge \alpha, \forall i \in K\} \neq \emptyset,\$$

the acceptance region of $\varphi_{mj}(v)$ is the intersection of E_g , g = c, t, and E_{mk} , hence it is of closed, convex and lower set.

Proof of Theorem 3.1. It follows from Lemma 2.1 that for each $\theta \in \Theta_{\epsilon}(\bar{\theta}_0, k) \setminus \{\gamma = 0\}$

$$\lim_{T \to \infty} E_{\theta} \varphi_{mj}(Y_T) = 1 - \int_E f(v; \gamma, \mathcal{G}_{\gamma\gamma}(\theta)) dv,$$
$$\lim_{T \to \infty} E_{\theta} \varphi'_{mj}(Y_T) = 1 - \int_{E'} f(v; \gamma, \mathcal{G}_{\gamma\gamma}(\theta)) dv,$$

for g = c, t. By Lemma A.2 the acceptance region E for the test $\varphi_{mj}(v)$ is a closed and convex lower set. Therefore, for any $\varepsilon > 0$ it follows from Lemma A.1 that there does not exist a distinct test $\varphi'_{mj}(Y_T)$ with the limiting acceptance region E' not being contained in $W^*(E_{mj})$ such that for all $\theta \in \Theta_{\epsilon}(\bar{\theta}_0, k) \setminus \{\gamma = 0\}$

$$\liminf_{T \to \infty} E_{\theta} \varphi'_{mj}(Y_T) \ge \limsup_{T \to \infty} E_{\theta} \varphi_{mj}(Y_T) + \varepsilon.$$

The result follows by letting $\varepsilon \to 0$.

Proof of Theorem 4.1. When $K_0 \neq \emptyset$ let \hat{i} be the (random) index i in Algorithm 2 such that $\tilde{p}_{\hat{i}}$ is

the smallest in $\{\tilde{p}_i, i \in K_0\}$ that is rejected. This implies that $K_i \supseteq K_0$. Without loss of generality let K_0 contains the first k_0 elements of K. Because the event that a false rejection occurs is the event

$$\min(\tilde{p}_i, i \in K_0) \le \tilde{c}_{|K_i|}(\alpha),$$

it follows that for any $\varepsilon > 0$

$$\limsup_{T \to \infty} \sup_{\theta \in \bar{\Theta}_{\epsilon}(\bar{\theta}_{0}, k_{0})} \Pr\{\min(\tilde{p}_{i}, i \in K_{0}) \leq \tilde{c}_{|K_{i}|}(\alpha)\}$$

$$\leq \limsup_{T \to \infty} \sup_{\theta \in \bar{\Theta}_{\epsilon}(\bar{\theta}_{0}, k_{0})} \Pr\{\min(\tilde{p}_{i}, i \in K_{0}) \leq c_{K_{0}}(\alpha) + \varepsilon\}$$
(A.9)

By the result of Lemma 2.1 the distribution of $\min(\tilde{p}_i, i \in K_0)$ under $\theta_0 \in \bar{\Theta}_{\epsilon}(\bar{\theta}_0, k_0)$ weakly converges to $\min(p_i, i \in K_0)$, where $p_i = 1 - \Phi(v_i)$ and $\{v_i, i \in K_0\}$ has the multivariate normal distribution with the mean 0 and the covariance matrix that is the correlation matrix of $\mathcal{G}(\theta_0)$ corresponding to the elements $i \in K_0$. Therefore, the right hand side of (A.9) is bounded above by α , thus we have $\limsup_{T \to \infty} \sup_{\theta \in \bar{\Theta}_{\epsilon}(\bar{\theta}_0, k_0)} FWER \leq \alpha$.

Proof of Theorem 4.2. The result stated in (i) suggests that the probability for any $i \in K_0$ being rejected is asymptotically locally bounded above by α . That requires $\limsup_{T\to\infty} \sup_{\theta\in\bar{\Theta}_{\epsilon}(\bar{\theta}_0,k_0)}$ FWER $\leq \alpha$. The result in (ii) follows because for those $i \in \bar{K}_0 \neq \emptyset$ H_{0i} is rejected with probability 1 as $T \to \infty$.

Proof of Proposition 1. Let $Y = (y_1, ..., y_T)'$, $Z = (z_1, ..., z_T)'$, $X = (x_1, ..., x_T)'$ and $\epsilon = (\epsilon_1, ..., \epsilon_T)'$. Let D = (Z, X) and $T^{-1}D'D \xrightarrow{p} \Omega$, where Ω is positive definite, implied by Assumption 1. Let Z further partitioned into $Z = (Z_1, Z_2)$ and rewrite the model as $Y = Z_1\gamma_1 + Z_2\gamma_2 + X\beta + \epsilon$. The constrained OLS estimate under $\overline{\Theta}(k_1)$ of $\theta = (\gamma'_1, \lambda')'$, where $\lambda = (\gamma'_2, \beta')'$, are $\tilde{\gamma}_1(k_1) = 0$, $\tilde{\lambda}(k_1) = (D'_2D_2)^{-1}D'_2Y$, where $D_2 = (Z_2, X)$ and the estimated variance of the error term is $\tilde{\sigma}^2(k_1) = T^{-1}(Y - D_2\tilde{\lambda}_2(k_1))'(Y - D_2\tilde{\lambda}_2(k_1))$. It follows $s_{T,\gamma_1}(\tilde{\theta}(k_1)) = \tilde{\sigma}^{-2}(k_1)Z'_1\{Y - D_2\tilde{\lambda}(k_1)\}$ and $\mathcal{J}_{T,\theta\theta}(\tilde{\theta}(k_1)) = \tilde{\sigma}^{-2}(k_1)D'D$. Thus, $U_{T,\gamma_1}(\tilde{\theta}(k_1)) = \mathcal{J}^{-1}_{T,\gamma_1}(\tilde{\theta}(k_1))s_{T,\gamma_1}(\tilde{\theta}(k_1))$, where $\mathcal{J}^{-1}_{T,\gamma_1}(\tilde{\theta}(k_1))$ is the block corresponding to γ_1 in $\mathcal{J}^{-1}_{T,\theta\theta}(\tilde{\theta}(k_1)) = \tilde{\sigma}^2(k_1)(D'D)^{-1}$.

Since $\tilde{\sigma}^2(k_1)$ can be viewed as unconstrained OLS estimate of $Y = Z_2 \gamma_2 + X\beta + \epsilon$ and $\tilde{\sigma}^2(k)$ as unconstrained OLS estimate of $Y = X\beta + \epsilon$, it is a well known result that $\tilde{\sigma}^2(k_1) \leq \tilde{\sigma}^2(k)$.

Suppose H_{0i} : $\gamma_i = 0$ is true for $i \in K_0 = K_1 \subseteq K$ and $\tilde{\sigma}^2(k_1) \xrightarrow{p} \sigma_0^2$. Following Lemma (2.1) the null distribution of $U_{T,\gamma_1}(\tilde{\theta}(k_1))$ weakly converges to the centred multivariate normal distribution with the covariance $\mathcal{J}_{\gamma_1}^{-1}(\theta_0)$ that is the block of $\sigma_0^2 \Omega^{-1}$ corresponding to γ_1 . Therefore, $\min(\tilde{p}_i(k_0), i \in K_0)$ weakly converges to $\min(p_i(k_0), i \in K_0)$, where $p_i = 1 - \Phi(v_i)$ and $\{v_i, i \in K_0\}$ has the multivariate normal distribution with the mean 0 and the covariance matrix that is the correlation matrix of $\mathcal{J}_{T,\gamma_1}^{-1}(\theta_0)$. Thus, the α th quantile denoted by $\tilde{c}_{m,K_0}(\alpha)$ of the distribution of $\min(\tilde{p}_i, i \in K_0)$ based on the empirical distribution $\{\tilde{p}_{m,K_0}^b, b = 1, ..., B\}$, where $\tilde{p}_{m,K_0}^b = \min(\tilde{p}_{mi}^b, i \in K_0)$ in Algorithm 2 converges to the α th quantile $c_{K_0}(\alpha)$ of the distribution of $\min(p_i, i \in K_0)$. That is,

$$\tilde{c}_{K_0}(\alpha) \to c_{K_0}(\alpha).$$

Because $\tilde{\sigma}^2(k_1) \leq \tilde{\sigma}^2(k)$ it is obvious that $\tilde{c}_{K_i}(\alpha) \leq \tilde{c}_{K_0}(\alpha)$ for $K_i \supseteq K_0$. Hence it suffices to prove the first part of the result.

Because $\tilde{\sigma}^2(k_1) \leq \tilde{\sigma}^2(k)$ holds true whenever $k_1 \leq k$, this implies that $\tilde{c}_{K_i}(\alpha)$ based on the estimate $\tilde{\theta}(k)$ is no greater than that based on the estimate $\tilde{\theta}(k_1)$. Hence it suffices to prove the second part of the result.

Lemma A.3. Let $v = (v_1, ..., v_k)'$ be the k-dimensional normal random variable with mean 0 and covariance $\{\rho_{ij}\}$ and let $u = (u_1, ..., u_k)'$ be the k-dimensional normal random variable with mean 0 and covariance $\{\xi_{ij}\}$. Let $\rho_{ii} = \xi_{ii} = 1$, $i \in K$. If $\rho_{ij} > \xi_{ij}$, $i, j \in K$, then

$$\Pr(v_1 > c_1, ..., v_k > c_k) \ge \Pr(u_1 > c_1, ..., u_k > c_k).$$

Proof. The result was established by Slepian (1962). See also Gupta (1963) for a proof. \Box

Lemma A.4. For v and u defined in Lemma A.3 let $p_{vi} = \Pr(v_i > c_i) = 1 - \Phi(c_i)$ and $p_{u_i} = \Pr(u_i > c_i) = 1 - \Phi(c_i)$, then

$$\Pr\{\min(p_{v_i}, i \in K) < c\} \ge \Pr\{\min(p_{u_i}, i \in K) < c\}$$

Proof. For v we have

$$\{v \in R^k : \min(p_{v_i}, i \in K) < c\} = \{v \in R^k : \min(p_{v_i}, i \in K) < c'\},\$$

hence

$$\Pr\{\min(p_{v_i}, i \in K) < c\} \\ = \Pr\{\max(v_i, i \in K) > c\} \\ = \Pr(v_1 > c, ..., v_k > c).$$

Similarly,

$$\Pr\{\min(p_{u_i}, i \in K) < c\} = \Pr(u_1 > c, ..., u_k > c).$$

The result then follows from Lemma A.3.

Proof of Proposition 2. If $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$ is a non-positive matrix, so is the correlation matrix corresponding to $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$. Let u be the k-dimensional centre normal distributions with the covariance matrix that is the correlation matrix corresponding to $\mathcal{G}_{\gamma_1\gamma_1}(\theta_0)$. Let v be the k-dimensional centre normal distributions with the covariance matrix that is the k-dimensional identity matrix. Let

$$\Pr\{\min(p_{v_i}, i \in K) < c_{m1}(\alpha)\} = \Pr\{\min(p_{u_i}, i \in K) < c_K(\alpha)\} = \alpha.$$

By Lemma A.4

$$c_{m1}(\alpha) \le c_K(\alpha) \tag{A.10}$$

For any $K_1 \subseteq K_2$ it is always true that $\min(p_{u_i}, i \in K_1) \ge \min(p_{u_i}, i \in K_2)$ for each value of u, hence, for $c_{K_1}(\alpha)$ and $c_{K_2}(\alpha)$ satisfying

$$\Pr\{\min(p_{u_i}, i \in K_1) < c_{K_1}(\alpha)\} = \Pr\{\min(p_{u_i}, i \in K_2) < c_{K_2}(\alpha)\}\$$

it follows

$$c_{K_1}(\alpha) \le c_{K_2}(\alpha). \tag{A.11}$$

Therefore, form (A.10) and (A.11) it follows for $K_0 \subseteq K_i \subseteq K$

$$c_{K_0}(\alpha) \le c_{K_i}(\alpha) \le c_{m1}(\alpha).$$

As $\tilde{c}_{K_i}(\alpha) \to c_{K_i}(\alpha)$, $K_i \subseteq K$, and $\tilde{c}_{m1}(\alpha) \leq \tilde{c}_{m2}(\alpha)$ as revealed in Remark 8 it suffices to prove the result.

Proof of Proposition 3. The proof is similar to the proof of Proposition 2. \Box

References

- Aitchison, J., Silvey, S.D., 1958. Maximum-likelihood estimation of parameters subject to restraints. The Annals of Mathematical Statistics 29, 813–828.
- Andrews, D.W.K., 1996. Admissibility of the likelihood ratio test when the parameter space is restricted under the alternative. Econometrica 64, 705–718.
- Andrews, D.W.K., 2001. Testing when a parameter is on the boundary of the maintained hypothesis. Econometrica 69, 683–734.
- Andrews, D.W.K., Moreira, M.J., Stock, J.H., 2006. Optimal two-sided invariant similar tests for instrumental variables regression. Econometrica 74, 715–752.
- Andrews, D.W.K., Ploberger, W., 1995. Admissibility of the likelihood ratio test when a nuisance parameter is present only under the alternative. The Annals of Statistics, 1609–1629.
- Birnbaum, A., 1955. Characterizations of complete classes of tests of some multiparametric hypotheses, with applications to likelihood ratio tests. The Annals of Mathematical Statistics , 21–36.

- Cavaliere, G., Nielsen, H.B., Pedersen, R.S., Rahbek, A., 2020. Bootstrap inference on the boundary of the parameter space, with application to conditional volatility models. Journal of Econometrics, available online.
- Chernozhukov, V., Hansen, C., Jansson, M., 2009. Admissible invariant similar tests for instrumental variables regression. Econometric Theory 25, 806–818.
- Demos, A., Sentana, E., 1998. Testing for GARCH effects: a one-sided approach. Journal of Econometrics 86, 97–127.
- Finner, H., Strassburger, K., 2002. The partitioning principle: a powerful tool in multiple decision theory. Annals of Statistics 30, 1194–1213.
- Francq, C., Zakoian, J.M., 2009. Testing the nullity of garch coefficients: Correction of the standard tests and relative efficiency comparisons. Journal of the American Statistical Association 104, 313–324.
- Gouriéroux, C., Holly, A., Monfort, A., 1982. Likelihood ratio test, Wald test, and Kuhn-Tucker test in linear models with inequality constraints on the regression parameters. Econometrica 50, 63–80.
- Gupta, S.S., 1963. Probability integrals of multivariate normal and multivariate t. The Annals of mathematical statistics 34, 792–828.
- Ketz, P., 2018. Subvector inference when the true parameter vector may be near or at the boundary. Journal of econometrics 207, 285–306.
- King, M., Wu, P., 1997. Locally optimal one-sided tests for multiparameter hypotheses. Econometric Reviews 16, 131–156.
- Lee, J.H.H., King, M.L., 1993. A locally most mean powerful based score test for arch and garch regression disturbances. Journal of Business & Economic Statistics 11, 17–27.
- Lehmann, E.L., Romano, J.P., 2005. Testing Statistical Hypotheses. Springer, New York. 3rd edition.

- Lu, Z.H., 2013. Halfline tests for multivariate one-sided alternatives. Computational Statistics & Data Analysis 57, 479–490.
- Lu, Z.H., 2016. Extended maxt tests of one-sided hypotheses. Journal of the American Statistical Association 111, 423–437.
- Marden, J.I., 1982. Combining independent noncentral chi squared or f tests. The Annals of Statistics 10, 266–277.
- Marden, J.I., 1985. Combining independent one-sided noncentral t or normal mean tests. The Annals of Statistics, 1535–1553.
- McKeague, I.W., Qian, M., 2015. An adaptive resampling test for detecting the presence of significant predictors. Journal of the American Statistical Association 110, 1422–1433.
- Rao, C.R., 1948. Large sample tests of statistical hypotheses concerning several parameters with applications to problems of estimation, in: Mathematical Proceedings of the Cambridge Philosophical Society, Cambridge Univ Press. pp. 50–57.
- Romano, J.P., Shaikh, A.M., Wolf, M., 2010. Hypothesis testing in econometrics. Annual Review of Economics 2, 75–104.
- Romano, J.P., Shaikh, A.M., Wolf, M., 2011. Consonance and the closure method in multiple testing. The International Journal of Biostatistics 7, 1–25.
- Romano, J.P., Wolf, M., 2005. Exact and approximate stepdown methods for multiple hypothesis testing. Journal of the American Statistical Association 100, 94–108.
- Silvapulle, M.J., Sen, P.K., 2005. Constrained Statistical Inference: Order, Inequality, and Shape Restrictions. Wiley, New York.
- Silvapulle, M.J., Silvapulle, P., 1995. A score test against one-sided alternatives. Journal of the American Statistical Association 90, 342–349.
- Silvey, S.D., 1959. The lagrangian multiplier test. The Annals of Mathematical Statistics 30, 389–407.

- Slepian, D., 1962. The one-sided barrier problem for gaussian noise. Bell System Technical Journal 41, 463–501.
- Stein, C., 1956. The admissibility of hotelling's t^2 -test. The Annals of Mathematical Statistics 27, 616–623.
- Westfall, P.H., Young, S.S., 1993. Resampling-based multiple testing: Examples and methods for p-value adjustment. John Wiley & Sons, Ltd., New York.
- Wolak, F.A., 1989a. Local and global testing of linear and nonlinear inequality constraints in nonlinear econometric models. Econometric Theory 5, 1–35.
- Wolak, F.A., 1989b. Testing inequality constraints in linear econometric models. Journal of Econometrics 41, 205–235.



Figure 1: Illustration of non-rejection region (E) and $W^*(E)$ for MinP-sc tests.

γ'	T		MinP	-sc			MinP	-st			MinF	P_{-S}		$\bar{\chi}^2$	t
		H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	H_0
							ρ	= -(0.45						
$(0 \ 0)$	60	4.7	3.9	2.3	1.7	4.8	3.4	2.1	1.4	4.5	4.5	2.6	2.0	5.0	5.6
	100	4.2	3.8	2.0	1.9	4.2	3.3	1.9	1.5	4.5	4.5	2.5	2.1	4.5	4.8
$(0.3 \ 0)$	60	68.0	2.7	51.7	2.7	68.7	2.4	48.8	2.4	57.9	3.1	55.6	3.1	73.8	69.5
	100	95.4	2.7	86.0	2.7	95.5	2.5	84.8	2.5	89.2	3.0	88.2	3.0	97.1	94.9
$(0.15 \ 0.15)$	60	52.3	0	13.7	17.3	56.9	0	12.6	16.2	34.1	0	15.5	19.7	65.1	68.1
	100	78.2	0	33.8	26.6	82.1	0	31.9	24.9	57.1	0	36.9	28.4	87.2	89.1
								$\rho =$	0						
$(0 \ 0)$	60	4.8	4.6	2.4	2.3	4.7	4.1	2.1	2.1	4.9	4.9	2.6	2.4	4.5	5.3
	100	4.6	4.4	1.9	2.7	5.0	4.0	1.8	2.4	4.7	4.7	2.1	2.8	4.3	5.0
$(0.3 \ 0)$	60	44.4	3.2	42.6	3.2	44.5	3.1	40.8	3.1	44.4	3.3	43.4	3.3	44.4	35.3
	100	78.2	3.9	77.9	3.9	77.9	3.9	76.0	3.9	78.9	4.0	78.6	4.0	78.4	57.2
$(0.15 \ 0.15)$	60	61.3	0	32.5	39.0	67.0	0	30.7	36.9	56.6	0	33.1	39.8	68.5	74.9
	100	71.6	0	52.1	48.6	75.9	0	50.0	47.0	70.8	0	52.8	49.5	77.0	81.9
							ĥ	p = 0	.45						
$(0 \ 0)$	60	6.3	5.8	3.6	3.2	6.0	5.9	3.6	3.3	6.1	6.1	3.8	3.4	6.0	5.5
	100	5.8	5.7	3.4	3.3	6.1	5.6	3.4	3.3	6.0	6.0	3.6	3.6	5.4	5.2
$(0.3 \ 0)$	60	66.3	4.1	61.4	4.1	61.7	4.1	61.2	4.1	63.1	4.2	62.9	4.2	70.7	37.1
	100	83.7	3.7	76.4	3.7	76.7	3.7	76.6	3.7	78.0	3.7	77.9	3.7	87.0	45.8
$(0.15 \ 0.15)$	60	58.4	0	34.8	46.1	61.2	0	32.9	43.5	58.2	0	35.1	46.6	60.1	66.5
·	100	69.7	0	55.4	57.1	73.4	0	54.8	56.6	70.7	0	55.8	57.9	70.1	79.7

Table 1: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in linear models with ϵ_n following N(0, 1).

γ'	T		MinP	-sc			MinP	-st			MinF	'- S		$\bar{\chi}^2$	t
		H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	H_0
							ρ	= -().45						
$(0 \ 0)$	60	5.1	4.3	2.4	2.0	5.0	3.6	1.9	1.8	5.2	5.2	3.0	2.3	5.0	5.1
	100	4.8	4.3	2.2	2.3	5.1	4.0	2.0	2.1	4.9	4.9	2.4	2.7	4.9	4.8
$(0.6 \ 0)$	60	98.2	2.2	95.8	2.2	98.3	2.1	95.3	2.1	96.7	2.4	96.5	2.4	98.8	96.9
	100	99.6	2.5	96.4	2.5	99.7	2.3	95.8	2.3	97.4	2.6	97.0	2.6	99.8	99.4
$(0.4 \ 0.4)$	60	98.7	0	64.9	73.2	99.3	0	63.8	71.9	92.5	0	66.5	75.2	99.6	99.8
	100	99.6	0	77.6	90.9	99.8	0	77.3	90.3	98.1	0	78.4	91.6	99.9	99.9
								$\rho =$	0						
$(0 \ 0)$	60	5.3	5.1	2.6	2.7	5.4	4.4	2.2	2.4	5.1	5.1	2.6	2.8	5.0	5.2
	100	5.3	4.8	2.8	2.2	5.4	4.1	2.4	1.9	5.0	5.0	2.9	2.3	5.4	5.7
$(0.6 \ 0)$	60	94.5	3.4	94.4	3.4	94.6	3.4	93.7	3.4	94.7	3.4	94.7	3.4	95.1	80.7
	100	99.0	3.6	98.8	3.6	99.0	3.6	98.5	3.6	99.0	3.6	98.9	3.6	99.1	96.3
$(0.4 \ 0.4)$	60	88.0	0	61.4	82.6	90.1	0	60.5	81.3	87.9	0	61.6	82.8	89.2	92.7
	100	97.7	0	86.5	88.4	98.3	0	86.1	87.9	96.6	0	86.9	88.5	98.5	99.3
							ĥ	p = 0	.45						
$(0 \ 0)$	60	4.8	4.2	2.7	2.9	4.8	4.6	3.0	3.1	5.0	5.0	3.2	3.3	5.0	5.5
	100	5.6	5.0	3.4	3.1	5.8	5.4	3.7	3.4	5.8	5.8	4.0	3.5	5.4	5.7
$(0.6 \ 0)$	60	96.5	3.3	94.1	3.3	94.2	3.3	94.2	3.3	94.5	3.3	94.5	3.3	97.9	63.3
	100	99.5	3.2	98.6	3.2	98.6	3.2	98.6	3.2	98.8	3.2	98.8	3.2	99.8	75.9
$(0.4 \ 0.4)$	60	78.2	0	68.0	70.7	80.9	0.0	68.0	71.0	79.9	0	69.0	72.0	75.8	85.2
	100	91.6	0	86.4	84.6	93.2	0.0	86.5	84.7	92.4	0	86.8	85.2	90.6	94.9

Table 2: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in linear models with ϵ_n following t_5 .

γ'	T		MinP	-sc			MinP	-st			MinF	°- s		DS	LK
		H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	H_0
							u_n	$\sim N$	(0, 1)						
$(0 \ 0)$	60	6.3	6.3	2.7	3.7	6.5	5.9	2.5	3.5	6.3	6.3	2.7	3.7	5.9	5.9
	100	5.3	5.2	2.6	2.8	5.3	4.1	2.2	2.0	5.3	5.3	2.7	2.8	6.1	5.3
$(0.6 \ 0)$	60	56.5	4.2	54.8	4.2	58.7	4.0	52.9	4.0	56.5	4.2	54.9	4.2	59.5	58.8
	100	79.1	6.4	78.2	6.4	80.1	6.2	75.9	6.2	78.9	6.4	78.2	6.4	78.4	81.0
$(0 \ 0.6)$	60	59.6	2.9	2.9	58.0	60.6	2.6	2.6	56.2	59.6	2.9	2.9	58.0	52.5	58.1
	100	81.3	3.9	3.9	80.4	81.3	3.8	3.8	79.1	81.2	3.9	3.9	80.4	73.1	79.9
$(0.5 \ 0.3)$	60	62.6	0	43.3	28.7	70.2	0	41.3	27.2	61.7	0	43.3	28.7	76.2	66.8
	100	84.5	0	67.1	41.2	89.9	0	65.9	40.1	83.7	0	67.1	41.2	92.7	88.0
$(0.3 \ 0.5)$	60	63.7	0	27.8	44.5	70.7	0	25.8	42.9	62.8	0	27.8	44.6	76.2	66.8
	100	86.8	0	48.9	64.7	90.8	0	47.8	63.3	86.3	0	48.9	64.7	92.2	89.4
$(0.4 \ 0.4)$	60	62.5	0	35.4	35.7	70.7	0	33.8	34.5	61.2	0	35.4	35.7	77.5	65.9
	100	84.9	0	59.2	52.1	89.5	0	57.6	51.0	83.9	0	59.2	52.1	92.5	87.9
							1	$u_n \sim$	t_5						
$(0 \ 0)$	60	4.5	4.5	2.0	2.6	4.6	4.0	1.8	2.2	4.5	4.5	2.0	2.6	4.5	4.3
	100	5.1	5.1	2.7	2.6	5.3	4.7	2.5	2.4	5.1	5.1	2.7	2.6	5.4	4.9
$(0.6 \ 0)$	60	56.1	5.0	53.3	5.0	60.1	4.7	51.6	4.7	55.5	5.0	53.3	5.0	63.9	59.5
	100	76.6	8.9	75.0	8.9	79.1	8.9	73.5	8.9	76.4	8.9	75.0	8.9	81.1	78.4
$(0 \ 0.6)$	60	58.8	2.2	2.2	57.5	58.7	2.0	2.0	55.7	58.7	2.2	2.2	57.5	54.7	56.4
	100	77.5	3.3	3.3	76.4	77.9	3.2	3.2	75.5	77.4	3.3	3.3	76.4	72.7	76.5
$(0.5 \ 0.3)$	60	66.2	0	45.1	31.1	72.5	0	43.3	29.7	65.4	0	45.1	31.1	79.1	69.4
	100	83.9	0	65.0	42.6	89.3	0	64.1	41.8	83.3	0	65.0	42.6	94.1	86.7
$(0.3 \ 0.5)$	60	64.5	0	32.6	42.6	72.6	0	31.2	41.2	64.0	0	32.6	42.6	78.8	68.1
	100	83.6	0	49.6	59.1	89.0	0	48.8	58.3	83.1	0	49.6	59.1	92.9	86.8
$(0.4 \ 0.4)$	60	64.5	0	39.4	36.0	72.4	0	37.7	34.7	64.1	0	39.4	36.0	80.3	68.8
	100	84.1	0	58.0	51.7	89.4	0	57.0	50.5	83.5	0	58.0	51.7	93.5	86.5

Table 3: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in ARCH(2) models.

γ'	Т		N	MinP	-sc				l	MinP	-st				•	MinF	P-8			DS	LK
		H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	H_0
										u_n	$\sim N$	(0, 1)									
$(0 \ 0 \ 0 \ 0)$	60	6.0	6.0	1.6	1.9	1.5	1.2	6.1	5.4	1.4	1.6	1.5	1.1	6.0	6.0	1.6	1.9	1.6	1.2	5.3	6.3
	100	5.2	5.1	1.1	1.3	1.4	1.6	5.7	4.6	1.1	1.3	1.3	1.3	5.1	5.1	1.1	1.3	1.4	1.6	5.6	5.5
$(0.6 \ 0 \ 0 \ 0)$	60	46.7	6.9	42.6	3.1	2.3	1.6	47.1	6.7	41.1	3.0	2.3	1.5	46.7	6.9	42.6	3.1	2.3	1.6	43.0	49.2
	100	69.1	7.9	66.8	3.7	3.0	1.5	70.0	7.7	65.8	3.6	3.0	1.5	69.0	7.9	66.8	3.7	3.0	1.5	60.4	70.6
$(0 \ 0.6 \ 0 \ 0)$	60	47.4	5.8	1.5	43.0	1.1	3.3	49.0	5.5	1.4	42.1	1.1	3.1	47.2	5.8	1.5	43.0	1.1	3.3	43.7	46.9
	100	70.0	7.4	2.0	66.6	1.6	4.0	70.9	7.3	1.9	65.8	1.6	4.0	69.8	7.4	2.0	66.6	1.6	4.0	60.2	70.0
$(0 \ 0 \ 0 \ 0.6)$	60	48.4	3.9	1.1	1.9	1.2	46.0	49.0	3.5	0.9	1.6	1.1	45.0	48.5	3.9	1.1	1.9	1.2	46.0	33.4	44.4
	100	71.2	5.0	1.7	2.1	1.7	69.4	71.7	4.8	1.6	2.0	1.6	68.3	71.2	5.0	1.7	2.1	1.7	69.4	51.8	68.5
$(0.4 \ 0.2 \ 0 \ 0)$	60	39.0	5.3	25.6	11.8	3.6	1.8	43.5	5.2	24.5	11.4	3.5	1.8	38.9	5.3	25.6	11.8	3.6	1.8	52.0	44.2
	100	61.0	7.1	46.2	18.2	4.8	2.4	66.1	7.0	45.0	17.6	4.8	2.4	60.7	7.1	46.2	18.2	4.8	2.4	70.7	66.8
$(0 \ 0.2 \ 0 \ 0.6)$	60	50.4	1.8	0.8	12.7	1.1	39.0	54.7	1.7	0.8	12.3	1.0	38.1	50.3	1.8	0.8	12.7	1.1	39.0	51.1	49.7
	100	72.6	2.7	1.4	19.5	1.6	60.1	76.1	2.7	1.4	19.2	1.6	58.9	72.5	2.7	1.4	19.5	1.6	60.1	72.5	73.8
$(0.3\ 0.2\ 0.1\ 0.05)$	60	38.2	0	19.2	11.4	8.9	4.4	46.6	0	18.3	11.3	8.4	4.3	38.0	0	19.2	11.4	8.9	4.5	57.1	44.1
	100	57.7	0	34.3	18.0	13.0	6.5	68.3	0	33.8	17.3	12.2	6.3	57.1	0	34.3	18.0	13.0	6.5	76.6	66.3
										,	$u_n \sim$	t_5									
$(0 \ 0 \ 0 \ 0)$	60	5.6	5.6	1.3	1.7	1.2	1.6	5.4	5.2	1.2	1.7	1.1	1.4	5.6	5.6	1.3	1.7	1.2	1.6	4.2	4.5
	100	6.2	6.1	1.6	1.5	2.0	1.3	6.2	5.7	1.6	1.4	1.9	1.1	6.1	6.1	1.6	1.5	2.0	1.3	5.6	5.6
$(0.6 \ 0 \ 0 \ 0)$	60	45.9	8.7	41.3	3.8	3.9	1.2	47.9	8.7	40.1	3.8	3.9	1.2	45.8	8.7	41.3	3.8	3.9	1.2	50.0	49.0
	100	63.8	12.1	59.4	5.7	5.2	1.5	65.4	11.9	58.8	5.5	5.1	1.5	63.7	12.1	59.4	5.7	5.2	1.5	66.0	65.4
$(0 \ 0.6 \ 0 \ 0)$	60	43.8	5.8	1.2	39.6	1.0	3.9	45.9	5.6	1.2	38.6	1.0	3.8	43.8	5.8	1.2	39.6	1.0	3.9	47.9	44.4
	100	65.2	7.3	1.9	61.2	1.3	4.3	66.9	7.2	1.9	60.4	1.3	4.3	64.9	7.3	1.9	61.2	1.3	4.3	65.9	65.9
$(0 \ 0 \ 0 \ 0.6)$	60	48.0	3.4	0.8	1.4	1.3	45.6	48.3	3.3	0.8	1.3	1.3	44.9	48.0	3.4	0.8	1.4	1.3	45.6	36.7	43.4
	100	68.8	4.4	1.4	1.9	1.6	66.3	69.8	4.3	1.4	1.8	1.5	65.4	68.7	4.4	1.4	1.9	1.6	66.3	56.6	66.3
$(0.4 \ 0.2 \ 0 \ 0)$	60	44.1	8.1	27.7	13.6	5.9	2.3	49.5	7.8	27.0	13.1	5.7	2.3	44.0	8.1	27.7	13.6	5.9	2.3	62.6	48.2
	100	62.8	9.8	40.8	23.0	7.9	2.0	69.2	9.5	40.1	22.4	7.7	1.9	62.2	9.8	40.8	23.0	7.9	2.0	78.8	69.2
$(0 \ 0.2 \ 0 \ 0.6)$	60	48.8	2.1	1.0	14.2	1.2	35.5	53.5	1.9	0.9	13.6	1.1	35.0	48.7	2.1	1.0	14.2	1.2	35.5	55.5	49.3
	100	71.5	2.2	1.1	24.3	1.4	53.2	75.8	2.2	1.1	23.8	1.4	52.6	71.5	2.2	1.1	24.3	1.4	53.2	79.3	73.7
$(0.3 \ 0.2 \ 0.1 \ 0.05)$	60	44.6	0	21.0	12.7	12.1	6.2	55.2	0	20.6	12.3	11.8	6.0	44.3	0	21.0	12.7	12.1	6.2	70.4	51.5
	100	64.9	0	31.6	22.7	17.1	8.0	76.8	0	31.1	22.3	16.7	7.7	64.1	0	31.6	22.7	17.1	8.0	87.8	73.3

Table 4: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in ARCH(4) models.

γ'	Т	MinP-sc						I	MinP	-st					MinF	P-S			DS	LK
	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_0	H_0
									u_n	$\sim N$	(0, 1))								
$(5\ 0\ 0\ 0)$	60 48.3	6.9	44.6	3.2	2.4	1.6	48.8	6.8	43.1	3.2	2.3	1.5	48.3	6.9	44.6	3.2	2.4	1.6	46.6	51.0
	100 62.7	6.6	60.5	2.8	2.6	1.4	63.6	6.6	59.4	2.8	2.6	1.4	62.7	6.6	60.5	2.8	2.6	1.4	52.0	64.1
$(0 \ 5 \ 0 \ 0)$	60 49.0	5.8	1.3	45.0	1.1	3.7	50.8	5.7	1.2	43.8	1.1	3.7	49.0	5.8	1.3	45.0	1.1	3.7	46.8	49.7
	$100 \ 63.6$	6.6	2.0	60.3	2.0	2.9	64.5	6.4	2.0	59.1	1.8	2.9	63.4	6.6	2.0	60.3	2.0	2.9	52.3	63.2
$(0 \ 0 \ 0 \ 5)$	$60 \ 50.4$	3.3	0.7	1.7	1.0	48.1	50.9	3.2	0.6	1.7	1.0	47.0	50.3	3.3	0.7	1.7	1.0	48.1	35.0	47.1
	100 64.2	4.9	1.5	2.0	1.6	62.0	65.1	4.8	1.5	2.0	1.6	61.0	64.1	4.9	1.5	2.0	1.6	62.0	46.2	61.0
$(3 \ 1.5 \ 0 \ 0)$	$60 \ 38.6$	4.9	25.5	11.2	3.2	1.8	41.9	4.8	24.3	10.6	3.2	1.7	38.3	4.9	25.5	11.2	3.2	1.8	50.6	43.8
	100 48.9	4.8	36.8	13.7	3.1	1.8	52.7	4.6	35.7	13.3	2.9	1.8	48.6	4.8	36.8	13.7	3.1	1.8	54.9	53.6
$(0 \ 1.5 \ 0 \ 5)$	60 52.4	2.0	1.0	12.4	1.1	41.9	56.1	2.0	1.0	12.0	1.1	40.7	52.3	2.0	1.0	12.4	1.1	41.9	53.0	51.8
	$100 \ 66.3$	3.0	1.8	14.7	1.4	56.7	69.3	3.0	1.8	14.5	1.4	56.1	66.2	3.0	1.8	14.7	1.4	56.7	63.7	67.3
$(2.5 \ 1.5 \ 0.7 \ 0.4)$	$60 \ 39.7$	0	20.3	11.4	8.7	5.1	47.8	0	20.0	11.2	8.4	4.9	39.3	0	20.3	11.4	8.7	5.1	56.9	44.8
	100 47.8	0	30.4	13.5	8.8	4.1	55.8	0	29.3	13.1	8.5	4.0	47.3	0	30.4	13.5	8.8	4.1	62.7	54.6
										$u_n \sim$	t_5									
$(5\ 0\ 0\ 0)$	60 47.5	9.1	42.6	4.0	4.3	1.3	49.7	8.8	41.6	3.8	4.2	1.3	47.4	9.1	42.6	4.0	4.3	1.3	52.2	49.8
	100 59.4	9.9	55.8	5.0	4.1	1.2	60.9	9.9	55.0	5.0	4.1	1.2	59.4	9.9	55.8	5.0	4.1	1.2	59.6	61.1
$(0\ 5\ 0\ 0)$	$60 \ 45.6$	6.4	1.6	40.9	1.0	4.1	48.0	6.1	1.5	39.8	0.9	4.0	45.6	6.4	1.6	40.9	1.0	4.1	50.6	45.1
	100 61.3	6.5	1.8	57.4	1.5	3.4	62.5	6.4	1.8	56.3	1.4	3.4	61.1	6.5	1.8	57.4	1.5	3.4	60.2	61.8
$(0 \ 0 \ 0 \ 5)$	60 49.7	3.3	0.9	1.1	1.4	47.4	50.3	3.2	0.8	1.1	1.4	46.7	49.7	3.3	0.9	1.1	1.4	47.4	38.0	45.2
	$100 \ 63.6$	4.8	1.6	1.9	1.7	60.8	64.9	4.6	1.6	1.7	1.7	60.4	63.7	4.8	1.6	1.9	1.7	60.9	51.3	61.1
$(3 \ 1.5 \ 0 \ 0)$	60 43.0	8.7	27.5	12.3	6.2	2.8	48.2	8.3	26.9	12.0	5.9	2.6	42.9	8.7	27.5	12.3	6.2	2.8	60.5	47.7
	$100\ 51.9$	7.0	34.4	17.7	5.4	1.6	57.7	6.9	33.6	17.1	5.3	1.6	51.5	7.0	34.4	17.7	5.4	1.6	67.6	58.2
$(0 \ 1.5 \ 0 \ 5)$	$60 \ 51.3$	1.8	0.8	14.6	1.0	38.1	54.9	1.7	0.8	14.2	1.0	37.3	51.3	1.8	0.8	14.6	1.0	38.1	55.5	50.6
	100 67.3	2.8	1.3	18.6	1.8	52.5	71.0	2.7	1.2	18.3	1.7	51.7	67.2	2.8	1.3	18.6	1.8	52.6	72.5	69.3
$(2.5 \ 1.5 \ 0.7 \ 0.4)$	60 44.9	0	22.7	13.0	11.1	6.2	55.8	0	21.8	12.6	10.9	6.0	44.7	0	22.7	13.0	11.1	6.2	70.0	51.4
	100 53.3	0	28.7	17.5	12.0	5.2	64.8	0	28.2	16.9	11.7	4.8	53.0	0	28.7	17.5	12.0	5.2	78.0	60.8

Table 5: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in ARCH(4) models under the local alternative γ/\sqrt{T} .

 $\gamma' = 0.95/k^{(1)} \times$ MinP-sc MinP-st MinP-s DS LK T $(1_{k^{(1)}}, 0_{k-k^{(1)}})$ H_0 FWER H_{01} H_{02} H_{03} H_{04} H_{05} H_{06} H_0 FWER H_{01} H_{02} H_{03} H_{04} H_{05} H_{06} H_0 FWER H_{01} H_{02} H_{03} H_{04} H_{05} H_{06} H_0 H_0 $u_n \sim N(0,1)$ 0_{6} 60 5.05.01.0 0.8 0.9 0.8 1.20.8 4.9 4.50.8 0.7 0.8 0.8 1.1 0.75.05.01.0 0.8 0.9 0.8 1.2 0.8 5.34.6 $100 \ 5.5$ 5.40.81.1 0.91.1 1.30.65.75.10.81.1 0.90.91.10.65.45.40.81.1 0.91.1 1.30.65.75.3 $k^{(1)} = 1$ 1.460 46.9 8.7 41.0 3.6 2.11.11.61.2 47.7 8.4 40.3 3.6 2.01.1 1.2 46.8 8.7 41.0 3.6 2.11.11.61.2 48.2 49.8 100 72.8 10.8 3.13.10.6 66.2 73.6 69.3 4.9 1.41.50.673.810.668.54.83.11.41.50.6 72.7 10.8 69.3 4.9 1.41.5 $k^{(2)} = 1$ 60 46.2 9.6 $22.0 \ 19.9 \ 4.5$ 2.32.01.651.49.5 $21.5 \ 19.5$ 4.52.31.9 $1.5 \ 46.0$ 9.622.0 19.9 4.5 2.32.01.6 62.7 53.2 $100\ 71.4$ 13.7 $37.6 \ 35.2 \ 6.9$ 3.42.71.475.8 13.5 $36.6 \ 34.3 \ 6.8$ 3.42.61.4 70.8 13.7 $37.6\ 35.2\ 6.9$ 3.42.71.4 80.8 77.8 $k^{(3)} = 1$ 60 44.2 7.7 $14.7 \ 13.5 \ 15.5 \ \ 3.2$ 3.02.253.07.414.3 13.1 15.1 3.02.9 $2.1 \ 44.1$ 7.714.7 13.5 15.5 3.23.02.2 67.1 52.2 $100\ 67.1$ 9.9 $27.4\ 23.5\ 25.4\ \ 4.1$ 3.72.677.9 $26.9\ 23.1\ 24.8\ 4.0$ 3.6 $2.6 \ 66.3$ 9.9 $27.4\ 23.5\ 25.4\ \ 4.1$ 3.72.6 85.3 77.8 9.7 $k^{(6)} = 1$ 60 37.6 0 6.8 6.5 7.7 8.0 7.88.350.10 $6.7 \ \ 6.3 \ \ 7.4$ 8.0 7.58.0 37.3 0 6.86.5 7.7 8.0 7.88.3 61.0 44.1 100 56.3 0 12.5 11.5 13.0 12.0 13.2 12.2 76.6 0 $12.2\ 11.3\ 12.6\ 11.7\ 13.0$ 11.9 55.1 0 12.5 11.5 13.0 12.0 13.2 12.2 84.5 69.7 $u_n \sim t_{10}$ 0.6 0.7 0_6 60 5.35.31.10.6 0.7 0.9 1.21.15.24.9 1.00.8 1.05.35.31.10.6 0.70.91.21.1 5.3 4.5 1.1 $100 \ 5.4$ 5.4 $1.2 \ 0.8$ 0.6 $1.0 \quad 0.9$ 1.15.85.11.10.80.60.9 $0.9 \ 1.0 \ 5.4$ 5.41.20.80.61.0 $0.9 \ 1.1 \ 4.4 \ 5.2$ $k^{(1)} = 1$ 60 45.8 9.0 40.6 3.4 2.61.1 1.71.2 47.0 8.6 $39.9 \ 3.3$ 2.51.01.71.1 45.8 9.0 $40.6 \ 3.4$ 2.61.11.71.2 48.9 49.7 100 66.7 14.2 $60.7 \ 6.1 \ 4.8$ 2.11.72.114.260.7 6.1 4.8 1.1 67.9 14.059.7 6.0 4.81.61.1 66.7 2.11.71.1 68.8 68.1 $k^{(2)} = 1$ 60 45.4 11.9 $21.5 \ 18.8 \ 4.9$ 3.03.8 $2.3 \ 50.9$ $21.1\ 18.5$ 4.93.03.72.2 45.2 11.9 $21.5 \ 18.8 \ 4.9$ 3.03.82.3 65.9 51.7 11.71.8 $100\ 68.1$ $33.9\ 33.1\ 7.3$ 4.32.875.8 $33.3\ 32.7\ 7.2$ 4.32.8 $1.8 \ 67.7$ $33.9\ 33.1\ 7.3$ 4.32.81.8 85.1 75.5 14.714.614.7 $k^{(3)} = 1$ 60 45.9 $15.4 \ 13.7 \ 16.4 \ \ 3.4$ 3.3 $14.9 \ 13.5 \ 15.9$ 3.45.43.2 45.7 15.4 13.7 16.4 3.45.43.3 71.4 54.4 11.05.454.010.911.0 $100 \ 69.3$ 13.0 $26.3\ 24.4\ 26.9\ 5.4$ 5.04.180.7 12.8 $25.7\ 24.0\ 26.4\ 5.2$ 5.0 $4.0 \ 68.8$ 13.0 $26.3\ 24.4\ 26.9\ 5.4$ 5.04.1 89.5 79.4 $k^{(6)} = 1$ 60 41.3 8.1 7.2 8.2 8.3 10.6 9.0 54.3 8.0 7.0 8.0 8.1 10.2 8.7 40.9 $8.1 \quad 7.2 \quad 8.2 \quad 8.3 \quad 10.6 \quad 9.0 \quad 66.3 \quad 48.0$ 0 0 0 100 59.9 0 $13.1\ 12.4\ 13.8\ 12.5\ 14.7\ 13.3\ 81.9$ 0 $12.9\ 12.1\ 13.4\ 12.4\ 14.4\ 13.0\ 58.6$ 0 $13.1\ 12.4\ 13.8\ 12.5\ 14.7\ 13.3\ 89.9\ 73.9$

Table 6: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in ARCH(6) models.

Table 7: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in ARCH(6) models under the local alternative γ/\sqrt{T} .

$\gamma' = 7/k^{(1)} \times$	T	MinP-sc									Ν	linP-	st						Ν	MinP	-s				DS	LK
$(1_{k^{(1)}}, 0_{k-k^{(1)}})$	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_{05}	H_{06}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_{05}	H_{06}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_{04}	H_{05}	H_{06}	H_0	H_0
												u_n	$\sim N($	(0, 1)												
$k^{(1)} = 1$	60 47.4	4 8.6	41.7	3.4	2.1	1.0	1.6	1.1	47.8	8.1	41.0	3.3	1.9	0.9	1.6	1.0	47.4	8.6	41.7	3.4	2.1	1.0	1.6	1.1	46.9	48.6
	100 67.	5 7.2	64.6	2.8	2.0	0.8	1.2	0.6	67.9	6.9	63.7	2.8	2.0	0.8	1.1	0.5	67.4	7.2	64.6	2.8	2.0	0.8	1.2	0.6	54.6	68.1
$k^{(2)} = 1$	60 43.9	9 8.5	20.2	19.7	4.0	2.1	2.0	1.1	48.2	8.2	19.6	19.2	3.9	2.0	1.9	1.1	43.6	8.5	20.2	19.7	4.0	2.1	2.0	1.1	60.2	50.3
	100 58.0	5 7.7	30.7	28.9	3.9	2.1	1.3	0.8	63.4	7.5	29.6	28.3	3.8	2.0	1.3	0.8	58.2	7.7	30.7	28.9	3.9	2.1	1.3	0.8	67.0	65.2
$k^{(3)} = 1$	60 41.8	8 6.9	13.6	12.5	15.0	3.0	2.6	1.9	50.3	6.7	13.4	12.1	14.6	2.9	2.6	1.8	41.5	6.9	13.6	12.5	15.0	3.0	2.6	1.9	64.2	50.3
	100 55.	l 5.5	21.5	19.2	20.0	2.7	2.4	0.6	62.2	5.5	20.9	18.5	19.5	2.7	2.4	0.6	54.2	5.5	21.5	19.2	20.0	2.7	2.4	0.6	70.8	63.7
$k^{(6)} = 1$	60 34.4	1 0	6.5	6.2	6.5	7.3	7.1	7.0	46.0	0	6.4	5.9	6.2	7.1	7.0	6.5	34.3	0	6.5	6.2	6.5	7.3	7.1	7.0	58.6	40.8
	100 43.	7 0	8.3	8.1	8.7	8.8	9.9	8.8	59.6	0	7.9	7.9	8.5	8.6	9.5	8.6	42.7	0	8.3	8.1	8.7	8.8	9.9	8.8	68.0	54.9
												u	$_n \sim t$	10												
$k^{(1)} = 1$	60 46.0) 8.8	41.1	3.4	2.1	1.1	1.6	1.4	47.0	8.6	40.3	3.4	2.0	1.0	1.5	1.4	45.8	8.8	41.1	3.4	2.1	1.1	1.6	1.4	47.2	48.2
	100 62.2	7 9.2	58.6	4.0	2.6	1.3	1.0	1.1	63.3	9.1	57.5	3.8	2.6	1.3	1.0	1.1	62.6	9.2	58.6	4.0	2.6	1.3	1.0	1.1	57.7	63.9
$k^{(2)} = 1$	60 42.0	5 10.4	20.3	18.4	4.4	3.2	3.0	1.8	47.8	10.3	20.0	18.0	4.4	3.1	2.9	1.8	42.5	10.4	20.3	18.4	4.4	3.2	3.0	1.8	63.5	49.4
	$100 \ 60.4$	4 10.3	29.7	29.1	5.0	2.6	2.2	1.3	66.9	10.0	29.2	28.1	4.9	2.6	2.1	1.2	60.0	10.3	29.7	29.1	5.0	2.6	2.2	1.3	74.3	67.7
$k^{(3)} = 1$	60 45.8	8 10.1	15.6	13.7	16.1	3.3	5.2	2.5	54.9	10.0	15.1	13.5	16.0	3.2	5.2	2.5	45.7	10.1	15.6	13.7	16.1	3.3	5.2	2.5	68.8	53.7
	100 58.	L 7.7	20.7	20.3	22.2	3.4	3.3	1.7	67.7	7.4	20.3	20.0	21.9	3.2	3.2	1.7	57.5	7.7	20.7	20.3	22.2	3.4	3.3	1.7	78.5	68.4
$k^{(6)} = 1$	60 37.9) ()	7.7	6.3	6.9	7.7	9.9	8.6	50.7	0	7.5	6.3	6.9	7.6	9.7	8.5	37.6	0	7.7	6.3	6.9	7.7	9.9	8.6	65.5	46.3
	100 47.8	8 0	9.3	9.4	10.0	9.8	9.9	8.8	65.6	0	9.2	9.2	9.7	9.5	9.7	8.3	46.8	0	9.3	9.4	10.0	9.8	9.9	8.8	77.6	58.3

Table 8: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in random coefficient models with k = 2.

γ'	T		MinP	-sc			MinP	-st			MinF	° -s		$\bar{\chi}^2$	t
		H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	FWER	H_{01}	H_{02}	H_0	H_0
							ϵ_n	$\sim N$	(0, 1)						
$(0 \ 0)$	100	5.1	4.6	3.0	1.8	4.9	3.3	2.1	1.3	4.8	4.8	3.0	1.9	5.1	5.1
	200	4.6	4.3	2.0	2.3	4.8	3.5	1.7	1.9	4.7	4.7	2.0	2.7	4.5	4.8
$(0.5 \ 0)$	100	64.1	4.5	58.9	4.5	55.6	4.1	54.1	4.1	60.6	4.5	59.7	4.5	67.8	18.7
	200	86.2	4.7	84.5	4.7	81.6	4.5	81.2	4.5	85.1	4.7	85.0	4.7	88.1	31.0
$(0.5 \ 0.5)$	100	66.6	0	44.8	51.2	58.1	0	41.0	46.7	65.2	0	44.9	51.4	70.0	11.6
	200	93.5	0	77.4	63.7	83.3	0	74.1	61.5	88.0	0	77.7	64.1	96.0	49.4
								$\epsilon_n \sim$	t_5						
$(0 \ 0)$	100	7.4	7.0	3.5	3.8	6.4	4.6	2.3	2.6	7.3	7.3	3.7	4.0	7.0	5.4
	200	6.7	6.5	3.2	4.0	5.7	4.6	2.3	2.9	6.8	6.8	3.3	4.2	5.9	3.7
$(0.5 \ 0)$	100	23.7	3.7	21.7	3.7	19.0	3.0	17.7	3.0	23.7	3.8	22.3	3.8	24.3	4.1
	200	52.8	4.7	48.3	4.7	44.8	4.4	43.5	4.4	50.6	4.9	49.2	4.9	54.2	11.4
$(0.5 \ 0.5)$	100	39.1	0	24.1	27.3	32.4	0	21.0	23.5	39.0	0	24.4	27.6	38.4	3.6
. ,	200	56.8	0	40.5	36.5	47.4	0	36.8	33.4	54.0	0	41.1	37.0	62.2	13.3

γ'	T		Mir	nP-so)			Mii	nP-st	-)			Mi	nP-s			$\bar{\chi}^2$	t
		H_0	FWER	H_{01}	H_{02}	H_{03}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_0	FWER	H_{01}	H_{02}	H_{03}	H_0	H_0
								e	$\epsilon_n \sim$	$\overline{N(0,}$	1)							
$(0 \ 0 \ 0)$	100	5.2	5.1	1.9	1.6	2.2	5.4	4.2	1.5	1.3	2.0	5.3	5.3	2.0	1.7	2.3	4.8	5.9
	200	5.7	5.4	2.2	1.5	1.8	5.8	4.9	2.0	1.4	1.7	5.7	5.7	2.3	1.7	2.0	5.1	4.2
(0.5 0 0)	100	50.6	4.0	46.7	2.3	2.0	44.5	3.5	42.9	2.2	1.7	48.7	4.1	47.2	2.4	2.1	52.5	9.9
	200	90.9	9.9	87.2	6.0	4.4	85.5	9.8	85.0	5.9	4.4	88.2	9.9	87.7	6.0	4.4	93.6	41.1
(0.5 0.5 0)	100	67.9	3.1	45.2	34.1	3.1	56.0	2.8	43.0	31.7	2.8	61.1	3.1	45.8	34.6	3.1	77.1	15.0
	200	84.1	5.0	49.0	54.8	5.0	69.4	4.8	47.6	52.4	4.8	73.2	5.1	49.2	55.0	5.1	91.0	24.5
$(0.4 \ 0.4 \ 0.4)$	100	62.5	0	33.3	21.6	31.2	51.6	0	31.1	20.0	29.0	57.6	0	33.9	22.1	31.6	72.0	7.6
	200	89.6	0	57.4	48.7	55.1	78.8	0	55.9	48.0	53.8	82.2	0	57.5	48.8	55.3	94.8	32.9
									ϵ_n	$\sim t_5$								
$(0 \ 0 \ 0)$	100	5.7	5.3	1.7	2.1	2.0	5.5	4.4	1.4	1.9	1.6	5.6	5.6	1.8	2.1	2.2	5.5	4.8
	200	5.5	5.1	1.8	1.7	1.9	5.5	4.1	1.4	1.4	1.7	5.4	5.4	1.9	1.8	2.1	5.7	4.4
(0.6 0 0)	100	27.6	5.0	22.0	2.3	2.9	23.5	4.6	20.0	2.1	2.7	26.7	5.3	23.2	2.4	3.1	29.7	6.0
	200	62.3	6.6	59.4	3.1	3.7	57.1	6.3	55.8	3.0	3.5	61.3	6.6	60.3	3.1	3.8	61.6	14.8
(0.5 0.5 0)	100	51.5	2.6	30.7	18.2	2.6	39.8	2.5	28.8	17.1	2.5	43.2	2.6	31.8	18.7	2.6	61.3	17.1
	200	66.3	3.4	44.8	46.8	3.4	59.8	3.3	42.9	44.8	3.3	64.1	3.4	45.2	47.2	3.4	71.7	18.8
$(0.4 \ 0.4 \ 0.4)$	100	48.3	0	20.6	18.1	17.3	36.6	0	19.2	16.4	15.8	41.6	0	21.0	18.3	17.6	57.0	5.0
	200	59.6	0	22.6	20.1	27.1	43.4	0	21.4	18.4	26.0	47.8	0	23.1	20.4	27.6	71.6	10.5

Table 9: Estimated probabilities in percentages of rejecting H_0 , FWER and H_{0i} in random coefficient models with k = 3.