

Lower Bound for Sculpture Garden Problem

M. Eskandari ^{*}, B. Sadeghi Bigham [†]

July 20, 2021

Abstract

The purpose of the current study is to investigate a special case of art gallery problem, namely Sculpture Garden Problem. In the said problem, for a given polygon P , the ultimate goal is to place the minimum number of guards to define the interior polygon P by applying a monotone Boolean formula composed of the guards. As the findings indicate, the conjecture about the issue that in the worst case, $n - 2$ guards are required to describe any n -gon (Eppstein et al. 2007) can be conclusively proved.

Key words: art galley, Boolean formula, computational geometry, prison yard problem, sculpture garden problem.

1 Introduction

A large and growing body of literature about computational geometry has explored the art gallery problem. The main goal in this problem is to place the minimum number of point guards inside a polygon P then, the set of guards can see the whole P . The number of guards that suffices and sometimes necessary for any arbitrary polygon with n vertices is $\lfloor n/3 \rfloor$ [1]. The main goal in our study is to find the minimum set of angle guards by which the geometry of the polygon can be defined through two operations $AND(\cdot)$ and $OR(+)$. An *angle guard* g views an infinite wedge of the plane (by going through the involved obstacles) and can be defined as a Boolean predicate, $B_g(p)$, which is

^{*}Corresponding author: Faculty of Mathematical Sciences, Alzahra University, Tehran, Iran. eskandari@alzahra.ac.ir

[†]Institute for Advanced Studies in Basic Sciences (IASBS), Zanjan, Iran. b_sadeghi_b@iasbs.ac.ir

The Authors' name are in alphabetical order.

True for a given point $p \in P$ if p is inside the view region of g , otherwise it is *False*. Given a polygon P , the aim is to place a set of angle guards on P in such a way that the monotone Boolean formula $F_P(p)$ is *True*, if and only if p is inside or on the polygon P , otherwise it is *False*:

$$F_P(p) = \begin{cases} \textit{True} & \text{if } p \in P \text{ or } p \in \partial P \\ \textit{False} & \text{otherwise} \end{cases}$$

An angle guard vertex placement is considered as *natural* if all the guards of P have the same view of their corresponding vertices [2]. As Eppstein et al. stated, a polygon P can be demonstrated in a way that a natural angle guard vertex placement cannot fully distinguish between points on the inside and outside of P which implies *Steiner-point* guards are sometimes necessary [2]. According to Figure 1.a, even the placement of a natural angle at every vertex of the pentagon is not able to distinguish between the points x and y and at least one unnatural guard is needed to define the polygon (Figure 1.b). Consequently the polygon is defined by $F = A.B.D$.

A variety of cases of problem is present in which the location and angel of view of the guards are different. We focus on a type of the problem that all the guards are placed on the vertices of P . It was a conjecture [3] that in the worst case, $n - 2$ guards are needed to describe n -gon. In this paper, we prove that the conjecture is true and present an algorithm to generate a polygon for a given n which needs at least $n - 2$ guards.

In the next section, the Sculpture guard problem is introduced and some applications are mentioned. Section 3 provides the main problem and presents an algorithm for generating the n -gon which needs exactly $n - 2$ guards to be defined. Finally, Section 4, presents the findings of the study and also some suggestions for further research.




Figure 1: (a) Natural angle guards do not suffice to define the polygon; (b) Coverage by three guards (formula A.B.D define the polygon) [3]

2 Sculpture Garden Problem and Applications

As mentioned previously, Sculpture Garden Problem can be considered as a special case of Art Gallery problem. There are various problems with similarities and differences with Sculpture garden problem. As Sculpture Garden Problem comes up from localization problems in wireless mobile computing, we wish to determine the position of some devices in a geometric environment.

Sculpture Garden Problem could be used in a localization problems in which a wireless device is used to prove that it belongs to a given polygonal environment. In this case, the locators would be simple and can broadcast information inside a certain angle. In this context, the Boolean predicates could be associated with secret keys. Therefore, each angle guard g could periodically broadcast a secret key K in its transmission angle and consequently the devices in range would have knowledge of this key. The wireless localization problem with natural and vertex guards is a NP-hard problem [4, 5].

In another application, namely constructive solid geometry (CSG), we wish to construct a geometric shape from simple combinations of primitive shapes [6]. Solutions to the Sculpture guard problem can be used to construct an efficient CSG formula that defines a given polygon P . Prison yard problem seeks a set of guards that can simultaneously see both the interior and exterior of a simple polygon, in which $\lfloor n/2 \rfloor$ guards are sufficient and sometimes necessary (tight bound) [7]. Another related problem is Floodlight illumination problem in which the vertex angle guards (called floodlights) with angles of 180 should see a simple polygon [8]. Likewise, studies have been conducted on the complexity of illuminating wedges with angle-restricted floodlights placed at a fixed set of points [9].

Since we are interested in more than simple observing the inside and outside of a polygon, solution to art gallery or prison yard problem would not change into solutions to sculpture garden problem. In other words, it is intended to establish the time when a point is inside a polygon using only the guards as witnesses.

Indeed, any polygon P can be triangulated and two of angel guards can be used to define each of the resulting $n + 2(h - 1)$ triangles, where h is the number of holes in P . This would give rise to a concise formula F defining P . However, it uses at least $2n + 4(h - 1)$ angle guards, which is more constant-depth formulas.

2.1 Upper and Lower Bounds

The Sculpture Garden Problem has different types due to the different restrictions as guards could be observed in varied forms including vertices, edges, interior, or exterior of polygon, the SGP can be manifested in different types, as well. However, in each case, finding the upper and lower bound is a problem that already was investigated.

An angle guard g with angle $\alpha \in (0, 360)$ is a pair (a, ω_α) of a point a and an infinitive wedge ω_α of aperture α at apex a and views ω_α . It can be shown as a Boolean predicate $B_g(p)$, in which for a point p in the plane, $B_g(p)$, is *True*, if p is inside the angle associated with g otherwise it is *False*. Given a polygon P with n vertices, we intend to allocate the minimum number of angle guards with arbitrary angles at vertices of P . Thus, a monotone Boolean formula, $F_P(\cdot)$, based on the angle guard predicates, $B_g(\cdot)$ is obtained as follows:

$$F_P(p) = \begin{cases} \textit{True} & \forall p \in \textit{int}(P) \\ \textit{False} & \textit{otherwise} \end{cases} \quad (1)$$

It is worth mentioning that $\textit{int}(P)$ is interior of polygon P . If $F_P(p)$ is a solution of sculpture garden problem for a given P , P is defined by F_P . The complement of an angle guard $g = (a, \omega_\alpha)$ is an angle guard g' at the point a with angle $2\pi - \alpha$. Hence, the wedge associated with g' is the *complement* of ω_α in plane. If formula F is a solution for sculpture garden problem for polygon P , then the complement of F which is denoted by F' defines the exterior of P . To obtain F' , initially, we replace every angle guard g by its complement, (i.e. g'), and then swap the operations *AND* and *OR*. In addition, we define , a *pocket* of a simple polygon as the areas outside of the polygon and inside of its convex hull.

As Eppstein et al. [2, 3] reported, for any polygon P , a set of $n + 2(h - 1)$ angle guards and an associated concise formula F are present solving the Sculpture Garden problem where h is the number of holes in P . So, a simple polygon can be defined with $n - 2$ guards. They have conjectured a class of simple n -gons that require at least $n - 2$ natural vertex guards. The main goal of this paper is to solve this open problem. They showed at least $\lceil n/2 \rceil$ guards are required to solve the Sculpture Garden problem for any polygon with no two edges lying on a same line. Besides, for any convex polygon P , a natural angel guard vertex placement is present whose $\lceil n/2 \rceil$ guards are sufficient. They showed that $\lceil n/2 \rceil + O(1)$ angle guards suffice to solve the Sculpture Garden problem for pseudo-

Table 1: Number of guards needed for a simple polygon with n vertices.

	<i>Natural Vertex</i>	<i>Natural</i>	<i>Vertex</i>	<i>General</i>
<i>Upper Bound</i>	<i>Not known</i>	$n - 2$ [11]	$n - 2$ [2]	$\lfloor \frac{(4n-2)}{5} \rfloor$ [11]
<i>Lower Bound</i>	<i>Not known</i>	$n - 2$ [11]	$\lfloor \frac{(2n)}{3} \rfloor - 1$ [10]	$\lceil \frac{(3n-4)}{5} \rceil$ [11]

triangles. Moreover, for any orthogonal polygon P (which is, probably the most likely real world applications) a set of $\lceil 3(n-2)/4 \rceil$ angle guards and an associated concise formula F are available to solve the Sculpture Garden problem using $\lceil n/2 \rceil$ natural angle guards. They gave an example of a class of simple polygons containing Sculpture Garden solutions are used $O(\sqrt{n})$ guards. Afterwards, they showed that the bound is optimal. On the contrary, some varied results are obtained for vertex guards. As Damian et al. demonstrated [10] a class of simple n -gons are presented that require at least $\lfloor 2n/3 \rfloor - 1$ guards placed at polygon vertices for localization. Through revealing the point that the maximum number of guards to describe any simple polygon on n vertices is roughly observable at $(3/5n, 4/5n)$, Hoffman et al. enhanced both upper and lower bounds for the general setting [11]. Eskandari et al. [12] improved the large upper bound $n + 2(h - 1)$ for an arbitrary n -gon. with h holes for placing guards and obtained a tight bound $(n - \lceil c/2 \rceil - h)$, where c is the number of vertices of convex hull of P . So in simple polygons, this bound is $n - \lceil c/2 \rceil$ which is tight too. To complete the first column of Table 1, a new class of polygons entitled *helix* is introduced in the next section.

3 Helix Polygon

In this section, we explore the special class of sculpture garden problem, where, the guards are natural. We manifest the point that the lower bound for the problem is $n - 2$. To do so, we commence with introducing a class of polygons demanding the exact number of $n - 2$ natural guards to be defined. In the next section, we introduce this class of polygons named *helix* (See Figure 2).

3.1 Construction Helix

An n -gon helix polygon (i.e. H_n) is constructed by an incremental method using an $n - 1$ -gon helix, H_{n-1} . A helix with three vertices is a triangle. By adding two new edges to H_{n-1} and also removing an edge of H_{n-1} , H_n is constructed on the basis of H_{n-1} where $n \geq 4$. The details are presented in Algorithm 1 and are illustrated in Figure 3.

Algorithm 1 Constructing Helix Polygon

Input: Integer number $n \geq 3$ as the number of vertices.

Output: The helix polygon H_n .

- 1: Construct $H_3 = \triangle v_1 v_2 v_3$, which is an equilateral triangle where $v_2 v_3$ is horizontal and the vertices are in counterclockwise order.
 - 2: Choose a positive real number δ so $0 < \delta < \frac{|v_2 v_3|}{2 \lfloor \frac{n-1}{3} \rfloor}$
 - 3: $p_1 = v_1; p_2 = v_2; p_3 = v_3$.
 - 4: **for** $i = 4; i \leq n; i++$ **do**
 - 5: $q_1 = p_1; p_1 =$ a point on $v_1 v_2$ such that $|p_1 q_1| = \delta, a = l_{13}(p_1);$
 - 6: $q_2 = p_2; p_2 =$ a point on $v_2 v_3$ such that $|p_2 q_2| = \delta, b = l_{12}(p_2);$
 - 7: $q_3 = p_3; p_3 =$ a point on $v_1 v_3$ such that $|p_3 q_3| = \delta, c = l_{23}(p_3);$
 - 8: **if** $i == 4$ **then**
 - 9: $l = b;$
 - 10: **else**
 - 11: $l = l_{24}(v_{i-2});$
 - 12: **end if**
 - 13: **if** $i == 5$ **then**
 - 14: $l' = c;$
 - 15: **else**
 - 16: $l' = l_{35}(v_{i-2});$
 - 17: **end if**
 - 18: **if** $i == 3k$ **then**
 - 19: $v_i = a \cap c;$
 - 20: **end if**
 - 21: **if** $i == 3k + 1$ **then**
 - 22: $v_i = c \cap l;$
 - 23: **end if**
 - 24: **if** $i == 3k + 2$ **then**
 - 25: $v_i = b \cap l';$
 - 26: **end if**
 - 27: Add edges $v_i v_{i-1}$ and $v_i v_{i-2}$.
 - 28: Remove $v_{i-1} v_{i-2}$ to obtain H_i
 - 29: **end for**
 - 30: Return H_n .
-

It is worth noting that the length of a line segment pq is denoted by $|pq|$ and for an arbitrary point p , $l_{ij}(p)$ denotes a line parallel to $v_i v_j$ which passes through p .




Figure 2: (a) A helix with 11 vertices. (b) A helix with 12 vertices.

3.2 Properties of Helix

Constructing the helix polygon sheds light that for $i > 2$, vertices v_{2i} are concave and for $i > 0$, vertices v_{2i+1} and v_2 are convex (Figure 2).

In fact, the pocket of a polygon P is defined as $CH(P) - P$ where $CH(P)$ is the convex hull of the vertices of P . The pocket of helix polygon with n vertices is a helix polygon with $n - 1$ vertices (Figure 4). The pocket of a polygon H_n is denoted by $P(H_n)$. For i , $1 \leq i \leq n - 1$, the vertices of $P(H_n)$ are called v'_i located on v_{i+1} . For $n > 4$, the angle $\widehat{v'_i}$ in $P(H_n)$ is obtained as follows:

$$\widehat{v'_i} = \begin{cases} \widehat{v_1 v_2 v_3} - \widehat{v_1 v_2 v_4} & i = 1 \\ \widehat{v_1 v_3 v_2} - \widehat{v_1 v_3 v_5} & i = 2 \\ 2\pi - \text{interior angle of } v_{i+1} \text{ in } H_n & i \geq 3. \end{cases} \quad (2)$$

We will show that polygon H_n can be defined by $n - 2$ natural vertices guards which are located on $v_1, v_2, \dots, v_{n-3}, v_n$. The Boolean formula, F_n , is as below:

$$F_n = \sum_{i=1}^{\lceil n/2 \rceil - 2} A_i \cdot g_{2i} + A_{\lceil n/2 \rceil - 1} \cdot g_n \quad (3)$$




Figure 3: Final step of Algorithm 1 for generating H_{12}

where $A_1 = g_1$, $A_{i+1} = A_i \cdot g_{2i+1}$ for all $1 \leq i \leq \lfloor n/2 \rfloor - 2$ and g_i is the natural vertex guard located on the vertex v_i of H_n . To clarify this point, F_8 can be written as follows:

$$F_8 = \sum_{i=1}^2 A_i \cdot g_{2i} + A_5 \cdot g_8 \quad (4)$$

where, $A_1 = g_1$, $A_2 = g_1 \cdot g_3$ and $A_3 = g_1 \cdot g_3 \cdot g_5$. Thus, we have:

$$F_8 = g_1 \cdot g_2 + g_1 \cdot g_3 \cdot g_4 + g_1 \cdot g_3 \cdot g_5 \cdot g_8$$

Generally, we expand F_n as follows:

$$F_n = \begin{cases} \sum_{i=1}^{k-1} (\prod_{j=0}^{i-1} g_{2j+1}) \cdot g_{2i} + (\prod_{j=0}^{k-2} g_{2j+1}) \cdot g_n & n = 2k + 1, k \in \mathbb{N} \\ \sum_{i=1}^{k-1} (\prod_{j=0}^{i-1} g_{2j+1}) \cdot g_{2i} + (\prod_{j=0}^{k-1} g_{2j+1}) \cdot g_n & n = 2k + 2, k \in \mathbb{N}. \end{cases} \quad (5)$$




Figure 4: (a) Helix H_{12} . (b) Pocket of H_{12} which is a helix with 11 vertices.

According to Lemma 1, F_n defined by Equation 5 describes H_n .

Lemma 1. *Helix polygon H_n can be defined by $n-2$ natural vertex guards g_i ($1 \leq i \leq n; i \neq n-1, n-2$) located on $v_1, v_2, \dots, v_{n-3}, v_n$ and the correspondent Boolean formula is Formula 5.*

Proof. We will prove the lemma by induction. When $n = 3$, $k = 1$ and $F_3 = g_1 \cdot g_3$ clearly defines triangle H_3 . For $n = 4$, $k = 1$ and $F_4 = g_1 \cdot g_4$ defines H_4 and it implies that the Lemma 1 holds for $n = 3, 4$. Now, for $n \geq 5$, without loss of generality, assume that $n = 2k + 2, k \in \mathbb{N}$. According to Property 2, $P(H_n)$ is a helix polygon with $2k + 1$ vertices. By induction hypothesis, $P(H_n)$ can be defined as follows:

$$F' = \sum_{i=1}^{k-1} \left(\prod_{j=0}^{i-1} g'_{2j+1} \right) \cdot g'_{2i} + \left(\prod_{j=0}^{k-2} g'_{2j+1} \right) \cdot g'_{2k+1} \quad (6)$$

where g'_i is a natural guard on the vertex v'_i of $P(H_n)$. According to correspondence between the vertices of H_n and $P(H_n)$, we have:

$$g'_i = \begin{cases} g_{i+1}^c & i \geq 3 \\ G_{i+1} \cdot g_{i+1}^c & i = 1, 2. \end{cases} \quad (7)$$

in which g^c is the complement of guard g and G_2 and G_3 are the guards located on v_2 and v_3 with the angles $\widehat{v_1 v_2 v_3}$ and $\widehat{v_1 v_3 v_2}$, respectively. So we have:

$$F' = g'_1 \cdot g'_2 + \sum_{i=2}^{k-1} \left(\prod_{j=0}^{i-1} g'_{2j+1} \right) \cdot g'_{2i} + \left(\prod_{j=0}^{k-2} g'_{2j+1} \right) \cdot g'_{2k+1}$$

$$\begin{aligned}
&= g'_1 \cdot g'_2 + \sum_{i=2}^{k-1} g'_1 \cdot \left(\prod_{j=1}^{i-1} g'_{2j+1} \right) \cdot g'_{2i} + g'_1 \cdot \left(\prod_{j=1}^{k-2} g'_{2j+1} \right) \cdot g'_{2k+1} \\
&= G_2 \cdot G_3 \cdot g_2^c \cdot g_3^c + \sum_{i=2}^{k-1} G_2 \cdot g_2^c \cdot \left(\prod_{j=1}^{i-1} g_{2j+2}^c \right) \cdot g_{2i+1}^c + G_2 \cdot g_2^c \cdot \left(\prod_{j=1}^{k-2} g_{2j+2}^c \right) \cdot g_{2k+2}^c \\
&= G_2 \cdot \left[G_3 \cdot g_2^c \cdot g_3^c + \sum_{i=2}^{k-1} \left(\prod_{j=0}^{i-1} g_{2j+2}^c \right) \cdot g_{2i+1}^c + \left(\prod_{j=0}^{k-2} g_{2j+2}^c \right) \cdot g_{2k+2}^c \right] \\
\implies (F')^c &= G_2^c + (G_3^c + g_2 + g_3) \cdot \left(\prod_{i=2}^{k-1} \left(\sum_{j=0}^{i-1} g_{2j+2} \right) + g_{2i+1} \right) \cdot \left(\sum_{j=0}^{k-2} g_{2j+2} + g_{2k+2} \right)
\end{aligned}$$

Consider the point that $F = (g_1 \cdot g_2) \cdot (F')^c + (g_1 \cdot g_3) \cdot (F')^c$. By replacing $(F')^c$ from the above relation, we obtain:

$$\begin{aligned}
F &= g_1 \cdot (g_2 + g_2 \cdot g_3) \cdot \left(\prod_{i=2}^{k-1} \left(\sum_{j=0}^{i-1} g_{2j+2} \right) + g_{2i+1} \right) \cdot \left(\sum_{j=0}^{k-2} g_{2j+2} + g_{2k+2} \right) \\
&\quad + g_1 \cdot (g_2 \cdot g_3 + g_3) \cdot \left(\prod_{i=2}^{k-1} \left(\sum_{j=0}^{i-1} g_{2j+2} \right) + g_{2i+1} \right) \cdot \left(\sum_{j=0}^{k-2} g_{2j+2} + g_{2k+2} \right)
\end{aligned}$$

Note that $g_1 \cdot g_2 \cdot G_2^c = g_1 \cdot g_3 \cdot G_2^c = g_1 \cdot g_2 \cdot G_3^c = g_1 \cdot g_3 \cdot G_3^c = \emptyset$. So we have:

$$F = \left(\prod_{i=2}^{k-1} \left(\sum_{j=0}^{i-1} g_{2j+2} \right) + g_{2i+1} \right) \cdot \left(\sum_{j=0}^{k-2} (g_{2j+2} + g_{2k+2}) \right) \cdot (g_2 + g_3) \cdot g_1$$

Now, we show that F can define H_n which contains exactly the natural guards $g_1, g_2, \dots, g_{n-3}, g_n$ and can be written in the form of F_n .

First, consider the definition of F which contains only natural guards. To prove that H_n can be defined by F , let x be an arbitrary point inside H_n . We have $g_1(x) = True$ and $(F')^c(x) = True$ ($x \in H_n \implies x \notin P(H_n) \implies F'(x) = False \implies (F')^c(x) = True$). There are two cases:

- $g_2(x) = True \implies F(x) = (g_1(x) \cdot g_2(x)) \cdot (F')^c(x) + (g_1(x) \cdot g_3(x)) \cdot (F')^c(x) = True$
- $g_2(x) = False \implies g_3(x) = True \implies (g_1(x) \cdot g_3(x)) \cdot (F')^c(x) = True \implies F(x) = True$.

Thus, F can distinguish the interior of H_n . Now, let $x \notin H_n$. There are two cases:

- $x \in P(H_n) \implies F'(x) = True, (F')^c(x) = False \implies F(x) = False$
- $x \notin P(H_n) \implies x \in Ext(\Delta v_1 v_2 v_3) \implies g_1(x) = False \implies F(x) = False$.

So, F can distinguish the exterior of H_n as well. Now, we show that F can be written in the form of F_n . Let

$$S = (g_2 + g_3) \cdot \left(\prod_{i=2}^{k-1} \left(\sum_{j=0}^{i-1} g_{2j+2} \right) + g_{2i+1} \right) \cdot \left(\sum_{j=0}^{k-2} g_{2j+2} + g_{2k+2} \right)$$

and

$$T = \sum_{i=1}^{k-1} \left(\prod_{j=1}^{i-1} (g_{2j+1}) \cdot g_{2i} \right) + \left(\prod_{j=1}^{k-1} (g_{2j+1}) \cdot g_{2k+2} \right)$$

Note that $F_n = g_1 \cdot T$ and $F = g_1 \cdot S$. To prove $F = F_n$, it is sufficient to show that $T = S$. For all integers r where $1 \leq r \leq k-1$, we define $S_i^{(r)}$ as follows:

$$S_i^{(r)} = \begin{cases} g_{2i+1} + \sum_{j=r}^{i-1} g_{2j+2} & r \leq i \leq k-1 \\ g_{2k+2} + \sum_{j=r}^{k-2} g_{2j+2} & i = k. \end{cases} \quad (8)$$

So, we have:

$$S_i^{(1)} = \begin{cases} g_{2i+1} + \sum_{j=1}^{i-1} g_{2j+2} & 1 \leq i \leq k-1 \\ g_{2k+2} + \sum_{j=1}^{k-2} g_{2j+2} & i = k. \end{cases} \quad (9)$$

By definition of $S_i^{(r)}$, it is clear that:

$$S = \prod_{i=1}^k (g_2 + S_i^{(1)}) = g_2 + \prod_{i=1}^k S_i^{(1)}$$

On the other hand, $S_i^{(r)} - S_i^{(r+1)} = g_{2r+2}$. Therefore we have:

$$\begin{aligned} \prod_{i=r}^k S_i^{(r)} &= S_r^{(r)} \cdot \prod_{i=r+1}^k S_i^{(r)} \\ &= g_{2r+1} \cdot \prod_{i=r+1}^k S_i^{(r)} = g_{2r+1} \cdot \prod_{i=r+1}^k (g_{2r+2} + S_i^{(r+1)}) = g_{2r+1} \cdot (g_{2r+2} + \prod_{i=r+1}^k S_i^{(r+1)}) \end{aligned}$$

By applying obtained recursive relation, $k-2$ times on $S = g_2 + \prod_{i=1}^k S_i^{(1)}$, $S = T$. In this respect, we have:

$$\begin{aligned} S &= g_2 + \prod_{i=1}^k S_i^{(1)} = g_2 + g_3 \cdot (g_4 + \prod_{i=2}^k S_i^{(2)}) = g_2 + g_3 \cdot g_4 + g_3 \cdot \prod_{i=2}^k S_i^{(2)} \\ &= g_2 + g_3 \cdot g_4 + g_3 \cdot g_5 \cdot g_6 + g_3 \cdot g_5 \cdot \prod_{i=3}^k S_i^{(3)} \end{aligned}$$

After t times, we have:

$$S = \left(\sum_{i=1}^{t+1} \left(\prod_{j=1}^{i-1} g_{2j+1} \right) \cdot g_{2i} \right) + \prod_{j=1}^t g_{2j+1} \cdot \prod_{i=t+1}^k S_i^{(t+1)}$$

So after $k-2$ times, we have:

$$S = \left(\sum_{i=1}^{k-1} \left(\prod_{j=1}^{i-1} g_{2j+1} \right) \cdot g_{2i} \right) + \prod_{j=1}^{k-2} g_{2j+1} \cdot \prod_{i=k-1}^k S_i^{(k-1)}$$

Note that $\prod_{i=k-1}^k S_i^{(k-1)} = S_{k-1}^{(k-1)} \cdot S_k^{(k-1)} = g_{2k-1} \cdot g_{2k+2}$, therefore,

$$S = \left(\sum_{i=1}^{k-1} \left(\prod_{j=1}^{i-1} g_{2j+1} \right) \cdot g_{2i} \right) + \prod_{j=1}^{k-1} g_{2j+1} \cdot g_{2k+2} = T.$$

$S = T$ implies $F = F_n$ which means that F can be written in the form of F_n and could define H_n . \square

3.3 Necessity of $n-2$ Natural Vertex Guards for Helix

In this section, we will prove that it is impossible to define helix polygon with fewer than $n-2$ natural vertex guards.

Lemma 2. *Every arbitrary set of natural vertex guards G which defines H_n contains g_1 , a natural vertex guard on v_1 . The final formula is in the form of $F = F_1 \cdot g_1$, where F_1 is a Boolean expression of $G - \{g_1\}$.*

Proof. Let G is an arbitrary set of natural guards which defines H_n by Boolean formula F . Suppose for a contradiction that g_1 does not belong to G . Since v_1v_2 and v_1v_3 are edges of H_n , G should contain two natural guards on v_2 and v_3 which are called g_2 and g_3 , respectively. So F can be written in the general form $F = g_2 \cdot g_3 \cdot T_1 + g_2 \cdot T_2 + g_3 \cdot T_3$ where T_i s are Boolean formulas which do not contain g_1 , g_2 and g_3 . This will result in a contradiction, mentioned below.

Consider two regions R_1 and R_2 as shown in Figure 5. Let x be an arbitrary point inside R_1 or R_2 . So we have: $g_2(x) = True$ and $g_3(x) = False$. So,

$$F(x) = T_2(x) \tag{10}$$

Also, note that we can expand T_2 in the following general form:

$$T_2 = T_2^{(1)} + T_2^{(2)} + \dots + T_2^{(l)} \quad (11)$$

where $T_2^{(i)}$ s are the multiplication of some natural guards in G . Let $x \in R_1$ be an arbitrary point. So from Equation 10, it is implied that:

$$x \in R_1 \implies T_2(x) = F(x) = True$$

Therefore, at least one of the expressions of T_2 should be True. Without loss of generality, it can be called $T_2^{(1)}$ and is written as $T_2^{(1)} = g_{i_1}.g_{i_2} \cdots .g_{i_m}$ where g_{i_j} is the natural guards of G and $i_j = 1, 2, 3$. Since $T_2^{(1)}(x) = True$, we have:

$$\forall j : 1 \leq j \leq m : g_{i_j}(x) = True$$

Regarding the structure of H_n , none of indices i_j can be odd. This is because we know that

$$\forall i \geq 1 : g_{2i+1}(x) = False$$

Now, let $y \in R_2$ be an arbitrary point. Due to the structure of H_n , it is inferred that for all $\forall y \in R_2$, we have:

$$\forall i \geq 2 : g_{2i}(y) = True$$

. So, $T_2^{(1)}(x) = True$ and from Equation 11, $T_2(y) = True$ is obtained and consequently $F(y) = T_2(y) = True$ (due to Equation 10). Nevertheless, $y \notin H_n$ which is a contradiction. With regard to existence of g_1 in G , F can be written in the form of $F = g_1.T_1 + T_2$ where T_2 does not contain g_1 . Indeed, $g_1.T_1 + g_1.T_2$ defines H_n as well. Let $x \in H_n$, then $g_1(x) = True$ and $g_1(x).T_1(x) + g_1(x).T_2(x) = g_1(x).T_1(x) + T_2(x) = F(x) = True$. If $y \notin H_n$ and $g_1(y) = True$, then $g_1(y).T_1(y) + g_1(y).T_2(y) = g_1(y).T_1(y) + T_2(y) = F(y) = False$. On the other hand, if $g_1(y) = False$, then $g_1(y).T_1(y) + g_1(y).T_2(y) = False$. So $F = g_1.(T_1 + T_2)$ defines H_n . In other words, F can be expressed as $F = g_1.F_1$.

□

Lemma 3. *Every arbitrary set of natural vertex guards G defining H_n contains g_2 (i.e. a natural guard on v_2). The final formula is in the form of $F = g_1.(g_2 + F_2)$, where F_2 is a Boolean expression of $G - \{g_1, g_2\}$.*




Figure 5: Regions R_1 and R_2 can be distinguishable without existence of g_1 in the formula.

Proof. Let G is an arbitrary set of natural guards which defines H_n by Boolean formula F , for $n \geq 4$. Suppose, a contradiction in which g_2 does not belong to G . From Lemma 2, we can write

$$F = g_1.(T_1 + T_2 + \dots + T_l)$$

where T_i s are Boolean expression of natural guards of G .

Consider two regions R_1 and R_3 as shown in Figure 6. Let $x \in R_1$ is an arbitrary point. We have $F(x) = g_1(x).(T_1(x) + T_2(x) + \dots + T_l(x)) = True$. Thus, at least one of the expressions T_i s is True. Without loss of generality, it can be named T_1 and is written as $T_1 = g_{i_1}.g_{i_2} \dots g_{i_m}$ where g_{i_j} s are some natural guards in G and $i_j \neq 1, 2$. Since $T_1(x) = True$, we have

$$\forall j, 1 \leq j \leq m : g_{i_j}(x) = True$$

Regarding the structure of H_n , i_j s are even, because $\forall i \geq 2 : g_{2i}(x) = True$.

Now, let $y \in R_3$ be an arbitrary point. Obviously, $\forall i \geq 2, g_{2i}(y) = True$ which implies $T_1(y) = True$. Then $F(y) = True$. However, $y \notin H_n$ which denotes a contradiction.

In addition, F can be manifested as below.

$$F = g_1.(g_2.T_1 + T_2)$$

Now note that for all points which are located in the interior (or exterior) of H_n , the above formula has the same value with the formula $g_1.(g_2 + T_2)$. This fact can be shown easily by considering all cases. Then $F = g_1.(g_2 + F_1)$.




Figure 6: Regions R_1 and R_3 can be distinguishable without existence of g_2 in the formula.

□

Corollary 4. *It is not possible to define H_5 with less than 3 natural vertex guards. The formula is $F = g_1.(g_2 + g_5)$.*

Proof. Regarding Lemmas 2 and 3, F can be written as $g_1.(g_2 + F_2)$. The edges v_4v_5 and v_3v_5 should have at least one guard on their endpoints. The optimal possibility is to locate a guard on v_5 as their intersection point. Clearly, $g_1.(g_2.g_5)$ defines H_5 .

□

Lemma 5. *Every arbitrary set of natural vertex guards G which defines H_n contains g_3 which is a natural guard on v_3 . The final formula is in the form of $F = g_1.(g_2 + g_3.F_3)$ where F_3 is a Boolean expression of $G - \{g_1, g_2, g_3\}$.*

Proof. Let G be an arbitrary set of natural guards which defines H_n by Boolean formula F for $n \geq 6$. Suppose contradiction in which g_3 does not belong to G . By Lemma 3, $F = g_1.(g_2 + F_2)$. Assume that $F_2 = T_1 + T_2 + \dots + T_l$ where T_i s are multiplication of natural guards in G .

Consider two regions R_3 and R_4 as shown in Figure 7. Let $x \in R_4$ be an arbitrary point. We have $F(x) = F_2(x) = True$. So at least one of T_i s is True. Without loss of generality, we call it T_1 which can be expressed as follows:

$$T_1 = g_{i_1} \cdot g_{i_2} \cdots g_{i_m}$$

where g_{i_j} s are natural guards in G and $i_j \neq 1, 2, 3$.

Since $T_1(x) = True$, we have:

$$\forall j, 1 \leq j \leq m : g_{i_j}(x) = True$$

Therefore i_j s are even. Now, let $y \in R_3$ be an arbitrary point. This point casts light that for all $i \geq 2$, $g_{2i}(y) = True$ which implies $T_1(y) = True$. Then, $F(y) = True$. However, $y \notin H_n$ showing a contradiction. So, $g_3 \in G$.

In addition, F can be written as bellow:

$$F + g_1 \cdot (g_2 + g_3 \cdot F_3)$$

It can be easily obtained from equivalency of $g_1 \cdot (g_2 + F_2)$ and $g_1 \cdot (g_2 + g_3 \cdot F_3)$ for all points with respect to H_n .




Figure 7: Regions R_3 and R_4 can be distinguishable without existence of g_3 in the formula.

□

Corollary 6. *It is not possible to define H_5 with fewer than 3 natural vertex guards. The formula is $F = g_1.(g_2 + g_3.g_6)$.*

Proof. Considering Lemma 5, H_6 can be defined by $F = g_1.(g_2 + g_3.F_3)$. Since v_4v_6 and v_6v_5 are two edges of H_6 , it is required to place at least one guard on one of the endpoints of these two edges. The optimal placement is to place a guard on v_6 . Obviously, H_6 can be defined by $F = g_1.(g_2 + g_3.g_6)$. \square

Lemma 7. *Every arbitrary set of natural vertex guards G which defines H_n , for all $n \geq 7$, contains g_4 , a natural guard on v_4 . The final formula is in the form of $F = g_1.(g_2 + g_3.(g_4 + F_4))$ where F_4 is a Boolean expression of $G - \{g_1, g_2, g_3, g_4\}$.*

Proof. Let G be an arbitrary set of natural guards defining H_n by Boolean formula F and $n \geq 7$. Suppose a contradiction in which g_4 does not belong to G . By Lemma 5, $F = g_1.(g_2 + g_3.F_3)$. It is intended to show that $F_3 = g_4 + F_4$. Assume that $F_3 = T_1 + T_2 + \dots + T_l$ where T_i s are multiplication of natural guards of G . Consider two regions R_5 and R_6 shown in Figure 8. Let $x \in R_5$ be an arbitrary point, then $F(x) = F_3(x) = True$. So, at least one of T_i s is True. Without loss of generality, it is called T_1 which can be expressed as bellow:

$$T_1 = g_{i_1}.g_{i_2} \dots g_{i_m}$$

where g_{i_j} s are natural guards and $i_j \neq 1, 2, 3, 4$. Since $T_1(x) = True$, for all $j : 1 \leq j \leq m$, $g_{i_j}(x) = True$. Therefore i_j s are even. Now let $y \in R_6$ be an arbitrary point (see Figure 8). It is clear that for all $i \geq 2$, $g_{2i}(y) = True$ which implies $T_1(y) = True$. Then, $F(y) = True$. However, $y \notin H_n$ indicating a contradiction. So $g_4 \in G$. In addition, F can be written as bellow:

$$F = g_1.(g_2 + g_3.(g_4 + F_4))$$

It can be easily shown that $g_1.(g_2 + g_3.F_3)$ is equivalent with $g_1.(g_2 + g_3.(g_4 + F_4))$ for all the points inside or outside of H_n . \square

Lemma 8. *Let H_n be defined by $F = g_1.(g_2 + g_3.(g_4 + F_4))$. Formula $F_P = g'_1.(g'_2 + g'_3.F_4^c)$ defines the pocket of H_n , $P(H_n)$ where g'_i s are defined in Equation 7.*

Proof. Let $x \in P(H_n)$ and $y \notin P(H_n)$ be two arbitrary points (see Figure 9). Then, it can be demonstrated that $F_P(x) = True$ and $F_P(y) = False$. Also, $g'_2 = G_3.g_3^c$ and $g'_3 = g_4^c$, so: $x \in P(H_n) \implies g'_1(x) = True$ and $G_3(x) = True$. So




Figure 8: Regions R_5 and R_6 can be distinguishable without existence of g_4 in the formula.

$$F_P(x) = g_3^c(x) + g_4^c(x).F_4^c(x) \quad (12)$$

On the other hand, $x \in P(H_n)$ implies that $F(x) = False$ and $g_1(x) = True$. Hence,

$$F(x) = g_2(x) + g_3(x).(g_4(x) + F_4(x)) = False \implies g_3(x).(g_4(x) + F_4(x)) = False$$

, so

$$g_3^c(x) + g_4^c(x).F_4^c(x) = True \quad (13)$$

From Equations 12 and 13, $F_P(x) = True$ can be obtained.

If $g'(y) = False$, for $y \notin P(H_n)$, $F_P(y) = False$ and the process of our calculations has been completed. Assume that $g'(y) = True$, then as $y \notin P(H_n)$, consequently $g'_2(y) = False$ (see Figure 9). If $g'_3(y) = False$, $F_P(y) = False$. Now, suppose that $g'_3(y) = True$ (i.e. $g_4(y) = False$). This assumption implies that $y \in H_n$ and we have $g_1(y) = True$, $g_2(y) = False$, $g_31(y) = True$ and $g_4(y) = False$. Since $y \in H_n$, $F(y) = F_4(y)$ and consequently, $F_4(y) = True$. So, we have $F_P(y) = False$. This means that F_P defines $P(H_n)$.

□




Figure 9: Natural vertex guards for packet of helix

Theorem 9. H_n requires at least $n - 2$ natural vertex guards.

Proof. We prove this theorem by induction. It is clear that for $n = 4$, H_4 is a tetragon and cannot be defined by fewer than two guards. We proved this in corollaries 4 and 6 for $n = 5$. Now, assume that this holds for $n - 1$, and we have to prove it for n where $n \geq 7$. Let F be a Boolean formula which defines H_n . With regard to Lemma 7, $F = g_1 \cdot (g_2 + (g_3 \cdot (g_4 + F_4)))$. Let m be the number of natural guards used in F . From Lemma 8, $P(H_n)$, a helix with $n - 1$ vertices, can be defined by $F_P = g'_1 \cdot (g'_2 + g'_3 \cdot F_4^c)$ which contains $m - 1$ guards. By induction hypothesis, $P(H_n)$ cannot be defined by fewer than $(n - 1) - 2$ natural guards. So $m - 1$ cannot be less than $n - 3$ and hence m cannot be fewer than $n - 2$. \square

Theorem 10. H_n requires exactly $n - 2$ natural vertex guards.

Proof. From Lemma 1 and Theorem 9, it is obviously implied that H_n requires exactly $n - 2$ natural vertex guards. \square

As we proved, there is an n -gon which needs exactly $n - 2$ natural vertex guards to be defined. This implies that $n - 2$ is the lower bound.

4 Conclusion:

Epstein et al. [3] in 2007 conjectured that for a given number n , at least one simple polygon is present that requires $n - 2$ natural vertex guards to describe the polygon. Introducing a new class of polygons named helix polygon, we proved the conjecture. Further research should be done to investigate the bounds for special cases of polygons (e.g. orthogonal polygons).

Acknowledgements

We deeply thank our colleagues in RoboCG (Robotics and Computational Geometry) lab for their insight and expertise that greatly assisted the research. We also appreciate Arya Falahi for the efforts he made to solve our Latex problems.

References

- [1] Mark De Berg, Marc Van Kreveld, Mark Overmars, and Otfried Cheong Schwarzkopf. *Computational geometry*. Springer, 2000.
- [2] David Eppstein, Michael T Goodrich, and Nodari Sitchinava. Guard placement for efficient point-in-polygon proofs. In *Proceedings of the twenty-third annual symposium on Computational geometry*, pages 27–36. ACM, 2007.
- [3] David Eppstein, Michael T Goodrich, and Nodari Sitchinava. Guard placement for wireless localization. *arXiv preprint cs/0603057*, 2006.
- [4] Tobias Christ, Michael Hoffmann, and Yoshio Okamoto. Natural wireless localization is np-hard. In *Abstracts 25th European Workshop Comput. Geom.*, pages 175–178. Citeseer, 2009.
- [5] Tobias Christ and Michael Hoffmann. Wireless localization with vertex guards is np-hard. In *CCCG*, pages 149–152. Citeseer, 2009.
- [6] David Dobkin, Leonidas Guibas, John Hershberger, and Jack Snoeyink. *An efficient algorithm for finding the CSG representation of a simple polygon*, volume 22. ACM, 1988.

- [7] Zoltán Füredi and Daniel J. Kleitman. The prison yard problem. *Combinatorica*, 14(3):287–300, 1994.
- [8] Vladimir Estivill-Castro, Joseph O’Rourke, Jorge Urrutia, and Dianna Xu. Illumination of polygons with vertex lights. *Information Processing Letters*, 56(1):9–13, 1995.
- [9] William Steiger and Ileana Streinu. Illumination by floodlights. *Computational Geometry*, 10(1):57–70, 1998.
- [10] Mirela Damian, Robin Flatland, Joseph O’Rourke, and Suneeta Ramaswami. A new lower bound on guard placement for wireless localization. *arXiv preprint arXiv:0709.3554*, 2007.
- [11] Tobias Christ, Michael Hoffmann, Yoshio Okamoto, and Takeaki Uno. Improved bounds for wireless localization. In *Algorithm Theory–SWAT 2008*, pages 77–89. Springer, 2008.
- [12] M Eskandari, A Mohades, and B Sadeghi Bigham. On the number of guards in sculpture garden problem. *World Applied Sciences Journal*, 10(10):1255–1263, 2010.