

FIBER DECOMPOSITION OF DETERMINISTIC REACTION NETWORKS WITH APPLICATIONS

CARSTEN WIUF AND CHUANG XU

ABSTRACT. Deterministic reaction networks (RNs) are tools to model diverse biological phenomena characterized by particle systems, when there are abundant number of particles. Examples include but are not limited to biochemistry, molecular biology, genetics, epidemiology, and social sciences. In this chapter we propose a new type of decomposition of RNs, called *fiber decomposition*. Using this decomposition, we establish lifting of mass-action RNs preserving stationary properties, including *multistationarity* and *absolute concentration robustness*. Such lifting scheme is simple and explicit which imposes little restriction on the reaction networks. We provide examples to illustrate how this lifting can be used to construct RNs preserving certain dynamical properties.

1. INTRODUCTION AND STATE OF THE ART

Reaction networks (RNs) can be regarded as a modelling machinery for many real-world dynamical systems. Examples include networks in epidemiology [4], pharmacology [6], ecology [18], and social sciences [31], as well as gene regulatory networks [9], biochemical reaction networks [21], signalling networks [29], and metabolic networks [32]. An RN is a finite nonempty set of reactions between complexes consisting of species. When there are abundant species and all species are homogeneously well mixed, an RN can be modelled deterministically by ordinary differential equations (ODEs), called *the rate equation*.

Multistationarity of deterministic reaction networks. A reaction network is *multistationary* if its rate equation admits multiple steady states (subject to the linear subspace the dynamics is confined to) [21]. Endowed with mass-action kinetics, the rate equation associated with an RN has a polynomial vector field. Hence to determine steady states of an RN amounts to determining zeros of a polynomial, which in general is challenging [21, 22]. Several approaches have been proposed to ensure the existence of multiple positive steady states (steady states with positive entries), e.g., based on deficiency theory [26, 19, 21], injectivity based tests using a Jacobian criterion [12, 13, 14], and homotopy and other approaches [15, 27, 10].

Lifting of RNs preserving multistationarity is well investigated in the literature [11, 27, 5]. The reason for studying lifting procedures is two-fold. First of all, whether an RN is multistationary might be solved for a smaller/simpler RN and if it is so, then the larger RN of interest is also multistationary by lifting. Secondly, lifting procedures might provide means to construct complex examples of RNs with the same properties as the simpler RN. The lifting scheme based on the so-called “atoms of multistationarity” [27] is valid for *fully open continuous-flow stirred-tank reactors* (CFSTRs), a network in which all chemical species enter the system at constant rates and are removed at rates proportional to their concentrations, that is, there are reactions $0 \rightleftharpoons S$ for all species in the RN.

An RN \mathcal{R} is *nondegenerately* multistationary provided a subnetwork $\tilde{\mathcal{R}} \subseteq \mathcal{R}$ with the same stoichiometric subspace as \mathcal{R} is so [27] (Here “nondegenerate” is in the sense of the

Date: 22nd July 2021.

Key words and phrases. Deterministic reaction networks, fiber decomposition, lifting, multistationarity, absolute concentration robustness.

Jacobian matrix of the vector field). This in particular implies two networks share the same set of species. The proof depends on constructing a mapping from every positive steady state (PSS) of the subnetwork $\tilde{\mathcal{R}}$ to a nearby point which is a PSS of \mathcal{R} . Based on the construction, the set of PSSs of $\tilde{\mathcal{R}}$ is not necessarily a subset of those of \mathcal{R} .

In contrast, our lifting scheme (Theorem 5.2) based on the fiber decomposition proposed in this chapter (i) does allow for two RNs to have different sets of species; (ii) under a certain assumption, the projection of the set of PSSs of the original reaction network \mathcal{R} onto the set of species of the subnetwork recovers precisely the set of PSSs of the subnetwork; (iii) the networks are not necessarily CFSTRs. Nevertheless, the specific construction of lifting does impose certain conditions on the reaction rate constants of the two RNs.

Absolute concentration robustness. One interesting property of RNs is absolute concentration robustness. An RN is *absolute concentration robust* (ACR) if the system has at least one PSS, and all PSSs projected to a given species S_i are identical (say, x_i^*). Such a species S_i is called an ACR species with x_i^* being the ACR value. Many biological systems have such ACR property, e.g., the EnvZ-OmpR osmoregulatory system, and double-phosphorylation systems of transcriptional regulatory proteins [34, 8]. This ACR property is closely related to a desirable property in bioengineering, called *robust perfect adaption*, which means that a biological system can adapt after an external stimulus has been applied, and be insensitive to variations in the biochemical parameters of the system [7]. We remark that ACR is also closely related to sensitivity analysis of parameters for biological systems [33].

We will show that our lifting procedure also works for ACR. For this we briefly review the literature on ACR. Based on linear algebra, a simple sufficient condition for a system of deficiency one to be ACR can be given [34] (see Proposition 5.10). This has recently been extended to a class of RNs with a weaker condition than deficiency one [8]. Also lifting of RNs preserving the ACR property under the former conditions is discussed therein. The notion of ACR has likewise been generalized to local ACR and necessary conditions for local ACR have been given [30]. An RN is *local ACR* with local ACR species S_i if the projection of the set of PSSs onto the i -th coordinate is nonempty and finite. We mention that results for a stochastic analogue of the ACR property are sparse [3, 2, 1, 20].

2. NOTATION

Let \mathbb{R} , \mathbb{R}_{\geq} , and \mathbb{Q} be the set of real numbers, nonnegative real numbers, and rational numbers, respectively. Given a finite index set $J \subseteq \mathbb{N}$, for any $x = (x_j)_{j \in J} \in \mathbb{R}^J$, denote $\text{supp } x := \{j \in J : x_j \neq 0\}$.

3. REACTION NETWORKS

In this section, we introduce reaction networks as well as elementary propositions, as prerequisites of the fiber decomposition of reaction networks.

A reaction network (RN) is composed of a triple $(\mathcal{S}, \mathcal{C}, \mathcal{R})$ of three non-empty finite sets:

- (i) $\mathcal{S} = \{S_i\}_{1 \leq i \leq d}$ is a set of symbols, termed *species*;
- (ii) $\mathcal{C} \subseteq \mathbb{N}_0^{\mathcal{S}}$ is a set of linear combinations of species, termed *complexes*, and
- (iii) $\mathcal{R} \subseteq \mathcal{C} \times \mathcal{C}$ is a set of *reactions*. A reaction (y, y') is denoted $y \longrightarrow y'$. The complex y is called the *reactant* and y' the *product*.

For convention, we assume every species is in some complex and every complex is in some reaction. Hence we also identify an RN with \mathcal{R} since \mathcal{S} and \mathcal{C} can be deduced from \mathcal{R} . We emphasize that \mathcal{R} is a set without multiplicity, and hence does not contain multiple identical reactions.

A reaction is *degenerate* if its reactant coincides with its product; otherwise it is non-degenerate. An RN is *degenerate* if it contains degenerate reactions; otherwise, it is *non-degenerate*. Given an RN \mathcal{R} , let $\mathcal{R}_* \subseteq \mathcal{R}$ be the subnetwork (with \mathcal{S}_* and \mathcal{C}_* being its sets

of species and complexes) only consisting of non-degenerate reactions. The concept of an RN herein is more general than the standard one in the literature of chemical reaction network theory (CRNT) [21] simply because we allow for degenerate reactions. Other definitions of RNs, similar to our definition, have also been explored in the literature [16, 17].

The pair $(\mathcal{C}, \mathcal{R})$ forms a (possibly non-simple) digraph referred to as the *reaction graph*. Hence the reaction graph is non-simple and contains a self-loop if and only if there exists a degenerate reaction in the RN. We adopt the convention that every node is strongly connected to itself. Any weakly connected component is called a *linkage class*. All nodes in a strongly connected component (called a *strong linkage class*) are *terminal* if there are no edges from any node in this component to a node in any other strongly connected component. Any node in a terminal strongly connected component is a *terminal complex*; otherwise it is a *non-terminal complex*. Let $\ell_{\mathcal{R}}$ be the number of linkage classes of \mathcal{R} .

Let $\Omega := \{y' - y: y \longrightarrow y' \in \mathcal{R}\}$ be the set of *reaction vectors* and $\Omega_* := \Omega \setminus \{0\} = \{y' - y: y \longrightarrow y' \in \mathcal{R}_*\}$. Let $\mathbf{S} := \{\sum_{\omega \in \Omega} c_{\omega} \omega: c_{\omega} \in \mathbb{R}\}$ be the span of Ω over \mathbb{R} , termed the *stoichiometric subspace* of \mathcal{R} . Note that $\mathbf{S} = \mathbf{S}_* := \{\sum_{\omega \in \Omega_*} c_{\omega} \omega: c_{\omega} \in \mathbb{R}\}$ and $\dim \mathbf{S} \leq d$. Define the *deficiency* of a reaction network \mathcal{R} :

$$\delta = \#\mathcal{C}_* - \ell_{\mathcal{R}_*} - \dim \mathbf{S}.$$

Then \mathcal{R} and \mathcal{R}_* share the same deficiency. For every $c \in \mathbb{R}_{\geq 0}^{\mathbf{S}}$, let $\mathbf{S}_c = (\mathbf{S} + c) \cap \mathbb{R}_{\geq 0}^{\mathbf{S}}$ be the *stoichiometric compatibility class* through c .

Given two RNs \mathcal{R}^1 and \mathcal{R}^2 . Let $y \longrightarrow y' \in \mathcal{R}^1$ and $\tilde{y} \longrightarrow \tilde{y}' \in \mathcal{R}^2$. We say $y \longrightarrow y'$ is representable by $\tilde{y} \longrightarrow \tilde{y}'$ and denoted $\{y \longrightarrow y'\} \preceq \{\tilde{y} \longrightarrow \tilde{y}'\}$ if

- i) $y' - y$ is a multiple of $\tilde{y}' - \tilde{y}$, that is, there exists $r \in \mathbb{R}$ such that $y' - y = r(\tilde{y}' - \tilde{y})$,
- ii) $\text{supp } \tilde{y} \subseteq \text{supp } y$.

A subnetwork $\mathcal{A} \subseteq \mathcal{R}^1$ is representable by $\tilde{y} \longrightarrow \tilde{y}'$, denoted $\mathcal{A} \preceq \tilde{y} \longrightarrow \tilde{y}'$ if every reaction in \mathcal{A} is so. \mathcal{R}^1 is representable by \mathcal{R}^2 , denoted by $\mathcal{R}^1 \preceq \mathcal{R}^2$ if every reaction in \mathcal{R}^1 is representable by one reaction in \mathcal{R}^2 .

A *deterministic reaction network* is a pair consisting of an RN and a *kinetics* $\mathcal{K} = (\lambda_{y \longrightarrow y'})_{y \longrightarrow y' \in \mathcal{R}}$, where $\lambda_{y \longrightarrow y'}: \mathbb{R}_{\geq 0}^{\mathbf{S}} \rightarrow \mathbb{R}_{\geq 0}$ is the *rate function* of $y \longrightarrow y'$, expressing the propensity of the reaction to occur. A special kinetics is *mass-action kinetics*,

$$(3.1) \quad \lambda_{y \longrightarrow y'}(x) = \kappa_{y \longrightarrow y'} x^y := \kappa_{y \longrightarrow y'} \prod_{i=1}^d x_i^{y_i},$$

where $\kappa_{y \longrightarrow y'}$ is referred to as the *reaction rate constant*. Hence, $\lambda_{y \longrightarrow y'}(x) > 0$ if and only if $\text{supp } x \supseteq \text{supp } y$ for $x \in \mathbb{R}^{\mathbf{S}}$.

For the ease of exposition rather than for generality, we assume throughout that all RNs are endowed with mass-action kinetics, and hence we also use \mathcal{R} to refer to the RN with mass-action kinetics.

The *rate equation* for a deterministic RN \mathcal{R} as well as for the corresponding \mathcal{R}_* , characterizing the change in species concentrations over time is then given by the ODE system

$$(3.2) \quad \dot{x} = \sum_{y \longrightarrow y' \in \mathcal{R}_*} \lambda_{y \longrightarrow y'}(x)(y' - y).$$

Hence for every $c \in \mathbb{R}_0^{\mathbf{S}}$, \mathbf{S}_c is an invariant subspace under the flow generated by (3.2).

4. FIBER DECOMPOSITION OF RNs

In this section, we define a fiber decomposition of an RN and use it to construct an explicit lifting scheme from one RN (*reference reaction network*) to another “larger” RN (with more species, complexes, and/or reactions) while preserving various stationary properties, including multistationarity and ACR property. We mention that converse to lifting, reduction of RNs

can be derived *mutatis mutandis*. Hence the main results can be potentially used to simplify large networks in biochemistry and synthetic biology.

Reference RN and base RN. Given an RN $(\mathcal{S}, \mathcal{C}, \mathcal{R})$, let $\mathcal{S}^1 \sqcup \mathcal{S}^2 = \mathcal{S}$ be a partition of \mathcal{S} into two disjoint sets \mathcal{S}^1 and \mathcal{S}^2 . We refer to \mathcal{S}^1 as the *reference* subset of species. Let \mathbf{P}_i is the natural projection from $\mathbb{R}^{\mathcal{S}}$ onto $\mathbb{R}^{\mathcal{S}^i}$ for $i = 1, 2$. Hence for $i = 1, 2$, for $y \in \mathcal{C}$, $\mathbf{P}_i y = 0$ if $\text{supp } y \cap \mathcal{S}^i = \emptyset$, and for $y \rightarrow y' \in \mathcal{R}$, $\mathbf{P}_i(y \rightarrow y') = \mathbf{P}_i y \rightarrow \mathbf{P}_i y'$ defines a reaction confined to the species set \mathcal{S}^i . Furthermore, let $\mathbf{P}_i \mathcal{R} = \{\mathbf{P}_i y \rightarrow \mathbf{P}_i y' : y \rightarrow y' \in \mathcal{R}\}$ (without multiplicity) and $\mathbf{P}_i \mathcal{C} = \{\mathbf{P}_i y : y \in \mathcal{C}\}$.

Hence $(\mathcal{S}^1, \mathcal{C}^b, \mathcal{R}^b)$ with $\mathcal{C}^b = \mathbf{P}_1 \mathcal{C}$ and $\mathcal{R}^b = \mathbf{P}_1 \mathcal{R}$ forms a new RN, called the *base reaction network* (BRN) of \mathcal{R} , denoted \mathcal{R}^b . Let $\mathcal{R}_*^b \subseteq \mathcal{R}^b$ be the subset of non-degenerate reactions.

In addition to \mathcal{R} , we consider another mass-action RN, termed the *reference* RN, given as $\mathcal{R}^\circ = \{y_k \rightarrow y'_k\}_{k \in K^\circ}$, where K° is an index set. Let $\mathcal{R}_*^\circ \subseteq \mathcal{R}^\circ$ be the RN consisting of the non-degenerate reactions of \mathcal{R}° and $K_*^\circ \subseteq K^\circ$ its index set.

Now we are ready to come up with a decomposition of \mathcal{R} w.r.t. the reference RN \mathcal{R}° . Assume

(A1) $\mathcal{R}_*^\circ \neq \emptyset$.

(A2) $\mathcal{R} = \sqcup_{k \in K^\circ} \mathcal{R}_k$ is a partition in disjoint RNs, such that for $k \in K_*^\circ$, $\mathbf{P}_1 \mathcal{R}_k \preceq y_k \rightarrow y'_k$, and for $k \in K^\circ \setminus K_*^\circ$, $\mathbf{P}_1 \mathcal{R}_k \subseteq \mathcal{R}^b$ consists of degenerate reactions.


For $y \rightarrow y' \in \mathcal{R}_k$, $k \in K_*^\circ$, let

$$(4.1) \quad a_k^{y \rightarrow y'} (y'_k - y_k) = \mathbf{P}_1 (y' - y), \quad a_k^{y \rightarrow y'} \in \mathbb{Q}.$$

Assumption (A2) implies $\mathcal{R}^b \preceq \mathcal{R}^\circ$. By definition of representability, a reaction in \mathcal{R}_*^b can only be representable by a reaction in \mathcal{R}_*° . Moreover, if \mathcal{R}° is non-degenerate, then so are \mathcal{R}^b and \mathcal{R} .

Nevertheless, a BRN of a non-degenerate RN can be degenerate.


Example 4.1. Consider the non-degenerate RN




Let $\mathcal{S}^1 = \{S_1\}$. Then its BRN $\{S_1 \rightarrow S_1\}$ is an RN consisting of a degenerate reaction. Hence \mathcal{R}° must be degenerate.

A reaction in \mathcal{R}^b can be representable by more than one reaction in \mathcal{R}° .

Example 4.2. Consider the RN \mathcal{R} :



Let \mathcal{R}° :



Hence $\mathcal{R}^b = \mathcal{R}^\circ$. Since either reaction in \mathcal{R}° is representable by the other, then either reaction in \mathcal{R}^b is representable by either reaction in \mathcal{R}° .

We further emphasize that such a decomposition given in (A2) may not be unique.

Example 4.3. Revisit Example 4.2. Then

$$\mathcal{R} = \mathcal{R}_1 \sqcup \mathcal{R}_2,$$

gives two decompositions with either

$$\mathcal{R}_1 = \{S_1 + S_2 + S_3 \rightarrow 2S_1 + S_2 + S_3\}, \quad \mathcal{R}_2 = \{2S_1 + S_2 + 2S_3 \rightarrow 3S_1 + S_2 + 3S_3\}$$

or

$$\mathcal{R}_1 = \{2S_1 + S_2 + 2S_3 \rightarrow 3S_1 + S_2 + 3S_3\}, \quad \mathcal{R}_2 = \{S_1 + S_2 + S_3 \rightarrow 2S_1 + S_2 + S_3\},$$

where $y_1 \rightarrow y'_1 = S_1 + S_2 \rightarrow 2S_1 + S_2$ and $y_2 \rightarrow y'_2 = 2S_1 + S_2 \rightarrow 3S_1 + S_2$.

We remark that decompositions of reaction networks are proposed in other contexts [24, 23, 25], e.g., for the pursuit of explicit formulas of stationary distributions [25].

Fiber decomposition. For a reaction $y \rightarrow y' \in \mathcal{R}$, we write $\mathbf{P}_1(y \rightarrow y') \oplus \mathbf{P}_2(y \rightarrow y')$ for the direct sum decomposition $y \rightarrow y' = \mathbf{P}_1 y \oplus \mathbf{P}_2 y \rightarrow \mathbf{P}_1 y' \oplus \mathbf{P}_2 y'$.

With a decomposition as in (A2), define the associated *fiber reaction network* (FRN) $(\mathcal{S}^2, \mathcal{C}_k, \mathcal{R}_k^f)$ on \mathcal{S}^2 at every reaction $y_k \rightarrow y'_k \in \mathcal{R}^\circ$ as

$$\mathcal{C}_k := \bigcup_{y \rightarrow y' \in \mathcal{R}_k} \{\mathbf{P}_2 y, \mathbf{P}_2 y'\}, \quad \mathcal{R}_k^f = \mathbf{P}_2 \mathcal{R}_k.$$

Let $\mathcal{R}_{k,*}^f \subseteq \mathcal{R}_k^f$ be the subset of non-degenerate reactions.

Let $\mathcal{R}_k^b = \mathbf{P}_1 \mathcal{R}_k$. Hence $\mathcal{R}^b = \bigcup_{k \in K^\circ} \mathcal{R}_k^b$. Recall by the definition of the decomposition in (A2), for every $k \in K_*$, $\mathcal{R}_k^b \preceq y \rightarrow y'_k$. Note that for two different reactions $y_k \rightarrow y'_k, y_{k'} \rightarrow y'_{k'} \in \mathcal{R}^\circ$, \mathcal{R}_k^b and $\mathcal{R}_{k'}^b$ may have a non-empty intersection or even coincide. Similarly, \mathcal{R}_k^f and $\mathcal{R}_{k'}^f$ may also have a non-empty intersection or coincide. Moreover, \mathcal{R} is non-degenerate if either (i) all FRNs are so or (ii) \mathcal{R}^b is so.

Finally we remark that depending on the choice of \mathcal{S}^1 , an RN can have different fiber decompositions in terms of the BRNs together with the FRNs.

Example 4.4. Consider the RN \mathcal{R} :



(i) Let $\mathcal{S}^1 = \{S_1\}$ and $\mathcal{R}^\circ = \{S_1 \rightarrow 0, 0 \rightarrow S_1, S_1 \rightarrow S_1\}$. Label the three reactions in \mathcal{R}° by 1-3 in the given order. Then $\mathcal{R}^b = \mathcal{R}^\circ$ is degenerate. There exists a unique decomposition (irrespective of the reaction rate constants) satisfying (A2) with $\mathcal{R}_1 = \{S_1 + S_2 \rightarrow S_2 + S_3\}$, $\mathcal{R}_2 = \{S_2 + S_3 \rightarrow S_1 + S_3\}$, and $\mathcal{R}_3 = \{S_1 + S_3 \rightarrow S_1 + S_2\}$. Hence the FRNs $\mathcal{R}_1^f = \{S_2 \rightarrow S_2 + S_3\}$, $\mathcal{R}_2^f = \{S_2 + S_3 \rightarrow S_3\}$, and $\mathcal{R}_3^f = \{S_3 \rightarrow S_2\}$ are all non-degenerate.

(ii) Let $\mathcal{S}^1 = \{S_1, S_2\}$ and $\mathcal{R}^\circ = \{S_1 + S_2 \rightarrow S_2, S_2 \rightarrow S_1, S_1 \rightarrow S_1 + S_2\}$. Label the three reactions in \mathcal{R}° by 1-3 in the given order. Hence $\mathcal{R}^b = \mathcal{R}^\circ$ is non-degenerate. However, with a unique decomposition satisfying (A2), $\mathcal{R}_1^f = \{0 \rightarrow S_3\}$, $\mathcal{R}_2^f = \{S_3 \rightarrow S_3\}$, and $\mathcal{R}_3^f = \{S_3 \rightarrow 0\}$ are *not* all non-degenerate.

5. LIFTING OF REACTION NETWORKS

Using the setup in the above two sections, we will construct a *larger* RN from a *smaller* RN (the reference RN), so that the set of *positive steady states* of the larger RN projected onto the species set of the smaller one coincides with the set of PSSs of the latter. Such reference RN plays a role as the core module of the larger RN. Specifically, in terms of a fiber decomposition of an RN, we look for an RN of the set E of PSSs with a prescribed reference RN of the set E° of PSSs such that $\emptyset \neq E \subseteq E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2}$.

Based on the fiber decomposition of an RN, given a reference RN \mathcal{R}° , there will be diverse ways to construct an RN \mathcal{R} with \mathcal{R}° as its prescribed BRN. In the following, we propose several ways to construct \mathcal{R} preserving the aforementioned stationary property.

Recall that we assume mass-action kinetics. For $y \rightarrow y' \in \mathcal{R}$, let $\kappa_{y \rightarrow y'}$ denote the corresponding rate constant, and for $y_k \rightarrow y'_k \in \mathcal{R}^\circ$, let κ_k° denote the corresponding rate constant.

Assume

$$\text{(A3)} \quad \sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} \text{ is independent of } k \in K_*^\circ.$$

Recall that mass-action kinetics of an RN is determined only by the reactants and the reaction rate constants. Hence there is no restriction on the products of \mathcal{R} (or equivalently, those of the FRNs). Assumption (A3) guarantees that the \mathbf{P}_1 -projection of the set E of PSSs of \mathcal{R} is a subset of the set E° of PSSs of \mathcal{R}° , due to (4.1).

In the light of (A3), assume

(A4) there exists $w \oplus v \in E^\circ \oplus \mathbb{R}_{>0}^{S^2}$ such that

$$\sum_{k \in K^\circ} \sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} \kappa_{y \rightarrow y'} w^{\mathbf{P}_1 y} v^{\mathbf{P}_2 y} \mathbf{P}_2(y' - y) = 0.$$

This assumption guarantees that $\emptyset \neq E \subseteq E^\circ \oplus \mathbb{R}_{>0}^{S^2}$. It is readily verified that the following assumption implies (A4) and ensures $E = E^\circ \oplus \mathbb{R}_{>0}^{S^2}$.

(A5) There exists a mapping $\sigma: \mathcal{S}^2 \rightarrow \mathcal{S}^1$ such that for $k \in K^\circ$ and $y \rightarrow y' \in \mathcal{R}_k$, there exists $b_k^{y \rightarrow y'} = (b_{k,i}^{y \rightarrow y'})_{i \in \mathcal{S}^2} \in (\mathbb{Q} \setminus \{0\})^{\mathcal{S}^2}$ such that for $i \in \mathcal{S}^2$,

$$\mathbf{P}_2(y'_i - y_i) = b_{k,i}^{y \rightarrow y'} ((y'_k)_{\sigma(i)} - (y_k)_{\sigma(i)})$$


and

$$\sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y}$$


is independent of $k \in K_*^\circ$.

From (A5) it follows that $\mathcal{R}_{k,*} = \emptyset$ for all $k \in K^\circ \setminus K_*^\circ$. In other words, all FRNs at degenerate reactions in the reference RN consist of degenerate reactions.

Example 5.1. Consider the mass-action RN \mathcal{R}° :



with $K^\circ = K_*^\circ = \{1, 2\}$ consistent with the indices of the reaction rate constants. Hence (A1) is satisfied. Consider its lifting \mathcal{R} :



where the labels κ_1° and $\kappa_2^\circ/2$ over the arrows are the rate constants associated with the reactions, which are ordered by the indices of κ_i° , $i = 1, 2$. Let $\mathcal{R} = \mathcal{R}_1 \sqcup \mathcal{R}_2$ with \mathcal{R}_i composed of the k -th reaction of \mathcal{R} for $k = 1, 2$. Hence for $k = 1, 2$ and $y \rightarrow y' \in \mathcal{R}_k$, $a_k^{y \rightarrow y'} = k$ and $\mathbf{P}_1 y - y_k = 1$ and $\mathbf{P}_2 y = 3$. Hence $\mathbf{P}_1 \mathcal{R}_k \preceq y_k \rightarrow y'_k$ for $k = 1, 2$, and (A2) is satisfied. Moreover, (A3) is satisfied with

$$\sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} = wv^2, \quad \text{for } k = 1, 2.$$

In addition, $\sigma(2) = 1$, and $b_{k,2}^{y \rightarrow y'} = -k$. Hence (A5) is satisfied with

$$\sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} = -wv^2.$$

It is easy to verify that $E^\circ = \{\kappa_2^\circ/\kappa_1^\circ\}$ and $E = \{\kappa_2^\circ/\kappa_1^\circ\} \oplus \mathbb{R}_{>0}$.

We remark that one can make a more general assumption than (A3), similar to (A5), by assuming the independence coordinate-wise. Such an assumption, however, will sacrifice the structure of the reference RN \mathcal{R}° as a core module of \mathcal{R} .

Theorem 5.2. *Given a non-degenerate mass-action RN \mathcal{R} . Let E and E° be the set of PSSs of \mathcal{R} and \mathcal{R}° , respectively. Assume (A1)-(A4). Then $E \subseteq E^\circ \oplus \mathbb{R}_{>0}^{S^2} \neq \emptyset$. In particular, assume (A5) additionally, then $E = E^\circ \oplus \mathbb{R}_{>0}^{S^2}$.*

Proof. Rewrite (3.2) as

$$(5.1) \quad \dot{x} = \sum_{k \in K^\circ} \sum_{y \rightarrow y' \in \mathcal{R}_{k \in K}} \kappa_{y \rightarrow y'} x^y (y' - y).$$

Let $x = w \oplus v$ with $w \in \mathbb{R}_{\geq 0}^{\mathcal{S}^1}$ and $v \in \mathbb{R}_{\geq 0}^{\mathcal{S}^2}$. In the light of (4.1) and

$$(5.2) \quad y'_k - y_k = 0, \quad \text{for all } k \in \mathbb{K}^\circ \setminus \mathbb{K}_*^\circ,$$

rewrite (5.1) as

$$(5.3) \quad \dot{w} = \sum_{k \in \mathbb{K}_*^\circ} \kappa_k^\circ w^{y_k} (y'_k - y_k) \cdot \sum_{y \rightarrow y' \in \mathcal{R}_k} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ} a_k^{y \rightarrow y'} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y}$$

$$(5.4) \quad \dot{v} = \sum_{k \in \mathbb{K}^\circ} \kappa_k^\circ w^{y_k} \cdot \sum_{y \rightarrow y' \in \mathcal{R}_k} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} (\mathbf{P}_2 y' - \mathbf{P}_2 y).$$

From (5.3) it follows that (A3) and (A4) together imply that $E \subseteq E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2} \neq \emptyset$.

Now assume (A5) additionally. For each $i \in \mathcal{S}^2$,

$$\begin{aligned} \dot{v}_i &= \sum_{k \in \mathbb{K}_*^\circ} \kappa_k^\circ w^{y_k} ((y'_k)_{\sigma(i)} - (y_k)_{\sigma(i)}) \sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} \\ &= \sum_{y \rightarrow y' \in \mathcal{R}_{k,*}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ} w^{\mathbf{P}_1 y - y_k} v^{\mathbf{P}_2 y} \sum_{k \in \mathbb{K}_*^\circ} \kappa_k^\circ w^{y_k} ((y'_k)_{\sigma(i)} - (y_k)_{\sigma(i)}). \end{aligned}$$

Therefore for all $x^* \in E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2} \neq \emptyset$, x^* is a PSS for \mathcal{R} . By (i), $E = E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2}$. \square

We propose more checkable assumptions than (A3) and (A5).

(A6) The sets $\mathcal{C}_1 = \{\mathbf{P}_1 y - y_k : y \rightarrow y' \in \mathcal{R}_k\}$ and $\mathcal{C}_2 = \{\mathbf{P}_2 y : y \rightarrow y' \in \mathcal{R}_k\}$ do not depend on $k \in \mathbb{K}^\circ$.

By (A6), for all $k \in \mathbb{K}^\circ$, \mathcal{R}_k can be decomposed as:

$$\mathcal{R}_k = \sqcup_{z_1 \in \mathcal{C}_1, z_2 \in \mathcal{C}_2} \mathcal{R}_{k, z_1, z_2},$$

where $\mathcal{R}_{k, z_1, z_2} = \{y \rightarrow y' \in \mathcal{R}_k : \mathbf{P}_1 y = z_1 + y_k, \mathbf{P}_2 y = z_2\}$.

(A7) There exists a mapping $\sigma : \mathcal{S}^2 \rightarrow \mathcal{S}^1$ such that for $k \in \mathbb{K}^\circ$ and $y \rightarrow y' \in \mathcal{R}_k$, there exists $b_k^{y \rightarrow y'} = (b_{k,i}^{y \rightarrow y'})_{i \in \mathcal{S}^2} \in (\mathbb{Q} \setminus \{0\})^{\mathcal{S}^2}$ such that for $i \in \mathcal{S}^2$,

$$\mathbf{P}_2 (y' - y)_i = b_{k,i}^{y \rightarrow y'} ((y'_k)_{\sigma(i)} - (y_k)_{\sigma(i)}),$$

both $\sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ}$ and $\sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ}$ are non-zero and independent of $k \in \mathbb{K}_*^\circ$.

Theorem 5.3. *Let \mathcal{R} be a non-degenerate mass-action RN. Let E and E° be the set of PSSs of \mathcal{R} and \mathcal{R}° , respectively. Assume (A1)-(A2) and (A6)-(A7). Then $E = E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2}$.*

Proof. By (A6) and (A7), we can rewrite (5.3) and (5.4) as

$$(5.5) \quad \begin{aligned} \dot{w} &= \sum_{k \in \mathbb{K}_*^\circ} \kappa_k^\circ w^{y_k} (y'_k - y_k) \sum_{z_1 \in \mathcal{C}_1} w^{z_1} \sum_{z_2 \in \mathcal{C}_2} v^{z_2} \\ &\cdot \sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ} \end{aligned}$$

$$(5.6) \quad \begin{aligned} \dot{v}_i &= \sum_{k \in \mathbb{K}_*^\circ} \kappa_k^\circ w^{y_k} ((y'_k)_{\sigma(i)} - (y_k)_{\sigma(i)}) \sum_{z_1 \in \mathcal{C}_1} w^{z_1} \sum_{z_2 \in \mathcal{C}_2} v^{z_2} \\ &\cdot \sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ}, \quad i \in \mathcal{S}^2. \end{aligned}$$

Since $\sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ}$ and $\sum_{y \rightarrow y' \in \mathcal{R}_{k, z_1, z_2}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}^{y \rightarrow y'}}{\kappa_k^\circ}$ are both non-zero and independent of $k \in \mathbb{K}_*^\circ$, we have $E = E^\circ \oplus \mathbb{R}_{>0}^{\mathcal{S}^2}$. \square





FIGURE 1. Streamlines for Example 5.5. Red circle: PSSs.

Definition 5.4. An RN is called *multistationary* if there exists a stoichiometric compatibility class with more than one PSS. Hence potentially an RN may admit multiple PSSs on some stoichiometric compatibility classes while admitting at most one PSS on other stoichiometric compatibility classes, as illustrated by Example 5.5 below. An RN is called *absolute concentration robust* (ACR) if the projection of all positive steady states onto a species S_i ($1 \leq i \leq d$) are identical [34].

Example 5.5. Consider the mass-action RN \mathcal{R}



It is readily verified that the rate equation for \mathcal{R} is

$$\begin{aligned} \dot{w} &= wv(w^2 - 4w + v^2 - 4v + 6) \\ \dot{v} &= -wv(w^2 - 4w + v^2 - 4v + 6). \end{aligned}$$

Moreover, \mathcal{R} is conservative with $w(t) + v(t) = w(0) + v(0)$ for all $t > 0$. Hence $E = \{(x, y) \in \mathbb{R}_{>0}^2 : (x - 2)^2 + (y - 2)^2 = 2\}$ is the set of PSSs. Hence for $c = (c_1, c_2) \in \mathbb{R}_{>0}^2$, S_c admits two PSSs (with the left one being an unstable node and the right one stable) if $2 < c_1 + c_2 < 6$, one PSS (saddle) if $c_1 + c_2 = 2$ or $c_1 + c_2 = 6$, and no PSS if $0 \leq c_1 + c_2 < 2$ or $c_1 + c_2 > 6$. See Figure 1.

The following two corollaries provide a lifting result pertaining to multistationarity and the ACR property. One can also view the results regarding reductions (of a larger RN), as opposed to lifting (of a smaller RN).

For $c \in \mathbb{R}_{>0}^{S_1}$, denote S° and S_c° the stoichiometric subspace and the stoichiometric compatibility class of the reference RN \mathcal{R}° , respectively.

Assume

(A8) For $k \in K_*^\circ$, $\mathcal{R}_k^* \neq \emptyset$.


Corollary 5.6. *Given a non-degenerate mass-action RN \mathcal{R} with non-degenerate mass-action reference RN \mathcal{R}° . Assume (A1)-(A2), (A8), and that \mathcal{R}^b is multistationary. Then \mathcal{R} is multistationary provided either (i) (A3) and (A5) or (ii) (A6) and (A7). In this case, \mathcal{R} is called a multistationarity lifting of \mathcal{R}^b .*

Proof. We only prove the conclusions under (i). The other case can be proved analogously. Let S° be the stoichiometric subspace of \mathcal{R}° . Then


$$S^\circ = \left\{ \sum_{k \in K_*^\circ} c_k (y'_k - y_k) : c_k \in \mathbb{R} \right\}.$$

By **(A1)**, **(A2)** and **(A8)**, we have $\mathbf{P}_1 S = S^\circ$. For every $c \in \mathbb{R}_{>0}^{S_1^1}$, let E_c° be the set of PSSs on the stoichiometric compatibility class S_c . Assume \mathcal{R}° is multistationary. Then $\#E_c^\circ > 1$ for some $c \in \mathbb{R}_{>0}^{S_1^1}$. By Theorem 5.3, we have $E = E^\circ \oplus \mathbb{R}_{>0}^{S_2^2}$. Choose a $\tilde{c} \in \mathbb{R}_{>0}^{S_2^2}$ with $\mathbf{P}_1 \tilde{c} = c$. Hence $E_{\tilde{c}} = E \cap S_{\tilde{c}}$. Since $\mathbf{P}_1 E_{\tilde{c}} = E_{\mathbf{P}_1 \tilde{c}}^\circ$, we have $E_{\tilde{c}} = E_{\mathbf{P}_1 \tilde{c}}^\circ \oplus (\mathbf{P}_2 S_{\tilde{c}} \cap \mathbb{R}_{>0}^{S_2^2})$. By **(A3)**, **(A5)** and **(A8)**, we have $\mathbf{P}_2 S_{\tilde{c}} \cap \mathbb{R}_{>0}^{S_2^2} \neq \emptyset$. Hence $\#E_{\tilde{c}} \geq \#E_{\mathbf{P}_1 \tilde{c}}^\circ > 1$, i.e., \mathcal{R} is also multistationary. \square

Example 5.7. Consider \mathcal{R}



Let $\mathcal{S}^1 = \{S_1\}$ and the reference RN \mathcal{R}° be




Assume $\kappa_2^2 > 4\kappa_1\kappa_3$. Then it is readily verified that $\#E^\circ = 2$. Moreover, in this case, $a_k^{y \rightarrow y'} \equiv 1$, $b_k^{y \rightarrow y'} \equiv -1$. All assumptions **(A1)**, **(A2)**, **(A6)** and **(A7)** are satisfied. By Corollary 5.6, $E = E^\circ \oplus \mathbb{R}$, and \mathcal{R} is a multistationarity lifting of \mathcal{R}^1 .

Corollary 5.8. *Given a non-degenerate mass-action RN \mathcal{R} with mass-action reference RN \mathcal{R}° . Assume **(A1)**-**(A2)**, and \mathcal{R}° is ACR in a species S_i , $i \in \mathcal{S}^1$. Then \mathcal{R} is also ACR in a species S_i with the same ACR value, provided either (i) **(A3)** and **(A4)** or (ii) **(A3)** and **(A5)** or (iii) **(A6)** and **(A7)**. In any case, \mathcal{R} is called an ACR lifting of \mathcal{R}° .*

Proof. The conclusions follow directly from Theorems 5.2 and 5.3. \square

Example 5.9. Consider the following RN \mathcal{R} [34],



Let $\mathcal{S}^1 = \{S_1\}$ and $\mathcal{R}^\circ = \{S_1 \xrightleftharpoons[\kappa_2]{\kappa_1} 0\}$ be the mass-action reference RN. This reaction network is the simple closed SIS epidemic contact network (S_1 represents the number of susceptibles and S_2 that of the infected). Hence **(A1)** is satisfied with $\mathcal{R}^b = \mathcal{R}^\circ$. Moreover, **(A2)** is satisfied with the decomposition $\mathcal{R} = \mathcal{R}_1 \sqcup \mathcal{R}_2$ where $\mathcal{R}_1 = \{S_2 \xrightarrow{1} 2S_2\}$ and $\mathcal{R}_2 = \{S_2 \xrightarrow{1} 0\}$. It is readily verified that **(A6)** is satisfied with $\mathcal{C}_1 = \{0\}$ and $\mathcal{C}_2 = \{S_2\}$, and **(A7)** is satisfied with $\sigma(2) = 1$, $b_{1,2}^{S_1+S_2 \rightarrow 2S_2} = b_{2,2}^{S_2 \rightarrow S_1} = -1$, and

$$\sum_{y \rightarrow y' \in \mathcal{R}_{k,z_1,z_2}} a_k^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} = 1, \quad \sum_{y \rightarrow y' \in \mathcal{R}_{k,z_1,z_2}} b_{k,i}^{y \rightarrow y'} \frac{\kappa_{y \rightarrow y'}}{\kappa_k^\circ} = -1.$$


Since \mathcal{R}^1 is ACR in species S_1 with ACR value $\frac{\kappa_2}{\kappa_1}$, we have by Corollary 5.8 that \mathcal{R} is also ACR in S_1 with the same ACR value.

A celebrated result provides a sufficient condition for ACR regardless of the reaction rate constants [34].

Proposition 5.10. *Let \mathcal{R} be a non-degenerate mass-action RN. Assume $E \neq \emptyset$ and \mathcal{R} has deficiency one. If there exist a pair of non-terminal complexes which differ only in species S_i , then \mathcal{R} is ACR in S_i .*

From the construction of the ACR lifting given in Corollary 5.8, the deficiency of the RN is generally *not* preserved. Hence one can combine Corollary 5.8 with Proposition 5.10 to generate RNs of *high deficiency* from deficiency one core modules, as illustrated by the two-species toy example below, which provides a different ACR lifting than given in Corollary 5.8.

Example 5.11. Consider the following mass-action RN \mathcal{R} :



(note that same reaction rates are identical to each other). The reaction graph associated with \mathcal{R} contains 10 nodes, 2 linkage classes and $\mathbf{S} = \mathbb{R}^2$. Hence the deficiency of \mathcal{R} is $10 - 2 - 2 = 6$.

Assume $\kappa_2 > \kappa_3$. Let $\mathcal{R}^\circ = \{0 \xrightleftharpoons[\kappa_2]{\kappa_1} S_1 \xrightarrow{\kappa_3} 2S_1\}$ be the reference RN. Label the reactions in \mathcal{R}° by the indices of the reaction rate constants. It is easy to show that the rate equation for \mathcal{R}° is

$$\dot{w} = \kappa_1 - (\kappa_2 - \kappa_3)w.$$

Hence the set of PSSs of \mathcal{R}° is given by $E^\circ = \{\frac{\kappa_2 - \kappa_3}{\kappa_1}\}$, and \mathcal{R}° is ACR (one can also deduce this from Proposition 5.10). Hence \mathcal{R} has the decomposition $\mathcal{R} = \sqcup_{i=1}^3 \mathcal{R}_i$. It is easy to verify that the rate equation for \mathcal{R} is

$$(5.7) \quad \begin{cases} \dot{w} = (v - 2)w^2v^3(\kappa_1 - (\kappa_2 - \kappa_3)w), \\ \dot{v} = (w - 3)w^2v^3(\kappa_1 - (\kappa_2 - \kappa_3)w), \end{cases}$$

which implies that

$$\frac{dw}{dv} = \frac{v - 2}{w - 3}, \quad \text{if } w \neq 3.$$

This gives the first integral of (5.8):

$$H(w, v) := (w - 3)^2 - (v - 2)^2,$$

and \mathcal{R} is an integrable system. Moreover, the set of PSSs $E = E^\circ \times \mathbb{R}_{>0} \cup \{(3, 2)\}$. Hence \mathcal{R} is ACR in species S_1 with ACR value 3 whenever $\frac{\kappa_2 - \kappa_3}{\kappa_1} = 3$.

Moreover, the equilibrium $(3, 2)$ is a saddle, $(3, v^*)$ are a stable node for $v^* > 2$ and an unstable node for $v^* < 2$. The invariant manifold through $(3, v^*)$ is $\{(w, v) \in \mathbb{R}_{>0}^2 : (w - 3)^2 - (v - 2)^2 = -(v - v^*)^2\}$, which is not a line (1-d hyperplane) but a hyperbola, for every $v^* \neq 2$. This is different from the conservative system in Example 5.9. See Figure 2.

Example 5.11 illustrates that even certain assumptions (e.g., (A7)) fail, similar lifting still is valid, and can produce stationary dynamics with different geometries. In particular, it demonstrates that the ACR system can have invariant manifolds (characterized in terms of first integrals) which are not hyperplanes (for conservative systems) but hyperbolas, different from what has been observed in the literature (e.g., Example 5.9). This may shed a new light on the study of ACR systems.

Now we present another RN whose invariant manifolds being ellipses.





FIGURE 2. Streamlines for Example 5.11 with $\kappa_1 = 3$, $\kappa_3 - \kappa_2 = 1$. Red line: PSSs.

Example 5.12. Consider the following mass-action RN \mathcal{R} :



(note that some reaction rates are identical). The reaction graph associated with \mathcal{R} contains 11 nodes, 2 linkage classes and $S = \mathbb{R}^2$. Hence the deficiency of \mathcal{R} is $11 - 2 - 2 = 7$.

Assume $\kappa_2 > \kappa_3$. Let \mathcal{R}° defined in Example 5.11 be the reference RN. Hence \mathcal{R}° is ACR. Moreover, \mathcal{R} has the decomposition $\mathcal{R} = \sqcup_{i=1}^3 \mathcal{R}_i$, and the rate equation for \mathcal{R} is

$$(5.8) \quad \begin{cases} \dot{w} = -(v-2)w^2v^3(\kappa_1 - (\kappa_2 - \kappa_3)w), \\ \dot{v} = (w-3)w^2v^3(\kappa_1 - (\kappa_2 - \kappa_3)w), \end{cases}$$

which implies that

$$\frac{dw}{dv} = -\frac{v-2}{w-3}, \quad \text{if } w \neq 3.$$

This gives the first integral of (5.8):

$$H(w, v) := (w-3)^2 + (v-2)^2,$$

and \mathcal{R} is an integrable system. Moreover, the set E of PSSs coincides with that in Example 5.11. Hence \mathcal{R} is ACR in species S_1 with ACR value 3 whenever $\frac{\kappa_2 - \kappa_3}{\kappa_1} = 3$.

Moreover, the equilibrium $(3, 2)$ is a saddle, $(3, v^*)$ are a stable node for $v^* < 2$ and an unstable node for $v^* > 2$. The invariant manifold through $(3, v^*)$ is $\{(w, v) \in \mathbb{R}_{>0}^2 : (w-3)^2 + (v-2)^2 = (v^*-2)^2\}$, which is an ellipse for every $v^* \neq 2$. This is also different from the conservative system in Example 5.9. See Figure 3.




FIGURE 3. Streamlines for Example 5.12 with $\kappa_1 = 3$, $\kappa_3 - \kappa_2 = 1$. Red line: PSSs.

6. OUTLOOKS

(i) In the main results, we provide conditions ensuring the projection of the set of PSSs of an RN \mathcal{R} coincides with that of the reference RN \mathcal{R}° . Nevertheless, it remains unknown if stability of the PSS of \mathcal{R} is consistent with that of PSS of \mathcal{R}° . This is the case for some known lifting schemes [27].


(ii) In this chapter, lifting of RNs preserving the ACR property is based on the fiber decomposition of a large RN. Such a decomposition is expected to preserve good properties of RNs. We list a few questions here:

- (a) Under what conditions can dynamical/algebraic properties of an RN be deduced barely from its BRN and the FRNs?
- (b) Are properties consistent for an RN and its BRN and FRNs? For instance, if all FRNs are of deficiency zero, is the original reaction network so? Or, if all FRNs are complex-balanced [26], are the original reaction network also complex-balanced?

(iii) The notion of decomposition of RNs has been applied to analysis of metabolic reaction networks and in bioinformatics [35, 24, 28]. We expect the fiber decomposition may play a role in these regards as well.

(iv) The fiber decomposition of RNs is readily adapted to *stochastic reaction networks* (by only adjusting the kinetics). In light of the applications of fiber decomposition in the deterministic setting, we believe the analogue for stochastic reaction networks might also play an important role on similar topics (i.e., lifting stochastic reaction networks preserving for example stationary properties [25] and *structural classification* [36]).


Example 6.1. Consider \mathcal{R}° :



where $\kappa_i > 0$, $\kappa_2 > \kappa_3$, $\kappa_4 < \kappa_5$ and $(\kappa_3 - \kappa_2)^2 = 4\kappa_1(\kappa_5 - \kappa_4)$. The rate equation for \mathcal{R}° is

$$\dot{w} = (\kappa_5 - \kappa_4)(w - w_*)^2,$$

where $w_* = \frac{\kappa_2 - \kappa_3}{2(\kappa_5 - \kappa_4)}$. Hence \mathcal{R}° is ACR. Consider the lifting \mathcal{R} of \mathcal{R}° :



The rate equation for \mathcal{R} is

$$\begin{aligned} \dot{w} &= v(\kappa_5 - \kappa_4)(w - w_*)^2, \\ \dot{v} &= -v(\kappa_5 - \kappa_4)(w - w_*)^2. \end{aligned}$$

It is readily verified that \mathcal{R} is conservative and an ACR-lifting of \mathcal{R}° . Nonetheless, by [36, Theorem 4.6], \mathcal{R} is positive recurrent on each of its finite compatibility classes while \mathcal{R}° is explosive a.s. on \mathbb{N}_0 .

This example reveals that an ACR lifting may *not* preserve the stochastic dynamics (e.g., explosivity) for the respective stochastic reaction networks.

ACKNOWLEDGEMENTS

CW acknowledges funding from the Novo Nordisk Foundation, Denmark. CX acknowledges the TUM Foundation Fellowship as well as the Alexander von Humboldt Fellowship funded by Alexander von Humboldt Foundation, Germany.

REFERENCES

- [1] ANDERSON, D.F. and CAPPELLETTI, D. Discrepancies between extinction events and boundary equilibria in reaction networks. *J. Math. Biol.*, **79**:1253–1277, 2019.
- [2] ANDERSON, D.F., CAPPELLETTI, D., and KURTZ, T.G. Finite time distributions of stochastically modeled chemical systems with absolute concentration robustness. *SIAM J. Appl. Dyn. Syst.*, **16**:1309–1339, 2017.
- [3] ANDERSON, D.F., ENCISO, G., and JOHNSTON, M.D. Stochastic analysis of biochemical reaction networks with absolute concentration robustness. *J. R. Soc. Interface*, **11**:20130943, 2014.
- [4] BALCAN, D., ET AL. Multiscale mobility networks and the spatial spreading of infectious diseases. *Proc. Natl. Acad. Sci. USA*, 106:21484–21489, 2009.
- [5] BANAJI, M. and PANTEA, C. The inheritance of nondegenerate multistationarity in chemical reaction networks. *SIAM J Appl Math*, **78**:1105–1130, 2018.
- [6] BERGER, S.I. and IYENGAR, R. Network analyses in systems pharmacology. *Bioinformatics*, 25:2466–2472, 2009.
- [7] BRIAT, C., GUPTA, A., and KHAMMASH, M. Antithetic integral feedback ensures robust perfect adaptation in noisy biomolecular networks. *Cell Syst.*, pages 15–26.
- [8] CAPPELLETTI, D., GUPTA, A., and KHAMMASH, M. A hidden integral structure endows absolute concentration robust systems with resilience to dynamical concentration disturbances. *J. Royal Soc. Interface*, page 20200437.
- [9] CHARLEBOIS, D.A. Multiscale effects of heating and cooling on genes and gene networks. *Proc. Natl. Acad. Sci. USA*, **115**:E10797–E10806, 2018.
- [10] CONRADI, C., FELIU, E., MINCHEVA, M., and WIUF, C. Identifying parameter regions for multistationarity. *PLOS Comp Biol*, 13:e1005751, 2017.
- [11] CONRADI, C., FLOCKERZI, D., RAISCH, J., and STELLING, J. Subnetwork analysis reveals dynamic features of complex (bio)chemical networks. *Proc. Natl. Acad. Sci. USA*, **104**:19175–19180, 2007.
- [12] CRACIUN, G. and FEINBERG, M. Multiple equilibria in complex chemical reaction networks. I. the injectivity property. *SIAM J. Appl. Math.*, **65**:1526–1546, 2005.
- [13] CRACIUN, G. and FEINBERG, M. Multiple equilibria in complex chemical reaction networks. II. the species-reaction graph. *SIAM J. Appl. Math.*, **66**:1321–1338, 2006.
- [14] CRACIUN, G. and FEINBERG, M. Multiple equilibria in complex chemical reaction networks: semiopen mass action systems. *SIAM J. Appl. Math.*, **70**:1859–1877, 2010.
- [15] CRACIUN, G., HELTON, J.W., and WILLIAMS, R.J. Homotopy methods for counting reaction network equilibria. *Math. Biosci.*, **216**:140–149, 2008.
- [16] DE FREITAS, M. M., FELIU, E., and WIUF, C. Intermediates, catalysts, persistence, and boundary steady states. *J. Math. Biol.*, **74**:887–932, 2017.
- [17] DE FREITAS, M. M., WIUF, C., and FELIU, E. Intermediates and generic convergence to equilibria. *Bull. Math. Biol.*, **79**:1662–1686, 2017.
- [18] DOMÍNGUEZ-GARCÍA, V., DAKOS, V., and KÉFI, S. Unveiling dimensions of stability in complex ecological networks. *Proc. Natl. Acad. Sci. USA*, **116**:25714–25720, 2019.
- [19] ELLISON, P. *The Advanced Deficiency Algorithm and Its Applications to Mechanism Discrimination*. Ph.D. thesis. University of Rochester, 1998.
- [20] ENCISO, G. and KIM, J. Absolutely robust controllers for chemical reaction networks. *J. Royal Soc. Interface*, **17**:20200031, 2020.
- [21] FEINBERG, M. *Foundations of Chemical Reaction Network Theory*. Applied Mathematical Sciences. Springer International Publishing, Cham, 2019.
- [22] FELIU, E. and WIUF, C. Enzyme sharing as a cause of multistationarity in signaling systems. *J. R. Soc. Interface*, **9**:1224–1232, 2011.
- [23] GROSS, E., HARRINGTON, H. MESHKAT, N., and SHIU, A. Joining and decomposing reaction networks. *J. Math. Biol.*, **80**:1683–1731, 2020.

- [24] HAUS, U.-U. and HEMMECKE, R. Decomposition of reaction networks: The initial phase of the permanganate/oxalic acid reaction. *J. Math. Chem.*, **48**:305–312, 2010.
- [25] HOESSLY, L. Stationary distributions via decomposition of stochastic reaction networks. <https://arxiv.org/abs/1910.02871>, 2019.
- [26] HORN, F. and JACKSON, R. General mass action kinetics. *Arch. Ration. Mech. Anal.*, **47**:81–116, 1972.
- [27] JOSHI, B. and SHIU, A. Atoms of multistationarity in chemical reaction networks. *J. Math. Chem.*, **51**:153–178, 2013.
- [28] KALTENBACH, H.M., CONSTANTINESCU, S., FEIGELMAN, J., and STELLING, J. Graph-based decomposition of biochemical reaction networks into monotone subsystems. In PRZYTYCKA T.M. and SAGOT M.F., editors, *Algorithms in Bioinformatics (WABI 2011)*, volume 6833 of *Lecture Notes in Computer Science*, pages 139–150. Springer-Verlag, Berlin, Heidelberg, 2011.
- [29] PAPIN, J.A., ET AL. Reconstruction of cellular signalling networks and analysis of their properties. *Nat. Rev. Mol. Cell Biol.*, **6**:99–111, 2005.
- [30] PASCUAL-ESCUERO, B. and FELIU, E. Local and global robustness in systems of polynomial equations. <https://arxiv.org/abs/2005.08796>, 2020.
- [31] PEEL, L., DELVENNE, J.-C., and LAMBIOTTE, R. Multiscale mixing patterns in networks. *Proc. Natl. Acad. Sci. USA*, **115**:4057–4062, 2018.
- [32] SCHUETZ, R., ET AL. Multidimensional optimality of microbial metabolism. *Science*, **336**:601–604, 2012.
- [33] SHINAR, G., ALON, U., and FEINBERG, M. Sensitivity and robustness in chemical reaction networks. *SIAM J. Appl. Math.*, **69**:977–998, 2009.
- [34] SHINAR, G. and FEINBERG, M. Structural sources of robustness in biochemical reaction networks. *Science*, **327**:1389–1391, 2010.
- [35] YOON, J., SI, Y., NOLAN, R., and LEE, K. Modular decomposition of metabolic reaction networks based on flux analysis and pathway projection. *Bioinformatics*, **23**:2433–2440, 2007.
- [36] C. Wiuf and C. Xu. Classification and threshold dynamics of stochastic reaction networks. <https://arxiv.org/abs/2012.07954>, 2020.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, COPENHAGEN, 2100, DENMARK

FACULTY OF MATHEMATICS, TECHNICAL UNIVERSITY OF MUNICH, MUNICH, GARCHING BEI MÜNCHEN, 85748, GERMANY.

Email address: wiuf@math.ku.dk

Email address: xuc@ma.tum.de