

# SPECIFYING A GAME-THEORETIC EXTENSIVE FORM AS AN ABSTRACT 5-ARY RELATION

Peter A. Streufert  
Economics Department  
Western University

ABSTRACT. This paper specifies an extensive form as a 5-ary relation (that is, as a set of quintuples) which satisfies eight abstract axioms. Each quintuple is understood to list a player, a situation (that is, a name for an information set), a decision node, an action, and a successor node. Accordingly, the axioms are understood to specify abstract relationships between players, situations, nodes, and actions. Such an extensive form is called a “pentaform”. Finally, a “pentaform game” is defined to be a pentaform together with utility functions.

To ground this new specification in the literature, the paper defines the concept of a “traditional game” to represent the literature’s many specifications of finite-horizon and infinite-horizon games. The paper’s main result is to construct an intuitive bijection between pentaform games and traditional games. Secondary results concern disaggregating pentaforms by subsets, constructing pentaforms by unions, and initial pentaform applications to Selten subgames and perfect-recall (an extensive application to dynamic programming is in Streufert 2023, [arXiv:2302.03855](#)).

## 1. INTRODUCTION

### 1.1. NEW CONCEPTS AND MAIN RESULT

A 5-ary relation is merely a set of quintuples. It is like a binary relation, which is a set of couples (ordered pairs), and also like a ternary relation, which is a set of triples.<sup>1</sup> In this paper, a quintuple is denoted  $\langle i, j, w, a, y \rangle$ . The first element is understood to be a player, the second a situation (that is, a name for an information set),<sup>2</sup> the third a decision node, the fourth an action, and the fifth a successor node. Thereby, a set

---

*Date:* July 5, 2024. Supersedes [arXiv:2107.10801v4](#) (the text’s results are essentially unchanged; the exposition and appendices are considerably improved). *Keywords:* extensive-form game, pentaform, subgame, perfect recall. *Classifications:* JEL C73, MSC 91A70. *Contact information:* pstreuf@uwo.ca, 519-661-3500, Economics Department, Western University, London, Ontario, N6A 5C2, Canada.

This paper has benefitted greatly from the extensive comments of an anonymous associate editor and two anonymous reviewers.

<sup>1</sup>This concept of a relation as a set accords with Halmos 1974, Section 7, and Enderton 1977, pages 41–42. In a similar vein, this paper regards functions (footnote 13) and correspondences (footnote 17) as sets of couples.

<sup>2</sup>In this paper, a “situation” can be either an information set (which is a set of nodes), or something else (such as a word like “tomorrow”). A situation can be interpreted as a name for an information set, or more abstractly, as a decision point. Regardless of interpretation, axiom [Pj+w] implies a bijection between situations and information sets (footnote 16). Myerson 1991 has a similar concept (footnote 8).

of quintuples is understood to specify relationships between players, situations, (two kinds of) nodes, and actions.

Such a quintuple set can specify a game-theoretic extensive form because an extensive form is, in essence, a collection of relationships between players, situations, nodes, and actions. Correspondingly, this paper specifies an extensive form as a “pentaform”, which is defined to be a quintuple set  $Q$  which obeys eight axioms. These axioms are formulated in terms of various projections of  $Q$ . For example, let  $\pi_{JI}(Q)$  denote the projection of  $Q$  onto its first two coordinates, with their order reversed. Then the first axiom requires that  $\pi_{JI}(Q)$  is a function. In other words, the axiom requires that each situation  $j$  is associated with exactly one player  $i$ . Intuitively, that player  $i$  is the one who controls the action at situation  $j$ . In a similar way, each of the other seven axioms formalizes one small independent feature of an extensive form.

Finally, a “pentaform game” is constructed by combining a pentaform with utility functions.<sup>3</sup> To ground this new specification in the literature, the paper defines the concept of a “traditional game” to represent the literature’s many specifications of finite-horizon and infinite-horizon games. Such a traditional game is defined in the usual way as a tree adorned with information sets, actions, players, and utility functions. The paper’s main result is Theorem 5.4, which shows that there is a constructive and intuitive bijection from the collection of traditional games to the collection of pentaform games with information-set situations. This suggests that pentaform games can equivalently formulate all discrete<sup>4</sup> extensive-form games in the literature.

## 1.2. MOTIVATION

Pentaforms are easy to manipulate because they are sets, and because the pentaform axioms are largely compatible with the concepts of subset and union.

More precisely, Section 4.1 shows that any subset of a pentaform satisfies six of the eight pentaform axioms (Proposition 4.1). This leads to a weak general condition under which a subset of a pentaform is itself a pentaform (Corollary 4.2). This result is a powerful tool for disaggregating a pentaform. First, Section 4.1 uses the result to characterize the subsets of a pentaform that correspond to Selten 1975 subgames (Proposition 4.3). Second, Section 4.1 explains how the same result, applied to other subsets of a pentaform, is the foundation for the generalized theory of dynamic programming in Streufert 2023. That sequel paper is the first paper to use value functions to characterize subgame perfection in arbitrary games.

---

<sup>3</sup>This paper does not formally consider probability. If a game has a finite number of nodes, mixed strategies and expected utilities can be derived by standard means (for example, Myerson 1991, Chapter 4). Meanwhile, if there are an infinite number of nodes, the very concepts of mixed strategy and expected utility can lead to subtle measurability issues.

<sup>4</sup>“Discreteness” means that each decision node has a finite number of predecessors. Non-discrete games include those in continuous time, as in Dockner, Jørgenson, Long, and Sorger 2000; and those yet more general, as in Alós-Ferrer and Ritzberger 2016, Chapters 1–5. [Discreteness is defined in terms of *decision* nodes because Alós-Ferrer and Ritzberger 2016, Section 6.2, admits a terminal node at the end of each infinite run (that is, each infinite play). Such terminal nodes do not appear in the present paper.]

In the opposite direction, Section 4.2 defines a “block” to be a quintuple set which satisfies all but one of the axioms. Then it essentially shows that the union of a “separated” collection of blocks is itself a block (Proposition 4.4), and that the union of an expanding sequence of pentaforms is itself a pentaform (Proposition 4.5). These are convenient tools for building new pentaforms from known components, as illustrated by the finite-horizon examples of Section 4.2 and the infinite-horizon example of Streufert 2023, Section 2.2. These techniques are complementary to those of Capucci, Ghani, Ledent, and Nordvall Forsberg 2022 in the computer-science literature (footnote 21).

In addition, pentaforms seem to have some less tangible benefits.

(a) Other axiomatic foundations, such as the well-known foundation for consumer preferences, have readily fostered new extensions and results. The same may occur with this paper’s axiomatic foundation for extensive forms. For example, the fine-grainedness of the eight axioms fostered the development of Propositions 4.1 and 4.4 for subsets and unions.

(b) The pentaform notation is distinctly new. Broadly, a pentaform is one high-dimensional relation, while a traditional extensive form is a list of low-dimensional relations. To suggest the value of this unification, Section 4.3 considers the concept of perfect-recall (Kuhn 1950, 1953), which simultaneously involves players, situations, nodes, and actions. It turns out that the concept can be efficiently reformulated in terms of pentaforms. The new formulation revolves around successor nodes rather than decision nodes, and is easily manipulated to produce a one-paragraph proof of the necessity of no-absentmindedness.

### 1.3. BACKGROUND

There have been many innovations in how to specify an extensive form. This paragraph discusses three broad strands in that literature. (1) Eighty years ago, von Neumann and Morgenstern 1944 (pages 73–74; 78) specified each node as a set of outcomes, and noted that set inclusion arranges these nodes in a tree. (2) Later, Kuhn 1953 specified each node as an abstract entity without any internal structure, and then separately specified the edges of the tree connecting the nodes. (3) Next, Osborne and Rubinstein 1994 (page 200) specified each node as a sequence of past actions, and thereby created a tractable specification of infinite-horizon extensive forms. More recently, Ritzberger 2002 (page 97) returned to the outcome-set formulation of von Neumann and Morgenstern 1944 and extended it to allow for an infinite horizon. Then Alós-Ferrer and Ritzberger 2016 (Section 6.3), Kline and Luckraz 2016, and Streufert 2019<sup>5</sup> formally connected the three strands of literature, by building bijections between their three classes of extensive forms. Infinite-horizon extensive forms are incorporated everywhere.

This paper’s main result (Theorem 5.4) uses a bijection to formally connect a generic extension (Definition 5.1) of Kuhn 1953 to the new pentaform specification (Definition 3.5). The pentaform specification has antecedents. In some fashion,

---

<sup>5</sup>For the purposes here, lump Streufert 2019’s choice-set form with its OR form, and lump its KS form with its simple form.

every extensive-form specification must determine a function which takes decision-node/feasible-action couples to successor nodes. Often this function is implicitly specified by assigning actions to the edges of a tree diagram. Alternatively, the function can be explicitly specified as a decision-node-indexed list of functions from actions to successor nodes (Kline and Luckraz 2016) or, in the reverse direction, as a pair of functions mapping each successor node to the immediately preceding node and the immediately preceding action (Mas-Colell, Whinston, and Green 1995, page 227).<sup>6</sup> Streufert 2018 departs by viewing this function in set-theoretic terms, as a set of triples. The present paper augments those triples with players and situations to form quintuples, and then defines an extensive form in terms of those quintuples.

#### 1.4. ORGANIZATION

Section 2 builds intuition through examples. Section 3 defines pentaform games. Section 4 discusses pentaform applications, and is unrelated to Section 5, which defines “traditional games” and relates them to pentaform games. Finally, Appendix A concerns graph-theoretic trees, and Appendices B–D provide lemmas and proofs for Sections 3–5, respectively.

## 2. INITIAL INTUITION

This brief section builds intuition through examples. The examples suggest how an extensive form can be expressed by a quintuple set (that is, by a 5-ary relation). This section presumes familiarity with tree diagrams.




FIGURE 2.1. By definition,  $Q^1$  is the set consisting of the table’s two rows, that is,  $\{ \langle \text{Alex}, \{0\}, 0, \text{left}, 1 \rangle, \langle \text{Alex}, \{0\}, 0, \text{right}, 2 \rangle \}$ . The tree diagram provides the same data. The set  $Q^1$  is a “pentaform” (Section 3.4).

Figure 2.1’s tree diagram has three nodes (0, 1, and 2), two actions (left and right), one player (Alex), and one information set ( $\{0\}$ ). The root node (0) is underlined, and there are two edges (that is, “arcs” or “twigs”). These edges can be denoted  $\langle 0, 1 \rangle$  and  $\langle 0, 2 \rangle$  (brackets  $\langle \rangle$  will appear only on tuples). Note that the action left labels the edge  $\langle 0, 1 \rangle$ , while the action right labels the edge  $\langle 0, 2 \rangle$ . This data can be encoded within the triples  $\langle 0, \text{left}, 1 \rangle$  and  $\langle 0, \text{right}, 2 \rangle$ . Next, the node 0 is in the information set  $\{0\}$ . This (self-evident) fact can be encoded within the quadruples

<sup>6</sup>A similar function from two variables to one can be found in the transition function of a stochastic game (Mertens 2002), and in the labelled transition system, i.e. state machine, of computer science (Blackburn, de Rijke, and Venema 2001, page 3).




FIGURE 2.2. By definition,  $Q^2$  is the set consisting of the table’s rows (expressed as quintuples). The tree diagram provides the same data. The set  $Q^2$  is a “pentaform” (Section 3.4).

$\langle \{0\}, 0, \text{left}, 1 \rangle$  and  $\langle \{0\}, 0, \text{right}, 2 \rangle$ . Finally, the player Alex makes the decision at information set  $\{0\}$ . This fact can be encoded within the quintuples  $\langle \text{Alex}, \{0\}, 0, \text{left}, 1 \rangle$  and  $\langle \text{Alex}, \{0\}, 0, \text{right}, 2 \rangle$ . In this sense, the set

$$Q^1 = \{ \langle \text{Alex}, \{0\}, 0, \text{left}, 1 \rangle, \langle \text{Alex}, \{0\}, 0, \text{right}, 2 \rangle \}$$

expresses Figure 2.1’s tree diagram. The set’s two quintuples correspond to the two (non-header) rows in the figure’s table (eventually the information set  $\{0\}$  will be regarded as a special kind of situation). Finally, the superscript on  $Q^1$  distinguishes this first example from future examples.

This process can be readily generalized. Essentially, each tree edge is changed into a quintuple. To be more specific, each tree edge is a couple  $\langle w, y \rangle$  consisting of a decision node  $w$  and a successor node  $y$ . This couple is changed into the quintuple  $\langle i, j, w, a, y \rangle$  in which  $a$  is the action labelling the edge  $\langle w, y \rangle$ ,  $j$  is the information set containing the decision node  $w$ , and  $i$  is the player making the decision at information set  $j$ . Eventually the information set  $j$  will be regarded as a special kind of situation.

Figure 2.2’s tree diagram is more complicated than the previous example because it has a nonsingleton information set.<sup>7</sup> Nonetheless it can be expressed as a quintuple set by the same process. The tree has eight edges. The edge  $\langle 3, 7 \rangle$  is changed to the quintuple  $\langle \text{Teacher}, \{2, 3\}, 3, f, 7 \rangle$  to encode the facts that (i) the action  $f$  labels the edge  $\langle 3, 7 \rangle$ , (ii) the decision node 3 is (self-evidently) in the information set  $\{2, 3\}$ , and (iii) the player Teacher controls the move at the information set  $\{2, 3\}$ . By similarly changing the other seven edges  $\langle w, y \rangle$  to quintuples  $\langle i, j, w, a, y \rangle$ , one obtains all eight rows in Figure 2.2’s table. Those rows define the set  $Q^2$ .

<sup>7</sup>Figures 2.2 and 2.3 (for examples  $Q^2$  and  $Q^3$ ) correspond to Selten 1975’s well-known “horse” game. To tell a story for these figures, suppose a Kid must decide, today, between the bad action of not doing her homework (called  $b$ ) and the correct action of doing her homework (called  $c$ ). Next, tonight, if the homework has been finished (node 1), a Dog must decide between the dumb action of eating the homework ( $d$ ) and the good action of going back to sleep ( $g$ ). Finally, tomorrow, without knowing whether the kid chose bad (node 2) or the kid chose correct and the dog chose dumb (node 3), the Teacher must decide between excusing the kid ( $e$ ) and failing the kid ( $f$ ).




FIGURE 2.3. By definition,  $Q^3$  is the set of the table’s rows (expressed as quintuples). The tree diagram provides the same data. In  $Q^3$ , situations are not information sets. The set  $Q^3$  is a “pentaform” (Section 3.4).

Figure 2.3’s tree diagram differs from the previous example to the extent that its information sets are named with the words “today”, “tonight”, and “tomorrow” (these words have meaning within the story of footnote 7). These three words are examples of situations. As footnote 2 explains, situations can be either information sets (as in the first two examples) or something else (such as the three words here). A situation can be interpreted as a name for an information set, or more abstractly, a decision point.<sup>8</sup>

This third example can be expressed as a quintuple set just as the previous two examples. Specifically, the edge  $\langle 3,7 \rangle$  becomes the quintuple  $\langle \text{Teacher, tomorrow, 3, f, 7} \rangle$  to encode the facts that (i) the action  $f$  labels the edge  $\langle 3,7 \rangle$ , (ii) the decision node 3 is associated with the situation **tomorrow**, and (iii) the player **Teacher** controls the move in the situation **tomorrow**. By similarly changing the other seven edges to quintuples, one obtains all eight rows in Figure 2.3’s table. These rows define the set  $Q^3$ .

For a fourth example, consider the quintuple set

$$(1) \quad Q^4 = \{ \langle 41,42,43,44,45 \rangle, \langle 46,47,48,49,50 \rangle \}.$$

Obviously, an arbitrary quintuple set like  $Q^4$  may or may not express a tree diagram.

Looking ahead, Definition 3.1 will designate which quintuple sets are to be called “pentaforms”, and Definition 3.5 will define a “pentaform game” to be a pentaform augmented with utility functions. In the end, Theorem 5.4 (the paper’s main result) will show that there is a bijection from the collection of “traditional games” to the collection of pentaform games with information-set situations. The forward direction of that bijection closely resembles this section’s informal process of expressing Figure 2.1 and 2.2’s tree diagrams as quintuple sets.

<sup>8</sup>A situation  $j$  is comparable to an information state  $s$  in Myerson 1991, Chapter 4 (there players are assumed to have disjoint sets of information states, unlike in Myerson 1991, Chapter 2).

## 3. PENTAFORM GAMES

This Section 3 defines pentaforms and pentaform games. The examples of Section 2 are used as illustrations.

## 3.1. THE COMPONENTS OF A QUINTUPLE

An arbitrary quintuple will be denoted  $\langle i, j, w, a, y \rangle$ . Call its first component  $i$  the *player*, call its second component  $j$  the *situation*, call its the third component  $w$  the *decision node*, call its fourth component  $a$  the *action*, and call its fifth component  $y$  the *successor node*. These five terms have no formal content. They merely name the five positions in a quintuple. For example, in the quintuple  $\langle 46, 47, 48, 49, 50 \rangle$ , the player is 46, the situation is 47, the decision node is 48, the action is 49, and the successor node is 50. Further, let the *nodes* of a quintuple be its decision node and its successor node. In other words, let the nodes of a quintuple be its third and fifth components. For example, the nodes of  $\langle 46, 47, 48, 49, 50 \rangle$  are 48 and 50.<sup>9</sup>

This concept of quintuple is extremely (and perhaps disturbingly) abstract in the sense that no structure is imposed on the components of a quintuple. For example, in the quintuple  $\langle 46, 47, 48, 49, 50 \rangle$ , each of the five components is an alphanumeric string (strings use the sans serif font). Similarly, in the quintuple  $\langle \text{Teacher}, \text{tomorrow}, 3, \text{f}, 7 \rangle$  from Figure 2.3, each of the five components is an alphanumeric string. Meanwhile, in the quintuple  $\langle \text{Teacher}, \{2, 3\}, 3, \text{f}, 7 \rangle$  from Figure 2.2, four of the five components are alphanumeric strings, but one of the components is a set of alphanumeric strings. In general, each component can be practically anything, including a number, a set of numbers, or a sequence of numbers or strings.

One advantage of this abstract formulation is that it is extremely flexible. One use of this flexibility is to directly accommodate the literature’s wide variety of notations for situations, nodes, and actions. For example, situations can be specified as sets of nodes (as in the information-set situation  $\{2, 3\}$  from Figure 2.2) or without any special structure (as in the situation *tomorrow* from Figure 2.3). Meanwhile, nodes can be specified as sequences of actions (Osborne and Rubinstein 1994), as sets of actions (Streufert 2019), as sets of outcomes (Alós-Ferrer and Ritzberger 2016, Section 6.2; see footnote 4 here), or without any special structure (as in the node 3 from Figures 2.2 and 2.3). Finally, actions can be specified as sets of nodes (van Damme 1991, page 103), as sets of edges (Selten 1975), as sets of outcomes (Alós-Ferrer and Ritzberger 2016, Section 6.2; see footnote 4 here), or without any special structure (as in the action *f* from Figures 2.2 and 2.3).

## 3.2. QUINTUPLE SETS AND THEIR SLICES

A set of quintuples will typically be denoted by the letter  $Q$ . Correspondingly, different quintuple sets will typically be distinguished from one another by means of markings around the letter  $Q$ . For instance, the examples in Section 2 are denoted  $Q^1$ ,  $Q^2$ ,  $Q^3$ , and  $Q^4$ . Similarly, Section 4.1 will consider subsets of a quintuple set  $Q$

<sup>9</sup>When necessary, a generic node will be denoted by  $x$ . Mnemonically,  $w$  is before  $x$  is before  $y$  in the alphabet, and correspondingly, decision nodes  $w$  are “early” nodes, and successor nodes  $y$  are “late” nodes.

denoted  $Q' \subseteq Q$  and  ${}^tQ \subseteq Q$ . Likewise, this section will consider other subsets of a quintuple set  $Q$  which will be denoted  $Q_j \subseteq Q$ .

Now consider an arbitrary quintuple set  $Q$ , and let  $J$  denote its set of situations  $j$ . In other words, let  $J$  be the projection of  $Q$  onto its second coordinate. Then, for each situation  $j \in J$ , define

$$(2) \quad Q_j = \{ \langle i_*, j, w_*, a_*, y_* \rangle \in Q \}.$$

Thus  $Q_j$  is the set of quintuples in  $Q$  that have situation  $j$ . Call  $Q_j$  the *slice of  $Q$  for situation  $j$* . By inspection,  $\langle Q_j \rangle_{j \in J}$  is an injectively indexed partition of  $Q$ . Call this the *slice partition of  $Q$* .

For example, consider example  $Q^3$  from Figure 2.3. In this example, the situation set  $J^3$  is  $\{\text{today, tonight, tomorrow}\}$ . Further, the slice partition  $\langle Q_j^3 \rangle_{j \in J^3}$  divides  $Q^3$  into the three quintuple sets

$$(3a) \quad Q_{\text{today}}^3 = \{ \langle \text{Kid, today, 0, c, 1} \rangle, \langle \text{Kid, today, 0, b, 2} \rangle \},$$

$$(3b) \quad Q_{\text{tonight}}^3 = \{ \langle \text{Dog, tonight, 1, g, 8} \rangle, \langle \text{Dog, tonight, 1, d, 3} \rangle \}, \text{ and}$$

$$(3c) \quad Q_{\text{tomorrow}}^3 = \{ \langle \text{Teacher, tomorrow, 2, e, 4} \rangle, \langle \text{Teacher, tomorrow, 2, f, 5} \rangle, \\ \langle \text{Teacher, tomorrow, 3, e, 6} \rangle, \langle \text{Teacher, tomorrow, 3, f, 7} \rangle \}.$$

These three sets are illustrated in Figure 3.0.<sup>10</sup>




FIGURE 3.0. The slice partition of  $Q^3$ , as illustrated by both the tree diagram and the table. The set  $Q^3$  is from Figure 2.3 and is a pentaform (Section 3.4). The partition's three slices are written explicitly in eq. (3).

<sup>10</sup>The unusual figure number “3.0” is intended to simplify example references. In the end, Figures 2.1 and 3.1 concern example  $Q^1$ ; Figures 2.2 and 3.2 concern example  $Q^2$ ; and Figures 2.3 and 3.0 concern example  $Q^3$ .



### 3.3. PROJECTIONS

Any quintuple set can be projected onto any sequence in  $\{I, J, W, A, Y\}$ . For example, Figure 2.1's table for  $Q^1$  implies<sup>11</sup>

$$(4) \quad \pi_Y(Q^1) = \{y \mid (\exists i, j, w, a) \langle i, j, w, a, y \rangle \in Q^1\} = \{1, 2\} \text{ and}$$

$$(5) \quad \pi_{JI}(Q^1) = \{\langle j, i \rangle \mid (\exists w, a, y) \langle i, j, w, a, y \rangle \in Q^1\} = \{\langle 0 \rangle, \text{Alex}\}.$$

For another example, equation (3c) for  $Q_{\text{tomorrow}}^3$  implies

$$(6) \quad \pi_A(Q_{\text{tomorrow}}^3) = \{a \mid (\exists i, j, w, y) \langle i, j, w, a, y \rangle \in Q_{\text{tomorrow}}^3\} = \{e, f\} \text{ and}$$

$$(7) \quad \begin{aligned} \pi_{WA}(Q_{\text{tomorrow}}^3) &= \{\langle w, a \rangle \mid (\exists i, j, y) \langle i, j, w, a, y \rangle \in Q_{\text{tomorrow}}^3\} \\ &= \{\langle 2, e \rangle, \langle 2, f \rangle, \langle 3, e \rangle, \langle 3, f \rangle\}. \end{aligned}$$

Note that projections, like  $\pi_{JI}(Q^1)$  in (5), can re-order the coordinates. Also note that projections of slices are well-defined simply because slices are quintuple sets. An example is the slice  $Q_{\text{tomorrow}}^3$  in (3c), (6), and (7).

Both slices and projections can be visualized by tables. Slices select rows and then projections select columns. For example, consider Figure 3.0's table for  $Q^3$ . Its last four rows constitute the slice  $Q_{\text{tomorrow}}^3$  in (3c), and this slice's fourth column determines the projection  $\pi_A(Q_{\text{tomorrow}}^3)$  in (6).

The notation for a single-coordinate projection will often be abbreviated by replacing the letter  $Q$  with the single coordinate. Specifically, define the five abbreviations  $I$ ,  $J$ ,  $W$ ,  $A$ , and  $Y$  by

$$(8) \quad I = \pi_I(Q), \quad J = \pi_J(Q), \quad W = \pi_W(Q), \quad A = \pi_A(Q), \quad \text{and} \quad Y = \pi_Y(Q).$$

These abbreviations inherit any markings on the letter  $Q$ . For example, (4) shows that  $Y^1 = \pi_Y(Q^1)$  is  $\{1, 2\}$ . Similarly, (6) shows that  $A_{\text{tomorrow}}^3 = \pi_A(Q_{\text{tomorrow}}^3)$  is  $\{e, f\}$ .

Two special cases will become important after Definition 3.1. Both concern a quintuple set  $Q$  and one of its situations  $j \in J$  (that is,  $j \in \pi_J(Q)$ ).

First,  $W_j$  (that is  $\pi_W(Q_j)$ ) is the decision-node set of situation  $j$ . In order to accord with standard terminology, call  $W_j$  the *information set* of situation  $j$ . For instance, in example  $Q^3$ , Figure 3.0's table shows that the information set of the situation  $j = \text{tomorrow}$  is

$$(9) \quad W_{\text{tomorrow}}^3 = \{2, 3\}.$$

Similarly, in example  $Q^2$ , Figure 2.2's table shows that the information set of the situation  $j = \{2, 3\}$  is

$$(10) \quad W_{\{2,3\}}^2 = \{2, 3\}.$$

Thus a situation  $j$  may or may not equal its information set  $W_j$ . This distinction will play a role in Sections 5.2–5.5, beginning with equation (32).

Second,  $A_j$  (that is  $\pi_A(Q_j)$ ) is the action set of situation  $j$ . For reasons associated with Proposition 3.3,  $A_j$  can also be called the *feasible* action set at situation  $j$ .

<sup>11</sup>When speaking aloud, it may be helpful to read  $\pi_Y(Q^1)$  as “the  $Y$  of  $Q^1$ ” (abbreviation (8) shortens this to  $Y^1$ ). Similarly, it may be helpful to read  $\pi_{JI}(Q^1)$  as “the  $JI$  of  $Q^1$ ”.

## 3.4. PENTAFORMS

**Definition 3.1 (Pentaform).** A pentaform is a (possibly infinite) set  $Q$  of quintuples  $\langle i, j, w, a, y \rangle$  such that

$$\begin{array}{ll}
[\text{Pi} \leftarrow j]^{12} & \pi_{JI}(Q) \text{ is a function,}^{13} \\
[\text{Pj} \leftarrow w] & \pi_{WJ}(Q) \text{ is a function,} \\
[\text{Pwa}] & (\forall j \in J) \pi_{WA}(Q_j) \text{ is a Cartesian product,} \\
[\text{Pwa} \rightarrow y] & \{ \langle \langle w, a \rangle, y \rangle \mid \langle w, a, y \rangle \in \pi_{WAY}(Q) \} \text{ is a function,}^{14} \\
[\text{Pw} \leftarrow y] & \pi_{YW}(Q) \text{ is a function,} \\
& \text{denoted by } p \text{ and called the immediate-predecessor function,}^{15} \\
[\text{Pa} \leftarrow y] & \pi_{YA}(Q) \text{ is a function,} \\
[\text{Py}] & (\forall y \in Y) (\exists m \geq 1) p^m(y) \notin Y, \text{ and} \\
[\text{Pr}] & W \setminus Y \text{ is a singleton}
\end{array}$$

(where (2) defines each  $Q_j$  and where (8) defines  $J$ ,  $W$ , and  $Y$ ).

The remainder of this Section 3.4 will discuss these eight axioms individually. En route, Section 2's examples will be reconsidered. It will be found that examples  $Q^1$ ,  $Q^2$ , and  $Q^3$  satisfy the axioms, and are therefore pentaforms. (An example of an infinite pentaform can be found in Streufert 2023, Section 2.2.)

*Axiom [Pi ← j].* This states that exactly one player  $i$  is assigned to each situation  $j$ . This is interpreted to mean that exactly one player controls the action at each situation. To be clear, [Pi ← j] states that  $\pi_{JI}(Q)$  is a function, which (by footnote 13) means that each  $j \in J$  is associated with exactly one  $i \in I$ . For example, (5) shows  $\pi_{JI}(Q^1) = \{ \langle \{0\}, \text{Alex} \rangle \}$ , and this is a function which maps  $\{0\}$  to Alex (here the only situation  $j$  is the information set  $\{0\}$ ). Similarly, examples  $Q^2$ ,  $Q^3$ , and  $Q^4$  satisfy axiom [Pi ← j].

*Axiom [Pj ← w].* This states that exactly one situation  $j$  is assigned to each decision node  $w$ . As Proposition 3.2(a ⇔ b) makes clear, this is equivalent to stating that distinct situations  $j_1$  and  $j_2$  have disjoint information sets  $W_{j_1}$  and  $W_{j_2}$ . For instance, the

<sup>12</sup>The label [Pi ← j] can be read as “ $i$  is a function of  $j$ ”. The labels [Pj ← w], [Pw ← y], and [Pa ← y] can be read similarly. Meanwhile, the label [Pwa → y] can be read as “ $w$  and  $a$  determine  $y$ ”. The arrows within the labels are visual crutches in the sense that they can be removed without introducing ambiguity.

<sup>13</sup>As in set theory (Halmos 1974, page 30; Enderton 1977, page 42), this paper identifies a function with its graph. For readers unfamiliar with set theory, [1] a *function*  $f$  is a set of couples such that  $(\forall x, y^A, y^B) \langle x, y^A \rangle \in f$  and  $\langle x, y^B \rangle \in f$  imply  $y^A = y^B$ , [2] the *domain* of a function  $f$  is  $\pi_1 f$ , and its *range* is  $\pi_2 f$ , [3] a *surjection from  $X$  to  $Y$*  is a function with domain  $X$  and range  $Y$ , [4] a *bijection from  $X$  to  $Y$*  is an injective surjection from  $X$  to  $Y$ , and [5] “ $f: X \rightarrow Z$ ” means  $f$  is a function with domain  $X$  and range included within  $Z$ .

<sup>14</sup>The set in axiom [Pwa → y] consists of couples of the form  $\langle \langle w, a \rangle, y \rangle$  whose first element is a couple of the form  $\langle w, a \rangle$ . The axiom requires that the set of “outer” couples is a function in the sense of footnote 13. Casually this means that  $\pi_{WAY}(Q)$  is a function from its first two coordinates.

<sup>15</sup>The statements  $w = p(y)$ ,  $\langle y, w \rangle \in \pi_{YW}(Q)$ , and  $\langle w, y \rangle \in \pi_{WY}(Q)$  are equivalent.

| Pentaform $Q$                       |  | [3.4]      |
|-------------------------------------|--|------------|
| $Q$                                 | set of quintuples $\langle i, j, w, a, y \rangle$                      | [3.1, 3.2] |
| $I = \pi_I(Q)$                      | $\hookrightarrow$ set of players $i$                                   | [3.1, 3.3] |
| $J = \pi_J(Q)$                      | $\hookrightarrow$ set of situations $j$                                | [3.1, 3.3] |
| $W = \pi_W(Q)$                      | $\hookrightarrow$ set of decision nodes $w$                            | [3.1, 3.3] |
| $A = \pi_A(Q)$                      | $\hookrightarrow$ set of actions $a$                                   | [3.1, 3.3] |
| $Y = \pi_Y(Q)$                      | $\hookrightarrow$ set of successor nodes $y$                           | [3.1, 3.3] |
| $Q_j \subseteq Q$                   | $\hookrightarrow$ situation $j$ 's slice of $Q$                        | [3.2]      |
| $W_j = \pi_W(Q_j)$                  | $\hookrightarrow$ situation $j$ 's decision-node set (information set) | [3.3]      |
| $A_j = \pi_A(Q_j)$                  | $\hookrightarrow$ situation $j$ 's (feasible) action set               | [3.3]      |
| $p = \pi_{YW}(Q)$                   | $\hookrightarrow$ immediate-predecessor function                       | [3.4]      |
| $F = \pi_{WA}(Q)$                   | $\hookrightarrow$ feasibility correspondence                           | [3.4]      |
| $W \cup Y$                          | $\hookrightarrow$ set of nodes $x$                                     | [3.5]      |
| $\pi_{WY}(Q)$                       | $\hookrightarrow$ set of edges $\langle w, y \rangle$                  | [3.5]      |
| $\{r\} = W \setminus Y$             | $\hookrightarrow$ root node $r$  | [3.5]      |
| $\preceq$                           | $\hookrightarrow$ weak precedence order                                | [3.5, A.1] |
| $<$                                 | $\hookrightarrow$ strict precedence order                              | [3.5, A.1] |
| $Y \setminus W$                     | $\hookrightarrow$ set of end nodes $y$                                 | [3.5, A.1] |
| $\mathcal{Z}$                       | $\hookrightarrow$ collection of runs $Z$                               | [3.5, A.1] |
| Pentaform Game $(Q, u)$             |  | [3.6]      |
| $u = \langle u_i \rangle_{i \in I}$ | profile with utility function $u_i$ for each player $i$                | [3.6]      |

TABLE 3.1. A pentaform is implicitly accompanied by its derivatives ( $\hookrightarrow$ ). Definitions are in the sections in brackets [].

situation set in example  $Q^3$  is  $J^3 = \{\text{today, tonight, tomorrow}\}$ . Further, (3) implies that the information sets are  $W_{\text{today}}^3 = \{0\}$ ,  $W_{\text{tonight}}^3 = \{1\}$ , and  $W_{\text{tomorrow}}^3 = \{2, 3\}$  (the third equality already appeared in (9)). Since these are disjoint, Proposition 3.2(a $\Leftarrow$ b) implies  $Q^3$  satisfies [Pj $\leftarrow$ w]. Similarly, examples  $Q^1$ ,  $Q^2$ , and  $Q^4$  satisfy [Pj $\leftarrow$ w].

**Proposition 3.2.** *Let  $Q$  be a quintuple set. Then the following are equivalent.*

- (a)  $Q$  satisfies [Pj $\leftarrow$ w].
- (b)  $(\forall j_1 \in J, j_2 \in J) j_1 \neq j_2$  implies  $W_{j_1} \cap W_{j_2} = \emptyset$ .
- (c)  $\langle W_j \rangle_{j \in J}$  is an injectively indexed partition of  $W$ .<sup>16</sup> (Proof B.3 in Appendix B.)

*Axiom [Pwa].* This states that for each situation  $j$ , the set  $\pi_{WA}(Q_j)$  is a Cartesian product. For instance, in example  $Q^3$ , axiom [Pwa] is satisfied at  $j = \text{tomorrow}$  because equation (7) implies  $\pi_{WA}(Q_{\text{tomorrow}}^3) = \{2, 3\} \times \{e, f\}$ . Cartesian products can also be found at the other two situations in  $Q^3$  and at all situations in  $Q^1$ ,  $Q^2$ , and  $Q^4$ . Hence all four examples satisfy [Pwa].

<sup>16</sup>To be clear, Proposition 3.2(c) means that [1]  $\{\langle j, W_j \rangle | j \in J\}$  is an injection and [2]  $\{W_j | j \in J\}$  partitions  $W$ . (Note that [1] implies that  $\{\langle j, W_j \rangle | j \in J\}$  is a bijection from the situation set  $J$  to the information-set collection  $\{W_j | j \in J\}$ .)

To interpret [Pwa], define the correspondence<sup>17</sup>

$$(11) \quad F = \pi_{WA}(Q).$$

Call  $F$  the *feasibility* correspondence, and call  $F(w) \subseteq A$  the set of *feasible* actions at decision node  $w \in W$ . For example, Figure 3.0's table implies

$$F^3 = \pi_{WA}(Q^3) = \{ \langle 0, \mathbf{b} \rangle, \langle 0, \mathbf{c} \rangle, \langle 1, \mathbf{d} \rangle, \langle 1, \mathbf{g} \rangle, \langle 2, \mathbf{e} \rangle, \langle 2, \mathbf{f} \rangle, \langle 3, \mathbf{e} \rangle, \langle 3, \mathbf{f} \rangle \},$$

which implies  $F^3(0) = \{\mathbf{b}, \mathbf{c}\}$ ,  $F^3(1) = \{\mathbf{d}, \mathbf{g}\}$ , and  $F^3(2) = F^3(3) = \{\mathbf{e}, \mathbf{f}\}$ . Below, Proposition 3.3(a $\Leftrightarrow$ d) characterizes [Pwa] by the property that the feasible set  $F(w)$  is constant across the nodes  $w$  in an information set. For instance, in the example  $Q^3$ , equation (9) implies that nodes 2 and 3 share the information set  $W_{\text{tomorrow}}^3 = \{2, 3\}$ , and the second-previous sentence shows that their feasible sets  $F^3(2)$  and  $F^3(3)$  are both equal to  $\{\mathbf{e}, \mathbf{f}\}$ . [Further, in accord with condition (c) of Proposition 3.3, this common feasible set is  $A_{\text{tomorrow}}^3 = \{\mathbf{e}, \mathbf{f}\}$ . For verification see (6).]

**Proposition 3.3.** *Suppose  $Q$  satisfies [Pj $\leftarrow$ w]. Then the following are equivalent.*

- (a)  $Q$  satisfies [Pwa].
- (b)  $(\forall j \in J) \pi_{WA}(Q_j) = W_j \times A_j$ .
- (c)  $(\forall j \in J, w \in W_j) F(w) = A_j$ .
- (d)  $(\forall j \in J, w_1 \in W_j, w_2 \in W_j) F(w_1) = F(w_2)$ . (Proof B.5 in Appendix B.)

*Axiom [Pwa $\rightarrow$ y].* This axiom states that each decision-node/feasible-action couple  $\langle w, a \rangle \in \pi_{WA}(Q)$  determines a successor node  $y$ . (Intuitively, selecting a feasible action at a decision node has the effect of selecting one of the decision node's successors.) To be clear, note that  $\pi_{WA}(Q)$  is the domain of the function in [Pwa $\rightarrow$ y], and that  $\pi_{WA}(Q)$  can be accurately called the set of decision-node/feasible-action couples because definition (11) implies  $\langle w, a \rangle \in \pi_{WA}(Q)$  iff  $a \in F(w)$ . In a different direction, it is sufficient for [Pwa $\rightarrow$ y] that each decision-node/feasible-action couple appears in exactly one quintuple in  $Q$ . By inspection, this holds in examples  $Q^1$ ,  $Q^2$ ,  $Q^3$ , and  $Q^4$ .

*Axioms [Pw $\leftarrow$ y] and [Pa $\leftarrow$ y].* These axioms are similar. Axiom [Pw $\leftarrow$ y] states that exactly one decision node is associated with each successor node. Similarly, [Pa $\leftarrow$ y] states that exactly one action is associated with each successor node. It is sufficient for both [Pw $\leftarrow$ y] and [Pa $\leftarrow$ y] that each successor node  $y \in Y$  appears in exactly one quintuple in  $Q$ . By inspection, this holds in examples  $Q^1$ ,  $Q^2$ ,  $Q^3$ , and  $Q^4$ .

*Axiom [Py].* This can be understood in the terms of a difference equation (Luenberger 1979, page 14). In particular, consider the difference equation  $y_{k-1} = p(y_k)$ , where the index  $k \in \{0, -1, -2, \dots\}$  runs backwards, and where  $p = \pi_{YW}(Q)$  is the immediate-predecessor function defined in axiom [Pw $\leftarrow$ y] of Definition 3.1. In this context, axiom [Py] states that the backward walk starting from any  $y_0 \in Y$  eventually leaves  $Y$ . In casual terms, the set  $Y$  is a "set-source". For instance, consider example  $Q^4$ . There definition (1) implies that  $Y^4 = \{45, 50\}$  and that  $p^4 = \pi_{YW}(Q^4) = \{ \langle 45, 43 \rangle, \langle 50, 48 \rangle \}$ . Hence  $Y^4$  is a "set-source" since  $(p^4)(45) = 43 \notin Y^4$  and since

<sup>17</sup>In this paper, a correspondence is simply a set of couples. Occasionally, the expression " $F: X \Rightarrow Z$ " is used to mean " $F$  is a correspondence such that  $\pi_1 F = X$  and  $\pi_2 F \subseteq Z$ ". (This paper does not apply the terms "domain" and "range" to correspondences.)

$(p^4)(50) = 48 \notin Y^4$ . The examples  $Q^1$ ,  $Q^2$ , and  $Q^3$  also satisfy [Py], though exhaustively showing so takes longer than for  $Q^4$  because there can be more successor nodes in  $Y$  and because the backward walks exiting  $Y$  can be longer.

*Axiom [Pr].* This states that there is exactly one decision node which is not also a successor node. For instance, consider example  $Q^3$ . There Figure 3.0's table implies that  $W^3 = \{0,1,2,3\}$  and that  $Y^3 = \{1,2,3,4,5,6,7,8\}$ , and these observations imply that  $W^3 \setminus Y^3 = \{0\}$ . Thus  $Q^3$  satisfies [Pr]. Similarly examples  $Q^1$  and  $Q^2$  satisfy [Pr]. Meanwhile, example  $Q^4$  violates [Pr] because definition (1) implies  $W^4 = \{43, 48\}$ ,  $Y^4 = \{45, 50\}$ , and  $W^4 \setminus Y^4 = \{43, 48\}$ .

*Summary.* In light of the preceding paragraphs, examples  $Q^1$ ,  $Q^2$ , and  $Q^3$  satisfy all eight axioms, and are consequently pentaforms. Meanwhile, example  $Q^4$  satisfies all axioms except [Pr], and is consequently not a pentaform.<sup>18</sup> Note that  $Q^4$  proves it is possible to violate [Pr] while satisfying the other seven axioms. In fact, Table B.1 in Appendix B shows that it is essentially possible to violate any one of the eight axioms while satisfying the other seven. In this sense the eight axioms are logically independent.

To verbally summarize the definition of a pentaform, recall from Section 3.1 that the terms “player”, “situation”, “decision node”, “action”, and “successor node” are defined to mean nothing but the five positions in a quintuple. Then a pentaform is a set of quintuples which satisfies the following five properties. (1) Exactly one player is assigned to each situation. (2) Exactly one situation is assigned to each decision node. (3) The set of actions assigned to a decision node is constant across the decision nodes assigned to each situation. (4) The assignment of a decision-node/action couple to a successor node is a bijection. (5) The assignment of decision nodes to successor nodes eventually takes every successor node to the unique decision node that is not a successor node. For details, note (1) paraphrases [Pi $\leftarrow$ j], (2) paraphrases [Pj $\leftarrow$ w], (3) paraphrases [Pwa] with the help of [Pj $\leftarrow$ w] and Proposition 3.3(a $\Leftrightarrow$ d), (4) paraphrases [Pwa $\rightarrow$ y], [Pw $\leftarrow$ y], and [Pa $\leftarrow$ y], and finally, (5) is based on [Pw $\leftarrow$ y] and paraphrases [Py] and [Pr].

### 3.5. THE OUT-TREE IMPLIED BY A PENTAFORM

An out-tree is a couple  $(X, E)$  consisting of a node set  $X$  and an edge set  $E$  which together satisfy the usual properties of a discrete game tree. The relevant definitions from graph theory are reviewed in the first paragraph of Section A.1 in Appendix A. The term “out-tree” is being used instead of “directed rooted tree” or “divergent arborescence” only because it is the shortest of these three synonymous terms.

**Proposition 3.4.** *Suppose  $Q$  is a quintuple set. Then (a)  $Q$  satisfies [Pw $\leftarrow$ y], [Py], and [Pr] iff  $(W \cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree. Further (b) if  $Q$  satisfies [Pw $\leftarrow$ y], [Py], and [Pr], then the unique element of  $W \setminus Y$  is the root of the out-tree  $(W \cup Y, \pi_{WY}(Q))$ . (Proof B.8.)*

<sup>18</sup>Here are some other examples: [1] the empty set is not a pentaform because it violates [Pr], [2] a singleton set  $\{\langle i, j, w, a, y \rangle\}$  with  $w \neq y$  is a pentaform, and [3] a singleton set  $\{\langle i, j, w, a, y \rangle\}$  with  $w = y$  is not a pentaform because it violates [Pr] (and also [Py]).

Consider a pentaform  $Q$ , or more narrowly, a quintuple set  $Q$  which satisfies  $[Pw\leftarrow y]$ ,  $[Py]$ , and  $[Pr]$ . Proposition 3.4 implies that  $(W\cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree, and that the unique element of  $W\setminus Y$  is the root node of this out-tree. Correspondingly, define  $Q$ 's *root node*  $r$  by<sup>19</sup>

$$(12) \quad \{r\} = W\setminus Y.$$

The remainder of this paragraph derives some familiar entities from  $Q$ 's out-tree  $(W\cup Y, \pi_{WY}(Q))$ . First, derive  $Q$ 's *weak precedence* relation  $\preceq$  and *strict precedence* relation  $\prec$  from the out-tree  $(W\cup Y, \pi_{WY}(Q))$  via (37) in Section A.1. Second, let  $Q$ 's set of *end nodes* be

$$(13) \quad Y\setminus W$$

(the same can be derived from  $(W\cup Y, \pi_{WY}(Q))$  via (38) in Section A.1). Third, derive  $Q$ 's *run* (or *play*) collection  $\mathcal{Z}$  from the out-tree  $(W\cup Y, \pi_{WY}(Q))$  via (39) in Section A.1.

### 3.6. PENTAFORM GAMES

Suppose  $Q$  is a pentaform with its player set  $I$  (from abbreviation (8)) and its run collection  $\mathcal{Z}$  (from the previous paragraph). A *utility function* for player  $i$  is a function of the form  $u_i: \mathcal{Z} \rightarrow \bar{\mathbb{R}}$ , where  $\bar{\mathbb{R}}$  is the extended real line  $\mathbb{R} \cup \{-\infty, \infty\}$ .<sup>20</sup> A *utility-function profile* is a  $u = \langle u_i \rangle_{i \in I}$  which lists a utility function  $u_i$  for each player  $i \in I$ . The following definition defines a pentaform game to be a couple listing a pentaform  $Q$  and a utility-function profile  $u$ . Figures 3.1 and 3.2 provide two relatively simple examples (Streufert 2023, Section 2.2 provides an infinite-horizon example).

**Definition 3.5 (Pentaform Game).** *A pentaform game is a couple  $(Q, u)$  such that  $Q$  is a pentaform (Definition 3.1) and  $u$  is of the form  $\langle u_i: \mathcal{Z} \rightarrow \bar{\mathbb{R}} \rangle_{i \in I}$  (where  $Q$  determines  $I$  and  $\mathcal{Z}$ , as summarized in Table 3.1).*

A less general formulation would be to augment each quintuple in a pentaform with a profile that specifies each player's reward for reaching the quintuple's successor node  $y$  (that is, for transiting the quintuple's edge  $\langle w, y \rangle$ ). Then a player's utility from a play  $Z$  could be calculated by summing their rewards from reaching the successor nodes in the play. In this formulation, the entire game is a set of tuples. This formulation is less general than Definition 3.5 when there is an infinite horizon (an infinite sum of rewards might be ill-defined, and further, there are infinite-horizon utility functions which cannot be expressed as sums).

<sup>19</sup>In another paper, one might define a special symbol (like “ $X$ ”) for  $Q$ 's node set  $W\cup Y$ . In this paper it seems better to avoid the additional notation.

<sup>20</sup>Infinite utility numbers are included because, in economics, many popular utility functions generate  $-\infty$  utility when some consumption level is zero. Such a utility function is often part of a consumer dynamic optimization problem, and such a problem can be specified as a one-player game.




FIGURE 3.1. The pentaform game  $(Q^1, u^1)$ . The set  $Q^1$  is from Figure 2.1, and consists of the upper table's rows (expressed as quintuples). The utility function  $u^1_{\text{Alex}}: \mathcal{Z}^1 \rightarrow \bar{\mathbb{R}}$  is defined by the lower table. The tree diagram provides the same data.




FIGURE 3.2. The pentaform game  $(Q^2, u^2)$ . The set  $Q^2$  is from Figure 2.2, and consists of the upper table's rows. The utility-function profile  $u^2 = \langle u^2_i: \mathcal{Z}^2 \rightarrow \bar{\mathbb{R}} \rangle_{i \in I^2}$  is defined by the lower table. The tree diagram provides the same data.

#### 4. SOME PENTAFORM TOOLS AND APPLICATIONS

Sections 4.1 and 4.2 develop tools for subsets and unions, and show their application to Selten subgames, dynamic programming, and pentaform construction. Separately, Section 4.3 applies pentaforms to perfect-recall and no-absentmindedness. (Section 5 will not depend on this Section 4.)

##### 4.1. SUBSETS OF PENTAFORMS

Part (a) of Proposition 4.1 shows that any subset of a pentaform satisfies six of the eight pentaform axioms. This and the definition of pentaform immediately imply part (b). Further, the proof of part (a) is intuitive. First consider the five axioms  $[P_i \leftarrow j]$ ,  $[P_j \leftarrow w]$ ,  $[P_{w \leftarrow y}]$ ,  $[P_{a \leftarrow y}]$ , and  $[P_{w \leftarrow y}]$ . Each states that some multidimensional projection of  $Q$  is a function, and any subset of a function is also a function (by the definition of function in footnote 13). So if one of these five axioms holds for  $Q$ , it almost obviously holds for any  $Q' \subseteq Q$ . Meanwhile, axiom  $[P_y]$  means that  $Y$  is a

“set-source” for the difference equation defined by  $p$ , as discussed in Section 3.4. If this holds for  $Y$ , it must also hold for  $Y' \subseteq Y$ . Thus it is intuitive that  $[Py]$  for  $Q$  implies  $[Py]$  for any  $Q' \subseteq Q$  (the proposition’s proof addresses some details about  $p'$ ).

**Proposition 4.1.** *Suppose  $Q$  is a pentaform and  $Q' \subseteq Q$ . Then (a)  $Q'$  satisfies  $[Pi^*j]$ ,  $[Pj^*w]$ ,  $[Pwa \rightarrow y]$ ,  $[Pa^*y]$ ,  $[Pw^*y]$ , and  $[Py]$ . Thus (b)  $Q'$  is a pentaform iff it satisfies  $[Pwa]$  and  $[Pr]$ . (Proof C.1.)*

As discussed in Section 3.2, each quintuple set  $Q$  is partitioned by the collection  $\{Q_j | j \in J\}$  of its slices  $Q_j$ . The following corollary states that the union of a subcollection of a pentaform’s slice partition satisfies the first seven pentaform axioms. This follows from Proposition 4.1(a) if it can be shown that such a union of slices satisfies  $[Pwa]$ . This holds because  $Q$  satisfies  $[Pwa]$  by assumption and because  $[Pwa]$  is defined in terms of individual slices.

**Corollary 4.2.** *Suppose  $Q$  is a pentaform and  $J' \subseteq J$ . Then (a)  $\cup\{Q_j | j \in J'\}$  satisfies  $[Pi^*j]$ ,  $[Pj^*w]$ ,  $[Pwa]$ ,  $[Pwa \rightarrow y]$ ,  $[Pa^*y]$ ,  $[Pw^*y]$ , and  $[Py]$ . Hence (b)  $\cup\{Q_j | j \in J'\}$  is a pentaform iff it satisfies  $[Pr]$ . (Proof above.)*

This remainder of this Section 4.1 uses Corollary 4.2 to construct the pentaforms for the Selten subgames of an arbitrary pentaform game. It also explains how the same Corollary 4.2 leads to Streufert 2023’s new results on dynamic programming.

Consider a pentaform  $Q$ . Then for any  $w \in W$ , define

$$(14) \quad {}^wQ = \{ \langle i_*, j_*, w_*, a_*, y_* \rangle \in Q \mid w \preceq w_* \}.$$

To put this in other words, say that a quintuple is *weakly after*  $w$  iff its decision node weakly succeeds  $w$ . Then  ${}^wQ$  is the set of quintuples that are weakly after  $w$ . A (*Selten*) *subroot* is a member of

$$(15) \quad T = \{ t \in W \mid {}^tJ \text{ and } \pi_J(Q \setminus {}^tQ) \text{ are disjoint} \},$$

where  ${}^tJ$  abbreviates  $\pi_J({}^tQ)$  by the sentence following (8). In other words, a decision node  $t \in W$  is a subroot iff each situation listed in a quintuple weakly after  $t$  is not listed in a quintuple anywhere else. Lemma C.3 shows this is equivalent to  ${}^tQ$  being the union of a subcollection of  $Q$ ’s slice partition.

Because of this, Corollary 4.2(b) can be applied to each  ${}^tQ$ . The result is the following proposition, which shows that the  ${}^tQ$  associated with each Selten subroot  $t \in T$  is a pentaform. As a consequence, the pentaform  ${}^tQ$  can serve as the extensive form of the Selten subgame starting at  $t$  (Streufert 2023, Section 4.2, completes the subgame by defining its utility functions).

**Proposition 4.3.** *Suppose  $Q$  is a pentaform and  $t \in T$ . Then  ${}^tQ$  is a pentaform. (Proof C.4.)*

Corollary 4.2 also plays a central role in the dynamic-programming theory of Streufert 2023, which is the first paper to use value functions to characterize subgame perfection in arbitrary games (only pure strategies are considered there). For an analogy, consider a repeated game. There, the whole-game extensive form (that is, the supergame’s extensive form) combines many replicas of a single-stage extensive



form (which is typically very simple). In this paper’s terminology, the “subroots” of the whole game are the nodes starting the replicas of the stage form. Now imagine that different subroots have different stage forms. In Streufert 2023’s terminology, these generalized stage forms are called “piece forms”. Each of the various piece forms has some combination of [a] finite runs, each of which terminates in a subsequent subroot or whole-form endnode, and [b] infinite runs, each of which fails to reach a subsequent subroot or whole-form endnode.

Streufert 2023 specifies an arbitrary pentaform game and shows [a] that the piece-form collection partitions the pentaform and [b] that this piece-form partition is coarser than the pentaform’s slice partition. Thus it can use Corollary 4.2(b) to show that each piece form is a pentaform. On this foundation the paper is able to build a notion of “piecewise Nashness” which generalizes dynamic-programming’s Bellman equation to arbitrary games.

#### 4.2. UNIONS OF BLOCKS

This Section 4.2 shows how to construct pentaforms as unions of “blocks”.<sup>21</sup> For the purposes of this section, let a *(penta)block* be a quintuple set  $Q$  satisfying the first seven pentaform axioms, namely

$$(16) \quad [Pi\leftarrow j], [Pj\leftarrow w], [Pwa], [Pwa\rightarrow y], [Pw\leftarrow y], [Pa\leftarrow y], \text{ and } [Py].$$

Thus a pentaform is equivalent to a block which satisfies the final pentaform axiom [Pr]. To put this in other words, consider an arbitrary quintuple set  $Q$  and call  $W \setminus Y$  the set of  $Q$ ’s *start nodes*. Then [Pr] is equivalent to  $Q$  having exactly one start node. Hence a pentaform is equivalent to a block with exactly one start node.

In this section’s terminology, Corollary 4.2(a) shows that if  $J' \subseteq J$  is a subset of a pentaform  $Q$ ’s situation set, then  $\cup\{Q_j | j \in J'\}$  is a block. This easily implies that each slice  $Q_j$  of a pentaform  $Q$  is a block. For example, reconsider Figure 3.0, which illustrates that  $\{Q_{\text{today}}^3, Q_{\text{tonight}}^3, Q_{\text{tomorrow}}^3\}$  is the slice partition of  $Q^3$ . Since  $Q^3$  is a pentaform, each of these three slices is a block.

Now consider an arbitrary collection  $\mathcal{Q}$  of quintuple sets  $Q$ . Then  $\mathcal{Q}$  is said to be *weakly separated* iff its member sets do not share situations,<sup>22</sup> decision nodes, or successor nodes. In other words,  $\mathcal{Q}$  is weakly separated iff, for all distinct  $Q^A$  and  $Q^B$  in  $\mathcal{Q}$ ,

$$(17) \quad J^A \cap J^B = \emptyset, W^A \cap W^B = \emptyset, \text{ and } Y^A \cap Y^B = \emptyset.$$

<sup>21</sup>This Section 4.2 is related to the ongoing work of Ghani, Kupke, Lambert, and Nordvall Forsberg 2018; Bolt, Hedges, and Zahn 2023; and Capucci, Ghani, Ledent, and Nordvall Forsberg 2022. Both that literature and this Section 4.2 seek to systematically construct games out of game fragments. A precise comparison is elusive because the mathematical foundations are very different. More is said there about utility. The relative advantages here include constructing general infinite-horizon games, using the relatively simple operation of union, and using relatively finely-grained axioms.

<sup>22</sup>Incidentally, suppose  $\mathcal{Q}$ ’s member sets have information-set situations (32). Then, in (17) and (18), the condition  $J^A \cap J^B = \emptyset$  is redundant. Specifically, the disjointness of  $W^A$  and  $W^B$  implies the disjointness of  $\mathcal{P}(W^A)$  and  $\mathcal{P}(W^B)$ , which implies the disjointness of the information-set collections  $\{W_j^A | j \in J^A\} \subseteq \mathcal{P}(W^A)$  and  $\{W_j^B | j \in J^B\} \subseteq \mathcal{P}(W^B)$ , which by (32) implies the disjointness of  $\{j | j \in J^A\}$  and  $\{j | j \in J^B\}$ , which by inspection implies the disjointness of  $J^A$  and  $J^B$ .

Further,  $\mathcal{Q}$  is said to be *strongly separated* iff its member sets do not share situations or nodes. In other words,  $\mathcal{Q}$  is strongly separated iff, for all distinct  $Q^A$  and  $Q^B$  in  $\mathcal{Q}$ ,

$$(18) \quad J^A \cap J^B = \emptyset \text{ and } (W^A \cup Y^A) \cap (W^B \cup Y^B) = \emptyset.$$

By inspection, (18) implies (17). Intuitively, not sharing nodes in general implies not sharing decision nodes in particular and also not sharing successor nodes in particular.

**Proposition 4.4.** (a) *Suppose  $\{Q^A, Q^B\}$  is a weakly separated collection of blocks such that the start-node set  $W^A \setminus Y^A$  is disjoint from the end-node set  $Y^B \setminus W^B$ . Then  $Q^A \cup Q^B$  is a block whose start-node set is the union of*

$$W^A \setminus Y^A \text{ and } (W^B \setminus Y^B) \setminus (Y^A \setminus W^A)$$

*and whose end-node set is the union of*

$$(Y^A \setminus W^A) \setminus (W^B \setminus Y^B) \text{ and } Y^B \setminus W^B.$$

(b) *Suppose  $\mathcal{Q}$  is a strongly separated collection of blocks. Then  $\cup \mathcal{Q}$  is a block with start-node set  $\cup_{Q \in \mathcal{Q}} (\pi_W(Q) \setminus \pi_Y(Q))$  and end-node set  $\cup_{Q \in \mathcal{Q}} (\pi_Y(Q) \setminus \pi_W(Q))$ . (Proofs C.6 and C.7.)<sup>23</sup>*

To explore this proposition, consider three quintuple sets: the example  $Q^3$  from Figure 3.0;

$$(19) \quad Q^{\text{guilty}} = \{ \langle \text{Kid, guilty, 4, s, 11} \rangle, \langle \text{Kid, guilty, 4, \tilde{s}, 12} \rangle, \\ \langle \text{Kid, guilty, 5, s, 13} \rangle, \langle \text{Kid, guilty, 5, \tilde{s}, 14} \rangle \}; \text{ and}$$

$$(20) \quad Q^{\text{innocent}} = \{ \langle \text{Kid, innocent, 6, s, 15} \rangle, \langle \text{Kid, innocent, 6, \tilde{s}, 16} \rangle, \\ \langle \text{Kid, innocent, 7, s, 17} \rangle, \langle \text{Kid, innocent, 7, \tilde{s}, 18} \rangle \}.$$
<sup>24</sup>

These three quintuple sets are illustrated in Figure 4.1. All three are blocks. In particular, (i)  $Q^3$  is a block because it is a pentaform and because every pentaform is a block by the block definition (16), (ii)  $Q^{\text{guilty}}$  is a block because it is like the slice  $Q^3_{\text{tomorrow}}$  from the pentaform  $Q^3$  and because every slice of every pentaform is a block by Corollary 4.2(a), and (iii)  $Q^{\text{innocent}}$  is a block because it also is like the slice  $Q^3_{\text{tomorrow}}$ . Further, the three blocks are weakly separated (17). To see this, note that the three blocks do not share situations, do not share decision nodes, and do not share successor nodes.

Proposition 4.4(a) assumes that  $Q^A$  and  $Q^B$  are weakly separated blocks and that no start node of  $Q^A$  is also an end node of  $Q^B$ . This admits the possibility that an

<sup>23</sup>Incidentally, Proposition 4.4 is proved via Lemma C.5, which shows that the union of an arbitrary weakly separated collection of blocks satisfies all axioms except [Py] and [Pr]. This relatively unstructured lemma appears to have independent value as an alternative to Proposition 4.4.

<sup>24</sup> $Q^{\text{guilty}}$  and  $Q^{\text{innocent}}$  can be interpreted as continuations of the story in footnote 7. In  $Q^{\text{guilty}}$ , the kid did not do her homework, the teacher has not yet reported a verdict, and the kid must decide whether to say something to influence her parents' reaction to the coming verdict (s denotes the action of saying something, and  $\tilde{s}$  denotes the opposite). Similarly, in  $Q^{\text{innocent}}$ , the dog ate the kid's homework, the teacher has not yet reported a verdict, and the kid must decide whether to say something to influence her parents' reaction.




FIGURE 4.1. Three blocks:  $Q^3$  from Figure 3.0,  $Q^{\text{guilty}}$  from equation (19), and  $Q^{\text{innocent}}$  from equation (20).

end node of  $Q^A$  is also a start node of  $Q^B$ . Roughly,  $Q^A$  can precede  $Q^B$ , but not vice versa. For example, consider  $(Q^A, Q^B) = (Q^3, Q^{\text{guilty}})$  and ignore  $Q^{\text{innocent}}$ . The only start node of  $Q^A = Q^3$  is 0, and this is not in  $\{11, 12, 13, 14\}$ , which is the end-node set of  $Q^B = Q^{\text{guilty}}$ . Thus Proposition 4.4(a) implies that  $Q^3 \cup Q^{\text{guilty}}$  is a block whose start nodes are

(21a) the start nodes of  $Q^A = Q^3$  together with

(21b) the start nodes of  $Q^B = Q^{\text{guilty}}$  that are not also end nodes of  $Q^A = Q^3$ .

In (21b), the start nodes of  $Q^{\text{guilty}}$  are 4 and 5, which are also end nodes of  $Q^3$ . Thus (21b) contributes nothing. Meanwhile, in (21a), the only start node of  $Q^3$  is 0. Therefore the only start node of  $Q^3 \cup Q^{\text{guilty}}$  is 0, which obviously implies that  $Q^3 \cup Q^{\text{guilty}}$  has exactly one start node, which by this Section 4.2's first paragraph implies that  $Q^3 \cup Q^{\text{guilty}}$  is a pentaform.

In a similar fashion, Proposition 4.4(a) can be applied at  $(Q^A, Q^B) = (Q^3, Q^{\text{innocent}})$  to show that  $Q^3 \cup Q^{\text{innocent}}$  is a pentaform. Further, using this technique twice shows that  $Q^3 \cup Q^{\text{guilty}} \cup Q^{\text{innocent}}$  is a pentaform. This can be accomplished by constructing the union as  $(Q^3 \cup Q^{\text{guilty}}) \cup Q^{\text{innocent}}$  or as  $(Q^3 \cup Q^{\text{innocent}}) \cup Q^{\text{guilty}}$ .

Proposition 4.4(b) provides another way to prove that  $Q^3 \cup Q^{\text{guilty}} \cup Q^{\text{innocent}}$  is a pentaform. Since  $Q^{\text{guilty}}$  and  $Q^{\text{innocent}}$  do not share nodes,  $\{Q^{\text{guilty}}, Q^{\text{innocent}}\}$  is strongly separated (18). Thus Proposition 4.4(b) implies that  $Q^{\text{guilty}} \cup Q^{\text{innocent}}$  is a block whose start-node set is

$$\bigcup_{Q \in \{Q^{\text{guilty}}, Q^{\text{innocent}}\}} (\pi_W(Q) \setminus \pi_Y(Q)) = \{4, 5\} \cup \{6, 7\}$$

and whose end-node set is

$$\bigcup_{Q \in \{Q^{\text{guilty}}, Q^{\text{innocent}}\}} (\pi_Y(Q) \setminus \pi_W(Q)) = \{11, 12, 13, 14\} \cup \{15, 16, 17, 18\}.$$

Then, as in the previous two paragraphs, Proposition 4.4(a) can be applied at  $(Q^A, Q^B) = (Q^3, Q^{\text{guilty}} \cup Q^{\text{innocent}})$  to show that  $Q^3 \cup (Q^{\text{guilty}} \cup Q^{\text{innocent}})$  is a block with exactly one start node. Thus the union is a pentaform.

Roughly, the previous paragraph showed how to augment the pentaform  $Q^3$  with a “layer”  $\mathcal{Q} = \{Q^{\text{guilty}}, Q^{\text{innocent}}\}$  of two additional blocks whose start nodes were among the end nodes of  $Q^3$ . This same technique can be used to augment any finite-horizon pentaform with any layer  $\mathcal{Q}$  of additional blocks whose start nodes are among the end nodes of the original pentaform. The layer’s blocks do not need to be similar to one another. Also, the layer can have arbitrarily many (and possibly uncountably many) blocks, which freely allows arbitrarily many end nodes of the original pentaform to be connected with successor nodes in the layer’s additional blocks.

Further, layer after layer can be added to generate an infinite expanding sequence of pentaforms. Proposition 4.5 shows that the union of such an expanding sequence is a pentaform. This provides a straightforward way to construct an infinite-horizon pentaform. For instance, Streufert 2023’s motivating example is a partially infinitely repeated game of cry-wolf. Its pentaform is built via Propositions 4.4 and 4.5 (details in Streufert 2023, Lemma A.6).

**Proposition 4.5.** *Suppose  $\langle Q^n \rangle_{n \geq 0}$  is an infinite sequence of pentaforms such that  $(\forall n \geq 1) Q^{n-1} \subseteq Q^n$  and  $r^n = r^0$ . Then  $\bigcup_{n \geq 0} Q^n$  is a pentaform with root  $r^0$ . (Proof C.8.)*

### 4.3. PERFECT-RECALL IN TERMS OF PENTAFORMS

Although perfect-recall will remain a relatively subtle concept, it seems helpful to re-express the concept in terms of pentaforms. This section does so, and is unconnected with Sections 4.1 and 4.2.

The concept of perfect-recall was introduced in two different but equivalent ways by Kuhn 1950, page 575, and Kuhn 1953, page 213. Later it was characterized in other ways by Okada 1987, page 87, Ritzberger 1999, Theorems 1 and 2, and Alós-Ferrer and Ritzberger 2017, Theorems 1 and 2. Two formulations especially close to the approach here are Kreps and Wilson 1982, equation (2.3), and Myerson 1991, page 43. Essentially, suppose that two decision nodes  $w_1$  and  $w_2$  are in the same information set, and that  $w_1$  has a predecessor controlled by the same player. Then perfect-recall requires that  $w_2$  has a predecessor [a] which is in the same information set as  $w_1$ ’s predecessor and [b] whose path to  $w_2$  starts with the same action as  $w_1$ ’s predecessor’s path to  $w_1$ . Unfortunately, it is notationally awkward to express the first action on the way to  $w_1$  and likewise the first action on the way to  $w_2$ . This is one reason that many people regard perfect-recall as a relatively subtle concept.

Pentaforms allow an alternative formulation which revolves around successor nodes rather than decision nodes. To begin, consider a pentaform  $Q$ , and note that the axioms  $[\text{Pi} \leftarrow j]$ ,  $[\text{Pj} \leftarrow w]$ ,  $[\text{Pw} \leftarrow y]$ , and  $[\text{Pa} \leftarrow y]$  imply that each successor node  $y \in Y$  determines its entire quintuple  $\langle i, j, w, a, y \rangle \in Q$ . Thus we may speak of  $y$ ’s player  $i_y$ ,




FIGURE 4.2. The definition (22) of perfect-recall assumes the solid lines and then requires the dashed lines.

$y$ 's situation  $jy$ , and  $y$ 's action  $a_y$ .<sup>25</sup> For intuitive purposes, it is important to be reminded that  $y$  is a successor node. Toward that end, say that a successor node  $y$  is obtained “by” player  $i_y$ , that  $y$  is obtained “from” situation  $j_y$ , and that  $y$  is obtained “via” action  $a_y$ .

Then say that a pentaform  $Q$  satisfies *perfect-recall* iff, for all successor nodes  $y_0$ ,  $y_1$ , and  $y_2$  in  $Y$ , the conditions

$$(22a) \quad jy_1 = jy_2, \quad y_0 \prec y_1, \quad \text{and} \quad iy_0 = iy_1,$$

imply the existence<sup>26</sup> of a successor node  $y_{00} \in Y$  such that

$$(22b) \quad y_{00} \prec y_2 \quad \text{and} \quad \langle jy_{00}, ay_{00} \rangle = \langle jy_0, ay_0 \rangle.$$

In English, suppose that  $y_1$  and  $y_2$  are obtained from the same situation, and that  $y_1$  has a predecessor  $y_0$  which is obtained by the same player.<sup>27</sup> Then perfect-recall requires that  $y_2$  has a predecessor  $y_{00}$  which is obtained from the same situation,<sup>28</sup> and via the same action, as  $y_0$ . Note that the actions  $a_{y_0}$  and  $a_{y_{00}}$  are easy to express because the new formulation revolves around successor nodes rather than decision nodes. This is an advantage over the traditional formulation discussed earlier.

For an example, modify the pentaform  $Q^3$  from Figure 3.0 by replacing both the player *Kid* and the player *Teacher* with a new player named *KidTeacher*. Here the new formulation of perfect-recall fails at  $(y_0, y_1, y_2) = (2, 5, 7)$ . In particular, 5 and 7 are obtained from the same situation (*tomorrow*), and 5 has a predecessor 2 which is obtained by the same player (*KidTeacher*). Yet, 7 does not have a predecessor which is obtained from the same situation, and via the same action, as 2. Specifically, 7 is preceded by 3 and 1 (the root node 0 is excluded because it is not a successor node). [a] The predecessor 3 is obtained from *tonight* while 2 is obtained from *today*. [b] The predecessor 1 is obtained from *today* as 2 is obtained from *today*, but, 1 is obtained

<sup>25</sup>To be explicit, let  $a_y = \pi_{YA}(Q)(y)$ , where  $\pi_{YA}(Q)$  is a function by  $[Pa \prec y]$ . Then let  $j_y = \pi_{WJ}(Q)(p(y))$  where  $p = \pi_{YW}(Q)$  is a function by  $[Pw \prec y]$  and  $\pi_{WJ}(Q)$  is a function by  $[Pj \prec w]$ . Finally, let  $i_y = \pi_{JI}(Q)(j_y)$  where  $\pi_{JI}(Q)$  is a function by  $[Pi \prec j]$ .

<sup>26</sup>By inspection, (22b) holds with  $y_{00}$  equaling  $y_0$  iff  $y_0 \prec y_2$ . Thus perfect-recall becomes interesting precisely when  $y_0$  does not precede  $y_2$ .

<sup>27</sup>Equation (22a)'s first equality and  $[Pi \prec j]$  imply that  $y_1$  and  $y_2$  are obtained by the same player, which by (22a)'s second equality implies that  $y_0$ ,  $y_1$ , and  $y_2$  are all obtained by the same player.

<sup>28</sup>This and axiom  $[Pi \prec j]$  imply that  $y_{00}$  and  $y_0$  are obtained by the same player, which by footnote 27 implies that all four nodes are obtained by the same player.

via the action  $c$  while 2 is obtained via the action  $b$ . Intuitively, when KidTeacher is at situation **tomorrow** (from which they obtain 5 and 7), they remember the situation **today** (from which they obtain predecessors of 5 and 7) but not the action which they took there (since they obtain the predecessors of 5 and 7 via different actions).

Lastly, the remainder of this section suggests that the new version of perfect-recall can be easily manipulated. It does so by considering the relationship between perfect-recall and no-absentmindedness. The concept of no-absentmindedness is [a] left unnamed and imposed as part of the definition of a game by Kuhn 1953, page 195, [b] called “linearity” by Isbell 1957, page 86, [c] called “nonrepetition” by Alpern 1988, page 470, and [d] called “no-absentmindedness” by Piccione and Rubinstein 1997, pages 9–10, and Ritzberger 1999, page 72. The concept is relatively simple. It says that one decision node cannot strictly precede another decision node in the same information set. Correspondingly, say that a pentaform  $Q$  satisfies *no-absentmindedness* iff

$$(23) \quad (\nexists y_1 \in Y, y_2 \in Y) \ y_1 \prec y_2 \text{ and } jy_1 = jy_2.$$

This says that no successor node strictly precedes another successor node from the same situation.

The proof of the following proposition is new. The result is stated without recorded proof in Piccione and Rubinstein 1997, pages 3–5, and Hillas and Kvasov 2020, Section 2 (a self-evident result in Isbell 1953, page 86, uses a different definition of perfect-recall). Meanwhile, there is a recorded proof in Ritzberger 1999, pages 73–74, which differs from the proposition’s proof by using actions, nontrivial feasible sets, and strategies.

**Proposition 4.6.** *Perfect-recall (22) implies no-absentmindedness (23).*

*Proof.* To prove the contrapositive, suppose that  $Q$  is a pentaform which violates no-absentmindedness. Then definition (23) implies there are successor nodes  $y_0$  and  $y_1$  such that  $y_0 \prec y_1$  and  $jy_0 = jy_1$ . The equality and [Pi<j] imply  $iy_0 = iy_1$ . Next, let  $y_2$  be the earliest weak predecessor of  $y_0$  and  $y_1$  which is from the same situation as  $y_0$  and  $y_1$  (it is irrelevant that  $y_2$  might equal  $y_0$ ). Note that  $jy_2 = jy_1$ . Thus  $y_0, y_1$ , and  $y_2$  satisfy the hypotheses of perfect-recall (22a). Yet the definition of  $y_2$  implies that there is no  $y_{00}$  such that  $y_{00} \prec y_2$  and  $jy_{00} = jy_0$ . Hence the conclusion of perfect-recall (22b) cannot be satisfied.  $\square$

## 5. EQUIVALENCE WITH TRADITIONAL GAMES

This Section 5 compares pentaform games with traditional games. It is unconnected with Section 4.

### 5.1. DEFINITION OF TRADITIONAL GAMES

Definition 5.1 will define a “traditional game” to be a tree that has been adorned with information sets, actions, players, and utility functions in a more-or-less usual way. The phrase “more-or-less” acknowledges that there is no generally accepted way

to define an extensive-form game. Rather, substantially different definitions appear in social science, mathematics, computer science, logic, and engineering; and further, there is often substantial variety within any one of these fields. Somewhere in the middle is Definition 5.1, which might best be called the definition of a “representative” traditional game. In developing this definition, the author’s guiding principles were to head for the middle, to avoid specialized assumptions, to use explicit notation, and to use standard mathematics whenever possible. (In the context of Section 1.3’s literature review, Definition 5.1 is a generic extension of Kuhn 1953, and fairly close to the “KS” formulations in Kline and Luckraz 2016 and Streufert 2019.)

The following five paragraphs define the symbols and terms that appear in Definition 5.1. The paragraphs have been given letters to facilitate referencing, and the material is summarized in the left-hand side of Table 5.1. Because of the diversity in the literature, almost every reader will be unfamiliar with something, and a number of specific hazards are discussed in footnotes 29–34. (Incidentally, it can be helpful to notice, from a mathematical perspective, that paragraphs c–e are almost independent of one another. The only link is that the players from paragraph d index the utility functions in paragraph e.)

*Paragraph a.* The first component of a traditional game (Definition 5.1) is a non-trivial out-tree  $(X, E)$ .<sup>29</sup> By definition, such an out-tree consists of a node set  $X$  and an edge set  $E$  which together satisfy the usual properties of a discrete game tree (full details in the first paragraph of Section A.1 in Appendix A). The term “out-tree” is being used instead of “directed rooted tree” or “divergent arborescence” only because it is the shortest of these three synonymous terms. By usual constructions, an out-tree determines its root node  $r$  (first paragraph of Section A.1), its decision-node set  $\pi_1 E$ ,<sup>30</sup> its successor-node set  $\pi_2 E$ , its immediate-predecessor function  $E^{-1}$  (Lemma A.1(40b)), its weak and strict precedence orders  $\preceq$  and  $\prec$  (equation (37)), its end-node set  $\pi_2 E \setminus \pi_1 E$  (equation (38)), and its collection  $\mathcal{Z}$  of runs or equivalently “plays” (equation (39)).

*Paragraph b.* The next component of a traditional game is a partition  $\mathcal{H}$  of the decision-node set  $\pi_1 E$ . The members (i.e. cells) of this partition are called *information sets*. (Intuitively, a player will be informed that they are in an information set, but not that they are at a particular node in that set.)

---

<sup>29</sup>This first step may be unfamiliar for any of several reasons. First, game trees can be specified as undirected rooted trees rather than out-trees. Second, game trees can be specified via order theory rather than graph theory. Third, some popular specifications use specialized notations which implicitly impose some of the properties of game trees. For example, some game-tree properties are implicitly imposed if nodes are specified as sequences of past actions, or sets of past actions, or sets of outcomes. Finally, it is common to specify game trees by diagrams rather than by notation.

<sup>30</sup>The reader might expect a special symbol (like “ $W$ ”) for the decision-node set  $\pi_1 E$ , and another special symbol (like “ $p$ ”) for the immediate-predecessor function  $E^{-1}$ . In this paper, it seems better to avoid the additional notation.

| Out-tree $(X, E)$  |  | [5.1a, A.1] | All in<br>pentaform terms   |
|--|--|-------------|---|
| $X$  | set of nodes $x$                                 | [5.1a, A.1] | $\hat{W} \cup \hat{Y}$  |
| $E$  | set of edges $\langle w, y \rangle$              | [5.1a, A.1] | $\pi_{WY}(\hat{Q})$   |
| $r$  | $\hookrightarrow$ root node                      | [5.1a, A.1] | $\{\hat{r}\} = \hat{W} \setminus \hat{Y}$   |
| $\pi_1 E$  | $\hookrightarrow$ set of decision nodes $w$      | [5.1a]      | $\hat{W} = \pi_W(\hat{Q})$  |
| $\pi_2 E$  | $\hookrightarrow$ set of successor nodes $y$     | [5.1a]      | $\hat{Y} = \pi_Y(\hat{Q})$  |
| $E^{-1}$   | $\hookrightarrow$ immediate-predecessor function | [5.1a]      | $\hat{p} = \pi_{YW}(\hat{Q})$   |
| $\preceq$  | $\hookrightarrow$ weak precedence order          | [5.1a, A.1] | $\hat{\preceq}$   |
| $\prec$  | $\hookrightarrow$ strict precedence order        | [5.1a, A.1] | $\hat{\prec}$   |
| $\mathcal{Z}$  | $\hookrightarrow$ collection of runs $Z$         | [5.1a, A.1] | $\hat{\mathcal{Z}}$   |
| Traditional Game $(X, E, \mathcal{H}, \alpha, \iota, u)$ |  | [5.1]       |   |
| $\mathcal{H}$  | collection of information sets $H$               | [5.1b]      | $\hat{J} = \pi_J(\hat{Q})$  |
| $\alpha$   | action-assigning function                        | [5.1c]      | $\{\langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(\hat{Q})\}$ |
| $A$  | $\hookrightarrow$ set of actions $a$             | [5.1c]      | $\hat{A} = \pi_A(\hat{Q})$  |
| $F$  | $\hookrightarrow$ feasibility correspondence     | [5.1c]      | $\hat{F} = \pi_{WA}(\hat{Q})$   |
| $\iota$  | player-assigning function                        | [5.1d]      | $\pi_{WI}(\hat{Q})$   |
| $I$  | $\hookrightarrow$ set of players $i$             | [5.1d]      | $\hat{I} = \pi_I(\hat{Q})$  |
| $u = \langle u_i \rangle_{i \in I}$                      | profile with utility function $u_i$ for each $i$ | [5.1e]      | $\hat{u} = \langle \hat{u}_i \rangle_{i \in \hat{I}}$   |

TABLE 5.1. *Left-hand Side:* Out-trees and traditional games are implicitly accompanied by their derivatives ( $\hookrightarrow$ ). Definitions are in the sections in brackets []. *Right-hand Side:* The same entities in terms of the pentaform game  $P(X, E, \mathcal{H}, \alpha, \iota, u) = (\hat{Q}, \hat{u})$  (Proposition 5.5).

*Paragraph c.* Next, each edge of the tree is assigned an action. Formally, this assignment is accomplished by a function  $\alpha$  from the edge set  $E$ . Call  $\alpha$  the *action-assigning function* (or “labelling function”).<sup>31</sup> Then let  $A$  be the range of  $\alpha$ , that is, let

$$(24) \quad A = \{ \alpha(w, y) \mid \langle w, y \rangle \in E \},$$

and call an element of  $A$  an *action*. (Intuitively, each edge’s action is what the controlling player would “do” to choose the edge.) It will be assumed that  $\alpha$  is *locally*

<sup>31</sup>Action-assigning functions may be unfamiliar for any of several reasons. First, actions might be assigned to successor nodes  $y \in \pi_2 E$  rather than to edges  $\langle w, y \rangle \in E$ . The two are equivalent because there is a bijection between  $\pi_2 E$  and  $E$  (this bijection follows from Lemma A.1(40b)). Second, some popular specifications use specialized notations which implicitly assign actions to edges or successor nodes. For example, if nodes are specified as sequences of past actions, then the action assigned to a successor node is simply the node’s most recent past action. Third, it is common to assign actions to edges in a tree diagram rather than through formal notation.



*injective* in the sense that, for any two edges of the form  $\langle w, y_1 \rangle \in E$  and  $\langle w, y_2 \rangle \in E$ ,

$$(25) \quad y_1 \neq y_2 \text{ implies } \alpha(w, y_1) \neq \alpha(w, y_2).$$

Thus local injectivity means that two different edges from one decision node  $w$  cannot be assigned the same action. Further, from the action-assigning function  $\alpha$ , derive the correspondence  $F: \pi_1 E \rightrightarrows A$  by

$$(26) \quad (\forall w \in \pi_1 E) F(w) = \{ a \mid (\exists y) \alpha(w, y) = a \}.$$

Thus each  $F(w)$  is the set of actions that label the edges leaving  $w$ . Call  $F(w)$  the set of actions that are *feasible* at  $w$ . Then let  $\langle F(w) \rangle_{w \in \pi_1 E}: \pi_1 E \rightarrow \mathcal{P}(A)$  be the associated set-valued function. It will be assumed that  $\langle F(w) \rangle_{w \in \pi_1 E}$  is *measurable*<sup>32</sup> in the sense that it is measurable as a function from  $\pi_1 E$  (endowed with the  $\sigma$ -algebra whose elements are arbitrary unions of cells from the information-set partition  $\mathcal{H}$ ) into the collection of the subsets of  $A$  (endowed with the discrete  $\sigma$ -algebra). This is equivalent to

$$(27) \quad (\forall H \in \mathcal{H}, w_1 \in H, w_2 \in H) F(w_1) = F(w_2),$$

which requires that two nodes in one information set have the same feasible-action set.

*Paragraph d.* Next, each decision node of the tree is assigned to a player. Formally, this assignment is accomplished by a function  $\iota$  from the decision-node set  $\pi_1 E$ .<sup>33</sup> Call  $\iota$  the *player-assigning function*. Then let  $I$  be the range of  $\iota$ , that is, let

$$(28) \quad I = \{ \iota(w) \mid w \in \pi_1 E \},$$

and call an element of  $I$  a *player*. (Intuitively, the role of  $\iota$  is to specify which player controls the action at each decision node.) It will be assumed that  $\iota$  is *measurable* (footnote 32) in the sense that it is measurable as a function from  $\pi_1 E$  (endowed with the  $\sigma$ -algebra whose elements are arbitrary unions of cells from the information-set partition  $\mathcal{H}$ ) into the player set  $I$  (endowed with the discrete  $\sigma$ -algebra). This is equivalent to

$$(29) \quad (\forall H \in \mathcal{H}, w_1 \in H, w_2 \in H) \iota(w_1) = \iota(w_2),$$

which requires that two nodes in one information set are controlled by same player.

*Paragraph e.* Finally, each player is given a utility function defined over the runs of the tree. Formally, this is accomplished by a profile  $u = \langle u_i \rangle_{i \in I}$  which lists a function  $u_i: \mathcal{Z} \rightarrow \bar{\mathbb{R}}$  for each player  $i$ . Here  $\mathcal{Z}$  is the run collection (39) of the out-tree  $(X, E)$  as defined in paragraph a, and  $\bar{\mathbb{R}}$  is the set of extended real numbers  $\mathbb{R} \cup \{-\infty, \infty\}$  as discussed in Section 3.6. Call  $u_i$  the *utility function* (or “payoff function”) of player  $i$ .<sup>34</sup>

<sup>32</sup>The term “measurability” is new in this context. The paper does not use measurability theory in any substantial way. Rather, “measurability” is just a mathematically ordinary and precise name for conditions (27) and (29). (The same constructions are called “continuity” in Streufert 2021.)

<sup>33</sup>An alternative is to assign players to information sets rather than to decision nodes. This would implicitly impose the measurability of (29).

<sup>34</sup>This specification of utility may be slightly unfamiliar. First, it is common, in finite-horizon games, to map end nodes to utility numbers. In this special case, there is a bijection between the end-node set  $\pi_2 E \setminus \pi_1 E$  and the run collection  $\mathcal{Z}$  (here  $\mathcal{Z}$  equals the collection  $\mathcal{Z}_{\text{ft}}$  of finite runs

**Definition 5.1 (Traditional Game).** A traditional game is a tuple  $(X, E, \mathcal{H}, \alpha, \iota, u)$  such that

[T1]  $(X, E)$  is a nontrivial out-tree (Section A.1),

[T2]  $\mathcal{H}$  is a partition of  $\pi_1 E$ ,

[T3]  $\alpha$  is a locally injective (25) function from  $E$ ,

[T4]  $\langle F(w) \rangle_{w \in \pi_1 E}$  is measurable (27),

[T5]  $\iota$  is a measurable (29) function from  $\pi_1 E$ , and

[T6]  $u$  is of the form  $\langle u_i: \mathcal{Z} \rightarrow \bar{\mathbb{R}} \rangle_{i \in I}$

(where  $\alpha$  determines  $F$  (26),  $\iota$  determines  $I$  (28), and  $(X, E)$  determines  $\mathcal{Z}$  (39)).

For an example, consider Figure 3.1's tree diagram. It has already served other purposes. Nonetheless, it also illustrates the traditional game  $(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5)$  defined by  $X^5 = \{0, 1, 2\}$ ,  $E^5 = \{\langle 0, 1 \rangle, \langle 0, 2 \rangle\}$ ,  $\mathcal{H}^5 = \{\{0\}\}$ ,

$$(30) \quad \begin{aligned} \alpha^5 &= \{ \langle \langle 0, 1 \rangle, \text{left} \rangle, \langle \langle 0, 2 \rangle, \text{right} \rangle \}, \\ \iota^5 &= \{ \langle 0, \text{Alex} \rangle \}, \text{ and} \\ u_{\text{Alex}}^5 &= \{ \langle \{0, 1\}, 2 \rangle, \langle \{0, 2\}, 4 \rangle \} \end{aligned}$$

(footnote 13 explains that, in this paper, a function is a set of couples). This definition implies that  $\pi_1 E^5 = \{0\}$ , that  $\mathcal{Z}^5 = \{\{0, 1\}, \{0, 2\}\}$ , that  $F^5(0) = \{\text{left}, \text{right}\}$ , that  $A^5 = \{\text{left}, \text{right}\}$ , and that  $I^5 = \{\text{Alex}\}$  (these derivative entities inherit the example's superscript <sup>5</sup>).

## 5.2. "PENTAFORMING" A TRADITIONAL GAME

The remainder of this Section 5 develops a bijection between traditional games (Definition 5.1) and certain kinds of pentaform games (Definition 3.5). To begin, this Section 5.2 constructs an operator  $\mathbf{P}$  which maps each traditional game to a pentaform game. Mnemonically,  $\mathbf{P}$  "pentaforms" a traditional game. To be specific, let  $\mathbf{P}$  be the operator (equivalently function) that takes each traditional game  $(X, E, \mathcal{H}, \alpha, \iota, u)$  to the  $\mathbf{P}(X, E, \mathcal{H}, \alpha, \iota, u) = (\hat{Q}, \hat{u})$  defined by

$$(31a) \quad \hat{Q} = \{ \langle \iota(w), H_w, w, \alpha(w, y), y \rangle \mid \langle w, y \rangle \in E \} \text{ and}$$

$$(31b) \quad \hat{u} = u,$$

where in (31a), for each decision node  $w \in \pi_1 E$ ,  $H_w$  is the cell of the information-set partition  $\mathcal{H}$  that contains  $w$  ( $\mathcal{H}$  partitions  $\pi_1 E$  by [T2] in Definition 5.1).

Say that a quintuple set  $Q$  has *information-set situations* iff

$$(32) \quad (\forall j \in J) W_j = j.$$

For example, equation (10) shows that the pentaform  $Q^2$  (Figure 2.2 or 3.2) has information-set situations, while equation (9) shows that the pentaform  $Q^3$  (Figure 2.3 or 3.0) does not. Thus the set of pentaform games with information-set situations is a

---

defined in Section A.1's last two paragraphs). Second, footnote 20 explains the slight benefit of allowing utility numbers in  $\bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$  rather than the more familiar  $\mathbb{R}$ .

proper subset of the set of pentaform games. Theorem 5.2 shows that the operator  $\mathsf{P}$  maps traditional games into this proper subset of pentaform games. This reflects the fact that the information sets of traditional games are less general than the situations of pentaform games.<sup>35</sup>

**Theorem 5.2.** *The operator  $\mathsf{P}$  takes each traditional game (Definition 5.1) to a pentaform game (Definition 3.5) with information-set situations (32). (Proof: Lemma D.1.)*

The definition (31) of  $\mathsf{P}$  accords with Section 2’s informal process of expanding edges into quintuples. To explore this, consider the traditional game  $(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5)$  defined in equation (30). It is illustrated by the tree diagram on Figure 3.1’s left-hand side. Then

$$(33) \quad \mathsf{P}(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5) = (Q^1, u^1),$$

where the couple  $(Q^1, u^1)$  shown in the figure’s right-hand side. Transparently, the lower table  $u^1$  is a mere rearrangement of the utilities in the tree diagram, in accord with (31b). Less transparently, the upper table  $Q^1$  is derived from the tree diagram, in accord with (31a). To see this, first note that the quintuples (rows) in the upper table can be indexed by the edges  $\langle w, y \rangle$  that they contain. This indexing accords with the dummy variable  $\langle w, y \rangle$  in (31a). Then consider, for example, the quintuple containing the edge  $\langle w, y \rangle = \langle 0, 2 \rangle$ . As in (31a), and as in Section 2’s informal process, this edge is expanded into the quintuple  $\langle \text{Alex}, \{0\}, 0, \text{right}, 2 \rangle$  by including its action  $\alpha^5(0, 2) = \text{right}$ , its decision node’s information set  $H_0 = \{0\} \in \mathcal{H}$ , and its decision node’s player  $\iota^5(0) = \text{Alex}$ . The other quintuple in the upper table is derived in the same way.

The practical import of Theorem 5.2 is that pentaform games are general enough to accommodate any discrete extensive-form game. Although Section 2’s examples are finite, pentaforms can have trees with arbitrary degree (which admits decision nodes with uncountably many immediate successors), and up to countably infinite height (which admits an infinite horizon, as in the example pentaform of Streufert 2023, Section 2.2).

### 5.3. “TRADITIONALIZING” A PENTAFORM GAME

This Section 5.3 constructs the operator  $\mathsf{T}$  from pentaform games (Definition 3.5) to traditional games (Definition 5.1). Mnemonically,  $\mathsf{T}$  “traditionalizes” a pentaform game. To be specific, let  $\mathsf{T}$  be the operator that takes a pentaform game  $(Q, u)$  to the  $\mathsf{T}(Q, u) = (\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u})$  defined by

---

<sup>35</sup>The lesser generality of traditional games is an artifact of Definition 5.1 rather than a disadvantage of the diverse literature that Definition 5.1 represents. For example, Myerson 1991 specifies a game with information states like the situations here (footnote 8), and such a construction has been left out of Definition 5.1 because it is unusual in the literature (other constructions have been left out as well, as explained in the first paragraph of Section 5.1).

$$\begin{aligned}
(34a) \quad & \bar{X} = W \cup Y, \\
(34b) \quad & \bar{E} = \pi_{WY}(Q), \\
(34c) \quad & \bar{\mathcal{H}} = \{W_j \mid j \in J\}, \\
(34d) \quad & \bar{\alpha} = \{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(Q) \},^{36} \\
(34e) \quad & \bar{t} = \pi_{WI}(Q), \text{ and} \\
(34f) \quad & \bar{u} = u
\end{aligned}$$

(where  $J$ ,  $W$ ,  $Y$ , and  $W_j$  abbreviate the projections  $\pi_J(Q)$ ,  $\pi_W(Q)$ ,  $\pi_Y(Q)$ , and  $\pi_W(Q_j)$ , in accord with (8) and the sentence thereafter).

**Theorem 5.3.** *The operator  $\mathbb{T}$  takes each pentaform game (Definition 3.5) to a traditional game (Definition 5.1). (Proof: Lemma D.2.)*

Broadly speaking, a pentaform  $Q$  is a high-dimensional relation, while a traditional extensive form is a list of low-dimensional relations. Correspondingly, a pentaform describes the relationships between players, situations, nodes, and actions all at once, while a traditional extensive form describes the same relationships one by one. In this light, it is reasonable that equations (34a)–(34e) would use various projections of a pentaform  $Q$  to build the first five components of a traditional game. To be completely precise, [a] definitions (34a), (34b), (34d), and (34e) use projections of  $Q$  itself, while [b] each information set  $W_j$  in (34c) is the projection of a *slice* of  $Q$  (but, if  $Q$  has information-set situations (32), then the right-hand side of (34c) reduces to  $\{W_j \mid j \in J\} = \{j \mid j \in J\} = J$ , which is yet another projection of  $Q$  itself).

The practical import of Theorem 5.3 is that each pentaform game can be interpreted as a traditional game. In other words, the pentaform formulation does not introduce anything new.

#### 5.4. BIJECTION

The bijection of the following theorem is the paper’s main result. The bijection is illustrated in Figure 5.1 by the two opposing arrows between the thick vertical bars. As claimed in Section 1.1, this bijection is constructive and intuitive. The construction is provided by equations (31) and (34). The intuition is provided by Section 5.2’s second-last paragraph, by Section 5.3’s second-last paragraph, and by the remainder of this Section 5.4.<sup>37</sup>

**Theorem 5.4 (Main Theorem).**  *$\mathbb{P}$  is a bijection from the collection of traditional games to the collection of pentaform games with information-set situations (32). Its inverse is the restriction of  $\mathbb{T}$  to the collection of pentaform games with information-set situations. (Proof D.5.)*

<sup>36</sup>Footnote 14 explains a similar construction. Intuitively, this is a function from two variables to one variable.

<sup>37</sup>Incidentally, Table 5.1, Figure 5.1, and all the equations of Sections 5.4 and 5.5 place traditional games on the left and pentaform games on the right. This consistency seems helpful.




FIGURE 5.1. The operators P and T

For example, consider the traditional game  $(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5)$  from (30) and the pentaform game  $(Q^1, u^1)$  from Figure 3.1's tables. Equation (33) shows that  $P(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5) = (Q^1, u^1)$ . By applying T to both sides, we have

$$TP(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5) = T(Q^1, u^1),$$

which by Theorem 5.4 implies  $(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5) = T(Q^1, u^1)$ . Thus the operators P and T toggle back and forth between  $(X^5, E^5, \mathcal{H}^5, \alpha^5, \iota^5, u^5)$  and  $(Q^1, u^1)$ . In this sense, the two games are equivalent in spite of their superficial differences.

For another example, consider the pentaform game  $(Q^2, u^2)$  in Figure 3.2. Since this game has information-set situations,<sup>38</sup> Theorem 5.4 implies  $PT(Q^2, u^2) = (Q^2, u^2)$ . In this sense, the traditional game  $T(Q^2, u^2)$  and the pentaform game  $(Q^2, u^2)$  are equivalent. Further, this  $T(Q^2, u^2)$  could be explicitly derived by applying the projections of definition (34) to  $(Q^2, u^2)$ .

### 5.5. DERIVATIVE ENTITIES

As discussed earlier, Table 5.1's left-hand side summarizes the notation and terminology of an arbitrary traditional game. More specifically, the table's left-hand side lists the 6 components of a traditional game, together with 10 of its derivatives. Now consider the table's right-hand side. It re-expresses the same 16 traditional entities using the terms of the traditional game's pentaform equivalent. To be precise, the 16 rows of the table reproduce the 16 conclusions of the following theorem.

**Proposition 5.5.** *Suppose  $(X, E, \mathcal{H}, \alpha, \iota, u)$  is a traditional game, and let  $P(X, E, \mathcal{H}, \alpha, \iota, u) = (\hat{Q}, \hat{u})$ . Then the following hold. [1]  $X = \hat{W} \cup \hat{Y}$ . [2]  $E = \pi_{WY}(\hat{Q})$ . [3]  $r = \hat{r}$ . [4]  $\pi_1 E = \hat{W}$ . [5]  $\pi_2 E = \hat{Y}$ . [6]  $E^{-1} = \hat{p}$ . [7]  $\preceq = \hat{\preceq}$ . [8]  $\prec = \hat{\prec}$ . [9]  $\mathcal{Z} = \hat{\mathcal{Z}}$ . [10]  $\mathcal{H} = \hat{J}$ . [11]  $\alpha = \{\langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(\hat{Q})\}$  (footnote 14). [12]  $A = \hat{A}$ . [13]  $F = \hat{F}$ . [14]  $\iota = \pi_{WI}(\hat{Q})$ . [15]  $I = \hat{I}$ . [16]  $u = \hat{u}$ . (Proof D.6.)*

<sup>38</sup>As Figure 5.1 suggests, the composition PT changes any pentaform game into a pentaform game with information-set situations. For example, consider  $(Q^3, u^2)$ , where  $Q^3$  is from Figure 3.0 and  $u^2$  is from Figure 3.2 (this mixture of examples is a well-defined pentaform game because  $I^3 = I^2$  and  $\mathcal{Z}^3 = \mathcal{Z}^2$ ). Definition (34) implies  $T(Q^3, u^2) = T(Q^2, u^2)$ , which implies  $PT(Q^3, u^2) = PT(Q^2, u^2)$ , which by Theorem 5.4 and  $Q^2$ 's information-set situations implies  $PT(Q^3, u^2) = (Q^2, u^2)$ . Hence the composition PT changes  $(Q^3, u^2)$ , which does not have information-set situations, into  $(Q^2, u^2)$ , which does.

Conversely, Proposition 5.6 interprets the derivatives of a pentaform game in terms of the pentaform game’s traditional equivalent. This proposition starts with an arbitrary pentaform game. This setting is more general than that of Proposition 5.5, because the range of  $\mathbf{P}$  (in Proposition 5.5) consists of the special pentaform games with information-set situations (this fact is from Theorem 5.4 and Figure 5.1). To reconcile the propositions’ conclusions, note that conclusion [10\*] of Proposition 5.6 reduces to conclusion [10] of Proposition 5.5 in the special case of information-set situations (32).

**Proposition 5.6.** *Suppose  $(\hat{Q}, \hat{u})$  is a pentaform game, and let  $(X, E, \mathcal{H}, \alpha, \iota, u) = \mathbf{T}(\hat{Q}, \hat{u})$ . Then all the conclusions of Proposition 5.5 hold, except that conclusion [10] is replaced by [10\*]  $\mathcal{H} = \{\hat{W}_j \mid j \in \hat{J}\}$ . (Proof D.6.)*

## APPENDIX A. OUT-TREES

### A.1. FAMILIAR DEFINITIONS

A *directed graph* is a couple  $(X, E)$  such that  $E \subseteq X^2$ . The elements of  $X$  are called *nodes*, and the elements of  $E$  are called *edges*. A *walk in a directed graph  $(X, E)$  from  $w$  to  $y$*  is a sequence  $\langle x_0, x_1, x_2, \dots, x_m \rangle$  in  $X$  such that

$$(35) \quad x_0 = w, \quad x_m = y, \quad \text{and } (\forall \ell < m) \langle x_\ell, x_{\ell+1} \rangle \in E.$$

Note that a walk<sup>39</sup> can visit a node more than once. An *out-tree*<sup>40</sup> is a directed graph  $(X, E)$  such that

$$(36) \quad (\exists r \in X)(\forall x \in X) \text{ there exists a unique walk in } (X, E) \text{ from } r \text{ to } x.$$

The term “out-tree” is synonymous with the terms “(outwardly) directed rooted tree” and “diverging arborescence”. It can be shown that each out-tree has exactly one node  $r$  that satisfies (36). Call this the *root node* of the out-tree. For example,  $(\{5\}, \emptyset)$  is a one-node out-tree because the one-node sequence  $\langle 5 \rangle$  is the unique walk in  $(\{5\}, \emptyset)$  from the root node  $r = 5$  to itself. An out-tree is said to be *nontrivial* iff it has more than one node.

Consider a nontrivial out-tree  $(X, E)$  with its root node  $r$ . The remainder of this section will define its precedence relation and its collection of runs (that is, plays).

<sup>39</sup>There is a related but distinct concept in the literature. Let a “path in a graph” be a walk in a graph which visits each of its nodes *exactly once*. Thus, within an arbitrary graph, a “path in the graph” is a special kind of a walk in the graph. Nonetheless, if the graph is an out-tree, a “path in the graph” is equivalent to a walk in the graph. This can be confusing: It is important that walks rather than “paths” appear in the definition of an out-tree, and at the same time, within an out-tree, the distinction between walks and “paths” is irrelevant.

<sup>40</sup>Similar definitions in the game-theory literature include Perea 2001, page 10; Maschler, Solon, and Zamir 2013, page 41; and Kline and Luckraz 2016, page 88. Similar definitions in the graph-theory literature include Tutte 1984, page 126; and Bang-Jensen and Gutin 2009, page 21. (Beware of variations in terminology.) Several of these definitions apply only to finite trees, and several definitions explicitly include antisymmetry or irreflexivity. In contrast, the definition here applies equally well to infinite trees, and the definition here implies antisymmetry and thus irreflexivity.

Let  $\preceq$  and  $\prec$  be the binary relations on  $X$  defined by

$$(37a) \quad x \preceq y \text{ iff there is a walk in } (X, E) \text{ from } x \text{ to } y, \text{ and}$$

$$(37b) \quad x \prec y \text{ iff } (x \preceq y \text{ and } x \neq y).$$

Note  $\preceq$  is reflexive because the right-hand side of (37a) admits the trivial walk  $(\{x\}, \emptyset)$  from a node  $x$  to itself. Call  $\preceq$  and  $\prec$  the *weak* and *strict precedence* orders, respectively. It can be shown that  $\preceq$  is a partial order on  $X$  and that  $\prec$  is the asymmetric part of  $\preceq$ .

Runs will be defined in three steps. First, let an *end node* (or leaf) be a member of

$$(38) \quad \pi_2 E \setminus \pi_1 E,$$

and let a *finite run* (or finite play) be a walk in  $(X, E)$  from the root node  $r$  to an end node. Second, let an *infinite run* (or infinite play) be a sequence  $\langle x_0, x_1, \dots \rangle$  in  $X$  such that  $x_0 = r$  and  $(\forall \ell) \langle x_\ell, x_{\ell+1} \rangle \in E$ . Third, let a *run* (or play) be a finite run or an infinite run.

Although each run is a *sequence* of nodes, each run can be characterized by the *set* of nodes that it visits.<sup>41</sup> Accordingly, a run will be regarded as a set  $Z \subseteq X$ . Let  $\mathcal{Z}_{\text{ft}}$  be the collection of finite runs, let  $\mathcal{Z}_{\text{inf}}$  be the collection of infinite runs, and let

$$(39) \quad \mathcal{Z} = \mathcal{Z}_{\text{ft}} \cup \mathcal{Z}_{\text{inf}}$$

be the collection of runs. There are three possibilities: (i)  $\mathcal{Z} = \mathcal{Z}_{\text{ft}}$  and  $\mathcal{Z}_{\text{inf}} = \emptyset$ , as in the out-trees of Section 2's figures, (ii)  $\mathcal{Z} = \mathcal{Z}_{\text{inf}}$  and  $\mathcal{Z}_{\text{ft}} = \emptyset$ , in which case there are no end nodes, and (iii) both  $\mathcal{Z}_{\text{ft}}$  and  $\mathcal{Z}_{\text{inf}}$  are nonempty, as in the out-tree of an infinite centipede game.

## A.2. TWO LEMMAS

Lemmas A.1 and A.2 depart slightly from the literature.<sup>42</sup> They are used to prove Proposition 3.4. Later, Proposition 3.4 is used to prove Theorems 5.2–5.5.

**Lemma A.1.** *Suppose  $(X, E)$  is a nontrivial out-tree and its root is  $r$ . Then*

$$(40a) \quad X = \pi_1 E \cup \pi_2 E,$$

$$(40b) \quad E^{-1} \text{ is a function,}$$

$$(40c) \quad (\forall y \in \pi_2 E) (\exists m \geq 1) E^{-m}(y) \notin \pi_2 E, \text{ and}$$

$$(40d) \quad \{r\} = \pi_1 E \setminus \pi_2 E.$$

*Proof.* See Claims 3–6.

<sup>41</sup>One proof of this result is built on the fact that each walk in an out-tree must also be a “path” in the out-tree (footnote 39).

<sup>42</sup>Lemmas A.1 and A.2 can be combined to characterize nontrivial out-trees: *Let  $E$  be a set of couples. Then  $(X, E)$  is a nontrivial out-tree iff (a)  $X = \pi_1 E \cup \pi_2 E$ , (b)  $E^{-1}$  is a function, (c)  $(\forall y \in \pi_2 E) (\exists m \geq 1) E^{-m}(y) \notin \pi_2 E$ , and (d)  $\pi_1 E \setminus \pi_2 E$  is a singleton.* This way of expressing an out-tree is unusual. A similar concept is in Streufert 2018, equation (1). A related but less similar concept is in Knuth 1997, page 373.

*Claim 1:*  $r \in \pi_1 E$ . Nontriviality implies there is a node  $x$  distinct from  $r$ . Thus (36) implies there is a walk of length greater than zero from  $r$  to  $x$ ; which by the walk definition (35) implies there is an  $x_1$  such that  $\langle r, x_1 \rangle \in E$ ; which implies  $r \in \pi_1 E$ .

*Claim 2:*  $r \notin \pi_2 E$ . Suppose  $r \in \pi_2 E$ . Then there is  $w \in X$  such that  $\langle w, r \rangle \in E$  (it is irrelevant whether  $w$  and  $r$  are distinct). Meanwhile, (36) implies there is a walk from  $r$  to  $w$ . Thus  $\langle w, r \rangle \in E$  implies there is a still longer walk from  $r$  to  $w$ , back to  $r$  via  $\langle w, r \rangle$ , and on to  $w$  for a second time. The previous two sentences contradict the uniqueness in (36).

*Claim 3:* (40a) holds. Since  $(X, E)$  is a directed graph, we have  $E \subseteq X^2$ , which implies  $X \supseteq \pi_1 E \cup \pi_2 E$ . To show  $X \subseteq \pi_1 E \cup \pi_2 E$ , take an arbitrary  $x \in X$ . On the one hand, if  $x = r$ , then Claim 1 implies  $x \in \pi_1 E$ . On the other hand, if  $x \neq r$ , then (36) implies there is a walk of length greater than zero from  $r$  to  $x$ ; which by the walk definition (35) implies there is a sequence  $\langle x_0, x_1, \dots, x_m \rangle$  with  $x_0 = r$ ,  $x_m = x$ ,  $m \geq 1$ , and  $(\forall \ell < m) \langle x_\ell, x_{\ell+1} \rangle \in E$ ; which implies  $\langle x_{m-1}, x \rangle \in E$ ; which implies  $x \in \pi_2 E$ .

*Claim 4:* (40b) holds. Since  $(X, E)$  is a directed graph, we have  $E \subseteq X^2$ , which implies  $E$  is a relation. Thus it suffices to show that there are not  $y$ ,  $w^1$ , and  $w^2$  such that  $\langle y, w^1 \rangle \in E^{-1}$  and  $\langle y, w^2 \rangle \in E^{-1}$  and  $w^1 \neq w^2$  (footnote 13). Suppose there were. Then we would have

$$(41a) \quad \langle w^1, y \rangle \in E,$$

$$(41b) \quad \langle w^2, y \rangle \in E, \text{ and}$$

$$(41c) \quad w^1 \neq w^2.$$

By (36) at  $x = w^1$ , and by (41a), there is a walk from  $r$  to  $w^1$  and then on to  $y$  in one step. Similarly, by (36) at  $x = w^2$ , and by (41b), there is a walk from  $r$  to  $w^2$  and then on to  $y$  in one step. Yet, by (36) at  $x = y$ , there is a unique walk from  $r$  to  $y$ , which contradicts the previous two sentences because  $w^1$  and  $w^2$  are distinct by (41c).

*Claim 5:* (40c) holds. Take  $y \in \pi_2 E$ . Then Claim 2 implies  $y \neq r$ ; which by (36) implies there is a walk of length greater than zero from  $r$  to  $y$ ; which implies there is a sequence  $\langle x_0, x_1, \dots, x_m \rangle$  such that

$$x_0 = r, \quad x_m = y, \quad m \geq 1, \quad \text{and} \quad (\forall \ell < m) \langle x_\ell, x_{\ell+1} \rangle \in E;$$

which by Claim 4 (40b) implies  $x_0 = r$ ,  $x_m = y$ ,  $m \geq 1$ , and  $(\forall \ell < m) x_\ell = E^{-1}(x_{\ell+1})$ ; which implies  $r = E^{-m}(y)$ ; which by Claim 2 implies  $E^{-m}(y) \notin \pi_2 E$ .

*Claim 6:* (40d) holds. For  $\{r\} \subseteq \pi_1 E \setminus \pi_2 E$ , Claim 1 implies  $r \in \pi_1 E$ , and Claim 2 implies  $r \notin \pi_2 E$ . For  $\{r\} \supseteq \pi_1 E \setminus \pi_2 E$ , it suffices to show

$$\pi_1 E \setminus \{r\} \subseteq \pi_2 E.$$

Take  $w \in \pi_1 E \setminus \{r\}$ . Then  $w \in \pi_1 E$  and Claim 3 (40a) imply  $w \in X$ . Thus  $w \neq r$  and (36) imply that there is a walk of length greater than zero from  $r$  to  $w$ ; which implies there is  $\langle x_0, x_1, \dots, x_m \rangle$  such that  $x_0 = r$ ,  $x_m = w$ ,  $m \geq 1$ , and  $(\forall \ell < m) \langle x_\ell, x_{\ell+1} \rangle \in E$ ; which implies  $\langle x_{m-1}, w \rangle \in E$ ; which implies  $w \in \pi_2 E$ .  $\square$



**Lemma A.2.** *Let  $E$  be a set of couples. Then  $(X, E)$  is a nontrivial out-tree if*

$$(42a) \quad X = \pi_1 E \cup \pi_2 E,$$

$$(42b) \quad E^{-1} \text{ is a function,}$$

$$(42c) \quad (\forall y \in \pi_2 E)(\exists m \geq 1) E^{-m}(y) \notin \pi_2 E, \text{ and}$$

$$(42d) \quad \pi_1 E \setminus \pi_2 E \text{ is a singleton.}$$

*Proof.* Suppose  $(X, E)$  satisfies (42a)–(42d). By (42d) define  $r$  by

$$(43) \quad \{r\} = \pi_1 E \setminus \pi_2 E.$$

For future reference, we argue that

$$(44) \quad (\forall x \in X, m \geq 0, n \geq 0) E^{-m}(x) = E^{-n}(x) = r \text{ implies } m = n.$$

Suppose  $E^{-m}(x) = E^{-n}(x) = r$  and yet  $m \neq n$ . Assume without loss of generality that  $m > n$ . Then the equalities and manipulation imply  $r = E^{-m}(x) = E^{-(m-n)}(E^{-n}(x)) = E^{-(m-n)}(r)$ , which by  $m > n$  implies  $\langle E^{-1}(r), r \rangle \in E$ , which implies  $r \in \pi_2 E$ , which violates (43).

Since  $E$  is assumed to be a set of couples, (42a) implies  $E \subseteq X^2$ , which implies  $(X, E)$  is a directed graph. Thus it suffices to show that  $r$  satisfies (36). Toward that end, take an arbitrary node  $x \in X$ .

On the one hand, suppose  $x = r$ . By inspection,  $\langle r \rangle$  is a one-node walk in  $(X, E)$  from  $x = r$  to itself. For uniqueness, note that any other walk in  $(X, E)$  from  $r$  to itself would be a sequence  $\langle y_0, y_1, \dots, y_n \rangle$  with  $y_0 = r$ ,  $y_n = r$ ,  $n \geq 1$ , and  $(\forall k < n) \langle y_k, y_{k+1} \rangle \in E$ . This implies  $\langle y_{n-1}, r \rangle \in E$ , which implies  $r \in \pi_2 E$ , which violates (43).

On the other hand, suppose  $x \neq r$ . Then (43) implies  $x \notin \pi_1 E \setminus \pi_2 E$ ; which by (42a) implies  $x \in \pi_2 E$ ; which by (42c) implies that there is an  $m \geq 1$  such that  $E^{-m}(x) \notin \pi_2 E$ ; which by (42a) implies  $E^{-m}(x) \in \pi_1 E \setminus \pi_2 E$ ; which by (43) implies

$$(45a) \quad E^{-m}(x) = r.$$

This implies that

$$(45b) \quad \langle x_\ell \rangle_{\ell=0}^m = \langle E^{-(m-\ell)}(x) \rangle_{\ell=0}^m$$

is a sequence such that  $x_0 = r$  and  $x_m = x$ . By (45b) and manipulation,

$$(\forall \ell < m) x_\ell = E^{-(m-\ell)}(x) = E^{-1}(E^{-(m-(\ell+1))}(x)) = E^{-1}(x_{\ell+1});$$

which implies  $(\forall \ell < m) \langle x_\ell, x_{\ell+1} \rangle \in E$ ; which by the previous sentence implies that  $\langle x_\ell \rangle_{\ell=0}^m$  is a walk in  $(X, E)$  from  $r$  to  $x$ .

For uniqueness, suppose that  $\langle y_k \rangle_{k=0}^n$  is an arbitrary walk in  $(X, E)$  from  $r$  to  $x$ . Then  $y_0 = r$ ,  $y_n = x$ , and  $(\forall k < n) \langle y_k, y_{k+1} \rangle \in E$ ; which implies  $y_0 = r$ ,  $y_n = x$ , and  $(\forall k < n) y_k = E^{-1}(y_{k+1})$ ; which implies

$$(46a) \quad y_0 = r \text{ and}$$

$$(46b) \quad \langle y_k \rangle_{k=0}^n = \langle E^{-(n-k)}(x) \rangle_{k=0}^n.$$

By (45b) and (46b), the walks  $\langle x_\ell \rangle_{\ell=0}^m$  and  $\langle y_k \rangle_{k=0}^n$  are equal provided that  $m = n$  (note  $\ell$  and  $k$  are just dummy variables). Thus it suffices to show that  $m = n$ . Toward that end, note that (46a) and (46b) at  $k = 0$  imply that  $E^{-n}(x) = r$ . This and (45a) imply that the hypothesis of (44) is met, which by (44) implies  $m = n$ .  $\square$

### APPENDIX B. FOR PENTAFORM DEFINITION (SECTION 3)

**Lemma B.1** (from set theory). *Suppose  $G$  is a set of couples  $\langle x, y \rangle$ . Then define  $X = \pi_1 G$ , define  $Y = \pi_2 G$ , and for each  $y \in Y$  define  $G^{-1}(y) = \{x \mid \langle x, y \rangle \in G\}$ . Then the following are equivalent.*

- (a)  $G$  is a function.
- (b)  $(\forall y_1 \in Y, y_2 \in Y) y_1 \neq y_2$  implies  $G^{-1}(y_1) \cap G^{-1}(y_2) = \emptyset$ .
- (c)  $\langle G^{-1}(y) \rangle_{y \in Y}$  is an injectively indexed partition of  $X$ .

**Lemma B.2.** *Suppose  $Q$  is a quintuple set. Then the following hold.*

- (a)  $(\forall j \in J) W_j = \{w \mid \langle w, j \rangle \in \pi_{WJ}(Q)\}$ .
- (b)  $(\forall j \in J) Y_j = \{y \mid \langle y, j \rangle \in \pi_{YJ}(Q)\}$ .
- (c)  $(\forall j \in J) \pi_{WA}(Q_j) = \{\langle w, a \rangle \mid \langle w, a, j \rangle \in \pi_{WAJ}(Q)\}$ .

*Proof.* (a). By  $Q_j$ 's definition (2),  $\pi_W(Q_j)$  equals  $\pi_W(\{\langle i_*, j, w_*, a_*, y_* \rangle \in Q\})$ , which by the definition of  $\pi_W$  equals

$$\{w \mid (\exists i_*, a_*, y_*) \langle i_*, j, w, a_*, y_* \rangle \in Q\},$$

which by the definition of  $\pi_{WJ}(Q)$  equals  $\{w \mid \langle w, j \rangle \in \pi_{WJ}(Q)\}$ .

(b). This is proved as part (a), except for reversing the roles of  $w$  and  $y$ .

(c). Definition (2) implies  $\pi_{WA}(Q_j)$  equals  $\pi_{WA}(\{\langle i_*, j, w_*, a_*, y_* \rangle \in Q\})$ , which by the definition of  $\pi_{WA}$  equals

$$\{\langle w, a \rangle \mid (\exists i_*, y_*) \langle i_*, j, w, a, y_* \rangle \in Q\},$$

which by the definition of  $\pi_{WAJ}(Q)$  equals  $\{\langle w, a \rangle \mid \langle w, a, j \rangle \in \pi_{WAJ}(Q)\}$ .  $\square$

### Proof B.3 (for Proposition 3.2).

*Claim 1:* For each  $j \in J$ , the inverse image  $(\pi_{WJ}(Q))^{-1}(j)$  equals the information set  $W_j$ . Fix a situation  $j \in J$ . Then its inverse image  $(\pi_{WJ}(Q))^{-1}(j)$  by definition equals  $\{w \mid \langle w, j \rangle \in \pi_{WJ}(Q)\}$ , which by Lemma B.2(a) equals the information set  $W_j$ .

*Conclusion.* Consider Lemma B.1 with  $G$  there equal to  $\pi_{WJ}(Q)$  here. Thus  $X, Y$ , and  $G^{-1}(y)$  there equal  $W, J$ , and  $(\pi_{WJ}(Q))^{-1}(j)$  here. So the lemma implies that the following are equivalent.

- (a\*)  $\pi_{WJ}(Q)$  is a function.
- (b\*)  $(\forall j_1 \in J, j_2 \in J) j_1 \neq j_2$  implies  $(\pi_{WJ}(Q))^{-1}(j_1) \cap (\pi_{WJ}(Q))^{-1}(j_2) = \emptyset$ .
- (c\*)  $\langle (\pi_{WJ}(Q))^{-1}(j) \rangle_{j \in J}$  is an injectively indexed partition of  $W$ .

Condition (a\*) is equivalent to the proposition's (a) by the definition of  $[Pj \leftarrow w]$  in Definition 3.1. Meanwhile, (b\*) is equivalent to the proposition's (b) by Claim 1. Similarly, (c\*) is equivalent to the proposition's (c) by Claim 1.  $\square$

**Lemma B.4.** *Suppose  $Q$  is a quintuple set which satisfies  $[Pj \leftarrow w]$  and  $[Pw \leftarrow y]$ . Then  $\langle Y_j \rangle_{j \in J}$  is an injectively indexed partition of  $Y$ .*

*Proof.* Consider Lemma B.1 with  $G$  there equal to  $\pi_{YJ}(Q)$  here. Then  $X$ ,  $Y$ , and  $G^{-1}(y)$  there equal  $Y$ ,  $J$ , and  $(\pi_{YJ}(Q))^{-1}(j)$  here. Therefore, since the lemma assumptions  $[Pj \leftarrow w]$  and  $[Pw \leftarrow y]$  imply that  $\pi_{YJ}(Q)$  is a function, Lemma B.1(a $\Rightarrow$ c) implies that  $\langle \pi_{YJ}(Q)^{-1}(j) \rangle_{j \in J}$  is an injectively indexed partition of  $Y$ . Each  $\pi_{YJ}(Q)^{-1}(j)$  by definition equals  $\{y \mid \langle y, j \rangle \in \pi_{YJ}(Q)\}$ , which by Lemma B.2(b) equals  $Y_j$ .  $\square$

**Proof B.5 (for Proposition 3.3).**

(a $\Rightarrow$ b). Assume (a), and take a situation  $j \in J$ . Then  $\pi_{WA}(Q_j)$  is a Cartesian product, which implies  $\pi_{WA}(Q_j) = \pi_W(\pi_{WA}(Q_j)) \times \pi_A(\pi_{WA}(Q_j))$ .<sup>43</sup> Note  $\pi_W(\pi_{WA}(Q_j))$  is  $\pi_W(Q_j)$ , which by abbreviation (8) is  $W_j$ . Similarly  $\pi_A(\pi_{WA}(Q_j))$  is  $\pi_A(Q_j)$ , which by abbreviation (8) is  $A_j$ .

(b $\Rightarrow$ c). Assume (b). Take a situation  $j \in J$  and a decision node  $w$  in  $j$ 's information set  $W_j$ . Express the node's feasible action set  $F(w)$  as  $\pi_A(F|_{W:w})$ .<sup>44</sup> By  $F$ 's definition (11), this is  $\pi_A(\pi_{WA}(Q)|_{W:w})$ , which by inspection is  $\pi_A(\pi_{WA}(Q|_{W:w}))$ , which by the assumption  $w \in W_j$  and the axiom  $[Pj \leftarrow w]$  is  $\pi_A(\pi_{WA}(Q_j|_{W:w}))$ , which by inspection is  $\pi_A(\pi_{WA}(Q_j)|_{W:w})$ , which by part (b) is  $\pi_A((W_j \times A_j)|_{W:w})$ , which by the assumption  $w \in W_j$  is  $\pi_A(\{w\} \times A_j)$ , which by inspection is  $A_j$ .

(c $\Rightarrow$ d). This holds by inspection.

(d $\Rightarrow$ a). Assume (d). Take a situation  $j \in J$ . It must be shown that  $\pi_{WA}(Q_j)$  is a Cartesian product. Note  $\pi_W(\pi_{WA}(Q_j))$  is  $\pi_W(Q_j)$ , which by abbreviation (8) is  $W_j$ . Thus it suffices to show that

$$(\forall w_1 \in W_j, w_2 \in W_j) \{a \mid \langle w_1, a \rangle \in \pi_{WA}(Q_j)\} = \{a \mid \langle w_2, a \rangle \in \pi_{WA}(Q_j)\}.$$

Toward that end, take decision nodes  $w_1 \in W_j$  and  $w_2 \in W_j$ . Then the left-hand side is  $\pi_A(\{\langle w_1, a \rangle \in \pi_{WA}(Q_j)\})$ , which can be expressed using the generalized slice operator of footnote 44 as  $\pi_A(\pi_{WA}(Q_j)|_{W:w_1})$ , which by assumption  $w_1 \in W_j$  and axiom  $[Pj \leftarrow w]$  is equal to  $\pi_A(\pi_{WA}(Q)|_{W:w_1})$ , which by  $F$ 's definition (11) is equal to  $\pi_A(F|_{W:w_1})$ , which is an alternative expression for  $F(w_1)$ . By the same reasoning, the right-hand side is  $F(w_2)$ . Finally,  $F(w_1) = F(w_2)$  by assumption (d).  $\square$

**Lemma B.6.** *Suppose  $Q$  is a quintuple set which satisfies  $[Pw \leftarrow y]$ ,  $[Py]$ , and  $[Pr]$ . Then  $(W \cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree.*

*Proof.*

*Claim 1:*  $W = \pi_1(\pi_{WY}(Q))$ . By abbreviation (8),  $W$  equals  $\pi_W(Q)$ , which by inspection equals  $\pi_1(\pi_{WY}(Q))$ .

*Claim 2:*  $Y = \pi_2(\pi_{WY}(Q))$ . By abbreviation (8),  $Y$  equals  $\pi_Y(Q)$ , which by inspection equals  $\pi_2(\pi_{WY}(Q))$ .

<sup>43</sup>To be clear, consider the expression  $\pi_W(\pi_{WA}(Q_j))$ . The inner projection is a projection of a set of quintuples, as discussed in Section 3.3. Meanwhile, the outer projection is a projection of a set of couples, which is taken to be self-explanatory. Similar self-explanatory projections appear throughout Proof B.5.

<sup>44</sup>Proof B.5 uses a general slice operator of the form  $T|_{C:c}$ . Specifically, suppose that  $T$  is a set of tuples with the same set of coordinates, and that  $C$  is one of those coordinates. Then let  $T|_{C:c}$  be the set consisting of those tuples in  $T$  that have the value  $c$  in the coordinate  $C$ . For example, any situation slice  $Q_j$  could be more explicitly expressed as  $Q|_{J:j}$ .

| $Q$  | $[Pi\leftarrow j]$ | $[Pj\leftarrow w]$ | $[Pwa]$ | $[Pwa\rightarrow y]$ | $[Pw\leftarrow y]$ | $[Pa\leftarrow y]$ | $[Py]$   | $[Pr]$  |
|--|--------------------|--------------------|---------|----------------------|--------------------|--------------------|----------|---------|
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Bob,now,0,b,1} \rangle\}$                                       | $\circ$            | $+$                | $+$     | $+$                  | $+$                | $+$                | $+$      | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,later,0,b,1} \rangle\}$                                     | $+$                | $\circ$            | $+$     | $+$                  | $+$                | $+$                | $+$      | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,0,c,2} \rangle, \langle \text{Ann,now,1,b,3} \rangle\}$ | $+$                | $+$                | $\circ$ | $+$                  | $+$                | $+$                | $+$      | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,0,b,2} \rangle\}$                                       | $+$                | $+$                | $+$     | $\circ$              | $+$                | $+$                | $+$      | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,1,b,1} \rangle\}$                                       | $+$                | $+$                | $+$     | $+$                  | $\circ$            | $+$                | $+^{45}$ | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,0,c,1} \rangle\}$                                       | $+$                | $+$                | $+$     | $+$                  | $+$                | $\circ$            | $+$      | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,2,b,3} \rangle, \langle \text{Ann,now,3,b,2} \rangle\}$ | $+$                | $+$                | $+$     | $+$                  | $+$                | $+$                | $\circ$  | $+$     |
| $\{\langle \text{Ann,now,0,b,1} \rangle, \langle \text{Ann,now,2,b,3} \rangle\}$                                       | $+$                | $+$                | $+$     | $+$                  | $+$                | $+$                | $+$      | $\circ$ |

TABLE B.1. Eight examples, each of which violates exactly one axiom. This can be efficiently verified by inspecting each column. (See footnote 45 about one exceptional cell.)

*Claim 3:* (a)  $\pi_{YW}(Q) = (\pi_{WY}(Q))^{-1}$ . (b)  $p = (\pi_{WY}(Q))^{-1}$ . Part (a) holds by inspection. Part (b) holds by (a) and the definition of  $p$  in axiom  $[Pw\leftarrow y]$ .

*Conclusion.* By inspection  $\pi_{WY}(Q)$  is a set of couples. Thus by Lemma A.2, it suffices to show that

$$(47a) \quad W \cup Y = \pi_1(\pi_{WY}(Q)) \cup \pi_2(\pi_{WY}(Q)),$$

$$(47b) \quad (\pi_{WY}(Q))^{-1} \text{ is a function,}$$

$$(47c) \quad (\forall y \in \pi_2(\pi_{WY}(Q))) (\exists m \geq 1) (\pi_{WY}(Q))^{-m}(y) \notin \pi_2(\pi_{WY}(Q)), \text{ and}$$

$$(47d) \quad \pi_1(\pi_{WY}(Q)) \setminus \pi_2(\pi_{WY}(Q)) \text{ is a singleton.}$$

Claims 1 and 2 imply (47a). Axiom  $[Pw\leftarrow y]$  and Claim 3(a) imply (47b). Axiom  $[Py]$  and Claims 2 and 3(b) imply (47c). Axiom  $[Pr]$  and Claims 1 and 2 imply (47d).  $\square$

**Lemma B.7.** *Suppose  $Q$  is a quintuple set such that  $(W \cup Y, \pi_{WY}(Q))$  is a non-trivial out-tree. Let  $r$  denote its root. Then  $Q$  satisfies  $[Pw\leftarrow y]$ ,  $[Py]$ ,  $[Pr]$ , and  $\{r\} = W \setminus Y$ .*

*Proof.*

<sup>45</sup>When axiom  $[Pw\leftarrow y]$  fails, the  $p = \pi_{YW}(Q)$  defined within  $[Pw\leftarrow y]$  fails to be a function, which then leaves axiom  $[Py]$  ill-defined. This affects one cell in Table B.1: in that context,  $p = \pi_{YW}(Q)$  is taken to be a correspondence, and  $[Py]$  is taken to mean that  $(\forall y \in Y) (\exists m \geq 1) p^m(y) \setminus Y \neq \emptyset$ .

*Claim 1:*  $\pi_1(\pi_{WY}(Q)) = W$ . By inspection,  $\pi_1(\pi_{WY}(Q))$  equals  $\pi_W(Q)$ , which by abbreviation (8) equals  $W$ .

*Claim 2:*  $\pi_2(\pi_{WY}(Q)) = Y$ . By inspection,  $\pi_2(\pi_{WY}(Q))$  equals  $\pi_Y(Q)$ , which by abbreviation (8) equals  $Y$ .

*Claim 3:*  $(\pi_{WY}(Q))^{-1} = \pi_{YW}(Q)$ . This holds by inspection.

*Conclusion.* Because  $(W \cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree by assumption, Lemma A.1(40b)–(40d) implies

$$(48a) \quad (\pi_{WY}(Q))^{-1} \text{ is a function,}$$

$$(48b) \quad (\forall y \in \pi_2(\pi_{WY}(Q))) (\exists m \geq 1) (\pi_{WY}(Q))^{-m}(y) \notin \pi_2(\pi_{WY}(Q)), \text{ and}$$

$$(48c) \quad \{r\} = \pi_1(\pi_{WY}(Q)) \setminus \pi_2(\pi_{WY}(Q)).$$

Equation (48a) and Claim 3 imply axiom  $[Pw \leftarrow y]$ . Axiom  $[Pw \leftarrow y]$  defines the immediate-predecessor function  $p$  to be  $\pi_{YW}(Q)$ , which by Claim 3 implies  $p = (\pi_{WY}(Q))^{-1}$ , so that equation (48b) and Claim 2 imply axiom  $[Py]$ . Equation (48c) and Claims 1 and 2 imply  $\{r\} = W \setminus Y$ . This implies axiom  $[Pr]$ .  $\square$

**Proof B.8 (for Proposition 3.4).** Part (a) follows by inspection from Lemmas B.6 and B.7. For part (b), suppose  $Q$  satisfies  $[Pj \leftarrow w]$ ,  $[Py]$ , and  $[Pr]$ . Then Lemma B.6 implies  $(W \cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree. Let  $r$  be its root. Then Lemma B.7 implies  $\{r\} = W \setminus Y$ .  $\square$

**Lemma B.9.** *Suppose  $Q$  is a quintuple set which satisfies  $[Pw \leftarrow y]$  and  $[Py]$ . Then the following hold.*

(a)  $(\forall y \in Y) (\exists \ell \geq 1) p^\ell(y) \in W \setminus Y$ .

(b) If  $Q$  also satisfies  $[Pr]$ , then  $(\forall y \in Y) (\exists \ell \geq 1) p^\ell(y) = r$ .

*Proof.* (a). Take a successor node  $y \in Y$ . Then  $[Py]$  implies there is  $\ell \geq 1$  such that  $p^\ell(y) \notin Y$ . Further, since  $[Pw \leftarrow y]$  defines  $p$  to be  $\pi_{YW}(Q)$ , the range of  $p$  is  $\pi_W(Q)$ , which by abbreviation (8) is  $W$ . Thus  $p^\ell(y) \in W \setminus Y$ .

(b). If  $[Pr]$  also holds, then (12) defines  $r$  to be the sole member of  $W \setminus Y$ . Hence (b) follows from (a).  $\square$

#### APPENDIX C. FOR PENTAFORM TOOLS AND APPLICATIONS (SECTION 4)

**Proof C.1 (for Proposition 4.1).** (a). First consider axiom  $[Pi \leftarrow j]$ . Since a function is being regarded as a set of couples (see footnote 13 on page 10), we have that [1] any subset of a function is itself a function. Meanwhile, the assumption  $Q' \subseteq Q$  implies [2]  $\pi_{JI}(Q') \subseteq \pi_{JI}(Q)$ . Then in steps,  $[Pi \leftarrow j]$  for  $Q$  implies  $\pi_{JI}(Q)$  is a function, which by [1] and [2] implies  $\pi_{JI}(Q')$  is a function, which implies  $[Pi \leftarrow j]$  for  $Q'$ .

Next consider each of the axioms  $[Pj \leftarrow w]$ ,  $[Pw \rightarrow y]$ ,  $[Pa \leftarrow y]$ , and  $[Pw \leftarrow y]$ . Here the axiom for  $Q$  implies the corresponding axiom for  $Q'$  by an argument similar to that of the previous paragraph for  $[Pi \leftarrow j]$ .

Finally consider axiom  $[Py]$ . First note that  $[Pw \leftarrow y]$  for  $Q'$  has already been derived, and that this implies  $p' = \pi_{YW}(Q')$  is a well-defined function with domain  $Y'$  and range  $W'$ . Now take  $y \in Y'$ . Then  $p'(y) \in W'$  is well-defined. [Step 1] If  $p'(y) \notin Y'$ , then  $p'(y) \in W' \setminus Y'$  and the argument is complete. Else  $p'(y) \in Y'$  so

$(p')^2(y) \in W'$  is well-defined. [Step 2] If  $(p')^2(y) \notin Y'$ , then  $(p')^2(y) \in W' \setminus Y'$  and the argument is complete. Else  $(p')^2(y) \in Y'$  so  $(p')^3(y) \in W'$  is well-defined. By repeating this process indefinitely, either the argument finishes at some step or  $(\forall m \geq 1) (p')^m(y) \in Y'$ .

To rule out the latter contingency, suppose it held. Note  $Q' \subseteq Q$  implies  $p' = \pi_{Y'W}(Q')$  is a restriction of  $p = \pi_{YW}(Q)$ . Hence the supposition implies that  $(\forall m \geq 1) p^m(y) \in Y'$ . Also note  $Q' \subseteq Q$  implies  $Y' \subseteq Y$ . Thus the second-previous sentence implies [1]  $(\forall m \geq 1) p^m(y) \in Y$ , and further, the definition of  $y$  in the previous paragraph implies [2]  $y \in Y$ . These two observations contradict [Py] for  $Q$ .

(b). This follows from part (a) and the pentaform definition (Definition 3.1).  $\square$

**Lemma C.2.** *Suppose  $Q$  is a pentaform and  $t \in T$  (where  $T$  is the subroot set (15)). Then the following hold.*

- (a)  ${}^tW = \{w \in W \mid t \preceq w\}$ .
- (b)  ${}^tY = \{y \in Y \mid t \prec y\}$ .
- (c)  $(\forall j \in {}^tJ) ({}^tQ)_j = Q_j$ .<sup>46</sup>
- (d)  ${}^tQ = \cup\{Q_j \mid j \in {}^tJ\}$ .

*Proof.* (a). In steps,  ${}^tW$  by abbreviation (8) is  $\pi_W({}^tQ)$ , which by definition (14) is  $\pi_W(\{\langle i, j, w, a, y \rangle \in Q \mid t \preceq w\})$ , which by inspection is  $\{w \in \pi_W(Q) \mid t \preceq w\}$ , which by abbreviation (8) is  $\{w \in W \mid t \preceq w\}$ .

(b). In steps,  ${}^tY$  abbreviates  $\pi_Y({}^tQ)$ , which by definition (14) is  $\pi_Y(\{\langle i, j, w, a, y \rangle \in Q \mid t \preceq w\})$ , which by inspection is

$$\{y \in Y \mid (\exists w) t \preceq w, \langle w, y \rangle \in \pi_{WY}(Q)\}.$$

Thus it suffices to show that this set's entrance requirement is equivalent to  $t \prec y$ . The entrance requirement implies  $t \preceq w$  and  $w \prec y$ , which implies  $t \prec y$ . Conversely,  $t \prec y$  implies that  $t \preceq p(y)$  and  $p(y) \prec y$ , which implies the entrance requirement at  $w = p(y)$ .

(c). Take a situation  $j \in {}^tJ$ . Simply,  ${}^tQ \subseteq Q$  implies  $({}^tQ)_j \subseteq Q_j$ . For the reverse inclusion, suppose there was a quintuple in  $Q_j$  that was not in  $({}^tQ)_j$ . Then the quintuple being in  $Q_j$  implies there is  $\langle i_*, w_*, a_*, y_* \rangle$  such that  $\langle i_*, j, w_*, a_*, y_* \rangle$  is in  $Q_j$  but not  $({}^tQ)_j$ . Hence two applications of the slice definition (2) implies  $\langle i_*, j, w_*, a_*, y_* \rangle$  is in  $Q$  but not  ${}^tQ$ , which implies  $\langle i_*, j, w_*, a_*, y_* \rangle \in Q \setminus {}^tQ$ , which implies  $j \in \pi_J(Q \setminus {}^tQ)$ . This and the assumption  $j \in {}^tJ$  imply that  ${}^tJ$  and  $\pi_J(Q \setminus {}^tQ)$  intersect. This contradicts the lemma's assumption that  $t \in T$ , by  $T$ 's definition (15).

(d). For the forward inclusion, take a quintuple  $\langle i, j, w, a, y \rangle \in {}^tQ$ . Then projection implies  $j \in \pi_J({}^tQ)$ , which by abbreviation (8) implies  $j \in {}^tJ$ . Further,  $\langle i, j, w, a, y \rangle \in Q$  and  $Q_j$ 's definition (2) imply  $\langle i, j, w, a, y \rangle \in Q_j$ . For the reverse inclusion, take a situation  $j \in {}^tJ$ . Then part (c) implies  $Q_j$  is equal to  $({}^tQ)_j$ , which by slice definition (2) is a subset of  ${}^tQ$ .  $\square$

**Lemma C.3.** *Suppose  $Q$  is a pentaform and  $w \in W$ . Then  $w \in T$  iff there is  $J' \subseteq J$  such that  ${}^wQ = \cup\{Q_j \mid j \in J'\}$ . (This lemma is cited in the text.)*

<sup>46</sup>To be clear,  $({}^tQ)_j$  is the situation- $j$  slice (definition (2)) of the quintuple set  ${}^tQ$ .

*Proof.* For the forward direction, suppose  $w \in T$  (that is, suppose  $w$  is a subroot). Then Lemma C.2(d) implies  ${}^wQ = \{Q_j | j \in J'\}$  for  $J' = {}^wJ$ .

For the reverse direction, suppose  $w \notin T$ . Then the assumption  $w \in W$  and  $T$ 's definition (15) imply there is a  $j_* \in J$  which is listed both in a quintuple of  ${}^wQ$  and in a quintuple of  $Q \setminus {}^wQ$ . Thus definition (2) implies that the slice  $Q_{j_*}$  intersects both  ${}^wQ$  and  $Q \setminus {}^wQ$ . Thus, since  $\{Q_j | j \in J\}$  is a partition, it does not have a subcollection  $\{Q_j | j \in J'\}$  whose union is  ${}^wQ$ .  $\square$

**Proof C.4 (for Proposition 4.3).**

*Claim 1:*  ${}^tQ$  satisfies [Pr]. It suffices to show  ${}^tW \setminus {}^tY = \{t\}$ .

For the forward inclusion, it suffices to show that  ${}^tW \setminus \{t\} \subseteq {}^tY$ . Toward that end, suppose  $w \in {}^tW \setminus \{t\}$ , that is, suppose  $w$  is a decision node in  ${}^tW$  other than  $t$  itself. Then Lemma C.2(a) implies  $t \preceq w$  and  $t \neq w$ , which by definition (37b) imply [a]  $t \prec w$ . Further, [a] with the order definitions (37a) and (37b) imply there is a nontrivial walk from  $t$  to  $w$ , which by the walk definition (35) implies [b]  $w \in Y$ . Finally, Lemma C.2(b) with [a] and [b] implies  $w \in {}^tY$ .

For the reverse inclusion, first note that  $T$ 's definition (15) implies  $t \in W$ , which by Lemma C.2(a) implies  $t \in {}^tW$ . Thus it remains to show  $t \notin {}^tY$ . If  $t \in {}^tY$  did hold, then Lemma C.2(b) would imply  $t \prec t$ , which is impossible by  $\prec$ 's definition (37b).

*Conclusion.* Lemma C.2(d) (or the forward direction of Lemma C.3) implies that there is  $J' \subseteq J$  such that  ${}^tQ = \cup\{Q_j | j \in J'\}$ . Thus Corollary 4.2(b) and Claim 1 imply that  ${}^tQ$  is a pentaform.  $\square$

**Lemma C.5.** *Suppose  $\mathcal{Q}$  is a weakly separated (17) collection of quintuple sets which satisfy [Pi $\prec$ j], [Pj $\prec$ w], [Pwa], [Pwa $\rightarrow$ y], [Pw $\prec$ y], and [Pa $\prec$ y]. Then  $\cup\mathcal{Q}$  satisfies the same six axioms.*

*Proof.* This holds by the following claims.

*Claim 1:*  $\cup\mathcal{Q}$  satisfies [Pi $\prec$ j]. By inspection, [1]  $\pi_{JI}(\cup\mathcal{Q}) = \cup_{Q \in \mathcal{Q}} \pi_{JI}(Q)$ . Also, each  $Q \in \mathcal{Q}$  satisfies [Pi $\prec$ j] by assumption, which implies that [2]  $(\forall Q \in \mathcal{Q}) \pi_{JI}(Q)$  is a function. Also, weak separation (17) implies that [3] the members  $Q$  of  $\mathcal{Q}$  have distinct situations  $j$ . These three facts imply that  $\pi_{JI}(\cup\mathcal{Q})$  is the union of a set of functions with disjoint domains, which implies that  $\pi_{JI}(\cup\mathcal{Q})$  is itself a function, which implies that  $\cup\mathcal{Q}$  satisfies [Pi $\prec$ j].

*Claim 2:*  $\cup\mathcal{Q}$  satisfies [Pj $\prec$ w], [Pwa $\rightarrow$ y], [Pw $\prec$ y], and [Pa $\prec$ y]. The arguments for these axioms mimic Claim 1's argument for [Pi $\prec$ j]. More precisely, the arguments for [Pj $\prec$ w] and [Pwa $\rightarrow$ y] rely on the members of  $\mathcal{Q}$  having distinct decision nodes  $w$ , and the arguments for [Pa $\prec$ y] and [Pw $\prec$ y] rely on the members of  $\mathcal{Q}$  having distinct successor nodes  $y$ .

*Claim 3:*  $\cup\mathcal{Q}$  satisfies [Pwa]. It suffices to show that

$$(\forall j \in \pi_J(\cup\mathcal{Q})) \pi_{WA}((\cup\mathcal{Q})_j) = \pi_W((\cup\mathcal{Q})_j) \times \pi_A((\cup\mathcal{Q})_j).$$

Toward that end, take a situation  $j^* \in \pi_J(\cup\mathcal{Q})$ . Weak separation (17) implies the members of  $\mathcal{Q}$  have distinct situations, which implies there is a unique member  $Q^* \in \mathcal{Q}$  such that  $j^* \in \pi_J(Q^*)$ . Hence the union's slice  $(\cup\mathcal{Q})_{j^*}$  equals the member's

slice  $Q_{j^*}^*$ . Thus it suffices to show that

$$\pi_{WA}(Q_{j^*}^*) = \pi_W(Q_{j^*}^*) \times \pi_A(Q_{j^*}^*).$$

This holds because  $Q^*$  satisfies [Pwa] by assumption.  $\square$

**Proof C.6 (for Proposition 4.4(a)).** The result follows from Claims 4–6.

*Claim 1:  $W^A$  and  $Y^B$  are disjoint.* Since  $W^A = (W^A \setminus Y^A) \cup (W^A \cap Y^A)$ ,<sup>47</sup> it suffices to show both (a)  $(W^A \setminus Y^A) \cap Y^B = \emptyset$  and (b)  $(W^A \cap Y^A) \cap Y^B = \emptyset$ . Note (b) holds because weak separation implies  $Y^A \cap Y^B = \emptyset$ . Now consider (a). Since  $Y^B = (Y^B \setminus W^B) \cup (Y^B \cap W^B)$ , it suffices to show both

$$(W^A \setminus Y^A) \cap (Y^B \setminus W^B) = \emptyset \text{ and } (W^A \setminus Y^A) \cap (Y^B \cap W^B) = \emptyset.$$

The former is assumed by part (a). The latter holds because weak separation implies  $W^A \cap W^B = \emptyset$ .

*Claim 2:  $Q^A \cup Q^B$  satisfies [Pi $\leftarrow$ j], [Pj $\leftarrow$ w], [Pwa], [Pwa $\rightarrow$ y], [Pw $\leftarrow$ y], and [Pa $\leftarrow$ y].* By the block definition (16), the blocks  $Q^A$  and  $Q^B$  satisfy these six axioms. Thus weak separability and Lemma C.5 imply that their union does.

*Claim 3:  $Q^A \cup Q^B$  satisfies [Py].* For notational ease, define  $\bar{p} = \pi_{YW}(Q^A \cup Q^B)$ . To be clear,  $\bar{p}$  is a function since  $Q^A \cup Q^B$  satisfies [Pw $\leftarrow$ y] by Claim 2. By inspection, it is a superset (equivalently an extension) of both  $p^A = \pi_{YW}(Q^A)$  and  $p^B = \pi_{YW}(Q^B)$  (these equalities are two instances of the definition of the immediate-predecessor function in axiom [Pw $\leftarrow$ y]). Further, Lemma B.9(a) applied to the block  $Q^A$  implies  $(\forall y \in Y^A)(\exists \ell^A \geq 1)(p^A)^{\ell^A}(y) \in W^A \setminus Y^A$ , which by Claim 1 implies

$$(49) \quad (\forall y \in Y^A)(\exists \ell^A \geq 1)(p^A)^{\ell^A}(y) \in W^A \setminus (Y^A \cup Y^B).$$

To show that  $Q^A \cup Q^B$  satisfies [Py], take an arbitrary successor node  $y \in Y^A \cup Y^B$ . It must be shown that

$$(50) \quad (\exists \ell \geq 1) \bar{p}^\ell(y) \notin Y^A \cup Y^B.$$

Obviously  $y \in Y^A$  or  $y \in Y^B$  (intuitively,  $Q^A$  “precedes”  $Q^B$ , so backward walks from nodes in  $Y^A$  will tend to be shorter than backward walks from nodes in  $Y^B$ ). First suppose  $y \in Y^A$ . Then (49) implies there is  $\ell^A \geq 1$  such that  $(p^A)^{\ell^A}(y) \notin Y^A \cup Y^B$ , which by  $p^A \subseteq \bar{p}$  implies (50). Second suppose  $y \in Y^B$ . Then [Py] for  $Q^B$  implies there is  $\ell^B \geq 1$  such that  $(p^B)^{\ell^B}(y) \notin Y^B$ . If  $(p^B)^{\ell^B}(y) \notin Y^A$ , the previous sentence implies  $(p^B)^{\ell^B}(y) \notin Y^A \cup Y^B$ , which by  $p^B \subseteq \bar{p}$  implies (50). Otherwise  $(p^B)^{\ell^B}(y) \in Y^A$ , which by (49) implies there is  $\ell^A \geq 1$  such that  $(p^A)^{\ell^A}((p^B)^{\ell^B}(y)) \notin Y^A \cup Y^B$ , which by  $p^A \subseteq \bar{p}$  and  $p^B \subseteq \bar{p}$  implies  $\bar{p}^{\ell^A + \ell^B}(y) \notin Y^A \cup Y^B$ , which implies (50).

<sup>47</sup>For intuition, the identity  $W = (W \setminus Y) \cup (W \cap Y)$  splits any decision-node set  $W$  into the start-node set  $W \setminus Y$  and the “middle”-node set  $W \cap Y$ . Similarly, the identity  $Y = (Y \setminus W) \cup (Y \cap W)$  splits any successor-node set  $Y$  into the end-node set  $Y \setminus W$  and the “middle”-node set  $Y \cap W$ . The proof of Claim 1 uses the first identity to split  $W^A$  and the second identity to split  $Y^B$ ; the proof of Claim 5 uses the second identity to split  $Y^A$ ; and the proof of Claim 6 uses the first identity to split  $W^B$ .



*Claim 4:*  $Q^A \cup Q^B$  is a block. This follows from Claims 2 and 3 and the block definition (16).

*Claim 5:*  $Q^A \cup Q^B$ 's start-node set is the union of

$$W^A \setminus Y^A \text{ and } (W^B \setminus Y^B) \setminus (Y^A \setminus W^A).$$

By inspection,  $\pi_W(Q^A \cup Q^B) = W^A \cup W^B$  and  $\pi_Y(Q^A \cup Q^B) = Y^A \cup Y^B$ . So  $Q^A \cup Q^B$ 's start-node set is  $(W^A \cup W^B) \setminus (Y^A \cup Y^B)$ , which by inspection is the union of

$$W^A \setminus (Y^A \cup Y^B) \text{ and } W^B \setminus (Y^A \cup Y^B).$$

The first set is equal to  $W^A \setminus Y^A$  by Claim 1. The second set is equal to  $(W^B \setminus Y^B) \setminus Y^A$ , which is equal to  $(W^B \setminus Y^B) \setminus [(Y^A \setminus W^A) \cup (Y^A \cap W^A)]$ , which by  $W^B \cap W^A = \emptyset$  (from weak separation) is equal to  $(W^B \setminus Y^B) \setminus (Y^A \setminus W^A)$ .

*Claim 6:*  $Q^A \cup Q^B$ 's end-node set is the union of

$$(Y^A \setminus W^A) \setminus (W^B \setminus Y^B) \text{ and } Y^B \setminus W^B.$$

By inspection,  $\pi_Y(Q^A \cup Q^B) = Y^A \cup Y^B$  and  $\pi_W(Q^A \cup Q^B) = W^A \cup W^B$ . Thus  $Q^A \cup Q^B$ 's end-node set is  $(Y^A \cup Y^B) \setminus (W^A \cup W^B)$ , which by inspection is the union of

$$Y^A \setminus (W^A \cup W^B) \text{ and } Y^B \setminus (W^A \cup W^B).$$

The second set is equal to  $Y^B \setminus W^B$  by Claim 1. The first set is equal to  $(Y^A \setminus W^A) \setminus W^B$ , which is equal to  $(Y^A \setminus W^A) \setminus [(W^B \setminus Y^B) \cup (W^B \cap Y^B)]$ , which by  $Y^A \cap Y^B = \emptyset$  (from weak separation) is equal to  $(Y^A \setminus W^A) \setminus (W^B \setminus Y^B)$ .  $\square$

**Proof C.7 (for Proposition 4.4(b)).** This follows from Claims 3–5.

*Claim 1:*  $\cup \mathcal{Q}$  satisfies  $[Pi \leftarrow j]$ ,  $[Pj \leftarrow w]$ ,  $[Pwa]$ ,  $[Pwa \rightarrow y]$ ,  $[Pw \leftarrow y]$ , and  $[Pa \leftarrow y]$ . Since strong separation (18) implies weak separation (17),  $\mathcal{Q}$  is weakly separated. Further, the block definition (16) implies that every block in  $\mathcal{Q}$  satisfies the six axioms. Thus Lemma C.5 implies that  $\cup \mathcal{Q}$  satisfies the six axioms.

*Claim 2:*  $\cup \mathcal{Q}$  satisfies  $[Py]$ . For notational ease, let  $\bar{p} = \pi_Y(\cup \mathcal{Q})$ . To be clear,  $\bar{p}$  is a function since  $\cup \mathcal{Q}$  satisfies  $[Pw \leftarrow y]$  by Claim 1. Thus it suffices to show

$$(\forall y \in \pi_Y(\cup \mathcal{Q})) (\exists \ell \geq 1) \bar{p}^\ell(y) \notin \pi_Y(\cup \mathcal{Q}).$$

Toward that end, take an arbitrary successor node  $y \in \pi_Y(\cup \mathcal{Q})$ . Since  $\pi_Y(\cup \mathcal{Q}) = \cup_{Q \in \mathcal{Q}} \pi_Y(Q)$  by inspection, there is a block  $Q^* \in \mathcal{Q}$  such that  $y \in Y^*$ . In accord with axiom  $[Pw \leftarrow y]$ , define  $p^* = \pi_{YW}(Q^*)$ , which by  $Q^* \in \mathcal{Q}$  implies  $p^* \subseteq \pi_{YW}(\cup \mathcal{Q})$ , which by  $\bar{p}$ 's definition above implies  $p^* \subseteq \bar{p}$  (equivalently  $p^*$  is a restriction of  $\bar{p}$ ). Further, Lemma B.9(a) applied to the block  $Q^*$  implies there is  $\ell \geq 1$  such that  $(p^*)^\ell(y) \in W^* \setminus Y^*$ , which by  $p^* \subseteq \bar{p}$  implies

$$(51) \quad \bar{p}^\ell(y) \in W^* \setminus Y^*.$$

Since (51) implies  $\bar{p}^\ell(y) \in W^*$ , this  $\bar{p}^\ell(y)$  is a node of  $Q^*$ , which by strong separation implies that  $\bar{p}^\ell(y)$  is not a node of  $\cup(\mathcal{Q} \setminus \{Q^*\})$ , which implies that  $\bar{p}^\ell(y)$  is not a successor node of  $\cup(\mathcal{Q} \setminus \{Q^*\})$ , which by  $\bar{p}^\ell(y) \notin Y^*$  from (51) implies that  $\bar{p}^\ell(y)$  is not a successor node of  $\cup \mathcal{Q}$ , which is equivalent to  $\bar{p}^\ell(y) \notin \pi_Y(\cup \mathcal{Q})$ .

*Claim 3:*  $\cup \mathcal{Q}$  is a block. This holds by Claims 1 and 2 and the block definition (16).

*Claim 4:*  $\cup \mathcal{Q}$ 's start-node set is  $\cup_{Q \in \mathcal{Q}} (\pi_W(Q) \setminus \pi_Y(Q))$ . Since  $\cup \mathcal{Q}$ 's start-node set is  $\pi_W(\cup \mathcal{Q}) \setminus \pi_Y(\cup \mathcal{Q})$ , it suffices to show

$$\pi_W(\cup \mathcal{Q}) \setminus \pi_Y(\cup \mathcal{Q}) = \cup_{Q \in \mathcal{Q}} (\pi_W(Q) \setminus \pi_Y(Q)).$$

For the forward inclusion, consider [a]  $w \in \pi_W(\cup \mathcal{Q}) \setminus \pi_Y(\cup \mathcal{Q})$ . Note [a] implies  $w \in \pi_W(\cup \mathcal{Q})$ , which implies there is a block  $Q^* \in \mathcal{Q}$  such that  $w \in \pi_W(Q^*)$ . Further, [a] implies  $w \notin \pi_Y(\cup \mathcal{Q})$ , which by  $Q^* \in \mathcal{Q}$  implies  $w \notin \pi_Y(Q^*)$ . The previous two sentences imply  $w \in \pi_W(Q^*) \setminus \pi_Y(Q^*)$ , which by  $Q^* \in \mathcal{Q}$  implies  $w \in \cup_{Q \in \mathcal{Q}} (\pi_W(Q) \setminus \pi_Y(Q))$ .

Conversely, for the reverse inclusion, suppose there is an individual block  $Q' \in \mathcal{Q}$  with start node [b]  $w \in \pi_W(Q') \setminus \pi_Y(Q')$ . Then [b] implies  $w \in \pi_W(Q')$ , which by  $Q' \in \mathcal{Q}$  implies [c]  $w \in \pi_W(\cup \mathcal{Q})$ . Also, [b] implies  $w \in \pi_W(Q')$ , which (in a different direction) implies that  $w$  is a node of  $Q'$ , which by strong separation implies that  $w$  is not a node of  $\cup(\mathcal{Q} \setminus \{Q'\})$ , which implies that  $w$  is not a successor node of  $\cup(\mathcal{Q} \setminus \{Q'\})$ , which is equivalent to  $w \notin \pi_Y(\cup \mathcal{Q} \setminus \{Q'\})$ , which by  $w \notin \pi_Y(Q')$  from [b] implies  $w \notin \pi_Y(\cup \mathcal{Q})$ . This and [c] imply  $w \in \pi_W(\cup \mathcal{Q}) \setminus \pi_Y(\cup \mathcal{Q})$ .

*Claim 5:*  $\cup \mathcal{Q}$ 's end-node set is  $\cup_{Q \in \mathcal{Q}} (\pi_Y(Q) \setminus \pi_W(Q))$ . This can be proved like Claim 4 was proved. Replace “start” with “end”, switch  $W$  and  $Y$ , replace  $w$  with  $y$ , and replace “successor node” with “decision node”.  $\square$

**Proof C.8 (for Proposition 4.5).** For notational ease, define

$$(52) \quad \bar{Q} = \cup_{n \geq 0} Q^n.$$

*Claim 1:*  $\bar{Q}$  satisfies  $[Pi \leftarrow j]$ . Suppose  $\bar{Q}$  violates  $[Pi \leftarrow j]$ . Then there exists a situation  $j \in \bar{J}$  and players  $i_1 \in \bar{I}$  and  $i_2 \in \bar{I}$  such that  $i_1 \neq i_2$  and both  $\langle j, i_1 \rangle$  and  $\langle j, i_2 \rangle$  are in  $\pi_{JI}(\bar{Q})$ . Note  $\langle j, i_1 \rangle$  being in  $\pi_{JI}(\bar{Q})$  implies there is  $\langle w_1, a_1, y_1 \rangle$  such that  $\langle i_1, j, w_1, a_1, y_1 \rangle \in \bar{Q}$ , which by  $\bar{Q}$ 's definition (52) implies there is  $n_1 \geq 0$  such that

$$\langle i_1, j, w_1, a_1, y_1 \rangle \in Q^{n_1}.$$

Similarly,  $\langle j, i_2 \rangle$  being in  $\pi_{JI}(\bar{Q})$  implies there is  $\langle w_2, a_2, y_2 \rangle$  such that  $\langle i_2, j, w_2, a_2, y_2 \rangle$  is in  $\bar{Q}$ , which by  $\bar{Q}$ 's definition (52) implies there is  $n_2 \geq 0$  such that

$$\langle i_2, j, w_2, a_2, y_2 \rangle \in Q^{n_2}.$$

Let  $n_* = \max\{n_1, n_2\}$ . Then the assumption  $(\forall n \geq 1) Q^{n-1} \subseteq Q^n$  implies that both  $\langle i_1, j, w_1, a_1, y_1 \rangle$  and  $\langle i_2, j, w_2, a_2, y_2 \rangle$  are in  $Q^{n_*}$ , which implies that both  $\langle j, i_1 \rangle$  and  $\langle j, i_2 \rangle$  are in  $\pi_{JI}(Q^{n_*})$ , which by  $i_1 \neq i_2$  implies  $Q^{n_*}$  violates  $[Pi \leftarrow j]$ , which violates the assumption that  $Q^{n_*}$  is a pentaform.

*Claim 2:*  $\bar{Q}$  satisfies  $[Pj \leftarrow w]$ ,  $[Pwa \rightarrow y]$ ,  $[Pw \leftarrow y]$ , and  $[Pa \leftarrow y]$ . Each of these four axioms is derived as Claim 1 derived  $[Pi \leftarrow j]$ .

*Claim 3:*  $\bar{Q}$  satisfies  $[Pwa]$ . Suppose  $\bar{Q}$  violates  $[Pwa]$ . Then there exists a situation  $j \in \bar{J}$  with a decision node  $w_1 \in \bar{W}_j$ , another decision node  $w_2 \in \bar{W}_j$ , and an action  $a$ , such that [a]  $\langle w_1, a \rangle \in \pi_{WA}(\bar{Q}_j)$  and [b]  $\langle w_2, a \rangle \notin \pi_{WA}(\bar{Q}_j)$ . We will find a

contradiction. (Looking ahead, [a] will be used immediately and [b] will be used at the end.)

Note that [a] and Lemma B.2(c) imply  $\langle j, w_1, a \rangle \in \pi_{JWA}(\bar{Q})$ , which implies there is  $\langle i_1, y_1 \rangle$  such that  $\langle i_1, j, w_1, a, y_1 \rangle \in \bar{Q}$ , which by  $\bar{Q}$ 's definition (52) implies there is  $n_1 \geq 0$  such that

$$\langle i_1, j, w_1, a, y_1 \rangle \in Q^{n_1}.$$

Meanwhile,  $w_2 \in \bar{W}_j$  and Lemma B.2(a) imply  $\langle j, w_2 \rangle \in \pi_{JW}(\bar{Q})$ , which implies there is  $\langle i_2, a_2, y_2 \rangle \in \bar{Q}$  such that  $\langle i_2, j, w_2, a_2, y_2 \rangle \in \bar{Q}$ , which by  $\bar{Q}$ 's definition (52) implies there is  $n_2 \geq 0$  such that

$$\langle i_2, j, w_2, a_2, y_2 \rangle \in Q^{n_2}.$$

Let  $n_* = \max\{n_1, n_2\}$ . Then the assumption  $(\forall n \geq 1) Q^{n-1} \subseteq Q^n$  implies that both  $\langle i_1, j, w_1, a, y_1 \rangle$  and  $\langle i_2, j, w_2, a_2, y_2 \rangle$  are in  $Q^{n_*}$ . The first implies  $\langle j, w_1, a \rangle \in \pi_{JWA}(Q^{n_*})$ , which by Lemma B.2(c) implies [c]  $\langle w_1, a \rangle \in \pi_{WA}(Q_j^{n_*})$ . The second implies  $\langle j, w_2 \rangle \in \pi_{JW}(Q^{n_*})$ , which by Lemma B.2(a) implies  $w_2 \in W_j^{n_*}$ , which by abbreviation (8) implies  $w_2 \in \pi_W(Q_j^{n_*})$ , which by inspection implies [d]  $w_2 \in \pi_W(\pi_{WA}(Q^{n_*}))$ .

By assumption,  $Q^{n_*}$  is a pentaform, which implies  $Q^{n_*}$  satisfies [Pwa], which implies that  $\pi_{WA}(Q_j^{n_*})$  is a rectangle, which by [c] and [d] implies  $\langle w_2, a \rangle \in \pi_{WA}(Q_j^{n_*})$ , which by Lemma B.2(c) implies  $\langle j, w_2, a \rangle \in \pi_{JWA}(Q^{n_*})$ , which implies there is  $\langle i_\Delta, y_\Delta \rangle$  such that  $\langle i_\Delta, j, w_2, a, y_\Delta \rangle \in Q^{n_*}$ , which by  $\bar{Q}$ 's definition (52) implies  $\langle i_\Delta, j, w_2, a, y_\Delta \rangle \in \bar{Q}$ , which implies  $\langle j, w_2, a \rangle \in \pi_{JWA}(\bar{Q})$ , which by Lemma B.2(c) implies  $\langle w_2, a \rangle \in \pi_{WA}(\bar{Q}_j)$ , which contradicts [b].

*Claim 4:* (a)  $\bar{W} = \bigcup_{n \geq 0} W^n$ . (b)  $\bar{Y} = \bigcup_{n \geq 0} Y^n$ . First consider (a). In steps,  $\bar{W}$  by abbreviation (8) is  $\pi_W(\bar{Q})$ , which by  $\bar{Q}$ 's definition (52) is  $\pi_W(\bigcup_{n \geq 0} Q^n)$ , which by inspection equals  $\bigcup_{n \geq 0} \pi_W(Q^n)$ , which by abbreviation (8) is  $\bigcup_{n \geq 0} W^n$ . A similar argument holds for (b).

*Claim 5:*  $\{r^0\} = \bar{W} \setminus \bar{Y}$ . For the forward inclusion  $\{r^0\} \subseteq \bar{W} \setminus \bar{Y}$ , it suffices to show  $r^0 \in \bar{W}$  and  $r^0 \notin \bar{Y}$ . First,  $r^0$ 's definition (12) implies  $r^0 \in W^0$ , which by Claim 4(a) implies  $r^0 \in \bar{W}$ . Second, to show  $r^0 \notin \bar{Y}$ , suppose  $r^0 \in \bar{Y}$ . Then Claim 4(b) implies there is  $n \geq 0$  such that  $r^0 \in Y^n$ , which implies  $\{r^0\} \neq W^n \setminus Y^n$ , which by  $r^n$ 's definition (12) implies  $r^0 \neq r^n$ , which contradicts the proposition's assumption that  $r^n = r^0$ .

For the reverse inclusion  $\bar{W} \setminus \bar{Y} \subseteq \{r^0\}$ , it suffices to show that  $(\forall w \in \bar{W} \setminus \{r^0\}) w \in \bar{Y}$ . In other words, it suffices to show that every decision node other than  $r^0$  is a successor node. Toward that end, take such a decision node  $w \in \bar{W} \setminus \{r^0\}$ . Then Claim 4(a) implies there is  $n \geq 0$  such that [a]  $w \in W^n \setminus \{r^0\}$ . Note  $r^n$ 's definition (12) implies  $W^n \setminus Y^n = \{r^n\}$ , which implies [b]  $W^n \setminus \{r^n\} \subseteq Y^n$ . Then in steps, [a] implies  $w \in W^n \setminus \{r^0\}$ , which by the assumption  $r^n = r^0$  implies  $w \in W^n \setminus \{r^n\}$ , which by [b] implies  $w \in Y^n$ , which by Claim 4(b) implies  $w \in \bar{Y}$ .

*Claim 6:*  $\bar{Q}$  satisfies [Py]. For any  $n$ , the assumption that  $Q^n$  is a pentaform implies that  $Q^n$  satisfies [Pw $\leftarrow$ y], which implies that the  $p^n = \pi_{YW}(Q)$  defined in [Pw $\leftarrow$ y] is a function. Meanwhile, Claim 2 implies that  $\bar{Q}$  satisfies [Pw $\leftarrow$ y], which implies that the  $\bar{p} = \pi_{YW}(\bar{Q})$  defined in [Pw $\leftarrow$ y] is a function. Further, for each  $n$ ,  $\bar{Q}$ 's definition

(52) implies  $Q^n \subseteq \bar{Q}$ , which implies  $\pi_{YW}(Q^n) \subseteq \pi_{YW}(\bar{Q})$ , which by the preceding definitions implies [a]  $p^n \subseteq \bar{p}$  (in other words,  $p^n$  is a restriction of  $\bar{p}$ ).

For [Py], it suffices to show that  $(\forall y \in \bar{Y})(\exists \ell \geq 1) \bar{p}^\ell(y) \notin \bar{Y}$ . Toward that end, take a successor node  $y \in \bar{Y}$ . By Claim 4(b), there is  $n \geq 0$  such that  $y \in \bar{Y}^n$ . Since  $Q^n$  is a pentaform by assumption, Lemma B.9(b) implies there is  $\ell \geq 0$  such that  $(p^n)^\ell(y) = r^n$ , which by the assumption  $r^n = r^0$  implies  $(p^n)^\ell(y) = r^0$ , which by [a] implies  $\bar{p}^\ell(y) = r^0$ , which by Claim 5 implies  $\bar{p}^\ell(y) \notin \bar{Y}$ .

*Conclusion.* Claim 5 implies that  $\bar{Q}$  satisfies [Pr]. Claims 1–3 and 6 show that  $\bar{Q}$  satisfies the other seven axioms. Thus  $\bar{Q}$  is a pentaform. Finally, definition (12) states that  $\bar{Q}$ 's root is the sole member of  $\bar{W} \setminus \bar{Y}$ , which by Claim 5 is  $r^0$ .  $\square$

#### APPENDIX D. FOR EQUIVALENCE (SECTION 5)

**Lemma D.1 (implies Theorem 5.2).** *Suppose  $(X, E, \mathcal{H}, \alpha, \iota, u)$  is a traditional game. Let  $(\hat{Q}, \hat{u}) = \mathbf{P}(X, E, \mathcal{H}, \alpha, \iota, u)$ . Then (a)  $(\hat{Q}, \hat{u})$  is a pentaform game with information-set situations (32). Further, (b)  $\hat{J} = \mathcal{H}$ , (c)  $\hat{W} \cup \hat{Y} = X$ , (d)  $\pi_{WY}(\hat{Q}) = E$ , (e)  $\{\langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(\hat{Q})\} = \alpha$ , and (f)  $\pi_{WI}(\hat{Q}) = \iota$ .*

*Proof.* Part (a) follows from Claim 22; part (b) from Claim 6; parts (c)–(d) from Claims 16–17; and parts (e)–(f) from Claims 23–24.

*Claim 1:*  $E \ni \langle w, y \rangle \mapsto \iota(w)$  is a surjection to  $I$ . This is the composition of two surjections. First,  $E \ni \langle w, y \rangle \mapsto w$  is a surjection to  $\pi_1 E$  by inspection. Second,  $\pi_1 E \ni w \mapsto \iota(w)$  is a surjection to  $I$  because of [T5] (Definition 5.1) and because  $I$  is the range of  $\iota$  by definition (28).

*Claim 2:*  $E \ni \langle w, y \rangle \mapsto H_w$  is a surjection to  $\mathcal{H}$ . This is the composition of two surjections. First,  $E \ni \langle w, y \rangle \mapsto w$  is a surjection to  $\pi_1 E$  by inspection. Second,  $\pi_1 E \ni w \mapsto H_w$  is a surjection to  $\mathcal{H}$  by [T2] (Definition 5.1) and by the definition of  $\langle H_w \rangle_{w \in \pi_1 E}$  (after definition (31)).

*Claim 3:*  $E \ni \langle w, y \rangle \mapsto \alpha(w, y)$  is a surjection to  $A$ . This holds because of [T3] and because  $A$  is the range of  $\alpha$  by definition (24).

*Claim 4:*  $\hat{Q}$  is well-defined. This follows from definition (31a) and Claims 1–3.

*Claim 5:*  $\hat{I} = I$ . By abbreviation (8),  $\hat{I}$  is the projection  $\pi_I(\hat{Q})$ , which by definition (31a) and the surjection of Claim 1 is equal to  $I$ .

*Claim 6:*  $\hat{J} = \mathcal{H}$ . By abbreviation (8),  $\hat{J}$  is the projection  $\pi_J(\hat{Q})$ , which by definition (31a) and the surjection of Claim 2 is equal to  $\mathcal{H}$ .

*Claim 7:*  $\hat{W} = \pi_1 E$ . By abbreviation (8),  $\hat{W}$  is the projection  $\pi_W(\hat{Q})$ , which by definition (31a) is equal to  $\pi_1 E$ .

*Claim 8:*  $\hat{Y} = \pi_2 E$ . By abbreviation (8),  $\hat{Y}$  is the projection  $\pi_Y(\hat{Q})$ , which by definition (31a) is equal to  $\pi_2 E$ .

*Claim 9:*  $\hat{Q}$  satisfies [Pi $\leftrightarrow$ j]. Note that  $\pi_{JI}(\hat{Q}) = \{\langle H_w, \iota(w) \rangle \mid \langle w, y \rangle \in E\}$  by definition (31a). Thus  $\pi_{JI}(\hat{Q}) = \{\langle H_w, \iota(w) \rangle \mid w \in \pi_1 E\}$  by inspection. To show that this is a function, it suffices to show that  $(\forall w_1 \in \pi_1 E, w_2 \in \pi_1 E)$  the condition  $H_{w_1} = H_{w_2}$  implies  $\iota(w_1) = \iota(w_2)$ . Toward that end, suppose  $w_1 \in \pi_1 E$  and  $w_2 \in \pi_1 E$  are two

decision nodes such that  $H_{w_1} = H_{w_2}$ . Then the definition of  $\langle H_w \rangle_{w \in \pi_1 E}$  (after definition (31)) implies that  $w_1$  and  $w_2$  belong to the same cell of the partition  $\mathcal{H}$ , which by the measurability (29) of [T5] implies  $\iota(w_1) = \iota(w_2)$ .

*Claim 10:*  $\hat{Q}$  satisfies [Pj\*w]. Note that  $\pi_{WJ}(\hat{Q}) = \{ \langle w, H_w \rangle \mid \langle w, y \rangle \in E \}$  by definition (31a). Thus  $\pi_{WJ}(\hat{Q}) = \{ \langle w, H_w \rangle \mid w \in \pi_1 E \}$  by inspection. This is the function  $\langle H_w \rangle_{w \in \pi_1 E}$  (defined after definition (31)).

*Claim 11:*  $\hat{F} = F$ . The correspondence  $F: \pi_1 E \rightrightarrows A$  is defined in equation (26) by  $F(w) = \{ a \mid (\exists y) \alpha(w, y) = a \}$ . Since a correspondence is a set of couples (footnote 17 on page 12), this implies that

$$F = \{ \langle w, a \rangle \mid w \in \pi_1 E, (\exists y) \alpha(w, y) = a \}.$$

Note [T3] implies that the domain of  $\alpha$  is  $E$ . Hence  $F$  is  $\{ \langle w, \alpha(w, y) \rangle \mid \langle w, y \rangle \in E \}$ , which by definition (31a) is  $\pi_{WA}(\hat{Q})$ , which by definition (11) is  $\hat{F}$ .

*Claim 12:*  $(\forall H \in \mathcal{H}) \{ w \in \pi_1 E \mid H_w = H \} = H$ . (For intuition, regard this as an identity which holds in every traditional game.) To prove this equality, note that  $\mathcal{H}$  partitions  $\pi_1 E$  (by [T2]), and that (for each  $w \in \pi_1 E$ )  $H_w$  is the cell that contains  $w$  (by the definition of  $H_w$  after (31)). Now consider a cell  $H \in \mathcal{H}$ . Then  $\{ w \in \pi_1 E \mid H_w = H \}$  consists of the  $w$  in the cell  $H$ , which is  $H$  itself.

*Claim 13:*  $\hat{Q}$  has information-set situations (32), that is,  $(\forall j \in \hat{J}) \hat{W}_j = j$ . Take a situation  $j \in \hat{J}$ . Definition (31a) implies that the slice  $\hat{Q}_j$  satisfies

$$\hat{Q}_j = \{ \langle \iota(w), H_w, w, \alpha(w, y), y \rangle \mid \langle w, y \rangle \in E, H_w = j \},$$

which by projection implies  $\pi_W(\hat{Q}_j) = \{ w \mid (\exists y) \langle w, y \rangle \in E, H_w = j \}$ , which by abbreviation (8) implies  $\hat{W}_j = \{ w \mid (\exists y) \langle w, y \rangle \in E, H_w = j \}$ , which by inspection implies  $\hat{W}_j = \{ w \mid w \in \pi_1 E, H_w = j \}$ , which by manipulation implies

$$(53) \quad \hat{W}_j = \{ w \in \pi_1 E \mid H_w = j \}.$$

Meanwhile, notice that  $j$  is a cell in the partition  $\mathcal{H}$  because  $j \in \hat{J}$  by assumption and because  $\hat{J} = \mathcal{H}$  by Claim 6. Thus Claim 12 with  $j$  replacing  $H$  implies  $\{ w \in \pi_1 E \mid H_w = j \} = j$ . This and (53) imply the result.

*Claim 14:*  $\hat{Q}$  satisfies [Pwa]. By Proposition 3.3(a $\Leftarrow$ d) and Claim 10, it suffices to show that

$$(\forall j \in \hat{J}, w_1 \in \hat{W}_j, w_2 \in \hat{W}_j) \hat{F}(w_1) = \hat{F}(w_2).$$

By replacing  $\hat{J}$  with  $\mathcal{H}$  via Claim 6, by replacing the two appearances of  $\hat{W}_j$  with  $j$  via Claim 13, and by replacing  $\hat{F}$  with  $F$  via Claim 11, this is equivalent to

$$(\forall j \in \mathcal{H}, w_1 \in j, w_2 \in j) F(w_1) = F(w_2).$$

By a change of variables, this is equivalent to  $(\forall H \in \mathcal{H}, w_1 \in H, w_2 \in H) F(w_1) = F(w_2)$ , which holds by the measurability (27) of [T4].

*Claim 15:*  $\hat{Q}$  satisfies  $[Pw \rightarrow y]$ . Note  $\pi_{WY}(\hat{Q}) = \{ \langle w, \alpha(w, y), y \rangle \mid \langle w, y \rangle \in E \}$  by Definition (31a). Thus it suffices to show, for all  $\langle w_1, y_1 \rangle \in E$  and  $\langle w_2, y_2 \rangle \in E$ , that

$$\langle w_1, \alpha(w_1, y_1) \rangle = \langle w_2, \alpha(w_2, y_2) \rangle \text{ implies } y_1 = y_2.$$

Toward that end, take  $\langle w_1, y_1 \rangle \in E$  and  $\langle w_2, y_2 \rangle \in E$  such that  $\langle w_1, \alpha(w_1, y_1) \rangle = \langle w_2, \alpha(w_2, y_2) \rangle$ . Since the equality's first coordinate implies  $w_1 = w_2$ , the rest of the previous sentence implies that we have two edges of the form  $\langle w, y_1 \rangle$  and  $\langle w, y_2 \rangle$  such that  $\alpha(w, y_1) = \alpha(w, y_2)$ . Thus the local injectivity (25) of [T3] implies  $y_1 = y_2$ .

*Claim 16:*  $\hat{W} \cup \hat{Y} = X$ . In steps,  $\hat{W} \cup \hat{Y}$  by Claims 7 and 8 is equal to  $\pi_1 E \cup \pi_2 E$ , which by Lemma A.1(40a) and [T1] is equal to  $X$ .

*Claim 17:*  $\pi_{WY}(\hat{Q}) = E$ . In steps,  $\pi_{WY}(\hat{Q})$  by (31a) is equal to  $\{ \langle w, y \rangle \mid \langle w, y \rangle \in E \}$ , which by inspection is equal to  $E$ .

*Claim 18:*  $\hat{Q}$  satisfies  $[Pw \leftarrow y]$ ,  $[Py]$ , and  $[Pr]$ . Since [T1] implies that  $(X, E)$  is a nontrivial out-tree, Claims 16 and 17 imply that  $(\hat{W} \cup \hat{Y}, \pi_{WY}(\hat{Q}))$  is a nontrivial out-tree. Thus the reverse direction of Proposition 3.4(a) implies that  $\hat{Q}$  satisfies  $[Pw \leftarrow y]$ ,  $[Py]$ , and  $[Pr]$ .

*Claim 19:*  $\hat{Q}$  satisfies  $[Pa \leftarrow y]$ . Note  $\pi_{YA}(\hat{Q}) = \{ \langle y, \alpha(w, y) \rangle \mid \langle w, y \rangle \in E \}$  by definition (31a). Thus it suffices to show, for all  $\langle w_1, y_1 \rangle \in E$  and  $\langle w_2, y_2 \rangle \in E$ , that

$$y_1 = y_2 \text{ implies } \alpha(w_1, y_1) = \alpha(w_2, y_2).$$

Toward that end, take  $\langle w_1, y_1 \rangle \in E$  and  $\langle w_2, y_2 \rangle \in E$  such that [1]  $y_1 = y_2$ . Claim 17 implies  $\langle w_1, y_1 \rangle \in \pi_{WY}(\hat{Q})$  and  $\langle w_2, y_2 \rangle \in \pi_{WY}(\hat{Q})$ . Further, Claim 18 implies  $[Pw \leftarrow y]$ , which implies  $\pi_{YW}(\hat{Q})$  is a function. Thus [1] implies [2]  $w_1 = w_2$ . Together, [1] and [2] imply  $\langle w_1, y_1 \rangle = \langle w_2, y_2 \rangle$ , which implies  $\alpha(w_1, y_1) = \alpha(w_2, y_2)$ .

*Claim 20:*  $\hat{Q}$  is a pentaform. This follows from Claims 9, 10, 14, 15, 18, and 19.

*Claim 21:*  $\hat{Z} = \mathcal{Z}$ . Claims 16 and 17 imply that  $(\hat{W} \cup \hat{Y}, \pi_{WY}(\hat{Q}))$  equals the out-tree  $(X, E)$  from [T1]. This suffices since both pentaform games (end of Section 3.5) and traditional games (Section 5.1 paragraph a) derive their run collections from their out-trees (via equation (39)).

*Claim 22:*  $(\hat{Q}, \hat{u})$  is a pentaform game with information-set situations. Claims 13 and 20 imply  $\hat{Q}$  is a pentaform with information-set situations. Thus by Definition 3.5 it suffices to show that  $\hat{u}$  is a utility-function profile for  $\hat{Q}$ . Definition (31b) implies  $\hat{u}$  is equal to  $u$ , which by [T6] has the form  $\langle u_i : \mathcal{Z} \rightarrow \bar{\mathbb{R}} \rangle_{i \in I}$ , which is identical to the form  $\langle u_i : \hat{\mathcal{Z}} \rightarrow \bar{\mathbb{R}} \rangle_{i \in \hat{I}}$  because  $I = \hat{I}$  by Claim 5 and because  $\mathcal{Z} = \hat{\mathcal{Z}}$  by Claim 21.

*Claim 23:*  $\{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(\hat{Q}) \} = \alpha$ . By definition (31a), the left-hand side is

$$\{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \{ \langle w, y, \alpha(w, y) \rangle \mid \langle w, y \rangle \in E \} \},$$

which by inspection is  $\{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y \rangle \in E, a = \alpha(w, y) \}$ , which by inspection is  $\{ \langle \langle w, y \rangle, \alpha(w, y) \rangle \mid \langle w, y \rangle \in E \}$ , which by [T3] is  $\alpha$ .

*Claim 24:*  $\pi_{WI}(\hat{Q}) = \iota$ . In steps,  $\pi_{WI}(\hat{Q})$  by (31a) is  $\{ \langle w, \iota(w) \rangle \mid (\exists y) \langle w, y \rangle \in E \}$ , which by inspection is  $\{ \langle w, \iota(w) \rangle \mid w \in \pi_1 E \}$ , which by [T5] is  $\iota$ .  $\square$

**Lemma D.2 (implies Theorem 5.3).** *Suppose  $(Q, u)$  is a pentaform game. Let  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{r}, \bar{u}) = \Gamma(Q, u)$ . Then, (a)  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{r}, \bar{u})$  is a traditional game. Further, (b)  $\bar{r} = r$ , (c)  $\pi_1 \bar{E} = W$ , (d)  $\pi_2 \bar{E} = Y$ , (e)  $\bar{E}^{-1} = p$ . (f)  $\bar{\succ} = \succ$ , (g)  $\bar{\prec} = \prec$ , (h)  $\bar{\mathcal{Z}} = \mathcal{Z}$ , (i)  $\bar{A} = A$ , (j)  $\bar{F} = F$ , and (k)  $\bar{I} = I$ .*

*Proof.* Part (a) follows from Claim 21; parts (b)–(h) from Claims 3–7; and parts (i), (j), and (k) from Claims 11, 12, and 19.

*Claim 1:*  $(\bar{X}, \bar{E})$  is equal to  $(W \cup Y, \pi_{WY}(Q))$  and is a nontrivial out-tree. Definitions (34a)–(34b) imply  $(\bar{X}, \bar{E})$  is equal to  $(W \cup Y, \pi_{WY}(Q))$ . Proposition 3.4(a)'s forward direction implies  $(W \cup Y, \pi_{WY}(Q))$  is a nontrivial out-tree.

*Claim 2:*  $[T1]$  holds.  $(\bar{X}, \bar{E})$  is a nontrivial out-tree by Claim 1.

*Claim 3:*  $\bar{r} = r$ . Since  $(\bar{X}, \bar{E})$  is an out-tree by Claim 1, Section 5.1 paragraph a defines  $\bar{r}$  to be the root of  $(\bar{X}, \bar{E})$ . By Claim 1 again, this  $\bar{r}$  is the root of the out-tree  $(W \cup Y, \pi_{WY}(Q))$ , which by Proposition 3.4(b) is the unique element of  $W \setminus Y$ , which by definition (12) is  $r$ .

*Claim 4:*  $\pi_1 \bar{E} = W$ . In steps,  $\pi_1 \bar{E}$  by  $\bar{E}$ 's definition (34b) equals  $\pi_1(\pi_{WY}(Q))$ , which by inspection equals  $\pi_W(Q)$ , which by abbreviation (8) equals  $W$ .

*Claim 5:*  $\pi_2 \bar{E} = Y$ . In steps,  $\pi_2 \bar{E}$  by  $\bar{E}$ 's definition (34b) equals  $\pi_2(\pi_{WY}(Q))$ , which by inspection equals  $\pi_Y(Q)$ , which by abbreviation (8) equals  $Y$ .

*Claim 6:*  $\bar{E}^{-1} = p$ . In steps,  $\bar{E}^{-1}$  by definition (34b) is  $(\pi_{WY}(Q))^{-1}$ , which by inspection is  $\pi_{YW}(Q)$ , which by the definition of  $p$  within  $[Pw \prec y]$  is  $p$ .

*Claim 7:* (a)  $\bar{\succ} = \succ$ . (b)  $\bar{\prec} = \prec$ . (c)  $\bar{\mathcal{Z}} = \mathcal{Z}$ . Claim 1 suffices because both traditional games (Section 5.1, paragraph a) and pentaform games (end of Section 3.5) derive their precedence relations and run collections from their out-trees (via (37) and (39) in Section A.1).

*Claim 8:*  $[T2]$  holds. Proposition 3.2(a $\Rightarrow$ c) and axiom  $[Pj \prec w]$  imply  $\{W_j | j \in J\}$  partitions  $W$ , which by  $\bar{\mathcal{H}}$ 's definition (34c) implies  $\bar{\mathcal{H}}$  partitions  $W$ , which by Claim 4 implies  $\bar{\mathcal{H}}$  partitions  $\pi_1 \bar{E}$ .

*Claim 9:*  $\bar{\alpha}$  is a function from  $\bar{E}$ . By  $\bar{\alpha}$ 's definition (34d),

$$(54) \quad \bar{\alpha} = \{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(Q) \}.$$

First, for functionhood, it suffices (by footnotes 13 and 14 on page 10) to show

$$\langle \langle w, y \rangle, a^1 \rangle \in \bar{\alpha} \text{ and } \langle \langle w, y \rangle, a^2 \rangle \in \bar{\alpha} \text{ imply } a^1 = a^2.$$

Toward that end, suppose  $\langle \langle w, y \rangle, a^1 \rangle$  and  $\langle \langle w, y \rangle, a^2 \rangle$  belong to  $\bar{\alpha}$ . Then (54) implies  $\langle w, y, a^1 \rangle$  and  $\langle w, y, a^2 \rangle$  belong to  $\pi_{WYA}(Q)$ , which implies  $\langle y, a^1 \rangle$  and  $\langle y, a^2 \rangle$  belong to  $\pi_{YA}(Q)$ , which by  $[Pa \prec y]$  implies  $a^1 = a^2$ . Second, the domain of the function  $\bar{\alpha}$  is  $\pi_1 \bar{\alpha}$ , which by (54) equals  $\pi_{WY}(Q)$ , which by  $\bar{E}$ 's definition (34b) is  $\bar{E}$ .

*Claim 10:*  $[T3]$  holds. Because of Claim 9, it suffices to show that  $\bar{\alpha}$  is locally injective (25). We will prove the contrapositive. Toward that end, consider two edges  $\langle w, y_1 \rangle \in E$  and  $\langle w, y_2 \rangle \in E$  from decision node  $w$  which are both assigned the action  $\bar{\alpha}(w, y_1) = \bar{\alpha}(w, y_2)$ . It suffices to show  $y_1 = y_2$ .

Let  $a$  denote the common action  $\bar{\alpha}(w, y_1) = \bar{\alpha}(w, y_2)$ . Then  $\bar{\alpha}$ 's definition (34d) implies that both  $\langle w, y_1, a \rangle$  and  $\langle w, y_2, a \rangle$  are in  $\pi_{WYA}(Q)$ . Thus by rearrangement, both  $\langle w, a, y_1 \rangle$  and  $\langle w, a, y_2 \rangle$  are in  $\pi_{WAY}(Q)$ . Hence axiom [Pwa $\rightarrow$ y] implies  $y_1 = y_2$ .

*Claim 11:*  $\bar{A} = A$ . In steps, the action set  $\bar{A}$  by its definition (24) is the range of  $\bar{\alpha}$ , which is  $\{\bar{\alpha}(w, y) \mid \langle w, y \rangle \in \bar{E}\}$ , which by Claim 9 and  $\bar{\alpha}$ 's definition (34d) is  $\pi_2(\{\langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(Q)\})$ , which by inspection is  $\pi_3(\pi_{WYA}(Q))$ , which by inspection is  $\pi_A(Q)$ , which by abbreviation (8) is  $A$ .

*Claim 12:*  $\bar{F} = F$ . The correspondence  $\bar{F}: \pi_1 \bar{E} \rightarrow \bar{A}$  is defined in (26) by  $\bar{F}(w) = \{a \mid (\exists y) \bar{\alpha}(w, y) = a\}$ . Thus  $\bar{F}$  (by footnote 17 on page 12) is equal to

$$\{\langle w, a \rangle \mid w \in \pi_1 \bar{E}, (\exists y) \bar{\alpha}(w, y) = a\},$$

which by  $\bar{E}$ 's definition (34b) and  $\bar{\alpha}$ 's definition (34d) is equal to

$$\{\langle w, a \rangle \mid w \in \pi_1(\pi_{WY}(Q)), (\exists y) \langle w, y, a \rangle \in \pi_{WYA}(Q)\},$$

which by inspection is equal to  $\{\langle w, a \rangle \mid w \in \pi_W(Q), (\exists y) \langle w, a, y \rangle \in \pi_{WAY}(Q)\}$ , which by inspection is equal to  $\pi_{WA}(Q)$ , which by  $F$ 's definition (11) is equal to  $F$ .

*Claim 13:* [T4] holds. Proposition 3.3(a $\Rightarrow$ d) and axioms [Pj $\leftarrow$ w] and [Pwa] imply that  $(\forall j \in J, w_1 \in W_j, w_2 \in W_j) F(w_1) = F(w_2)$ . Since Proposition 3.2(a $\Rightarrow$ c) and axiom [Pj $\leftarrow$ w] imply that  $\langle W_j \rangle_{j \in J}$  is an injectively indexed partition, this is equivalent to

$$(\forall H \in \{W_j \mid j \in J\}, w_1 \in H, w_2 \in H) F(w_1) = F(w_2),$$

which by  $\bar{H}$ 's definition (34c) and Claim 12 implies

$$(\forall H \in \bar{\mathcal{H}}, w_1 \in H, w_2 \in H) \bar{F}(w_1) = \bar{F}(w_2),$$

which implies the measurability (27) of  $\bar{F}$ .

*Claim 14:* (a)  $\pi_{WJ}(Q)$  is a surjection from  $W$  to  $J$ . (b)  $\pi_{JI}(Q)$  is a surjection from  $J$  to  $I$ . Part (a) holds by axiom [Pj $\leftarrow$ w], and because  $W = \pi_W(Q)$  and  $J = \pi_J(Q)$  by abbreviation (8). Similarly, (b) holds by axiom [Pi $\leftarrow$ j], and because  $J = \pi_J(Q)$  and  $I = \pi_I(Q)$  by abbreviation (8).

*Claim 15:*  $\pi_{JI}(Q) \circ \pi_{WJ}(Q) = \pi_{WI}(Q)$ . By definition,  $\pi_{JI}(Q) \circ \pi_{WJ}(Q)$  is equal to  $\{\langle w, i \rangle \mid (\exists j) \langle w, j \rangle \in \pi_{WJ}(Q), \langle j, i \rangle \in \pi_{JI}(Q)\}$ , which by inspection is equal to

$$\{\langle w, i \rangle \mid (\exists j, i_\Delta) \langle w, j, i_\Delta \rangle \in \pi_{WJI}(Q), \langle j, i \rangle \in \pi_{JI}(Q)\}.$$

Consider the conditions  $\{w, y, i_\Delta\} \in \pi_{WJI}(Q)$  and  $\langle j, i \rangle \in \pi_{JI}(Q)$ . With axiom [Pi $\leftarrow$ j], they imply  $i_\Delta = i$ . Thus the set is equal to

$$\{\langle w, i \rangle \mid (\exists j) \langle w, j, i \rangle \in \pi_{WJI}(Q), \langle j, i \rangle \in \pi_{JI}(Q)\},$$

which by inspection is equal to  $\{\langle w, i \rangle \mid (\exists j) \langle w, j, i \rangle \in \pi_{WJI}(Q)\}$ , which by inspection is equal to  $\pi_{WI}(Q)$ .

*Claim 16:*  $\pi_{WI}(Q)$  is a surjection from  $W$  to  $I$ . This holds by Claims 14 and 15.

*Claim 17:*  $\bar{v}$  is a function from  $\pi_1 \bar{E}$ . Claim 16 implies that  $\pi_{WI}(Q)$  is a function from  $W$ , which by Claim 4 implies that  $\pi_{WI}(Q)$  is a function from  $\pi_1 \bar{E}$ , which by  $\bar{v}$ 's definition (34e) implies that  $\bar{v}$  is a function from  $\pi_1 \bar{E}$ .



*Claim 18:* [T5] holds. Because of Claim 17, it suffices to show  $\bar{\iota}$  is measurable (29). In other words, it suffices to show that

$$(\forall H \in \bar{\mathcal{H}}, w_1 \in H, w_2 \in H) \bar{\iota}(w_1) = \bar{\iota}(w_2).$$

By  $\bar{\mathcal{H}}$ 's definition (34c) and  $\bar{\iota}$ 's definition (34e), this is equivalent to

$$(\forall H \in \{W_j | j \in J\}, w_1 \in H, w_2 \in H) \pi_{WI}(Q)(w_1) = \pi_{WI}(Q)(w_2),$$

which, because  $\langle W_j \rangle_{j \in J}$  is an injectively indexed partition by Proposition 3.2(a $\Rightarrow$ c) and axiom [Pj $\leftarrow$ w], is equivalent to

$$(\forall j \in J, w_1 \in W_j, w_2 \in W_j) \pi_{WI}(Q)(w_1) = \pi_{WI}(Q)(w_2),$$

which by Claim 15 is equivalent to

$$(\forall j \in J, w_1 \in W_j, w_2 \in W_j) \pi_{JI}(Q) \circ \pi_{WJ}(Q)(w_1) = \pi_{JI}(Q) \circ \pi_{WJ}(Q)(w_2).$$

Now, to show this, take a situation  $j \in J$  and two decision nodes  $w_1$  and  $w_2$  in its information set  $W_j$ . Then  $w_1 \in W_j$  and Lemma B.2(a) imply  $\langle w_1, j \rangle \in \pi_{WJ}(Q)$ , which by axiom [Pj $\leftarrow$ w] implies  $\pi_{WJ}(Q)(w_1) = j$ . By the same reasoning,  $\pi_{WJ}(Q)(w_2) = j$ . Hence  $\pi_{WJ}(Q)(w_1) = \pi_{WJ}(Q)(w_2)$ , which by axiom [Pi $\leftarrow$ j] implies the desired equality.

*Claim 19:*  $\bar{I} = I$ . In steps,  $\bar{I}$  by its definition (28) is equal to the range of  $\bar{\iota}$ , which by  $\bar{\iota}$ 's definition (34e) is equal to the range of  $\pi_{WI}(Q)$ , which by Claim 16 is equal to  $I$ .

*Claim 20:* [T6] holds. Definition (34f) sets  $\bar{u}$  equal to  $u$ , which by the definition of a pentaform game (Definition 3.5) has the form  $\langle u_i: \mathcal{Z} \rightarrow \mathbb{R} \rangle_{i \in I}$ , which is identical to the form  $\langle u_i: \bar{\mathcal{Z}} \rightarrow \mathbb{R} \rangle_{i \in \bar{I}}$  because  $\bar{\mathcal{Z}} = \mathcal{Z}$  by Claim 7(c) and because  $\bar{I} = I$  by Claim 19.

*Claim 21:*  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u})$  is a traditional game. This follows from Definition 5.1 and Claims 2, 8, 10, 13, 18, and 20.  $\square$

**Lemma D.3.** *Let  $(X, E, \mathcal{H}, \alpha, \iota, u)$  be a traditional game. Then  $\text{TP}(X, E, \mathcal{H}, \alpha, \iota, u) = (X, E, \mathcal{H}, \alpha, \iota, u)$ .*

*Proof.* Let  $(\hat{Q}, \hat{u}) = \text{P}(X, E, \mathcal{H}, \alpha, \iota, u)$ , which by Theorem 5.2 is a pentaform game with information-set situations. Next let  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u}) = \text{T}(\hat{Q}, \hat{u})$ . Since  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u}) = \text{TP}(X, E, \mathcal{H}, \alpha, \iota, u)$  by inspection, it suffices to show  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u}) = (X, E, \mathcal{H}, \alpha, \iota, U)$ . This is done, one component at a time, by Claims 1–6.

*Claim 1:*  $\bar{X} = X$ . Note  $\bar{X}$  by its definition (34a) is  $\hat{W} \cup \hat{Y}$ , which by Lemma D.1(c) is  $X$ .

*Claim 2:*  $\bar{E} = E$ . Note  $\bar{E}$  by its definition (34b) is  $\pi_{WY}(\hat{Q})$ , which by Lemma D.1(d) is  $E$ .

*Claim 3:*  $\bar{\mathcal{H}} = \mathcal{H}$ . By the proof's first sentence,  $(\hat{Q}, \hat{u})$  has information-set situations (32). In other words,  $(\forall j \in \hat{J}) \hat{W}_j = j$ . Then in steps,  $\bar{\mathcal{H}}$  by its definition (34c) is  $\{\hat{W}_j | j \in \hat{J}\}$ , which by the previous sentence is  $\{j | j \in \hat{J}\}$ , which reduces to  $\hat{J}$ , which by Lemma D.1(b) is  $\mathcal{H}$ .

*Claim 4:*  $\bar{\alpha} = \alpha$ . Note  $\bar{\alpha}$  by its definition (34d) is  $\{ \langle \langle w, y \rangle, a \rangle \mid \langle w, y, a \rangle \in \pi_{WYA}(\hat{Q}) \}$ , which by Lemma D.1(e) is  $\alpha$ .

*Claim 5:*  $\bar{\iota} = \iota$ . Note  $\bar{\iota}$  by its definition (34e) is  $\pi_{WI}(\hat{Q})$ , which by Lemma D.1(f) is  $\iota$ .

*Claim 6:*  $\bar{u} = u$ . Note  $\bar{u}$  by its definition (34f) is  $\hat{u}$ , which by its definition (31b) is  $u$ .  $\square$

**Lemma D.4.** *Suppose  $(Q, u)$  is a pentaform game with information-set situations. Then  $\text{PT}(Q, u) = (Q, u)$ .*

*Proof.* Let  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u}) = \text{T}(Q, u)$ , which by Theorem 5.3 is a traditional game. Next let  $(\hat{Q}, \hat{u}) = \text{P}(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u})$ . Since  $(\hat{Q}, \hat{u}) = \text{PT}(Q, u)$  by inspection, it suffices to show that  $(\hat{Q}, \hat{u}) = (Q, u)$ . Definitions (31b) and (34f) imply  $\hat{u} = \bar{u} = u$ . Thus it suffices to show  $\hat{Q} = Q$ . This is Claim 4 below. (Looking ahead, claims 1–3 concern the starting point  $Q$  and the midpoint  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota})$  but not end point  $\hat{Q}$ .)

*Claim 1:*  $(\forall \langle j, w \rangle \in \pi_{JW}(Q)) j = \bar{H}_w$  (where  $\bar{H}_w$  is the information set in  $\bar{\mathcal{H}}$  that contains  $w$ , as defined below (31)). Take an original situation/decision-node couple  $\langle j, w \rangle \in \pi_{JW}(Q)$ . Easily, projection implies  $j \in \pi_J(Q)$ , which by abbreviation (8) implies [a]  $j \in J$ . Further, the assumption  $\langle j, w \rangle \in \pi_{JW}(Q)$  and Lemma B.2(a) imply [b]  $w \in W_j$ .

Note that [a] and  $\bar{\mathcal{H}}$ 's definition (34c) imply  $W_j \in \bar{\mathcal{H}}$ , that is, that the original pentaform information set  $W_j$  is a traditional information set in  $\bar{\mathcal{H}}$ . Thus since  $\bar{\mathcal{H}}$  is a partition by [T2], [b] implies that  $W_j$  is the member of  $\bar{\mathcal{H}}$  that contains  $w$ . In other words, [c]  $W_j = \bar{H}_w$ . Meanwhile, since  $(Q, u)$  has information-set situations (32) by assumption,  $W_j = j$ . This and [c] imply  $j = \bar{H}_w$ .

*Claim 2:*  $Q \subseteq \{ \langle \bar{\iota}(w), \bar{H}_w, w, \bar{\alpha}(w, y), y \rangle \mid \langle w, y \rangle \in \bar{E} \}$ . (The following argument will letter its observations in an unusual way.) Take an original quintuple [q]  $\langle i, j, w, a, y \rangle \in Q$ . We begin with four preliminary observations. First, [q] implies  $\langle w, i \rangle \in \pi_{WI}(Q)$ , which by  $\bar{\iota}$ 's definition (34e) implies [i]  $i = \bar{\iota}(w)$ . Second, [q] implies  $\langle w, j \rangle \in \pi_{WJ}(Q)$ , which by Claim 1 implies [j]  $j = \bar{H}_w$ . Third, [q] implies  $\langle w, y, a \rangle \in \pi_{WYA}(Q)$ , which by  $\bar{\alpha}$ 's definition (34d) implies [a]  $a = \bar{\alpha}(w, y)$ . Fourth, [q] implies  $\langle w, y \rangle \in \pi_{WY}(Q)$ , which by  $\bar{E}$ 's definition (34b) implies [e]  $\langle w, y \rangle \in \bar{E}$ . In conclusion, [i], [j], and [a] imply  $\langle i, j, w, a, y \rangle = \langle \bar{\iota}(w), \bar{H}_w, w, \bar{\alpha}(w, y), y \rangle$ , which with [e] completes the argument.

*Claim 3:*  $Q = \{ \langle \bar{\iota}(w), \bar{H}_w, w, \bar{\alpha}(w, y), y \rangle \mid \langle w, y \rangle \in \bar{E} \}$ . Claim 2 shows the forward inclusion. From another perspective, Claim 2 shows that the set  $Q$  is a subset of a function from  $\bar{E}$  (to be clear, the function's argument  $\langle w, y \rangle \in \bar{E}$  appears in the third and fifth coordinates, and the function takes each  $\langle w, y \rangle \in \bar{E}$  to the triple  $\langle \bar{\iota}(w), \bar{H}_w, \bar{\alpha}(w, y) \rangle$  which appears in the first, second, and fourth coordinates). Thus the set  $Q$  and the function are equal if the projection  $\pi_{WY}(Q)$  is equal to the domain  $\bar{E}$  (as opposed to merely being a subset of  $\bar{E}$ ). This holds by  $\bar{E}$ 's definition (34b).

*Claim 4:*  $Q = \hat{Q}$ . The right-hand side of Claim 3's equality is equal to  $\hat{Q}$  by  $\hat{Q}$ 's definition (31a).  $\square$

**Proof D.5 (for Theorem 5.4).** Theorem 5.2 shows that  $\mathsf{P}$  maps traditional games to pentaform games with information-set situations. Hence the theorem follows from Lemmas D.3 and D.4.  $\square$

**Proof D.6 (for Propositions 5.5 and 5.6).** As suggested in the text, Proposition 5.6 is more general than Proposition 5.5.

*Proposition 5.6.* By assumption,  $(X, E, \mathcal{H}, \alpha, \iota, u) = \mathsf{T}(\hat{Q}, \hat{u})$ . Parts [1], [2], [10\*], [11], [14], and [16] are implied by definitions (34a)–(34f), respectively. Parts [3]–[9], [12]–[13], and [15] are implied by Lemma D.2(b)–(k), respectively. (Note that  $(X, E, \mathcal{H}, \alpha, \iota, u)$  and  $(\hat{Q}, \hat{u})$  here and in Proposition 5.6 correspond to  $(\bar{X}, \bar{E}, \bar{\mathcal{H}}, \bar{\alpha}, \bar{\iota}, \bar{u})$  and  $(Q, u)$  in definition (34) and Lemma D.2.)

*Proposition 5.5.* By assumption,  $\mathsf{P}(X, E, \mathcal{H}, \alpha, \iota, u) = (\hat{Q}, \hat{u})$ . Thus Theorem 5.4 implies that  $(\hat{Q}, \hat{u})$  is a pentaform game with information-set situations and that  $(X, E, \mathcal{H}, \alpha, \iota, u) = \mathsf{T}(\hat{Q}, \hat{u})$ . Hence the previous paragraph (that is, Proposition 5.6) implies conclusions [1]–[9], [10\*], and [11]–[16]. Since  $(\hat{Q}, \hat{u})$  has information-set situations (32), conclusion [10\*] implies conclusion [10].  $\square$

## REFERENCES

- ALÓS-FERRER, C., AND K. RITZBERGER (2016): *The Theory of Extensive Form Games*. Springer.
- (2017): “Characterizations of Perfect Recall,” *International Journal of Game Theory*, 46, 311–326.
- ALPERN, S. (1988): “Games with Repeated Decisions,” *SIAM Journal of Optimization and Control*, 26, 468–477.
- BANG-JENSEN, J., AND G. Z. GUTIN (2009): *Digraphs: Theory, Algorithms and Applications, Second Edition*. Springer.
- BLACKBURN, P., M. DE RIJKE, AND Y. VENEMA (2001): *Modal Logic*. Cambridge.
- BOLT, J., J. HEDGES, AND P. ZAHN (2023): “Bayesian Open Games,” *Compositionality*, 5(9), 46 pages, arXiv:1910.03656v2.
- CAPUCCI, M., N. GHANI, J. LEDENT, AND F. NORDVALL FORSBERG (2022): “Translating Extensive Form Games to Open Games with Agency,” in *Fourth International Conference on Applied Category Theory (ACT 2021)*, ed. by K. Kishida, EPTCS, 372, pp. 221–234. doi:10.4204/EPTCS.372.16.
- DOCKNER, E., S. JØRGENSEN, N. V. LONG, AND G. SORGER (2000): *Differential Games in Economics and Management Science*. Cambridge.
- ENDERTON, H. B. (1977): *Elements of Set Theory*. Academic Press.
- GHANI, N., C. KUPKE, A. LAMBERT, AND F. NORDVALL FORSBERG (2018): “A Composition Treatment of Iterated Open Games,” *Theoretical Computer Science*, 741, 48–57.
- HALMOS, P. R. (1974): *Naive Set Theory*. Springer.
- HILLAS, J., AND D. KVASOV (2020): “Backward Induction in Games without Perfect Recall,” *Games and Economic Behavior*, 124, 207–218.
- ISELL, J. R. (1957): “Finitary Games,” in *Contributions to the Theory of Games, Volume III*, ed. by M. Dresher, A. W. Tucker, and P. Wolfe, pp. 79–96. Princeton.

- KLINE, J. J., AND S. LUCKRAZ (2016): “Equivalence between Graph-Based and Sequence-Based Extensive Form Games,” *Economic Theory Bulletin*, 4, 85–94.
- KNUTH, D. E. (1997): *The Art of Computer Programming: Volume 1 Fundamental Algorithms, Third Edition*. Addison-Wesley.
- KREPS, D. M., AND R. WILSON (1982): “Sequential Equilibria,” *Econometrica*, 50, 863–894.
- KUHN, H. W. (1950): “Extensive Games,” *Proceedings of the National Academy of Sciences*, 36(10), 570–576.
- (1953): “Extensive Games and the Problem of Information,” in *Contributions to the Theory of Games, Volume II*, ed. by H. W. Kuhn, and A. W. Tucker, pp. 193–216. Princeton, reprinted in *Classics in Game Theory*, ed. by H. W. Kuhn, pp. 46–68, Princeton (1997).
- LUENBERGER, D. G. (1979): *Introduction to Dynamic Systems*. Wiley.
- MAS-COLELL, A., M. D. WHINSTON, AND J. R. GREEN (1995): *Microeconomic Theory*. Oxford.
- MASCHLER, M., E. SOLAN, AND S. ZAMIR (2013): *Game Theory*. Cambridge University Press.
- MERTENS, J.-F. (2002): “Stochastic Games,” in *Handbook of Game Theory, Volume 3*, ed. by R. J. Aumann, and S. Hart, pp. 1809–1832. North-Holland.
- MYERSON, R. B. (1991): *Game Theory: Analysis of Conflict*. Harvard.
- OKADA, A. (1987): “Complete Inflation and Perfect Recall in Extensive Games,” *International Journal of Game Theory*, 16, 85–91.
- OSBORNE, M. J., AND A. RUBINSTEIN (1994): *A Course in Game Theory*. MIT.
- PEREA, A. (2001): *Rationality in Extensive Form Games*. Kluwer.
- PICCIONE, M., AND A. RUBINSTEIN (1997): “On the Interpretation of Decision Problems with Imperfect Recall,” *Games and Economic Behavior*, 20, 3–24.
- RITZBERGER, K. (1999): “Recall in Extensive Form Games,” *International Journal of Game Theory*, 28, 69–87.
- (2002): *Foundations of Non-Cooperative Game Theory*. Oxford.
- SELTEN, R. (1975): “Reexamination of the Perfectness Concept for Equilibrium Points in Extensive Games,” *International Journal of Game Theory*, 4, 25–55.
- STREUFERT, P. A. (2018): “The Category of Node-and-Choice Preforms for Extensive-Form Games,” *Studia Logica (Special Issue: Logics for Social Behaviour)*, 106, 1001–1064.
- (2019): “Equivalences among Five Game Specifications, including a New Specification whose Nodes are Sets of Past Choices,” *International Journal of Game Theory*, 48, 1–32.
- (2021): “A Category for Extensive-Form Games,” arXiv:2105.11398, also Western University, Department of Economics Research Report Series 2021-2, 60 pages.
- (2023): “Dynamic Programming for Pure-Strategy Subgame Perfection in an Arbitrary Game,” arXiv:2302.03855v3, 56 pages.
- TUTTE, W. T. (1984): *Graph Theory*. Addison-Wesley.

- VAN DAMME, E. (1991): *Stability and Perfection of Nash Equilibria, Second, Revised and Enlarged Edition*. Springer-Verlag.
- VON NEUMANN, J., AND O. MORGENSTERN (1944): *Theory of Games and Economic Behavior*. Princeton.