

Nonlinear first order PDEs reducible to autonomous form polynomially homogeneous in the derivatives

M. Gorgone and F. Oliveri

MIFT Department, University of Messina
Viale F. Stagno d'Alcontres 31, 98166 Messina, Italy
mgorgone@unime.it; foliveri@unime.it

Published in *J. Geom. Phys.* **113**, 53–64 (2017).

Abstract

It is proved a theorem providing necessary and sufficient conditions enabling one to map a nonlinear system of first order partial differential equations, polynomial in the derivatives, to an equivalent autonomous first order system polynomially homogeneous in the derivatives. The result is intimately related to the symmetry properties of the source system, and the proof, involving the use of the canonical variables associated to the admitted Lie point symmetries, is constructive. First order Monge–Ampère systems, either with constant coefficients or with coefficients depending on the field variables, where the theorem can be successfully applied, are considered.

Keywords. Lie symmetries; First order Monge–Ampère systems; Transformation to quasilinear form.

1 Introduction

Lie group analysis [1, 2, 3, 4, 5, 6, 7, 8] provides a unified and elegant algorithmic framework to a deep understanding and fruitful handling of differential

equations. It is known that Lie point symmetries admitted by ordinary differential equations allow for their order lowering and possibly reducing them to quadrature, whereas in the case of partial differential equations they can be used for the determination of special (invariant) solutions of initial and boundary value problems. Also, the Lie symmetries are important ingredients in the derivation of conserved quantities, or in the construction of relations between different differential equations that turn out to be equivalent [8, 9, 10, 11, 12, 13, 14, 15, 16]. Lie point symmetries of differential equations, in fact, can be used to construct a mapping from a given (source) system of differential equations to another (target) suitable system; if we consider one-to-one (invertible) point mappings, then a one-to-one correspondence between Lie point symmetries admitted by the source and target system of differential equations arises. In other words, the Lie algebra of infinitesimal operators of the target system of differential equations has to be isomorphic to the Lie algebra of infinitesimal operators of the source system of differential equations. This property has been used to give necessary and sufficient conditions for reducing a system of partial differential equations to autonomous form [12, 13], a system of first order nonlinear partial differential equations to linear form [9, 10, 11, 13], nonautonomous and/or nonhomogeneous quasilinear systems of partial differential equations to autonomous and homogeneous form [14, 16]. In particular, in [16], it has been proved a theorem providing the necessary and sufficient conditions in order to map a general first order quasilinear system of partial differential equations, say

$$\sum_{i=1}^n A^i(\mathbf{x}, \mathbf{u}) \frac{\partial \mathbf{u}}{\partial x_i} = \mathbf{B}(\mathbf{x}, \mathbf{u}), \quad (1)$$

where $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{u} \in \mathbb{R}^m$, A^i are $m \times m$ matrices with entries depending at most on \mathbf{x} and \mathbf{u} , and the source term $\mathbf{B} \in \mathbb{R}^m$ depends at most on \mathbf{x} and \mathbf{u} too, into a first order quasilinear homogeneous and autonomous system. This reduction, when it is possible, is performed by an invertible point transformation like

$$\mathbf{z} = \mathbf{Z}(\mathbf{x}), \quad \mathbf{w} = \mathbf{W}(\mathbf{x}, \mathbf{u}), \quad (2)$$

which preserves the quasilinear structure of the system, and whose construction is algorithmically suggested by the Lie symmetries admitted by (1).

In this paper, we consider a general nonlinear system of first order partial differential equations involving the derivatives of the unknown variables in

polynomial (of degree greater than 1) form, and establish a theorem giving necessary and sufficient conditions in order to map it to an autonomous system which is polynomially homogeneous in the derivatives.

In some relevant situations, *e.g.*, Monge–Ampère systems, the target system results to be quasilinear, but there are cases where the system we obtain is polynomially homogeneous in the derivatives but not quasilinear. This means that the conditions of the theorem are only necessary for the reduction of a *nonlinear* first order system to autonomous and homogeneous *quasilinear* form [17].

The main difference of the theorem presented in this paper with the similar one proved in [16] (concerned with the transformation of a general first order quasilinear system of partial differential equations into a first order quasilinear homogeneous and autonomous system) consists in the possibility of admitting now an invertible point transformation like

$$\mathbf{z} = \mathbf{Z}(\mathbf{x}, \mathbf{u}), \quad \mathbf{w} = \mathbf{W}(\mathbf{x}, \mathbf{u}), \quad (3)$$

i.e., a mapping where the new independent variables \mathbf{z} are allowed to depend also on the old dependent ones.

The plan of the paper is the following. In Section 2, the theorem giving necessary and sufficient conditions for the existence of an invertible mapping linking a nonlinear system of first order partial differential equations which is polynomial in the derivatives to an autonomous system polynomially homogeneous in the derivatives is proved. In Section 3, the theorem is applied to various general first order Monge–Ampère systems. Finally, Section 4 contains some concluding remarks.

2 Main result

Let us consider a general system of first order partial differential equations

$$\Delta(\mathbf{x}, \mathbf{u}, \mathbf{u}^{(1)}) = 0, \quad (4)$$

where $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{u}^{(1)} \in \mathbb{R}^{mn}$ are the independent variables, the dependent variables, and the first order partial derivatives, respectively. In particular, in the following we consider systems (4) composed by equations which are polynomial in the derivatives, with coefficients depending at most

on \mathbf{x} and \mathbf{u} , *i.e.* systems made by equations of the form

$$\sum_{|\alpha|, |\mathbf{j}|=1}^{N_s} A_{\alpha\mathbf{j}}^s(\mathbf{x}, \mathbf{u}) \prod_{k=1}^{|\alpha|} \frac{\partial u_{\alpha_k}}{\partial x_{j_k}} + B^s(\mathbf{x}, \mathbf{u}) = 0, \quad s = 1, \dots, m, \quad (5)$$

where α is the multi-index $(\alpha_1, \dots, \alpha_r)$, \mathbf{j} the multi-index (j_1, \dots, j_r) , $\alpha_k = 1, \dots, m$, $j_k = 1, \dots, n$, N_s are integers, and $A_{\alpha\mathbf{j}}^s(\mathbf{x}, \mathbf{u})$, $B^s(\mathbf{x}, \mathbf{u})$ smooth functions of their arguments.

The aim is to determine necessary and sufficient conditions for the construction of an invertible point transformation

$$\mathbf{z} = \mathbf{Z}(\mathbf{x}, \mathbf{u}), \quad \mathbf{w} = \mathbf{W}(\mathbf{x}, \mathbf{u}), \quad (6)$$

mapping (5) into an equivalent autonomous system which is homogeneous polynomial in the derivatives $\mathbf{w}^{(1)}$, *i.e.*, made by equations of the form

$$\sum_{|\alpha|, |\mathbf{j}|=\overline{N}_s} \tilde{A}_{\alpha\mathbf{j}}^s(\mathbf{w}) \prod_{k=1}^{\overline{N}_s} \frac{\partial w_{\alpha_k}}{\partial z_{j_k}} = 0, \quad s = 1, \dots, m, \quad (7)$$

for some integers \overline{N}_s ; of course, it may occur that the target system turns out to be linear in the derivatives, *i.e.*, $\overline{N}_s = 1$ ($s = 1, \dots, m$), whereupon we have an autonomous and homogeneous quasilinear system.

The following lemma, guarantees that an invertible point transformation like (6) preserves the polynomial structure in the derivatives.

Lemma 1 *Given a first order system of partial differential equations like (5) which is polynomial in the derivatives, then an invertible point transformation like (6) produces a first order system which is still polynomial in the derivatives.*

Proof. Straightforward, by using the chain rule. □

Theorem 1 *The nonlinear first order system of partial differential equations polynomial in the derivatives*

$$\sum_{|\alpha|, |\mathbf{j}|=1}^{N_s} A_{\alpha\mathbf{j}}^s(\mathbf{x}, \mathbf{u}) \prod_{k=1}^{|\alpha|} \frac{\partial u_{\alpha_k}}{\partial x_{j_k}} + B^s(\mathbf{x}, \mathbf{u}) = 0, \quad s = 1, \dots, m, \quad (8)$$

is mapped by an invertible point transformation, say

$$\mathbf{z} = \mathbf{Z}(\mathbf{x}, \mathbf{u}), \quad \mathbf{w} = \mathbf{W}(\mathbf{x}, \mathbf{u}), \quad (9)$$

to the equivalent nonlinear first order autonomous system having homogeneous polynomial form, say

$$\sum_{|\boldsymbol{\alpha}|, |\mathbf{j}| = \overline{N}_s} \tilde{A}_{\boldsymbol{\alpha}\mathbf{j}}^s(\mathbf{w}) \prod_{k=1}^{\overline{N}_s} \frac{\partial w_{\alpha_k}}{\partial z_{j_k}} = 0, \quad s = 1, \dots, m, \quad (10)$$

for some integers \overline{N}_s , if and only if there exists an $(n + 1)$ -dimensional subalgebra of the Lie algebra of point symmetries, admitted by system (8), spanned by the vector fields

$$\Xi_i = \sum_{j=1}^n \xi_i^j(\mathbf{x}, \mathbf{u}) \frac{\partial}{\partial x_j} + \sum_{\alpha=1}^m \eta_i^\alpha(\mathbf{x}, \mathbf{u}) \frac{\partial}{\partial u_\alpha}, \quad i = 1, \dots, n + 1, \quad (11)$$

generating a distribution of rank $(n + 1)$, and such that

$$\begin{aligned} [\Xi_i, \Xi_j] &= 0, & i = 1, \dots, n - 1, & \quad i < j \leq n, \\ [\Xi_i, \Xi_{n+1}] &= \Xi_i, & i = 1, \dots, n. \end{aligned} \quad (12)$$

Moreover, the variables \mathbf{w} , which by construction are invariants of Ξ_1, \dots, Ξ_n , have to be invariant with respect to Ξ_{n+1} too.

Proof. Suppose the conditions of Theorem 1 are satisfied, and so the system (8) admits an $(n + 1)$ -dimensional algebra as subalgebra of the algebra of its Lie point symmetries generating a distribution of rank $(n + 1)$ and verifying the structure conditions (12). Let us introduce a set of canonical variables for the vector field Ξ_1 , say

$$y_i^1 \quad (i = 1, \dots, n), \quad v_\alpha^1 \quad (\alpha = 1, \dots, m),$$

such that

$$\Xi_1 y_1^1 = 1, \quad \Xi_1 y_{i_1}^1 = 0, \quad \Xi_1 v_\alpha^1 = 0$$

($i_1 = 2, \dots, n$); as a consequence, Ξ_1 takes the form

$$\Xi_1 = \frac{\partial}{\partial y_1^1},$$

i.e., it corresponds to a translation in the variable y_1^1 .

Since $[\Xi_1, \Xi_2] = 0$, it is

$$\Xi_1(\Xi_2 y_i^1) = \Xi_2(\Xi_1 y_i^1) = 0, \quad \Xi_1(\Xi_2 v_\alpha^1) = \Xi_2(\Xi_1 v_\alpha^1) = 0, \quad (13)$$

for $i = 1, \dots, n$ and $\alpha = 1, \dots, m$. Thus, the infinitesimals of Ξ_2 , represented in terms of the canonical variables of Ξ_1 , will depend upon the invariants of Ξ_1 only, *i.e.*, Ξ_2 writes as

$$\Xi_2 = \sum_{i=1}^n \Theta_i^2(y_{j_1}^1, v_\beta^1) \frac{\partial}{\partial y_i^1} + \sum_{\alpha=1}^m \Lambda_\alpha^2(y_{j_1}^1, v_\beta^1) \frac{\partial}{\partial v_\alpha^1} \quad (14)$$

($j_1 = 2, \dots, n, \beta = 1, \dots, m$).

If $\Theta_1^2 \neq 0$ we need to replace y_1^1 with

$$y_1^1 + \varphi_1^1(y_{j_1}^1, v_\beta^1),$$

where the function φ_1^1 satisfies

$$\Theta_1^2(y_{j_1}^1, v_\beta^1) + \sum_{i_1=2}^n \Theta_{i_1}^2(y_{j_1}^1, v_\beta^1) \frac{\partial \varphi_1^1}{\partial y_{i_1}^1} + \sum_{\alpha=1}^m \Lambda_\alpha^2(y_{j_1}^1, v_\beta^1) \frac{\partial \varphi_1^1}{\partial v_\alpha^1} = 0.$$

That enables us to write Ξ_1 and Ξ_2 as follows

$$\Xi_1 = \frac{\partial}{\partial y_1^1}, \quad \Xi_2 = \sum_{i_1=2}^n \Theta_{i_1}^2(y_{j_1}^1, v_\beta^1) \frac{\partial}{\partial y_{i_1}^1} + \sum_{\alpha=1}^m \Lambda_\alpha^2(y_{j_1}^1, v_\beta^1) \frac{\partial}{\partial v_\alpha^1}, \quad (15)$$

where $j_1 = 2, \dots, n$.

Introducing the canonical variables

$$y_1^2 = y_1^1, \quad y_2^2, \quad y_{i_2}^2 \quad (i_2 = 3, \dots, n), \quad v_\alpha^2 \quad (\alpha = 1, \dots, m),$$

such that

$$\Xi_2 y_2^2 = 1, \quad \Xi_2 y_{i_2}^2 = 0, \quad \Xi_2 v_\alpha^2 = 0$$

($i_1 = 2, \dots, n$), it is obtained

$$\Xi_2 = \frac{\partial}{\partial y_2^2}.$$

Continuing inductively for $k = 2, \dots, n - 1$, since Ξ_{k+1} commutes with Ξ_1, \dots, Ξ_k , in terms of the canonical variables

$$y_i^k \quad (i = 1, \dots, n), \quad v_\alpha^k \quad (\alpha = 1, \dots, m),$$

we have

$$\begin{aligned} \Xi_1 &= \frac{\partial}{\partial y_1^k}, & \Xi_2 &= \frac{\partial}{\partial y_2^k}, & \dots, & & \Xi_k &= \frac{\partial}{\partial y_k^k}, \\ \Xi_{k+1} &= \sum_{i=1}^n \Theta_i^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial}{\partial y_i^k} + \sum_{\alpha=1}^m \Lambda_\alpha^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial}{\partial v_\alpha^k}, \end{aligned} \quad (16)$$

where $j_k = k + 1, \dots, n$. If $\Theta_\ell^{k+1} \neq 0$, for $\ell = 1, \dots, k$, we need to replace the variable y_ℓ^k with

$$y_\ell^k + \varphi_\ell^k(y_{j_k}^k, v_\beta^k),$$

where the function φ_ℓ^k satisfies

$$\Theta_\ell^{k+1}(y_{j_k}^k, v_\beta^k) + \sum_{i_k=k+1}^n \Theta_{i_k}^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial \varphi_\ell^k}{\partial y_{i_k}^k} + \sum_{\alpha=1}^m \Lambda_\alpha^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial \varphi_\ell^k}{\partial v_\alpha^k} = 0,$$

so that Ξ_{k+1} writes as

$$\Xi_{k+1} = \sum_{i_k=k+1}^n \Theta_{i_k}^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial}{\partial y_{i_k}^k} + \sum_{\alpha=1}^m \Lambda_\alpha^{k+1}(y_{j_k}^k, v_\beta^k) \frac{\partial}{\partial v_\alpha^k}; \quad (17)$$

hence, we may construct the canonical variables

$$y_1^{k+1} = y_1^k, \dots, y_k^{k+1} = y_k^k, \quad y_{i_k}^{k+1} \quad (i_k = k+1, \dots, n), \quad v_\alpha^{k+1} \quad (\alpha = 1, \dots, m),$$

related to the operator Ξ_{k+1} , such that the latter writes as

$$\Xi_{k+1} = \frac{\partial}{\partial y_{k+1}^{k+1}}.$$

The complete application of the described algorithm enables us to write each operator Ξ_i in the form

$$\Xi_i = \frac{\partial}{\partial z_i}, \quad i = 1, \dots, n,$$

and the new independent and dependent variables are $z_i = y_i^n$ ($i = 1, \dots, n$), $w_\alpha = v_\alpha^n$ ($\alpha = 1, \dots, m$), respectively.

Therefore, what we have obtained is a variable transformation like (9) allowing to write the system (8) in autonomous form.

Finally, since $[\Xi_i, \Xi_{n+1}] = \Xi_i$ ($i = 1, \dots, n$), it is

$$\begin{aligned}\Xi_i(\Xi_{n+1}z_j) &= \Xi_{n+1}(\Xi_i z_j) + \Xi_i z_j = \delta_{ij}, \\ \Xi_i(\Xi_{n+1}w_\alpha) &= \Xi_{n+1}(\Xi_i w_\alpha) + \Xi_i w_\alpha = 0;\end{aligned}\tag{18}$$

where δ_{ij} is the Kronecker symbol; these relations, together with the hypothesis that the variables w_α ($\alpha = 1, \dots, m$) are invariant with respect to Ξ_{n+1} , allow the vector field Ξ_{n+1} to gain the representation

$$\Xi_{n+1} = \sum_{j=1}^n z_j \frac{\partial}{\partial z_j}.\tag{19}$$

As a consequence, since the resulting system, written in the variables \mathbf{z} and \mathbf{w} , is autonomous and polynomial in the derivatives, and is invariant with respect to a uniform scaling of all independent variables, then it necessarily must be polynomially homogeneous in the derivatives, *i.e.*, it has the form (10).

The condition that the symmetries generate a distribution of rank $(n+1)$, whence the vector fields spanning the n -dimensional Abelian Lie subalgebra generate a distribution of rank n , ensures that we may construct the complete set of the new independent variables \mathbf{z} .

Conversely, if the nonautonomous and/or nonhomogeneous system (8) can be mapped by the invertible point transformation (9) to the autonomous system polynomially homogeneous in the derivatives (10), then, since the latter admits the n vector fields $\frac{\partial}{\partial z_i}$, spanning an n -dimensional Abelian Lie algebra, and the vector field $\sum_{j=1}^n z_j \frac{\partial}{\partial z_j}$, then it follows that also the system (8) must admit $(n+1)$ Lie point symmetries with the requested algebraic structure. \square

Example 1 Let us consider the first order system made by the equations

$$\begin{aligned}
& \frac{\partial u_1}{\partial x_2} - \frac{\partial u_2}{\partial x_1} = 0, \\
& \kappa_1 \left(\frac{\partial u_1}{\partial x_2} \right)^4 + \left(\kappa_2 \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \kappa_3 \left(\frac{\partial u_1}{\partial x_2} \right)^2 \right) \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} \\
& + \left(\kappa_4 \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \kappa_5 \left(\frac{\partial u_1}{\partial x_2} \right)^2 \right) \frac{\partial u_1}{\partial x_1} + \left(\kappa_6 \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \kappa_7 \left(\frac{\partial u_1}{\partial x_2} \right)^2 \right) \frac{\partial u_1}{\partial x_2} \\
& + \left(\kappa_8 \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \kappa_9 \left(\frac{\partial u_1}{\partial x_2} \right)^2 \right) \frac{\partial u_2}{\partial x_2} + \kappa_{10} \left(\frac{\partial u_1}{\partial x_1} \right)^2 + \kappa_{11} \frac{\partial u_1}{\partial x_1} \frac{\partial u_1}{\partial x_2} \\
& + \kappa_{12} \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \kappa_{13} \left(\frac{\partial u_1}{\partial x_2} \right)^2 + \kappa_{14} \frac{\partial u_1}{\partial x_2} \frac{\partial u_2}{\partial x_2} + \kappa_{15} \left(\frac{\partial u_2}{\partial x_2} \right)^2 = 0,
\end{aligned} \tag{20}$$

with $u_1(x_1, x_2)$, $u_2(x_1, x_2)$ scalar functions, and $\kappa_i(u_1, u_2)$ ($i = 1, \dots, 15$) arbitrary smooth functions of the indicated arguments.

It can be easily ascertained that system (20) admits the Lie point symmetries spanned by the operators

$$\begin{aligned}
& \Xi_1 = \frac{\partial}{\partial x_1}, \quad \Xi_2 = \frac{\partial}{\partial x_2}, \\
& \Xi_3 = (x_1 - au_1 - bu_2) \frac{\partial}{\partial x_1} + (x_2 - bu_1 - cu_2) \frac{\partial}{\partial x_2},
\end{aligned} \tag{21}$$

a, b, c being constants, provided that the conditions

$$\begin{aligned}
& \kappa_1 - c^2 \kappa_{10} + bck_{11} - ack_{12} - b^2 \kappa_{13} + ab\kappa_{14} - a^2 \kappa_{15} = 0, \\
& \kappa_2 - c^2 \kappa_{10} + bck_{11} - ack_{12} - b^2 \kappa_{13} + ab\kappa_{14} - a^2 \kappa_{15} = 0, \\
& \kappa_3 + 2(c^2 \kappa_{10} - bck_{11} + ack_{12} + b^2 \kappa_{13} - ab\kappa_{14} + a^2 \kappa_{15}) = 0, \\
& \kappa_4 + 2c\kappa_{10} - b\kappa_{11} + a\kappa_{12} = 0, \\
& \kappa_5 - 2c\kappa_{10} + b\kappa_{11} - a\kappa_{12} = 0, \\
& \kappa_6 + c\kappa_{11} - 2b\kappa_{13} + a\kappa_{14} = 0, \\
& \kappa_7 - c\kappa_{11} + 2b\kappa_{13} - a\kappa_{14} = 0, \\
& \kappa_8 + c\kappa_{12} - b\kappa_{14} + 2a\kappa_{15} = 0, \\
& \kappa_9 - c\kappa_{12} + b\kappa_{14} - 2a\kappa_{15} = 0
\end{aligned} \tag{22}$$

are satisfied. Since

$$[\Xi_1, \Xi_2] = 0, \quad [\Xi_1, \Xi_3] = \Xi_1, \quad [\Xi_2, \Xi_3] = \Xi_2, \tag{23}$$

applying the theorem, we introduce the new independent and dependent variables

$$\begin{aligned} z_1 &= x_1 - au_1 - bu_2, & z_2 &= x_2 - bu_1 - cu_2, \\ w_1 &= u_1, & w_2 &= u_2, \end{aligned} \quad (24)$$

and the nonlinear system (20) reduces to

$$\begin{aligned} \frac{\partial w_1}{\partial z_2} - \frac{\partial w_2}{\partial z_1} &= 0, \\ \kappa_{10} \left(\frac{\partial w_1}{\partial z_1} \right)^2 + \kappa_{11} \frac{\partial w_1}{\partial z_1} \frac{\partial w_1}{\partial z_2} + \kappa_{12} \frac{\partial w_1}{\partial z_1} \frac{\partial w_2}{\partial z_2} \\ &+ \kappa_{13} \left(\frac{\partial w_1}{\partial z_2} \right)^2 + \kappa_{14} \frac{\partial w_1}{\partial z_2} \frac{\partial w_2}{\partial z_2} + \kappa_{15} \left(\frac{\partial w_2}{\partial z_2} \right)^2 = 0, \end{aligned} \quad (25)$$

i.e., reads as an autonomous system polynomially homogeneous in the derivatives.

We notice that system (25), by specializing the functions $\kappa_{10}, \dots, \kappa_{15}$ as follows,

$$\begin{aligned} \kappa_{10} &= -\kappa(1 + w_2^2)^2, \\ \kappa_{11} &= 4\kappa w_1 w_2 (1 + w_2^2), \\ \kappa_{12} &= 2((2 - \kappa)(1 + w_1^2 + w_2^2) - \kappa w_1^2 w_2^2), \\ \kappa_{13} &= -4(1 + w_1^2 + w_2^2 + \kappa w_1^2 w_2^2), \\ \kappa_{14} &= 4\kappa w_1 w_2 (1 + w_1^2), \\ \kappa_{15} &= -\kappa(1 + w_1^2)^2, \end{aligned} \quad (26)$$

is equivalent to the second order partial differential equation

$$\begin{aligned} &\kappa (1 + w_{z_2}^2)^2 w_{z_1 z_1}^2 - 4\kappa w_{z_1} w_{z_2} (1 + w_{z_2}^2) w_{z_1 z_1} w_{z_1 z_2} \\ &- 2((2 - \kappa)(1 + w_{z_1}^2 + w_{z_2}^2) - \kappa w_{z_1}^2 w_{z_2}^2) w_{z_1 z_1} w_{z_2 z_2} \\ &+ 4(1 + w_{z_1}^2 + w_{z_2}^2 + \kappa w_{z_1}^2 w_{z_2}^2) w_{z_1 z_2}^2 - 4\kappa w_{z_1} w_{z_2} (1 + w_{z_1}^2) w_{z_1 z_2} w_{z_2 z_2} \\ &+ \kappa (1 + w_{z_1}^2)^2 w_{z_2 z_2}^2 = 0, \end{aligned} \quad (27)$$

where $w_{z_1} = \frac{\partial w}{\partial z_1} = w_1$, $w_{z_2} = \frac{\partial w}{\partial z_2} = w_2$, and κ is an arbitrary function of w_{z_1} and w_{z_2} .

Considering a smooth surface in \mathbb{R}^3 with the metric $ds^2 = dz_1^2 + dz_2^2 + dw^2$,

and its Gaussian and mean curvature,

$$\begin{aligned} G &= \frac{w_{z_1 z_1} w_{z_2 z_2} - w_{z_1 z_2}^2}{(1 + w_{z_1}^2 + w_{z_2}^2)^2}, \\ H &= \frac{1}{2} \frac{(1 + w_{z_2}^2) w_{z_1 z_1} - 2w_{z_1} w_{z_2} w_{z_1 z_2} + (1 + w_{z_1}^2) w_{z_2 z_2}}{(1 + w_{z_1}^2 + w_{z_2}^2)^{3/2}}, \end{aligned} \quad (28)$$

respectively, equation (27) can be written as

$$G = \kappa H^2, \quad (29)$$

whereupon it should be $\kappa(w_{z_1}, w_{z_2}) \leq 1$. In the limit case $\kappa \equiv 1$, Eq. (29) characterizes a surface with all its points umbilic; it is known that a surface with all its point umbilic is a (open) domain of a plane or a sphere [18]. It is worth of being remarked that Eq. (29) with $\kappa \equiv 1$ is strongly Lie remarkable [19], since it is the unique second order partial differential equation uniquely characterized by the conformal Lie algebra in \mathbb{R}^3 [20].

Remark 1 Notice that the Example 1 provides a system polynomial in the derivatives which is transformed into a system polynomially homogeneous of degree 2 in the derivatives. In Section 3, we will analyze various first order Monge–Ampère systems, and show that they can be transformed to quasilinear form.

3 Applications

In this section, we provide some examples of first order nonlinear systems polynomial in the derivatives whose Lie symmetries satisfy the conditions of Theorem 1, and prove that they can be transformed under suitable conditions to autonomous first order systems having homogeneous polynomial form; the systems that will be considered are of Monge–Ampère type, and, remarkably, they are reduced to quasilinear (or linear) form.

In particular, we are concerned with the nonlinear first order systems of Monge–Ampère equations for the unknowns $u_\alpha(x_i)$ ($\alpha = 1, \dots, m$, $i = 1, \dots, n$). These systems have been characterized by Boillat in 1997 [21] by looking for the nonlinear first order systems possessing, as the quasilinear systems, the property of the linearity of the Cauchy problem. These systems are also completely exceptional [22, 23], and are made by equations which

are expressed as linear combinations (with coefficients depending at most on the independent and the dependent variables) of all minors extracted from the gradient matrix of $u_\alpha = u_\alpha(x_i)$.

Hereafter, to shorten the formulas, we denote with $u_{\alpha,i}$ the first order partial derivative of $u_\alpha(x_i)$ with respect to x_i , and with $w_{\alpha,i}$ the first order partial derivative of $w_\alpha(z_i)$ with respect to z_i ; moreover, we denote with $f_{i;\alpha}$ the first order partial derivative of the function f_i with respect to u_α (or w_α). In the following we limit ourselves to consider the coefficients of the Monge–Ampère systems at most functions of the field variables.

3.1 Case $m = n = 2$

Let us consider the nonlinear first order system of Monge–Ampère made by the equations

$$\kappa_0^i (u_{1,1}u_{2,2} - u_{1,2}u_{2,1}) + \kappa_1^i u_{1,1} + \kappa_2^i u_{1,2} + \kappa_3^i u_{2,1} + \kappa_4^i u_{2,2} + \kappa_5^i = 0 \quad (30)$$

($i = 1, 2$), with $u_1(x_1, x_2), u_2(x_1, x_2)$ scalar functions, and $\kappa_j^i(u_1, u_2)$ ($i = 1, 2; j = 0, \dots, 5$) arbitrary smooth functions of the indicated arguments.

The substitutions

$$u_1 \rightarrow u_1 + \alpha_{11}x_1 + \alpha_{12}x_2, \quad u_2 \rightarrow u_2 + \alpha_{21}x_1 + \alpha_{22}x_2, \quad (31)$$

where α_{ij} are arbitrary constants, produce a system with $\kappa_5^i = 0$ ($i = 1, 2$) provided that

$$\kappa_0^i (\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21}) + \kappa_1^i \alpha_{11} + \kappa_2^i \alpha_{12} + \kappa_3^i \alpha_{21} + \kappa_4^i \alpha_{22} + \kappa_5^i = 0 \quad (32)$$

($i = 1, 2$). Conditions (32) provide two constraints on the functional form of the coefficients so that not all systems can be written in a form where $\kappa_5^i = 0$; however, if the coefficients κ_j^i are constant, due to the arbitrariness of the constants α_{ij} , then (32) can always be satisfied whatever the values of the coefficients are.

It is easily recognized that system (30), now taken with $\kappa_5^i = 0$, admits the Lie point symmetries spanned by the operators

$$\Xi_1 = \frac{\partial}{\partial x_1}, \quad \Xi_2 = \frac{\partial}{\partial x_2}, \quad \Xi_3 = (x_1 - f_1) \frac{\partial}{\partial x_1} + (x_2 - f_2) \frac{\partial}{\partial x_2}, \quad (33)$$

where $f_i(u_1, u_2)$ ($i = 1, 2$) are arbitrary smooth functions of their arguments, provided that

$$\kappa_0^i + \kappa_1^i f_{2;2} - \kappa_2^i f_{1;2} - \kappa_3^i f_{2;1} + \kappa_4^i f_{1;1} = 0, \quad i = 1, 2. \quad (34)$$

The constraints (34), once we assign the 10 functions $k_j^i(u_1, u_2)$ ($i = 1, 2$, $j = 0, \dots, 4$), are the differential equations providing us the functional form of $f_1(u_1, u_2)$ and $f_2(u_1, u_2)$.

In the case where all the coefficients κ_j^i are constant, then the functions f_1 and f_2 are forced to be linear, *i.e.*,

$$f_1 = \beta_{11}u_1 + \beta_{12}u_2, \quad f_2 = \beta_{21}u_1 + \beta_{22}u_2, \quad (35)$$

β_{ij} being constants whose value is determined by the coefficients κ_j^i .

Since

$$[\Xi_1, \Xi_2] = 0, \quad [\Xi_1, \Xi_3] = \Xi_1, \quad [\Xi_2, \Xi_3] = \Xi_2, \quad (36)$$

we introduce the new variables

$$z_1 = x_1 - f_1, \quad z_2 = x_2 - f_2, \quad w_1 = u_1, \quad w_2 = u_2, \quad (37)$$

and the generators of the point symmetries write as

$$\Xi_1 = \frac{\partial}{\partial z_1}, \quad \Xi_2 = \frac{\partial}{\partial z_2}, \quad \Xi_3 = z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2}. \quad (38)$$

In terms of the new variables (37), the nonlinear system (30) becomes

$$\kappa_1^i w_{1,1} + \kappa_2^i w_{1,2} + \kappa_3^i w_{2,1} + \kappa_4^i w_{2,2} = 0, \quad (39)$$

i.e., reads as an autonomous and homogeneous quasilinear system. This system is linear if all the coefficients κ_j^i are constant; nevertheless, since it is a 2×2 homogeneous and autonomous quasilinear system, it can be written in linear form by means of the hodograph transformation also when the coefficients κ_j^i depend on u_1 and u_2 .

In conclusion, all Monge–Ampère systems with $m = n = 2$ can be reduced to a linear system when the coefficients κ_j^i are constant; on the contrary, when the coefficients depend upon u_1 and u_2 the reduction to the linear form is possible provided that the constraints (32) are satisfied.

3.2 Case $m = 2$, $n = 3$

By considering the gradient matrix of $u_\alpha(x_i)$ ($\alpha = 1, 2$, $i = 1, \dots, 3$)

$$H = \begin{pmatrix} u_{1,1} & u_{1,2} & u_{1,3} \\ u_{2,1} & u_{2,2} & u_{2,3} \end{pmatrix} \quad (40)$$

and its extracted minors

$$H^1 = \begin{vmatrix} u_{1,1} & u_{1,2} \\ u_{2,1} & u_{2,2} \end{vmatrix}, \quad H^2 = \begin{vmatrix} u_{1,1} & u_{1,3} \\ u_{2,1} & u_{2,3} \end{vmatrix}, \quad H^3 = \begin{vmatrix} u_{1,2} & u_{1,3} \\ u_{2,2} & u_{2,3} \end{vmatrix}, \quad (41)$$

the nonlinear first order system of Monge–Ampère is made by equations like

$$\begin{aligned} &\kappa_1^i H^1 + \kappa_2^i H^2 + \kappa_3^i H^3 \\ &+ \kappa_4^i u_{1,1} + \kappa_5^i u_{1,2} + \kappa_6^i u_{1,3} + \kappa_7^i u_{2,1} + \kappa_8^i u_{2,2} + \kappa_9^i u_{2,3} + \kappa_{10}^i = 0, \end{aligned} \quad (42)$$

$\kappa_j^i(u_\alpha)$ ($i = 1, 2$, $j = 1, \dots, 10$) arbitrary smooth functions of the indicated arguments.

The substitutions

$$u_1 \rightarrow u_1 + \alpha_{11}x_1 + \alpha_{12}x_2 + \alpha_{13}x_3, \quad u_2 \rightarrow u_2 + \alpha_{21}x_1 + \alpha_{22}x_2 + \alpha_{23}x_3, \quad (43)$$

where α_{ij} are arbitrary constants, produce a system with $\kappa_{10}^i = 0$ ($i = 1, 2$) provided that

$$\begin{aligned} &\kappa_1^i(\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21}) + \kappa_2^i(\alpha_{11}\alpha_{23} - \alpha_{13}\alpha_{21}) + \kappa_3^i(\alpha_{12}\alpha_{23} - \alpha_{13}\alpha_{22}) \\ &+ \kappa_4^i\alpha_{11} + \kappa_5^i\alpha_{12} + \kappa_6^i\alpha_{13} + \kappa_7^i\alpha_{21} + \kappa_8^i\alpha_{22} + \kappa_9^i\alpha_{23} + \kappa_{10}^i = 0 \end{aligned} \quad (44)$$

($i = 1, 2$). Actually, conditions (44) can always be satisfied when the coefficients κ_j^i are constant because of the arbitrariness of the constants α_{ij} .

This nonlinear system (42), now taken with $\kappa_{10}^i = 0$, admits the Lie point symmetries spanned by the operators

$$\begin{aligned} \Xi_1 &= \frac{\partial}{\partial x_1}, & \Xi_2 &= \frac{\partial}{\partial x_2}, & \Xi_3 &= \frac{\partial}{\partial x_3}, \\ \Xi_4 &= (x_1 - f_1) \frac{\partial}{\partial x_1} + (x_2 - f_2) \frac{\partial}{\partial x_2} + (x_3 - f_3) \frac{\partial}{\partial x_3}, \end{aligned} \quad (45)$$

where $f_i(u_1, u_2)$ ($i = 1, \dots, 3$) are arbitrary smooth functions of their arguments, provided that

$$\begin{aligned} &\kappa_1^i + \kappa_4^i f_{2;2} - \kappa_5^i f_{1;2} - \kappa_7^i f_{2;1} + \kappa_8^i f_{1;1} = 0, \\ &\kappa_2^i + \kappa_4^i f_{3;2} - \kappa_6^i f_{1;2} - \kappa_7^i f_{3;1} + \kappa_9^i f_{1;1} = 0, \\ &\kappa_3^i + \kappa_5^i f_{3;2} - \kappa_6^i f_{2;2} - \kappa_8^i f_{3;1} + \kappa_9^i f_{2;1} = 0. \end{aligned} \quad (46)$$

The six conditions (46) cannot be fulfilled for an arbitrary choice of the coefficients κ_j^i . In the simplest case, where the coefficients κ_j^i are constant, they can be always satisfied and the functions f_i must be linear:

$$f_1 = \beta_{11}u_1 + \beta_{12}u_2, \quad f_2 = \beta_{21}u_1 + \beta_{22}u_2, \quad f_3 = \beta_{31}u_1 + \beta_{32}u_2, \quad (47)$$

β_{ij} being arbitrary constants.

The symmetries (45) generate a 4–dimensional solvable Lie algebra,

$$[\Xi_i, \Xi_j] = 0, \quad [\Xi_i, \Xi_4] = \Xi_i, \quad (i, j = 1, 2, 3), \quad (48)$$

whereupon we may introduce the new variables

$$\begin{aligned} z_1 &= x_1 - f_1, & z_2 &= x_2 - f_2, & z_3 &= x_3 - f_3, \\ w_1 &= u_1, & w_2 &= u_2, \end{aligned} \quad (49)$$

and the generators of the point symmetries write as

$$\Xi_1 = \frac{\partial}{\partial z_1}, \quad \Xi_2 = \frac{\partial}{\partial z_2}, \quad \Xi_3 = \frac{\partial}{\partial z_3}, \quad \Xi_4 = z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3}. \quad (50)$$

In terms of the new variables (49), the nonlinear system (42) reduces to

$$\kappa_4^i w_{1,1} + \kappa_5^i w_{1,2} + \kappa_6^i w_{1,3} + \kappa_7^i w_{2,1} + \kappa_8^i w_{2,2} + \kappa_9^i w_{2,3} = 0, \quad (51)$$

i.e., reads as an autonomous and homogeneous quasilinear system.

3.3 Case $m = 3, n = 2$

By considering the gradient matrix of $u_\alpha(x_i)$ ($\alpha = 1, \dots, 3, i = 1, 2$)

$$H = \begin{pmatrix} u_{1,1} & u_{1,2} \\ u_{2,1} & u_{2,2} \\ u_{3,1} & u_{3,2} \end{pmatrix} \quad (52)$$

and its extracted minors

$$H^1 = \begin{vmatrix} u_{1,1} & u_{1,2} \\ u_{2,1} & u_{2,2} \end{vmatrix}, \quad H^2 = \begin{vmatrix} u_{1,1} & u_{1,2} \\ u_{3,1} & u_{3,2} \end{vmatrix}, \quad H^3 = \begin{vmatrix} u_{2,1} & u_{2,2} \\ u_{3,1} & u_{3,2} \end{vmatrix}, \quad (53)$$

the nonlinear first order system of Monge–Ampère is made by equations like

$$\begin{aligned} \kappa_1^i H^1 + \kappa_2^i H^2 + \kappa_3^i H^3 \\ + \kappa_4^i u_{1,1} + \kappa_5^i u_{1,2} + \kappa_6^i u_{2,1} + \kappa_7^i u_{2,2} + \kappa_8^i u_{3,1} + \kappa_9^i u_{3,2} + \kappa_{10}^i = 0, \end{aligned} \quad (54)$$

$\kappa_j^i(u_\alpha)$ ($i = 1, \dots, 3$, $j = 1, \dots, 10$) arbitrary smooth functions of the indicated arguments.

The substitutions

$$\begin{aligned} u_1 &\rightarrow u_1 + \alpha_{11}x_1 + \alpha_{12}x_2, \\ u_2 &\rightarrow u_2 + \alpha_{21}x_1 + \alpha_{22}x_2, \\ u_3 &\rightarrow u_3 + \alpha_{31}x_1 + \alpha_{32}x_2, \end{aligned} \quad (55)$$

where α_{ij} are arbitrary constants, produce a system with $\kappa_{10}^i = 0$ ($i = 1, \dots, 3$) provided that

$$\begin{aligned} &\kappa_1^i(\alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21}) + \kappa_2^i(\alpha_{11}\alpha_{32} - \alpha_{12}\alpha_{31}) + \kappa_3^i(\alpha_{21}\alpha_{32} - \alpha_{22}\alpha_{31}) \\ &+ \kappa_4^i\alpha_{11} + \kappa_5^i\alpha_{12} + \kappa_6^i\alpha_{21} + \kappa_7^i\alpha_{22} + \kappa_8^i\alpha_{31} + \kappa_9^i\alpha_{32} + \kappa_{10}^i = 0 \end{aligned} \quad (56)$$

($i = 1, \dots, 3$). Also in this case, conditions (56) can always be satisfied when the coefficients κ_j^i are constant because of the arbitrariness of the constants α_{ij} .

This nonlinear system (54), now taken with $\kappa_{10}^i = 0$, admits the Lie point symmetries spanned by the operators

$$\Xi_1 = \frac{\partial}{\partial x_1}, \quad \Xi_2 = \frac{\partial}{\partial x_2}, \quad \Xi_3 = (x_1 - f_1) \frac{\partial}{\partial x_1} + (x_2 - f_2) \frac{\partial}{\partial x_2}, \quad (57)$$

where $f_i(u_1, u_2, u_3)$ ($i = 1, 2$) are arbitrary smooth functions of their arguments, provided that

$$\begin{aligned} \kappa_1^i + \kappa_4^i f_{2;2} - \kappa_5^i f_{1;2} - \kappa_6^i f_{2;1} + \kappa_7^i f_{1;1} &= 0, \\ \kappa_2^i + \kappa_4^i f_{2;3} - \kappa_5^i f_{1;3} - \kappa_8^i f_{2;1} + \kappa_9^i f_{1;1} &= 0, \\ \kappa_3^i + \kappa_6^i f_{2;3} - \kappa_7^i f_{1;3} - \kappa_8^i f_{2;2} + \kappa_9^i f_{1;2} &= 0. \end{aligned} \quad (58)$$

The six conditions (58) cannot be fulfilled for an arbitrary choice of the coefficients κ_j^i . In the simplest case, where the coefficients κ_j^i are constant, they can be always satisfied and the functions f_i must be linear:

$$f_1 = \beta_{11}u_1 + \beta_{12}u_2 + \beta_{13}u_3, \quad f_2 = \beta_{21}u_1 + \beta_{22}u_2 + \beta_{23}u_3, \quad (59)$$

β_{ij} being arbitrary constants.

The symmetries (57) generate a 3-dimensional solvable Lie algebra,

$$[\Xi_i, \Xi_j] = 0, \quad [\Xi_i, \Xi_3] = \Xi_i, \quad (i, j = 1, 2), \quad (60)$$

whereupon we may introduce the new variables

$$\begin{aligned} z_1 &= x_1 - f_1, & z_2 &= x_2 - f_2, \\ w_1 &= u_1, & w_2 &= u_2, & w_3 &= u_3, \end{aligned} \quad (61)$$

and the generators of the point symmetries write as

$$\Xi_1 = \frac{\partial}{\partial z_1}, \quad \Xi_2 = \frac{\partial}{\partial z_2}, \quad \Xi_4 = z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2}. \quad (62)$$

In terms of the new variables (61), the nonlinear system (54) reduces to

$$\kappa_4^i w_{1,1} + \kappa_5^i w_{1,2} + \kappa_6^i w_{2,1} + \kappa_7^i w_{2,2} + \kappa_8^i w_{3,1} + \kappa_9^i w_{3,2} = 0, \quad (63)$$

i.e., reads as an autonomous and homogeneous quasilinear system.

3.4 Case $m = n = 3$

By considering the gradient matrix of $u_\alpha(x_i)$ ($i = 1, \dots, 3$, $\alpha = 1, \dots, 3$)

$$H = \begin{pmatrix} u_{1,1} & u_{1,2} & u_{1,3} \\ u_{2,1} & u_{2,2} & u_{2,3} \\ u_{3,1} & u_{3,2} & u_{3,3} \end{pmatrix} \quad (64)$$

and its extracted minors of order 2

$$\begin{aligned} H^1 &= \begin{vmatrix} u_{2,2} & u_{2,3} \\ u_{3,2} & u_{3,3} \end{vmatrix}, & H^2 &= \begin{vmatrix} u_{2,1} & u_{2,3} \\ u_{3,1} & u_{3,3} \end{vmatrix}, & H^3 &= \begin{vmatrix} u_{2,1} & u_{2,2} \\ u_{3,1} & u_{3,2} \end{vmatrix}, \\ H^4 &= \begin{vmatrix} u_{1,2} & u_{1,3} \\ u_{3,2} & u_{3,3} \end{vmatrix}, & H^5 &= \begin{vmatrix} u_{1,1} & u_{1,3} \\ u_{3,1} & u_{3,3} \end{vmatrix}, & H^6 &= \begin{vmatrix} u_{1,1} & u_{1,2} \\ u_{3,1} & u_{3,2} \end{vmatrix}, \\ H^7 &= \begin{vmatrix} u_{1,2} & u_{1,3} \\ u_{2,2} & u_{2,3} \end{vmatrix}, & H^8 &= \begin{vmatrix} u_{1,1} & u_{1,3} \\ u_{2,1} & u_{2,3} \end{vmatrix}, & H^9 &= \begin{vmatrix} u_{1,1} & u_{1,2} \\ u_{2,1} & u_{2,2} \end{vmatrix}, \end{aligned}$$

the nonlinear first order system of Monge–Ampère results composed by equations like

$$\begin{aligned} \kappa_0^i \det(H) + \kappa_1^i H^1 + \kappa_2^i H^2 + \kappa_3^i H^3 + \kappa_4^i H^4 + \kappa_5^i H^5 + \kappa_6^i H^6 \\ + \kappa_7^i H^7 + \kappa_8^i H^8 + \kappa_9^i H^9 + \kappa_{10}^i u_{1,1} + \kappa_{11}^i u_{1,2} + \kappa_{12}^i u_{1,3} + \kappa_{13}^i u_{2,1} \\ + \kappa_{14}^i u_{2,2} + \kappa_{15}^i u_{2,3} + \kappa_{16}^i u_{3,1} + \kappa_{17}^i u_{3,2} + \kappa_{18}^i u_{3,3} + \kappa_{19}^i = 0, \end{aligned} \quad (65)$$

$\kappa_j^i(u_\alpha)$ ($i = 1, \dots, 3$, $j = 0, \dots, 19$, $\alpha = 1, \dots, 3$) arbitrary smooth functions of the indicated arguments.

Also in this case, the substitutions

$$\begin{aligned} u_1 &\rightarrow u_1 + \alpha_{11}x_1 + \alpha_{12}x_2 + \alpha_{13}x_3, \\ u_2 &\rightarrow u_2 + \alpha_{21}x_1 + \alpha_{22}x_2 + \alpha_{23}x_3, \\ u_3 &\rightarrow u_3 + \alpha_{31}x_1 + \alpha_{32}x_2 + \alpha_{33}x_3, \end{aligned} \tag{66}$$

where α_{ij} are arbitrary constants, allow us to obtain a system with $\kappa_{19}^i = 0$ provided that $u_{i,j} = \alpha_{ij}$ is a solution of equations (65). This requirement implies some constraints on the coefficients κ_j^i in the general case; on the contrary, no limitation to the values of the coefficients exists if they are assumed to be constant.

The system (65), with $\kappa_{19}^i = 0$, admits the Lie point symmetries spanned by the operators

$$\begin{aligned} \Xi_1 &= \frac{\partial}{\partial x_1}, & \Xi_2 &= \frac{\partial}{\partial x_2}, & \Xi_3 &= \frac{\partial}{\partial x_3}, \\ \Xi_4 &= (x_1 - f_1) \frac{\partial}{\partial x_1} + (x_2 - f_2) \frac{\partial}{\partial x_2} + (x_3 - f_3) \frac{\partial}{\partial x_3}, \end{aligned} \tag{67}$$

where $f_i(u_1, u_2, u_3)$ ($i = 1, \dots, 3$) are arbitrary smooth functions of their

arguments, provided that

$$\begin{aligned}
& \kappa_0^i - (f_{2;2}f_{3;3} - f_{2;3}f_{3;2})\kappa_{10}^i - (f_{1;3}f_{3;2} - f_{1;2}f_{3;3})\kappa_{11}^i \\
& - (f_{1;2}f_{2;3} - f_{1;3}f_{2;2})\kappa_{12}^i - (f_{2;3}f_{3;1} - f_{2;1}f_{3;3})\kappa_{13}^i \\
& - (f_{1;1}f_{3;3} - f_{1;3}f_{3;1})\kappa_{14}^i - (f_{1;3}f_{2;1} - f_{1;1}f_{2;3})\kappa_{15}^i \\
& - (f_{2;1}f_{3;2} - f_{2;2}f_{3;1})\kappa_{16}^i - (f_{1;2}f_{3;1} - f_{1;1}f_{3;2})\kappa_{17}^i \\
& - (f_{1;1}f_{2;2} - f_{1;2}f_{2;1})\kappa_{18}^i = 0, \\
& \kappa_1^i + \kappa_{14}^i f_{3;3} - \kappa_{15}^i f_{2;3} - \kappa_{17}^i f_{3;2} + \kappa_{18}^i f_{2;2} = 0, \\
& \kappa_2^i + \kappa_{13}^i f_{3;3} - \kappa_{15}^i f_{1;3} - \kappa_{16}^i f_{3;2} + \kappa_{18}^i f_{1;2} = 0, \\
& \kappa_3^i + \kappa_{13}^i f_{2;3} - \kappa_{14}^i f_{1;3} - \kappa_{16}^i f_{2;2} + \kappa_{17}^i f_{1;2} = 0, \\
& \kappa_4^i + \kappa_{11}^i f_{3;3} - \kappa_{12}^i f_{2;3} - \kappa_{17}^i f_{3;1} + \kappa_{18}^i f_{2;1} = 0, \\
& \kappa_5^i + \kappa_{10}^i f_{3;3} - \kappa_{12}^i f_{1;3} - \kappa_{16}^i f_{3;1} + \kappa_{18}^i f_{1;1} = 0, \\
& \kappa_6^i + \kappa_{10}^i f_{2;3} - \kappa_{11}^i f_{1;3} - \kappa_{16}^i f_{2;1} + \kappa_{17}^i f_{1;1} = 0, \\
& \kappa_7^i + \kappa_{11}^i f_{3;2} - \kappa_{12}^i f_{2;2} - \kappa_{14}^i f_{3;1} + \kappa_{15}^i f_{2;1} = 0, \\
& \kappa_8^i + \kappa_{10}^i f_{3;2} - \kappa_{12}^i f_{1;2} - \kappa_{13}^i f_{3;1} + \kappa_{15}^i f_{1;1} = 0, \\
& \kappa_9^i + \kappa_{10}^i f_{2;2} - \kappa_{11}^i f_{1;2} - \kappa_{13}^i f_{2;1} + \kappa_{14}^i f_{1;1} = 0.
\end{aligned} \tag{68}$$

The vector fields (67) span a 4-dimensional solvable Lie algebra,

$$[\Xi_i, \Xi_j] = 0, \quad [\Xi_i, \Xi_4] = \Xi_i, \quad (i, j = 1, \dots, 3), \tag{69}$$

whereupon we may introduce the new variables

$$\begin{aligned}
z_1 &= x_1 - f_1, & z_2 &= x_2 - f_2, & z_3 &= x_3 - f_3, \\
w_1 &= u_1, & w_2 &= u_2, & w_3 &= u_3,
\end{aligned} \tag{70}$$

and the generators of the point symmetries write as

$$\begin{aligned}
\Xi_1 &= \frac{\partial}{\partial z_1}, & \Xi_2 &= \frac{\partial}{\partial z_2}, & \Xi_3 &= \frac{\partial}{\partial z_3}, \\
\Xi_4 &= z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3}.
\end{aligned} \tag{71}$$

In terms of the new variables (70), Eqs. (65) write as

$$\begin{aligned}
& \kappa_{10}^i w_{1,1} + \kappa_{11}^i w_{1,2} + \kappa_{12}^i w_{1,3} + \kappa_{13}^i w_{2,1} + \kappa_{14}^i w_{2,2} + \kappa_{15}^i w_{2,3} \\
& + \kappa_{16}^i w_{3,1} + \kappa_{17}^i w_{3,2} + \kappa_{18}^i w_{3,3} = 0,
\end{aligned} \tag{72}$$

i.e., they are in autonomous and homogeneous quasilinear (linear, if the coefficients are constant) form.

Conditions (68) play severe restrictions to the expression of the coefficients κ_j^i . When these coefficients are assumed to be constant, we are forced to take

$$\begin{aligned} f_1 &= \beta_{11}u_1 + \beta_{12}u_2 + \beta_{13}u_3, \\ f_2 &= \beta_{21}u_1 + \beta_{22}u_2 + \beta_{23}u_3, \\ f_3 &= \beta_{31}u_1 + \beta_{32}u_2 + \beta_{33}u_3, \end{aligned} \tag{73}$$

where β_{ij} are arbitrary constants; also in such simple case, the reduction to linear form is not always possible due to (68).

3.5 Case m and n arbitrary

It is easily recognized that, a general Monge–Ampère system with m dependent variables and n independent variables, provided that some suitable conditions on the coefficients (at most depending on the field variables) are satisfied, is invariant with respect to the Lie groups generated by the vector fields

$$\Xi_i = \frac{\partial}{\partial x_i} \quad (i = 1, \dots, n), \quad \Xi_{n+1} = \sum_{i=1}^n (x_i - f_i(u_\alpha)) \frac{\partial}{\partial x_i}, \tag{74}$$

where $f_i(u_\alpha)$ are smooth functions of (u_1, \dots, u_m) which have to be linear in their arguments when the coefficients of the Monge–Ampère system are constant.

As one expects, for $m > 3$ or $n > 3$, we have a situation similar to the case $m = n = 3$, *i.e.*, even in the case of constant coefficients, not all Monge–Ampère systems can be reduced to (quasi)linear form.

4 Conclusions

In this paper, we proved a theorem giving necessary and sufficient conditions for transforming a nonlinear first order system of partial differential equations involving the derivatives in polynomial form to an equivalent autonomous system polynomially homogeneous in the derivatives. The theorem is based on the Lie point symmetries admitted by the nonlinear system, and the proof

is constructive, in the sense that it leads to the algorithmic construction of the invertible mapping performing the task.

The theorem is applied to a class of first order nonlinear systems belonging to the family of Monge–Ampère systems that have been characterized by Boillat in 1997. These systems share with the quasilinear systems the property of the linearity of the Cauchy problem. These systems are also completely exceptional [22, 23], and are made by equations which are expressed as linear combinations (with coefficients depending at most on the independent and the dependent variables) of all minors extracted from the gradient matrix of $u_\alpha = u_\alpha(x_i)$ ($\alpha = 1, \dots, m$, $i = 1, \dots, n$). We considered explicitly either the case of constant coefficients or the case of coefficients depending on the field variables, for $m = 2, 3$ and $n = 2, 3$. If $m = 2$ and $n = 2, 3$, or $n = 2$ and $m = 2, 3$, and the coefficients are assumed to be constant, we proved that the Monge–Ampère systems can always be transformed to linear form.

Nevertheless, for arbitrary m and n , Monge–Ampère systems, provided that the coefficients entering their equations satisfy some constraints, can be mapped to first order quasilinear autonomous and homogeneous systems. This, in some sense, casts new light on the fact, underlined by Boillat [21], that Monge–Ampère systems, because of the linearity of the Cauchy problem, are the closest to quasilinear systems, which are Monge systems.

Moreover, an example of a first order system polynomial in the derivatives that can be reduced to a system polynomially homogeneous in the derivatives (equivalent to a second order partial differential equation for a surface in \mathbb{R}^3 such that its Gaussian curvature is proportional to the square of its mean curvature), is provided.

Acknowledgments

The authors thank the Referee for the helpful comments leading to improve the quality of the paper. Work supported by G.N.F.M. of the Istituto Nazionale di Alta Matematica.

References

- [1] L. V. Ovsiannikov. Group analysis of differential equations. Academic Press, New York, 1982.
- [2] N. H. Ibragimov. Transformation groups applied to mathematical physics. D. Reidel Publishing Company, Dordrecht, 1985.
- [3] P. J. Olver. Applications of Lie groups to differential equations. Springer, New York, 1986.
- [4] P. J. Olver. Equivalence, Invariants, and Symmetry. Cambridge University Press, Cambridge, 1995.
- [5] G. Baumann. Symmetry Analysis of Differential Equations with Mathematica. Springer, New York, 2000.
- [6] G. W. Bluman, S. C. Anco. Symmetry and Integration Methods for Differential Equations. Springer, New York, 2002.
- [7] S. V. Meleshko. Methods for Constructing Exact Solutions of Partial Differential Equations. Springer, New York, 2005.
- [8] G. W. Bluman, A. F. Cheviakov, S. C. Anco. Applications of Symmetry Methods to Partial Differential Equations. Springer, New York, 2009.
- [9] S. Kumei, G. W. Bluman. When nonlinear differential equations are equivalent to linear differential equations. *SIAM J. Appl. Math.*, **42**, 1157–1173, 1982.
- [10] G. W. Bluman, S. Kumei. Symmetries and differential equations. Springer, New York, 1989.
- [11] A. Donato, F. Oliveri. Linearization procedure of nonlinear first order systems of PDE's by means of canonical variables related to Lie groups of point transformations. *J. Math. Anal. Appl.*, **188**, 552–568, 1994.
- [12] A. Donato, F. Oliveri. When nonautonomous equations are equivalent to autonomous ones. *Appl. Anal.*, **58**, 313–323, 1995.

- [13] A. Donato, F. Oliveri. How to build up variable transformations allowing one to map nonlinear hyperbolic equations into autonomous or linear ones. *Transp. Th. Stat. Phys.*, **25**, 303–322, 1996.
- [14] C. Currò, F. Oliveri. Reduction of nonhomogeneous quasilinear 2×2 systems to homogeneous and autonomous form. *J. Math. Phys.*, **49**, 103504-1–103504-11, 2008.
- [15] F. Oliveri. Lie symmetries of differential equations: classical results and recent contributions. *Symmetry*, **2**, 658–706, 2010.
- [16] F. Oliveri. General dynamical systems described by first order quasilinear PDEs reducible to homogeneous and autonomous form. *Int. J. Non-linear Mech.*, **47**, 53–60, 2012.
- [17] M. Gorgone, F. Oliveri. Nonlinear first order partial differential equations reducible to first order homogeneous and autonomous quasilinear ones. *Ricerche di Matematica*, 1–13, doi: 10.1007/s11587-016-0286-8, 2016.
- [18] A. Pressley. *Elementary differential geometry*. Springer–Verlag, London, 2010.
- [19] G. Manno, F. Oliveri, R. Vitolo. On differential equations characterized by their Lie point symmetries. *J. Math. Anal. Appl.*, **332**, 767–786, 2007.
- [20] G. Manno, F. Oliveri, R. Vitolo. Differential equations uniquely determined by algebras of point symmetries. *Theor. Math. Phys.*, **151**, 843–850, 2007.
- [21] G. Boillat. Sur la forme générale du système de Monge–Ampère. *C. R. Acad. Sci. Paris*, **325**, Série I, 339–342, 1997.
- [22] P. D. Lax. Hyperbolic systems of conservation laws, II. *Comm. Pure Appl. Math.*, **10**, 537–566, 1957.
- [23] G. Boillat. *La propagation des ondes*. Gauthier–Villars, Paris, 1965.