

# Representations of the Bondi–Metzner–Sachs group in three space–time dimensions in the Hilbert topology I. Determination of the representations

---

Evangelos Melas <sup>a</sup>

<sup>a</sup> *University of Thessaly,  
Department of Mathematics,  
Lamia 35100,  
Greece*

*E-mail:* [evanmelas@uth.gr](mailto:evanmelas@uth.gr)

ABSTRACT: The original Bondi–Metzner–Sachs (BMS) group  $B$  is the common asymptotic symmetry group of all asymptotically flat Lorentzian 4–dim space–times. As such,  $B$  is the best candidate for the universal symmetry group of General Relativity (G.R.). Here, the analogue  $B(2,1)$  of  $B$  in 3 space–time dimensions is properly defined. We study its representation theory in the Hilbert topology by using an infinite–dimensional extension of Wigner–Mackey theory. We obtain the necessary data in order to construct the strongly continuous irreducible unitary representations (IRS) of  $B(2,1)$ . The main results of the representation theory are: The IRS are induced from “little groups” which are compact. There is one infinite connected “little group”, the special orthogonal group  $SO(2)$ . There are infinite non–connected finite discrete “little groups”, the cyclic groups  $C_n$  of even order. The inducing construction is exhaustive notwithstanding the fact that  $B(2,1)$  is not locally compact in the employed Hilbert topology.  $B(2,1)$  is also derived, in Klein’s sense, as the automorphism group of the “strong conformal geometry” of  $\mathcal{S}^+$ . Besides the Hilbert topology other reasonable topologies are given to  $B(2,1)$  and their physical relevance is discussed. Connections of the IRS of  $B(2,1)$  with geometry are outlined.

KEYWORDS: Space–Time Symmetries Models of Quantum Gravity Classical Theories of Gravity

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The group <math>B^{2,1}(N^+)</math></b>	<b>3</b>
<b>3</b>	<b>The double cover <math>B^{2,1}(N^+)_c</math></b>	<b>4</b>
<b>4</b>	<b>The group <math>B^{2,1}(N^+)_c</math></b>	<b>4</b>
4.1	The projective space of $N^+$	5
4.2	The bijections $F(P(N^+)) \leftrightarrow F_1(N^+) \leftrightarrow F_1(\widetilde{N^+}) \leftrightarrow F_2(\mathcal{I})$	7
4.3	The spinor version of $B^{2,1}(N^+)_c$	9
4.4	Unconstrained supertranslations	9
4.5	Almost unconstrained supertranslations	11
<b>5</b>	<b>The group <math>B(2, 1)</math></b>	<b>12</b>
<b>6</b>	<b><math>B(2, 1)</math> as the automorphism group of the strong conformal geometry of <math>\mathcal{I}^+</math></b>	<b>14</b>
6.1	“Future null infinity” $\mathcal{I}^+$ of 3–dim Minkowski space–time	15
6.2	The Newman–Unti group	17
6.3	Self–diffeomorphisms of $\mathcal{I}^+$ which preserve its strong conformal geometry	18
6.4	“Null angles”	19
<b>7</b>	<b>The classical action <math>B(2, 1) \times \mathcal{I}^+ \longrightarrow \mathcal{I}^+</math></b>	<b>20</b>
<b>8</b>	<b>The supertranslation subgroup</b>	<b>21</b>
<b>9</b>	<b>On the subgroups of <math>B(2, 1)</math></b>	<b>22</b>
<b>10</b>	<b>The spaces <math>\mathcal{A}</math>, <math>\mathcal{A}'</math>, <math>\hat{\mathcal{A}}</math></b>	<b>24</b>
<b>11</b>	<b>The dual action <math>T'</math> and the supermomentum space <math>\mathcal{A}'</math></b>	<b>25</b>
<b>12</b>	<b>Representation theory and its extension</b>	<b>26</b>
12.1	Wigner–Mackey’s theory	27
12.2	Extension of Wigner–Mackey’s theory	29
<b>13</b>	<b>Topologies in <math>\mathcal{A}</math></b>	<b>30</b>
13.1	$\mathcal{A}$ is not unique	30
13.2	Topologies in $\mathcal{A}$	31
<b>14</b>	<b>“Little groups” are compact</b>	<b>31</b>
14.1	“Little groups” depend on the topology	31
14.2	Proof	32

<b>15</b>	<b>The “Little groups” of <math>B(2, 1)</math></b>	<b>33</b>
15.1	The compact subgroups of $SO(2)$	34
15.2	The “Little groups” of $B(2, 1)$	35
<b>16</b>	<b>Exhaustivity of Induced Representations</b>	<b>35</b>
16.1	Properties of the dual action	35
16.2	Sufficient conditions are satisfied	37
<b>17</b>	<b>Construction of the IRS of <math>B(2, 1)</math></b>	<b>37</b>
<b>18</b>	<b>Connections to Geometry</b>	<b>39</b>

---

## 1 Introduction

The BMS group  $B$  is the common asymptotic group of all curved real Lorentzian space–times which are asymptotically flat in future null directions [1, 2], and is the best candidate for the universal symmetry group of G.R..

The BMS group  $B$  of general relativity was first discovered [1, 2] not as a transformation group of (exact) global diffeomorphisms of a fixed manifold, but as a pseudo–group of local diffeomorphisms (“asymptotic isometries”) of the asymptotic region of Lorentzian space–times which are asymptotically flat in lightlike future directions.

However, Penrose [3–5] showed that, by “going to infinity”,  $B$  could be interpreted as an (exact) global transformation group  $B \times \mathfrak{S}^+ \rightarrow \mathfrak{S}^+$  of the “future null boundary”  $\mathfrak{S}^+$  of the space–times concerned. Furthermore, he gave [3, 4] a geometric structure to  $\mathfrak{S}^+$ , the “strong conformal geometry” for which the transformation group is precisely the group of automorphisms [5].

In 1939 Wigner laid the foundations of special relativistic quantum mechanics [6] and relativistic quantum field theory by constructing the Hilbert space IRS of the (universal cover) of the Poincare group  $P$ .

The universal property of  $B$  for G.R. makes it reasonable to attempt to lay a similarly firm foundation for quantum gravity by following through the analogue of Wigner’s programme with  $B$  replacing  $P$ . Some years ago McCarthy constructed explicitly [7–14] the IRS of  $B$  for exactly this purpose. This work was based on G. W. Mackey’s pioneering work on group representations [15–20]; in particular it was based on an extension [11, 21] of G. W. Mackey’s work to the relevant infinite dimensional case.

It is difficult to overemphasize the importance of Piard’s results [22, 23] who soon afterwards proved that *all* the IRS of  $B$ , when this is equipped with the Hilbert topology, are derivable by the inducing construction. This proves the exhaustivity of McCarthy’s list of representations and renders his results even more important.

However, in quantum gravity, complexified or euclidean versions of G.R. are frequently considered and the question arises: Are there similar symmetry groups for these versions of

the theory? McCarthy constructed [24], in abstract form, all possible analogues of  $B$ , both real and in any signature, or complex, with all possible notions of asymptotic flatness “near infinity”. There are, in fact, forty–one such groups. One of them is  $B(2, 2)$ , the asymptotic symmetry group of all curved real ultrahyperbolic space–times which are asymptotically flat in null directions.

These abstract constructions were given in a quantum setting; the paper [24] was concerned with finding the IRS of these groups in Hilbert spaces (especially for the complexification  $\mathcal{CB}$  of  $B$  itself). It has been argued [24, 25] that these Hilbert space representations are related to elementary particles and quantum gravity (via gravitational instantons). The IRS of  $B(2, 2)$  were constructed in [26, 27]. The representation theory of HB, which is a subgroup of  $B(2, 2)$ , was initiated in [28–30].

Here, we follow this programme for 3–dim G.R. We properly define the analogue  $B(2, 1)$  of  $B$  in three space–time dimensions. We construct the IRS of  $B(2, 1)$  by endowing it with the Hilbert topology. IRS in the Hilbert topology describe *bounded* sources [13].

$B(2, 1)$  was first introduced in [21]. However, the treatment of  $B(2, 1)$  in [21] was rather discursive. Only the main features of  $B(2, 1)$  were given without any relevant proofs. [21] dealt primarily with the needed infinite–dimensional extension of Wigner–Mackey theory required to study the representation theory  $B$  and all its variants in 3 space–time dimensions and higher in all signatures. In this paper, the extension of Wigner–Mackey theory given in [21], is particularized in the case of  $B(2, 1)$ . Moreover, further particulars of the necessary extension of Wigner–Mackey theory appropriate to the case of  $B(2, 1)$  are given mainly regarding the various reasonable topologies  $B(2, 1)$  can be equipped with and their physical underpinnings.

The main results of the representation theory are: The IRS of  $B(2, 1)$  are induced from IRS of *compact* “little groups”. The “little groups” are of two types: 1. Infinite connected Lie groups. 2. Non–connected finite discrete groups. The infinite connected “little group” is the special orthogonal group  $SO(2)$ . The non–connected finite discrete “little groups” are cyclic groups of even order, which are symmetry groups of regular polygons in ordinary euclidean 2–space. The inducing construction is exhaustive notwithstanding the fact that  $B(2, 1)$  is *not* locally compact in the employed Hilbert topology. We give all the necessary information in order to construct the operators of the induced representations of  $B(2, 1)$ .

$B(2, 1)$  is the semidirect product of  $SL(2, R)$  with the additive group of real–valued functions defined on the circle. The asymptotic symmetry group in 3 space–time dimensions, to the best of our knowledge, was first considered in [31], a work which is entirely classical in aims and scope. Later in [32] the authors defined, in three space–time dimensions,  $BMS_3$ , the semidirect product of  $Diff(S^1)$ , the group of the orientation preserving diffeomorphisms of the circle, with the additive group of real–valued functions defined on the circle. In [33], [34] the projective representations of  $BMS_3$  were studied. The main difference between this paper and [31], [32], [33], [34] is that in [31], [32], [33], [34],  $SL(2, R)$  has been enlarged to the full group  $Diff(S^1)$ . As a result there is no overlap between this paper and [31], [32], [33], [34].

This paper is organised as follows: In section 2 the group  $B^{2,1}(N^+)$  is introduced. In section 3 the double cover  $B^{2,1}(N^+)_c$  of  $B^{2,1}(N^+)$  is defined. In section 4 the group

$B^{2,1}(N^+)_c$  is studied. In section 5 the group  $B(2,1)$  is defined. In section 6 the group  $B(2,1)$  is rederived as automorphism group of the strong conformal geometry of  $\mathcal{S}^+$ . In section 7 the classical action of  $B(2,1)$  on  $\mathcal{S}^+$  is given. In section 8 the subgroup of supertranslations of  $B(2,1)$  is studied. In section 9 the subgroup structure of  $B(2,1)$  is analyzed. In section 10 the spaces  $\mathcal{A}$ ,  $\mathcal{A}'$ ,  $\hat{\mathcal{A}}$  are defined. In section 11 the dual action of the action which defines  $B(2,1)$  on  $\mathcal{A}'$  is given. In section 12 Wigner–Mackey’s theory is outlined and its necessary extension is presented. In section 13 a class of “reasonable” topologies for  $\mathcal{A}$  is given. In section 14 it is proved that all “Little groups” are compact. In section 15 the “Little groups” of  $B(2,1)$  are given. In section 16 it is proved that *all* IRS of  $B(2,1)$  are obtained by the Wigner–Mackey’s inducing construction. In section 17 all the necessary data in order to construct the IRS of  $B(2,1)$  are given. In section 18 connections of the IRS of  $B(2,1)$  with geometry are discussed.

## 2 The group $B^{2,1}(N^+)$

We turn now to the study of  $B(2,1)$ , the analogue of  $B$  in three space–time dimensions. Recall that the  $2+1$  Minkowski space is the vector space  $R^3$  of row vectors with 3 real components, with inner product defined as follows. Let  $x, y \in R^3$  have components  $x^\mu$  and  $y^\mu$  respectively, where  $\mu = 0, 1, 2$ . Define the inner product  $x \cdot y$  between  $x$  and  $y$  by

$$x \cdot y = x^0 y^0 - x^1 y^1 - x^2 y^2. \quad (2.1)$$

Then the  $2+1$  Minkowski space, sometimes written  $R^{2,1}$ , is just  $R^3$  with this inner product. The “2,1” refers to the one plus and two minus signs in the inner product. Let  $SO(2,1)$  be the (connected component of the identity element of the) group of linear transformations preserving the inner product. Matrices  $\Lambda \in SO(2,1)$  are taken as acting by matrix multiplication from the right,  $x \mapsto x\Lambda$ , on row vectors  $x \in R^{2,1}$ .

The future null cone  $N^+ \subset R^{2,1}$  is just the set of nonzero vectors with zero length and  $x^0 > 0$ :

$$N^+ = \{x \in R^{2,1} \mid x \cdot x = 0, x^0 > 0\}. \quad (2.2)$$

Let  $R^+$  denote the multiplicative group of all *positive* real numbers. Obviously, if  $x \in N^+$ , then  $tx \in N^+$  for any  $t \in R^+$ . Let  $F_1(N^+)$  denote the vector space (under pointwise addition) of all functions  $f : N^+ \rightarrow R$  satisfying the homogeneity condition

$$f(tx) = tf(x) \quad (2.3)$$

for all  $x \in N^+$  and  $t \in R^+$ . Define a representation  $T$  of  $SO(2,1)$  on  $F_1(N^+)$  by setting, for each  $x \in N^+$  and  $\Lambda \in SO(2,1)$ ,

$$(T(\Lambda)f)(x) = f(x\Lambda). \quad (2.4)$$

Now let  $B^{2,1}(N^+)$  be the semi–direct product

$$B^{2,1}(N^+) = F_1(N^+) \circledast_T SO(2,1). \quad (2.5)$$

That is to say,  $B^{2,1}(N^+)$  is, as a set, just the product  $F_1(N^+) \times SO(2,1)$ , and the group multiplication law for pairs is

$$(f_1, \Lambda_1)(f_2, \Lambda_2) = (f_1 + T(\Lambda_1)f_2, \Lambda_1\Lambda_2). \quad (2.6)$$

### 3 The double cover $B^{2,1}(N^+)_c$

Let  $SL(2, R)$  be the group of all real  $2 \times 2$  matrices with determinant one.  $SL(2, R)$  is sometimes denoted  $G$  below. Let  $M_s(2, R)$  be the set of all  $2 \times 2$  symmetric real matrices. We define a right action of  $G$  on  $M_s(2, R)$  by  $M_s(2, R) \times G \rightarrow M_s(2, R)$  with

$$(m, g) \mapsto g^\top m g, \quad (3.1)$$

where the superscript  $\top$  means transpose. Clearly any element  $\mu \in M_s(2, R)$  can be parameterized as follows

$$\mu = \begin{bmatrix} x^0 - x^1 & x^2 \\ x^2 & x^0 + x^1 \end{bmatrix}$$

where  $x^0, x^1, x^2 \in R$ .

We now consider the map  $b : R^3 \rightarrow M_s(2, R)$  defined by

$$b(x) = \begin{bmatrix} x^0 - x^1 & x^2 \\ x^2 & x^0 + x^1 \end{bmatrix}, \quad (3.2)$$

where the  $x^\mu$  are the components of  $x \in R^3$ . This map is a linear bijection, so the right action of  $G$  on  $M_s(2, R)$  induces a linear right action of  $G$  on  $R^3$ . Since

$$\det(b(x)) = x \cdot x \quad (3.3)$$

and the  $G$  action preserves determinants (indeed  $\det g = 1$ ) in  $M_s(2, R)$ ,  $G$  acts as transformations from  $SO(2, 1)$ . In fact, this construction gives an homomorphism

$$\gamma : G \rightarrow SO(2, 1) \quad (3.4)$$

which is onto, and has kernel  $Z_2 = \{Id, -Id\}$  in  $G$ ,  $Id$  denotes the identity element of  $G$ .

Thus  $\gamma$  identifies  $G$  as the double cover of  $SO(2, 1)$

$$G = SO(2, 1)_c. \quad (3.5)$$

Therefore, the double cover of the group  $B^{2,1}(N^+)$ , given in (2.5), has the form

$$B^{2,1}(N^+)_c = F_1(N^+) \otimes_T G. \quad (3.6)$$

Strictly speaking, “ $T$ ” should read “ $T\gamma$ ”, but the notation is simpler as above.

### 4 The group $B^{2,1}(N^+)_c$

We describe now the structure of the group  $B^{2,1}(N^+)_c$ .

#### 4.1 The projective space of $N^+$

The null cone  $N^+$  becomes, under identification (3.2),  $\widetilde{N}^+ = b(N^+)$  with

$$\widetilde{N}^+ = \{\mu \in M_s(2, R), | x^0 > 0, \det \mu = 0\}. \quad (4.1)$$

Thus  $\mu \in \widetilde{N}^+$  if and only if  $\mu$  has rank exactly 1 and  $x^0 > 0$ . Let  $\mathcal{S}$  be the set of all non-zero real two-component row vectors  $\sigma$ ;  $\mathcal{S} = R^2 - 0$ .  $\mathcal{S}$  is the real section of Penrose's "spin space" appropriate to  $N^+ \subset R^{2,1}$ . From the rank condition and the requirement  $x^0 > 0$ , it follows that

$$\mu \in \widetilde{N}^+ \quad \Leftrightarrow \quad \mu = \sigma^\top \sigma \quad (4.2)$$

where  $\sigma \in \mathcal{S}$ . Equation (4.2) gives a parametrisation of the null cone  $\widetilde{N}^+$  by means of the non-zero real two-component row vectors  $\sigma$ . However, this parametrisation, though (by construction) onto, is not one-one.

To make this explicit we define an equivalence relation in  $\mathcal{S}$ . Two elements  $\sigma, \sigma'$  of  $\mathcal{S}$  are equivalent if and only if

$$\sigma' = \kappa \sigma, \quad \kappa = 1, -1. \quad (4.3)$$

Therefore if  $[\sigma]$  is the equivalence class of  $\sigma$ ,  $[\sigma] = \{\sigma, -\sigma\}$ .

If  $\sigma'$  and  $\sigma$  belong to the same equivalence class, then, by Eq. (4.3),  $\sigma' = \kappa \sigma$ ,  $\kappa = 1, -1$ . Therefore,

$$\sigma'^\top \sigma' = (\kappa \sigma)^\top (\kappa \sigma) = \kappa^2 \sigma^\top \sigma = \sigma^\top \sigma.$$

Consequently,  $\sigma'$  and  $\sigma$  define the same  $\mu$ .

Conversely, if  $\sigma'$  and  $\sigma$  define the same  $\mu$ , then

$$\sigma'^\top \sigma' = \sigma^\top \sigma. \quad (4.4)$$

Postmultiplying (4.4) by the transpose  $\sigma^\top$  of  $\sigma$  gives

$$\sigma'^\top (\sigma' \sigma^\top) = \sigma^\top (\sigma \sigma^\top). \quad (4.5)$$

Since  $\sigma \sigma^\top \neq 0$ ,  $\sigma' \sigma^\top$  cannot be zero. Hence  $\sigma'$  is a non zero real multiple of  $\sigma$ ,  $\sigma' = \lambda \sigma$ . Substituting this into equation (4.5) gives  $\lambda^2 = 1$ . Consequently,  $\sigma'$  and  $\sigma$  belong to the same equivalent class. We therefore arrive at the Proposition which follows.

**Proposition 1** *Every  $\mu \in \widetilde{N}^+$  corresponds exactly to one equivalence class  $[\sigma]$  and vice versa.*

Let  $\pi$  be the projection  $\pi : \mathcal{S} \rightarrow \widetilde{N}^+$  given by

$$\pi(\sigma) = \sigma^\top \sigma. \quad (4.6)$$

Let  $Z_2 = \{-1, 1\}$ . Define a left action  $Z_2 \times \mathcal{S} \rightarrow \mathcal{S}$  by

$$(\kappa, \sigma) \mapsto \kappa \sigma. \quad (4.7)$$

Then this  $Z_2$  action is fixed point free. Indeed Let  $\sigma = (\sigma_1 \ \sigma_2) \in \mathcal{I}$ . If  $\lambda\sigma = \sigma$ , then,  $\lambda\sigma_1 = \sigma_1$  and  $\lambda\sigma_2 = \sigma_2$ . But  $\sigma_1$  and  $\sigma_2$  do not both vanish, so  $\lambda = 1$  and the action (4.7) is fixed point free.

According to Proposition 1 the  $Z_2$  orbits are precisely the fibres of the projection

$$\pi : \mathcal{I} \rightarrow \widetilde{N^+}. \quad (4.8)$$

We conclude that

**Proposition 2** *The left free action of  $Z_2$  on  $\mathcal{I}$*

$$Z_2 \times \mathcal{I} \rightarrow \mathcal{I}, \quad (\kappa, \sigma) \mapsto \kappa\sigma,$$

*gives rise to a principal  $Z_2$ -bundle  $(\mathcal{I}, \pi, \widetilde{N^+})$  with projection*

$$\pi : \mathcal{I} \rightarrow \widetilde{N^+}, \quad \pi(\sigma) = \sigma^\top \sigma,$$

*and base space diffeomorphic to  $\widetilde{N^+}$ .*

Let  $R^+$  be the multiplicative group of positive real numbers. Define a left action  $R^+ \times \widetilde{N^+} \rightarrow \widetilde{N^+}$  by

$$(t, \mu) \mapsto t\mu. \quad (4.9)$$

This action is fixed point free and defines a principal bundle  $\widetilde{N^+} \rightarrow P(\widetilde{N^+})$ .

Combining the projections

$$\mathcal{I} \rightarrow \widetilde{N^+} \rightarrow P(\widetilde{N^+}) \quad (4.10)$$

evidently corresponds to combining the  $Z_2$  and  $R^+$  actions to give a  $R^+ \times Z_2$  action on  $\mathcal{I}$ . Indeed, we can take the combined action as

$$((t, \lambda), \sigma) \mapsto t\lambda\sigma. \quad (4.11)$$

But,  $t\lambda = \kappa \in R^*$ , where  $R^*$  is the multiplicative group of non-zero real numbers.

Consequently, the action (4.11) is equivalent to the  $R^*$  action given by

$$(\kappa, \sigma) \mapsto \kappa\sigma. \quad (4.12)$$

Hence the base  $P(\widetilde{N^+})$  is the set of orbits for this last action, namely

$$P(\widetilde{N^+}) \simeq S^1/Z_2 \equiv P_1(R). \quad (4.13)$$

Here  $P_1(R)$  is the one-dimensional real projective space. Hence we have

**Proposition 3** *The left free action of  $R^*$  on  $\mathcal{I}$*

$$R^* \times \mathcal{I} \rightarrow \mathcal{I}, \quad (\kappa, \sigma) \mapsto \kappa\sigma,$$



gives rise to a principal  $R^*$ -bundle  $(\mathcal{I}, \pi, P_1(R))$  with projection

$$\begin{aligned} \pi &: \mathcal{I} \rightarrow P_1(R), \\ \pi(\sigma) = \pi(\sigma_1 \ \sigma_2) &= \left( \frac{\rho}{w(\rho)}, \frac{1}{w(\rho)} \right), \\ \rho &= \frac{\sigma_1}{\sigma_2}, \quad w(\rho) = (\rho^2 + 1)^{1/2}, \\ &\text{when } \sigma_2 \neq 0, \text{ and,} \\ \pi(\sigma) = \pi(\sigma_1 \ 0) &= (1 \ 0), \text{ when } \sigma_2 = 0, \end{aligned} \tag{4.14}$$

and base space diffeomorphic to  $P_1(R)$ .

Noticing that  $\widetilde{N^+} = b(N^+)$  we also have

$$P(N^+) \simeq P(\widetilde{N^+}) \simeq P_1(R). \tag{4.15}$$

#### 4.2 The bijections $F(P(N^+)) \leftrightarrow F_1(N^+) \leftrightarrow F_1(\widetilde{N^+}) \leftrightarrow F_2(\mathcal{I})$

Let

$$F_1(\widetilde{N^+}) = \{ \tilde{f} : \widetilde{N^+} \rightarrow R \mid \tilde{f}(t\mu) = t\tilde{f}(\mu) \} \tag{4.16}$$

for all  $t \in R^+$  and  $\mu \in \widetilde{N^+}$ .

We have the following

**Proposition 4** *There are the bijections*

$$F(P(N^+)) \leftrightarrow F_1(N^+) \leftrightarrow F_1(\widetilde{N^+}).$$

**Proof** *The orbits of the dilatation action  $R^+ \times N^+ \rightarrow N^+$ ;  $(t, x) \mapsto tx$  are open half-lines in  $N^+$  from the origin of  $R^{2,1}$ . The projective null cone  $P(N^+)$  is the space of these half lines. Since the homogeneity constraint (2.3) fixes the behaviour of functions  $f \in F_1(N^+)$  along these half lines,  $F_1(N^+)$  may also be realized as the set  $F(P(N^+))$  of all arbitrary (unconstrained or “free”) real valued functions defined on  $P(N^+)$ :*

$$F_1(N^+) \simeq F(P(N^+)). \tag{4.17}$$

Let  $x = (x^1, x^2, x^0) \in N^+$  and let the unit circle

$$S_m^1 = \{ x \in N^+ \mid x^0 = 1 \}$$

be our model for  $P(N^+)$ . The orbit  $R^+x$  intersects  $S_m^1$  at exactly one point

$$\tilde{x} = \frac{x}{\sqrt{(x^1)^2 + (x^2)^2}}. \tag{4.18}$$

With each  $f \in F_1(N^+)$  we associate a function  $\alpha \in F(P(N^+))$  by restricting  $f$  to the circle  $S_m^1$ :

$$\alpha(\tilde{x}) = f(\tilde{x}). \tag{4.19}$$

Using the homogeneity condition (2.3) we have

$$\alpha(\tilde{x}) = \frac{f(x)}{\sqrt{(x^1)^2 + (x^2)^2}}. \quad (4.20)$$

Formula (4.52) associates not only  $f$  with the corresponding  $\alpha$ , but also vice versa. It explicitly gives the bijection

$$F_1(N^+) \leftrightarrow F(P(N^+)). \quad (4.21)$$

Evidently, the bijection between  $F_1(N^+)$  and  $F_1(\widetilde{N^+})$  is given by

$$f(x) = \tilde{f}(b(x)). \quad (4.22)$$

This completes the proof.

Now suppose that  $\psi : \mathcal{S} \rightarrow R$  is any function which satisfies

$$\psi(-\sigma) = \psi(\sigma) \quad (4.23)$$

for all  $\sigma \in \mathcal{S}$ . This just means that  $\psi$  is constant on each  $Z_2$  orbit of Proposition 2. That is,  $\psi$  only depends on the points  $\mu = \sigma^\top \sigma \in \widetilde{N^+}$ , and so can be written as a function  $\tilde{f} : \widetilde{N^+} \rightarrow R$ :

$$\psi(\sigma) = \tilde{f}(\sigma^\top \sigma). \quad (4.24)$$

Conversely, any function  $\tilde{f} : \widetilde{N^+} \rightarrow R$  defines a function  $\psi$ , given by (4.24), which satisfies (4.23).

We can now use (4.24) to identify  $F_1(\widetilde{N^+})$  with a subset of the space of all functions  $\psi : \mathcal{S} \rightarrow R$ . Let  $F_2(\mathcal{S})$  be the set of all such functions which are homogeneous of degree 2, i.e., of all functions satisfying

$$\psi(q\sigma) = q^2\psi(\sigma) \quad (4.25)$$

for all  $q \in R^*$  and all  $\sigma \in \mathcal{S}$ . Then we have the following

**Proposition 5** *There is a bijection*

$$F_1(\widetilde{N^+}) \leftrightarrow F_2(\mathcal{S}).$$

**Proof** *Given any  $\psi \in F_2(\mathcal{S})$ , we have, using (4.25),  $\psi(-\sigma) = \psi(\sigma)$ , so  $\psi$  satisfies (4.23). So, as in (4.24), we can write  $\psi(\sigma) = \tilde{f}(\sigma^\top \sigma)$  for some function  $\tilde{f} : \widetilde{N^+} \rightarrow R$ . Writing  $\mu = \sigma^\top \sigma$ , we then have*

$$\tilde{f}(t\mu) = \tilde{f}(t(\sigma^\top \sigma)) = \tilde{f}((\sqrt{t}\sigma)^\top (\sqrt{t}\sigma)) = \psi(\sqrt{t}\sigma) = t\psi(\sigma) = t\tilde{f}(\sigma^\top \sigma) = t\tilde{f}(\mu),$$

for all  $t \in R^+$ , so  $\tilde{f} \in F_1(\widetilde{N^+})$ .

Conversely, given any  $\tilde{f} \in F_1(\widetilde{N^+})$ , define  $\psi(\sigma) = \tilde{f}(\sigma^\top \sigma)$ . Then, for any  $q \in R^*$ , we have

$$\psi(q\sigma) = \tilde{f}((q\sigma)^\top (q\sigma)) = \tilde{f}(q^2(\sigma^\top \sigma)) = q^2\tilde{f}(\sigma^\top \sigma) = q^2\psi(\sigma). \quad (4.26)$$

Thus  $\psi$  satisfies (4.25), so  $\psi \in F_2(\mathcal{S})$ . This completes the proof.

Propositions (4) and (5) lead to

**Proposition 6** *There are the bijections*

$$F(P(N^+)) \leftrightarrow F_1(N^+) \leftrightarrow F_1(\widetilde{N^+}) \leftrightarrow F_2(\mathcal{S}).$$

### 4.3 The spinor version of $B^{2,1}(N^+)_c$

The relationship between  $g \in G$  and the image  $\Lambda \in SO(2,1)$  is given by

$$g^\top b(x)g = b(x\Lambda). \quad (4.27)$$

The action (3.1) restricts to an action of  $G$  on  $\widetilde{N^+}$ . For  $x \in N^+$ ,  $b(x) = \sigma^\top \sigma$ ,  $\sigma \in \mathcal{I}$ . Then equation (4.27) reads

$$g^\top \sigma^\top \sigma g = b(x\Lambda). \quad (4.28)$$

Thus

$$b(x\Lambda) = (\sigma g)^\top (\sigma g). \quad (4.29)$$

Since the identification between  $F_1(N^+)$  and  $F_1(\widetilde{N^+})$  is given by

$$f(x) = \tilde{f}(b(x)), \quad (4.30)$$

we may now compute the representation  $T(g)$  in terms of  $F_1(\widetilde{N^+})$  as follows. We have

$$(T(\gamma(g))f)(x) = (T(\Lambda)f)(x) = f(x\Lambda) = \tilde{f}(b(x\Lambda)) = \tilde{f}((\sigma g)^\top (\sigma g)) = \psi(\sigma g). \quad (4.31)$$

Now define a new representation  $T$  of  $G$  in  $F_2(\mathcal{I})$  by the following formula

$$(T(g)\psi)(\sigma) = \psi(\sigma g). \quad (4.32)$$

This representation reduces to the identity on the subgroup  $\{Id, -Id\}$  of  $G$  (that is on the kernel of the homomorphism from  $G$  onto  $SO(2,1)$ ). We have just seen that it corresponds to the representation  $T(g)$  of  $G$  on  $F_1(N^+)$  defining  $B^{2,1}(N^+)_c$ . Hence

**Theorem 1** *The spinor version of  $B^{2,1}(N^+)_c$  is defined by*

$$B_s^{2,1}(N^+)_c = F_2(\mathcal{I}) \otimes_T G, \quad (4.33)$$

where the representation  $T$  is given by

$$(T(g)\psi)(\sigma) = \psi(\sigma g).$$

### 4.4 Unconstrained supertranslations

In considering representation theory, it is convenient to work in terms of unconstrained functions. To this end we find a new realization of  $B_s^{2,1}(N^+)_c$  with  $F_2(\mathcal{I})$  realized as  $F(P(N^+))$ .

First we introduce convenient local coordinates into  $\mathcal{I}$ . Given  $\sigma = (\sigma_1, \sigma_2) \in \mathcal{I}$ , assume that  $\sigma_2 \neq 0$ . Let  $r(\sigma)$  be the length of  $\sigma$  and let  $\mathcal{S}(\sigma_2)$  be the sign of  $\sigma_2$ :

$$r(\sigma) = (\sigma_1^2 + \sigma_2^2)^{1/2}, \quad \mathcal{S}(\sigma_2) = \frac{\sigma_2}{|\sigma_2|}. \quad (4.34)$$

Further define

$$\rho = \frac{\sigma_1}{\sigma_2}, \quad w = w(\rho) = \frac{r}{|\sigma_2|} = (\rho^2 + 1)^{1/2}. \quad (4.35)$$

Thus  $\sigma$  can be written as

$$\sigma = (\sigma_1, \sigma_2) = r \left( \frac{\sigma_1}{r}, \frac{\sigma_2}{r} \right) = r \mathcal{S}(\sigma_2) \left( \frac{\rho}{w(\rho)}, \frac{1}{w(\rho)} \right), \quad (4.36)$$

and so, writing the unit vector in the final brackets as  $e(\rho)$ , we have

$$\sigma = r(\sigma) \mathcal{S}(\sigma_2) e(\rho). \quad (4.37)$$

The vector  $e(\rho)$  has the second component positive. Evidently  $\rho$  is a local projective coordinate for  $P(N^+)$ .

Using the homogeneity condition (4.25) we have

$$\psi(\sigma) = \psi(r(\sigma) \mathcal{S}(\sigma_2) e(\rho)) = r^2(\sigma) \mathcal{S}^2(\sigma_2) \psi(e(\rho)). \quad (4.38)$$

Writing  $\psi(e(\rho)) = \alpha(\rho)$  this expresses (locally) every  $\psi \in F_2(\mathcal{S})$  in terms of a function  $\alpha \in F(P(N^+))$  by

$$\psi(\sigma) = r^2(\sigma) \alpha(\rho). \quad (4.39)$$

To find the expression for the  $T(g)$  operators in terms of the  $\alpha$ s note first that if  $g \in G$  is

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

then the components  $\sigma_1, \sigma_2$  of  $\sigma$  transform linearly, so that the ratio  $\rho = \frac{\sigma_1}{\sigma_2}$  transforms fraction linearly. Writing  $\rho g$  for the transformed ratio,

$$\rho g = \frac{(\sigma g)_1}{(\sigma g)_2} = \frac{\sigma_1 a + \sigma_2 c}{\sigma_1 b + \sigma_2 d} = \frac{\rho a + c}{\rho b + d}. \quad (4.40)$$

Hence we have

$$\psi(\sigma g) = r^2(\sigma g) \alpha(\rho g). \quad (4.41)$$

The action  $T$  of  $G$  on the  $\psi$ s in Theorem 1 induces an action, also denoted  $T$ , of  $G$  on the  $\alpha$ s defined by

$$(T(g)\psi)(\sigma) = r^2(\sigma) (T(g)\alpha)(\rho). \quad (4.42)$$

The last two equations give

$$(T(g)\alpha)(\rho) = \kappa_g(\rho) \alpha(\rho g), \quad (4.43)$$

where the factor  $\kappa_g(\rho)$  on the right is defined by

$$\kappa_g(\rho) = \frac{r^2(\sigma g)}{r^2(\sigma)} = \frac{(a\sigma_1 + c\sigma_2)^2 + (b\sigma_1 + d\sigma_2)^2}{\sigma_1^2 + \sigma_2^2} = \frac{(\rho a + c)^2 + (\rho b + d)^2}{\rho^2 + 1}. \quad (4.44)$$

Note that the last two formulae are expressed entirely in terms of the local coordinate  $\rho$  for  $P_1(R)$ . Strictly speaking one more local chart is needed to cover all of  $P_1(R)$  related to the above via  $\frac{1}{\rho}$ , but the single one used above will be sufficient here.

Summarising, we have

**Theorem 2** *The group  $B_s^{2,1}(N^+)_c$  can be realized as*

$$B_s^{2,1}(N^+)_c = F(P(N^+)) \otimes_{TG} \quad (4.45)$$

*with semidirect product specified by*

$$(T(g)\alpha)(\rho) = \kappa_g(\rho) \alpha(\rho g). \quad (4.46)$$

## 4.5 Almost unconstrained supertranslations

We now find a new model for  $F(P(N^+))$  on the double cover of  $P_1(R)$ . With each vector  $\sigma = (\sigma_1, \sigma_2) \in \mathcal{S}$  we associate the unit length vector  $e_\sigma = \sigma/|\sigma|$  having the same direction as  $\sigma$ . Thus we have

$$\sigma = r(\sigma)e_\sigma, \quad r(\sigma) = |\sigma|, \quad e_\sigma = \sigma/|\sigma|. \quad (4.47)$$

Let  $S^1 \subset \mathcal{S}$  be the set of vectors of unit length in  $\mathcal{S}$ :

$$S_m^1 = \{\sigma \in \mathcal{S} \mid |\sigma| = 1\}. \quad (4.48)$$

Let  $F_e(S^1)$  be the set of all real valued *even* functions  $\beta : S^1 \rightarrow R$ . That is these functions satisfy the even-ness condition

$$\beta(-e_\sigma) = \beta(e_\sigma). \quad (4.49)$$

With each  $\psi \in F_2(\mathcal{S})$  we associate a function  $\beta \in F_e(S^1)$  by restricting  $\psi$  to the circle  $S^1$ :

$$\beta(e_\sigma) = \psi(e_\sigma). \quad (4.50)$$

Using the homogeneity condition (4.25), we have

$$\psi(\sigma) = \psi(r(\sigma)e_\sigma) = r^2(\sigma)\psi(e_\sigma) = r^2(\sigma)\beta(e_\sigma). \quad (4.51)$$

Hence we can write

$$\beta(e_\sigma) = \frac{\psi(\sigma)}{r^2(\sigma)}. \quad (4.52)$$

Formula (4.52) associates not only any  $\psi$  with the corresponding  $\beta$ , but also vice versa. So we have a bijection

$$F_2(\mathcal{S}) \leftrightarrow F_e(S^1), \quad (4.53)$$

giving us a new realisation of the supertranslation space  $F_2(\mathcal{S})$ . In this new realisation, the supertranslations are ‘‘almost unconstrained’’, they are subject to the condition that they must be even.

Hence we can now calculate the representation  $T(g)$  in this new realisation  $F_e(S^1)$ . The transformation  $\sigma \mapsto \sigma' = \sigma g$  becomes, in the coordinates defined by (4.47),  $r(\sigma) \mapsto r(\sigma')$  and  $e_\sigma \mapsto e_{\sigma'}$  where

$$r(\sigma') = |\sigma'| = |\sigma g| = |r(\sigma)e_\sigma g| = r(\sigma)|e_\sigma g|, \quad \text{and}, \quad (4.54)$$

$$e_{\sigma'} = e_{\sigma g} = (\sigma g)/|\sigma g| = (r(\sigma)e_\sigma g)/|r(\sigma)e_\sigma g| = (e_\sigma g)/|e_\sigma g|. \quad (4.55)$$

Let us denote  $T(g)\psi$  by  $\psi'$ , and, correspondingly,  $T(g)\beta$  by  $\beta'$ . Then, using (5.12), (4.50), (4.52), and (4.54), (4.55), we get

$$\begin{aligned} (T(g)\beta)(e_\sigma) &= \beta'(e_\sigma) = \frac{\psi'(\sigma)}{r^2(\sigma)} = \frac{\psi(\sigma g)}{r^2(\sigma)} = \frac{\psi(\sigma')}{r^2(\sigma)} = \frac{\psi(r(\sigma')e_{\sigma'})}{r^2(\sigma)} = \frac{r^2(\sigma')}{r^2(\sigma)}\psi(e_{\sigma'}) \\ &= \frac{r^2(\sigma')}{r^2(\sigma)}\beta(e_{\sigma'}) = \kappa_g(e_\sigma)\beta(e_{\sigma'}) = \kappa_g(e_\sigma)\beta(e_{\sigma g}), \end{aligned} \quad (4.56)$$

where the factor  $\kappa_g(e_\sigma)$  on the right is defined by

$$\kappa_g(e_\sigma) = \frac{r^2(\sigma')}{r^2(\sigma)} = \frac{|\sigma g|^2}{|\sigma|^2} = |e_\sigma g|^2. \quad (4.57)$$

From (4.44) and (4.57) we have

$$\kappa_g(e_\sigma) = \kappa_g(\rho). \quad (4.58)$$

We conclude that

**Theorem 3** *The group  $B_s^{2,1}(N^+)_c$  can be realized as*

$$B_s^{2,1}(N^+)_c = F_e(S^1) \otimes_T G \quad (4.59)$$

with semidirect product given by

$$(T(g)\beta)(e_\sigma) = \kappa_g(e_\sigma)\beta(e_\sigma g). \quad (4.60)$$

We can use either of the realizations given in Theorems 2 and 3 to develop the representation theory of  $B_s^{2,1}(N^+)_c$ . Here we use the realization given in Theorem 2.

## 5 The group $B(2, 1)$

So far, the supertranslation space  $F(P(N^+))$  has been defined as a space of truly arbitrary functions. This has been merely for clarity; for physical applications it is necessary to give to this space additional structure. To this end we now restrict  $F(P(N^+))$  to be the Hilbert space of square integrable functions on  $P(N^+)$  with respect to the standard (Lebesgue) measure.

It will be convenient then to use as coordinates for  $P(N^+)$  the two components of a unit length vector  $\mathbf{m}$  in  $R^2$

$$P(N^+) = \{\mathbf{m} \in R^2 \mid |\mathbf{m}| = 1\}. \quad (5.1)$$

The  $\mathbf{m}$  coordinates are globally defined. The usual angular coordinate  $\theta$  for  $P(N^+)$  is defined by

$$\mathbf{m} = (\sin\theta, \cos\theta) = (\mathbf{m}_1, \mathbf{m}_2), \quad (5.2)$$

$\theta \in [0, 2\pi)$ . The angular coordinate  $\theta$  is related to the projective coordinate  $\rho$  for  $P(N^+)$  by stereographic projection

$$\rho = \cot\left(\frac{\theta}{2}\right). \quad (5.3)$$

The required measure  $\lambda$  on  $P(N^+)$  is the usual (normalized) length of an arc segment for  $S^1$ . The explicit form is the 1-form

$$d\lambda(\mathbf{m}) = \frac{d\mathbf{m}_1}{2\pi\mathbf{m}_2}, \quad (5.4)$$

where  $\mathbf{m}_1, \mathbf{m}_2$  are the components of a unit vector  $\mathbf{m} \in R^2$ . This formula is only valid for  $\mathbf{m}_2 \neq 0$ , but a permutation of the indices in the components gives another valid expression for  $d\lambda(\mathbf{m})$ . In terms of  $\theta$  the normalized Lebesgue measure is given by

$$d\lambda(\theta) = \frac{d\theta}{2\pi} \quad (5.5)$$

which evidently is  $SO(2)$ -invariant. In terms of  $\rho$  the corresponding expression is

$$d\lambda(\rho) = -\frac{d\rho}{\pi(1+\rho^2)}. \quad (5.6)$$

Letting  $\mathbf{m}g$  be the transform of  $\mathbf{m}$  by  $g \in G$  (so that  $\rho \mapsto \rho g$  corresponds to  $\mathbf{m} \mapsto \mathbf{m}g$ ), a simple calculation gives

$$\frac{d\lambda(\mathbf{m})}{d\lambda(\mathbf{m}g)} = \kappa_g(\rho) = \kappa_g(\mathbf{m}). \quad (5.7)$$

Here  $\kappa_g(\mathbf{m})$  denotes  $\kappa_g(\rho)$ , where  $\mathbf{m}$  corresponds to  $\rho$ . While the expression given for  $d\lambda(\mathbf{m})$  is local,  $\kappa_g(\mathbf{m})$  is defined globally.

Let  $L^2(P_1(R), \lambda, R)$  be the space of all functions  $\alpha : P_1(R) \rightarrow R$  which are Lebesgue measurable and satisfy

$$\int_{P_1(R)} |\alpha(\rho)|^2 d\lambda(\rho) < +\infty. \quad (5.8)$$

Introduce an inner product into  $L^2(P_1(R), \lambda, R)$  by setting, for each  $\alpha_1, \alpha_2 \in L^2(P_1(R), \lambda, R)$ ,

$$\langle \alpha_1, \alpha_2 \rangle = \int_{P_1(R)} \alpha_1(\rho)\alpha_2(\rho) d\lambda(\rho). \quad (5.9)$$

Then, as is well known,  $L^2(P_1(R), \lambda, R)$  becomes a real Hilbert space with respect to this inner product in the usual way. We denote the resulting Hilbert space by  $\mathcal{H}(P_1(R))$ . In  $\mathcal{H}(P_1(R))$  functions are identified whenever they differ, at most, on a set of measure zero. We use the homeomorphism  $P(N^+) \simeq P_1(R)$  to identify the spaces  $L^2(P(N^+), \lambda, R)$  and  $L^2(P_1(R), \lambda, R)$ , and consequently, the ensuing Hilbert spaces  $\mathcal{H}(P(N^+))$  and  $\mathcal{H}(P_1(R))$  respectively.

We now replace in the definition (4.45) the supertranslation space  $F(P(N^+))$  by  $\mathcal{H}(P_1(R))$ . Note that the representation  $T$ , given by (4.43), restricts to the subspace  $\mathcal{H}(P_1(R))$  and it is given by the same formula as in (4.43). So, our final definition of our group, which we will, now and henceforth, denote by  $B(2, 1)$  (to distinguish it from previous definitions), is the group

**Definition 1**

$$B(2, 1) = \mathcal{H}(P_1(R)) \otimes_T G. \quad (5.10)$$

with semidirect product specified by

$$(T(g)\alpha)(\rho) = \kappa_g(\rho)\alpha(\rho g).$$

We can also use the bijections between all the different models for supertranslation space to restrict all of them to functions corresponding to  $L^2(P_1(R), \lambda, R)$  and to define corresponding Hilbert space structures for all of them. Denoting the resulting spaces by  $\mathcal{H}$ , and using subscripts, as before, to indicate constraints, we now have

$$\mathcal{H}(P(N^+)) \leftrightarrow \mathcal{H}_1(N^+) \leftrightarrow \mathcal{H}_1(\widetilde{N^+}) \leftrightarrow \mathcal{H}_2(\mathcal{I}). \quad (5.11)$$

We note that there are two distinct transitive actions of  $G$  on  $S^1$ .

1. The action (4.40) on  $P_1(R)$  induced by the usual linear action of  $G$  on  $R^2$ . The stabilizer of  $\infty$  is the subgroup  $\mathcal{U}$  of  $G$  of upper triangular matrices. Therefore,  $P_1(R)$  is in bijective correspondence with and homeomorphic to the quotient space  $G/\mathcal{U}$ . In this case the action (4.40) of  $G$  on  $P_1(R)$  is not faithful; the kernel of the action is the group  $\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \right\} \simeq Z_2$ . This action was used to define the representation  $T$  in (4.43) which specifies the semi-direct product  $B(2, 1)$  in (5.10).
2. The action on the circle  $(R^2 - \{0\})/R_{>0}$  inferred from (4.55) is

$$e_\sigma g = e_{\sigma'} = (e_\sigma g)/|e_\sigma g|. \quad (5.12)$$

If  $e_\sigma = (x \ y)$ ,  $x$  and  $y$  are the cartesian coordinates of  $e_\sigma$ , action (5.12) reads

$$(x \ y) \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \frac{1}{\sqrt{(ax + cy)^2 + (bx + dy)^2}} (ax + cy \ bx + dy). \quad (5.13)$$

The stabilizer of  $(1 \ 0)$  for this action is  $\mathcal{U}^+$ , the subgroup of index 2 of  $\mathcal{U}$  with positive entries in the main diagonal. Therefore the circle  $(R^2 - \{0\})/R_{>0}$  is in bijective correspondence with and homeomorphic to the quotient space  $G/\mathcal{U}^+$ . In this case the action of  $G$  on  $G/\mathcal{U}^+$  is faithful. This action was used to define the representation  $T$  in (4.60) which specifies the semi-direct product  $B_s^{2,1}(N^+)_c$  in (4.59).

There is a continuous  $G$ -equivariant surjection  $G/\mathcal{U}^+ \rightarrow G/\mathcal{U}$  which is a connected 2-fold covering.

## 6 $B(2, 1)$ as the automorphism group of the strong conformal geometry of $\mathcal{I}^+$

According to Klein's Erlangen programme, one may (indirectly) specify a geometry by giving a group action. Conversely, given a group action, one may ask for the corresponding geometry. Penrose [3], [4], [5] interpreted the original BMS group  $B$  as a transformation group  $B \times \mathcal{I}^+ \rightarrow \mathcal{I}^+$  of the "future null infinity" of conformally compactified weakly asymptotically simple space-times. Furthermore, he gave a geometric structure to  $\mathcal{I}^+$ , the "strong conformal geometry", such that this classical action  $B \times \mathcal{I}^+ \rightarrow \mathcal{I}^+$  is the group of automorphisms of the geometry. In this section we rederive  $B(2, 1)$  as the automorphism group of the "strong conformal geometry" of the "future null infinity"  $\mathcal{I}^+$  of 3-dim Minkowski space-time.



In dealing with  $B(2, 1)$  it has been convenient to specify the action  $T$  of  $G$  on  $\mathcal{S}$  from the right. In order to agree with notations used elsewhere (e.g. [4], [5]), it now becomes convenient to use the action of  $G$  on  $\mathcal{S}$  from the left. So, for this and the next section only, we redefine the groups and quantities needed. Let  $g \in G$  and write

$$g = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \quad \sigma = \begin{bmatrix} \sigma_1 \\ \sigma_2 \end{bmatrix}, \quad g\sigma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} \sigma_1 \\ \sigma_2 \end{bmatrix}, \quad (6.1)$$

$$\rho = \frac{\sigma_1}{\sigma_2}, \quad g\rho = \frac{a\rho + b}{c\rho + d}. \quad (6.2)$$

### 6.1 “Future null infinity” $\mathcal{S}^+$ of 3–dim Minkowski space–time

The following definition will be useful

**Definition 2** *The 3–dim Einstein universe is a solution to Einstein’s equations with zero cosmological constant in 3 space–time dimensions with metric  $\mathcal{M}$  given by*

$$ds^2 = -dT^2 + d\Psi^2 + \sin^2\Psi d\phi^2, \quad (6.3)$$

$$\text{where, } -\infty < T < \infty, \quad 0 \leq \Psi < \pi, \quad 0 \leq \phi < 2\pi. \quad (6.4)$$

Einstein’s 3–dim universe contains dust with constant proper energy density  $\rho$ ,  $\rho = \frac{1}{\kappa}$ ,  $\kappa = \frac{8\pi G}{c^4}$ . It has four killing vector fields

$$\xi_1 = \partial_T, \quad \xi_2 = \partial_\phi, \quad \xi_3 = -(\cos\phi\partial_\Psi - \cot\Psi\sin\phi\partial_\phi), \quad (6.5)$$

$$\xi_4 = \sin\phi\partial_\Psi + \cot\Psi\cos\phi\partial_\phi. \quad (6.6)$$

$\xi_2, \xi_3, \xi_4$  are the three killing vector fields of the 2–sphere.  $\xi_1$  is a global timelike Killing vector field, orthogonal to the level hypersurfaces of  $T$  which are 2–spheres. Therefore, 3–dim Einstein universe is static with topology  $R \times S^2$ . 3–dim Einstein universe will be denoted by  $Ein_3$ ,  $Ein_3 = (R \times S^2, \mathcal{M})$ .

We obtain

$$\mathcal{M} = \Omega^2\eta, \quad \text{where,} \quad (6.7)$$

$$\Omega = \frac{2}{\sqrt{(1+u^2)(1+v^2)}}, \quad (6.8)$$

$$\begin{aligned} \eta &= \frac{(1+u^2)(1+v^2)}{4} (-dT^2 + d\Psi^2) + \frac{(u-v)^2}{4} d\phi^2 \\ &= -dudv + \frac{(u-v)^2}{4} d\phi^2 \\ &= -dt^2 + dr^2 + r^2 d\phi^2, \end{aligned} \quad (6.9)$$

$$T = \arctanu + \arctanv, \quad \Psi = \arctanv - \arctanu, \quad (6.10)$$

$$u = t - r, \quad v = t + r, \quad (6.11)$$

$$r \geq 0, \quad -\infty < t, \quad u, \quad v < \infty. \quad (6.12)$$

The metric  $\eta$  is the metric of the 3–dim Minkowski space–time  $Min_3$ ,  $Min_3 = (R^3, \eta)$ .

By using equations (6.7)-(6.12) we are easily led to the following

**Proposition 7** *There exists a conformal isometry of  $Min_3$  into the open region  $Min_3$  of  $Ein_3$  given by the coordinate restrictions*

$$-\pi < T < \pi, \quad 0 \leq \Psi < \pi, \quad 0 \leq \phi < 2\pi. \quad (6.13)$$

**Definition 3** *The conformal infinity of  $Min_3$  is the boundary  $\mathcal{B}_{Min_3}$  of  $Min_3$  in  $Ein_3$ .*

Let  $\overline{Min_3}$  be the closure of  $Min_3$ , i.e.,  $\overline{Min_3} = Min_3 \cup \mathcal{B}_{Min_3}$ .

We can easily prove the following

**Lemma 1** *The 2-dim surface*

$$\mathcal{I}^+ = \{(T, \Psi, \phi) : T = \pi - \Psi, \quad 0 < \Psi < \pi, \quad 0 \leq \phi < 2\pi\}$$

*in  $\mathcal{B}_{Min_3}$  is smooth null hypersurface for  $\mathcal{M}$  (hence the terminology “null infinity”). Its null generators are the integral curves of the vector field  $\partial_T - \partial_\Psi$ .*

Conformal “compactification” is not always possible. The property for a space–time to admit a smooth conformal compactification can be characterized in terms of decay of the Weyl curvature at infinity. When such a compactification exists the boundary, which is  $\mathcal{B}_{Min_3}$  in the case under consideration, will have a structure: different parts corresponding to different ways of going to infinity in the physical space–time.  $\mathcal{I}^+$  is the subset of the boundary, i.e., of  $\mathcal{B}_{Min_3}$  in our case, with elements the end points of all future directed null geodesics of  $Min_3$  (hence the terminology “future infinity”).

Conformal “compactification” when possible is not unique: There is a freedom [5] in the choice of  $\Omega$  in (6.7). Indeed,  $\Omega$  can be replaced by any

$$\Omega' = \Theta\Omega, \quad (6.14)$$

$\Theta$  being any smooth positive real valued function on  $\overline{Min_3}$ . As a result of this freedom, a conformal class  $\mathcal{C}_{Min_3}$  of metrics is defined on  $Min_3$ . Thus  $\mathcal{C}_{Min_3}$  consists of all metrics  $\Omega^2\mathcal{M}$ , where  $\Omega$  is a smooth positive real valued function on  $Min_3$ , which vanishes on  $\mathcal{I}^+$ , and has non–zero divergence on  $\mathcal{I}^+$ .

By restricting  $\mathcal{M}$  to  $\mathcal{I}^+$  we obtain that the induced metric on  $\mathcal{I}^+$  is given by

$$dl^2 = -0d\Psi^2 + \sin^2\Psi d\phi^2, \quad (6.15)$$

$$\text{where,} \quad 0 \leq \Psi < \pi, \quad 0 \leq \phi < 2\pi. \quad (6.16)$$

We can use the freedom in the choice of  $\Omega$  to scale the metric (6.15) so that the divergence of the null generators of  $\mathcal{I}^+$  vanishes. In fact one can easily prove the following

**Lemma 2** *With a proper choice of  $\Omega$  the induced metric on  $\mathcal{I}^+$  can be cast into the form  $\mathcal{M}_{\mathcal{I}^+}$  given by*

$$dl^2 = 0du^2 + d\phi^2 = 0du^2 + \frac{4}{(1+\rho^2)^2}d\rho^2 \quad (6.17)$$

$$\text{where,} \quad \Psi = \frac{\pi}{2} - \arctan u, \quad -\infty < u < \infty, \quad (6.18)$$

$$\rho = \frac{\cot\phi}{2}, \quad -\infty < \rho < \infty. \quad (6.19)$$

Therefore, in the  $(u, \phi)$  coordinates, the null generators of  $\mathcal{I}^+$ , affinely parameterized by  $u$ , are the integrals curves of the vector field  $\frac{\partial}{\partial u}$ . The part of the metric (6.17) that is transverse to  $\frac{\partial}{\partial u}$  is the metric of a circle with unit radius. Therefore,  $\mathcal{I}^+$  has topology  $R \times S^1$ , “ $R$ ” being the null generators of  $\mathcal{I}^+$ , “ $S^1$ ” being the circle of unit radius.

$\mathcal{C}_{Min_3}$  induces, via (6.7), a conformal class  $\mathcal{C}_{\mathcal{I}^+}$  of metrics on  $\mathcal{I}^+$ .  $\mathcal{C}_{\mathcal{I}^+}$  consists of all metrics  $\frac{1}{\Omega^2} \mathcal{M}_{\mathcal{I}^+}$ , where  $\Omega$ , as in  $\mathcal{C}_{Min_3}$ , is a smooth positive real valued function on  $Min_3$ , which vanishes on  $\mathcal{I}^+$ , and has non-zero divergence on  $\mathcal{I}^+$ .

## 6.2 The Newman–Unti group

The *local* orientation-preserving conformal maps of the plane onto itself are given by

$$\zeta \longrightarrow \bar{\zeta} = f(\zeta)$$

where  $f$  is holomorphic and  $\zeta = \sigma_1 + i\sigma_2$ ,  $\sigma_1, \sigma_2$  being Cartesian coordinates on the plane.

However, for a *global* map, we would require that both  $f$  and its inverse  $f^{-1}$  be non-singular over the whole plane. This restricts  $f$  to be a linear function

$$f(\zeta) = w\zeta + q. \quad (6.20)$$

Therefore, the group (6.20) of *global* orientation-preserving conformal maps of the plane onto itself is a four-real-parameter group. Writing  $w = |w|e^{i\vartheta}$  in polar coordinates, we see that the affine transformations (6.20) can be viewed as the composition of a rotation, followed by a scaling, followed by a translation.

The linear action (6.20) induces a fraction linear action on  $P_1(R)$

$$\rho \longrightarrow \bar{\rho} = g(\rho) = \frac{a\rho + b}{c\rho + d}, \quad (6.21)$$

$\rho = \frac{\sigma_1}{\sigma_2}$ ,  $a, b, c, d \in R$ . The three real ratios  $a : b : c : d$  define  $g$ . We can normalize  $a, b, c, d$  by

$$ad - bc = 1. \quad (6.22)$$

Thus every map  $g$  corresponds to an element of  $PSL(2, R)$  and vice versa.

One can easily prove the following

**Proposition 8** *The group of non-reflective motions of  $\mathcal{I}^+$  preserving its intrinsic degenerate metric*

$$dl^2 = 0du^2 + \frac{4}{(1 + \rho^2)^2} d\rho^2$$

$$\text{where, } \quad -\infty < u < \infty, \quad -\infty < \rho < \infty,$$

up to conformal rescalings is given by

$$\rho \longrightarrow \bar{\rho} = \frac{a\rho + b}{c\rho + d}, \quad (6.23)$$

$$u \longrightarrow \bar{u} = F(u, \rho), \quad (6.24)$$

$$\text{where, } \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, R), \quad \frac{\partial F}{\partial u} > 0. \quad (6.25)$$

$F$  is appropriately smooth on the whole of  $\mathcal{S}^+ \simeq R \times S^1$ .  $F$  is monotonic increasing in  $u$  for each  $\rho$ , mapping the entire range for  $u$  for each  $\rho$ -generator to itself, and with non-vanishing  $u$ -derivative so that the inverse transformation is also smooth. The condition  $\frac{\partial F}{\partial u} > 0$  ensures that the transformation (6.24) preserves the orientation on each generator of  $\mathcal{S}^+$ . The group of transformations (6.23), (6.24), (6.25) is the Newman–Unti (NU) group in 3 space–time dimensions. The analogue of the NU group in 4 space–time dimensions has been defined in [4], [5], [35].

$\mathcal{C}_{\mathcal{S}^+}$  is determined, via (6.7), by  $\mathcal{C}_{Min_3}$  and that is certainly preserved by  $P(2, 1)$ , the Poincare group in 3 space–time dimensions. Thus  $P(2, 1)$  is a subgroup of the NU group.

### 6.3 Self–diffeomorphisms of $\mathcal{S}^+$ which preserve its strong conformal geometry

Following Penrose [4] it is possible to assign more geometric structure to  $\mathcal{S}^+$  than we have given hitherto. The preservation of this additional structure will reduce the freedom in  $F(u, \rho)$  in (6.24) to that of a function of  $\alpha(\rho)$  only. This will give us the BMS group  $B(2, 1)$ .

To this end it is natural to seek for quantities which are invariants of the class  $\mathcal{C}_{\mathcal{S}^+}$ , i.e., are the same for all members of the class  $\mathcal{C}_{\mathcal{S}^+}$ . In fact we can easily prove the following

**Proposition 9** *The ratio  $\frac{du}{dl}$  is independent of the choice of  $\Omega$ .*

The infinitesimal increments  $du$  and  $dl$  were defined respectively in Lemma 2 and in (6.15), (6.17).

We can use this invariance to define the concept of “null angle”  $\nu$  [4] between two tangent directions at a point  $P$  of  $\mathcal{S}^+$ , whose span contains the null normal direction to  $\mathcal{S}^+$ , by

$$\nu = \frac{du}{dl}. \quad (6.26)$$

The induced degenerate metric (6.17) supplemented by the concept of null angle, provides the “strong conformal geometry” [4] at  $\mathcal{S}^+$ .

We seek to find the group of self–diffeomorphisms of  $\mathcal{S}^+$  which preserves the “strong conformal geometry” at  $\mathcal{S}^+$ , i.e., the group of diffeomorphisms of  $\mathcal{S}^+$  to itself which preserves angles and leaves invariant the ratio  $\frac{du}{dl}$ .

The allowed diffeomorphisms must have the form (6.23), (6.24), (6.25) since these are the ones preserving the inner conformal metric but the function  $F$  must now have the special form that allows the ratio  $\frac{du}{dl}$  to remain invariant.

It is an easy exercise to prove that under (6.23), (6.24)

$$dl \longrightarrow Kdl, \quad \text{where} \quad K = \frac{1 + \rho^2}{(a\rho + b)^2 + (c\rho + d)^2}. \quad (6.27)$$

We must therefore also have

$$du \longrightarrow Kdu. \quad (6.28)$$

Since  $K$  is independent of  $u$ , this means  $u$  transforms according to

$$u \longrightarrow K(u + \alpha(\rho)), \quad (6.29)$$

where  $\alpha(\rho)$  is some real-valued function defined on  $S^1$ .

Let  $\mathcal{S}(S^1)$  be the space of real-valued functions defined on  $S^1$ . The Theorem which follows summarizes the previous results.

**Theorem 4** *The group of self-diffeomorphisms of  $\mathcal{I}^+$  which preserves the “strong conformal geometry” at  $\mathcal{I}^+$  is given by*

$$\rho \longrightarrow \bar{\rho} = \frac{a\rho + b}{c\rho + d}, \quad (6.30)$$

$$u \longrightarrow \bar{u} = \frac{1 + \rho^2}{(a\rho + b)^2 + (c\rho + d)^2} (u + \alpha(\rho)), \quad (6.31)$$

$$\text{where, } \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, R), \quad \alpha(\rho) \in \mathcal{S}(S^1). \quad (6.32)$$

Two remarks are in order regarding Theorem 4:

1. Historically, the analogue of the classical action (6.30), (6.31) on  $\mathcal{I}^+$  in the case of the original BMS group  $B$  was given in [2] Equation (3.12), [36] Equation (II.23), [3] Equation (3.14).
2. Since truly arbitrary supertranslation functions  $\alpha(\rho)$  in (6.31) describe symmetry transformations in Penrose’s sense, supertranslations need not have some minimum degree of smoothness.

Let  $\mathcal{B}$  be the set of pairs  $(\alpha(\rho), g)$ ,  $\alpha(\rho) \in \mathcal{S}(S^1)$ ,  $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, R)$ . Application of two consecutive self-diffeomorphisms

$$\left\{ \begin{array}{l} \rho \longrightarrow \bar{\rho} \\ u \longrightarrow \bar{u} \end{array} \right\}, \quad \left\{ \begin{array}{l} \bar{\rho} \longrightarrow \bar{\bar{\rho}} \\ \bar{u} \longrightarrow \bar{\bar{u}} \end{array} \right\}, \quad (6.33)$$

defines an operation between the elements of  $\mathcal{B}$ . It appears that  $\mathcal{B}$  cannot be identified with  $B(2, 1)$  since the action  $T_{\mathcal{B}}$ , resulting from the two consecutive self-diffeomorphisms (6.33), which defines the operation between the elements of  $\mathcal{B}$ , is not identical to the action  $T$ , given by (4.43), (4.44), which defines  $B(2, 1)$ .

In order to bring home the identity of  $\mathcal{B}$  with  $B(2, 1)$  a convenient reformulation of (6.30), (6.31) is needed. This is given in section 7. Not surprisingly, the reformulation needed, is the analogue of Equation (50) in [1], where  $B$  firstly appeared as a pseudo-group of local diffeomorphisms.

#### 6.4 “Null angles”

Let  $P$  be a point of  $\mathcal{I}^+$  and consider two non-null (spacelike) tangent oriented directions  $\gamma$  and  $\delta$  to  $\mathcal{I}^+$  at  $P$ . Let

$$N_{\alpha} = -\nabla_{\alpha}\Omega \quad (6.34)$$

near  $\mathcal{I}^+$ . The vector field  $N_\alpha$  is normal to the null hypersurface  $\mathcal{I}^+$ . Because  $N_\alpha$  is orthogonal to itself ( $N^\alpha N_\alpha = 0$ ), this vector field is also tangent to  $\mathcal{I}^+$ . The sign in (6.34) is chosen so that  $N^\alpha$  is future-directed when  $\Omega$  increases toward the future.

In 4 space-time dimensions there are two types of angle [5] that arise between  $\gamma$  and  $\delta$ . The first type occurs when the plane defined by  $\gamma$  and  $\delta$  does not contain  $N$ . The angle between  $\gamma$  and  $\delta$  is measured with the intrinsic degenerate metric on  $\mathcal{I}^+$ . The second type arises when the plane defined by  $\gamma$  and  $\delta$  contains  $N$  (neither of  $\gamma$  and  $\delta$  being actually in the direction of  $N$ ). One can easily prove that in this case the angle between  $\gamma$  and  $\delta$  measured with the intrinsic degenerate metric on  $\mathcal{I}^+$  is always zero.

Although the angle between  $\gamma$  and  $\delta$  in this case is always zero, there is still an invariant concept of separation between  $\gamma$  and  $\delta$ . This concept is precisely the concept of “null angle”.  $B$  is the group of self-diffeomorphisms of  $\mathcal{I}^+$  which leaves conformally invariant the intrinsic degenerate metric on  $\mathcal{I}^+$  and the “null angle” between any two non-null tangent directions at a point  $P$  of  $\mathcal{I}^+$  which are coplanar with  $N$ .

In three space-time dimensions the situation is different. One can prove the Proposition which follows.

**Proposition 10** *In three space-time dimensions there is only one type of angle that arises between  $\gamma$  and  $\delta$ . The plane spanned by  $\gamma$  and  $\delta$  always contains  $N$ . The angle between  $\gamma$  and  $\delta$  measured with the intrinsic degenerate metric (6.17) on  $\mathcal{I}^+$  is always zero. The “null angle”*

$$\nu = \frac{du}{dl}$$

*gives an invariant concept of separation between  $\gamma$  and  $\delta$ .*

**Proof** *The proof can be given with standard Linear Algebra and Differential Geometry and as such it is omitted.*

$B(2, 1)$  is the group of self-diffeomorphisms of  $\mathcal{I}^+$  which leaves conformally invariant the intrinsic degenerate metric (6.17) on  $\mathcal{I}^+$  and the “null angle” between any two non-null tangent directions  $\gamma$  and  $\delta$  at a point  $P$  of  $\mathcal{I}^+$ .

## 7 The classical action $B(2, 1) \times \mathcal{I}^+ \longrightarrow \mathcal{I}^+$

In order to agree with notations used elsewhere (e.g. [4], [5]) for the classical action  $B \times \mathcal{I}^+ \longrightarrow \mathcal{I}^+$  from the left, it now becomes convenient to use the action of  $G$  on  $\mathcal{I}$  from the left. Now, we use (6.1), (6.2), and

$$r^2(\sigma) = \sigma_1^2 + \sigma_2^2, \quad r^2(g\sigma) = (a\sigma_1 + b\sigma_2)^2 + (c\sigma_1 + d\sigma_2)^2, \quad (7.1)$$

and we redefine

$$\kappa_g(\rho) = \frac{r^2(g\sigma)}{r^2(\sigma)} = \frac{(\rho a + b)^2 + (\rho c + d)^2}{\rho^2 + 1}. \quad (7.2)$$

With these new definitions adapted to the left action of  $G$  on  $\mathcal{I}$  one can now redefine

$$B(2, 1) = \mathcal{H}(P_1(R)) \otimes_T G, \quad (7.3)$$

$$(T(g)\alpha)(\rho) = \kappa_{g^{-1}}(\rho)\alpha(g^{-1}\rho). \quad (7.4)$$

$\mathcal{I}^+$  can be identified with the product  $R \times P_1(R)$ . Let  $[u, \rho] \in R \times P_1(R) = \mathcal{I}^+$ . Then  $(\alpha, g) \in B(2, 1)$  may be obtained as the product  $(\alpha, g) = (\alpha, Id)(0, g)$  (pure Lorentz followed by pure supertranslations). The separate actions are, in order,

$$(0, g)[u, \rho] = [\kappa_g^{-1}(\rho)u, g\rho] \in \mathcal{I}^+, \quad (7.5)$$

$$(\alpha, Id)[u, \rho] = [u + \alpha(\rho), \rho] \in \mathcal{I}^+. \quad (7.6)$$

Combining the two gives

$$(\alpha, g)[u, \rho] = [\kappa_g^{-1}(\rho)u + \alpha(g\rho), g\rho]. \quad (7.7)$$

This last formula, then gives the classical action  $B(2, 1) \times \mathcal{I}^+ \rightarrow \mathcal{I}^+$ . Applying a further  $(\beta, h)$  gives

$$(\beta, h)(\alpha, g) = (\beta + T(h)\alpha, hg), \quad (7.8)$$

where  $T(h)\alpha$  is easily checked to be given by the action (7.4) defining  $B(2, 1)$  in (7.3). We note that  $K = \kappa_g^{-1}(\rho)$ ,  $K, \kappa_g(\rho)$  are given respectively by (6.27) and (7.2). Moreover, (7.7) is a convenient reformulation of (6.30), (6.31). An easy calculation shows

$$(T(g)\alpha, g)[u, \rho] = [\kappa_g^{-1}(\rho)(u + \alpha(\rho)), g\rho]. \quad (7.9)$$

The right-hand side of (7.9) is identical to (6.30), (6.31). We turn now to the study of  $B(2, 1)$ .

## 8 The supertranslation subgroup

For notational convenience, hereafter, we use the symbol  $\mathcal{A}$  for  $\mathcal{H}(P_1(R))$ . As is well known, the set of functions

$$e_0 = \frac{1}{\sqrt{2\pi}}, e_n = \frac{1}{\sqrt{\pi}} \cos n\theta, \varepsilon_n = \frac{1}{\sqrt{\pi}} \sin n\theta, \quad n = 1, 2, \dots, \quad (8.1)$$

is a complete orthonormal basis for the Hilbert space  $\mathcal{A}$ . Therefore, any function  $\alpha \in \mathcal{A}$  may be written as

$$\alpha(\theta) = \sum_{n=0}^{\infty} (c_n e_n + s_n \varepsilon_n). \quad (8.2)$$

The inner product space  $\mathcal{A}$  is an orthogonal direct sum of the subspaces  $\mathcal{T}$  and  $\mathcal{S}$ , the translation and proper supertranslation subspaces respectively,

$$\mathcal{A} = \mathcal{T} \oplus \mathcal{S}, \quad (8.3)$$

these subspaces consisting of the following functions:

$$\mathcal{T} : t(\theta) = \sum_{n=0}^1 (c_n e_n + s_n \varepsilon_n) = a_0 + a_1 \cos \theta + a_2 \sin \theta, \quad (8.4)$$

where the real numbers  $a_0, a_1, a_2$  may be easily found from  $c_0, c_1, s_1$  by using the formulae for the  $e_0, e_1, \varepsilon_1$ ,

$$\mathcal{S} : s(\theta) = \sum_{n=2}^{\infty} (c_n e_n + s_n \varepsilon_n). \quad (8.5)$$

Therefore, every element  $\alpha \in \mathcal{A}$  is uniquely decomposed as

$$\alpha = t + s \quad (8.6)$$

where  $t \in \mathcal{T}$  and  $s \in \mathcal{S}$ . The elements  $t$  and  $s$  are the orthogonal projections of  $\alpha$  on the spaces  $\mathcal{T}$  and  $\mathcal{S}$  respectively:

$$t = P_{\mathcal{T}}(\alpha), \quad s = P_{\mathcal{S}}(\alpha). \quad (8.7)$$

$\mathcal{T}$  is invariant under the action  $T$  of  $G$  on  $\mathcal{A}$ , given by (4.43), whereas  $\mathcal{S}$  is not. Thus the orthogonal direct sum decomposition  $\mathcal{A} = \mathcal{T} \oplus \mathcal{S}$  is not preserved by the  $G$  action. Restricting  $\mathcal{A}$  to the  $G$  invariant subspace  $\mathcal{T}$  restricts  $B(2, 1) = \mathcal{A} \otimes_T G$  to  $P(2, 1) = \mathcal{T} \otimes_T G$ , the double cover of the Poincare group in three space–time dimensions. In fact the 3–tuple  $(a_0, a_1, a_2)$  in the formula for  $t(\theta) \in \mathcal{T}$  is the translation vector of  $P(2, 1)$ , and one quickly checks that the action  $T$  of  $G$  on these vectors gives the natural action of  $G/Z_2 \approx L_+^\uparrow, L_+^\uparrow$  is the connected component of the identity of the Lorentz group in three dimensions, on three–dimensional Minkowski space vectors.

## 9 On the subgroups of $B(2, 1)$

The following Proposition clarifies the relation of  $B(2, 1)$  with  $\mathcal{T}$  and  $\mathcal{A}$ .

### Proposition 11

*The group  $B(2, 1)$  has two normal subgroups and both are abelian:*

1.  $\mathcal{T}$  which is its unique 3–dimensional normal subgroup, and,
2.  $\mathcal{A}$  which is infinite–dimensional.

**Proof** *The proof is similar to the one given by Sachs [36] for the original BMS group  $B$  and as such it is omitted.*

Property 1 identifies  $\mathcal{T}$  uniquely. Obviously, the factor group  $B(2, 1)/\mathcal{A}$  is isomorphic to  $G$ .  $B(2, 1)$  has an infinite–parameter family of Poincare subgroups, all of them on equal footing. Not even the restricted Poincare group  $P(2, 1)$  itself arises naturally as a subgroup of  $B(2, 1)$  in general. It is only in Minkowski space  $\mathcal{M}$  that  $P(2, 1)$  is naturally singled out. In the case of  $\mathcal{M}$  we have some additional structure at future null infinity  $\mathfrak{J}^+$ , namely the notion of which cuts are to be labelled as “good cuts”. These “good cuts”, which form a three–parameter family, are obtained from one another by translations; but a proper supertranslation takes a “good cut” into a “bad cut”. For such a supertranslation  $s \in \mathcal{S}$ , the



six-parameter subgroup of  $B(2, 1)$ ,  $\mathcal{P} = s^{-1}P(2, 1)s$ , will be distinct from  $P(2, 1)$ , though isomorphic with it. For a general  $s$ , the subgroups  $P(2, 1)$  and  $\mathcal{P}$  have only the group of translations  $\mathcal{T}$  in common.

In the case of the original BMS group  $B$ , the infinite-parameter family of Poincare subgroups  $P$  contained in  $B$  [37] underlies a new resolution [38] of the black hole information paradox. In 2+1-dimensional gravity with zero cosmological constant no black holes with event horizons exist [39]. However, as in the four-dimensional case, the infinite-parameter family of Poincare subgroups contained in  $B(2, 1)$ , implies that in 2+1-dimensional quantum gravity with zero cosmological constant the vacuum is not unique. In fact there is an infinite-parameter family of inequivalent vacua which have zero energy but different angular momenta. None of these vacua are preferred, and each is annihilated by a different Poincare subgroup of  $B(2, 1)$ .

$\mathcal{T}$  is a normal subgroup of  $\mathcal{A}$ . The quotient group  $PST = \mathcal{A}/\mathcal{T}$  is naturally isomorphic to  $\mathcal{S}$ , the isomorphism being given by

$$i : PST \rightarrow \mathcal{S}, \quad i(\alpha + \mathcal{T}) = P_{\mathcal{S}}(\alpha), \quad (9.1)$$

where  $\alpha \in \mathcal{A}$ , and  $P_{\mathcal{S}}(\alpha)$  is the projection of  $\alpha$  onto  $\mathcal{S}$ .  $PST$  has as elements the proper supertranslations. Isomorphism (9.1) identifies the infinite-dimensional group

$$PST(2, 1) = PST \otimes_T G, \quad (9.2)$$

as a subgroup of  $B(2, 1)$ , with semidirect product specified by

$$T(g)(\alpha + \mathcal{T}) = T(g)\alpha + \mathcal{T}, \quad (9.3)$$

where  $\alpha \in \mathcal{A}$ .

An immediate consequence of the definition (6), (9.3) is the Corollary which follows. Normal subgroup and group isomorphism are denoted respectively by  $\trianglelefteq$  and  $\simeq$ .

**Corollary 1** *Regarding the group  $PST(2, 1)$  we have the following*

1.  $PST(2, 1) \not\trianglelefteq B(2, 1)$
2.  $PST \trianglelefteq PST(2, 1)$
3.  $PST(2, 1)/PST \simeq G$
4.  $PST(2, 1) \simeq B(2, 1)/\mathcal{T}$

The IRS of  $PST(2, 1)$  describe quantum systems invariant under all rigid translations in  $R^3$ . Therefore, the IRS of  $B$  are not associated with elementary free particles. It seems plausible to suggest that the carrier spaces of the IRS of  $PST(2, 1)$  have as elements quantum states of three-dimensional Chern–Simons gravity. In view of the relation  $PST(2, 1)/PST \simeq G$ , it is tempting to propose that  $G$ , which double covers  $SO(2, 1)$ , contains within its structure a preferred way of representing a spin-like quantity which can be assigned to these quantum states.

## 10 The spaces $\mathcal{A}$ , $\mathcal{A}'$ , $\hat{\mathcal{A}}$

A character  $\chi$  of  $\mathcal{A}$  is a continuous homomorphism

$$\chi : \mathcal{A} \rightarrow U(1) \quad (10.1)$$

of  $\mathcal{A}$  into the multiplicative group of complex numbers of unit modulus. A composition law

$$(\chi_1\chi_2)(\alpha) = \chi_1(\alpha)\chi_2(\alpha) \quad (10.2)$$

on the characters turns them into an abelian group  $\hat{\mathcal{A}}$ . Besides, a multiplication by scalars

$$(\kappa\chi)(\alpha) = \kappa\chi(\alpha), \quad \kappa \in R, \quad (10.3)$$

turns the characters into a vector space over the reals. By writing the character  $\chi$  as

$$\chi(\alpha) = e^{if(\alpha)}, \quad (10.4)$$

one easily finds that the real-valued function  $f$  1. Is continuous 2. Is additive, i.e.,  $f(\alpha_1 + \alpha_2) = f(\alpha_1) + f(\alpha_2)$ , and, 3. It satisfies  $f(\kappa\alpha) = \kappa f(\alpha)$ ,  $\kappa \in R$ . Therefore  $f$  is a continuous linear functional on  $\mathcal{A}$ , that is,  $f$  belongs to the topological dual  $\mathcal{A}'$  of  $\mathcal{A}$ . Thus the map

$$j : \mathcal{A}' \rightarrow \hat{\mathcal{A}}, \quad (j(f))(\alpha) = e^{if(\alpha)}, \quad (10.5)$$

$\alpha \in \mathcal{A}$ , provides an isomorphism between  $\hat{\mathcal{A}}$  and  $\mathcal{A}'$ , as real vector spaces, and in particular as abelian groups.

From the Reisz–Fréchet theorem for Hilbert spaces, one concludes that *every* element  $f \in \mathcal{A}'$  can be written uniquely in the form

$$f(\alpha) = \langle \beta, \alpha \rangle, \quad (10.6)$$

where  $\beta \in \mathcal{A}$  and  $\langle \cdot, \cdot \rangle$  denotes the inner product of the Hilbert space  $\mathcal{A}$ . Thus the map

$$l : \mathcal{A} \rightarrow \mathcal{A}', \quad (l(\beta))(\alpha) = \langle \beta, \alpha \rangle, \quad (10.7)$$

$\beta, \alpha \in \mathcal{A}$ , provides an isomorphism between  $\mathcal{A}$  and  $\mathcal{A}'$ , as real vector spaces, and in particular as abelian groups. Therefore,  $\mathcal{A}' \simeq \mathcal{H}(P_1(R))$ . Hereafter we will use the same symbol  $\beta$  to denote both an element of  $\mathcal{A}$  and its associated functional in  $\mathcal{A}'$ .

We conclude that

**Proposition 12** *There are the isomorphisms*

$$\mathcal{A} \xrightarrow{l} \mathcal{A}' \xrightarrow{j} \hat{\mathcal{A}}, \quad (10.8)$$

given by

$$(l(\beta))(\alpha) = \langle \beta, \alpha \rangle,$$

and,

$$(j(f))(\alpha) = e^{if(\alpha)}.$$

## 11 The dual action $T'$ and the supermomentum space $\mathcal{A}'$

Requiring pointwise preservation of the isomorphism  $\mathcal{A}' \xrightarrow{j} \hat{\mathcal{A}}$  by the  $G$  action determines uniquely a dual action  $T'$  of  $G$  on  $\mathcal{A}'$ . We have the following

**Corollary 2** *The dual action  $T'$  of  $G$  on  $\mathcal{A}'$  is given by*

$$\langle T'(g)\beta, \alpha \rangle = \langle \beta, T(g^{-1})\alpha \rangle. \quad (11.1)$$

**Proof** *The action  $T$  of  $G$  on  $\mathcal{A}$  induces, in the usual way, an action  $\hat{T}$  of  $G$  on  $\hat{\mathcal{A}}$ , defined by*

$$(\hat{T}\chi)(\alpha) = \chi(T(g^{-1})\alpha). \quad (11.2)$$

*From (10.5) and (10.7) we infer that every character is of the form*

$$\chi(\alpha) = e^{i\langle \beta, \alpha \rangle}, \quad (11.3)$$

*with  $\beta$  uniquely determined by the sequence of isomorphisms  $l$  and  $j$  given respectively by (10.7) and (10.5). Equation (11.2) implies that the isomorphism  $\mathcal{A}' \xrightarrow{j} \hat{\mathcal{A}}$  is preserved pointwise by the  $G$  action, i.e.,*

$$(\hat{T}(g)\chi)(\alpha) = e^{i\langle T'(g)\beta, \alpha \rangle}, \quad (11.4)$$

*if and only if the action  $T'$  of  $G$  on  $\mathcal{A}'$  is given by*

$$\langle T'(g)\beta, \alpha \rangle = \langle \beta, T(g^{-1})\alpha \rangle.$$

*This completes the proof.*

We readily find the dual action  $T'$  of  $G$  on  $\mathcal{A}'$  as follows: By using (4.43) we obtain

$$\begin{aligned} \langle T'(g)\beta, \alpha \rangle &= \langle \beta, T(g^{-1})\alpha \rangle = \int \beta(\rho)(T(g^{-1})\alpha)(\rho) d\lambda(\rho) \\ &= \int \beta(\rho)\kappa_{g^{-1}}(\rho)\alpha(\rho g^{-1}) d\lambda(\rho). \end{aligned} \quad (11.5)$$

By using (5.7) we find

$$\langle T'(g)\beta, \alpha \rangle = \int \kappa_g^{-2}(\rho)\beta(\rho g)\alpha(\rho) d\lambda(\rho). \quad (11.6)$$

This holds for all  $\alpha \in \mathcal{A}$ , so we conclude

$$(T'(g)\beta)(\rho) = \kappa_g^{-2}(\rho)\beta(\rho g). \quad (11.7)$$

It is this dual action  $T'$  of  $G$  on  $\mathcal{A}'$  which determines the structure of the IRS of  $B(2, 1)$ .

$\mathcal{A}'$  is isomorphic to  $\mathcal{H}(P_1(R))$ . This implies that the orthogonal direct sum decomposition  $\mathcal{A} = \mathcal{T} \oplus \mathcal{S}$  dualizes to

$$\mathcal{A}' = \mathcal{T}' \oplus \mathcal{S}', \quad (11.8)$$

where  $\mathcal{T}'$  and  $\mathcal{S}'$  are the dual spaces of  $\mathcal{T}$  and  $\mathcal{S}$  respectively. Obviously we have

$$\mathcal{T}' = \mathcal{S}^\circ, \quad \mathcal{S}' = \mathcal{T}^\circ, \quad (11.9)$$

where  $\mathcal{S}^\circ$  is the annihilator of  $\mathcal{S}$ , consisting of all linear functionals which vanish on  $\mathcal{S}$ .  $\mathcal{T}^\circ$  is the annihilator of  $\mathcal{T}$ . We have the isomorphisms

$$\mathcal{A}'/\mathcal{T}' \approx \mathcal{T}^\circ \quad \text{and} \quad \mathcal{A}'/\mathcal{T}^\circ \approx \mathcal{T}'. \quad (11.10)$$

It is of outmost significance that on passing to the dual action  $T'$  of  $G$  on  $\mathcal{A}'$ , the subspace  $\mathcal{S}'$  becomes invariant whereas the subspace  $\mathcal{T}'$  becomes *noninvariant* (as it is easily inferred from the dual action formula (11.7) had the space  $\mathcal{S}$  been invariant the space  $\mathcal{T}'$  would also have been invariant). This explains in particular that there is no reason to expect that the  $B(2,1)$  little groups should include or be the same as the  $P(2,1)$  little groups: Since  $\mathcal{T}'$  is not  $T'$ -invariant the usual action of  $G$  on Poincare momentum space is not “contained” in the action of  $G$  in  $\mathcal{A}'$ . In fact, as we will prove in Section 14, all the little groups of  $B(2,1)$  are compact, whereas the little groups of  $P(2,1)$  are not all compact.

By using the isomorphism  $\mathcal{A}' \simeq \mathcal{A} = \mathcal{H}(P_1(R))$ , elements  $\beta$  of  $\mathcal{A}'$  can be expanded, as the elements of  $\mathcal{A}$  do, in the complete orthonormal basis (8.1):

$$\beta(\theta) = p_0 + p_1 \cos \theta + p_2 \sin \theta + \sum_{n=2}^{\infty} (p_n e_n + p_n \varepsilon_n), \quad (11.11)$$

where the decomposition  $\mathcal{A}' = \mathcal{T}' \oplus \mathcal{S}'$  has already been indicated: The set  $\{1, \cos \theta, \sin \theta\}$  is a spanning set of  $\mathcal{T}'$  and the set  $\{e_n, \varepsilon_n, n = 2, 3, \dots\}$  is a spanning set of  $\mathcal{S}'$ . The isomorphism  $\mathcal{A}'/\mathcal{T}^\circ \approx \mathcal{T}'$  suggests that the triplet  $(p_0, p_1, p_2)$  may be thought of as the components of a “momentum” vector with respect to the Poincare group  $P(2,1)$ . As a complement to this observation we note that given that  $p = (p_0, p_1, p_2)$  is a “momentum” vector then the character  $\chi(\alpha) = e^{i\langle p, \alpha \rangle}$  associated to  $p$  becomes a plane wave as the wave function of a free particle with a well defined momentum  $p$  in the momentum representation we expect to be.

## 12 Representation theory and its extension

Our construction of the IRS of  $B(2,1)$  is based on G. W. Mackey’s pioneering work on group representations (see, for example, [18], [19], [20], [40], [41]); in particular, is based on an extension to the relevant infinite-dimensional case [10], [21], [27], [42] of his semidirect product theory. It is interesting that the main sources of Mackey’s work were his extension of the Stone–von Neumann theorem [43], which is the cornerstone of quantum mechanics, Von Neuman’s theory of direct integral decompositions of operator algebras which he developed in the 1930s, and Wigner’s classic paper [6] in which he broke new grounds in mathematics by giving the first explicit treatment of infinite dimensional representations of a Lie group.

## 12.1 Wigner–Mackey’s theory

Let  $\mathcal{A}$  and  $\mathcal{G}$  be topological groups, and let  $T$  be a given homomorphism from  $\mathcal{G}$  into the group of automorphisms  $\text{Aut}(\mathcal{A})$  of  $\mathcal{A}$ . Suppose  $\mathcal{A}$  is abelian and  $\mathcal{B} = \mathcal{A} \rtimes_T \mathcal{G}$  is the semi-direct product of  $\mathcal{A}$  and  $\mathcal{G}$ , specified by the continuous action  $T : \mathcal{G} \rightarrow \text{Aut}(\mathcal{A})$ . In the product topology of  $\mathcal{A} \times \mathcal{G}$ ,  $\mathcal{B}$  then becomes a topological group. It is assumed that it becomes a separable locally compact topological group.

The action  $T$  of  $\mathcal{G}$  on  $\mathcal{A}$  induces a dual action  $T'$  of  $\mathcal{G}$  on  $\mathcal{A}'$  given by (11.7)

$$\langle T'(g)\beta, \alpha \rangle = \langle \beta, T(g^{-1})\alpha \rangle .$$

For a given functional  $\beta \in \mathcal{A}'$ , the largest subgroup  $L_\beta$  of  $\mathcal{G}$  which leaves  $\beta$  fixed is called the “little group” of  $\beta$ , and the set of functionals which can be reached from  $\beta$  by the  $\mathcal{G}$  action is called the orbit of  $\beta$ , denoted  $\mathcal{G}\beta$ .

Let  $U$  be a unitary representation of the “little group”  $L_\beta$  on a Hilbert space  $D$ . Let  $D_\mu$  be the space of functions  $\psi : \mathcal{G} \rightarrow D$  which satisfy the conditions

$$(a) \quad \psi(gl) = U(l^{-1})\psi(g), \quad g \in \mathcal{G}, \quad l \in L_\beta,$$

$$(b) \quad \int_{\mathcal{G}\beta} \langle \psi(q), \psi(q) \rangle d\mu(q) < \infty,$$

where the scalar product under the integral sign is that of  $D$ . Note, that the constraint (a) implies that  $\langle \psi(gl), \psi(gl) \rangle = \langle \psi(g), \psi(g) \rangle$ , and therefore the inner product  $\langle \psi(g), \psi(g) \rangle$ ,  $g \in \mathcal{G}$ , is constant along every element  $q$  of the coset space  $\mathcal{G}/L_\beta \approx \mathcal{G}\beta$ . This allows to assign a meaning to  $\langle \psi(q), \psi(q) \rangle$ , where  $q = gL_\beta$ , by defining  $\langle \psi(q), \psi(q) \rangle := \langle \psi(g), \psi(g) \rangle$ . Thus the integrand in (b) becomes meaningful due to the condition (a).

A pre-Hilbert space structure can now be given to  $D_\mu$  by defining the scalar product

$$\langle \psi_1, \psi_2 \rangle = \int_{\mathcal{G}\beta} \langle \psi_1(q), \psi_2(q) \rangle d\mu(q), \quad (12.1)$$

where  $\psi_1, \psi_2 \in D_\mu$ . It is convenient to complete the space  $D_\mu$  with respect to the norm defined by the scalar product (12.1). In the resulting Hilbert space, functions are identified whenever they differ, at most, on a set of  $\mu$ -measure zero. Thus our Hilbert space is

$$D_\mu = L^2(\mathcal{G}\beta, \mu, D). \quad (12.2)$$

We note the following. Let  $\mathcal{T}$  be a representation of  $\mathcal{B}$  and let  $Q$  and  $V$  be its restriction to  $\mathcal{A}$  and  $\mathcal{G}$  respectively, i.e.,

$$Q(\alpha) \equiv \mathcal{T}(a), \quad V(g) \equiv \mathcal{T}(g), \quad (12.3)$$

where,  $a = (\alpha, I)$ ,  $g = (i, g)$ , and  $I, i$  are the identity elements of  $\mathcal{G}$  and  $\mathcal{A}$  respectively. Since  $(\alpha, g) = (\alpha, I)(i, g)$ , for all  $\alpha \in \mathcal{A}$  and  $g \in \mathcal{G}$ , for any  $h = (\alpha, g) \in \mathcal{B}$  we have

$$\mathcal{T}(h) \equiv \mathcal{T}(\alpha, g) = Q(\alpha)V(g). \quad (12.4)$$

Thus a representation  $\mathcal{T}$  of  $\mathcal{B}$  is completely determined by its restriction  $Q$  and  $V$  to the subgroups  $\mathcal{A}$  and  $\mathcal{G}$  respectively. Conversely representations  $Q$  and  $V$  of  $\mathcal{A}$  and  $\mathcal{G}$  correspondingly, define a unique representation  $\mathcal{T}$  of  $\mathcal{B}$  by (12.4).  $\mathcal{T}$  is unitary when  $Q$  and  $V$  are unitary.

With this remark in mind we define now actions of  $\mathcal{A}$  and  $\mathcal{G}$  on  $D_\mu$  by

$$(Q(\alpha)\psi)(q) = e^{i\langle g\beta, \alpha \rangle} \psi(q), \text{ and} \quad (12.5)$$

$$(V(g)\psi)(q) = \sqrt{\frac{d\mu_g}{d\mu}}(q) \psi(g^{-1}q), \quad (12.6)$$

respectively, where  $g \in \mathcal{G}$ ,  $q = gL_\beta \in \mathcal{G}/L_\beta \approx \mathcal{G}\beta$ , and  $\alpha \in \mathcal{A}$ . It is a straightforward exercise to verify that these actions are in fact representations of  $\mathcal{A}$  and  $\mathcal{G}$  on  $D_\mu$ . The ‘‘Jacobian’’  $\frac{d\mu_g}{d\mu}$  is known as the *Radon–Nikodym* derivative of  $\mu_g$  with respect to  $\mu$  and ensures that the representation  $V$  is unitary. Actions (12.5) and (12.6) define a unitary representation of  $\mathcal{B}$  on  $D_\mu$  by

$$\begin{aligned} (\mathcal{T}(h)\psi)(q) &= (\mathcal{T}(\alpha, g)\psi)(q) = ((Q(\alpha)V(g))\psi)(q) = (Q(\alpha)(V(g)\psi))(q) \\ &= e^{i\langle g\beta, \alpha \rangle} (V(g)\psi)(q) = \sqrt{\frac{d\mu_g}{d\mu}}(q) e^{i\langle g\beta, \alpha \rangle} \psi(g^{-1}q). \end{aligned} \quad (12.7)$$

The central results of induced representation theory ([6], [18], [19], [20], [40], [41], and references therein) are the following:

1. Given the topological restrictions on  $\mathcal{B} = \mathcal{A} \rtimes_{\mathcal{T}} \mathcal{G}$  (separability and local compactness), any representation  $\mathcal{T}$  of  $\mathcal{B}$ , constructed by the method above, is irreducible if the representation  $U$  of  $L_\beta$  on  $D$  is irreducible. Thus an irreducible representation of  $\mathcal{B}$  is induced for *each*  $\beta \in \mathcal{A}'$  and *each* irreducible representation  $U$  of  $L_\beta$ .
2. If  $\mathcal{B} = \mathcal{A} \rtimes_{\mathcal{T}} \mathcal{G}$  is a regular semi–direct product (i.e., if  $\mathcal{A}'$  contains a Borel subset which meets each  $\mathcal{G}$ –orbit in  $\mathcal{A}'$  at just one point) then *all* of its irreducible representations are obtained by this inducing construction.

It follows from the previous discussion that in order to give the operators  $\mathcal{T}$  of the IRS of  $\mathcal{B}$  explicitly it is necessary to give the following information

1. An irreducible unitary representation  $U$  of  $L_\beta$  on a Hilbert space  $D$  for each ‘‘little group’’  $L_\beta$ .
2. A  $\mathcal{G}$ –quasi–invariant measure  $\mu$  on each orbit  $\mathcal{G}\beta \approx \mathcal{G}/L_\beta$ , where  $L_\beta$  denotes the ‘‘little group’’ of the base point  $\beta \in \mathcal{A}'$  of the orbit  $\mathcal{G}\beta$ .

In the case of  $B(2, 1)$  then, in order to find its IRS  $\mathcal{T}$ , it is enough to provide the information cited in 1 and 2 for each of the orbit types.

## 12.2 Extension of Wigner–Mackey’s theory

The key feature of  $B(2,1)$  is the enlargement of the three–dimensional vector space of rigid translations in  $R^{2,1}$  to the infinite–dimensional vector space of supertranslations  $\mathcal{H}(P_1(R))$ . This turns  $B(2,1)$  into an infinite–dimensional Lie group. In fact  $\mathcal{H}(P_1(R))$  equipped with the standard inner product (5.9) is a separable *infinite–dimensional* Hilbert space. It is precisely this key feature which calls for an extension of Wigner–Mackey’s theory.

Mackey assumes [17] that both  $\mathcal{A}$  and  $\mathcal{G}$  in  $\mathcal{B} = \mathcal{A} \rtimes_T \mathcal{G}$  are locally compact. In the applications to Physics it is assumed that  $\mathcal{A}$  has an additional structure, it is also a vector space, with vector addition being the group multiplication. It is well known that every Hausdorff topological vector space is locally compact if and only if it is finite. This implies in particular that a Hausdorff infinite–dimensional topological vector space is never locally compact. In the case of  $B(2,1)$   $\mathcal{A}$  is an infinite–dimensional vector space. In the whole class of topologies, physically motivated,  $\mathcal{A}$  is Hausdorff and therefore not locally compact. This renders Mackey’s Theorems 14.1 and 14.2 in [17] inapplicable.

In this paper  $B(2,1)$  is equipped with the Hilbert topology.  $\mathcal{H}(P_1(R))$  is endowed with the topology whose open sets are the balls determined by the metric induced by the scalar product (5.9). Moreover, the group  $G$ , considered as a subset of  $R^4$ , is endowed with the induced topology. In the product topology of  $\mathcal{H}(P_1(R)) \times G$ , the Hilbert topology,  $B(2,1)$  becomes a topological group; in particular it becomes a non–locally compact group. The proof is similar to the corresponding proof [44] for the original BMS group  $B$ .

Since  $B(2,1)$  is not locally compact in the Hilbert topology, Mackey’s Theorem 14.2 in [17] which proves irreducibility of the representations obtained by inducing does not no longer applies. However, when  $B(2,1)$  is employed with the Hilbert topology we can prove that the representations of  $B(2,1)$  obtained by inducing are irreducible notwithstanding the fact that  $B(2,1)$  is not locally compact in the employed Hilbert topology. The proof is practically the same with the corresponding proof given in [45] for the original BMS group  $B$  when  $B$  is endowed with the Hilbert topology.

Mackey proved, Theorem 14.1 in [17], that only for regular semi–direct products  $\mathcal{B} = \mathcal{A} \rtimes_T \mathcal{G}$  Wigner–Mackey inducing construction is exhaustive, i.e., that only for regular  $\mathcal{B}$  all IRS of  $\mathcal{B}$  are obtained by inducing. Regular  $\mathcal{B}$  means that  $\mathcal{A}'$  can have no  $\mathcal{G}$ –quasi–invariant measures  $\mu$  such that the action of  $\mathcal{G}$  is strictly ergodic with respect to  $\mu$ . Whenever such measures  $\mu$  do exist, an irreducible representation of  $\mathcal{B}$  may be associated with each that is not obtained by inducing. A sufficient condition which ensures the regularity of  $\mathcal{B}$  [46] (p. 43 and p. 141) is the existence of a Borel set in  $\mathcal{A}'$  that meets each  $\mathcal{G}$ –orbit exactly once.

When the abelian normal subgroup  $\mathcal{A}$  is also a finite–dimensional vector space finding a Borel cross section which intersects each  $\mathcal{G}$ –orbit in  $\mathcal{A}'$  exactly once is fairly easy (e.g. [19]). However, when the abelian normal subgroup  $\mathcal{A}$  is also an infinite–dimensional vector space,  $\mathcal{A}'$  is also an infinite–dimensional vector space, it is not generally feasible [9], [21] to classify completely the  $\mathcal{G}$ –orbits in  $\mathcal{A}'$ , let alone to search for a Borel set in  $\mathcal{A}'$  which meets each  $\mathcal{G}$ –orbit in  $\mathcal{A}'$  exactly once.

Consequently, the existence of a Borel cross section which meets each  $\mathcal{G}$ –orbit in  $\mathcal{A}'$

exactly once cannot be used as a sufficient condition which can be practically implemented when  $\mathcal{A}'$  is infinite-dimensional and new sufficient conditions which insure the exhaustivity of the IRS of  $\mathcal{B}$  constructed by Wigner-Mackey's "little group" method need to be introduced.

It is expected that it is the properties of the dual action of  $\mathcal{G}$  on  $\mathcal{A}'$  which will determine if the list of IRS obtained by inducing is exhaustive. As a result we expect that the properties of the dual action of  $\mathcal{G}$  on  $\mathcal{A}'$  will imply the validity of the sought new sufficient conditions. In conclusion, when  $\mathcal{A}$  is an infinite-dimensional vector space, proving the exhaustivity of the IRS obtained by inducing is reduced to the solution of the following two problems:

1. Find sufficient conditions which ensure that  $\mathcal{A}'$  can have no  $\mathcal{G}$ -quasi-invariant measures  $\mu$  such that the action of  $\mathcal{G}$  is strictly ergodic with respect to  $\mu$ .
2. Find those properties of the dual action of  $\mathcal{G}$  on  $\mathcal{A}'$  which imply the validity of the sufficient conditions stated in Problem 1.

Problems 1 and 2 were solved in two remarkable papers [22], [23] by Piard, in the case where  $\mathcal{B}$  is endowed with the Hilbert topology.

### 13 Topologies in $\mathcal{A}$

The class of functions in  $\mathcal{A}$  is not uniquely determined. This allows a wide range of "reasonable" topologies for  $\mathcal{A}$ .

#### 13.1 $\mathcal{A}$ is not unique

The key observation is that Sachs's original derivation [36] was superseded by that of Penrose [3-5, 47]. As it was shown in subsection 6.3, since truly arbitrary supertranslation functions describe symmetry transformations in Penrose's sense, *supertranslations need not have some minimum degree of smoothness*.

Therefore, there is to a certain extent an arbitrariness in the choice of the class of functions for  $\mathcal{A}$ . Sachs's original derivation [36] of the group  $B$  required that the supertranslations be twice differentiable. McCarthy widened [7, 9]  $C^2(S^2, R)$  to  $L^2(S^2, R)$ , the square-integrable real-valued functions defined on the two sphere, in order to simplify the treatment.

Other choices for  $\mathcal{A}$  are the function spaces  $C^k(S^2, R)$ ,  $k = 0, 1, 2, \dots$ , [48], [49], the  $k$ -times continuously differentiable functions on  $S^2$ , which are physically plausible in a gravitational radiation context because they are adapted to initial-value problems for hyperbolic differential equations, the function space  $C^\infty(S^2, R)$  of smooth functions on  $S^2$  [10], [48], [49], the function space  $C^\omega(S^2, R)$  of analytic functions on  $S^2$  [50].

As it was shown in section 6  $B(2, 1)$  can also be derived in Penrose's sense, i.e., as that group of *exact* conformal motions of the future (or past) null boundary  $\mathcal{I}^+$  (or  $\mathcal{I}^-$ ) of 3-dim Minkowski space-time which is the automorphism group of the "strong conformal geometry" at  $\mathcal{I}^+$  (or  $\mathcal{I}^-$ ). Consequently similar remarks for the available freedom in the choice of the degree of smoothness of the supertranslations apply also in the case of  $B(2, 1)$ .



Thus besides  $\mathcal{H}_1(P_1(R))$ , other physically plausible choices for  $\mathcal{A}$  are the function spaces  $C^k(S^1, R)$ ,  $k = 0, 1, 2, \dots$ ,  $C^\infty(S^1, R)$ , and  $C^\omega(S^1, R)$ .

### 13.2 Topologies in $\mathcal{A}$

It is precisely the arbitrariness in the class of functions allowed in  $\mathcal{A}$  which permits a wide range of choices of “reasonable” topologies for  $\mathcal{A}$ . The only way to interpret the topologies is to look at the structure of IRS themselves.

The class of functions in  $\mathcal{A}$  does not determine uniquely the topology in  $\mathcal{A}$ . For the reasons set out in [7], [9], [10], [48], [49], in the case of the original BMS group  $B$  the Hilbert topology gives rise to IRS which are identified with bound states, corresponding to elementary particles. For the same reasons the Hilbert topology given in  $B(2, 1)$  in this paper leads to IRS which describe elementary particles.

If we choose for  $\mathcal{A}$  the function spaces  $C^k(S^1, R)$ ,  $k = 0, 1, 2, \dots$ , as in the case of  $B$  [49], [51], a physically reasonable class of topologies are the Sobolev topologies. If we take  $\mathcal{A}$  to be  $C^\infty(S^1, R)$ ,  $\mathcal{A}'$  is enlarged to the space of generalised functions  $\mathcal{F}(S^1)$  on  $S^1$ , and a natural choice, as in the case of  $B$  [10], [48], [49], is the nuclear topology. The expectation is [49] that in the nuclear topology new IRS are going to arise which describe scattering states. If we choose for  $\mathcal{A}$  the function space  $C^\omega(S^1, R)$  rather than the function space  $C^\infty(S^1, R)$ , then, with an appropriately fine topology,  $\mathcal{A}'$  becomes enlarged to the space  $\mathcal{Y}(S^1)$  of real hyperfunctions on  $S^1$ .  $\mathcal{Y}(S^1)$  is larger than  $\mathcal{F}(S^1)$ . It seems [50] that the category of hyperfunctions is more appropriate than that of distributions for discussing  $S$ -matrix theory.

## 14 “Little groups” are compact

Before giving the proof that the “little groups” of  $B(2, 1)$  are compact in the Hilbert topology, we make some comments on the dependence of the “little groups” on the topology  $B(2, 1)$  comes equipped with.

### 14.1 “Little groups” depend on the topology

“Little groups” might change when the topology of  $\mathcal{A}$  is modified: The induced representations are associated with the existence of invariant characters, i.e., of elements in  $\mathcal{A}'$  which are left invariant by the action of some “little group”. “Little groups” are then connected, via  $\mathcal{A}'$ , to the topology of  $\mathcal{A}$ . Since, e.g., a refinement of the topology may broaden  $\mathcal{A}'$ , new invariant elements with associated little groups may come into existence.

In fact this is precisely what happens in the case of the original BMS group  $B$ . In the Hilbert topology all little groups of  $B$  are compact [7], [9]. The effect of passing from the Hilbert (coarser) topology to the nuclear (finer) topology is to enlarge  $\mathcal{A}'$  from  $L^2(S^2, R)$  to the space of generalised functions  $\mathcal{F}(S^2)$  on  $S^2$ , i.e., the distributions  $\mathcal{D}(S^2) = \mathcal{F}(S^2) - L^2(S^2, R)$  as well as “regular”  $L^2(S^2, R)$  functions. Non-compact little groups arise in the nuclear topology and are associated with distributional supermomenta  $\chi \in \mathcal{D}(S^2)$ . We expect that the same will happen in the case of  $B(2, 1)$ . In this section we prove that all “little groups” of  $B(2, 1)$  are compact in the Hilbert topology. We expect that if we pass

from the Hilbert to the nuclear topology non–compact little groups are going to arise in the nuclear topology which will leave invariant distributional supermomenta  $\varphi \in \mathcal{D}(S^1)$ . We also expect that if we choose for  $\mathcal{A}$  the function space  $C^\omega(S^1, R)$  non–compact little groups are going to arise which leave invariant hyperfunctional supermomenta.

## 14.2 Proof

Recall that the little group  $L_\beta$  of any  $\beta \in \mathcal{A}'$  is the set

$$L_\beta = \{g \in G \mid T'(g)\beta = \beta\}. \quad (14.1)$$

By continuity of the representation  $T'$ ,  $L_\beta$  is always a closed subgroup of  $G$ . Evidently, if  $\beta = 0$ ,  $L_\beta = G$ . Representations of  $B(2, 1)$  that are induced from  $\beta = 0$ , and  $L_\beta = G$ , are unfaithful on the *whole* supertranslation subgroup, and in particular describe quantum systems invariant under all rigid translations in  $R^{2,1}$ . Clearly they have no physical significance in the context of  $B(2, 1)$  which is concerned with isolated sources. So, hereafter, we always assume that  $\beta \neq 0$ .

We now prove that all little groups of  $B(2, 1)$  are in fact compact in the Hilbert topology. The standard topology on  $G$  is given by the metric derived from the norm

$$|g| = (|a|^2 + |b|^2 + |c|^2 + |d|^2)^{1/2}, \quad g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in G. \quad (14.2)$$

The following result will be useful.

**Theorem 5** *If  $L \subset G$  is a non–compact subgroup of  $G$ , and has a fixed point in  $\mathcal{A}'$ , there is an infinite sequence  $g_n$ ,  $n = 1, 2, 3, \dots$ ,  $g_n \in L$  such that, for large enough  $n$ , the function  $\kappa_{g_n}(\rho)$  is greater than any given  $M > 0$  over a region  $R_1$  of  $S^1$ , and  $0 < \kappa_{g_n}(\rho) \leq M$  over a region  $R_2$  of  $S^1$ , where  $S^1 = R_1 \cup R_2$  (disjoint union), and the measure  $\lambda(R_2)$  is not greater than any given  $\varepsilon > 0$ .*

**Proof** *The proof is similar to the corresponding proof for  $B(2, 2)$  [42] and as such it is omitted.*

The  $\kappa$  factor corresponds physically to a “redshift factor”. Theorem 5 amounts to the anticipated fact that a sequence of Lorentz transformations which contains a sequence of pure boosts with velocity parameter tending arbitrary close to the speed of light subjects a region of the one–dimensional “selestial sphere” with arbitrary small complement to arbitrary large “blue–shifts” (the complement is “redshifted”).

We now prove the following:

**Theorem 6** *Every little group of  $B(2, 1)$  is compact.*

**Proof** *The proof proceeds by contradiction. Let  $\beta \in \mathcal{A}' - \mathbf{0}$  have little group  $L_\beta$ . It will be shown that if  $L_\beta$  is non–compact and  $\beta$  is fixed under  $L_\beta$ , then  $\beta = 0$  or  $\beta$  cannot be*

square integrable. A somewhat stronger result will, in fact, be proved, namely that  $\beta = 0$  or  $|\beta|$  cannot be integrable. This implies the desired result since, from  $(1 - \beta)^2 \geq 0$ , one gets  $|\beta| \leq \frac{1}{2}(1 + \beta^2)$ .

Assume, contrary to the theorem, that  $L_\beta$  is non-compact for some  $\beta \neq 0$ . Then  $L_\beta \subset G$  must be unbounded with respect to the standard metric (14.2) for  $G$ . Indeed,  $L_\beta$  is closed, so if it were bounded, it would be compact, contrary to our starting assumption. By definition  $(T'(g)\beta)(\rho) = \beta(\rho)$  for all  $g \in L_\beta$ . Hence, for all  $g \in L_\beta$ ,

$$\int_{S^1} |(T'(g)\beta)(\rho)| d\lambda(\rho) = \int_{S^1} |\beta(\rho)| d\lambda(\rho) \quad (14.3)$$

for all  $g \in L_\beta$ . Using (11.7) to substitute for  $T'(g)\beta$ , changing the variable  $\rho \mapsto \rho g^{-1}$ , and using (5.7), gives

$$\int_{S^1} \kappa_{g^{-1}}(\rho) |\beta(\rho)| d\lambda(\rho) = \int_{S^1} |\beta(\rho)| d\lambda(\rho) \quad (14.4)$$

for all  $g \in L_\beta$ . Since  $L_\beta$  is a group, the same equation applies with  $g^{-1}$  replaced by  $g$ . Then the previous equation becomes

$$\int_{S^1} (\kappa_g(\rho) - 1) |\beta(\rho)| d\lambda(\rho) = 0 \quad (14.5)$$

for all  $g \in L_\beta$ . Then using Theorem 5 for  $\kappa_g(\rho)$ , the rest of the proof is essentially the same as the corresponding proof for  $B(2, 2)$  in [42]. This completes the proof.

Theorem 6 intuitively can be understood intuitively as follows: The dual action (11.7)  $T'$  of  $G$  on  $\mathcal{H}(P_1(R))$  is similar [21] to the transformation law for light intensity distributions on distant  $1 - \dim$  “celestial” sphere. A non-compact little group  $L_\beta$  contains a sequence of boosts with velocity which becomes arbitrary close to the speed of light. For an intensity distribution  $\beta$  to be fixed under  $L_\beta$  it must be such that when subjected to arbitrary large “blueshifts” over a region of the  $1 - \dim$  “celestial” sphere with arbitrary small complement (the complement is “redshifted”), it remains the same. It is not surprising that there are no such non-zero  $\beta$ 's, so that there cannot be any non-compact little groups. However, by the same reasoning, there may exist *distributional intensity distributions*  $\beta$  invariant under non-compact little groups. This is precisely what we expect to happen if we take  $\mathcal{A}$  to be  $C^\infty(S^1, R)$  and equip it with the nuclear topology.

## 15 The “Little groups” of $B(2, 1)$

In section 14 it was shown that the “little groups” of  $B(2, 1)$  are compact subgroups of  $SL(2, R)$ . The maximal compact subgroup of  $G = SL(2, R)$  is just the subgroup  $SO(2)$ . That is to say, if  $H$  is a compact subgroup of  $G$ , then some conjugate  $gHg^{-1}$ ,  $g \in G$ , of  $H$  is a compact subgroup of  $SO(2)$ . In the representation theory of  $B(2, 1)$ , the “little groups” are only significant up to conjugacy. So, after a possible conjugation, we may take every “little group” to be a compact (or equivalently closed) subgroup of  $SO(2)$ . Therefore, we need to find the compact subgroups of  $SO(2)$ .

### 15.1 The compact subgroups of $SO(2)$

We prove now the following

**Proposition 13** *The compact subgroups of  $SO(2)$  are classified as follows:*

1.  $SO(2)$  is the only one-dimensional compact subgroup.
2.  $C_n$ ,  $n \geq 1$ , the finite cyclic groups of order  $n$ , are the only zero-dimensional compact subgroups.

**Proof**  $SO(2)$  is the set of matrices of the form

$$g(\omega) = g = \begin{bmatrix} \cos \omega & \sin \omega \\ -\sin \omega & \cos \omega \end{bmatrix} \in G. \quad (15.1)$$

where  $\omega$  is any real number, determined mod  $2\pi$ , and may be taken in the range  $0 \leq \omega \leq 2\pi$ .

The infinitesimal generator of  $SO(2)$  is

$$\left. \frac{dg(\omega)}{d\omega} \right|_{\omega=0} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = J. \quad (15.2)$$

Hence the Lie algebra  $\mathcal{L}K$  of  $K = SO(2)$  is the vector space of all matrices of the form

$$\alpha J = \begin{bmatrix} 0 & \alpha \\ -\alpha & 0 \end{bmatrix}, \quad (15.3)$$

where  $\alpha$  is any real number. The exponential map  $exp : \mathcal{L}K \rightarrow K$  here is the usual exponential for matrices. This is given by

$$exp(\alpha J) = g(\alpha). \quad (15.4)$$

This map is obviously onto but it is not one-to-one.

We now wish to describe the compact (equivalently closed) subgroups of  $K$ . Every closed subgroup of a Lie group is a Lie subgroup (see e.g. [52]). Let  $H$  be any closed subgroup of  $K$ ,  $H \subset K$ , with dimension 1. The Lie algebra  $\mathcal{L}H$  of  $H$  has dimension 1, so must coincide with  $\mathcal{L}K$ ;  $\mathcal{L}H = \mathcal{L}K$ .  $K$  is a connected topological group, so *any* neighborhood of the identity of  $K$  generates the whole of  $K$  (see e.g. [53]). But  $exp(\mathcal{L}H) = exp(\mathcal{L}K)$  contains a neighborhood of the identity of  $H$ , and also of  $K$ . The intersection of these is a neighborhood of the identity of both groups. This intersection belongs to  $H$ , and generates the whole of  $K$ . So  $K \subset H$ . But  $H \subset K$ . Therefore,  $H = K$ .

The only remaining case to consider is that in which  $H$  is zero-dimensional. Then  $H$  is a discrete closed subgroup of the compact group  $K$  and so, necessarily, a finite subgroup  $F \subset K$ . But the only finite discrete subgroups of  $K$  are the finite cyclic groups  $C_n$  of order  $n$ ,  $n \geq 1$ . Therefore  $F = C_n$ ,  $n \geq 1$ . This completes the proof.

## 15.2 The “Little groups” of $B(2, 1)$

In [54] we prove that not all the compact subgroups of  $SO(2)$ , given in Proposition 13, qualify as little groups. We prove in particular the following

**Theorem 7** *The “little groups” of  $B(2, 1)$  are*

1.  $SO(2)$  itself.
2. The cyclic groups  $C_n$  of even order.

## 16 Exhaustivity of Induced Representations

Proving exhaustivity opens a whole new chapter in the theory of dynamical systems. The dynamical system at hand is the triple  $(\mathcal{A}', \mathcal{M}, \mathcal{G})$  where  $\mathcal{M}$  is a class of  $\mathcal{G}$ -quasi-invariant measures on  $\mathcal{A}'$ . It is the properties of the  $\mathcal{G}$  action on  $\mathcal{A}'$  which determine if the list of IRS of  $\mathcal{B} = \mathcal{A} \rtimes_T \mathcal{G}$  obtained by inducing is exhaustive.

Piard with a view to proving exhaustivity of McCarthy’s list of representations [7], [9], studied class of semi-direct products of the form  $\mathcal{B} = \mathcal{A} \rtimes_T \mathcal{G}$ , where  $\mathcal{A}$  is either  $C^\infty(S^1, R)$  endowed with the nuclear topology, or  $\mathcal{A}$  is the Hilbert space of a Gelfand triple  $\mathcal{A}_o \subset \mathcal{A} \subset \mathcal{A}'_o$  ( $\mathcal{A}_o$  is nuclear dense in  $\mathcal{A}$ ).  $\mathcal{A}$  in the latter case is a separable countable infinite-dimensional Hilbert space endowed with the strong topology which is also known in the literature as metric or norm topology.  $\mathcal{G}$  in both cases is assumed to be locally compact. Both the original BMS group  $B$  in the Hilbert topology and  $B(2, 1)$  in the Hilbert topology fall into this latter class of groups.

In a nutshell Piard associates a cylindrical measure on  $\mathcal{A}'$  (respectively  $\mathcal{A}'_o$  in which case the measure is concentrated on  $\mathcal{A}'_o \cap \mathcal{A}'$ ). When the representation of  $\mathcal{B}$  is irreducible this measure is ergodic for the dual action of  $\mathcal{G}$  on  $\mathcal{A}'$  and under the assumption that the measure is concentrated on an orbit Piard proves that the representation of  $\mathcal{B}$  is induced. When  $\mathcal{A}$  is a Hilbert space Piard gives sufficient conditions that all these ergodic measures be concentrated on an orbit and hence that all IRS of  $\mathcal{B}$  be induced.

In [23] Piard by using the properties of the  $\mathcal{G}$  action in  $\mathcal{A}'$  proved that the sufficient conditions introduced in [22] are verified by  $B$ . He thus solved, in the case of  $B$ , problems 1 and 2 stated in subsection 12.2. In the case of  $B$ ,  $\mathcal{G} = SL(2, C)$  and  $\mathcal{A}' \simeq L^2(S^2, R)$ . Therefore, Piard concluded, McCarthy’s list of IRS in the Hilbert topology [7], [9] is exhaustive. In this section, by using the properties of the dual action of  $SL(2, R)$  on  $\mathcal{A}' \simeq \mathcal{H}(P_1(R))$ , we prove that the sufficient conditions introduced in [22] are also verified by  $B(2, 1)$ .

### 16.1 Properties of the dual action

Three are the properties of the dual action of  $SL(2, R)$  on  $\mathcal{A}'$  which suffice to insure the validity of the sufficient conditions introduced in [22]. These properties are stated in the three Propositions which follow.

**Proposition 14** *Let  $g_n, n = 1, 2, 3, \dots$ , be a sequence in  $G$  such that  $|g_n| \rightarrow \infty$ . Then there exists a subsequence  $g_{n_p}$  such that  $\kappa_{g_{n_p}}(\rho) \rightarrow \infty$  almost uniformly as  $p \rightarrow \infty$ .*

**Proof** The proof is the same as the corresponding proof for the original BMS group  $B$  given in [23].

Proposition 14 is an immediate consequence of Theorem 5 and asserts that every sequence of Lorentz transformations which subjects a region of the one-dimensional “selestial sphere” with arbitrary small complement to arbitrary large “blueshifts” contains a subsequence of Lorentz transformations which subjects a region of the one-dimensional “selestial sphere” with complement of measure zero to infinite “blueshift”.

**Proposition 15** Let  $g_n, n = 1, 2, 3, \dots$ , be a sequence in  $G$  such that  $|g_n| \rightarrow \infty$ . Then there exists a subsequence  $g_{n_p}$  such that  $\|T'(g_{n_p})\beta\| \rightarrow \infty$  almost uniformly as  $p \rightarrow \infty$  for all  $\beta \in \mathcal{A}'$ .

**Proof** From (11.7) we obtain

$$\|(T'(g_n)\beta)\|^2 = \int_{S^1} \kappa_{g_n}^{-4}(\rho)\beta^2(\rho g_n)d\lambda(\rho). \quad (16.1)$$

Using (5.7) and changing the variable  $\rho \mapsto \xi = \rho g_n$  yields

$$\|(T'(g_n)\beta)\|^2 = \int_{S^1} \kappa_{g_n}^{-3}(\xi g_n^{-1})\beta^2(\xi)d\lambda(\xi). \quad (16.2)$$

From (4.43) we have

$$\begin{aligned} [(T(g_n^{-1})T(g_n)\alpha)](\xi) &= [T(g_n^{-1})(T(g_n)\alpha)](\xi) = \kappa_{g_n^{-1}}(\xi)[T(g_n)\alpha](\xi g_n^{-1}) \\ &= \kappa_{g_n^{-1}}(\xi)\kappa_{g_n}(\xi g_n^{-1})\alpha(\xi) \end{aligned} \quad (16.3)$$

for all  $\alpha \in \mathcal{A}$ . Therefore,  $\kappa_{g_n^{-1}}(\xi)\kappa_{g_n}(\xi g_n^{-1}) = 1$  almost everywhere. Henceforth we obtain

$$\|(T'(g_n)\beta)\|^2 = \int_{S^1} \kappa_{g_n^{-1}}^3(\xi)\beta^2(\xi)d\lambda(\xi). \quad (16.4)$$

The rest of the proof is essentially the same as the corresponding proof for the original BMS group  $B$  given in [23].

Proposition 15 is a consequence of Proposition 14 and states that every sequence of Lorentz transformations which subjects a region of the one-dimensional “selestial sphere” with arbitrary small complement to arbitrary large “blueshifts” contains a subsequence of Lorentz transformations which transforms any square integrable intensity distribution to a distribution of infinite intensity in a region of the one-dimensional “selestial sphere” with complement of measure zero.

A consequence of Proposition 15 which refers to any sequence of square integrable intensity distributions on the one-dimensional “selestial sphere” is given in the Proposition which follows.

**Proposition 16** Let  $\beta_n, n = 1, 2, 3, \dots, \beta_n \in \mathcal{A}'$ , be a sequence weakly convergent to  $\beta_0$  and  $g_n, n = 1, 2, 3, \dots$ , a sequence in  $G$  such that  $|g_n| \rightarrow \infty$ . Then there exists a subsequence  $g_{n_p}$  such that  $\|T'(g_{n_p})\beta_{n_p}\| \rightarrow \infty$  almost uniformly as  $p \rightarrow \infty$ .

**Proof** The proof is practically the same as the corresponding proof for the original BMS group  $B$  given in [23].

## 16.2 Sufficient conditions are satisfied

Piard's sufficient conditions introduced in [22] are:

1. The  $\mathcal{G}$ -orbits in  $\mathcal{A}'$  are open relatively to their strong closure.
2. For any two minimal invariant cones  $C_{\beta_1}$  and  $C_{\beta_2}$  such that  $C_{\beta_1} \cap C_{\beta_2} = \{0\}$ , their intersections  $F_{\beta_1}$  and  $F_{\beta_2}$  with the unit ball are such that  $\bar{F}_{\beta_1} \neq \bar{F}_{\beta_2}$  where  $\bar{F}_{\beta_i}$  is the weak closure of  $F_{\beta_i}$ .

The minimal invariant cone  $C_\beta$  is the smallest cone invariant under the  $\mathcal{G}$ -action in  $\mathcal{A}'$  which contains  $\beta$  :

$$C_\beta = \{\lambda(T'(g)\beta), g \in \mathcal{G}, \lambda \in \mathbb{R}^+\}.$$

$\mathcal{A}'$  is equipped with the weak topology. In the case at hand  $\mathcal{G} = SL(2, \mathbb{R})$ ,  $\mathcal{A} = \mathcal{H}(P_1(\mathbb{R}))$ , and  $\mathcal{A}' \simeq \mathcal{H}(P_1(\mathbb{R}))$ .

By using Propositions 15 and 16 we arrive at the following

**Proposition 17** *Piard's sufficient conditions 1 and 2 are satisfied in the case of  $B(2, 1)$ .*

**Proof** *The proof is essentially the same as the corresponding proof for the original BMS group  $B$  given in [23].*

Consequently all the cylindrical ergodic measures on  $\mathcal{A}'$  associated to the IRS of  $B(2, 1)$  are concentrated on the  $SL(2, \mathbb{R})$ -orbits on  $\mathcal{A}'$ . These measures correspond to subsets of  $\mathcal{A}'$  which are "indecomposable" in the set theoretic sense. We conclude [22], [23] that the following Theorem holds:

**Theorem 8** *All IRS of  $B(2, 1)$  are induced.*

Thus all IRS of  $B(2, 1)$  are obtained by the Wigner–Mackey's inducing construction given in subsection 12.1.

## 17 Construction of the IRS of $B(2, 1)$

To find explicitly the operators of the induced representations of  $B(2, 1)$ , it suffices to provide the information cited in (i) and (ii) in subsection 12.1 for each of the orbit types. We note that all the little groups are abelian. All IRS of an abelian group are one-dimensional.

(i) The IRS of the "Little groups", i.e., of  $SO(2)$  and of  $C_n$ ,  $n$  even, have been given in [21].

(ii) A  $\mathcal{G}$ -quasi-invariant measure  $\mu$  on each orbit  $\mathcal{G}\phi \approx \mathcal{G}/L_\phi$  is required, however, a  $\mathcal{G}$ -invariant measure  $\mu$  on each orbit  $\mathcal{G}\phi \approx \mathcal{G}/L_\phi$  will be given in all cases. We note that when  $\mu$  is  $\mathcal{G}$ -invariant then

$$\frac{d\mu_g}{d\mu}(q) = 1, \quad g \in \mathcal{G}, \quad q \in \mathcal{G}\phi. \quad (17.1)$$

This is precisely what happens in the case of  $B(2, 1)$  where  $\mathcal{G} = G = \mathrm{SL}(2, R)$ , and  $L_\phi$  is either  $SO(2)$  or  $C_n$ ,  $n$  even. (17.1) applies in particular in (12.6). In [21]  $G$ -invariant measures on each orbit  $G\phi \approx G/L_\phi$  were obtained by projecting the unique (up to a multiplicative constant) Haar measure of  $G$  to the orbits  $G\phi \approx G/L_\phi$  via the canonical map

$$G \longrightarrow G/L_\phi.$$

However, explicit expressions for the  $G$ -invariant measures on the orbits  $G/L_\phi$  were not given in [21]. Here we give explicit expressions for these measures. Apart from their own interest, these measures will be particularly useful when we make contact with other approaches to quantum gravity, e.g. with canonical quantum gravity [55].

(ii $_\alpha$ ) The orbit  $01 \equiv G/L_\phi$ ,  $L_\phi = K = SO(2)$ , is the coset space  $\mathcal{S} = \mathrm{SL}(2, R)/SO(2)$ . It is well known [56] that every element  $\gamma$  of  $\mathcal{S}$  can be uniquely written in the form

$$\gamma = na(na)^T, \quad (17.2)$$

where,

$$n \in N = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \cong R, \quad a \in A = \begin{pmatrix} y & 0 \\ 0 & \frac{1}{y} \end{pmatrix} \cong R_{>0}, \quad (17.3)$$

$x \in R$ ,  $y > 0$ , and  $T$  denotes transposition.  $\mathcal{S}$  can be identified with the Poincare half-plane

$$H = \{(x, y) | y > 0, x, y \in R\} \quad (17.4)$$

via the map

$$(x, y) \longrightarrow \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} y & 0 \\ 0 & \frac{1}{y} \end{pmatrix} \left( \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} y & 0 \\ 0 & \frac{1}{y} \end{pmatrix} \right)^T. \quad (17.5)$$

A  $G$ -invariant measure on  $H$  is given by

$$d\mu = \frac{dx \wedge dy}{y^2}. \quad (17.6)$$

This is also, given (17.5), a  $G$ -invariant measure on  $\mathcal{S}$ .

(ii $_\beta$ ) A  $G$ -invariant measure on

$$G = \mathrm{SL}(2, R) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad a, b, c, d \in R, \quad ad - bc = 1 \right\},$$

is given by

$$dg = \frac{da \wedge db \wedge dc}{a}. \quad (17.7)$$



The orbits  $02 \equiv G/L_\phi$ ,  $L_\phi = C_n$ , where  $n$  is even, can be endowed with the  $G$ -invariant measure on  $G$  given in (17.7). Indeed, for a given little group  $L_\phi$ , the orbit  $02 \equiv G/L_\phi$  is the space of orbits of the right action  $T : G \times L_\phi \rightarrow G$  of the group  $L_\phi$  on  $G$  given by

$$g \star c := g \cdot c, \tag{17.8}$$

where  $g \in G$ , and  $c \in L_\phi$ . Thus the action  $T$  denoted by  $\star$  is identical to the group multiplication in  $G$ . Since the group  $L_\phi$  is finite and since the action (17.8) is fixed point free the coset spaces  $G/L_\phi$  inherit the measure on  $G$ .

This completes the necessary information in order to construct the induced representations of  $B(2, 1)$ . To conclude, in this paper *all* IRS of  $B(2, 1)$  have been constructed in the Hilbert topology.

## 18 Connections to Geometry

In three space–time dimensions with zero cosmological constant when the Einstein tensor vanishes so does the four–index Riemann–Christoffel curvature tensor. As a result, all three–dimensional Einstein spaces with zero cosmological constant are flat. Thus there are no local gravitational degrees of freedom but a finite number of global degrees of freedom which arise from the topology and matter content of the space–time.

In the case of the four–dimensional asymptotically flat space–times Foster showed [57], [58] that when Penrose’s future null infinity  $\mathfrak{J}^+$  is homeomorphic to  $R \times S^2$  there exists an asymptotic symmetry group, the original BMS group  $B$ , and vice versa. For other homeomorphic types of  $\mathfrak{J}^+$  there is no in general asymptotic symmetry group and when it exists the asymptotic symmetry group neither is it determined uniquely nor is it  $B$ .

By following through Foster’s analysis one can easily show that in the case of three–dimensional flat space–times a similar result holds:  $B(2, 1)$  is the asymptotic symmetry group if and only if  $\mathfrak{J}^+$  is homeomorphic to  $R \times S^1$ . This implies in particular that  $B(2, 1)$  cannot be the asymptotic symmetry group if a few generators of  $\mathfrak{J}^+$  are “missing” or if one has “only a piece of  $\mathfrak{J}^+$ ”.

Therefore in order to interpret the IRS of  $B(2, 1)$  and relate their properties with the properties of the underlying space–times we need to answer the question which of the non–compact three–manifolds which admit flat Lorentzian metrics also admit, when they are conformally compactified, future null infinity  $\mathfrak{J}^+$  homeomorphic to  $R \times S^1$ . A complete answer to this question cannot be given as these space–times have not yet been studied in much detail [59].

A second class of well–known (2+1)–space–times are those solutions to the Einstein equations with particle matter content. They were first investigated in the Euclidean context by Staruszkiewicz [60] and in the Lorentzian signature by Deser, Jackiw and ’t Hooft [61]. A lot of work has been done on this class of space–times since then, e.g., [62], [63], [64], [65], [66].

The simplest space–time in this class contains a single spinless point particle with mass  $m$  in  $R^3$  at rest at the origin. The corresponding metric was constructed in [61]. The space

is flat, but a wedge with opening angle  $2\pi(1 - a)$ ,  $a = 1 - 4Gm$ , is cut out, and its edges are identified. Thus the space is a cone, the unique two–space which is flat except at one point, its vertex. The angular defect  $2\pi(1 - a) = 8\pi Gm$  characterises the mass.

The cone geometry has the following consequence: The metric admits local solutions of the killing equation corresponding to the full Poincare group, most of the resulting would–be killing vectors are not preserved under the identification of the edges of the wedge. Only the killing vectors corresponding to rotations and time translation are globally defined.

Space–times with multiple point particles were introduced in [60], [61], [62]. Curvature is created by the particles, but only locally at their position; elsewhere space–time remains flat. Their physical properties have been studied in [60], [61], [62], [63], [64], [65], [66], for an overview see e.g. [59]. Their mathematical structure is more involved than that of vacuum space–times and its study was initiated in [67], [68], [69].

Space–times with multiple point particles define manifolds with conical singularities or, in the case where the masses of all particles are rational multiples of  $2\pi$ , orbifolds. With the exception of the orbifold case, they cannot be obtained as quotients of regions in three–dimensional Minkowski space.

Outside point particles, space–time is flat. All effects of localized sources are on the global geometry, which is fixed by singularities of the worldlines of the particles; these are arbitrary flat space geodesics. This means in particular that the conserved quantities, total energy–momentum and angular momentum, are related to topological invariants. There is for example a static  $N$ –body solution [61] with conical spatial geometry, whose total energy is additive and determines the Euler invariant of the spatial surface.

We await that the IRS of  $B(2, 1)$  in the Hilbert topology will make contact with both the empty space–times and the space–times with particle matter content via the set of invariants which classify the  $SL(2, R)$ –orbits on  $\mathcal{A}' \simeq \mathcal{H}(P_1(R))$ .

For example the invariant which corresponds to the Poincare “mass squared” invariant  $m^2$  is identified with the mass of the particle matter content of the associated space–time, and subsequently, at least in the case of the static  $N$ –body solution [61], this invariant determines the Euler invariant of the spatial surface of this space–time.

In the case of the simplest space–time of a single spinless point particle at rest at the origin in  $R^3$  the mass  $m$  also determines the holonomy group of the space–time manifold which is either a finite or a countable dense subgroup of  $SO(2)$  according to whether  $1 - 4Gm$  is rational or not [70]. Holonomy groups serve as tools for the global classification of space–times [71] in a manner similar to but distinct from the local Petrov and Segre type classifications.

Two–body quantum–mechanical scattering in 2+1 gravity has been studied in [62], [72], [73]. Since there are no gravitons forces are not mediated by graviton exchange, rather, they are geometrical/topological in origin, arising from global properties of space–time, which is not Minkowskian in the large, even when it is locally flat.

It is expected that the IRS of  $B(2, 1)$  in the nuclear topology associated with non–compact little groups which leave invariant distributional supermomenta will make contact with the two–body and more generally with the  $N$ –body quantum–mechanical scattering in 2+1

gravity. This also it is expected to happen when  $\mathcal{A}'$  is enlarged to the space  $\mathcal{Y}(S^1)$  of real hyperfunctions on  $S^1$ .

One of the most interesting features of the representation theory of  $B$  and of its generalizations in all signatures and in all space–time dimensions is the appearance of discrete (finite and infinite) groups as “little groups” [9], [11], [24], [25], [26], [27], [30].

In earlier papers [24], [42], [74] (and more forcibly in [25]) it has been argued that the IRS of  $B$  and of its generalizations in complex space–times as well as in space–times with Euclidean or Ultrahyperbolic signature induced from finite “little groups” are what really lie behind the full description of (unconstrained) moduli spaces of (generalised) gravitational instantons. It is an interesting open question to find if the IRS of  $B(2,1)$  induced from the cyclic “little groups”  $C_n$  of even order correspond to gravitational instantons in three space–time dimensions.

The holographic description of bulk physics in terms of a theory living on the boundary has been concretely realised by the AdS/CFT correspondence for space–times with global negative curvature. It remains an important outstanding problem to understand suitable formulations of holography for flat space–time, a goal that has elicited a considerable amount of work from several complementary approaches [75], [76], [77], [78], [79].  $B(2,1)$  is going to be relevant for exploring holography in three–dimensional Minkowski space–time.

## References

- [1] Bondi H. & Van Der Berg M. G. J. & Metzner A. W. K. *Gravitational waves in general relativity VII. Waves from axi–symmetric isolated sources*. Proc. R. Soc. Lond. A. **269**, 21–51 (1962).
- [2] Sachs R. K. *Gravitational waves in general relativity VIII. Waves in asymptotically flat space–time*. Proc. R. Soc. Lond. A. **270**, 103–126 (1962).
- [3] Penrose R. *Asymptotic Properties of Fields and Space–Times*. . Phys. Rev. Lett. **10**, 66 (1963).
- [4] Penrose R. *Relativistic symmetry groups, in: Group theory and nonlinear problems*. Ed. A.O. Barut. Reidel, Dordrecht (1976).
- [5] Penrose, R. & Rindler W. *Spinors and spacetime, Vol. 2*. Cambridge Univ. Press, Cambridge (1986).
- [6] Wigner E. *On unitary representations of the Inhomogeneous Lorentz Group*. Annals of Mathematics **40**, 149–204 (1939).
- [7] McCarthy P. J. *Representations of the Bondi–Metzner–Sachs group I. Determination of the representations*. Proc. R. Soc. Lond. A. **330**, 517–535 (1972).
- [8] McCarthy P. J. *Structure of the Bondi–Metzner–Sachs group*. J. Math. Phys. **13**, 1837–1842 (1972).
- [9] McCarthy P. J. *Representations of the Bondi–Metzner–Sachs group II. Properties and classification of the of the representations*. Proc. R. Soc. Lond. A. **333**, 317–336 (1973).
- [10] McCarthy P. J. & Crampin M. *Representations of the Bondi–Metzner–Sachs group III. Poincare spin multiplicities and irreducibility*. Proc. R. Soc. Lond. A. **335**, 301–311 (1973).

- [11] McCarthy P. J. *The Bondi–Metzner–Sachs group in the nuclear topology*. Proc. R. Soc. Lond. A. **343**, 489–523 (1975).
- [12] McCarthy P. J. *Lifting of projective representations of the Bondi–Metzner–Sachs group in the nuclear topology*. Proc. R. Soc. Lond. A. **358**, 141–171 (1978).
- [13] McCarthy P. J. & Crampin M. *Physical significance of the topology of the Bondi–Metzner–Sachs group*. Phys. Rev. Letters. **33**, 547–550 (1974).
- [14] McCarthy P. J. *Asymptotically flat Space–Times and Elementary Particles*. . Phys. Rev. Lett. **29**, 817–819 (1972).
- [15] Mackey G. W. *Induced Representations of Locally Compact Groups I*. Annals of Mathematics **55**, 101–139 (1952).
- [16] Mackey G. W. *Induced Representations of Locally Compact Groups II. The Frobenius Reciprocity Theorem*. Annals of Mathematics **58**, 193–221 (1953).
- [17] Mackey G. W. *Induced representations of groups and quantum mechanics*. Benjamin, New York, Amsterdam and Torino (1968).
- [18] Mackey G. W. 1955 *The theory of group representations*. The University of Chicago Press, Chicago (1955).
- [19] Simms D. J. *Lie groups and quantum mechanics*. Bonn notes Springer, Berlin, Heidelberg and New York (1968).
- [20] Isham C. J. *Topological and quantum aspects of quantum theory*. in Relativity groups and topology. North-Holland, Amsterdam (1984).
- [21] Melas E. *On the representation theory of the Bondi–Metzner–Sachs group and its variants in three space–time dimensions* JMP **58**, 071705 (2017).
- [22] Piard A. *UNITARY REPRESENTATIONS OF SEMIDIRECT PRODUCT GROUPS WITH INFINITE DIMENSIONAL ABELIAN NORMAL SUBGROUP*. Rep. Math. Phys. **11**, 259–278 (1977).
- [23] Piard A. *REPRESENTATIONS OF THE BONDI–METZNER–SACHS GROUP WITH THE HILBERT TOPOLOGY*. Rep. Math. Phys. **11**, 279–283 (1977).
- [24] McCarthy P. J. *Real and Complex Asymptotic Symmetries in Quantum Gravity, Irreducible Representations, Polygons, Polyhedra, and the A, D, E Series*. Phil.Trans. R. Soc. Lond. A. **338**, 271–299 (1992).
- [25] Melas E. *The BMS group and generalized gravitational instantons*. J. Math. Phys. **45** 3, 996–1002 (2004).
- [26] McCarthy P. J. & Melas E. *On irreducible representations of the ultrahyperbolic BMS group*. Nuc. Phys. B. **653**, 369–399 (2003).
- [27] Melas E. *Construction of the irreducibles of  $B(2,2)$* . J. Phys. A: Math. Gen. **39**, 3341–3366 (2006).
- [28] Melas E. *Representations of the Ultrahyperbolic BMS HB. I. General Results* (unpublished)
- [29] Melas E. *Representations of the ultrahyperbolic BMS group HB. II. Determination of the representations induced from infinite "little groups"*. (2013) arXiv:1312.0532v1 (2013).
- [30] Melas E. *Representations of the ultrahyperbolic BMS group HB. III. Determination of the representations induced from finite "little groups"*. arXiv:1402.1428v1 (2014).

- [31] Ashtekhar A. & Bičák J. & Schmidt B. G. *Asymptotic structure of symmetry reduced general relativity*. Phys. Rev. D. **55** 2, 669–686 (1997).
- [32] Barnich G. & Troessaert C. *Aspects of the BMS/CFT correspondence*. J. High Energy Phys. **062** (2010).
- [33] Barnich G. & Oblak B. *Notes on the BMS group in three dimensions: I. Induced representations*. J. High Energy Phys. **129** (2014).
- [34] Barnich G. & Oblak B. *Notes on the BMS group in three dimensions: II. Coadjoint representation*. J. High Energy Phys. **33** (2015).
- [35] Newman E. T. & Unti W. J. *Behavior of Asymptotically Flat Empty Spaces*. JMP **3**, 891–901 (1962).
- [36] Sachs R. K. *Asymptotic Symmetries in Gravitational Theory*. Phys. Rev. **128**, 2851–2864 (1962).
- [37] Penrose R. *Spinors and space–time Volume 2*. Cambridge University Press (1986).
- [38] Hawking S. W. & Perry M. J. & Strominger A. *Soft hair on black holes*. arXiv:1601.00921v1
- [39] Giddings S. Abbot J. & Kuchar K. *Einstein’s Theory in a Three-Dimensional Space–Time*. Gen. Rel. Grav. **16**, No 8, 751–775 (1984).
- [40] Mackey G. W. *Induced representations of groups and quantum mechanics* (New york : Benjamin) (1968).
- [41] Mackey G. W. *Unitary group representations in physics, probability and number theory*. London. Benjamin. (1978).
- [42] McCarthy P. J. & Melas E. *On irreducible representations of the ultrahyperbolic BMS group*. Nucl. Phys. B. **653**, 369–399 (2003).
- [43] Stone M. H. *Linear transformations in Hilbert space III, operational methods and group theory* Proc. Nat. Acad. Sci. U.S.A. **16**, 172–175 (1930).
- [44] Cantoni V. *On the representations of the Bondi–Metzner–Sachs group* Ph.D. thesis, University of London (1967).
- [45] McCarthy P. J. & Crampin M. *Representations of the Bondi–Metzner–Sachs group III. Poincare Spin Multiplicities and Irreducibility* Proc. R. Soc. Lond. A. **335**, 301–311 (1973).
- [46] Mackey G. W. *Induced Representations of Groups and Quantum Mechanics* Benjamin, Amsterdam, Torino, New York, (1968).
- [47] Penrose R. *Zero Rest–Mass Fields Including Gravitation: Asymptotic Behaviour*. Proc. R. Soc. Lond. A. **284**, 159–203 (1965).
- [48] Girardello L. & Parravicini G. *Continuous Spins in the Bondi–Metzner–Sachs Group of Asymptotic Symmetry in General Relativity*. Phys. Rev. Let. **32**, 10 565–568 (1974).
- [49] Crampin M. & McCarthy P. J. *Physical Significance of the Topology of the Bondi–Metzner–Sachs Group*. Phys. Rev. Let. **33**, 9 547–550 (1974).
- [50] McCarthy P. J. *Hyperfunctions and asymptotic symmetries*. Proc. R. Soc. Lond. A. **358**, 495–498 (1978).
- [51] Sobolev S. L. *Partial Differential Equations of Mathematical Physics*. Pergamon New York (1964).

- [52] Helgason S. *Differential geometry and symmetric spaces*. Academic Press. (1962).
- [53] Hausner M. & Schwartz J. T. *Lie groups, Lie algebras*. Thomas Nelson and Sons Ltd London (1968).
- [54] E. Melas Representations of the Bondi–Metzner–Sachs group in three space–time dimensions in the Hilbert topology II. Properties and Classification of the Representations (under preparation)
- [55] Martinec E. *Soluble systems in quantum gravity* Phys. Rev. D **30**,6, 1198–1204 (1984).
- [56] Helgason S. *Lie Groups and Symmetric Spaces* Battelle Recontres, p. 1, edited by C. M. DeWitt and J. A. Wheeler, Benjamin, New York (1968).
- [57] Foster J. *Conformal structure of  $\mathfrak{I}^+$  and asymptotic symmetry I. Definitions and local theory*. J. Phys. A: Math. Gen. **11**, 1, 93–102 (1978).
- [58] Foster J. *Asymptotic Symmetry and the Global Structure of Future Null Infinity*. Int. J. of Theor. Phys. **26**, 11, 1107–1124 (1987).
- [59] Carlip S. *Quantum Gravity in 2+1 Dimensions*. Cambridge University Press (1998).
- [60] Staruszkiewicz A. *Gravitation theory in three–dimensional space* . Acta Phys. Pol. **24**, 735–740 (1963).
- [61] Deser S. & Jackiw R. & 't Hooft G. *Three–dimensional Einstein gravity: dynamics of flat space* . Ann. Phys. **152**, 220–235 (1984).
- [62] Deser S. & Jackiw R. *Classical and quantum scattering on a cone* . Commun. Math. Phys. **118**, 495–509 (1988).
- [63] de Sousa Gerbert P. *On spin and (quantum) gravity in 2+ 1 dimensions* . Nucl. Phys. B **346**, 440–472 (1990).
- [64] 't Hooft G. *The evolution of gravitating point particles in 2+1 dimensions* . Class. Quantum Grav. **10**, 1023–1038 (1993).
- [65] 't Hooft G. *Canonical quantization of gravitating point particles in 2+1 dimensions*. Class. Quantum Grav. **10**, 1653–1664 (1993).
- [66] 't Hooft G. *Quantization of point particles in (2+1)-dimensional gravity and spacetime discreteness*. Class. Quantum Grav. **13**, 1023–1039 (1996).
- [67] Krasnov K. & Schlenker J. *Minimal surfaces and particles in 3–manifolds* Geometriae Dedicata **126** 187–254 (2007) (arXiv:math/0511441).
- [68] Bonsante F. & Krasnov K. & Schlenker J. *Multi–black holes and earthquakes on Riemann surfaces with boundaries* Int. Math. Res. Not. **2011** 487–552 (2010) (arXiv:math/0610429).
- [69] Barbot T. & Bonsante F. & Schlenker J. *Collisions of particles in locally AdS space–times: I. Local description and global examples* (2010) (arxiv.org/abs/1010.3602v1).
- [70] Oliveira–Neto G. *Identifying conical singularities* J. Math. Phys. **37** 4716–4723 (1996).
- [71] Hall G. S. & Lonie D. P. *Holonomy groups and spacetimes*. Class. Quantum Grav. **17**, 1369–1382 (2000).
- [72] 't Hooft G. *Non–Perturbative 2 Particle Scattering Amplitudes in 2+1 Dimensional Quantum Gravity*. Commun. Math. Phys. **117**, 685–700 (1988).

- [73] de Sousa Gerbert P. & Jackiw R. *Classical and Quantum Scattering on a Spinning Cone*. Commun. Math. Phys. **124**, 229–260 (1989).
- [74] Melas E. *Approaching quantum gravity via the representation theory of the BMS group and its variants. Some conceptual issues and a conjecture* Nucl. Phys. B. (Proc. Suppl.) **104**, 212–216 (2002).
- [75] J. de Boer & Solodukhin S. N. *A Holographic reduction of Minkowski space–time*. Nucl. Phys. B. **665**, 545 (2003).
- [76] Cheung C. A. de la Fuente & Sundrum R. *4D scattering amplitudes and asymptotic symmetries from 2D CFT*. JHEP **1701**, 112 (2017).
- [77] Kapec D. Mitra P. Raclariu A. M. & Strominger A. *2D Stress Tensor for 4D Gravity*. Phys. Rev. Lett. **119**, 12, 121601 (2017).
- [78] Kapec D. Lysov V. Pasterski S. & Strominger A. *Semiclassical Virasoro symmetry of the quantum gravity S–matrix*. JHEP **1408**, 058 (2014).
- [79] Cachazo F. & Strominger A. *Evidence for a New Soft Graviton Theorem*. arXiv: 1404.4091.