

Planar graphs with the maximum number of induced 4-cycles or 5-cycles

Michael Savery

Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK, and
the Heilbronn Institute for Mathematical Research, Bristol, UK
savery@maths.ox.ac.uk

September 28, 2021

Abstract

For large n we determine exactly the maximum numbers of induced C_4 and C_5 subgraphs that a planar graph on n vertices can contain. We show that $K_{2,n-2}$ uniquely achieves this maximum in the C_4 case, and we identify the graphs which achieve the maximum in the C_5 case. This extends work in a paper by Hakimi and Schmeichel and a paper by Ghosh, Győri, Janzer, Paulos, Salia, and Zamora which together determine both maxima asymptotically.

1 Introduction

An important class of problems in extremal graph theory concerns determining the maximum number of induced copies of a small graph H that can be contained in a graph on n vertices. These questions were first considered by Pippenger and Golumbic in [19] where they showed among other things that for every k -vertex graph H , the maximum number of induced copies of H in an n -vertex graph is asymptotically at least $\frac{n^k}{k^{k-k}}$. The maximum is now known asymptotically for all graphs H on at most four vertices except the path of length 3, as well as for certain complete partite graphs. See [6] and its references for a good summary of these results, and see [15] and [22] for some results for other graphs H .

The case where H is a cycle has received particular attention. Pippenger and Golumbic conjectured in their paper that the lower bound stated above is asymptotically tight for cycles of length at least 5. This conjecture was verified for 5-cycles by Balogh, Hu, Lidický, and Pfender in [2], where they determined exactly the maximum number of induced 5-cycles in a graph on n vertices when n is large. The best upper bound known for general cycles is due to Král', Norin, and Volec in [17], who showed that for all n and $k \geq 5$, every n -vertex graph contains at most $2n^k/k^k$ induced k -cycles.

A closely related problem is to determine the maximum number of induced copies of H that can be contained in a planar graph on n vertices. We will write $f_I(n, H)$ for this quantity and $f(n, H)$ for the corresponding quantity when the copies of H do not have to be induced.

We consider in particular the case where H is a small cycle. Hakimi and Schmeichel showed in [14] that $f(n, C_3) = 3n - 8$ for $n \geq 3$, and $f(n, C_4) = \frac{1}{2}(n^2 + 3n - 22)$ for $n \geq 4$. Since a 3-cycle in a graph is always induced, this gives $f_I(n, C_3) = 3n - 8$. Also, it is straightforward to see that the complete bipartite graph $K_{2, n-2}$ contains $\frac{1}{2}(n^2 - 5n + 6)$ induced 4-cycles, so $f_I(n, C_4) = \frac{1}{2}n^2 + O(n)$, as observed in [7]. In [12] Győri, Paulos, Salia, Tompkins, and Zamora determined $f(n, C_5)$ exactly for all $n \geq 5$, and in [7] Ghosh, Győri, Janzer, Paulos, Salia, and Zamora showed that $f_I(n, C_5) = \frac{1}{3}n^2 + O(n)$.

In [3] and [4] Cox and Martin determined $f(n, H)$ asymptotically when H is a small even cycle, showing that $f(n, C_{2k}) = \binom{n}{k}^k + o(n^k)$ for $k \in \{3, 4, 5, 6\}$. For $n \equiv 0 \pmod{k}$, the graph $G_{n,k}$ defined by replacing every second vertex in a $2k$ -cycle with $\frac{n}{k} - 1$ copies of that vertex contains $(\frac{n}{k} - 1)^k$ induced $2k$ -cycles, hence also $f_I(n, C_{2k}) = \binom{n}{k}^k + o(n^k)$ for $k \in \{3, 4, 5, 6\}$. Cox and Martin go on to conjecture that $f(n, C_{2k}) = \binom{n}{k}^k + o(n^k)$ for all $k \geq 7$, which would similarly determine $f_I(n, C_{2k})$ asymptotically for all $k \geq 7$.

Much less is known for odd cycles of length greater than 5. In the same spirit as above, by evenly blowing up k pairwise non-adjacent vertices in a $(2k+1)$ -cycle until the graph has approximately n vertices, one can obtain a lower bound of $\binom{n}{k}^k + o(n^k)$ for $f_I(n, C_{2k+1})$ for all $k \geq 1$. One can obtain a slightly better lower bound for $f(n, C_{2k+1})$ by adding a path through each of the blown up sets of vertices in the graph $G_{n,k}$ defined above to obtain a planar graph containing $2k \binom{n}{k}^k + o(n^k)$ non-induced copies of C_{2k+1} . These blow-up constructions were first given in [10] and [13], and were also considered in [3] and [4]. Hakimi and Schmeichel showed in [14] that $f(n, C_{2k+1}) = O(n^k)$, and this seems to be the best known upper bound.

For results on $f(n, H)$ for various other graphs H see [1], [3], [5], [10], [11], [20], and [21]. In the general case, Huynh, Joret, and Wood proved a far-reaching result in [16] which gives the order of magnitude of $f(n, H)$ for all H in terms of a graph parameter called the ‘flap-number’ of H . Very recently, in [18], Liu showed (Corollary 6.1 in that paper) that for all graphs H , $f_I(n, H) = \Theta(f(n, H))$, thus determining the order of magnitude of $f_I(n, H)$ for all H .

In this paper we will determine $f_I(n, C_4)$ and $f_I(n, C_5)$ exactly for large n , and will identify the graphs which exhibit these maxima. Our result for 4-cycles is the following.

Theorem 1. *For large n , $f_I(n, C_4) = \frac{1}{2}(n^2 - 5n + 6)$. Moreover, for large n , the only n -vertex planar graph which contains $f_I(n, C_4)$ induced 4-cycles is the complete bipartite graph $K_{2, n-2}$.*

Turning to the 5-cycle case, we first define a family of graphs which we will show to be the only n -vertex planar graphs containing $f_I(n, C_5)$ induced 5-cycles for large n .

Definition 1. A graph G on $n \geq 19$ vertices is *of the required form* if it contains distinct vertices u_1, u_2, u_3 , and w such that the remainder of its vertices can be partitioned into

sets $A = \{a_1, \dots, a_{|A|}\}$, $B = \{b_1, \dots, b_{|B|}\}$, $C = \{c_1, \dots, c_{|C|}\}$, and $Z = \{z_1, \dots, z_6\}$ such that all of the following hold:

1. (a) if $n \equiv 1 \pmod{3}$, then $|A| = |B| = \frac{n-7}{3}$ and $|C| = \frac{n-16}{3}$,
 (b) if $n \equiv 2 \pmod{3}$, then $|A| = \frac{n-5}{3}$, $|B| = \frac{n-8}{3}$, and $|C| = \frac{n-17}{3}$, or $|A| = |B| = \frac{n-8}{3}$ and $|C| = \frac{n-14}{3}$,
 (c) if $n \equiv 0 \pmod{3}$, then $|A| = \frac{n}{3} - 2$, $|B| = \frac{n}{3} - 3$, and $|C| = \frac{n}{3} - 5$, or $|A| = |B| = \frac{n}{3} - 2$ and $|C| = \frac{n}{3} - 6$,
2. the edge set of G contains the edges u_1u_2 , u_1u_3 , u_2u_3 , u_1z_1 , u_2z_1 , u_1z_2 , u_2z_4 , z_1z_2 , z_1z_3 , z_1z_4 , z_2z_3 , z_3z_4 , z_2w , z_3w , z_4w , u_2z_5 , u_3z_5 , z_5w , u_1z_6 , u_3z_6 , and z_6w , and the edges in the three complete bipartite graphs with vertex classes $\{u_1, w\}$ and A , $\{u_2, w\}$ and B , and $\{u_3, w\}$ and C , and
3. the remaining edges of G are taken from $a_{|A|}z_2$, z_4b_1 , $b_{|B|}z_5$, z_5c_1 , $c_{|C|}z_6$, z_6a_1 , $a_i a_{i+1}$ for $1 \leq i \leq |A| - 1$, $b_i b_{i+1}$ for $1 \leq i \leq |B| - 1$, and $c_i c_{i+1}$ for $1 \leq i \leq |C| - 1$.

We refer to the edges in point 3 of Definition 1 as *optional* edges of a graph of the required form, and if none of these edges are present then we say that G is a *principal graph of the required form*. An illustration of a general graph of the required form is given in Figure 1. In this illustration red lines represent optional edges. A graph of the required form is clearly planar. It is interesting to note that a graph of the required form in which all optional edges are present is a maximal planar graph.

Roughly speaking, a large graph of the required form is built from a smaller graph of the required form with $n \equiv 1 \pmod{3}$ vertices by “adding vertices to A , B , and C as evenly as possible”. If the number of vertices in the smaller graph has some other value modulo 3, then we need to add the first one or two vertices to particular classes before adding the rest as evenly as possible.

In a large graph of the required form almost all of the vertices are in A , B , or C . For each $a \in A$ and $b \in B$ the cycle u_1awbu_2 is an induced 5-cycle in the graph. Similarly, there is an induced 5-cycle containing each pair of vertices $b \in B$ and $c \in C$, and each pair $a \in A$ and $c \in C$. Since these 5-cycles are different for different pairs of vertices, and each of A , B , and C has size $\frac{1}{3}n - O(1)$, this accounts for $3(\frac{1}{3}n - O(1))^2 = \frac{1}{3}n^2 + O(n)$ induced 5-cycles in G .

Theorem 2. *For large n ,*

$$f_I(n, C_5) = \begin{cases} \frac{1}{3}(n^2 - 8n + 22), & \text{if } n \equiv 1 \pmod{3} \\ \frac{1}{3}(n^2 - 8n + 21), & \text{if } n \equiv 0, 2 \pmod{3}. \end{cases}$$

Moreover, for large n , if G is a planar graph on n vertices, then G contains $f_I(n, C_5)$ induced 5-cycles if and only if G is of the required form.

The conclusion of Theorem 2 does not hold for all n . Indeed, for $n = 10$ Theorem 2 would assert that no planar graph on 10 vertices contains more than 14 induced 5-cycles,




Figure 1: A graph of the required form

but the graph shown in Figure 2 contains 16. In this figure, the colours are only intended to highlight the structure of the graph.

For longer cycles we make the following conjecture based on the constructions given above.

Conjecture 3. *For $k \geq 6$ and n sufficiently large relative to k , the n -vertex graph obtained by blowing up $\lfloor \frac{k}{2} \rfloor$ pairwise non-adjacent vertices in a k -cycle to sets of as equal size as possible contains the most induced k -cycles of any planar graph.*




Figure 2: A planar graph with 10 vertices and 16 induced 5-cycles

It would also be interesting to know how many induced k -cycles can be contained in an n -vertex graph which can be embedded in a fixed surface for surfaces other than the sphere.

Question 4. *For each $k \geq 3$ and each surface Σ other than the sphere, what is the maximum number of induced k -cycles which can be contained in an n -vertex graph which can be embedded in Σ ?*

The results of Huynh, Joret, and Wood [16] and Liu [18] described above extend to all surfaces Σ . Combined, these results show that the answer to Question 4 is $\Theta(n^{\lfloor k/2 \rfloor})$ for all $k \geq 3$ and all surfaces Σ .

1.1 Notation and organisation of the paper

In this paper all graphs are simple, and we use the following standard graph theoretic notation. For a graph G we write $V(G)$ and $E(G)$ for the vertex and edge sets of G respectively. For a vertex v of a graph G we write $N_G(v)$ for the set of neighbours of v in G , also called the neighbourhood of v in G , and $d_G(v)$ for the degree of v in G . In both cases we drop the subscript if the graph in question is clear. For a graph G and a set $S \subseteq V(G)$, we write $G[S]$ for the induced subgraph of G with vertex set S , and $G - S$ for the induced subgraph with vertex set $V(G) \setminus S$. We write $K_{a,b}$ for the complete bipartite graph with parts of size a and b , and we write C_k for the k -cycle graph.

In Section 2 we prove Theorem 1 after stating and proving two preliminary lemmas. Section 3 contains the proof of Theorem 2, which is broken down into three small preliminary lemmas and two larger lemmas which are proved in Sections 4 and 5 respectively. The proofs of these two lemmas contain the bulk of the work in proving the theorem. One of these proofs involves some repetitive case checking, which is handled in Appendix A.

1.2 Update

Very recently, the authors of [7] updated their paper and independently proved Theorem 2 [8].

2 Proof of Theorem 1

In this section we give two preliminary lemmas and then prove Theorem 1.

Lemma 5. *Let n be large, and suppose that G is an n -vertex planar graph in which every vertex of G is in at least $n - 3$ induced 4-cycles. If G contains a vertex of degree 2, then G is isomorphic to $K_{2,n-2}$.*

Proof. Suppose that $v \in V(G)$ has degree 2. Let the neighbours of v be u and w . Since v is in at least $n - 3$ induced 4-cycles, the remaining $n - 3$ vertices of G are all adjacent to both u and w , so u and w are adjacent to all vertices of G except each other.

Suppose that $x_1x_2x_3x_4$ is a 4-cycle in G which doesn't contain u or w . Then $\{u, x_1, x_3\}$ and $\{w, x_2, x_4\}$ form the partite sets of a subdivision of $K_{3,3}$ in G , which contradicts the planarity of G by Kuratowski's theorem. Hence every 4-cycle in G contains u or w . Any induced 4-cycle in G containing u must also contain w , and vice versa, since no vertex in an induced 4-cycle can be a neighbour of all the others in the cycle. Hence every induced 4-cycle in G has the form $uxwy$ for some $x, y \in V(G) \setminus \{u, w\}$.

Let $z \in V(G) \setminus \{u, w\}$. Then every induced 4-cycle in G containing z is of the form $uzwy$ where $y \in V(G) \setminus \{u, w, z\}$. By assumption, there are at least $n - 3$ such cycles, so z is not adjacent to any vertex in G other than u and w . Therefore G is isomorphic to $K_{2,n-2}$ as required. \square

Lemma 6. *Let n be large, and suppose that G is an n -vertex planar graph in which every vertex of G is in at least $n - 3$ induced 4-cycles. Then G contains no vertices of degree 3, 4, or 5.*

Proof. Suppose for a contradiction that $v \in V(G)$ has degree 3, 4, or 5. Let the neighbours of v in G be $x_1, \dots, x_{d(v)}$. The number of induced 4-cycles in G containing v is at most

$$\sum_{1 \leq i < j \leq d(v)} |(N(x_i) \cap N(x_j)) \setminus (\{v\} \cup N(v))|. \quad (1)$$

For each set of three neighbours of v , there is at most one common neighbour of all three other than v , otherwise G has an obvious $K_{3,3}$ subgraph. So for each distinct pair of terms in sum (1), of which there are at most

$$\binom{\binom{5}{2}}{2} = 45,$$

there is at most one vertex contributing to both terms, i.e. there are at most 45 vertices which contribute to more than one term in the sum. Each of these could contribute up

to 10 in total, since there are at most 10 terms in the sum, so the total contribution to the sum from these vertices is at most 450.

By assumption v is in at least $n - 3$ induced 4-cycles, so there are at least $n - 453$ induced 4-cycles containing v in which the vertex opposite v in the cycle is not in any other induced 4-cycle containing v . This implies that there are at least $n - 453$ vertices in $V(G) \setminus (\{v\} \cup N(v))$ which are adjacent to at least two neighbours of v . So there are at most 449 vertices in $V(G) \setminus (\{v\} \cup N(v))$ which are not adjacent to at least two neighbours of v .

By the pigeonhole principle, we may assume that $|(N(x_1) \cap N(x_2)) \setminus (\{v\} \cup N(v))| \geq (n - 3)/10$. Draw G and consider the induced drawing of the complete bipartite graph with parts $\{x_1, x_2\}$ and $(N(x_1) \cap N(x_2)) \setminus (\{v\} \cup N(v))$. Label the vertices in the latter set as y_1, \dots, y_k in natural order, where $k \geq (n - 3)/10$. The drawing of the complete bipartite graph splits the plane into k regions R_1, \dots, R_k , where for each i , R_i is bounded by the cycle $x_1 y_i x_2 y_{i+1}$, where here and henceforth we take subscript addition to be modulo k .

Let R_j be the region with v in its interior. Then every vertex in $N(v)$ is in R_j (including its boundary). Hence every vertex with a neighbour in $N(v) \setminus \{x_1, x_2\}$ is in $R_{j-2} \cup R_{j-1} \cdots \cup R_{j+2}$ (including its boundary). There are at most 449 vertices in $V(G) \setminus (\{v\} \cup N(v))$ which are not adjacent to at least two neighbours of v , so there are at most 449 values of i other than $j - 2, \dots, j + 2$ for which the interior of R_i contains a vertex. Thus since n is large, there exists i such that R_{i-1} and R_i have no vertices in their interiors.

By assumption y_i is in at least $n - 3$ induced 4-cycles. Since $N(y_i) \subseteq \{x_1, x_2, y_{i-1}, y_{i+1}\}$, every such 4-cycle either contains the path $y_{i-1} y_i y_{i+1}$ or the path $x_1 y_i x_2$. However since n is large (so $k \geq 5$), the existence of y_{i-2} and y_{i+2} implies there is no common neighbour of y_{i-1} and y_{i+1} in G other than x_1 and x_2 , which are both also neighbours of y_i . Hence there are no induced 4-cycles containing the path $y_{i-1} y_i y_{i+1}$. Therefore there are $n - 3$ common neighbours of x_1 and x_2 besides y_i , and y_i is not adjacent to any of them. Hence y_i has degree 2 in G , so by Lemma 5 G is isomorphic to $K_{2, n-2}$. But for large n this has no vertex of degree 3, 4, or 5, which gives the required contradiction and completes the proof of the lemma. \square

Proof of Theorem 1. First, it is straightforward to see that every induced 4-cycle in $K_{2, n-2}$ contains exactly two vertices from the part of size $n - 2$, and that there is a unique, distinct induced 4-cycle containing each such pair, so the number of induced 4-cycles in $K_{2, n-2}$ is $\binom{n-2}{2} = \frac{1}{2}(n^2 - 5n + 6)$.

By Lemmas 5 and 6, to prove the theorem it is sufficient to show that for large n any n -vertex planar graph containing $f_I(n, C_4)$ induced 4-cycles has no vertex in fewer than $n - 3$ induced 4-cycles. Indeed, if this is the case, then if n is large and G is an n -vertex planar graph containing $f_I(n, C_4)$ induced 4-cycles, then the minimum degree of G is at most 5 by the planarity of G and at least 2 since every vertex is in at least one induced 4-cycle. So by the lemmas G is isomorphic $K_{2, n-2}$.

Let n be large and suppose that G is an n -vertex planar graph in which every vertex of G is in more than $n - 3$ induced 4-cycles. Then as above the minimum degree of G

is at least 2 but at most 5, so by Lemmas 5 and 6 G is isomorphic to $K_{2,n-2}$. But this contains vertices in at most $n - 3$ induced 4-cycles, which is a contradiction. Thus for large n , every n -vertex planar graph has a vertex in at most $n - 3$ induced 4-cycles.

For large n , deleting a vertex in at most $n - 3$ induced 4-cycles from an n -vertex planar graph containing $f_I(n, C_4)$ induced 4-cycles yields an $(n - 1)$ -vertex planar graph containing at least $f_I(n, C_4) - n + 3$ induced 4-cycles. This shows that $f_I(n, C_4) \leq f_I(n - 1, C_4) + n - 3$ for large n .

We have seen that there are exactly $\frac{1}{2}(n^2 - 5n + 6)$ induced 4-cycles in $K_{2,n-2}$ for each n , so there are $n - 3$ more induced 4-cycles in $K_{2,n-2}$ than there are in $K_{2,n-3}$. If $f_I(n, C_4) < f_I(n - 1, C_4) + n - 3$ for infinitely many values of n , then for large enough n we have $\frac{1}{2}(n^2 - 5n + 6) > f_I(n, C_4)$, which is a contradiction. Hence for large n , $f_I(n, C_4) = f_I(n - 1, C_4) + n - 3$. Therefore if n is large enough that this holds, and G is an n -vertex planar graph containing $f_I(n, C_4)$ induced 4-cycles but also containing a vertex in fewer than $n - 3$ induced 4-cycles, then we can delete this vertex to obtain an $(n - 1)$ -vertex planar graph containing more than $f_I(n - 1, C_4)$ induced 4-cycles, which is a contradiction. This completes the proof of the theorem. \square

3 Proof of Theorem 2

3.1 Preliminaries to the proof of Theorem 2

We start with the following definition.

Definition 2. Two vertices in a graph G are *principal neighbours* if they are adjacent and there is an induced 5-cycle in G containing both of them.

We will use the following notation and result adapted from [7]. Let v be a vertex of a planar graph G with distinct neighbours u and w . Let $X_{uvw}^0 = N(u) \setminus (N(w) \cup \{w\})$ and let $Y_{uvw}^0 = N(w) \setminus (N(u) \cup \{u\})$. Then let X_{uvw} be the set of vertices in X_{uvw}^0 which have a neighbour in Y_{uvw}^0 , and similarly let Y_{uvw} be the set of vertices in Y_{uvw}^0 which have a neighbour in X_{uvw}^0 . The following important lemma is a small adaptation of Lemma 1 in [7]. We include the proof here since it is short, provides intuition, and illustrates a method we will use repeatedly.

Lemma 7 ([7]). *Let G be a planar graph with $v \in V(G)$ and $u, w \in N(v)$ such that $u \neq w$. Suppose that there is an induced 5-cycle in G containing the path uvw . Define the sets X_{uvw} and Y_{uvw} as above. Then G' , the bipartite subgraph of G induced by vertex classes X_{uvw} and Y_{uvw} , is a non-empty forest. Moreover the number of induced 5-cycles in G containing the path uvw is at most $|E(G')|$, so in particular, there are at most $|X_{uvw}| + |Y_{uvw}| - 1$ such cycles.*

Proof. Certainly any induced 5-cycle containing the path uvw contains an edge of G' , and each of these edges is in at most one such cycle. Since there is an induced 5-cycle in G containing the path uvw , the graph G' is non-empty. Hence it's enough to show

that G' is acyclic. Indeed, suppose G' contains a cycle $x_1y_1x_2 \dots x_ky_k$ for some $k \geq 2$, $x_i \in X_{uvw}$, and $y_i \in Y_{uvw}$. Then G contains a subdivision of $K_{3,3}$ with vertex classes $\{u, y_1, y_2\}$ and $\{w, x_1, x_2\}$ which is impossible since G is planar. \square

Our second preliminary lemma says that every drawing of a principal graph of the required form has a particular structure.

Lemma 8. *Let $n \geq 19$, and let G be an n -vertex principal graph of the required form. Given a drawing of G and a labelling of the vertices of G consistent with Definition 1, we may assume that the boundaries of the faces in the drawing consist of a fixed set of cycles, namely $u_1u_2u_3$, $u_1a_iwa_{i+1}$ for $1 \leq i \leq |A| - 1$, $u_2b_iwb_{i+1}$ for $1 \leq i \leq |B| - 1$, $u_3c_iwc_{i+1}$ for $1 \leq i \leq |C| - 1$, $u_1a_{|A|}wz_2$, $u_1z_1z_2$, $z_1z_2z_3$, z_2z_3w , $u_1u_2z_1$, $u_2z_1z_4$, $z_1z_3z_4$, z_3z_4w , $u_2z_4wb_1$, $u_2b_{|B|}wz_5$, $u_3z_5wc_1$, $u_2u_3z_5$, $u_3c_{|C|}wz_6$, $u_1z_6wa_1$, and $u_1u_3z_6$.*

Proof. Consider the drawing of $G - Z$ induced by the drawing of G . It is straightforward to see that we can relabel the vertices within each of A , B , and C such the faces of this drawing are bounded by the cycles $u_1u_2u_3$, $u_1a_iwa_{i+1}$ for $1 \leq i \leq |A| - 1$, $u_2b_iwb_{i+1}$ for $1 \leq i \leq |B| - 1$, $u_3c_iwc_{i+1}$ for $1 \leq i \leq |C| - 1$, $u_1a_{|A|}wb_1u_2$, $u_2b_{|B|}wc_1u_3$, and $u_3c_{|C|}wa_1u_1$.

It is clear which of these faces contains each of the vertices in Z in the drawing of G . One by one, add the vertices in Z (and their edges to the current graph) back to the drawing of $G - Z$ in the order z_1, \dots, z_6 , keeping track of the faces of the drawing and their boundaries at each step. At each step there is a unique face to which the next vertex can be added, and the cycles forming the boundaries of the faces in the resulting drawing are fixed. After adding z_6 , the cycles forming the boundaries of the faces are those in the list in the statement of the lemma. Clearly this new labeling of the vertices is still consistent with Definition 1. \square

Finally, we count the induced 5-cycles in a principal graph of the required form.

Lemma 9. *Let $n \geq 19$. Every principal n -vertex graph of the required form contains exactly $\frac{1}{3}(n^2 - 8n + 22)$ induced 5-cycles if $n \equiv 1 \pmod{3}$, and exactly $\frac{1}{3}(n^2 - 8n + 21)$ otherwise. Moreover, every n -vertex graph of the required form contains at least as many induced 5-cycles as a principal n -vertex graph of the required form.*

Proof. Let G be a principal n -vertex graph of the required form, and label its vertices as in Definition 1. Let $a \in A$, then the only neighbours of a are u_1 and w . Define $X = X_{u_1aw}$ and $Y = Y_{u_1aw}$ as above, then the number of induced 5-cycles in G containing a is equal to the number of edges between these two sets. We see that $X = \{u_2, u_3, z_1\}$ and $Y = B \cup C \cup \{z_3, z_4, z_5\}$, and the number of edges between these sets is $|B| + |C| + 5$. None of these 5-cycles contain another vertex in A , so there are $|A|(|B| + |C| + 5)$ induced 5-cycles containing a vertex in A .

Now consider $G - A$. By the same method, we see that each vertex in B is in $|C| + 5$ induced 5-cycles in $G - A$, and none of these use another vertex in B , so there are $|B|(|C| + 5)$ induced 5-cycles in $G - A$ containing a vertex in B . Applying the method once more to the graph $G - (A \cup B)$ and a vertex in C , we find that there are $2|C|$ induced 5-cycles in that graph containing a vertex in C .

It remains to count the number of induced 5-cycles in $G[\{u_1, u_2, u_3, w\} \cup Z]$. Let Γ be an induced 5-cycle in this graph and suppose that it does not contain w . Then the only available neighbours of z_5 and z_6 are u_2 and u_3 , and u_1 and u_3 respectively, but both these pairs are adjacent, so neither z_5 nor z_6 are in Γ . Then similarly u_3 is not in Γ . Now z_1 is adjacent to all the remaining vertices, so it too is not in Γ . Thus Γ is the induced 5-cycle $u_1z_2z_3z_4u_2$.

Now suppose Γ contains w and z_5 . There are exactly five such cycles (two containing the path $wz_5u_2u_1$, two containing $wz_5u_2z_1$, one containing $wz_5u_3u_1$, and none containing any other path of length 4 extending wz_5 beyond z_5). By a similar count, there are exactly four induced 5-cycles containing w and z_6 but not z_5 . If Γ contains w but not z_5 or z_6 , then it must contain z_2 and z_4 , and we see the only such cycle is $u_1z_2wz_4u_2$.

So in total there are exactly $|A|\frac{1}{3}(n^2 - 8n + 21) + |B| + |A||C| + |B||C| + 5(|A| + |B|) + 2|C| + 11$ induced 5-cycles in G , which is $\frac{1}{3}(n^2 - 8n + 22)$ if $n \equiv 1 \pmod{3}$ and $\frac{1}{3}(n^2 - 8n + 21)$ otherwise. Finally, we see from this count that no induced 5-cycle in a principal graph of the required form contains two vertices which are the endpoints of an optional edge. Hence every n -vertex graph of the required form contains at least as many induced 5-cycles as the principal n -vertex graph of the required form that it contains. \square

3.2 Two key lemmas in the proof of Theorem 2

The following two lemmas are the two key steps in the proof of Theorem 2.

Lemma 10. *Let n be large, and suppose that G is an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles. Suppose also that every vertex of G is in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles. Then it contains distinct vertices u_1, u_2, u_3, a, b, c , and w , such that*

1. u_1, u_2 , and u_3 are all adjacent to one another but none are adjacent to w , and
2. the principal neighbours of a, b , and c are exactly u_1 and w, u_2 and w , and u_3 and w respectively.

Lemma 11. *Let $n \geq 19$, and suppose that G is an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles. Suppose also that it contains distinct vertices u_1, u_2, u_3, a, b, c , and w satisfying the conditions in Lemma 10. Then there exists a vertex of G in at most $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles. If moreover $n \equiv 1 \pmod{3}$ and every vertex of G is in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2 = \frac{2n-8}{3}$ induced 5-cycles, then G is of the required form and contains exactly $\frac{1}{3}(n^2 - 8n + 22)$ induced 5-cycles.*

3.3 Proof of Theorem 2 given the key lemmas

Before proving Lemmas 10 and 11 we use them to prove Theorem 2.

Proof of Theorem 2. Let n be large, and let G be an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles. By Lemmas 10 and 11, some vertex of G is in at most

$\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles. By deleting such a vertex from G we find that $f_I(n, C_5) \leq f_I(n-1, C_5) + \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ for large n .

Let $c(n)$ denote the number of induced 5-cycles in an n -vertex principal graph of the required form. By Lemma 9 this is well-defined and satisfies $c(n) = c(n-1) + \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ for all $n \geq 20$. Suppose that the inequality at the end of the last paragraph is strict for infinitely many values of n . Then for large enough n , we have $c(n) > f_I(n, C_5)$ which is a contradiction since graphs of the required form are planar. Hence for large n , $f_I(n, C_5) = f_I(n-1, C_5) + \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$.

Therefore, for large n , if G is an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles, then every vertex of G is in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles, otherwise we could delete the vertex in the fewest induced 5-cycles to obtain an $(n-1)$ -vertex planar graph containing more than $f_I(n-1, C_5)$ induced 5-cycles. Hence by Lemmas 10 and 11, if n is large with $n \equiv 1 \pmod{3}$, and G is an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles, then G is of the required form. Moreover, for such n we have $f_I(n, C_5) = \frac{1}{3}(n^2 - 8n + 22)$. Hence for large n with $n \equiv 0, 2 \pmod{3}$ we have $f_I(n, C_5) = \frac{1}{3}(n^2 - 8n + 21)$, and so by Lemma 9, for large enough n every n -vertex graph of the required form contains $f_I(n, C_5)$ induced 5-cycles. To complete the proof of the theorem it is sufficient to show that if n is large with $n \equiv 0, 2 \pmod{3}$, and G is an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles, then G is of the required form.

Claim 1. *Let n be large, with $n \equiv 0 \pmod{3}$ or $n \equiv 2 \pmod{3}$, and let G be an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles. Suppose that G contains an $(n-1)$ -vertex graph of the required form, H , as an induced subgraph. Then G is of the required form.*

Proof. Let v be the vertex we delete from G to obtain H . Let H_0 be a principal graph of the required form on $n-1$ vertices contained in H . Fix a drawing of G and consider the induced drawings of H and H_0 . By Lemma 8 we can label the vertices of H according to Definition 1 such that the boundaries of the faces in the drawing of H_0 consist of the cycles listed in that lemma.

The only non-triangular faces in the drawing of H have a boundary consisting of a cycle from the list in Lemma 8 of the form $u_i y w y'$ for some $i \in \{1, 2, 3\}$ and $y, y' \in A \cup B \cup C \cup Z$. We know that v is contained in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles in G so it cannot be contained in a triangular face of H . Let the face of H which contains v be bounded by cycle $u_i y w y'$, with i, y , and y' as above. Both y and y' have degree at most 5 in H so by Lemma 7, since n is large, these cannot be the only principal neighbours of v in G . Hence v is adjacent to u_i and w . Its only other possible neighbours are y and y' , so by the planarity of G it is now clear that if $n \equiv 2 \pmod{3}$, then G is of the required form.

If $n \equiv 0 \pmod{3}$, then there is some $i \in \{1, 2, 3\}$ for which this does not imply that G is of the required form. If $n \equiv 0 \pmod{3}$, then by the proof of Lemma 9,

one of the vertex classes A , B , and C in H is such that all the vertices in that class are in exactly $\left\lfloor \frac{2(n-2)}{3} \right\rfloor - 2 = \frac{2n}{3} - 4$ induced 5-cycles in H . This is strictly less than $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2 = \frac{2n}{3} - 3$ and hence each vertex in that class must be in an induced 5-cycle containing v in G . Suppose that A is the class to which this applies (similar arguments hold for B and C). Then it is enough to show that $i \neq 1$ in the argument above.

Suppose for a contradiction that v is adjacent to u_1 and w in G . Vertex a_3 is in an induced 5-cycle containing v in G , and this 5-cycle cannot contain both u_1 and w , so there is a path of length at most 3 from a_3 to v which avoids u_1 and w . Using the list of cycles forming the boundaries of the faces in the drawing of H_0 , we can deduce that the only vertices to which there is a path in H of length at most 2 which avoids u_1 and w are a_1, a_2, a_3, a_4 , and a_5 . Thus v is adjacent to one of these five vertices in G . Similarly, since a_9 is in an induced 5-cycle containing v in G , v is adjacent to one of a_7, a_8, a_9, a_{10} , and a_{11} . However using the list of the boundaries of the faces in the drawing of H_0 again, we see that this is impossible. Hence $i \neq 1$ as required, and if $n \equiv 0 \pmod{3}$, then G is of the required form. \square

To conclude, let n be large with $n \equiv 2 \pmod{3}$, and let G be an n -vertex planar graph containing $f_I(n, C_5)$ induced 5-cycles. Then there is a vertex in exactly $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles in G , and deleting this vertex gives an $(n-1)$ -vertex planar graph containing $f_I(n-1, C_5)$ induced 5-cycles. Since $n-1 \equiv 1 \pmod{3}$, this graph is of the required form. Hence by the claim, G is of the required form. Repeating this argument, we can extend this to $n \equiv 0 \pmod{3}$. This completes the proof of the theorem. \square

4 Proof of Lemma 10

Following the authors of [7], we say that in a drawing of a planar graph G , an *empty* $K_{2,7}$ is a $K_{2,7}$ subgraph of G with parts $\{a_1, a_2\}$ and $\{b_1, \dots, b_7\}$, with b_1, \dots, b_7 labelled in a natural order in the drawing, such that in the drawing of G the bounded region with boundary $a_1b_1a_2b_7$ contains exactly the vertices b_2, \dots, b_6 . Empty $K_{2,7}$'s will be useful in the proof of Lemma 10 since, with notation as above, a_1 and a_2 are the only principal neighbours of b_4 in the graph. Indeed, the only other possible neighbours of b_4 are b_3 and b_5 , but there is no path of length 3 from b_3 to b_5 avoiding a_1, a_2 , and b_4 , so no induced 5-cycle contains the path $b_3b_4b_5$. Also, no induced 5-cycle contains the path $a_i b_4 b_j$ for $i \in \{1, 2\}$ and $j \in \{3, 5\}$ since a_i and b_j are neighbours. Similarly, a_1 and a_2 are the only principal neighbours of b_2, b_3, b_5 , and b_6 .

The proof of Lemma 10 will use the following three results, proved in [7] as Lemma 2, Lemma 4, and Corollary 1 respectively (a small correction to their Corollary 1 was given in an updated version of their paper, see [9]).

Lemma 12 ([7]). *Let n be large and let G be an n -vertex plane graph in which every vertex is in more than $\frac{11n}{20}$ induced 5-cycles. Then G contains an empty $K_{2,7}$.*

Lemma 13 ([7]). *Let n be large and let G be an n -vertex plane graph in which every vertex is in more than $\frac{11n}{20}$ induced 5-cycles. Let u and w be distinct vertices of G , and let v_1, \dots, v_6 be some of their common neighbours, labelled in a natural order in the drawing of G . Suppose that the interior of the bounded region with boundary formed of the cycle uv_3wv_4 contains no common neighbours of u and w . Then if this region contains a vertex, then it contains at least $n^{1/5}$ vertices.*

Lemma 14 ([7], [9]). *Let n be large and let G be an n -vertex plane graph in which every vertex is in more than $\frac{11n}{20}$ induced 5-cycles. If u and w are distinct vertices of G with $|N(u) \cap N(w)| \geq 7n^{4/5}$, then G contains an empty $K_{2,7}$ whose part of size 2 is $\{u, w\}$.*

We now prove Lemma 10. The proof is a very slight adaptation of that of Lemmas 5 and 6 from [7], but we repeat it here for completeness.

Proof of Lemma 10. Fix a drawing of G . By Lemma 12 this drawing contains an empty $K_{2,7}$. Let u and w be the vertices in the part of size 2, and let v be the ‘central’ vertex of the seven in the other part. Then u and w are the only principal neighbours of v , so in particular u and w are not adjacent. Let $X = X_{uvw}$ and $Y = Y_{uvw}$ be defined as in Section 3.1. Then by Lemma 7 we have $|X| + |Y| \geq \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 1$. Let G' be the induced bipartite graph between X and Y .

Claim 2. *No vertex in $V(G) \setminus (Y \cup \{u\})$ is adjacent to more than two vertices in X , and no vertex in $V(G) \setminus (X \cup \{w\})$ is adjacent to more than two vertices in Y .*

Proof. Suppose for a contradiction that $a \in V(G) \setminus (Y \cup \{u\})$ is adjacent to three vertices in X , say x_1, x_2 , and x_3 . By the definition of X , $a \neq w$ and $x_1, x_2, x_3 \in N(u)$. Moreover, there is a path of length 2 to each of x_1, x_2 , and x_3 from w , where the middle vertex in the path is in Y . Hence there exists a vertex $b \in Y \cup \{w\}$ such that G contains a subdivision of $K_{3,3}$ with parts $\{x_1, x_2, x_3\}$ and $\{u, a, b\}$, contradicting Kuratowski’s theorem. This proves the first statement, and the second is similar. \square

Claim 3. *The maximum degree of G' is at least $n^{5/6}$.*

Proof. Suppose otherwise. We will start by showing that u and w have a common neighbourhood of size at least $\frac{1}{3}n + o(n)$. Let $x \in X$. By assumption, x is in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles in G . If an induced 5-cycle contains u, x , and w , then since neither u nor x is adjacent to w , and by the definition of Y , the 5-cycle must be of the form $uxywa$ for some $y \in Y$ and $a \in N(u) \cap N(w)$. Hence the number of such 5-cycles is at most $d_{G'}(x)|N(u) \cap N(w)|$.

If an induced 5-cycle contains x but not both u and w , then each vertex in the cycle which is not adjacent to x is either w or has a path of length at most 3 to x which avoids both u and w . For a vertex $z \in X \cup Y$, let $T(z)$ be the number of vertices of G from which there is a path to z of length at most 3 which avoids both u and w . Then by Lemma 7 there are at most $\binom{d_G(x)}{2}T(x)$ induced 5-cycles in G containing x but not both u and w .

Similar arguments hold if we replace x with a vertex in Y , so for all $z \in X \cup Y$ we have

$$\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2 \leq d_{G'}(z)|N(u) \cap N(w)| + \binom{d_G(z)}{2} T(z). \quad (2)$$

To show that $|N(u) \cap N(w)|$ is ‘large’ we will therefore aim to show that there exists $z \in X \cup Y$ such that $d_{G'}(z)$, $d_G(z)$, and $T(z)$ are ‘small’.

First, recall from the proof of Lemma 7 that G' is planar and hence has at most $|X| + |Y| - 1$ edges. So if l_1 is the number of vertices of degree at least 3 in G' , then $3l_1 \leq 2(|X| + |Y| - 1)$, which gives $|X| + |Y| - l_1 \geq \frac{1}{3}(|X| + |Y| + 2) \geq \frac{1}{3} \left(\left\lfloor \frac{2(n-1)}{3} \right\rfloor + 1 \right) \geq \frac{n}{5}$. Hence there are at least $\frac{n}{5}$ vertices in G' with degree at most 2.

Next, since G is planar it has at most $3n$ edges. If l_2 is the number of vertices of degree at least 61 in G , then $61l_2 \leq 6n$, so $l_2 \leq \frac{n}{10}$. Thus there are at least $\frac{9n}{10}$ vertices in G with degree at most 60.

Finally we turn to $T(z)$. Take a maximal matching in G' between X and Y . Label the vertices in this matching as x_1, \dots, x_k and y_1, \dots, y_k where $x_i \in X$ and $y_i \in Y$ for all i , where edge $x_i y_i$ is in the matching for all i , and where the vertices x_1, \dots, x_k are arranged in that order around u in the drawing of G . This matching, along with the edges $u x_i$ and $w y_i$ for each i split the plane into k regions R_1, \dots, R_k , where R_i has boundary $u x_i y_i w y_{i+1} x_{i+1}$ for each i , with the addition in the subscripts taken modulo k . Since the matching is maximal, every vertex in $X \cup Y$ in the interior of region R_i is adjacent to one of the vertices in $X \cup Y$ on its boundary. We have assumed that the maximum degree of G' is less than $n^{5/6}$, so the total number of vertices in $X \cup Y$ in each region R_i (including the vertices on its boundary) is at most $4n^{5/6}$.

Let q be any vertex of G which is not u or w . Then there is some $1 \leq i \leq k$ such that q is in R_i (including possibly on its boundary). If z is a vertex in $X \cup Y$ such that there is a path in G of length at most 3 from q to z avoiding u and w , then z is in $R_{i-4} \cup \dots \cup R_{i+4}$ (including on its boundary) where again the addition in the subscripts is modulo k . Hence there are at most $36n^{5/6}$ vertices z for which this can hold, and at most $36n^{11/6}$ pairs (q, z) for which this holds. Let l_3 be the number of vertices in $X \cup Y$ with $T(z) \geq 1000n^{5/6}$. Then there are at least $1000l_3n^{5/6}$ pairs (q, z) as above, so $1000l_3n^{5/6} \leq 36n^{11/6}$. Rearranging, we find $l_3 \leq \frac{36n}{1000} \leq \frac{n}{20}$.

Combining these three calculations, there are at least $\frac{n}{5}$ vertices $z \in X \cup Y$ with $d_{G'}(z) \leq 2$, of which at most $\frac{n}{10}$ have degree greater than 60 in G and at most $\frac{n}{20}$ have $T(z) \geq 1000n^{5/6}$. So there exists $z \in X \cup Y$ with $d_{G'}(z) \leq 2$, $d_G(z) \leq 60$, and $T(z) < 1000n^{5/6}$. Hence by (2) we have

$$|N(u) \cap N(w)| \geq \frac{1}{2} \left(\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2 - \binom{60}{2} \cdot 1000n^{5/6} \right) = \frac{1}{3}n + o(n).$$

To complete the proof of the claim we will show that G contains at most $\frac{2}{9}n^2 + o(n^2)$ induced 5-cycles. By Lemma 9, an n -vertex principal graph of the required form contains $\frac{1}{3}n^2 + O(n)$ induced 5-cycles, so since n is large this contradicts G containing $f_I(n, C_5)$ induced 5-cycles.

Label the vertices in $N(u) \cap N(w)$ in the order they're arranged around u as v_1, \dots, v_t . The drawing of the bipartite graph with parts $\{u, w\}$ and $\{v_1, \dots, v_t\}$ contained in the drawing of G splits the plane into t regions R_1, \dots, R_t , where R_i has boundary uv_iwv_{i+1} , with the addition in the subscript taken modulo t . By Lemma 13, if any of the interiors of the regions R_3, \dots, R_{t-3} contain a vertex, then they contain at least $n^{1/5}$ vertices, so at most $n^{4/5}$ of these interiors contain a vertex.

Thus for all but $o(n)$ values of i , none of the regions $R_{i-3}, R_{i-2}, \dots, R_{i+2}$ have any vertices in their interior, and hence the only principal neighbours of v_i are u and w . This implies that v_i is in at most $|X| + |Y| - 1 \leq n - |N(u) \cap N(w)| \leq \frac{2}{3}n + o(n)$ induced 5-cycles in G . Hence deleting all $\frac{1}{3}n + o(n)$ of these vertices v_i removes at most $\frac{2}{9}n^2 + o(n^2)$ induced 5-cycles.

Let H be the graph which remains after these vertices have been deleted. Then H contains $\frac{2}{3}n + o(n)$ vertices, and it's enough to show that it contains $o(n^2)$ induced 5-cycles. First, let $S = V(H) \setminus (X \cup Y \cup \{u, w\})$. Then since $|X| + |Y| \geq \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 1$, S has size $o(n)$. By Claim 2, no vertex in S is adjacent to more than six vertices in $X \cup Y \cup \{u, w\}$. Since $H[S]$ is planar, there is therefore a vertex in S of degree at most 11 in H . By Lemma 7 this vertex is in at most $\binom{11}{2}n$ induced 5-cycles in H . Hence removing the vertices in S from H one by one, at each stage removing the vertex with the smallest degree, we find that deleting S from H removes only $o(n^2)$ induced 5-cycles.

It remains to show that $H' = G[X \cup Y \cup \{u, w\}]$ contains only $o(n^2)$ induced 5-cycles. Let Γ be an induced 5-cycle in H' , and suppose it does not contain u or w . If Γ contains a path that goes from X to Y , then back X , then back to Y , then in a similar way to in the proof of Lemma 7, this implies the existence of a subdivision of $K_{3,3}$ in G . So there are no induced 5-cycles of this form.

Suppose instead that Γ is of the form $x_1x_2x_3y_1y_2$ where $x_1, x_2, x_3 \in X$ and $y_1, y_2 \in Y$. Suppose we are given x_2 . Then by Claim 2, x_2 has at most two neighbours in X , so there are two choices for x_1 , and then x_3 is determined. By assumption each vertex in X has at most $n^{5/6}$ neighbours in Y , so for each choice of x_1 there are at most $n^{5/6}$ choices for y_2 . Finally, each such y_2 has at most two neighbours in Y . Hence there are only $o(n^2)$ such cycles Γ . Similarly, there are $o(n^2)$ induced 5-cycles with two vertices in X and three in Y .

The final step in the proof of the claim will be to show that there are only $o(n^2)$ induced 5-cycles in H' containing u or w . We do these by deleting the vertices in $X \cup Y$ one by one in such an order that at each stage the vertex we delete is in $o(n)$ induced 5-cycles containing u or w . The vertex we delete at each step is a vertex in what remains of $X \cup Y$ which has the lowest degree in what remains of G' . Since G' is a forest, this vertex has degree at most 1 in G' .

Consider a particular step in the deletion process where without loss of generality we are deleting $x \in X$. If Γ is an induced 5-cycle in H' containing x and one of u and w , then it must be of the form $uxyy'x'$ or $wyxx'y'$ for $x' \in X$ and $y, y' \in Y$. If x has degree 0 in what remains of G' , then clearly there are no such cycles. Otherwise let y be the unique neighbour of x in what remains of Y . By Claim 2, y has at most two neighbours in Y , and by assumption each of these has at most $n^{5/6}$ neighbours in X . Similarly, x has

at most two neighbours in X and each of these has at most $n^{5/6}$ neighbours in Y . Thus x is in $o(n)$ induced 5-cycles containing u or w in the current graph, as required. \square

Relabel v and u as a and u_1 respectively. By Claim 3, without loss of generality there is a vertex $u_2 \in X$ with at least $n^{5/6}$ neighbours in Y . So the complete bipartite graph with parts $\{u_2, w\}$ and $N(u_2) \cap N(w)$ contains an empty $K_{2,7}$ by Lemma 14. Let $b_1, b_2,$ and b_3 be the central three vertices of the part of size 7, then u_2 and w are the only principal neighbours of each of these vertices. None of $b_1, b_2,$ and b_3 are equal to u_1 , since u_1 and w are not neighbours, and moreover at most two of these vertices are adjacent to u_1 , otherwise G has a $K_{3,3}$ subgraph. Let b be one of the three which is not adjacent to u_1 , then $b \in Y$ by the definition of Y .

Let $X' = X_{u_2bw}$ and $Y' = Y_{u_2bw}$. Then by Lemma 7, $|X'| + |Y'| \geq \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 1$, so $|(X \cup Y) \cap (X' \cup Y')| \geq \frac{1}{4}n$. Clearly $X \cap Y'$ and $X' \cap Y$ are empty, so $|X \cap X'| + |Y \cap Y'| \geq \frac{1}{4}n$. If $z \in X \cap X'$, then z is a vertex in X which is adjacent to $u_2 \in X$, so $|X \cap X'| \leq 2$ by Claim 2. Thus $|Y \cap Y'| \geq \frac{1}{5}n$.

Let $z_1, z_2,$ and z_3 be distinct vertices in $Y \cap Y'$. Let $t_1, t_2,$ and t_3 be (not necessarily distinct) vertices in X' such that z_i and t_i are neighbours for each $i = 1, 2, 3$. Without loss of generality, the cycle $u_2 t_2 z_2 w b$ splits the plane into two regions, one of which contains z_1 and the other of which contains z_3 . Again without loss of generality, u_1 is in the same region as z_1 . Since $z_3 \in Y$, there is a path of length 2 from u_1 to z_3 , so u_1 and z_3 have a common neighbour among the vertices in the cycle $u_2 t_2 z_2 w b$. However, $u_2 \notin N(z_3)$ since $z_3 \in Y'$, $z_2 \notin N(u_1)$ since $z_2 \in Y$, $w \notin N(u_1)$ since there is an induced 5-cycle in G containing the path $u_1 a w$, and $b \notin N(u_1)$ since $b \in Y$. Hence $t_2 \in N(u_1)$, and consequently $t_2 \in X$.

Therefore all but at most two vertices in $Y \cap Y'$ have a neighbour in $X \cap X'$, but we know there are at most two vertices in $X \cap X'$, so there exists one, say u_3 , with at least $\frac{1}{12}n$ neighbours in $Y \cap Y'$. Then u_3 and w have a common neighbourhood of size at least $\frac{1}{12}n$ in G , so by Lemma 14 G contains an empty $K_{2,7}$ whose part of size 2 is $\{u_3, w\}$. Let c be the central vertex in the part of size seven. Since $u_2 \in X$ and $u_3 \in X \cap X'$, we have that $u_1, u_2,$ and u_3 are all mutually adjacent and none are adjacent to w . The principal neighbours of $a, b,$ and c are exactly u_1 and w, u_2 and $w,$ and u_3 and w respectively, and thus the lemma is proved. \square

5 Proof of Lemma 11

We now prove Lemma 11.

Proof of Lemma 11. As in the statement of the lemma, let $n \geq 19$, and let G be an n -vertex planar graph containing $f_1(n, C_5)$ induced 5-cycles. Suppose that G contains distinct vertices $u_1, u_2, u_3, a, b, c,$ and w satisfying the conditions in the statement of Lemma 10. Let A be the set of vertices of G whose principal neighbours are exactly u_1 and w . Let B and C be the corresponding sets for u_2 and $w,$ and u_3 and w respectively. Note that $a \in A, b \in B,$ and $c \in C$ but $u_1, u_2, u_3, w \notin A \cup B \cup C$.

Define H to be the graph obtained by deleting from G any edges with one endpoint in A and the other endpoint not in $\{u_1, w\}$, and also any analogous edges for B and C . Then H is a planar n -vertex graph which, by the definitions of A , B , and C , has the same set of induced 5-cycles as G .

Let $Z = V(H) \setminus (A \cup B \cup C \cup \{u_1, u_2, u_3, w\})$. Then the graph $H - Z$ consists of the triangle $u_1u_2u_3$ and the three complete bipartite graphs with parts $\{u_1, w\}$ and A , $\{u_2, w\}$ and B , and $\{u_3, w\}$ and C (see Figure 3). It is straightforward to see that a is in exactly $|B| + |C|$ induced 5-cycles in H which avoid Z , and analogously for b and c . In fact, we see that there are exactly $|A||B| + |A||C| + |B||C|$ induced 5-cycles in H which avoid Z .

Let k_1, k_2 , and k_3 be the non-negative constants such that a, b , and c are in exactly $|B| + |C| + k_1$, $|A| + |C| + k_2$, and $|A| + |B| + k_3$ induced 5-cycles in H respectively. Let $T = 2(|A| + |B| + |C|) + k_1 + k_2 + k_3$ be the sum of these numbers. Since $|A| + |B| + |C| = n - |Z| - 4$, we have

$$T - 2n + 2|Z| + 8 = k_1 + k_2 + k_3. \quad (3)$$

Vertices a, b , and c are respectively in k_1, k_2 , and k_3 induced 5-cycles in H containing a vertex of Z . If an induced 5-cycle in H contains two vertices from a, b , and c , then it does not contain any vertices in Z , so the number of induced 5-cycles in H which contain a vertex in Z and one of a, b , and c is $k_1 + k_2 + k_3$.

Using a similar approach to that used in the proof of Lemma 8, it is straightforward to see that for any drawing of $H - Z$ there exists a labeling of the vertices of A as $a_1, \dots, a_{|A|}$ and similarly for B and C such that the faces of the drawing have boundaries given by the cycles $u_1u_2u_3$, $u_1a_{|A|}wb_1u_2$, $u_2b_{|B|}wc_1u_3$, $u_3c_{|C|}wa_1u_1$, $u_1a_iwa_{i+1}$ for $1 \leq i \leq |A| - 1$, $u_2b_iwb_{i+1}$ for $1 \leq i \leq |B| - 1$, and $u_3c_iwc_{i+1}$ for $1 \leq i \leq |C| - 1$. Fix a drawing of H , consider the induced drawing of $H - Z$, and label the vertices of A, B , and C as above. We will refer to the faces whose boundaries are the second, third, and fourth of the cycles listed as F_1, F_2 , and F_3 respectively (see Figure 3).

We now consider which faces of the drawing of $H - Z$ contain vertices of Z in the drawing of H . Suppose $|A| \geq 2$, and let $Z_0 \subseteq Z$ be the set of vertices in the face with boundary $u_1a_1wa_2$. Then the only vertices in $V(H) \setminus Z_0$ which could be adjacent to a vertex in Z_0 are u_1 and w . The only neighbours of $a_{|A|}$ in H are u_1 and w , so we can redraw H with no vertices in the face with boundary $u_1a_1wa_2$ by moving the vertices in Z_0 into F_1 . Repeating this process for the other faces of $H - Z$ whose boundaries are 4-cycles, we see that way may assume that all vertices of Z are in F_1, F_2, F_3 , or the triangular face.

If $z \in Z$ is in an induced 5-cycle in H containing one of a, b , and c , then z is at distance 1 or 2 from w . Therefore z is not in the triangular face of $H - Z$, so it is in one of F_1, F_2 , or F_3 . Note that no induced 5-cycle in H containing one of a, b , and c can contain vertices in Z from two of F_1, F_2 , and F_3 . Hence the number of induced 5-cycles in H containing one of a, b , and c and a vertex in Z is the sum over F_1, F_2 , and F_3 of the number of such cycles containing a vertex of Z in that face.

Let $Z_1 \subseteq Z$ be the set of vertices in F_1 . For each $S \subseteq \{u_1, u_2, w\}$, let L_S be the set of vertices in Z_1 whose neighbours among u_1, u_2 , and w are exactly the vertices in S ,




Figure 3: Illustration of the drawing of $H - Z$, with faces F_1 , F_2 , and F_3 labelled

and let $l_S = |L_S|$. The number of induced 5-cycles containing c and a vertex in Z_1 is $2l_{\{u_1, u_2, w\}} + l_{\{u_1, w\}} + l_{\{u_2, w\}}$. The number of induced 5-cycles containing b and a vertex in Z_1 is $l_{\{u_1, w\}}$ plus the number of paths $u_2 z_1 z_2 w$ where $z_1, z_2 \in Z_1$ are such that $z_1 \notin N(w)$ and $z_2 \notin N(u_2)$. The number of such paths is the number of edges between $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$ and $L_{\{w\}} \cup L_{\{u_1, w\}}$. By a similar argument to that used in the proof of Lemma 7, the bipartite graph between these two sets is a forest. Moreover, it is straightforward to use the planarity of H to show that there can be at most one vertex in $L_{\{u_1, w\}}$ which has a neighbour in $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$.

Hence the number of paths $u_2 z_1 z_2 w$ is at most $l_{\{u_2\}} + l_{\{u_1, u_2\}} + l_{\{w\}}$, with equality only if one of the following holds:

- (i) there is a vertex in $L_{\{u_1, w\}}$ which has a neighbour in $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$, every vertex in $L_{\{u_1, u_2\}}$ has a neighbour in $L_{\{w\}} \cup L_{\{u_1, w\}}$, and every vertex in $L_{\{w\}}$ has a neighbour

in $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$, or

$$(ii) \ l_{\{u_2\}} = l_{\{u_1, u_2\}} = l_{\{w\}} = 0.$$

Indeed, if there is no vertex in $L_{\{u_1, w\}}$ with a neighbour in $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$, then since the bipartite graph between $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$ and $L_{\{w\}}$ is a forest it contains at most $l_{\{u_2\}} + l_{\{u_1, u_2\}} + l_{\{w\}}$ edges, with equality if and only if it has no vertices. If instead there is a vertex $p \in L_{\{u_1, w\}}$ which has a neighbour in $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$, then the number of edges in the bipartite graph between $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$ and $L_{\{w\}} \cup L_{\{u_1, w\}}$ is the same as the number of edges between $L_{\{u_2\}} \cup L_{\{u_1, u_2\}}$ and $L_{\{w\}} \cup \{p\}$. Again, this is at most $l_{\{u_2\}} + l_{\{u_1, u_2\}} + l_{\{w\}}$ with equality if and only if the forest is connected, which in particular implies that (i) is satisfied.

So the number of induced 5-cycles containing b and a vertex in Z_1 is at most $l_{\{u_1, w\}} + l_{\{u_2\}} + l_{\{u_1, u_2\}} + l_{\{w\}}$ with equality only if one of (i) or (ii) holds. We can similarly bound the number of induced 5-cycles containing a and a vertex in Z_1 by swapping the roles of u_1 and u_2 . Let the conditions corresponding to (i) and (ii) in this setting be called (i)' and (ii)'. Summing, we find that the total number of induced 5-cycles that contain one of a , b , and c and a vertex in Z_1 is at most $2l_{\{u_1, u_2, w\}} + 2l_{\{u_1, w\}} + 2l_{\{u_2, w\}} + l_{\{u_2\}} + 2l_{\{u_1, u_2\}} + 2l_{\{w\}} + l_{\{u_1\}}$, with equality only if one of each of (i) and (ii), and (i)' and (ii)' hold.

Repeating for F_2 and F_3 , and recalling that no vertices in the triangular face of $H - Z$ are in an induced 5-cycle in H containing any of a , b , and c , we find that there are at most twice as many induced 5-cycles in H containing one of a , b , and c and a vertex in Z as there are vertices in Z . Recall that the number of such cycles is $k_1 + k_2 + k_3$, which is equal to $T - 2n + 2|Z| + 8$ by equation (3). Hence

$$T \leq 2n - 8. \tag{4}$$

Suppose that all the vertices in G , so in particular a , b , and c , are in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 1$ induced 5-cycles. Then this is also true in H , so

$$T \geq 3 \left\lfloor \frac{2(n-1)}{3} \right\rfloor - 3 = \begin{cases} 2n - 6, & \text{if } n \equiv 0 \pmod{3} \\ 2n - 5, & \text{if } n \equiv 1 \pmod{3} \\ 2n - 7, & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

This is a contradiction, so there is a vertex of G in at most $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2$ induced 5-cycles, which proves the first part of the lemma.

We now prove the second part of the lemma, so assume that $n \equiv 1 \pmod{3}$ and every vertex of G is in at least $\left\lfloor \frac{2(n-1)}{3} \right\rfloor - 2 = \frac{2n-8}{3}$ induced 5-cycles. Then the same is true in H , so $T \geq 2n - 8$. So in fact we have $T = 2n - 8$, and each of a , b , and c are in exactly $\frac{2n-8}{3}$ induced 5-cycles. Moreover, the fact that we have equality in (4) implies that there are exactly twice as many induced 5-cycles in H containing one of a , b , and c and a vertex in Z as there are vertices in Z . Hence there are no vertices of Z in the triangular face of $H - Z$, and in F_1 we must have $l_\emptyset = l_{\{u_2\}} = l_{\{u_1\}} = 0$ and one of each of conditions (i) and (ii), and (i)' and (ii)' must hold (and analogously in F_2 and F_3).




Figure 4: The possible configurations of vertices in F_1

We now use these conditions on the vertices in Z_1 to show that $H[Z_1 \cup \{u_1, u_2, w\}]$ must belong to one of three families of graphs. The same will hold for faces F_2 and F_3 by symmetry. The first possibility is that $Z_1 = \emptyset$, in which case we will say that F_1 is of *type 1*. The second possibility is that Z_1 contains a single vertex z which is adjacent to all of u_1 , u_2 , and w , in which case we will say that F_1 is of *type 2*.

The final possibility is that Z_1 contains $m \geq 3$ vertices z_1, \dots, z_m and $H[Z_1 \cup \{u_1, u_2, w\}]$ contains the edges u_1u_2 , u_1z_1 , u_2z_1 , u_1z_2 , and u_2z_m , and the edges z_iz_i and z_iw for all $2 \leq i \leq m$. The remaining edges of $H[Z_1 \cup \{u_1, u_2, w\}]$ form a subset of $\{z_iz_{i+1} : 2 \leq i \leq m-1\}$. In this case we say that F_1 is of *type 3*. The three possibilities for F_1 are illustrated in Figure 4, where red lines indicate edges each of whose presence does not affect whether the face is of that type.

Claim 4. *Face F_1 is of type 1, 2, or 3. By symmetry, the same is true for F_2 and F_3 .*

Proof. We know that $l_\emptyset = l_{\{u_2\}} = l_{\{u_1\}} = 0$ and one of each of conditions (i) and (ii), and (i)' and (ii)' hold. First, if condition (ii) holds, then $l_{\{u_1, u_2\}} = l_{\{w\}} = 0$. So all vertices in Z_1 are adjacent to exactly the vertices in one of $\{u_1, w\}$, $\{u_2, w\}$, and $\{u_1, u_2, w\}$ among u_1 , u_2 , and w . In particular all the vertices are adjacent to w . Suppose there exists $z \in L_{\{u_1, w\}}$. Then since z is not in A , there is some $z' \in Z_1$ which is a principal neighbour of z .

Let Γ be an induced 5-cycle in H containing both z and z' . Then Γ does not contain w since it's a common neighbour of z and z' . If every vertex in Γ is in Z_1 , then all its vertices are adjacent to u_1 or u_2 so without loss of generality, some two non-adjacent vertices in Γ , say p_1 and p_2 , are both adjacent to u_1 . Let q_1 and q_2 be two other distinct non-adjacent vertices of Γ . Then $\{w, p_1, p_2\}$ and $\{u_1, q_1, q_2\}$ form the partite sets of a subdivision of $K_{3,3}$ in G , which is a contradiction.

Thus Γ must contain a vertex which is not in Z_1 , so without loss of generality it contains u_1 . If it doesn't contain u_2 , then the rest of its vertices are in Z_1 , and the two which are not adjacent to u_1 , say p_1 and p_2 , must be adjacent to u_2 . Let q_1 and q_2 be the other two vertices in Γ . Then $\{u_1, w, p_1\}$ and $\{u_2, q_1, q_2\}$ form the partite sets of a subdivision of $K_{3,3}$ in H , giving a contradiction. If Γ does contain u_2 , then since u_1 and

u_2 are neighbours, the other three vertices in Γ are in Z_1 . One of these three must be adjacent to neither u_1 nor u_2 , but no such vertex exists. Hence $l_{\{u_1, w\}} = 0$. Similarly $l_{\{u_2, w\}} = 0$. If $l_{\{u_1, u_2, w\}} = 0$ then F_1 is of type 1. It is straightforward to use the planarity of H to show that $l_{\{u_1, u_2, w\}} \leq 1$, so if $l_{\{u_1, u_2, w\}} \neq 0$, then F_1 is of type 2.

If condition (ii) doesn't hold, then neither does condition (ii)', so conditions (i) and (i)' both hold. In particular,

- (a) there is a vertex in $L_{\{u_1, w\}}$ which has a neighbour in $L_{\{u_1, u_2\}}$,
- (b) there is a vertex in $L_{\{u_2, w\}}$ which has a neighbour in $L_{\{u_1, u_2\}}$,
- (c) every vertex in $L_{\{u_1, u_2\}}$ has a neighbour in $L_{\{w\}} \cup L_{\{u_1, w\}}$, and
- (d) every vertex in $L_{\{w\}}$ has a neighbour in $L_{\{u_1, u_2\}}$.

Condition (a) implies that $l_{\{u_1, u_2\}} \geq 1$, and it is straightforward to deduce from (c) and the planarity of H that $l_{\{u_1, u_2\}} \leq 1$. It is also straightforward to use (a) and the planarity of H to deduce that $l_{\{u_1, u_2, w\}} = 0$. It remains to determine $l_{\{u_1, w\}}$, $l_{\{u_2, w\}}$ and $l_{\{w\}}$.

Let z_1 be the unique vertex in $L_{\{u_1, u_2\}}$ and let z_2 be a vertex in $L_{\{u_1, w\}}$ which is adjacent to z_1 . Let $m = |L_{\{w\}}| + 3$ and let z_m be a vertex in $L_{\{u_2, w\}}$ which is adjacent to z_1 . Consider $H[\{u_1, u_2, w, a_{|A|}, b_1, z_1, z_2, z_m\}]$ with the edge $z_2 z_m$ deleted if present, and the drawing of this induced by our drawing of H . Face F_1 is split into six faces whose boundaries are the cycles in a fixed list. Let E_1 , E_2 , and E_3 respectively be the faces whose boundaries are formed of the cycles $u_1 a_{|A|} w z_2$, $u_2 b_1 w z_m$, and $z_1 z_2 w z_m$ (see Figure 5). Any vertices in $L_{\{u_1, w\}} \setminus \{z_2\}$ must be in E_1 and any vertices in $L_{\{u_2, w\}} \setminus \{z_m\}$ must be in E_2 . By (d), every vertex in $L_{\{w\}}$ is in E_3 .

Suppose there exists some $z \in L_{\{u_1, w\}} \setminus \{z_2\}$. Then since $z \notin A$, z has a principal neighbour other than u_1 and w , say z' . Then z' is either z_2 or is in the interior of E_1 , so in particular $z' \in L_{\{u_1, w\}}$. Let Γ be an induced 5-cycle in G containing z and z' . Then Γ does not contain u_1 or w since these are common neighbours of z and z' . Hence every vertex in Γ is either z_2 or is in the interior of E_1 , so they are all adjacent to u_1 and w . By an argument similar to that used in the proof of Lemma 7, this implies the existence of a subdivision of $K_{3,3}$ in H , which is a contradiction. Therefore $L_{\{u_1, w\}} = \{z_2\}$ and similarly $L_{\{u_2, w\}} = \{z_m\}$.

Consider adding the vertices in $L_{\{w\}}$ and their edges to w and z_1 to the drawing of $H[\{u_1, u_2, w, a_{|A|}, b_1, z_1, z_2, z_m\}]$. We see that we can label them as z_3, \dots, z_{m-1} such that to preserve planarity the only other edges which could be present in F_1 are of the form $z_i z_{i+1}$ for $2 \leq i \leq m-1$. Hence F_1 is of type 3. This completes the proof of the claim. \square

To summarise, we know that:

- there are exactly $|A||B| + |A||C| + |B||C|$ induced 5-cycles in $H - Z$,
- a , b , and c are in $|B| + |C|$, $|A| + |C|$, and $|A| + |B|$ induced 5-cycles in $H - Z$ respectively,




Figure 5: Illustration of the drawing of $H[\{u_1, u_2, w, a_{|A|}, b_1, z_1, z_2, z_m\}]$ with the edge z_2z_m deleted if present

- each of a , b , and c is in exactly $\frac{2n-8}{3}$ induced 5-cycles in H ,
- every vertex of Z is in F_1 , F_2 , or F_3 ,
- each of F_1 , F_2 , and F_3 is of type 1, 2, or 3,
- there are no induced 5-cycles in H containing two vertices in $A \cup B \cup C$ and a vertex in Z , and
- there are no induced 5-cycles in H containing one of a , b , and c and vertices in Z from two of F_1 , F_2 , and F_3 .

For each assignment of types 1, 2, and 3 to faces F_1 , F_2 , and F_3 and for each $x \in \{a, b, c\}$, we now count how many induced 5-cycles there are in H containing x and a vertex in Z . After that, we will count how many induced 5-cycles there are in $H[\{u_1, u_2, u_3, w\} \cup Z]$ for each of these assignments. We will then use the second and third points above to calculate the sizes of A , B , and C in each case. We can then use all of this information to determine the total number of induced 5-cycles in H in each case.

For the first of these steps it is sufficient by symmetry to count the induced 5-cycles containing x and a vertex in F_1 for each $x \in \{a, b, c\}$ and each assignment of a type to F_1 . If F_1 is of type 1, then there are clearly no induced 5-cycles containing a and a vertex in F_1 , and similarly for b and c . If F_1 is of type 2, containing a single vertex z , then $u_1zwc u_3$ and $u_2zwc u_3$ are induced 5-cycles in H . We know there are at most twice as many induced 5-cycles in H containing one of a , b , and c and a vertex in F_1 as there are vertices in F_1 , so there are exactly two induced 5-cycles containing c and z , and none containing a and z or b and z .

Finally, if F_1 is of type 3 with $m \geq 3$ vertices labelled as in the definition of a type 3 face (and Figure 4c), then H contains the induced 5-cycles $u_1z_2wbu_2$, $u_1z_2wcu_3$, $u_2z_mwau_1$, $u_2z_mwcu_3$, $u_1z_1z_iwa$ for $3 \leq i \leq m$, and $u_2z_1z_iwb$ for $2 \leq i \leq m-1$. We have identified $2m$ suitable induced 5-cycles, so as above we know this is all of them. So a and b are each in exactly $m-1$ induced 5-cycles in H containing a vertex in F_1 , and c is exactly in two.

We now determine the number of induced 5-cycles in $H[\{u_1, u_2, u_3, w\} \cup Z]$ in each case. Note that every induced 5-cycle in this graph contains a vertex of Z , and no induced 5-cycle can contain a vertex of Z from each of F_1 , F_2 , and F_3 . Let $Z_1 \subseteq Z$ be the set of vertices in F_1 as before, and let $Z_2 \subseteq Z$ be the set of vertices in F_2 . Then by symmetry it is sufficient to calculate the number of induced 5-cycles in $H[\{u_1, u_2, u_3, w\} \cup Z_1]$ in each case, and the number of induced 5-cycles in $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ which contain vertices from both Z_1 and Z_2 in each case.

We first count the number of induced 5-cycles in $H[\{u_1, u_2, u_3, w\} \cup Z_1]$. Clearly there are none if F_1 is of type 1 or 2. Suppose F_1 is of type 3 with $m \geq 3$ vertices labelled as in the definition of a face of type 3. The graph $H[\{u_1, u_2, u_3, w\} \cup Z_1]$ is illustrated in Figure 6a, where a red line indicates an edge which may or may not be present.


In this graph u_3 is only adjacent to u_1 and u_2 and these are themselves neighbours, so u_3 is not in an induced 5-cycle in this graph. Next, z_1 is adjacent to all the remaining vertices except w , so it also cannot be in an induced 5-cycle in this graph. If an induced 5-cycle contains w , then it contains exactly two of the vertices z_2, \dots, z_m , and these two must not be neighbours. The remaining two vertices must be u_1 and u_2 , and hence the only possible induced 5-cycle containing w is $u_1z_2wz_mu_2$. This induced 5-cycle is realised if and only if $m \geq 4$, or $m = 3$ and z_2 and z_3 are not neighbours.

Now suppose Γ is an induced 5-cycle which does not contain w . Then it must contain at least 3 vertices from z_2, \dots, z_m . Clearly it cannot only contain vertices from z_2, \dots, z_m , so it contains at least one of u_1 and u_2 . Each of these only has one neighbour in $\{z_2, \dots, z_m\}$, so in fact Γ contains both u_1 and u_2 . Hence it also contains z_2 and z_m , and the remaining vertex is a common neighbour of these two among z_3, \dots, z_{m-1} . So for such a cycle to appear we must have $m = 4$, and the edges z_2z_3 and z_3z_4 must be present. This condition is also sufficient for the induced cycle $u_1z_2z_3z_4u_2$ to appear.


For each assignment of types 1, 2, and 3 to faces F_1 and F_2 , we now count the number of induced 5-cycles in $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ which contain vertices from both Z_1 and Z_2 . If either face is of type 1, then clearly there are no such cycles. If both faces are of type 2, then label the vertex in F_1 as z and the vertex in F_2 as z' . The graph $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ is illustrated in Figure 6b. It is straightforward to see (for example by considering whether or not the cycle contains w) that there is exactly one induced 5-cycle of the required form, namely $u_1zwz'u_3$.

Now suppose that F_1 is of type 3 with $m \geq 3$ vertices and F_2 is of type 2. Label the vertices in F_1 as z_1, \dots, z_m in the usual way, and the vertex in F_2 as z . The graph $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ is illustrated in Figure 6c, where a red line indicates an edge which may or may not be present.


Let Γ be a 5-cycle in the graph containing z and one of z_1, \dots, z_m . Then since Γ




(a) $H[\{u_1, u_2, u_3, w\} \cup Z_1]$ if F_1 is of type 3



(b) $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ if F_1 and F_2 are both of type 2



(c) $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ if F_1 is of type 3 and F_2 is of type 2



(d) $H[\{u_1, u_2, u_3, w\} \cup Z_1 \cup Z_2]$ if F_1 and F_2 are both of type 3

Figure 6: Certain subgraphs of H

contains z it must also contain w . Suppose Γ contains z_1 , then since z_1 is not adjacent to w the cycle must also contain a common neighbour of z_1 and w , i.e. one of z_2, \dots, z_m . So Γ contains the path $zwz_i z_1$ for some $2 \leq i \leq m$. The only common neighbour of z and z_1 is u_2 , so Γ is $zwz_i z_1 u_2$ for some $2 \leq i \leq m$. This induced 5-cycle is realised if and only if $i \neq m$, so there are exactly $m - 2$ such 5-cycles.

Now suppose Γ does not contain z_1 . Then it must contain one of z_2, \dots, z_m , and since these are all neighbours of w it must in fact contain exactly one of them. For $3 \leq i \leq m - 1$, z_i has no neighbours outside $\{w, z_1, \dots, z_m\}$, so Γ must contain z_2 or z_m . If it contains z_m , then it also contains u_2 , but this is a neighbour of z . Hence Γ contains the path $zwz_2 u_1$. The only common neighbours of u_1 and z are u_2 and u_3 , and the induced 5-cycle is realised in both cases. So in total there are m induced 5-cycles in the graph containing a vertex from Z_1 and a vertex from Z_2 .

Finally, suppose F_1 and F_2 are both of type 3, containing $m \geq 3$ and $m' \geq 3$ vertices respectively. Label the vertices in F_1 as z_1, \dots, z_m in the usual way, and the vertices in F_2 as $z'_1, \dots, z'_{m'}$ in the analogous way, as illustrated in Figure 6d. As usual, red lines in the figure indicate edges which may or may not be present.

Let Γ be a 5-cycle in the graph containing a vertex from Z_1 and a vertex from Z_2 . Since Γ contains vertices in both F_1 and F_2 , it must contain two vertices on the boundaries of each. Therefore it contains a vertex which is on the boundary of both faces, i.e. w or u_2 . If it does not contain w , then in order to contain two vertices on the boundary of each face it must contain u_1, u_2 , and u_3 which is impossible. Hence Γ contains w .

Suppose Γ contains z_1 . If it does not also contain one of z_2, \dots, z_m , then it contains u_1 and u_2 as these are the only other vertices adjacent to z_1 . But then Γ contains u_1, u_2 , and z_1 , which form a triangle, giving a contradiction. So Γ contains one of z_2, \dots, z_m . It cannot contain more than one of z_2, \dots, z_m since these are all common neighbours of z_1 and w .

Hence Γ contains a path $u_i z_1 z_j w$ for some $i \in \{1, 2\}$ and $j \in \{2, \dots, m\}$. There are no common neighbours of u_1 and w which are not adjacent to z_1 , and the only common neighbour of u_2 and w which is not adjacent to z_1 is z'_2 . Hence if Γ contains z_1 , then it is of the form $u_2 z_1 z_j w z'_2$ for some $j \in \{2, \dots, m\}$. This induced 5-cycle is realised if and only if $j \in \{2, \dots, m - 1\}$, so there are exactly $m - 2$ induced cycles of the required form containing z_1 . Similarly there are exactly $m' - 2$ containing z'_1 .

Now suppose Γ contains neither z_1 nor z'_1 . Then it contains one vertex from z_2, \dots, z_m and one from $z'_2, \dots, z'_{m'}$. All of these vertices are neighbours of w , so in fact Γ must contain exactly one from each list. The remaining two vertices in Γ must be picked from u_1, u_2 , and u_3 . None of z_3, \dots, z_{m-1} or $z'_3, \dots, z'_{m'-1}$ are adjacent to any of these vertices, so the neighbours of w in the cycle must be one of z_2 and z_m and one of z'_2 and $z'_{m'}$. For each such combination except z_m and z'_2 , we see there is a single induced 5-cycle of the required form, and for z_m and z'_2 there are none. So in total there are $m + m' - 1$ induced 5-cycles in the graph containing a vertex in Z_1 and a vertex in Z_2 .

To summarise our findings:

- if F_1 is of type 1, then there are no induced 5-cycles containing a vertex in F_1 ,

- if F_1 is of type 2, then there are no induced 5-cycles containing a or b and a vertex in F_1 , and there are exactly two containing c and a vertex in F_1 ,
- if F_1 is of type 3 with $m \geq 3$ vertices, then for $x \in \{a, b\}$ there are exactly $m - 1$ induced 5-cycles containing x and a vertex in F_1 , and there are exactly two containing c and a vertex in F_1 ,
- $H[\{u_1, u_2, u_3, w\} \cup Z]$ contains exactly
 - one induced 5-cycle per face of type 3 with $m \geq 5$ vertices, four vertices but not both optional edges present, or three vertices with the optional edge not present,
 - two induced 5-cycles per face of type 3 with four vertices and both optional edges present,
 - one induced 5-cycle per pair of faces of type 2,
 - m induced 5-cycles per pair of faces where one is of type 2 and the other is of type 3 with m vertices,
 - $m+m'-1$ induced 5-cycles per pair of faces of type 3, where one has m vertices and the other has m' ,

and no further induced 5-cycles.

For each assignment of types to F_1 , F_2 , and F_3 , we can now calculate the sizes of A , B , and C and thence can determine the total number of induced 5-cycles in H in each case. The details of this process are laid out in Appendix A, where the following claim is proved.

Claim 5. *H is the principal graph of the required form on n vertices.*

Label the vertices of H according to Definition 1. Fix a drawing of G , then by Lemma 8 we may assume that the boundaries of the faces in the induced drawing of H consist of the cycles listed in the statement of that lemma. Consider adding the edges in $E(G) \setminus E(H)$ back to this drawing of H . Clearly no edge is added to a triangular face. The remaining faces' boundaries are cycles of the form $u_i y w y'$ for some $i \in \{1, 2, 3\}$ and $y, y' \in A \cup B \cup C \cup Z$. Hence every edge in $E(G) \setminus E(H)$ is of the form $u_i w$ for some $i \in \{1, 2, 3\}$, or yy' for some $y, y' \in A \cup B \cup C \cup Z$ such that there exists $i \in \{1, 2, 3\}$ such that cycle $u_i y w y'$ appears in the list in Lemma 8. The sets of induced 5-cycles in G and H are the same, and there is an induced 5-cycle in H containing u_i and w for each $i \in \{1, 2, 3\}$, so $u_i w \notin E(G) \setminus E(H)$ for all $i \in \{1, 2, 3\}$. Analysing the list of cycles in Lemma 8, we see that G is of the required form.

By Lemma 9, H contains exactly $\frac{1}{3}(n^2 - 8n + 22)$ induced 5-cycles. We know that G has the same set of induced 5-cycles as H , so G contains exactly $\frac{1}{3}(n^2 - 8n + 22)$ induced 5-cycles too. \square

Acknowledgements

Thank you to Tom Johnston, Emil Powierski, and Jane Tan for helpful discussions. Particular thanks to Tom for finding the graph in Figure 2, and to Emil and Jane for spotting that an argument to show $f_I(n, C_4) = \frac{1}{2}(n^2 - 5n + 6)$ for large n could be extended to show that $K_{2,n-2}$ uniquely achieves the maximum. Thank you also to Alex Scott for support and helpful comments. The author is grateful to the Heilbronn Institute for Mathematical Research for their support.

References

- [1] N. Alon and Y. Caro. On the number of subgraphs of prescribed type of planar graphs with a given number of vertices. In M. Rosenfeld and J. Zaks, editors, *Annals of Discrete Mathematics (20): Convexity and Graph Theory*, volume 87 of *North-Holland Mathematics Studies*, pages 25–36. North-Holland, 1984.
- [2] J. Balogh, P. Hu, B. Lidický and F. Pfender. Maximum density of induced 5-cycle is achieved by an iterated blow-up of 5-cycle. *European Journal of Combinatorics*, **52**:47–58, 2016.
- [3] C. Cox and R. R. Martin. Counting paths, cycles and blow-ups in planar graphs. *arXiv:2101.05911* preprint, 2021.
- [4] C. Cox and R. R. Martin. The maximum number of 10- and 12-cycles in a planar graph. *arXiv:2106.02966* preprint, 2021.
- [5] D. Eppstein. Connectivity, graph minors, and subgraph multiplicity. *Journal of Graph Theory*, **17**(3):409–416, 1993.
- [6] C. Even-Zohar and N. Linial. A note on the inducibility of 4-vertex graphs. *Graphs and Combinatorics*, **31**:1367–1380, 2015.
- [7] D. Ghosh, E. Győri, O. Janzer, A. Paulos, N. Salia and O. Zamora. The maximum number of induced C_5 's in a planar graph. *arXiv:2004.01162v1* preprint, 2020.
- [8] D. Ghosh, E. Győri, O. Janzer, A. Paulos, N. Salia and O. Zamora. The maximum number of induced C_5 's in a planar graph. *arXiv:2004.01162v2* preprint, 2021.
- [9] D. Ghosh, E. Győri, O. Janzer, A. Paulos, N. Salia and O. Zamora. The maximum number of induced C_5 's in a planar graph. *arXiv:2004.01162v3* preprint, 2021.
- [10] D. Ghosh, E. Győri, R. R. Martin, A. Paulos, N. Salia, C. Xiao and O. Zamora. The maximum number of paths of length four in a planar graph. *Discrete Mathematics*, **344**(5):112317, 2021.
- [11] A. Grzesik, E. Győri, A. Paulos, N. Salia, C. Tompkins and O. Zamora. The maximum number of paths of length three in a planar graph. *arXiv:1909.13539* preprint, 2021.

- [12] E. Győri, A. Paulos, N. Salia, C. Tompkins and O. Zamora. The maximum number of pentagons in a planar graph. *arXiv:1909.13532* preprint, 2019.
- [13] E. Győri, A. Paulos, N. Salia, C. Tompkins and O. Zamora. Generalized planar Turán numbers. *arXiv:2002.04579* preprint, 2020.
- [14] S. L. Hakimi and E. F. Schmeichel. On the number of cycles of length k in a maximal planar graph. *Journal of Graph Theory*, **3**(1):69–86, 1979.
- [15] H. Hatami, J. Hirst and S. Norine. The inducibility of blow-up graphs. *Journal of Combinatorial Theory, Series B*, **109**:196–212, 2014.
- [16] T. Huynh, G. Joret and D. R. Wood. Subgraph densities in a surface. *arXiv:2003.13777* preprint, 2021.
- [17] D. Král', S. Norin and J. Volec. A bound on the inducibility of cycles. *Journal of Combinatorial Theory, Series A*, **161**:359–363, 2019.
- [18] C.-H. Liu. Homomorphism counts in robustly sparse graphs. *arXiv:2107.00874* preprint, 2021.
- [19] N. Pippenger and M. C. Golumbic. The inducibility of graphs. *Journal of Combinatorial Theory, Series B*, **19**(3):189–203, 1975.
- [20] D. R. Wood. On the maximum number of cliques in a graph. *Graphs and Combinatorics*, **23**:337–352, 2007.
- [21] N. C. Wormald. On the frequency of 3-connected subgraphs of planar graphs. *Bulletin of the Australian Mathematical Society*, **34**(2):309–317, 1986.
- [22] R. Yuster. On the exact maximum induced density of almost all graphs and their inducibility. *Journal of Combinatorial Theory, Series B*, **136**:81–109, 2019.

A Proof of Claim 5

Let $K_1 = \frac{n}{3} - |A|$, $K_2 = \frac{n}{3} - |B|$, and $K_3 = \frac{n}{3} - |C|$. For each assignment of types to F_1 , F_2 , and F_3 (ignoring symmetric cases), we will assume that H follows this assignment and go through the following process. We know that each of a , b , and c is in exactly $\frac{2n-8}{3}$ induced 5-cycles in H . We also know that a is in exactly $|B| + |C|$ induced 5-cycles which avoid Z , and for each assignment of types to faces we know how many induced 5-cycles there are which contain a and a vertex of Z . Repeating for b and c gives three linear equations which we can solve to find K_1 , K_2 , and K_3 .

Once we know the sizes of A , B , and C , we can use our earlier findings to determine the total number of induced 5-cycles in H in each case. There are $|A||B| + |A||C| + |B||C|$ induced 5-cycles avoiding Z , $|A|(\frac{2n-8}{3} - |B| - |C|) + |B|(\frac{2n-8}{3} - |A| - |C|) + |C|(\frac{2n-8}{3} - |A| - |B|)$ induced 5-cycles containing a vertex of $A \cup B \cup C$ and a vertex of Z , and in each case we know how many induced 5-cycles there are avoiding $A \cup B \cup C$.

Note that by our earlier findings, for a face of type 3 with three vertices the number of induced 5-cycles in H can only increase if the optional edge is not present, so we may assume this is always the case. Similarly, for a face of type 3 with four vertices we may assume that both optional edges are present.

Of the below cases, case 7 with $m = 4$ uniquely gives the most induced 5-cycles, so since H contains $f_I(n, C_5)$ induced 5-cycles we have $f_I(n, C_5) \leq \frac{1}{3}(n^2 - 8n + 22)$ for this n . By Lemma 9, in fact $f_I(n, C_5) = \frac{1}{3}(n^2 - 8n + 22)$ for this n , and hence case 7 with $m = 4$ must occur. In other words, H is the principal graph of the required form on n vertices.

Case 1: F_1, F_2 , and F_3 type 1.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3$, $\frac{2n}{3} - K_1 - K_3$, and $\frac{2n}{3} - K_1 - K_2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_2 = K_3 = \frac{4}{3}$. There are no induced 5-cycles avoiding $A \cup B \cup C$. Hence there are $3(\frac{n-4}{3})^2 = \frac{1}{3}(n^2 - 8n + 16)$ induced 5-cycles in H .

Case 2: F_1 type 2, F_2 and F_3 type 1.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3$, $\frac{2n}{3} - K_1 - K_3$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_2 = \frac{7}{3}$ and $K_3 = \frac{1}{3}$. There are no induced 5-cycles avoiding $A \cup B \cup C$. Hence there are $2(\frac{n-7}{3})(\frac{n-1}{3}) + (\frac{n-7}{3})^2 + 2(\frac{n-1}{3}) = \frac{1}{3}(n^2 - 8n + 19)$ induced 5-cycles in H .

Case 3: F_1 and F_2 type 2, F_3 type 1.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3 + 2$, $\frac{2n}{3} - K_1 - K_3$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_3 = \frac{4}{3}$ and $K_2 = \frac{10}{3}$. There is exactly one induced 5-cycle avoiding $A \cup B \cup C$. Hence there are $2(\frac{n-4}{3})(\frac{n-10}{3}) + (\frac{n-4}{3})^2 + 4(\frac{n-4}{3}) + 1 = \frac{1}{3}(n^2 - 8n + 19)$ induced 5-cycles in H .

Case 4: F_1, F_2 , and F_3 type 2.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3 + 2$, $\frac{2n}{3} - K_1 - K_3 + 2$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_2 = K_3 = \frac{7}{3}$. There are exactly 3 induced 5-cycles avoiding $A \cup B \cup C$. Hence there are $3(\frac{n-7}{3})^2 + 6(\frac{n-7}{3}) + 3 = \frac{1}{3}(n^2 - 8n + 16)$ induced 5-cycles in H .

Case 5: F_1 type 3 with m vertices, F_2 and F_3 type 1.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3 + m - 1$, $\frac{2n}{3} - K_1 - K_3 + m - 1$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_2 = \frac{7}{3}$ and $K_3 = m - \frac{2}{3}$. There is exactly one induced 5-cycle avoiding $A \cup B \cup C$ for $m \neq 4$. Hence there are $2(\frac{n-7}{3})(\frac{n+2}{3} - m) + (\frac{n-7}{3})^2 + 2(m-1)(\frac{n-7}{3}) + 2(\frac{n+2}{3} - m) + 1 = \frac{1}{3}(n^2 - 8n + 28 - 6m)$ induced 5-cycles in H for $m \neq 4$. Since $m \geq 3$, this is at most $\frac{1}{3}(n^2 - 8n + 10)$. For $m = 4$ there is one more induced 5-cycle, so there are $\frac{1}{3}(n^2 - 8n + 7)$ in total.

Case 6: F_1 type 3 with m vertices, F_2 type 2, F_3 type 1.

Vertices a, b , and c are in $\frac{2n}{3} - K_2 - K_3 + m + 1$, $\frac{2n}{3} - K_1 - K_3 + m - 1$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = \frac{4}{3}$, $K_2 = \frac{10}{3}$, and $K_3 = m + \frac{1}{3}$. There are exactly $m + 1$ induced 5-cycles avoiding $A \cup B \cup C$ for $m \neq 4$. Hence there are $(\frac{n-4}{3})(\frac{n-10}{3}) + (\frac{n-4}{3})(\frac{n-1}{3} - m) + (\frac{n-10}{3})(\frac{n-1}{3} - m) + (m+1)(\frac{n-4}{3}) + (m-1)(\frac{n-10}{3}) +$

$2(\frac{n-1}{3} - m) + m + 1 = \frac{1}{3}(n^2 - 8n + 25 - 3m)$ induced 5-cycles in H for $m \neq 4$. Since $m \geq 3$, this is at most $\frac{1}{3}(n^2 - 8n + 16)$. For $m = 4$ there is one more induced 5-cycle, so there are $\frac{1}{3}(n^2 - 8n + 16)$ in total.

Case 7: F_1 type 3 with m vertices, F_2 and F_3 type 2.

Vertices a , b , and c are in $\frac{2n}{3} - K_2 - K_3 + m + 1$, $\frac{2n}{3} - K_1 - K_3 + m + 1$, and $\frac{2n}{3} - K_1 - K_2 + 2$ induced 5-cycles respectively. Solving, we obtain $K_1 = K_2 = \frac{7}{3}$ and $K_3 = m + \frac{4}{3}$. There are exactly $2m + 2$ induced 5-cycles avoiding $A \cup B \cup C$ for $m \neq 4$. Hence there are $2(\frac{n-7}{3})(\frac{n-4}{3} - m) + (\frac{n-7}{3})^2 + 2(m+1)(\frac{n-7}{3}) + 2(\frac{n-4}{3} - m) + 2m + 2 = \frac{1}{3}(n^2 - 8n + 19)$ induced 5-cycles in H for $m \neq 4$. For $m = 4$ there is one more, so there are $\frac{1}{3}(n^2 - 8n + 22)$.

Case 8: F_1 type 3 with m vertices, F_2 type 3 with m' vertices, F_3 type 1.

Vertices a , b , and c are in $\frac{2n}{3} - K_2 - K_3 + m + 1$, $\frac{2n}{3} - K_1 - K_3 + m + m' - 2$, and $\frac{2n}{3} - K_1 - K_2 + m' + 1$ induced 5-cycles respectively. Solving, we obtain $K_1 = m' + \frac{1}{3}$, $K_2 = \frac{10}{3}$, and $K_3 = m + \frac{1}{3}$. There are exactly $m + m' + 1$ induced 5-cycles avoiding $A \cup B \cup C$ if $m, m' \neq 4$. Hence there are $(\frac{n-1}{3} - m')(\frac{n-10}{3}) + (\frac{n-1}{3} - m')(\frac{n-1}{3} - m) + (\frac{n-10}{3})(\frac{n-1}{3} - m) + (m+1)(\frac{n-1}{3} - m') + (m+m'-2)(\frac{n-10}{3}) + (m'+1)(\frac{n-1}{3} - m) + m + m' + 1 = \frac{1}{3}(n^2 - 8n + 28 - 3mm')$ induced 5-cycles in H for $m, m' \neq 4$. Since $m, m' \geq 3$, this is at most $\frac{1}{3}(n^2 - 8n + 1)$. If one or both of m and m' is 4, then this increases by at most 2, to give a total of at most $\frac{1}{3}(n^2 - 8n + 7)$.

Case 9: F_1 type 3 with m vertices, F_2 type 3 with m' vertices, F_3 type 2.

Vertices a , b , and c are in $\frac{2n}{3} - K_2 - K_3 + m + 1$, $\frac{2n}{3} - K_1 - K_3 + m + m'$, and $\frac{2n}{3} - K_1 - K_2 + m' + 1$ induced 5-cycles respectively. Solving, we obtain $K_1 = m' + \frac{4}{3}$, $K_2 = \frac{7}{3}$, and $K_3 = m + \frac{4}{3}$. There are exactly $2m + 2m' + 1$ induced 5-cycles avoiding $A \cup B \cup C$ if $m, m' \neq 4$. Hence there are $(\frac{n-4}{3} - m')(\frac{n-7}{3}) + (\frac{n-4}{3} - m')(\frac{n-4}{3} - m) + (\frac{n-7}{3})(\frac{n-4}{3} - m) + (m+1)(\frac{n-4}{3} - m') + (m+m')(\frac{n-7}{3}) + (m'+1)(\frac{n-4}{3} - m) + 2m + 2m' + 1 = \frac{1}{3}(n^2 - 8n + 19 + 3(m+m' - mm'))$ induced 5-cycles in H for $m, m' \neq 4$. For $m, m' \geq 3$ we see that $m + m' - mm'$ is decreasing in m and m' , so this is at most $\frac{1}{3}(n^2 - 8n + 10)$. If one or both of m and m' is 4, then this increases by at most 2, to give a total of at most $\frac{1}{3}(n^2 - 8n + 16)$.

Case 10: F_1 , F_2 , and F_3 type 3 with m , m' , and m'' vertices respectively.

Vertices a , b , and c are in $\frac{2n}{3} - K_2 - K_3 + m + m''$, $\frac{2n}{3} - K_1 - K_3 + m + m'$, and $\frac{2n}{3} - K_1 - K_2 + m' + m''$, induced 5-cycles respectively. Solving, we obtain $K_1 = m' + \frac{4}{3}$, $K_2 = m'' + \frac{4}{3}$, and $K_3 = m + \frac{4}{3}$. There are exactly $2m + 2m' + 2m''$ induced 5-cycles avoiding $A \cup B \cup C$ if $m, m', m'' \neq 4$. Hence there are $(\frac{n-4}{3} - m')(\frac{n-4}{3} - m'') + (\frac{n-4}{3} - m')(\frac{n-4}{3} - m) + (\frac{n-4}{3} - m'')(\frac{n-4}{3} - m) + (m+m'')(\frac{n-4}{3} - m') + (m+m')(\frac{n-4}{3} - m'') + (m'+m'')(\frac{n-4}{3} - m) + 2m + 2m' + 2m'' = \frac{1}{3}(n^2 - 8n + 16 + 3(2m + 2m' + 2m'' - mm' - m'm'' - mm''))$ induced 5-cycles in H for $m, m', m'' \neq 4$. For $m, m', m'' \geq 3$ we see that $2m + 2m' + 2m'' - mm' - m'm'' - mm''$ is decreasing in m , m' , and m'' , so this is at most $\frac{1}{3}(n^2 - 8n - 11)$. If any of m , m' , and m'' are 4, then this increases by at most 3, to give a total of at most $\frac{1}{3}(n^2 - 8n - 2)$.