

# ON BIRKHOFF-JAMES ORTHOGONALITY PRESERVERS BETWEEN REAL NON-ISOMETRIC BANACH SPACES

RYOTARO TANAKA

ABSTRACT. Real smooth three-dimensional or higher Banach spaces are isomorphic with respect to the nonlinear structure of Birkhoff-James orthogonality if and only if they are isometrically isomorphic. Moreover, using smooth Radon planes and non-smooth direct sums, in arbitrary dimensions, we construct examples of non-isometric pairs of real Banach spaces that admit norm-preserving homogeneous bicontinuous Birkhoff-James orthogonality preservers among them.

## 1. INTRODUCTION

Throughout this paper, the term “Banach space” indicates a *real* Banach space. The present paper is concerned with nonlinear classification of Banach spaces based on Birkhoff-James orthogonality. Let  $X$  be a Banach space, and let  $x, y \in X$ . Then,  $x$  is said to be *Birkhoff-James* orthogonal to  $y$ , which is denoted as  $x \perp_{BJ} y$ , if  $\|x + \lambda y\| \geq \|x\|$  for each  $\lambda \in \mathbb{R}$ . This generalized orthogonality relation in Banach spaces was introduced by Birkhoff [5] and deeply studied by James [9, 10], and is known as one of the most important generalized orthogonality relations since it is closely related to the geometric structure of Banach spaces. For example, if  $x, y \in X$  and  $x \perp_{BJ} y$ , then  $x$  is a nearest point from the origin in the line  $L = \{x + \lambda y : \lambda \in \mathbb{K}\}$ . Further, if  $x$  is a unit vector,  $L$  becomes a tangent line to the unit ball of  $X$ . This last statement also implies a relationship between Birkhoff-James orthogonality and support functionals for the unit ball  $B_X$  of  $X$ . Indeed, studies on Birkhoff-James orthogonality have frequently begun with recalling the following useful result by James [10]:

**Lemma 1.1** (James, 1947). *Let  $X$  be a Banach space, and let  $x, y \in X$ . Then,  $x \perp_{BJ} y$  if and only if  $f(y) = 0$  for some  $f \in \nu(x)$ , where  $\nu(x)$  is the set of all elements of  $B_{X^*}$  that support  $B_X$  at  $x$ ; that is,  $\nu(x) = \{f \in B_{X^*} : f(x) = \|x\|\}$ .*

It is known that Birkhoff-James orthogonality is *non-degenerate* (that is,  $x \perp_{BJ} x$  implies  $x = 0$ ), and *homogeneous* (that is,  $x \perp_{BJ} y$  implies  $\alpha x \perp_{BJ} \beta y$  for each  $\alpha, \beta \in \mathbb{R}$ ). However, it is *asymmetric* in general. Indeed, as was independently shown by Day [7, Theorem 6.4] and James [9, Theorem 1], if  $\dim X \geq 3$  and  $x \perp_{BJ} y$  implies  $y \perp_{BJ} x$  in  $X$ , then  $X$  is a Hilbert space. Here, it should be noted that the assumption on the dimension of  $X$  cannot be omitted. We can construct non-Hilbert two-dimensional real Banach spaces in which Birkhoff-James orthogonality is symmetric (called *Radon planes*); see, for example, [5] and [7], and [14] and [16], for various characterizations of Radon planes. For further

---

2010 *Mathematics Subject Classification.* 46B20; 46B80.

*Key words and phrases.* Banach space, Birkhoff-James orthogonality, Nonlinear classification.

information about generalized orthogonality relations, the readers are referred to Amir [3], who described many characterizations of inner product spaces, and a comprehensive survey on Birkhoff-James orthogonality and isosceles orthogonality by Alonso, Martini, and Wu [2].

As an interesting application of Birkhoff-James orthogonality to the theory of classification of Banach spaces, Koldobsky [12] showed the following theorem.

**Theorem 1.2** (Koldobsky, 1993). *Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$  be linear. Then,  $T$  is a scalar multiple of a linear isometry if and only if  $x, y \in X$  and  $x \perp_{BJ} y$  imply  $Tx \perp_{BJ} Ty$ .*

Later, the complex version of the preceding theorem was proved by Blanco and Turnšek [6]. Moreover, in 2019, Wójcik [21] gave a simpler proof of Theorem 1.2, and improved it as follows.

**Theorem 1.3** (Wójcik, 2019). *Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$  be additive. Then,  $T$  is a scalar multiple of a linear isometry if and only if  $x, y \in X$  and  $x \perp_{BJ} y$  imply  $Tx \perp_{BJ} Ty$ .*

An important consequence of Theorem 1.3 is the following: For Banach spaces  $X, Y$ , if there exists an additive bijection  $T : X \rightarrow Y$  that preserves Birkhoff-James orthogonality in *one direction*, then  $X = Y$  (that is,  $X$  is *isometrically isomorphic* to  $Y$ ). This means that the combination of structures of addition and Birkhoff-James orthogonality determines the entire structure of a Banach space. In other words, with the aid of additivity, Banach spaces are fully classified by their structure of Birkhoff-James orthogonality. A natural question then arises: What happens if the additivity is omitted? In this direction, the author [19] studied the nonlinear equivalence of Banach spaces based on Birkhoff-James orthogonality. Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$  be a (possibly non-additive) bijection. Then,  $T$  is called a *Birkhoff-James orthogonality preserver* if it preserves the Birkhoff-James orthogonality in *both directions* (that is,  $x \perp_{BJ} y$  if and only if  $Tx \perp_{BJ} Ty$ ); in addition,  $X$  is said to be isomorphic to  $Y$  *with respect to the structure of Birkhoff-James orthogonality*, denoted by  $X \sim_{BJ} Y$ , a Birkhoff-James orthogonality preserver exists between  $X$  and  $Y$ . Here, the bidirectionality of a Birkhoff-James orthogonality preserver is required to ensure the symmetry of “ $\sim_{BJ}$ ,” whereas the bijectivity is assumed to guarantee that  $T(X)$  is also a Banach space. It was shown in [19, Corollary 3.7 and Theorem 3.16] that if  $X, Y$  are finite-dimensional, or reflexive and smooth, then  $X \sim_{BJ} Y$  implies  $X \cong Y$  (that is,  $X$  is *linearly isomorphic* to  $Y$ ). Moreover, [19, Theorem 4.3] states that if  $X$  is a Banach space with  $\dim X \geq 3$ , and  $H$  is a Hilbert space, then  $X \sim_{BJ} H$  implies  $X = H$ . By contrast, an example of a non-Hilbert smooth Radon plane  $X$  such that  $X \sim_{BJ} \ell_2^2$  was constructed in [19, Example 4.8], where  $\ell_2^2$  denotes the two-dimensional Euclidean space. This last result shows that, at least in the two-dimensional setting, a Birkhoff-James orthogonality preserver is not necessarily a scalar multiple of an isometric isomorphism. Hence, in general, the additivity assumption cannot be omitted in Theorem 1.3.

In this paper, we present two theorems that further develop the theory of non-linear classification of Banach spaces through Birkhoff-James orthogonality. The first one states that if  $X, Y$  are smooth Banach spaces, then  $X \sim_{BJ} Y$  if and only if  $X = Y$ . The second gives non-isometric pairs of Banach spaces of arbitrary dimensions that admit norm-preserving homogeneous bicontinuous Birkhoff-James orthogonality preservers between them.

2. RESULTS IN SMOOTH BANACH SPACES

Here in, we show that Birkhoff-James orthogonality preservers between non-isometric Banach spaces cannot be constructed under a smooth setting, where a Banach space  $X$  is said to be *smooth* if  $\nu(x)$  is a singleton for each nonzero  $x \in X$ . To see this, we recall the following results described in [19, Theorems 3.6 and 3.9].

**Theorem 2.1.** *Let  $X, Y$  be Banach spaces. Suppose that either  $X$  or  $Y$  is finite-dimensional. Then,  $X \sim_{BJ} Y$  implies that  $\dim X = \dim Y$*

**Theorem 2.2.** *Let  $X, Y$  be smooth Banach spaces, and let  $T : X \rightarrow Y$  be a Birkhoff-James orthogonality preserver. If  $M$  is a reflexive subspace of  $X$ , then  $T(M)$  is a closed subspace of  $Y$ .*

The argument herein is essentially based on the fundamental theorem of projective geometry. To be precise, we make use of the following form given by Mackey [15, Lemma A].

**Lemma 2.3** (Mackey, 1942). *Let  $X, Y$  be Banach spaces, and let  $\mathcal{M}_1(X)$  and  $\mathcal{M}_1(Y)$  be the families of all one-dimensional subspaces of  $X$  and  $Y$ , respectively. If  $\rho : \mathcal{M}_1(X) \rightarrow \mathcal{M}_1(Y)$  is a bijection that preserves the linear independence of the finite elements of  $\mathcal{M}_1(X)$  and  $\mathcal{M}_1(Y)$ . Then, there exists a linear bijection  $T : X \rightarrow Y$  such that  $T(M) = \rho(M)$  for each  $M \in \mathcal{M}_1(X)$ .*

Furthermore, the following auxiliary result is needed. In the following, if  $x_1, \dots, x_n$  are elements of a Banach space, then  $[x_1, \dots, x_n]$  denotes the linear span of  $x_1, \dots, x_n$ .

**Lemma 2.4.** *Let  $X, Y$  be Banach spaces, and let  $\mathcal{F}(X)$  and  $\mathcal{F}(Y)$  be the families of all finite-dimensional subspaces of  $X$  and  $Y$ , respectively. If  $\rho : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$  is an order isomorphism, then  $\rho$  is restricted to a bijection from  $\mathcal{M}_1(X)$  onto  $\mathcal{M}_1(Y)$  that preserves the linear independence of the finite elements of  $\mathcal{M}_1(X)$  and  $\mathcal{M}_1(Y)$ .*

*Proof.* We first note that  $\rho(\{0_X\}) = \{0_Y\}$ . Indeed, we have  $\{0_Y\} = \rho(M)$  for some  $M \in \mathcal{F}(X)$ . Since  $\{0_X\} \subset M$ , it follows that  $\rho(\{0_X\}) \subset \rho(M) = \{0_Y\}$ . This proves that  $\rho(\{0_X\}) = \{0_Y\}$ .

Next, suppose that  $x \in X \setminus \{0\}$ . Then,  $\rho([x]) \neq \{0_Y\}$  as indicated in the preceding paragraph. Take an arbitrary  $y \in \rho([x]) \setminus \{0\}$ . It follows from  $[y] \subset \rho([x])$  that  $\rho^{-1}([y]) \subset [x]$ ; by  $\rho^{-1}([y]) \neq \{0_X\}$ , it implies that  $\rho^{-1}([y]) = [x]$ . Hence, we obtain  $\rho([x]) = [y]$ . In particular,  $\rho(\mathcal{M}_1(X)) \subset \mathcal{M}_1(Y)$ . Since  $\rho^{-1}$  has the same property as  $\rho$ , we also have  $\rho^{-1}(\mathcal{M}_1(Y)) \subset \mathcal{M}_1(X)$ . Therefore,  $\rho$  is restricted to a bijection from  $\mathcal{M}_1(X)$  onto  $\mathcal{M}_1(Y)$ .

Now, let  $x_1, \dots, x_n \in X$  occur such that  $\{x_1, \dots, x_n\}$  is linearly independent. For each  $j$ , choose a  $y_j \in \rho([x_j]) \setminus \{0\}$ . Then,  $\rho([x_j]) = [y_j]$  as indicated in the preceding paragraph. If  $\{y_1, \dots, y_n\}$  is linearly dependent, we have  $[y_i] \subset [y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n]$  for some  $i$ . Meanwhile, if  $j \neq i$ , it follows from  $[x_j] \subset [x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]$  that  $[y_j] \subset \rho([x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n])$ , which implies that

$$[y_i] \subset [y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n] \subset \rho([x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]).$$

Thus, we obtain  $[x_i] \subset [x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n]$ . This contradicts the linear independence of  $\{x_1, \dots, x_n\}$ . Hence,  $\{y_1, \dots, y_n\}$  must be linearly dependent. The exact same argument is also valid for  $\rho^{-1}$ . Consequently,  $\rho$  preserves the linear independence of the finite elements of  $\mathcal{M}_1(X)$  and  $\mathcal{M}_1(Y)$ .  $\square$

Combining these results with Theorem 1.2, we have an improvement of [19, Theorem 3.16]. The following is the first main theorem of this paper.

**Theorem 2.5.** *Let  $X, Y$  be smooth Banach spaces. Suppose that  $\dim X \geq 3$  or  $\dim Y \geq 3$ . Then,  $X \sim_{BJ} Y$  if and only if  $X = Y$ .*

*Proof.* It is sufficient to show the “only if” part. Suppose that  $X \sim_{BJ} Y$ . Then, there exists a Birkhoff-James orthogonality preserver  $T : X \rightarrow Y$ . We first note that  $\dim X \geq 3$  and  $\dim Y \geq 3$  by Theorem 2.1. Let  $M$  be a finite-dimensional subspace of  $X$ . Since  $M$  is reflexive, by Theorem 2.2,  $T(M)$  is a closed subspace of  $Y$ . In particular, the restriction of  $T$  to  $M$  gives a Birkhoff-James orthogonality preserver from  $M$  onto  $T(M)$ . Hence,  $\dim M = \dim T(M)$  by Theorem 2.1. This proves that  $T$  maps a finite-dimensional subspace of  $X$  to a subspace of  $Y$  having the same dimension. Since  $T^{-1}$  has the same property as  $T$ , we can define an order isomorphism  $\rho : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$  by  $\rho(M) = T(M)$ . Moreover, by Lemma 2.4,  $\rho$  is restricted to a bijection from  $\mathcal{M}_1(X)$  onto  $\mathcal{M}_1(Y)$  that preserves the linear independence of the finite elements of  $\mathcal{M}_1(X)$  and  $\mathcal{M}_1(Y)$ . From this, by Lemma 2.3, we have a linear bijection  $S : X \rightarrow Y$  such that  $S(M) = T(M)$  for each  $M \in \mathcal{M}_1(X)$ . Now, suppose that  $x, y \in X$ , and  $x \perp_{BJ} y$ . Then,  $Tx \perp_{BJ} Ty$ . Based on the homogeneity of Birkhoff-James orthogonality, and  $[Sx] = [Tx]$  and  $[Sy] = [Ty]$ , it is implied that  $Sx \perp_{BJ} Sy$ . Thus, by Theorem 1.2,  $S$  is a scalar multiple of an isometric isomorphism from  $X$  onto  $Y$ . This proves that  $X = Y$ .  $\square$

**Remark 2.6.** In contrast to the previous version of the preceding theorem [19, Theorem 3.16], the proof of Theorem 2.5 no longer relies on the main results of Mackey [15, Theorem in p. 246]. The property of the existing linear operator  $S$  follows from Theorem 1.2.

**Remark 2.7.** The assumption that  $\dim X \geq 3$  or  $\dim Y \geq 3$  cannot be omitted. Indeed, it was noted in [19, Example 4.8] that a two-dimensional non-Hilbert smooth real Banach space  $X$  exists that satisfies  $X \sim_{BJ} \ell_2^2$ .

### 3. TOOLS FOR CONSTRUCTION

To prove the second main theorem of this paper, preliminary works are needed. We first recall the acute and obtuse angles based on Birkhoff-James orthogonality.

**Definition 3.1.** Let  $X$  be a Banach space, and let  $x, y \in X$ . Then,  $x$  is *at an acute angle* to  $y$ , as denoted by  $x \perp_{BJ}^+ y$ , if  $\|x + \lambda y\| \geq \|x\|$  for each  $\lambda \geq 0$ . If  $x \perp_{BJ}^+ y$  and  $x \not\perp_{BJ} y$ , then  $x$  is *at a strictly acute angle* to  $y$ , which we write as  $x \perp_{BJ}^{++} y$ .

**Definition 3.2.** Let  $X$  be a Banach space, and let  $x, y \in X$ . Then,  $x$  is *at an obtuse angle* to  $y$ , as denoted by  $x \perp_{BJ}^- y$ , if  $\|x + \lambda y\| \geq \|x\|$  for each  $\lambda \leq 0$ . If  $x \perp_{BJ}^- y$  and  $x \not\perp_{BJ} y$ , then  $x$  is *at a strictly obtuse angle* to  $y$ , which we write as  $x \perp_{BJ}^{--} y$ .

**Remark 3.3.** The notions of Birkhoff-James acute and obtuse angles were already considered by Sain [17]. In particular, their basic properties were stated in [17, Proposition 2.1]. For example, we have the following properties of “ $\perp_{BJ}^+$ ” and “ $\perp_{BJ}^-$ ”:

- (i)  $x \perp_{BJ} y$  if and only if  $x \perp_{BJ}^+ y$  and  $x \perp_{BJ}^- y$ ,
- (ii)  $x \perp_{BJ}^+ y$  implies  $\alpha x \perp_{BJ}^+ \beta y$  for each nonnegative real numbers  $\alpha, \beta$ ,

- (iii)  $x \perp_{B,J}^- y$  implies  $\alpha x \perp_{B,J}^- \beta y$  for each nonnegative real numbers  $\alpha, \beta$ , and
- (iv)  $x \perp_{B,J}^+ y$  if and only if  $x \perp_{B,J}^- (-y)$ .

Now, using support functionals, we present the following characterizations of Birkhoff-James acute and strictly acute angles. An idea is derived from Lemma 1.1.

**Lemma 3.4.** *Let  $X$  be a Banach space, and let  $x, y \in X$ . Then, the following hold:*

- (i)  $x \perp_{B,J}^+ y$  if and only if  $f(y) \geq 0$  for some  $f \in \nu(x)$ ,
- (ii)  $x \perp_{B,J}^{++} y$  if and only if  $f(y) > 0$  for each  $f \in \nu(x)$ .

*Proof.* (i) Suppose that  $x \perp_{B,J}^+ y$ . Then, for each  $n \in \mathbb{N}$ , there exists an  $f_n \in S_{X^*}$  such that

$$f_n(x + n^{-1}y) = \|x + n^{-1}y\| \geq \|x\|.$$

In particular, we have

$$f_n(x) \geq \|x\| - \frac{1}{n}f_n(y) \geq \|x\| - \frac{1}{n}\|y\|$$

and

$$f_n(y) \geq n(\|x\| - f_n(x)) \geq 0$$

for each  $n \in \mathbb{N}$ . Since  $B_{X^*}$  is weakly\* compact by the Banach-Alaoglu theorem, there exists a subnet  $(f_{n_a})_a$  of  $(f_n)_n$  that converges weakly\* to some  $f_0 \in B_{X^*}$ . It follows that

$$f_0(x) = \lim_a f_{n_a}(x) = \lim_n f_n(x) = \|x\|,$$

and  $f_0(y) = \lim_a f_{n_a}(y) \geq 0$ ; that is,  $f_0 \in \nu(x)$  and  $f_0(y) \geq 0$ .

Conversely, if  $f(y) \geq 0$  for some  $f \in \nu(x)$ , then

$$\|x + \lambda y\| \geq f(x + \lambda y) = \|x\| + \lambda f(y) \geq \|x\|$$

for each  $\lambda \geq 0$ . Hence,  $x \perp_{B,J}^+ y$  holds.

(ii) If  $x \perp_{B,J}^{++} y$ , then  $x \perp_{B,J}^+ y$  and  $x \not\perp_{B,J} y$ . By (i), we obtain  $f_1(y) \geq 0$  for some  $f_1 \in \nu(x)$ . Moreover, by Lemma 1.1,  $f(y) \neq 0$  for each  $f \in \nu(x)$ . It follows that  $f_1(y) > 0$ . Now, suppose that  $f_2(y) < 0$  for some  $f_2 \in \nu(x)$ . Since  $\nu(x)$  is convex, we derive

$$g = \frac{-f_2(y)}{f_1(y) - f_2(y)}f_1 + \frac{f_1(y)}{f_1(y) - f_2(y)}f_2 \in \nu(x),$$

and  $g(y) = 0$ . However, this contradicts  $x \not\perp_{B,J} y$  by Lemma 1.1. Thus, there is no  $f \in \nu(x)$  satisfying  $f(y) \leq 0$ ; that is, we have  $f(y) > 0$  for each  $f \in \nu(x)$ .

The converse follows from (i) and Lemma 1.1. □

In the following, let  $R_x = \{y \in X : x \perp_{B,J} y\}$  and  $R_x^{++} = \{y \in X : x \perp_{B,J}^{++} y\}$  for each  $x \in X$ . Below, we show that  $R_x^{++}$  is an open convex subset of  $X$  which is maximal as a connected subset of  $X \setminus R_x$ .

**Lemma 3.5.** *Let  $X$  be a Banach space, and let  $x \in X \setminus \{0\}$ . Then,  $R_x^{++}$  is an open convex subset of  $X \setminus R_x$  containing  $x$ .*

*Proof.* Set  $\Pi^{++}(f) = \{y \in X : f(y) > 0\}$  for each  $f \in X^*$ . We then have  $R_x^{++} = \bigcap \{\Pi^{++}(f) : f \in \nu(x)\}$  by Lemma 3.4 (ii). Since  $\Pi^{++}(f)$  is open, convex, and contains  $x$  whenever  $f \in \nu(x)$ , it follows that  $R_x^{++}$  is an open convex subset of  $X \setminus R_x$  containing  $x$ . □

**Lemma 3.6.** *Let  $X$  be a Banach space, and let  $x \in X \setminus \{0\}$ . Suppose that  $A$  is a connected subset of  $X \setminus R_x$  containing  $x$ . Then,  $A \subset R_x^{++}$ .*

*Proof.* To show  $A \subset R_x^{++}$ , suppose to the contrary that  $A \not\subset R_x^{++}$ . Let  $y \in A \setminus R_x^{++}$ . Then,  $f_0(y) \leq 0$  for some  $f_0 \in \nu(x)$  by Lemma 3.4. However, since  $A$  is connected and  $f_0(x) = \|x\| > 0$ , the intermediate value theorem ensures that  $f_0(z) = 0$  for some  $z \in A$ . It follows from Lemma 1.1 that  $x \perp_{BJ} z$ . This contradicts  $A \subset X \setminus R_x$ . Therefore,  $A \subset R_x^{++}$  holds.  $\square$

Based on these lemmas, it turns out that if a Birkhoff-James orthogonality preserver is continuous, it then also preserves the Birkhoff-James acute angles.

**Theorem 3.7.** *Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$  be a Birkhoff-James orthogonality preserver. Suppose that  $T$  is continuous. Then,  $x, y \in X$  and  $x \perp_{BJ}^+ y$  implies  $Tx \perp_{BJ}^+ Ty$ . Consequently, if  $T$  is bicontinuous, then  $T$  is a Birkhoff-James acute angle preserver.*

*Proof.* We first note that  $T(R_x) = R_{Tx}$  holds for each  $x \in X$ . Take an arbitrary  $x \in X \setminus \{0\}$ . Then, by Lemma 3.5,  $R_x^{++}$  is an open convex subset of  $X \setminus R_x$  containing  $x$ . Since  $T$  is continuous, it follows that  $T(R_x^{++})$  is a connected subset of  $Y \setminus R_{Tx}$  containing  $Tx$ . Hence, by Lemma 3.6, we obtain  $T(R_x^{++}) \subset R_{Tx}^{++}$ .

Now, suppose that  $x, y \in X$ , and  $x \perp_{BJ}^+ y$ . If  $x \perp_{BJ} y$ , then  $Tx \perp_{BJ} Ty$ , whereas if  $x \not\perp_{BJ} y$ , it then follows that  $y \in R_x^{++}$ , and  $Ty \in R_{Tx}^{++}$ . Therefore, in either case, we obtain  $Tx \perp_{BJ}^+ Ty$ .  $\square$

Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$ . Then,  $T$  is said to be *norm-preserving* if  $\|Tx\| = \|x\|$  for each  $x \in X$ , and *homogeneous* if  $T(cx) = cTx$  for each  $x \in X$  and  $c \in \mathbb{R}$ . We now introduce a new notion of the equivalence of Banach spaces that is stronger than “ $\sim_{BJ}$ .”

**Definition 3.8.** Let  $X, Y$  be Banach spaces. Then,  $X$  is said to be *strongly isomorphic* to  $Y$  with respect to the structure of Birkhoff-James orthogonality, denoted by  $X =_{BJ} Y$ , if there exists a norm-preserving homogeneous bicontinuous Birkhoff-James orthogonality preserver  $T : X \rightarrow Y$ .

**Remark 3.9.** Obviously,  $X = Y$  implies  $X =_{BJ} Y$ , and  $X =_{BJ} Y$  implies  $X \sim_{BJ} Y$ .

The rest of this section is devoted to showing that if  $X =_{BJ} Z$  and  $Y =_{BJ} W$ , then  $X \oplus_\infty Y =_{BJ} Z \oplus_\infty W$ , where  $X \oplus_\infty Y$  is the Banach space  $X \times Y$  endowed with the norm  $\|(x, y)\|_\infty = \max\{\|x\|, \|y\|\}$ . We begin with a further analysis of Birkhoff-James acute angles.

**Lemma 3.10.** *Let  $X$  be a Banach space, and let  $x, y \in X$ . Then,  $x \perp_{BJ}^+ y$  if and only if there exists a sequence of positive real numbers  $(\lambda_n)_n$  such that  $\lim_n \lambda_n = 0$  and  $\|x + \lambda_n y\| \geq \|x\|$  for each  $n$ .*

*Proof.* It is sufficient to show the “if” part. Suppose that  $(\lambda_n)_n$  is a sequence of positive real numbers such that  $\lim_n \lambda_n = 0$  and  $\|x + \lambda_n y\| \geq \|x\|$  for each  $n$ . Then, for each  $n$ , there exists an  $f_n \in S_{X^*}$  such that

$$f_n(x + \lambda_n y) = \|x + \lambda_n y\| \geq \|x\|$$

for each  $n$ . By the Banach-Alaoglu theorem, we have a subnet  $(f_{n_a})_a$  of  $(f_n)_n$  that converges weakly\* to some  $f_0 \in B_{X^*}$ . Since

$$f_n(x) \geq \|x\| - \lambda_n f_n(y) \geq \|x\| - \lambda_n \|y\|$$

and

$$f_n(y) \geq \lambda_n^{-1}(\|x\| - f_n(x)) \geq 0$$

for each  $n$ , it follows that  $f_0(x) = \|x\|$  and  $f_0(y) \geq 0$ . Therefore,  $x \perp_{B,J}^+ y$  by Lemma 3.4 (i).  $\square$

**Lemma 3.11.** *Let  $X, Y$  be Banach spaces, and let  $(x_1, y_1), (x_2, y_2) \in X \oplus_\infty Y$ . Then,  $(x_1, y_1) \perp_{B,J}^+ (x_2, y_2)$  if and only if one of the following holds:*

- (i)  $\|x_1\| > \|y_1\|$ , and  $x_1 \perp_{B,J}^+ x_2$ ;
- (ii)  $\|x_1\| = \|y_1\|$ , and  $x_1 \perp_{B,J}^+ x_2$  or  $y_1 \perp_{B,J}^+ y_2$ ;
- (iii)  $\|x_1\| < \|y_1\|$ , and  $y_1 \perp_{B,J}^+ y_2$ .

*Proof.* Suppose that  $(x_1, y_1) \perp_{B,J}^+ (x_2, y_2)$ . Then, either  $\|x_1\| > \|y_1\|$ ,  $\|x_1\| = \|y_1\|$ , or  $\|x_1\| < \|y_1\|$  holds. If  $\|x_1\| > \|y_1\|$ , we have  $\|(x_1, y_1)\|_\infty = \|x_1\|$ . Moreover,  $\|x_1 + n^{-1}x_1\| > \|y_1 + n^{-1}y_2\|$  for a sufficiently large  $n$ , and thus

$$\|x_1 + n^{-1}x_2\| = \|(x_1, y_1) + n^{-1}(x_2, y_2)\|_\infty \geq \|(x_1, y_1)\|_\infty = \|x_1\|.$$

Hence, by Lemma 3.10, we obtain  $x_1 \perp_{B,J}^+ x_2$ . In the case of  $\|x_1\| < \|y_1\|$ , we know  $y_1 \perp_{B,J}^+ y_2$  in a similar way.

Finally, we consider the case of  $\|x_1\| = \|y_1\|$ . Then,  $\|(x_1, y_1)\|_\infty = \|x_1\| = \|y_1\|$ . Set

$$N_1 = \{n \in \mathbb{N} : \|x_1 + n^{-1}x_2\| \geq \|y_1 + n^{-1}y_2\|\}$$

$$N_2 = \{n \in \mathbb{N} : \|x_1 + n^{-1}x_2\| < \|y_1 + n^{-1}y_2\|\}.$$

Since  $N_1 \cup N_2 = \mathbb{N}$ , either  $N_1$  or  $N_2$  is an infinite set. If  $N_1$  is an infinite set, as in the preceding paragraph, we have  $x_1 \perp_{B,J}^+ x_2$ . Similarly, if  $N_2$  is an infinite set,  $y_1 \perp_{B,J}^+ y_2$  holds.

Conversely, suppose that (i), (ii), or (iii) holds. If (i) holds, then

$$\|(x_1, y_1) + \lambda(x_2, y_2)\|_\infty \geq \|x_1 + \lambda x_2\| \geq \|x_1\| = \|(x_1, y_1)\|_\infty$$

for each  $\lambda \geq 0$ . Hence,  $(x_1, y_1) \perp_{B,J}^+ (x_2, y_2)$ . A similar argument shows the same conclusion in the case of (iii). Assume that (ii) holds. Then, either

$$\|x_1 + \lambda x_2\| \geq \|x_1\| = \|(x_1, y_1)\|_\infty$$

for each  $\lambda \geq 0$ , or

$$\|y_1 + \lambda y_2\| \geq \|y_1\| = \|(x_1, y_1)\|_\infty$$

for each  $\lambda \geq 0$ . Thus, in either case, it follows that

$$\|(x_1, y_1) + \lambda(x_2, y_2)\|_\infty \geq \|y_1\| = \|(x_1, y_1)\|_\infty$$

for each  $\lambda \geq 0$ . This proves that  $(x_1, y_1) \perp_{B,J}^+ (x_2, y_2)$ .  $\square$

Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$ . Then,  $T$  is called a *Birkhoff-James acute angle preserver* if it is bijective and  $x \perp_{B,J}^+ y$  if and only if  $Tx \perp_{B,J}^+ Ty$ . For homogeneous Birkhoff-James acute angle preservers, we have the following result:

**Proposition 3.12.** *Let  $X, Y$  be Banach spaces, and let  $T : X \rightarrow Y$  be a homogeneous Birkhoff-James acute angle preserver. Then,  $T$  is a Birkhoff-James orthogonality preserver.*

*Proof.* Suppose that  $x, y \in X$  and  $x \perp_{BJ} y$ . Then,  $x \perp_{BJ}^+ y$  and  $x \perp_{BJ}^+ (-y)$ ; in addition, by  $T(-y) = -Ty$ , this implies that  $Tx \perp_{BJ}^+ Ty$  and  $Tx \perp_{BJ}^+ (-Ty)$ . Hence, it follows that  $Tx \perp_{BJ} Ty$ . This completes the proof since the exact same argument is valid for  $T^{-1}$ .  $\square$

We are now ready to prove the following theorem.

**Theorem 3.13.** *Let  $X, Y, Z, W$  be Banach spaces such that  $X =_{BJ} Z$  and  $Y =_{BJ} W$ . Then,  $X \oplus_{\infty} Y =_{BJ} Z \oplus_{\infty} W$ .*

*Proof.* Let  $S : X \rightarrow Z$  and  $T : Y \rightarrow W$  be norm-preserving homogeneous bicontinuous Birkhoff-James preservers, and let

$$R(x, y) = (Sx, Ty)$$

for each  $(x, y) \in X \oplus_{\infty} \mathbb{R}$ . Then,  $R : X \oplus_{\infty} Y \rightarrow Z \oplus_{\infty} W$  is a norm-preserving homogeneous bicontinuous bijection because  $R^{-1}$  is given by  $R^{-1}(z, w) = (S^{-1}z, T^{-1}w)$  for each  $(z, w) \in Z \oplus_{\infty} W$ . Moreover,  $R, T$  preserve the Birkhoff-James acute angle by Theorem 3.7. Now, suppose that  $(x_1, y_1), (x_2, y_2) \in X \oplus_{\infty} Y$ , and  $(x_1, y_1) \perp_{BJ}^+ (x_2, y_2)$ . From Lemma 3.11, one of the following holds:

- (i)  $\|x_1\| > \|y_1\|$ , and  $x_1 \perp_{BJ}^+ x_2$ ;
- (ii)  $\|x_1\| = \|y_1\|$ , and  $x_1 \perp_{BJ}^+ x_2$  or  $y_1 \perp_{BJ}^+ y_2$ ;
- (iii)  $\|x_1\| < \|y_1\|$ , and  $y_1 \perp_{BJ}^+ y_2$ .

If (i) holds, then  $\|Sx_1\| = \|x_1\| > \|y_1\| = \|Ty_1\|$  and  $Sx_1 \perp_{BJ}^+ Sx_2$ ; hence, from Lemma 3.11 again,  $R(x_1, y_1) \perp_{BJ}^+ R(x_2, y_2)$ . Similarly, we obtain the same conclusion for the other cases. Thus,  $R$  preserves the Birkhoff-James acute angle. Since the same is true for  $R^{-1}$ , by Proposition 3.12, we can conclude that  $R$  is a Birkhoff-James orthogonality preserver. Therefore,  $X \oplus_{\infty} Y =_{BJ} Z \oplus_{\infty} W$ .  $\square$

**Corollary 3.14.** *Let  $X, Y, Z$  be Banach spaces such that  $X =_{BJ} Y$ . Then,  $X \oplus_{\infty} Z =_{BJ} Y \oplus_{\infty} Z$ .*

#### 4. EXAMPLES

In this section, we construct example pairs of real Banach spaces  $(X, Y)$  of arbitrary dimensions such that  $X =_{BJ} Y$  and  $X \neq Y$ . First, we improve [19, Theorem 4.7] by slightly modifying its proof.

**Theorem 4.1.** *Let  $X$  be a smooth Radon plane. Then,  $X =_{BJ} \ell_2^2$ .*

*Proof.* Let  $x(\theta) = (\cos \theta, \sin \theta)$  and  $y(\theta) = \|x(\theta)\|_X^{-1} x(\theta)$  for each  $\theta \in [0, 2\pi]$ . Then, as in the proof of [19, Theorem 4.7], there exists a bijection  $\eta : [0, \pi/2] \rightarrow [\pi/2, \pi]$  such that  $\eta(0) = \pi/2$ ,  $\eta(\pi/2) = \pi$ , and  $y(\theta) \perp_{BJ} y(\eta(\theta))$  for each  $\theta \in [0, \pi/2]$ . We first show that this  $\eta$  is automatically continuous. Let  $\theta_0 \in [0, \pi/2]$ . Suppose that  $(\theta_n)_n$  is a sequence in  $[0, \pi/2]$  that converges to  $\theta_0$ . By the definition of  $\eta$ , we have  $y(\theta_n) \perp_{BJ} y(\eta(\theta_n))$ , which implies that

$$\|y(\theta_n) + \lambda y(\eta(\theta_n))\| \geq \|y(\theta_n)\| = 1$$

for each  $\lambda \in \mathbb{R}$ . Now, take an arbitrary subsequence  $(\eta(\theta_{k_n}))_n$  of  $(\eta(\theta_n))_n$ . Then, there exists a subsequence  $(\eta(\theta_{l_{k_n}}))_n$  of  $(\eta(\theta_{k_n}))_n$  that converges to some  $\theta'_0 \in$



$[\pi/2, \pi]$ . Since the mapping  $\theta \mapsto y(\theta)$  is a homeomorphism from  $[0, \pi]$  to  $S_X^+$ , we derive

$$\|y(\theta_0) + \lambda y(\theta'_0)\| = \lim_n \|y(\theta_{l_{k_n}}) + \lambda y(\eta(\theta_{l_{k_n}}))\| \geq 1$$

for each  $\lambda \in \mathbb{R}$ . This means that  $y(\theta_0) \perp_{BJ} y(\theta'_0)$ ; hence,  $\theta'_0 = \eta(\theta_0)$  by the smoothness of  $X$ . Therefore,  $\eta(\theta_n) \rightarrow \eta(\theta_0)$ ; that is,  $\eta$  is continuous on  $[0, \pi/2]$ .

Next, set

$$T(x(\theta)) = \begin{cases} y(\theta) & (\theta \in [0, \pi/2]) \\ y(\eta(\theta - \pi/2)) & (\theta \in [\pi/2, \pi]) \end{cases} .$$

Then, as was shown in [19, Theorem 4.7],  $T : S_{\ell_2^+} \rightarrow S_X$  is a bijection such that  $x(\theta) \perp x(\theta')$  in  $\ell_2^2$  if and only if  $Tx(\theta) \perp_{BJ} Tx(\theta')$  in  $X$ . We note that the mapping  $\theta \mapsto x(\theta)$  is a homeomorphism from  $[0, \pi/2]$  onto  $S_{\ell_2^+}$ . Since  $y(\pi/2) = y(\eta(0))$ , and  $\eta$  is continuous as indicated in the preceding paragraph,  $T$  is also continuous.

We extend  $T$  by setting  $T^=(x) = -T(-x)$  for each  $-S_{\ell_2^+}$ . Then,  $T^= : S_{\ell_2^+} \rightarrow S_X$  is a bijection such that  $x \perp_{BJ} y$  in  $\ell_2^2$  if and only if  $T^=x \perp_{BJ} T^=y$ . It is easy to check whether  $T^=$  is continuous. In fact,  $T^=$  is a homeomorphism since it is a continuous bijection from a compact space  $S_{\ell_2^+}$  onto a Hausdorff space  $S_X$ .

Finally, we define the full extension  $T^\sim$  of  $T^=$  to  $\ell_2^2$  by

$$T^\sim x = \begin{cases} 0 & (x = 0) \\ \|x\|_2 T^= \left( \frac{1}{\|x\|_2} x \right) & (x \neq 0) \end{cases} .$$

This gives rise to a norm-preserving homogeneous continuous Birkhoff-James orthogonality preserver from  $\ell_2^2$  onto  $X$ . Moreover, since

$$(T^\sim)^{-1} y = \begin{cases} 0 & (y = 0) \\ \|y\|_X (T^=)^{-1} \left( \frac{1}{\|y\|_X} y \right) & (y \neq 0) \end{cases} ,$$

$(T^\sim)^{-1}$  is also continuous. Thus,  $X =_{BJ} \ell_2^2$  holds. □

For a nonempty set  $I$ , the symbol  $c_0(I)$  denotes the Banach space of all systems  $(a_n)_{n \in I}$  such that  $\{n \in I : |a_n| \geq \varepsilon\}$  is finite for each  $\varepsilon > 0$ . To prove the non-isometric clause of the examples, we need the following technical lemma.

**Lemma 4.2.** *Let  $X$  be a Banach space, and let  $I$  be a nonempty set. If  $X$  does not contain a two-dimensional Hilbert subspace, the same is then true for  $X \oplus_\infty c_0(I)$ .*

*Proof.* We first prove the theorem under the case in which  $I$  is a singleton. Suppose that  $X \oplus_\infty \mathbb{R}$  contains a two-dimensional Hilbert subspace  $M$ . Let  $\{(x, r), (y, s)\} \subset M$  be an orthonormal basis for  $M$ . We then have

$$\|\alpha(x, r) + \beta(y, s)\|_\infty = (\alpha^2 + \beta^2)^{1/2}$$

for each  $\alpha, \beta \in \mathbb{R}$ . In particular, setting  $(\alpha, \beta) = (r, s)$ , we obtain

$$r^2 + s^2 \leq \|r(x, r) + s(y, s)\|_\infty = (r^2 + s^2)^{1/2},$$

which implies that  $(r^2 + s^2)^{1/2} \leq 1$ . Combining this with the Cauchy-Schwartz inequality yields

$$|\alpha r + \beta s| \leq (\alpha^2 + \beta^2)^{1/2}$$

for each  $(\alpha, \beta)$ . Moreover, the equality holds only if  $(\alpha, \beta) = c(r, s)$  for some  $c \in \mathbb{R}$ . This means that

$$\|\alpha x + \beta y\| = (\alpha^2 + \beta^2)^{1/2}$$

for each  $(\alpha, \beta) \in \mathbb{R}^2 \setminus [(r, s)]$ . Since the set  $\mathbb{R}^2 \setminus [(r, s)]$  is dense in  $\mathbb{R}^2$  with respect to the product topology on  $\mathbb{R}^2$ , it follows that

$$\|\alpha x + \beta y\| = (\alpha^2 + \beta^2)^{1/2}$$

for all  $(\alpha, \beta) \in \mathbb{R}^2$ . Therefore,  $[x, y]$  is a two-dimensional Hilbert subspace of  $X$ .

Next, suppose that  $I$  is finite, and that  $X \oplus_\infty c_0(I)$  contains a two-dimensional Hilbert subspace. We may write  $I = \{1, \dots, n\}$ . Since

$$X \oplus_\infty c_0(\{1, \dots, n\}) = (X \oplus_\infty c_0(\{1, \dots, n-1\})) \oplus_\infty \mathbb{R},$$

as indicated in the first paragraph of the proof,  $X \oplus_\infty c_0(\{1, \dots, n-1\})$  contains a two-dimensional Hilbert subspace. Continuing this process, we infer that  $X$  contains a two-dimensional Hilbert subspace.

Finally, assume that  $I$  is infinite. Suppose that  $\{(x, (a_n)_{n \in I}), (y, (b_n)_{n \in I})\} \subset X \oplus_\infty c_0(I)$  is an orthonormal basis for two-dimensional Hilbert subspace. Since  $(a_n)_{n \in I}, (b_n)_{n \in I} \in c_0(I)$ , the sets

$$I_1 = \left\{ n \in I : |a_n| \geq \frac{1}{\sqrt{2}} \right\}$$

$$I_2 = \left\{ n \in I : |b_n| \geq \frac{1}{\sqrt{2}} \right\}$$

are both finite. Set  $I_0 = I_1 \cup I_2$ . Then,  $I_0$  is also finite, and

$$|\alpha a_n + \beta b_n| \leq (\alpha^2 + \beta^2)^{1/2} (a_n^2 + b_n^2)^{1/2} < (\alpha^2 + \beta^2)^{1/2}$$

for each  $n \notin I_0$  and each nonzero  $(\alpha, \beta) \in \mathbb{R}^2$ . Hence, it follows that

$$\begin{aligned} (\alpha^2 + \beta^2)^{1/2} &= \|\alpha(x, (a_n)_{n \in I}) + \beta(y, (b_n)_{n \in I})\|_\infty \\ &= \max \left\{ \|\alpha x + \beta y\|, \max_{n \in I} |\alpha a_n + \beta b_n| \right\} \\ &= \max \left\{ \|\alpha x + \beta y\|, \max_{n \in I_0} |\alpha a_n + \beta b_n| \right\} \end{aligned}$$

for each  $(\alpha, \beta) \in \mathbb{R}^2$ . This means that  $X \oplus_\infty c_0(I_0)$  contains a two-dimensional Hilbert subspace. Thus, as described in the preceding paragraph, we can conclude that  $X$  contains a two-dimensional Hilbert subspace.  $\square$

Now, we prove the second main theorem of this paper.

**Theorem 4.3.** *Let  $X$  be a smooth Radon plane, and let  $I$  be a nonempty set. Suppose that  $X \neq \ell_2^2$ . Then,  $X \oplus_\infty c_0(I) =_{BJ} \ell_2^2 \oplus_\infty c_0(I)$ , and  $X \oplus_\infty c_0(I) \neq \ell_2^2 \oplus_\infty c_0(I)$ .*

*Proof.* By Corollary 3.14 and Theorem 4.1, we have  $X \oplus_\infty c_0(I) =_{BJ} \ell_2^2 \oplus_\infty c_0(I)$ . Moreover, since  $X \neq \ell_2^2$ , by Lemma 4.2,  $X \oplus_\infty c_0(I)$  does not contain two-dimensional Hilbert subspace, whereas  $\ell_2^2 \subset \ell_2^2 \oplus_\infty c_0(I)$ . Therefore,  $X \oplus_\infty c_0(I) \neq \ell_2^2 \oplus_\infty c_0(I)$ .  $\square$

**Example 4.4.** As was shown in [19, Example 4.8], the Day-James space  $\ell_{p,q}^2$  with  $p^{-1} + q^{-1} = 1$  is a non-Hilbert smooth Radon plane, where  $\ell_{p,q}^2$  is the Banach space  $\mathbb{R}^2$  endowed with the norm

$$\|(a, b)\|_{p,q} = \begin{cases} (|a|^p + |b|^p)^{1/p} & (ab \geq 0) \\ (|a|^q + |b|^q)^{1/q} & (ab \leq 0) \end{cases} .$$

Hence, we have an infinitely many examples of  $X$  that satisfy the assumption of the preceding theorem.

### 5. REMARKS

The proofs of two main theorems of this paper are only valid for real Banach spaces. Indeed, in a complex case, the operator  $S$  constructed in the proof of Theorem 2.5 is generally *semilinear*. To obtain the linearity or antilinearity of  $S$ , we need some arguments related to the continuity. For example, it was shown by [11, Lemma 2 and Corollary] that a semilinear bijection between infinite-dimensional complex Banach spaces is linear or antilinear if it preserves closed hyperplanes in both directions. Moreover, such an operator is automatically continuous; see Fillmore and Longstaff [8, Lemma 2 and Lemma 3]. The finite-dimensional case is known to be more complicated. At this stage, we do not know if a Birkhoff-James orthogonality preserver between smooth complex Banach spaces satisfies some continuity conditions. At a minimum, if the considered complex Banach spaces are reflexive, smooth, and infinite-dimensional, we may thus overcome this difficulty by using [19, Corollary 3.10] and [8, Theorem 1] instead of Theorem 2.2 and Lemma 2.3. However, there is no idea on how to fill in the gaps if the spaces are non-reflexive, or finite-dimensional.

The validity of the second main theorem for complex spaces is under a more serious condition. The construction of examples described herein relies heavily on the existence of non-Hilbert Radon planes. However, to the best of the author's knowledge, it is not known if a complex Radon plane exists.

Finally, we note that all known pairs of real Banach spaces  $(X, Y)$  satisfying  $X \sim_{BJ} Y$  are mutually isomorphic. Hence, we wonder if  $X \sim_{BJ} Y$  implies  $X \cong Y$  without any assumption. Meanwhile, by [19, Remarks 3.14 and 3.15], we have a complex Banach space  $X$  that satisfies (automatically)  $X \sim_{BJ} \overline{X}$ , and  $X \not\cong \overline{X}$ . Therefore, the corresponding problem in the complex case would be the following: Does  $X \sim_{BJ} Y$  imply  $X \cong Y$  or  $X \cong \overline{Y}$ ?

We conclude this paper with some useful tools for analyzing the structure of Birkhoff-James orthogonality in Banach spaces. In 2005, Turnšek [20] considered a local symmetry condition to Birkhoff-James orthogonality, with was then applied to determine the forms of the Birkhoff-James orthogonality preservers on  $B(H)$ , where  $B(H)$  is the Banach space of all bounded linear operators on a complex Hilbert space  $H$ . It was shown accurately in [20, Theorem 2.5] that if  $B \in B(H)$ , and if  $A \perp_{BJ} B$  always implies  $B \perp_{BJ} A$  in  $B(H)$ , then  $B$  is a scalar multiple of an isometry or a coisometry. Moreover, this result was generalized to a characterization of extreme points of  $B_A$  for a von Neumann algebra  $A$ ; see [13, Theorem 4.7]. Now, such a property of  $B$  is known as *right symmetry* for Birkhoff-James orthogonality. Similarly, an element  $x$  of a Banach space  $X$  is called a *left symmetric point* for Birkhoff-James orthogonality if  $x \perp_{BJ} y$  always implies  $y \perp_{BJ} x$  in  $X$ . The terms “left symmetric” and “right symmetric” were introduced by Sain [17],

and Sain, Ghosh, and Paul [18], respectively. These locally symmetric points for Birkhoff-James orthogonality reflect the geometric features of Banach spaces, and are preserved under Birkhoff-James orthogonality preservers. Hence, they may play important roles in classifying general Banach spaces based on the structure of Birkhoff-James orthogonality.

Another interesting tool is *orthographs* of *symmetrized* Birkhoff-James orthogonality that were recently introduced and studied by Arambašić et al. [4]. We define  $x \perp y$  by  $x \perp_{BJ} y$  and  $y \perp_{BJ} x$ . Then, “ $\perp$ ” is called *mutual* Birkhoff-James orthogonality. The orthograph  $\Gamma(X)$  of a Banach space  $X$  induced by “ $\perp$ ” is defined as follows: The vertex set  $V(\Gamma(X))$  is the set of all one-dimensional subspaces of  $X$ , and the vertices  $[x], [y]$  are *adjacent* if  $x \perp y$ . Naturally, the properties of the orthographs depend on the geometric structure of the Banach spaces. The author believes that orthographs are preserved under “ $\sim_{BJ}$ ” in a reasonable manner, and are useful in some important circumstances.

#### ACKNOWLEDGMENT

The author would like to thank Editage (www.editage.com) for English language editing. This work was supported by JSPS KAKENHI Grant Number JP19K14561.

#### REFERENCES

- [1] J. Alonso, *Any two-dimensional normed space is a generalized Day-James space*, J. Inequal. Appl., **2011**, 2011:2, 3 pp.
- [2] J. Alonso, H. Martini and S. Wu, *On Birkhoff orthogonality and isosceles orthogonality in normed linear spaces*, Aequationes Math., **83** (2012), 153–189.
- [3] D. Amir, *Characterizations of inner product spaces*, Birkhäuser Verlag, Basel, 1986.
- [4] L. Arambašić, A. Guterman, B. Kuzma, R. Rajić and S. Zhilina, *Symmetrized Birkhoff-James orthogonality in arbitrary normed spaces*, J. Math. Anal. Appl., **502**, (2021), 125203, 16 pp.
- [5] G. Birkhoff, *Orthogonality in linear metric spaces*, Duke Math. J., **1** (1935), 169–172.
- [6] A. Blanco and A. Turnšek, *On maps that preserve orthogonality in normed spaces*, Proc. R. Soc. Edinburgh Sect. A, **136** (2006), 709–716.
- [7] M. M. Day, *Some characterization of inner-product spaces*, Trans. Amer. Math. Soc., **47** (1947), 320–337.
- [8] P. A. Fillmore and W. E. Longstaff, *On isomorphisms of lattice of closed subspaces*, Canad. J. Math., **36** (1984), 820–829.
- [9] R. C. James, *Inner product in normed linear spaces*, Bull. Amer. Math. Soc., **53** (1947), 559–566.
- [10] R. C. James, *Orthogonality and linear functionals in normed linear spaces*, Trans. Amer. Math. Soc., **61** (1947), 265–292.
- [11] S. Kakutani and G. W. Mackey, *Ring and lattice characterization of complex Hilbert space*, Bull. Amer. Math. Soc., **52** (1946), 727–733.
- [12] A. Koldobsky, *Operators preserving orthogonality are isometries*, Proc. Roy. Soc. Edinburgh Sect. A, **123** (1993), 835–837.
- [13] N. Komuro, K.-S. Saito and R. Tanaka, *Symmetric points for (strong) Birkhoff-James orthogonality in von Neumann algebras with applications to preserver problems*, J. Math. Anal. Appl., **463** (2018), 1109–1131.
- [14] N. Komuro, K.-S. Saito and R. Tanaka, *A characterization of Radon planes using generalized Day-James spaces*, Ann. Funct. Anal., **11** (2020), 62–74.
- [15] G. W. Mackey, *Isomorphisms of normed linear spaces*, Ann. of Math. (2), **43** (1942), 244–260.
- [16] H. Martini and K. J. Swanepoel, *Antinorms and Radon curves*, Aequationes Math., **72** (2006), 110–138.
- [17] D. Sain, *Birkhoff-James orthogonality of linear operators on finite dimensional Banach spaces*, J. Math. Anal. Appl., **447** (2017), 860–866.
- [18] D. Sain, P. Ghosh and K. Paul, *On symmetry of Birkhoff-James orthogonality of linear operators on finite-dimensional real Banach spaces*, Oper. Matrices, **11** (2017), 1087–1095.

- [19] R. Tanaka, *Nonlinear equivalence of Banach spaces based on Birkhoff-James orthogonality*, J. Math. Anal. Appl., **505** (2022), 125444.
- [20] A. Turnšek, *On operators preserving James' orthogonality*, Linear Algebra Appl., **407** (2005), 189–195.
- [21] P. Wójcik, *Mappings preserving B-orthogonality*, Indag. Math. (N.S.), **30** (2019), 197–200.

KATSUSHIKA DIVISION, INSTITUTE OF ARTS AND SCIENCES, TOKYO UNIVERSITY OF SCIENCE,  
TOKYO 125-8585, JAPAN

*Email address:* `r-tanaka@rs.tus.ac.jp`