

# A local estimate for the mean curvature flow

Zhen Wang

August 3, 2021

## Abstract

We establish a pointwise estimate of  $A$  along the mean curvature flow in terms of the initial geometry and the  $|HA|$  bound. As corollaries we obtain the extension theorem of  $HA$  and the blowup rate estimate of  $HA$ .

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Preliminaries</b>	<b>3</b>
<b>3</b>	<b><math>L^p</math> estimate</b>	<b>4</b>
<b>4</b>	<b><math>L^\infty</math> estimate and extension theorem</b>	<b>11</b>
<b>5</b>	<b>Blowup estimate of <math>HA</math></b>	<b>14</b>

## 1 Introduction

Let  $\mathbf{x}_0 : \Sigma^n \rightarrow \mathbb{R}^{n+1}$  be a complete smooth immersed hypersurface without boundary and a family of immersions  $\mathbf{x}(x, t) : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a solution to the equation

$$\partial_t \mathbf{x} = -H\mathbf{n}, \quad \mathbf{x}(0) = \mathbf{x}_0,$$

which is called a mean curvature flow with the maximal time  $T \leq \infty$ . When  $\Sigma$  is compact,  $T < \infty$  by the comparison principle for mean curvature flow and  $\sup_{\Sigma_t} |A| \rightarrow \infty$  as  $t \rightarrow T$  by Huisken in [4].

Since the finite-time singularity for a compact mean curvature flow is characterized by the blowup of the second fundamental form, it is of great interest to express this criterion in terms of some simpler quantity. Let  $n \geq 2$ . A natural conjecture is the blowup of the mean curvature  $H$ , which was confirmed in dimension two by Li-Wang [6]. In [2] Cooper proved the  $HA$  tensor blows up at time  $T$ . In [2, 7, 9], Cooper and Le-Sesum proved that the mean curvature blows up assuming some blowup rate of the second fundamental form. Some extension results under integral conditions can be seen in Le-Sesum [8] and Xu-Ye-Zhao [16].

Similar blowup problems and extension problems have been studied for Ricci flow as well. In [3] Hamilton proved that the Riemann curvature tensor blows up at the finite singular time. In [13] Sesum proved the blowup of the Ricci curvature. In [15, 1, 5], Wang, Chen-Wang and Kotschwar-Munteanu-Wang arrived at estimates on curvature growth in terms of the Ricci curvature.

The explicit local estimate in Kotschwar-Munteanu-Wang [5] has some precedent on a gradient shrinking soliton in [11], a bound on  $Ric$  implies a polynomial growth bound on  $Rm$ . The feasibility lies in the observation that the second order derivatives of  $Ric$  appear as time-derivative of  $Rm$ , i.e.,

$$\partial_t Rm = c\nabla^2 Ric,$$

which helps to yield a differential inequality on spatial integrations. This equation follows from the fact that  $Ric$  describes the metric evolution along a Ricci flow. Now, for a mean curvature flow  $HA$  describes the metric evolution and plays the role of  $Ric$  by

$$\partial_t |A|^2 = 2(\nabla^2 H \cdot A + H|A|^3).$$

In the present paper, we follow the techniques on integration estimates from [5] and establish the following local  $L^\infty$  estimate of  $A$  in terms of the initial geometry and the  $|HA|$  bound along the flow.

**Theorem 1.1.** (Theorem 4.2) Fix  $x_0 \in \mathbb{R}^{n+1}$  and  $r > 0$ . Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a complete smooth mean curvature flow satisfying the uniform bound

$$\sup_{B(x_0, r) \cap \Sigma_t} |HA|(\cdot, t) \leq K(t), \quad \forall t \in [t_0, t_1].$$

Then for any  $q > n + 2$  there exist positive constants  $C = C(n, r, t_1 - t_0, q, K(t_0))$  and  $c = c(n, q)$  such that for any  $t \in [t_0, t_1]$

$$\sup_{B(x_0, r/2) \cap \Sigma_t} |A| \leq C \left(1 + \|A\|_{L^q(B_{2r})}^q\right)^c \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^c \left(\int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + cK) ds} ds\right)^c,$$

where  $B_{2r} = B(x_0, 2r + n^{1/4} \int_{t_0}^t \sqrt{K}) \cap \Sigma_{t_0}$ .

This result provides a new proof of the blowup of  $HA$  in [2] and extends the estimates for the Ricci flow in terms of  $Ric$  in [15, 1, 5] to an estimate for the mean curvature flow in terms of  $HA$ . One of its direct corollaries is the following extension theorem in terms of  $HA$ .

**Theorem 1.2.** (Corollary 4.3) Let  $\mathbf{x} : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow. Each time slice  $\Sigma_t$  has bounded  $HA$ . There exists a positive constant  $C = C(n, T, K, V, E, q)$  such that if

(1) the  $|HA|$  bound satisfies

$$\sup_{t \in [0, T)} \sup_{\Sigma_t} |HA|(\cdot, t) \leq K < \infty;$$

(2) the initial data satisfies a uniform volume bound

$$\sup_{x \in \Sigma_0} \text{Vol}_{g(0)}(B(x, 1 + n^{1/4} T \sqrt{K}) \cap \Sigma_0) \leq V < \infty;$$

(3) the initial data satisfies an integral bound

$$\sup_{x \in \Sigma_0} \|A\|_{L^q(B(x,1) \cap \Sigma_0)} \leq E < \infty$$

for some  $q > n + 2$ ,

then

$$\limsup_{t \rightarrow T} \sup_{\Sigma_t} |A|(\cdot, t) \leq C < \infty.$$

In particular, the flow can be extended past time  $T$ .

Combining the  $L^\infty$  estimate and the blowup estimate of  $A$  in [4], we also obtain the following estimate of the blowup rate of  $HA$  at the first finite singularity. This result generalizes Theorem 1.2 of [9] and Theorem 5.1 of [2] and can be seen as another version of Theorem 1.1 of [15] and Theorem 2 of [5].

**Theorem 1.3.** (Theorem 5.1) Let  $\mathbf{x} : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a complete smooth mean curvature flow with a finite maximal time  $T$ . Each time slice  $\Sigma_t$  has bounded second fundamental form. Then there exists a positive constant  $\epsilon = \epsilon(n)$  such that

$$\limsup_{t \rightarrow T} \left( |T - t| \sup_{\Sigma_t} |HA| \right) \geq \epsilon.$$

The organization of this paper is as follows. In Sect.2 we recall some basic results on mean curvature flow. In Sect.3 we develop  $L^p$  estimate in terms of initial data and  $|HA|$  bound, following the argument in [5]. In Sect.4 we establish the  $L^\infty$  estimate by Moser iteration as in [8] and then derive the extension theorem. In Sect.5 we estimate the blowup rate of  $HA$ , using the  $L^\infty$  estimate and the blowup estimate of  $A$ .

**Acknowledgements:** The author would like to thank H.Z.Li for insightful discussions.

## 2 Preliminaries

Let  $\mathbf{x}(p, t) : \Sigma^n \rightarrow \mathbb{R}^{n+1}$  be a family of smooth immersions.  $\{(\Sigma^n, \mathbf{x}(\cdot, t)), 0 \leq t < T\}$  is called a mean curvature flow if  $\mathbf{x}$  satisfies

$$\partial_t \mathbf{x} = -H \mathbf{n}, \quad \forall t \in [0, T), \quad (2.1)$$

where  $A = (h_{ij})$  denotes the second fundamental form and  $H = g^{ij} h_{ij}$  denotes the mean curvature.  $\Sigma_t := \mathbf{x}_t(\Sigma)$  denotes the time slice of the flow for  $t \in [0, T)$ .

Some equations are listed here for later calculations. See [4] or [10] for details.

**Lemma 2.1.** (Sect.3 of [4]) Along the mean curvature flow,

$$\begin{aligned} \partial_t d\mu &= -H^2 d\mu, \\ \partial_t |A|^2 &= 2(\nabla^2 H \cdot A + H|A^3|), \end{aligned} \quad (2.2)$$

$$2|\nabla H|^2 = (\Delta - \partial_t)H^2 + 2H^2|A|^2, \quad (2.3)$$

$$2|\nabla A|^2 = (\Delta - \partial_t)|A|^2 + 2|A|^4. \quad (2.4)$$

where

$$\nabla^2 H \cdot A = H_{ij} A_{ij}, \quad |A^3| = A_{ij} A_{jk} A_{ki}.$$

Observe that the equation (2.1) is invariant under the rescaling

$$\tilde{\mathbf{x}}(p, t) = \lambda \left( \mathbf{x}(p, t_0 + \frac{t}{\lambda^2}) - x_0 \right), \quad \forall (p, t) \in \Sigma \times [-\lambda^2 t_0, \lambda^2 T - \lambda^2 t_0],$$

for some  $(x_0, t_0) \in \mathbb{R}^{n+1} \times [0, T)$  and  $\lambda > 0$ . Under the rescaling we have

$$\tilde{A} = \lambda A, \quad \tilde{g} = \lambda^2 g, \quad |\tilde{A}| = \lambda^{-1} |A|, \quad \tilde{H} = \lambda^{-1} H,$$

where  $\tilde{A}$ ,  $\tilde{g}$  and  $\tilde{H}$  denote the second fundamental form, the induced metric and the mean curvature of the hypersurface  $\tilde{\mathbf{x}}(\Sigma)$  respectively. Moreover, we have

**Lemma 2.2.** *Under the rescaling,*

$$\begin{aligned} \text{Vol}_{\tilde{g}}(B(x_0, \tilde{r}) \cap \tilde{\Sigma}_{\tilde{t}}) &= \lambda^n \text{Vol}_g(B(x_0, r) \cap \Sigma_t), \\ \int_{B(x_0, \tilde{r}) \cap \tilde{\Sigma}_{\tilde{t}}} |\tilde{A}|^p d\tilde{\mu} &= \lambda^{n-p} \int_{B(x_0, r) \cap \Sigma_t} |A|^p d\mu, \\ \int_{\tilde{t}_1}^{\tilde{t}_2} \int_{B(x_0, \tilde{r}) \cap \tilde{M}_{\tilde{t}}} |\tilde{A}|^p d\tilde{\mu} d\tilde{t} &= \lambda^{n+2-p} \int_{t_1}^{t_2} \int_{B(x_0, r) \cap \Sigma_t} |A|^p d\mu dt, \end{aligned}$$

where  $\tilde{r} = \lambda r$  and  $\tilde{t} = t_0 + t/\lambda^2$ .

The standard Moser iteration in Sect.4 depends on the Michael-Simon inequality. See [4, 12].

**Lemma 2.3.** (Lemma 5.7 of [4]) *For any Lipschitz function  $f$  with compact support on a hypersurface  $\Sigma^n \subset \mathbb{R}^{n+1}$ ,*

$$\left( \int_{\Sigma} |f|^{\frac{n}{n-1}} d\mu \right)^{\frac{n-1}{n}} \leq c_n \int_{\Sigma} (|\nabla f| + H|f|) d\mu. \quad (2.5)$$

By maximum principle the second fundamental form blows up at least at a rate of  $1/2$ , which holds for noncompact cases as well.

**Lemma 2.4.** (Proposition 2.4.6 of [10]) *Suppose each time slice  $\Sigma_t$  has bounded second fundamental form. Then*

$$\sup_{\Sigma_t} |A| \geq \frac{1}{\sqrt{2(T-t)}}.$$

### 3 $L^p$ estimate

Throughout this paper  $c$  denotes nonnegative constants depending only on  $n$  and  $p$  while  $c_n$  denotes nonnegative constants depending only on  $n$ , which may change from line to line.

**Theorem 3.1.** *Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a closed smooth mean curvature flow satisfying the uniform bound*

$$\sup_{\Sigma_t} |HA|(\cdot, t) \leq K(t), \quad \forall t \in [t_0, t_1]. \quad (3.1)$$

*Then for any  $p \geq 4$  there exist positive constants  $c = c(n, p)$  such that for any  $t \in [t_0, t_1]$ ,*

$$\int_{\Sigma_t} |A|^p \leq \left( \int_{\Sigma_{t_0}} |A|^p(t_0) + cK(t_0) \int_{\Sigma_{t_0}} |A|^{p-2}(t_0) \right) e^{\int_{t_0}^t (|K'|/K + cK)}.$$

*Proof.* Using equation (2.2) from Lemma 2.1, we get

$$\begin{aligned}
\partial_t \int_{\Sigma_t} |A|^p &= p \int_{\Sigma_t} |A|^{p-2} (\nabla^2 HA + H|A^3|) - \int_{\Sigma_t} |A|^p H^2 \\
&\leq p(p-1) \int_{\Sigma_t} |A|^{p-2} |\nabla H| |\nabla A| + p \int_{\Sigma_t} |H| |A|^{p+1} \\
&\leq \frac{1}{4K} \int_{\Sigma_t} |A|^p |\nabla H|^2 + p^4 K \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 + pK \int_{\Sigma_t} |A|^p. \tag{3.2}
\end{aligned}$$

Using equation (2.3), we get

$$\begin{aligned}
2 \int_{\Sigma_t} |A|^p |\nabla H|^2 &= \int_{\Sigma_t} |A|^p (\Delta - \partial_t) H^2 + 2 \int_{\Sigma_t} H^2 |A|^{p+2} \\
&\leq 2p \int_{\Sigma_t} |H| |A|^{p-1} |\nabla H| |\nabla A| - \partial_t \int_{\Sigma_t} H^2 |A|^p + \int_{\Sigma_t} H^2 \partial_t (|A|^p) \\
&\quad - \int_{\Sigma_t} H^4 |A|^p + 2 \int_{\Sigma_t} H^2 |A|^{p+2}.
\end{aligned}$$

Since

$$\begin{aligned}
2p \int_{\Sigma_t} |H| |A|^{p-1} |\nabla H| |\nabla A| &\leq \int_{\Sigma_t} |A|^p |\nabla H|^2 + p^2 \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla A|^2, \\
\partial_t (|A|^p) &= p |A|^{p-2} (\nabla^2 HA + H|A^3|), \\
p |H|^3 |A|^{p+1} &\leq H^4 |A|^p + \frac{p^2}{4} H^2 |A|^{p+2},
\end{aligned}$$

we have

$$\begin{aligned}
\int_{\Sigma_t} |A|^p |\nabla H|^2 &\leq p^2 \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla A|^2 - \partial_t \int_{\Sigma_t} H^2 |A|^p + p \int_{\Sigma_t} H^2 |A|^{p-2} \nabla^2 HA \\
&\quad + p \int_{\Sigma_t} |H|^3 |A|^{p+1} - \int_{\Sigma_t} H^4 |A|^p + 2 \int_{\Sigma_t} H^2 |A|^{p+2} \\
&\leq -\partial_t \int_{\Sigma_t} H^2 |A|^p + p^2 K^2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 + p \int_{\Sigma_t} H^2 |A|^{p-2} \nabla^2 HA \\
&\quad + \frac{p^2}{2} K^2 \int_{\Sigma_t} |A|^p. \tag{3.3}
\end{aligned}$$

For the third term on the right of (3.3),

$$\begin{aligned}
p \int_{\Sigma_t} H^2 |A|^{p-2} \nabla^2 HA &\leq p(p-1) \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla H| |\nabla A| + 2p \int_{\Sigma_t} |H| |A|^{p-1} |\nabla H|^2 \\
&\leq \frac{1}{2} \int_{\Sigma_t} |A|^p |\nabla H|^2 + \frac{p^4}{2} \int_{\Sigma_t} H^4 |A|^{p-4} |\nabla A|^2 \\
&\quad + \frac{1}{4} \int_{\Sigma_t} |A|^p |\nabla H|^2 + 4p^2 \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla H|^2 \\
&\leq \frac{3}{4} \int_{\Sigma_t} |A|^p |\nabla H|^2 + \left(\frac{np^4}{2} + 4np^2\right) \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla A|^2 \\
&\leq \frac{3}{4} \int_{\Sigma_t} |A|^p |\nabla H|^2 + np^2 \left(\frac{p^2}{2} + 4\right) K^2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2. \tag{3.4}
\end{aligned}$$

Substituting (3.4) into (3.3), we get

$$\frac{1}{4} \int_{\Sigma_t} |A|^p |\nabla H|^2 \leq -\partial_t \int_{\Sigma_t} H^2 |A|^p + np^4 K^2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 + \frac{p^2}{2} K^2 \int_{\Sigma_t} |A|^p.$$

Back to (3.2), we obtain

$$\partial_t \int_{\Sigma_t} |A|^p \leq -\frac{1}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p + (n+1)p^4 K \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 + \left(\frac{p^2}{2} + p\right) K \int_{\Sigma_t} |A|^p.$$

By equation (2.4) from Lemma 2.1, for  $p \geq 4$ ,

$$\begin{aligned} 2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 &= \int_{\Sigma_t} |A|^{p-4} (\Delta - \partial_t) |A|^2 + 2 \int_{\Sigma_t} |A|^p \\ &= -2(p-4) \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 - \frac{2}{p-2} \int_{\Sigma_t} \partial_t (|A|^{p-2}) + 2 \int_{\Sigma_t} |A|^p, \end{aligned}$$

i.e.,

$$\int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 = -\frac{1}{(p-2)(p-3)} \partial_t \int_{\Sigma_t} |A|^{p-2} + \frac{1}{p-3} \int_{\Sigma_t} |A|^p.$$

Hence we have

$$\begin{aligned} &\partial_t \int_{\Sigma_t} |A|^p \\ &\leq -\frac{1}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p - \frac{(n+1)p^4}{(p-2)(p-3)} K \partial_t \int_{\Sigma_t} |A|^{p-2} + \left(\frac{(n+1)p^4}{p-3} + \frac{p^2}{2} + p\right) K \int_{\Sigma_t} |A|^p \\ &\leq -\frac{1}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p - cK \partial_t \int_{\Sigma_t} |A|^{p-2} + cK \int_{\Sigma_t} |A|^p. \end{aligned} \quad (3.5)$$

If we set

$$U(t) = \int_{\Sigma_t} |A|^p + \frac{1}{K} \int_{\Sigma_t} H^2 |A|^p + cK \int_{\Sigma_t} |A|^{p-2},$$

then actually (3.5) becomes

$$\begin{aligned} U' &\leq -\frac{K'}{K^2} \int_{\Sigma_t} H^2 |A|^p + cK' \int_{\Sigma_t} |A|^{p-2} + cK \int_{\Sigma_t} |A|^p \\ &\leq (|K'|/K + cK)U, \end{aligned}$$

which implies

$$U(t) \leq e^{\int_{t_0}^t (|K'|/K + cK)} U(t_0).$$

In particular,

$$\int_{\Sigma_t} |A|^p \leq \left( \int_{\Sigma_{t_0}} |A|^p(t_0) + cK(t_0) \int_{\Sigma_{t_0}} |A|^{p-2}(t_0) \right) e^{\int_{t_0}^t (|K'|/K + cK)}.$$

□

**Corollary 3.2.** *Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a closed smooth mean curvature flow satisfying the uniform bound*

$$\sup_{s \in [t_0, t]} \sup_{\Sigma_s} |HA|(\cdot, s) \leq K(t), \quad \forall t \in [t_0, t_1].$$

*Then for any  $p \geq 4$  there exist positive constants  $c = c(n, p)$  such that for any  $t \in [t_0, t_1]$*

$$\begin{aligned} \int_{\Sigma_t} |A|^p &\leq \left( \int_{\Sigma_{t_0}} |A|^p(t_0) + cK(t) \int_{\Sigma_{t_0}} |A|^{p-2}(t_0) \right) e^{c(t-t_0)K(t)} \\ &\leq \text{Vol}_{g(t_0)}(\Sigma_{t_0}) \left( (\sup_{\Sigma_{t_0}} |A|)^p + cK(t) (\sup_{\Sigma_{t_0}} |A|)^{p-2} \right) e^{c(t-t_0)K(t)}. \end{aligned}$$

*Proof.* Fix  $\tilde{t} \in [t_0, t_1]$  and  $\tilde{K} := K(\tilde{t})$ . Consider instead the rescaled flow  $\tilde{\mathbf{x}} : \Sigma \times [\tilde{K}t_0, \tilde{K}\tilde{t}] \rightarrow \mathbb{R}^{n+1}$  with  $\tilde{\mathbf{x}}(\cdot, t) = \sqrt{\tilde{K}} \mathbf{x}(\cdot, t/\tilde{K})$ . One sees

$$\sup_{\tilde{\Sigma}_t} |\tilde{H}\tilde{A}| \leq 1, \quad \forall t \in [\tilde{K}t_0, \tilde{K}\tilde{t}].$$

Provided the uniform bound  $\hat{K} \equiv 1$ , applying Theorem 3.1 yields

$$\begin{aligned} \tilde{K}^{\frac{n-p}{2}} \int_{\Sigma_{\tilde{t}}} |A|^p &= \int_{\tilde{\Sigma}_{\tilde{K}\tilde{t}}} |\tilde{A}|^p \\ &\leq \left( \int_{\tilde{\Sigma}_{\tilde{K}t_0}} |\tilde{A}|^p(\tilde{K}t_0) + c \int_{\tilde{\Sigma}_{\tilde{K}t_0}} |\tilde{A}|^{p-2}(\tilde{K}t_0) \right) e^{c\tilde{K}(\tilde{t}-t_0)} \\ &= \left( \tilde{K}^{\frac{n-p}{2}} \int_{\Sigma_{t_0}} |A|^p(t_0) + c\tilde{K}^{\frac{n+2-p}{2}} \int_{\Sigma_{t_0}} |A|^{p-2}(t_0) \right) e^{c\tilde{K}(\tilde{t}-t_0)}, \end{aligned}$$

which implies

$$\int_{\Sigma_{\tilde{t}}} |A|^p \leq \left( \int_{\Sigma_{t_0}} |A|^p(t_0) + c\tilde{K} \int_{\Sigma_{t_0}} |A|^{p-2}(t_0) \right) e^{c\tilde{K}(\tilde{t}-t_0)}.$$

Since the choice of  $\tilde{t}$  is arbitrary, we actually obtain the result.  $\square$

By the use of some cutoff function, Theorem 3.1 can be generalized to a local version.

**Theorem 3.3.** *Fix  $x_0 \in \mathbb{R}^{n+1}$  and  $r > 0$ . Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow satisfying the uniform bound*

$$\sup_{B(x_0, r) \cap \Sigma_t} |HA|(\cdot, s) \leq K(t), \quad \forall t \in [t_0, t_1].$$

*Then for any  $p \geq 4$  there exist positive constants  $c = c(n, p)$  such that for any  $t \in [t_0, t_1]$*

$$\int_{B(x_0, r/2) \cap \Sigma_t} |A|^p \leq \left( \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^p(t_0) + cK(t_0) \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^{p-2}(t_0) + r^{-p} \text{Vol}_{g(t_0)}(B_r) \right) \cdot e^{\int_{t_0}^t (|K'|/K + cK)},$$

where  $B_r = B(x_0, r + n^{1/4} \int_{t_0}^t \sqrt{K}) \cap \Sigma_{t_0}$ .

*Proof.* Note that  $c$  always denotes nonnegative constants depending on  $n$  and  $p$ , which may vary from line to line. Let  $\phi(x, t)$  be a smooth function with compact support which will be determined later. Following the steps in the proof of Theorem 3.1, we have

$$\begin{aligned}
\partial_t \int_{\Sigma_t} |A|^p \phi &\leq \frac{c}{K} \int_{\Sigma_t} |A|^p |\nabla H|^2 \phi + cK \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi + cK \int_{\Sigma_t} |A|^p \phi \\
&\quad + c \int_{\Sigma_t} |A|^{p-1} |\nabla H| |\nabla \phi| + \int_{\Sigma_t} |A|^p \partial_t \phi \\
&\leq \frac{c}{K} \int_{\Sigma_t} |A|^p |\nabla H|^2 \phi + cK \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi + cK \int_{\Sigma_t} |A|^p \phi \\
&\quad + cK \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + \int_{\Sigma_t} |A|^p \partial_t \phi,
\end{aligned}$$

$$\begin{aligned}
\int_{\Sigma_t} |A|^p |\nabla H|^2 \phi &\leq -\partial_t \int_{\Sigma_t} H^2 |A|^p \phi + c \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla A|^2 \phi + c \int_{\Sigma_t} H^2 |A|^{p+2} \phi \\
&\quad + 2 \int_{\Sigma_t} |H| |A|^p |\nabla H| |\nabla \phi| + p \int_{\Sigma_t} H^2 |A|^{p-1} |\nabla H| |\nabla \phi| + \int_{\Sigma_t} H^2 |A|^p \partial_t \phi \\
&\leq -\partial_t \int_{\Sigma_t} H^2 |A|^p \phi + c \int_{\Sigma_t} H^2 |A|^{p-2} |\nabla A|^2 \phi + c \int_{\Sigma_t} H^2 |A|^{p+2} \phi \\
&\quad + \frac{1}{2} \int_{\Sigma_t} |A|^p |\nabla H|^2 \phi + c \int_{\Sigma_t} H^2 |A|^p \phi^{-1} |\nabla \phi|^2 + \int_{\Sigma_t} H^2 |A|^p \partial_t \phi \\
&\leq -\partial_t \int_{\Sigma_t} H^2 |A|^p \phi + cK^2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi + cK^2 \int_{\Sigma_t} |A|^p \phi \\
&\quad + \frac{1}{2} \int_{\Sigma_t} |A|^p |\nabla H|^2 \phi + cK^2 \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + \int_{\Sigma_t} H^2 |A|^p \partial_t \phi,
\end{aligned}$$

$$\begin{aligned}
\frac{c}{K} \int_{\Sigma_t} |A|^p |\nabla H|^2 \phi &\leq -\frac{c}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p \phi + cK \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi + cK \int_{\Sigma_t} |A|^p \phi \\
&\quad + cK \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + \frac{c}{K} \int_{\Sigma_t} H^2 |A|^p \partial_t \phi,
\end{aligned}$$

$$\begin{aligned}
2 \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi &\leq -c \partial_t \int_{\Sigma_t} |A|^{p-2} \phi + 2 \int_{\Sigma_t} |A|^p \phi + 2 \int_{\Sigma_t} |A|^{p-3} |\nabla A| |\nabla \phi| \\
&\quad + c \int_{\Sigma_t} |A|^{p-2} \partial_t \phi \\
&\leq -c \partial_t \int_{\Sigma_t} |A|^{p-2} \phi + 2 \int_{\Sigma_t} |A|^p \phi + \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi \\
&\quad + c \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + c \int_{\Sigma_t} |A|^{p-2} \partial_t \phi,
\end{aligned}$$

$$\begin{aligned}
K \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi &\leq -cK \partial_t \int_{\Sigma_t} |A|^{p-2} \phi + 2K \int_{\Sigma_t} |A|^p \phi + cK \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 \\
&\quad + cK \int_{\Sigma_t} |A|^{p-2} \partial_t \phi,
\end{aligned}$$



$$\begin{aligned}
\partial_t \int_{\Sigma_t} |A|^p \phi &\leq -\frac{c}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p \phi + cK \int_{\Sigma_t} |A|^{p-4} |\nabla A|^2 \phi + cK \int_{\Sigma_t} |A|^p \phi \\
&\quad + cK \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + \frac{c}{K} \int_{\Sigma_t} H^2 |A|^p \partial_t \phi + \int_{\Sigma_t} |A|^p \partial_t \phi \\
&\leq -\frac{c}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p \phi - cK \partial_t \int_{\Sigma_t} |A|^{p-2} \phi + cK \int_{\Sigma_t} |A|^p \phi \\
&\quad + cK \int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 + \int_{\Sigma_t} |A|^p |\partial_t \phi| + cK \int_{\Sigma_t} |A|^{p-2} |\partial_t \phi|. \quad (3.6)
\end{aligned}$$

Consider a smooth decreasing function  $\eta$ , which equals 1 on  $[0, r/2]$  and vanishes on  $[r, \infty)$ , satisfying  $|\eta'| \leq 3/r$ . For any  $0 < \delta < 1$  we set  $\psi := \eta^{1/\delta}$  such that

$$|\psi'| \leq \frac{3}{\delta r} \psi^{1-\delta}.$$

Now we choose  $\phi := \psi(|x - x_0|)$ . Then

$$\phi^{-1} |\nabla \phi|^2 \leq n \psi^{-1} (\psi')^2 \leq \frac{9n}{\delta^2 r^2} \phi^{1-2\delta},$$

$$|\partial_t \phi| = |\psi'| |\partial_t |x - x_0|| \leq |\psi'| |H| \leq \frac{3}{\delta r} |H| \phi^{1-\delta}.$$

Take  $\delta = \frac{1}{p}$ . By Young's inequality,

$$\begin{aligned}
\int_{\Sigma_t} |A|^{p-2} \phi^{-1} |\nabla \phi|^2 &\leq \frac{c}{\delta^2 r^2} \int_{\Sigma_t} |A|^{p-2} \phi^{1-2\delta} \leq c \int_{\Sigma_t} |A|^p \phi + \frac{c}{\delta^p r^p} \int_{\Sigma_t} \phi^{1-p\delta} \\
&\leq c \int_{\Sigma_t} |A|^p \phi + \frac{c}{r^p} \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t),
\end{aligned}$$

$$\begin{aligned}
\int_{\Sigma_t} |A|^{p-2} |\partial_t \phi| &\leq \frac{c}{\delta r} \int_{\Sigma_t} |A|^{p-1} \phi^{1-\delta} \leq c \int_{\Sigma_t} |A|^p \phi + \frac{c}{\delta^p r^p} \int_{\Sigma_t} \phi^{1-p\delta} \\
&\leq c \int_{\Sigma_t} |A|^p \phi + \frac{c}{r^p} \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t),
\end{aligned}$$

$$\begin{aligned}
\int_{\Sigma_t} |A|^p |\partial_t \phi| &\leq \frac{c}{\delta r} \int_{\Sigma_t} |H| |A|^p \phi^{1-\delta} \leq \frac{c}{\delta r} K \int_{\Sigma_t} |A|^{p-1} \phi^{1-\delta} \\
&\leq cK \int_{\Sigma_t} |A|^p \phi + \frac{c}{r^p} K \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t).
\end{aligned}$$

Back to (3.6), we obtain

$$\begin{aligned}
&\partial_t \int_{\Sigma_t} |A|^p \phi + \frac{c}{K} \partial_t \int_{\Sigma_t} H^2 |A|^p \phi + cK \partial_t \int_{\Sigma_t} |A|^{p-2} \phi \\
&\leq cK \int_{\Sigma_t} |A|^p \phi + \frac{cK}{r^p} \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t). \quad (3.7)
\end{aligned}$$

If we set

$$U(t) = \int_{\Sigma_t} |A|^p \phi + \frac{c}{K} \int_{\Sigma_t} H^2 |A|^p \phi + cK \int_{\Sigma_t} |A|^{p-2} \phi,$$

then actually it becomes

$$\begin{aligned} U' &\leq -\frac{cK'}{K^2} \int_{\Sigma_t} H^2 |A|^p \phi + cK' \int_{\Sigma_t} |A|^{p-2} \phi + cK \int_{\Sigma_t} |A|^p \phi + \frac{cK}{r^p} \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t) \\ &\leq (|K'|/K + cK)U + cr^{-p}K \text{Vol}_{g(t)}(B(x_0, r) \cap \Sigma_t). \end{aligned}$$

Since

$$\partial_t |x - x_0| \leq |H| \leq n^{1/4} \sqrt{K}$$

and

$$\partial_t d\mu = -H^2 d\mu,$$

we know

$$\begin{aligned} B(x_0, r) \cap \Sigma_t &\subset B(x_0, r + n^{1/4} \int_{t_0}^t \sqrt{K}) \cap \Sigma_{t_0} := B_r, \\ \text{Vol}_{g(s)}(B(x_0, r) \cap \Sigma_s) &\leq \text{Vol}_{g(t_0)}(B_r), \quad \forall s \in [t_0, t]. \end{aligned}$$

Then for any  $s \in [t_0, t]$ ,

$$\begin{aligned} \partial_s \left( e^{-\int_{t_0}^s (|K'|/K + cK)} U(s) \right) &\leq cr^{-p}K e^{-\int_{t_0}^s (|K'|/K + cK)} \text{Vol}_{g(t_0)}(B_r) \\ &\leq r^{-p}(cK + |K'|/K) e^{-\int_{t_0}^s (|K'|/K + cK)} \text{Vol}_{g(t_0)}(B_r) \\ &= \partial_s \left( -e^{-\int_{t_0}^s (|K'|/K + cK)} \right) r^{-p} \text{Vol}_{g(t_0)}(B_r), \end{aligned}$$

i.e.,

$$\partial_s \left( e^{-\int_{t_0}^s (|K'|/K + cK)} (U(s) + r^{-p} \text{Vol}_{g(t_0)}(B_r)) \right) \leq 0,$$

which implies

$$U(s) \leq e^{\int_{t_0}^s (|K'|/K + cK)} \left( U(t_0) + r^{-p} \text{Vol}_{g(t_0)}(B_r) \right), \quad \forall s \in [t_0, t]. \quad (3.8)$$

In particular,

$$\int_{B(x_0, r/2) \cap \Sigma_t} |A|^p \leq \left( \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^p(t_0) + cK(t_0) \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^{p-2}(t_0) + r^{-p} \text{Vol}_{g(t_0)}(B_r) \right) \cdot e^{\int_{t_0}^t (|K'|/K + cK)}.$$

□

**Corollary 3.4.** Fix  $x_0 \in \mathbb{R}^{n+1}$  and  $r > 0$ . Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow satisfying the uniform bound

$$\sup_{s \in [t_0, t]} \sup_{B(x_0, r) \cap \Sigma_s} |HA|(\cdot, s) \leq K(t), \quad \forall t \in [t_0, t_1].$$

Then for any  $p \geq 4$  there exist positive constants  $c = c(n, p)$  such that for any  $t \in [t_0, t_1]$

$$\int_{B(x_0, r/2) \cap \Sigma_t} |A|^p \leq \left( \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^p(t_0) + cK(t) \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^{p-2}(t_0) + r^{-p} \text{Vol}_{g(t_0)}(\tilde{B}_r) \right) \cdot e^{c(t-t_0)K(t)},$$

where  $\tilde{B}_r = B(x_0, r + n^{1/4}(t - t_0)\sqrt{K(t)}) \cap \Sigma_{t_0}$ .

*Proof.* As in the proof of Theorem 3.2, by rescaling argument and Theorem 3.1 one has

$$\begin{aligned} K^{\frac{n-p}{2}} \int_{B(x_0, \frac{r}{2}) \cap \Sigma_t} |A|^p &= \int_{B(x_0, \frac{\sqrt{K}r}{2}) \cap \tilde{\Sigma}_{Kt}} |\tilde{A}|^p \\ &\leq \left( \int_{B(x_0, \sqrt{K}r) \cap \tilde{\Sigma}_{Kt_0}} |\tilde{A}|^p(Kt_0) + c \int_{B(x_0, \sqrt{K}r) \cap \tilde{\Sigma}_{Kt_0}} |\tilde{A}|^{p-2}(Kt_0) + (\sqrt{K}r)^{-p} \text{Vol}_{\tilde{g}(Kt_0)}(\tilde{B}) \right) e^{c(t-t_0)K} \\ &\leq \left( K^{\frac{n-p}{2}} \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^p(t_0) + cK^{\frac{n+2-p}{2}} \int_{B(x_0, r) \cap \Sigma_{t_0}} |A|^{p-2}(t_0) + K^{\frac{n-p}{2}} r^{-p} \text{Vol}_{g(t_0)}(\tilde{B}) \right) e^{c(t-t_0)K}, \end{aligned}$$

where

$$\begin{aligned} \tilde{B} &= B(x_0, \sqrt{K}r + n^{1/4}(t - t_0)K) \cap \tilde{\Sigma}_{Kt_0} \\ &= B(x_0, r + n^{1/4}(t - t_0)\sqrt{K}) \cap \Sigma_{t_0}. \end{aligned}$$

Thus we obtain the result.  $\square$

## 4 $L^\infty$ estimate and extension theorem

From  $L^p$  estimate to  $L^\infty$  estimate we require the process of Moser iteration using the Michael-Simon inequality, i.e, Lemma 2.3, which relies on mean curvature. We conclude the following result from Lemma 5.2 in [8].

**Lemma 4.1 (Moser iteration).** *(Lemma 5.2 of [8]) Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a smooth mean curvature flow. Consider the differential inequality*

$$(\partial_t - \Delta)v \leq fv, \quad v \geq 0.$$

Fix  $x_0 \in \mathbb{R}^{n+1}$  and  $r > 0$ . For any  $q > n + 2$  and  $\beta \geq 2$  there exists a constant  $C = C(n, r, t_1 - t_0, q, \beta)$  such that for any  $t \in [t_0, t_1]$

$$\|v\|_{L^\infty(D'_{t,r})} \leq C \|f\|_{L^{q/2}(D_{t,r})}^{\frac{qn^2}{\beta(q-n-2)}} \left(1 + \|H\|_{L^{n+2}(D_{t,r})}^{n+2}\right)^{\frac{qn^3}{\beta(n+2)(q-n-2)}} \|v\|_{L^\beta(D_{t,r})},$$

where

$$\begin{aligned} D_{t,r} &:= \bigcup_{t_0 \leq s \leq t} (B(x_0, r) \cap \Sigma_s), \\ D'_{t,r} &:= \bigcup_{(t_0+t)/2 \leq s \leq t} (B(x_0, r/2) \cap \Sigma_s). \end{aligned}$$

Combining Theorem 3.3 and Lemma 4.1 we obtain the following local estimate.

**Theorem 4.2** ( $L^\infty$  estimate). *Fix  $x_0 \in \mathbb{R}^{n+1}$  and  $r > 0$ . Let  $\mathbf{x} : \Sigma^n \times [t_0, t_1] \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow satisfying the uniform bound*

$$\sup_{B(x_0, r) \cap \Sigma_t} |HA|(\cdot, t) \leq K(t), \quad \forall t \in [t_0, t_1].$$

*Then for any  $q > n + 2$  there exist positive constants  $C = C(n, r, t_1 - t_0, q, K(t_0))$  and  $c = c(n, q)$  such that for any  $t \in [t_0, t_1]$*

$$\sup_{D'_{t,r}} |A| \leq C \left(1 + \|A\|_{L^q(B_{2r})}^q\right)^c \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^c \left(\int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + cK)} ds\right)^{\frac{1}{q} + \frac{n^2}{q-n-2}},$$

*where  $B_{2r} = B(x_0, 2r + n^{1/4} \int_{t_0}^t \sqrt{K}) \cap \Sigma_{t_0}$ . In particular, there exist positive constants  $C_0 = C_0(n, r, t_1 - t_0, \Sigma_0, \int_{t_0}^{t_1} \sqrt{K})$  and  $c_n$  such that for any  $t \in [t_0, t_1]$*

$$\sup_{B(x_0, r/2) \cap \Sigma_t} |A| \leq C_0 \left(\int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + c_n K)} ds\right)^{c_n} \leq C_0 e^{c_n \int_{t_0}^t (|K'|/K + K)}.$$

*Proof.* Take  $\beta = \frac{n+2}{2}$ . Applying Lemma 4.1 to

$$(\partial_t - \Delta)|A|^2 = -2|\nabla A|^2 + 2|A|^4 \leq 2|A|^2 \cdot |A|^2$$

yields

$$\begin{aligned} \sup_{D'_{t,r}} |A| &\leq C' \|A\|_{L^q(D_{t,r})}^{\frac{qn^2}{\beta(q-n-2)}} \left(1 + \|A\|_{L^{n+2}(D_{t,r})}^{n+2}\right)^{\frac{qn^3}{2\beta(n+2)(q-n-2)}} \|A\|_{L^{2\beta}(D_{t,r})} \\ &\leq C' \|A\|_{L^q(D_{t,r})}^{\frac{2qn^2}{(n+2)(q-n-2)}} \left(1 + \|A\|_{L^{n+2}(D_{t,r})}^{1 + \frac{qn^3}{(n+2)(q-n-2)}}\right) \\ &\leq C' \|A\|_{L^q(D_{t,r})}^{\frac{2qn^2}{(n+2)(q-n-2)}} \left(1 + \|A\|_{L^q(D_{t,r})}^{1 + \frac{qn^3}{(n+2)(q-n-2)}} \text{Vol}(D_{t,r})^{\frac{q-n-2}{q(n+2)} + \frac{n^3}{(n+2)^2}}\right) \\ &\leq C' \|A\|_{L^q(D_{t,r})}^{\frac{2qn^2}{(n+2)(q-n-2)}} \left(1 + \|A\|_{L^q(D_{t,r})}^{1 + \frac{qn^3}{(n+2)(q-n-2)}} \text{Vol}_{g(t_0)}(B_{2r})^{\frac{q-n-2}{q(n+2)} + \frac{n^3}{(n+2)^2}}\right) \\ &\leq C' \left(1 + \|A\|_{L^q(D_{t,r})}^q\right)^{\frac{1}{q} + \frac{n^2}{q-n-2}} \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^{\frac{q-n-2}{q(n+2)} + \frac{n^3}{(n+2)^2}}, \end{aligned} \quad (4.1)$$

where

$$C' = C'(n, r, t_1 - t_0, q),$$

$$\text{Vol}(D_{t,r}) := \int_{t_0}^t \text{Vol}_{g(s)}(B(x_0, r) \cap \Sigma_s) ds,$$

$$B_{2r} = B(x_0, 2r + n^{1/4} \int_{t_0}^t \sqrt{K}) \cap \Sigma_{t_0} \leq B(x_0, 2r + n^{1/4} \int_{t_0}^{t_1} \sqrt{K}) \cap \Sigma_{t_0} := B'_{2r}.$$

It is derived from Theorem 3.3 that for  $q > n + 2$

$$\begin{aligned} \|A\|_{L^q(D_{t,r})}^q &\leq \left(\|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}^q + cK(t_0) \|A\|_{L^{q-2}(B(x_0, 2r) \cap \Sigma_{t_0})}^{q-2} + (2r)^{-q} \text{Vol}_{g(t_0)}(B_{2r})\right) \\ &\quad \cdot \int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + cK)} ds, \end{aligned}$$

where  $c = c(n, q)$ . Note that

$$\begin{aligned}
& \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}^q + cK(t_0) \|A\|_{L^{q-2}(B(x_0, 2r) \cap \Sigma_{t_0})}^{q-2} + (2r)^{-q} \text{Vol}_{g(t_0)}(B_{2r}) \\
& \leq \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}^q + cK(t_0) \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}^{q-2} \text{Vol}_{g(t_0)}(B_{2r})^{\frac{2}{q}} + (2r)^{-q} \text{Vol}_{g(t_0)}(B_{2r}) \\
& \leq C(n, r, q, K(t_0)) \left(1 + \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}\right)^q \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right).
\end{aligned}$$

The final coefficient is

$$\begin{aligned}
& C \left(1 + \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_0)}\right)^{q \left(\frac{1}{q} + \frac{n^2}{q-n-2}\right)} \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^{\frac{1}{q} + \frac{n^2}{q-n-2} + \frac{q-n-2}{q(n+2)} + \frac{n^3}{(n+2)^2}} \\
& \leq C \left(1 + \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}\right)^{1 + \frac{qn^2}{q-n-2}} \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^{\frac{1}{n+2} + \frac{n^2}{q-n-2} + \frac{n^3}{(n+2)^2}},
\end{aligned}$$

where  $C = C(n, r, t_1 - t_0, q, K(t_0))$ . Back to (4.1), we have the local  $L^\infty$  estimate

$$\sup_{D'_{t,r}} |A| \leq C \left(1 + \|A\|_{L^q(B(x_0, 2r) \cap \Sigma_{t_0})}\right)^c \left(1 + \text{Vol}_{g(t_0)}(B_{2r})\right)^c \left(\int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + cK) ds}\right)^{\frac{1}{q} + \frac{n^2}{q-n-2}}.$$

The second result follows from taking some  $q = q(n) > n + 2$ .  $\square$

As an application of the local estimate above, one immediately gets the following extension theorem about  $HA$ .

**Corollary 4.3 (HA-extension).** *Let  $\mathbf{x} : \Sigma^n \times [0, T] \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow. Each time slice  $\Sigma_t$  has bounded  $HA$ . There exists a positive constant  $C = C(n, T, K, V, E, q)$  such that if*

(1) *the  $|HA|$  bound satisfies*

$$\sup_{t \in [0, T]} \sup_{\Sigma_t} |HA|(\cdot, t) \leq K < \infty;$$

(2) *the initial data satisfies a uniform volume bound*

$$\sup_{x \in \Sigma_0} \text{Vol}_{g(0)}(B(x, 1 + n^{1/4} T \sqrt{K}) \cap \Sigma_0) \leq V < \infty;$$

(3) *the initial data satisfies an integral bound*

$$\sup_{x \in \Sigma_0} \|A\|_{L^q(B(x, 1) \cap \Sigma_0)} \leq E < \infty$$

for some  $q > n + 2$ ,

then

$$\limsup_{t \rightarrow T} \sup_{\Sigma_t} |A|(\cdot, t) \leq C < \infty.$$

In particular, the flow can be extended past time  $T$ .

*Proof.* It suffices to take  $r = 1$  in Theorem 4.2.  $\square$

## 5 Blowup estimate of $HA$

In this section we derive a blowup estimate of  $HA$  from Theorem 4.2 and Lemma 2.4, which also implies a blowup estimate of mean curvature.

**Theorem 5.1 (HA-blowup).** *Let  $\mathbf{x} : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow with a finite maximal time  $T$ . Each time slice  $\Sigma_t$  has bounded second fundamental form. Then there exists a positive constant  $\epsilon = \epsilon(n)$  such that*

$$\limsup_{t \rightarrow T} \left( |T - t| \sup_{\Sigma_t} |HA| \right) \geq \epsilon.$$

*Proof.* For otherwise one finds  $t_0 \in [0, T)$  and  $\epsilon > 0$  such that

$$\sup_{\Sigma_t} |HA| < \frac{\epsilon}{T - t}, \quad \forall t \in [t_0, T).$$

Actually we find a smooth mean curvature flow  $\mathbf{x} : \Sigma^n \times [t_0, T) \rightarrow \mathbb{R}^{n+1}$  with a  $|HA|$  bound

$$K(t) = \frac{\epsilon}{T - t}.$$

For  $t$  close to  $T$ ,

$$\int_{t_0}^t (|K'|/K + c_n K) = \int_{t_0}^t \frac{1 + c_n \epsilon}{T - s} ds = (1 + c_n \epsilon) \log\left(\frac{T - t_0}{T - t}\right),$$

$$\int_{t_0}^t e^{\int_{t_0}^s (|K'|/K + c_n K)} ds = \int_{t_0}^t \left(\frac{T - t_0}{T - s}\right)^{1 + c_n \epsilon} ds = \frac{(T - t_0)^{1 + c_n \epsilon}}{c_n \epsilon} \left( (T - t)^{-c_n \epsilon} - (T - t_0)^{-c_n \epsilon} \right).$$

On the other hand, by Lemma 2.4 we know

$$\sup_{\Sigma_t} |A| \geq \frac{1}{\sqrt{2}} (T - t)^{-\frac{1}{2}}.$$

Note that  $\Sigma_{t_0}$  has bounded geometry. Combining the inequalities and applying Theorem 4.2 imply that as  $t \rightarrow T$ ,

$$(T - t)^{-\frac{1}{2}} \leq C(n, T, \Sigma_{t_0}, \epsilon) (T - t)^{-c_n \epsilon}.$$

Letting  $c_n \epsilon < \frac{1}{2}$  causes a contradiction.  $\square$

Remark that Theorem 5.1 certainly works for the closed cases. The type-I blowup is optimal since the standard sphere  $S^n \hookrightarrow \mathbb{R}^{n+1}$  satisfies  $|HA| = \frac{n}{2(T-t)}$ .

**Corollary 5.2 (H-blowup).** *Let  $\mathbf{x} : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow with a finite maximal time  $T$ . Each time slice  $\Sigma_t$  has bounded second fundamental form. Assume that*

$$\limsup_{t \rightarrow T} \left( |T - t|^\lambda \sup_{\Sigma_t} |A| \right) < \infty$$

for some  $\lambda \in [\frac{1}{2}, 1)$ . Then we have the blowup estimate of mean curvature

$$\limsup_{t \rightarrow T} \left( |T - t|^{1-\lambda} \sup_{\Sigma_t} |H| \right) > 0.$$

*Proof.* For otherwise for any  $\varepsilon > 0$  one finds  $t_\varepsilon$  such that

$$\sup_{\Sigma_t} |H| \leq \varepsilon(T-t)^{\lambda-1}, \quad \forall t \in [t_\varepsilon, T).$$

By the assumption there exist nonnegative constants  $t_1 \in [0, T)$  and

$$C := \limsup_{t \rightarrow T} \left( |T-t|^\lambda \sup_{\Sigma_t} |A| \right) < \infty$$

such that

$$\sup_{\Sigma_t} |A| \leq C(T-t)^{-\lambda}, \quad \forall t \in [t_1, T).$$

Hence we have

$$\sup_{\Sigma_t} |HA| \leq C\varepsilon(T-t)^{-1}, \quad \forall t \in [\max\{t_\varepsilon, t_1\}, T).$$

Recall the constant  $\varepsilon = \varepsilon(n)$  in Theorem 5.1. Letting  $C\varepsilon < \varepsilon$  contradicts Theorem 5.1.  $\square$

Remark that by Theorem 5.1 of [2] Cooper proved the blowup of mean curvature under the same assumption in Corollary 5.2 and by Theorem 1.2 of [9] Le-Sesum proved the case of  $\lambda = \frac{1}{2}$ . Hence Theorem 5.1 and Corollary 5.2 can be seen as generalizations of these results.

**Corollary 5.3.** *Let  $x : \Sigma^n \times [0, T) \rightarrow \mathbb{R}^{n+1}$  be a complete noncompact smooth mean curvature flow with a finite maximal time  $T$ . Each time slice  $\Sigma_t$  has bounded second fundamental form. Assume that*

$$\limsup_{t \rightarrow T} \left( |T-t|^\lambda \sup_{\Sigma_t} |H| \right) < \infty$$

for some  $\lambda \in [0, \frac{1}{2})$ . Then we have the blowup estimate

$$\limsup_{t \rightarrow T} \left( |T-t|^{1-\lambda} \sup_{\Sigma_t} |A| \right) > 0.$$

In particular,  $t = T$  is a type-II singularity.

*Proof.* By the same argument used in the proof of Corollary 5.2 we obtain the result.  $\square$

## References

- [1] X. X. Chen, B. Wang, *On the conditions to extend Ricci flow(III)*. Int. Math. Res. Not. IMRN 2013, no. 10, 2349-2367.
- [2] A. Cooper, *A characterization of the singular time of the mean curvature flow*. Proc. Amer. Math. Soc. 139 (2011), no. 8, 2933-2942.
- [3] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. 17 (1982), no. 2, 255-306.
- [4] G. Huisken, *Flow by mean curvature of convex surfaces into spheres*, J. Differential Geom. 20 (1984), no. 1, 237-266.

- [5] B. Kotschwar, O. Munteanu, J. P. Wang, *A local curvature estimate for the Ricci flow*, J. Funct. Anal. 271 (2016), no. 9, 2604–2630.
- [6] H. Z. Li, B. Wang, *The extension problem of the mean curvature flow(I)*, Invent. Math. 218, 721-777 (2019).
- [7] N. Q. Le, N. Sesum, *The mean curvature at the first singular time of the mean curvature flow*. Ann. Inst. H. Poincaré Anal. Non Linéaire 27 (2010), no. 6, 1441-1459.
- [8] N. Q. Le, N. Sesum, *On the extension of the mean curvature flow*, Math. Z. 267, 583–604 (2011).
- [9] N. Q. Le, N. Sesum, *Blow-up rate of the mean curvature during the mean curvature flow and a gap theorem for self-shrinkers*, Commun. Anal. Geom. 19(4), 633-659 (2011).
- [10] C. Mantegazza, *Lecture notes on mean curvature flow*, Progress in Mathematics, 290. Birkhäuser/Springer Basel AG, Basel, 2011. xii+166 pp.
- [11] O. Munteanu, M. T. Wang, *The curvature of gradient Ricci solitons*. Math. Res. Lett. 18 (2011), no. 6, 1051–1069. (Reviewer: Bo Yang) 53C21 (53C25).
- [12] J. H. Michael, L. M. Simon, *Sobolev and mean value inequalities on generalized submanifolds of  $\mathbb{R}^{n+1}$* , Comm. Pure Appl. Math. 26 (1973), 316-379.
- [13] N. Sesum, *Curvature tensor under the Ricci flow*, Amer. J. Math. 127 (2005), no. 6, 1315-1324.
- [14] B. Wang, *On the conditions to extend Ricci flow*, Int. Math. Res. Not. IMRN 2008, no. 8, Art. ID rnn012, 30 pp.
- [15] B. Wang, *On the conditions to extend Ricci flow(II)*. Int. Math. Res. Not. IMRN 2012, no. 14, 3192-3223.
- [16] H. W. Xu, F. Ye, E. T. Zhao, *Extend mean curvature flow with finite integral curvature*. Asian J. Math. 15 (2011), no. 4, 549-556.