

Isometries of ultrametric normed spaces

Javier Cabello Sánchez, José Navarro Garmendia*

ABSTRACT. We show that the group of isometries of an ultrametric normed space can be seen as a kind of a fractal. Then, we apply this description to study ultrametric counterparts of some classical problems in Archimedean analysis, such as the so called *Problème des rotations de Mazur* or Tingley’s problem.

In particular, it turns out that, in contrast with the case of real normed spaces, isometries between ultrametric normed spaces can be very far from being linear.

1. Introduction

The study of isometries dates back to the classification of them as translation, reflections and rotations in the two-dimensional Euclidean space and comprises a huge variety of results, that include one of the most ancient and celebrated results in the theory of Banach spaces: the Mazur–Ulam Theorem states that every onto isometry between real Banach spaces is affine.

In [8], there was an attempt to prove an ultrametric version of the Mazur–Ulam Theorem introducing the notion of non-Archimedean strictly convex space. Nevertheless, Professor A. Kubzdela observed some years later that non-Archimedean strictly convex spaces are a *rarity* (see [7]).

Quite recently, the authors of the present work have shown that any attempt to obtain ultrametric versions of the Mazur–Ulam Theorem via strictly convex spaces is doomed to fail (see [4]).

The aim of this paper is to go a little further and analyse to which extent the behaviour of isometries in ultrametric analysis is far from their behaviour in real analysis. Our main result is Theorem 2.3, whose proof is elementary, and that exhibits a fractal structure of the group of isometries of an ultrametric normed space.

As a consequence, we prove in Corollary 3.8 that the only ultrametric normed spaces that possess a Mazur–Ulam like property are the trivial examples whose bijections are always affine; namely, the one-dimensional normed spaces over $\mathbb{Z}/2\mathbb{Z}$ or $\mathbb{Z}/3\mathbb{Z}$, and the two-dimensional normed spaces over $\mathbb{Z}/2\mathbb{Z}$.

1.1. Preliminaries. The first definitions are a commonplace in ultrametric analysis, and we reflect them just for the sake of completeness, see [8].

Keywords: Mazur–Ulam Theorem; ultrametric normed spaces; isometries.

Mathematics Subject Classification: 46S10, 26E30.

DEFINITION 1.1. A non-Archimedean (or ultrametric) field is a field \mathbb{K} equipped with a function $|\cdot| : \mathbb{K} \rightarrow [0, \infty)$ such that

- i) $|\lambda| = 0$ if and only if $\lambda = 0$,
- ii) $|\lambda\mu| = |\lambda||\mu|$,
- iii) $|\lambda + \mu| \leq \max\{|\lambda|, |\mu|\}$ for all $\lambda, \mu \in \mathbb{K}$.

DEFINITION 1.2. An ultrametric normed space is a linear space X over a non-Archimedean field $(\mathbb{K}, |\cdot|)$ that is endowed with an ultrametric norm; that is to say, endowed with a function $\|\cdot\| : X \rightarrow [0, \infty)$ such that

- iv) $\|x\| = 0$ if and only if $x = 0$,
- v) $\|\lambda x\| = |\lambda|\|x\|$, for all $\lambda \in \mathbb{K}, x \in X$.
- vi) $\|x + y\| \leq \max\{\|x\|, \|y\|\}$ for all $x, y \in X$.

Definition 1.2 endows X with a structure of metric space, thus allowing us to study the isometries between (subsets of) ultrametric normed spaces:

DEFINITION 1.3. Let $A \subseteq X$ and $B \subseteq Y$ be subsets of two ultrametric spaces X, Y . A map $f : A \rightarrow B$ is an isometry if it is bijective and, for any $x, y \in A$,

$$\|f(y) - f(x)\| = \|y - x\|.$$

The inversion through the origin, $x \mapsto -x$, is always an isometry $X \rightarrow X$, as so they are translations by a vector $x \mapsto x + z$.

Finally, let us also recall the following well-known result:

LEMMA 1.4 ([10], 8.C). On an ultrametric normed space X , every triangle is isosceles in the big, i.e., for every triplet of points $x, y, z \in X$, $\|x - z\| < \|y - z\|$ implies $\|y - x\| = \|y - z\|$.

2. Fractality of isometries

Let X be an ultrametric normed space. Let us denote the open ball and the sphere of radius $r \in (0, \infty)$ as

$$B(x, r) := \{y \in X : \|y - x\| < r\} \quad , \quad S_X(r) := \{x \in X : \|x\| = r\} ,$$

and let us also convene that

$$B(x, \infty) := X .$$

PROPOSITION 2.1. Let $r \in (0, \infty]$, and consider an arbitrary family:

$$f_{r'} : S_X(r') \rightarrow S_X(r')$$

of isometries on each sphere $S_X(r')$, for $r' \in (0, r)$.

Then, the map:

$$f : B(0, r) \longrightarrow B(0, r) \quad , \quad f(x) := \begin{cases} f_{r'}(x), & \text{if } x \in S_X(r') \\ 0, & \text{if } x = 0 \end{cases}$$

is a centred isometry (i.e., an isometry such that $f(0) = 0$).

PROOF. Observe that $\|x\| = \|f(x)\|$, for any $x \in B(0, r)$.

Thus, for any pair of points $x, y \in B(0, r)$,

- if $\|x\| = \|y\| = r'$, then

$$\|y - x\| = \|f_{r'}(y) - f_{r'}(x)\| = \|f(y) - f(x)\| .$$

- if $\|x\| < \|y\|$, then $\|f(x)\| < \|f(y)\|$ and

$$\|f(y) - f(x)\| = \|f(y)\| = \|y\| = \|y - x\| .$$

- if $\|x\| > \|y\|$, then $\|f(x)\| > \|f(y)\|$ and

$$\|f(y) - f(x)\| = \|f(x)\| = \|x\| = \|y - x\| .$$

□

The following conditions define equivalence relations on X , for any $r \in (0, \infty]$,

$$x \sim_r y \iff \|y - x\| < r .$$

The equivalence classes for these relations are precisely the open balls $B(x, r)$. Therefore, an ultrametric normed space X , as well as any other subset $A \subset X$, canonically decomposes as a disjoint union of open balls of any fixed radius r .

In particular, if we write S/\sim_r to abbreviate the quotient space $S_X(r)/\sim_r$, and we denote

$$B_{\bar{x}}(r) := B(x, r) , \text{ where } x \in X \text{ is any representative of } \bar{x} \in S/\sim_r ,$$

any sphere decomposes as a disjoint union of open balls:

$$(1) \quad S_X(r) = \bigsqcup_{\bar{x} \in S/\sim_r} B_{\bar{x}}(r) .$$

PROPOSITION 2.2. *Let $r \in (0, \infty)$ be a positive radius.*

If σ is a permutation of the quotient set S/\sim_r and $\{\varphi_{\bar{x}}: B_{\bar{x}}(r) \rightarrow B_{\sigma(\bar{x})}(r)\}_{\bar{x} \in S/\sim_r}$ is an arbitrary family of isometries, then the disjoint union of these isometries

$$S_X(r) \xrightarrow{f} S_X(r) \quad , \quad f(x) := \varphi_{\bar{x}}(x) ,$$

is an isometry of the sphere $S_X(r)$.

PROOF. Any two points of the sphere $x, y \in S_X(r)$ satisfy $\|y - x\| \leq r$ (because of Lemma 1.4), so that the distance between points $x, y \in S_X(r)$ in different open balls of (1) is exactly r .

Therefore, for any pair of points $x, y \in S_X(r)$,

- if $\|y - x\| < r$, then $\bar{x} = \bar{y}$ and

$$\|f(y) - f(x)\| = \|\varphi_{\bar{x}}(y) - \varphi_{\bar{x}}(x)\| = \|y - x\| .$$

- if $\|y - x\| = r$, then $\bar{x} \neq \bar{y}$. Hence, $\sigma(\bar{x}) \neq \sigma(\bar{y})$, so that $B_{\sigma(\bar{x})}(r) \neq B_{\sigma(\bar{y})}(r)$ and, in particular, $f(x)$ and $f(y)$ lie in different open balls of (1); that is to say, $\|f(y) - f(x)\| = r$.

□

We can summarize these results as follows:

THEOREM 2.3. *The following statements hold:*

- (1) *(Isometries between balls) Let B_1 and B_2 be open balls with the same (possibly infinite) radius $r \in (0, \infty]$. A map $f: B_1 \rightarrow B_2$ is an isometry if and only if there exist translations τ, τ' and a centred isometry $f_c: B(0, r) \rightarrow B(0, r)$ such that*

$$f = \tau' \circ f_c \circ \tau .$$

- (2) *(Centred isometries between balls) A centred map $f: B(0, r) \rightarrow B(0, r)$, with $r \in (0, \infty]$, is an isometry if and only if*
- (a) *It preserves norms: $\|f(x)\| = \|x\|$, for any $x \in B(0, r)$.*
 - (b) *It is an isometry on each sphere:*

$$f: S_X(r') \rightarrow S_X(r') \text{ is an isometry, for any } r' \in (0, r).$$

- (3) *(Isometries between spheres) A map $f: S_X(r) \rightarrow S_X(r)$, with $r \in (0, \infty)$, is an isometry if and only if there exist a permutation σ of the quotient set S/\sim_r and isometries $\varphi_{\bar{x}}: B_{\bar{x}}(r) \rightarrow B_{\sigma(\bar{x})}(r)$, for $\bar{x} \in S/\sim_r$, such that f is the disjoint union of the $\varphi_{\bar{x}}$:*

$$S_X(r) = \bigsqcup_{\bar{x} \in S/\sim_r} B_{\bar{x}}(r) \xrightarrow{f = \sqcup \varphi_{\bar{x}}} \bigsqcup_{\bar{x} \in S/\sim_r} B_{\bar{x}}(r) = S_X(r) .$$

PROOF. For the first statement, assume $f: B_1 \rightarrow B_2$ is an isometry between open balls of the same radius $r \in (0, \infty]$.

For any pair of points $x \in B_1$ and $y \in B_2$, the translation by the vector $y - x$ is an isometry $\tau_{y-x}: B_1 \rightarrow B_2$: in fact, for any $z \in B_1$, $\|\tau_{y-x}(z) - y\| = \|z - x\| < r$, so that $\tau_{y-x}(z) \in B_2$.

Therefore, if we take any point $x \in B_1$, the translations $\tau_{-x}: B_1 \rightarrow B(0, r)$ and $\tau_{f(x)}: B(0, r) \rightarrow B_2$ are well-defined, and f_c is the unique isometry that makes the square commutative:

$$\begin{array}{ccc} B_1 & \xrightarrow[\sim]{f} & B_2 \\ \tau_{-x} \downarrow & & \uparrow \tau_{f(x)} \\ B(0, r) & \xrightarrow{\dots\dots\dots f_c \dots\dots\dots} & B(0, r) . \end{array}$$

Finally, both the second and third statements are reformulations of Proposition 2.1 and Proposition 2.2, respectively. \square

EXAMPLE 2.4. In order to give an explanation of the name of this Section, let us describe the group of centred autoisometries of a quite simple space as $X = \mathbb{Q}_3^2$ when we consider the following valuation and norm:

$$\left| 3^n \frac{a}{b} \right| = 3^{-n}; \quad \|(\lambda, \mu)\| = \max\{|\lambda|, |\mu|\},$$

for every $n, a, b \in \mathbb{Z}$ such that a and b are coprime and $\lambda, \mu \in \mathbb{Q}_3$.

The norm $\|\cdot\|$ only takes values in $\{0\} \cup \{3^n : n \in \mathbb{Z}\}$, so the second part of Theorem 2.3 implies that we only need to study the autoisometries $\tau_n: S_X(3^n) \rightarrow S_X(3^n)$ for each $n \in \mathbb{Z}$. Later, we can glue these isometries to obtain $\tau: X \rightarrow X$ defined as $\tau(x) = \tau_n(x)$ when $\|x\| = 3^n$, $\tau(0) = 0$.

It is clear that any isometry $\tau_n : S_X(3^n) \rightarrow S_X(3^n)$ is associated with $\tau_0 : S_X \rightarrow S_X$ via the dilations $S_X(3^n) \rightarrow S_X$ and $S_X \rightarrow S_X(3^n)$, so the structure of Iso_X is determined by the structure of Iso_{S_X} : we only need to find one isometry $S_X \rightarrow S_X$ for each integer n and then glue them all in their corresponding spheres $S_X(3^n)$. But we can also take some bijection $\mathbb{Z} \rightarrow \mathbb{N} \cup \{0\}$, like the one defined as $n \mapsto 2n$, $-n \mapsto (-2n - 1)$ for every $n \in \mathbb{N}$ and $0 \mapsto 0$. This way, we have a correspondence between Iso_X and Iso_{B_X} thanks to Iso_{S_X} . So, let us analyse what happens with Iso_{S_X} . We can decompose S_X as in (1), but we have the nice feature that $B_X(x, 1) = B_X[x, 1/3]$ for every $x \in X$ –because $\|y\| < 1$ is equivalent to $\|y\| \leq 1/3$ – and in particular we get

$$(2) \quad S_X = \bigsqcup_{\bar{x} \in S/\sim_1} B_X[x, 1/3].$$

The decomposition given in (2) is much more simple than it seems. Namely, as the space that we are dealing with is $X = \mathbb{Q}_3^2$, the quotient S/\sim_1 contains only a finite number k of equivalence classes. So, we only need to choose some permutation $\sigma \in S_k$ (S_k stands for the symmetric group) and k isometries $\varphi_i : B_X[0, 1/3] \rightarrow B_X[0, 1/3]$. Of course, the structure of $\text{Iso}_{B_X[0, 1/3]}$ and the one of Iso_{B_X} are the same, so we can restrict ourselves to the study of $\text{Iso}_{B_X} \dots$

3. Some consequences

3.1. Isotropy of ultrametric spaces. One of the most famous open problems in functional analysis is the following:

3.1 (Problème des rotations de Mazur). Let X be a separable real or complex Banach space and suppose that, for every $x, y \in S_X$ there is an onto isometry $f : X \rightarrow X$ such that $f(x) = y$. Does this imply that X is an inner product space?

This was solved in the positive for finite-dimensional spaces and in the negative for non-separable spaces, but it is still open in the above form.

In the ultrametric setting, though, the group of isometries always acts transitively on the spheres:

COROLLARY 3.2. *Let X be an ultrametric normed space. For any pair of vectors $x, y \in X$ with the same norm, there exists a centred isometry $f : X \rightarrow X$ such that $f(x) = y$.*

PROOF. For any $x, y \in S_X(r)$, there exists an isometry $\varphi : S_X(r) \rightarrow S_X(r)$ such that $\varphi(x) = y$ (Proposition 2.2).

A global isometry is a gluing of isometries on each sphere (Proposition 2.1) so we are done. \square

REMARK 3.3. We distinguish *non-Archimedean* from *ultrametric*, although both terminologies are sometimes seen as equivalent. But they do not need to be. Actually, the definition of normed space over a valuated field that the reader can find in Bourbaki's monumental work [1] does not involve the ultrametric inequality. Both concepts *are* equivalent when one refers to valuated fields (because a multiplicative norm $|\cdot| : \mathbb{K} \rightarrow [0, \infty)$ satisfies the ultrametric inequality if and only if the image of the natural map $\mathbb{N} \rightarrow \mathbb{K}$ is bounded, [[10], 8.2]), but we have found no reason whatsoever to limit the non-Archimedean norms to ultrametric norms.

In fact, recent work ([3]) points out that it may be interesting to study the structure that usual (not ultrametric) norms give to linear spaces over non-Archimedean fields.

3.2. Tingley’s Problem. Due to the Mazur–Ulam Theorem and Mankiewicz’s Theorem, that states that every onto isometry $\tau : B_X \rightarrow B_Y$ between the unit balls of two real Banach spaces extends to a linear isometry $\tilde{\tau} : X \rightarrow Y$, it stands to reason to wonder whether every onto isometry $\tau : S_X \rightarrow S_Y$ between the unit spheres of real Banach spaces extends to a linear isometry, too. This is known as Tingley’s Problem, and is receiving a lot of attention nowadays, see, e.g., [5, 9] and the question can be generalised to *does every onto isometry between the boundaries of open convex subsets extend to a linear isometry between the spaces?*, see [2]; and to finding minimal requirements on the subset where the isometry is defined, see [6]. One key in this question is that τ extends linearly if and only if it extends to an isometry, so we have that the question is equivalent, in real Banach spaces, to any of the following:

- 1 Does every onto isometry $\tau : S_X \rightarrow S_Y$ extend to an isometry $\tilde{\tau} : X \rightarrow Y$?
- 2 Does every onto isometry $\tau : S_X \rightarrow S_Y$ extend to a linear map $\tilde{\tau} : X \rightarrow Y$?

In ultrametric normed spaces, it is clear that the answer to the first question is *yes, with conditions* and the answer to the second one is *not even close*. Please be aware that, for the question to make sense we need to choose a nonempty sphere instead of the unit sphere.

PROPOSITION 3.4. *Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be ultrametric normed spaces over some valued field \mathbb{K} , $r, r' > 0$ such that $rS_X \neq \emptyset$ and $\tau : rS_X \rightarrow r'S_Y$ an isometry. If the valuation is not trivial, then $r' = r$ and τ extends to an isometry $\tilde{\tau} : X \rightarrow Y$.*

PROOF. Let $x_0 \in rS_X$. The ultrametric inequality implies that $B_X(x_0, r) \subseteq S_X(0, r)$. It is clear that $\tau : B_X(x_0, r) \rightarrow B_Y(\tau(x_0), r)$ is also an onto isometry, so we may compose τ with translations to get a (centred) isometry $\tilde{\tau} : B_X(0, r) \rightarrow B_Y(0, r)$ defined as $\tilde{\tau}(x) = \tau(x+x_0) - \tau(x_0)$. Gluing τ with $\tilde{\tau}$ we obtain an isometry, which we will not rename,

$$\tilde{\tau} : B_X[0, r] \rightarrow B_Y[0, r].$$

Take some $\alpha \in \mathbb{K}$ such that $a = |\alpha| > 1$ – it exists because the valuation is not trivial. For any $x \in B_X[0, ra] \setminus B_X[0, r]$ (equivalently, $r < \|x\|_X \leq ra$), define $\tilde{\tau}(x) = \alpha\tau(x/\alpha)$. It is clear that $\tilde{\tau} : S_X(0, t) \rightarrow S_Y(0, t)$ is a well-defined onto isometry for every $t \in (r, ra]$. This implies that $\tilde{\tau} : B_X[0, ra] \rightarrow B_Y[0, ra]$ is a centred onto isometry. The same way we can define $\tilde{\tau}(x) = \alpha^n\tau(x/\alpha^n)$ whenever $x \in B_X[0, ra^n] \setminus B_X[0, ra^{n-1}]$, thus finishing the proof. \square

REMARK 3.5. If the valuation of \mathbb{K} is trivial, then we can find at least two pathological behaviours. Namely, if $(\mathbb{Z}/2\mathbb{Z})^2$ is endowed with the ultrametric norms

$$\|(a, b)\|_X = \max\{|a|, 2|b|\}, \|(a, b)\|_Y = \max\{|a|, 3|b|\}$$

then both unit spheres consist in the singleton $\{(1, 0)\}$ but the isometry $\tau : S_X \rightarrow S_Y$ defined as $\tau(1, 0) = (1, 0)$ does not extend. The same happens if we define the same norm over any other two-dimensional space over a field whose valuation is trivial.

Moreover, if we endow $(\mathbb{Z}/2\mathbb{Z})^2$ with

$$\|(a, b)\|_X = \max\{|a|, 2|b|\}, \|(a, b)\|_Y = \max\{2|a|, 3|b|\}$$

then both S_X and $2S_Y$ contain exactly one point. This means that they are trivially isometric, but, with the notations of Proposition 3.4, we have $r' \neq r$.

3.3. Non-existence of Mazur-Ulam type theorems.

EXAMPLE 3.6. The map $x \rightarrow -x$ is always a centred isometry $X \rightarrow X$ and, for any fixed x_0 , the map $x \mapsto x + x_0$ is an isometry of any sphere $S_X(r)$ with $r > \|x_0\|$.

Therefore, for any $r > \|x_0\|$ such that $S_X(r) \neq \emptyset$, the maps $f, \tilde{f}: X \rightarrow X$ defined as

$$f(x) = \begin{cases} -x, & \text{if } x \in S_X(r) \\ x, & \text{if } x \notin S_X(r) \end{cases}, \quad \tilde{f}(x) = \begin{cases} x + x_0, & \text{if } x \in S_X(r) \\ x, & \text{if } x \notin S_X(r) \end{cases}$$

are centred isometries.

Observe that, whenever they are distinct from the identity, f and \tilde{f} are non-linear maps.

EXAMPLE 3.7 (Isometries of one-dimensional spaces over finite fields). Any finite field \mathbb{F}_q is a non-Archimedean field in a unique way: equipped with the trivial valuation $|\lambda| = 1$, for any non-zero element $\lambda \in \mathbb{F}_q$. As a result, any bijective map $f: \mathbb{F}_q \rightarrow \mathbb{F}_q$ is an isometry.

In the same way, every ultrametric norm on a one-dimensional linear space X_1 over \mathbb{F}_q is trivial; i.e., for any such norm there exists $a \in (0, \infty)$ such that $\|x\| = a$ for any non-zero $x \in X_1$. Consequently, any bijective map $f: X_1 \rightarrow X_1$ is an isometry.

COROLLARY 3.8. *Let X be an ultrametric normed space. If every centred isometry $f: X \rightarrow X$ is a linear map, then one of the following options holds:*

- (1) $X = 0$.
- (2) X is a one-dimensional linear space over $\mathbb{Z}/2\mathbb{Z}$.
- (3) X is a one-dimensional linear space over $\mathbb{Z}/3\mathbb{Z}$.
- (4) X is a two-dimensional linear space over $\mathbb{Z}/2\mathbb{Z}$.

PROOF. If the norm on X is trivial, then any bijection $f: X \rightarrow X$ is an isometry. If X has more than four elements, then we may take different $x_1, x_2 \in X \setminus \{0\}$ such that $x_3 = x_1 + x_2 \neq 0$ and $x_4 \notin \{x_1, x_2, x_3, 0\}$ and define $f: X \rightarrow X$ as $f(x) = x$ for every $x \notin \{x_3, x_4\}$, $f(x_3) = x_4$, $f(x_4) = x_3$. This is obviously a nonlinear onto isometry from X to X .

If $\|\cdot\|$ is not trivial, then we have two options: apart from the value 0, either $\|\cdot\|$ takes exactly two values or it takes at least three.

If there are x_0, x_1, x_2 such that $0 < \|x_0\| < \|x_1\| < \|x_2\|$, then we may define the mapping $f: X \rightarrow X$ as

$$f(x) = \begin{cases} x, & \text{if } \|x\| \neq \|x_1\| \\ x + x_0, & \text{if } \|x\| = \|x_1\| \end{cases}.$$

Proposition 2.1 implies that f is an onto isometry and it is not linear because $f(x_2 + x_1) = x_2 + x_1$, $f(x_2) = x_2$, $f(x_1) = x_1 + x_0$.

The only option left is that $\|\cdot\|$ takes exactly two positive values, say there are $x_0, x_1 \in X$ such that $0 < \|x_0\| < \|x_1\|$.

Consider the map

$$f: X \rightarrow X, \quad f(x) := \begin{cases} x, & \text{if } \|x\| = \|x_1\| \\ -x, & \text{if } \|x\| = \|x_0\| \end{cases}.$$

This map is a bijection such that $f(0) = 0$ and it is clear that fulfils the conditions in Proposition 2.1, so it is an isometry.

Moreover, this map is not linear:

$$f(x_1 + x_0) - f(x_1) - f(x_0) = x_1 + x_0 - x_1 + x_0 = 2x_0 \neq 0 \dots$$

unless $2 = 0$ in \mathbb{K} . So, the only problem we have right now is that \mathbb{K} has characteristic 2, its valuation is trivial (if it is not, then the valuation takes infinitely many values, and so does the norm), and the norm takes exactly two positive values.

If X fulfils everything, then take $0 < \|x_1\| < \|x_2\|$. As every triangle is *isosceles in the big*, for every pair x, y such that $\|x\| = \|y\| = \|x_1\|$ we have $\|x + y\| \leq \|x_1\|$. Furthermore, the valuation of \mathbb{K} is trivial, so

$$X_1 = \{x \in X : \|x\| \leq \|x_1\|\} = \{x \in X : \|x\| = \|x_1\|\} \cup \{0\}$$

is closed under addition and scalar multiplication and this means that it is a vector subspace of X with trivial norm, so if it has only affine isometries it must have 4 elements or less. If there is some nonlinear isometry $X_1 \rightarrow X_1$ that sends 0 to 0 then we just need to extend it to X by identity, so our only problem is that X_1 has at most four elements. In this case, $\mathbb{K} = \mathbb{Z}/2\mathbb{Z}$ implies that X is (linearly) isometric to either $\mathbb{Z}/2\mathbb{Z}$ or $(\mathbb{Z}/2\mathbb{Z})^2$ and $\mathbb{K} = \mathbb{F}_4$ implies that X is (linearly) isometric to \mathbb{K} .

In the first case, it is easy to see that a bijection $f : X \rightarrow X$ is an isometry if and only if $f(x + x_1) = f(x) + x_1$ for every $x \in X$. As the dimension of X is at least 3, there are $x, y, z \notin X_1$ such that $z = x + y$. Now, the map $f : X \rightarrow X$ defined as $f(x) = x$ whenever $x \notin \{z, z + x_1\}$, $f(z) = z + x_1$, $f(z + x_1) = z$ is an isometry and it is not affine.

In the second case, we only need to consider some linear isometry $\tau : X_1 \rightarrow X_1$ that does not send x_1 to x_1 —it exists because X_1 contains four elements— and define $f : X \rightarrow X$ as

$$f(x) = \begin{cases} x & \text{if } x \notin X_1 \\ \tau(x) & \text{if } x \in X_1 \end{cases} .$$

This is an isometry by Proposition 2.1. Furthermore it is not linear because $f(x_2 + x_1) = x_2 + x_1$, $f(x_1) = \tau(x_1) \neq x_1$ and $f(x_2) = x_2$.

To finish the proof, we must observe that if we consider the four element field \mathbb{F}_4 , that can be seen as $\mathbb{F}_4 = ((\mathbb{Z}/2\mathbb{Z})[x])/(x^2 + x + 1)$, then every centred bijection $\mathbb{F}_4 \rightarrow \mathbb{F}_4$ is an additive isometry. But it is clear that the map $f : \mathbb{F}_4 \rightarrow \mathbb{F}_4$ defined as

$$f([0]) = [0], f([1]) = [1], f([x]) = [x^2], f([x^2]) = [x]$$

is not (\mathbb{F}_4) -linear. □

Acknowledgments

Supported in part by DGICYT project PID2019-103961GB-C21 (Spain), Feder funds and Junta de Extremadura program IB-18087.

References

- [1] N. Bourbaki. *Espaces vectoriels topologiques sur un corps valué*, pages 1–29. Springer Berlin Heidelberg, Berlin, Heidelberg, 2007.
- [2] J. Cabello Sánchez. Linearity of isometries between convex Jordan curves. *Linear Algebra and its Applications*, 621:1–17, 2021.
- [3] J. Cabello Sánchez and F. Carmona Fuertes. On the concept of non-ultrametric non-Archimedean analysis. *Preprint*, 2021.
- [4] J. Cabello Sánchez and J. Navarro Garmendia. A precision on the concept of strict convexity in non-Archimedean analysis. *Nonlinear Analysis*, 207:112304, 2021.
- [5] J. Cabello Sánchez. A reflection on Tingley’s problem and some applications. *Journal of Mathematical Analysis and Applications*, 476(2):319–336, 2019.
- [6] M. Cueto-Avellaneda and A. M. Peralta. Can one identify two unital JB*-algebras by the metric spaces determined by their sets of unitaries? *Preprint*, 2021.
- [7] A. Kubzdela. Isometries, Mazur–Ulam theorem and Aleksandrov problem for non-Archimedean normed spaces. *Nonlinear Analysis: Theory, Methods & Applications*, 75(4):2060–2068, 2012.
- [8] M. S. Moslehian and G. A. Sadeghi. Mazur-Ulam Theorem in non-Archimedean normed spaces. *Nonlinear Anal.*, 69(10):3405–3408, 2008.
- [9] A. M. Peralta. A survey on Tingley’s problem for operator algebras. *Acta Scientiarum Mathematicarum*, 84(1–2):81–123, 2018.
- [10] W. Slichhoff. *Ultrametric Calculus, an introduction to p-adic analysis*. Cambridge studies in advanced mathematics, 1984.

*CORRESPONDING AUTHOR: DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS S/N, 06006; BADAJOZ. SPAIN

Email address: `coco@unex.es`, `navarrogarmendia@unex.es`