

EXTREMAL PRODUCT-ONE FREE SEQUENCES OVER $C_n \rtimes_s C_2$

F. E. BROCHERO MARTÍNEZ AND S. RIBAS

ABSTRACT. Let G be a finite group multiplicatively written. The small Davenport constant of G is the maximum positive integer $d(G)$ such that there exists a sequence S of length $d(G)$ for which every subsequence of S is product-one free. Let $s^2 \equiv 1 \pmod{n}$, where $s \not\equiv \pm 1 \pmod{n}$. It has been proven that $d(C_n \rtimes_s C_2) = n$ (see [12, Lemma 6]). In this paper, we determine all sequences over $C_n \rtimes_s C_2$ of length n which are product-one free. It completes the classification of all product-one free sequences over every group of the form $C_n \rtimes_s C_2$, including the quasidihedral groups and the modular maximal-cyclic groups.

1. INTRODUCTION

Given a finite group G multiplicatively written, the *zero-sum problems* study conditions to ensure that a given sequence over G has a non-empty subsequence with prescribed properties (such as length, repetitions, weights) such that the product of its terms in some order equals 1.

1.1. Definitions and notations. A sequence over a finite group G is a finite and unordered element S of the free abelian monoid $\mathcal{F}(G)$, equipped with the sequence concatenation product denoted by \cdot . The form of S is

$$S = g_1 \cdot \dots \cdot g_k = \prod_{1 \leq i \leq k}^{\bullet} g_i = \prod_{1 \leq i \leq k}^{\bullet} g_{\tau(i)} = \prod_{g \in G}^{\bullet} g^{v_g(S)} \in \mathcal{F}(G),$$

where $g_1, \dots, g_k \in G$ are the *terms* of S , $k = |S| \geq 0$ is the *length* of S , $\tau : \{1, 2, \dots, k\} \rightarrow \{1, 2, \dots, k\}$ is a permutation, and $v_g(S) = \#\{i \in \{1, 2, \dots, k\}; g_i = g\}$ is the *multiplicity* of g in S . We say that a sequence T is a *subsequence* of S if $v_g(T) \leq v_g(S)$ for all $g \in G$, that is, if $T \mid S$ in $\mathcal{F}(G)$; in this case, we write $S \cdot T^{[-1]} = \prod_{g \in G}^{\bullet} g^{[v_g(S) - v_g(T)]}$. We also define:

$$\pi(S) = \{g_{\tau(1)} \cdot \dots \cdot g_{\tau(k)} \in G; \tau \text{ is a permutation of } \{1, 2, \dots, k\}\} \quad (\text{the set of products of } S);$$

$$\Pi(S) = \bigcup_{\substack{T \mid S \\ |T| \geq 1}} \{\pi(T)\} \subset G \quad (\text{the set of subproducts of } S);$$

$$S \cap K = \prod_{\substack{g \mid S \\ g \in K}}^{\bullet} g \quad (\text{the subsequence of } S \text{ that lie in a subset } K \text{ of } G).$$

The sequence S is called *product-one free* if $1 \notin \Pi(S)$, and *product-one sequence* if $1 \in \pi(S)$.

1.2. The small Davenport constant. An important type of zero-sum problem is the *small Davenport constant* of G . Denoted by $d(G)$, this constant is the maximal integer such that there exists a sequence over G (repetition allowed) of length $d(G)$ which is product-one free, i.e.,

$$d(G) = \sup\{|S| > 0; S \in \mathcal{F}(G) \text{ is product-one free}\}.$$

Date: August 3, 2021.

2010 Mathematics Subject Classification. 11P70 (primary), 11B50 (secondary).

Key words and phrases. Zero-sum problem, small Davenport constant, inverse zero-sum problem, semidirect product.

The first author was partially supported by FAPEMIG APQ-02973-17, Brazil.

Denote by C_n the cyclic group of order n . It is known that $d(C_n) = n - 1$. Zhuang & Gao [12] showed that $d(G) = n$ for every non-abelian group G of the form $C_n \rtimes_s C_2$. The conditions $s^2 \equiv 1 \pmod{n}$ and $s \not\equiv 1 \pmod{n}$ are required in order to obtain non-abelian groups. The case $s \equiv -1 \pmod{n}$ yields D_{2n} , the Dihedral Group of order $2n$.

1.3. Inverse zero-sum problems. Let G be a finite group. By definition, there exist sequences $S \in \mathcal{F}(G)$ with $d(G)$ terms that are product-one free. The *inverse zero-sum problems* study the structure of these extremal length sequences; see [5, 10, 4] for an overview on inverse problems over abelian groups. In [1] (see also Lemma 2.1 below) and [10, Theorem 3.2], one can find the solution of the inverse problem associated with small Davenport constant over cyclic groups and over abelian groups of rank two, respectively. For non-abelian groups, the first results of this kind have emerged in recent years; see [2, 3, 8, 9]. In particular, the authors solved in [3] the inverse problem associated with small Davenport constant over Dihedral Groups.

Let n be a positive integer such that there exists an integer s satisfying

$$s^2 \equiv 1 \pmod{n}, \quad \text{but} \quad s \not\equiv \pm 1 \pmod{n}, \quad (1.1)$$

and consider the group $C_n \rtimes_s C_2$ with presentation

$$\langle x, y \mid x^2 = y^n = 1, yx = xy^s \rangle, \quad \text{where the pair } (n, s) \text{ satisfies the conditions (1.1)}. \quad (1.2)$$

The conditions (1.1) are necessary, since the semidirect product $C_n \rtimes_s C_2$ should yield a non-abelian group. Indeed, $s \equiv 1 \pmod{n}$ means that $C_n \rtimes_s C_2$ is an abelian group (C_{2n} or $C_2 \times C_n$, depending on whether n is odd or even, respectively), while $s \equiv -1 \pmod{n}$ means that $C_n \rtimes_s C_2$ is the Dihedral Group of order $2n$, for which the inverse problems are already solved. It is worth mentioning that this family of metacyclic groups includes the quasidihedral groups, when $(n, s) = (2^t, 2^{t-1} - 1)$, $t \geq 3$, and the modular maximal-cyclic groups, when $(n, s) = (2^t, 2^{t-1} + 1)$, $t \geq 3$.

In this paper, we solve the inverse problem over $C_n \rtimes_s C_2$ related to small Davenport constant, proving the following result:

Theorem 1.1. *Let $G \simeq C_n \rtimes_s C_2$ be the group with presentation given by (1.2). Let $S \in \mathcal{F}(G)$ be a sequence of length $|S| = n$. Then S is product-one free if and only if*

- (i) *For any (n, s) , $S = (y^u)^{[n-1]} \cdot xy^v$ for some integers u, v with $\gcd(u, n) = 1$, or*
- (ii) *For $(n, s) = (2^t, 2^{t-1} + 1)$ where $t \geq 3$, $S = \begin{cases} (xy^u)^{[n-1]} \cdot xy^v & \text{for } u \text{ odd and } v \text{ even, or} \\ (xy^u)^{[n-1]} \cdot y^v & \text{for } u, v \text{ odd.} \end{cases}$*

In particular, together with [3, Theorem 1.3] and [10, Theorem 3.2], the previous theorem completes the classification of all the extremal length product-one free sequences over all groups of the form $C_n \rtimes_s C_2$. The case (ii) of previous theorem appears to have some extra extremal product-one free sequences, but this is not actually true. It turns out that $C_n \rtimes_s C_2 = \langle x, xy^v \mid v \text{ is odd} \rangle$ for $(n, s) = (2^t, 2^{t-1} + 1)$ and $t \geq 3$, that is, the extra sequences arise only because of this change of generators.

The paper is organized as follows: In Section 2, we present two lemmas which will be used later. In Section 3, we prove the main result by splitting the proof into cases according to the number of terms of S that belong to the normal subgroup of order n , and in some cases we further split into subcases according to either the value of s or the factorization of n .

2. PRELIMINARY RESULTS

In this section, we present two preliminary results that will be used in the proof of the main theorem. The first can be seen as a generalization of the inverse zero-sum problem over cyclic groups, dealing with product-one free sequences of large length and their subsequence sums.

Lemma 2.1 (See [7, Theorem 11.1]). *Let $m \geq 3$ be an integer and fix a generator y of C_m . Let $S = \prod_{1 \leq i \leq |S|} y^{a_i} \in \mathcal{F}(C_m)$ be a product-one free sequence of length $|S| > m/2$. Then there exists $y^{a_i} \mid S$ such that $v_{y^{a_i}}(S) \geq \max\{m - 2|S| + 1, |S| - \lfloor \frac{m-1}{3} \rfloor\}$. In addition, for a residue class $r \pmod{m}$, denote by \bar{r} the integer such that $0 \leq \bar{r} \leq m - 1$ and $r \equiv \bar{r} \pmod{m}$. Then there exists an integer t with $\gcd(t, m) = 1$ and $\sum_{1 \leq i \leq |S|} \overline{a_i t} < m$. Moreover, for every $1 \leq k \leq \sum_{1 \leq i \leq |S|} \overline{a_i t}$, there exists $T_k \mid S$ such that $\sum_{y^{a_i} \mid T_k} \overline{a_i t} = k$.*

The second lemma is related to the factorization of n and has essentially been considered in [6, Section 5], nevertheless we summarize it here for convenience. We notice that conditions (1.1) guarantee that n can be neither an odd prime power nor twice an odd prime power, otherwise we would have $s \equiv \pm 1 \pmod{n}$. Assuming that n is not a power of 2, we are going to show that these conditions suffice to factor n in a helpful way.

Lemma 2.2. *Let $n \geq 8$ and s be positive integers satisfying the conditions (1.1).*

- *If both $n \neq p^t$ and $n \neq 2p^t$ for every prime p and every integer $t \geq 1$, then there exist coprime integers $n_1, n_2 \geq 3$ such that $s \equiv -1 \pmod{n_1}$, $s \equiv 1 \pmod{n_2}$, and either (A) $n = n_1 n_2$ or (B) $n = 2n_1 n_2$.*
- *If $n = 2^t$ for some $t \geq 3$, then (B) $n = 2n_1 n_2$, where either $(n_1, n_2) = (1, 2^{t-1})$ satisfies $s \equiv 1 \pmod{n_2}$ or $(n_1, n_2) = (2^{t-1}, 1)$ satisfies $s \equiv -1 \pmod{n_1}$.*

Proof: Let $n = 2^t m$, where m is odd and $t \geq 0$ is an integer. Since m divides $s^2 - 1$ and $\gcd(s - 1, s + 1) \in \{1, 2\}$ (depending on s is even or odd), each prime power factor of m divides either $s + 1$ or $s - 1$. Let $m_1 = \gcd(m, s + 1)$ and $m_2 = \gcd(m, s - 1)$, so that $m = m_1 m_2$. In addition, $s^2 \equiv 1 \pmod{2^t}$ implies that either $s \equiv \pm 1 \pmod{2^t}$ or $s \equiv 2^{t-1} \pm 1 \pmod{2^t}$ (whether $t \geq 3$). We consider some cases:

- (i) **CASE** $t = 0$. In this case, $n = m = m_1 m_2$. We set $n_1 = m_1$ and $n_2 = m_2$, hence $n = n_1 n_2$ is the desired factorization, as in (A). This is the only case where s can be even; in the following, s must be odd.
- (ii) **CASE** $t = 1$. It is possible to set either $n_1 = 2m_1$ and $n_2 = m_2$ or $n_1 = m_1$ and $n_2 = 2m_2$, hence $n = n_1 n_2$ is the desired factorization, as in (A).
- (iii) **CASE** $t \geq 2$ **and** $m \geq 3$. If $s \equiv -1 \pmod{2^t}$, then we set $n_1 = 2^t m_1$ and $n_2 = m_2$. If $s \equiv 1 \pmod{2^t}$, then we set $n_1 = m_1$ and $n_2 = 2^t m_2$. Therefore, $n = n_1 n_2$ is the desired factorization, as in (A).

In the case that $t \geq 3$, it is possible that $s \equiv 2^{t-1} \pm 1 \pmod{2^t}$. If $s \equiv 2^{t-1} - 1 \pmod{2^t}$, then we set $n_1 = 2^{t-1} m_1$ and $n_2 = m_2$. If $s \equiv 2^{t-1} + 1 \pmod{2^t}$, then we set $n_1 = m_1$ and $n_2 = 2^{t-1} m_2$. Thus, $n = 2n_1 n_2$ is the desired factorization as in (B).

- (iv) **CASE** $t \geq 3$ **and** $m = 1$. In this case, $s \equiv 2^{t-1} \pm 1 \pmod{2^t}$, which implies that $s \equiv \pm 1 \pmod{2^{t-1}}$. For the negative sign, it follows that $(n_1, n_2) = (2^{t-1}, 1)$, and for the positive sign, it follows that $(n_1, n_2) = (1, 2^{t-1})$. Therefore, $n = 2n_1 n_2$ is the factorization as in (B).

□

We highlight that the case (iv) of previous lemma yields the quasidihedral group provided $s \equiv 2^{t-1} - 1 \pmod{2^t}$ and the modular maximal-cyclic group provided $s \equiv 2^{t-1} + 1 \pmod{2^t}$.

From the previous lemma and Chinese Remainder Theorem, there exists a natural projection

$$\Psi : C_n \rightarrow C_{n_1} \oplus C_{n_2} \tag{2.1}$$

satisfying $\Psi(e) = (e_1, e_2)$ and $\Psi(s \cdot e) = (-e_1, e_2)$, where $C_{n_1} = \langle e_1 \rangle$, $C_{n_2} = \langle e_2 \rangle$ and $C_n = \langle e \rangle$. In case (A) of previous lemma, Ψ is an isomorphism.

3. PROOF OF THEOREM 1.1

The sequences of the form $(y^u)^{[n-1]} \cdot xy^v \in \mathcal{F}(C_n \rtimes_s C_2)$, where $\gcd(u, n) = 1$, are product-one free. In addition, if $(n, s) = (2^t, 2^{t-1} + 1)$ for some $t \geq 3$, then the sequences $(xy^u)^{[n-1]} \cdot xy^v \in \mathcal{F}(C_n \rtimes_s C_2)$, for u odd and v even, and $(xy^u)^{[n-1]} \cdot y^v \in \mathcal{F}(C_n \rtimes_s C_2)$, for u, v odd, are product-one free. In fact:

- If $(xy^u)^a = 1$ for some $a \in \{1, 2, \dots, n-1\}$, then a must be even, thus

$$1 = (y^{2u+n/2})^{a/2} = y^{(2u+n/2)a/2} = y^{(u+n/4)a},$$

which implies that $n = 2^t$ divides $a(u + n/4)$. Since $\gcd(u + n/4, n) = 1$, n divides a , a contradiction.

- If $(xy^u)^a(xy^v) = 1$, then a must be odd, thus

$$1 = [(xy^u)^2]^{(a-1)/2} \cdot (xy^u \cdot xy^v) = y^{(2u+n/2)(a-1)/2} \cdot y^{u+n/2+v} = y^{(2u+n/2)(a-1)/2+u+v+n/2}.$$

It implies that $n = 2^t$ divides $(2u + n/2)(a-1)/2 + u + v + n/2$, but this is a contradiction since $(2u + n/2)(a-1)/2 + u + v + n/2$ is odd.

- If $(xy^u)^a y^v = 1$, then a must be even, thus

$$1 = [(xy^u)^2]^{a/2} \cdot y^u = y^{(2u+n/2)a/2} \cdot y^u = y^{(2u+n/2)a+v}.$$

It implies that $n = 2^t$ divides $(2u + n/2)a/2 + v$, but this is a contradiction since $(2u + n/2)a/2 + v$ is odd.

Therefore it only remains to verify the converse. Let $S \in \mathcal{F}(C_n \rtimes_s C_2)$ be a product-one free sequence of length $|S| = n$. Let $H = \langle y \rangle \simeq C_n$ be the normal subgroup of $C_n \rtimes_s C_2$ and let $N = xH$. We split into some cases and subcases:

- (a) **CASE $|S \cap H| = n$.** In this case, S is contained in the cyclic subgroup of order n . Since $d(C_n) = n-1$, S cannot be product-one free.
- (b) **CASE $|S \cap H| = n-1$.** In this case, by Lemma 2.1, the terms of $S \cap H$ must all be equal to the same fixed generator of H . Therefore, there exist integers t and u with $\gcd(t, n) = 1$ such that $S = (y^t)^{[n-1]} \cdot xy^u$.
- (c) **CASE $|S \cap H| = n-k$, where $2 \leq k < n/2$.** By Lemma 2.1, we may assume without loss of generality that $S \cap H = \prod_{1 \leq i \leq n-k}^\bullet y^{a_i}$, where $n-k \leq \sum_{1 \leq i \leq n-k} a_i \leq n-1$. Moreover, Lemma 2.1 also ensures that $y^\gamma \in \Pi(S \cap H)$ for every $1 \leq \gamma \leq n-k$. If there exists a subsequence $xy^\alpha \cdot xy^\beta \mid S \cap N$ such that $\alpha s + \beta$ has a representant in the range $[k, n-1]$ modulo n , then there exists $y^\gamma \in \Pi(S \cap H)$, $1 \leq \gamma \leq n-k$, such that $xy^\alpha \cdot xy^\beta \cdot y^\gamma = 1$. Therefore, we assume that $\alpha s + \beta$ has a representant in $[1, k-1]$ for all α, β such that $xy^\alpha \cdot xy^\beta \mid S$. Since $s^2 \equiv 1 \pmod{n}$, we may assume that s has an integer representant in the range $[\sqrt{n+1}, n - \sqrt{n+1}]$. The elements from $S \cap N$ are paired forming more $[k/2]$ terms in H , say y^{a_i} for $n-k+1 \leq i \leq n-k+[k/2]$, and it yields a new sequence $T \in \mathcal{F}(H)$ of length $|T| = n-k+[k/2] \geq 3n/4$, that is, $T = \prod_{1 \leq i \leq n-k+[k/2]}^\bullet y^{a_i}$ is the concatenation of $S \cap H$ and the pairs of $S \cap N$. If T is product-one free, then the average of the exponents of the terms other than y is at least two, therefore

$$\frac{n-1 - v_y(T)}{3n/4 - v_y(T)} \geq \frac{\sum_{1 \leq i \leq |T|} a_i - v_y(T)}{n-k+[k/2] - v_y(T)} \geq 2,$$

which implies $v_y(T) \geq [n/2] + 1$.

We are going to show that at least one of the $y \mid T$ comes from a pair of $S \cap N$, that is, $v_y(T) > v_y(S \cap H) = v_y(S)$. Otherwise, the average of the $\lfloor k/2 \rfloor$ terms of T coming from $S \cap N$ is at most

$$\frac{\sum_{y^{a_i} \mid T} a_i - \sum_{y^{a_i} \mid S \cap H} a_i}{\lfloor k/2 \rfloor} \leq \frac{n-1-(n-k)}{\lfloor k/2 \rfloor} = \begin{cases} \frac{2(k-1)}{k} < 2 & \text{if } k \text{ is even,} \\ 2 & \text{if } k \text{ is odd.} \end{cases}$$

The Pigeonhole Principle implies that $\Pi(S \cap N) = \{y^2\}$ and k is odd. Let $xy^\alpha \cdot xy^\beta \mid S \cap N$ such that $xy^\alpha \cdot xy^\beta = y^2$, thus $xy^\beta \cdot xy^\alpha = y^{2s}$. In general, it increases the exponent and contradicts the average at most two. In fact, if $2s \equiv 1 \pmod{n}$, then we would be done since $y^{2s} = y \mid T$. Moreover, $2s \equiv 2 \pmod{n}$ if and only if either $s \equiv 1 \pmod{n}$ if n is odd, a contradiction, or $s \equiv n/2 + 1 \pmod{n}$ if n is even. In the case that n is even, it follows that n is divisible by 4 since $\gcd(s, n) = 1$. In addition, $S \cap N = (xy^\alpha)^{\lfloor k \rfloor}$, where $\alpha s + \alpha \equiv 2 \pmod{n}$. In fact, if $k \geq 3$, then let $xy^\alpha \cdot xy^\beta \cdot xy^\gamma \mid S \cap N$. It follows that $\alpha s + \beta \equiv 2 \pmod{n}$ and $\alpha s + \gamma \equiv 2 \pmod{n}$, from where we obtain $\beta \equiv \gamma \pmod{n}$ and similarly we obtain $S \cap N = (xy^\alpha)^{\lfloor k \rfloor}$. If $k = 2$, then $S \in \{y^{\lfloor n-2 \rfloor} \cdot xy^\alpha \cdot xy^\beta, y^{\lfloor n-3 \rfloor} \cdot y^2 \cdot xy^\alpha \cdot xy^\beta\}$, and in both cases it is easy to verify that S is not product-one free since either $\alpha s + \beta \not\equiv 1 \pmod{n}$ or $\beta s + \alpha \not\equiv 1 \pmod{n}$. Therefore, $S \cap N = (xy^\alpha)^{\lfloor k \rfloor}$. If α is even, then $\alpha s + \alpha \equiv 2 \pmod{n}$ and $s \equiv n/2 + 1 \pmod{n}$ imply that $2\alpha \equiv 2 \pmod{n}$, therefore $\alpha \equiv 1 \pmod{n/2}$ and α is odd, contradiction. If α is odd, then $n/2 + 2\alpha \equiv 2 \pmod{n}$, therefore $\alpha \equiv n/4 + 1 \pmod{n/2}$ and hence $\alpha \in \{n/4 + 1, 3n/4 + 1\}$. For each value of α , we obtain a contradiction using the following products:

$$\begin{cases} xy^{n/4+1} \cdot y \cdot xy^{n/4+1} \cdot y^{n/2-3} = 1 & \text{if } \alpha = n/4 + 1 \text{ and } n \equiv 0 \pmod{8}, \\ xy^{n/4+1} \cdot y^2 \cdot xy^{n/4+1} \cdot y^{n/2-4} = 1 & \text{if } \alpha = n/4 + 1 \text{ and } n \equiv 4 \pmod{8}, \\ xy^{3n/4+1} \cdot y \cdot xy^{3n/4+1} \cdot y^{n/2-3} = 1 & \text{if } \alpha = 3n/4 + 1 \text{ and } n \equiv 0 \pmod{8}, \\ xy^{3n/4+1} \cdot y^2 \cdot xy^{3n/4+1} \cdot y^{n/2-4} = 1 & \text{if } \alpha = 3n/4 + 1 \text{ and } n \equiv 4 \pmod{8}, \end{cases}$$

which implies that $v_y(T) > v_y(S)$ so far.

Let $xy^\alpha \cdot xy^\beta \mid S \cap N$ with $y = xy^\alpha \cdot xy^\beta$, so that $y^s = xy^\beta \cdot xy^\alpha$. We consider the following subcases:

(c.1) **Subcase** $n/2 < s \leq n - \sqrt{n+1}$. Since $1 \leq n-s < n/2$ and $|S \cap H| > n/2$, Lemma 2.1 ensures that $y^{n-s} \in \Pi(S \cap H)$. Therefore $xy^\beta \cdot xy^\alpha \cdot y^{n-s} = 1$, and we are done.

(c.2) **Subcase** $\sqrt{n+1} \leq s < n/2$. We proceed similarly to the baby-step giant-step method for the calculus of discrete logarithm [11]. Let $\ell \geq 2$ be the greatest integer such that $\ell s \leq n$. It follows that $\ell \leq \lfloor \frac{n}{\sqrt{n+1}} \rfloor = \lfloor \sqrt{n} \rfloor$ and $n-s < \ell s \leq n$. Since $v_y(T) \geq \lfloor n/2 \rfloor + 1$, we consider the following product

$$xy^\beta \cdot \underbrace{y \cdots y}_{\ell-1 \text{ times}} \cdot xy^\alpha = y^{\ell s}. \quad (3.1)$$

If $\sqrt{n+1} \leq s \leq \lfloor n/2 \rfloor + 1 - \lfloor \sqrt{n} \rfloor$, then $\ell + s \leq \lfloor n/2 \rfloor + 1$ and we complete the product in (3.1) as $y^{\ell s} \cdot \underbrace{y \cdots y}_{n-\ell s \text{ times}} = 1$ since $1 \leq n - \ell s < s$.

Otherwise, $\lfloor n/2 \rfloor - \lfloor \sqrt{n} \rfloor + 2 \leq s < n/2$ implies that $\ell = 2$, therefore $2 + s \leq \lfloor n/2 \rfloor + 1$ and in the same way we obtain $y^{2s} \cdot y^{n-2s} = 1$, since $n - 2s \leq \lfloor n/2 \rfloor - 1$. Hence we are done.

(d) **CASE** $|S \cap H| = n - k$, where $n/2 \leq k \leq n$.

(d.1) **Subcase** $n = n_1 n_2$. Group the terms of $S \cap N = xy^{\beta_1} \cdots xy^{\beta_k}$ in pairs $xy^{\beta_i} \cdot xy^{\beta_j}$ in such way that $\beta_i s + \beta_j \equiv 0 \pmod{n_1}$. This is possible always that $\beta_i \equiv \beta_j \pmod{n_1}$ because $s \equiv -1 \pmod{n_1}$, hence we group these pairs until remains $\ell \leq n_1$ terms, each of them having the exponents of y in distinct residue classes modulo n_1 . The number of pairs formed is $\lceil \frac{k-\ell}{2} \rceil$. The ℓ remaining terms are grouped into $\lfloor \ell/2 \rfloor$ pairs with product in H . We also group these

$\lfloor \ell/2 \rfloor$ terms and the terms of $S \cap H = y^{\alpha_1} \cdot \dots \cdot y^{\alpha_{n-k}}$ into $\lfloor \frac{n-k+\lfloor \ell/2 \rfloor}{n_1} \rfloor$ subsequences, each of them is not product-one free in the quotient $C_n/C_{n_1} \simeq \langle y^{n_1} \rangle$. Thus, we obtain $\frac{k-\ell}{2} + \lfloor \frac{n-k+\lfloor \ell/2 \rfloor}{n_1} \rfloor$ elements in $\langle y^{n_1} \rangle \simeq C_{n_2}$. Since $\ell \leq n_1$, we have that

$$\begin{aligned} \frac{k-\ell}{2} + \left\lfloor \frac{n-k+\lfloor \ell/2 \rfloor}{n_1} \right\rfloor &\geq \frac{k-\ell}{2} + \frac{n-k+\frac{\ell-1}{2} - (n_1-1)}{n_1} \\ &\geq \frac{k}{2} - \frac{n_1}{2} + n_2 - \frac{k}{n_1} - \frac{1}{2} + \frac{1}{2n_1}, \end{aligned}$$

and if the latter is bigger than $d(C_{n_2}) = n_2 - 1$, then we are done, since we obtain more than $d(C_{n_2})$ elements in $\langle y^{n_1} \rangle$, which yields a product-one sequence over C_n . But this assertion is equivalent to $k > n_1 + 1 + \frac{3}{n_1-2}$. Since $k \geq \frac{n}{2} = \frac{n_1 n_2}{2}$, if $\frac{n_1 n_2}{2} > n_1 + 1 + \frac{3}{n_1-2}$, then we are done. It implies that we are done unless either $(n_1, n_2) \in (3, 4)$ and $6 \leq k \leq 7$ or $(n_1, n_2) = (4, 3)$ and $k = 6$, since otherwise $k > n_1 + 1 + \frac{3}{n_1-2}$. But these cases follows directly from the inequality

$$\frac{k-\ell}{2} + \left\lfloor \frac{n-k+\lfloor \ell/2 \rfloor}{n_1} \right\rfloor \geq n_2.$$

(d.2) **Subcase** $n = 2n_1 n_2$, n_1 even and $n_2 \geq 3$. We slightly modify the previous argument to obtain $\lfloor \frac{n-k}{2n_1} \rfloor + \lfloor \frac{k-2n_1}{2} \rfloor$ elements in $\langle y^{2n_1} \rangle \simeq C_n/C_{2n_1} \simeq C_{n_2}$. If

$$\left\lfloor \frac{n-k}{2n_1} \right\rfloor + \left\lfloor \frac{k-2n_1}{2} \right\rfloor \geq \frac{n-k-2n_1+1}{2n_1} + \frac{k-2n_1-1}{2} > n_2 - 1 = d(C_{n_2}),$$

then we are done, since in this case we obtain more than $d(C_{n_2})$ elements in $\langle y^{2n_1} \rangle$ which yields a product-one subsequence over C_n . But the last inequality above is equivalent to $k > 2n_1 + 3 + \frac{2}{n_1-1}$.

Since $k \geq \frac{n}{2} = n_1 n_2$, if $n_1 n_2 > 2n_1 + 3 + \frac{2}{n_1-1}$, then we are done. Since $n_1 \geq 4$ and $n_2 \geq 3$, it follows that $n_1 n_2 \geq 3n_1 > 2n_1 + 4 \geq 2n_1 + 3 + \frac{2}{n_1-1}$, hence we are done.

(d.3) **Subcase** $n = 2n_1 n_2$, n_2 even and $n_1 \geq 3$. Similar to the previous case, we obtain $\lfloor \frac{n-k}{n_1} \rfloor + \lfloor \frac{k-n_1}{2} \rfloor$ elements in $\langle y^{n_1} \rangle \simeq C_n/C_{n_1} \simeq C_{2n_2}$. If

$$\left\lfloor \frac{n-k}{n_1} \right\rfloor + \left\lfloor \frac{k-n_1}{2} \right\rfloor \geq \frac{n-k-n_1+1}{n_1} + \frac{k-n_1-1}{2} > 2n_2 - 1 = d(C_{2n_2}),$$

then we are done, since in this case we obtain more than $d(C_{2n_2})$ elements in $\langle y^{n_1} \rangle$ which yields a product-one subsequence over C_n . But the last inequality above is equivalent to $k > n_1 + 3 + \frac{4}{n_1-2}$.

Since $k \geq \frac{n}{2} = n_1 n_2$, if $n_1 n_2 > n_1 + 3 + \frac{4}{n_1-2}$, then we are done. Since $n_1 \geq 3$ and $n_2 \geq 4$, it follows that $n_1 n_2 \geq 4n_1 > n_1 + 7 \geq n_1 + 3 + \frac{4}{n_1-2}$, hence we are done.

(d.4) **Subcase** $n = 2^t = 2n_1 n_2$, $n_2 = 1$. In this case, $s \equiv n_1 - 1 \pmod{n}$, thus $s \equiv -1 \pmod{n_1}$. Let $S = \prod_{1 \leq i \leq k}^{\bullet} (xy^{\alpha_i}) \cdot \prod_{1 \leq i \leq n-k}^{\bullet} (y^{\beta_i})$. We are going to look at the α_i 's modulo $n/2$ and, for this, we consider the following values of k :

(d.4.i) Sub-subcase $k \geq n/2 + 3$. Since $k - 2 \geq n/2 + 1$, there exist exponents (say) $\alpha_1 \equiv \alpha_2 \pmod{n/2}$ and $\alpha_3 \equiv \alpha_4 \pmod{n/2}$. It implies that either

$$xy^{\alpha_1} \cdot xy^{\alpha_2} = 1 \quad \text{or} \quad xy^{\alpha_3} \cdot xy^{\alpha_4} = 1 \quad \text{or} \quad xy^{\alpha_1} \cdot xy^{\alpha_2} \cdot xy^{\alpha_3} \cdot xy^{\alpha_4} = 1.$$

In fact, the previous first two products are $y^{\alpha_2 - \alpha_1} \in \{1, y^{n/2}\}$ and $y^{\alpha_4 - \alpha_3} \in \{1, y^{n/2}\}$. If both of them are distinct than 1, then the third product is $y^{\alpha_2 - \alpha_1 + \alpha_4 - \alpha_3} = 1$, and we are done.

(d.4.ii) Sub-subcase $k = n/2 + 2$. Since $k - 2 = n/2$, there exist exponents (say) $\alpha_1 \equiv \alpha_2 \pmod{n/2}$ and either $[\alpha_3 \equiv \alpha_4 \pmod{n/2}]$ or $[\alpha_{i+3} \equiv i \pmod{n/2}]$ for $0 \leq i \leq$

$n/2 - 1$]. If $\alpha_3 \equiv \alpha_4 \pmod{n/2}$, then we use the same argument than Sub-subcase (d.4.i). Otherwise, $\alpha_{i+3} \equiv i \pmod{n/2}$ for $0 \leq i \leq n/2 - 1$ implies that

$$xy^{\alpha_{i+3}} \cdot xy^{\alpha_{i+5}} \cdot xy^{\alpha_{i+6}} \cdot xy^{\alpha_{i+4}} \in \{1, y^{n/2}\},$$

therefore either

$$\begin{aligned} xy^{\alpha_1} \cdot xy^{\alpha_2} &= 1 \quad \text{or} \\ xy^{\alpha_{i+3}} \cdot xy^{\alpha_{i+5}} \cdot xy^{\alpha_{i+6}} \cdot xy^{\alpha_{i+4}} &= 1 \quad \text{or} \\ xy^{\alpha_1} \cdot xy^{\alpha_2} \cdot xy^{\alpha_{i+3}} \cdot xy^{\alpha_{i+5}} \cdot xy^{\alpha_{i+6}} \cdot xy^{\alpha_{i+4}} &= 1, \end{aligned}$$

and we are done.

(d.4.iii) Sub-subcase $k = n/2 + 1$. Since $k - 2 = n/2 - 1$, there exist exponents (say) $\alpha_1 \equiv \alpha_2 \pmod{n/2}$ and either $[\alpha_3 \equiv \alpha_4 \pmod{n/2}]$ or [there exist $0 \leq \ell \leq n/2 - 1$ such that the sets $\{\alpha_{i+3}; 0 \leq i \leq n/2 - 2\}$ and $\{0, 1, 2, \dots, n/2 - 1\} \setminus \{\ell\}$ are the same modulo $n/2$]. If $\alpha_3 \equiv \alpha_4 \pmod{n/2}$, then we use the same argument than Sub-subcase (d.4.i). Otherwise, there exist $0 \leq \ell \leq n/2 - 1$ such that the sets $\{\alpha_{i+3}; 0 \leq i \leq n/2 - 2\}$ and $\{0, 1, 2, \dots, n/2 - 1\} \setminus \{\ell\}$ are the same modulo $n/2$. If $t \geq 4$ (that is, $n \geq 16$), then the set $\{\alpha_{i+3}; 0 \leq i \leq n/2 - 2\}$ has four consecutive elements, therefore we can apply the same argument as Subcase (d.4.ii). If $t = 3$ (that is, $n = 8$), then $|S \cap H| = n - k = 3$ and $xy^{\alpha_3} \cdot xy^{\alpha_4}$ generates another product in $C_n = C_8$. These four elements in C_8 yields a subsequence $T \mid (S \cap H) \cdot xy^{\alpha_3} \cdot xy^{\alpha_4}$ such that either $1 \in \pi(T)$ or $y^4 \in \pi(T)$. Therefore either

$$xy^{\alpha_1} \cdot xy^{\alpha_2} = 1 \quad \text{or} \quad 1 \in \pi(T) \quad \text{or} \quad 1 \in \pi(T \cdot xy^{\alpha_1} \cdot xy^{\alpha_2}),$$

thus we are done.

(d.4.iv) Sub-subcase $k = n/2$. Since $|S \cap H| = n/2$, there exists $T \mid S \cap H$ such that either $1 \in \pi(T)$ or $y^{n/2} \in \pi(T)$. Since $|S \cap N| = n/2$, either there exist exponents (say) $\alpha_1 \equiv \alpha_2 \pmod{n/2}$ or $[\alpha_{i+1} \equiv i \pmod{n/2}$ for $0 \leq i \leq n/2 - 1$]. If $\alpha_1 \equiv \alpha_2 \pmod{n/2}$, then either

$$xy^{\alpha_1} \cdot xy^{\alpha_2} = 1 \quad \text{or} \quad 1 \in \pi(T) \quad \text{or} \quad 1 \in \pi(T \cdot xy^{\alpha_1} \cdot xy^{\alpha_2}),$$

thus we are done. Otherwise, $\alpha_{i+1} \equiv i \pmod{n/2}$ for $0 \leq i \leq n/2 - 1$ implies that $xy^{\alpha_1} \cdot xy^{\alpha_3} \cdot xy^{\alpha_4} \cdot xy^{\alpha_2} \in \{1, y^{n/2}\}$. Therefore either

$$1 \in \pi(T) \quad \text{or} \quad xy^{\alpha_1} \cdot xy^{\alpha_3} \cdot xy^{\alpha_4} \cdot xy^{\alpha_2} = 1 \quad \text{or} \quad 1 \in \pi \left(T \cdot \prod_{1 \leq i \leq 4}^{\bullet} (xy^{\alpha_i}) \right),$$

hence we are done.

(d.5) **Subcase** $n = 2^t = 2n_1n_2$, $n_1 = 1$. In this case, $s \equiv n_2 + 1 \pmod{n}$. Let

$$S = S_1 \cdot S_2 \cdot S_3 \cdot S_4,$$

where

$$S_1 = \prod_{\substack{y^\alpha \mid S \\ \alpha \text{ odd}}}^{\bullet} y^\alpha, \quad S_2 = \prod_{\substack{y^\alpha \mid S \\ \alpha \text{ even}}}^{\bullet} y^\alpha, \quad S_3 = \prod_{\substack{xy^\alpha \mid S \\ \alpha \text{ odd}}}^{\bullet} xy^\alpha, \quad S_4 = \prod_{\substack{xy^\alpha \mid S \\ \alpha \text{ even}}}^{\bullet} xy^\alpha.$$

We have that

$$xy^\alpha \cdot xy^\beta = y^{\alpha(n/2+1)+\beta} = \begin{cases} y^{\alpha+\beta} & \text{if } \alpha \text{ is even,} \\ y^{\alpha+\beta+n/2} & \text{if } \alpha \text{ is odd.} \end{cases}$$

Let $C_{n/2} \simeq \langle y^2 \rangle$. We see that $S_2 \in \mathcal{F}(C_{n/2})$. Furthermore, for $i \in \{1, 3, 4\}$, the product of any two terms of S_i belongs to $C_{n/2}$. Therefore the terms can be grouped into pairs whose products belong to $C_{n/2}$. In this way, we obtain

$$L = \left\lfloor \frac{|S_1|}{2} \right\rfloor + |S_2| + \left\lfloor \frac{|S_3|}{2} \right\rfloor + \left\lfloor \frac{|S_4|}{2} \right\rfloor \geq \frac{n + |S_2| - 3}{2} \geq \frac{n-3}{2}$$

disjoint products in $C_{n/2}$. If $L > d(C_{n/2}) = n/2 - 1$, then S is not product-one free. Therefore we may assume that $L = n/2 - 1$. In particular, $|S_2| \leq 1$. If $|S_2| = 1$ and $L = n/2 - 1$, then $|S_1|$, $|S_3|$ and $|S_4|$ are all odd. It implies that there is one term left for each S_1 , S_3 and S_4 , and their product (in any order) lies in $C_{n/2}$, hence we obtain one element more in $C_{n/2}$.

From now on, we assume that $|S_2| = 0$. By Lemma 2.1, these $n/2 - 1$ terms of $C_{n/2}$ must comprise the same fixed generator of $C_{n/2}$, say equal to y^2 without loss of generality.

Suppose that $|S_3| \geq 3$ and let $xy^{\alpha_1} \cdot xy^{\alpha_2} \cdot xy^{\alpha_3} \in S_3$. By changing the order in which the elements are obtained, we can show that all terms of S_3 are equal. In fact,

$$xy^{\alpha_1} \cdot xy^{\alpha_2} = y^2 = xy^{\alpha_1} \cdot xy^{\alpha_3}$$

implies that $\alpha_2 \equiv \alpha_3 \pmod{n}$, and similarly $\alpha_1 \equiv \alpha_2 \equiv \alpha_3 \pmod{n}$. In particular, all terms of S_3 are equal. A similar result can be obtained if $|S_4| \geq 3$.

Since $n/2 \leq k = |S_3| + |S_4|$, it follows that $\max\{|S_3|, |S_4|\} \geq n/4 \geq 2$. If $|S_3| = |S_4| = 2$ and $n = 8$, then $L = 4 = n/2$, a contradiction. Therefore we assume that $\max\{|S_3|, |S_4|\} \geq 3$. If this maximum is $|S_4|$ and $xy^\alpha \in S_4$ (α is even), then $xy^\alpha \cdot xy^\alpha = y^{2\alpha} \in C_{n/4} = \langle y^4 \rangle \neq y^2$, a contradiction. Therefore, $|S_4| \leq 2$ and $|S_3| \geq 3$.

We have so far $|S_1| = n - k$, $|S_2| = 0$, $|S_3| \geq 3$, $|S_4| \leq 2$ and $|S_3| + |S_4| = k$. Consider the values of $|S_4|$:

(d.5.i) Sub-subcase $|S_4| = 2$. Let $S_4 = xy^{\alpha_1} \cdot xy^{\alpha_2}$, and let $(xy^\beta)^{[2]} \in S_3$ (that is, α_1 and α_2 are even, and β is odd). It follows that

$$y^{2\beta+n/2} = xy^\beta \cdot xy^\beta = y^2 = xy^{\alpha_1} \cdot xy^{\alpha_2} = y^{\alpha_1+\alpha_2}.$$

We reorder only these four terms, obtaining the product

$$xy^\beta \cdot xy^{\alpha_1} \cdot xy^\beta \cdot xy^{\alpha_2} = y^{\alpha_1+\beta+n/2} \cdot y^{\alpha_2+\beta+n/2} = y^{\alpha_1+\alpha_2+2\beta} = y^{n/2+4},$$

and we complete the previous product with the subsequence $(y^2)^{[n/4-2]}$ over $C_{n/2}$, obtaining a product-one subsequence.

(d.5.ii) Sub-subcase $|S_4| = 1$. We may assume without loss of generality that $S = y^{[n-k]} \cdot (xy^\alpha)^{[k-1]} \cdot xy^\beta$, where α is odd and β is even. Since $y^2 = (xy^\alpha)^2 = y^{2\alpha+n/2}$, we have that $\alpha \in \{n/4 + 1, 3n/4 + 1\}$. Suppose $\alpha = n/4 + 1$ (the case $\alpha = 3n/4 + 1$ is completely similar). If $k < n$, then it is possible to obtain both products

$$xy^\beta \cdot xy^{n/4+1} \cdot y = y^{n/4+\beta+2} \quad \text{and} \quad xy^{n/4+1} \cdot xy^\beta \cdot y = y^{3n/4+\beta+2}.$$

Both exponents are even numbers and, modulo n , some of them is at least $n/2$, therefore it is possible to complete such product with some y^2 's, obtaining a product-one subsequence. It implies that $k = n$ and $S = (xy^\alpha)^{[n-1]} \cdot xy^\beta$, where α is odd and β is even.

- (d.5.iii) Sub-subcase $|S_4| = 0$. We may assume without loss of generality that $S = y^{[n-k]} \cdot (xy^{n/4+1})^{[k]}$. If $k \leq n - 2$, then

$$xy^{n/4+1} \cdot y \cdot xy^{n/4+1} \cdot y = y^{n/2+2},$$

therefore it is possible to complete such product with $(y^2)^{[n/4-1]}$, thus S is not product-one free. If $k = n$, then $S = (xy^{n/4+1})^{[n]}$ and $(xy^{n/4+1})^n = 1$, thus S is not product-one free. Hence, $k = n - 1$.

It completes the proof. □

REFERENCES

- [1] J.D. Bovey, P. Erdős, I. Niven; *Conditions for a zero sum modulo n*. Canad. Math. Bull. Vol. **18** (1), (1975), 27-29.
- [2] F.E. Brochero Martínez, S. Ribas; *Extremal product-one free sequences in $C_q \rtimes_s C_m$* . J. Number Theory **204** (2019), 334-353.
- [3] F.E. Brochero Martínez, S. Ribas; *Extremal product-one free sequences in Dihedral and Dicyclic Groups*. Disc. Math. **341** (2018), 570-578.
- [4] W.D. Gao, A. Geroldinger, D.J. Grynkiewicz; *Inverse zero-sum problems III*. Acta Arith. **141.2** (2010), 103-152.
- [5] W.D. Gao, A. Geroldinger, W.A. Schmid; *Inverse zero-sum problems*. Acta Arith. **128.3** (2007), 245-279.
- [6] A. Geroldinger, D.J. Grynkiewicz; *The large Davenport constant I: Groups with a cyclic, index 2 subgroup*. J. Pure Appl. Algebra **217** (2013), 863-885.
- [7] D.J. Grynkiewicz; *Structural Additive Theory*. Springer (2013).
- [8] J.S. Oh, Q. Zhong; *On Erdős-Ginzburg-Ziv inverse theorems for dihedral and dicyclic groups*. Isr. J. Math. **238** (2020), 715-743.
- [9] J.S. Oh, Q. Zhong; *On minimal product-one sequences of maximal length over Dihedral and Dicyclic groups*. Commun. Korean Math. Soc. **35**, no. 1 (2020), 83-116.
- [10] W.A. Schmid; *Inverse zero-sum problems II*. Acta Arith. **143**, no. 4 (2010), 333-343.
- [11] D. Shanks; *Class number, a theory of factorization and genera*. In Proc. Symp. Pure Math. **20**, Providence, R.I.: AMS (1971), 415-440.
- [12] J.J. Zhuang, W.D. Gao; *Erdős-Ginzburg-Ziv theorem for dihedral groups of large prime index*. Europ. J. Combin. **26** (2005), 1053-1059.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, UFMG, BELO HORIZONTE, MG, 31270-901, BRAZIL,

Email address: `fbrocher@mat.ufmg.br`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE OURO PRETO, UFOP, OURO PRETO, MG, 35400-000, BRAZIL,

Email address: `savio.ribas@ufop.edu.br`