

ASYMPTOTIC EXPANSIONS FOR THE TRUNCATION ERROR IN RAMANUJAN-TYPE SERIES

LORENZ MILLA, 05/2022

ABSTRACT. Many of the fastest known algorithms to compute π involve generalized hypergeometric series, such as the Ramanujan-Sato series. In this paper, we investigate the rates of convergence for several such series and we give asymptotic expansions for the error of finite approximation.

For example, when using the first n terms of the Chudnovskys' series, we obtain the finite approximation $\pi_n \approx \pi$. It is known that the truncation error satisfies $|\pi_n - \pi| \approx 53360^{-3n}$. In this paper, we prove that the asymptotic expansion for the truncation error in the Chudnovskys' series is

$$|\pi_n - \pi| = 53360^{-3n} \cdot \frac{106\,720\sqrt{10\,005}\pi}{1\,672\,209\sqrt{n}} \cdot \exp\left(\frac{A_1}{n} + \frac{A_2}{n^2} + \frac{\delta_n}{n^3}\right),$$

with $0.006\,907 < \delta_n < 0.008\,429$ and the exact rational values of A_1 and A_2 :

$$A_1 = -\frac{1\,781\,843\,197\,433}{7\,456\,754\,505\,816},$$

$$A_2 = -\frac{1\,080\,096\,011\,925\,710\,088\,395}{3\,475\,199\,235\,000\,451\,148\,614\,116}.$$

Thus we demonstrate how to establish precise error bounds for the approximations for π obtained through Ramanujan-like series for $1/\pi$.

We also give asymptotic expansions for all known rational hypergeometric series for $1/\pi$ in the appendix.

Keywords: Chudnovsky Algorithm · error bounds · inequalities · hypergeometric series · approximations · asymptotic expansions · Stirling's formula

Mathematics Subject Classification (2010): 26D15 · 33C20 · 33B15

INTRODUCTION

In 1914, Srinivasa Ramanujan gave 17 series for the number π . The fastest among them (see [8, Eq. (44)]) is

$$(1a) \quad \frac{1}{\pi} = \frac{\sqrt{8}}{9801} \sum_{k=0}^{\infty} \frac{(4k)!}{(k!)^4} \frac{1103 + 26390k}{396^{4k}}.$$

The brothers David and Gregory Chudnovsky found an even faster series in 1988 (see [2, Eq. (1.5)]), which was used in most recent computations of π – it reads

$$(1b) \quad \frac{\sqrt{640320^3}}{12\pi} = \sum_{k=0}^{\infty} \frac{(-1)^k (6k)!}{(3k)! (k!)^3} \frac{13591409 + 545140134k}{640320^{3k}}.$$

More details can be found in [1]. In [4], we gave a detailed proof of this series using elliptic curves and the Picard Fuchs differential equation. In [5], we calculated the coefficients in the Chudnovskys' series using the integrality of certain non-holomorphic modular functions.

To calculate digits of π , one has to decide how many terms to use:

Definition 1. *The first n terms of the Chudnovskys' series (1b) yield the finite approximation $\pi_n \approx \pi$*

$$(2a) \quad \frac{1}{\pi_n} = \frac{12 \cdot 545\,140\,134}{\sqrt{640\,320^3}} \sum_{k=0}^{n-1} (-1)^k s_k,$$

$$(2b) \quad \text{with } s_k = \frac{(6k)!}{(3k)! (k!)^3} \frac{k+S}{640\,320^{3k}}$$

$$(2c) \quad \text{and } S = \frac{13\,591\,409}{545\,140\,134}.$$

Throughout the paper we denote the convergence rate of s_k by

$$(2d) \quad \varepsilon = \frac{6^6/3^3}{640\,320^3} = \frac{1\,728}{640\,320^3} = 53\,360^{-3}.$$

In Section 1, we prove the following asymptotic expansion of the truncation error in the Chudnovskys' series:

Theorem 1. *For all $n \geq 1$, the omitted terms of the Chudnovskys' series (1b) admit the asymptotic expansion*

$$(3a) \quad \left| \sum_{k=n}^{\infty} (-1)^k s_k \right| = \frac{s_n}{1+\varepsilon} \left(1 + \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n \varepsilon}{n^3} \right).$$

For the definition of s_k , S and ε see (2b), (2c) and (2d). The coefficients of this expansion are the rational numbers

$$(3b) \quad a_1 = \frac{\varepsilon}{2(1+\varepsilon)} = \frac{1}{303\,862\,746\,112\,002}$$

$$(3c) \quad \text{and } a_2 = \frac{S\varepsilon}{1+\varepsilon} - \frac{23\varepsilon - 4\varepsilon^2}{36(1+\varepsilon)^2} = \frac{-62\,186\,213\,362\,465}{15\,388\,761\,412\,454\,497\,761\,254\,741\,334}.$$

The error term e_n in this expansion satisfies

$$(3d) \quad 0.3216 < e_n < 0.6704.$$

In Section 2, we derive the recursion formula (11) which allows one to compute the rational coefficients a_k of the asymptotic expansion (3a) more efficiently and to higher orders. But this involves a formal power series (9) which is divergent, thus the results from Section 2 do not readily yield rigorous error bounds like (3d) in Theorem 1.

In Section 3, we use Stirling's approximation and Theorem 1 to prove the following asymptotic expansion for $|\pi_n - \pi|$ mentioned in the abstract:

Theorem 2. *If we denote by π_n the approximation we obtain by using the first n terms in the Chudnovskys' series as in (2a), then we have the asymptotic expansion*

$$(4a) \quad |\pi_n - \pi| = 53360^{-3n} \cdot \frac{A_0}{\sqrt{n}} \cdot \exp\left(\frac{A_1}{n} + \frac{A_2}{n^2} + \frac{\delta_n}{n^3}\right).$$

Here, the error term δ_n satisfies

$$(4b) \quad 0.006\,907 < \delta_n < 0.008\,429$$

for all $n \geq 1$ and the coefficients A_0 , A_1 and A_2 can be expressed in terms of S from (2c), ε from (2d), a_1 from (3b) and a_2 from (3c):

$$(4c) \quad A_0 = \frac{6\sqrt{\pi} \cdot 545\,140\,134}{\sqrt{640\,320^3}(1+\varepsilon)} = \frac{106\,720 \cdot \sqrt{10\,005\pi}}{1\,672\,209},$$

$$(4d) \quad A_1 = a_1 - \frac{19}{72} + S = -\frac{1\,781\,843\,197\,433}{7\,456\,754\,505\,816},$$

$$(4e) \quad A_2 = a_2 - \frac{a_1^2}{2} - \frac{S^2}{2} = -\frac{1\,080\,096\,011\,925\,710\,088\,395}{3\,475\,199\,235\,000\,451\,148\,614\,116}.$$

We have chosen to calculate all terms exactly up to order n^{-2} , with upper and lower bounds on the remainders r like $\frac{a}{n^3} < r(n) < \frac{b}{n^3}$. All parts of this paper can be verified without specialized software.

In Section 4, we will do a numerical analysis showing that the bounds we have proven are close to optimal. We will also generalize our results to other hypergeometric series and we will give an asymptotic expansion for Ramanujan's series (1a).

In Appendix A, we prove an asymptotic expansion of a product of Pochhammer symbols. This enables us to give asymptotic expansions for all known rational hypergeometric series for $1/\pi$ in Appendix B.

1. PROOF OF THEOREM 1

In this section, we prove Theorem 1 using only basic methods of analysis. For the definition of s_n , S and ε keep (2b), (2c) and (2d) from Definition 1 in mind.

Lemma 1. *For all $n \geq 1$ there is φ_n with $0.05 < \varphi_n < 2.19$ and*

$$\frac{s_n}{\varepsilon \cdot s_{n-1}} = \sum_{j=0}^{\infty} \frac{b_j}{n^j} = 1 - \frac{1}{2n} + \frac{\frac{5}{36} - S}{n^2} + \frac{\frac{5}{72} - \frac{S}{2} + S^2}{n^3} + \frac{\varphi_n}{n^4}.$$

If $n \geq 2$, we have that $\varphi_n < 0.11$ and

$$\frac{s_n}{\varepsilon \cdot s_{n-1}} < \exp\left(-\frac{1}{2n} - \frac{S - \frac{1}{72}}{n^2} + \frac{0.67}{n^3}\right).$$

Proof. The definition of $s_n = \frac{(6n)!}{(3n)!(n!)^3} \frac{n+S}{640 \cdot 320^{3n}}$ in (2b) yields for all $n \geq 1$:

$$\begin{aligned} (5) \quad \frac{s_n}{\varepsilon \cdot s_{n-1}} &= \left(1 - \frac{1}{6n}\right) \left(1 - \frac{1}{2n}\right) \left(1 - \frac{5}{6n}\right) \cdot \frac{S+n}{S+n-1} \\ &= \left(1 - \frac{3}{2n} + \frac{23}{36n^2} - \frac{5}{72n^3}\right) \cdot \left(1 + \sum_{k=0}^{\infty} \frac{(1-S)^k}{n^{k+1}}\right) \\ &= 1 - \frac{1}{2n} + \frac{\frac{5}{36} - S}{n^2} + \frac{\frac{5}{72} - \frac{S}{2} + S^2}{n^3} + \frac{\varphi_n}{n^4}, \end{aligned}$$

$$\text{with } \frac{\varphi_n}{n^4} = \sum_{k=4}^{\infty} \frac{\left(\frac{5}{72} - \frac{23S}{36} + \frac{3S^2}{2} - S^3\right)(1-S)^{k-4}}{n^k} = \frac{\frac{5}{72} - \frac{23S}{36} + \frac{3S^2}{2} - S^3}{n^4 \left(1 - \frac{1-S}{n}\right)}.$$

This proves $0.05 < \varphi_n < 2.19$ for $n \geq 1$ and $\varphi_n < 0.11$ for $n \geq 2$. Next, (5) yields

$$\begin{aligned} \ln\left(\frac{s_n}{\varepsilon \cdot s_{n-1}}\right) &= \ln\left(1 - \frac{1}{6n}\right) + \ln\left(1 - \frac{1}{2n}\right) + \ln\left(1 - \frac{5}{6n}\right) \\ &\quad + \ln\left(1 + \frac{S}{n}\right) - \ln\left(1 - \frac{1-S}{n}\right). \end{aligned}$$

Using $\ln(1-x) = -\sum_{k=1}^{\infty} \frac{x^k}{k}$ we obtain:

$$\begin{aligned} \ln\left(\frac{s_n}{\varepsilon \cdot s_{n-1}}\right) &= \sum_{k=1}^{\infty} \frac{(1-S)^k - (-S)^k - \left(\frac{1}{6}\right)^k - \left(\frac{1}{2}\right)^k - \left(\frac{5}{6}\right)^k}{k \cdot n^k} \\ &< -\frac{1}{2n} - \frac{S - \frac{1}{72}}{n^2} + \sum_{k=3}^{\infty} \frac{1}{k \cdot n^k}. \end{aligned}$$

For $n \geq 2$ we have $\sum_{k=3}^{\infty} \frac{1}{k \cdot n^k} < \frac{1}{3n^3} \sum_{k=0}^{\infty} \frac{1}{2^k} = \frac{2}{3n^3}$ which proves the lemma. \square

Lemma 2. *For $n \geq 1$ and $k \geq 1$ we have that*

$$\frac{s_{n+k}}{\varepsilon^k \cdot s_n} < 1 - \frac{k}{2n} + \frac{\frac{3}{8}k^2 + \left(\frac{19}{72} - S\right)k}{n^2} + \frac{1.2k^6}{n^3}.$$

Proof. Lemma 1 yields

$$\frac{s_{n+k}}{\varepsilon^k \cdot s_n} = \prod_{j=1}^k \frac{s_{n+j}}{\varepsilon \cdot s_{n+j-1}} < \exp \left(\sum_{j=1}^k \left(-\frac{1}{2(n+j)} - \frac{S - \frac{1}{72}}{(n+j)^2} + \frac{0.67}{(n+j)^3} \right) \right).$$

For $x < 0$, we have that $\exp(x) < 1 + x + \frac{1}{2}x^2$. Since the sum is negative, we obtain

$$\begin{aligned} \frac{s_{n+k}}{\varepsilon^k \cdot s_n} &< 1 + \sum_{j=1}^k \left(-\frac{1}{2(n+j)} - \frac{S - \frac{1}{72}}{(n+j)^2} + \frac{0.67}{(n+j)^3} \right) \\ &\quad + \frac{1}{2} \left(\sum_{j=1}^k \left(\frac{1}{2(n+j)} + \frac{S - \frac{1}{72}}{(n+j)^2} - \frac{0.67}{(n+j)^3} \right) \right)^2. \end{aligned}$$

We estimate $\frac{1}{n} - \frac{j}{n^2} < \frac{1}{n+j} < \frac{1}{n} - \frac{j}{n^2} + \frac{j^2}{n^3}$ as well as $\frac{1}{n^2} - \frac{2j}{n^3} < \frac{1}{(n+j)^2} < \frac{1}{n^2}$ and obtain

$$\begin{aligned} \frac{s_{n+k}}{\varepsilon^k \cdot s_n} &< 1 + \sum_{j=1}^k \left(-\frac{1}{2n} + \frac{j - S + \frac{1}{72}}{n^2} + \frac{2j(S - \frac{1}{72}) + 0.67}{n^3} \right) \\ &\quad + \frac{1}{2} \left(\sum_{j=1}^k \left(\frac{1}{2n} - \frac{j}{2n^2} + \frac{j^2}{2n^3} + \frac{S - \frac{1}{72}}{n^2} \right) \right)^2 \\ &= 1 - \frac{k}{2n} + \frac{\frac{k(k+1)}{4} - Sk + \frac{k}{72}}{n^2} + \frac{k(k+1)(S - \frac{1}{72}) + 0.67k}{n^3} \\ &\quad + \frac{1}{2} \left(\frac{k}{2n} - \frac{k(k+1)}{4n^2} + \frac{k(k+1)(2k+1)}{12n^3} + \frac{k(S - \frac{1}{72})}{n^2} \right)^2 \\ &\leq 1 - \frac{k}{2n} + \frac{\frac{1}{4}k^2 + (\frac{19}{72} - S)k}{n^2} + \frac{0.7k^2}{n^3} + \frac{1}{2} \left(\frac{k}{2n} + \frac{k^3}{2n^3} \right)^2 \\ &\leq 1 - \frac{k}{2n} + \frac{\frac{1}{4}k^2 + (\frac{19}{72} - S)k}{n^2} + \frac{0.7k^2}{n^3} + \frac{k^2}{8n^2} + \frac{k^4}{4n^4} + \frac{k^6}{8n^6} \\ &\leq 1 - \frac{k}{2n} + \frac{\frac{3}{8}k^2 + (\frac{19}{72} - S)k}{n^2} + \frac{1.2k^6}{n^3}. \quad \square \end{aligned}$$

Proof of upper bound in Theorem 1: Since s_n decreases monotonically to zero, we obtain

$$\begin{aligned} \left| \sum_{k=n}^{\infty} (-1)^k s_k \right| &= \sum_{k=0}^{\infty} (s_{n+2k} - s_{n+2k+1}) = \sum_{k=0}^{\infty} s_{n+2k} \cdot \left(1 - \varepsilon \cdot \frac{s_{n+2k+1}}{\varepsilon \cdot s_{n+2k}} \right) \\ (6) \quad &= s_n \cdot \sum_{k=0}^{\infty} \varepsilon^{2k} \cdot \underbrace{\frac{s_{n+2k}}{\varepsilon^{2k} \cdot s_n}}_{=Q_{n,k}} \cdot \underbrace{\left(1 - \varepsilon \cdot \frac{s_{n+2k+1}}{\varepsilon \cdot s_{n+2k}} \right)}_{=P_{n+2k}}. \end{aligned}$$

Lemma 2 tells us that

$$\begin{aligned} Q_{n,k} = \frac{s_{n+2k}}{\varepsilon^{2k} \cdot s_n} &< 1 - \frac{2k}{2n} + \frac{\frac{3}{8}(2k)^2 + (\frac{19}{72} - S) \cdot 2k}{n^2} + \frac{1.2(2k)^6}{n^3} \\ &= 1 - \frac{k}{n} + \frac{\frac{3}{2}k^2 + (\frac{19}{36} - 2S)k}{n^2} + \frac{76.8k^6}{n^3}. \end{aligned}$$

From Lemma 1 we deduce

$$\begin{aligned} P_{n+2k} &= 1 - \varepsilon \cdot \frac{s_{n+2k+1}}{\varepsilon \cdot s_{n+2k}} \\ &< 1 - \varepsilon \cdot \left(1 - \frac{1}{2(n+2k+1)} + \frac{\frac{5}{36} - S}{(n+2k+1)^2} + \frac{0.0576}{(n+2k+1)^3} \right) \\ &= 1 - \varepsilon + \frac{\varepsilon}{2(n+2k+1)} - \frac{(\frac{5}{36} - S)\varepsilon}{(n+2k+1)^2} - \frac{0.0576\varepsilon}{(n+2k+1)^3}. \end{aligned}$$

Here we write $K = 2k + 1$ and estimate $\frac{1}{n+K} < \frac{1}{n} - \frac{K}{n^2} + \frac{K^2}{n^3}$, as well as $\frac{1}{(n+K)^2} > \frac{1}{n^2} - \frac{2K}{n^3}$ and $\frac{1}{(n+K)^3} > \frac{1}{n^3}$:

$$\begin{aligned} P_{n+2k} &< 1 - \varepsilon + \frac{\varepsilon}{2n} - \frac{\varepsilon K}{2n^2} + \frac{\varepsilon K^2}{2n^3} - \frac{(\frac{5}{36} - S)\varepsilon}{n^2} + \frac{2(\frac{5}{36} - S)\varepsilon K}{n^3} - \frac{0.0576\varepsilon}{n^3} \\ &= 1 - \varepsilon + \frac{\varepsilon}{2n} - \frac{\frac{\varepsilon K}{2} + (\frac{5}{36} - S)\varepsilon}{n^2} + \frac{\frac{\varepsilon}{2}K^2 + 2(\frac{5}{36} - S)\varepsilon K - 0.0576\varepsilon}{n^3} \\ &< 1 - \varepsilon + \frac{\varepsilon}{2n} - \frac{(k + \frac{23}{36} - S)\varepsilon}{n^2} + \frac{(2k^2 + (\frac{23}{9} - 4S)k + 0.67032)\varepsilon}{n^3} \\ &\leq 1 - \varepsilon + \frac{\varepsilon}{2n} - \frac{(k + \frac{23}{36} - S)\varepsilon}{n^2} + \frac{(4.456k^2 + 0.67032)\varepsilon}{n^3}. \end{aligned}$$

Since both estimates are positive, we may multiply them and obtain

$$\begin{aligned} Q_{n,k} \cdot P_{n+2k} &< \left(1 - \frac{k}{n} + \frac{\frac{3}{2}k^2 + (\frac{19}{36} - 2S)k}{n^2} + \frac{76.8k^6}{n^3} \right) \\ &\quad \cdot \left(1 - \varepsilon + \frac{\varepsilon}{2n} - \frac{(k + \frac{23}{36} - S)\varepsilon}{n^2} + \frac{(4.456k^2 + 0.67032)\varepsilon}{n^3} \right) \\ &= 1 - \varepsilon + \frac{\frac{\varepsilon}{2} - (1 - \varepsilon)k}{n} + \frac{\alpha_2(k)}{n^2} + R_3(n, k), \end{aligned}$$

$$\begin{aligned} \text{with } \alpha_2(k) &= \left(\frac{3}{2}k^2 + \left(\frac{19}{36} - 2S \right) k \right) \cdot (1 - \varepsilon) - k \cdot \frac{\varepsilon}{2} - \left(k + \frac{23}{36} - S \right) \varepsilon \\ &= \frac{3}{2}(1 - \varepsilon)k^2 + \left[\left(\frac{19}{36} - 2S \right) (1 - \varepsilon) - \frac{3}{2}\varepsilon \right] k - \left(\frac{23}{36} - S \right) \varepsilon. \end{aligned}$$

To estimate the remaining term $R_3(n, k)$, we first omit all negative terms, then we use $\frac{1}{n^l} \leq \frac{1}{n^3}$ for $l \geq 3$ and $k^l \leq k^8$ for $1 \leq l \leq 8$. Since $\varepsilon < 10^{-14}$, this yields

$$R_3(n, k) < \frac{76.9k^8 + 0.67032\varepsilon}{n^3}.$$

From (6) we obtain

$$\begin{aligned} \frac{|\sum_{k=n}^{\infty} (-1)^k s_k|}{s_n} &= \sum_{k=0}^{\infty} \varepsilon^{2k} \cdot Q_{n,k} \cdot P_{n+2k} \\ &< \sum_{k=0}^{\infty} \varepsilon^{2k} \left(1 - \varepsilon + \frac{\frac{\varepsilon}{2} - (1 - \varepsilon)k}{n} + \frac{\alpha_2(k)}{n^2} + \frac{76.9k^8 + 0.67032\varepsilon}{n^3} \right) \\ &=: c_0 + \frac{c_1}{n} + \frac{c_2}{n^2} + \frac{c_3}{n^3}. \end{aligned}$$

The numbers c_0 , c_1 and c_2 can be computed using the geometric series and its moments $\sum_{k=0}^{\infty} k \varepsilon^{2k} = \frac{\varepsilon^2}{(1-\varepsilon^2)^2}$ and $\sum_{k=0}^{\infty} k^2 \varepsilon^{2k} = \frac{\varepsilon^2(1+\varepsilon^2)}{(1-\varepsilon^2)^3}$:

$$c_0 = (1 - \varepsilon) \cdot \frac{1}{1 - \varepsilon^2} = \frac{1}{1 + \varepsilon},$$

$$c_1 = \frac{\varepsilon}{2} \cdot \frac{1}{1-\varepsilon^2} - (1-\varepsilon) \cdot \frac{\varepsilon^2}{(1-\varepsilon^2)^2} = \frac{\varepsilon}{2(1+\varepsilon)^2},$$

$$c_2 = \frac{3}{2}(1-\varepsilon) \cdot \frac{\varepsilon^2(1+\varepsilon^2)}{(1-\varepsilon^2)^3} + \left[\left(\frac{19}{36} - 2S \right) (1-\varepsilon) - \frac{3}{2}\varepsilon \right] \cdot \frac{\varepsilon^2}{(1-\varepsilon^2)^2} - \frac{\left(\frac{23}{36} - S \right) \varepsilon}{1-\varepsilon^2}$$

$$= \frac{\varepsilon S}{(1+\varepsilon)^2} + \frac{4\varepsilon^2 - 23\varepsilon}{36(1+\varepsilon)^3}.$$

Here we recognize the values of $a_1 = (1+\varepsilon)c_1$ and $a_2 = (1+\varepsilon)c_2$ from Theorem 1. To estimate c_3 , we use $\varepsilon^k \cdot k^8 \leq \varepsilon$ (valid for $k \geq 1$) and thus

$$\sum_{k=0}^{\infty} \varepsilon^{2k} \cdot k^8 = \sum_{k=1}^{\infty} \varepsilon^{2k} \cdot k^8 < \sum_{k=1}^{\infty} \varepsilon^k \cdot \varepsilon = \frac{\varepsilon^2}{1-\varepsilon},$$

which implies that

$$c_3 < 76.9 \cdot \frac{\varepsilon^2}{1-\varepsilon} + 0.67032\varepsilon \cdot \frac{1}{1-\varepsilon^2} < \frac{0.6704\varepsilon}{1+\varepsilon}.$$

This proves the upper bound, $e_n < 0.6704$. □

Proof of lower bound in Theorem 1: Since s_n decreases monotonically to zero, we obtain

$$\left| \sum_{k=n-1}^{\infty} (-1)^k s_k \right| = s_{n-1} - \left| \sum_{k=n}^{\infty} (-1)^k s_k \right|,$$

which implies that

$$(7a) \quad \underbrace{\frac{1+\varepsilon}{s_{n-1}} \cdot \left| \sum_{k=n-1}^{\infty} (-1)^k s_k \right|}_{=f(n-1)} = 1 + \varepsilon - \underbrace{\frac{1+\varepsilon}{s_n} \cdot \left| \sum_{k=n}^{\infty} (-1)^k s_k \right|}_{=f(n)} \cdot \frac{s_n}{s_{n-1}}.$$

Here we observe a relation between $f(n-1)$ and $-f(n)$, i.e.

$$(7b) \quad f(n-1) = 1 + \varepsilon - f(n) \cdot \frac{s_n}{\varepsilon \cdot s_{n-1}} \cdot \varepsilon,$$

which allows us to transform the already proven upper bound $e_n < 0.6704$ from Theorem 1 into a lower bound: Using $e_n < 0.6704$ and Lemma 1 yields for $n \geq 2$

$$f(n) \cdot \frac{s_n}{\varepsilon \cdot s_{n-1}} < \left(1 + \frac{\varepsilon}{2(1+\varepsilon)n} + \frac{\frac{\varepsilon S}{(1+\varepsilon)} + \frac{4\varepsilon^2 - 23\varepsilon}{36(1+\varepsilon)^2}}{n^2} + \frac{0.6704\varepsilon}{n^3} \right)$$

$$\cdot \left(1 - \frac{1}{2n} + \frac{\frac{5}{36} - S}{n^2} + \frac{0.0576}{n^3} + \frac{0.11}{n^4} \right)$$

$$< 1 - \frac{1}{2(1+\varepsilon)n} + \frac{C}{n^2} + \frac{0.113}{n^3},$$

$$(8) \quad \text{with } C = \left(\frac{\varepsilon S}{1+\varepsilon} + \frac{4\varepsilon^2 - 23\varepsilon}{36(1+\varepsilon)^2} \right) + \left(\frac{5}{36} - S \right) - \frac{\varepsilon}{4(1+\varepsilon)}$$

$$= \frac{5 - 22\varepsilon}{36(1+\varepsilon)^2} - \frac{S}{1+\varepsilon}.$$

This proves the lower bound on $f(n-1)$ for $n \geq 2$:

$$f(n-1) = 1 + \varepsilon - \varepsilon \cdot f(n) \cdot \frac{s_n}{\varepsilon \cdot s_{n-1}}$$

$$> 1 + \frac{\varepsilon}{2(1+\varepsilon)n} - \frac{C\varepsilon}{n^2} - \frac{0.113\varepsilon}{n^3}.$$

For $n \geq 2$ we have that $\frac{1}{(n+1)^2} < \frac{1}{n^2} - \frac{1}{n^3}$. This shows that for $n \geq 2$ we have that

$$\begin{aligned} f(n) &> 1 + \frac{\varepsilon}{2(1+\varepsilon)(n+1)} - \frac{C\varepsilon}{(n+1)^2} - \frac{0.113\varepsilon}{(n+1)^3} \\ &> 1 + \frac{\varepsilon}{2(1+\varepsilon)n} - \frac{\varepsilon}{2(1+\varepsilon)n^2} + \frac{\varepsilon}{2(1+\varepsilon)n^3(1+1/n)} - \frac{C\varepsilon}{n^2} + \frac{C\varepsilon}{n^3} - \frac{0.113\varepsilon}{n^3} \\ &> 1 + \frac{\varepsilon}{2(1+\varepsilon)n} - \frac{\frac{\varepsilon}{2(1+\varepsilon)} + C\varepsilon}{n^2} + \frac{0.334\varepsilon}{n^3} = 1 + \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{0.334\varepsilon}{n^3}. \end{aligned}$$

Using the value of C from (8), we recognize a_1 and a_2 as in Theorem 1 and we have proven $e_n > 0.334$ for $n \geq 2$. In the case $n = 1$ we have:

$$\frac{\sqrt{640320^3}}{12\pi \cdot 545\,140\,134} = \sum_{k=0}^{\infty} (-1)^k s_k = s_0 - \left| \sum_{k=1}^{\infty} (-1)^k s_k \right|,$$

which implies that

$$\ln \left(\left| \sum_{k=1}^{\infty} (-1)^k s_k \right| \cdot \frac{1+\varepsilon}{s_1} \right) = \ln \left(\frac{\left(S - \frac{\sqrt{640320^3}}{12\pi \cdot 545\,140\,134} \right) (1+\varepsilon)}{\frac{720}{6 \cdot 1^3} \cdot \frac{1+S}{640320^3}} \right) > 0.20767\varepsilon.$$

This yields $a_1 + a_2 + e_1\varepsilon > 0.20767\varepsilon$ and $e_1 > 0.3216$, which finishes the proof of the lower bound on e_n and of Theorem 1. \square

2. SHARPER ESTIMATES IN HIGHER ORDERS

Using the terms s_n of the Chudnovskys' series from Definition 1, we denote the truncated part of the series by $f(n)$ as in (7a). This can be written as a formal power series

$$(9) \quad f(n) = \frac{1+\varepsilon}{s_n} \cdot \left| \sum_{k=n}^{\infty} (-1)^k s_k \right| =: \sum_{k=0}^{\infty} \frac{a_k}{n^k},$$

which we use to define the coefficients a_k . For $k \leq 2$, this definition is consistent with the values of a_k from Theorem 1. In this section, we explain a method how to compute the rational coefficients a_k efficiently to higher orders (see (11)), but without error bounds like (3d).

In Lemma 1 we proved

$$(10) \quad \begin{aligned} \frac{s_n}{\varepsilon \cdot s_{n-1}} &= 1 - \frac{1}{2n} + \frac{\frac{5}{36} - S}{n^2} + \frac{\frac{5}{72} - \frac{S}{2} + S^2}{n^3} \\ &+ \sum_{k=4}^{\infty} \frac{\left(\frac{5}{72} - \frac{23S}{36} + \frac{3S^2}{2} - S^3 \right) (1-S)^{k-4}}{n^k} =: \sum_{k=0}^{\infty} \frac{b_k}{n^k}, \end{aligned}$$

which gives $\frac{s_n}{s_{n-1}} = \varepsilon \cdot \sum_{k=0}^{\infty} \frac{b_k}{n^k}$ as a power series. Thus (7b) reads

$$f(n-1) = 1 + \varepsilon - f(n) \cdot \frac{s_n}{s_{n-1}},$$

$$\text{which yields} \quad \sum_{k=0}^{\infty} \frac{a_k}{(n-1)^k} = 1 + \varepsilon - \sum_{k=0}^{\infty} \frac{a_k}{n^k} \cdot \varepsilon \cdot \sum_{k=0}^{\infty} \frac{b_k}{n^k}.$$

Using $\frac{1}{(1-x)^k} = \sum_{l=0}^{\infty} \binom{l+k-1}{k-1} \cdot x^l$ with $x = \frac{1}{n}$ yields

$$a_0 + \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \frac{a_k \cdot \binom{l+k-1}{k-1}}{n^{k+l}} = 1 + \varepsilon - \varepsilon \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{a_k \cdot b_l}{n^{k+l}}.$$

Here, $b_0 = 1$ yields $a_0 = 1$ and equating the coefficients for $N = k + l \geq 1$ gives the recursion formula:

$$(11) \quad -(1 + \varepsilon) \cdot a_N = \varepsilon \cdot b_N + \sum_{k=1}^{N-1} a_k \cdot \left(\binom{N-1}{k-1} + \varepsilon \cdot b_{N-k} \right).$$

Using recursion (11) and the values of b_k from (10), we compute further terms of the sequence a_k :

$$\begin{aligned} a_1 &= \frac{1}{303\,862\,746\,112\,002} \approx 3.2910 \cdot 10^{-15}, \\ a_2 &= \frac{-62\,186\,213\,362\,465}{15\,388\,761\,412\,454\,497\,761\,254\,741\,334} \approx -4.0410 \cdot 10^{-15}, \\ a_3 &= \frac{20\,630\,598\,257\,083\,699\,331\,942\,595\,295}{4\,676\,071\,302\,050\,834\,345\,503\,446\,765\,524\,996\,702\,890\,668} \approx 4.4120 \cdot 10^{-15}, \\ a_4 &= \frac{-1\,933\,230\,018\,398\,723\,806\,508\,418\,321\,549\,998\,750\,727\,169}{405\,966\,819\,101\,911\,798\,225\,167\,895\,714\,948\,764\,118\,181\,484\,848\,996\,742\,096} \\ &\approx -4.7620 \cdot 10^{-15} \\ a_{10} &\approx -6.6157 \cdot 10^{-15}, \\ a_{100} &\approx -1.4296 \cdot 10^5, \\ a_{1000} &\approx -1.3214 \cdot 10^{1049}. \end{aligned}$$

Figure 1 shows the values of $|a_k|$ for $k \leq 200$. We observe numerically that the a_k have alternating signs for most k and that they grow roughly like $|a_k| \approx k! / q^k$ with $q = \ln(53360^3)$, thus (9) diverges for all $n \in \mathbb{N}$.




FIGURE 1. Values of $|a_k|$. Simple dot (\cdot): $a_k a_{k+1} < 0$. Dot with circle (\odot): $a_k a_{k+1} > 0$. Gray line: rough approximations $|a_k| \approx k! / q^k$ with $q = \ln(53360^3)$.

3. PROOF OF THEOREM 2

From $\frac{1}{\pi} - \frac{1}{\pi_n} = \frac{\pi_n - \pi}{\pi \pi_n}$ we deduce using Definition 1:

$$(12) \quad |\pi_n - \pi| = |\pi \pi_n| \cdot \left| \frac{1}{\pi} - \frac{1}{\pi_n} \right| = \pi \pi_n \frac{12 \cdot 545\,140\,134}{\sqrt{640320^3}} \left| \sum_{k=n}^{\infty} (-1)^k s_k \right|$$

$$(13) \quad = \frac{12 \cdot 545\,140\,134 \pi^2}{\sqrt{640320^3} (1 + \varepsilon)} \cdot \frac{\pi_n}{\pi} \cdot s_n \cdot \underbrace{\frac{1 + \varepsilon}{s_n} \left| \sum_{k=n}^{\infty} (-1)^k s_k \right|}_{\text{Theorem 1}}.$$

Most of the work towards proving Theorem 2 consists in establishing the error bound in Theorem 1. We still have to find approximations for s_n (Lemma 3) and for $\frac{\pi_n}{\pi}$ (Lemma 4).

Lemma 3. For all $n \geq 1$ there is h_n with $-0.0024 < h_n < -0.0005$ and

$$s_n = \frac{\varepsilon^n}{2 \pi^{3/2} \sqrt{n}} \cdot \exp \left(\frac{S - \frac{19}{72}}{n} + \frac{-\frac{1}{2} S^2}{n^2} + \frac{\frac{131}{15552} + \frac{1}{3} S^3}{n^3} + \frac{h_n}{n^4} \right).$$

Proof. We start with Stirling's formula $n! = \sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^n \cdot e^{r_n}$ with the enveloping series $r_n = \sum_{j=1}^{\infty} \frac{B_{2j}}{2j(2j-1)n^{2j-1}}$ proved in [9] which tells us that:

$$\frac{1}{12n} - \frac{1}{360n^3} + \frac{1}{1260n^5} - \frac{1}{1680n^7} < r_n < \frac{1}{12n} - \frac{1}{360n^3} + \frac{1}{1260n^5}.$$

This yields $\frac{(6n)!}{(3n)!(n!)^3} = \frac{1728^n}{2(\pi n)^{3/2}} \cdot e^{y_n}$ with the following bounds for $y_n = r_{6n} - r_{3n} - 3r_n$:

$$y_n < -\frac{19}{72n} + \frac{131}{15552n^3} - \frac{3337}{1399680n^5} + \frac{3281}{1837080n^7},$$

$$y_n > -\frac{19}{72n} + \frac{131}{15552n^3} - \frac{3337}{1399680n^5} - \frac{1}{470292480n^7}.$$

To approximate $S + n = n \cdot \exp \ln(1 + x)$ with $x = \frac{S}{n} > 0$, we use

$$x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 < \ln(1 + x) < x - \frac{1}{2}x^2 + \frac{1}{3}x^3.$$

Combining these two error bounds, we obtain

$$s_n = \frac{(6n)!}{(3n)!(n!)^3} \cdot \frac{S + n}{640320^{3n}} = \frac{1728^n}{2(\pi n)^{3/2}} \cdot e^{y_n} \cdot \frac{n}{640320^{3n}} \cdot \left(1 + \frac{S}{n}\right)$$

$$= \frac{53360^{-3n}}{2 \pi^{3/2} \sqrt{n}} \cdot \exp \left(-\frac{19}{72n} + \frac{131}{15552n^3} + \frac{S}{n} - \frac{\frac{1}{2} S^2}{n^2} + \frac{\frac{1}{3} S^3}{n^3} + \frac{h_n}{n^4} \right),$$

where $\frac{h_n}{n^4}$ contains the terms of higher order

$$-\frac{3337}{1399680n^5} - \frac{1}{470292480n^7} - \frac{\frac{1}{4} S^4}{n^4} < \frac{h_n}{n^4} < -\frac{3337}{1399680n^5} + \frac{3281}{1837080n^7},$$

which proves $-0.0024 < h_n < -0.0005$ and the lemma \square

Lemma 4. For $n \geq 1$ we have the really weak estimate $\pi_n = \pi \exp \left(\frac{f_n \varepsilon}{n^{10}} \right)$ with $-4 < f_n < 4$.

Proof. Since s_n decreases monotonically to zero, (12) yields

$$|\pi_n - \pi| < \pi \pi_n \frac{12 \cdot 545\,140\,134}{\sqrt{640320^3}} s_n.$$

Using $s_n < \frac{\varepsilon^n}{2 \pi^{3/2}}$ (see Lemma 3) and $\pi_n < 3.2$ yields

$$\frac{|\pi_n - \pi|}{\pi} < \frac{12 \cdot 545\,140\,134}{\sqrt{640320^3}} \pi_n s_n < 3.7 \varepsilon^n \leq \frac{3.7 \varepsilon}{n^{10}},$$

and thus $\pi \left(1 - \frac{3.7\varepsilon}{n^{10}}\right) < \pi_n < \pi \left(1 + \frac{3.7\varepsilon}{n^{10}}\right)$, which proves the lemma. \square

Lemma 5. *For $n \geq 1$ there is d_n with $0.321 < d_n < 0.671$ and*

$$\left| \sum_{k=n}^{\infty} (-1)^k s_k \right| = \frac{s_n}{1+\varepsilon} \cdot \exp\left(\frac{a_1}{n} + \frac{a_2 - \frac{1}{2}a_1^2}{n^2} + \frac{d_n\varepsilon}{n^3}\right),$$

where a_1 and a_2 are the coefficients from Theorem 1.

Proof. Theorem 1 implies that $\left|\sum_{k=n}^{\infty} (-1)^k s_k\right| = \frac{s_n}{1+\varepsilon} \cdot (1+x_n)$ with $x_n = \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3}$. Since $x_n > 0$ for all $n \geq 1$, we can use $x - \frac{1}{2}x^2 < \ln(1+x) < x - \frac{1}{2}x^2 + \frac{1}{3}x^3$.

Here, $e_n < 0.6704$ yields the upper bound on d_n :

$$\begin{aligned} \ln(1+x_n) &< \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3} - \frac{1}{2} \left(\frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3}\right)^2 + \frac{1}{3} \left(\frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3}\right)^3 \\ &< \frac{a_1}{n} + \frac{a_2 - \frac{1}{2}a_1^2}{n^2} + \frac{0.671\varepsilon}{n^3}. \end{aligned}$$

In the last step, we used that $|a_1|, |a_2|$ and ε are smaller than 10^{-14} .

For the lower bound on d_n , we use $e_n > 0.3216$:

$$\begin{aligned} \ln(1+x_n) &> \frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3} - \frac{1}{2} \left(\frac{a_1}{n} + \frac{a_2}{n^2} + \frac{e_n\varepsilon}{n^3}\right)^2 \\ &> \frac{a_1}{n} + \frac{a_2 - \frac{1}{2}a_1^2}{n^2} + \frac{0.321\varepsilon}{n^3}. \end{aligned}$$

This proves $0.321 < d_n < 0.671$, which finishes the proof of Lemma 5. \square

Proof of Theorem 2: We use Lemma 3, Lemma 4 and Lemma 5 in equation (13) and obtain

$$\begin{aligned} |\pi_n - \pi| &= \frac{12 \cdot 545\,140\,134 \pi^2}{\sqrt{640\,320^3}(1+\varepsilon)} \cdot \frac{\pi_n}{\pi} \cdot s_n \cdot \frac{1+\varepsilon}{s_n} \left| \sum_{k=n}^{\infty} (-1)^k s_k \right| \\ &= \frac{12 \cdot 545\,140\,134 \pi^2}{\sqrt{640\,320^3}(1+\varepsilon)} \cdot \frac{\varepsilon^n}{2\pi^{3/2}\sqrt{n}} \cdot \exp(R_n). \end{aligned}$$

The terms in the exponent R_n of this expression are

$$R_n = \frac{a_1}{n} + \frac{a_2 - \frac{1}{2}a_1^2}{n^2} + \frac{d_n\varepsilon}{n^3} + \frac{S - \frac{19}{72}}{n} + \frac{-\frac{1}{2}S^2}{n^2} + \frac{\frac{131}{15552} + \frac{1}{3}S^3}{n^3} + \frac{h_n}{n^4} + \frac{f_n\varepsilon}{n^{10}}.$$

Here we sort the terms as stated in Theorem 2:

$$\begin{aligned} A_0 &= \frac{6\sqrt{\pi} \cdot 545\,140\,134}{\sqrt{640\,320^3}(1+\varepsilon)} = \frac{106\,720 \cdot \sqrt{10\,005\pi}}{1\,672\,209}, \\ A_1 &= a_1 + S - \frac{19}{72} = -\frac{1\,781\,843\,197\,433}{7\,456\,754\,505\,816}, \\ A_2 &= a_2 - \frac{1}{2}a_1^2 - \frac{1}{2}S^2 = -\frac{1\,080\,096\,011\,925\,710\,088\,395}{3\,475\,199\,235\,000\,451\,148\,614\,116}, \end{aligned}$$

and the error term

$$\frac{\delta_n}{n^3} = \frac{d_n\varepsilon}{n^3} + \frac{\frac{131}{15552} + \frac{1}{3}S^3}{n^3} + \frac{h_n}{n^4} + \frac{f_n\varepsilon}{n^{10}}.$$

This proves $0.007 < \delta_n < 0.008\,429$ for $n \geq 2$. Using $\delta_1 = \ln\left(|\pi_1 - \pi| \cdot \frac{53360^3}{A_0}\right) - A_1 - A_2$ and $\pi_1 = \frac{\sqrt{640\,320^3}}{12 \cdot 13\,591\,409}$ yields $0.006\,907 < \delta_1 < 0.007$. \square

4. NUMERICAL VERIFICATION AND FURTHER RESULTS

For a numerical analysis, we computed the partial sums $\sum_{k=0}^{n-1} (-1)^k s_k$. Using these, we calculated the numerical values of e_n from Theorem 1 and of δ_n from Theorem 2. Figure 2 shows the results and that the bounds we have proven are close to optimal.




FIGURE 2. Numerically computed values of e_n (left) and δ_n (right) along with the proven bounds (dashed lines).

In [3], H. Cohen and J. Guillera list all 36 known rational hypergeometric series for $1/\pi$. They can be written with Pochhammer symbols $(a)_n = \Gamma(a+n)/\Gamma(n)$ and $R \in \{1/2; 1/3; 1/4; 1/6\}$:

$$(14) \quad \frac{1}{\pi} = P \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} (n+S) Z^n.$$

To generalize our results to formulae of this type, we can use the asymptotic estimate $\frac{(R)_n}{n!} \approx \frac{n^{R-1}}{\Gamma(R)}$ and Euler's reflection formula $\Gamma(R)\Gamma(1-R) = \pi/\sin(\pi R)$ to obtain

$$(15) \quad \frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} \approx \frac{\sin(\pi R)}{(\pi n)^{3/2}}.$$

In Appendix A, we can use the method presented by C. Mortici in [7] to prove a generalized version of Lemma 3: For $n \geq 1$ and $0 < R < 1$ we have that

$$(16) \quad \exp\left(\frac{\sigma_1}{n} + \frac{\sigma_3}{n^3} + \frac{\sigma_5}{n^5}\right) < \frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} \cdot \frac{(\pi n)^{3/2}}{\sin(\pi R)} < \exp\left(\frac{\sigma_1}{n} + \frac{\sigma_3}{n^3}\right),$$

with $\sigma_1 = -\frac{1}{8} - r$ and $\sigma_3 = \frac{1}{192} + \frac{1}{6}r^2$ and $\sigma_5 = \frac{-1}{640} - \frac{1}{30}r^2 - \frac{1}{15}r^3$ and $r = R(1-R)$. Here, $R = 1/6$ yields $\sigma_1 = -19/72$, $\sigma_3 = 131/15552$ and $\sigma_5 = -3337/1399680$.

Truncating one of the 36 rational hypergeometric series for $1/\pi$ of the form (14) results in approximations $\hat{\pi}_n$ of π . Using (16) instead of Lemma 3 gives the following estimate of the truncation error:

$$(17) \quad |\hat{\pi}_n - \pi| \approx \frac{P \sin(\pi R) \sqrt{\pi}}{(1-Z)\sqrt{n}} \cdot |Z|^n.$$

For asymptotic expansions of these 36 series see Appendix B.

We conclude our paper with another example – Ramanujan's series (1a) from the introduction: For this series, we obtain the asymptotic expansion

$$(18) \quad \sum_{k=n}^{\infty} \tilde{s}_k = \frac{\tilde{s}_n}{1-99^{-4}} \cdot \left(1 - \frac{1}{192119200n} + \frac{248104223}{3690978700864000n^2} - \frac{\tilde{e}_n}{n^3}\right),$$

with $3.558 \cdot 10^{-9} < \tilde{e}_n < 7.463 \cdot 10^{-9}$. Thus Ramanujan’s famous series (1a) results in approximations $\tilde{\pi}_n$ of π with an asymptotic expansion

$$(19) \quad |\tilde{\pi}_n - \pi| = 99^{-4n} \frac{9801\sqrt{\pi}}{1820\sqrt{n}} \exp\left(-\frac{1793359}{6624800n} - \frac{15333610991}{17555190016000n^2} + \frac{\tilde{\delta}_n}{n^3}\right),$$

where $0.0090 < \tilde{\delta}_n < 0.0111$. In [6, Theorem 3], C. Mortici has proven similar bounds for another of Ramanujan’s series.

Finally, we observe that the exponent $A_1/n + A_2/n^2 + \delta_n/n^3$ in Theorem 2 is negative, which implies the weaker estimate for the truncation error in the Chudnovskys’ series (1b)

$$(20) \quad |\pi_n - \pi| < \frac{106720\sqrt{10005}\pi}{1672209\sqrt{n}} \cdot 53360^{-3n} < \frac{11.315}{\sqrt{n}} \cdot 53360^{-3n},$$

which is valid for all $n \geq 1$. This much weaker estimate is sufficient to determine the number of terms needed to compute a given number of digits of π – even when computing trillions of digits. But the methods described in this paper demonstrate how to establish precise error bounds for the approximation of π through Ramanujan-like series for $1/\pi$ – and they can be applied to various other generalized hypergeometric series.

REFERENCES

- [1] Berndt, B., Chan, H.: Eisenstein series and approximations to π . *Illinois J. Math.* **45**, 75-90 (2001). <https://doi.org/10.1215/ijm/1258138255>.
- [2] Chudnovsky, D., Chudnovsky, G.: Approximation and complex multiplication according to Ramanujan. In: Andrews, G., Askey, R., Berndt, B., Ramanathan, K., Rankin, R. (eds.) *Ramanujan Revisited*, pp. 375-472. Academic Press, Boston (1988)
- [3] Cohen, H., Guillera, J.: Rational Hypergeometric Ramanujan Identities for $1/\pi^c$: Survey and Generalizations (2021). arXiv:2101.12592.
- [4] Milla, L.: A detailed proof of the Chudnovsky formula with means of basic complex analysis (2019). arXiv:1809.00533.
- [5] Milla, L.: An efficient determination of the coefficients in the Chudnovskys’ series for $1/\pi$. *Ramanujan J.* **57**, 803-809 (2022). <https://doi.org/10.1007/s11139-020-00330-6>.
- [6] Mortici, C.: Error estimates of Ramanujan-type series. *Ramanujan J.* **27**, 169-179 (2012). <https://doi.org/10.1007/s11139-011-9300-7>.
- [7] Mortici, C.: Product Approximations via Asymptotic Integration. *Amer. Math. Monthly* **117**, 434-441 (2010). <https://doi.org/10.4169/000298910X485950>.
- [8] Ramanujan, S.: Modular equations and approximations to π . *Q. J. Math.* **45**, 350-372 (1914).
- [9] Sasvari, Z.: An elementary proof of Binet’s formula for the gamma function. *Amer. Math. Monthly* **106**(2), 156-158 (1999). <https://doi.org/10.2307/2589052>.

APPENDIX A. ASYMPTOTIC EXPANSION OF A POCHHAMMER PRODUCT

In this appendix, we prove (16) using the method presented by C. Mortici in [7].

Lemma 6. *For all R with $0 < R < 1$ it holds*

$$\frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} = \frac{\sin(\pi R)}{(\pi n)^{3/2}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right).$$

Proof. The definition $(a)_n = \Gamma(n+a)/\Gamma(a)$ yields

$$\frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} = \underbrace{\frac{\Gamma(n+\frac{1}{2})\Gamma(n+R)\Gamma(n+1-R)}{\Gamma(n+1)^3}}_{=A} \cdot \underbrace{\frac{1}{\Gamma(\frac{1}{2})\Gamma(R)\Gamma(1-R)}}_{=B}.$$

From Stirling's formula $\Gamma(n+1) = n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ we obtain

$$\frac{\Gamma(n+1+x)}{n^x \Gamma(n+1)} \approx \frac{\sqrt{2\pi(n+x)} \left(\frac{n+x}{e}\right)^{n+x}}{n^x \sqrt{2\pi n} \left(\frac{n}{e}\right)^n} = \left(1 + \frac{x}{n}\right)^n e^{-x} \left(1 + \frac{x}{n}\right)^{x+\frac{1}{2}} \approx 1$$

which proves $A \approx n^{-3/2}$. Using the reflection formula $\Gamma(R)\Gamma(1-R) = \frac{\pi}{\sin(\pi R)}$ of the Γ function yields $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ and $B = \frac{\sin(\pi R)}{\pi^{3/2}}$ which proves the Lemma. \square

Lemma 7. For all $n \geq 1$ and all $0 < R < 1$ it holds

$$\exp\left(\frac{\sigma_1}{n} + \frac{\sigma_3}{n^3} + \frac{\sigma_5}{n^5}\right) < \frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} \cdot \frac{(\pi n)^{3/2}}{\sin(\pi R)} < \exp\left(\frac{\sigma_1}{n} + \frac{\sigma_3}{n^3}\right)$$

with $\sigma_1 = -\frac{1}{8} - r$ and $\sigma_3 = \frac{1}{192} + \frac{1}{6}r^2$ and $\sigma_5 = \frac{-1}{640} - \frac{1}{30}r^2 - \frac{1}{15}r^3$ and $r = R(1-R)$.

Proof. First we prove the upper bound and define r_n as follows:

$$\begin{aligned} \frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} &= \frac{\sin(\pi R)}{(\pi n)^{3/2}} \exp\left(\frac{\sigma_1}{n} + \frac{\sigma_3}{n^3} + r_n\right) \\ \implies r_n &= \ln\left(\frac{\left(\frac{1}{2}\right)_n (R)_n (1-R)_n}{n!^3} \cdot \frac{(\pi n)^{3/2}}{\sin(\pi R)}\right) - \frac{\sigma_1}{n} - \frac{\sigma_3}{n^3}. \end{aligned}$$

Here, Lemma 6 proves that r_n approaches 0 for $n \rightarrow \infty$.

Next, we compute $r_n - r_{n-1}$ as a power series $\sum_{k=1}^{\infty} \varrho_k/n^k$:

$$\begin{aligned} r_n - r_{n-1} &= \ln\left(\frac{\left(n - \frac{1}{2}\right) (n+R-1) (n-R)}{n^3} \cdot \frac{n^{3/2}}{(n-1)^{3/2}}\right) \\ &\quad - \frac{\sigma_1}{n} - \frac{\sigma_3}{n^3} + \frac{\sigma_1}{n-1} + \frac{\sigma_3}{(n-1)^3} \\ &= \ln\left(\frac{\left(1 - \frac{1}{2n}\right) \left(1 - \frac{1-R}{n}\right) \left(1 - \frac{R}{n}\right)}{\left(1 - \frac{1}{n}\right)^{3/2}}\right) + \sum_{k=2}^{\infty} \frac{\sigma_1}{n^k} + \sum_{k=4}^{\infty} \frac{\binom{k-1}{2} \sigma_3}{n^k} \\ &= \sum_{k=1}^{\infty} \frac{\frac{3}{2} - \left(\frac{1}{2}\right)^k - (1-R)^k - R^k}{kn^k} + \sum_{k=2}^{\infty} \frac{\sigma_1}{n^k} + \sum_{k=4}^{\infty} \frac{\binom{k-1}{2} \sigma_3}{n^k} =: \sum_{k=1}^{\infty} \frac{\varrho_k}{n^k} \end{aligned}$$

Here we observe $\varrho_1 = 0$ and $\varrho_2 = \sigma_1 + \frac{1}{8} + R(1-R)$. This explains our choice of σ_1 and proves $\varrho_2 = \varrho_3 = 0$ and $\varrho_4 = -\frac{1}{64} - \frac{1}{2}r^2 + 3 \cdot \sigma_3$. By our choice of σ_3 , we have $\varrho_4 = \varrho_5 = 0$. This proves

$$r_n - r_{n-1} = \sum_{k=6}^{\infty} \frac{\varrho_k}{n^k} \quad \text{with} \quad \varrho_6 = \frac{1}{128} + \frac{1}{6}r^2 + \frac{1}{3}r^3.$$

In Lemma 8 we prove $\varrho_k > 0$ for all $k \geq 6$ and thus $r_n - r_{n-1} > 0$. This proves that r_n is *increasing* to zero, thus $r_n < 0$.

For the lower bound, we use $\tilde{r}_n = r_n - \sigma_5/n^5$:

$$\tilde{r}_n - \tilde{r}_{n-1} = \frac{\sigma_5}{(n-1)^5} - \frac{\sigma_5}{n^5} + \sum_{k=6}^{\infty} \frac{\varrho_k}{n^k} = \sum_{k=6}^{\infty} \frac{\varrho_k + \binom{k-1}{4} \sigma_5}{n^k} =: \sum_{k=6}^{\infty} \frac{\tilde{\varrho}_k}{n^k}$$

Here, our choice of $\sigma_5 = -\varrho_6/5$ yields $\tilde{\varrho}_6 = \tilde{\varrho}_7 = 0$ and

$$\tilde{r}_n - \tilde{r}_{n-1} = \sum_{k=8}^{\infty} \frac{\tilde{\varrho}_k}{n^k} \quad \text{with} \quad \tilde{\varrho}_8 = -\frac{17}{2048} - \frac{1}{6}r^2 - \frac{1}{3}r^3 - \frac{1}{4}r^4.$$

In Lemma 9 we prove $\tilde{\varrho}_k < 0$ for all $k \geq 8$ and thus $r_n - r_{n-1} < 0$. This proves that \tilde{r}_n is *decreasing* to zero, which proves $\tilde{r}_n > 0$. \square

Lemma 8. For $k \geq 6$ and for all $0 \leq R \leq 1$ it holds $\varrho_k > 0$ with

$$\varrho_k = \frac{\frac{1}{2} - \left(\frac{1}{2}\right)^k}{k} + \frac{R - R^k}{k} + \frac{(1 - R) - (1 - R)^k}{k} + \sigma_1 + \binom{k-1}{2}\sigma_3$$

and $\sigma_1 = -\frac{1}{8} - r$ and $\sigma_3 = \frac{1}{192} + \frac{1}{6}r^2$ and $r = R(1 - R)$.

Proof. The first three terms are positive. For $k \geq 10$ it holds $\binom{k-1}{2} \geq 36$ and

$$\sigma_1 + \binom{k-1}{2}\sigma_3 \geq -\frac{1}{8} - r + 36\left(\frac{1}{192} + \frac{1}{6}r^2\right) = \frac{1}{16} + 6r^2 - r > 0.$$

It remains to check $\varrho_k > 0$ for $k < 10$: $\varrho_6 = \frac{1}{128} + \frac{1}{6}r^2 + \frac{1}{3}r^3$, $\varrho_7 = \frac{3}{128} + \frac{1}{2}r^2 + r^3$, $\varrho_8 = \frac{95}{2048} + r^2 + 2r^3 - \frac{1}{4}r^4$ and $\varrho_9 = \frac{39}{512} + \frac{5}{3}r^2 + \frac{10}{3}r^3 - r^4$ are also positive. \square

Lemma 9. For $k \geq 8$ and for all $0 \leq R \leq 1$ it holds $\tilde{\varrho}_k < 0$ with

$$\tilde{\varrho}_k = \frac{\frac{1}{2} - \left(\frac{1}{2}\right)^k}{k} + \frac{R - R^k}{k} + \frac{(1 - R) - (1 - R)^k}{k} + \sigma_1 + \binom{k-1}{2}\sigma_3 + \binom{k-1}{4}\sigma_5$$

and $\sigma_1 = -\frac{1}{8} - r$ and $\sigma_3 = \frac{1}{192} + \frac{1}{6}r^2$ and $\sigma_5 = \frac{-1}{640} - \frac{1}{30}r^2 - \frac{1}{15}r^3$ and $r = R(1 - R)$.

Proof. Since $\binom{k-1}{4} = \frac{(k-3)(k-4)}{12}\binom{k-1}{2}$, for $k \geq 12$ we have $\binom{k-1}{4} \geq 6\binom{k-1}{2}$ and

$$\binom{k-1}{2}\sigma_3 + \binom{k-1}{4}\sigma_5 \leq \binom{k-1}{2}(\sigma_3 + 6\sigma_5) \leq 55\left(-\frac{1}{240} - \frac{1}{30}r^2 - \frac{2}{5}r^3\right) \leq -\frac{11}{48}$$

The first three terms in $\tilde{\varrho}_k$ are less than $\frac{3/2}{k}$. This proves $\tilde{\varrho}_k \leq \frac{3/2}{12} + \sigma_1 - \frac{11}{48} < 0$ (if $k \geq 12$).

It remains to check $\tilde{\varrho}_k < 0$ for $k < 12$:

$$\begin{aligned} \tilde{\varrho}_8 &= \varrho_8 + 35\sigma_5 = -\frac{17}{2048} - \frac{1}{6}r^2 - \frac{1}{3}r^3 - \frac{1}{4}r^4 \\ \tilde{\varrho}_9 &= \varrho_9 + 70\sigma_5 = -\frac{17}{512} - \frac{2}{3}r^2 - \frac{4}{3}r^3 - r^4 \\ \tilde{\varrho}_{10} &= \frac{2r^5 - 25r^4 + 50r^3 - 35r^2 + 10r + \frac{511}{1024}}{10} + \sigma_1 + 36\sigma_3 + 126\sigma_5 \\ &= \frac{-173}{2048} - \frac{17}{10}r^2 - \frac{17}{5}r^3 - \frac{5}{2}r^4 + \frac{1}{5}r^5 \\ \tilde{\varrho}_{11} &= \frac{11r^5 - 55r^4 + 77r^3 - 44r^2 + 11r + \frac{1023}{2048}}{11} + \sigma_1 + 45\sigma_3 + 210\sigma_5 \\ &= \frac{-355}{2048} - \frac{7}{2}r^2 - 7r^3 - 5r^4 + r^5 \end{aligned}$$

Since we have $0 \leq R \leq 1$, we have $0 \leq r \leq 1$, which proves $\tilde{\varrho}_k < 0$ for all k . \square

APPENDIX B. ASYMPTOTIC EXPANSIONS FOR FURTHER SERIES

We have used Lemma 4 to estimate $\pi_n \approx \pi$ in

$$\pi_n - \pi = \pi \pi_n \left(\frac{1}{\pi} - \frac{1}{\pi_n}\right) \approx \pi^2 \left(\frac{1}{\pi} - \frac{1}{\pi_n}\right).$$

This has worked well, because the Chudnovskys' Series 7 and Ramanujan's Series 23 converge so fast. For most other series, this estimate is valid only for larger n . For example in Series 1, we need $n \geq 25$ (see Fig. 3).

In Series 33, the convergence is too slow and we can not use this estimate at all. This is why we only give an expansion for $\frac{1}{\pi} - \frac{1}{\pi_{33}(n)}$ at Series 33.

#	R	S	Z	P
1	1/6	8/63	-64/125	21√15/25
2	1/6	15/154	-27/512	77√2/32
3	1/6	25/342	-1/512	57√6/32
4	1/6	31/506	-9/64000	759√30/800
5	1/6	263/5418	-1/512000	2709√15/1600
6	1/6	10177/261702	-1/85184000	43617√330/96800
7	1/6	13591409/545140134	-1/53360 ³	90856689√10005/711822400
8	1/6	3/28	27/125	28√5/25
9	1/6	1/11	4/125	22√15/25
10	1/6	5/63	8/1331	84√33/121
11	1/6	8/133	64/614125	2394√255/7225
12	1/4	3/20	-1/4	5/2
13	1/4	23/260	-1/324	65/18
14	1/4	1123/21460	-1/777924	5365/882
15	1/4	8/65	-256/3969	65√7/63
16	1/4	3/28	-1/48	7√3/4
17	1/4	41/644	-1/25920	161√5/72
18	1/4	1/7	32/81	14/9
19	1/4	1/8	1/9	4√3/3
20	1/4	1/10	1/81	20√2/9
21	1/4	3/40	1/2401	120√3/49
22	1/4	19/280	1/9801	140√11/99
23	1/4	1103/26390	1/96059601	52780√2/9801
24	1/3	7/51	-1/16	17√3/12
25	1/3	53/615	-1/1024	205√3/96
26	1/3	827/14151	-1/250000	4717√3/1500
27	1/3	1/5	-9/16	5√3/4
28	1/3	1/9	-1/80	3√15/4
29	1/3	13/165	-1/3024	55√7/36
30	1/3	1/6	1/2	2√3/3
31	1/3	2/15	2/27	20/9
32	1/3	4/33	4/125	22√3/15
33	1/2	1/4	-1	2
34	1/2	1/6	-1/8	3√2/2
35	1/2	1/6	1/4	3/2
36	1/2	5/42	1/64	21/8

TABLE 1. Coefficients (R, S, Z, P) in formulae of type (14).

Series 1. If we denote by $1/\pi_1(n)$ the truncated series

$$\frac{1}{\pi_1(n)} = \frac{21\sqrt{15}}{25} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{8}{63}\right) \left(\frac{-64}{125}\right)^k,$$

then we have that $\pi_1(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_1(n) - \pi = \left(\frac{-64}{125}\right)^n \frac{5\sqrt{15}\pi}{18\sqrt{n}} \cdot \exp\left(\frac{7}{216n} + \frac{-80}{729n^2} + \frac{23129}{3779136n^3} + \frac{47168}{531441n^4} \Theta_1(n)\right).$$

For all $n \geq 25$ we have that $0 < \Theta_1(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_1(n) \rightarrow 1$.




FIGURE 3. Values of $\Theta_1(n)$ from Series 1 showing strong oscillation for $n < 25$ (left) and damped oscillation for $n \geq 25$ (right).

Series 2. If we denote by $1/\pi_2(n)$ the truncated series

$$\frac{1}{\pi_2(n)} = \frac{77\sqrt{2}}{32} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{15}{154}\right) \left(\frac{-27}{512}\right)^k,$$

then we have that $\pi_2(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_2(n) - \pi &= \left(\frac{-27}{512}\right)^n \frac{8\sqrt{2\pi}}{7\sqrt{n}} \\ &\cdot \exp\left(\frac{-499}{3528n} + \frac{-291}{9604n^2} + \frac{57332435}{1829677248n^3} + \frac{-2919207}{184473632n^4} \Theta_2(n)\right). \end{aligned}$$

For all $n \geq 4$ we have that $0 < \Theta_2(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_2(n) \rightarrow 1$.

Series 3. If we denote by $1/\pi_3(n)$ the truncated series

$$\frac{1}{\pi_3(n)} = \frac{57\sqrt{6}}{32} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{25}{342}\right) \left(\frac{-1}{512}\right)^k,$$

then we have that $\pi_3(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_3(n) - \pi = \left(\frac{-1}{512}\right)^n \frac{8\sqrt{6\pi}}{9\sqrt{n}} \cdot \exp\left(\frac{-41}{216n} + \frac{-11}{2916n^2} + \frac{36665}{3779136n^3} \Theta_3(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_3(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_3(n) \rightarrow 1$.

Series 4. If we denote by $1/\pi_4(n)$ the truncated series

$$\frac{1}{\pi_4(n)} = \frac{759\sqrt{30}}{800} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{31}{506}\right) \left(\frac{-9}{64000}\right)^k,$$

then we have that $\pi_4(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_4(n) - \pi &= \left(\frac{-9}{64000}\right)^n \frac{120\sqrt{30\pi}}{253\sqrt{n}} \\ &\cdot \exp\left(\frac{-933499}{4608648n} + \frac{-32087155}{16388608324n^2} + \frac{35019155487209387}{4078583656660041408n^3} \Theta_4(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_4(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_4(n) \rightarrow 1$.

Series 5. If we denote by $1/\pi_5(n)$ the truncated series

$$\frac{1}{\pi_5(n)} = \frac{2709\sqrt{15}}{1600} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{263}{5418}\right) \left(\frac{-1}{512000}\right)^k,$$

then we have that $\pi_5(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_5(n) - \pi &= \left(\frac{-1}{512000}\right)^n \frac{160\sqrt{15}\pi}{189\sqrt{n}} \\ &\cdot \exp\left(\frac{-20513}{95256n} + \frac{-668795}{567106596n^2} + \frac{2742951440609}{324121835451456n^3} \Theta_5(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_5(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_5(n) \rightarrow 1$.

Series 6. If we denote by $1/\pi_6(n)$ the truncated series

$$\frac{1}{\pi_6(n)} = \frac{43617\sqrt{330}}{96800} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{10177}{261702}\right) \left(\frac{-1}{440^3}\right)^k,$$

then we have that $\pi_6(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_6(n) - \pi &= \left(\frac{-1}{440^3}\right)^n \frac{440\sqrt{330}\pi}{1953\sqrt{n}} \cdot \exp\left(\frac{-2288537}{10171224n} + \frac{-4889066795}{6465862353636n^2}\right. \\ &\quad \left. + \frac{3331546471820293001}{394594406111993822784n^3} \Theta_6(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_6(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_6(n) \rightarrow 1$.

Series 7. If we denote by $1/\pi_7(n)$ the truncated series

$$\frac{1}{\pi_7(n)} = \frac{90856689\sqrt{10005}}{711822400} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{13591409}{545140134}\right) \left(\frac{-1}{53360^3}\right)^k,$$

then we have that $\pi_7(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_7(n) - \pi &= \left(\frac{-1}{53360^3}\right)^n \frac{106720\sqrt{10005}\pi}{1672209\sqrt{n}} \\ &\cdot \exp\left(\frac{-1781843197433}{7456754505816n} + \frac{-1080096011925710088395}{3475199235000451148614116n^2}\right. \\ &\quad \left. + \frac{1310485187935583963485460802564780329}{155482245325187582131326612761170191936n^3} \Theta_7(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_7(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_7(n) \rightarrow 1$.

Series 8. If we denote by $1/\pi_8(n)$ the truncated series

$$\frac{1}{\pi_8(n)} = \frac{28\sqrt{5}}{25} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{3}{28}\right) \left(\frac{27}{125}\right)^k,$$

then we have that $\pi_8(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_8(n) - \pi &= \left(\frac{27}{125}\right)^n \frac{5\sqrt{5}\pi}{7\sqrt{n}} \cdot \exp\left(\frac{-1039}{3528n} + \frac{3615}{19208n^2} + \frac{-601881853}{1829677248n^3}\right. \\ &\quad \left. + \frac{288589047}{368947264n^4} + \frac{-132250237090171}{56482136645760n^5} \Theta_8(n)\right). \end{aligned}$$

For all $n \geq 4$ we have that $0 < \Theta_8(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_8(n) \rightarrow 1$.

Series 9. If we denote by $1/\pi_9(n)$ the truncated series

$$\frac{1}{\pi_9(n)} = \frac{22\sqrt{15}}{25} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{1}{11}\right) \left(\frac{4}{125}\right)^k,$$

then we have that $\pi_9(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_9(n) - \pi = & \left(\frac{4}{125}\right)^n \frac{5\sqrt{15}\pi}{11\sqrt{n}} \cdot \exp\left(\frac{-1651}{8712n} + \frac{3865}{263538n^2} + \frac{-339804517}{27551316672n^3}\right. \\ & \left. + \frac{1739500937}{69452277444n^4} + \frac{-119121688877123}{3300372224138880n^5} \Theta_9(n)\right). \end{aligned}$$

For all $n \geq 2$ we have that $0 < \Theta_9(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_9(n) \rightarrow 1$.

Series 10. If we denote by $1/\pi_{10}(n)$ the truncated series

$$\frac{1}{\pi_{10}(n)} = \frac{84\sqrt{33}}{121} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{5}{63}\right) \left(\frac{8}{1331}\right)^k,$$

then we have that $\pi_{10}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{10}(n) - \pi = & \left(\frac{8}{1331}\right)^n \frac{22\sqrt{33}\pi}{63\sqrt{n}} \cdot \exp\left(\frac{-1985}{10584n} + \frac{899}{3500658n^2} + \frac{2216760425}{444611571264n^3}\right. \\ & \left. + \frac{46674125279}{12254606432964n^4} + \frac{-483110468975635}{74115859706566272n^5} \Theta_{10}(n)\right). \end{aligned}$$

For all $n \geq 2$ we have that $0 < \Theta_{10}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{10}(n) \rightarrow 1$.

Series 11. If we denote by $1/\pi_{11}(n)$ the truncated series

$$\frac{1}{\pi_{11}(n)} = \frac{2394\sqrt{255}}{7225} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(k!)^3} \left(k + \frac{8}{133}\right) \left(\frac{64}{85^3}\right)^k,$$

then we have that $\pi_{11}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{11}(n) - \pi = & \left(\frac{64}{85^3}\right)^n \frac{85\sqrt{255}\pi}{513\sqrt{n}} \cdot \exp\left(\frac{-143017}{701784n} + \frac{-13456880}{7695324729n^2}\right. \\ & \left. + \frac{1092857149734121}{129610938470796864n^3} + \frac{3781366699871168}{59218022684758923441n^4}\right. \\ & \left. + \frac{-8157754659852064155749017}{3324660866544068506249499520n^5} \Theta_{11}(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_{11}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{11}(n) \rightarrow 1$.

Series 12. If we denote by $1/\pi_{12}(n)$ the truncated series

$$\frac{1}{\pi_{12}(n)} = \frac{5}{2} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{3}{20}\right) \left(\frac{-1}{4}\right)^k,$$

then we have that $\pi_{12}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{12}(n) - \pi = & \left(\frac{-1}{4}\right)^n \frac{\sqrt{2\pi}}{\sqrt{n}} \\ & \cdot \exp\left(\frac{-1}{16n} + \frac{-3}{32n^2} + \frac{73}{1536n^3} + \frac{33}{1024n^4} + \frac{-1861}{20480n^5} \Theta_{12}(n)\right). \end{aligned}$$

For all $n \geq 12$ we have that $0 < \Theta_{12}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{12}(n) \rightarrow 1$.

Series 13. If we denote by $1/\pi_{13}(n)$ the truncated series

$$\frac{1}{\pi_{13}(n)} = \frac{65}{18} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{23}{260}\right) \left(\frac{-1}{324}\right)^k,$$

then we have that $\pi_{13}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{13}(n) - \pi = \left(\frac{-1}{324}\right)^n \frac{9\sqrt{2\pi}}{5\sqrt{n}} \cdot \exp\left(\frac{-89}{400n} + \frac{-23}{4000n^2} + \frac{317969}{24000000n^3} \Theta_{13}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{13}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{13}(n) \rightarrow 1$.

Series 14. If we denote by $1/\pi_{14}(n)$ the truncated series

$$\frac{1}{\pi_{14}(n)} = \frac{5365}{882} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{1123}{21460}\right) \left(\frac{-1}{777924}\right)^k,$$

then we have that $\pi_{14}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{14}(n) - \pi &= \left(\frac{-1}{777924}\right)^n \frac{441\sqrt{2\pi}}{145\sqrt{n}} \\ &\cdot \exp\left(\frac{-87521}{336400n} + \frac{-155039}{113164960n^2} + \frac{158694702263081}{14275759704000000n^3} \Theta_{14}(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_{14}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{14}(n) \rightarrow 1$.

Series 15. If we denote by $1/\pi_{15}(n)$ the truncated series

$$\frac{1}{\pi_{15}(n)} = \frac{65\sqrt{7}}{63} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{8}{65}\right) \left(\frac{-256}{3969}\right)^k,$$

then we have that $\pi_{15}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{15}(n) - \pi &= \left(\frac{-256}{3969}\right)^n \frac{63\sqrt{14\pi}}{130\sqrt{n}} \cdot \exp\left(\frac{-10757}{67600n} + \frac{-140944}{3570125n^2} + \frac{4556280871313}{115843416000000n^3}\right. \\ &\quad \left.+ \frac{-5541878772448}{318644812890625n^4} + \frac{-123624675437230027457}{275716983698000000000n^5} \Theta_{15}(n)\right). \end{aligned}$$

For all $n \geq 5$ we have that $0 < \Theta_{15}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{15}(n) \rightarrow 1$.

Series 16. If we denote by $1/\pi_{16}(n)$ the truncated series

$$\frac{1}{\pi_{16}(n)} = \frac{7\sqrt{3}}{4} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{3}{28}\right) \left(\frac{-1}{48}\right)^k,$$

then we have that $\pi_{16}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{16}(n) - \pi &= \left(\frac{-1}{48}\right)^n \frac{6\sqrt{6\pi}}{7\sqrt{n}} \cdot \exp\left(\frac{-153}{784n} + \frac{-1331}{76832n^2} + \frac{1385667}{60236288n^3}\right. \\ &\quad \left.+ \frac{-61756703}{5903156224n^4} + \frac{3777837381}{826441871360n^5} \Theta_{16}(n)\right). \end{aligned}$$

For all $n \geq 3$ we have that $0 < \Theta_{16}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{16}(n) \rightarrow 1$.

Series 17. If we denote by $1/\pi_{17}(n)$ the truncated series

$$\frac{1}{\pi_{17}(n)} = \frac{161\sqrt{5}}{72} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{41}{644}\right) \left(\frac{-1}{25920}\right)^k,$$

then we have that $\pi_{17}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{17}(n) - \pi = \left(\frac{-1}{25920}\right)^n \frac{180\sqrt{10}\pi}{161\sqrt{n}} \cdot \exp\left(\frac{-103193}{414736n} + \frac{-44090635}{21500743712n^2} + \frac{299081123487449}{26751397332420096n^3}\Theta_{17}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{17}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{17}(n) \rightarrow 1$.

Series 18. If we denote by $1/\pi_{18}(n)$ the truncated series

$$\frac{1}{\pi_{18}(n)} = \frac{14}{9} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{1}{7}\right) \left(\frac{32}{81}\right)^k,$$

then we have that $\pi_{18}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{18}(n) - \pi = \left(\frac{32}{81}\right)^n \frac{9\sqrt{2}\pi}{7\sqrt{n}} \cdot \exp\left(\frac{-389}{784n} + \frac{2939}{4802n^2} + \frac{-302662495}{180708864n^3} + \frac{148922927}{23059204n^4} + \frac{-26488619802887}{826441871360n^5}\Theta_{18}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{18}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{18}(n) \rightarrow 1$.

Series 19. If we denote by $1/\pi_{19}(n)$ the truncated series

$$\frac{1}{\pi_{19}(n)} = \frac{4\sqrt{3}}{3} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{1}{8}\right) \left(\frac{1}{9}\right)^k,$$

then we have that $\pi_{19}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{19}(n) - \pi = \left(\frac{1}{9}\right)^n \frac{3\sqrt{6}\pi}{4\sqrt{n}} \cdot \exp\left(\frac{-1}{4n} + \frac{37}{512n^2} + \frac{-1195}{12288n^3} + \frac{46945}{262144n^4} + \frac{-1956919}{5242880n^5}\Theta_{19}(n)\right).$$

For all $n \geq 3$ we have that $0 < \Theta_{19}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{19}(n) \rightarrow 1$.

Series 20. If we denote by $1/\pi_{20}(n)$ the truncated series

$$\frac{1}{\pi_{20}(n)} = \frac{20\sqrt{2}}{9} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{1}{10}\right) \left(\frac{1}{81}\right)^k,$$

then we have that $\pi_{20}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{20}(n) - \pi = \left(\frac{1}{81}\right)^n \frac{9\sqrt{\pi}}{4\sqrt{n}} \cdot \exp\left(\frac{-7}{32n} + \frac{5}{2048n^2} + \frac{323}{98304n^3} + \frac{37313}{4194304n^4} + \frac{-2253817}{167772160n^5}\Theta_{20}(n)\right).$$

For all $n \geq 2$ we have that $0 < \Theta_{20}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{20}(n) \rightarrow 1$.

Series 21. If we denote by $1/\pi_{21}(n)$ the truncated series

$$\frac{1}{\pi_{21}(n)} = \frac{120\sqrt{3}}{49} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{3}{40}\right) \left(\frac{1}{2401}\right)^k,$$

then we have that $\pi_{21}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{21}(n) - \pi = \left(\frac{1}{2401}\right)^n \frac{49\sqrt{6\pi}}{40\sqrt{n}} \cdot \exp\left(\frac{-1141}{4800n} + \frac{-23567}{9216000n^2} + \frac{3626854163}{331776000000n^3} + \frac{606576592193}{2123366400000000n^4} + \frac{-44351997882926281}{1274019840000000000n^5}\Theta_{21}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{21}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{21}(n) \rightarrow 1$.

Series 22. If we denote by $1/\pi_{22}(n)$ the truncated series

$$\frac{1}{\pi_{22}(n)} = \frac{140\sqrt{11}}{99} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{19}{280}\right) \left(\frac{1}{9801}\right)^k,$$

then we have that $\pi_{22}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{22}(n) - \pi = \left(\frac{1}{9801}\right)^n \frac{99\sqrt{22\pi}}{140\sqrt{n}} \cdot \exp\left(\frac{-1199}{4900n} + \frac{-344063}{153664000n^2} + \frac{250800696613}{22588608000000n^3} + \frac{40239008483873}{590315622400000000n^4} + \frac{-959530669459838519}{29515781120000000000n^5}\Theta_{22}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{22}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{22}(n) \rightarrow 1$.

Series 23. If we denote by $1/\pi_{23}(n)$ the truncated series

$$\frac{1}{\pi_{23}(n)} = \frac{52780\sqrt{2}}{9801} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k}{(k!)^3} \left(k + \frac{1103}{26390}\right) \left(\frac{1}{99^4}\right)^k,$$

then we have that $\pi_{23}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{23}(n) - \pi = \left(\frac{1}{99^4}\right)^n \frac{9801\sqrt{\pi}}{1820\sqrt{n}} \cdot \exp\left(\frac{-1793359}{6624800n} + \frac{-15333610991}{17555190016000n^2} + \frac{9674999636298154667}{872247171134976000000n^3} + \frac{-5815449554312741373727}{7704617412446652006400000000n^4} + \frac{-28927901748784728363322170328567}{911456239892438932357120000000000n^5}\Theta_{23}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{23}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{23}(n) \rightarrow 1$.

Series 24. If we denote by $1/\pi_{24}(n)$ the truncated series

$$\frac{1}{\pi_{24}(n)} = \frac{17\sqrt{3}}{12} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{7}{51}\right) \left(\frac{-1}{16}\right)^k,$$

then we have that $\pi_{24}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{24}(n) - \pi = \left(\frac{-1}{16}\right)^n \frac{2\sqrt{\pi}}{\sqrt{n}} \cdot \exp\left(\frac{-13}{72n} + \frac{-1}{24n^2} + \frac{665}{15552n^3} + \frac{-95}{5184n^4} + \frac{-6559}{1399680n^5}\Theta_{24}(n)\right).$$

For all $n \geq 5$ we have that $0 < \Theta_{24}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{24}(n) \rightarrow 1$.

Series 25. If we denote by $1/\pi_{25}(n)$ the truncated series

$$\frac{1}{\pi_{25}(n)} = \frac{205\sqrt{3}}{96} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{53}{615}\right) \left(\frac{-1}{1024}\right)^k,$$

then we have that $\pi_{25}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{25}(n) - \pi = \left(\frac{-1}{1024}\right)^n \frac{16\sqrt{\pi}}{5\sqrt{n}} \cdot \exp\left(\frac{-469}{1800n} + \frac{-13}{3000n^2} + \frac{3482081}{243000000n^3} \Theta_{25}(n)\right).$$

For all $n \geq 1$ we have that $0 < \Theta_{25}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{25}(n) \rightarrow 1$.

Series 26. If we denote by $1/\pi_{26}(n)$ the truncated series

$$\frac{1}{\pi_{26}(n)} = \frac{4717\sqrt{3}}{1500} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{827}{14151}\right) \left(\frac{-1}{250000}\right)^k,$$

then we have that $\pi_{26}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{26}(n) - \pi &= \left(\frac{-1}{250000}\right)^n \frac{250\sqrt{\pi}}{53\sqrt{n}} \\ &\cdot \exp\left(\frac{-58405}{202248n} + \frac{-107963}{63123848n^2} + \frac{4656308646401}{344700144278208n^3} \Theta_{26}(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_{26}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{26}(n) \rightarrow 1$.

Series 27. If we denote by $1/\pi_{27}(n)$ the truncated series

$$\frac{1}{\pi_{27}(n)} = \frac{5\sqrt{3}}{4} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{1}{5}\right) \left(\frac{-9}{16}\right)^k,$$

then we have that $\pi_{27}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{27}(n) - \pi &= \left(\frac{-9}{16}\right)^n \frac{6\sqrt{\pi}}{5\sqrt{n}} \\ &\cdot \exp\left(\frac{59}{1800n} + \frac{-127}{1000n^2} + \frac{-186271}{243000000n^3} + \frac{2740273}{25000000n^4} \Theta_{27}(n)\right). \end{aligned}$$

For all $n \geq 31$ we have that $0 < \Theta_{27}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{27}(n) \rightarrow 1$.

Series 28. If we denote by $1/\pi_{28}(n)$ the truncated series

$$\frac{1}{\pi_{28}(n)} = \frac{3\sqrt{15}}{4} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{1}{9}\right) \left(\frac{-1}{80}\right)^k,$$

then we have that $\pi_{28}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{28}(n) - \pi &= \left(\frac{-1}{80}\right)^n \frac{10\sqrt{5}\pi}{9\sqrt{n}} \cdot \exp\left(\frac{-149}{648n} + \frac{-715}{52488n^2} + \frac{2207033}{102036672n^3} \right. \\ &\quad \left. + \frac{-20699503}{2754990144n^4} + \frac{5625196531}{2231542016640n^5} \Theta_{28}(n)\right). \end{aligned}$$

For all $n \geq 3$ we have that $0 < \Theta_{28}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{28}(n) \rightarrow 1$.

Series 29. If we denote by $1/\pi_{29}(n)$ the truncated series

$$\frac{1}{\pi_{29}(n)} = \frac{55\sqrt{7}}{36} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{13}{165}\right) \left(\frac{-1}{3024}\right)^k,$$

then we have that $\pi_{29}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{29}(n) - \pi &= \left(\frac{-1}{3024}\right)^n \frac{42\sqrt{21}\pi}{55\sqrt{n}} \\ &\cdot \exp\left(\frac{-58429}{217800n} + \frac{-436999}{131769000n^2} + \frac{5956214110601}{430489323000000n^3} \Theta_{29}(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_{29}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{29}(n) \rightarrow 1$.

Series 30. If we denote by $1/\pi_{30}(n)$ the truncated series

$$\frac{1}{\pi_{30}(n)} = \frac{2\sqrt{3}}{3} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{1}{6}\right) \left(\frac{1}{2}\right)^k,$$

then we have that $\pi_{30}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{30}(n) - \pi &= \left(\frac{1}{2}\right)^n \frac{2\sqrt{\pi}}{\sqrt{n}} \\ &\cdot \exp\left(\frac{-49}{72n} + \frac{7}{6n^2} + \frac{-65647}{15552n^3} + \frac{7057}{324n^4} + \frac{-202977547}{1399680n^5} \Theta_{30}(n)\right). \end{aligned}$$

For all $n \geq 1$ we have that $0 < \Theta_{30}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{30}(n) \rightarrow 1$.

Series 31. If we denote by $1/\pi_{31}(n)$ the truncated series

$$\frac{1}{\pi_{31}(n)} = \frac{20}{9} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{2}{15}\right) \left(\frac{2}{27}\right)^k,$$

then we have that $\pi_{31}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{31}(n) - \pi &= \left(\frac{2}{27}\right)^n \frac{6\sqrt{3\pi}}{5\sqrt{n}} \cdot \exp\left(\frac{-457}{1800n} + \frac{19}{450n^2} + \frac{-12156583}{243000000n^3} + \frac{11619823}{126562500n^4} \right. \\ &\quad \left. + \frac{-2235858759091}{1366875000000n^5} \Theta_{31}(n)\right). \end{aligned}$$

For all $n \geq 3$ we have that $0 < \Theta_{31}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{31}(n) \rightarrow 1$.

Series 32. If we denote by $1/\pi_{32}(n)$ the truncated series

$$\frac{1}{\pi_{32}(n)} = \frac{22\sqrt{3}}{15} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(k!)^3} \left(k + \frac{4}{33}\right) \left(\frac{4}{125}\right)^k,$$

then we have that $\pi_{32}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\begin{aligned} \pi_{32}(n) - \pi &= \left(\frac{4}{125}\right)^n \frac{25\sqrt{\pi}}{11\sqrt{n}} \cdot \exp\left(\frac{-2113}{8712n} + \frac{580}{43923n^2} + \frac{-262753543}{27551316672n^3} \right. \\ &\quad \left. + \frac{491042788}{17363069361n^4} + \frac{-138715488606809}{3300372224138880n^5} \Theta_{32}(n)\right). \end{aligned}$$

For all $n \geq 3$ we have that $0 < \Theta_{32}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{32}(n) \rightarrow 1$.

Series 33. If we denote by $1/\pi_{33}(n)$ the truncated series

$$\frac{1}{\pi_{33}(n)} = 2 \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k}{(k!)^3} \left(k + \frac{1}{4}\right) (-1)^k,$$

then we have that $\pi_{33}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\frac{1}{\pi} - \frac{1}{\pi_{33}(n)} = (-1)^n \frac{1}{\sqrt{\pi^3 n}} \cdot \exp\left(\frac{1}{8n} + \frac{-1}{8n^2} + \frac{-13}{192n^3} + \frac{7}{64n^4} + \frac{81}{640n^5} \Theta_{33}(n)\right).$$

For all $n \geq 2$ we have that $0 < \Theta_{33}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{33}(n) \rightarrow 1$.

Series 34. If we denote by $1/\pi_{34}(n)$ the truncated series

$$\frac{1}{\pi_{34}(n)} = \frac{3\sqrt{2}}{2} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k}{(k!)^3} \left(k + \frac{1}{6}\right) \left(\frac{-1}{8}\right)^k,$$

then we have that $\pi_{34}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{34}(n) - \pi = \left(\frac{-1}{8}\right)^n \frac{4\sqrt{2\pi}}{3\sqrt{n}} \cdot \exp\left(\frac{-11}{72n} + \frac{-23}{324n^2} + \frac{8171}{139968n^3} + \frac{-1823}{209952n^4} + \frac{-1668011}{37791360n^5} \Theta_{34}(n)\right).$$

For all $n \geq 6$ we have that $0 < \Theta_{34}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{34}(n) \rightarrow 1$.

Series 35. If we denote by $1/\pi_{35}(n)$ the truncated series

$$\frac{1}{\pi_{35}(n)} = \frac{3}{2} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k}{(k!)^3} \left(k + \frac{1}{6}\right) \left(\frac{1}{4}\right)^k,$$

then we have that $\pi_{35}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{35}(n) - \pi = \left(\frac{1}{4}\right)^n \frac{2\sqrt{\pi}}{\sqrt{n}} \cdot \exp\left(\frac{-3}{8n} + \frac{1}{4n^2} + \frac{-31}{64n^3} + \frac{41}{32n^4} + \frac{-2723}{640n^5} \Theta_{35}(n)\right).$$

For all $n \geq 4$ we have that $0 < \Theta_{35}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{35}(n) \rightarrow 1$.

Series 36. If we denote by $1/\pi_{36}(n)$ the truncated series

$$\frac{1}{\pi_{36}(n)} = \frac{21}{8} \sum_{k=0}^{n-1} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k \left(\frac{1}{2}\right)_k}{(k!)^3} \left(k + \frac{5}{42}\right) \left(\frac{1}{64}\right)^k,$$

then we have that $\pi_{36}(n) \rightarrow \pi$ as $n \rightarrow \infty$ and we have the asymptotic expansion

$$\pi_{36}(n) - \pi = \left(\frac{1}{64}\right)^n \frac{8\sqrt{\pi}}{3\sqrt{n}} \cdot \exp\left(\frac{-19}{72n} + \frac{1}{324n^2} + \frac{667}{139968n^3} + \frac{2689}{209952n^4} + \frac{-753139}{37791360n^5} \Theta_{36}(n)\right).$$

For all $n \geq 2$ we have that $0 < \Theta_{36}(n) < 1$ and for $n \rightarrow \infty$ we have that $\Theta_{36}(n) \rightarrow 1$.