

Estimates for Interpolation Projectors and Related Problems in Computational Geometry

Mikhail Nevskii* and Alexey Ukhalov†

August 2, 2021

This paper contains a survey of results obtained by the authors mostly during the past few years and published by 2021. In particular, we present the best of known estimates of numerical characteristics related to the research theme.

Sections: 1. Introduction. 2. The case when $n + 1$ is an Hadamard number. 3. Estimates for the minimal absorption index of a cube by a simplex. 4. Estimates for the minimal norm of a projector in linear interpolation on a cube in \mathbb{R}^n . 5. Estimates of numbers ξ'_n and θ'_n . 6. Simplices satisfying the inclusions $S \subset Q_n \subset nS$. 7. Perfect simplices. 8. Equisecting simplices. 9. Properties of $(0, 1)$ -matrices of order n having maximal determinant. 10. Problems for a simplex and a Euclidean ball. 11. Linear interpolation on a Euclidean ball.

Bibliography: 56 titles.

Keywords: simplex, cube, Euclidean ball, homothety, axial diameter, absorption index, Hadamard number, interpolation, projector, norm, estimate.

1 Introduction

This paper provides a survey of some results obtained by the authors mainly in the last seven years. The subject of our consideration is polynomial interpolation in several variables and also application in this area of geometric constructions and methods. As analysis in general, approximation theory is closely related to geometry.

*Department of Mathematics, P. G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia, mnevsk55@yandex.ru, orcid.org/0000-0002-6392-7618

†Department of Mathematics, P. G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia, alex-uhalov@yandex.ru, orcid.org/0000-0001-6551-5118

It is well known that many fundamental results in approximation theory essentially have geometric character. In detail, this connection is discovered in literature.

In polynomial interpolation of functions in several variables, the important geometric questions arise in connection with the structure of the basic domain and the choice of the nodes set. In particular, when interpolating functions of n variables using the space of polynomials of degree ≤ 1 , the interpolation nodes coincide with vertices of an n -dimensional nondegenerate simplex. By this, it occurs to be possible to provide estimates of the interpolation projector's norm through some geometric or numerical characteristics of the corresponding simplex. This is where we can use the properties of some numeric characteristics of a simplex and another convex body (volume, axial diameters, minimal positive homothety coefficients related to absorption a convex body by a homothet of a simplex, with or without translation). This approach can be carried over also to interpolation with wider spaces of polynomials.

The basic theoretical apparatus in this area was developed by M. Nevsky approximately in 2000–2014. Here we note the monograph [26] containing many results and links, the papers [14]–[15], [17]–[23], [25], [27]–[29], and also the dissertation [30]. A. Ukhalov joined the research starting around 2015. In particular, he very successfully applied computer methods in this area. The is written mainly on the basis of papers published by the authors or their students in 2016–2020. In order to make the text more understandable, we also cite the works written in previous years.

This paper continues the series of review articles in the anniversary collections issued for the 25th, 30th, 35th, and 40th anniversaries of Department of Mathematics of P. G. Demidov Yaroslavl State University (see [8], [16], [24], [31]). The original Russian text was prepared for a collection dedicated to 45th anniversary of the faculty which took place in 2021.

We start with main definitions and denotations. Everywhere further $n \in \mathbb{N}$. An element $x \in \mathbb{R}^n$ is written in the form $x = (x_1, \dots, x_n)$. By e_1, \dots, e_n we denote the standard basis in \mathbb{R}^n . By definition,

$$\|x\| := \sqrt{(x, x)} = \left(\sum_{i=1}^n x_i^2 \right)^{1/2},$$

$$B(x^{(0)}; R) := \{x \in \mathbb{R}^n : \|x - x^{(0)}\| \leq R\} \quad (x^{(0)} \in \mathbb{R}^n, R > 0),$$

$$B_n := B(0; 1), \quad Q_n := [0, 1]^n, \quad Q'_n := [-1, 1]^n.$$

The notation $L(n) \asymp M(n)$ means that there exist $c_1, c_2 > 0$ not depending on n such that $c_1 M(n) \leq L(n) \leq c_2 M(n)$.

Let Ω be a convex body in \mathbb{R}^n , i. e., a compact convex subset of \mathbb{R}^n with nonempty interior. By $\sigma\Omega$ we denote a homothetic copy of C with the center of homothety in the center of gravity of C and the ratio of homothety σ . The symbol $\text{vol}(\Omega)$ denotes the volume of Ω . If Ω is a convex polytope, then $\text{ver}(\Omega)$ is a set of vertices of Ω . By a *translate* we mean the result of a parallel shift.

An n -dimensional simplex S is circumscribed around a convex body Ω if $\Omega \subset S$ and each $(n - 1)$ -dimensional face of S contains a point of Ω . A convex polytope is inscribed into Ω if every vertex of this polytope belongs to the boundary of Ω .

Define $d_i(\Omega)$ as the minimal length of a segment contained in Ω and parallel to the x_i -axis. We call $d_i(\Omega)$ *the i th axial diameter of Ω* . The notion *axial diameter of a convex body* was introduced by P. Scott [48],[49].

For convex bodies Ω_1, Ω_2 , let us denote by $\xi(\Omega_1; \Omega_2)$ the minimal $\sigma \geq 1$ having the property $\Omega_1 \subset \sigma\Omega_2$. An equality $\xi(\Omega_1; \Omega_2) = 1$ is equivalent to the inclusion $\Omega_1 \subset \Omega_2$. We call $\xi(\Omega_1, \Omega_2)$ *the absorption index of Ω_1 by Ω_2* . Define $\alpha(\Omega_1, \Omega_2)$ as minimal $\sigma > 0$ such that Ω_1 is a subset of a translate of $\sigma\Omega_2$. Clearly, $\alpha(\Omega_1, \Omega_2) \leq \xi(\Omega_1, \Omega_2)$. We put $\xi(\Omega) := \xi(Q_n; \Omega)$, $\alpha(\Omega) := \alpha(Q_n; \Omega)$.

The symbol $C(\Omega)$ denotes a space of continuous functions $f : \Omega \rightarrow \mathbb{R}$ with the uniform norm

$$\|f\|_{C(\Omega)} := \max_{x \in \Omega} |f(x)|.$$

By $\Pi_1(\mathbb{R}^n)$ we mean a set of polynomials in n variables of degree ≤ 1 , i. e., a set of linear functions upon \mathbb{R}^n .

Let S be a nondegenerate simplex in \mathbb{R}^n with vertices $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)})$, $1 \leq j \leq n + 1$. *The vertices matrix* of this simplex

$$\mathbf{A} := \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}$$

is nondegenerate and $\text{vol}(S) = \frac{1}{n!} |\det(\mathbf{A})|$. Assume $\mathbf{A}^{-1} = (l_{ij})$. Define $\lambda_j \in \Pi_1(\mathbb{R}^n)$ as polynomials with coefficients forming the columns of \mathbf{A}^{-1} :

$$\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}.$$

These polynomials have the property $\lambda_j(x^{(k)}) = \delta_j^k$, where δ_j^k is the Kronecker delta. We call λ_j *the basic Lagrange polynomials corresponding to S* . For $x \in \mathbb{R}^n$, the numbers $\lambda_1(x), \dots, \lambda_{n+1}(x)$ are *the barycentric coordinates of x with respect to S* . Equations $\lambda_j(x) = 0$ give the $(n - 1)$ -dimensional hyperplanes containing the faces of S . We have

$$S = \{x \in \mathbb{R}^n : \lambda_j(x) \geq 0, j = 1, \dots, n + 1\}.$$

For more information on λ_j , see [26], [46].

The first author proved that

$$\xi(\Omega; S) = (n + 1) \max_{1 \leq k \leq n+1} \max_{x \in \Omega} (-\lambda_k(x)) + 1 \quad (\Omega \not\subset S), \quad (1)$$

$$\alpha(\Omega; S) = \sum_{j=1}^{n+1} \max_{x \in \Omega} (-\lambda_j(x)) + 1. \quad (2)$$

(see [23]; the proofs are given also in [26]). In the present text, we consider mainly the cases $\Omega = Q_n$ and $\Omega = B_n$.

If $\Omega = Q_n$, then (1) and (2) are equivalent correspondingly to the equalities

$$\xi(S) = (n+1) \max_{1 \leq k \leq n+1} \max_{x \in \text{ver}(Q_n)} (-\lambda_k(x)) + 1, \quad (3)$$

$$\alpha(S) = \sum_{j=1}^{n+1} \max_{x \in \text{ver}(Q_n)} (-\lambda_j(x)) + 1. \quad (4)$$

The equality $\xi(S) = \alpha(S)$ holds true if and only if the simplex $\xi(S)S$ is circumscribed around Q_n . This is also equivalent to the relation

$$\max_{x \in \text{ver}(Q_n)} (-\lambda_1(x)) = \dots = \max_{x \in \text{ver}(Q_n)} (-\lambda_{n+1}(x)). \quad (5)$$

Very important result is the following formula for $\alpha(S)$ obtained in [23]:

$$\alpha(S) = \sum_{i=1}^n \frac{1}{d_i(S)}. \quad (6)$$

It was proved in [21] that

$$\frac{1}{d_i(S)} = \frac{1}{2} \sum_{j=1}^{n+1} |l_{ij}|. \quad (7)$$

From (6) and (7), we have

$$\alpha(S) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^{n+1} |l_{ij}|. \quad (8)$$

Obviously, $\alpha(\Omega_1; \Omega_2)$ is invariant under the set translations and $\alpha(\tau\Omega_1; \Omega_2) = \tau\alpha(\Omega_1; \Omega_2)$, $\tau > 0$. Since $Q'_n = [-1, 1]^n$ is a translate of $2Q_n$, equality (8) gives

$$\alpha(Q'_n; S) = \sum_{i=1}^n \sum_{j=1}^{n+1} |l_{ij}|. \quad (9)$$

Relation (6) has a lot of useful corollaries given in [23] and [26]. For instance, from (6) it follows that for a simplex $S \subset Q_n$

$$\xi(S) \geq \alpha(S) = \sum_{i=1}^n \frac{1}{d_i(S)} \geq n, \quad (10)$$

since $d_i(S) \leq 1$. If $\xi(S) = n$, then $\alpha(S) = n$ и $d_i(S) = 1$.

Let us note also a result obtained on another way by M. Lassak [12]. *If a simplex $S \subset Q_n$ has the maximum volume, then $d_i(S) = 1$.* Indeed, for such an S holds $S \subset -nS$. If this was not so, then some vertex of S may be moved in Q_n in such

a way that its distance from the opposite face of the simplex increases. In this case the volume of S was not maximum. The inclusion $Q_n \subset -nS$ means that $-Q_n$ is a subset of a translate of nS . Since $-Q_n = Q_n$, we have $\alpha(S) \leq n$. As it was marked, if $S \subset Q_n$, then $\alpha(S) \geq n$. Therefore, $\alpha(S) = n$ and (6) gives $d_i(S) = 1$. This property of a maximum volume simplex essentially was used in [44].

Computational formulæ for $\xi(B_n; S)$ and $\alpha(B_n; S)$ were obtained by M. Nevskii in [32] (see Section 8).

Define the value

$$\xi_n(\Omega) := \min\{\xi(\Omega; S) : S \text{ is an } n\text{-dimensional simplex, } S \subset \Omega, \text{vol}(S) \neq 0\}.$$

In other words, $\xi_n(\Omega)$ is the minimal absorption index of a convex body Ω by an inner nondegenerate simplex. The results of the present survey are connected mainly with the case $\Omega = Q_n$.

Let $\xi_n := \xi_n(Q_n)$. If $S \subset Q_n$, then $\xi(S) \geq n$, see (10). This estimate occurs to be exact in order of n . Really, consider for $n > 2$ the simplex with the zero-vertex and other n vertices coinciding with the vertices of Q_n adjacent to $(1, \dots, 1)$. For this simplex, $\xi(S) = \frac{n^2-3}{n-1}$. Hence, for $n > 2$,

$$\xi_n \leq \frac{n^2 - 3}{n - 1}. \quad (11)$$

If $n > 1$, the right-hand part of (11) is strictly smaller than $n + 1$. The inequality $\xi_n < n + 1$ holds true also for $n = 1, 2$. If a simplex $S \subset Q_n$ has the maximum volume in Q_n , then $\xi(S) \leq n + 2$. For details, see [26].

Thus, always $n \leq \xi_n < n + 1$, i. e., $\xi_n - n \in [0, 1)$. However, the exact values of ξ_n still are known only for $n = 2$, $n = 5$, $n = 9$ and also for infinite set of n when there exists an Hadamard matrix of order $n + 1$. In all these cases except $n = 2$, holds $\xi_n = n$ while $\xi_2 = \frac{3\sqrt{5}}{5} + 1 = 2.34\dots$. Dimension $n = 2$ still remains a unique from even n when ξ_n was calculated exactly.

The accurate value of $\xi_n(B_n)$ is given in [32]. If $S \subset B_n$, then $\xi(B_n; S) \geq n$, with an equality only for a regular simplex inscribed into the boundary sphere. (For brevity, further we will say that such a simplex is inscribed into the ball.) Consequently, $\xi_n(B_n) = n$.

Let $x^{(j)} \in \Omega$ be the vertices of an n -dimensional nondegenerate simplex S . We say that an interpolation projector $P : C(\Omega) \rightarrow \Pi_1(\mathbb{R}^n)$ corresponds to S if its interpolation nodes coincides with the points $x^{(j)}$. This projector is given by the equalities $Pf(x^{(j)}) = f_j := f(x^{(j)})$. The following analogue of the Lagrange interpolation formula takes place:

$$Pf(x) = \sum_{j=1}^{n+1} f(x^{(j)}) \lambda_j(x). \quad (12)$$

Let us denote by $\|P\|_\Omega$ the norm of P as an operator from $C(\Omega)$ to $C(\Omega)$. It follows from (12) that

$$\|P\|_\Omega = \sup_{\|f\|_{C(\Omega)}=1} \|Pf\|_{C(\Omega)} = \sup_{-1 \leq f_j \leq 1} \max_{x \in \Omega} \sum_{j=1}^{n+1} f_j \lambda_j(x).$$

The expression $\sum f_j \lambda_j(x)$ is linear in x and f_1, \dots, f_{n+1} , hence,

$$\|P\|_\Omega = \max_{f_j = \pm 1} \max_{x \in \Omega} \sum_{j=1}^{n+1} f_j \lambda_j(x) = \max_{x \in \Omega} \max_{f_j = \pm 1} \sum_{j=1}^{n+1} f_j \lambda_j(x) = \max_{x \in \Omega} \sum_{j=1}^{n+1} |\lambda_j(x)|. \quad (13)$$

If Ω is a convex polytope in \mathbb{R}^n (e. g., $\Omega = Q_n$), then we have also a simpler equality

$$\|P\|_\Omega = \max_{x \in \text{ver}(\Omega)} \sum_{j=1}^{n+1} |\lambda_j(x)|.$$

By $\theta_n(\Omega)$ define the minimal value of $\|P\|_\Omega$ under the condition $x^{(j)} \in \Omega$. Put $\theta_n := \theta_n(Q_n)$.

We say that a point $x \in \Omega$ is an *1-point of Ω with respect to S* , if the projector $P : C(\Omega) \rightarrow \Pi_1(\mathbb{R}^n)$ with nodes in the vertices of S satisfies an equality $\|P\| = \sum |\lambda_j(x)|$ and among numbers $\lambda_j(x)$ there is a unique negative. It was proved in [19] that for any P and the corresponding S

$$\frac{n+1}{2n} (\|P\|_\Omega - 1) + 1 \leq \xi(\Omega; S) \leq \frac{n+1}{2} (\|P\|_\Omega - 1) + 1. \quad (14)$$

If in Ω there exists an 1-point with respect to S , then the right-hand inequality becomes an equality. The latter property in equivalent form was proved in [19]; the notion *an 1-vertex of the cube* was introduced later in [20].

Suppose $\Omega = Q_n$ or $\Omega = B_n$. If $S \subset \Omega$, then $\xi(\Omega; S) \geq n$. Applying a right-hand inequality from (14), we see that for any projector having nodes in Ω

$$\|P\|_\Omega \geq 3 - \frac{4}{n+1}.$$

This implies, for all n ,

$$\theta_n \geq 3 - \frac{4}{n+1}, \quad \theta_n(B_n) \geq 3 - \frac{4}{n+1}.$$

More advanced estimates of numbers θ_n and $\theta_n(B_n)$ were obtained with the use of the Legendre polynomials. *The standardized Legendre polynomial of degree n* is a function

$$\chi_n(t) := \frac{1}{2^n n!} [(t^2 - 1)^n]^{(n)}$$

(Rodrigues formula). For properties of χ_n see, e. g., [50]. The Legendre polynomials are orthogonal upon the segment $[-1, 1]$ with the weight $w(t) = 1$. As is known, $\chi_n(1) = 1$; if $n \geq 1$, then $\chi_n(t)$ increases for $t \geq 1$. By χ_n^{-1} , we denote a function inverse to χ_n on the half-axis $[1, +\infty)$.

The appearance of the Legendre polynomials in our questions is connected with the following their property. For $\gamma \geq 1$, let us consider the set

$$E_{n,\gamma} := \left\{ x \in \mathbb{R}^n : \sum_{j=1}^n |x_j| + \left| 1 - \sum_{j=1}^n x_j \right| \leq \gamma \right\}.$$

In 2003, M. Nevskii [14] established the equality

$$\text{mes}_n(E_{n,\gamma}) = \frac{\chi_n(\gamma)}{n!} \quad (15)$$

(the proof is given also in [26]). Applying this equality, the first author obtained a relation

$$\theta_n \geq \chi_n^{-1} \left(\frac{1}{\nu_n} \right) = \chi_n^{-1} \left(\frac{n!}{h_n} \right). \quad (16)$$

Here ν_n is the maximum volume of a simplex contained in Q_n , h_n is the maximum value of a determinant of order n consisting of 0's and 1's. From (16), some more visible inequalities were obtained. This led to a relation $\theta_n \asymp \sqrt{n}$. If $P : C(Q_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is an interpolation projector having the nodes in vertices of a maximum volume simplex in Q_n , then $\|P\|_{Q_n} \asymp \theta_n$ (with constants not depending on n).

Let us mark here some inequalities (links and proofs are available in [26]). For $n \neq 2$,

$$\frac{1}{e} \sqrt{n-1} < \theta_n \leq \min \left(\frac{n+1}{2}, \frac{4\sqrt{e}}{3} \sqrt{n+2} + 1 \right).$$

The left-hand inequality is valid for an arbitrary n . Also

$$\frac{1}{4} \sqrt{n} < \theta_n < 3\sqrt{n}, \quad n \in \mathbb{N}.$$

For a projector corresponding to a maximum volume simplex, we have

$$\|P\|_{Q_n} \leq \min \left(n+1, \frac{4\sqrt{e}}{3} \sqrt{n+2} + 1 \right),$$

If additionally $n+1$ is an Hadamard number, then $\|P\|_{Q_n} \leq \sqrt{n+1}$.

An analogue of (16) holds true also for interpolation by linear functions on a Euclidean ball. It is proved in [33] that for each n

$$\theta_n(B_n) \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\sigma_n} \right). \quad (17)$$

Here $\varkappa_n := \text{vol}(B_n)$ and σ_n denotes the volume of an n -dimensional regular simplex inscribed into B_n . Utilizing (17), we obtain a relation $\theta_n(B_n) \asymp \sqrt{n}$. If $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is an interpolation projector with the nodes in vertices of a regular simplex inscribed into B_n , then $\sqrt{n} \leq \|P\|_{B_n} \leq \sqrt{n+1}$. In other words, for such a projector $\|P\|_{B_n} \asymp \theta_n(B_n)$ (see [33]). Moreover, at least for $1 \leq n \leq 4$ we have $\|P\|_{B_n} = \theta_n(B_n)$ (this is proved in [43] and [34] by different methods).

Nowadays, the exact values of θ_n are known only for $n = 1, 2, 3$, and $n = 7$ (see [26]). Namely,

$$\theta_1 = 1, \quad \theta_2 = \frac{2\sqrt{5}}{5} + 1 = 1.89\dots, \quad \theta_3 = 2, \quad \theta_7 = \frac{5}{2}.$$

For all these dimensions,

$$\xi_n = \frac{n+1}{2}(\theta_n - 1) + 1.$$

Moreover, if $n = 1, 3, 7$, then

$$\theta_n = 3 - \frac{4}{n+1}.$$

The exact values of $\theta_n(B_n)$ are known for $1 \leq n \leq 4$:

$$\theta_1(B_1) = 1, \quad \theta_2(B_2) = \frac{5}{3}, \quad \theta_3(B_3) = 2, \quad \theta_4(B_4) = \frac{11}{5}.$$

In all these cases

$$\theta_n(B_n) = 3 - \frac{4}{n+1}.$$

If $n \geq 5$, then $\theta_n(B_n) > 3 - \frac{4}{n+1}$. These results are obtained in [43].

In the present survey we will also discuss some estimates of numbers ξ'_n и θ'_n which are the analogues of ξ_n and θ_n under the condition when vertices of simplices coincide with vertices of the cube. By definition,

$$\xi'_n := \min\{\xi(S) : \text{ver}(S) \subset \text{ver}(Q_n)\}, \quad \theta'_n := \min\{\|P\|_{Q_n} : \text{ver}(S) \subset \text{ver}(Q_n)\}.$$

In the second equality S is a simplex with vertices in the nodes of a projector $P : C(Q_n) \rightarrow \Pi_1(\mathbb{R}^n)$. Obviously,

$$\xi'_n \geq \xi_n, \quad \theta'_n \geq \theta_n,$$

This leads to the corresponding inequalities for upper and lower bounds of these values. The numbers θ'_n were introduced in paper [14] which is one of the very first M. Nevskii's works in this field. The estimates of ξ'_n and θ'_n are collected in [42]; we will describe these results in Section 4.

For calculation and graph drawing, the authors often utilized the Wolfram Mathematica system (see, e. g., [55], [10], [52]).

2 The Case when $n + 1$ is an Hadamard Number

An *Hadamard matrix* of order m is a nondegenerate $(m \times m)$ -matrix \mathbf{H}_m with entries ± 1 which satisfies

$$\mathbf{H}_m^{-1} = \frac{1}{m} \mathbf{H}_m^T.$$

Some information on Hadamard matrices is given in M. Hall's book [6]. If \mathbf{H}_m exists, then $m = 1$, $m = 2$ or m is a multiple of 4. For an infinite set of numbers having form $m = 4k$, including $m = 2^l$, the existence of \mathbf{H}_m is long ago established. By 2008, the smallest m , for which it is not known whether there exists an Hadamard matrix of order m , was equal to 668. We will call m an *Hadamard number* if there exists an Hadamard matrix of order m .

When and only when $n + 1$ is an Hadamard number, there exists a regular simplex S which is inscribed into Q_n in such a way that the vertices of S coincide with vertices of the cube (see [7, Theorem 4.5]). In [22] and [26, § 3.2], it was proved by various methods that $\xi(S) = n$, so $\xi_n = n$. Here we give another proof given in [40] and directly related to Hadamard matrices.

Theorem 2.1. *Let $n + 1$ be an Hadamard number. Then there exists a regular simplex S with the following properties:*

- 1) $\text{ver}(S) \subset \text{ver}(Q_n)$;
- 2) $\xi_n = \xi(S) = \alpha(S) = n$;
- 3) $d_1(S) = \dots = d_n(S) = 1$;
- 4) *the simplex nS is circumscribed around Q_n in such a way that each $(n - 1)$ -dimensional face of nS contains the only one vertex of Q_n .*

Proof. Taking into consideration similarity, we can prove the statement for the cube $Q'_n = [-1, 1]^n$. Since $n + 1$ is an Hadamard number, there exists *the normalized Hadamard matrix* of order $n + 1$, i. e., the Hadamard matrix with the first row and the first column consisting of 1's (see [6, Chapter 14]). Let us write the rows of this matrix in the inverse order:

$$\mathbf{H} = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ \dots & \dots & \dots & \dots & 1 \\ \dots & \dots & \dots & \dots & 1 \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & 1 \end{pmatrix}.$$

The obtained matrix \mathbf{H} also is an Hadamard matrix of order $n + 1$. Consider the simplex S' with vertices formed by the first n numbers in rows of \mathbf{H} .

All the vertices of S' are also vertices of Q'_n , therefore, the simplex is inscribed into the cube. Let us show that S' is a regular simplex and the length of its edge is equal to $\sqrt{2(n + 1)}$. Let a, b be two different rows of \mathbf{H} . All elements of \mathbf{H} are ± 1 , hence, we have $\|a\|^2 = \|b\|^2 = n + 1$. Rows of an Hadamard matrix are mutually orthogonal, therefore,

$$\|a - b\|^2 = (a - b, a - b) = \|a\|^2 + \|b\|^2 - 2(a, b) = 2(n + 1).$$

Denote by u and w the vertices of S' obtained from a and b respectively by removing the last component. This last component is equal to 1. It follows that n -dimensional length of vector $u - w$ is equal to $(n + 1)$ -dimensional length of vector $a - b$, i. e., $\|u - w\| = \sqrt{2(n + 1)}$.

Denote by λ_j the basic Lagrange polynomials of S' . Since $\mathbf{H}^{-1} = \frac{1}{n+1}\mathbf{H}^T$, the coefficients of $(n+1)\lambda_j$ are situated in rows of \mathbf{H} . All the constant terms of these polynomials stand in the last column of \mathbf{H} . Consequently, they are equal to 1. This means that the constant terms of polynomials $-(n+1)\lambda_j$ are equal to -1 . Hence, for any $j = 1, \dots, n+1$,

$$(n+1) \max_{x \in \text{ver}(Q'_n)} (-\lambda_j(x)) = n-1.$$

The coefficients of polynomials $-(n+1)\lambda_j$ are equal to ± 1 , therefore, for any j , the vertex v of Q'_n , such that $(n+1)(-\lambda_j(v)) = n-1$, is unique. Indeed, $v = (v_1, \dots, v_n)$ is defined by equalities $v_i = -\text{sign } l_{ij}$, where l_{ij} are the coefficients of λ_j .

Now let us find $\xi(Q'_n; S')$ using a formula analogous to (3):

$$\xi(Q'_n; S') = (n+1) \max_{1 \leq j \leq n+1} \max_{x \in \text{ver}(Q'_n)} (-\lambda_j(x)) + 1 = n-1 + 1 = n.$$

Consider the similarity transformation which maps Q'_n into Q_n . This transformation also maps S' into a simplex inscribed into Q_n . Denote by S the image of S' . Obviously, $\xi(S) = \xi(S'; Q'_n) = n$. It follows that, if $n+1$ is an Hadamard number, then $\xi_n \leq n$. As we know, for any n , $\xi_n \geq n$. Hence, $\xi_n = \xi(S) = n$. Since the coefficient of similarity for a mapping Q'_n to Q_n equals $\frac{1}{2}$, the length of any edge of S equals $\sqrt{\frac{n+1}{2}}$.

The condition

$$\max_{x \in \text{ver}(Q'_n)} (-\lambda_1(x)) = \dots = \max_{x \in \text{ver}(Q'_n)} (-\lambda_{n+1}(x)) = \frac{n-1}{n+1}$$

and (5) mean that the simplex nS' is circumscribed around the cube Q'_n . From the above it also follows that each $(n-1)$ -face of nS' contains the only one vertex of Q'_n . Hence, the simplex nS is circumscribed around the cube Q_n in the same way. Since nS is circumscribed around Q_n , we have $\alpha(S) = \xi(S) = n$ and $d_i(S) = 1$. These equalities could be also obtained from inequalities $n \leq \alpha(S) \leq \xi(S)$ and the equality $\xi(S) = n$.

The equalities $d_i(S) = 1$ and $\alpha(S) = n$ can be also derived in another way. Note that a regular simplex inscribed into Q_n has the maximum volume among all the simplices contained in this cube (see [7, Theorem 2.4]). As we marked, each axial diameter of any maximum volume simplex in Q_n is equal to 1.

The theorem is proved. □

3 Estimates for the Minimal Absorption Index of a Cube by a Simplex

The above estimates

$$n \leq \xi_n < n+1, \quad n \in \mathbb{N} \tag{18}$$

and

$$\xi_n \leq \frac{n^2 - 3}{n - 1}, \quad n > 2$$

were recently specified for some dimensions n . Here we give all the exact values and the best estimates of ξ_n known for today. For $n = 1, \dots, 10$, see Table 1.

Table 1: Estimates of ξ_n ($1 \leq n \leq 10$)

n	ξ_n	n	ξ_n
1	1	6	$6 \leq \xi_6 < 6.0166$
2	$\frac{3\sqrt{5}}{5} + 1 = 2.3416\dots$	7	7
3	3	8	$8 \leq \xi_8 < 8.1355$
4	$4 \leq \xi_4 \leq \frac{19+5\sqrt{13}}{9} = 4.1141\dots$	9	9
5	5	10	$10 \leq \xi_{10} < 10.2342$

Clearly, $\xi_1 = 1$. The exact values of ξ_2 , ξ_3 , and ξ_7 were delivered by M. Nevskii (see [26]). Note that ξ_1 , ξ_3 , and ξ_7 can be also calculated as a corollary of Theorem 2.1, since 2, 4, and 8 are Hadamard numbers. As is noted, $n = 2$ remains a unique even n for which we know the exact ξ_n .

The given estimate for ξ_4 was obtained in [36]. In this paper, also an assumption was made that $\xi_4 = \frac{19+5\sqrt{13}}{9}$. The authors have not yet succeeded in proving or refuting this conjecture.

The value of ξ_5 and the clarified estimate for ξ_6 were firstly given in [37]. It was shown that there exists a one-parameter family of simplices $R = R(t)$ with a property $\xi(R) = 5$. In the same way, in [40] it was established that $\xi_9 = 9$.

For today, $n = 5$ is the minimal n for which an infinite set of extremal simplices was found. Later such the families were built also for $n = 7$ and $n = 9$ (see [40]). For $n = 5$, in [40] a two-parameter family of simplices $R = R(s, t)$ satisfying $\xi(R) = 5$ was built. In addition to being extremal, simplices from this family also have some other interesting properties (see Sections 6–8).

The most precise upper bounds for ξ_8 and ξ_{10} were obtained by A. Ukhalov and D. Fedulov in [4].

The given estimates for ξ_6 , ξ_8 , and ξ_{10} were discovered by numerical minimization of $\xi(S)$ as a function of vertices coordinates of a simplex.

Tables 2, 3, and 4 contain the most precise upper bounds of ξ_n for $11 \leq n \leq 118$. These estimates are obtained in [39]. In order to provide the estimates, we used the following approach. M. Nevskii [26] showed that if $S \subset Q_n$ is a maximum volume

Table 2: Estimates of ξ_n ($11 \leq n \leq 58$)

n	$\xi_n \leq$		n	$\xi_n \leq$		n	$\xi_n \leq$	
11	11	H	27	27	H	43	43	H
12	$\frac{184}{15} = 12.267$	B	28	$\frac{781}{27} = 28.926$	A	44	$\frac{1933}{43} = 44.953$	A
13	$\frac{529}{39} = 13.564$	B	29	$\frac{5963}{203} = 29.374$	B	45	$\frac{2493}{55} = 45.327$	B
14	$\frac{193}{13} = 14.846$	A	30	$\frac{2981}{98} = 30.418$	B	46	$\frac{2113}{45} = 46.956$	A
15	15	H	31	31	H	47	47	H
16	$\frac{163}{10} = 16.300$	B	32	$\frac{1600}{49} = 32.653$	B	48	$\frac{2301}{47} = 48.957$	A
17	$\frac{296}{17} = 17.412$	B	33	$\frac{543}{16} = 33.938$	A	49	$\frac{689}{14} = 49.214$	B
18	$\frac{321}{17} = 18.882$	AB	34	$\frac{5237}{152} = 34.454$	B	50	$\frac{1162}{23} = 50.522$	B
19	19	H	35	35	H	51	51	H
20	$\frac{596}{29} = 20.552$	B	36	$\frac{875}{24} = 36.458$	B	52	$\frac{2701}{51} = 52.961$	A
21	$\frac{219}{10} = 21.900$	A	37	$\frac{4139}{111} = 37.288$	B	53	$\frac{36707}{689} = 53.276$	B
22	$\frac{481}{21} = 22.905$	A	38	$\frac{2309}{60} = 38.483$	B	54	$\frac{2017}{37} = 54.514$	B
23	23	H	39	39	H	55	55	H
24	$\frac{339}{14} = 24.214$	B	40	$\frac{1808}{45} = 40.178$	B	56	$\frac{394}{7} = 56.286$	B
25	$\frac{379}{15} = 25.267$	B	41	$\frac{8479}{205} = 41.361$	B	57	$q = 57.618$	B
26	$\frac{4853}{182} = 26.665$	B	42	$\frac{40204}{945} = 42.544$	B	58	$\frac{31455}{539} = 58.358$	B

Table 3: Estimates of ξ_n ($59 \leq n \leq 106$)

n	$\xi_n \leq$		n	$\xi_n \leq$		n	$\xi_n \leq$	
59	59	H	75	75	H	91	91	H
60	$\frac{1985}{33} = 60.152$	B	76	$\frac{5773}{75} = 76.973$	A	92	$\frac{8461}{91} = 92.978$	A
61	$\frac{11219}{183} = 61.306$	B	77	$\frac{2963}{38} = 77.974$	A	93	$\frac{4323}{46} = 93.978$	A
62	$\frac{21299}{341} = 62.460$	B	78	$\frac{4081}{52} = 78.481$	B	94	$\frac{5857}{62} = 94.468$	B
63	63	H	79	79	H	95	95	H
64	$\frac{4093}{63} = 64.968$	A	80	$\frac{3937}{49} = 80.347$	B	96	$\frac{9213}{95} = 96.979$	A
65	$\frac{33949}{520} = 65.287$	B	81	$\frac{3653}{45} = 81.178$	B	97	$\frac{18863}{194} = 97.232$	B
66	$\frac{5918}{89} = 66.494$	B	82	$\frac{8990}{109} = 82.477$	B	98	$\frac{4234}{43} = 98.465$	B
67	67	H	83	83	H	99	99	H
68	$\frac{4621}{67} = 68.970$	A	84	$\frac{42406}{501} = 84.643$	B	100	$\frac{714817}{7110} = 100.537$	B
69	$\frac{2379}{34} = 69.971$	A	85	$\frac{152113}{1785} = 85.217$	B	101	$\frac{51097}{505} = 101.182$	B
70	$\frac{3313}{47} = 70.489$	B	86	$\frac{1643}{19} = 86.474$	B	102	$\frac{19555}{191} = 102.382$	B
71	71	H	87	87	H	103	103	H
72	$\frac{5181}{71} = 72.972$	A	88	$\frac{7741}{87} = 88.977$	A*	104	$\frac{10813}{103} = 104.981$	A*
73	$\frac{48128}{657} = 73.254$	B	89	$\frac{3959}{44} = 89.977$	A*	105	$\frac{5511}{52} = 105.981$	A
74	$\frac{2458}{33} = 74.485$	B	90	$\frac{1538}{17} = 90.471$	B	106	$\frac{7021}{66} = 106.379$	B

Table 4: Estimates of ξ_n ($107 \leq n \leq 118$)

n	$\xi_n \leq$		n	$\xi_n \leq$	
107	107	H	114	$\frac{24247}{212} = 114.373$	B
109	$\frac{107131}{981} = 109.206$	B	115	115	H
110	$\frac{22627}{205} = 110.376$	B	116	$\frac{13453}{115} = 116.983$	A*
111	111	H	117	$\frac{132583}{1131} = 117.226$	B
112	$\frac{6727}{60} = 112.117$	B	118	$\frac{8641}{73} = 118.370$	B
113	$\frac{25591}{226} = 113.235$	B			

simplex, then $\xi(S) \asymp \xi_n$. Calculation of $\xi(S)$ for maximum volume simplices often makes it possible to refine the upper bound in the double inequality

$$n \leq \xi_n \leq \frac{n^2 - 3}{n - 1} \quad (19)$$

(see (11) and (18)). For constructing maximum volume simplices in Q_n , we utilized some data from website <http://www.indiana.edu/~maxdet/> related to maximum $(-1, 1)$ -determinants. This site contained maximum $(-1, 1)$ -determinants up to the order 119. Among the listed determinants there were also those whose maximality has not been established. For some orders, the determinants were not given at all.

Along with upper bounds of ξ_n , the tables also provide information about methods of obtaining the estimates. We use the following designations:

- A — The estimate is obtained from inequality (19). The using of a maximum volume simplex did not improve this estimate.
- B — We give the smallest value $\xi(S)$ obtained from considering one or more simplices of maximum volume in Q_n . This value is less than the right-hand side of (19).
- AB — The value of $\xi(S)$ obtained for simplices of maximum volume coincides with the right-hand side of (19).
- H — $n + 1$ is an Hadamard number. We give the exact value $\xi_n = n$.

- A^* — We use inequality (19), since for this n no maximum determinant is given on the site.

For $n = 57$, we use denotation $q = \frac{34018994107517188105}{590424166322794597}$.

4 Estimates for the Minimal Norm of a Projector in Linear Interpolation on a Cube in \mathbb{R}^n

Evidently, $\theta_1 = 1$. The exact values of θ_2 , θ_3 , and θ_7 were obtained by M. Nevskii (see [26]). For today, $n = 1$, $n = 2$, $n = 3$, and $n = 7$ are the only cases when we know θ_n exactly. For other n , we know common estimates (see Section 1). In some cases, these estimates were clarified with the use of computer. Here we mostly give the material from [41].

Table 5 contains the most precise upper estimates of θ_n for $1 \leq n \leq 27$. If we know the exact value, we write this value. For providing estimates, the following approach described in [26] may be used. The upper bound of θ_n appears from consideration of an projector having the nodes in vertices of a maximum volume simplex in Q_n . For the first time, this method was applied by M. Nevskii and I. Hlestkova in [35].

A significant part of the given estimates was obtained using actually this approach. For constructing maximum volume simplices for a given n , we utilized maximal $(-1, 1)$ -determinants of order $n + 1$ from the site <http://www.indiana.edu/~maxdet/>. Hadamard matrices were imported from website <http://www.maths.gla.ac.uk/~es/hadamard/hadamard.php>.

The maximality of $(-1, 1)$ -determinants of order $1, \dots, 21$ listed on website www.indiana.edu/~maxdet is proved. Moreover, the website contains completely all the maximum determinants of order ≤ 21 (up to equivalence of matrices). While getting the estimates, we used all these data. The maximality of determinants of order 22, 23, and 27 has not been established. Such situations are marked in the table with a question mark. It is unclear whether is exhaustive the set of maximum determinants of order 25 and 26 given on the site. To refine the upper estimate of θ_n , we examined all the known maximum $(-1, 1)$ -determinants of order $n + 1$, since the corresponding projectors can have different norms.

Calculations using the formula

$$\|P\|_{Q_n} = \max_{x \in \text{ver}(Q_n)} \sum_{j=1}^{n+1} |\lambda_j(x)|. \quad (20)$$

reduce to looking through all 2^n vertices of Q_n . For large n , these calculations become rather time-consuming. This is the reason why we limited by investigation of cases $1 \leq n \leq 27$.

In all the situations when $n + 1$ is an Hadamard number, we used the complete set of Hadamard matrices of the appropriate order. In particular, to obtain the estimate

Table 5: Upper estimates of θ_n ($1 \leq n \leq 27$)

n	min P 	N	$\theta_n \leq$	
1	1	1	$\theta_1 = 1$	N
2	3	1	$\theta_2 = \frac{2\sqrt{5}}{5} + 1 = 1.8944 \dots$	N
3	2	1	$\theta_3 = 2$	N
4	$\frac{7}{3} = 2.3333 \dots$	1	$\frac{3(4+\sqrt{2})}{7} = 2.3203 \dots$	K
5	$\frac{13}{5} = 2.6$	1	2.448804	NU
6	3	1	2.60004...	L
7	$\frac{5}{2} = 2.5$	1	$\theta_7 = \frac{5}{2} = 2.5$	N
8	$\frac{22}{7} = 3.1428 \dots$	1	$\frac{22}{7} = 3.1428 \dots$	M
9	3	1	3	M
10	$\frac{19}{5} = 3.8$	3	$\frac{19}{5} = 3.5186 \dots$	L
11	3	1	3	H
12	$\frac{17}{5} = 3.4$	1	$\frac{17}{5} = 3.4$	M
13	$\frac{49}{13} = 3.7692 \dots$	1	$\frac{49}{13} = 3.7692 \dots$	M
14	$\frac{21}{5} = 4.2$	1	$\frac{21}{5} = 4.0156 \dots$	L
15	$\frac{7}{2} = 3.5$	5	$\frac{7}{2} = 3.5$	H
16	$\frac{21}{5} = 4.2$	3	$\frac{21}{5} = 4.2$	M
17	$\frac{139}{34} = 4.0882 \dots$	3	$\frac{139}{34} = 4.0882 \dots$	M
18	$\frac{95}{17} = 5.5882 \dots$	3	5.14006...	L
19	4	3	4	H
20	$\frac{137}{29} = 4.7241 \dots$	7	4.68879...	L
21	$\frac{251}{50} = 5.02$	1	$\frac{251}{50} = 5.02$	M?
22	$\frac{1817}{335} = 5.4238 \dots$	1	$\frac{1817}{335} = 5.4238 \dots$	M?
23	$\frac{9}{2} = 4.5$	60	$\frac{9}{2} = 4.5$	O
24	$\frac{103}{21} = 4.9047 \dots$	2	$\frac{103}{21} = 4.9047 \dots$	M
25	5	3	5	M
26	$\frac{474}{91} = 5.2087 \dots$	1	$\frac{474}{91} = 5.2087 \dots$	M?
27	5.0	487	5.0	U

of θ_{23} , we considered all the existing 60 Hadamard matrices of order 24. This case was studied by A. Ukhalov and E. Ozerova [11]. The estimate of θ_{27} was obtained by A. Ukhalov and E. Udovenko [51], [53]. To get the result, it was necessary to consider 487 Hadamard matrices of order 28.

For $n = 4, 5, 6, 10, 14, 18$, and 20, the estimates of θ_n were made via minimization of the function standing in the right-hand part (20). The links are given in the description of the table.

The authors conjecture that $\theta_4 = \frac{3(4+\sqrt{2})}{7} = 2.320377\dots$

In the first column of Table 5, we put dimension n . The second column contains the minimal projector norm obtained via consideration of maximum $(-1, 1)$ -determinants of order $n + 1$. The third column shows number N of matrices of order $n + 1$ considered for obtaining the result written in the second column. In the fourth column, the best nowadays upper bound of θ_n is given. The last column of the table contains a comment on the way to get the top boundary of θ_n indicated in the fourth column. Here we use the following notation.

- H — The case when $n + 1$ is an Hadamard number. There was used a regular simplex built from the corresponding Hadamard matrix.
- K — The estimate is obtained by I. Kudryavcev (see [11]).
- L — The estimate is obtained by A. Luterkov in [13].
- M — There was used a maximum volume simplex in Q_n built from maximum $(-1, 1)$ -determinant of order $n + 1$.
- M? — There was used a simplex in Q_n built from determinant of order $n + 1$ which is maximum from known nowadays. The global maximality of such a determinant yet is not proved.
- N — The exact value of θ_n is obtained by M. Nevskii (see [26]).
- NU — The upper estimate is obtained by M. Nevskii and A. Ukhalov in [37].
- O — The upper estimate is obtained by E. Ozerova (see [11]).
- U — The upper estimate is obtained by E. Udovenko in [51].

For $n = 5$, E. Spasova (2020, bachelor's thesis under the guidance of M. Nevskii) constructed a projector having the norm $\|P\|_{Q_5} = \frac{5}{2} = 2.5$. This value exceeds the smallest known upper bound $2.4488\dots$ for θ_5 established in [37]. Nevertheless, the nodes of P have a rather simple form — they are located at the vertices or at midpoints of the edges of Q_5 . For example, the following points are suitable: $(0, 0, 0, 0, 0)$, $(0, 0, \frac{1}{2}, 1, 1)$, $(0, 1, \frac{1}{2}, 0, 1)$, $(0, 1, 1, 1, 0)$, $(1, 0, 1, 0, \frac{1}{2})$, $(1, 1, 0, 1, \frac{1}{2})$. Back in 2006, a projector with nodes in the vertices of Q_5 and the norm 2.6 was built (see, e. g., [15]), and for at least 10 years the estimate $\theta_5 \leq 2.6$ was the best from the known.

Table 6: Lower estimates of θ_n ($1 \leq n \leq 54$)

n	$\chi_n^{-1}(\frac{1}{\nu_n})$	$\mathbf{3} - \frac{4}{n+1}$	Max	n	$\chi_n^{-1}(\frac{1}{\nu_n})$	$\mathbf{3} - \frac{4}{n+1}$	Max
1	1	1	1	28	2.2768	2.8621	2.8621
2	1.291	1.6667	1.6667	29	2.3074	2.8667	2.8667
3	1.2492	2	2	30	2.3487	2.871	2.871
4	1.3478	2.2	2.2	31	2.3452	2.875	2.875
5	1.4284	2.3333	2.3333	32	2.3955	2.8788	2.8788
6	1.5018	2.4286	2.4286	33	2.4259	2.8824	2.8824
7	1.4678	2.5	2.5	34	2.4642	2.8857	2.8857
8	1.5626	2.5556	2.5556	35	2.4601	2.8889	2.8889
9	1.6034	2.6	2.6	36	2.5019	2.8919	2.8919
10	1.6699	2.6364	2.6364	37	2.5348	2.8947	2.8947
11	1.6488	2.6667	2.6667	38	2.5722	2.8974	2.8974
12	1.7086	2.6923	2.6923	39	2.5697	2.9	2.9
13	1.7659	2.7143	2.7143	40	2.6056	2.9024	2.9024
14	1.8211	2.7333	2.7333	41	2.641	2.9048	2.9048
15	1.8108	2.75	2.75	42	2.6759	2.907	2.907
16	1.8778	2.7647	2.7647	43	2.6747	2.9091	2.9091
17	1.9156	2.7778	2.7778	44	2.7179	2.9111	2.9111
18	1.965	2.7895	2.7895	45	2.743	2.913	2.913
19	1.9587	2.8	2.8	46	2.7791	2.9149	2.9149
20	2.0159	2.8095	2.8095	47	2.7756	2.9167	2.9167
21	2.0588	2.8182	2.8182	48	2.8201	2.9184	2.9184
22	2.1039	2.8261	2.8261	49	2.8413	2.92	2.92
23	2.0958	2.8333	2.8333	50	2.8805	2.9216	2.9216
24	2.1408	2.84	2.84	51	2.8729	2.9231	2.9231
25	2.1847	2.8462	2.8462	52	2.9173	2.9245	2.9245
26	2.2278	2.8519	2.8519	53	2.9362	2.9259	2.9362
27	2.2242	2.8571	2.8571	54	2.9735	2.9273	2.9735

Now let us pass to the lower bounds of θ_n . Tables 6, 7, and 8 present the largest known lower bounds for these numbers. To obtain the estimates, we used the inequality

$$\theta_n \geq \max \left[3 - \frac{4}{n+1}, \chi_n^{-1} \left(\frac{1}{\nu_n} \right) \right]. \quad (21)$$

For each n , we give the values $\chi_n^{-1} \left(\frac{1}{\nu_n} \right)$ and $3 - \frac{4}{n+1}$, as well as the maximum of them. This maximum is the most accurate lower bound for θ_n . The tables are missing $n = 104$ and $n = 116$, since website www.indiana.edu/~maxdet does not show any maximum determinants of orders 105 and 117.

Graphs of the functions $\chi_n^{-1} \left(\frac{1}{\nu_n} \right)$, $3 - \frac{4}{n+1}$ and $\frac{\sqrt{n-1}}{e}$ are shown in Fig. 1. These functions are defined for an integer argument, but when constructing the graphs, we use continuous lines of various type. The inequality

$$\chi_n^{-1} \left(\frac{1}{\nu_n} \right) > 3 - \frac{4}{n+1}$$

holds for $n \geq 53$.




Figure 1: Graphs of the functions $\chi_n^{-1} \left(\frac{1}{\nu_n} \right)$, $3 - \frac{4}{n+1}$, and $\frac{\sqrt{n-1}}{e}$

Table 7: Lower estimates of θ_n ($55 \leq n \leq 109$)

n	$\chi_n^{-1}\left(\frac{1}{\nu_n}\right)$	$\mathbf{3} - \frac{4}{n+1}$	Max	n	$\chi_n^{-1}\left(\frac{1}{\nu_n}\right)$	$\mathbf{3} - \frac{4}{n+1}$	Max
55	2.9669	2.9286	2.9669	82	3.5595	2.9518	3.5595
56	3.0034	2.9298	3.0034	83	3.5546	2.9524	3.5546
57	3.0315	2.931	3.0315	84	3.5899	2.9529	3.5899
58	3.0583	2.9322	3.0583	85	3.6053	2.9535	3.6053
59	3.058	2.9333	3.058	86	3.6355	2.954	3.6355
60	3.0877	2.9344	3.0877	87	3.6307	2.9545	3.6307
61	3.1173	2.9355	3.1173	88	3.667	2.9551	3.667
62	3.1465	2.9365	3.1465	89	3.6803	2.9556	3.6803
63	3.1463	2.9375	3.1463	90	3.7099	2.956	3.7099
64	3.1849	2.9385	3.1849	91	3.7052	2.9565	3.7052
65	3.2039	2.9394	3.2039	92	3.7401	2.957	3.7401
66	3.2375	2.9403	3.2375	93	3.755	2.9574	3.755
67	3.2322	2.9412	3.2322	94	3.783	2.9579	3.783
68	3.2717	2.942	3.2717	95	3.7781	2.9583	3.7781
69	3.2888	2.9429	3.2888	96	3.8121	2.9588	3.8121
70	3.32	2.9437	3.32	97	3.8258	2.9592	3.8258
71	3.3158	2.9444	3.3158	98	3.8542	2.9596	3.8542
72	3.3529	2.9452	3.3529	99	3.8497	2.96	3.8497
73	3.3703	2.9459	3.3703	100	3.8814	2.9604	3.8814
74	3.4023	2.9467	3.4023	101	3.8964	2.9608	3.8964
75	3.3973	2.9474	3.3973	102	3.9246	2.9612	3.9246
76	3.4348	2.9481	3.4348	103	3.92	2.9615	3.92
77	3.4521	2.9487	3.4521	105	3.9662	2.9623	3.9662
78	3.4818	2.9494	3.4818	106	3.9936	2.9626	3.9936
79	3.4769	2.95	3.4769	107	3.989	2.963	3.989
80	3.5138	2.9506	3.5138	108	4.0841	2.9633	4.0841
81	3.5288	2.9512	3.5288	109	4.034	2.9636	4.034

Table 8: Lower estimates of θ_n ($110 \leq n \leq 118$)

n	$\chi_n^{-1}(\frac{1}{\nu_n})$	$3 - \frac{4}{n+1}$	Max	n	$\chi_n^{-1}(\frac{1}{\nu_n})$	$3 - \frac{4}{n+1}$	Max
110	4.0615	2.964	4.0615	114	4.1281	2.9652	4.1281
111	4.0568	2.9643	4.0568	115	4.1234	2.9655	4.1234
112	4.0789	2.9646	4.0789	117	4.167	2.9661	4.167
113	4.101	2.9649	4.101	118	4.1937	2.9664	4.1937

5 Estimates of Numbers ξ'_n and θ'_n

Let us give estimates of numbers ξ'_n and θ'_n related to the situation when vertices of simplices located at vertices of the cube. By our definition,

$$\xi'_n := \min_{\text{ver}(S) \subset \text{ver}(Q_n)} \xi(S), \quad \theta'_n := \min_{\text{ver}(S) \subset \text{ver}(Q_n)} \|P\|_{Q_n}. \quad (22)$$

In the right-hand equality, S is the simplex having vertices at the nodes of a projector $P : C(Q_n) \rightarrow \Pi_1(\mathbb{R}^n)$. The sequence θ'_n was introduced by M. Nevskii in [14]. He proved that $\theta'_n \geq \frac{1}{2e}\sqrt{n}$, and if $n+1$ is an Hadamard number, then $\theta'_n \leq \sqrt{n+1}$. More systematically the numbers (22) were studied by the authors in [42]. Here we describe some results of this work.

First note general estimates for ξ'_n и θ'_n . Many of them are also estimates for ξ_n and θ_n . The inequality holds

$$\theta'_n > \frac{1}{e}\sqrt{n-1}.$$

If $n \neq 2$, then

$$\theta'_n \leq \min\left(\frac{n+1}{2}, \frac{4\sqrt{e}}{3}\sqrt{n+2}+1\right).$$

For each n , we have $\xi'_n \geq n$. If n is even, then $\xi'_n > n$. If $n+1$ is an Hadamard number, then $\xi'_n = n$. For $n > 2$,

$$\xi'_n \leq \frac{n^2-3}{n-1}.$$

Thus, $\theta'_n \asymp n^{1/2}$, $\xi'_n \asymp n$.

Assume S is an n -dimensional $(0,1)$ -simplex having the maximum volume in Q_n and P is the corresponding interpolation projector. Then with constants not depending on n we have $\|P\|_{Q_n} \asymp \theta'_n$, $\xi(S) \asymp \xi'_n$.

For $n \geq 3$, the sequence $\{\xi'_n\}$ strictly increases.

Consider the n -dimensional simplex S^* for which the first n vertices are the vertices of the cube adjacent to $(1, \dots, 1)$ and the $(n+1)$ th vertex coincides with 0:

$$x^{(1)} = (0, 1, \dots, 1, 1), \dots, x^{(n)} = (1, 1, \dots, 1, 0), x^{(n+1)} = (0, \dots, 0).$$

The lengths of n edges of S^* containing $x^{(n+1)}$ are equal to $\sqrt{n-1}$; the lengths of the rest edges are $\sqrt{2}$. For $n = 3$ (and only in this case) S^* is a regular simplex. Starting $n = 3$, the simplex S^* has the following property (see [7, Lemma 3.3]): while replacing any vertex of S^* by any point of Q_n , the volume of the simplex strictly decreases. That is why for $n \geq 3$ the simplex S^* is called in [7] *rigid*. This property holds also for $n = 1$. In the two-dimensional case the volume of the simplex in such a way does not increase. If and only if $1 \leq n \leq 4$, the volume of S^* is the maximum possible for a simplex contained in Q_n .

It was proved in [26] that

$$\xi(S^*) = \begin{cases} 1, & n = 1; \\ 4, & n = 2; \\ \frac{n^2-3}{n-1}, & n \geq 3. \end{cases}$$

For each n , we have $d_1(S^*) = \dots = d_n(S^*) = 1$, $\alpha(S^*) = n$. The one-dimensional case is the single when $\alpha(S^*) = \xi(S^*)$. This means that the simplex $\xi(S^*)S^*$ is circumscribed around Q_n only for $n = 1$.

It turns out that for $1 \leq n \leq 5$ the simplex S^* delivers the exact values of ξ'_n , i. e., in these cases $\xi(S^*) = \xi'_n$. The minimal n satisfying $\xi(S^*) > \xi'_n$ is equal to 6. Let $P^* : C(Q_n) \rightarrow \Pi_1(\mathbb{R}^n)$ be the interpolation projector corresponding to S^* . The projector P^* is extremal with respect to θ'_n , if $1 \leq n \leq 4$. In these dimensions, $\|P^*\|_{Q_n} = \theta'_n$. The minimal n with a property $\|P^*\| > \theta'_n$ equals 5. Let us describe these cases in more details.

Suppose $n = 1$. We have $Q_1 = S^* = [0, 1]$, $\|P^*\|_{Q_1} = \xi(S^*) = 1$, hence, $\theta_1 = \xi_1 = \theta'_1 = \xi'_1 = 1$.

In the case $n = 2$ there is only one (up to similarity) $(0, 1)$ -simplex, namely, the simplex S^* with vertices $(0, 1)$, $(1, 0)$, $(0, 0)$. It follows that $\theta'_2 = 3$, $\xi'_2 = 4$.

The case $n = 3$ was analyzed by the first author (see, e. g., [26]). The minima of $\xi(S)$ and $\|P\|_{Q_3}$ are delivered by the simplex S^* with vertices $(0, 1, 1)$, $(1, 0, 1)$, $(1, 1, 0)$, $(0, 0, 0)$. We have $\xi_3 = \xi'_3 = 3$, $\theta_3 = \theta'_3 = 2$. The same values of $\xi(S)$ and $\|P\|_{Q_3}$ are given also by the simplex S^{**} with vertices $(\frac{1}{2}, 0, 0)$, $(\frac{1}{2}, 1, 0)$, $(0, \frac{1}{2}, 1)$, $(1, \frac{1}{2}, 1)$ (and by similar simplices). But vertices of S^{**} are not $(0, 1)$ -points. There are no another extremal simplices in the three-dimensional case.

For $n = 4$ and $n = 5$, we have performed the full search of n -dimensional simplices satisfying the condition $\text{ver}(S) \subset \text{ver}(Q_n)$. It was discovered that S^* and P^* give both the minima of $\xi(S)$ and $\|P\|_{Q_n}$. The following equalities hold: $\theta'_4 = \frac{7}{3}$, $\xi'_4 = \frac{13}{3}$. The simplex S^* remains an extremal also with respect to ξ_5 , i. e., $\xi_5 = \xi(S^*) = \frac{11}{2}$. It was obtained that $\theta_5 = \frac{13}{5}$. The inequality $\theta'_5 \leq \frac{13}{5}$ was established earlier in [26, п. 3.8.3].

Let us turn to the case $n = 6$. From consideration S^* , it follows that $6 \leq \xi'_6 \leq \frac{33}{5}$. The upper estimate may be improved. We have made the full search of six-dimensional simplices satisfying the condition $\text{ver}(S) \subset \text{ver}(Q_6)$. For each simplex, one vertex was fixed and the rest six vertices were chosen from the rest $2^6 - 1$

vertices of the cube. Thus, we were needed to look through $\binom{2^6-1}{6} = 67\,945\,521$ simplices. Calculations were performed with the use of a special programme in Wolfram Mathematica system (see, e. g., [55], [10]). The minimum of $\xi(S)$ equal to $\frac{25}{4}$ is delivered by the simplex with vertices $(0, 0, 0, 0, 1, 1)$, $(0, 0, 0, 1, 0, 1)$, $(0, 1, 1, 1, 1, 0)$, $(1, 0, 1, 1, 1, 0)$, $(1, 1, 0, 1, 1, 0)$, $(1, 1, 1, 0, 0, 1)$, $(0, 0, 0, 0, 0, 0)$. Therefore, $\xi'_6 = \frac{25}{4}$. Simultaneously, we detected that the minimal n , for which S^* does not give an exact value of ξ'_n , is equal to 6.

From the results of [26, п.3.8.3], it follows that $\theta'_6 \leq 3$. The full search of $(0, 1)$ -simplices in \mathbb{R}^6 have shown that the above simplex minimizes also the norm of an interpolation projector. For the corresponding projector, $\|P\|_{Q_6} = 3$. Thus, the exact value of θ'_6 equals 3. Note that $\|P^*\|_{Q_6} = \frac{17}{5} > \theta'_6$.

Table 9: The numbers ξ_n , ξ'_n , θ_n , and θ'_n for $1 \leq n \leq 7$

n	ξ_n	ξ'_n	θ_n	θ'_n
1	1	1	1	1
2	$\frac{3\sqrt{5}}{5} + 1 = 2.34\dots$	4	$\frac{2\sqrt{5}}{5} + 1 = 1.89\dots$	3
3	3	3	2	2
4	$4 \leq \xi_4 \leq \frac{19+5\sqrt{13}}{9} = 4.11\dots$	$\frac{13}{3}$	$\frac{11}{5} \leq \theta_4 \leq \frac{3(4+\sqrt{2})}{7} = 2.3203\dots$	$\frac{7}{3}$
5	5	$\frac{11}{2}$	$\frac{7}{3} \leq \theta_5 < 2.448804$	$\frac{13}{5}$
6	$6 \leq \xi_6 < 6.0166$	$\frac{25}{4}$	$\frac{17}{7} \leq \theta_6 \leq 2.60014$	3
7	7	7	$\frac{5}{2}$	$\frac{5}{2}$

Finally, consider the case $n = 7$. Since 8 is an Hadamard number, there exists a seven-dimensional regular simplex having vertices at vertices of Q_7 . This is, for example, the simplex S with vertices $(1, 1, 1, 1, 1, 1, 1)$, $(0, 1, 0, 1, 0, 1, 0)$, $(0, 0, 1, 1, 0, 0, 1)$, $(1, 0, 0, 1, 1, 0, 0)$, $(0, 0, 0, 0, 1, 1, 1)$, $(1, 0, 1, 0, 0, 1, 0)$, $(1, 1, 0, 0, 0, 0, 1)$, $(0, 1, 1, 0, 1, 0, 0)$. We have $\xi(S) = 7$. From inequalities $\xi'_n \geq \xi_n \geq n$, it follows that $\xi'_7 = \xi_7 = 7$. Now let us apply the estimate

$$\xi_n \leq \frac{n+1}{2} (\theta_n - 1) + 1 \quad (23)$$

which appears from the right-hand inequality in (14) for $\Omega = Q_n$. Since $\xi_7 = 7$, (23) yields $\theta_7 \geq \frac{5}{2}$. But the projector corresponding to S has the norm $\|P\|_{Q_7} = \frac{5}{2}$. Therefore, $\theta'_7 = \theta_7 = \frac{5}{2}$.

A collection of the known results related to the numbers ξ_n , ξ'_n , θ_n , and θ'_n for $1 \leq n \leq 7$ is given in Table 9.

6 Simplicies Satisfying the Inclusions $S \subset Q_n \subset nS$

As is noted, the absorption index of the unit cube by a simplex contained in this cube is not less than the dimension of the space. This is equivalent to the inequality $\xi_n \geq n$ which follows from (10). If for some $S \subset Q_n$ we have $\xi(S) = n$, then $\alpha(S) = \xi(S) = n$ and $\xi_n = n$. Moreover, for each i , holds $d_i(S) = 1$. On the contrary, the equality $\xi_n = n$ means that there exists a simplex $S \subset Q_n$ with the property $\alpha(S) = \xi(S) = n$, i. e., $S \subset nS$. However, simplicies satisfying $S \subset Q_n \subset nS$ do exist not for any n . The first such value is $n = 2$. Namely, $\xi_2 = 1 + \frac{3\sqrt{5}}{5} = 2.34\dots > 2$; nontrivial proof is given in [26]. Below we note another way to see that $\xi_2 > 2$.

Nowadays the equality $\xi_n = n$ is proved for $n = 5$, $n = 9$ and for an infinite set of n such that $n + 1$ is an Hadamard number. In particular, such are all odd n from an interval $1 \leq n \leq 11$. No one even n with the condition $\xi_n = n$ has not yet been found. Moreover, it remains unclear does there exist at least one such even n .

Simplicies satisfying the inclusions $S \subset Q_n \subset nS$, if exist, have a number of remarkable properties. Such simplicies were studied by the authors in [38].

Further $x^{(j)}$ are the vertices and $\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$ are the basic Lagrange polynomials of n -dimensional simplex S . By $c(\Omega)$ we denote the center of gravity of a convex body Ω . By definition,

$$D_j := \left\{ x \in \mathbb{R}^n : -\frac{n-1}{n+1} \leq \lambda_j(x) \leq 1 \right\}.$$

The following theorem combines the results of Theorem 1 and a number of corollaries from [38].

Theorem 6.1. *Suppose $S \subset Q_n \subset nS$. Then the following relation hold:*

$$\max_{x \in \text{ver}(Q_n)} \lambda_1(x) = \dots = \max_{x \in \text{ver}(Q_n)} \lambda_{n+1}(x) = 1, \quad (24)$$

$$\max_{x \in \text{ver}(Q_n)} (-\lambda_1(x)) = \dots = \max_{x \in \text{ver}(Q_n)} (-\lambda_{n+1}(x)) = \frac{n-1}{n+1}, \quad (25)$$

$$c(S) = c(Q_n) = \left(\frac{1}{2}, \dots, \frac{1}{2} \right), \quad (26)$$

$$\dots \subset \frac{1}{n^2} S \subset \frac{1}{n} Q_n \subset S \subset Q_n \subset nS \subset n^2 Q_n \subset n^3 S \subset \dots,$$

$$\sum_{j=1}^{n+1} |l_{ij}| = 2 \quad (1 \leq i \leq n), \quad \sum_{i=1}^n |l_{ij}| = \frac{2n}{n+1} \quad (1 \leq j \leq n+1),$$

$$\sum_{i \leq n: l_{ij} \geq 0} l_{ij} = 1 - l_{n+1,j}, \quad \sum_{i \leq n: l_{ij} < 0} |l_{ij}| = \frac{n-1}{n+1} + l_{n+1,j} \quad (1 \leq j \leq n+1),$$

$$Q_n \subset \bigcap_{j=1}^{n+1} D_j.$$

The equalities similar to (24)–(25) hold also in the case when maxima are taken over $x \in Q_n$.

Applying (26), it is rather easy to prove that $\xi_2 > 2$. Earlier the exact value of ξ_2 was obtained by M. Nevskii (see [26]). Let us note some other corollaries of Theorem 6.1.

Let n be an even number. Then there does not exist a simplex satisfying simultaneously the condition $S \subset Q \subset nS$ and $\text{ver}(S) \subset \text{ver}(Q_n)$.

Assume the converse: there exists a simplex S such that the vertices $x^{(j)}$ coincide with vertices of Q_n and $S \subset Q_n \subset nS$. By (26),

$$c(S) = \frac{1}{n+1} \sum_{j=1}^{n+1} x^{(j)} = \left(\frac{1}{2}, \dots, \frac{1}{2} \right),$$

hence,

$$u := \sum_{j=1}^{n+1} x^{(j)} = \left(\frac{n+1}{2}, \dots, \frac{n+1}{2} \right).$$

Since $x^{(j)} \in \text{ver}(Q_n)$, coordinates of u are integer. If n is even, the latter equality cannot be valid. Thus, we have a contradiction.

Suppose $S \subset Q_n \subset nS$. Then this simplex S has the following property: while replacing any vertex of S by an arbitrary point of Q_n , the volume of the simplex does not increase.

Let us prove it. Assume $1 \leq j \leq n+1$ and $y \in Q_n$. Since

$$-\frac{n-1}{n+1} \leq \lambda_j(y) \leq 1, \quad \lambda_j(x^{(j)}) = 1,$$

we have

$$\text{dist}(y; B_j) \leq \text{dist}(x^{(j)}; B_j). \quad (27)$$

Here B_j is the $(n-1)$ -dimensional hyperplane given by the equation $\lambda_j(x) = 0$ and

$$\text{dist}(y; B_j) := \min_{z \in B_j} \|y - z\|$$

is the distance from y to B_j . Denote by S' a simplex with vertices $x^{(1)}, \dots, x^{(j-1)}, y, x^{(j+1)}, \dots, x^{(n+1)}$, i. e., the convex hull of the points $x^{(k)}$ for $k \neq j$ and the point y . From (27), it follows that $\text{vol}(S') \leq \text{vol}(S)$.

By the terminology of [7], a simplex $S \subset Q_n$, for which replacing any vertex by any point of Q_n decreases its volume, is called *rigid*. According to the above result, a simplex S satisfying the inclusions $S \subset Q_n \subset nS$, may be called *quasi-rigid*.

7 Perfect Simplices

This section is based on the authors' paper [40], see also [46].

By our definition, S is a perfect simplex with respect to a cube Q if $S \subset Q \subset \xi_n S$ and the cube Q is inscribed into the simplex $\xi_n S$, i. e., the boundary of $\xi_n S$ contains all the vertices of Q . A simplex perfect with respect to the unit cube Q_n is called simply *perfect*. Today we know the only three dimensions n when there exist perfect simplices in \mathbb{R}^n . These are $n = 1$, $n = 2$, and $n = 5$; in all the cases $\xi_n = n$.

Consider the question on existence of perfect simplices in \mathbb{R}^n for various n . We need the following proposition proved in [40].

Theorem 7.1. *Suppose $S \subset Q_n$. A vertex v of Q_n belongs to the j th face of the simplex $\xi(S)S$ when and only when*

$$-\lambda_j(v) = \max_{1 \leq k \leq n+1, x \in \text{ver}(Q_n)} (-\lambda_k(x)). \quad (28)$$

By the j th face of the simplex $\xi(S)S$, we mean the $(n-1)$ -dimensional face contained in the hyperplane $\lambda_j(x) = \text{const}$. As above, λ_j is the j th basic Lagrange polynomial related to S .

1) $n = 1$.

This case is very simple. For the segment $S = [0, 1]$, we have $S = Q_1$. Hence, $\xi_1 = 1$ and S is a unique perfect simplex. The equality $\xi_1 = 1$ also follows from Theorem 2.1 since 2 is an Hadamard number.

2) $n = 2$.

There are no perfect simplices in \mathbb{R}^2 . A unique (up to rotations) simplex S satisfying $\xi(S) = \xi_2$ is described in [26, §2.4]. Obviously, S is not perfect, since the zero-vertex of Q_2 is not contained in the boundary of $\xi(S)S$.

3) $n = 3$.

The exact value $\xi_3 = 3$ is given in [26, §2.4]. The equality $\xi(S) = 3$ holds for two (up to equivalence) tetrahedrons. Let us consider these cases.

Since for $n = 3$ the conditions of Theorem 2.1 are fulfilled, we have $\xi_3 = 3$. There exists a regular simplex S inscribed into Q_3 and satisfying $Q_3 \subset 3S$. We may take the simplex S_1 with vertices $(0, 0, 0)$, $(1, 1, 0)$, $(1, 0, 1)$, $(0, 1, 1)$. For this simplex,

$$\mathbf{A} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix}, \quad \mathbf{A}^{-1} = \frac{1}{2} \begin{pmatrix} -1 & 1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ -1 & -1 & 1 & 1 \\ 2 & 0 & 0 & 0 \end{pmatrix}.$$

Formulae (6)–(7) give $d_1(S_1) = d_2(S_1) = d_3(S_1) = 1$, $\alpha(S_1) = 3$. The basic Lagrange polynomials are

$$\begin{aligned} \lambda_1(x) &= \frac{1}{2}(-x_1 - x_2 - x_3 + 2), & \lambda_2(x) &= \frac{1}{2}(x_1 + x_2 - x_3), \\ \lambda_3(x) &= \frac{1}{2}(x_1 - x_2 + x_3), & \lambda_4(x) &= \frac{1}{2}(-x_1 + x_2 + x_3). \end{aligned}$$

Calculation shows that

$$\xi(S_1) = 4 \max_{1 \leq k \leq 4} \max_{x \in \text{ver}(Q_3)} (-\lambda_k(x)) + 1 = 4 \cdot \frac{1}{2} + 1 = 3. \quad (29)$$

For each k , maximum in (29) is delivered by the only one vertex of Q_3 :

$$-\lambda_1(1, 1, 1) = -\lambda_2(0, 0, 1) = -\lambda_3(0, 1, 0) = -\lambda_4(1, 0, 0) = \frac{1}{2}. \quad (30)$$

By Theorem 7.1 and (30), the extremal vertices of the cube, i. e., the vertices $(1, 1, 1)$, $(0, 0, 1)$, $(0, 1, 0)$, $(1, 0, 0)$, belong to the faces of the simplex $3S_1$. Moreover, each face contains a unique vertex. The rest four vertices of the cube are inside the simplex $3S_1$. Therefore, S_1 , though satisfies the inclusions $S \subset Q_3 \subset 3S_1$, is not a perfect simplex. Largest segments in S_1 parallel to the axes intersect at the center of Q_3 .

Now consider the simplex S_2 with vertices $(\frac{1}{2}, 0, 0)$, $(\frac{1}{2}, 1, 0)$, $(0, \frac{1}{2}, 1)$, $(1, \frac{1}{2}, 1)$. For S_2 ,

$$\mathbf{A} = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 1 \\ \frac{1}{2} & 1 & 0 & 1 \\ 0 & \frac{1}{2} & 1 & 1 \\ 1 & \frac{1}{2} & 1 & 1 \end{pmatrix}, \quad \mathbf{A}^{-1} = \frac{1}{2} \begin{pmatrix} 0 & 0 & -1 & 1 \\ -1 & 1 & 0 & 0 \\ -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 1 & 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix}.$$

By our formulae, $d_1(S_2) = d_2(S_2) = d_3(S_2) = 1$, $\alpha(S_2) = 3$. We have

$$\begin{aligned} \lambda_1(x) &= -x_2 - \frac{1}{2}x_3 + 1, & \lambda_2(x) &= x_2 - \frac{1}{2}x_3, \\ \lambda_3(x) &= \frac{1}{2}(-2x_1 + x_3 + 1), & \lambda_4(x) &= \frac{1}{2}(2x_1 + x_3 - 1), \\ -\lambda_1(0, 1, 1) &= -\lambda_1(1, 1, 1) = -\lambda_2(0, 0, 1) = -\lambda_2(1, 0, 1) = \\ & & & \\ -\lambda_3(1, 0, 0) &= -\lambda_3(1, 1, 0) = -\lambda_4(0, 0, 0) = -\lambda_4(0, 1, 0) = \frac{1}{2}. \end{aligned} \quad (31)$$

Hence,

$$\xi(S_2) = 4 \max_{1 \leq k \leq 4} \max_{x \in \text{ver}(Q_3)} (-\lambda_k(x)) + 1 = 4 \cdot \frac{1}{2} + 1 = 3.$$

In this case, any vertex of the cube is an extremal. By Theorem 7.1 and (31), all the vertices of Q_3 belong to the faces of $3S_2$. This means that S_2 , differently from S_1 , is a perfect simplex.

4) $n = 5$.

As we marked, $\xi_5 = 5$. Let us consider the following family of five-dimensional simplices $V = V(s, t)$. This family was introduced and studied by the authors in [40]. The vertices of V are the points $(s, 1, \frac{1}{3}, 1, 1)$, $(s, 0, \frac{1}{3}, 1, 1)$, $(s, 2 - 3t, \frac{1}{3}, 0, 1)$,

$(2 - 3s, t, 0, \frac{1}{3}, 0)$, $(0, t, 1, \frac{1}{3}, 0)$, $(1, t, 1, \frac{1}{3}, 0)$. The vertices matrix is

$$\mathbf{A} = \mathbf{A}(s, t) = \begin{pmatrix} s & 1 & \frac{1}{3} & 1 & 1 & 1 \\ s & 0 & \frac{1}{3} & 1 & 1 & 1 \\ s & 2 - 3t & \frac{1}{3} & 0 & 1 & 1 \\ 2 - 3s & t & 0 & \frac{1}{3} & 0 & 1 \\ 0 & t & 1 & \frac{1}{3} & 0 & 1 \\ 1 & t & 1 & \frac{1}{3} & 0 & 1 \end{pmatrix}.$$

For any $s, t \in \mathbb{R}$, the simplex V is nondegenerate and $\text{vol}(V) = \frac{1}{120}$. The center of gravity of the simplex coincides with the center of the cube. The matrix \mathbf{A}^{-1} has the form

$$\mathbf{A}^{-1} = \mathbf{A}^{-1}(s, t) = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & 1 \\ 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 3s - 1 & 2 - 3s \\ 2 - 3t & 3t - 1 & -1 & 0 & 0 & 0 \\ 3t - \frac{4}{3} & \frac{5}{3} - 3t & \frac{2}{3} & -\frac{2}{3} & 3s - \frac{5}{3} & \frac{4}{3} - 3s \\ -\frac{2}{3} & \frac{1}{3} & \frac{1}{3} & 1 & 2 - 3s & 3s - 2 \end{pmatrix}.$$

The inclusion $V \subset Q_5$ takes places if and only if $\frac{1}{3} \leq s \leq \frac{2}{3}$, $\frac{1}{3} \leq t \leq \frac{2}{3}$. Calculations described in [40] lead to the following conclusions.

Under the condition

$$s \in \left[\frac{4}{9}, \frac{5}{9} \right], \quad t \in \left[\frac{4}{9}, \frac{5}{9} \right], \quad (32)$$

we have the equalities $d_i(V) = 1$, $\alpha(V) = \xi(V) = 5$. Moreover, applying Theorem 7.1, we see that in this case all the vertices of the cube Q_5 belong to the boundary of the simplex $5V$. This means that *under the key condition (32) each simplex $V(s, t)$ is perfect*.

Note that if $s \notin \left[\frac{4}{9}, \frac{5}{9} \right]$ or $t \notin \left[\frac{4}{9}, \frac{5}{9} \right]$, then $\xi(V(s, t)) > 5$.

The fact that there are perfect simplices in \mathbb{R}^5 is the basic result of the authors' paper [40]. It is still unknown whether there exist perfect simplices for odd $n > 5$. Also it remains unclear whether there exist perfect simplices for at least one even n .

8 Equisecting Simplices

The notion *equisection property* was introduced by the authors in [38] and also was studied by E. Esipova and A. Ukhalov in [1]–[2].

We call $S \subset Q_n$ an *equisecting simplex* (or a *simplex having the equisecting property*), if all the $(n - 1)$ -dimensional hyperplanes containing the faces of S cut off

from the cube (outward from the simplex) the parts of equal volumes. A simplex $S \subset Q_n$ is called here *an extremal* if $\xi(S) = \xi_n$.

The equations of $(n-1)$ -dimensional hyperplanes containing the faces of S have the form $\lambda_j(x) = 0$. The j th hyperplane contains all vertices of the simplex except the j th one. Denote by v_j the volume of a convex body cut off from the cube by the j th hyperplane and not containing the simplex. Thus,

$$v_j = \text{vol}(Q_n \cap \{x \in \mathbb{R}^n : \lambda_j(x) \leq 0\}).$$

The simplex is an equisectioning if and only if $v_1 = v_2 = \dots = v_{n+1}$.

We have noticed that some extremal simplices are also equisectioning. Obviously, the extremal one-dimensional simplex $S = [0, 1]$ has this property, since the faces of S cut off from $Q_1 = [0, 1]$ the zero-length segments. The triangle with vertices $(0, 0)$, $(1, \tau)$, $(\tau, 1)$, where $\tau = \frac{3-\sqrt{5}}{2} = 0.38196601\dots$, is an equisectioning simplex on plane. M. Nevskii showed (see [26]) that this is a unique (up to rotations) extremal simplex for $n = 2$. The area of any triangle cut off from Q_2 equals $\frac{3-\sqrt{5}}{4} = 0.1909\dots$. It is easy to see that both extremal simplices for $n = 3$ described in Section 7 also are equisectioning. Therefore, for $1 \leq n \leq 3$, all the extremal simplices have the equisection property.

Note that the equisection property does not imply an extremality. One can construct equisectioning simplices being not extremal. A problem of connection between equisection and extremality is rather interesting. It was discovered that the authors' first conjecture that all the extremal simplices are also equisectioning is not true (see [38]). In particular, for $n = 5$, $n = 7$, and $n = 9$, A. Ukhalov has built extremal simplices without the equisection property. Yet it was discovered that for all extremal simplices the volumes v_j obey some regularities. Namely, the set of these values is always limited to only two quantities. Further note that equisectioning simplices do exist also for $n > 3$. In [2], it was proved that the five-dimensional extremal simplex $V = V(s, t)$ described in Section 7 has the equisection property. Let us give the corresponding formulae from [2] for the cut-off volumes:

$$v_1(t) = \begin{cases} \frac{81t-35}{1458t^2-1620t+432} + \frac{1}{2}, & -\infty < t < \frac{1}{3}, \\ -\frac{t}{2} + \frac{2}{54-81t} + \frac{4}{9}, & \frac{1}{3} < t < \frac{4}{9}, \\ \frac{1}{3}, & \frac{4}{9} < t < \frac{2}{3}, \\ \frac{t}{2} + \frac{2}{81t-36} - \frac{1}{9}, & \frac{2}{3} < t < \frac{7}{9}, \\ \frac{55-81t}{1458t^2-1620t+432} + \frac{1}{2}, & \frac{7}{9} < t < +\infty, \end{cases}$$

$$v_2(t) = \begin{cases} \frac{81t-26}{1458t^2-1296t+270} + \frac{1}{2}, & -\infty < t < \frac{2}{9}, \\ -\frac{t}{2} + \frac{2}{45-81t} + \frac{7}{18}, & \frac{2}{9} < t < \frac{1}{3}, \\ \frac{1}{3}, & \frac{1}{3} < t < \frac{5}{9}, \\ \frac{81t^2-36t+7}{162t-54}, & \frac{5}{9} < t < \frac{2}{3}, \\ \frac{46-81t}{1458t^2-1296t+270} + \frac{1}{2}, & \frac{2}{3} < t < +\infty, \end{cases}$$

$$v_3 = \frac{1}{3}, \quad v_4 = \frac{1}{3}, \quad v_5(s) = v_2(s), \quad v_6(s) = v_1(s).$$

The expressions for $v_5(s)$ and $v_6(s)$ are the same as for $v_2(t)$ and $v_1(t)$ relatively. An analysis of the expressions for $v_1(t)$ and $v_2(t)$ shows that $v_2(t) = v_1(t + \frac{1}{9})$. Therefore, the behaviour of v_j for $j = 1, 2, 5, 6$ is described, in fact, by the same function.

In Fig. 2, we present graphs of the functions $v_1(t)$ and $v_2(t)$ on the segment $[-\frac{1}{9}, \frac{10}{9}]$.




Figure 2: Graphs of the functions $v_1(t)$ and $v_2(t)$

By the above formulae, if $s, t \in [\frac{4}{9}, \frac{5}{9}]$, then $v_j = \frac{1}{3}$, $j = 1, \dots, 6$. It is for those s and t for which $V(s, t)$ is perfect, this simplex is also equisectioning.

We see that all known today perfect simplices are also equisectioning. This allows the authors to conjecture the following: *all the perfect simplices have the equisection property.*

9 Properties of $(0, 1)$ -Matrices of Order n Having Maximal Determinant

In this section, we describe the results presented in [44]. Complementary materials to this paper are given in [45].

As we mark in Section 1, if a simplex $S \subset Q_n$ has the maximum volume, then the axial diameters of S are equal to 1. For this simplex $\alpha(S) = n$. By (6), this is equivalent to the equalities $d_i(S) = 1$. These are the basic facts for our analysis.

Denote by h_n and g_n maximal determinants of $(0, 1)$ - and $(-1, 1)$ -matrices of order n respectively. Define ν_n as the maximum volume of an n -dimensional simplex contained in Q_n . We have $g_{n+1} = 2^n h_n$, $h_n = n! \nu_n$ (see [7]).

If a $(0, 1)$ -matrix of order n having maximal determinant is known, it is possible to construct a maximal volume simplex in Q_n . Let us enlarge the row set of such a determinant by the row $(0, \dots, 0)$. Then simplex S with these vertices is contained in Q_n and has the maximal possible volume. Indeed, consider for S the vertices matrix \mathbf{A} . Then

$$\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!} = \frac{h_n}{n!} = \nu_n.$$

One can also obtain nonzero vertices of an n -dimensional simplex in Q_n with maximal volume in the same way using the columns of a $(0, 1)$ -matrix with maximal determinant of order n . On the other hand, if an n -dimensional simplex $S \subset Q_n$ with the 0-vertex has the maximal possible volume, then the coordinates of its nonzero vertices, being written in rows or in columns, form $(0, 1)$ -matrices with maximal determinant of order n .

Theorem 9.1. *Suppose \mathbf{M} is an arbitrary nondegenerate $(0, 1)$ -matrix of order n . Let \mathbf{A} be the matrix of order $n + 1$ which is obtained from \mathbf{M} by adding the $(n + 1)$ th row $(0, 0, \dots, 0, 1)$ and the $(n + 1)$ th column consisting of 1's. Denote $\mathbf{A}^{-1} = (l_{i,j})$. Then for each $i = 1, \dots, n$*

$$\sum_{j=1}^{n+1} |l_{i,j}| \geq 2. \quad (33)$$

If $|\det(\mathbf{M})| = h_n$, then for all $i = 1, \dots, n$

$$\sum_{j=1}^{n+1} |l_{i,j}| = 2. \quad (34)$$

If for some i we have the strong inequality

$$\sum_{j=1}^{n+1} |l_{i,j}| > 2, \quad (35)$$

then $|\det(\mathbf{M})| < h_n$.

Proof. Consider the simplex S with the 0-vertex and the rest vertices coinciding in the coordinate form with the rows of \mathbf{M} . Obviously, \mathbf{A} is the vertices matrix of S . Since $\det(\mathbf{A}) = \det(\mathbf{M}) \neq 0$, the simplex S is nondegenerate. The inclusion $S \subset Q_n$ means that $d_i(S) \leq 1$. Being combined with (7), this gives (33).

Now assume $|\det(\mathbf{M})| = h_n$. Then we have

$$\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!} = \frac{|\det(\mathbf{M})|}{n!} = \frac{h_n}{n!} = \nu_n.$$

Consequently, S is a simplex in Q_n with the maximal possible volume. All the axial diameters $d_i(S)$ are equal to 1. Applying (7), we obtain the conditions (34). Finally, if for some i the strong inequality (35) holds then $d_i(S) < 1$. In this case, the volume of S is not maximal and $|\det(\mathbf{M})| < h_n$. \square

Note that a $(0, 1)$ -matrix \mathbf{T} of order n having maximal determinant h_n can be obtained from a $(-1, 1)$ -matrix \mathbf{U} of order $n + 1$ with maximal determinant g_{n+1} . It is sufficient to apply to \mathbf{U} a special procedure described in [39]. If at the exit will turn out $|\det(\mathbf{T})| < h_n$, then at the entrance we had $|\det(\mathbf{U})| < g_{n+1}$. Using the above theorem, in some cases, it is possible to show that the corresponding determinants are non-maximal.




Figure 3: $(-1, 1)$ -matrix matrix \mathbf{U} of order 101. White squares denote -1 's, black squares denote 1 's




Figure 4: $(0,1)$ -matrix \mathbf{T} of order 100. White squares denote 0's, black squares denote 1's

A collection of matrices having maximal known determinants of orders $1, \dots, 119$ can be found on the site [47]. Consider the matrix of order 101 constructed in 2003 by W. Orrick and B. Solomon. They don't claim that the given $(-1, 1)$ -matrix has the largest possible determinant but only state that its determinant «surpasses the previous record». In graphical form, the original $(-1, 1)$ -matrix \mathbf{U} of order 101 and the resulting $(0, 1)$ -matrix \mathbf{T} of order 100 are shown in Fig. 3–4.

Our calculations performed by the use of the Wolfram Mathematica system show that a huge in absolute value 81-digit determinant of \mathbf{T} :

$$|\det(\mathbf{T})| = 2 \cdot 3^2 \cdot 5^{97} \cdot 79 =$$

89740816577942302983638241586569761487623964058002457022666931152343750,

yet is not maximal. Applying Theorem 9.1 to $\mathbf{M} = \mathbf{T}$, we have

$$\sum_{j=1}^{101} |l_{ij}| = \begin{cases} \frac{1438}{711} = 2.0225 \dots, & i = 1, 2, 3, \\ \frac{490}{237} = 2.0675 \dots, & i = 4, \\ 2, & i = 5, \dots, 100. \end{cases}$$

For $i \leq 4$, the strict inequality (35) holds. This yields $|\det(\mathbf{T})| < h_{100}$. Consequently, $|\det(\mathbf{U})| < g_{101}$. For the corresponding simplex S , by (7),

$$d_i(S) = \begin{cases} \frac{711}{719} = 0.9888 \dots, & i = 1, 2, 3, \\ \frac{237}{245} = 0.9673 \dots, & i = 4, \\ 1, & i = 5, \dots, 100. \end{cases}$$

This means that S is not a maximum volume simplex in Q_{100} .

10 Problems for a Simplex and a Euclidean Ball

In this section, we will discuss the results of [32] for characteristics $\alpha(B_n; S)$ and $\xi(B_n; S)$ defined for a simplex and the unit Euclidean ball in \mathbb{R}^n . Replacing a cube with a ball makes many questions much more simpler. Nevertheless, geometric interpretation of general results has a certain interest also in this particular case. Moreover, we will note some additional applications of the basic Lagrange polynomials.

The *inradius* of an n -dimensional simplex S is the maximum of the radii of balls contained within S . The center of this unique maximum ball is called the *incenter* of S . The boundary of the maximum ball is a sphere that has a single common point with each $(n - 1)$ -dimensional face of S . By the *circumradius* of S we mean the minimum of the radii of balls containing S . The boundary of this unique minimal ball does not necessarily contain all the vertices of S . Namely, this is only when the center of the minimal ball lies inside the simplex.

The inradius r and the circumradius R of a simplex S satisfy the so-called *Euler inequality*

$$R \geq nr. \tag{36}$$

Equality in (36) takes place if and only if S is a regular simplex. For history, proofs, and generalizations of Euler inequality, see, e. g., [9], [56], [54].

Let $x^{(1)}, \dots, x^{(n+1)}$ be the vertices and let $\lambda_1, \dots, \lambda_{n+1}$ be the basic Lagrange polynomials of a nondegenerate simplex $S \subset \mathbb{R}^n$. Suppose Γ_j is an $(n - 1)$ -dimensional hyperplane given by the equation $\lambda_j(x) = 0$. By Σ_j we mean an $(n - 1)$ -dimensional face of S contained in Γ_j . Symbol h_j denotes the height of S conducted from the vertex $x^{(j)}$ onto Γ_j ; r denotes the inradius of S . Define σ_j as

the $(n-1)$ -measure of Σ_j . Put $\sigma := \sum_{j=1}^{n+1} \sigma_j$. Consider the vector $a_j := \{l_{1j}, \dots, l_{nj}\}$ orthogonal to Γ_j and directed into a subspace containing $x^{(j)}$. Obviously,

$$\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j} = (a_j, x) + l_{n+1,j} = (a_j, x) + \lambda_j(0).$$

Theorem 10.1. *The following equalities are true:*

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}, \quad (37)$$

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \frac{1}{h_j}, \quad (38)$$

$$\alpha(B_n; S) = \frac{1}{r}, \quad (39)$$

$$\alpha(B_n; S) = \frac{\sigma}{n \text{vol}(S)}. \quad (40)$$

Let us note some corollaries of Theorem 10.1. We have

$$\frac{1}{r} = \sum_{j=1}^{n+1} \frac{1}{h_j}.$$

For proving, it is sufficient to apply (38) and (39). It seems to be interesting that this relation (which evidently can be obtained also in a direct way) occurs to be equivalent to the general formula (2) for $\alpha(\Omega; S)$ in the case when a convex body Ω coincide with the unit ball.

The inradius r and the incenter z of S can be calculated by formulae

$$r = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}}, \quad (41)$$

$$z = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}} \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} x^{(j)}. \quad (42)$$

The tangent point of the ball $B(z; r)$ and the face Σ_k has the form

$$y^{(k)} = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}} \left[\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} x^{(j)} - \frac{1}{\left(\sum_{i=1}^n l_{ik}^2 \right)^{1/2}} (l_{1k}, \dots, l_{nk}) \right]. \quad (43)$$

Equality (41) follows immediately from (37) and (39). To obtain (42), let us remark that

$$r = \text{dist}(z; \Gamma_j) = \frac{|\lambda_j(z)|}{\|a_j\|}.$$

Since z lies inside S , each barycentric coordinate of this point $\lambda_j(z)$ is positive, i. e., $\lambda_j(z) = r\|a_j\|$. Consequently,

$$z = \sum_{j=1}^{n+1} \lambda_j(z) x^{(j)} = r \sum_{j=1}^{n+1} \|a_j\| x^{(j)}.$$

This coincides with (42). Finally, since vector $a_k = \{l_{1k}, \dots, l_{nk}\}$ is orthogonal to Σ_k and is directed from this face inside the simplex, a unique common point of $B(z; r)$ and Σ_k has the form

$$y^{(k)} = z - \frac{r}{\|a_k\|} a_k = r \left(\sum_{j=1}^{n+1} \|a_j\| x^{(j)} - \frac{1}{\|a_k\|} a_k \right).$$

The latter is equivalent to (43).

It is interesting to compare (37) with formula (9) for $\alpha(Q'_n; S)$. Since B_n is a subset of $Q'_n = [-1, 1]^n$, we have $\alpha(B_n; S) \leq \alpha(Q'_n; S)$. Analytically, this inequality also follows from the estimate

$$\left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} \leq \sum_{i=1}^n |l_{ij}|.$$

For arbitrary $x^{(0)}$ and $\varrho > 0$, the value $\alpha(B(x^{(0)}; \varrho); S)$ can be calculated with the use of Theorem 10.1 and the equality $\alpha(B(x^{(0)}; \varrho); S) = \varrho \alpha(B_n; S)$.

If $S \subset Q_n$, then $d_i(S) \leq 1$. As it was noted, (6) immediately gives $\alpha(Q_n; S) \geq n$. The equality $\alpha(Q_n; S) = n$ holds when and only when each axial diameter of S is equal to 1. The following proposition expresses the analogues of these properties for simplices contained in a ball. This theorem is equivalent to the Euler inequality (36).

Theorem 10.2. *If $S \subset B_n$, then $\alpha(B_n; S) \geq n$. The equality $\alpha(B_n; S) = n$ holds true if and only if S is a regular simplex inscribed into B_n .*

Now let us give a formula from [32] for the absorption index of a Euclidean ball by a simplex.

Theorem 10.3. *Suppose S is a nondegenerate simplex in \mathbb{R}^n , $x^{(0)} \in \mathbb{R}^n$, $\varrho > 0$. If $B(x^{(0)}; \varrho) \not\subset S$, we have*

$$\xi(B(x^{(0)}; \varrho); S) = (n+1) \max_{1 \leq j \leq n+1} \left[\varrho \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - \sum_{i=1}^n l_{ij} x_i^{(0)} - l_{n+1,j} \right] + 1. \quad (44)$$

In particular, if $B_n \not\subset S$, then

$$\xi(B_n; S) = (n+1) \max_{1 \leq j \leq n+1} \left[\left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - l_{n+1,j} \right] + 1. \quad (45)$$

Theorem 10.4. *If $S \subset B_n$, then $\xi(B_n; S) \geq n$. The equality $\xi(B_n; S) = n$ takes place if and only if S is a regular simplex inscribed into B_n .*

Theorem 10.4 follows immediately from Theorem 10.2 and the inequality $\xi(B_n; S) \geq \alpha(B_n; S)$. Therefore, $\xi_n(B_n) = n$. For proofs and commentaries, see [32].

11 Linear Interpolation on a Euclidean Ball

In the authors' paper [43] and in the paper [33] by M. Nevskii, some questions related to linear interpolation on a Euclidean ball in \mathbb{R}^n are considered. Let us describe the main results of these papers.

An interpolation projector $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is called *minimal* if $\|P\|_{B_n} = \theta_n(B_n)$. The existence of a minimal projector follows from the continuity of $\|P\|_{B_n}$ as the nodes function which can be considered on a closed bounded subset of \mathbb{R}^m , $m = n(n+1)$ given by the conditions $x^{(j)} \in B_n$, $\det \mathbf{A} \geq \varepsilon_n > 0$. It is showed that there exists a minimal projector with nodes belonging to the boundary sphere $\|x\| = 1$.

Suppose $B = B(x^{(0)}; R)$, $P : C(B) \rightarrow \Pi_1(\mathbb{R}^n)$ is an interpolation projector with nodes $x^{(j)} \in B$, $\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$ are the basic polynomials of the simplex S having the vertices $x^{(j)}$. Then

$$\|P\|_B = \max_{x \in B} \sum_{j=1}^{n+1} |\lambda_j(x)|, \quad (46)$$

see (13) for $\Omega = B$. In [43] formula (46) is supplemented with another expression for the projector norm.

Theorem 11.1. *The following equality holds:*

$$\begin{aligned} \|P\|_B &= \max_{f_j = \pm 1} \left[R \left(\sum_{i=1}^n \left(\sum_{j=1}^{n+1} f_j l_{ij} \right)^2 \right)^{1/2} + \left| \sum_{j=1}^{n+1} f_j \left(\sum_{i=1}^n l_{ij} x_i^{(0)} + l_{n+1,j} \right) \right| \right] = \\ &= \max_{f_j = \pm 1} \left[R \left(\sum_{i=1}^n \left(\sum_{j=1}^{n+1} f_j l_{ij} \right)^2 \right)^{1/2} + \left| \sum_{j=1}^{n+1} f_j \lambda_j(x^{(0)}) \right| \right]. \end{aligned} \quad (47)$$

In the case when the center of gravity $c(S)$ of the simplex is coincided with the center of the ball, formula (47) became noticeably simpler. Indeed, if $c(S) = x^{(0)}$, then

$$\|P\|_B = \max_{f_j = \pm 1} \left[R \left(\sum_{i=1}^n \left(\sum_{j=1}^{n+1} f_j l_{ij} \right)^2 \right)^{1/2} + \frac{1}{n+1} \left| \sum_{j=1}^{n+1} f_j \right| \right].$$

Let S^* be a regular simplex inscribed into the ball $B = B(x^{(0)}; R)$ and let $P^* : C(B) \rightarrow \Pi_1(\mathbb{R}^n)$ be the corresponding interpolation projector. It is easy to see that $\|P^*\|_B$ depends not on the center $x^{(0)}$, nor on radius R of the ball, nor on the choice of a regular simplex inscribed into that ball. In other words, $\|P^*\|_B$ is a function of only dimension n . For $0 \leq t \leq n+1$, consider the function

$$\psi(t) := \frac{2\sqrt{n}}{n+1} \left(t(n+1-t) \right)^{1/2} + \left| 1 - \frac{2t}{n+1} \right|.$$

By definition, put $a := \left\lfloor \frac{n+1}{2} - \frac{\sqrt{n+1}}{2} \right\rfloor$.

Theorem 11.2. *The following relations are true:*

$$\begin{aligned} \|P^*\|_B &= \max\{\psi(a), \psi(a+1)\}, \\ \sqrt{n} &\leq \|P^*\|_B \leq \sqrt{n+1}. \end{aligned} \tag{48}$$

Moreover, $\|P^*\|_B = \sqrt{n}$ only for $n = 1$; the equality $\|P^*\|_B = \sqrt{n+1}$ holds if and only if $\sqrt{n+1}$ is an integer.

Recall that $\theta_n(B)$ denotes the minimal norm $\|P\|_B$ of an interpolation projector with nodes in B . Clearly, $\theta_n(B)$ doesn't depend on the center and the radius of a ball; this is a function only of n . From (48), it follows immediately that $\theta_n(B) \geq c\sqrt{n}$. As we will see, this estimate is exact in n . Further we take $B = B_n$.

With the use of Theorem 11.2 and arguments based on (14) for $\Omega = B_n$, it was proved in [43] that

$$\theta_1(B_1) = 1, \quad \theta_2(B_2) = \frac{5}{3}, \quad \theta_3(B_3) = 2, \quad \theta_4(B_4) = \frac{11}{5}.$$

For any $n \leq 4$, the equality $\|P\|_{B_n} = \theta_n(B_n)$ takes place only for projectors corresponding to regular inscribed simplices.

Another approach to get $\|P^*\|_{B_n} = \theta_n(B_n)$, was suggested by M. Nevskii in [34]. This approach is based on some geometric conjecture for a ball and an ellipsoid that are circumscribed around a simplex. However, also on this way, the equality $\|P^*\|_{B_n} = \theta_n(B_n)$ so far is managed to be proved only for $n \leq 4$.

Let us present some illustrations, the results of numerical analysis and some comments given in [43]. Graphs of the function

$$\psi(t) = \frac{2\sqrt{n}}{n+1} \left(t(n+1-t) \right)^{1/2} + \left| 1 - \frac{2t}{n+1} \right|, \quad 0 \leq t \leq n+1,$$

for $n = 3, 4, 5, 15$ are shown in Fig. 5–8. We mark the maximum points of this function $t_- := \frac{n+1}{2} - \frac{\sqrt{n+1}}{2}$, $t_+ := \frac{n+1}{2} + \frac{\sqrt{n+1}}{2}$ and also the points $a = \left\lfloor \frac{n+1}{2} - \frac{\sqrt{n+1}}{2} \right\rfloor$ and $a+1$. One of the points a and $a+1$ maximizes $\psi(k)$ for integer $1 \leq k \leq \frac{n+1}{2}$.

In Fig. 9–10, the values $d_n = \sqrt{n+1} - \|P^*\|_{B_n}$ for $n > 23$ are presented. As we have established, $\|P^*\|_{B_n} \leq \sqrt{n+1}$, with an equality only for $n = 3, 8, 15, 24, 35, 48, 63, 80, \dots$, i. e., for n having the form $m^2 - 1$. It is at these points that $d_n = 0$, as can




Figure 5: The graph of $\psi(t)$ for $n = 3$. Here $n + 1 = 4$, $t_- = a = 1$




Figure 6: The graph of $\psi(t)$ for $n = 4$. Here $n + 1 = 5$, $a = 1$, $t_- = \frac{5-\sqrt{5}}{2}$




Figure 7: The graph of $\psi(t)$ for $n = 5$. Here $n + 1 = 5$, $a = 1$, $t_- = \frac{6-\sqrt{6}}{2}$




Figure 8: The graph of $\psi(t)$ for $n = 15$. Here $n + 1 = 16$, $t_- = a = 6$

be seen in Fig. 9. The dashed line denotes the graph of the linear interpolation spline $l(n)$ constructed by the nodes $n = m^2 - 2$, $n = m^2$ and by the values d_n in these nodes. Always $d_n \leq l(n)$; the equality comes only for $n = m^2 - 2$ and $n = m^2$. We have $l(n) \rightarrow 0$ monotonically as $n \rightarrow \infty$. Thus, the two-side estimate

$$\sqrt{n+1} - l(n) \leq \|P^*\|_{B_n} \leq \sqrt{n+1}$$

takes place. Both the right and left inequalities turn into equalities for an infinite set of n . In [43], we also give the data related to the calculation of $\|P^*\|_{B_n}$ for $1 \leq n \leq 15$, $n = 50$, $n = 100$, and $n = 1000$.




Figure 9: The numbers $d_n = \sqrt{n+1} - \|P^*\|_{B_n}$ for $23 \leq n \leq 160$

In [33], M. Nevskii developed an approach wherein the norm of an interpolation projector $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ can be estimated from below through the volume of the corresponding simplex. The essential feature of this approach is the application of the classical Legendre polynomials. On this way, it was proved that $\theta_n(B_n) \asymp \sqrt{n}$. Let us describe these results in more detail.

The standardized Legendre polynomial of degree n is the function

$$\chi_n(t) := \frac{1}{2^{n_n!}} [(t^2 - 1)^n]^{(n)}.$$

The reasons why these polynomials appeared in the range of our questions are explained in Section 1. There we also gave the estimates for minimal norms of interpolation projectors on a cube and on a ball with the participation of the function χ_n^{-1} inverse to χ_n on the half-axis $[1, +\infty)$. Let us present estimates for interpolation on a ball proved in [33]. The central result of this paper is the following theorem. Denote $\varkappa_n := \text{vol}(B_n)$.




Figure 10: The numbers $d_n = \sqrt{n+1} - \|P^*\|_{B_n}$ for $23 \leq n \leq 300$

Theorem 11.3. *Suppose $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is an arbitrary interpolation projector. Then for the corresponding simplex $S \subset B_n$ and the vertices matrix \mathbf{A} we have*

$$\|P\|_{B_n} \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\text{vol}(S)} \right) = \chi_n^{-1} \left(\frac{n! \varkappa_n}{|\det(\mathbf{A})|} \right). \quad (49)$$

Denote by σ_n the volume of a regular simplex inscribed into B_n . From (49), it follows immediately for each n

$$\theta_n(B_n) \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\sigma_n} \right). \quad (50)$$

It is known that

$$\varkappa_n = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right)}, \quad \sigma_n = \frac{1}{n!} \sqrt{n+1} \left(\frac{n+1}{n} \right)^{\frac{n}{2}},$$

$$\varkappa_{2k} = \frac{\pi^k}{k!}, \quad \varkappa_{2k+1} = \frac{2^{k+1} \pi^k}{(2k+1)!!} = \frac{2(k!)(4\pi)^k}{(2k+1)!}$$

(see, e. g., [5], [26]). Therefore, the estimate (50) can be made more concrete:

$$\theta_n(B_n) \geq \chi_n^{-1} \left(\frac{\pi^{\frac{n}{2}} n!}{\Gamma\left(\frac{n}{2} + 1\right) \sqrt{n+1} \left(\frac{n+1}{n}\right)^{\frac{n}{2}}} \right). \quad (51)$$

If $n = 2k$, then (51) is equivalent to the inequality

$$\theta_{2k}(B_{2k}) \geq \chi_{2k}^{-1} \left(\frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k} \right). \quad (52)$$

For $n = 2k + 1$, we have

$$\theta_{2k+1}(B_{2k+1}) \geq \chi_{2k+1}^{-1} \left(\frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left(\frac{2k+2}{2k+1}\right)^{\frac{2k+1}{2}}} \right). \quad (53)$$

The Stirling formula $n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{\zeta_n}{12n}}$, $0 < \zeta_n < 1$, yields

$$\sqrt{2\pi n} \left(\frac{n}{e}\right)^n < n! < \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{1}{12n}}. \quad (54)$$

It was proved in [26, п. 3.4.2] that

$$\chi_{2k}^{-1}(s) > \left(\frac{(k!)^2 s}{(2k)!}\right)^{\frac{1}{2k}}, \quad \chi_{2k+1}^{-1}(s) > \left(\frac{(k+1)!k!s}{(2k+1)!}\right)^{\frac{1}{2k+1}}. \quad (55)$$

With the use of the estimates (52)–(55), the following result was obtained in [33].

Theorem 11.4. *There exists a constant $c > 0$ not depending on n such that*

$$\theta_n(B_n) > c\sqrt{n}. \quad (56)$$

The inequality (56) takes place, e. g., with the constant

$$c = \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} = 0.2135\dots$$

From Theorems 11.2 and 11.4, we obtain $\theta_n(B_n) \asymp \sqrt{n}$. Our results also mean that any interpolation projector P^* corresponding to an inscribed regular simplex has the norm equivalent to the minimal possible. In other words, with constants not depending on n , we have $\|P^*\|_{B_n} \asymp \theta_n(B_n)$. The equality $\|P^*\|_{B_n} = \theta_n(B_n)$ remains proved only for $1 \leq n \leq 4$.

References

- [1] E. M. Esipova, Geometric characteristics of simplices having the equicutting property, In: *Sovremennye Problemy Matematiki i Informatiki. Vyp. 17*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2017, 49–61 (in Russian).
- [2] E. M. Esipova and A. Yu. Ukhalov, Properties of extremum simplices related to the cut-off volumes, In: *Sovremennye Problemy Matematiki i Informatiki. Vyp. 18*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2018, 4–17 (in Russian).
- [3] D. D. Fedulov, The numerical estimation of geometric characteristics of an n -dimensional simplex, In: *The Way to Science. Materials of the Conference. Mathematics*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2019, 36–38 (in Russian).

- [4] D. D. Fedulov and A. Yu. Ukhalov, Estimates of minimal absorption index of an n -dimensional simplex, In: *Sovremennyye Problemy Matematiki i Informatiki. Vyp. 19*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2019, 53–59 (in Russian).
- [5] G. M. Fikhtengol'ts, *The Course in Differential and Integral Calculation. Vol.3*, Fizmatlit, Moscow, 2001, 662 p. (in Russian).
- [6] M. Hall, Jr. *Combinatorial Theory*, Blaisdall Publishing Company, Waltham (Massachusetts) – Toronto – London, 1967.
- [7] M. Hudelson, V. Klee, and D. Larman, Largest j -simplices in d -cubes: some relatives of the Hadamard maximum determinant problem, *Linear Algebra Appl.*, **241–243** (2019), 519–598.
- [8] I. P. Irodova and M. V. Nevskii, Dyadic Besov spaces and other questions in approximation theory, In: *Mathematics in Yaroslavl University. Collection of Survey Articles. Dedicated to the 25th Anniversary of Mathematical Department*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2001, 115–131 (in Russian).
- [9] M. S. Klamkin and G. A. Tsifinis, Circumradius-inradius inequality for a simplex, *Mathematics Magazine*, **52**:1 (1979), 20–22.
- [10] V. S. Klimov and A. Yu. Ukhalov, *Solving the Mathematical Analysis Problems with Systems of Computer Mathematics*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2014, 96 p. (in Russian).
- [11] I. S. Kudryavcev, E. A. Ozerova, and A. Yu. Ukhalov, New estimates for the norms of minimal projectors, In: *Sovremennyye Problemy Matematiki i Informatiki. Vyp. 17*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2017, 74–81 (in Russian).
- [12] M. Lassak, Parallelotopes of maximum volume in a simplex, *Discr. Comput. Geom.*, **21** (1999), 449–462.
- [13] A. V. Lutentkov, Minimal norms of interpolation projectors, In: *Sovremennyye Problemy Matematiki i Informatiki. Vyp. 18*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2018, 51–63 (in Russian).
- [14] M. V. Nevskii, Estimates for the minimum norm of a projection in linear interpolation over the vertices of an n -dimensional cube, *Model. Anal. Inform. Sist.*, **10**:1 (2003), 9–19 (in Russian).
- [15] M. V. Nevskii, Geometric methods in the minimal projection problem, *Model. Anal. Inform. Sist.*, **13**:2 (2006), 16–29 (in Russian).

- [16] M. V. Nevskii, Inequalities for the norms of projections in interpolation over verices of of an n -dimensional cube, In: *Mathematics in Yaroslavl University. Collection of Survey Articles. Dedicated to the 30th Anniversary of Mathematical Department*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2006, 308–330 (in Russian).
- [17] M. V. Nevskii, Minimal projections and largest simplices, *Model. Anal. Inform. Sist.*, **14**:1 (2007), 3–10 (in Russian).
- [18] M. V. Nevskii, Orthogonal projection and minimal linear interpolation on an n -dimensional cube, *Model. Anal. Inform. Sist.*, **14**:3 (2007), 8–28 (in Russian).
- [19] M. V. Nevskii, Inequalities for the norms of interpolating projections, *Model. Anal. Inform. Sist.*, **15**:3 (2008), 28–37 (in Russian).
- [20] M. V. Nevskii, On a certain relation for the minimal norm of an interpolational projection, *Model. Anal. Inform. Sist.*, **16**:1 (2009), 24–43 (in Russian).
- [21] M. V. Nevskii, On a property of n -dimensional simplices, *Matem. zametki*, **87**:4 (2010), 580–593 (in Russian). English translation: *Math. Notes*, **87**:4 (2010), 543–555.
- [22] M. V. Nevskii, On geometric characteristics of an n -dimensional simplex, *Model. Anal. Inform. Sist.*, **18**:2 (2011), 52–64 (in Russian).
- [23] M. Nevskii, Properties of axial diameters of a simplex, *Discr. Comput. Geom.*, **46**:2 (2011), 301–312.
- [24] M. V. Nevskii, Geometric inequalities for interpolation projectors, In: *Mathematics in Yaroslavl University. Collection of Survey Articles. Dedicated to the 35th Anniversary of Mathematical Department*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2011, 143–154 (in Russian).
- [25] M. V. Nevskii, On the axial diameters of a convex body, *Matem. zametki*, **90**:2 (2011), 313–315 (in Russian). English translation: *Math. Notes*, **90**:2 (2011), 295–298.
- [26] M. V. Nevskii, *Geometric Estimates in Polynomial Interpolation*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2012, 218 p. (in Russian).
- [27] M. V. Nevskii, On the minimal positive homothetic image of a simplex which contains a convex body, *Matem. zametki*, **93**:3 (2013), 295–298 (in Russian). English translation: *Math. Notes*, **93**:3–4 (2011), 470–478.
- [28] M. V. Nevskii, On some problem for a simplex and a cube in \mathbb{R}^n , *Model. Anal. Inform. Sist.*, **20**:3 (2013), 77–85 (in Russian). English translation: *Aut. Control Comp. Sci.*, **48**:7 (2014), 521–527.

- [29] M. V. Nevskii, Computation of the longest segment of a given direction in a Simplex, *Fundam. Prikl. Mat.*, **18**:2 (2013), 147–152 (in Russian). English translation: *J. Math. Sci.*, **203**:6 (2014), 851–854.
- [30] M. V. Nevskii, *Geometric Estimates in Polynomial Interpolation*, Doct. Diss., Yaroslavl, 2014, 294 p. (in Russian).
- [31] M. V. Nevskii, Polynomial interpolation and geometry of convex bodies, In: *Mathematics in Yaroslavl University, Collection of Survey Articles, Dedicated to the 40th Anniversary of Mathematical Department*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2016, 162–190 (in Russian).
- [32] M. V. Nevskii, On some problems for a simplex and a ball in \mathbb{R}^n , *Model. Anal. Inform. Sist.*, **25**:6 (2018), 680–691 (in Russian). English translation: *Aut. Control Comp. Sci.*, **53**:7 (2019), 644–652.
- [33] M. V. Nevskii, Geometric estimates in interpolation on an n -dimensional ball, *Model. Anal. Inform. Sist.*, **26**:3 (2019), 441–449 (in Russian). English translation: *Aut. Control Comp. Sci.*, **54**:7 (2020), 712–718.
- [34] M. V. Nevskii, On properties of a regular simplex inscribed into a ball, *Model. Anal. Inform. Sist.*, **28**:2 (2021), 186–197 (in Russian).
- [35] M. V. Nevskii and I. V. Hlestkova, On minimal linear interpolation, In: *Sovremennyye Problemy Matematiki i Informatiki, Vyp. 9*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2008, 31–37 (in Russian).
- [36] M. V. Nevskii and A. Yu. Ukhalov, On numerical characteristics of a simplex and their estimates, *Model. Anal. Inform. Sist.*, **23**:5 (2016), 602–618 (in Russian). English translation: *Aut. Control Comp. Sci.*, **51**:7 (2017), 757–769.
- [37] M. V. Nevskii and A. Yu. Ukhalov, New estimates of numerical values related to a simplex, *Model. Anal. Inform. Sist.*, **24**:1 (2017), 94–110 (in Russian). English translation: *Aut. Control Comp. Sci.*, **51**:7 (2017), 770–782.
- [38] M. V. Nevskii and A. Yu. Ukhalov, On n -dimensional simplices satisfying inclusions $S \subset [0, 1]^n \subset nS$, *Model. Anal. Inform. Sist.*, **24**:5 (2017), 578–595 (in Russian). English translation: *Aut. Control Comp. Sci.*, **52**:7 (2018), 667–679.
- [39] M. V. Nevskii and A. Yu. Ukhalov, On minimal absorption index for an n -dimensional simplex, *Model. Anal. Inform. Sist.*, **25**:1 (2018), 140–150 (in Russian). English translation: *Aut. Control Comp. Sci.*, **52**:7 (2018), 680–687.
- [40] M. Nevskii and A. Ukhalov, Perfect Simplices in \mathbb{R}^5 , *Beitr. Algebra Geom.*, **59**:3 (2018), 501–521.

- [41] M. V. Nevskii and A. Yu. Ukhalov, On optimal interpolation by linear functions on an n -dimensional cube, *Model. Anal. Inform. Sist.*, **25**:3 (2018), 291–311 (in Russian). English translation: *Aut. Control Comp. Sci.*, **52**:7 (2018), 828–842.
- [42] M. V. Nevskii and A. Yu. Ukhalov, Some properties of 0/1-simplices, *Izv. Saratov Univ. (N. S.), Ser. Math. Mech. Inform.*, **18**:3 (2018), 305–315 (in Russian).
- [43] M. V. Nevskii and A. Yu. Ukhalov, Linear interpolation on a Euclidean ball in \mathbb{R}^n , *Model. Anal. Inform. Sist.*, **26**:2 (2019), 279–296 (in Russian). English translation: *Aut. Control Comp. Sci.*, **54**:7 (2020), 601–614.
- [44] M. Nevskii and A. Ukhalov, Properties of $(0, 1)$ -matrices of order n having maximal determinant, *Matem. zametki SVFU*, **26**:2 (2019), 109–115 (in Russian).
- [45] M. Nevskii and A. Ukhalov, Functions for checking necessary conditions for maximality of 0/1-determinant and example // doi: 10.17632/sm3x4xrb42.1 url: <http://dx.doi.org/10.17632/sm3x4xrb42.1>
Complementary materials to the paper: M. Nevskii and A. Ukhalov, Properties of $(0, 1)$ -matrices of order n having maximal determinant, *Math. Notes of NEFU*, **26**:2 (2019), 109–115.
- [46] M. V. Nevskii and A. Yu. Ukhalov, *Selected Problems in Analysis and Computational Geometry. Part 1*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2020, 98 p. (in Russian).
- [47] W. P. Orrick and B. Solomon, The Hadamard maximal determinant problem (website), <http://www.indiana.edu/~maxdet/~>.
- [48] P. R. Scott, Lattices and convex sets in space, *Quart. J. Math. Oxford Ser. (2)*, **36** (1985), 359–362.
- [49] P. R. Scott, Properties of axial diameters, *Bull. Austral. Math. Soc.*, **39** (1989), 329–333.
- [50] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society, New York, 1959, 421 p.
- [51] E. D. Udovenko, The norms of interpolation projectors, In: *The Way to Science. Materials of the Conference. Mathematics*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2019, 35–36 (in Russian).
- [52] A. Yu. Ukhalov, *Practicum in Wolfram Mathematica*, P. G. Demidov Yaroslavl State University, Yaroslavl, 2020, 40 p. (in Russian).

- [53] A. Ukhalov and E. Udovenko, Hadamard matrices of order 28 in machine readable form, *Mendeley Data*, V2, 2020. doi: 10.17632/tw66ksdfhh.2.
<https://data.mendeley.com/datasets/tw66ksdfhh/2>
- [54] A. Vince, A simplex contained in a sphere, *J. Geom.*, **89**:2 (2008), 169–178.
- [55] S. Wolfram, *Essentials of Programming in Mathematica*, Cambridge University Press, 2016, 422 p.
- [56] S. Yang and J. Wang, Improvements of n -dimensional Euler inequality, *J. Geom.*, **51** (1994), 190–195.