

# THE INCREASING FAMILIES OF SETS GENERATED BY SELF-DUAL CLUTTERS

ANDREY O. MATVEEV

ABSTRACT. Using the KKS inequalities, we establish bounds on the numbers of  $k$ -sets in the increasing families generated by self-dual clutters (i.e., clutters  $\mathcal{A}$  that coincide with the blockers  $\mathfrak{B}(\mathcal{A})$ ) on their ground set of even cardinality.

## 1. INTRODUCTION

We denote by  $E_t$ , where  $t \geq 3$ , the set of integers  $[t] := [1, \dots, t] := \{1, \dots, t\}$ . A finite collection of sets  $\mathcal{F}$  is called a *family*. The family  $\mathbf{2}^{[t]}$  of all subsets of the set  $E_t$  is called the *power set* of  $E_t$ . The *empty set* is denoted by  $\hat{0}$ , and the *empty family* containing *no sets* is denoted by  $\emptyset$ . We denote by  $|\cdot|$  the cardinalities of sets, while the numbers of sets in families are denoted by  $\#\cdot$ . Sometimes we say that the cardinality of a set  $A$  is the *size* of  $A$ .

If  $\mathcal{F}$  is a set family such that  $\emptyset \neq \mathcal{F} \neq \{\hat{0}\}$ , then the union  $V(\mathcal{F}) := \bigcup_{F \in \mathcal{F}} F$  is called the *vertex set* of  $\mathcal{F}$ . Any finite set  $S$  such that  $S \supseteq V(\mathcal{F})$ , fixed for someone's research purposes, is called the *ground set* of the family  $\mathcal{F}$ .

Given a nonempty family  $\mathcal{F} \subseteq \mathbf{2}^{[t]}$  on its ground set  $E_t$ , we let  $\mathcal{F}^c := \{F^c : F \in \mathcal{F}\}$  denote the family of their *complements*, where  $F^c := E_t - F$ . We also associate with the family  $\mathcal{F}$  the family

$$\mathcal{F}^* := \{G^c : G \in \mathbf{2}^{[t]} - \mathcal{F}\}.$$

A family of sets  $\mathcal{A}$ , such that  $\emptyset \neq \mathcal{A} \neq \{\hat{0}\}$ , is called a *nontrivial clutter* if for any indices  $i \neq j$  of members of the family  $\{A_1, \dots, A_\alpha\} =: \mathcal{A}$  we have  $A_i \not\subseteq A_j$ .

For a subset  $A \subseteq E_t$ , the *principal increasing family* of sets  $\{A\}^\nabla \subseteq \mathbf{2}^{[t]}$ , generated by the *one-member clutter*  $\{A\}$  on its *ground set*  $E_t$ , is defined by  $\{A\}^\nabla := \{B \subseteq E_t : B \supseteq A\}$ . In particular, we have  $\{\hat{0}\}^\nabla = \mathbf{2}^{[t]}$ , and  $\{E_t\}^\nabla = \{E_t\}$ .

Given a nonempty clutter  $\mathcal{A} := \{A_1, \dots, A_\alpha\} \subset \mathbf{2}^{[t]}$ , the *increasing family* of sets  $\mathcal{A}^\nabla \subseteq \mathbf{2}^{[t]}$ , generated by  $\mathcal{A}$  on its *ground set*  $E_t$ , is defined as the union  $\mathcal{A}^\nabla := \bigcup_{A \in \mathcal{A}} \{A\}^\nabla$  of the principal increasing families  $\{A\}^\nabla$ .

A subset  $B \subseteq E_t$  is a *blocking set* of a nontrivial clutter  $\mathcal{A} \subset \mathbf{2}^{[t]}$  if it holds  $|B \cap A| > 0$ , for each member  $A \in \mathcal{A}$ . The *blocker*  $\mathfrak{B}(\mathcal{A})$  of the clutter  $\mathcal{A}$  is defined to be the family of all *inclusion-minimal blocking sets* of  $\mathcal{A}$ ; see, e.g., the monographs [4, 7, 9, 10, 11, 12, 13, 15, 16, 19, 20, 21, 22, 24, 26,

31, 32, 33, 35, 36, 38]. Additional references can be found in [27, Pt 2]. The increasing family  $\mathfrak{B}(\mathcal{A})^\nabla$  is by definition the family of *all* blocking sets of the clutter  $\mathcal{A}$ .

Recall that for a nontrivial clutter  $\mathcal{A}$  on its ground set  $E_t$  we have

$$\mathfrak{B}(\mathcal{A})^\nabla = (\mathcal{A}^\nabla)^* .$$

Clutters  $\mathcal{A}$  with the property

$$\mathfrak{B}(\mathcal{A}) = \mathcal{A} ,$$

or, equivalently,

$$\mathfrak{B}(\mathcal{A})^\nabla = \mathcal{A}^\nabla ,$$

are called *self-dual* or *identically self-blocking*; see, e.g., [1, 2][4, §2.1][24, Ch. 9][26, §5.7] on such clutters.

A clutter  $\mathcal{A}$  on its ground set  $E_t$  is self-dual if and only if we have

$$(\mathcal{A}^\nabla)^* = \mathcal{A}^\nabla , \tag{1.1}$$

and although the (lack of) self-duality of clutters does not depend structurally on the cardinalities of their vertex sets and ground sets, relation (1.1) suggests the following criterion:

**Remark 1.1.** (see [26, Cor. 5.28(i)]) *A clutter  $\mathcal{A} \subset \mathbf{2}^{[t]}$  on its ground set  $E_t$  is self-dual if and only if*

$$\#\mathcal{A}^\nabla = 2^{t-1} . \tag{1.2}$$

In Theorem 4.1 we consider condition (1.2) from a perspective of constraints that are satisfied by the numbers of  $k$ -sets in the increasing families  $\mathcal{A}^\nabla$  generated by self-dual clutters  $\mathcal{A} \subset \mathbf{2}^{[t]}$  on their ground set  $E_t$  of even cardinality  $t$ .

## 2. LONG $f$ - AND $h$ -VECTORS OF SET FAMILIES

Let us associate with each family  $\mathcal{F} \subseteq \mathbf{2}^{[t]}$  on the ground set  $E_t$  its *long  $f$ -vector*

$$\mathbf{f}(\mathcal{F}; t) := (f_0(\mathcal{F}; t), f_1(\mathcal{F}; t), \dots, f_t(\mathcal{F}; t)) \in \mathbb{N}^{t+1} ,$$

where

$$f_k(\mathcal{F}; t) := \#\{F \in \mathcal{F} : |F| = k\} , \quad 0 \leq k \leq t .$$

If ‘ $\mathbf{x}$ ’ is a formal variable, then the *long  $h$ -vector*

$$\mathbf{h}(\mathcal{F}; t) := (h_0(\mathcal{F}; t), h_1(\mathcal{F}; t), \dots, h_t(\mathcal{F}; t)) \in \mathbb{Z}^{t+1}$$

of the family  $\mathcal{F}$  is defined by means of the relation

$$\sum_{i=0}^t h_i(\mathcal{F}; t) \cdot \mathbf{x}^{t-i} := \sum_{i=0}^t f_i(\mathcal{F}; t) \cdot (\mathbf{x} - 1)^{t-i} .$$

**Example 2.1.** (cf. Example 3.1) For any element  $a \in E_t$ , the principal increasing family  $\{\{a\}\}^\nabla \subset \mathbf{2}^{[t]}$ , generated by the self-dual clutter  $\{\{a\}\}$  on its ground set  $E_t$ , is described by the vectors

$$\mathbf{f}(\{\{a\}\}^\nabla; t) = (0, \binom{t-1}{1-1}, \binom{t-1}{2-1}, \dots, \binom{t-1}{t-1})$$

and

$$\mathbf{h}(\{\{a\}\}^\nabla; t) = (0, 1, 0, \dots, 0).$$

**Example 2.2.** (i) (a) Given the self-dual clutter

$$\mathcal{A} := \{\{1, 2\}, \{1, 3\}, \{2, 3\}\} \quad (2.1)$$

that generates on its ground set  $E_3 = V(\mathcal{A})$  the increasing family

$$\mathcal{A}^\nabla = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, E_3\},$$

we have

$$\begin{aligned} \mathbf{f}(\mathcal{A}^\nabla; 3) &= (0, 0, 3, 1), \\ \mathbf{h}(\mathcal{A}^\nabla; 3) &= (0, 0, 3, -2). \end{aligned}$$

(b) If we choose the set  $E_4 \supsetneq V(\mathcal{A})$  as the ground set of the self-dual clutter (2.1), instead of the set  $E_3$ , then  $\mathcal{A}$  generates on  $E_4$  the increasing family

$$\mathcal{A}^\nabla = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}, \{2, 3, 4\}, E_4\},$$

and we have

$$\begin{aligned} \mathbf{f}(\mathcal{A}^\nabla; 4) &= (0, 0, 3, 4, 1), \\ \mathbf{h}(\mathcal{A}^\nabla; 4) &= (0, 0, 3, -2, 0). \end{aligned}$$

(ii) (a) For the self-dual clutter

$$\mathcal{A} := \{\{2\}\} \quad (2.2)$$

that generates on its ground set  $E_3 \supsetneq V(\mathcal{A})$  the principal increasing family

$$\mathcal{A}^\nabla = \{\{2\}, \{1, 2\}, \{2, 3\}, E_3\},$$

we have

$$\begin{aligned} \mathbf{f}(\mathcal{A}^\nabla; 3) &= (0, 1, 2, 1), \\ \mathbf{h}(\mathcal{A}^\nabla; 3) &= (0, 1, 0, 0). \end{aligned}$$

(b) For the self-dual clutter (2.2) that generates on its ground set  $E_4 \supsetneq V(\mathcal{A})$  the principal increasing family

$$\mathcal{A}^\nabla = \{\{2\}, \{1, 2\}, \{2, 3\}, \{2, 4\}, \{1, 2, 3\}, \{1, 2, 4\}, \{2, 3, 4\}, E_4\},$$

we have

$$\begin{aligned} \mathbf{f}(\mathcal{A}^\nabla; 4) &= (0, 1, 3, 3, 1), \\ \mathbf{h}(\mathcal{A}^\nabla; 4) &= (0, 1, 0, 0, 0). \end{aligned}$$

(iii) *The self-dual clutter*

$$\mathcal{A} := \{\{1, 2, 3, 4\}, \{1, 5\}, \{2, 5\}, \{3, 5\}, \{4, 5\}\},$$

that generates on its ground set  $E_5 = V(\mathcal{A})$  the increasing family

$$\begin{aligned} \mathcal{A}^\nabla = \{ & \{1, 5\}, \{2, 5\}, \{3, 5\}, \{4, 5\}, \{1, 2, 5\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 5\}, \\ & \{2, 4, 5\}, \{3, 4, 5\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 4, 5\}, \\ & \{1, 3, 4, 5\}, \{2, 3, 4, 5\}, E_5 \}, \end{aligned}$$

is described by the vectors

$$\begin{aligned} \mathbf{f}(\mathcal{A}^\nabla; 5) &= (0, 0, 4, 6, 5, 1), \\ \mathbf{h}(\mathcal{A}^\nabla; 5) &= (0, 0, 4, -6, 5, -2). \end{aligned}$$

**Remark 2.3.** Given a family  $\mathcal{F} \subseteq \mathbf{2}^{[t]}$  on its ground set  $E_t$ , we have:

(i) (see [26, Prop. 2.1(iii)(a)])

$$h_\ell(\mathcal{F}; t) = (-1)^\ell \sum_{k=0}^{\ell} (-1)^k \binom{t-k}{t-\ell} f_k(\mathcal{F}; t), \quad 0 \leq \ell \leq t; \quad (2.3)$$

$$f_\ell(\mathcal{F}; t) = \sum_{k=0}^{\ell} \binom{t-k}{t-\ell} h_k(\mathcal{F}; t), \quad 0 \leq \ell \leq t. \quad (2.4)$$

(ii) (see [26, Prop. 2.1(iii)(b)])

$$\begin{aligned} h_0(\mathcal{F}; t) &= f_0(\mathcal{F}; t); \\ h_1(\mathcal{F}; t) &= f_1(\mathcal{F}; t) - t f_0(\mathcal{F}; t); \\ h_{t-1}(\mathcal{F}; t) &= (-1)^{t-1} \sum_{k=0}^{t-1} (-1)^k (t-k) f_k(\mathcal{F}; t); \\ h_t(\mathcal{F}; t) &= (-1)^t \sum_{k=0}^t (-1)^k f_k(\mathcal{F}; t); \end{aligned} \quad (2.5)$$

$$\sum_{k=0}^t h_k(\mathcal{F}; t) = f_t(\mathcal{F}; t).$$

(iii) (see [26, Prop. 2.1(iii)(c)])

$$\sum_{k=0}^t f_k(\mathcal{F}; t) = \sum_{k=0}^t 2^{t-k} h_k(\mathcal{F}; t) = \#\mathcal{F}.$$

(iv)

$$\mathbf{h}(\mathcal{F}; t) + \mathbf{h}(\mathbf{2}^{[t]} - \mathcal{F}; t) = (1, 0, 0, \dots, 0) = \mathbf{h}(\mathbf{2}^{[t]}; t).$$

Being *redundant* analogues of *standard*  $h$ -vectors of abstract simplicial complexes [8, 23, 28, 34, 37, 40], *long*  $h$ -vectors (see, e.g., [29, p. 170][30, p. 265]) demonstrate their usefulness below in relations (2.6), where they describe a curious enumerative connection between the families  $\mathcal{F}$  and  $\mathcal{F}^*$ .

**Remark 2.4.** Given a family  $\mathcal{F} \subseteq \mathbf{2}^{[t]}$  on its ground set  $E_t$ , we have:

(i)

$$\#\mathcal{F}^* + \#\mathcal{F} = 2^t .$$

More precisely,

$$f_\ell(\mathcal{F}^*; t) + f_{t-\ell}(\mathcal{F}; t) = \binom{t}{\ell} , \quad 0 \leq \ell \leq t .$$

(ii) (see [26, Prop. 2.1(iii)(d)])

$$h_\ell(\mathcal{F}^*; t) + (-1)^\ell \sum_{k=\ell}^t \binom{k}{\ell} h_k(\mathcal{F}; t) = \delta_{\ell,0} , \quad 0 \leq \ell \leq t , \quad (2.6)$$

where  $\delta_{\ell,0} := 1$  if  $\ell = 0$ , and  $\delta_{\ell,0} := 0$  otherwise. Note that

$$h_t(\mathcal{F}^*; t) = (-1)^{t+1} h_t(\mathcal{F}; t) .$$

### 3. ABSTRACT SIMPLICIAL COMPLEXES $\Delta$ , SUCH THAT $\Delta^* = \Delta$ , ON THEIR GROUND SETS $E_t$ OF EVEN CARDINALITY $t$

Although in the present paper we are concerned with enumerative properties of specific increasing families of sets, in this section we turn to specific *abstract simplicial complexes* (i.e., “*decreasing*” families of “*faces*”) because the numbers of faces of size  $k$  in complexes are known to satisfy the classical *Kruskal–Katona–Schützenberger (KKS) inequalities*; see, e.g., [34, §10.3] and [3, 4, 6, 8, 14, 17, 18, 23, 24, 25, 37, 38, 39, 40].

Let  $\Delta \subseteq \mathbf{2}^{[t]}$ , where  $\emptyset \neq \Delta \neq \{\hat{0}\}$ , be an abstract simplicial complex on its ground set  $E_t$ , with its *vertex set*  $V(\Delta) \subseteq E_t$ . By definition of complex, the following implications hold:  $(B \in \Delta, A \subset B) \implies A \in \Delta$ . The *inclusion-maximal* faces of  $\Delta$  are called its *facets*. Sometimes one says that a complex with *one* facet (i.e., the power set of a set) is a *simplex*. The long  $f$ -vector of the complex  $\Delta$  has the form

$$\mathbf{f}(\Delta; t) := \left( 1, \underbrace{f_1(\Delta; t)}_{:=|V(\Delta)|>0}, \dots, \underbrace{f_{d(\Delta)}(\Delta; t)}_{>0}, 0, \dots, 0 \right) ,$$

where  $d(\Delta) := \max\{k \in [t]: f_k(\Delta; t) > 0\}$ .

**Example 3.1.** (cf. Example 2.1) For any element  $a \in E_t$ , the simplex  $\Delta := \{F: F \subseteq (E_t - \{a\})\} = \Delta^*$ , whose facet is the subset  $(E_t - \{a\}) \subset E_t$  of size  $(t-1)$ , is described by the vectors

$$\mathbf{f}(\Delta; t) = \left( \binom{t-1}{0}, \dots, \binom{t-1}{t-2}, \binom{t-1}{t-1}, 0 \right)$$

and

$$\mathbf{h}(\Delta; t) = (1, -1, 0, \dots, 0) .$$

**Remark 3.2.** An abstract simplicial complex  $\Delta$  with vertex set  $V(\Delta)$ , such that  $\Delta$  coincides with its Alexander dual complex  $\Delta^\vee$ , defined by

$$\Delta^\vee := \{V(\Delta) - F : F \in \mathbf{2}^{V(\Delta)} - \Delta\}$$

when  $V(\Delta^\vee) = V(\Delta)$ , is known as an Alexander self-dual complex. Note that

$$\Delta = \Delta^\vee \iff \#\Delta = 2^{|V(\Delta)|-1}.$$

See, e.g., [38] and [5] on *combinatorial Alexander duality*.

**Remark 3.3.** Suppose that  $\Delta \subset \mathbf{2}^{[t]}$  is an abstract simplicial complex, such that  $\Delta^* = \Delta$ , on its ground set  $E_t$ . Then we have:

$$(i) \quad \#\Delta = \sum_{k=0}^t f_k(\Delta; t) = \sum_{k=0}^{d(\Delta)} f_k(\Delta; t) = \sum_{k=0}^t 2^{t-k} h_k(\Delta; t) = 2^{t-1}.$$

$$(ii) \quad f_\ell(\Delta; t) + f_{t-\ell}(\Delta; t) = \binom{t}{\ell}, \quad 0 \leq \ell \leq t. \quad (3.1)$$

In particular, if  $t$  is even, then

$$f_{t/2}(\Delta; t) = \frac{1}{2} \binom{t}{t/2}.$$

**Example 3.4.** (i) The abstract simplicial complex

$$\Delta := \{\hat{0}, \{1\}, \{2\}, \{3\}, \{4\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\},$$

such that  $\Delta^* = \Delta$ , on its ground set  $E_4 = V(\Delta)$ , is described by the vectors

$$\mathbf{f}(\Delta; 4) = (1, 4, 3, 0, 0),$$

$$\mathbf{h}(\Delta; 4) = (1, 0, -3, 2, 0).$$

The complex  $\Delta$  is an Alexander self-dual complex, that is,  $\Delta = \Delta^\vee$ , because  $8 = \#\Delta = 2^{|V(\Delta)|-1} = 2^{4-1}$ .

(ii) The simplex

$$\Delta := \{\hat{0}, \{2\}, \{3\}, \{4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{2, 3, 4\}\},$$

such that  $\Delta^* = \Delta$ , on its ground set  $E_4 \supsetneq V(\Delta)$ , is described by the vectors

$$\mathbf{f}(\Delta; 4) = (1, 3, 3, 1, 0),$$

$$\mathbf{h}(\Delta; 4) = (1, -1, 0, 0, 0).$$

• Suppose again that  $\Delta \subset \mathbf{2}^{[t]}$  is an abstract simplicial complex, such that  $\Delta^* = \Delta$ , on its ground set  $E_t$  of *even* cardinality  $t$ . We would like to apply the KKS inequalities to the complex  $\Delta$ .

◦ Since  $\frac{1}{2} \binom{t}{t/2} = \binom{t-1}{t/2}$ , we have the simplest  $(t/2)$ -binomial expansion

$$f_{t/2}(\Delta; t) = \binom{\alpha_{t/2}(t/2)}{t/2}$$

of the middle component of the long  $f$ -vector  $\mathbf{f}(\Delta; t)$ , where

$$\alpha_{t/2}(t/2) = t - 1.$$

Then relations (3.1) and the KKS inequalities (lower and upper bound versions) [34, Cor. 10.1, Cor. 10.2] imply that

$$\begin{aligned} \binom{t}{(t/2)-1} - f_{(t/2)+1}(\Delta; t) &= f_{(t/2)-1}(\Delta; t) \geq \binom{\alpha_{t/2}(t/2)}{(t/2)-1} = \binom{t-1}{(t/2)-1} = \binom{t-1}{t/2}, \\ \binom{t}{(t/2)+1} - f_{(t/2)-1}(\Delta; t) &= f_{(t/2)+1}(\Delta; t) \leq \binom{\alpha_{t/2}(t/2)}{(t/2)+1} = \binom{t-1}{(t/2)+1} = \binom{t-1}{(t/2)-2}. \end{aligned}$$

Further, note that if our complex  $\Delta$  had the minimum possible number of its faces of size  $((t/2)-1)$ , namely  $\binom{t-1}{(t/2)-1}$  faces, then the following implication would hold:

$$\begin{aligned} f_{(t/2)-1}(\Delta; t) &:= \binom{t-1}{(t/2)-1} = \binom{t-1}{t/2} \\ \implies f_{(t/2)+1}(\Delta; t) &= \binom{t}{(t/2)+1} - \binom{t-1}{(t/2)-1} \\ &= \binom{t}{(t/2)-1} - \binom{t-1}{(t/2)-1} = \binom{t-1}{(t/2)-2}. \end{aligned}$$

Note also that  $\binom{t}{(t/2)-1} - \binom{t-1}{(t/2)-1} = \binom{t-1}{(t/2)-2} = \binom{t-1}{(t/2)+1}$ .

◦ If  $((t/2) - 1) > 1$ , then suppose that our complex  $\Delta$  indeed has the *minimum possible* number of faces of size  $((t/2) - 1)$ , that is,  $f_{(t/2)-1}(\Delta; t) := \binom{t-1}{(t/2)-1}$ . In this case we have the  $((t/2) - 1)$ -binomial expansion

$$f_{(t/2)-1}(\Delta; t) = \binom{\alpha_{(t/2)-1}((t/2)-1)}{(t/2)-1}$$

of the component  $f_{(t/2)-1}(\Delta; t)$  of the vector  $\mathbf{f}(\Delta; t)$ , where

$$\alpha_{(t/2)-1}((t/2) - 1) = t - 1.$$

Relations (3.1) and the KKS inequalities (lower bound version) [34, Cor. 10.1] imply that

$$\binom{t}{(t/2)-2} - f_{(t/2)+2}(\Delta; t) = f_{(t/2)-2}(\Delta; t) \geq \binom{\alpha_{(t/2)-1}((t/2)-1)}{(t/2)-2} = \binom{t-1}{(t/2)-2}.$$

On the other hand, suppose that the complex  $\Delta$  has the *maximum possible* number of faces of size  $((t/2) + 1)$ , that is,  $f_{(t/2)+1}(\Delta; t) := \binom{t-1}{(t/2)+1}$ . We have the  $((t/2) + 1)$ -binomial expansion

$$f_{(t/2)+1}(\Delta; t) = \binom{\alpha_{(t/2)+1}((t/2)+1)}{(t/2)+1}$$

of the component  $f_{(t/2)+1}(\Delta; t)$  of the vector  $\mathbf{f}(\Delta; t)$ , where

$$\alpha_{(t/2)+1}((t/2) + 1) = t - 1.$$

Now relations (3.1) and the KKS inequalities (upper bound version) [34, Cor. 10.2] imply that

$$\binom{t}{(t/2)+2} - f_{(t/2)-2}(\Delta; t) = f_{(t/2)+2}(\Delta; t) \leq \binom{\alpha_{(t/2)+1}((t/2)+1)}{(t/2)+2} = \binom{t-1}{(t/2)+2}.$$

Note that  $\binom{t}{(t/2)-2} - \binom{t-1}{(t/2)-2} = \binom{t-1}{(t/2)-3} = \binom{t-1}{(t/2)+2}$ .

◦ Proceeding by induction, we arrive at the following counterpart of Theorem 4.1:

**Lemma 3.5.** (see also Example 3.1) Let  $\Delta \subset \mathbf{2}^{[t]}$  be an abstract simplicial complex on its ground set  $E_t$  of even cardinality  $t$ . If

$$\Delta^* = \Delta ,$$

then we have

$$\begin{aligned} f_0(\Delta; t) &= 1 , \quad \text{and} \quad f_t(\Delta; t) = 0 ; \\ f_{t/2}(\Delta; t) &= \frac{1}{2} \binom{t}{t/2} = \binom{t-1}{t/2} ; \\ f_{(t/2)-i}(\Delta; t) &\geq \binom{t-1}{(t/2)-i} = \binom{t-1}{(t/2)+i-1} , \quad 1 \leq i \leq (t/2) - 1 ; \\ f_{(t/2)+j}(\Delta; t) &\leq \binom{t-1}{(t/2)+j} = \binom{t-1}{(t/2)-j-1} , \quad 1 \leq j \leq (t/2) - 1 ; \\ f_{(t/2)-k}(\Delta; t) + f_{(t/2)+k}(\Delta; t) &= \binom{t}{(t/2)-k} = \binom{t}{(t/2)+k} , \quad 1 \leq k \leq (t/2) - 1 . \end{aligned}$$

#### 4. THE INCREASING FAMILIES OF SETS GENERATED BY SELF-DUAL CLUTTERS ON THEIR GROUND SETS $E_t$ OF EVEN CARDINALITY $t$

Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family on its ground set  $E_t$ . The implication

$$\mathcal{F}^* = \mathcal{F} \quad \Longleftrightarrow \quad (\mathbf{2}^{[t]} - \mathcal{F})^* = \mathbf{2}^{[t]} - \mathcal{F} ,$$

allows us to obtain from Lemma 3.5 the following result:

**Theorem 4.1.** (see also Example 2.1) Let  $\mathcal{A} \subset \mathbf{2}^{[t]}$  be a clutter on its ground set  $E_t$  of even cardinality  $t$ . If the clutter  $\mathcal{A}$  is self-dual, that is,

$$\mathfrak{B}(\mathcal{A}) = \mathcal{A} ,$$

then we have

$$\begin{aligned} f_0(\mathcal{A}^\nabla; t) &= 0 , \quad \text{and} \quad f_t(\mathcal{A}^\nabla; t) = 1 ; \\ f_{t/2}(\mathcal{A}^\nabla; t) &= \frac{1}{2} \binom{t}{t/2} = \binom{t-1}{t/2} ; \\ f_{(t/2)-i}(\mathcal{A}^\nabla; t) &\leq \binom{t-1}{(t/2)-i-1} = \binom{t-1}{(t/2)+i} , \quad 1 \leq i \leq (t/2) - 1 ; \\ f_{(t/2)+j}(\mathcal{A}^\nabla; t) &\geq \binom{t-1}{(t/2)+j-1} = \binom{t-1}{(t/2)-j} , \quad 1 \leq j \leq (t/2) - 1 ; \\ f_{(t/2)-k}(\mathcal{A}^\nabla; t) + f_{(t/2)+k}(\mathcal{A}^\nabla; t) &= \binom{t}{(t/2)-k} = \binom{t}{(t/2)+k} , \quad 1 \leq k \leq (t/2) - 1 . \end{aligned}$$

#### 5. APPENDIX: REMARKS ON SET FAMILIES $\mathcal{F}$ SUCH THAT $\mathcal{F}^* = \mathcal{F}$

**Remark 5.1.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$ .

(i) We have

$$\#\mathcal{F} = 2^{t-1} .$$

More precisely,

$$f_\ell(\mathcal{F}; t) + f_{t-\ell}(\mathcal{F}; t) = \binom{t}{\ell} , \quad 0 \leq \ell \leq t . \quad (5.1)$$

(ii) Relations (5.1) and (2.4) imply that

$$\sum_{k=0}^{\ell} \binom{t-k}{t-\ell} h_k(\mathcal{F}; t) + \sum_{j=0}^{t-\ell} \binom{t-j}{\ell} h_j(\mathcal{F}; t) = \binom{t}{\ell} , \quad 0 \leq \ell \leq t .$$



Given a family  $\mathcal{F} \subset \mathbf{2}^{[t]}$ , such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$ , from (2.3) we have for any  $\ell$ , where  $0 \leq \ell \leq t$ :

$$\begin{aligned} h_\ell(\mathcal{F}; t) &= (-1)^\ell \sum_{k=0}^{\ell} (-1)^k \binom{t-k}{t-\ell} f_k(\mathcal{F}; t) \\ &= (-1)^\ell \sum_{k=0}^{\ell} (-1)^k \binom{t-k}{t-\ell} \left( \binom{t}{k} - f_{t-k}(\mathcal{F}; t) \right). \end{aligned}$$

**Remark 5.2.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$ .

(i) We have

$$h_\ell(\mathcal{F}; t) = \delta_{\ell,0} - (-1)^{t-\ell} \sum_{j=t-\ell}^t (-1)^j \binom{j}{t-\ell} f_j(\mathcal{F}; t), \quad 0 \leq \ell \leq t; \quad (5.2)$$

$$f_\ell(\mathcal{F}; t) = \binom{t}{\ell} - \sum_{j=0}^{t-\ell} \binom{t-j}{\ell} h_j(\mathcal{F}; t), \quad 0 \leq \ell \leq t. \quad (5.3)$$

(ii) Relations (5.2) and (2.3) yield

$$\begin{aligned} &(-1)^\ell \sum_{k=0}^{\ell} (-1)^k \binom{t-k}{t-\ell} f_k(\mathcal{F}; t) \\ &+ (-1)^{t-\ell} \sum_{j=t-\ell}^t (-1)^j \binom{j}{t-\ell} f_j(\mathcal{F}; t) = \delta_{\ell,0}, \quad 0 \leq \ell \leq t. \end{aligned}$$

**Remark 5.3.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$  of odd cardinality  $t$ . From (2.5) we have

$$\begin{aligned} h_t(\mathcal{F}; t) &= (-1)^t \sum_{k=0}^t (-1)^k f_k(\mathcal{F}; t) \\ &= - \sum_{k=0}^{\lfloor t/2 \rfloor} (-1)^k \left( f_k(\mathcal{F}; t) - \left( \binom{t}{k} - f_k(\mathcal{F}; t) \right) \right) \\ &= \binom{t-1}{(t-1)/2} - 2 \sum_{k=0}^{(t-1)/2} (-1)^k f_k(\mathcal{F}; t), \end{aligned}$$

and

$$\begin{aligned}
h_t(\mathcal{F}; t) &= (-1)^t \sum_{j=0}^t (-1)^j f_j(\mathcal{F}; t) \\
&= - \sum_{j=\lceil t/2 \rceil}^t (-1)^j \left( f_j(\mathcal{F}; t) - \binom{t}{j} f_j(\mathcal{F}; t) \right) \\
&= - \binom{t-1}{(t-1)/2} - 2 \sum_{j=(t+1)/2}^t (-1)^j f_j(\mathcal{F}; t) .
\end{aligned}$$

As a consequence, we see that

$$\sum_{k=0}^{(t-1)/2} (-1)^k f_k(\mathcal{F}; t) = \binom{t-1}{(t-1)/2} + \sum_{j=(t+1)/2}^t (-1)^j f_j(\mathcal{F}; t) .$$

**Remark 5.4.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$ .

(i) From (2.6) we have

$$h_\ell(\mathcal{F}; t) = \delta_{\ell,0} + (-1)^{\ell+1} \sum_{k=\ell}^t \binom{k}{\ell} h_k(\mathcal{F}; t) , \quad 0 \leq \ell \leq t . \quad (5.4)$$

If  $t$  is even, then we see that

$$h_t(\mathcal{F}; t) = 0 . \quad (5.5)$$

(ii) For even indices  $\ell$ , where  $2 \leq \ell \leq t$ , relations (5.4) imply that

$$\sum_{k=\ell}^t \binom{k}{\ell-1} h_k(\mathcal{F}; t) = 0 , \quad (5.6)$$

that is,

$$\ell h_\ell(\mathcal{F}; t) + \sum_{k=\ell+1}^t \binom{k}{\ell-1} h_k(\mathcal{F}; t) = 0 ;$$

we also have

$$2h_\ell(\mathcal{F}; t) + \sum_{k=\ell+1}^t \binom{k}{\ell} h_k(\mathcal{F}; t) = 0 .$$

If  $t$  is odd, then we have

$$(t-1)h_{t-1}(\mathcal{F}; t) + \binom{t}{2} h_t(\mathcal{F}; t) = 0 ,$$

that is,

$$2h_{t-1}(\mathcal{F}; t) + t h_t(\mathcal{F}; t) = 0 .$$

If  $t$  is even, then we have

$$\frac{t}{2} h_{t/2}(\mathcal{F}; t) + \sum_{k=(t/2)+1}^{t-1} \binom{k}{(t/2)-1} h_k(\mathcal{F}; t) = 0 ,$$

and

$$h_{t/2}(\mathcal{F}; t) + \frac{1}{2} \sum_{k=(t/2)+1}^{t-1} \binom{k}{t/2} h_k(\mathcal{F}; t) = 0 . \quad (5.7)$$

**Remark 5.5.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$ .

If  $t$  is even, then relations (5.5) and (5.6) in the case  $\ell := 2$  yield

$$h_t(\mathcal{F}; t) = 0 , \quad \text{and} \quad \sum_{k=2}^{t-1} k h_k(\mathcal{F}; t) = 0 .$$

Similarly, if  $t$  is odd, then from (5.6) we have

$$\sum_{k=2}^t k h_k(\mathcal{F}; t) = 0 .$$

**Remark 5.6.** Let  $\mathcal{F} \subset \mathbf{2}^{[t]}$  be a family, such that  $\mathcal{F}^* = \mathcal{F}$ , on its ground set  $E_t$  of even cardinality  $t$ .

(i) On the one hand, since

$$f_{t/2}(\mathcal{F}; t) = \frac{1}{2} \binom{t}{t/2} , \quad (5.8)$$

relations (2.4) imply that

$$\begin{aligned} & \sum_{k=0}^{t/2} \binom{t-k}{t/2} h_k(\mathcal{F}; t) = f_{t/2}(\mathcal{F}; t) \\ & = h_{t/2}(\mathcal{F}; t) + \sum_{k=0}^{(t/2)-1} \binom{t-k}{t/2} h_k(\mathcal{F}; t) = \frac{1}{2} \binom{t}{t/2} . \end{aligned} \quad (5.9)$$

On the other hand, (5.8) and relations (5.3) imply that

$$\binom{t}{t/2} - \sum_{j=0}^{t/2} \binom{t-j}{t/2} h_{t/2}(\mathcal{F}; t) = f_{t/2}(\mathcal{F}; t) = \frac{1}{2} \binom{t}{t/2} ,$$

that is,

$$\sum_{j=0}^{t/2} \binom{t-j}{t/2} h_{t/2}(\mathcal{F}; t) = \frac{1}{2} \binom{t}{t/2} .$$

(ii) Relations (5.7) and (5.9) imply that

$$\sum_{k=0}^{(t/2)-1} \binom{t-k}{t/2} h_k(\mathcal{F}; t) - \frac{1}{2} \sum_{j=(t/2)+1}^{t-1} \binom{j}{t/2} h_j(\mathcal{F}; t) = \frac{1}{2} \binom{t}{t/2},$$

and

$$h_{t/2}(\mathcal{F}; t) = \frac{1}{4} \binom{t}{t/2} - \frac{1}{2} \sum_{k=0}^{(t/2)-1} \binom{t-k}{t/2} h_k(\mathcal{F}; t) - \frac{1}{4} \sum_{j=(t/2)+1}^{t-1} \binom{j}{t/2} h_j(\mathcal{F}; t).$$

## REFERENCES

- [1] *Abdi A., Cornuéjols G., Lee D.* Identically self-blocking clutters. 20th International Conference Integer Programming and Combinatorial Optimization IPCO 2019 held in Ann Arbor, MI, USA, May 22–24, 2019. *A. Lodi and V. Nagarajan* (eds.). Lecture Notes in Computer Science, 11480. Cham: Springer, 2019, pp. 1–12.
- [2] *Abdi A., Pashkovich K.* Delta minors, delta free clutters, and entanglement. *SIAM Journal on Discrete Mathematics*, 2018, 32, no. 3, pp. 1750–1774.
- [3] *Anderson I.* Combinatorics of finite sets. Corrected reprint of the 1989 edition. Mineola, NY: Dover Publications, Inc., 2002.
- [4] *Berge C.* Hypergraphs. Combinatorics of finite sets. Translated from the French. North-Holland Mathematical Library, 45. Amsterdam: North-Holland Publishing Co., 1989.
- [5] *Björner A., Tancer M.* Note: Combinatorial Alexander duality—A short and elementary proof. *Discrete and Computational Geometry*, 2009, 42, no. 4, pp. 586–593.
- [6] *Bollobás B.* Combinatorics. Set systems, hypergraphs, families of vectors and combinatorial probability. Cambridge: Cambridge University Press, 1986.
- [7] *Bretto A.* Hypergraph theory. An introduction. Mathematical Engineering. Cham: Springer, 2013.
- [8] *Bruns W., Herzog J.* Cohen–Macaulay rings. Second edition. Cambridge Studies in Advanced Mathematics, 39. Cambridge: Cambridge University Press, 1998.
- [9] *Conforti M., Cornuéjols G., Zambelli G.* Integer programming. Cham: Springer, 2014.
- [10] *Cornuéjols G.* Combinatorial optimization. Packing and covering. CBMS–NSF Regional Conference Series in Applied Mathematics, 74. Philadelphia, PA: Society for Industrial and Applied Mathematics (SIAM), 2001.
- [11] *Crama Y., Hammer P.L.* Boolean functions. Theory, algorithms, and applications. Encyclopedia of Mathematics and its Applications, 142. Cambridge: Cambridge University Press, 2011.
- [12] *Cygan M., Fomin F.V., Kowalik L., Lokshtanov D., Marx D., Pilipczuk M., Pilipczuk M., Saurabh S.* Parameterized algorithms. Cham: Springer, 2015.
- [13] *Fomin F.V., Lokshtanov D., Saurabh S., Zehavi M.* Kernelization. Theory of parameterized preprocessing. Cambridge: Cambridge University Press, 2019.
- [14] *Frankl P., Tokushige N.* Extremal problems for finite sets. Student Mathematical Library, 86. Providence, RI: American Mathematical Society, 2018.
- [15] *Gainanov D.N.* Graphs for pattern recognition. Infeasible systems of linear inequalities. Berlin: De Gruyter, 2016.
- [16] *Garey M.R., Johnson D.S.* Computers and intractability. A guide to the theory of NP-completeness. A Series of Books in the Mathematical Sciences. San Francisco, CA: W.H. Freeman and Co., 1979.
- [17] *Gerbner D., Patkós B.* Extremal finite set theory. Discrete Mathematics and its Applications (Boca Raton). Boca Raton, FL: CRC Press, 2019.

- [18] *Godsil C., Meagher K.* Erdős–Ko–Rado theorems: Algebraic approaches. Cambridge Studies in Advanced Mathematics, 149. Cambridge: Cambridge University Press, 2016.
- [19] *Grötschel M., Lovász L., Schrijver A.* Geometric algorithms and combinatorial optimization. Second edition. Algorithms and Combinatorics, 2. Berlin: Springer-Verlag, 1993.
- [20] *Haynes T.W., Hedetniemi S.T., Henning M.A.* (eds.). Structures of domination in graphs. Developments in Mathematics, 66. Cham: Springer, 2021.
- [21] *Henning M.A., Yeo A.* Total domination in graphs. Springer Monographs in Mathematics. New York: Springer, 2013.
- [22] *Henning M.A., Yeo A.* Transversals in linear uniform hypergraphs. Developments in Mathematics, 63. Cham: Springer, 2020.
- [23] *Herzog J., Hibi T.* Monomial ideals. Graduate Texts in Mathematics, 260. London: Springer-Verlag London, Ltd., 2011.
- [24] *Jukna S.* Extremal combinatorics. With applications in computer science. Second edition. Texts in Theoretical Computer Science. An EATCS Series. Heidelberg: Springer, 2011.
- [25] *Knuth D.E.* The art of computer programming. Vol. 4A. Combinatorial algorithms. Part 1. Upper Saddle River, NJ: Addison-Wesley, 2011.
- [26] *Matveev A.O.* Pattern recognition on oriented matroids. Berlin: De Gruyter, 2017.
- [27] *Matveev A.O.* Pattern recognition on oriented matroids: Symmetric cycles in the hypercube graphs. V. Preprint [arXiv:2106.03832], 2021.
- [28] *McMullen P.* The numbers of faces of simplicial polytopes. Israel Journal of Mathematics, 1971, 9, 559–570.
- [29] *McMullen P., Shephard G.C.* Convex polytopes and the upper bound conjecture. Prepared in collaboration with *J.E. Reeve* and *A.A. Ball*. London Mathematical Society Lecture Note Series, vol. 3. London: Cambridge University Press, 1971.
- [30] *McMullen P., Walkup D.W.* A generalized lower-bound conjecture for simplicial polytopes. Mathematika, 1971, 18, pp. 264–273.
- [31] *Mirsky L.* Transversal theory. An account of some aspects of combinatorial mathematics. Mathematics in Science and Engineering, 75. New York: Academic Press, 1971.
- [32] *Nagamochi H., Ibaraki T.* Algorithmic aspects of graph connectivity. Encyclopedia of Mathematics and its Applications, 123. Cambridge: Cambridge University Press, 2008.
- [33] *Nemhauser G.L., Wolsey L.A.* Integer and combinatorial optimization. Reprint of the 1988 original. Wiley-Interscience Series in Discrete Mathematics and Optimization. New York: John Wiley & Sons, Inc., 1999.
- [34] *Petersen T.K.* Eulerian numbers. With a foreword by *R.P. Stanley*. Birkhäuser Advanced Texts: Basler Lehrbücher. New York: Birkhäuser/Springer, 2015.
- [35] *Scheinerman E.R., Ullman D.H.* Fractional graph theory. A rational approach to the theory of graphs. With a foreword by *C. Berge*. Reprint of the 1997 original. Mineola, NY: Dover Publications, Inc., 2011.
- [36] *Schrijver A.* Combinatorial optimization. Polyhedra and efficiency. Vol. C. Disjoint paths, hypergraphs. Chapters 70–83. Algorithms and Combinatorics, 24,C. Berlin: Springer-Verlag, 2003.
- [37] *Stanley R.P.* Combinatorics and commutative algebra. Second edition. Progress in Mathematics, 41. Boston, MA: Birkhäuser, 1996.
- [38] *Villarreal R.H.* Monomial algebras. Second edition. Monographs and Research Notes in Mathematics. Boca Raton, FL: CRC Press, 2015.
- [39] *West D.B.* Combinatorial mathematics. Cambridge: Cambridge University Press, 2021.

- [40] *Ziegler G.M.* Lectures on polytopes. Revised edition. Graduate Texts in Mathematics, 152. Berlin: Springer-Verlag, 1998.

*Email address:* `andrey.o.matveev@gmail.com`