

Open and free EEG datasets for epilepsy diagnosis

Palak Handa · Monika Mathur · Nidhi Goel

Received: date / Accepted: date

Abstract The Epilepsies are a common, chronic neurological disorder affecting more than 50 million individuals across the globe. It is characterized by unprovoked, recurring (similar or different type) seizures which are commonly diagnosed through clinical EEGs. Good-quality, open-access and free EEG data can act as a catalyst for on-going state-of-the-art (SOTA) research works for detection, prediction and management of epilepsy and seizures. They can also aid in improving the quality of life (QOL) of these diseased individuals and contribute research in healthcare multimedia, data analytics and Artificial Intelligence (AI) in personalized medicine. This paper presents widely used, available, open and free EEG datasets available for epilepsy and seizure diagnosis. A brief comparison and discussion of open and private datasets has also been done. Such datasets will help in development and evaluation of automatic computer-aided system in healthcare.

Keywords Open datasets · Biomedical multimedia · EEG signals · Epilepsy diagnosis · Seizure · Performance

1 Introduction

The Epilepsies are a chronic neurological disorder characterized by unprovoked, recurring (similar or differ-

ent type) seizures. A clinical EEG setting is used by doctors to observe different types of epileptic activity as it leaves distinct impressions in the form of interictal epileptiform discharges, peri-ictal activities and high frequency oscillations *etc* [1].

The annual economic burden of epilepsy is enormous in developing countries like India where it is estimated to be 88.2% of gross national product (GNP) per capita and 0.5% of the overall GNP [1]. Hence, early diagnosis through recent technologies like Artificial Intelligence (AI), feature engineering, data analytics and multimedia is vital and can aid in Quality Of Life (QOF) of patients and their associated caretakers.

Secured, reproducible AI algorithm, good quality data and efficient computing horse power are the major elements for development of early detection and prediction of epileptic wave forms through EEG signals. There are several types of EEGs such as intracranial, scalp, ambulatory, *etc*. They are recorded in video, image and signal format depending on their use and application in hospitals.

There is a huge demand of real-time biomedical multimedia tools for data analysis, and pattern recognition of such formats. Bonn EEG time series database [2] was the first EEG dataset to be publically available for research applications in this field. It remains as the benchmark dataset for most research works due to its availability. Several datasets discussed in this paper encourage scientific advances in this field.

The data quality of biomedical datasets is measured through various factors such as presence of artefacts and noise, missing values, descriptive information, annotations by health experts, pre-fined data structure, processing and robustness to outliers *etc*.

The datasets mentioned in the section 2 are freely available (except temple university which requires a lo-

P. Handa
Dept. of ECE, DTU, Delhi
E-mail: palakhanda97@gmail.com

M. Mathur
Dept. of ECE, IGDTUW, Delhi
E-mail: mathur.monika2007@gmail.com

N. Goel
Dept. of ECE, IGDTUW, Delhi
E-mail: nidhi.iitr1@gmail.com

gin), re-distributable for research purposes, previously freely available and or became private or removed from the portal based on adult and paediatric population. Table 1 shows a comparison of publically available, open-access (except temple university which requires a login) human EEG datasets for epilepsy diagnosis.

The source and availability of these were verified on 26-07-2021, which may change in the future. They were found using different keywords like ‘EEG datasets for epilepsy’, ‘datasets for epilepsy detection’, ‘EEG based epilepsy diagnosis’, and ‘open EEG datasets’ on Pub med and google scholar search engine.

2 EEG datasets for epilepsy diagnosis

There are several EEG datasets for epilepsy diagnosis which are freely available and private due to various reasons such as lack of ethical clearance. This section lists all the existing EEG datasets with their URLs for adult and paediatric population.

2.1 Adult datasets

The adult population consists of affected individuals above 20 years of age. There are several databases like American Epilepsy Society Seizure Prediction Challenge database [3], dataset of EEG recordings of pediatric patients with epilepsy based on the 10-20 system [4] and Karunya University [5] which contain both adult and paediatric EEG data. They are mentioned in section 2.2.

2.1.1 Bonn EEG time series database[2]

This database comprises of 100 single channels EEG of 23.6 seconds with sampling rate of 173.61 Hz. Its spectral bandwidth range is between 0.5 Hz and 85 Hz. It was taken from a 128 channel acquisition system. Five patients EEG sets were cut out from a multi-channel EEG recording and named A, B, C, D and E. Set A and B are the surface EEG recorded during eyes closed and open situation of healthy patients respectively. Set C and D are the intracranial EEG recorded during a seizure free from within seizure generating area and from outside seizure generating area of epileptic patients respectively. Set E is the intracranial EEG of an epileptic patient during epileptic seizures. Each set contains 100 text files wherein each text file has 4097 samples of 1 EEG time series in ASCII code. A band pass filter with cut off frequency as 0.53 Hz and 40 Hz has been applied on the data. It is an artifact free data and hence no prior pre-processing is required for the

classification of healthy (non-epileptic) and un-healthy (epileptic) signals. The strong eye movement’s artefacts were omitted. It was made available in 2001. The extended version of this data is now a part of EPILEPSIA project. Available link: [a](#) and [b](#)

2.1.2 Bern-Barcelona EEG database [6]

This multi channel EEG database was recorded using specialized electrodes and consists of five patients with longstanding pharmacoresistant temporal lobe epilepsy. The patients underwent epilepsy surgery. The sampling rate was either 512 or 1024 Hz based upon whether they were recorded with less or more than 64 channels of EEG system. Three out of five attained complete seizure freedom. Two types of EEG are present in the data *i.e.*, focal and non-focal. Each file has about 10240 samples for a time duration of 20 seconds. Available link: [c](#) and [d](#)

2.1.3 Temple University EEG corpus [7]

Temple University EEG corpus is the largest free EEG data available for epilepsy and seizure types diagnosis till date. It consists of data acquired from 2000 to 2013 using different EEG clinical settings for about 10,874 patients. This community has developed various software products such as annotation tools, toolboxes for seizure detection, and EDF browser for data analysis of EEG, EMG, and ECG *etc* signals. EDF browser helps to view EEG recording in a video form. There are various datasets available such as IBM Features For Seizure Detection (IBMFT), the TUH EEG epilepsy corpus, seizure corpus, slowing corpus, and events corpus *etc*. A user ID and password is required to get access to these datasets. [Available link.](#)

2.1.4 Neurology and sleep center, New Delhi EEG dataset [8]

This database comprises of 5.12 seconds EEG data. It was recorded using 57 EEG channel Grass Tele-factor Comet AS40 Amplification System; sampled at 200 Hz. It’s spectral bandwidth range is between 0.5 Hz and 70 Hz. Time series EEG datasets are categorized into three major MATLAB file folder namely ictal, pre-ictal and inter-ictal stages. Each MAT file has 1024 samples. A subset of this database is publically available. [Available link.](#)

2.1.5 Epileptic Seizure Recognition Data Set [2]

The time series EEG dataset consists of 11500 instances of EEGs of 4 subjects suffering from epilepsy. This data

has been removed from the UCI machine learning repository recently and was released in 2017. It is a simplified version of the original data released by [2]. It consists of 5 subjects (4 unhealthy and 1 healthy) performing different activities and experience epileptic seizures except subject 1. The time duration for each EEG was 23.5 seconds.

2.1.6 Siena Scalp EEG Database [9, 10]

This multi-channel EEG database of 14 epileptic patients (9 males and 5 females) was recorded using specialized amplifiers, and reusable electrodes. The signals were recorded with a sampling rate of 512 Hz and stored in EDF files. The data has been acquired from Unit of Neurology and Neurophysiology of the University of Siena, Italy and focuses on seizure prediction. It is an integral part of national interdisciplinary research project PANACEE. This data contains 47 seizures from 128 hours of video EEG recording. The start and end time of a seizure was also recorded and contains the list of electrodes present on the scalp of a patient during event recording. Three types of seizures namely focal onset with and without impaired awareness, and focal to bilateral tonic-clonic (FBTC) were found and recorded in the diseased patients. [Available link](#).

2.1.7 Single electrode EEG data of healthy and epileptic patients [11, 12]

This dataset was generated with a motive to build predictive epilepsy diagnosis model and publically available since 2020. It was generated on a similar acquisition and settings *i.e.*, sampling frequency, bandpass filtering and number of signals and time duration as of University of Bonn. It has overcome the limitations faced by University of Bonn dataset such as different EEG recording (inter-cranial and scalp) for healthy and epileptic patients [11]. All the data were taken exclusively using surface EEG electrodes for 15 healthy and epileptic patients. [Available link](#).

2.1.8 Epileptic EEG Dataset [13]

This multi-channel, long term EEG database was recorded for 6 patients suffering from focal epilepsy. They were undergoing pre-surgical evaluation for possible epilepsy surgery. Different EEG segments of a seizure like ictal, pre-ictal, inter-ictal and its onsets have been included in the data. The signals were recorded with a sampling rate of 500 Hz and stored in EDF files. Labelled and classified data points (train and test set) have been mentioned for complex partial electrographic, and video-detected seizures. All the EEG signals underwent band

pass filtering of range 1-70 Hz where 50 Hz (utility frequency) was also removed. [Available link](#).

2.2 Paediatric datasets

The paediatric EEG database consists of affected individuals from age 1 month - 20 years.

2.2.1 Children's hospital Boston-MIT database [14]

This database comprises of 844 hour continuous EEG. 23 pediatric patients from age 1.5-19 who underwent scalp multi-channel EEG recording. It is the first paediatric EEG database available for epilepsy and seizure diagnosis. The patients were given anti-seizure medications. About 200 seizures were recorded in a universal bio-polar montage with about 24-27 EEG channels. Sampling frequency was kept to be 256 Hz. Each EEG segment is called as a record which usually is for duration of one hour. There are 9-42 edf files from a single subject. Additional vagal nerve stimulus signals are also present. Separate file name and montages have been mentioned for seizure v/s non seizure episode in EEG segments. [Available link](#).

2.2.2 Karunya University [5]

This database comprises of 18 channel EEG data with segments of normal, focal and generalized epileptic seizure activities from 1-107 years of patients. It was released in 2014 but the website is not available for research use now. Each segment has 2056 sample points. Sampling frequency was kept to be 256 Hz. The EEG recordings vary from 40 minutes to one hour. It was collected from a diagnostic center based in Coimbatore, India. [Available link](#).

2.2.3 A dataset of neonatal EEG recordings with seizures annotations [15]

This database consists of multi-channel, good quality EEG recordings of 79 term neonates where 39 of them suffered from neonatal seizures in the NICU of Helsinki University Hospital, Finland. The recordings were captured with NicOne EEG amplifier, and 19 EEG channel cap. The signals were recorded with a sampling rate of 256 Hz and stored in EDF files. It consists of seizure annotations by healthcare experts for seizure detection purpose. The data was pre-processed using butterworth high-pass filtering. The data also contains natural artefacts. [Available link](#).

Table 1 Comparison of existing EEG datasets for epilepsy diagnosis

Ref.	Availability	Type	Source	Year	Size	No. of channels	No. of patients	Sampling frequency	EEG segments
[2]	Freely available	Adult	e-repositori upf.	2001	3.05 MB	100 single	5	173.61 Hz	seizure states, healthy
[14]	Freely available	Paediatric	PhysioNet repository	2010	40 GB	23-26	22	256 Hz	Intractable seizures
[6]	Freely available	Adult	e-repositori upf.	2012	814 MB	64	5	512 Hz	Focal, Non-focal
[3]	Freely available	Dog and human	Kaggle	2014	105 GB	-	-	-	different types
[5]	Not available	Adult and Paediatric	Website	2014	-	-	-	256 Hz	normal, focal and generalized epileptic seizures different types
[7]	Free but requires login	Adult	Website	2015	572 GB	20-31	10,874	250, 256, 512 Hz	
[8]	Freely available	Adult	Researchgate	2016	604 KB	57	10	200 Hz	Ictal, inter-ictal, pre-ictal EEGs
[2]	Removed	Adult	UCI repository	2017	3 MB	100 single	5	173.61 Hz	seizure related, healthy
[15]	Freely available	Paediatric (neonates)	Zenedo	2018	4.3 GB	19	79	256 Hz	seizure onset
[16]	Private	Adult	-	2019	-	19	115	128 Hz	epileptic and healthy
[11]	Private	Adult	-	2019	-	-	50	250, 256 Hz	generalized and focal epilepsies
[17]	Private	Adult	-	2019	-	21	5	500 Hz	focal and tonic-clonic
[18]	Private	Pediatric	-	2019	-	-	29	200, 500 Hz	typical absence seizures
[19]	Private	Adult	-	2019	-	-	12	256 Hz	seizure events
[20]	Private	-	-	2019	-	21	25	200 Hz	seizure events
[21]	Private	-	-	2019	-	18	10	256 Hz	seizure states
[22]	Private	-	-	2019	-	22	22	250 Hz	ictal, non-ictal
[12]	Freely available	Adult	Zenedo	2020	20 MB	-	15	173.61 Hz	Inter-ictal
[23]	Private	-	-	2020	-	21	-	250 Hz	seizure onsets
[24]	Private	Adult	-	2020	-	21	150	256 Hz	seizure and normal
[10]	Freely available	Adult	PhysioNet repository	2020	20 GB	29	14	512 Hz	Epileptic seizures (focal onset, tonic-clonic)
[13]	Freely available	Adult	Mendeley repository	2021	3133 MB	21	6	500 Hz	Complex partial, electrographic and video-detected seizures
[4]	Freely available	Paediatric and Adult	Open neuro repository	2021	15 GB	52	30	2000 Hz	HFO markings

2.2.4 Dataset of EEG recordings of pediatric patients with epilepsy based on the 10-20 system [4]

This dataset consists of scalp EEG recordings to study the impact age on observed High Frequency Oscillations (HFO) in pediatric epileptic patients. Three hours of pediatric and adult EEG sleep data was recorded for 30 focal or generalised epileptic patients. The signals were recorded with a sampling rate of 2000 Hz and stored in EDF files. Different sleep stage annotations are available in this database. [Available link.](#)

2.3 Others

2.3.1 American Epilepsy Society Seizure Prediction Challenge database [3]

This database consists of intra-cranial EEG segments from dogs and humans with different acquisitions of sampling rate, duration of EEGs, and no. of electrodes *etc.* It was released as a part of the kaggle challenge hosted by the American Epilepsy Society in 2014 for development of seizure forecasting systems and witnessed about 504 teams. Different seizure segments of ictal, pre-ictal, post-ictal, inter-ictal were provided in MATLAB files. The data storage was about 105 GB. Additional annotated EEG data was also provided by the University of Pennsylvania and the Mayo Clinic. [Available link.](#)

2.3.2 Private databases

Several private databases have also been recorded for epilepsy diagnosis using EEG signals [11, 24, 16, 17, 18, 22, 23, 19, 20, 21]. The European Epilepsy database [25] is a private database which consists of high quality, annotated EEG signals from University of Bonn [2], Freiburg [26], Flint hills, and many multi-modal like MRI imaging data. The website of [5] is not available. EEG data in [27] was freely available till 2015 when its portal crashed.

3 Conclusion

Diagnosis, treatment and management of epilepsy is still a challenging task for the scientific and health-care community. It's detection by visual introspection of long hour EEG is not only time taking but a very tedious and subjective task. Artificial intelligence can help in escalating this process and lead to successful detection of different types of epilepsies through efficient, high quality and annotated EEG data. This paper has

presented all the existing EEG datasets for epilepsy diagnosis with its availability and brief comparisons. Such datasets motivate scientific research in early diagnosis of epilepsy through robust techniques.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. SV Thomas, PS Sarma, M Alexander, L Pandit, L Shekhar, C Trivedi, and B Vengamma. Economic burden of epilepsy in india. *Epilepsia*, 42(8):1052–1060, 2001.
2. Ralph G Andrzejak, Klaus Lehnertz, Florian Mormann, Christoph Rieke, Peter David, and Christian E Elger. Indications of nonlinear deterministic and finite-dimensional structures in time series of brain electrical activity: Dependence on recording region and brain state. *Physical Review E*, 64(6):061907, 2001.
3. J Jeffrey Howbert, Edward E Patterson, S Matt Stead, Ben Brinkmann, Vincent Vasoli, Daniel Crepeau, Charles H Vite, Beverly Sturges, Vanessa Ruedebusch, Jaideep Mavoori, et al. Forecasting seizures in dogs with naturally occurring epilepsy. *PloS one*, 9(1):e81920, 2014.
4. Dorottya Cserpan et al. Dataset of eeg recordings of pediatric patients with epilepsy based on the 10-20 system, 2021.
5. Thomas George Selvaraj, Balakrishnan Ramasamy, Stanly Johnson Jeyaraj, and Easter Selvan Suvishamuthu. Eeg database of seizure disorders for experts and application developers. *Clinical EEG and neuroscience*, 45(4):304–309, 2014.
6. Ralph G Andrzejak, Kaspar Schindler, and Christian Rummel. Nonrandomness, nonlinear dependence, and nonstationarity of electroencephalographic recordings from epilepsy patients. *Physical Review E*, 86(4):046206, 2012.
7. Iyad Obeid and Joseph Picone. The temple university hospital eeg data corpus. *Frontiers in neuroscience*, 10:196, 2016.
8. P Swami, B Panigrahi, S Nara, M Bhatia, and T Gandhi. Eeg epilepsy datasets, 2016.
9. Paolo Detti, Giampaolo Vatti, and Garazi Zabalo Manrique de Lara. Eeg synchronization analysis for seizure prediction: A study on data of noninvasive recordings. *Processes*, 8(7):846, 2020.

10. P. Detti. Siena scalp eeg database (version 1.0.0), 2020. URL <https://doi.org/10.13026/5d4a-j060>.
11. Siddharth Panwar, Shiv Dutt Joshi, Anubha Gupta, and Puneet Agarwal. Automated epilepsy diagnosis using eeg with test set evaluation. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 27(6):1106–1116, 2019.
12. Siddharth Panwar. Single electrode EEG data of healthy and epileptic patients, February 2020. URL <https://doi.org/10.5281/zenodo.3684992>.
13. Wassim Nasreddine. Epileptic eeg dataset, 2021. URL <https://doi.org/10.17632/5pc2j46cbc.1>.
14. Ary L Goldberger, Luis AN Amaral, Leon Glass, Jeffrey M Hausdorff, Plamen Ch Ivanov, Roger G Mark, Joseph E Mietus, George B Moody, Chung-Kang Peng, and H Eugene Stanley. Physiobank, physiotoolkit, and physionet: components of a new research resource for complex physiologic signals. *circulation*, 101(23):e215–e220, 2000.
15. Nathan Stevenson, Karoliina Tapani, Leena Lauronen, and Sampsa Vanhatalo. A dataset of neonatal EEG recordings with seizures annotations, June 2018. URL <https://doi.org/10.5281/zenodo.2547147>.
16. Shivarudhrappa Raghu, Natarajan Sriraam, Yasin Temel, Shyam Vasudeva Rao, Alangar Satyaranjandas Hegde, and Pieter L Kubben. Performance evaluation of dwt based sigmoid entropy in time and frequency domains for automated detection of epileptic seizures using svm classifier. *Computers in biology and medicine*, 110:127–143, 2019.
17. Duanpo Wu, Zimeng Wang, Lurong Jiang, Fang Dong, Xunyi Wu, Shuang Wang, and Yao Ding. Automatic epileptic seizures joint detection algorithm based on improved multi-domain feature of ceeg and spike feature of aeeg. *IEEE Access*, 7: 41551–41564, 2019.
18. Mustafa Talha Avcu, Zhuo Zhang, and Derrick Wei Shih Chan. Seizure detection using least eeg channels by deep convolutional neural network. In *ICASSP 2019-2019 IEEE international conference on acoustics, speech and signal processing (ICASSP)*, pages 1120–1124. IEEE, 2019.
19. Peter Z Yan, Fei Wang, Nathaniel Kwok, Baxter B Allen, Sotirios Keros, and Zachary Grinspan. Automated spectrographic seizure detection using convolutional neural networks. *Seizure*, 71:124–131, 2019.
20. Gwangho Choi, Chulkyun Park, Junkyung Kim, Kyoungin Cho, Tae-Joon Kim, HwangSik Bae, Kyeongyuk Min, Ki-Young Jung, and Jongwha Chong. A novel multi-scale 3d cnn with deep neural network for epileptic seizure detection. In *2019 IEEE International Conference on Consumer Electronics (ICCE)*, pages 1–2. IEEE, 2019.
21. Jiuwen Cao, Jiahua Zhu, Wenbin Hu, and Anton Kummert. Epileptic signal classification with deep eeg features by stacked cnns. *IEEE Transactions on Cognitive and Developmental Systems*, 12(4):709–722, 2019.
22. Muhammad Bilal, Muhammad Rizwan, Sajid Saleem, Muhammad Murtaza Khan, Mohammed Saeed Alkathir, and Mohammed Alqarni. Automatic seizure detection using multi-resolution dynamic mode decomposition. *IEEE Access*, 7:61180–61194, 2019.
23. Dhanalekshmi P Yedurkar and Shilpa P Metkar. Multiresolution approach for artifacts removal and localization of seizure onset zone in epileptic eeg signal. *Biomedical Signal Processing and Control*, 57:101794, 2020.
24. Khakon Das, Debashis Daschakladar, Partha Pratim Roy, Atri Chatterjee, and Shankar Prasad Saha. Epileptic seizure prediction by the detection of seizure waveform from the pre-ictal phase of eeg signal. *Biomedical Signal Processing and Control*, 57:101720, 2020.
25. Matthias Ihle, Hinnerk Feldwisch-Drentrup, César A Teixeira, Adrien Witon, Björn Schelter, Jens Timmer, and Andreas Schulze-Bonhage. Epilepsiae—a european epilepsy database. *Computer methods and programs in biomedicine*, 106(3):127–138, 2012.
26. Björn Schelter, Matthias Winterhalder, Thomas Maiwald, Armin Brandt, Ariane Schad, Jens Timmer, and Andreas Schulze-Bonhage. Do false predictions of seizures depend on the state of vigilance? a report from two seizure-prediction methods and proposed remedies. *Epilepsia*, 47(12):2058–2070, 2006.
27. Piotr Zwoliński, Marcin Roszkowski, Jarosław Żygierewicz, Stefan Haufe, Guido Nolte, and Piotr J Durka. Open database of epileptic eeg with mri and postoperational assessment of foci—a real world verification for the eeg inverse solutions. *Neuroinformatics*, 8(4):285–299, 2010.