KK-like relations of α' corrections to disk amplitudes

Carlos R. Mafra

Mathematical Sciences and STAG Research Centre, University of Southampton, Highfield, Southampton, SO17 1BJ, UK

Inspired by the definition of color-dressed amplitudes in string theory, we define analogous color-dressed permutations replacing the color-ordered string amplitudes by their corresponding permutations. Decomposing the color traces into symmetrized traces and structure constants, the color-dressed permutations define BRST-invariant permutations, which we show are elements of the inverse Solomon descent algebra and we find a closed formula for them. We then present evidence that these permutations encode KK-like relations among the different α' corrections to disk amplitudes refined by their motivic MZV content. In particular, the number of linearly independent amplitudes at a given α' order and motivic MZV content is given by (sums of) Stirling cycle numbers. In addition, we show how the superfield expansion of BRST invariants of the pure spinor formalism corresponding to $\alpha'^2 f_2$ corrections is encoded in the descent algebra.

arXiv:2108.01081v3 [hep-th] 26 Aug 2022 [arXiv:2108.01081v3 \[hep-th\] 26 Aug 2022](http://arxiv.org/abs/2108.01081v3)

August 2021

[†] email: c.r.mafra@soton.ac.uk

Contents

1. Introduction

It is well-known that superstring n-point color-ordered disk amplitudes satisfy monodromy relations which imply that the number of linearly independent amplitudes is $(n-3)!$, for all α' corrections [1,2]. These relations involve coefficients that depend on Mandelstam variables [3] and are famously related to the Bern-Carrasco-Johansson (BCJ) color-kinematics amplitude relations in the field-theory limit. In this paper, we investigate a weaker set of relations, called KK-like relations [4], of higher α' corrections to disk amplitudes [5] refined by their motivic MZV content written in the f-alphabet of [6]. More precisely, writing the motivic superstring color-ordered disk amplitude as¹

$$
\phi\big(A^{\text{string}}(1,2,\ldots,n)\big) = A^{\text{SYM}}(1,2,\ldots,n) + \sum_{m=0}^{\infty} \sum_{M} A_{f_2^m f_M}(1,2,\ldots,n) f_2^m f_M \qquad (1.1)
$$

where M runs over all words composed of odd positive integers $\{2k+1 \mid k \geq 1\}$ and $f_M = f_{m_1} f_{m_2} \dots f_{m_p}$ for words of length p (e.g. $f_{3,5,3} = f_3 f_5 f_3$), we will focus on relations obeyed by the *motivic MZV amplitudes* $A_{f_2^m f_M}$ of the form

$$
\sum_{\sigma} c_{\sigma} A_{f_2^m f_M}(\sigma) = 0,
$$
\n(1.2)

where the coefficients c_{σ} are rational numbers independent of Mandelstam invariants. In addition, the map ϕ in (1.1) converts the motivic MZVs to the f-alphabet as described in [6,5]. It is well-known that (1.1) is cyclically symmetric and satisfies $A^{string}(1, 2, \ldots, n) =$ $(-1)^n A^{\text{string}}(n, n-1, \ldots, 1)$, so the basis dimensions of motivic MZV amplitudes is at most $\frac{1}{2}(n-1)!$. When the superstring amplitude (1.1) is restricted to its field-theory limit A^{SYM} , the KK-like relations correspond to the famous Kleiss-Kuijf (KK) relations [7], under which there are only $(n-2)!$ independent amplitudes. However, it is not yet known the general form of the KK-like relations and the corresponding basis dimensions for the $f_2^m f_M$ components of (1.1) with $m \geq 2$. For instance, a brute-force search indicates that the upper bound $\frac{1}{2}(n-1)!$ is saturated for $A_{f_2^2 f_M}(1, 2, \ldots, n)$ when $n = 4, 5, 6$ and 7, but the $n = 8$ case is different. In fact,

$$
#(A_{f_2^2 f_M}(1,2,\ldots,8)) = 2519 = \frac{1}{2}7! - 1.
$$
\n(1.3)

¹ For more information on motivic amplitudes and the f -alphabet expansion, see [5,6].

In this paper, this result and its generalization will be obtained as

$$
\#(A_{f_M}(1,2,\ldots,n)) = {n-1 \brack 1} = (n-2)! \qquad (1.4)
$$

$$
\#(A_{f_2f_M}(1,2,\ldots,n)) = {n-1 \brack 3}
$$

$$
\#(A_{f_2^m f_M}(1,2,\ldots,n)) = {n-1 \brack 1} + {n-1 \brack 3} + \cdots + {n-1 \brack 2m+1}, \quad m \ge 2.
$$

We will see that the general KK-like relations are closely related to the mathematical framework of the Solomon descent algebra [8,9,10,11,12,13,14]. To see this we will define the color-dressed permutation

$$
P_n = \sum_{\sigma \in S_n, \sigma(1)=1} \text{Tr}(T^{\sigma}) \sigma, \qquad T^{\sigma} := T^{\sigma(1)} T^{\sigma(2)} \cdots T^{\sigma(n)} \qquad (1.5)
$$

which is inspired by the expression of the color-dressed disk amplitudes, where $T^i := T^{a_i}$ denotes a Chan-Paton factor. When the closed formula from [15] for the color-trace decomposition [16] is plugged into (1.5), the permutations appearing as coefficients with respect to a basis of color factors define what we call BRST invariant permutations $\gamma_{1|P_1,...,P_k}$ with $1 \leq k \leq n-1$. We conjecture the following closed formula

$$
\gamma_{1|P_1,\dots,P_k} = 1.\mathcal{E}(P_1)\sqcup \mathcal{E}(P_2)\sqcup \dots \sqcup \mathcal{E}(P_k) \tag{1.6}
$$

where $\mathcal{E}(P)$ satisfying $\mathcal{E}(R \sqcup S) = 0$ for $R, S \neq \emptyset$ is the *Berends-Giele idempotent*, defined in section 2.2 from mapping the permutations of the Solomon idempotent [17] into their inverses. Then in section 3.1 we will find evidence that these BRST-invariant permutations encode the general KK-like relations as

$$
A_{f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 1,
$$

\n
$$
A_{f_2f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 3
$$

\n
$$
A_{f_2^m f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 1, 3, 5,..., 2m + 1, \quad m \ge 2.
$$
\n(1.7)

Studying the color-dressed string disk amplitude we will obtain, following the results of [18], a correspondence between the above permutations and kinematics from the string disk amplitudes. More precisely, the duality from [18] is generalized to

$$
A^{\text{SYM}}(1,2,\ldots,n) \longleftrightarrow \gamma_{123\ldots n}^{(1)}, \qquad A_{f_2}(1,2,\ldots,n) \iff \gamma_{123\ldots n}^{(3)}, \qquad (1.8)
$$

where $\gamma^{(1)}$ and $\gamma^{(3)}$ are orthogonal idempotents of the (inverse) descent algebra constructed from linear combinations of $\gamma_{1|P_1,...,P_k}$ with $k=1$ and $k=3$ respectively. This interpretation is important because we can use a theorem from the work of Garsia and Reutenauer [9] to prove results conjectured in [18].

In section 3.3, the BRST-invariant permutations will be used to define higher-mass BRST-invariant superfields via

$$
\gamma_{1|P_1, P_2, \dots, P_k} \leftrightarrow A^{\text{string}}(\gamma_{1|P_1, P_2, \dots, P_k}), \qquad A_{f_2^m f_M}(\gamma_{1|P_1, \dots, P_k}) = k! C_{1|P_1, \dots, P_k}^{f_2^m f_M}. \qquad (1.9)
$$

As shown in section 3.4, the case for f_2 reduces to the BRST-invariants studied in [18,19]. In the appendices we review the descent algebra and collect various proofs and explicit expansions omitted from the main text.

1.0.1. Conventions

Words from the alphabet $\mathbb{N} = \{1, 2, \ldots\}$ will be denoted either by capital Latin letters or, especially when viewed as elements of the permutation group, by lower case Greek letters. The symmetric group S_n acts on words of length n via the right-action multiplication [14]

$$
P \circ \sigma = p_{\sigma(1)} p_{\sigma(2)} \dots p_{\sigma(n)}, \tag{1.10}
$$

where p_i denotes the *i*th letter of P. For example $abcd \circ 3124 = cabd$. The inverse σ^{-1} of a permutation σ of length n is such that $\sigma \circ \sigma^{-1} = \sigma^{-1} \circ \sigma = 12 \dots n$. For example, $(2314)^{-1} = 3124$. For typographical convenience, we write a permutation σ as W_{σ} .

2. The combinatorics of color-dressed permutations

2.1. Color-dressed permutations

In this section we will investigate the combinatorics of the *color-dressed permutations* P_n

$$
P_n = \sum_{\sigma \in S_n, \sigma(1)=1} \text{Tr}(T^{\sigma}) W_{\sigma}, \qquad T^{\sigma} := T^{\sigma(1)} T^{\sigma(2)} \cdots T^{\sigma(n)}, \qquad (2.1)
$$

arising from decomposing [16] the traces of color factors into symmetrized traces $d^{12...k}$ and structure constants f^{abc} of the gauge group [15]

$$
\text{Tr}(T^0 T^1 \cdots T^{n-1}) = \sum_{S_{n-1}\ni\sigma=\sigma_1\cdots\sigma_k} i^{n-1-k} \kappa_{\sigma_1} \cdots \kappa_{\sigma_k} d^{0a_1\cdots a_k} F^{\sigma_1}_{a_1} \cdots F^{\sigma_k}_{a_k}.
$$
 (2.2)

where $\sigma = \sigma_1 \cdot \sigma_2 \cdot \ldots \cdot \sigma_k$ denotes the decreasing Lyndon factorization of σ to be defined below, F_a^{σ} for a word σ and a letter a is defined recursively by $F_a^{Pj} = F_b^P f^{bja}$ with base case $F_a^i = \delta_a^i$. The coefficients κ_{σ} are defined in (A.20). In addition, the symmetrized trace and the structure constant are given by

$$
d^{12...k} := \frac{1}{k!} \sum_{\sigma \in S_k} \text{Tr}(T^{\sigma}), \qquad d^{12} := \frac{1}{2} \delta^{12}, \quad [T^a, T^b] = i f^{abc} T^c. \tag{2.3}
$$

The decreasing Lyndon factorization (dLf) of σ is defined as [20,9] $\sigma = \sigma_1 \cdot \sigma_2 \ldots \sigma_k$ representing the unique deconcatenation of σ into Lyndon subwords $\sigma_1, \ldots, \sigma_k$ such that $\sigma_1 > \cdots > \sigma_k$ in the lexicographical order of the alphabet $\mathbb{N} = \{1, 2, \ldots, \}$. Representing the concatenation by a dot to distinguish the subwords σ_i in the dLF factorization of σ , we have, for example $1432 = 1432$, $2134 = 2.134$, $54132 = 5.4.132$, $42671835 = 4.267.1835$.

Definition (BRST-invariant permutations). The BRST-invariant permutations are the coefficients with respect to the basis of color factors $d^{1a_1...a_k}F^{\sigma_1}_{a_1} \dots F^{\sigma_k}_{a_k}$, where $\sigma_1 \cdots \sigma_k$ is the decreasing Lyndon factorization of σ , in the color-dressed permutation (2.1), i.e.

$$
P_n = \sum_{\sigma \in S_{n-1}} i^{n-1-k} d^{1a_1 \dots a_k} F_{a_1}^{\sigma_1} \cdots F_{a_k}^{\sigma_k} \gamma_{1 | \sigma_1, \dots, \sigma_k}, \qquad (2.4)
$$

The reason for this terminology will become clear in (3.23) when $\gamma_{1|\sigma_1,\sigma_2,\sigma_3}$ will be associated to BRST invariants superfields in the pure spinor formalism. For example, plugging $\text{Tr}(T^1T^2T^3) = d^{1a_1a_2}F^2_{a_1}F^3_{a_2} + \frac{i}{2}$ $\frac{i}{2}d^{1a}F_a^{23}$ into (2.1) yields

$$
P_3 = d^{1ab} F_a^2 F_b^3 \gamma_{1|2,3} + i d^{1a} F_a^{23} \gamma_{1|23} , \qquad (2.5)
$$

with

$$
\gamma_{1|2,3} = W_{123} + W_{132}, \quad \gamma_{1|23} = \frac{1}{2} W_{123} - \frac{1}{2} W_{132}.
$$
 (2.6)

Repeating the same exercise for $n = 4$ using (2.2) we obtain

$$
P_4 = d^{1abc} F_a^2 F_b^3 F_c^4 \gamma_{1|2,3,4}
$$

+ $i d^{1ab} F_a^{23} F_b^4 \gamma_{1|23,4} + i d^{1ab} F_a^{24} F_b^3 \gamma_{1|24,3} + i d^{1ab} F_a^2 F_b^{34} \gamma_{1|2,34}$
+ $i^2 d^{1a} F_a^{234} \gamma_{1|234} + i^2 d^{1a} F_a^{243} \gamma_{1|243}$ (2.7)

where the BRST-invariant permutations are given by

$$
\gamma_{1|2,3,4} = W_{1234} + W_{1243} + W_{1324} + W_{1342} + W_{1423} + W_{1432},
$$
\n
$$
\gamma_{1|23,4} = \frac{1}{2}W_{1234} + \frac{1}{2}W_{1243} - \frac{1}{2}W_{1324} - \frac{1}{2}W_{1342} + \frac{1}{2}W_{1423} - \frac{1}{2}W_{1432},
$$
\n
$$
\gamma_{1|2,34} = \frac{1}{2}W_{1234} - \frac{1}{2}W_{1243} + \frac{1}{2}W_{1324} + \frac{1}{2}W_{1342} - \frac{1}{2}W_{1423} - \frac{1}{2}W_{1432},
$$
\n
$$
\gamma_{1|24,3} = \frac{1}{2}W_{1234} + \frac{1}{2}W_{1243} + \frac{1}{2}W_{1324} - \frac{1}{2}W_{1342} - \frac{1}{2}W_{1423} - \frac{1}{2}W_{1432},
$$
\n
$$
\gamma_{1|234} = \frac{1}{3}W_{1234} - \frac{1}{6}W_{1243} - \frac{1}{6}W_{1324} - \frac{1}{6}W_{1342} - \frac{1}{6}W_{1423} + \frac{1}{3}W_{1432},
$$
\n
$$
\gamma_{1|243} = -\frac{1}{6}W_{1234} + \frac{1}{3}W_{1243} - \frac{1}{6}W_{1324} + \frac{1}{3}W_{1342} - \frac{1}{6}W_{1423} - \frac{1}{6}W_{1432}.
$$
\n(2.8)

For $n = 5$ we obtain

$$
P_{5} = d^{1abcd}F_{a}^{2}F_{b}^{3}F_{c}^{4}F_{d}^{5}\gamma_{1|2,3,4,5}
$$
\n
$$
+ id^{1abc}F_{a}^{23}F_{b}^{4}F_{c}^{5}\gamma_{1|23,4,5} + id^{1abc}F_{a}^{24}F_{b}^{3}F_{c}^{5}\gamma_{1|24,3,5} + id^{1abc}F_{a}^{25}F_{b}^{3}F_{c}^{4}\gamma_{1|25,3,4}
$$
\n
$$
+ id^{1abc}F_{a}^{2}F_{b}^{34}F_{c}^{5}\gamma_{1|2,34,5} + id^{1abc}F_{a}^{2}F_{b}^{35}F_{c}^{4}\gamma_{1|2,35,4} + id^{1abc}F_{a}^{2}F_{b}^{3}F_{c}^{45}\gamma_{1|2,3,45}
$$
\n
$$
+ i^{2}d^{1ab}F_{a}^{234}F_{b}^{5}\gamma_{1|234,5} + i^{2}d^{1ab}F_{a}^{243}F_{b}^{5}\gamma_{1|243,5} + i^{2}d^{1ab}F_{a}^{235}F_{b}^{4}\gamma_{1|235,4}
$$
\n
$$
+ i^{2}d^{1ab}F_{a}^{245}F_{b}^{3}\gamma_{1|245,3} + i^{2}d^{1ab}F_{a}^{253}F_{b}^{4}\gamma_{1|253,4} + i^{2}d^{1ab}F_{a}^{254}F_{b}^{3}\gamma_{1|254,3}
$$
\n
$$
+ i^{2}d^{1ab}F_{a}^{345}F_{b}^{2}\gamma_{1|345,2} + i^{2}d^{1ab}F_{a}^{354}F_{b}^{2}\gamma_{1|354,2}
$$
\n
$$
+ i^{2}d^{1ab}F_{a}^{23}F_{b}^{45}\gamma_{1|23,45} + i^{2}d^{1ab}F_{a}^{24}F_{b}^{35}\gamma_{1|24,35} + i^{2}d^{1ab}F_{a}^{25}F_{b}^{34}\gamma_{1|25,34}
$$
\n
$$
+ i^{3}d^{1a}F_{a}^{2
$$

where the various $\gamma_{1|A_1,A_2,...,A_k}$ are listed in the appendix C.

2.1.1. Relating the BRST-invariant permutations with the descent algebra

The permutations in each $\gamma_{1|A_1,A_2,...,A_k}$ turn out to be related to the descent algebra reviewed in the appendix A. To see this consider $\gamma_{1|23,4}$ from (2.8), relabel $i \rightarrow i - 1$ and strip off the leading "0" (denoted by \times) to obtain

$$
\gamma_{\times|12,3} = \frac{1}{2} \underbrace{W_{123}}_{\in D_0} + \frac{1}{2} \underbrace{W_{132}}_{\in D_{\{2\}}} - \frac{1}{2} \underbrace{W_{213}}_{\in D_{\{1\}}} - \frac{1}{2} \underbrace{W_{231}}_{\in D_{\{2\}}} + \frac{1}{2} \underbrace{W_{312}}_{\in D_{\{1\}}} - \frac{1}{2} \underbrace{W_{321}}_{\in D_{\{1,2\}}},
$$
\n(2.10)

The above permutations are *not* in the descent algebra \mathcal{D}_3 since permutations in the same descent class have different coefficients (see also proposition 2.1 from [21]). However, the inverse permutations in $\theta(\gamma_{\times 12.3})$ do belong to the same descent classes:

$$
\theta(\gamma_{\times|12,3}) = \frac{1}{2} \underbrace{W_{123}}_{\in D_{\emptyset}} + \frac{1}{2} \underbrace{W_{132}}_{\in D_{\{2\}}} - \frac{1}{2} \underbrace{W_{213}}_{\in D_{\{1\}}} - \frac{1}{2} \underbrace{W_{312}}_{\in D_{\{1\}}} + \frac{1}{2} \underbrace{W_{231}}_{\in D_{\{2\}}} - \frac{1}{2} \underbrace{W_{321}}_{\in D_{\{1,2\}}}.
$$
\n(2.11)\n
$$
= \frac{1}{2} D_{\emptyset} - \frac{1}{2} D_{\{1\}} + \frac{1}{2} D_{\{2\}} - \frac{1}{2} D_{\{1,2\}}.
$$

as can be verified using (A.3). This suggests that the BRST-invariant permutations belong to the *inverse* descent algebra $\mathcal{D}'_n := \theta(\mathcal{D}_n)$. To find an algorithm that generates these permutations, we will consider the inverse of the Eulerian idempotent (A.20).

2.2. The Berends-Giele idempotent

Define the Berends-Giele idempotent² \mathcal{E}_n as the inverse of the Eulerian idempotent (A.20):

$$
\mathcal{E}_n = \sum_{\sigma \in S_n} \kappa_{\sigma^{-1}} \sigma, \qquad \mathcal{E}(P) = \mathcal{E}_P := P \circ \mathcal{E}_n, \quad |P| = n. \tag{2.12}
$$

The reason for this terminology is the correspondence with the standard Berends-Giele current of Yang-Mills theory [22], see section 3.2. A few examples of (2.12) are

$$
\mathcal{E}(1) = W_1, \qquad \mathcal{E}(12) = \frac{1}{2}(W_{12} - W_{21}), \tag{2.13}
$$
\n
$$
\mathcal{E}(123) = \frac{1}{3}W_{123} - \frac{1}{6}W_{132} - \frac{1}{6}W_{213} - \frac{1}{6}W_{231} - \frac{1}{6}W_{312} + \frac{1}{3}W_{321},
$$

while the expansion of $\mathcal{E}(1234)$ can be found in the appendix C.1.

Proposition (Shuffle Symmetry). The Berends-Giele idempotent (2.12) satisfies

$$
\mathcal{E}(R \sqcup S) = 0, \qquad R, S \neq \emptyset. \tag{2.14}
$$

*Proof.*³ Since the sum in (2.12) is over all permutations we rename $P \circ \sigma = \tau$ and sum over τ. Notice that $\sigma^{-1} = \tau^{-1} \circ P$, so $\mathcal{E}(P) = \sum_{\sigma} \kappa_{\sigma^{-1}} P \circ \sigma = \sum_{\tau} \kappa_{(\tau^{-1} \circ P)} \tau$ and therefore

$$
\mathcal{E}(R \sqcup S) = \sum_{\tau} \kappa_{(\tau^{-1} \circ (R \sqcup S))} \tau = \sum_{\tau} \kappa_{(\tau^{-1}(R) \sqcup \tau^{-1}(S))} \tau = 0 \tag{2.15}
$$

where the last equality follows from $(A.22)$ and the crucial observation in (1.5) of [9] that $\sigma^{-1} \circ (R \sqcup S) = \sigma^{-1}(R) \sqcup \sigma^{-1}(S)$, where $\sigma^{-1}(R)$ denotes the word obtained by replacing each letter in R by its image under σ^{-1} . \Box

² The inverse of an idempotent is also idempotent.

³ We know from [23] that any Lie polynomial can be expanded as $\sum_{\sigma} M_{\sigma} \sigma$ with $M_{R \sqcup S} = 0$ for nonempty R, S, so it follows that if Γ is a Lie polynomial then the word function $\mathcal{F}(P) := P \circ \theta(\Gamma)$ satisfies the shuffle symmetry $\mathcal{F}(R \sqcup S) = 0$ for $R, S \neq \emptyset$.

2.3. Inverse idempotent basis and BRST-invariant permutations

Following the realization in section 2.1.1 that the BRST-invariant permutations are related to the inverse of the descent algebra, we define the inverse of the idempotent basis I_p as

$$
\mathcal{I}_{p_1p_2...p_k}(P_1, P_2,..., P_k) := \theta(I_{p_1p_2...p_k})(P), \quad |P_i| = p_i, \quad P_1P_2...P_k = P \qquad (2.16)
$$

where the map θ is defined in (A.14). For example $\mathcal{I}_{21}(12,3) = \frac{1}{2}(W_{123} + W_{132} - W_{213} - W_{123})$ $W_{231} + W_{312} - W_{321}$. See (B.4) for the explicit permutations in $\mathcal{I}_{22}(12, 34)$.

An alternative representation is proven in the appendix B

$$
\mathcal{I}_{p_1p_2\ldots p_k}(P_1, P_2, \ldots, P_k) = \mathcal{E}(P_1) \sqcup \mathcal{E}(P_2) \sqcup \ldots \sqcup \mathcal{E}(P_k), \quad |P_i| = p_i, \qquad (2.17)
$$

from which it follows that $(\tilde{P}$ denotes the reversal⁴ of P)

$$
\mathcal{I}_{\ldots p_i \ldots (s-1, R \sqcup S, \ldots)} = 0, \quad R, S \neq \emptyset, \quad |R| + |S| = p_i, \tag{2.18}
$$
\n
$$
\mathcal{I}_{\ldots p_i \ldots p_j \ldots (s-1, P_i, \ldots, P_j, \ldots)} = \mathcal{I}_{\ldots p_j \ldots p_i \ldots (s-1, P_j, \ldots, P_i, \ldots)},
$$
\n
$$
\tilde{\mathcal{I}}_{p_1 \ldots p_k}(P_1, \ldots, P_k) = (-1)^{\# \text{even}(p)} \mathcal{I}_{p_1 \ldots p_k}(P_1, \ldots, P_k), \tag{2.19}
$$

where $\#\text{even}(p)$ denotes the number of even parts in the composition p. To see this note that if a function satisfies the shuffle symmetry or, in other words, belongs to the dual space of Lie polynomials [24], then $\tilde{F}(P) = (-1)^{|P|-1}F(P)$ and (2.19) follows from (2.17).

Finally, based on multiple examples we conjecture that

$$
\gamma_{1|P_1,...,P_k} = 1.\mathcal{I}_{p_1...p_k}(P_1,...,P_k) = 1.\mathcal{E}(P_1) \sqcup \mathcal{E}(P_2) \sqcup ... \sqcup \mathcal{E}(P_k), \quad p_i = |P_i|.
$$
 (2.20)

The shuffle symmetry of $\mathcal{E}(P_i)$ can be used to fix the first letter of P_i and the commutativity of the shuffle product implies total symmetry in word exchanges, so the basis dimension of $\gamma_{1|P_1,...,P_k}$ with n letters is given by the Stirling cycle numbers [25]

$$
\#(\gamma_{1|P_1,P_2,...,P_k}) = \begin{bmatrix} n-1 \\ k \end{bmatrix}, \qquad \sum_{i=1}^k |P_i| = n-1, \qquad (2.21)
$$

which explains the total number of terms in P_n as $\sum_{k=1}^{n-1} \binom{n-1}{k}$ $\begin{bmatrix} -1 \\ k \end{bmatrix} = (n-1)!$.

⁴ If $Q = q_1 \dots q_n$ then its reversal is the word $\tilde{Q} := q_n \dots q_1$.

2.4. BRST-invariant permutations and orthogonal idempotents

Since the BRST-invariant permutations have been related to the idempotent basis of the (inverse) descent algebra in (2.20) we may construct orthogonal idempotents as in section A.3 by taking the inverse of the Reutenauer idempotents (A.32) $\gamma_{12...n}^{(i)} := 1.\theta(E^{(i)}),$ where the labels in $\theta(E^{(i)})$ must be shifted as $i \to i+1$ prior to the left concatenation with the letter 1. Equivalently, from $(A.30)$, $(A.32)$, and (2.20) we obtain

$$
\gamma_{12...n}^{(k)} = \sum_{23...n = P_1...P_k} \frac{1}{k!} \gamma_{1|P_1,...,P_k}.
$$
\n(2.22)

From the discussion of section A.3 it follows that (2.22) are orthogonal idempotents in the inverse descent algebra \mathcal{D}'_n satisfying (δ^{ij} is the Kronecker delta)

$$
\sum_{k=1}^{n-1} \gamma_{12...n}^{(k)} = W_{12...n}, \qquad \gamma_{12...n}^{(i)} \circ \gamma_{12...n}^{(j)} = \delta^{ij} \gamma_{12...n}^{(i)}
$$
(2.23)

For example, from the BRST-invariant permutations in (2.6) we get

$$
\gamma_{123}^{(1)} \equiv \gamma_{1|23} = \frac{1}{2} W_{123} - \frac{1}{2} W_{132}, \quad \gamma_{123}^{(2)} \equiv \frac{1}{2} \gamma_{1|2,3} = \frac{1}{2} W_{123} + \frac{1}{2} W_{132}, \tag{2.24}
$$

which clearly satisfies (2.23). Similarly, at multiplicity four the definition (2.22)

$$
\gamma_{1234}^{(1)} = \gamma_{1|234}, \qquad \gamma_{1234}^{(2)} = \frac{1}{2} \left(\gamma_{1|23,4} + \gamma_{1|2,34} \right), \qquad \gamma_{1234}^{(3)} = \frac{1}{3!} \gamma_{1|2,3,4}, \tag{2.25}
$$

yields

$$
\gamma_{1234}^{(1)} = \frac{1}{3} W_{1234} - \frac{1}{6} W_{1243} - \frac{1}{6} W_{1324} - \frac{1}{6} W_{1342} - \frac{1}{6} W_{1423} + \frac{1}{3} W_{1432},
$$
\n(2.26)\n
$$
\gamma_{1234}^{(2)} = \frac{1}{2} W_{1234} - \frac{1}{2} W_{1432},
$$
\n
$$
\gamma_{1234}^{(3)} = \frac{1}{6} W_{1234} + \frac{1}{6} W_{1243} + \frac{1}{6} W_{1324} + \frac{1}{6} W_{1342} + \frac{1}{6} W_{1423} + \frac{1}{6} W_{1432}.
$$

It is straightforward but tedious [26] to check that the above satisfy (2.23). At multiplicity five, the orthogonal idempotents are given by

$$
\gamma_{12345}^{(1)} = \gamma_{1|2345},
$$
\n
$$
\gamma_{12345}^{(3)} = \frac{1}{3!} \left(\gamma_{1|23,4,5} + \gamma_{1|2,3,45} \right),
$$
\n
$$
\gamma_{12345}^{(2)} = \frac{1}{2} \left(\gamma_{1|234,5} + \gamma_{1|23,45} + \gamma_{1|2,345} \right),
$$
\n
$$
\gamma_{12345}^{(4)} = \frac{1}{4!} \gamma_{1|2,3,4,5},
$$
\n
$$
\gamma_{12345}^{(4)} = \frac{1}{4!} \gamma_{1|2,3,4,5},
$$
\n
$$
(2.27)
$$

whose expansions can be found in the appendix C and can be checked to obey (2.23).

3. The descent algebra and string disk amplitudes

3.1. KK-like relations of α' corrections to disk amplitudes

The color-dressed string motivic disk amplitude

$$
M_n = \sum_{\sigma \in S_{n-1}} \text{Tr}\left(T^1 T^{\sigma(2)} \cdots T^{\sigma(n)}\right) \phi\left(A^{\text{string}}(1, \sigma(2), \ldots, \sigma(n))\right),\tag{3.1}
$$

is a sum over disk orderings of the $(\phi \text{ map of})$ open string color-ordered amplitude weighted by traces of Chan-Paton factors. The explicit form of the disk amplitudes is a linear combination of field-theory amplitudes A^{SYM} of ten-dimensional super-Yang-Mills [27] given by [28,29]

$$
A^{\text{string}}(P) = \sum_{Q, R \in S_{n-3}} Z(P|1, R, n, n-1) S[R|Q]_1 A^{\text{SYM}}(1, Q, n-1, n) \tag{3.2}
$$

where $S(P|Q)$ ₁ is the field-theory KLT kernel [30,31,32] conveniently computed recursively by $S[A, j|B, j, C]_i = (k_{iB} \cdot k_j)S[A|B, C]_i$, with $S[\emptyset|\emptyset]_i := 1$ [33,34,24]. In addition, $Z(P|Q)$ are the non-abelian Z-theory amplitudes of [35,29]. The color-ordered motivic amplitudes [5] are decomposed as

$$
\phi(A^{\text{string}}(1,2,\ldots,n)) = A^{\text{SYM}}(1,2,\ldots,n) + \sum_{m=0}^{\infty} \sum_{M} A_{f_2^m f_M}(1,2,\ldots,n) f_2^m f_M \qquad (3.3)
$$

where M runs over all words composed of odd positive integers ≥ 3 and f_M = $f_{m_1}f_{m_2}\dots f_{m_p}$ in the f-alphabet of [6] for words of length p. From now on we will use the shorthand *motivic MZV amplitudes* for the $A_{f_2^m f_M}$ components of (3.3), and we will see below that components with the same α' but different motivic MZV content satisfy different relations.

It is well-known that the color-ordered string disk amplitudes are cyclically symmetric and parity invariant [36]

$$
Astring(1, 2, ..., n) = Astring(2, 3, ..., n, 1)
$$
\n(3.4)
\n
$$
Astring(1, 2, ..., n) = (-1)n Astring(n, n - 1, ..., 2, 1).
$$

These relations imply an upper bound on the number of linearly-independent motivic MZV amplitudes $A_{f_2^m f_M}(1, 2, \ldots, n)$ of $\frac{1}{2}(n-1)!$. We want to know whether the different motivic MZV amplitudes in (3.3) satisfy additional KK-like relations and their corresponding basis dimensions, where:

Definition (KK-like relation). An identity of the form

$$
\sum_{\sigma} c_{\sigma} A_{f_2^m f_M}(\sigma) = 0, \tag{3.5}
$$

where the coefficients c_{σ} are rational numbers is said to be KK-like [4].

For an example KK-like relation, one can verify that the f_2 amplitudes satisfy [18],

$$
A_{f_2}(1,2,3,4,5) + \text{perm}(2,4,5) = 0.
$$
\n(3.6)

A central observation of this paper is that the BRST-invariant permutations $\gamma_{1|P_1,...,P_k}$ seem to encode all the KK-like relations among the motivic MZV amplitudes. While the cyclicity relation in (3.4) was used in the definition (3.1) to fix the label in $\gamma_{1|...}$, the parity and cyclicity relations (3.4) are encoded by compositions of $n-1$ with even number of parts:

$$
Astring(\gamma_{1|P_1,...,P_k}) = 0, \quad k \text{ even.}
$$
\n(3.7)

See the appendix D for a proof. The identity $\sum_{k \text{ even}} \binom{n-1}{k}$ $\left[\frac{-1}{k} \right] = \frac{1}{2}$ $\frac{1}{2}(n-1)!$ agrees with the counting of linearly independent amplitudes following from (3.4). Note that (3.7) implies the parity relation in (3.4) as

$$
\sum_{k \text{ even}} \frac{1}{k!} A^{\text{string}}(\gamma_{1|P_1,...,P_k}) = A^{\text{string}}(1, 2, ..., n) - (-1)^n A^{\text{string}}(1, n, n-1, ..., 2).
$$
 (3.8)

Based on explicit computations, we find that motivic MZV amplitudes with different f_2^m content^5 satisfy the following KK-like relations:

$$
A_{f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 1,
$$

\n
$$
A_{f_2f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 3
$$

\n
$$
A_{f_2^m f_M}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 1, 3, 5,..., 2m + 1, \quad m \ge 2,
$$
\n
$$
(3.9)
$$

which constitute the *descent algebra decomposition of KK-like relations*. To count the basis dimensions implied by (3.9) we recall that $\#(\gamma_{1|P_1,...,P_k}) = \begin{bmatrix} n-1 \\ k \end{bmatrix}$ $\begin{bmatrix} -1 \\ k \end{bmatrix}$ and $\sum_{k=1}^{n-1} \begin{bmatrix} n-1 \\ k \end{bmatrix}$ $\left[\begin{smallmatrix} -1 \ k \end{smallmatrix}\right]=$

 5 The string monodromy relations [1,2] give rise to deformations of the field-theory BCJ relations by powers of $\alpha'^{2m} f_2^m$ [33,37]. Since the BCJ-satisfying f_M components lead to the minimum $(n-3)!$ dimension, they are not expected to modify the dimensions within a given $f_2^m f_M$ class.

 $(n-1)!$. Thus, subtracting the dimensions of the BRST-invariant permutations from the number of cyclically symmetric *n*-point amplitudes⁶ leads to

$$
#(A_{f_M}(\gamma_{1|P_1,\ldots,P_k})) = \begin{bmatrix} n-1 \\ 1 \end{bmatrix} = (n-2)!,
$$
\n(3.10)

$$
#(A_{f_2f_M}(\gamma_{1|P_1,\ldots,P_k})) = \begin{bmatrix} n-1 \\ 3 \end{bmatrix},
$$
\n(3.11)

$$
\#(A_{f_2^m f_M}(\gamma_{1|P_1,\ldots,P_k})) = \begin{bmatrix} n-1 \\ 1 \end{bmatrix} + \begin{bmatrix} n-1 \\ 3 \end{bmatrix} + \cdots + \begin{bmatrix} n-1 \\ 2m+1 \end{bmatrix}.
$$
 (3.12)

The dimension (3.10) corresponds to the number of independent amplitudes under the Kleiss-Kuijf relations $A^{SYM}(P1Q, n) - (-1)^{|P|}A^{SYM}(1, \tilde{P}\sqcup Q, n) = 0$, which is also valid for the BCJ-satisfying f_M corrections. The dimension (3.11) was obtained in [18] following a similar discussion of the all-plus one-loop amplitudes of [4] (see also [38]).

The basis dimension formula (3.12) and the corresponding amplitude relations in (3.9) are new. Interestingly, they imply that the basis dimension of the $f_2^m f_M$ components with $m \geq 2$ is given by $\frac{1}{2}(n-1)!$ only when $n \leq 2m+3$. More explicitly,

$$
#(A_{f_2^m f_M}(1,2,\ldots,n)) = \frac{1}{2}(n-1)!, \quad n = 4,5,\ldots,2m+3.
$$
 (3.13)

For a deviation from the $\frac{1}{2}(n-1)!$ dimension, we have, for example with $m=2$

$$
\#(A_{f_2^2 f_M}(1,2,\ldots,8)) = \begin{bmatrix} 7 \\ 1 \end{bmatrix} + \begin{bmatrix} 7 \\ 3 \end{bmatrix} + \begin{bmatrix} 7 \\ 5 \end{bmatrix} = 2519 = \frac{1}{2}7! - 1, \tag{3.14}
$$

which was confirmed by a long brute-force search using **FORM** [26]. The additional relation on top of (3.7) is seen to be $A_{f_2^2}(\gamma_{1|2,3,4,5,6,7,8}) = A_{f_2^2}(1,2 \sqcup 3 \sqcup \ldots \sqcup 8) = 0$, in agreement with the analysis of [33,39]. At nine points and $m = 2$, the formula (3.12) predicts a mismatch due to $\begin{bmatrix} 8 \\ 7 \end{bmatrix}$ $\binom{8}{7} = 28$ additional relations etc.

The evidence for the descent algebra decomposition of KK-like relations in (3.9) was collected from explicit calculations of $A_{f_2^n f_M}(\gamma_{1|P_1,\ldots,P_k})$ for various compositions $p = (p_1, \ldots, p_k) \models n-1$ with $|P_i| = p_i$. The results are summarized in Table 1⁷. The number of checks of $A_{f_2^m f_M}(\gamma_{1|P_1,...,P_k})$ at n points would appear to grow rapidly, but

⁶ Note that $\frac{1}{2}(n-1)!$ is equal to the sum over $\binom{n-1}{k}$ $\binom{-1}{k}$ with even k. Since k even is included in (3.9), the Stirling cycle numbers are subtracted from $(n-1)!$.

⁷ This data was collected using the α' corrections to disk amplitudes obtained in [28,40,41,29] (see also [42,43,44,45] and references therein for earlier work and [46] for a discussion on MZVs).

κ	$(A^{\text{string}}(\gamma_{1 P_1,,P_k}))=0$	$\Lambda^{\text{string}}(\gamma_{1 P_1,,P_k})\neq 0$
$\overline{7}$	$f_7, f_5, f_3, f_3^2, f_2, f_2f_5, f_2f_3, f_2^2, f_2^2f_3$	f_2^3
6	$\forall f_M$	
5	$f_7, f_5, f_3, f_3^2, f_2, f_2f_5, f_2f_3$	$f_2^2, f_2^2 f_3, f_2^3$
4	$\forall f_M$	
3	f_7, f_5, f_3, f_3^2	$f_2, f_2f_5, f_2f_3, f_2^2, f_2^2f_3, f_2^3$
$\mathcal{D}_{\mathcal{L}}$	$\forall f_M$	
	f_2, f_2f_3, f_2f_5	$f_7, f_5, f_3, f_3^2, f_2^2, f_2^2 f_3, f_2^3$

Table 1. Overview of the descent algebra symmetries of higher α' corrections to string disk amplitudes of up to $n = 8$ points displayed by their motivic MZV content of weight $w \le 7$. The entries depend only on the number of parts k of the composition of $n-1$. However, a partition with k parts cannot be probed by disk amplitudes with fewer than $k+1$ points.

luckily a vanishing outcome was observed to depend only on the number of parts k of the composition, independently of n (with data up to $n = 8$). Thus it suffices to test the single case $k = n-1$ for each additional external leg. Since $\gamma_{1|2,3,...,n} = 1.(2 \sqcup 3 \sqcup ... \sqcup n)$ by (2.20) , we get a sum over all cyclic orderings of the *n*-point string disk amplitude (3.2) , leading to the α' -corrected abelian Z-theory amplitudes $A^{\text{string}}(\gamma_{1|2,3,...,n}) \sim A^{\text{NLSM}}(1,2,...,n)$ of [33,39]. As a consistency check, the proof (D.1) implying that $A_{f_2^m f_M}(\gamma_{1|2,\ldots,k})$ vanishes when k is even (so n is odd) agrees with the vanishing of NLSM odd-point amplitudes.

3.2. The field-theory and α'^2 corrections

The SYM amplitudes are computed in pure spinor superspace from the expression M_1E_P , where E_P is a superfield satisfying the same shuffle symmetry $E_{R\sqcup S} = 0$ for $R, S \neq \emptyset$ of the standard Berends-Giele current J_P^m [47,48]. The A_{f_2} amplitudes [49] can be computed in pure spinor superspace [18] using BRST-closed combinations of superfields $C_{1|X,Y,Z}$ symmetric under exchanges of any pairs $X \leftrightarrow Y, Z$ and satisfying $C_{1|R\sqcup S,Y,Z} = 0$ for $R, S \neq \emptyset$. For convenience, define the BRST-closed combination $C_{1|P} := M_1E_P$ so that⁸

$$
A^{\text{SYM}}(1,2,\ldots,n) = \langle C_{1|2\ldots n} \rangle, \qquad (3.15)
$$

$$
A_{f_2}(1,2,\ldots,n) = \sum_{2\ldots n} \langle C_{1|X,Y,Z} \rangle, \qquad (3.16)
$$

where $\sum_{2...n}$ is a shorthand for the sum over the deconcatenations of $2...n = XYZ$. In terms of these BRST invariants, the color-dressed amplitudes at four and five points can

⁸ The angular brackets $\langle \ldots \rangle$ denotes the pure spinor zero-mode integration of [50], but it plays no role in the subsequent discussions.

be written as [18]

$$
M_{4} = 6f_{2}\alpha'^{2}d^{1234}C_{1|2,3,4}
$$
\n
$$
+ i^{2}d^{1a}F_{a}^{234}C_{1|234} + i^{2}d^{1a}F_{a}^{243}C_{1|243} + \mathcal{O}(\alpha'^{3}),
$$
\n
$$
M_{5} = 6f_{2}\alpha'^{2}id^{1abc}\left(F_{a}^{23}F_{b}^{4}F_{c}^{5}C_{1|23,4,5} + F_{a}^{24}F_{b}^{3}F_{c}^{5}C_{1|24,3,5} + F_{a}^{25}F_{b}^{3}F_{c}^{4}C_{1|25,3,4}\right)
$$
\n
$$
+ F_{a}^{2}F_{b}^{34}F_{c}^{5}C_{1|2,34,5} + F_{a}^{2}F_{b}^{35}F_{c}^{4}C_{1|2,35,4} + F_{a}^{2}F_{b}^{3}F_{c}^{45}C_{1|2,3,45}\right)
$$
\n
$$
+ i^{3}d^{1a}F_{a}^{2345}C_{1|2345} + i^{3}d^{1a}F_{a}^{2354}C_{1|2354} + i^{3}d^{1a}F_{a}^{2435}C_{1|2435}
$$
\n
$$
+ i^{3}d^{1a}F_{a}^{2453}C_{1|2453} + i^{3}d^{1a}F_{a}^{2534}C_{1|2534} + i^{3}d^{1a}F_{a}^{2543}C_{1|2543} + \mathcal{O}(\alpha'^{3}),
$$
\n
$$
(3.17)
$$
\n
$$
(3.17)
$$

with similar expansions at higher points. Comparing (3.17) with the color-dressed permutations P_4 and P_5 in (2.7) and (2.9) implies the correspondences⁹ at the given α' order,

$$
C_{1|X} \longleftrightarrow \gamma_{1|X}, \qquad C_{1|X,Y,Z} \longleftrightarrow \frac{1}{6} \gamma_{1|X,Y,Z}. \tag{3.18}
$$

So the A^{SYM} and A_{f_2} amplitudes correspond to idempotents of the descent algebra

$$
A^{\text{SYM}}(1,2,\ldots,n) \longleftrightarrow \gamma_{123\ldots n}^{(1)}, \qquad A_{f_2}(1,2,\ldots,n) \iff \gamma_{123\ldots n}^{(3)}.\tag{3.19}
$$

where the deconcatenations (3.16) and (2.22) have been used in the last line. The correspondences in (3.19) can be used to justify the first two relations in (3.4) from the theorem 4.2 of [9], namely $E_{\mu} \circ I_{p} = 0$ if $\lambda(p) \neq \mu$, where E_{μ} for a partition μ is reviewed in (A.30), p is a composition and $\lambda(p)$ is its shape as defined in the beginning of section A.3. Under the dualities (3.19) this identity implies the relations¹⁰

$$
A^{\text{SYM}}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 1, \qquad A_{f_2}(\gamma_{1|P_1,...,P_k}) = 0, \quad k \neq 3. \tag{3.20}
$$

⁹ The first correspondence in (3.18) implies $M_1E_P \leftrightarrow 1 \cdot \mathcal{E}_P$ and suggests the duality $E_P \leftrightarrow \mathcal{E}_P$ (E_P) here is the superfield of [51], not the Eulerian idempotent). Since the superfield E_P is related to the Berends-Giele current M_P [51], this motivates the terminology of \mathcal{E}_P in (2.12).

¹⁰ To see this we use that $P \cdot (\sigma \circ \tau) = ((P \cdot \sigma) \circ (P \cdot \tau))$ if P has no common letters with σ and τ and $\theta(\sigma \circ \tau) = \theta(\tau) \circ \theta(\sigma)$ to show that the theorem 4.2 of [9] implies $(1 \cdot \mathcal{I}_p) \circ (1 \cdot E_\mu^{\theta}) = 0$ for $\lambda(p) \neq \mu$ leading to (3.20) since A_{f_2} corresponds to a sum of $\theta(E_\mu)$ with partitions with three parts $k(\mu) = 3$ and A^{SYM} to $\theta(E_{\mu})$ with $k(\mu) = 1$.

3.3. BRST-invariant permutations and BRST-invariant superfields

The linearity condition $M_n = \phi(A^{\text{string}}(P_n))$ implies that BRST-invariant permutations in P_n are mapped to kinematics in M_n leading to a correspondence between the BRSTinvariant permutations (2.4) and a series of motivic MZV corrections from the motivic string disk amplitude in the f-alphabet

$$
\gamma_{1|P_1, P_2, \dots, P_k} \leftrightarrow \phi\big(A^{\text{string}}(\gamma_{1|P_1, P_2, \dots, P_k})\big). \tag{3.21}
$$

whose precise content follows from Table 1 and (3.2). For example,

$$
\gamma_{1|234} \leftrightarrow A^{\text{SYM}}(1,2,3,4) + f_3 \alpha'^3 s_{12} s_{23} (s_{12} + s_{23}) A^{\text{SYM}}(1,2,3,4) + \cdots \qquad (3.22)
$$

\n
$$
\gamma_{1|2,3,4} \leftrightarrow -6 f_2 \alpha'^2 s_{12} s_{23} A^{\text{SYM}}(1,2,3,4) -3 f_2^2 \alpha'^4 (s_{12}^3 s_{23} + s_{12}^2 s_{23}^2 + s_{12} s_{23}^3) A^{\text{SYM}}(1,2,3,4) + \cdots
$$

with similar expansions at higher points. The data presented in Table 1 and the discussion in section 3.2 suggest that the BRST-invariant permutations can be associated to a series of higher-mass BRST-invariant superfields by defining

$$
A_{f_2^m f_M}(\gamma_{1|P_1,\dots,P_k}) = k! C_{1|P_1,\dots,P_k}^{f_2^m f_M}.
$$
\n(3.23)

The motivic MZV amplitudes then follow from

$$
A_{f_2^m f_M}(1,2,3,\ldots,n) = \sum_{k=1}^{2m+1} \sum_{23\ldots n} C_{1|P_1,\ldots,P_k}^{f_2^m f_M},
$$
\n(3.24)

where $\sum_{23...n}$ is a shorthand notation for the deconcatenations of $P_1P_2...P_k = 23...n$. For example,

$$
A_{f_2^2}(1,2,3,4) = C_{1|234}^{f_2^2} + C_{1|2,3,4}^{f_2^2} = -\frac{2}{5}(s_{12}s_{23}^3 + \frac{1}{4}s_{12}^2s_{23}^2 + s_{12}^3s_{23})A^{\text{SYM}}(1,2,3,4),
$$
 (3.25)

where the BRST invariants $C_{1|234}^{f_2^2} = 1/10(s_{12}s_{23}^3 + 4s_{12}^2s_{23}^2 + s_{12}^3s_{23})A^{\text{SYM}}(1,2,3,4)$ and $C_{1[2,3,4}^{f_2^2} = -1/2(s_{12}s_{23}^3 + s_{12}^2s_{23}^2 + s_{12}^3s_{23})A^{\text{SYM}}(1,2,3,4)$ were used. The superfield representation of the above BRST invariants is known only for the two simplest cases, $C_{1|P} = M_1 E_P$ [51] and $C_{1}^{f_2}$ $I_{1|P_1,P_2,P_3}^{f_2}$ [19,52]. Note that $C_{1|P_1} = A^{\text{SYM}}(1,P_1)$ while the S-map algorithm of [19] gives rise to a purely combinatorial translation between $C_{1}^{f_2}$ $1|P_1,P_2,P_3|$ and sums of $s_{ij}^2 A^{\text{SYM}}$. It remains to be seen whether there exists a general algorithm to rewrite $C^{f_2^m f_M}$ in terms of SYM amplitudes.

3.4. BRST invariants from A_{f_2}

Another consequence of the duality (3.18) is that the representation of A_{f_2} in terms of $C_{1|P_1,P_2,P_3}$ given in (3.16) is invertible, as argued indirectly in [18]. This follows from Theorem 4.2 of [9], $E_{\mu} \circ I_{p} = I_{p}$ if $\lambda(p) = \mu$ where $\lambda(p)$ is the shape of the composition p and E_{μ} is defined in (A.30). This implies $(1 \cdot \mathcal{I}_{p}) \circ (1 \cdot \theta(E_{\mu})) = (1 \cdot \mathcal{I}_{p})$ for $\lambda(p) = \mu$ or, using the function interpretation of the right action $\sigma \circ F := F(\sigma)$ with a partition with three parts $k(\mu) = 3$

$$
A_{f_2}(\gamma_{1|P_1, P_2, P_3}) = 6C_{1|P_1, P_2, P_3}, \quad |P_i| = p_i, \quad P_1 P_2 P_3 = 23...n, \tag{3.26}
$$

where we used the identifications (2.20) , (2.22) and duality (3.19) on the left-hand side and the duality (3.18) on the right-hand side. For example, from $\gamma_{1|23,4,5}$ of (C.1) we get

$$
6C_{1|23,4,5} = A_{12345}^{f_2} + A_{12354}^{f_2} + A_{12435}^{f_2} + A_{12453}^{f_2} + A_{12534}^{f_2} + A_{12543}^{f_2}
$$
\n
$$
- A_{13245}^{f_2} - A_{13254}^{f_2} - A_{13425}^{f_2} - A_{13524}^{f_2} + A_{14235}^{f_2} - A_{14325}^{f_2},
$$
\n
$$
(3.27)
$$

where we used the parity relation (3.4) in the RHS.

3.5. The superfield expansion of $C_{1|P,Q,R}$ from BRST-invariant permutations

The so-called BRST invariants $C_{1|P,Q,R}$ of the pure spinor formalism [50] play an important role in the mapping between BRST-invariant permutations and kinematics, see section 3.3. They were firstly derived at low multiplicities in [18] and were subsequently studied in different contexts and given general recursive algorithms, see [19,52,53]. Their superfield expansions in terms of Berends-Giele currents follow from

$$
C_{i|P,Q,R} = M_i M_{P,Q,R} + M_i \cdot \left[C_{p_1|p_2...p_{|P|},Q,R} - C_{p_{|P|}|p_1...p_{|P|-1},Q,R} + (P \leftrightarrow Q,R) \right] \tag{3.28}
$$

starting from $C_{i|j,k,l} = M_i M_{j,k,l}$ with the dot representing concatenation, $M_i \cdot M_A := M_{iA}$. For example, the first few expansions are given by

$$
C_{1|2,3,4} = M_1 M_{2,3,4},
$$
\n
$$
C_{1|23,4,5} = M_1 M_{23,4,5} + M_{12} M_{3,4,5} - M_{13} M_{2,4,5},
$$
\n
$$
C_{1|234,5,6} = M_1 M_{234,5,6} + M_{12} M_{34,5,6} + M_{123} M_{4,5,6} - M_{124} M_{3,5,6}
$$
\n
$$
- M_{14} M_{23,5,6} - M_{142} M_{3,5,6} + M_{143} M_{2,5,6},
$$
\n
$$
C_{1|23,45,6} = M_1 M_{23,45,6} + M_{12} M_{45,3,6} - M_{13} M_{45,2,6} + M_{14} M_{23,5,6} - M_{15} M_{23,4,6}
$$
\n
$$
+ M_{124} M_{3,5,6} - M_{134} M_{2,5,6} + M_{142} M_{3,5,6} - M_{152} M_{3,4,6}
$$
\n
$$
- M_{125} M_{3,4,6} + M_{135} M_{2,4,6} - M_{143} M_{2,5,6} + M_{153} M_{2,4,6}.
$$
\n(3.29)

These terms can be extracted from the permutations of the BRST-invariant permutations $\gamma_{1|P,Q,R}$ of the inverse descent algebra as follows:

1. Sum over the cyclic permutations of all permutations in $\gamma_{1|P,Q,R}$:

$$
W_{\sigma} \to W_{\sigma} + \text{cyclic}(\sigma) \tag{3.30}
$$

2. Decompose W_{σ} into all possible four-word deconcatenations:

$$
W_{\sigma} = \sum_{XYZW = \sigma} W_X . W_Y . W_Z . W_W \tag{3.31}
$$

3. Move label 1 to the front by repeatedly commuting $W_C.W_{A1B} = W_{A1B}.W_C$ if necessary and write the result in terms of Berends-Giele superfields:

$$
W_{A1B}.W_C.W_D.W_E := \frac{1}{4!} M_{A1B} M_{C,D,E}
$$
\n(3.32)

The resulting expressions have been explicitly checked¹¹ for all topologies of BRST invariants up to eight points. In addition, using the descent duality (3.18) one may also derive the change of basis identities for $C_{i\neq1|...} = \sum C_{1|...}$ from [52,53] by choosing a different label to be singled-out in the color-dressed permutation (2.1) [54].

4. Conclusion

In this paper we investigated the combinatorial properties of the permutations appearing in the *color-dressed permutations* (2.1) using the tools from the descent algebra. In particular, we defined BRST-invariant permutations, found a closed formula, and related them to orthogonal idempotents which sum to the identity permutation [10,55,9].

We then considered the color-dressed motivic string disk amplitudes of [5] within this framework. This led to the discovery of the relations (3.9) obeyed by the motivic MZV amplitudes, dubbed the *descent algebra decomposition of KK-like relations*. The basis dimensions of linearly independent motivic MZV amplitudes are given by sums of Stirling cycle numbers. These claims have been explicitly checked using various data points in string theory up to $n = 8$ and ${\alpha'}^7$.

¹¹ The shuffle symmetry $AiB = (-1)^{|A|} i\tilde{A} \sqcup B$ [21] is needed to rewrite words in a Lyndon basis.

Inspired by [18], we proposed a correspondence between the permutations from the (inverse) descent algebra and kinematics from the motivic string disk amplitudes in terms of higher-mass BRST invariants. In the particular case of α'^2 , we exploited a theorem from the mathematics literature on descent algebra to prove certain claims in [18] and to systematically express BRST invariants as linear combinations of A_{f_2} corrections to motivic disk amplitudes in (3.26).

And finally, we found an algorithm to extract the superfield content of the BRST invariants in the pure spinor formalism from the BRST-invariant permutations in the inverse descent algebra. It will be interesting to obtain superfield realizations of the higher-mass BRST invariants defined in section 3.3. They can probably be extracted from a perturbiner series of amplitudes at the appropriate mass level. For instance, the superfields in $C_{1|P}^{\text{SYM}} = M_1 E_P$ are related to the series Tr(VVV) [56,27]. The superfields in $C_{1|P}^{f_2}$ $^{1/2}_{1|P_1,P_2,P_3}$ are related to the series $\text{Tr}(\mathbb{V}_1(\lambda \gamma^m \mathbb{W})(\lambda \gamma^n \mathbb{W})\mathbb{F}_{mn})$, while the superfields in $C_{1\perp}^{fs}$ $l_{1|P}^{J3}$ should follow from $\text{Tr}((\lambda \gamma^{mnpqr} \lambda)(\lambda \gamma^{s} \mathbb{W}) \mathbb{F}_{mn} \mathbb{F}_{pq} \mathbb{F}_{rs})$. It would also be interesting to find combinatorial algorithms to directly translate the higher-mass BRST invariants $C_{1|P_1}^{f_2^m f_M}$ $\prod_{1|P_1,...,P_k}^{J_2 J M}$ into linear combinations of super-Yang–Mills amplitudes and powers of Mandelstam invariants, generalizing the S-map algorithm of [19] for $C_{1}^{f_2}$ $\frac{J_{2}}{1|P_{1},P_{2},P_{3}}$. This would give rise to a combinatorial description of the P_n and M_n matrices of [5].

Acknowledgements: I thank Ruggero Bandiera for discussions during an early stage of this project. I also thank Oliver Schlotterer for sharing notes containing intriguing observations about the permutations in the color-dressed amplitude that served as the motivation for this paper, for discussions, for collaboration on related topics, and for comments on the draft (including suggesting a v3 to be uploaded). CRM is supported by a University Research Fellowship from the Royal Society.

Appendix A. The Solomon descent algebra

We review the salient features of the Solomon descent algebra [8,9,10,11,12,13,14]. In particular, we discuss different bases and highlight the orthogonal idempotents discovered by Reutenauer, as they will be related to α' corrections to string amplitudes.

A.1. Descent classes and the Solomon descent algebra

The descent set $D(\sigma)$ and the and the descent number d_{σ} of a permutation $\sigma = \sigma_1 \sigma_2 \ldots \sigma_n$ in S_n are defined by

$$
D(\sigma) = \{ i \in \{1, 2, \dots, n-1\} \mid \sigma_i > \sigma_{i+1} \}, \qquad d_{\sigma} = \#(D(\sigma)). \tag{A.1}
$$

For example, the permutation $\sigma = 546132$ has descent set $D(\sigma) = \{1, 3, 5\}$ and descent number $d_{\sigma} = 3$. The collection of permutations with a given descent set S is called a descent class,

$$
D_S = \sum_{D(\sigma) = S} \sigma \,. \tag{A.2}
$$

For example, the permutations in S_3 are distributed into four descent classes,

$$
D_{\emptyset} = W_{123}, \quad D_{\{1\}} = W_{213} + W_{312}, \quad D_{\{2\}} = W_{132} + W_{231}, \quad D_{\{1,2\}} = W_{321}. \tag{A.3}
$$

In general, the permutations of S_n decompose into 2^{n-1} distinct descent classes; all the subsets in the powerset of $\{1, 2, \ldots, n-1\}$ since the last *n*-th position is never a descent. Solomon showed the remarkable property that descent classes are closed under the right action (1.10)

$$
D_S \circ D_T = \sum_{U \subseteq \{1, 2, \dots, n-1\}} c_{S, T, U} D_U \tag{A.4}
$$

where the coefficients $c_{S,T,U}$ are non-negative integers [8]. The descent classes therefore form a 2^{n-1} dimensional algebra, the so-called Solomon's descent algebra \mathcal{D}_n $[8,9,11,12,13,14]$.

As an example of $(A.4)$, consider the permutations in S_4 . Its 24 elements are organized into 8 descent classes as follows

$$
D_{\{0\}} = W_{1234},
$$
\n
$$
D_{\{1,2\}} = W_{3214} + W_{4213} + W_{4312},
$$
\n
$$
D_{\{1\}} = W_{2134} + W_{3124} + W_{4123},
$$
\n
$$
D_{\{2\}} = W_{1324} + W_{1423} + W_{2314} + W_{2413} + W_{3412},
$$
\n
$$
D_{\{2,3\}} = W_{1432} + W_{2431} + W_{3421},
$$
\n
$$
D_{\{1,3\}} = W_{2143} + W_{3142} + W_{3241} + W_{4132} + W_{4231},
$$
\n
$$
D_{\{1,2,3\}} = W_{4321}.
$$
\n(A.5)

It is straightforward to multiply the permutations among these descent classes using the right-action of the symmetric group (1.10). For example,

$$
D_{\{1\}} \circ D_{\{2\}} = W_{1234} + W_{1243} + W_{1324} + W_{1342} + W_{1423} + W_{1432} + W_{2314}
$$
\n
$$
+ W_{2341} + W_{2413} + W_{2431} + W_{3214} + W_{3412} + W_{3421} + W_{4213} + W_{4312}
$$
\n
$$
= D_{\{\emptyset\}} + D_{\{1,2\}} + D_{\{2\}} + D_{\{2,3\}} + D_{\{3\}},
$$
\n(A.6)

where the last line follows from the remarkable property (A.4) which ensures that the permutations in (A.6) are themselves a sum of descent classes.

A.2. Bases of the descent algebra

Apart from the descent classes D_S indexed by descent sets S, there are other convenient bases of the descent algebra [9].

A.2.1. Composition basis B_p

The composition p of n, denoted $p \models n$, is a k-tuple of positive integers with sum n,

$$
p = (p_1, p_2, \dots, p_k), \quad p_1 + p_2 + \dots + p_k = n.
$$
 (A.7)

There is a bijection between compositions $p \models n$ and subsets S of $\{1, 2, \ldots, n-1\}$

$$
p = (p_1, p_2, \dots, p_k) \mapsto \{p_1, p_1 + p_2, \dots, p_1 + p_2 + \dots + p_{k-1}\} := S(p), \quad (A.8)
$$

$$
S = \{i_1, i_2, \dots, i_k\} \mapsto (i_1, i_2 - i_1, \dots, i_k - i_{k-1}, n - i_k) := C_n(S).
$$
 (A.9)

Thus the total number of compositions of n is 2^{n-1} , the cardinality of the powerset of $\{1, 2, \ldots, n-1\}$. Note that the map $C_n(S) = p$ depends on the order n of the permutation group S_n for $C_3({1, 2}) = (1, 1, 1)$ but $C_4({1, 2}) = (1, 1, 2)$. In particular, $C_n(\emptyset) = (n)$.

The basis B_p is indexed by compositions p rather than subsets and is defined by [9]

$$
B_p = D_{\subseteq S(p)}\,,\tag{A.10}
$$

with $S(p)$ given by (A.8). For example, the D_S basis elements (A.5) become

$$
B_{1111} = D_{\emptyset} + D_{\{1\}} + D_{\{2\}} + D_{\{3\}} + D_{\{1,2\}} \t B_{13} = D_{\emptyset} + D_{\{1\}},
$$

\n
$$
+ D_{\{1,3\}} + D_{\{2,3\}} + D_{\{1,2,3\}},
$$

\n
$$
B_{112} = D_{\emptyset} + D_{\{1\}} + D_{\{2\}} + D_{\{1,2\}},
$$

\n
$$
B_{121} = D_{\emptyset} + D_{\{1\}} + D_{\{3\}} + D_{\{1,3\}},
$$

\n
$$
B_{21} = D_{\emptyset} + D_{\{1\}} + D_{\{3\}} + D_{\{1,3\}},
$$

\n
$$
B_{31} = D_{\emptyset} + D_{\{3\}},
$$

\n
$$
B_{4} = D_{\emptyset}.
$$

\n(A.11)

The inverse of (A.10) is given by Lemma 8.18 in [14]

$$
D_S = \sum_{T \subseteq S} (-1)^{|S| - |T|} D_{\subseteq T} . \tag{A.12}
$$

For example (in S_4), $D_{\{1,2\}} = B_{112} - B_{13} - B_{22} + B_4$, $D_{\{1\}} = B_{13} - B_4$, $D_{\{2\}} = B_{22} - B_4$, and $D_{\emptyset} = B_4$, from which we verify that $D_{\emptyset} + D_{\{1\}} + D_{\{2\}} + D_{\{1,2\}} = B_{112}$.

The permutations within a basis element B_p can be found via [55,9]

$$
B_{p_1p_2...p_k} = \theta(X_1 \sqcup X_2 \sqcup ... \sqcup X_k), \quad 12...n = X_1...X_k, \quad |X_i| = p_i.
$$
 (A.13)

where the inverse map θ is given by

$$
\theta(\sigma) \mapsto \sigma^{-1} \,. \tag{A.14}
$$

For example, if $p = (1, 1, 2)$ then $X_1 = 1$, $X_2 = 2$ and $X_3 = 34$ and we get

$$
B_{112} = \theta(1 \sqcup 2 \sqcup 34) = W_{1234} + W_{1324} + W_{1423} + W_{2134} + W_{2314} + W_{2413} \quad (A.15)
$$

$$
+ W_{3124} + W_{3214} + W_{3412} + W_{4123} + W_{4213} + W_{4312}.
$$

A.2.2. Multiplication table for $B_p \circ B_q$

There is a closed formula for the multiplication of $B_p \circ B_q$ [12,9,57]. Let M be a matrix with non-negative integer entries m_{ij} whose row sum $r(M)$ and column sum $c(M)$ are vectors defined by

$$
r(M)_i := \sum_j m_{ij}, \qquad c(M)_j := \sum_i m_{ij}.
$$
 (A.16)

Then

$$
B_p \circ B_q = \sum_{\substack{c(M) = p \\ r(M) = q}} B_{\text{co}(M)} \tag{A.17}
$$

where $\text{co}(M)$ denotes the composition obtained by reading the matrix M row by row from top to bottom while excluding the zero entries $m_{ij} = 0$. This product is associative and B_n is a multiplicative identity for compositions of n [57].

For example, let us recover the result (A.6) for $D_{\{1\}} \circ D_{\{2\}}$ using the above multiplication table (A.17) in S_4 . Given that $D_{\{1\}} = B_{13} - B_4$ and $D_{\{2\}} = B_{22} - B_4$, the only non-trivial product we need is $B_{13} \circ B_{22}$ since B_4 is the identity for compositions of $n = 4$. The set of integer matrices M with $c(M) = (1, 3)$ and $r(M) = (2, 2)$ is given by

$$
\begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}, \qquad \begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix}. \tag{A.18}
$$

Thus $B_{13} \circ B_{22} = B_{112} + B_{211}$ and $D_{\{1\}} \circ D_{\{2\}} = (B_{13} - B_4) \circ (B_{22} - B_4)$ implies

$$
D_{\{1\}} \circ D_{\{2\}} = B_{112} + B_{211} - B_{13} - B_{22} + B_4 = D_{\emptyset} + D_{\{1,2\}} + D_{\{2\}} + D_{\{2,3\}} + D_{\{3\}} \quad (A.19)
$$

where we used the conversions $(A.11)$.

A.2.3. The Eulerian idempotent

The Eulerian (or Solomon) idempotent is defined by [17,10,55,58] (see also [59])

$$
E_n = \sum_{\sigma \in S_n} \kappa_{\sigma} \sigma, \qquad \kappa_{\sigma} = \frac{(-1)^{d_{\sigma}}}{|\sigma| \binom{|\sigma| - 1}{d_{\sigma}}}
$$
(A.20)

where d_{σ} denotes the descent number (A.1) of the permutation σ . For example,

$$
E_2 = \frac{1}{2}(W_{12} - W_{21}), \quad E_3 = \frac{1}{3}W_{123} - \frac{1}{6}W_{132} - \frac{1}{6}W_{213} - \frac{1}{6}W_{231} - \frac{1}{6}W_{312} + \frac{1}{3}W_{321}.
$$
 (A.21)

Apart from being an idempotent satisfying $E_n \circ E_n = E_n$, the definition (A.20) is also a Lie polynomial [10]. Therefore its coefficients κ_{σ} must satisfy the shuffle symmetry [23]

$$
\kappa_{R \sqcup S} = 0, \quad R, S \neq \emptyset. \tag{A.22}
$$

As usual, the definition (A.20) in terms of the fixed alphabet N in S_n can be turned into a function of an arbitrary word P by the right action (1.10) of the symmetric group $[14,60]$,

$$
E(P) = EP := P \circ E_n, \quad |P| = n.
$$
 (A.23)

For example, $E(i, j, k) = ijk \circ E_3 = \frac{1}{3}W_{ijk} - \frac{1}{6}W_{ikj} - \frac{1}{6}W_{jik} - \frac{1}{6}W_{jki} - \frac{1}{6}W_{kij} + \frac{1}{3}W_{kji}$.

A.2.4. The idempotent basis I_p

The idempotent basis I_p of the descent algebra \mathcal{D}_n satisfying $I_p \circ I_p = I_p$ was introduced in [9] and it is indexed by the compositions of n

$$
I_{p_1p_2...p_k}(P) = \sum_{\substack{x_1,\ldots,x_k\\|x_i|=p_i}} \langle P, X_1 \sqcup X_2 \sqcup \ldots \sqcup X_k \rangle E^{X_1} E^{X_2} \ldots E^{X_k},
$$
 (A.24)

where the sum is constrained by the length of X_i being equal to the corresponding p_i in the composition p and E^{X_i} denote the Eulerian idempotent function (A.23). For example, with canonical $P = 12...n$ we have

$$
I_{11} = W_{12} + W_{21}, \t I_2 = \frac{1}{2}(W_{12} - W_{21}),
$$
\n
$$
I_{111} = W_{123} + W_{132} + W_{213} + W_{231} + W_{312} + W_{321},
$$
\n
$$
I_{21} = \frac{1}{2}W_{123} + \frac{1}{2}W_{132} - \frac{1}{2}W_{213} + \frac{1}{2}W_{231} - \frac{1}{2}W_{312} - \frac{1}{2}W_{321},
$$
\n
$$
I_{12} = \frac{1}{2}W_{123} - \frac{1}{2}W_{132} + \frac{1}{2}W_{213} - \frac{1}{2}W_{231} + \frac{1}{2}W_{312} - \frac{1}{2}W_{321},
$$
\n
$$
I_3 = \frac{1}{3}W_{123} - \frac{1}{6}W_{132} - \frac{1}{6}W_{213} - \frac{1}{6}W_{231} - \frac{1}{6}W_{312} + \frac{1}{3}W_{321}.
$$
\n(A.25)

A.2.5. I_p to B_p

The idempotent basis elements I_p for $p = p_1p_2 \ldots p_k$ can be expanded in terms of compositions B_q using an algorithm discussed in [9]. First one defines *moments* e_m as a polynomial in non-commuting variables t_i for $i = 1, 2, \ldots$ from the generating series

$$
\sum x^n e_n = \log(1 + \sum t_i x^i)
$$
\n(A.26)

where x is a commuting parameter. For example, from $(A.26)$ it follows that

$$
e_1 = t_1, \quad e_2 = t_2 - \frac{1}{2}t_1^2, \quad e_3 = t_3 - \frac{1}{2}(t_1t_2 + t_2t_1) + \frac{1}{3}t_1^3
$$
\n
$$
e_4 = -\frac{1}{4}t_1^4 + \frac{1}{3}t_1^2t_2 + \frac{1}{3}t_1t_2t_1 - \frac{1}{2}t_1t_3 + \frac{1}{3}t_2t_1^2 - \frac{1}{2}t_2^2 - \frac{1}{2}t_3t_1 + t_4
$$
\n(A.27)

Then to convert the I_p basis elements to the composition basis B_q one uses [9]

$$
I_p = \delta(e_{p_1}e_{p_2}\dots e_{p_k}), \text{ with } \delta(t_{i_1}t_{i_2}\dots t_{i_k}) := B_{i_1i_2\dots i_k}.
$$
 (A.28)

For example,

$$
I_4 = -\frac{1}{4}B_{1111} + \frac{1}{3}B_{112} + \frac{1}{3}B_{121} - \frac{1}{2}B_{13} + \frac{1}{3}B_{211} - \frac{1}{2}B_{22} - \frac{1}{2}B_{31} + B_4.
$$
 (A.29)

A.3. Reutenauer orthogonal idempotents

A partition λ of n, denoted $\lambda \vdash n$, is a k-tuple of positive integers with sum n satisfying $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_k$. If $p \models n$ is a composition of n, the shape $\lambda(p)$ of p is the partition of n obtained by rearranging the parts of p in decreasing order. Also, $k(p)$ is the number of parts of the composition p. For example, $p = (2, 3, 1, 2)$ implies $\lambda(p) = 3221$ and $k(p) = 4$.

Given a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ into k parts, theorem 3.1 of [9] shows that

$$
E_{\lambda} := \frac{1}{k!} \sum_{\lambda(p)=\lambda} I_p, \qquad \sum_{\lambda \vdash n} E_{\lambda} = W_{12...n}.
$$
 (A.30)

Note that when the partition λ of n has only one part, $E_{\lambda} = I_n$ coincides with the Eulerian idempotent E_n (A.20), so this notation is not ambiguous. For example, $E_1 = I_1$ and

$$
E_2 = I_2, \t E_3 = I_3, \t E_{111} = \frac{1}{3!} I_{111},
$$

\n
$$
E_{11} = \frac{1}{2} I_{11}, \t E_{21} = \frac{1}{2} (I_{12} + I_{21}), \t E_{211} = \frac{1}{3!} (I_{112} + I_{121} + I_{211}).
$$
\n(A.31)

one can readily verify $E_3 + E_{21} + E_{111} = W_{123}$ using the expansions listed in the appendix C.

The Reutenauer idempotents $E^{(m)}$ are defined in the alphabet $\{1, 2, ...\}$ as the sum over all permutations of E_{λ} from (A.30) such that λ is a partition of n with m parts, i.e.,

$$
E^{(m)} = \sum_{\substack{\lambda \vdash n \\ k(\lambda) = m}} E_{\lambda} \tag{A.32}
$$

For example,

$$
n = 2 \t E^{(1)} = E_2, \t E^{(2)} = E_{11}
$$

\n
$$
n = 3 \t E^{(1)} = E_3, \t E^{(2)} = E_{21}, \t E^{(3)} = E_{111}
$$

\n
$$
n = 4 \t E^{(1)} = E_4, \t E^{(2)} = E_{31} + E_{22}, \t E^{(3)} = E_{211}, \t E^{(4)} = E_{1111}
$$
\n(A.33)

It was shown in [9,10] that (A.32) are orthogonal idempotents which sum to the identity permutation

$$
\sum_{i=1}^{n} E^{(i)} = W_{123...n}, \qquad E^{(i)} \circ E^{(j)} = \begin{cases} E^{(i)} & \text{if } i = j; \\ 0 & \text{otherwise.} \end{cases}
$$
 (A.34)

An alternative definition of the Reutenauer idempotents in terms of a generating function can be found in [14].

Appendix B. The inverse of the idempotent basis

In this appendix we will prove (2.17), that is:

Proposition. The inverse of the idempotent basis (2.16) satisfies

$$
\mathcal{I}_{p_1p_2\ldots p_k}(P_1, P_2, \ldots, P_k) = \mathcal{E}(P_1) \sqcup \mathcal{E}(P_2) \sqcup \ldots \sqcup \mathcal{E}(P_k), \quad |P_i| = p_i \qquad (B.1)
$$

where $P = P_1 \dots P_k$ is the factorization of P with P_i of length p_i .

Proof. The proof will be based on the following observations collected from [14], which should be consulted for more details as the equation numbers below refer to it. First, the adjoint of an arbitrary function $F(P) = P \circ F$ of a word P is given by $\theta(F)(P) = P \circ \theta(F)$, see (3.3.5). Second, the adjoint of $F_{p_1} \star F_{p_2} \ldots \star F_{p_k}$ is $\theta(F_{p_1}) \star' \theta(F_{p_2}) \ldots \star' \theta(F_{p_k})$ where \star and \star' are the convolution operators defined in (1.5.7) and (1.5.8) and $\theta(F_j)$ is the adjoint of F_j when viewed as a function by the right-action (1.10), see proof of Lemma 3.13. Third, for permutations F_{p_i} of length p_i one can show (by adapting the proof of Lemma 3.13)

$$
(F_{p_1} \star' \ldots \star' F_{p_k})(P) = F_{p_1}(P_1) \sqcup \ldots \sqcup F_{p_k}(P_k)
$$
\n(B.2)

where the functions are defined via a right action as $F_{p_i}(P_i) := P_i \circ F_{p_i}$. The proof of (B.1) then follows from the observation by (1.5.4) and (1.5.7) that the idempotent basis I_p (A.24) can be rewritten as a convolution $I_{p_1...p_k}(P) = (E_{p_1} \star ... \star E_{p_k})(P)$ where E_p is the Eulerian idempotent (A.20). Therefore its adjoint $\theta(I_{p_1...p_k})(P)$ is given by

$$
\theta(I_{p_1\ldots p_k})(P) = \theta(E_{p_1})(P_1) \star' \ldots \star' \theta(E_{p_k})(P_k), \quad P = P_1 \ldots P_k, \quad |P_i| = p_i
$$

= $\mathcal{E}(P_1) \sqcup \ldots \sqcup \mathcal{E}(P_k)$ (B.3)

where we used (B.2) and $\mathcal{E}(P_i) = \theta(E_{p_i})(P_i)$.

As a multiplicity-four example of the inverse idempotent basis of (2.16) we have

$$
\mathcal{I}_{22}(12,34) = \frac{1}{4}W_{1234} - \frac{1}{4}W_{1243} + \frac{1}{4}W_{1324} + \frac{1}{4}W_{1342} - \frac{1}{4}W_{1423} - \frac{1}{4}W_{1432}
$$
\n
$$
- \frac{1}{4}W_{2134} + \frac{1}{4}W_{2143} - \frac{1}{4}W_{2314} - \frac{1}{4}W_{2341} + \frac{1}{4}W_{2413} + \frac{1}{4}W_{2431}
$$
\n
$$
+ \frac{1}{4}W_{3124} + \frac{1}{4}W_{3142} - \frac{1}{4}W_{3214} - \frac{1}{4}W_{3241} + \frac{1}{4}W_{3412} - \frac{1}{4}W_{3421}
$$
\n
$$
- \frac{1}{4}W_{4123} - \frac{1}{4}W_{4132} + \frac{1}{4}W_{4213} + \frac{1}{4}W_{4231} - \frac{1}{4}W_{4312} + \frac{1}{4}W_{4321}.
$$
\n(B.4)

Appendix C. Explicit permutations at low multiplicities

The multiplicity-five BRST-invariant permutations (defined in (2.4)) are given by

$$
\gamma_{1|2,3,4,5} = W_{1(2 \text{u}3 \text{u}4 \text{u}5)} \qquad (C.1)
$$
\n
$$
\gamma_{1|23,4,5} = \frac{1}{2} W_{12345} + \frac{1}{2} W_{12354} + \frac{1}{2} W_{12435} + \frac{1}{2} W_{12453} + \frac{1}{2} W_{12534} + \frac{1}{2} W_{12543} - \frac{1}{2} W_{13245} - \frac{1}{2} W_{13254} - \frac{1}{2} W_{13425} - \frac{1}{2} W_{13452} - \frac{1}{2} W_{13524} - \frac{1}{2} W_{13542} + \frac{1}{2} W_{14235} + \frac{1}{2} W_{14253} - \frac{1}{2} W_{14325} - \frac{1}{2} W_{14352} + \frac{1}{2} W_{14523} - \frac{1}{2} W_{14532} + \frac{1}{2} W_{15234} + \frac{1}{2} W_{15243} - \frac{1}{2} W_{15324} - \frac{1}{2} W_{15342} + \frac{1}{2} W_{15423} - \frac{1}{2} W_{15432} - \frac{1}{6} W_{12435} - \frac{1}{6} W_{12435} + \frac{1}{3} W_{12534} - \frac{1}{6} W_{13245} - \frac{1}{6} W_{13245} - \frac{1}{6} W_{13452} - \frac{1}{6} W_{13524} - \frac{1}{6} W_{13542} - \frac{1}{6} W_{14235} - \frac{1}{6} W_{14235} - \frac{1}{6} W_{14235} + \frac{1}{3} W_{14325} + \frac{1}{3} W_{14352} - \frac{1}{6} W_{14523} + \frac{1}{3} W_{14532} + \frac{1}{
$$

 \Box

$$
\begin{split} \gamma_{1|23,45} &= \frac{1}{4}W_{12345} - \frac{1}{4}W_{12354} + \frac{1}{4}W_{12435} + \frac{1}{4}W_{12453} - \frac{1}{4}W_{12534} - \frac{1}{4}W_{12543} \\ & - \frac{1}{4}W_{13245} + \frac{1}{4}W_{13254} - \frac{1}{4}W_{13425} - \frac{1}{4}W_{13452} + \frac{1}{4}W_{13524} + \frac{1}{4}W_{13542} \\ & + \frac{1}{4}W_{14235} + \frac{1}{4}W_{14253} - \frac{1}{4}W_{14325} - \frac{1}{4}W_{14352} + \frac{1}{4}W_{14523} - \frac{1}{4}W_{14532} \\ & - \frac{1}{4}W_{15234} - \frac{1}{4}W_{15243} + \frac{1}{4}W_{15324} + \frac{1}{4}W_{15342} - \frac{1}{4}W_{15423} + \frac{1}{4}W_{15432} \\ \gamma_{1|2345} &= \frac{1}{4}W_{12345} - \frac{1}{12}W_{12354} - \frac{1}{12}W_{12435} - \frac{1}{12}W_{12453} - \frac{1}{12}W_{12534} + \frac{1}{12}W_{12543} \\ & - \frac{1}{12}W_{13245} + \frac{1}{12}W_{13254} - \frac{1}{12}W_{13425} - \frac{1}{12}W_{13452} + \frac{1}{12}W_{13524} + \frac{1}{12}W_{13542} \\ & - \frac{1}{12}W_{14235} - \frac{1}{12}W_{14253} + \frac{1}{12}W_{14325} + \frac{1}{12}W_{14352} - \frac{1}{12}W_{14523} + \frac{1}{12}W_{14532} \\ & - \frac{1}{12}W_{15234} + \frac{1}{12}W_{1524
$$

According to the deconcatenation (2.22) these BRST-invariant permutations give rise to the following orthogonal idempotents:

$$
\begin{split} \gamma_{12345}^{(1)}&=\frac{1}{4}W_{12345}-\frac{1}{12}W_{12354}-\frac{1}{12}W_{12435}-\frac{1}{12}W_{12453}-\frac{1}{12}W_{12534}+\frac{1}{12}W_{12543}\\ &-\frac{1}{12}W_{13245}+\frac{1}{12}W_{13254}-\frac{1}{12}W_{13425}-\frac{1}{12}W_{13452}+\frac{1}{12}W_{13524}+\frac{1}{12}W_{13542}\\ &-\frac{1}{12}W_{14235}-\frac{1}{12}W_{14253}+\frac{1}{12}W_{14325}+\frac{1}{12}W_{14352}-\frac{1}{12}W_{14523}+\frac{1}{12}W_{14532}\\ &-\frac{1}{12}W_{15234}+\frac{1}{12}W_{15243}+\frac{1}{12}W_{15324}+\frac{1}{12}W_{15342}+\frac{1}{12}W_{15423}-\frac{1}{4}W_{15432}\\ &-\frac{1}{24}W_{12345}-\frac{1}{24}W_{12354}-\frac{1}{24}W_{12435}-\frac{1}{24}W_{12453}-\frac{1}{24}W_{12534}-\frac{1}{24}W_{12543}\\ &-\frac{1}{24}W_{13245}-\frac{1}{24}W_{13254}-\frac{1}{24}W_{13425}-\frac{1}{24}W_{13452}-\frac{1}{24}W_{13524}-\frac{1}{24}W_{13542}\\ &-\frac{1}{24}W_{14235}-\frac{1}{24}W_{14253}-\frac{1}{24}W_{14325}-\frac{1}{24}W_{14352}-\frac{1}{24}W_{14523}-\frac{1}{24}W_{14532}\\ &-\frac{1}{24}W_{15234}-\frac{1}{24}W_{15243}-\frac{1}{24}W_{15324}-\frac{1}{24}W_{15342}-
$$

Here we list the first few expansions of E_{λ} defined in $(A.30)$:

$$
E_2 = \frac{1}{2}W_{12} - \frac{1}{2}W_{21}, \qquad E_{11} = \frac{1}{2}W_{12} + \frac{1}{2}W_{21}
$$
(C.2)
\n
$$
E_3 = \frac{1}{3}W_{123} - \frac{1}{6}W_{132} - \frac{1}{6}W_{213} - \frac{1}{6}W_{231} - \frac{1}{6}W_{312} + \frac{1}{3}W_{321}
$$

\n
$$
E_{21} = \frac{1}{2}W_{123} - \frac{1}{2}W_{321}, \qquad E_{111} = \frac{1}{6}W_{123} + \text{perm}(1, 2, 3)
$$

\n
$$
E_{211} = \frac{1}{4}W_{1234} + \frac{1}{12}W_{1243} + \frac{1}{12}W_{1324} + \frac{1}{12}W_{1342} + \frac{1}{12}W_{1423} - \frac{1}{12}W_{1432}
$$

\n
$$
+ \frac{1}{12}W_{2134} - \frac{1}{12}W_{2143} + \frac{1}{12}W_{2314} + \frac{1}{12}W_{2341} + \frac{1}{12}W_{2413} - \frac{1}{12}W_{2431}
$$

\n
$$
+ \frac{1}{12}W_{3124} - \frac{1}{12}W_{3142} - \frac{1}{12}W_{3214} - \frac{1}{12}W_{3241} + \frac{1}{12}W_{3412} - \frac{1}{12}W_{3421}
$$

\n
$$
+ \frac{1}{12}W_{4123} - \frac{1}{12}W_{4132} - \frac{1}{12}W_{4213} - \frac{1}{12}W_{4231} - \frac{1}{12}W_{4312} - \frac{1}{4}W_{4321}
$$

C.1. The Berends-Giele idempotents

The Berends-Giele idempotents $\mathcal{E}(P)$ are defined in section 2.2 as the inverse $\theta(E(P))$ of the Eulerian idempotent (A.20). Their expansions up to multiplicity three were given in (2.13) and now we write down the multiplicity four:

$$
\mathcal{E}(1234) = \frac{1}{4}W_{1234} - \frac{1}{12}W_{1243} - \frac{1}{12}W_{1324} - \frac{1}{12}W_{1342} - \frac{1}{12}W_{1423} + \frac{1}{12}W_{1432}
$$
 (C.3)
\n
$$
- \frac{1}{12}W_{2134} + \frac{1}{12}W_{2143} - \frac{1}{12}W_{2314} - \frac{1}{12}W_{2341} + \frac{1}{12}W_{2413} + \frac{1}{12}W_{2431}
$$
\n
$$
- \frac{1}{12}W_{3124} - \frac{1}{12}W_{3142} + \frac{1}{12}W_{3214} + \frac{1}{12}W_{3241} - \frac{1}{12}W_{3412} + \frac{1}{12}W_{3421}
$$
\n
$$
- \frac{1}{12}W_{4123} + \frac{1}{12}W_{4132} + \frac{1}{12}W_{4213} + \frac{1}{12}W_{4231} + \frac{1}{12}W_{4312} - \frac{1}{4}W_{4321}
$$

As a curiosity, noting that $E_4 = I_4$ one can derive these permutations using the conversion (A.28) together with (A.13) for the permutations in $\theta(B_p)$ (note $\theta^2 = 1$). So (A.29) yields the permutations in $\mathcal{E}_{1234} = \theta(I_4)$ as

$$
\mathcal{E}(1234) = -\frac{1}{4}1 \sqcup 2 \sqcup 3 \sqcup 4 + \frac{1}{3}1 \sqcup 2 \sqcup 34 + \frac{1}{3}1 \sqcup 23 \sqcup 4 - \frac{1}{2}1 \sqcup 234
$$
\n
$$
+ \frac{1}{3}12 \sqcup 3 \sqcup 4 - \frac{1}{2}12 \sqcup 34 - \frac{1}{2}123 \sqcup 4 + 1234.
$$
\n(C.4)

Appendix D. Parity of the disk amplitude and even partitions

Parity of the amplitude $A^{\text{string}}(1,\ldots,n) = (-1)^n A^{\text{string}}(n,\ldots,1)$ explains the vanishing of $A_{f_2^m f_M}(\gamma_{1|P_1,...,P_k})$ for even k as observed in Table 1. A quick counting argument suggests why this is so as $\sum_{k} \begin{bmatrix} n-1 \\ 2k \end{bmatrix}$ $\binom{k-1}{2k} = \frac{1}{2}$ $\frac{1}{2}(n-1)!$ is the upper bound in the dimension of string disk amplitudes from properties of the string worldsheet alone [2]. More precisely:

Proposition. If k is even then the *n*-point disk amplitude satisfies

$$
A^{\text{string}}(\gamma_{1|P_1,\dots,P_k}) = 0,\tag{D.1}
$$

where $\gamma_{1|P_1,...,P_k}$ is the BRST-invariant permutation (2.4).

Proof. The parity of A^{string} at n points can be written as $A^{string}(1, \sigma) = (-1)^n A^{string}(1, \tilde{\sigma})$ by cyclicity. This means, by (2.20) , that $A^{\text{string}}(\gamma_{1|P_1,...,P_k})$ will vanish whenever the parity of A^{string} at n points is opposite to the parity of \mathcal{I}_p for $p \models n-1$. To see why this is true consider the example of $A^{\text{string}}(\gamma_{1|23,4})$ with the expression for the BRST-invariant permutation in (2.8). The terms can be rearranged as

$$
A^{\text{string}}(\gamma_{1|23,4}) = \frac{1}{2} \left(A^{\text{string}}_{1234} - A^{\text{string}}_{1432} \right) + \frac{1}{2} \left(A^{\text{string}}_{1243} - A^{\text{string}}_{1342} \right) + \frac{1}{2} \left(A^{\text{string}}_{1423} - A^{\text{string}}_{1324} \right) = 0 \tag{D.2}
$$

which vanishes by parity $A_{1234}^{string} = A_{1432}^{string}$. Notice that this happens because the parity of $\mathcal{I}_{21}(23,4)$ from (2.19) is the opposite of the string disk amplitude; $\tilde{\mathcal{I}}_{21}(23,4) = -\mathcal{I}_{21}(23,4)$. The proposition can now be proven by considering the two cases when n is even or odd.

For *n* even the parity of the *n*-point disk amplitude is + so $A^{\text{string}}(\gamma_{1|P_1,...,P_k})$ vanishes if the parity of \mathcal{I}_p is – for a composition p of n–1. By (2.19) this means that there must be an odd number of even parts in the composition p (which sum to even). But since $n-1$ is odd, there must be an odd number of odd parts in p (which sum to odd). Therefore the number of parts $k(p)$ is even (= odd + odd). Similarly, when n is odd the number of parts $k(p)$ in the composition of p is also even (from even + even). This finishes the proof. ⊔

References

- [1] N.E.J. Bjerrum-Bohr, P.H. Damgaard and P. Vanhove, "Minimal Basis for Gauge Theory Amplitudes," Phys. Rev. Lett. 103, 161602 (2009). [arXiv:0907.1425].
- [2] S. Stieberger, "Open & Closed vs. Pure Open String Disk Amplitudes," [arXiv:0907.2211].
- [3] S. Mizera, "Inverse of the String Theory KLT Kernel," JHEP 06, 084 (2017) [arXiv:1610.04230].
- [4] N. E. J. Bjerrum-Bohr, P. H. Damgaard, H. Johansson and T. Sondergaard, "Monodromy–like Relations for Finite Loop Amplitudes," JHEP 1105, 039 (2011). [arXiv:1103.6190].
- [5] O. Schlotterer and S. Stieberger, "Motivic Multiple Zeta Values and Superstring Amplitudes," J. Phys. A 46, 475401 (2013). [arXiv:1205.1516].
- [6] F. Brown, "On the decomposition of motivic multiple zeta values." In GaloisTeichmller theory and arithmetic geometry, pp. 31-58. Mathematical Society of Japan, 2012 [arXiv:1102.1310 [math.NT]]
- [7] R. Kleiss and H. Kuijf, "Multi Gluon Cross-sections and Five Jet Production at Hadron Colliders," Nucl. Phys. B 312, 616 (1989)..
- [8] L. Solomon, "A Mackey formula in the group ring of a Coxeter group," Journal of Algebra, 41(2), (1976) 255-264.
- [9] A.M. Garsia, C. Reutenauer, "A decomposition of Solomon's descent algebra". Advances in Mathematics, 77(2) (1989), 189-262
- [10] C. Reutenauer, "Theorem of Poincaré-Birkhoff-Witt, logarithm and symmetric group representations of degrees equal to Stirling numbers". In Combinatoire énumérative (pp. 267-284) 1986. Springer, Berlin, Heidelberg.
- [11] M. Schocker, "The descent algebra of the symmetric group," Representations of finite dimensional algebras and related topics in Lie theory and geometry 40 (2004): 145-161.
- [12] A.M.Garsia and J.Remmel, "Shuffles of permutations and the Kronecker product," Graphs Combin. 1 (1985), 217-263.
- [13] J.-Y. Thibon, "Lie idempotents in descent algebras" (lecture notes), Workshop on Hopf Algebras and Props, Boston, March 5 - 9, 2007 Clay Mathematics Institute.
- [14] C. Reutenauer, "Free Lie Algebras," London Mathematical Society Monographs, 1993
- [15] R. Bandiera and C.R. Mafra, "A closed-formula solution to the color-trace decomposition problem," [arXiv:2009.02534 [math.CO]].
- [16] T. van Ritbergen, A. N. Schellekens and J. A. M. Vermaseren, "Group theory factors for Feynman diagrams," Int. J. Mod. Phys. A 14, 41 (1999). [hep-ph/9802376].
- [17] L. Solomon "On the Poincaré-Birkhoff-Witt theorem". Journal of Combinatorial Theory. 1968 May 1;4(4):363-75.
- [18] C.R. Mafra and O. Schlotterer, "The Structure of n-Point One-Loop Open Superstring Amplitudes," JHEP 1408, 099 (2014). [arXiv:1203.6215].
- [19] C.R. Mafra and O. Schlotterer, "Multiparticle SYM equations of motion and pure spinor BRST blocks," JHEP 1407, 153 (2014). [arXiv:1404.4986].
- [20] Lothaire, M., "Combinatorics on Words", (Cambridge Mathematical Library), Cambridge University Press (1997).
- [21] M. Schocker, "Lie elements and Knuth relations," Canad. J. Math. 56 (2004), 871-882. [math/0209327].
- [22] F.A. Berends and W.T. Giele, "Recursive Calculations for Processes with n Gluons," Nucl. Phys. B 306, 759 (1988).
- [23] R. Ree, "Lie elements and an algebra associated with shuffles", Ann.Math. 62, No.2 (1958), 210–220.
- [24] H. Frost, C.R. Mafra and L. Mason, "A Lie bracket for the momentum kernel," [arXiv:2012.00519].
- [25] D.E. Knuth, "Two notes on notation", Amer. Math. Monthly 99 (1992), no. 5, 403–422 [math/9205211].
- [26] J.A.M. Vermaseren, "New features of FORM," arXiv:math-ph/0010025. M. Tentyukov and J.A.M. Vermaseren, "The multithreaded version of FORM," arXiv:hep-ph/0702279.
- [27] L. Brink, J. H. Schwarz and J. Scherk, "Supersymmetric Yang-Mills Theories," Nucl. Phys. B 121, 77-92 (1977)
- [28] C.R. Mafra, O. Schlotterer and S. Stieberger, "Complete N-Point Superstring Disk Amplitude I. Pure Spinor Computation," [arXiv:1106.2645].
- [29] C.R. Mafra and O. Schlotterer, "Non-abelian Z-theory: Berends-Giele recursion for the α' expansion of disk integrals," [arXiv:1609.07078]. http://repo.or.cz/BGap.git
- [30] Z. Bern, L. J. Dixon, M. Perelstein and J. S. Rozowsky, "Multileg one loop gravity amplitudes from gauge theory," Nucl. Phys. B 546, 423 (1999). [hep-th/9811140].
- [31] N. E. J. Bjerrum-Bohr, P. H. Damgaard, B. Feng and T. Sondergaard, "Gravity and Yang-Mills Amplitude Relations," Phys. Rev. D 82, 107702 (2010). [arXiv:1005.4367].
- [32] N. E. J. Bjerrum-Bohr, P. H. Damgaard, T. Sondergaard and P. Vanhove, "The Momentum Kernel of Gauge and Gravity Theories," JHEP 1101, 001 (2011)=. [arXiv:1010.3933].
- [33] J.J.M. Carrasco, C.R. Mafra and O. Schlotterer, "Abelian Z-theory: NLSM amplitudes and α' corrections from the open string," JHEP 06, 093 (2017) [arXiv:1608.02569].
- [34] C.R. Mafra, "Planar binary trees in scattering amplitudes." Algebraic Combinatorics, Resurgence, Moulds and Applications (CARMA) (2020): 349-365. [arXiv:2011.14413 [math.CO]].
- [35] J. Broedel, O. Schlotterer and S. Stieberger, "Polylogarithms, Multiple Zeta Values and Superstring Amplitudes," Fortsch. Phys. 61, 812 (2013). [arXiv:1304.7267].
- [36] J.H. Schwarz, "Superstring Theory," Phys. Rept. 89, 223-322 (1982)
- [37] J. Broedel and L. J. Dixon, "Color-kinematics duality and double-copy construction for amplitudes from higher-dimension operators," JHEP 1210, 091 (2012). [arXiv:1208.0876].
- [38] R.M. Schabinger, "One-Loop $N = 4$ Super Yang-Mills Scattering Amplitudes to All Orders in the Dimensional Regularization Parameter," [arXiv:1103.2769 [hep-th]].
- [39] J. J. M. Carrasco, C.R. Mafra and O. Schlotterer, "Semi-abelian Z-theory: $NLSM + \phi^3$ from the open string," JHEP 08, 135 (2017) [arXiv:1612.06446].
- [40] C.R. Mafra, O. Schlotterer and S. Stieberger, "Complete N-Point Superstring Disk Amplitude II. Amplitude and Hypergeometric Function Structure," [arXiv:1106.2646].
- [41] J. Broedel, O. Schlotterer, S. Stieberger and T. Terasoma, "All order α' -expansion of superstring trees from the Drinfeld associator," Phys. Rev. D 89, no. 6, 066014 (2014). [arXiv:1304.7304]. http://mzv.mpp.mpg.de
- [42] R. Medina, F. T. Brandt and F. R. Machado, "The Open superstring five point amplitude revisited," JHEP 0207, 071 (2002). [hep-th/0208121]. ; L. A. Barreiro and R. Medina, "5-field terms in the open superstring effective action," JHEP 0503, 055 (2005). [hep-th/0503182].
- [43] D. Oprisa and S. Stieberger, "Six gluon open superstring disk amplitude, multiple hypergeometric series and Euler-Zagier sums," [hep-th/0509042].
- [44] S. Stieberger and T. R. Taylor, "Multi-Gluon Scattering in Open Superstring Theory," Phys. Rev. D 74, 126007 (2006). [hep-th/0609175].
- [45] J. M. Drummond and E. Ragoucy, "Superstring amplitudes and the associator," JHEP 1308, 135 (2013). [arXiv:1301.0794].
- [46] J. Blumlein, D. J. Broadhurst and J. A. M. Vermaseren, "The Multiple Zeta Value Data Mine," Comput. Phys. Commun. 181, 582 (2010). [arXiv:0907.2557 [math-ph]].
- [47] F.A. Berends and W.T. Giele, "Multiple Soft Gluon Radiation in Parton Processes," Nucl. Phys. B 313, 595 (1989).
- [48] S. Lee, C.R. Mafra and O. Schlotterer, "Non-linear gauge transformations in $D = 10$ SYM theory and the BCJ duality," JHEP 1603, 090 (2016). [arXiv:1510.08843].
- [49] M. B. Green and J. H. Schwarz, "Supersymmetrical Dual String Theory. 2. Vertices and Trees," Nucl. Phys. B 198, 252-268 (1982)
- [50] N. Berkovits, "Super-Poincare covariant quantization of the superstring," JHEP 0004, 018 (2000) [arXiv:hep-th/0001035].
- [51] C.R. Mafra, O. Schlotterer, S. Stieberger and D. Tsimpis, "A recursive method for SYM n-point tree amplitudes," Phys. Rev. D 83, 126012 (2011). [arXiv:1012.3981].
- [52] C.R. Mafra and O. Schlotterer, "Cohomology foundations of one-loop amplitudes in pure spinor superspace," [arXiv:1408.3605].
- [53] C.R. Mafra and O. Schlotterer, "Towards the n-point one-loop superstring amplitude. Part I. Pure spinors and superfield kinematics," JHEP 08, 090 (2019) [arXiv:1812.10969].
- [54] R. Bandiera, private communication.
- [55] A.M. Garsia, "Combinatorics of the Free Lie Algebra and the Symmetric Group", In Analysis, et Cetera, edited by Paul H. Rabinowitz and Eduard Zehnder, Academic Press, (1990) 309-382
- [56] C.R. Mafra and O. Schlotterer, "Berends-Giele recursions and the BCJ duality in superspace and components," JHEP 1603, 097 (2016). [arXiv:1510.08846 [hep-th]].
- [57] Atkinson, M. D. "Solomon's Descent Algebra Revisited," Bulletin of the London Mathematical Society Bull London Math Soc (1992)
- [58] J.L. Loday, "Série de Hausdorff, idempotents Eulériens et algebres de Hopf". Exp. Math. 12 (1994), 165-178.
- [59] R. Bandiera, F. Schaetz, "Eulerian idempotent, pre-Lie logarithm and combinatorics of trees". arXiv:1702.08907.
- [60] C. Malvenuto, and C. Reutenauer, "Duality between quasi-symmetrical functions and the solomon descent algebra", Journal of Algebra, 177(3), (1995) pp.967-982.