A note on extremal intersecting linear Ryser systems

Adrián Vázquez-Ávila*

Subdirección de Ingeniería y Posgrado Universidad Aeronáutica en Querétaro

Abstract

A famous conjecture of Ryser states that any r-partite set system has transversal number at most r-1 times their matching number. This conjecture is only known to be true for $r \leq 3$ in general, for $r \leq 5$ if the set system is intersecting, and for $r \leq 9$ if the intersecting set system is linear. In this note, we deal with Ryser's Conjecture for intersecting r-partite linear systems; that is, if τ is the transversal number for an intersecting r-partite linear system, then Ryser's Conjecture states that $\tau \leq r-1$. If this conjecture is true, this is known to be sharp for r for which there exists a projective plane of order r-1. There has also been considerable effort to find intersecting r-partite set systems whose transversal number is r-1.

In this note, the following is proved: if $r \ge 4$ is an even integer, then $f_l(r) \ge 3(r-2)+1$, where $f_l(r)$ is the minimum number of lines of an intersecting *r*-partite linear system whose transversal number is r-1. This lower bound gives an exact value for $f_l(r)$, for some small values of *r*. Also, we prove that any *r*-partite linear system satisfies $\tau \le r-1$ if $\nu_2 \le r$ for all $r \ge 3$ odd integer and $\nu_2 \le r-1$ for all $r \ge 4$ even integer, where ν_2 is the maximum cardinality of a subset of lines $R \subseteq \mathcal{L}$ such that every triplet of different elements of *R* does not have a common point.

^{*}adrian.vazquez@unaq.mx

Keywords. Ryser's Conjecture, linear systems, transversal number, 2-packing number, projective planes

1 Introduction

A set system is a pair (X, \mathcal{F}) where \mathcal{F} is a finite family of subsets on a ground set X. A set system can be also thought of as a hypergraph, where the elements of X and \mathcal{F} are called *vertices* and *hyperedges* respectively. The set system (X, \mathcal{F}) is called *r-uniform*, when all subsets of \mathcal{F} has size *r*. The set system (X, \mathcal{F}) is *r-partite* if the elements of X can be partitioned into *r* sets X_1, \ldots, X_r , called the *sides*, such that each element of \mathcal{F} contain exactly one element of X_i , for all $i = 1, \ldots, r$, that is $|F \cap X_i| = 1$, where $F \in \mathcal{F}$. Thus, an *r*-partite set system is an *r*-uniform set system.

Let (X, \mathcal{F}) be a set system. A subset $T \subseteq X$ is a *transversal* of (X, \mathcal{F}) if for any element $F \in \mathcal{F}$ contain at least an element of T, that is, $T \cap F \neq \emptyset$, for every $F \in \mathcal{F}$. The *transversal number* of (X, \mathcal{F}) , $\tau = \tau(X, \mathcal{F})$, is the smallest possible cardinality of a transversal of (X, \mathcal{F}) . The transversal number has been studied in the literature in many different contexts and names. For example, with the name of *piercing number* and *covering number*, see for instance [8, 9, 7, 11, 14, 16, 17, 18, 19].

Let (X, \mathcal{F}) be a set system. A subset $\mathcal{E} \subseteq \mathcal{F}$ of a set system (X, \mathcal{F}) is called a *matching* if $F \cap \hat{F} = \emptyset$ for every $F, \hat{F} \in \mathcal{E}$. The matching number of $(X, \mathcal{F}), \nu = \nu(X, \mathcal{F})$, is the cardinality of the largest matching of (X, \mathcal{F}) . A set system is called *intersecting* if $\nu = 1$; that is, $F \cap \hat{F} \neq \emptyset$, for every $F, \hat{F} \in \mathcal{F}$.

It is not hard to see that any r-uniform set system (X, \mathcal{F}) satisfies the inequality $\tau \leq r\nu$. It is well-known this bound is sharp, as shown by the family of all subsets of size r in a ground set of size kr - 1, which has $\nu = k - 1$ and $\tau = (k - 1)r$. On the other hand, if $\nu = 1$, any projective plane of order r - 1 (denoted by Π_{r-1} in this paper), where r - 1 is a prime power, satisfies $\tau = r$. However, for r-partite set systems, Ryser conjectured in the 1960's that the upper bound could be improved. **Ryser's Conjecture:** Any *r*-partite set system satisfies $\tau \leq (r-1)\nu$.

For the special case r = 2, Ryser's Conjecture is equivalent to Kőnig's Theorem. The only other known general case of the conjecture is when r = 3, see [2]. However, Ryser's conjecture is also known to be true in some special cases. Tuza verified [21] Ryser's Conjecture for $r \leq 5$ if the set system is assumed intersecting. Furthermore, Francetić et al. [13] verified Ryser's Conjecture for $r \leq 9$ if the set system is a linear system, that is, a set system (X, \mathcal{F}) is *linear* if it satisfies $|E \cap F| \leq 1$, for every pair of distinct subsets $E, F \in \mathcal{F}$. In this note every linear system is denoted by (P, \mathcal{L}) , where the elements of P and \mathcal{L} are called *points* and *lines*, respectively. In the rest of this paper, only linear systems are considered. Most of the definitions can be generalized for set systems. Thus, we deal with Ryser's Conjecture for intersecting r-partite linear systems.

Intersecting linear Ryser's Conjecture: Every intersecting rpartite linear system satisfies $\tau \leq r - 1$.

In case the conjecture would be true, it is tight in the sense that for infinitely many r's there are constructions of intersecting r-partite linear systems with $\tau = r - 1$. For example, if r - 1 is a prime power, consider the finite projective plane of order r-1 as a linear system, Π_{r-1} . This linear system is r-uniform and intersecting. To make it r-partite, one just needs to delete one point from the projective plane. This truncated projective plane gives an intersecting r-partite linear system with $\tau \geq r-1$, and r(r-1) points and $(r-1)^2$ lines, and it will be denoted by Π'_{r-1} . However, the construction obtained from the projective plane is not the "optimal" extremal. Although the projective plane construction only contains r(r-1)points (which is an optimal number of points), it has a lot of lines. Let f(r) be the minimum integer so that there exists an intersecting r-partite set system (X, \mathcal{F}) with $\tau = r - 1$ and $|\mathcal{F}| = f(r)$ lines. Analogously, let $f_l(r)$ be the minimum integer so that there exists an intersecting r-partite linear system (P, \mathcal{L}) with $\tau = r - 1$ and $|\mathcal{L}| = f_l(r)$ lines. It is probably that $f_l(r)$ does not exist for some values of r (if r-1 is a prime power, then Π'_{r-1} is known to exist, providing proof that $f_l(r)$ is well-defined). Hence, if $f_l(r)$ does exist, for some r, then $f(r) \leq f_l(r)$. It is not difficult to prove that $f_l(2) = 1$ and $f_l(3) = 3$, see [15]. Furthermore, Mansour et al. [15]

proved that $f_l(4) = 6$ and $f_l(5) = 9$. On the other hand, Aharoni et al. [3] proved that $f_l(6) = 13$ and f(7) = 17 (even when the truncated projective plane does not exist, since it has been proved that finite projective planes of order six do not exist, see [20]); however Francetić proved [13] that $f_l(7)$ does not exist, that is, there is no intersecting 7-partite linear system such that $\tau = 6$. Abu-Khazneha et al. [1] constructed new infinite family of intersecting *r*-partite set systems extremal to Ryser's Conjecture, which exist whenever a projective plane of order r - 2 exists. That construction produces a large number of non-isomorphic extremal set systems. Finally, Aharoni et al. [3] gave a lower bound on f(r) when $r \to \infty$, showing that $f(r) \geq 3.052r + O(1)$, this lower bound is an improvement since Mansour proved [15] that $f(r) \geq (3 - \frac{1}{\sqrt{18}})r(1-o(1)) \approx 2.764(1-o(1))$, when $r \to \infty$. In this paper, a lower bound for $f_l(r)$ is given for $r \geq 4$ an even integer.

Theorem 1.1. If $r \ge 4$ is an even integer, then $3(r-2) + 1 \le f_l(r)$.

Notice that, if $r \in \{2, 4, 6\}$, then $f_l(r) = 3(r-2) + 1$. Aharoni et al. [3] proved $18 \le f(8)$ and $24 \le f(10)$. Hence, by Theorem 1.1 we have: $19 \le f_l(8)$ and $25 \le f_l(10)$.

2 Main Results

Let (P, \mathcal{L}) be a linear system and $p \in P$ be a point. It is denoted by \mathcal{L}_p to be the set of lines incident to p. The *degree* of p is defined as $deg(p) = |\mathcal{L}_p|$ and the maximum degree over all points of the linear system is denoted by $\Delta = \Delta(P, \mathcal{L}).$

A subset R of lines of a linear system (P, \mathcal{L}) is a 2-packing of (P, \mathcal{L}) if the elements of R are triplewise disjoint, that is, if any three elements are chosen in R, then they have not a common point. The 2-packing number of $(P, \mathcal{L}), \nu = \nu_2(P, \mathcal{L})$, is the maximum cardinality of a 2-packing of (P, \mathcal{L}) . There are some works that study this new parameter, see [4, 5, 10, 6, 22, 24, 23, 25, 26].

Theorem 2.1. [5] Let (P, \mathcal{L}) be a linear system with $p, q \in P$ be two points such that $\Delta = deg(p)$ and $\Delta' = \max\{deg(x) : x \in P \setminus \{p\}\}$. If $|\mathcal{L}| \leq \Delta + \Delta' + \nu_2 - 3$, then $\tau \leq \nu_2 - 1$.

Lemma 2.2. [22] Let (P, \mathcal{L}) be an intersecting r-uniform linear system, with $r \geq 3$ be an odd integer. If $\tau = r$, then $\nu_2 = r + 1$.

Corollary 2.3. Let (P, \mathcal{L}) be an intersecting r-partite linear system, with $r \geq 3$ be an odd integer. If $\nu_2 \leq r$ then $\tau \leq r - 1$.

Now, let deal with the case when r is an even integer.

Lemma 2.4. [24] Let (P, \mathcal{L}) be an r-uniform intersecting linear system with $r \geq 2$ be an even integer. If $\nu_2 = r + 1$ then $\tau = \frac{r+2}{2}$.

Hence, if $\tau > \frac{r+2}{2}$ then $\nu_2 \leq r$ and $r \geq 4$.

Lemma 2.5. [24] Let (P, \mathcal{L}) be an intersecting r-uniform linear system, with $r \geq 4$ be an even integer. If $\tau = r$, then $\nu_2 = r$.

Lemma 2.6. Let (P, \mathcal{L}) be an intersecting r-uniform linear system with $r \geq 4$ be an even integer. If $\tau = r - 1$, then $\nu_2 = r$.

Proof. Let (P, \mathcal{L}) be an intersecting *r*-partite linear system. Let $p, q \in P$ be two points such that $\Delta = deg(p)$ and $\Delta' = \max\{deg(x) : x \in P \setminus \{p\}\}$. By Theorem 2.1 if $|\mathcal{L}| \leq \Delta + \Delta' + \nu_2 - 3 \leq 3(r-2) + 1$, then $\tau \leq \nu_2 - 1$, which implies that $\nu_2 = r$.

Corollary 2.7. Let (P, \mathcal{L}) be an intersecting *r*-partite linear system with $r \geq 4$ be an even integer. If $\nu_2 \leq r-1$ then $\tau \leq r-2$.

By Lemmas 2.5 and 2.6, we have:

Corollary 2.8. Let (P, \mathcal{L}) be an intersecting r-uniform linear system with $r \geq 4$ be an even number. If $\tau \in \{r-1, r\}$, then $\nu_2 = r$.

By Corollaries 2.3 and 2.7, and Lemma 2.5, we have:

Theorem 2.9. Let $r \ge 3$ be an integer. Then every intersecting r-partite linear system satisfies

- 1. $\tau \leq r-1$ If $\nu_2 \leq r$ and $r \geq 3$ be an odd integer; and
- 2. $\tau \leq r-2$ if $\nu_2 \leq r-1$ and $r \geq 4$ be an even integer.

The intersecting linear Ryser's Conjecture is almost proved; it only remains to analyzes two cases concerning the 2-packing number:

Conjecture 2.10. Let $r \ge 3$ be an integer. Then every intersecting *r*-partite linear system satisfies:

- 1. $\tau \leq r-1$ if $\nu_2 = r+1$, with $r \geq 3$ an odd number.
- 2. $\tau \leq r-1$ if $\nu_2 = r$, with $r \geq 4$ an even number.

Theorem 2.11. Let $r \ge 4$ be an even integer, then $3(r-2) + 1 \le f_l(r)$.

Proof. Suppose that $\nu_2 \leq r-1$ (by Corollary 2.8). Let $p, q \in P$ be two points such that $deg(p) = \Delta$ and $\Delta' = deg(q) = max\{deg(x) : x \in P \setminus \{p\}\}$. By Theorem 2.1 if $|\mathcal{L}| \leq \Delta + \Delta' + \nu_2 - 3 \leq 3(r-2)$, then $\tau \leq \nu_2 - 1 \leq r-2$. Therefore, $3(r-2) + 1 \leq f_l(r)$.

Acknowledgment

Research was partially supported by SNI and CONACyT.

References

- A. Abu-Khazneha, J. Barát, A. Pokrovskiy and T. Szabó, A family of extremal hypergraphs for Ryser's conjecture, J. Combin. Theory Ser. A 161, 164–177 (2019)
- [2] R. Aharoni, Ryser's conjecture for 3-graphs, Combinatorica 21 (1), 1–4 (2001)
- [3] R. Aharoni, J. Barát and I.M. Wanless, Multipartite hypergraphs achieving equality in Ryser's conjecture, Graphs Combin. 32, 1–15 (2016)

- [4] C. Alfaro and A. Vázquez-Ávila, A note on a problem of Henning and Yeo about the transversal number of uniform linear systems whose 2packing number is fixed, Discrete Math. Lett., 3, 61–66 (2020)
- [5] C. Alfaro, G. Araujo-Pardo, C. Rubio-Montiel and A. Vázquez-Ávila, On transversal and 2-packing numbers in uniform linear systems, AKCE J. Graphs Comb., 17 (1), 335-3341 (2020)
- [6] C. Alfaro, C. Rubio-Montiel and A. Vázquez-Ávila, Covering and 2degree-packing number in graphs, J. Comb. Math. Comb. Comput. Accepted.
- [7] N. Alon, G. Kalai, J. Matoušek, R. Meshulam, Transversal numbers for hypergraphs arising in geometry, Advances in Applied Math. 29, 79-101 (2002)
- [8] N. Alon, D.J. Kleitman, Piercing convex sets and the Hadwiger Debrunner (p,q)-problem, Advances in Math. 96, 103-112 (1992)
- [9] N. Alon, D.J. Kleitman, *Piercing convex sets*, Bull. Of the Amér. Math. Soc. 29, 252-256 (1992)
- [10] G. Araujo-Pardo, A. Montejano, L. Montejano and A. Vázquez-Ávila, On transversal and 2-packing numbers in straight line systems on ℝ², Util. Math. 105, 317–336 (2017)
- [11] J. Eckhoff, A Survey of Hadwiger-Debrunner (p,q)-problem, Discrete and Computational Geometry, The Goodman-Pollack Festschrift Algorithms and Combinatorics, 25, 347-377 (2003)
- [12] A. Eustis, and J. Verstraëte, On the independence number of Steiner systems, Combin. Probab. Comput. 22 (2), 241–252 (2013)
- [13] N. Francetić, S. Herke, B. D. McKay and I. M. Wanless, On Ryser's conjecture for linear intersecting multipartite hypergraphs, Eur. J. Comb. 61, 91–105 (2017)
- [14] M. Huicochea, J. Jerónimo-Castro, L. Montejano and D. Oliveros, About the piercing number of a family of intervals, Discrete Math., 338 (12), 2545-2548 (2015)

- [15] T. Mansour, C. Song and R. Yuster, A comment on Ryser's conjecture for intersecting hypergraphs, Graphs Combin. 25, 101–109 (2009)
- [16] L. Montejano, P. Soberón, Piercing numbers for balanced and unbalanced families, Discrete Computational Geometry 45 (2), 358–364 (2011)
- [17] A. Noga, K. Gil, J. Matousek and R. Meshulam, *Transversal numbers for hypergraphs arising in geometry*, Adv. in Appl. Math. **29** (1), 79–101 (2002)
- [18] A. Noga and D. J. Kleitman, *Piercing convex sets*, Bull. Amer. Math. Soc. (N.S.) **27** (2), 252–256 (1992)
- [19] D. Oliveros, C. O'Neill and S. Zerbib, The geometry and combinatorics of discrete line segment hypergraphs, Discrete Math. 343 (6), 111825 (2020)
- [20] G. Tarry, Le probléme de 36 officiers, Compte Rendu de l'Assoc. Francais Avanc. Sci. Naturel 2, 170–203 (1901)
- [21] Z. Tuza, Ryser's conjecture on transversals of r-partite hypergraphs, Ars. Combin. 16, 201–209 (1983)
- [22] A. Vázquez-Ávila, A note on domination and 2-packing numbers in intersecting linear systems, Appl. Math. E-Notes 19, 310–314 (2019)
- [23] A. Vázquez-Ávila, Domination and 2-degree-packing number in graphs,J. Comb. Math. Comb. Comput. Accepted.
- [24] A. Vázquez-Ávila, On domination and 2-packing numbers in intersecting linear systems, Ars. Comb. Accepted.
- [25] A. Vázquez-Ávila, On intersecting straight line systems, J. Discret. Math. Sci. Cryptogr. Accepted.
- [26] A. Vázquez-Ávila, On transversal numbers of intersecting straight line systems and intersecting segment systems. Bol. Soc. Mat. Mex. Accepted.