On $(1, C_4)$ one-factorization and two orthogonal $(2, C_4)$ one-factorization of complete graphs

Adrián Vázquez-Ávila^{*}

Subdirección de Ingeniería y Posgrado Universidad Aeronáutica en Querétaro

Abstract

An one-factorization \mathcal{F} of the complete graph K_n is (l, C_k) , where $l \geq 0$ and $k \geq 4$ are integers, if the union $F \cup G$, for any $F, G \in \mathcal{F}$, includes exactly l (edge-disjoint) cycles of length k ($lk \leq n$). Moreover, a pair of orthogonal one-factorizations \mathcal{F} and \mathcal{G} of the complete graph K_n is (l, C_k) if the union $F \cup G$, for any $F \in \mathcal{F}$ and $G \in \mathcal{G}$, includes exactly l cycles of length k.

In this paper, we prove the following: if $q \equiv 11 \pmod{24}$ is an odd prime power, then there is a $(1, C_4)$ one-factorization of K_{q+1} . Also, there is a pair of orthogonal $(2, C_4)$ one-factorization of K_{q+1} .

Keywords. One-factorization, orthogonal one-factorization, Strong starters.

1 Introduction

A one-factor of a graph G is a regular spanning subgraph of degree one. A one-factorization of a graph G is a set $\mathcal{F} = \{F_1, F_2, \dots, F_n\}$ of edge disjoint

^{*}adrian.vazquez@unaq.mx

one-factors such that $E(G) = \bigcup_{i=1}^{n} E(F_i)$. Two one-factorizations $\mathcal{F} = \{F_1, F_2, \ldots, F_n\}$ and $\mathcal{H} = \{H_1, H_2, \ldots, H_n\}$ of a graph G are orthogonal if $|F \cap H| \leq 1$, for every $F \in \mathcal{F}$ and $H \in \mathcal{H}$. A one-factorization $\mathcal{F} = \{F_1, F_2, \ldots, F_n\}$ of a graph G is (l, C_k) , where $l \geq 0$ and $k \geq 4$ integers, if $F_i \cup F_j$ include exactly l cycles of length k ($lk \leq n$). In particular, if l = 0, we said that the one factorization is C_k -free, see for example [4, 18, 30].

A pair of orthogonal one-factorizations \mathcal{F} and \mathcal{H} of the complete graph K_n is (l, C_k) , where $l \geq 0$ and $k \geq 4$ integers, if $F \cup H$ include exactly l cycles of length k, for all $F \in \mathcal{F}$ and $H \in \mathcal{H}$.

An interesting way of constructing one-factorizations of the complete graph is by using starters of additive Abelian groups of odd order: Let Γ be a finite additive Abelian group of odd order n = 2k + 1. And let $\Gamma^* = \Gamma \setminus \{0\}$ be the set of non-zero elements of Γ . A starter for Γ is a set $S = \{\{x_1, y_1\}, \ldots, \{x_k, y_k\}\}$ such that $\{x_1, \ldots, x_k, y_1, \ldots, y_k\} = \Gamma^*$ and $\{\pm (x_i - y_i) : i = 1, \ldots, k\} = \Gamma^*$. Moreover, if $\{x_i + y_i : i = 1, \ldots, k\} \subseteq$ Γ^* with $|\{x_i + y_i : i = 1, \ldots, k\}| = k$, then S is called strong starter for Γ . To see some works related to strong starters, the reader may consult [1, 7, 8, 11, 12, 16, 17, 19, 20, 21, 27, 29, 28].

Strong starters were first introduced by Mullin and Stanton in [25] in constructing of Room squares. Starters and strong starters have been useful to built many combinatorial designs such as Room cubes [10], Howell designs [3], Kirkman triple systems [17, 22], Kirkman squares and cubes [24, 26], and factorizations of complete graphs [2, 4, 9, 11, 15, 30].

Let Γ be a finite additive Abelian group of odd order n = 2k + 1. It is well known that $F_{\gamma} = \{\{\infty, \gamma\}\} \cup \{\{x_i + \gamma, y_i + \gamma\} : 1 \leq i \leq k\}$, for all $\gamma \in \Gamma$, forms a one-factorization of the complete graph on $\Gamma \cup \{\infty\}$. Hence, if $F_0 = \{\{\infty, 0\}\} \cup \{\{x_i, y_i\} : 1 \leq i \leq k\}$, then $F_{\gamma} = F_0 + \gamma$, for all $\gamma \in \Gamma^*$. On the other hand, let $S = \{\{x_i, y_i\} : 1 \leq i \leq k\}$ and $T = \{\{u_i, v_i\} : 1 \leq i \leq k\}$ two starters for Γ . Without loss of generality, we assume that $x_i - y_i = u_i - v_i$, for all $i = 1, \ldots, k$. Then S and T are orthogonal starters if $u_i - x_i = u_j - x_j$ implies i = j, and if $u_i \neq x_i$, for all $i = 1, \ldots, k$.

Let $S = \{\{x_i, y_i\} : i = 1, ..., k\}$ be a starter for a finite additive

Abelian group Γ of odd order n = 2k + 1. It is not hard to see that $-S = \{\{-x_i, -y_i\} : i = 1, \dots, k\}$ is also a starter for Γ .

Theorem 1.1. [14] If there is a strong starter S in an additive Abelian group of odd order, then S and -S are pairwise orthogonal starters.

Let q be an odd prime power. An element $x \in \mathbb{F}_q^*$ is called a *quadratic* residue if there exists an element $y \in \mathbb{F}_q^*$ such that $y^2 = x$. If there is no such y, then x is called a *non-quadratic residue*. The set of quadratic residues of \mathbb{F}_q^* is denoted by QR(q) and the set of non-quadratic residues is denoted by NQR(q). It is well known that QR(q) is a cyclic subgroup of \mathbb{F}_q^* of order $\frac{q-1}{2}$ (see for example [13]). Also, it is well known that if either $x, y \in QR(q)$ or $x, y \in NQR(q)$, then $xy \in QR(q)$. Also, if $x \in QR(q)$ and $y \in NQR(q)$, then $xy \in NQR(q)$. For more details of this kind of results the reader may consult [5, 13].

Horton in [14] proved the following (see too [1]):

Proposition 1.2. [14] If $q \equiv 3 \pmod{4}$ is an odd prime power $(q \neq 3)$ and $\beta \in NQR(q) \setminus \{-1\}$, then

$$S_{\beta} = \{\{x, x\beta\} : x \in QR(q)\},\$$

is a strong starter for \mathbb{F}_q .

Theorem 1.3. [14] Let $q \equiv 3 \pmod{4}$ be a prime power $(q \neq 3)$. If $\beta_1, \beta_2 \in NQR(q) \setminus \{-1\}$, with $\beta_1 \neq \beta_2$, then S_{β_1} and S_{β_2} are orthogonal.

In this paper, we prove the following:

Theorem 1.4. If $q \equiv 3 \pmod{8}$ is an odd prime power such that $q \equiv -1 \pmod{12}$, then there is $(1, C_4)$ one-factorization of K_{q+1} .

Theorem 1.5. If $q \equiv 3 \pmod{8}$ is an odd prime power such that $q \equiv -1 \pmod{12}$, then there is a pair of orthogonal $(2, C_4)$ one-factorization of K_{q+1} .

2 $(1, C_4)$ one-factorizations of complete graph

Let $n \geq 11$ be an odd number. A one-factorization $\mathcal{F} = \{F_1, F_2 \dots, F_n\}$ of K_{n+1} is uniform (also called semi-regular), if for any $F_i, F_j, F_k, F_m \in \mathcal{F}$ such that $i \neq j$ and $k \neq m$ satisfy that $F_i \cup F_j$ and $F_k \cup F_m$ generate the same cycle structure. Rosa in [23] shows several examples of infinite families of uniform one-factorizations of the complete graph K_{2n} . An important example of uniform one-factorization of complete graph K_{2n} is the onefactorization generated by the Mullin-Nemeth strong starters [19]: Let pbe an odd prime and let m be a positive integer such that $3 < p^m = 2t + 1$, where t is odd such that $p^m \equiv 3 \pmod{4}$. If r is a primitive root in \mathbb{F}_{p^m} , then

$$S = \{\{r^0, r\}, \{r^2, r^3\}, \dots, \{r^{2t-2}, r^{2t-1}\}\}\$$

is a strong starter for \mathbb{F}_{p^m} , see for instance [1, 19].

The following proof is completely analogous to the proof of Theorem 2.1 given Anderson in [2].

Theorem 2.1. Let $q \equiv 3 \pmod{4}$ be an odd prime power $(q \neq 3)$ and let $y \in QR(q)$ be a generator of QR(q). If $\beta \in NQR(q) \setminus \{-1\}$ then the cycle structure of $F_0 \cup F_0 + y$ is the same as the cycle structure of $(F_0 + g) \cup (F_0 + h)$, for every different elements $g, h \in \mathbb{F}_q$. That is, the one-factorization generated by the starter S_β is uniform.

Lemma 2.2. Let $q \equiv 3 \pmod{4}$ be an odd prime power with $q \geq 11$ and $\beta \in NQR(q) \setminus \{-1\}$. If $\beta^2 - \beta + 1 = 0$, then the one-factorization generated by the starter S_β include a cycle of length four, C_4 , with $\infty \in V(C_4)$.

Proof. Let $F_i = \{\{\infty, i\}\} \cup \{\{x + i, x\beta + i\} : x \in QR(q)\}$, for $i \in \mathbb{F}_q$. Then $\mathcal{F} = \{F_i : i \in \mathbb{F}_q\}$ is a one-factorization of the complete graph on $\mathbb{F}_q \cup \{\infty\}$. By Theorem 2.1 it is sufficient to prove that $F_0 \cup F_i$, where $i \in QR(q)$ is a generator of QR(q), include a cycle of length four, C_4 , such that $\infty \in V(C_4)$.

Hence, $\{\infty, 0\}, \{\infty, i\} \in E(C_4)$, where $\{\infty, 0\} \in F_0$ and $\{\infty, i\} \in F_i$. Then $\{-i + i, -i\beta^{-1} + i\}, \{i, i\beta\} \in E(C_4)$. Therefore, $F_0 \cup F_i$ include the following cycle of length 4:

$$\{\infty, 0\}\{0, -i\beta^{-1} + i\}\{i, i\beta\}\{\infty, i\} = C_4,$$

which implies that $i\beta^{-1} + i = i\beta$. Hence, $\beta^2 - \beta + 1 = 0$.

Lemma 2.3. If $q \equiv -1 \pmod{12}$ is an odd prime power with $q \geq 11$ and $\beta \in NQR(q) \setminus \{-1\}$, then $\beta^2 - \beta + 1 \neq 0$.

Proof. Suppose that there exists $\beta \in NQR(q) \setminus \{-1\}$ such that $\beta^2 - \beta + 1 = 0$. Notice that

$$\beta(\beta - 2) = \beta^2 - 2\beta = -(\beta + 1).$$

Given that $\beta \in NQR(q)$ and $-1 \in NQR(q)$, then $(\beta - 2)(\beta + 1) \in QR(q)$. Furthermore, $(\beta - 2)(\beta + 1) = -3$, which implies that $(\beta - 2)(\beta + 1) \in NQR(q)$ (since $q \equiv -1 \mod 12$ implies that $3 \in QR(q)$), which is a contradiction.

It is not hard to check that, if $q \equiv 1 \pmod{3}$ is an odd prime power with $q \geq 7$, then there exist $\beta \in NQR(q) \setminus \{-1\}$ such that $\beta^2 - \beta + 1 = 0$. Hence, we have the following:

Theorem 2.4. Let $q \equiv 3 \pmod{8}$ be an odd prime power with $q \geq 11$ such that $3 \in NQR(q)$ (such that $q \equiv 1 \pmod{3}$). If $\beta \in NQR(q) \setminus \{-1\}$ is such that $\beta^2 - \beta + 1 = 0$, then the one-factorization generated by the starter S_β is $(1, C_4)$.

Proof. Let $F_i = \{\{\infty, i\}\} \cup \{\{x+i, x\beta+i\} : x \in QR(q)\}$, for $i \in \mathbb{F}_q$. Then $\mathcal{F} = \{F_i : i \in \mathbb{F}_q\}$ is a one-factorization of the complete graph on $\mathbb{F}_q \cup \{\infty\}$. By Theorem 2.1 it is sufficient to prove that $F_0 \cup F_i$, where $i \in QR(q)$ is a generator of QR(q), include an unique cycle of length four, C_4 .

Suppose that there is a $\beta \in NQR(q) \setminus \{-1\}$ such that $\beta^2 - \beta + 1 = 0$ and S_β include a cycle of length four, C_4 , with $\infty \neq V(C_4)$ (by Lemma ??). Given that $E(C_4) \cap F_0 \neq \emptyset$, then there is an $a \in V(C_4) \cap QR(q)$ such that

$$\{a, a\beta\} \in F_0 \text{ and } \{a\beta, (a\beta - i)\beta^{t_1} + i\} \in F_i,$$

with $t_1 = 1$ if $a\beta - i \in QR(q)$, and $t_1 = -1$ if $a\beta - i \in NQR(q)$. On the other hand

$$\{a, (a-i)\beta^{t_2} + i\} \in F_i \text{ and } \{(a-i)\beta^{t_2} + i, ((a-i)\beta^{t_2} + i)\beta^{t_3}\} \in F_0,$$

whit $t_2 = 1$ if $a - i \in QR(q)$, and $t_2 = -1$ if $a - i \in NQR(q)$, and $t_3 = 1$ if $(a - i)\beta^{t_2} + i \in QR(q)$, and $t_3 = -1$ if $(a - i)\beta^{t_2} + i \in NQR(q)$. Therefore

$$a(\beta^{t_1+1} - \beta^{t_2+t_3}) + i(\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1) = 0,$$

where $t_1, t_2, t_3 \in \{-1, 1\}$. Notice that, if $\{a, b\}\{b, c\}\{a, d\}\{d, c\} = C_4$, then $|V(C_4) \cap QR(q)| = 2$. It is not hard to check that, if $t_1 + 1 = t_2 + t_3$, where $t_1, t_2, t_3 \in \{-1, 1\}$, then $\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1 \neq 0$, which implies that the one-factorization generated by the starter S_β is C_4 -free. Suppose that $t_1 + 1 \neq t_2 + t_3$.

- case (i) Suppose that $t_1 = t_2 = 1$ and $t_3 = -1$. In this case $(a i)\beta + i \in QR(q)$ and $(a\beta i)\beta + i \in NQR(q)$. Since $\{(a i)\beta + i, (a\beta i)\beta + i\} \in F_0$, then $(a\beta i)\beta + i ((a i)\beta + i) = x(\beta 1)$, where $x = -a\beta$. Hence $(a i)\beta + i = -a\beta$ and $(a\beta 1)\beta + i = -a\beta^2$, since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$. But $(a\beta 1)\beta + i = -a\beta^2$ implies that $2a\beta^2 = i(\beta 1)$, which is a contradiction, since $2a\beta^2 \in NQR(q)$ (given that $2 \in NQR(q)$) and $i(\beta 1) \in QR(q)$ (given that $\beta^2 \beta + 1 = 0$). Hence, the one-factorization generated by the starter S_β is $(0, C_4)$, C_4 -free.
- case (ii) Suppose that $t_1 = t_3 = 1$ and $t_2 = -1$. In this case $(a i)\beta^{-1} + i \in QR(q)$ and $(a\beta i)\beta + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a i)\beta^{-1} + i$ and $x\beta = (a\beta i)\beta + i$. Hence, $(a i) + i\beta = (a\beta i)\beta + i$, which implies that $2i\beta = a(\beta^2 + \beta) = a(2\beta 1)$ (since $\beta^2 \beta + 1 = 0$), which implies that $2\beta(a i) = a$, a contradiction (since $a i \in NQR(q)$). Hence, the one-factorization generated by the starter S_β is C_4 -free.

case (iii) Suppose that $t_2 = t_3 = 1$ and $t_1 = -1$. In this case $(a - i)\beta + i \in QR(q)$ and $(a\beta - i)\beta^{-1} + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a - i)\beta + i$ and $x\beta = (a\beta - i)\beta^{-1} + i$. Hence, $(a - i)\beta^2 + i\beta = (a\beta - i)\beta^{-1} + i$, which implies that $a(\beta - 1) = a\beta - i$, a contradiction (since $a\beta - i \in NQR(q)$). Hence, the one-factorization generated by the starter S_β is C_4 -free.

case (iv) Suppose that $t_1 = 1$ and $t_2 = t_3 = -1$. Then

$$a(\beta^2 - \beta^{-2}) + i(\beta^{-2} - \beta^{-1} - \beta + 1) = 0$$

Given that $\beta^2 - \beta + 1 = 0$ (implying that $\beta + \beta^{-1} = 1$), then $a(\beta^2 - \beta^{-2}) + i\beta^{-2} = 0$, which implies that $a\beta^2 = \beta^{-2}(a-i)$, a contradiction (since $a - i \in NQR(q)$). Hence, the one-factorization generated by the starter S_{β} is C_4 -free.

case (v) Suppose that $t_1 = t_2 = t_3 = -1$. Then

 $a(1-\beta^{-2}) + i(\beta^{-2} - 2\beta^{-1} + 1) = 0,$

which implies that $a(\beta^2 - 1) + i(\beta^2 - 2\beta + 1) = a(\beta - 2) - i\beta = 0$ (since $\beta^2 - \beta + 1 = 0$). Hence, $\beta(a - i) = 2a$, which is a contradiction (since $a - i \in NQR(q)$ and $2 \in NQR(q)$). Hence, the one-factorization generated by the starter S_{β} is C_4 -free.

Hence, by Lemma 2.2 the one-factorization generated by the starter S_{β} is $(1, C_4)$.

Example 1. Let q = 19. A primitive root of \mathbb{F}_{19} is r = 2 and $i = r^2 = 4$ is a generator of QR(19). Furthermore, the set of non-quadratic residues of \mathbb{F}_{19} is:

$$\{2, 3, 8, 10, 12, 13, 14, 15, 18\}.$$

Hence, if $\beta = 8$, then $\beta^2 - \beta + 1 = 0$, which implies that the one-factorization generated by the starter S_8 include an unique cycle C_4 , with $V(C_4) = \{\infty, 0, 4, 13\}$.

Lemma 2.5. Let $q \equiv 3 \pmod{8}$ be an odd prime power such that $q \equiv -1 \pmod{12}$. If there exist $\beta \in NQR(q) \setminus \{-1\}$ such that $\beta - 1 \in NQR(q)$ and $\beta^2 + 1 \in QR(q)$, then the one-factorization generated by the starter S_β is $(1, C_4)$.

Proof. Let $F_i = \{\{\infty, i\}\} \cup \{\{x+i, x\beta+i\} : x \in QR(q)\}$, for $i \in \mathbb{F}_q$. Then $\mathcal{F} = \{F_i : i \in \mathbb{F}_q\}$ is a one-factorization of the complete graph on $\mathbb{F}_q \cup \{\infty\}$. Then the one-factorization generated by the starter S_β include a cycle of length four, C_4 , with $\infty \notin V(C_4)$ (by Lemma 2.3 and Theorem 2.4).

By Theorem 2.1, if $i \in QR(q)$ is a generator of QR(q), then the cycle structure $F_0 \cup F_i$ is the same as the cycle structure of the union of any two distinct one-factors, say $F_g \cup F_h$. Given that $E(C_4) \cap F_0 \neq \emptyset$, then there is an $a \in V(C_4) \cap QR(q)$ such that

$$\{a, a\beta\} \in F_0 \text{ and } \{a\beta, (a\beta - i)\beta^{t_1} + i\} \in F_i$$

with $t_1 = 1$ if $a\beta - i \in QR(q)$, and $t_1 = -1$ if $a\beta - i \in NQR(q)$. On the other hand

$$\{a, (a-i)\beta^{t_2}+i\} \in F_i \text{ and } \{(a-i)\beta^{t_2}+i, ((a-i)\beta^{t_2}+i)\beta^{t_3}\} \in F_0,$$

whit $t_2 = 1$ if $a - i \in QR(q)$, and $t_2 = -1$ if $a - i \in NQR(q)$, and $t_3 = 1$ if $(a - i)\beta^{t_2} + i \in QR(q)$, and $t_3 = -1$ if $(a - i)\beta^{t_2} + i \in NQR(q)$. Therefore

$$a(\beta^{t_1+1} - \beta^{t_2+t_3}) + i(\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1) = 0,$$

where $t_1, t_2, t_3 \in \{-1, 1\}$. Notice that, if $\{a, b\}\{b, c\}\{a, d\}\{d, c\} = C_4$, then $|V(C_4) \cap QR(q)| = 2$. It is not hard to check that, if $t_1 + 1 = t_2 + t_3$, where $t_1, t_2, t_3 \in \{-1, 1\}$, then $\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1 \neq 0$, which implies that the one-factorization generated by the starter S_β is C_4 -free. Suppose that $t_1 + 1 \neq t_2 + t_3$.

Case (i): Suppose that $t_1 = 1$ and $-t_2 = t_3 = 1$. In this case $(a - i)\beta^{-1} + i \in QR(q)$ and $(a\beta - i)\beta + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a - i)\beta^{-1} + i$ and $x\beta = (a\beta - i)\beta + i$. Hence,

 $(a\beta - i)\beta + i = (a - i) + i\beta$, which implies that $2i = a(\beta + 1)$. Therefore, if $\beta + 1 \in NQR(q)$, then the one-factorization generated by the starter S_{β} does include a cycle of length four; otherwise don't.

Case (ii): Suppose that $t_1 = 1$ and $t_2 = t_3 = -1$. In this case $(a - i)\beta^{-1} + i \in NQR(q)$ and $(a\beta - i)\beta + i \in QR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta - i)\beta + i$ and $x\beta = (a - i)\beta^{-1} + i$. Hence, $(a - i)\beta^{-1} + i = (a\beta - i)\beta^2 + i\beta$, which implies that

$$i(\beta^3 - \beta^2 + \beta - 1) = a(\beta^4 - 1) = a(\beta + 1)(\beta^3 - \beta^2 + \beta - 1)$$

implying that $a - i = -a\beta$, which is a contradiction, given that $t_2 = -1$ implies that $a - i \in NQR(q)$. the one-factorization generated by the starter S_β is C_4 -free.

- Case (iii): Suppose that $t_1 = 1$ and $t_2 = -t_3 = 1$. In this case $(a i)\beta + i \in NQR(q)$ and $(a\beta i)\beta + i \in QR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta i)\beta + i$ and $x\beta = (a i)\beta + i$. Hence, $(a i)\beta + i = (a\beta i)\beta^2 + i\beta$, which implies that $i(\beta 1) = a\beta(\beta + 1)$. If $\beta + 1 \in QR(q)$, then the one-factorization generated by the starter S_β does include a cycle of length four; otherwise don't.
- Case (iv): Suppose that $t_1 = -1$ and $t_2 = t_3 = 1$. In this case $(a i)\beta + i \in QR(q)$ and $(a\beta i)\beta^{-1} + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a i)\beta + i$ and $x\beta = (a\beta i)\beta^{-1} + i$. Hence, $(a\beta i)\beta^{-1} + i = (a i)\beta^2 + i\beta$, which implies that $a\beta(\beta + 1) = i(\beta^2 + 1)$. If $\beta + 1 \in NQR(q)$, then the one-factorization generated by the starter S_β does include a cycle of length four; otherwise don't.
- Case (v): Suppose that $t_1 = t_2 = t_3 = -1$. In this case $(a i)\beta^{-1} + i \in NQR(q)$ and $(a\beta i)\beta^{-1} + i \in QR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta i)\beta^{-1} + i$ and $x\beta = (a i)\beta^{-1} + i$. Hence, $(a i)\beta^{-1} + i = (a\beta i) + i\beta$, which implies that $i(\beta 1)^2 = 0$, which is a contradiction.

It is not hard to check that Case (i) and Case (iv) generate the same (unique) cycle of length four. On the other hand, Case (iii) generate an unique cycle of length four. Therefore, the pair of orthogonal one-factorizations generated by the starter S_{β} and $-S_{\beta}$ does include exactly one cycle of length four.

Example 2. Let q = 59. A primitive root of \mathbb{F}_{59} is r = 2 and $i = r^2 = 4$ is a generator of QR(59). Hence, if $\beta = 32$, then $\beta - 1 \in NQR(59)$ and $\beta^2 + 1 \in QR(q)$. Given that $a = 36 \in QR(59)$ is such that $2i = a(\beta - 1)$, then the one-factorization generated by the starter S_{32} include an unique cycle C_4 , with $V(C_4) = \{5, 31, 36, 42\}$.

Remark 1. Let q = ef + 1 be a prime power and let H be the subgroup of \mathbb{F}_q^* of order f with $\{H = C_0, \ldots, C_{e-1}\}$ the set of (multiplicative) cosets of H in \mathbb{F}_q^* (that is, $C_i = g^i C_0$, where g is the least primitive element of \mathbb{F}_q). The cyclotomic number (i, j) is defined as $|\{x \in C_i : x + 1 \in C_j\}|$. In particular, if e = 2 and f is odd, then $(0, 0) = \frac{f-1}{2}$, $(0, 1) = \frac{f+1}{2}$, $(1, 0) = \frac{f-1}{2}$ and $(1, 1) = \frac{f-1}{2}$. On the other hand, if e = 2 and f is even, then $(0, 0) = \frac{f-2}{2}$, $(0, 1) = \frac{f}{2}$, $(1, 0) = \frac{f}{2}$ and $(1, 1) = \frac{f}{2}$, see [6], Table VII.8.50.

Hence, if e = 2, then $C_0 = QR(q)$ and $C_1 = NQR(q)$ are the cosets of QR(q) in \mathbb{F}_q^* .

Lemma 2.6. Let $q \equiv 3 \pmod{8}$ be an odd prime power such that $q \equiv -1 \pmod{12}$. Then the set

$$M = \{\beta \in NQR(q) \setminus \{-1\} : \beta - 1 \in NQR(q), \beta^2 + 1 \in QR(q)\} \neq \emptyset$$

Proof. Given that $-1 \in NQR(q)$, then $\beta \to \beta^2$, for all $\in \beta \in NQR(q)$, is a bijection between NQR(q) and QR(q). Let $A = \{\alpha \in QR(q) : \alpha^2 + 1, \alpha^2 - 1 \in QR(q)\}$ (see Remark 1). Assume that $\alpha - 1 \in QR(q)$ (which implies that $\alpha + 1 \in QR(q)$), otherwise, if $\alpha - 1 \in NQR(q)$, then $M \neq \emptyset$. Moreover, assume that $\alpha^4 + 1 \in NQR(q)$, otherwise $\beta = -\alpha^2 \in M$. Given that $\alpha - 1, \alpha + 1 \in QR(q)$, then $\alpha^2 - 1 \in QR(q)$, which implies that $(\alpha^2 - 1)(\alpha^2 + 1) \in QR(q)$ and $\alpha^4 + 1 \in NQR(q)$, a contradiction. Hence, by Theorem 2.4, Lemma 2.5 and Lemma 2.6, we have:

Theorem 2.7. Let $q \equiv 3 \pmod{8}$ be an odd prime power such that $q \equiv -1 \pmod{12}$. Then there exist a one-factorization $(1, C_4)$ of K_{q+1} .

3 $(2, C_4)$ two orthogonal one-factorizations of complete graph

In this section we will prove that the strong starter given by Horton (see Proposition 1.2) doesn't generate two orthogonal $(1, C_4)$ one-factorization of the complete graph; however, does generate two orthogonal C_4 -free (see for instance [4, 30]) and $(2, C_4)$ one-factorizations of complete graphs.

Lemma 3.1. Let $q \equiv 3 \pmod{8}$ be an odd prime power and $\beta \in NQR(q) \setminus \{-1\}$. If $\beta \notin \{2, 2^{-1}\}$, then the pair of orthogonal one-factorizations generated by the starter S_{β} and $-S_{\beta}$ doesn't include a cycle of length four, C_4 , with $\infty \notin V(C_4)$.

Proof. Let $F_i = \{\{\infty, i\}\} \cup \{\{x + i, x\beta + i\} : x \in QR(q)\}$ and $G_i = \{\{\infty, i\}\} \cup \{\{y + i, y\beta + i\} : y \in NQR(q)\}$, for $i \in \mathbb{F}_q$. Then $\mathcal{F} = \{F_i : i \in \mathbb{F}_q\}$ and $\mathcal{G} = \{G_i : i \in \mathbb{F}_q\}$ are orthogonal one-factorizations of the complete graph on $\mathbb{F}_q \cup \{\infty\}$. Given that $F_i = F_0 + i$ and $G_i = G_0 + i$, without loss of generality we can suppose that $F_0 \cup G_i$, where $i \in \mathbb{F}(q)$, doesn't include a cycle of length 4, C_4 , such that $\infty \notin V(C_4)$.

Assume to the contrary that $F_0 \cup G_i$, for some $i \in \mathbb{F}_q$, does include a cycle of length 4, C_4 , such that $\infty \in V(C_4)$. Consequently $\{\infty, 0\}, \{\infty, i\} \in E(C_4)$, where $\{\infty, 0\} \in F_0$ and $\{\infty, i\} \in G_i$. Then $\{0, -i\beta^{t_1}+i\}, \{i, i\beta^{t_2}\} \in E(C_4)$, where $t_1 = 1$ if $-i \in NQR(q)$ and $t_1 = -1$ if $-i \in QR(q)$, and $t_2 = 1$ if $i \in QR(q)$ and $t_2 = -1$ if $i \in NQR(q)$. Given that, if $i \in QR(q)$ then $t_2 = 1$ and $t_1 = 1$, and if $i \in NQR(q)$ then $t_2 = -1$ and $t_1 = -1$. Therefore $-i\beta^{t_1} + i = i\beta^{t_2}$, where $t_1, t_2 \in \{-1, 1\}$ with $t_1 = t_2 = t$. Hence, $2\beta^t = 1$ with $t \in \{-1, 1\}$, which is a contradiction to the hypothesis.

Lemma 3.2. Let $q \equiv 3 \pmod{8}$ be an odd prime power with $q \equiv -1 \pmod{8}$

12) and $\beta \in NQR(q) \setminus \{-1\}$. If $\beta \notin \{2^{-1}, 2\}$ is such that $\beta + 1, \beta - 1 \in NQR(q)$ and $\beta^2 + 1 \in QR(q)$, then the pair of orthogonal one-factorizations generated by the starter S_{β} and $-S_{\beta}$ does include exactly two cycles of length four.

Proof. Let $F_i = \{\{\infty, i\}\} \cup \{\{x + i, x\beta + i\} : x \in QR(q)\}$ and $G_i = \{\{\infty, i\}\} \cup \{\{y + i, y\beta + i\} : y \in NQR(q)\}$, for $i \in \mathbb{F}_q$. Then $\mathcal{F} = \{F_i : i \in \mathbb{F}_q\}$ and $\mathcal{G} = \{G_i : i \in \mathbb{F}_q\}$ are orthogonal one-factorizations of the complete graph on $\mathbb{F}_q \cup \{\infty\}$. Given that $F_i = F_0 + i$ and $G_i = G_0 + i$, without loss of generality we can suppose that $F_0 \cup G_i$, where $i \in \mathbb{F}(q)$, does include one cycle of length four, C_4 , such that $\infty \notin V(C_4)$ (see Lemma 3.1). Given that $E(C_4) \cap F_0 \neq \emptyset$, then there is an $a \in V(C_4) \cap QR(q)$ such that

$$\{a, a\beta\} \in F_0 \text{ and } \{a\beta, (a\beta - i)\beta^{t_1} + i\} \in G_i,$$

whit $t_1 = 1$ if $a\beta - i \in NQR(q)$, and $t_1 = -1$ if $a\beta - i \in QR(q)$. On the other hand,

$$\{a, (a-i)\beta^{t_2} + i\} \in G_i \text{ and } \{(a-i)\beta^{t_2} + i, ((a-i)\beta^{t_2} + i)\beta^{t_3}\} \in F_0,$$

with $t_2 = 1$ if $a - i \in NQR(q)$ and $t_2 = -1$ if $a - i \in QR(q)$, and $t_3 = 1$ if $(a - i)\beta^{t_2} + i \in QR(q)$ and $t_3 = -1$ if $(a - i)\beta^{t_2} + i \in NQR(q)$. Therefore

$$a(\beta^{t_1+1} - \beta^{t_2+t_3}) + i(\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1) = 0$$

for all $t_1, t_2, t_3 \in \{-1, 1\}$. Notice that, if $\{a, b\}\{b, c\}\{a, d\}\{d, c\} = C_4$, then $|V(C_4) \cap QR(q)| = 2$. It is not hard to check that, if $t_1 + 1 = t_2 + t_3$, where $t_1, t_2, t_3 \in \{-1, 1\}$, then $\beta^{t_2+t_3} - \beta^{t_3} - \beta^{t_1} + 1 \neq 0$, which implies that the pair of orthogonal one-factorizations generated by the starter S_β and $-S_\beta$ doesn't include a cycle of length four. Suppose that $t_1 + 1 \neq t_2 + t_3$.

Case (i): Suppose that $t_1 = t_2 = 1$ and $t_3 = -1$. In this case $(a\beta - i)\beta + i \in QR(q)$ and $(a - i)\beta + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta - i)\beta + i$ and $x\beta = (a - i)\beta + i$. Hence, $(a\beta - i)\beta^2 + i\beta = (a - i)\beta + i$, which implies that $a\beta(\beta + 1) = i(\beta - 1)$. If $i \in NQR(q)$, then the pair of orthogonal one-factorizations

generated by the starter S_{β} and $-S_{\beta}$ does include a cycle of length four; otherwise don't.

- Case (ii): Suppose that $t_1 = t_3 = 1$ and $t_2 = -1$. In this case $(a\beta i)\beta + i \in NQR(q)$ and $(a i)\beta^{-1} + i \in QR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a i)\beta^{-1} + i$ and $x\beta = (a\beta i)\beta + i$. Hence, $(a\beta i)\beta + i = (a i) + i\beta$, which implies that $2i = a(\beta + 1)$. If $i \in QR(q)$, then the pair of orthogonal one-factorizations generated by the starter S_β and $-S_\beta$ does include a cycle of length four; otherwise don't.
- Case (iii): Suppose that $t_1 = 1$ and $t_2 = t_3 = -1$. In this case $(a\beta i)\beta + i \in QR(q)$ and $(a i)\beta^{-1} + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta i)\beta + i$ and $x\beta = (a i)\beta^{-1} + i$. Hence, $(a\beta i)\beta^2 + i\beta = (a i)\beta^{-1} + i$, which implies that $i = a(\beta + 1)$. If $i \in NQR(q)$, then the pair of orthogonal one-factorizations generated by the starter S_β and $-S_\beta$ does include a cycle of length four; otherwise don't.
- Case (iv): Suppose that $t_2 = t_3 = 1$ and $t_1 = -1$. In this case $(a i)\beta + i \in QR(q)$ and $(a\beta i)\beta^{-1} + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a i)\beta + i$ and $x\beta = (a\beta i)\beta^{-1} + i$. Hence, $(a i)\beta^2 + i\beta = (a\beta i)\beta^{-1} + i$, which implies that $a\beta(\beta + 1) = i(\beta^2 + 1)$. If $i \in QR(q)$, then the pair of orthogonal one-factorizations generated by the starter S_β and $-S_\beta$ does include a cycle of length four; otherwise don't.
- Case (v): Suppose that $t_1 = t_2 = t_3 = -1$. In this case $(a\beta i)\beta^{-1} + i \in QR(q)$ and $(a - i)\beta^{-1} + i \in NQR(q)$. Since $\{x, x\beta\} \in F_0$, where $x \in QR(q)$, then $x = (a\beta - i)\beta^{-1} + i$ and $x\beta = (a - i)\beta^{-1} + i$. Hence, $(a\beta - i) + i\beta = (a - i)\beta^{-1} + i$, which implies that $a(\beta + 1) = -i(\beta - 1)$. If $i \in NQR(q)$, then the pair of orthogonal one-factorizations generated by the starter S_β and $-S_\beta$ does include a cycle of length four; otherwise don't.

It is not hard to check that Case (iii) and Case (v) generate different cycles of length four. On the other hand, Case (iii) generate one cycle of length four and Case (i) and (v) generate the other cycle of length four. Therefore, the pair of orthogonal one-factorizations generated by the starter S_{β} and $-S_{\beta}$ does include exactly two cycles of length four.

Theorem 3.3. Let $q \equiv 3 \pmod{8}$ be an odd prime power such that $q \equiv -1 \pmod{12}$. Then there is a pair of orthogonal $(2, C_4)$ one-factorization of K_{q+1} .

Proof. Let

 $M=\{\beta\in NQR(q)\backslash \{-2,-1,2\}:\beta-1,\beta+1\in NQR(q),\beta^2+1\in QR(q)\}.$

By Lemma 3.1 and Lemma 3.2, we only need to show that M is not the empty set: see proof of Lemma 2.6.

Hence, by Theorem 2.7 and Theorem 3.3, we have the following:

Corollary 3.4. Let $q \equiv 3 \pmod{8}$ be an odd prime power such that $q \equiv -1 \pmod{12}$. Then there are a pair of orthogonal one-factorization of K_{q+1} , \mathcal{F} and \mathcal{G} , such that $F \cup G$ include at most two cycles of length four, for every $F, G \in \mathcal{F} \cup \mathcal{G}$.

Acknowledgment

Research was partially supported by SNI and CONACyT.

References

 C. A. Alfaro, C. Rubio-Montiel and A. Vázquez-Ávila, On two-quotient strong starters for F_q, Util. Math. **112** (2019), 287–302.

- [2] B. A. Anderson, A class of starter induced 1-factorizations, Graphs and combinatorics (Proc. Capital Conf., George Washington Univ., Washington, D.C., 1973), Springer, Berlin, 1974, pp. 180–185. Lecture Notes in Math., Vol. 406.
- [3] B. A. Anderson, P. J. Schellenberg and D. R. Stinson, *The existence of Howell designs of even side*, J. Combin. Theory Ser. A **36** (1984), no. 1, 23–55.
- [4] J. Bao and L. Ji, Two Orthogonal 4-Cycle-Free One-Factorizations of Complete Graphs, Graphs Combin. 35 (2019), no. 2, 373–392.
- [5] D. M. Burton, *Elementary number theory*, sixth ed., McGraw-Hill, 2007.
- [6] C. J. Colbourn and J. H. Dinitz, Handbook of Combinatorial Designs, 2nd ed., Discrete mathematics and its applications, Chapman & Hall/Taylor & Francis, 2007.
- [7] J. F. Dillon and R. A. Morris, A skew Room square of side 257, Utilitas Math. 4 (1973), 187–192.
- [8] J. H. Dinitz, Room n-cubes of low order, J. Austral. Math. Soc. Ser. A 36 (1984), 237–252.
- [9] J. H. Dinitz and P. Dukes, On the structure of uniform onefactorizations from starters in finite fields, Finite Fields Appl. 12 (2006), no. 2, 283–300.
- [10] J. H. Dinitz and D. R. Stinson, *The spectrum of Room cubes*, European J. Combin. 2 (1981), no. 3, 221–230.
- [11] J. H. Dinitz and D. R. Stinson, Some new perfect one-factorizations from starters in finite fields, J. Graph Theory, 13 (1989), no. 4, 405–415.
- [12] K. B. Gross, Some new classes of strong starters, Discrete Math. 12 (1975), no. 3, 225–243.
- [13] G. H. Hardy and E. M. Wright, An introduction to the theory of numbers, sixth ed., Oxford University Press, Oxford, 2008, Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles.

- [14] J. D. Horton, Room designs and one-factorizations, Aequationes Math.
 12 (1981), no. 1, 56–63.
- [15] J. D. Horton, The construction of Kotzig factorizations, Discrete Math.
 43 (1983), no. 2-3, 199–206.
- [16] J. D. Horton, Orthogonal starters in finite abelian groups, Discrete Math. 79 (1990), no. 3, 265–278.
- [17] W. L. Kocay, D. R. Stinson and S. A. Vanstone, On strong starters in cyclic groups, Discrete Math. 56 (1985), no. 1, 45–60.
- [18] M. Meszka, k-cycle free one-factorizations of complete graphs, Electron. J. Combin. 16 (2009), no. 1, 255–258.
- [19] R. C. Mullin and E. Nemeth, An existence theorem for room squares, Canad. Math. Bull. 12 (1969), 493–497.
- [20] R. C. Mullin and E. Nemeth, On furnishing Room squares, J. Combinatorial Theory 7 (1969), 266–272.
- [21] O. Ogandzhanyants, M. Kondratieva and N. Shalaby, Strong Skolem starters, J. Combin. Des., 27 (2019), no. 1, 5–21.
- [22] D. K. Ray-Chaudhuri and R. M. Wilson, Solution of Kirkman's schoolgirl problem, Combinatorics (Proc. Sympos. Pure Math., Vol. XIX, Univ. California, Los Angeles, Calif., 1968), Amer. Math. Soc., Providence, R.I. (1971), 187–203.
- [23] A. Rosa, Perfect 1-factorizations, Math. Slovaca. 69 (2019), 479–496.
- [24] A. Rosa and S. A. Vanstone, On the existence of strong Kirkman cubes of order 39 and block size 3, Algorithms in combinatorial design theory, North-Holland Math. Stud., vol. 114, North-Holland, Amsterdam (1985), no. 1, 309–319.
- [25] R. G. Stanton and R.C. Mullin, Construction of Room squares, Ann. Math. Statist. 39 (1968), 1540–1548.
- [26] D. R. Stinson and S. A. Vanstone, A Kirkman square of order 51 and block size 3, Discrete Math. 55 (1985), no. 1, 107–111.

- [27] A. Vázquez-Ávila, A note on strong Skolem starters, Discrete Math. 343 (2020), no. 2, 111672.
- [28] A. Vázquez-Ávila, On strong Skolem starters for Z_{pq}, AKCE J. Graphs Comb. 17 (2020), No. 3, 988–991.
- [29] A. Vázquez-Ávila, On strong Skolem starters, J. Discret. Math. Sci. Cryptogr. Accepted.
- [30] A. Vázquez-Ávila, A note on two orthogonal totally C₄-free onefactorizations of complete graphs, Bol. Soc. Mat. Mex. 27 (2021), no. 1, 5.