Twist polynomials of delta-matroids

Qi Yan

School of Mathematics China University of Mining and Technology P. R. China Xian'an Jin¹ School of Mathematical Sciences Xiamen University P. R. China

Email:qiyan@cumt.edu.cn; xajin@xmu.edu.cn

Abstract

Recently, Gross, Mansour and Tucker introduced the partial duality polynomial of a ribbon graph and posed a conjecture that there is no orientable ribbon graph whose partial duality polynomial has only one non-constant term. We found an infinite family of counterexamples for the conjecture and showed that essentially these are the only counterexamples. This is also obtained independently by Chumutov and Vignes-Tourneret and they posed a problem: it would be interesting to know whether the partial duality polynomial and the related conjectures would make sence for general delta-matroids. In this paper, we show that partial duality polynomials have delta-matroid analogues. We introduce the twist polynomials of delta-matroids and discuss its basic properties for delta-matroids. We give a characterization of even normal binary delta-matroids whose twist polynomials have only one term and then prove that the twist polynomial of a normal binary deltamatroid contains non-zero constant term if and only if its intersection graph is bipartite.

Keywords: delta-matroid, binary, twist, polynomial

2020 MSC: 05B35, 05C10, 05C31

¹Corresponding author.

1. Introduction

For any ribbon graph G, there is a natural dual ribbon graph G^* , also called *geometric dual*. Chmutov [4] introduced an extension of geometric duality called *partial duality*. Roughly speaking, a partial dual G^A is obtained by forming the geometric dual with respect to only a subset $A \subseteq E(G)$ of a ribbon graph G.

In [10], Gross, Mansour and Tucker introduced the enumeration of the partial duals G^A of a ribbon graph G, by Euler genus ε , over all edge subsets $A \subseteq E(G)$. The associated generating functions, denoted as $\partial \varepsilon_G(z)$, are called *partial duality polynomials* of G. They formulated the following conjecture.

Conjecture 1. [10] There is no orientable ribbon graph having a nonconstant partial duality polynomial with only one non-zero coefficient.

The conjecture is not true. In [13] we found an infinite family of counterexamples. Furthermore, we [14] proved that essentially these are the only counterexamples. This is also obtained independently by Chumutov and Vignes-Tourneret in [5] and they also posed the following question:

Question 2. [5] Ribbon graphs may be considered from the point of view of delta-matroid. In this way the concepts of partial duality and genus can be interpreted in terms of delta-matroids [6, 7]. It would be interesting to know whether the partial duality polynomial and the related conjectures would make sence for general delta-matroids.

In this paper, we show that partial duality polynomials have delta-matroid analogues. We introduce the twist polynomials of delta-matroids and discuss its basic properties and consider Conjecture 1 for delta-matroids. We give a characterization of even normal binary delta-matroids whose twist polynomials have only one term and then prove that the twist polynomial of a normal binary delta-matroid contains non-zero constant term if and only if its intersection graph is bipartite.

2. Preliminaries

2.1. Delta-matroids

A set system is a pair $D = (E, \mathcal{F})$, where E or E(D), is a finite set, called the ground set, and \mathcal{F} , or $\mathcal{F}(D)$, is a collection of subsets of E, called *feasible* sets. A set system D is proper if $\mathcal{F} \neq \emptyset$ and D is trivial if $E = \emptyset$. As introduced by Bouchet in [1], a *delta-matroid* is a proper set system $D = (E, \mathcal{F})$ for which satisfies the symmetric exchange axiom: for all triples (X, Y, u) with $X, Y \in \mathcal{F}$ and $u \in X\Delta Y$, there is a $v \in X\Delta Y$ (possibly v = u) such that $X\Delta\{u, v\} \in \mathcal{F}$. Here $X\Delta Y = (X \cup Y) \setminus (X \cap Y)$ is the usual symmetric difference of sets.

Let $D = (E, \mathcal{F})$ be a delta-matroid. If for any $A_1, A_2 \in \mathcal{F}$, we have $|A_1| = |A_2|$. Then D is said to be a *matroid* and we refer to \mathcal{F} as its *bases*. If a set system forms a matroid M, then we usually denote M by (E, \mathcal{B}) . The rank function of M takes any subset $A \subseteq E(M)$ to the number

$$\max\{|A \cap B| : B \in \mathcal{B}\}.$$

This is written as $r_M(A)$. We say that the rank of M, written r(M), is equal to |B| for any $B \in \mathcal{B}(M)$. It is clear that the rank function of a matroid M on a set E has the following properties [12]:

- 1. If $X \subseteq Y \subseteq E$, then $r_M(X) \leq r_M(Y)$;
- 2. If X and Y are subsets of E, then

$$r_M(X \cup Y) + r_M(X \cap Y) \le r_M(X) + r_M(Y).$$

The nullity of A, written $n_M(A)$, is $|A| - r_M(A)$.

Bouchet [2] defined an analogue of the rank function for delta-matroids. Let $D = (E, \mathcal{F})$ be a delta-matroid. For $A \subseteq E$, define

$$\rho_D(A) := |E| - \min\{|A\Delta F| : F \in \mathcal{F}\}.$$

A delta-matroid is *even* if for every pair F and \tilde{F} of its feasible sets $|F\Delta \tilde{F}|$ is even. Otherwise, we call the delta-matroid *odd*. A delta-matroid is *normal* if the empty set is feasible.

For a delta-matroid $D = (E, \mathcal{F})$, let $\mathcal{F}_{max}(D)$ and $\mathcal{F}_{min}(D)$ be the collection of maximum and minimum cardinality feasible sets of D, respectively. Bouchet [3] showed that the set systems $D_{max} = (E, \mathcal{F}_{max})$ and $D_{min} = (E, \mathcal{F}_{min})$ are matroids. The width of D, denote by w(D), is defined by

$$v(D) := r(D_{max}) - r(D_{min}).$$

Particularly, D is a matroid if and only if w(D) = 0.

A fundamental operation on delta-matroids, introduced by Bouchet in [1], is the twist. Let $D = (E, \mathcal{F})$ be a set system. For $A \subseteq E$, the *twist* of D with respect to A, denoted by D * A, is given by

$$(E, \{A\Delta X : X \in \mathcal{F}\}).$$

The dual of D, written D^* , is equal to D * E. Using the identity

$$(A\Delta C)\Delta(B\Delta C) = A\Delta B$$

it is straightforward to show that the twist of a delta-matroid is a deltamatroid [1]. Note that being even is preserved under taking twists.

Definition 3. The *twist polynomial* of any delta-matroid $D = (E, \mathcal{F})$ is the generating function

$${}^{\partial}w_D(z) := \sum_{A \subseteq E} z^{w(D*A)}$$

that enumerates all twists of D by width.

Definition 4. [7] For delta-matroids $D = (E, \mathcal{F})$ and $\widetilde{D} = (\widetilde{E}, \widetilde{\mathcal{F}})$ with $E \cap \widetilde{E} = \emptyset$, the *direct sum* of D and \widetilde{D} , written $D \oplus \widetilde{D}$, is the delta-matroid defined as

$$D \oplus \widetilde{D} := (E \cup \widetilde{E}, \{F \cup \widetilde{F} : F \in \mathcal{F} \text{ and } \widetilde{F} \in \widetilde{\mathcal{F}}\}).$$

A delta-matroid is *disconnected* if it can be written as $D \oplus \widetilde{D}$ for some non-trivial delta-matroids D and \widetilde{D} , and *connected* otherwise.

Let $D = (E, \mathcal{F})$ be a proper set system. An element $e \in E$ contained in every feasible set of D is said to be a *coloop*, while an element $e \in E$ contained in no feasible set of D is said to be a *loop*.

Let $D = (E, \mathcal{F})$ be a proper set system and $e \in E$. Then D delete by e, denoted $D \setminus e$, is defined as $D \setminus e := (E \setminus e, \mathcal{F}')$, where

$$\mathcal{F}' := \begin{cases} \{F : F \in \mathcal{F}, F \subseteq E \setminus e\}, & \text{if } e \text{ is not a coloop} \\ \{F \setminus e : F \in \mathcal{F}\}, & \text{if } e \text{ is a coloop.} \end{cases}$$

Bouchet [1] has shown that the order in which deletions are performed does not matter. Let $A \subseteq E$. We define $D \setminus A$ as the result of deleting every element of A in any order. The *restriction* of D to A, written $D|_A$, is the delta-matroid $D \setminus (E \setminus A)$. Throughout the paper, we will often omit the set brackets in the case of a single element set. For example, we write D * einstead of $D * \{e\}$, or $D|_e$ instead of $D|_{\{e\}}$.

2.2. Ribbon graphs

We give a brief review of ribbon graphs referring the reader to [8, 9] for further details.

Definition 5 ([9]). A ribbon graph G = (V(G), E(G)) is a (possibly nonorientable) surface with boundary represented as the union of two sets of discs, a set V(G) of vertices, and a set E(G) of edges such that

- 1. The vertices and edges intersect in disjoint line segments;
- 2. Each such line segment lies on the boundary of precisely one vertex and precisely one edge;
- 3. Every edge contains exactly two such line segments.

A bouquet is a ribbon graph with only one vertex. An edge e of a ribbon graph is a *loop* if it is incident with exactly one vertex. A loop is *non-orientable* if together with its incident vertex it forms a Möbius band, and is *orientable* otherwise. A *signed rotation* of a bouquet is a cyclic ordering of the half-edges at the vertex and if the edge is an orientable loop, then we give the same sign + to the corresponding two half-edges, and give the different signs (one +, the other -) otherwise. The sign + is always omitted. See Figure 1 for an example.



Figure 1: The signed rotation of the bouquet is (-1, -2, 3, 4, 2, 1, 3, 4).

Definition 6. [7] Let G = (V, E) be a ribbon graph and let

 $\mathcal{F} := \{F \subseteq E(G) : F \text{ is the edge set of a spanning quasi-tree of } G\}.$ We call $D(G) =: (E, \mathcal{F})$ the delta-matroid of G.

2.3. Binary and intersection graphs

For a finite set E, let C be a symmetric |E| by |E| matrix over GF(2), with rows and columns indexed, in the same order, by the elements of E. Let C[A] be the principal submatrix of C induced by the set $A \subseteq E$. We define the set system $D(C) = (E, \mathcal{F})$ with

$$\mathcal{F} := \{ A \subseteq E : C[A] \text{ is non-singular} \}.$$

By convention $C[\emptyset]$ is non-singular. Then D(C) is a delta-matroid [2]. A delta-matroid is said to be *binary* if it has a twist that is isomorphic to D(C) for some symmetric matrix C over GF(2).

Let $D = (E, \mathcal{F})$ be a delta-matroid. If D = D(C) for some symmetric matrix C over GF(2), that is, D is a normal binary delta-matroid, then we can get C as following [11]:

- 1. Set $C_{v,v} = 1$ if and only if $\{v\} \in \mathcal{F}$. This determines the diagonal entries of C;
- 2. Set $C_{u,v} = 1$ if and only if $\{u\}, \{v\} \in \mathcal{F}$ but $\{u, v\} \notin \mathcal{F}$, or $\{u, v\} \in \mathcal{F}$ but $\{u\}$ and $\{v\}$ are not both in \mathcal{F} . Then the feasible sets of size two determine the off-diagonal entries of C.

Let $D = (E, \mathcal{F})$ be a normal binary delta-matroid. Then there exists a symmetric |E| by |E| matrix C over GF(2) such that D = D(C). The *intersection graph* G_D of D is the graph with vertex set E and in which two vertices u and v of G_D are adjacent if and only if $C_{u,v} = 1$. Recall that a *looped simple graph* is a graph obtained from a simple graph by adding (exactly) one loop to some of its vertices. If D is odd, then G_D is a looped simple graph, and if D is even, then G_D is a simple graph. Note that D is connected if and only if G_D is connected.

Conversely, the adjacency matrix A(G) of a looped simple graph G is the matrix over GF(2) whose rows and columns correspond to the vertices of G; and where, $A(G)_{u,v} = 1$ if and only if u and v are adjacent in G and $A(G)_{u,u} = 1$ if and only if there is a loop at u. Let D be a normal binary delta-matroid. It obvious that $D = D(A(G_D))$.

3. Main results

Proposition 7. Let $D = (E, \mathcal{F})$ and $\widetilde{D} = (\widetilde{E}, \widetilde{\mathcal{F}})$ be two delta-matroids and $A \subseteq E$. Then

1. $\partial w_D(1) = 2^{|E|};$

2.
$${}^{\partial}w_D(z) = {}^{\partial}w_{D*A}(z);$$

3. ${}^{\partial}w_{D\oplus\widetilde{D}}(z) = {}^{\overline{\partial}}w_D(z) {}^{\partial}w_{\widetilde{D}}(z).$

Proof. For (1), the evaluation ${}^{\partial}w_D(1)$ counts the total number of twists, which is $2^{|E|}$. For (2), this is because the sets of all twists of D and D * A are the same. For (3), the underlying phenomenon is the additivity of width over the direct sum. It follows immediately that for any subset $B \subseteq E \cup \widetilde{E}$, we have

$$(D \oplus \widetilde{D}) * B = D * (B \cap E) \oplus \widetilde{D} * (B \cap \widetilde{E}),$$

from which formula (3) follows.

Remark 8. By Proposition 7, we can observe that analyzing the twist polynomials of all delta-matroids is equivalent to analyzing normal delta-matroids. Consequently, it is natural to focus on normal delta-matroids.

Lemma 9. [7] Let G = (V, E) be a ribbon graph, $A \subseteq E$ and $e \in E$. Then $D(G^A) = D(G) * A$ and $\varepsilon(G) = w(D(G))$.

Lemma 10. Let G = (V, E) be a ribbon graph. Then $\partial w_{D(G)}(z) = \partial \varepsilon_G(z)$.

Proof. By Lemma 9, for any $A \subseteq E$,

$$w(D(G) * A) = w(D(G^A)) = \varepsilon(G^A).$$

Hence $\partial w_{D(G)}(z) = \partial \varepsilon_G(z)$.

Theorem 11. If two normal binary delta-matroids D and D have the same intersection graph, then ${}^{\partial}w_D(z) = {}^{\partial}w_{\widetilde{D}}(z)$.

Proof. Since $G_D = G_{\widetilde{D}}$, $D = D(A_{G_D})$ and $\widetilde{D} = D(A_{G_{\widetilde{D}}})$, it follows that $D = \widetilde{D}$. Therefore ${}^{\partial}w_D(z) = {}^{\partial}w_{\widetilde{D}}(z)$.

Theorem 12. Let *B* and \widetilde{B} be two bouquets. If $G_{D(B)} = G_{D(\widetilde{B})}$, then ${}^{\partial} \varepsilon_B(z) = {}^{\partial} \varepsilon_{\widetilde{B}}(z)$.

Proof. If $G_{D(B)} = G_{D(\widetilde{B})}$, then $D(B) = D(\widetilde{B})$. For any $A \subseteq E(B)$, we denoted its corresponding subset of $E(\widetilde{B})$ by \widetilde{A} . By Lemma 9,

$$D(B^A) = D(B) * A = D(\widetilde{B}) * \widetilde{A} = D(\widetilde{B}^A).$$

We have $w(D(B^A)) = w(D(\widetilde{B}^{\widetilde{A}}))$. Since $w(D(B^A)) = \varepsilon(B^A)$ and $w(D(\widetilde{B}^{\widetilde{A}})) = \varepsilon(\widetilde{B}^{\widetilde{A}})$, it follows that $\varepsilon(B^A) = \varepsilon(\widetilde{B}^{\widetilde{A}})$. Thus ${}^{\partial}\varepsilon_B(z) = {}^{\partial}\varepsilon_{\widetilde{B}}(z)$.

Lemma 13. [13] Let B_t be a bouquet with the signed rotation

We have

$${}^{\partial}\varepsilon_{B_t}(z) = \begin{cases} 2^t z^{t-1}, & \text{if } t \text{ is odd} \\ 2^{t-1} z^t + 2^{t-1} z^{t-2}, & \text{if } t \text{ is even.} \end{cases}$$

Proposition 14. Let D be a normal binary delta-matroid and let v be the number of vertices of G_D . If G_D is a complete graph, then

$${}^{\partial}w_D(z) = \begin{cases} 2^v z^{v-1}, & \text{if } v \text{ is odd} \\ 2^{v-1} z^v + 2^{v-1} z^{v-2}, & \text{if } v \text{ is even}. \end{cases}$$

Proof. By Theorem 11, we just need to find a normal binary delta-matroid \widetilde{D} such that $G_{\widetilde{D}} = G_D$ and the evaluation ${}^{\partial}w_{\widetilde{D}}(z)$ is easy to obtained. Let B_v be a bouquet with the signed rotation (1, 2, 3, ..., v, 1, 2, 3, ..., v). Obviously, $G_{D(B_v)} = G_D = K_v$. By Lemma 10 and Theorem 11,

$${}^{\partial}w_D(z) = {}^{\partial}w_{D(B_v)}(z) = {}^{\partial}\varepsilon_{B_v}(z).$$

Therefore $\partial w_D(z)$ can be obtained by Lemma 13.

Theorem 15. Let $D = (E, \mathcal{F})$ be a delta-matroid. Then ${}^{\partial}w_D(z) = k$ for some integer k if and only if $|\mathcal{F}| = 1$.

Proof. The sufficiency is straightforward. To prove the necessity, suppose that $|\mathcal{F}| \geq 2$, then there exist $A_1, A_2 \in \mathcal{F}$ such that $A_1 \neq A_2$. Since $\partial w_D(z) = k$, we have w(D) = 0. Thus $|A_1| = |A_2|$. Then for any $x \in A_1 \setminus A_2$ we have $A_1 \setminus x, A_2 \cup x \in \mathcal{F}(D * x)$. Obviously, $|A_1 \setminus x| \neq |A_2 \cup x|$, a contradiction, since $\partial w_D(z) = \partial w_{D*x}(z) = k$ by Proposition 7.

Lemma 16. [7] Let $D = (E, \mathcal{F})$ be a delta-matroid and $A \subseteq E$. Then

$$w(D|_A) = \rho_D(A) - r_{D_{min}}(A) - n_{D_{min}}(E) + n_{D_{min}}(A).$$

Lemma 17. Let $D = (E, \mathcal{F})$ be a normal delta-matroid and $A \subseteq E$. Then

$$w(D * A) = w(D|_A) + w(D|_{A^c})$$

Proof. Since D is a normal delta-matroid, it follows that $D_{min} = (E, \{\emptyset\})$. We have $r_{D_{min}}(A) = r_{D_{min}}(A^c) = 0$, $n_{D_{min}}(E) = |E|, n_{D_{min}}(A) = |A|$ and $n_{D_{min}}(A^c) = |A^c|$. Then by Lemma 16,

$$\begin{split} w(D|_{A}) + w(D|_{A^{c}}) \\ &= \rho_{D}(A) - r_{D_{min}}(A) - n_{D_{min}}(E) + n_{D_{min}}(A) + \\ \rho_{D}(A^{c}) - r_{D_{min}}(A^{c}) - n_{D_{min}}(E) + n_{D_{min}}(A^{c}) \\ &= \rho_{D}(A) + \rho_{D}(A^{c}) - |E| \\ &= |E| - min\{|A\Delta F| : F \in \mathcal{F}\} + \\ |E| - min\{|A^{c}\Delta F| : F \in \mathcal{F}\} - |E| \\ &= |E| - min\{|A^{c}\Delta F| : F \in \mathcal{F}\} - min\{|A\Delta F| : F \in \mathcal{F}\} \\ &= |E| - r((D * A^{c})_{min}) - r((D * A)_{min}) \\ &= |E| - r((D * A)_{min}^{*}) - r((D * A)_{min}) \\ &= r((D * A)_{max}) - r((D * A)_{min}) = w(D * A). \end{split}$$

Remark 18. This is not right for non-normal delta-matroids. For example, let $D = (\{1,2\},\{\{1\},\{2\}\})$. It is easy to check that $D * 1 = (\{1,2\},\{\emptyset,\{1,2\}\})$, $D|_1 = (\{1\},\{\{1\}\})$ and $D|_2 = (\{2\},\{\{2\}\})$. Obviously, w(D * 1) = 2 and $w(D|_1) = w(D|_2) = 0$. Note that $w(D * 1) \neq w(D|_1) + w(D|_2)$.

Theorem 19. Let $D = (E, \mathcal{F})$ be a binary normal delta-matroid. Then ${}^{\partial}w_D(z)$ contains non-zero constant term if and only if G_D is a bipartite graph.

Proof. Since ${}^{\partial}w_D(z)$ contains non-zero constant term, it follows that D is a twist of a matroid. On account of the property that twist preserving evenness, we have that D is even and hence G_D is a simple graph. Suppose that G_D is not bipartite. Then G_D contains an odd cycle P of length more than or equal to 3. We denote by A the subset of E corresponding to the vertices of P. It is obvious that deleting can not increase the width. Then for any subset B of E, we have $w(D|_{B\cap A}) \leq w(D|_B)$ and $w(D|_{B^c\cap A}) \leq w(D|_{B^c})$. Since D is a normal binary delta-matroid, we know that D = D(C) for some symmetric matrix C over GF(2). Note that there are $e, f \in B \cap A$ or $e, f \in B^c \cap A$ such that

$$C[\{e,f\}] = \begin{array}{c} e & f \\ f \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then

$$w(D|_B) + w(D|_{B^c}) \ge w(D|_{B \cap A}) + w(D|_{B^c \cap A}) > 0.$$

Thus by Lemma 17 $w(D * B) = w(D|_B) + w(D|_{B^c}) > 0$, a contradiction.

Conversely, if G_D is bipartite and non-trivial, then its vertex set can be partitioned into two subsets X and Y so that every edge of G_D has one end in X and the other end in Y. For these two subsets X and Y of the vertex set of G_D , we denoted these two corresponding subset of E also by X and Y. Obviously, $X \cup Y = E$, $X \cap Y = \emptyset$ and $w(D|_X) = w(D|_Y) = 0$. Thus $w(D * X) = w(D|_X) + w(D|_Y) = 0$ by Lemma 17. Hence $\partial w_D(z)$ contains non-zero constant term.

Theorem 20. Let $D = (E, \mathcal{F})$ be a connected even normal binary deltamatroid. Then ${}^{\partial}w_D(z) = mz^k$ if and only if G_D is a complete graph of odd order.

Proof. The sufficiency is easily verified by Proposition 14. For necessity,

$$D = \begin{cases} (\{e\}, \{\emptyset\}), & \text{if } |E| = 1\\ (\{e, f\}, \{\emptyset, \{e, f\}\}), & \text{if } |E| = 2. \end{cases}$$

Then the result is easily verified when $|E| \in \{1, 2\}$. Assume that $|E| \ge 3$. Let $e, f, g \in E$. We consider three claims:

Claim 1. If $\{e, f\} \in \mathcal{F}$, we have $r_{D^*_{min}}(\{e, f\}) = 1$ and $r_{D^*_{min}}(e) = 1$.

Proof of Claim 1. For any $A \in \mathcal{F}_{max}$, we observe that $A \cap \{e, f\} \neq \emptyset$. Otherwise, $\emptyset, A \cup \{e, f\} \in \mathcal{F}(D * \{e, f\})$. Since $\{e, f\} \in \mathcal{F}$, it follows that $\emptyset \in \mathcal{F}(D * \{e, f\})$. Then $w(D * \{e, f\}) > w(D)$, this contradicts $\partial w_D(z) = mz^k$. Furthermore, we observe that there exists $B \in \mathcal{F}_{max}$ such that $e \notin B$. Otherwise, $r(D * e_{max}) = r(D_{max}) - 1$. Since $e \notin \mathcal{F}$ and $\emptyset \in \mathcal{F}$, we have $r(D * e_{min}) = 1$. Then w(D * e) = w(D) - 2, this also contradicts $\partial w_D(z) = mz^k$. Consequently, for any $A \in \mathcal{F}^*_{min}$, $A \cap \{e, f\} \neq \{e, f\}$ and there exists $B \in \mathcal{F}^*_{min}$ such that $e \in B$. Thus $r_{D^*_{min}}(\{e, f\}) = 1$ and $r_{D^*_{min}}(e) = 1$.

Claim 2. If $\{e, f\} \notin \mathcal{F}$, we have $r_{D^*_{min}}(\{e, f\}) = 2$.

Proof of Claim 2. There exists $A \in \mathcal{F}_{max}$ such that $\{e, f\} \cap A = \emptyset$. Otherwise, $r(D * \{e, f\}_{max}) \leq r(D_{max})$. Since $\{e, f\} \notin \mathcal{F}$ and $\emptyset \in \mathcal{F}$, we have $r(D * \{e, f\}_{min}) = 2$. Then $w(D * \{e, f\}) \leq w(D) - 2$, a contradiction. Consequently, there exists $A \in \mathcal{F}^*_{min}$ such that $\{e, f\} \in A$. Thus $r_{D^*_{min}}(\{e, f\}) = 2$.

Claim 3. E does not contain e, f, g such that $\{e, f\}, \{e, g\} \in \mathcal{F}$ and $\{f, g\} \notin \mathcal{F}$.

Proof of Claim 3. Assume that Claim 3 is not true. Since $\{e, f\}, \{e, g\} \in \mathcal{F}$, it follows that $r_{D^*_{min}}(\{e, f\}) = 1$ and $r_{D^*_{min}}(\{e, g\}) = 1$ by Claim 1. Then

$$r_{D^*_{min}}(\{e, f\} \cup \{e, g\}) + r_{D^*_{min}}(\{e, f\} \cap \{e, g\})$$

$$= r_{D^*_{min}}(\{e, f, g\}) + r_{D^*_{min}}(e)$$

$$\leq r_{D^*_{min}}(\{e, f\}) + r_{D^*_{min}}(\{e, g\})$$

$$= 2.$$

Hence $r_{D^*_{min}}(\{e, f, g\}) \leq 1$. Since $\{f, g\} \notin \mathcal{F}$, we have $r_{D^*_{min}}(\{f, g\}) = 2$ by Claim 2. But

$$r_{D^*_{min}}(\{f,g\}) \le r_{D^*_{min}}(\{e,f,g\}) \le 1,$$

a contradiction.

Suppose that G_D is not a complete graph. Note that G_D is connected. Then there is a vertex set v_e, v_f, v_g of G_D such that the induced subgraph $G_D(\{v_e, v_f, v_g\})$ is a 2-path. We may assume without loss of generality that the degree of v_e is 2 in $G_D(\{v_e, v_f, v_g\})$. Since D is a normal binary deltamatroid, we know that D = D(C) for some symmetric matrix C over GF(2). Then

$$C[\{e, f, g\}] = \begin{array}{c} e & f & g \\ e \\ f \\ g \\ \end{array} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \\ \end{pmatrix}.$$

Thus $\{e, f\}, \{e, g\} \in \mathcal{F}$ and $\{f, g\} \notin \mathcal{F}$, this contradicts Claim 3. Therefore G_D is a complete graph.

Corollary 21. Let $D = (E, \mathcal{F})$ be an even normal binary delta-matroid. Then ${}^{\partial}w_D(z) = mz^k$ if and only if each connected component of G_D is a complete graph of odd order.

Proof. The proof is straightforward by Proposition 7 and Theorem 20. \Box

Acknowledgements

This work is supported by NSFC (No. 11671336) and the Fundamental Research Funds for the Central Universities (Nos. 20720190062, 2021QN1037).

References

- A. Bouchet, Greedy algorithm and symmetric matroids, *Math. Program.* 38 (1987) 147–159.
- [2] A. Bouchet, Representability of Δ-matroids, Colloq. Math. Soc. János Bolyai (1987) 167–182.
- [3] A. Bouchet, Maps and delta-matroids, *Discrete Math.* **78** (1989) 59–71.
- [4] S. Chmutov, Generalized duality for graphs on surfaces and the signed Bollobás-Riordan polynomial, J. Combin. Theory Ser. B 99 (2009) 617– 638.
- [5] S. Chmutov and F. Vignes-Tourneret, On a conjecture of Gross, Mansour and Tucker, *European J. Combin.* 97 (2021) 103368.
- [6] C. Chun, I. Moffatt, S. D. Noble and R. Rueckriemen, On the interplay between embedded graphs and delta-matroids, *Proc. London Math. Soc.* 118 (2019) 3: 675–700.
- [7] C. Chun, I. Moffatt, S. D. Noble and R. Rueckriemen, Matroids, deltamatroids and embedded graphs, J. Combin. Theory Ser. A 167 (2019) 7–59.
- [8] J. A. Ellis-Monaghan and I. Moffatt, Twisted duality for embedded graphs, Trans. Amer. Math. Soc. 364 (2012) 1529–1569.
- [9] J. A. Ellis-Monaghan and I. Moffatt, Graphs on surfaces, Springer New York, 2013.
- [10] J. L. Gross, T. Mansour and T. W. Tucker, Partial duality for ribbon graphs, I: Distributions, *European J. Combin.* 86 (2020) 103084.
- [11] I. Moffatt, Surveys in Combinatorics, 2019: Delta-matroids for graph theorists, 2019.
- [12] J. Oxley, Matroid theory, 2nd edn, Oxford University Press, New York, 2011.
- [13] Q. Yan and X. Jin, Counterexamples to a conjecture by Gross, Mansour and Tucker on partial-dual genus polynomials of ribbon graphs, *European J. Combin.* **93** (2021) 103285.

[14] Q. Yan and X. Jin, Partial-dual genus polynomials and signed intersection graphs, Preprint arXiv:math.CO/2102.01823.