Turk J Math (2023) 47: 794 – 815 © TÜBİTAK doi:10.55730/1300-0098.3394

# Clairaut Riemannian maps

## Kiran MEENA, Akhilesh YADAV

Department of Mathematics, Institute of Science, Banaras Hindu University, Varanasi, India

Received: 31.08.2022 • Accepted/Published Online: 14.02.2023 • Final Version: 09.03.2023

Abstract: In this paper, first we define Clairaut Riemannian map between Riemannian manifolds by using a geodesic curve on the base space and find necessary and sufficient conditions for a Riemannian map to be Clairaut with a non-trivial example. We also obtain necessary and sufficient condition for a Clairaut Riemannian map to be harmonic. Thereafter, we study Clairaut Riemannian map from Riemannian manifold to Ricci soliton with a non-trivial example. We obtain scalar curvatures of  $rangeF_*$  and  $(rangeF_*)^{\perp}$  by using Ricci soliton. Further, we obtain necessary conditions for the leaves of  $rangeF_*$  to be almost Ricci soliton and Einstein. We also obtain necessary condition for the vector field  $\dot{\beta}$  to be Killing on  $(rangeF_*)^{\perp}$ , where  $\beta$  is a geodesic curve on the base space of Clairaut Riemannian map. Also, we obtain necessary condition for the mean curvature vector field of  $rangeF_*$  to be constant. Finally, we introduce Clairaut anti-invariant Riemannian map from Riemannian manifold to Kähler manifold, and obtain necessary and sufficient condition for an anti-invariant Riemannian map to be Clairaut with a non-trivial example. Further, we find necessary condition for  $rangeF_*$  to be minimal and totally geodesic. We also obtain necessary and sufficient condition for Clairaut anti-invariant Riemannian maps to be harmonic.

**Key words:** Riemannian manifold, Kähler manifold, Riemannian map, Clairaut Riemannian map, anti-invariant Riemannian map, Ricci soliton

### 1. Introduction

The geometry of Riemannian submersions has been discussed widely in [8]. In 1992, Fischer introduced Riemannian map between Riemannian manifolds as a generalization of an isometric immersion and Riemannian submersion that satisfies the well-known generalized eikonal equation  $||F_*||^2 = rankF$ , which is a bridge between geometric optics and physical optics [9]. Further, the geometry of Riemannian maps was investigated in [2, 3, 20–26].

An important Clairaut's relation states that  $\tilde{r}sin\theta$  is constant, where  $\theta$  is the angle between the velocity vector of a geodesic and a meridian, and  $\tilde{r}$  is the distance to the axis of a surface of revolution. In 1972, Bishop defined Clairaut Riemannian submersion with connected fibers and gave a necessary and sufficient condition for a Riemannian submersion to be Clairaut Riemannian submersion [5]. Further, Clairaut submersions were studied in [1, 12, 14]. In [25], Şahin introduced Clairaut Riemannian map by using a geodesic curve on the total space and obtained necessary and sufficient conditions for Riemannian map to be Clairaut Riemannian map. Further, Şahin gave an open problem to find characterizations for Clairaut Riemannian maps (see [26], page

<sup>\*</sup>Correspondence: kirankapishmeena@gmail.com 2010 AMS Mathematics Subject Classification: 53B20, 53B35, 53C25.

165, open problem 2). In Section 3, we introduce a new type of Clairaut Riemannian map by using a geodesic curve on the base space and obtain necessary and sufficient conditions for a Riemannian map to be Clairaut Riemannian map.

A Riemannian manifold  $(N, g_2)$  is called a Ricci soliton [11] if there exists a smooth vector field  $Z_1$  (called potential vector field) on N such that  $\frac{1}{2}(L_{Z_1}g_2)(X_1,Y_1) + Ric(X_1,Y_1) + \lambda g_2(X_1,Y_1) = 0$ , where  $L_{Z_1}$  is the Lie derivative of the metric tensor of  $g_2$  with respect to  $Z_1$ , Ric is the Ricci tensor of  $(N, g_2)$ ,  $\lambda$  is a constant function and  $X_1$ ,  $Y_1$  are arbitrary vector fields on N. We shall denote a Ricci soliton by  $(N, g_2, Z_1, \lambda)$ . The Ricci soliton  $(N, g_2, Z_1, \lambda)$  is said to be shrinking, steady or expanding accordingly as  $\lambda < 0$ ,  $\lambda = 0$  or  $\lambda > 0$ , respectively. It is obvious that a trivial Ricci soliton is an Einstein manifold [4] with  $Z_1$  zero or Killing (Lie derivative of metric tensor  $g_2$  with respect to  $Z_1$  is vanishes). Ricci soliton can be used to solve the Poincaré conjecture [17]. A Ricci soliton  $(N, g_2, Z_1, \lambda)$  becomes an almost Ricci soliton [18] if the function  $\lambda$  is a variable. The Ricci soliton  $(N, g_2, Z_1, \lambda)$  is said to be a gradient Ricci soliton if the potential vector field  $Z_1$  is the gradient of some smooth function f on N, which is denoted by  $(N, g_2, f, \lambda)$ . Moreover, a non-Killing tangent vector field  $Z_1$  on a Riemannian manifold  $(N, g_2)$  is called conformal [7] if it satisfies  $L_{Z_1}g_2 = 2fg_2$ , where f is called the potential function of  $Z_1$ . The submersions and Riemannian maps from a Ricci soliton to a Riemannian manifold were studied in [10, 13, 15, 29, 30]. In [32], present authors introduced Riemannian map from a Riemannian manifold to a Ricci soliton.

In [28], Watson studied almost Hermitian submersions. In [23], Şahin introduced holomorphic Riemannian map as generalization of holomorphic submersion and holomorphic submanifold. In [2, 3, 20, 22] invariant, anti-invariant and semi-invariant Riemannian maps were studied from a Riemannian manifold to a Kähler manifold. Recently, present authors introduced Clairaut invariant Riemannian map from a Riemannian manifold to a Kähler manifold in [31]. In Section 5, we introduce Clairaut anti-invariant Riemannian map from a Riemannian manifold to a Kähler manifold.

## 2. Preliminaries

In this section, we recall the notion of Riemannian map between Riemannian manifolds and give a brief review of basic facts.

Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a smooth map between Riemannian manifolds such that  $0< rank F\le min\{m,n\}$ , where dim(M)=m and dim(N)=n. We denote the kernel space of  $F_*$  by  $\nu_p=ker F_{*p}$  at  $p\in M$  and consider the orthogonal complementary space  $\mathcal{H}_p=(ker F_{*p})^\perp$  to  $ker F_{*p}$  in  $T_pM$ . Then the tangent space  $T_pM$  of M at p has the decomposition  $T_pM=(ker F_{*p})\oplus (ker F_{*p})^\perp=\nu_p\oplus \mathcal{H}_p$ . We denote the range of  $F_*$  by  $range F_*$  at  $p\in M$  and consider the orthogonal complementary space  $(range F_{*p})^\perp$  to  $range F_{*p}$  in the tangent space  $T_{F(p)}N$  of N at  $F(p)\in N$ . Since  $rank F\leq min\{m,n\}$ , we have  $(range F_{*p})^\perp\neq\{0\}$ . Thus the tangent space  $T_{F(p)}N$  of N at  $F(p)\in N$  has the decomposition  $T_{F(p)}N=(range F_{*p})\oplus (range F_{*p})^\perp$ . Then F is called Riemannian map at  $p\in M$  if the horizontal restriction  $F_{*p}^h:(ker F_{*p})^\perp\to (range F_{*p})$  is a linear isometry between the spaces  $((ker F_{*p})^\perp,g_{1(p)}|_{(ker F_{*p})^\perp})$  and  $(range F_{*p},g_{2(p_1)}|_{(range F_{*p})})$ , where  $F(p)=p_1$ . In other words,  $F_*$  satisfies

$$g_2(F_*X, F_*Y) = g_1(X, Y), \tag{2.1}$$

for all X, Y vector field tangent to  $\Gamma(ker F_{*p})^{\perp}$ . It follows that isometric immersions and Riemannian submer-

sions are particular Riemannian maps with  $ker F_* = \{0\}$  and  $(range F_*)^{\perp} = \{0\}$ , respectively. The differential map  $F_*$  of F can be viewed as a section of bundle  $Hom(TM, F^{-1}TN) \to M$ , where  $F^{-1}TN$  is the pullback bundle whose fibers at  $p \in M$  is  $(F^{-1}TN)_p = T_{F(p)}N$ ,  $p \in M$ . The bundle  $Hom(TM, F^{-1}TN)$  has a con-

nection  $\nabla$  induced from the Levi-Civita connection  $\nabla^M$  and the pullback connection  $\overset{N}{\nabla^F}$ . Then the second fundamental form of F is given by [16]

$$(\nabla F_*)(X,Y) = \nabla_X^F F_* Y - F_* (\nabla_X^M Y), \tag{2.2}$$

for all  $X,Y \in \Gamma(TM)$ , where  $\overset{N}{\nabla_X^F}F_*Y \circ F = \overset{N}{\nabla_{F_*X}}F_*Y$ . It is known that the second fundamental form is symmetric. In [20] Şahin proved that  $(\nabla F_*)(X,Y)$  has no component in  $rangeF_*$ , for all  $X,Y \in \Gamma(kerF_*)^{\perp}$ . More precisely, we have

$$(\nabla F_*)(X,Y) \in \Gamma(rangeF_*)^{\perp}. \tag{2.3}$$

The tension field of F is defined to be the trace of the second fundamental form of F, i.e.  $\tau(F) = trace(\nabla F_*) = \sum_{i=1}^{m} (\nabla F_*)(e_i, e_i)$ , where m = dim(M) and  $\{e_1, e_2, ..., e_m\}$  is the orthonormal frame on M. Moreover, a map  $F: (M^m, g_1) \to (N^n, g_2)$  between Riemannian manifolds is harmonic if and only if the tension field of F vanishes at each point  $p \in M$ .

**Lemma 2.1** [21] Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a Riemannian map between Riemannian manifolds. Then the tension field of F is given by  $\tau(F)=-rF_*(H)+(m-r)H_2$ , where  $r=\dim(\ker F_*)$ ,  $(m-r)=\operatorname{rank} F$ , H and  $H_2$  are the mean curvature vector fields of the distribution  $\ker F_*$  and  $\operatorname{range} F_*$ , respectively.

**Lemma 2.2** [22] Let  $F:(M,g_1) \to (N,g_2)$  be a Riemannian map between Riemannian manifolds. Then F is umbilical Riemannian map if and only if

$$(\nabla F_*)(X,Y) = q_1(X,Y)H_2,$$

for  $X,Y\in\Gamma(ker F_*)^{\perp}$  and  $H_2$  is the mean curvature vector field of range  $F_*$  .

For any vector field X on M and any section V of  $(rangeF_*)^{\perp}$ , we have  $\nabla_X^{F\perp}V$ , which is the orthogonal projection of  $\nabla_X^N V$  on  $(rangeF_*)^{\perp}$ , where  $\nabla^{F\perp}$  is linear connection on  $(rangeF_*)^{\perp}$  such that  $\nabla^{F\perp}g_2 = 0$ . Now, for a Riemannian map F we define  $\mathcal{S}_V$  as ([24], p. 188)

$$\nabla_{F_*X}^N V = -\mathcal{S}_V F_* X + \nabla_X^{F\perp} V, \tag{2.4}$$

where  $\nabla^N$  is Levi-Civita connection on N,  $\mathcal{S}_V F_* X$  is the tangential component (a vector field along F) of  $\nabla^N_{F_* X} V$ . Thus at  $p \in M$ , we have  $\nabla^N_{F_* X} V(p) \in T_{F(p)} N$ ,  $\mathcal{S}_V F_* X \in F_{*p}(T_p M)$  and  $\nabla^{F^{\perp}}_X V(p) \in (F_{*p}(T_p M))^{\perp}$ . It is easy to see that  $\mathcal{S}_V F_* X$  is bilinear in V, and  $F_* X$  at p depends only on  $V_p$  and  $F_{*p} X_p$ . Hence from (2.2) and (2.4), we obtain

$$g_2(S_V F_* X, F_* Y) = g_2(V, (\nabla F_*)(X, Y)), \tag{2.5}$$

for  $X, Y \in \Gamma(kerF_*)^{\perp}$  and  $V \in \Gamma(rangeF_*)^{\perp}$ , where  $S_V$  is self-adjoint operator.

### 3. Clairaut Riemannian map between Riemannian manifolds

In this section, we define Clairaut Riemannian map between Riemannian manifolds by using a geodesic curve [6] on the base space and investigate geometry.

The notion of Clairaut Riemannian map was defined by Şahin in [25]. According to the definition, a Riemannian map  $F:(M,g_1)\to (N,g_2)$  between Riemannian manifolds is called Clairaut Riemannian map if there is a function  $\tilde{r}:M\to\mathbb{R}^+$  such that for every geodesic  $\alpha$  on M, the function  $(\tilde{r}\circ\alpha)\sin\theta$  is constant, where, for all t,  $\theta(t)$  is the angle between  $\dot{\alpha}(t)$  and the horizontal space at  $\alpha(t)$ .

Thus, the notion of Clairaut Riemannian map comes from a geodesic curve on a surface of revolution. Therefore, we are going to give a definition of Clairaut Riemannian map by using geodesic curve on the base space.

**Definition 3.1** A Riemannian map  $F:(M,g_1)\to (N,g_2)$  between Riemannian manifolds is called Clairaut Riemannian map if there is a function  $\tilde{s}:N\to\mathbb{R}^+$  such that for every geodesic  $\beta$  on N, the function  $(\tilde{s}\circ\beta)\sin\omega(t)$  is constant, where,  $F_*X\in\Gamma(rangeF_*)$  for  $X\in\Gamma(kerF_*)^\perp$  and  $V\in\Gamma(rangeF_*)^\perp$  are components of  $\dot{\beta}(t)$ , and  $\omega(t)$  is the angle between  $\dot{\beta}(t)$  and V for all t.

**Note:** For all  $U, V \in \Gamma(rangeF_*)^{\perp}$  we define

$$\nabla_U^N V = \mathcal{R}(\nabla_U^N V) + \nabla_U^{F\perp} V,$$

where  $\mathcal{R}(\nabla_U^N V)$  and  $\nabla_U^{F\perp} V$  denote  $range F_*$  and  $(range F_*)^{\perp}$  part of  $\nabla_U^N V$ , respectively. Therefore  $(range F_*)^{\perp}$  is totally geodesic if and only if

$$\nabla_{U}^{N}V = \nabla_{U}^{F\perp}V.$$

Note that from now, throughout the paper, we are assuming  $(rangeF_*)^{\perp}$  is totally geodesic.

**Lemma 3.2** Let  $F:(M,g_1)\to (N,g_2)$  be a Riemannian map between Riemannian manifolds and  $\alpha:I\to M$  be a geodesic curve on M. Then the curve  $\beta=F\circ\alpha$  is geodesic curve on N if and only if

$$(\nabla F_*)(X,X) + \nabla_X^{F\perp} V + \nabla_V^{F\perp} V = 0, \tag{3.1}$$

$$-S_V F_* X + F_* (\nabla_X^M X) + \nabla_V^N F_* X = 0, \tag{3.2}$$

where  $F_*X \in \Gamma(rangeF_*)$ ,  $V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$  and  $\nabla^N$  is Levi-Civita connection on N and  $\nabla^{F\perp}$  is a linear connection on  $(rangeF_*)^{\perp}$ .

**Proof** Let  $\alpha: I \to M$  be a geodesic on M with  $U(t) = \nu \dot{\alpha}(t)$  and  $X(t) = \mathcal{H}\dot{\alpha}(t)$ . Let  $\beta = F \circ \alpha$  be a geodesic on N with  $F_*X \in \Gamma(rangeF_*)$  and  $V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$ . Now,

$$\nabla^{N}_{\dot{\beta}}\dot{\beta} = \nabla^{N}_{F_{*}X+V}(F_{*}X+V),$$

which implies

$$\nabla^N_{\dot{\beta}}\dot{\beta} = \nabla^N_{F_*X}F_*X + \nabla^N_{F_*X}V + \nabla^N_VF_*X + \nabla^N_VV.$$

Using (2.4) in above equation, we get

$$\nabla_{\dot{\beta}}^{N}\dot{\beta} = \nabla_{X}^{F}F_{*}X \circ F + (-\mathcal{S}_{V}F_{*}X + \nabla_{X}^{F\perp}V) + \nabla_{V}^{N}F_{*}X + \nabla_{V}^{N}V.$$

Using (2.2) in above equation, we get

$$\nabla^{N}_{\dot{\beta}}\dot{\beta} = (\nabla F_{*})(X,X) + F_{*}(\nabla^{M}_{X}X) - \mathcal{S}_{V}F_{*}X + \nabla^{F\perp}_{X}V + \nabla^{N}_{V}F_{*}X + \nabla^{N}_{V}V. \tag{3.3}$$

Since  $(rangeF_*)^{\perp}$  is totally geodesic, (3.3) can be written as

$$\nabla^{N}_{\dot{\beta}}\dot{\beta} = (\nabla F_{*})(X,X) + F_{*}(\nabla^{M}_{X}X) - \mathcal{S}_{V}F_{*}X + \nabla^{F\perp}_{X}V + \nabla^{N}_{V}F_{*}X + \nabla^{F\perp}_{V}V. \tag{3.4}$$

Now  $\beta$  is geodesic on N if and only if  $\nabla^N_{\dot{\beta}}\dot{\beta}=0$ . Then (3.4) implies  $(\nabla F_*)(X,X)+F_*(\nabla^M_XX)-\mathcal{S}_VF_*X+\nabla^{F\perp}_XV+\nabla^N_VF_*X+\nabla^{F\perp}_VV=0$ , which completes the proof.

**Theorem 3.3** Let  $F:(M,g_1) \to (N,g_2)$  be a Riemannian map between Riemannian manifolds such that  $rangeF_*$  is connected and  $\alpha$ ,  $\beta = F \circ \alpha$  are geodesic curves on M and N, respectively. Then F is Clairaut Riemannian map with  $\tilde{s} = e^g$  if and only if any one of the following conditions holds:

- (i)  $S_V F_* X = -V(g) F_* X$ , where  $F_* X \in \Gamma(range F_*), V \in \Gamma(range F_*)^{\perp}$  are components of  $\dot{\beta}(t)$ .
- (ii) F is umbilical map, and has  $H_2 = -\nabla^N g$ , where g is a smooth function on N and  $H_2$  is the mean curvature vector field of range  $F_*$ .

**Proof** First we prove F is a Clairaut Riemannian map with  $\tilde{s} = e^g$  if and only if for any geodesic  $\beta: I \to N$  with tangential components  $F_*X \in \Gamma(rangeF_*)$  and  $V \in \Gamma(rangeF_*)^{\perp}$ ,  $t \in I$  the equation

$$g_{2\beta(t)}(F_*X(t), F_*X(t))g_2(\dot{\beta}(t), (\nabla^N g)) + g_2(\mathcal{S}_V F_*X(t), F_*X(t)) = 0,$$
(3.5)

is satisfied. To prove this, let  $\beta$  be a geodesic on N with  $\dot{\beta}(t) = F_*X(t) + V(t)$  and let  $\omega(t) \in [0, \pi]$  denote the angle between  $\dot{\beta}(t)$  and V(t). If  $\dot{\beta}(t) \in \Gamma(rangeF_*)^{\perp}$ , then we have  $F_*X(t_0) = 0$  (i.e. (3.5) is satisfied), which implies  $\sin \omega(t) = 0$  at point  $\beta(t_0)$ . Thus for any function  $\tilde{s} = e^g$  on M,  $(\tilde{s}(\beta(t)))\sin \omega(t)$  identically vanishes. Therefore, the statement holds trivially in this case. Now, we consider the case  $\sin \omega(t) \neq 0$ , i.e.  $\dot{\beta}(t)$  does not belongs only in  $\Gamma(rangeF_*)^{\perp}$ . Since  $\beta$  is geodesic, its speed is constant  $b = ||\dot{\beta}||^2$  (say). Then

$$g_{2\beta(t)}(V,V) = b\cos^2(\omega(t)), \tag{3.6}$$

$$g_{2\beta(t)}(F_*X, F_*X) = bsin^2\omega(t). \tag{3.7}$$

Now differentiating (3.7) along  $\beta$ , we get

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2bsin\omega cos\omega \frac{d\omega}{dt}.$$
(3.8)

On the other hand,

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2g_2(\nabla^N_{\dot{\beta}}F_*X, F_*X).$$

By putting  $\dot{\beta} = F_*X + V$  in above equation, we get

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2g_2(\nabla^N_{F_*X}F_*X + \nabla^N_V F_*X, F_*X),$$

which implies

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2g_2(\nabla_X^F F_*X \circ F + \nabla_V^N F_*X, F_*X). \tag{3.9}$$

Using (2.2) and (3.2) in (3.9), we get

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2g_2((\nabla F_*)(X, X) + F_*(\nabla_X^M X) + \mathcal{S}_V F_*X - F_*(\nabla_X^M X), F_*X).$$

Using (2.3) in above equation, we get

$$\frac{d}{dt}g_2(F_*X, F_*X) = 2g_2(\mathcal{S}_V F_*X, F_*X). \tag{3.10}$$

Now from (3.8) and (3.10), we get

$$g_2(S_V F_* X, F_* X) = b \sin \omega \cos \omega \frac{d\omega}{dt}.$$
 (3.11)

Moreover, F is a Clairaut Riemannian map with  $\tilde{s} = e^g$  if and only if  $\frac{d}{dt}(e^{g\circ\beta}sin\omega) = 0$ , that is,  $e^{g\circ\beta}sin\omega\frac{d(g\circ\beta)}{dt} + e^{g\circ\beta}cos\omega\frac{d\omega}{dt} = 0$ . By multiplying this with nonzero factor  $bsin\omega$  and using (3.7), we get

$$g_2(F_*X, F_*X)\frac{d(g \circ \beta)}{dt} = -bsin\omega cos\omega \frac{d\omega}{dt}.$$
 (3.12)

Now from (3.11) and (3.12), we get

$$g_2(\mathcal{S}_V F_* X, F_* X) = -g_2(F_* X, F_* X) \frac{d(g \circ \beta)}{dt},$$

which means

$$g_2(S_V F_* X, F_* X) = -g_2(F_* X, F_* X) g_2(\nabla^N g, \dot{\beta}). \tag{3.13}$$

Indeed assuming (3.5) and considering any geodesic  $\beta$  on N with initial tangent vector which belongs in  $\Gamma(rangeF_*)$ , then by using  $V(t_0) = 0$  in (3.13), we get g is constant on  $rangeF_*$  and since  $rangeF_*$  is connected,  $\nabla^N g \in \Gamma(rangeF_*)^{\perp}$ . Then by (3.13), we get

$$g_2(S_V F_* X, F_* X) = -g_2(F_* X, F_* X) g_2(\nabla^N g, V).$$
(3.14)

Thus  $S_V F_* X = -V(g) F_* X$ , where V(g) is a smooth function on N, which implies the proof of (i). Now, by using (2.5) in (3.14), we get

$$g_2(V, (\nabla F_*)(X, X)) = -g_2(F_*X, F_*X)g_2(\nabla^N g, V), \tag{3.15}$$

for  $F_*X \in \Gamma(rangeF_*)$  and  $V \in \Gamma(rangeF_*)^{\perp}$ . Now using (2.2) in (3.15), we get

$$g_2(V, \nabla_X^F F_* X) = -g_2(\nabla^N g, V)g_2(F_* X, F_* X).$$

Thus by comparing, we get

$$\nabla_X^F F_* X = -(\nabla^N g) g_2(F_* X, F_* X). \tag{3.16}$$

Taking trace of (3.16), we get

$$\sum_{j=r+1}^{m} \nabla_{X_j}^F F_* X_j = -(\nabla^N g)(m-r), \tag{3.17}$$

where  $\{X_{r+1}, X_{r+2}, ..., X_m\}$  and  $\{F_*X_{r+1}, F_*X_{r+2}, ..., F_*X_m\}$  are orthonormal bases of  $(kerF_*)^{\perp}$  and  $rangeF_*$ , respectively.

Moreover, the mean curvature vector field of  $rangeF_*$  is defined by ([21], [24] page 199)

$$H_2 = \frac{1}{m-r} \sum_{j=r+1}^{m} \nabla_{X_j}^F F_* X_j, \tag{3.18}$$

where  $\{X_j\}_{r+1\leq j\leq m}$  is an orthonormal basis of  $(ker F_*)^{\perp}$ . Then from (3.17) and (3.18), we get

$$H_2 = -\nabla^N g. (3.19)$$

Also, by (3.15), we get

$$(\nabla F_*)(X, X) = -g_2(F_*X, F_*X)(\nabla^N g). \tag{3.20}$$

Since F is Riemannian map, using (2.1) in (3.20), we get

$$(\nabla F_*)(X, X) = -g_1(X, X)(\nabla^N g). \tag{3.21}$$

From (3.19) and (3.21), we get

$$(\nabla F_*)(X, X) = g_1(X, X)H_2.$$

Thus by Lemma 2.2 F is umbilical map, which completes the proof.

**Remark 3.4** In [25], Sahin considered geodesic curve on the total manifold of a Riemannian map F, then by using Clairaut relation fibers of F are totally umbilical. On the other hand, in Definition 3.1, we considered geodesic curve on the base manifold of F, then by using Clairaut's relation F becomes totally umbilical.

**Theorem 3.5** Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a Clairaut Riemannian map with  $\tilde{s}=e^g$  between Riemannian manifolds such that  $kerF_*$  is minimal. Then F is harmonic if and only if g is constant function on N.

**Proof** Since H = 0, then by Lemma 2.1 F is harmonic if and only if  $H_2 = 0$  if and only if  $\nabla^N g = 0$ , which completes the proof.

**Theorem 3.6** Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a Clairaut Riemannian map with  $\tilde{s}=e^g$  between Riemannian manifolds. Then  $N=N_{(rangeF_*)^{\perp}}\times_f N_{rangeF_*}$  is a twisted product manifold.

**Proof** By (3.20), (3.21) and Theorem 3.3, we have  $\nabla_X^F F_* Y = g_1(X,Y) H_2$  for  $X,Y \in \Gamma(ker F_*)^{\perp}$ , which implies  $range F_*$  is totally umbilical. Then proof follows by [19].

**Example 3.7** Let  $M=\{(x_1,x_2)\in\mathbb{R}^2: x_1\neq 0, x_2\neq 0\}$  be a Riemannian manifold with Riemannian metric  $g_1=e^{2x_2}dx_1^2+dx_2^2$  on M. Let  $N=\{(y_1,y_2)\in\mathbb{R}^2\}$  be a Riemannian manifold with Riemannian metric  $g_2=e^{2x_2}dy_1^2+dy_2^2$  on N. Consider a map  $F:(M,g_1)\to (N,g_2)$  defined by

$$F(x_1, x_2) = (x_1, 0).$$

Then, we get

$$ker F_* = span\{U = e_2\} \text{ and } (ker F_*)^{\perp} = span\{X = e_1\},$$

where  $\left\{e_1=e^{-x_2}\frac{\partial}{\partial x_1},e_2=\frac{\partial}{\partial x_2}\right\}$  and  $\left\{e_1'=e^{-x_2}\frac{\partial}{\partial y_1},e_2'=\frac{\partial}{\partial y_2}\right\}$  are bases on  $T_pM$  and  $T_{F(p)}N$ , respectively, for all  $p\in M$ . By easy computations, we see that  $F_*(X)=e_1'$  and  $g_1(X,X)=g_2(F_*X,F_*X)$  for  $X\in \Gamma(\ker F_*)^\perp$ . Thus F is Riemannian map with range  $F_*=\operatorname{span}\{F_*(X)=e_1'\}$  and  $(\operatorname{range}F_*)^\perp=\operatorname{span}\{e_2'\}$ . Now to show F is Clairaut Riemannian map we will verify Theorem 3.3, for this we will verify (3.14). Since V and  $(\nabla F_*)(X,X)\in \Gamma(\operatorname{range}F_*)^\perp$ , e here we can write  $V=\operatorname{ae}_2'$  and  $(\nabla F_*)(X,X)=\operatorname{be}_2'$  for some  $a,b\in\mathbb{R}$ . Then we get

$$g_2(V, (\nabla F_*)(X, X)) = g_2(ae_2', be_2') = ab,$$
 (3.22)

and

$$g_2(F_*X, F_*X) = g_2(e'_1, e'_1) = 1.$$
 (3.23)

Since  $\nabla^N g = \sum_{i,j=1}^2 g_2^{ij} \frac{\partial g}{\partial y_i} \frac{\partial}{\partial y_j}$ . Therefore for the function  $g = -by_2$ 

$$g_2(\nabla^N g, V) = -ab. (3.24)$$

Thus by using (2.5), (3.22), (3.23) and (3.24) we see that (3.14) holds. Thus F is a Clairaut Riemannian map.

# 4. Clairaut Riemannian map from Riemannian manifold to Ricci soliton

In this section, we study Clairaut Riemannian map  $F:(M,g_1)\to (N,g_2)$  from a Riemannian manifold to a Ricci soliton and give some characterizations.

**Lemma 4.1** [32] Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a Riemannian map between Riemannian manifolds. Then the Ricci tensor on  $(N,g_2)$  given by

$$Ric(F_*X, F_*Y) = Ric^{rangeF_*}(F_*X, F_*Y) - \sum_{k=1}^{n_1} \left\{ g_2(\mathcal{S}_{\nabla_{e_k}^{F_\perp} e_k} F_*X, F_*Y) - g_2(\nabla_{e_k}^N \mathcal{S}_{e_k} F_*X, F_*Y) + g_2(\mathcal{S}_{e_k} F_*X, \mathcal{S}_{e_k} F_*Y) + g_2(\nabla_{e_k}^N F_*X, \mathcal{S}_{e_k} F_*Y) \right\},$$

$$(4.1)$$

$$Ric(V,W) = Ric^{(rangeF_*)^{\perp}}(V,W) - \sum_{j=r+1}^{m} \left\{ g_2(\mathcal{S}_{\nabla_{V}^{F_{\perp}}W}F_*X_j, F_*X_j) + g_2(\mathcal{S}_{V}F_*X_j, \mathcal{S}_{W}F_*X_j) - \nabla_{V}^{N}(g_2(\mathcal{S}_{W}F_*X_j, F_*X_j)) + 2g_2(\mathcal{S}_{W}F_*X_j, \nabla_{V}^{N}F_*X_j) \right\},$$

$$(4.2)$$

$$Ric(F_*X, V) = \sum_{j=r+1}^{m} \left\{ g_2((\tilde{\nabla}_X \mathcal{S})_V F_* X_j, F_* X_j) - g_2((\tilde{\nabla}_{X_j} \mathcal{S})_V F_* X, F_* X_j) \right\} - \sum_{k=1}^{n_1} g_2(R^{F\perp}(F_* X, e_k) V, e_k), \tag{4.3}$$

for  $X,Y \in \Gamma(kerF_*)^{\perp}$ ,  $V,W \in \Gamma(rangeF_*)^{\perp}$  and  $F_*X,F_*Y \in \Gamma(rangeF_*)$ , where  $\{F_*X_j\}_{r+1 \leq j \leq m}$  and  $\{e_k\}_{1 \leq k \leq n_1}$  are orthonormal bases of range $F_*$  and  $(rangeF_*)^{\perp}$ , respectively.

**Theorem 4.2** Let  $F:(M^m,g_1) \to (N^n,g_2)$  be a Clairaut Riemannian map with  $\tilde{s}=e^g$  between Riemannian manifolds. Then the Ricci tensor on  $(N,g_2)$  given by

$$Ric(F_*X, F_*Y) = Ric^{rangeF_*}(F_*X, F_*Y) - \sum_{k=1}^{n_1} (e_k(g))^2 g_2(F_*X, F_*Y) + \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F_\perp} e_k, \nabla^N g) g_2(F_*X, F_*Y) - \sum_{k=1}^{n_1} (\nabla_{e_k}^N e_k(g)) g_2(F_*X, F_*Y),$$

$$(4.4)$$

$$Ric(V,W) = Ric^{(rangeF_*)^{\perp}}(V,W) + (m-r)g_2(\nabla^N g, \nabla_V^{F_{\perp}}W) - (m-r)V(g)W(g) - (m-r)\nabla_V^N W(g),$$
(4.5)

$$Ric(F_*X, V) = \sum_{j=r+1}^{m} g_2((\tilde{\nabla}_X \mathcal{S})_V F_*X_j, F_*X_j) - \sum_{j=r+1}^{m} g_2((\tilde{\nabla}_{X_j} \mathcal{S})_V F_*X_j, F_*X_j) - \sum_{k=1}^{n_1} g_2(R^{F\perp}(F_*X, e_k)V, e_k), (4.6)$$

for  $X,Y \in \Gamma(kerF_*)^{\perp}$ ,  $V,W \in \Gamma(rangeF_*)^{\perp}$  and  $F_*X,F_*Y \in \Gamma(rangeF_*)$ , where  $\{F_*X_j\}_{r+1 \leq j \leq m}$  and  $\{e_k\}_{1 \leq k \leq n_1}$  are orthonormal bases of range $F_*$  and  $(rangeF_*)^{\perp}$ , respectively.

**Proof** Using Theorem 3.3 and (3.14) in (4.1), we get

$$Ric(F_*X, F_*Y) = Ric^{rangeF_*}(F_*X, F_*Y) - \sum_{k=1}^{n_1} (e_k(g))^2 g_2(F_*X, F_*Y) + \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F_\perp} e_k, \nabla^N g) g_2(F_*X, F_*Y) - \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^N (e_k(g)F_*X), F_*Y) + \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^N F_*X, e_k(g)F_*Y),$$

which implies (4.4). Also using Theorem 3.3 and (3.14) in (4.2), we get

$$\begin{aligned} Ric(V,W) & = Ric^{(rangeF_*)^{\perp}}(V,W) + \sum_{j=r+1}^{m} g_2(\nabla_V^{F_{\perp}}W,\nabla^N g)g_2(F_*X_j,F_*X_j) - \sum_{j=r+1}^{m} g_2(V(g)F_*X_j,W(g)F_*X_j) \\ & - \sum_{j=r+1}^{m} \nabla_V^N (g_2(W(g)F_*X_j,F_*X_j)) + 2\sum_{j=r+1}^{m} g_2(W(g)F_*X_j,\nabla_V^N F_*X_j), \end{aligned}$$

which implies (4.5). Also the proof of (4.3) and (4.6) is same.

**Theorem 4.3** Let  $(N, g_2, H_2, \lambda)$  be a Ricci soliton with the potential vector field  $H_2 \in \Gamma(rangeF_*)^{\perp}$  and  $F: (M, g_1) \to (N, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then

$$s^{rangeF_*} = -\lambda (m-r) + (m-r)\Delta g - (m-r)(m-r-2) \|\nabla^N g\|^2,$$

where  $s^{rangeF_*}$  is the scalar curvature of  $rangeF_*$  and  $(m-r) = \dim(rangeF_*)$ .

**Proof** Since  $(N, g_2, H_2, \lambda)$  admit Ricci soliton with the potential vector field  $H_2 \in \Gamma(rangeF_*)^{\perp}$  then, we have

$$\frac{1}{2}(L_{H_2}g_2)(F_*X, F_*Y) + Ric(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0,$$

for  $F_*X, F_*Y \in \Gamma(rangeF_*)$ , which implies

$$\frac{1}{2}\{g_2(\nabla^N_{F_*X}H_2,F_*Y)+g_2(\nabla^N_{F_*Y}H_2,F_*X)\}+Ric(F_*X,F_*Y)+\lambda g_2(F_*X,F_*Y)=0.$$

Using (2.4) in above equation, we get

$$\frac{1}{2}\{g_2(-\mathcal{S}_{H_2}F_*X,F_*Y)+g_2(-\mathcal{S}_{H_2}F_*Y,F_*X)\}+Ric(F_*X,F_*Y)+\lambda g_2(F_*X,F_*Y)=0.$$

Since  $\mathcal{S}_{H_2}$  is self-adjoint, above equation can be written as

$$-g_2(S_{H_2}F_*X, F_*Y) + Ric(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0.$$
(4.7)

Using (3.14), (3.19) and (4.4) in (4.7), we get

$$-g_{2}(\nabla^{N}g, \nabla^{N}g)g_{2}(F_{*}X, F_{*}Y) + Ric^{rangeF_{*}}(F_{*}X, F_{*}Y) - \sum_{k=1}^{n_{1}} (e_{k}(g))^{2}g_{2}(F_{*}X, F_{*}Y) + \sum_{k=1}^{n_{1}} g_{2}(\nabla^{F_{\perp}}e_{k}, \nabla^{N}g)g_{2}(F_{*}X, F_{*}Y) - \sum_{k=1}^{n_{1}} \nabla^{N}e_{k}(g)g_{2}(F_{*}X, F_{*}Y) + \lambda g_{2}(F_{*}X, F_{*}Y) = 0,$$

where  $\{e_k\}_{1\leq k\leq n_1}$  is an orthonormal basis of  $(rangeF_*)^{\perp}$ . This implies

$$-2\|\nabla^{N}g\|^{2}g_{2}(F_{*}X, F_{*}Y) + Ric^{rangeF_{*}}(F_{*}X, F_{*}Y) - \sum_{k=1}^{n_{1}} g_{2}(e_{k}, \nabla^{N}_{e_{k}}\nabla^{N}g)g_{2}(F_{*}X, F_{*}Y) + \lambda g_{2}(F_{*}X, F_{*}Y) = 0.$$

$$(4.8)$$

Taking trace of (4.8) for  $rangeF_*$ , we get

$$s^{rangeF_*} - 2(m-r)\|\nabla^N g\|^2 - (m-r)\sum_{k=1}^{n_1} g_2(\nabla^N_{e_k}\nabla^N g, e_k) + \lambda(m-r) = 0.$$

Using definition of Hessian form of g (i.e.  $H^g(X_1, Y_1) = g_2(\nabla_{X_1}^N \nabla^N g, Y_1)$  for all  $X_1, Y_1 \in \Gamma(TN)$ ) from [8] in above equation, we get

$$s^{rangeF_*} + (m-r)\{-2\|\nabla^N g\|^2 - \sum_{k=1}^{n_1} H^g(e_k, e_k) + \lambda\} = 0.$$
 (4.9)

Since we know that

$$\Delta g = \sum_{j=r+1}^{m} H^g(F_* X_j, F_* X_j) + \sum_{k=1}^{n_1} H^g(e_k, e_k), \tag{4.10}$$

where  $\{F_*X_j\}_{r+1\leq j\leq m}$  and  $\{e_k\}_{1\leq k\leq n_1}$  are orthonormal bases of  $rangeF_*$  and  $(rangeF_*)^{\perp}$ , respectively. Then by using definition of Hessian form of g in (4.10), we get

$$\Delta g = \sum_{j=r+1}^{m} g_2(\nabla_{F_*X_j}^N \nabla^N g, F_*X_j) + \sum_{k=1}^{n_1} H^g(e_k, e_k). \tag{4.11}$$

Using (2.4) in (4.11), we get

$$\Delta g = -\sum_{j=r+1}^{m} g_2(\mathcal{S}_{\nabla^N g} F_* X_j, F_* X_j) + \sum_{k=1}^{n_1} H^g(e_k, e_k).$$

Using Theorem 3.3 in above equation, we get

$$\Delta g - (m - r) \|\nabla^N g\|^2 = \sum_{k=1}^{n_1} H^g(e_k, e_k). \tag{4.12}$$

Thus (4.9) and (4.12) implies the proof.

**Theorem 4.4** Let  $(N, g_2, H_2, \lambda)$  be a Ricci soliton with the potential vector field  $H_2 \in \Gamma(rangeF_*)^{\perp}$  and  $F: (M^m, g_1) \to (N^n, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then

$$s^{(rangeF_*)^{\perp}} = -\lambda n_1 + (m-r+1)\Delta g - (m-r)^2 \|\nabla^N g\|^2$$

where  $s^{(rangeF_*)^{\perp}}$  denotes the scalar curvature of  $(rangeF_*)^{\perp}$  and  $(m-r) = \dim(rangeF_*)$ ,  $n_1 = \dim((rangeF_*)^{\perp})$ .

**Proof** Since  $(N, g_2, H_2, \lambda)$  admit Ricci soliton with the potential vector field  $H_2 \in \Gamma(rangeF_*)^{\perp}$  then, we have

$$\frac{1}{2}(L_{H_2}g_2)(V,W) + Ric(V,W) + \lambda g_2(V,W) = 0,$$

for  $V, W \in \Gamma(rangeF_*)^{\perp}$ , which implies

$$\frac{1}{2}\{g_2(\nabla_V^N H_2, W) + g_2(\nabla_W^N H_2, V)\} + Ric(V, W) + \lambda g_2(V, W) = 0.$$

Putting  $H_2 = -\nabla^N g$  in above equation, we get

$$-\frac{1}{2}\{g_2(\nabla_V^N \nabla^N g, W) + g_2(\nabla_W^N \nabla^N g, V)\} + Ric(V, W) + \lambda g_2(V, W) = 0.$$
(4.13)

Using definition of Hessian form of g and (4.5) in (4.13), we get

$$-H^{g}(V,W) + Ric^{(rangeF_{*})^{\perp}}(V,W) + (m-r)g_{2}(\nabla^{N}g,\nabla_{V}^{F\perp}W) - (m-r)V(g)W(g) - (m-r)\nabla_{V}^{N}W(g) + \lambda g_{2}(V,W) = 0.$$
(4.14)

Taking trace of (4.14) for  $(rangeF_*)^{\perp}$ , we get

$$-\sum_{k=1}^{n_1} H^g(e_k, e_k) + s^{(rangeF_*)^{\perp}} + \sum_{k=1}^{n_1} (m-r)g_2(\nabla^N g, \nabla^{F_{\perp}}_{e_k} e_k) - (m-r)\sum_{k=1}^{n_1} (e_k(g))^2 - (m-r)\sum_{k=1}^{n_1} \nabla^N_{e_k} e_k(g) + \lambda n_1 = 0,$$

where  $\{e_k\}_{1\leq k\leq n_1}$  is an orthonormal basis of  $(rangeF_*)^{\perp}$ , which implies

$$s^{(rangeF_*)^{\perp}} + \lambda n_1 - (m-r) \sum_{k=1}^{n_1} (e_k(g))^2 - (m-r+1) \sum_{k=1}^{n_1} H^g(e_k, e_k) = 0.$$

Using (4.12) and  $(e_k(g))^2 = g_2(\nabla^N g, e_k)^2 = g_2(\nabla^N g, \nabla^N g)$  in above equation, we get the proof.

**Remark 4.5** Since  $rangeF_*$  and  $(rangeF_*)^{\perp}$  are subbundles of TN, they define distributions on N. Then for  $F_*X, F_*Y \in \Gamma(rangeF_*)$ , we have

$$[F_*X, F_*Y] = \nabla^N_{F_*X} F_*Y - \nabla^N_{F_*Y} F_*X$$
$$= \nabla^N_{F_*Y} F_*Y \circ F - \nabla^P_{Y} F_*X \circ F.$$

Using (2.2) in above equation, we get

$$[F_*X, F_*Y] = F_*(\nabla_X Y) - F_*(\nabla_Y X) = F_*(\nabla_X Y - \nabla_Y X) \in \Gamma(rangeF_*).$$

Thus range  $F_*$  is an integrable distribution. Then for any point  $F(p) \in N$  there exists maximal integral manifold or a leaf of range  $F_*$  containing F(p).

**Theorem 4.6** Let  $(N, g_2, F_*Z, \lambda)$  be a Ricci soliton with the potential vector field  $F_*Z \in \Gamma(rangeF_*)$  and  $F: (M, g_1) \to (N, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then a leaf of  $rangeF_*$  is an almost Ricci soliton.

**Proof** Since  $(N, g_2, F_*Z, \lambda)$  admit Ricci soliton with the potential vector field  $F_*Z \in \Gamma(rangeF_*)$  then, we have

$$\frac{1}{2}(L_{F_*Z}g_2)(F_*X, F_*Y) + Ric(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0, \tag{4.15}$$

for  $F_*X, F_*Y, F_*Z \in \Gamma(rangeF_*)$ . Using (4.4) in (4.15), we get

$$\begin{split} &\frac{1}{2}(L_{F_*Z}g_2)(F_*X,F_*Y) + Ric^{rangeF_*}(F_*X,F_*Y) - \sum_{k=1}^{n_1}(e_k(g))^2g_2(F_*X,F_*Y) \\ &+ \sum_{k=1}^{n_1}g_2(\nabla_{e_k}^{F_\perp}e_k,\nabla^Ng)g_2(F_*X,F_*Y) - \sum_{k=1}^{n_1}\nabla_{e_k}^Ne_k(g)g_2(F_*X,F_*Y) + \lambda g_2(F_*X,F_*Y) = 0, \end{split}$$

where  $\{e_k\}_{1\leq k\leq n_1}$  is an orthonormal basis of  $(rangeF_*)^{\perp}$ , which implies

$$\frac{1}{2}(L_{F_*Z}q_2)(F_*X, F_*Y) + Ric^{rangeF_*}(F_*X, F_*Y) + \tilde{\lambda}q_2(F_*X, F_*Y) = 0,$$

where  $\tilde{\lambda} = -\sum_{k=1}^{n_1} (e_k(g))^2 + \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F\perp} e_k, \nabla^N g) - \sum_{k=1}^{n_1} e_k(e_k(g)) + \lambda$  is a smooth function on N. Thus a leaf of  $rangeF_*$  is an almost Ricci soliton, which completes the proof.

**Theorem 4.7** Let  $(N, g_2, V, \lambda)$  be a Ricci soliton with the potential vector field  $V \in \Gamma(rangeF_*)^{\perp}$  and  $F : (M, g_1) \to (N, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then a leaf of  $rangeF_*$  is an Einstein.

**Proof** Since  $(N, g_2, F_*Z, \lambda)$  admit Ricci soliton with the potential vector field  $F_*Z \in \Gamma(rangeF_*)$  then, we have

$$\frac{1}{2}(L_V g_2)(F_* X, F_* Y) + Ric(F_* X, F_* Y) + \lambda g_2(F_* X, F_* Y) = 0,$$

for  $F_*X, F_*Y \in \Gamma(rangeF_*)$ , which implies

$$\frac{1}{2}\{g_2(\nabla^N_{F_*X}V, F_*Y) + g_2(\nabla^N_{F_*Y}V, F_*X)\} + Ric(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0.$$

Using (2.4) in above equation, we get

$$\frac{1}{2}\{g_2(-\mathcal{S}_V F_* X, F_* Y) + g_2(-\mathcal{S}_V F_* Y, F_* X)\} + Ric(F_* X, F_* Y) + \lambda g_2(F_* X, F_* Y) = 0.$$

Since  $S_V$  is self-adjoint, above equation can be written as

$$-g_2(S_V F_* X, F_* Y) + Ric(F_* X, F_* Y) + \lambda g_2(F_* X, F_* Y) = 0.$$
(4.16)

Since F is Clairaut Riemannian map, using  $S_V F_* X = -V(g) F_* X$  and (4.4) in (4.16), we get

$$V(g)g_{2}(F_{*}X, F_{*}Y) + Ric^{rangeF_{*}}(F_{*}X, F_{*}Y) - \sum_{k=1}^{n_{1}} (e_{k}(g))^{2}g_{2}(F_{*}X, F_{*}Y) + \sum_{k=1}^{n_{1}} g_{2}(\nabla_{e_{k}}^{F\perp}e_{k}, \nabla^{N}g)g_{2}(F_{*}X, F_{*}Y) - \sum_{k=1}^{n_{1}} \nabla_{e_{k}}^{N}e_{k}(g)g_{2}(F_{*}X, F_{*}Y) + \lambda g_{2}(F_{*}X, F_{*}Y) = 0,$$

where  $\{e_k\}_{1 \leq k \leq n_1}$  is an orthonormal basis of  $(rangeF_*)^{\perp}$ , which implies

$$Ric^{rangeF_*}(F_*X, F_*Y) = \lambda' g_2(F_*X, F_*Y),$$

where 
$$\lambda' = \sum_{k=1}^{n_1} (e_k(g))^2 - \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F\perp} e_k, \nabla^N g) + \sum_{k=1}^{n_1} e_k(e_k(g)) - \lambda - V(g)$$
 is a smooth function on  $N$ . Thus a leaf of  $rangeF_*$  is an Einstein, which completes the proof.

**Theorem 4.8** Let  $\beta$  be a geodesic curve on N and  $(N, g_2, \dot{\beta}, \lambda)$  be a Ricci soliton with the potential vector field  $\dot{\beta} \in \Gamma(TN)$ . Let  $F: (M, g_1) \to (N, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  from a Riemannian manifold M to an Einstein manifold N. Then the following statements are true:

- (i)  $\dot{\beta}$  is a conformal vector field on range  $F_*$ .
- (ii)  $\dot{\beta}$  is Killing vector field on  $(rangeF_*)^{\perp}$  if and only if  $V(g)W(g) = -H^g(V,W)$  for all  $V,W \in \Gamma(rangeF_*)^{\perp}$ .

**Proof** Since  $(N, g_2, \dot{\beta}, \lambda)$  is a Ricci soliton then, we have

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(F_*X, F_*Y) + Ric(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0, \tag{4.17}$$

for  $F_*X, F_*Y \in \Gamma(rangeF_*)$ . Using (4.4) in (4.17), we get

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(F_*X, F_*Y) + Ric^{rangeF_*}(F_*X, F_*Y) - \sum_{k=1}^{n_1} (e_k(g))^2 g_2(F_*X, F_*Y) 
+ \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F_\perp} e_k, \nabla^N g) g_2(F_*X, F_*Y) - \sum_{k=1}^{n_1} \nabla_{e_k}^N e_k(g) g_2(F_*X, F_*Y) + \lambda g_2(F_*X, F_*Y) = 0,$$
(4.18)

where  $\{e_k\}_{1\leq k\leq n_1}$  is an orthonormal basis of  $(rangeF_*)^{\perp}$ . Since N is Einstein, putting  $Ric^{rangeF_*}(F_*X, F_*Y) = -\lambda g_2(F_*X, F_*Y)$  in (4.18), we get

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(F_*X, F_*Y) + \mu g_2(F_*X, F_*Y) = 0,$$

where  $\mu = -\sum_{k=1}^{n_1} (e_k(g))^2 + \sum_{k=1}^{n_1} g_2(\nabla_{e_k}^{F\perp} e_k, \nabla^N g) - \sum_{k=1}^{n_1} e_k(e_k(g))$  is a smooth function on N. Thus  $\dot{\beta}$  is a conformal vector field on  $rangeF_*$ . On the other hand, since  $(N, g_2, \dot{\beta}, \lambda)$  is a Ricci soliton then, we have

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(V,W) + Ric(V,W) + \lambda g_2(V,W) = 0, \tag{4.19}$$

for any  $V, W \in \Gamma(rangeF_*)^{\perp}$ . Using (4.5) in (4.19), we get

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(V,W) + Ric^{(rangeF_*)^{\perp}}(V,W) + (m-r)g_2(\nabla^N g, \nabla_V^{F^{\perp}}W) 
-(m-r)V(g)W(g) - (m-r)\nabla_V^N W(g) + \lambda g_2(V,W) = 0.$$
(4.20)

Since N is Einstein, putting  $Ric^{(rangeF_*)^{\perp}}(V,W) = -\lambda g_2(V,W)$  in (4.20), we get

$$\frac{1}{2}(L_{\dot{\beta}}g_2)(V,W) + \{g_2(\nabla^N g, \nabla_V^{F\perp} W) - V(g)W(g) - \nabla_V^N W(g)\}(m-r) = 0.$$

Then by using  $\nabla_V^N W(g) = \nabla_V^N (g_2(W, \nabla^N g)) = g_2(\nabla_V^N W, \nabla^N g) + H^g(V, W) = g_2(\nabla_V^{F\perp} W, \nabla^N g) + H^g(V, W)$  in above equation, we get  $\frac{1}{2}(L_{\dot{\beta}}g_2)(V, W) = 0$  if and only if  $V(g)W(g) = -H^g(V, W)$ . This completes the proof.

**Lemma 4.9** Let  $(N, g_2, X_1, \lambda)$  be a Ricci soliton with the potential vector field  $X_1 \in \Gamma(TN)$  and  $F: (M^m, g_1) \to (N^n, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then

$$s = -\lambda n, (4.21)$$

where s denotes the scalar curvature of N.

**Proof** The proof is similar to remark 9 of [30]; therefore, we are omitting it.  $\Box$ 

**Theorem 4.10** Let  $(N, g_2, -H_2, \lambda)$  be a Ricci soliton with the potential vector field  $-H_2 \in \Gamma(rangeF_*)^{\perp}$  and  $F: (M, g_1) \to (N, g_2)$  be a Clairaut Riemannian map with  $\tilde{s} = e^g$  between Riemannian manifolds. Then following statements are true:

- (i) N admits a gradient Ricci soliton.
- (ii) The mean curvature vector field of range  $F_*$  is constant.

**Proof** By similar proof as theorem 10 of [30], we get

$$\Delta q = 0.$$

Hence  $\nabla^N(\nabla^N g) = 0$ , i.e.  $\nabla^N H_2 = 0$ , which means  $H_2$  is constant. This completes the proof.

**Example 4.11** The map  $F: M \to N$  given in Example 3.7 is Clairaut Riemannian map. Now, we will show that N admits a Ricci soliton, i.e.

$$\frac{1}{2}(L_{Z_1}g_2)(X_1, Y_1) + Ric(X_1, Y_1) + \lambda g_2(X_1, Y_1) = 0, \tag{4.22}$$

for any  $X_1, Y_1, Z_1 \in \Gamma(TN)$ . By similar computations as example 6.1 of [32], we get

$$\frac{1}{2}(L_{Z_1}g_2)(X_1, Y_1) = 0, (4.23)$$

$$g_2(X_1, Y_1) = (a_1 a_3 + a_2 a_4), (4.24)$$

and

$$Ric(X_1, Y_1) = a_1 a_3 Ric(e'_1, e'_1) + (a_1 a_4 + a_2 a_3) Ric(e'_1, e'_2) + a_2 a_4 Ric(e'_2, e'_2).$$

$$(4.25)$$

By (4.4), we get

$$Ric(e'_1,e'_1) = Ric^{rangeF_*}(e'_1,e'_1) - (g_2(\nabla^N g,e'_2))^2 + g_2(\nabla^{F\perp}_{e'_2}e'_2,\nabla^N g) - \nabla^N_{e'_2}(g_2(e'_2,\nabla^N g)).$$

Since dimension of range  $F_*$  is one,  $Ric^{range}F_*(e_1',e_1')=0$  and we have  $\nabla^N g=-be_2'$  for some  $b\in\mathbb{R}$ . So

$$Ric(e'_1, e'_1) = -b^2,$$
 (4.26)

By (4.5), we get

$$Ric(e_2', e_2') = Ric^{(rangeF_*)^{\perp}}(e_2', e_2') + g_2(\nabla^N g, \nabla^{F_\perp}_{e_2'} e_2') - e_2'(g)e_2'(g) - \nabla^N_{e_2'}(e_2'(g)).$$

Since dimension of  $(rangeF_*)^{\perp}$  is one,  $Ric^{(rangeF_*)^{\perp}}(e_2',e_2')=0$  and putting  $\nabla^N g=-be_2'$  for some  $b\in\mathbb{R}$ , we get

$$Ric(e_2', e_2') = -b^2.$$
 (4.27)

And by similar computation as example 6.1 of [32], we get

$$Ric(e'_1, e'_2) = 0.$$
 (4.28)

Using (4.26), (4.27) and (4.28) in (4.25), we get

$$Ric(X_1, Y_1) = -(a_1 a_3 + a_2 a_4)b^2. (4.29)$$

Now, using (4.23), (4.24) and (4.29) in (4.22), we obtain that metric  $g_2$  admits Ricci soliton for

$$\lambda = b^2$$
.

Since  $b \in \mathbb{R}$ , for some choices of b Ricci soliton  $(N, g_2)$  will be expanding or steady according to  $\lambda > 0$  or  $\lambda = 0$ .

# 5. Clairaut anti-invariant Riemannian map from Riemannian manifold to Kähler manifold

In this section, we introduce Clairaut anti-invariant Riemannian map from a Riemannian manifold to a Kähler manifold and investigate the geometry with a non-trivial example.

Let  $(N, g_2)$  be an almost Hermitian manifold [33], then N admits a tensor J of type (1, 1) on N such that  $J^2 = -I$  and

$$g_2(JX_1, JY_1) = g_2(X_1, Y_1), (5.1)$$

for all  $X_1, Y_1 \in \Gamma(TN)$ . An almost Hermitian manifold N is called Kähler manifold if

$$(\nabla_{X_1}^N J)Y_1 = 0,$$

for all  $X_1, Y_1 \in \Gamma(TN)$ , where  $\nabla^N$  is the Levi-Civita connection on N.

**Definition 5.1** [20] Let  $F:(M,g_1) \to (N,g_2)$  be a proper Riemannian map from a Riemannian manifold M to an almost Hermitian manifold N with almost complex structure J. We say that F is an anti-invariant Riemannian map at  $p \in M$  if  $J(rangeF_{*p}) \subset (rangeF_{*p})^{\perp}$ . If F is an anti-invariant Riemannian map for every  $p \in M$  then F is called an anti-invariant Riemannian map.

In this case we denote the orthogonal subbundle to  $J(rangeF_*)$  in  $(rangeF_*)^{\perp}$  by  $\mu$ , i.e.  $(rangeF_*)^{\perp} = J(rangeF_*) \oplus \mu$ . For any  $V \in \Gamma(rangeF_*)^{\perp}$ , we have

$$JV = BV + CV, (5.2)$$

where  $BV \in \Gamma(rangeF_*)$  and  $CV \in \mu$ . Note that if  $\mu = 0$  then F is called Lagrangian Riemannian map [27].

**Lemma 5.2** Let  $F:(M,g_1) \to (N,g_2,J)$  be an anti-invariant Riemannian map from a Riemannian manifold M to a Kähler manifold N and  $\alpha:I \to M$  be a geodesic curve on M. Then the curve  $\beta=F\circ\alpha$  is geodesic on N if and only if

$$-S_{JF_*X}F_*X - S_{CV}F_*X + \nabla_V^N BV + F_*(\nabla_X^{M*}F_*BV) = 0,$$
(5.3)

$$(\nabla F_*)(X, {}^*F_*BV) + \nabla_X^{F\perp} J F_* X + \nabla_V^{F\perp} J F_* X + \nabla_X^{F\perp} CV + \nabla_V^{F\perp} CV = 0, \tag{5.4}$$

where  $F_*X \in \Gamma(rangeF_*), V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$  and  $^*F_*$  is the adjoint map of  $F_*$ , and  $\nabla^N$  is the Levi-Civita connection on N, and  $\nabla^{F\perp}$  is a linear connection on  $(rangeF_*)^{\perp}$ .

**Proof** Let  $\alpha: I \to M$  be a geodesic on M and let  $\beta = F \circ \alpha$  be a geodesic on N with  $F_*X \in \Gamma(rangeF_*)$  and  $V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$ . Since N is Kähler manifold,  $\nabla^N_{\dot{\beta}}\dot{\beta} = -J\nabla^N_{\dot{\beta}}J\dot{\beta}$ . Thus

$$\nabla^N_{\dot{\beta}}\dot{\beta} = -J\nabla^N_{\dot{\beta}}J\dot{\beta} = -J\nabla^N_{F_*X+V}J(F_*X+V),$$

which implies

$$\nabla^{N}_{\dot{\beta}}\dot{\beta} = -J(\nabla^{N}_{F_{*}X}JF_{*}X + \nabla^{N}_{F_{*}X}JV + \nabla^{N}_{V}JF_{*}X + \nabla^{N}_{V}JV). \tag{5.5}$$

Using (2.4) and (5.2) in (5.5), we get

$$\nabla^{N}_{\dot{\beta}}\dot{\beta} = -J(-\mathcal{S}_{JF_{*}X}F_{*}X - \mathcal{S}_{CV}F_{*}X + \nabla^{N}_{V}BV + \nabla^{N}_{F_{*}X}BV + \nabla^{F}_{Y}^{\perp}JF_{*}X + \nabla^{F}_{V}^{\perp}JF_{*}X + \nabla^{F}_{X}^{\perp}CV + \nabla^{F}_{V}^{\perp}CV).$$

$$(5.6)$$

Since  $\nabla^N$  is Levi-Civita connection on N and  $g_2(\nabla^N_V BV, U) = 0$  for any  $U \in \Gamma(rangeF_*)^{\perp}$ ,  $\nabla^N_V BV \in \Gamma(rangeF_*)$  and using (2.2), we get  $\nabla^N_{F_*X} BV = \nabla^F_X BV \circ F = (\nabla F_*)(X, {}^*F_*BV) + F_*(\nabla^M_X {}^*F_*BV)$ . Then by (5.6), we get

$$\begin{array}{ll} \nabla^N_{\dot{\beta}} \dot{\beta} &= -J(-\mathcal{S}_{JF_*X}F_*X - \mathcal{S}_{CV}F_*X + \nabla^N_VBV + (\nabla F_*)(X, {}^*F_*BV) \\ &+ F_*(\nabla^M_X {}^*F_*BV) + \nabla^{F\perp}_XJF_*X + \nabla^{F\perp}_VJF_*X + \nabla^{F\perp}_XCV + \nabla^{F\perp}_VCV). \end{array}$$

Now  $\beta$  is geodesic on  $N \iff \nabla^N_{\dot{\beta}}\dot{\beta} = 0 \iff -\mathcal{S}_{JF_*X}F_*X - \mathcal{S}_{CV}F_*X + \nabla^N_VBV + (\nabla F_*)(X, {}^*F_*BV) + F_*(\nabla^M_X {}^*F_*BV) + \nabla^{F\perp}_XJF_*X + \nabla^{F\perp}_VJF_*X + \nabla^{F\perp}_XCV + \nabla^{F\perp}_VCV = 0$ , which completes the proof.

**Definition 5.3** An anti-invariant Riemannian map from a Riemannian manifold to a Kähler manifold is called Clairaut anti-invariant Riemannian map if it satisfies the condition of Clairaut Riemannian map.

**Theorem 5.4** Let  $F:(M,g_1) \to (N,g_2,J)$  be an anti-invariant Riemannian map from a Riemannian manifold M to a Kähler manifold N and  $\alpha$ ,  $\beta = F \circ \alpha$  are geodesic curves on M and N, respectively. Then F is Clairaut anti-invariant Riemannian map with  $\tilde{s} = e^g$  if and only if  $g_2(S_{JF_*X}F_*X + S_{CV}F_*X, BV) - g_2((\nabla F_*)(X, {}^*F_*BV) + \nabla_X^{F\perp}JF_*X + \nabla_V^{F\perp}JF_*X, CV) - g_2(F_*X, F_*X) \frac{d(g \circ \beta)}{dt} = 0$ , where g is a smooth function on N and  $F_*X \in \Gamma(rangeF_*)$ ,  $V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$ .

**Proof** Let  $\alpha: I \to M$  be a geodesic on M and let  $\beta = F \circ \alpha$  be a geodesic on N with  $F_*X \in \Gamma(rangeF_*)$  and  $V \in \Gamma(rangeF_*)^{\perp}$  are components of  $\dot{\beta}(t)$  and  $\omega(t)$  denote the angle in  $[0, \pi]$  between  $\dot{\beta}$  and V. Assuming  $b = \|\dot{\beta}(t)\|^2$ , then we get

$$g_{2\beta(t)}(V,V) = b\cos^2 \omega(t), \tag{5.7}$$

$$g_{2\beta(t)}(F_*X, F_*X) = b\sin^2(\omega(t)). \tag{5.8}$$

Now differentiating (5.7) along  $\beta$ , we get

$$\frac{d}{dt}g_2(V,V) = -2bsin\omega(t)cos\omega(t)\frac{d\omega}{dt}.$$
(5.9)

On the other hand by (5.1), we get

$$\frac{d}{dt}g_2(V,V) = \frac{d}{dt}g_2(JV,JV).$$

Using (5.2) in above equation, we get

$$\frac{d}{dt}g_2(V,V) = \frac{d}{dt}\Big(g_2(BV,BV) + g_2(CV,CV)\Big),$$

which implies

$$\frac{d}{dt}g_2(V,V) = 2g_2(\nabla^N_{\dot{\beta}}BV,BV) + 2g_2(\nabla^N_{\dot{\beta}}CV,CV). \tag{5.10}$$

Putting  $\dot{\beta} = F_*X + V$  in (5.10), we get

$$\tfrac{d}{dt} g_2(V,V) = 2g_2(\nabla^N_{F_*X}BV,BV) + 2g_2(\nabla^N_{F_*X}CV,CV) + 2g_2(\nabla^N_VBV,BV) + 2g_2(\nabla^N_VCV,CV).$$

Since  $(rangeF_*)^{\perp}$  is totally geodesic, above equation can be written as

$$\frac{d}{dt}g_2(V,V) = 2g_2(\nabla_V^R BV \circ F, BV) + 2g_2(\nabla_{F_*X}^N CV, CV) + 2g_2(\nabla_V^N BV, BV) + 2g_2(\nabla_V^{F^{\perp}} CV, CV). \tag{5.11}$$

Using (2.2), (2.3) and (2.4) in (5.11), we get

$$\frac{d}{dt}g_2(V,V) = 2g_2(F_*(\nabla_X^{M*}F_*BV) + \nabla_V^{N}BV,BV) + 2g_2(\nabla_X^{F\perp}CV + \nabla_V^{F\perp}CV,CV). \tag{5.12}$$

Using (5.3) and (5.4) in (5.12), we get

$$\frac{d}{dt}g_2(V,V) = 2g_2(S_{JF_*X}F_*X + S_{CV}F_*X,BV) - 2g_2\Big((\nabla F_*)(X, F_*BV) + \nabla_X^{F_\perp}JF_*X + \nabla_V^{F_\perp}JF_*X,CV\Big).$$
(5.13)

Now from (5.9) and (5.13), we get

$$g_2(\mathcal{S}_{JF_*X}F_*X + \mathcal{S}_{CV}F_*X, BV) - g_2\Big((\nabla F_*)(X, {^*F_*BV}) + \nabla_X^{F\perp}JF_*X + \nabla_V^{F\perp}JF_*X, CV\Big) = -bsin\omega cos\omega \frac{d\omega}{dt}.$$

$$(5.14)$$

Moreover, F is a Clairaut Riemannian map with  $\tilde{s} = e^g$  if and only if  $\frac{d}{dt}(e^{g\circ\beta}sin\omega) = 0$ , that is,  $e^{g\circ\beta}sin\omega\frac{d(g\circ\beta)}{dt} + e^{g\circ\beta}cos\omega\frac{d\omega}{dt} = 0$ . By multiplying this with nonzero factor  $bsin\omega$  and using (5.8), we get

$$g_2(F_*X, F_*X)\frac{d(g \circ \beta)}{dt} = -bsin\omega cos\omega \frac{d\omega}{dt}.$$
 (5.15)

Thus (5.14) and (5.15) complete the proof.

**Theorem 5.5** Let  $F:(M^m,g_1)\to (N^n,g_2,J)$  be a Clairaut anti-invariant Riemannian map with  $\tilde{s}=e^g$  from a Riemannian manifold M to a Kähler manifold N. Then at least one of the following statement is true:

(i)  $\dim(rangeF_*)=1$ ,

(ii) g is constant on  $J(rangeF_*)$ , where g is a smooth function on N.

**Proof** Since F is Clairaut Riemannian map with  $\tilde{s} = e^g$  then using (2.2) in (3.21), we get

$$\nabla_{X}^{F} F_{*} Y - F_{*} (\nabla_{X}^{M} Y) = -g_{1}(X, Y) \nabla^{N} g, \qquad (5.16)$$

for  $F_*Y \in \Gamma(rangeF_*)$  and  $X,Y \in \Gamma(kerF_*)^{\perp}$ . Taking inner product of (5.16) with  $JF_*Z \in \Gamma(rangeF_*)^{\perp}$ , we get

$$g_2(\nabla_X^F F_* Y - F_*(\nabla_X^M Y), JF_* Z) = -g_1(X, Y)g_2(\nabla^N g, JF_* Z).$$
(5.17)

Since  $\nabla^F$  is pullback connection of the Levi-Civita connection  $\nabla^N$ . Therefore  $\nabla^F$  is also Levi-Civita connection. Then using metric compatibility condition in (5.17), we get

$$-g_2(\nabla_X^N J F_* Z, F_* Y) = -g_1(X, Y)g_2(\nabla^N g, J F_* Z),$$

which implies

$$g_2(J\nabla_X^F F_* Z, F_* Y) = g_1(X, Y)g_2(\nabla^N g, JF_* Z).$$
(5.18)

Using (5.1) in (5.18), we get

$$-g_2(\nabla_X^F F_* Z, JF_* Y) = g_1(X, Y)g_2(\nabla^N g, JF_* Z).$$

Using (5.16) in above equation, we get

$$g_1(X, Z)g_2(\nabla^N g, JF_*Y) = g_1(X, Y)g_2(\nabla^N g, JF_*Z).$$
(5.19)

Now putting X = Y in (5.19), we get

$$g_1(X, Z)g_2(\nabla^N g, JF_*X) = g_1(X, X)g_2(\nabla^N g, JF_*Z).$$
 (5.20)

Now interchanging X and Z in (5.20), we get

$$g_1(X, Z)g_2(\nabla^N g, JF_*Z) = g_1(Z, Z)g_2(\nabla^N g, JF_*X).$$
(5.21)

From (5.20) and (5.21), we get

$$g_2(\nabla^N g, JF_*X)\left(1 - \frac{g_1(X, X)g_1(Z, Z)}{g_1(X, Z)g_1(X, Z)}\right) = 0,$$

which implies either dim $((ker F_*)^{\perp}) = 1$  or  $g_2(\nabla^N g, JF_*X) = 0$ , which means  $(JF_*X)(g) = 0$ , which completes the proof.

**Theorem 5.6** Let  $F:(M^m,g_1)\to (N^n,g_2,J)$  be a Clairaut Lagrangian Riemannian map with  $\tilde{s}=e^g$  from a Riemannian manifold M to a Kähler manifold N such that  $\dim(rangeF_*)>1$ . Then following statements are true:

- (i)  $rangeF_*$  is minimal.
- (ii)  $rangeF_*$  is totally geodesic.

**Proof** Since F is Clairaut Riemannian map then from (3.21) and Theorem 3.3, we have

$$(\nabla F_*)(X, X) = g_1(X, X)H_2,$$

for  $X \in \Gamma(kerF_*)^{\perp}$  and  $H_2$  is the mean curvature vector field of  $rangeF_*$ . Now multiply above equation by  $U \in \Gamma(rangeF_*)^{\perp}$ , we get

$$g_2((\nabla F_*)(X,X),U) = g_1(X,X)g_2(H_2,U). \tag{5.22}$$

Using (2.2) in (5.22), we get

$$g_2(\nabla_X^F F_* X, U) = g_1(X, X)g_2(H_2, U). \tag{5.23}$$

Since N is Kähler manifold, using (5.1) in (5.23), we get

$$g_2(\nabla_X^F J F_* X, J U) = g_1(X, X) g_2(H_2, U).$$
(5.24)

Since  $\nabla^N$  is Levi-Civita connection on N, using metric compatibility condition in (5.24), we get

$$-g_2(JF_*X, \nabla_X^F JU) = g_1(X, X)g_2(H_2, U). \tag{5.25}$$

Using (5.23) in (5.25), we get

$$-g_2(JF_*X, g_1(X, F_*JU)H_2) = g_1(X, X)g_2(H_2, U),$$
(5.26)

where  ${}^*F_*$  is the adjoint map of  $F_*$ . Now using  $H_2 = -\nabla^N g$  in (5.26), we get

$$g_1(X, {}^*F_*JU)g_2(JF_*X, \nabla^N g) = g_1(X, X)g_2(H_2, U),$$

which implies

$$g_1(X, F_*JU)JF_*X(g) = g_1(X, X)g_2(H_2, U).$$
 (5.27)

Since  $\dim(rangeF_*) > 1$  then by Theorem 5.5, g is constant on  $J(rangeF_*)$ , which means  $JF_*X(g) = 0$ . Then (5.27) implies  $g_2(H_2, U) = 0$ . Thus

$$H_2 = 0,$$
 (5.28)

which implies (i).

Since 
$$H_2 = trace \ (\nabla_X^F F_* Y)$$
. Then by (5.28), we get  $\nabla_X^F F_* Y = 0$ , which implies (ii).

**Theorem 5.7** Let  $F:(M^m,g_1) \to (N^n,g_2,J)$  be a Clairaut Lagrangian Riemannian map with  $\tilde{s}=e^g$  from a Riemannian manifold M to a Kähler manifold N such that  $\dim(rangeF_*) > 1$ . Then F is harmonic if and only if mean curvature vector field of  $\ker F_*$  is constant.

**Proof** Let  $F:(M^m,g_1)\to (N^n,g_2)$  be a smooth map between Riemannian manifolds. Then F is harmonic if and only if the tension field  $\tau(F)$  of map F vanishes. Then proof follows by Lemma 2.1 and Theorem 5.6.  $\square$ 

**Theorem 5.8** Let  $F:(M^m,g_1)\to (N^n,g_2,J)$  be a Clairaut Lagrangian Riemannian map with  $\tilde{s}=e^g$  from a Riemannian manifold M to a Kähler manifold N such that  $\dim(rangeF_*)>1$ . Then  $N=N_{rangeF_*}\times N_{(rangeF_*)^{\perp}}$  is a usual product manifold.

**Proof** The proof follows by [19] and Theorem 5.6.

**Example 5.9** Let  $M = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \neq 0, x_2 \neq 0\}$  be a Riemannian manifold with Riemannian metric  $g_1 = e^{2x_2} dx_1^2 + e^{2x_2} dx_2^2$  on M. Let  $N = \{(y_1, y_2) \in \mathbb{R}^2\}$  be a Riemannian manifold with Riemannian metric  $g_2 = e^{2x_2} dy_1^2 + dy_2^2$  on N and the complex structure J on N defined as  $J(y_1, y_2) = (-y_2, y_1)$ . Consider a map  $F: (M, g_1) \to (N, g_2, J)$  defined by

$$F(x_1, x_2) = \left(\frac{x_1 - x_2}{\sqrt{2}}, 0\right).$$

Then

$$ker F_* = span \Big\{ U = \frac{e_1 + e_2}{\sqrt{2}} \Big\} \text{ and } (ker F_*)^{\perp} = span \Big\{ X = \frac{e_1 - e_2}{\sqrt{2}} \Big\},$$

where  $\left\{e_1 = e^{-x_2} \frac{\partial}{\partial x_1}, e_2 = e^{-x_2} \frac{\partial}{\partial x_2}\right\}$  and  $\left\{e_1' = e^{-x_2} \frac{\partial}{\partial y_1}, e_2' = \frac{\partial}{\partial y_2}\right\}$  are bases on  $T_pM$  and  $T_{F(p)}N$  respectively, for  $p \in M$ . By easy computations, we see that  $F_*(X) = e_1'$  and  $g_1(X, X) = g_2(F_*X, F_*X)$  for  $X \in \Gamma(\ker F_*)^{\perp}$ . Thus F is Riemannian map with  $\operatorname{range} F_* = \operatorname{span} \{F_*(X) = e_1'\}$  and  $(\operatorname{range} F_*)^{\perp} = \operatorname{span} \{e_2'\}$ . Moreover it is easy to see that  $JF_*X = Je_1' = -e_2'$ . Thus F is an anti-invariant Riemannian map.

Now to show F is Clairaut Riemannian map we will find a smooth function g on N satisfying  $(\nabla F_*)(X,X) = -g_1(X,X)\nabla^N g$  for  $X \in \Gamma(\ker F_*)^{\perp}$ . Since  $(\nabla F_*)(X,X) \in \Gamma(\operatorname{range} F_*)^{\perp}$  for any  $X \in \Gamma(\ker F_*)^{\perp}$ . So here we can write  $(\nabla F_*)(X,X) = \operatorname{ae}'_2$ , for some  $a \in \mathbb{R}$ . Since  $\nabla^N g = e^{-2x_2} \frac{\partial g}{\partial y_1} \frac{\partial}{\partial y_1} + \frac{\partial g}{\partial y_2} \frac{\partial}{\partial y_2}$ . Hence  $\nabla^N g = -a\frac{\partial}{\partial y_2} = -\operatorname{ae}'_2$  for the function  $g = -\operatorname{ay}_2$ . Then it is easy to verify that  $(\nabla F_*)(X,X) = -g_1(X,X)\nabla^N g$ , where  $g_1(X,X) = 1$ , for vector field  $X \in \Gamma(\ker F_*)^{\perp}$  and we can easily see that  $\nabla^N_{e'_2} e'_2 = 0$ . Thus by Theorem 3.3, F is Clairaut anti-invariant Riemannian map.

### Acknowledgment

We would like to thank all the anonymous referees for his/her valuable comments and suggestions towards the improvement of quality of the paper. The first author, Kiran Meena gratefully acknowledges the financial support provided by the Human Resource Development Group - Council of Scientific and Industrial Research (HRDG-CSIR), New Delhi, India [File No.: 09/013(0887)/2019-EMR-I].

#### References

- [1] Allison D. Lorentzian Clairaut submersions. Geometriae Dedicata 1996; 63: 309-319. https://doi.org/10.1007/BF00181419
- [2] Akyol MA, Şahin B. Conformal anti-invariant Riemannian maps to Kähler manifolds. UPB Scientific Bulletin, Series A: Applied Mathematics and Physics 2018; 80 (4): 187-198.
- [3] Akyol MA, Şahin B. Conformal semi-invariant Riemannian maps to Kähler manifolds. Revista de la Union Matematica Argentina 2019; 60 (2): 459-468. https://doi.org/10.33044/revuma.v60n2a12
- [4] Besse AL. Einstein Manifolds. Springer-Verlag, Berlin, Heidelberg, New York, 1987.
- [5] Bishop RL. Clairaut submersions. Differential geometry (in Honor of K-Yano), Kinokuniya, Tokyo, 1972; 21-31.
- [6] Carmo MP. Differential geometry of curves and surfaces. Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1976.
- [7] Deshmukh S, Al-Solamy FR. Conformal vector fields on a Riemannian manifold. Balkan Journal of Geometry and its Applications 2014; 19 (2): 86-93.
- [8] Falcitelli M, Ianus S, Pastore AM. Riemannian Submersions and Related Topics. River Edge, NJ: World Scientific, 2004.
- [9] Fischer AE. Riemannian maps between Riemannian manifolds. Contemporary Mathematics 1992; 132: 331-366. https://doi.org/10.1090/conm/132/1188447
- [10] Gupta G, Sachdeva R, Kumar R, Rani R. On conformal Riemannian maps whose total manifolds admit a Ricci soliton. Journal of Geometry and Physics 2022; 178: 1-19. https://doi.org/10.1016/j.geomphys.2022.104539
- [11] Hamilton RS. The Ricci flow on surfaces, mathematics and general relativity. Contemporary Mathematics 1966; 71: 237-262. https://doi.org/10.1090/conm/071/954419
- [12] Lee J, Park J, Şahin B, Song D. Einstein conditions for the base space of anti-invariant Riemannian submersions and Clairaut submersions. Taiwanese Journal of Mathematics 2015; 19 (4): 1145-1160. https://doi.org/10.11650/tjm.19.2015.5283
- [13] Meena K, Yadav A. Conformal submersions whose total manifolds admit a Ricci soliton. Mediterranean Journal of Mathematics 2023; To appear.
- [14] Meena K, Zawadzki T. Clairaut conformal submersions. preprint, 2022, arXiv:2202.00393 [math.DG]. https://doi.org/10.48550/arXiv.2202.00393
- [15] Meriç SE, Kiliç E. Riemannian submersions whose total manifolds admit a Ricci soliton. International Journal of Geometric Methods in Modern Physics 2019; 16: 1950196-1-1950196-12. https://doi.org/10.1142/S0219887819501962
- [16] Nore T. Second fundamental form of a map. Annali di Matematica pura ed applicata 1986; 146: 281-310. https://doi.org/10.1007/BF/01762368
- [17] Perelman G. The Entropy formula for the Ricci flow and its geometric applications. preprint, 2002, arxiv: math/0211159 [math.DG].
- [18] Pigola S, Rigoli M, Rimoldi M, Setti AG. Ricci almost solitons. Annali della Scuola Normale Superiore di Pisa Classe di Scienze 2011; 10 (4): 757-799.

# Meena and Yadav/Turk J Math

- [19] Ponge R, Reckziegel H. Twisted products in pseudo-Riemannian geometry. Geometriae Dedicata 1993; 48 (1): 15-25. https://doi.org/10.1007/BF01265674
- [20] Şahin B. Invariant and anti-invariant Riemannian maps to Kähler manifolds. International Journal of Geometric Methods in Modern Physics 2010; 7 (3): 337-355. https://doi.org/10.1142/S0219887810004324
- [21] Şahin B. Conformal Riemannian maps between Riemannian manifolds, their harmonicity and decomposition theorems. Acta Applicandae Mathematicae 2010; 109: 829-847. https://doi.org/10.1007/s10440-008-9348-6
- [22] Şahin B. Semi-invariant Riemannian maps to Kähler manifolds. International Journal of Geometric Methods in Modern Physics 2011; 8 (7): 1439-1454. https://doi.org/10.1142/S0219887811005725
- [23] Şahin B. Holomorphic Riemannian maps. Journal of Mathematical Physics, Analysis, Geometry 2014; 10 (4): 422-430
- [24] Şahin B. Riemannian submersions, Riemannian maps in Hermitian geometry, and their applications. Academic Press: Elsevier, 2017.
- [25] Şahin B. Circles along a Riemannian map and Clairaut Riemannian maps. Bulletin of the Korean Mathematical Society 2017; 54 (1): 253-264. https://doi.org/10.4134/BKMS.b160082
- [26] Şahin B. A survey on differential geometry of Riemannian maps between Riemannian manifolds. Analele Stiintifice ale Universitatii Al I Cuza din Iasi Matematica 2017; 63: 151-167.
- [27] Taştan HM. On Lagrangian submersions. Hacettepe Journal of Mathematics and Statistics 2014; 43 (6): 993-1000. https://doi.org/10.15672/HJMS.2014437529
- [28] Watson B. Almost Hermitian submersions. Journal of Differential Geometry 1976; 11: 147-165.
- [29] Yadav A, Meena K. Riemannian maps whose total manifolds admit a Ricci soliton. Journal of Geometry and Physics 2021; 168: 1-13. https://doi.org/10.1016/j.geomphys.2021.104317
- [30] Yadav A, Meena K. Clairaut Riemannian maps whose total manifolds admit a Ricci soliton. International Journal of Geometric Methods in Modern Physics 2022; 19 (2): 2250024-1-2250024-17. https://doi.org/10.1142/S0219887822500244
- [31] Yadav A, Meena K. Clairaut invariant Riemannian maps with Kähler structure. Turkish Journal of Mathematics 2022; 46 (3): 1020-1035. https://doi.org/10.55730/1300-0098.3139
- [32] Yadav A, Meena K. Riemannian maps whose base manifolds admit a Ricci soliton. Publicationes Mathematicae Debrecen 2023; To appear. https://doi.org/10.5486/PMD.2023.9413
- [33] Yano K, Kon M. Structure on Manifolds. Singapore: World Scientific, 1984.