

Anisotropic curvature measures and uniqueness of convex bodies

Mario Santilli

Abstract

We prove that an arbitrary convex body $C \subseteq \mathbf{R}^{n+1}$, whose k -th anisotropic curvature measure (for $k = 0, \dots, n-1$) is a multiple constant of the anisotropic perimeter of C , must be a rescaled and translated Wulff shape. This result provides a generalization of a theorem of Schneider (1979) and resolves a conjecture of Andrews and Wei (2017).

1 Introduction

Jellett (1853) proved that a compact embedded star-shaped hypersurface of the Euclidean space with constant mean curvature must be a round sphere. A century later Hsiung (see [Hsi54]), extending a result of Süß (1929), proved that the same conclusion holds for compact embedded star-shaped hypersurfaces with constant higher-order mean curvatures. The fundamental results of Alexandrov (see [Ale62]), Ros (see [Ros87]), Korevaar-Ros (see [Ros88]) and Montiel-Ros (see [MR91]) guarantee that the same results hold without assuming that the hypersurfaces are star-shaped. These theorems, as well as their method of proof, lie at the core of modern differential geometry and its applications.

It is a natural question to extend these type of results to arbitrary convex bodies. Of course, in this setting one has to carefully choose the notion of curvature, in order to handle the unavoidable singular set. An insightful notion of curvature has been introduced by Federer in [Fed59] with the concept of *curvature measures*. If $C \subseteq \mathbf{R}^{n+1}$ is an arbitrary convex body, we denote with δ_C the distance function from C and with ξ_C the metric projection onto C . Then there exist uniquely determined Radon measures $\mathcal{C}_0(K, \cdot), \dots, \mathcal{C}_n(C, \cdot)$ supported on the boundary ∂C of C such that

$$\mathcal{L}^{n+1}(\{x \in \mathbf{R}^{n+1} : 0 < \delta_C(x) \leq \rho, \xi_C(x) \in B\}) = \sum_{m=0}^n \rho^{n+1-m} \mathcal{C}_m(C, B)$$

for every Borel subset $B \subseteq \partial C$. For an arbitrary convex body the equality $\mathcal{C}_n(C, \cdot) = \mathcal{H}^n \llcorner \partial C$ always holds. If ∂C is a \mathcal{C}^2 -hypersurface, then

$$\mathcal{C}_m(C, B) = \frac{1}{n+1-m} \int_B H_{n-m} d\mathcal{H}^n \quad \text{for } m = 0, \dots, n \text{ and } B \subseteq \partial C,$$

where

$$H_k = \sum_{1 \leq l_1 < \dots < l_k \leq n} \kappa_{l_1} \cdots \kappa_{l_k}$$

denotes the k -th mean curvature of ∂C (with $H_0 \equiv 1$). The aforementioned uniqueness results for hypersurfaces with constant k -th mean curvatures have been extended to arbitrary convex bodies by Schneider in 1979.

1.1 Theorem (cf. [Sch79]). *If $C \subseteq \mathbf{R}^{n+1}$ is an arbitrary convex body, $m = 0, \dots, n-1$, $\lambda > 0$ and $\mathcal{C}_m(C, \cdot) = \lambda \mathcal{C}_n(C, \cdot)$, then C is a round ball.*

Related characterizations of the round ball among arbitrary convex bodies can be found in [Koh95], [Koh98] and [San20b]. A very detailed account on many other uniqueness results for convex bodies can be found in [Sch14].

Besides of being a central result in the theory of convex bodies, Theorem 1.1 has recently emerged in [AW21] as an important tool to study the asymptotic behaviour of mixed-volume preserving flows in \mathbf{R}^{n+1} . If $\Omega \subseteq \mathbf{R}^{n+1}$ is a compact domain with \mathcal{C}^2 -boundary, we define for $k = 0, \dots, n$ the $n+1-k$ mixed volume $V_{n+1-k}(\Omega)$ as

$$V_{n+1-k}(\Omega) := \int_{\partial\Omega_t} E_{k-1}(x) d\mathcal{H}^n(x),$$

where $V_{n+1}(\Omega) = (n+1)\mathcal{L}^{n+1}(\Omega)$ and E_k is the normalized k -th mean curvature of $\partial\Omega$, namely $E_k = \binom{n}{k}^{-1} H_k$. Consider a smooth embedding $X_0 : M \rightarrow \mathbf{R}^{n+1}$ of a closed n -dimensional manifold M such that $X_0(M) = \partial\Omega_0$ is a smooth strictly convex hypersurface and a smooth flow $X : M^n \times [0, T) \rightarrow \mathbf{R}^{n+1}$ of the form

$$\begin{cases} \frac{\partial X}{\partial t}(x, t) = (\mu(t) - E_k(x, t)^{\alpha/k})\nu(x, t) \\ X(\cdot, 0) = X_0, \end{cases} \quad (1)$$

where $\alpha > 0$, $\nu(\cdot, t)$ is the outward unit-normal of the hypersurface $X(M, t) = \partial\Omega_t$, $k \in \{1, \dots, n\}$ and $\mu(t)$ is chosen to keep constant a general monotone function of $V_{n+1-k}(\Omega_t)$ and $V_{n+1}(\Omega_t)$ along the flow (in particular allowing to keep constant along the flow either only $V_{n+1-k}(\Omega_t)$ or only $V_{n+1}(\Omega_t)$); see [AW21, pag. 194]. The following bubbling theorem for the geometric flow in (1) is proved in [AW21].

1.2 Theorem (cf. [AW21, Theorem 1.1]). *The flow in (1) has a smooth strictly convex solution $\partial\Omega_t$ defined for all $t \geq 0$ and $\partial\Omega_t$ smoothly converges to a round sphere as $t \rightarrow \infty$.*

Special cases of this result were known before; see [AW21, pp. 195-196] for a detailed account. In particular, Sinestrari in [Sin15] treats the case $k = 1$ and $\alpha > 0$ (see also [BS18]). The main novelty of Theorem 1.2 is the treatment of the case $k > 1$ (for any exponent $\alpha > 0$). As explained in [AW21], the asymptotic analysis for the case $k > 1$ cannot be done in this case with the previously known approaches and the authors introduce a new method based on Theorem 1.1.

It is a natural question to extend Theorem 1.2 to the anisotropic setting and in [AW21] the authors make a conjecture in this direction. Before introducing the problem, let us briefly review few definitions. Let ϕ be a uniformly convex smooth norm on \mathbf{R}^{n+1} and we denote with ϕ^* its dual norm. The set $\mathcal{W}^\phi = \{x \in \mathbf{R}^{n+1} : \phi^*(x) = 1\}$ is called Wulff shape of ϕ . If $M = \partial\Omega \subseteq \mathbf{R}^{n+1}$ is a \mathcal{C}^2 -hypersurface and $\eta : M \rightarrow \mathbf{S}^n$ is the outward unit-normal vector field, then we define the (outward) anisotropic normal $\nu : M \rightarrow \mathcal{W}^\phi$ setting $\nu(x) = \nabla\phi(\eta(x))$ for $x \in M$. One can prove that $D\nu(x) \in \text{Hom}(\text{Tan}(M, x), \text{Tan}(M, x))$ and $D\nu(x)$ is diagonalizable with real eigenvalues $\kappa_1^\phi(x) \leq \dots \leq \kappa_n^\phi(x)$. We refer to these eigenvalues as the anisotropic principal curvatures of M at x . Then we define the anisotropic k -th man curvature of M as

$$H_k^\phi(x) = \sum_{1 \leq l_1 < \dots < l_k \leq n} \kappa_{l_1}^\phi(x) \cdots \kappa_{l_k}^\phi(x) \quad \text{for } k = 0, \dots, n,$$

where $H_0^\phi \equiv 1$. We define the renormalized k -th mean curvature $E_k^\phi = \binom{n}{k}^{-1} H_k^\phi$ and the anisotropic mixed-volume

$$V_{n+1-k}^\phi(\Omega) = \int_{\partial\Omega} \phi(\eta(x)) E_{k-1}^\phi(x) d\mathcal{H}^n(x) \quad \text{for } k = 0, \dots, n,$$

where η is the outward unit-normal of Ω and $V_{n+1}^\phi(\Omega) = (n+1)\mathcal{L}^{n+1}(\Omega)$. Using these definitions we can naturally formulate a general anisotropic version of (1). Given a closed strictly convex hypersurface $X_0 : M \rightarrow \mathbf{R}^{n+1}$, $\alpha > 0$, $k \in \{1, \dots, n\}$, we consider the flow of the form

$$\begin{cases} \frac{\partial X}{\partial t}(x, t) = (\mu^\phi(t) - E_k^\phi(x, t)^{\alpha/k})\nu^\phi(x, t) \\ X(\cdot, 0) = X_0, \end{cases} \quad (2)$$

where $\nu^\phi(\cdot, t)$ is the outward anisotropic normal of $X(M, t) = \partial\Omega_t$ and the term $\mu^\phi(t)$ is chosen to keep constant along the flow a monotone function $V_{n+1-k}^\phi(\Omega_t)$ and $V_{n+1}^\phi(\Omega_t)$. It is remarked in [AW21, pag. 219] that *if* Theorem 1.1 can be extended to the anisotropic setting, *then* the method of [AW21] carries through directly to prove the following Conjecture on the asymptotic behaviour of the anisotropic flow in (2).

1.3 Conjecture (cf. [AW21]). *Suppose $k \in \{2, \dots, n-1\}$. The flow in (1) has a smooth strictly convex solution $\partial\Omega_t$ defined for all $t \geq 0$ and $\partial\Omega_t$ smoothly converges to a scaled and translated Wulff shape of ϕ as $t \rightarrow \infty$.*

We remark that the cases $k = 1$ and $k = n$ can be proved with other techniques (see [AW21, pag. 219]).

To formulate the anisotropic version of Theorem 1.1, we first need to introduce the anisotropic analogous of curvature measures. To the best of our knowledge this notion has been first studied in [HL00]. If $C \subseteq \mathbf{R}^{n+1}$ is an arbitrary convex body, we denote with δ_C^ϕ the distance function from C with respect to the norm ϕ and with ξ_C^ϕ the metric projection onto C with respect to ϕ . Then there exist uniquely determined Radon measures $\mathcal{C}_0^\phi(C, \cdot), \dots, \mathcal{C}_n^\phi(C, \cdot)$ supported on the boundary ∂C of C such that

$$\mathcal{L}^{n+1}(\{x \in \mathbf{R}^{n+1} : 0 < \delta_C^\phi(x) \leq \rho, \xi_C^\phi(x) \in B\}) = \sum_{m=0}^n \rho^{n+1-m} \mathcal{C}_m^\phi(C, B)$$

for every Borel subset $B \subseteq \partial C$; cf. [HL00, Theorem 2.3]. The following conjecture is explicitly formulated in [AW21].

1.4 Conjecture (cf. [AW21, 8.2]). *If $C \subseteq \mathbf{R}^{n+1}$ is an arbitrary convex body, $m = 0, \dots, n-1$, $\lambda > 0$ and $\mathcal{C}_m^\phi(C, \cdot) = \lambda \mathcal{C}_m^\phi(C, \cdot)$, then C is a scaled and translated Wulff shape of ϕ .*

In this paper we prove Conjecture 1.4. As already pointed out, Conjecture 1.3 follows from Conjecture 1.4 with a direct extension of the method in [AW21] to the anisotropic setting.

We now describe the content of the paper. In the preliminary section 2, besides introducing the notation and recalling few facts on the geometry of the Wulff shapes, we introduce a notion of anisotropic normal bundle for closed sets and we recall from the recent work [KS21] the Lipschitz and differentiability properties of the anisotropic nearest point projection onto a closed set. The nearest point projection onto a closed set K might not be single-valued at some points of $\mathbf{R}^{n+1} \setminus K$ (if it is everywhere single-valued then the set K is convex); indeed the set of points where it fails to be single-valued might even be dense in $\mathbf{R}^{n+1} \setminus K$, and even if K is the complementary of a convex body with \mathcal{C}^1 -boundary; see [San21]. Therefore in our proofs we use an approach based on the theory of multivalued functions.

In Section 3 we introduce the anisotropic higher-order mean curvatures for an arbitrary closed set K as functions on the anisotropic normal bundle. We use them to find a (local) formula for the anisotropic tubular neighbourhood around K in Theorem 3.12. This formula generalizes the isotropic Steiner formula for arbitrary closed sets proved in [HLW04]. Our motivation to develop a theory for arbitrary (non-convex) closed sets is given by the fact that in our proof of Theorem 5.2 we need to work with the complementary of the convex body C . In fact, the proof of the inequality employs the general Steiner formula of Theorem 3.12 with $K = \mathbf{R}^{n+1} \setminus C$, while the analysis of the equality case uses Corollary 3.16 of Theorem 3.12 always for the complementary of C .

In Section 4, specializing the Steiner formula of section 3 to convex sets, we obtain an integral representation of the anisotropic curvature measures. This result extends to the anisotropic setting the classical integral representation in [Zäh86] for the isotropic curvature measures. Then we prove an anisotropic version of the Minkowski formulae for arbitrary convex bodies and we use them to study the k -th mean curvature of an arbitrary convex body $C \subseteq \mathbf{R}^{n+1}$, whose $(n-k)$ -th anisotropic curvature measure satisfies $C_{n-k}^\phi(K, \cdot) = \lambda C_n^\phi(K, \cdot)$ for some constant $\lambda > 0$. In particular we obtain a lower bound for λ in terms of the anisotropic isoperimetric ratio of C .

Finally in Section 5 we prove an optimal geometric inequality for arbitrary convex bodies. This inequality is inspired by an inequality originally found by Heintze-Karcher in [HK78] and used by Montiel-Ros in [MR91] to prove the uniqueness of compact smooth hypersurfaces with constant higher-order mean curvature. Combining the optimal geometric inequality with the lower bound for the constant λ in section 4 we finally obtain the proof of conjecture 1.4.

2 Preliminaries

In general, but with few exceptions explained below, we follow the notation and terminology of [Fed69] (see [Fed69, pp. 669-676]).

We denote by \bullet a fixed scalar product on \mathbf{R}^{n+1} and by $|\cdot|$ its associated norm. We denote with \mathbf{S}^n the unit sphere; i.e. $\mathbf{S}^n = \{x \in \mathbf{R}^{n+1} : |x| = 1\}$. The map $\mathbf{p} : \mathbf{R}^{n+1} \times \mathbf{R}^{n+1} \rightarrow \mathbf{R}^{n+1}$ is the projection onto the first component; i.e. $\mathbf{p}(x, \eta) = x$. If $S \subseteq \mathbf{R}^k$ and $a \in \mathbf{R}^p$, then we denote with $\text{Tan}(S, a)$ the tangent cone of S at a (see [Fed69, 3.1.21]) and with $\text{Tan}^m(\mathcal{H}^m \lrcorner S, a)$ the cone of all $(\mathcal{H}^m \lrcorner S, m)$ approximate tangent vectors at a (see [Fed69, 3.2.16]). For an (\mathcal{H}^m, m) rectifiable and \mathcal{H}^m -measurable set $S \subseteq \mathbf{R}^p$, the cone $\text{Tan}^m(\mathcal{H}^m \lrcorner S, a)$ is an m -dimensional plane for \mathcal{H}^m a.e. $a \in S$ (see [Fed69, 3.2.19]); in this case, if $f : S \rightarrow \mathbf{R}^q$ is a Lipschitzian function and $k \in \{1, \dots, m\}$, then we denote with $\text{ap} J_k^S f$ its $(\mathcal{H}^m \lrcorner S, m)$ -approximate k -dimensional Jacobian. See [Fed69, 3.2.19, 3.2.10 and 3.2.22] for details about this definition and applications to area/coarea formula that will be used in this paper.

If X is a topological space and $S \subseteq X$ then its topological boundary is ∂S and the characteristic map of S is $\mathbf{1}_S$. If $Q \subseteq X \times Y$ and $S \subseteq X$, we define $Q|_S = \{(x, y) \in Q : x \in S\}$.

2.1 Multivalued maps

A map T defined on a set X is called Y -multivalued, if $T(x)$ is a subset of Y for every $x \in X$. If $T(x)$ is a singleton, with a little abuse of notation we denote with $T(x)$ the unique element of the set $T(x) \subseteq Y$. Suppose $(X, \|\cdot\|)$ and $(Y, \|\cdot\|)$ are finite dimensional normed vectorspaces, T is a Y -multivalued map such that $T(u) \neq \emptyset$ for every $u \in X$ and $x \in X$.

- (1) We say that T is *weakly continuous at x* if and only if for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$T(y) \subseteq T(x) + \{v \in Y : \|v\| < \epsilon\} \quad \text{whenever } \|y - x\| < \delta;$$

if, additionally, $T(x)$ is a singleton, then we say that T is continuous at x .

- (2) We say that T is *strongly differentiable at $x \in X$* if and only if $T(x)$ is a singleton and there exists a linear map $L : X \rightarrow Y$ such that for any $\epsilon > 0$ there exists $\delta > 0$ satisfying

$$\|w - T(x) - L(y - x)\| \leq \epsilon \|y - x\| \quad \text{whenever } \|x - y\| \leq \delta \text{ and } w \in T(y);$$

cf. [KS21, Definition 2.21]. The linear map L is unique (cf. [KS21, Remark 2.22]) and we denote it with $\text{DT}(x)$. Moreover we denote with dmn DT the set of points $x \in X$ where T is strongly differentiable.

The following general fact on the Borel measurability of the differential of a multivalued map will be useful.

2.1 Lemma. *Suppose $(X, \|\cdot\|)$ and $(Y, \|\cdot\|)$ are finite dimensional normed vectorspaces, T is a Y -multivalued weakly continuous map such that $T(u) \neq \emptyset$ for every $u \in X$.*

Then $\{x \in X : T(x) \text{ is a singleton}\}$ and dmn DT are Borel subsets of X and $\text{DT} : \text{dmn DT} \rightarrow \text{Hom}(X, Y)$ is a Borel function.

Proof. We define $U = \{x \in X : T(x) \text{ is a singleton}\}$ and the function $\text{diam} : 2^Y \setminus \{\emptyset\} \rightarrow [0, \infty]$ as $\text{diam } S = \sup\{\|y_1 - y_2\| : y_1, y_2 \in S\}$ for every $S \in 2^Y \setminus \{\emptyset\}$. Noting that $\text{diam} \circ T : X \rightarrow [0, +\infty]$ is upper-semicontinuous, we conclude that $U = \{x \in X : \text{diam}(T(x)) = 0\}$ is a Borel subset of X .

For the positive integers i, j we define

$$C_{ij} = \left\{ (x, L) \in U \times \text{Hom}(X, Y) : \|w - T(x) - L(h)\| \leq \frac{1}{i} \|h\| \quad \text{for } \|h\| \leq \frac{1}{j} \text{ and } w \in T(x+h) \right\}.$$

We prove that C_{ij} is relatively closed in $U \times \text{Hom}(X, Y)$. By contradiction assume that C_{ij} is not closed. Then there exists $(x_0, L_0) \in (U \times \text{Hom}(X, Y)) \setminus C_{ij}$ and a sequence $(x_k, L_k) \in C_{ij}$ converging to (x_0, L_0) . We notice that there exist $h_0 \in X$ with $\|h_0\| \leq \frac{1}{j}$ and $w_0 \in T(x_0 + h_0)$ such that

$$\|w_0 - T(x_0) - L_0(h_0)\| > \frac{1}{i} \|h_0\|,$$

we define $h_k = x_0 + h_0 - x_k$ for every $k \geq 1$ and we select $k_0 \geq 1$ so that $\|h_k\| \leq \frac{1}{j}$ for every $k \geq k_0$. Since $x_0 + h_0 = x_k + h_k$ and $w_0 \in T(x_k + h_k)$ for every $k \geq 1$, we infer that

$$\|w_0 - T(x_k) - L_k(h_k)\| \leq \frac{1}{i} \|h_k\| \quad \text{for every } k \geq k_0.$$

Noting that $T(x_k) \rightarrow T(x_0)$ and $h_k \rightarrow h_0$ as $k \rightarrow \infty$, we deduce that

$$\|w_0 - T(x_0) - L_0(h_0)\| \leq \frac{1}{i} \|h_0\|$$

and we obtain a contradiction.

Let $G := \{(x, DT(x)) : x \in \text{dmn } DT\}$ and $\pi_X : X \times \text{Hom}(X, Y) \rightarrow X$, $\pi_X(x, T) = x$ for every $(x, T) \in X \times \text{Hom}(X, Y)$. Noting that

$$G = \bigcap_{i=1}^{\infty} \bigcup_{j=1}^{\infty} C_{ij},$$

we infer that G is a Borel subset of $U \times \text{Hom}(X, Y)$. Since $\pi|_G$ is injective, we obtain the conclusion from [Fed69, 2.2.10]. \square

2.2 Norms and Wulff shapes.

Let ϕ be a norm on \mathbf{R}^{n+1} . We say that ϕ is a C^k -norm if and only if $\phi \in C^k(\mathbf{R}^{n+1} \setminus \{0\})$. We say that ϕ is *uniformly convex* if and only if there exists a constant $\gamma > 0$ (ellipticity constant) such that the function $\mathbf{R}^{n+1} \ni u \rightarrow \phi(u) - \gamma|u|$ is convex. If ϕ is a uniformly convex C^2 -norm then

$$D^2 \phi(u)(v, v) \geq \gamma|v|^2$$

for all $u \in \mathbf{R}^{n+1}$ with $|u| = 1$ and for all $v \in \mathbf{R}^{n+1}$ perpendicular to u .

For any norm ϕ we denote by ϕ^* the *conjugate norm* of ϕ ; namely if $u \in \mathbf{R}^n$ then $\phi^*(u) = \sup\{v \bullet u : \phi(v) = 1\}$. It is well known that if ϕ is a uniformly convex C^2 -norm then ϕ^* is a uniformly convex C^2 -norm. We refer to [DRKS20, Lemma 2.32] for this and other basic facts on ϕ and ϕ^* . These facts will be tacitly used through the paper. If $B = \{x \in \mathbf{R}^{n+1} : \phi^*(x) \leq 1\}$ we define the *Wulff shape* (or *Wulff crystal*) of ϕ as

$$\mathcal{W}^\phi = \partial B.$$

If ϕ is a uniformly convex C^2 -norm then the Wulff shape of ϕ is a uniformly convex C^2 -hypersurface. In this case we denote the exterior unit-normal of B with $\mathbf{n}^\phi : \mathcal{W}^\phi \rightarrow \mathbf{S}^n$; we remark (see [DRKS20, 2.32]) that \mathbf{n}^ϕ is a C^1 -diffeomorphism onto \mathbf{S}^n and

$$\nabla \phi(\mathbf{n}^\phi(u)) = u \quad \text{for } u \in \mathcal{W}^\phi, \quad \mathbf{n}^\phi(\nabla \phi(\eta)) = \eta \quad \text{for } \eta \in \mathbf{S}^n. \quad (3)$$

2.3 Distance function and normal bundle.

Warning. In this paper sometimes we refer to [KS21]. Notice that in this paper we use the same symbols with a different meaning; compare the definitions below with those given in [KS21, Section 2].

Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed and ϕ is a uniformly convex C^2 -norm on \mathbf{R}^{n+1} . If ϕ is the Euclidean norm the dependence on ϕ is omitted in all the symbols introduced below.

The ϕ -distance function $\delta_K^\phi : \mathbf{R}^{n+1} \rightarrow \mathbf{R}$ is defined by

$$\delta_K^\phi(x) = \inf\{\phi^*(x - c) : c \in K\} \quad \text{for every } x \in \mathbf{R}^{n+1}$$

and

$$S^\phi(K, r) = \{x : \delta_K^\phi(x) = r\} \quad \text{for } r > 0.$$

The set $\text{Unp}^\phi(K)$ is the set of $x \in \mathbf{R}^{n+1} \setminus K$ such that there exists a unique $c \in K$ with $\phi^*(x - c) = \delta_K^\phi(x)$. Since δ_K^ϕ is a Lipschitz map, it follows that $\mathcal{L}^{n+1}(\mathbf{R}^{n+1} \setminus (K \cup \text{Unp}^\phi(K))) = 0$. The *nearest ϕ -projection* $\xi_K^\phi : \mathbf{R}^{n+1} \rightarrow \mathbf{2}^K$ is the K -multivalued map characterized by

$$\xi_K^\phi(x) = \{c \in K : \delta_K^\phi(x) = \phi^*(x - c)\} \quad \text{for every } x \in \mathbf{R}^{n+1}.$$

This is a weakly continuous by [KS21, Lemma 2.30(f)]; moreover notice that $\text{Unp}^\phi(K)$ is a Borel subset of \mathbf{R}^{n+1} by Lemma 2.1. The ϕ -Cahn-Hoffman map of K is the \mathcal{W}^ϕ -multivalued function $\nu_K^\phi : \mathbf{R}^{n+1} \setminus K \rightarrow \mathbf{2}^{\mathcal{W}^\phi}$ defined by

$$\nu_K^\phi(x) = \delta_K^\phi(x)^{-1}(x - \xi_K^\phi(x)) \quad \text{for } x \in \mathbf{R}^{n+1} \setminus K.$$

Finally we set $\psi_K^\phi : \mathbf{R}^{n+1} \setminus K \rightarrow \mathbf{2}^K \times \mathbf{2}^{\mathcal{W}^\phi}$ by

$$\psi_K^\phi(x) = (\xi_K^\phi(x), \nu_K^\phi(x)) \quad \text{for } x \in \mathbf{R}^{n+1} \setminus K.$$

We define the ϕ -unit normal bundle of K as

$$N^\phi(K) = \{(x, \eta) : x \in K, \eta \in \mathcal{W}^\phi, \delta_K^\phi(x + r\eta) = r \text{ for some } r > 0\}$$

and we set $N^\phi(K, x) = N^\phi(K)|_{\{x\}}$. We recall (cf. [DRKS20, Lemma 5.2]) that $N^\phi(K)$ is Borel and countably $(n-1)$ -rectifiable (in the sense of [Fed69, 3.2.14(2)]) subset of $\mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$; moreover

$$N^\phi(K) = \{(a, \nabla\phi(\eta)) : (a, \eta) \in N(K)\}. \quad (4)$$

The normal ϕ -distance function to the cut locus of K is the upper-semicontinuous function $r_K^\phi : N^\phi(K) \rightarrow (0, +\infty]$ given by

$$r_K^\phi(a, \eta) = \sup\{s : \delta_K^\phi(a + s\eta) = s\} \quad \text{for } (a, \eta) \in N^\phi(K)$$

and the ϕ -cut locus of K is given by

$$\text{Cut}^\phi(K) = \{a + r_K^\phi(a, \eta)\eta : (a, \eta) \in N^\phi(K)\}.$$

We recall that $\mathcal{L}^{n+1}(\text{Cut}^\phi(K)) = 0$; cf. [DRKS20, Remark 5.11]; if K is convex then $\text{Cut}^\phi(K) = \emptyset$. A related function which will be useful in the sequel is defined as

$$\rho_K^\phi(x) = \sup\{s \geq 0 : \delta_K^\phi(a + s(x-a)) = s\delta_K^\phi(x)\} \quad \text{for } x \in \mathbf{R}^{n+1} \setminus K \text{ and } a \in \xi_K^\phi(x).$$

This definition does not depend on the choice of $a \in \xi_K^\phi(x)$ and the function $\rho_K^\phi : \mathbf{R}^{n+1} \setminus K \rightarrow [1, +\infty]$ is upper-semicontinuous; cf. [KS21, Remark 2.29]. Notice that $\{x : \rho_K^\phi(x) > 1\} \subseteq \text{Unp}^\phi(K)$ and

$$r_K^\phi(a, u) = r\rho_K^\phi(a + ru) \quad \text{for every } (a, u) \in N^\phi(K) \text{ and } 0 < r < r_K^\phi(a, u) \quad (5)$$

Let $\Sigma^\phi(K)$ be the set of non-differentiability points of δ_K^ϕ in $\mathbf{R}^{n+1} \setminus K$. It is well known (see [KS21] and references therein) that

$$\Sigma^\phi(K) \subseteq \text{Cut}^\phi(K) \subseteq \text{Clos} \Sigma^\phi(K) \quad \text{and} \quad \Sigma^\phi(K) = \mathbf{R}^{n+1} \setminus (K \cup \text{Unp}^\phi(K));$$

moreover for $x \in \text{Unp}^\phi(K)$

$$\nabla\delta_K^\phi(x) = \nabla\phi^*(x - \xi_K^\phi(x)) \in \mathcal{W}^{\phi^*} \quad \text{and} \quad \nabla\phi(\nabla\delta_K^\phi(x)) = \nu_K^\phi(x) \in \mathcal{W}^\phi \quad (6)$$

cf. [KS21, Lemma 2.30(c)]. It follows from [DRKS20, Lemma 2.32] that $\nu_K^\phi(x) \bullet \nabla\delta_K^\phi(x) = \phi(\nabla\delta_K^\phi(x)) = 1$ for $x \in \text{Unp}^\phi(K)$; in particular $\nu_K^\phi(x)$ and $\nabla\delta_K^\phi(x)$ are linearly independent.

2.2 Theorem (cf. [KS21, Corollary 3.10]). *Suppose ϕ is a uniformly convex \mathcal{C}^2 -norm on \mathbf{R}^{n+1} , $K \subseteq \mathbf{R}^{n+1}$ is closed, $1 < \lambda < \infty$, $0 < s < t < \infty$, and*

$$K_{\lambda,s,t} = \{x \in \mathbf{R}^{n+1} \setminus K : \rho_K^\phi(x) \geq \lambda, s \leq \delta_K^\phi(x) \leq t\}.$$

Then $\xi_K^\phi|_{K_{\lambda,s,t}}$ is Lipschitz continuous.

2.3 Theorem (cf. [KS21, Theorem 1.4]). *Suppose ϕ is a uniformly convex \mathcal{C}^2 -norm on \mathbf{R}^{n+1} and $K \subseteq \mathbf{R}^{n+1}$ is closed.*

Then $\mathcal{L}^{n+1}(\mathbf{R}^{n+1} \setminus (K \cup \text{dnn}(\text{D}\nu_K^\phi))) = 0$ and

$$\{a + r\eta : 0 < r < r_K^\phi(a, \eta)\} \subseteq \text{dnn}(\text{D}\nu_K^\phi)$$

for \mathcal{H}^n almost all $(a, \eta) \in N^\phi(K)$.

3 Anisotropic Steiner formula for arbitrary closed sets

In this section we assume that ϕ is a uniformly convex \mathcal{C}^2 norm. We start recalling few known facts.

3.1 Remark. Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed. If $x \in \text{Unp}^\phi(K)$, $r = \delta_K^\phi(x)$, $0 < t < 1$ and $y = \xi_K^\phi(x) + t\nu_K^\phi(x)$, then $y \in \text{Unp}^\phi(K)$ and

$$\text{Tan}(S^\phi(K, r), x) = \{v \in \mathbf{R}^{n+1} : v \bullet \nabla \delta_K^\phi(x) = 0\},$$

$$\nabla \delta_K^\phi(x) = \nabla \delta_K^\phi(y), \quad \text{Tan}(S^\phi(K, r), x) = \text{Tan}(S^\phi(K, tr), y).$$

cf. [KS21, Lemma 2.30, Lemma 2.33].

3.2 Remark. For every $(a, u) \in N^\phi(K)$ and for every $0 < r < r_K^\phi(a, u)$ we have that

$$\text{Tan}(S^\phi(K, r), a + ru) = \text{Tan}(\mathcal{W}^\phi, u).$$

Setting $\eta = \frac{\nabla \delta_K^\phi(a+ru)}{|\nabla \delta_K^\phi(a+ru)|}$, this assertion follows from Remark 3.1, noting that (see (3) and (6))

$$\nabla \phi(\eta) = \nabla \phi(\nabla \delta_K^\phi(a + ru)) = u, \quad \mathbf{n}^\phi(u) = \eta.$$

3.3 Lemma (cf. [KS21, Lemma 4.8]). *Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed, $x \in \text{dmn}(\text{D}\nu_K^\phi)$ and $r = \delta_K^\phi(x)$.*

Then $\rho_K^\phi(x) > 1$, $\text{D}\nu_K^\phi(x)(\nu_K^\phi(x)) = 0$, $\text{D}\nu_K^\phi(x)[\text{Tan}(S^\phi(K, r), x)] \subseteq \text{Tan}(S^\phi(K, r), x)$ and the linear automorphism $\text{D}\nu_K^\phi(x)|_{\text{Tan}(S^\phi(K, r), x)}$ is diagonalizable with real eigenvalues. Moreover for every eigenvalue χ of $\text{D}\nu_K^\phi(x)|_{\text{Tan}(S^\phi(K, r), x)}$ holds that

$$\frac{1}{(1 - \rho_K^\phi(x))r} \leq \chi \leq \frac{1}{r}.$$

Proof. All the assertions of this statement with the exception of the diagonalizability property of $\text{D}\nu_K^\phi(x)|_{\text{Tan}(S^\phi(K, r), x)}$, are contained in [KS21, Lemma 4.8]. Indeed, the first part of the proof of this lemma in combination with [DRKS20, Remark 2.25] shows that $\text{D}\nu_K^\phi(x)|_{\text{Tan}(S^\phi(K, r), x)}$ is diagonalizable. \square

3.4 Definition. Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed, $x \in \text{dmn}(\text{D}\nu_K^\phi)$ and $r = \delta_K^\phi(x)$. We define

$$\chi_{K,1}^\phi(x) \leq \dots \leq \chi_{K,n}^\phi(x)$$

to be the eigenvalues (counted with their algebraic multiplicity) of $\text{D}\nu_K^\phi(x)|_{\text{Tan}(S^\phi(K, r), x)}$.

3.5 Lemma. *The set $\text{dmn D}\nu_K^\phi$ is a Borel subset of \mathbf{R}^{n+1} and the functions $\chi_{K,i}^\phi : \text{dmn D}\nu_K^\phi \rightarrow \mathbf{R}$ are Borel functions.*

Proof. Let X be the set of all automorphisms $T \in \text{Hom}(\mathbf{R}^{n+1}, \mathbf{R}^{n+1})$ with real eigenvalues. For each $T \in X$ we define $\lambda_0(T) \leq \dots \leq \lambda_n(T)$ to be the eigenvalues of T counted with their algebraic multiplicity and we define the map $\lambda : X \rightarrow \mathbf{R}^{n+1}$ as

$$\lambda(T) = (\lambda_0(T), \dots, \lambda_n(T)) \quad \text{for } T \in X.$$

We observe that λ is a continuous map by [HM87, Theorem B]. Moreover we notice that $\text{dmn D}\nu_K^\phi = \text{dmn D}\xi_K^\phi$ and that this is a Borel subset of \mathbf{R}^{n+1} by Lemma 2.1. For each $x \in \text{dmn D}\xi_K^\phi$, since $\text{D}\xi_K^\phi(x)(\nu_K^\phi(x)) = 0$, we have that

$$\lambda_0(\text{D}\xi_K^\phi(x)) = 0 \quad \text{and} \quad \lambda_i(\text{D}\xi_K^\phi(x)) = 1 - \delta_K^\phi(x)\chi_{K,n+1-i}^\phi(x) \geq 0 \quad \text{for } i = 1, \dots, n.$$

Since the map $\text{D}\xi_K^\phi : \text{dmn D}\xi_K^\phi \rightarrow X$ is a Borel function, we obtain the conclusion. \square

3.6 Lemma. Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed, $(a, u) \in N^\phi(K)$, $0 < r < s < \mathbf{r}_K^\phi(a, u)$ so that $a + ru, a + su \in \text{dmn } D\nu_K^\phi$ and $\tau_1, \dots, \tau_n \in \text{Tan}(\mathcal{W}^\phi, u)$.

Then $D\nu_K^\phi(a + ru)\tau_i = \chi_{K,i}^\phi(a + ru)\tau_i$ for every $i = 1, \dots, n$ if and only if $D\nu_K^\phi(a + su)\tau_i = \chi_{K,i}^\phi(a + su)\tau_i$ for every $i = 1, \dots, n$, in which case holds that

$$\frac{\chi_{K,i}^\phi(a + ru)}{1 - r\chi_{K,i}^\phi(a + ru)} = \frac{\chi_{K,i}^\phi(a + su)}{1 - s\chi_{K,i}^\phi(a + su)}.$$

Proof. We define $x = a + ru$, $y = a + su$ and $t = \frac{r}{s}$. We notice that ξ_K^ϕ is differentiable at y and

$$D\xi_K^\phi(y)|\text{Tan}(\mathcal{W}^\phi, u) = \text{Id}_{\text{Tan}(\mathcal{W}^\phi, u)} - sD\nu_K^\phi(y)|\text{Tan}(\mathcal{W}^\phi, u).$$

Let $\xi : \mathbf{R}^{n+1} \setminus K \rightarrow K$ and $\nu : \mathbf{R}^{n+1} \setminus K \rightarrow \mathcal{W}^\phi$ such that $\xi(z) \in \xi_K^\phi(z)$ and $\nu(z) = \delta_K^\phi(z)^{-1}(z - \xi(z))$ for each $z \in \mathbf{R}^{n+1} \setminus K$. It follows from the strict convexity of ϕ (see Remark [KS21, 2.11]) that

$$\nu(\xi(z) + t(z - \xi(z))) = \nu(z) \quad \text{for } z \in \mathbf{R}^{n+1} \setminus K;$$

differentiating this equality in y we obtain

$$D\nu(x) \circ [D\xi(y) + t(\text{Id}_{\mathbf{R}^{n+1}} - D\xi(y))] = D\nu(y).$$

Assume now $D\nu(y)\tau_i = \chi_{K,i}^\phi(y)\tau_i$ for every $i = 1, \dots, n$. Then we get

$$\chi_{K,i}^\phi(y)\tau_i = [1 - (s - r)\chi_{K,i}^\phi(y)]D\nu(x)\tau_i \quad \text{for every } i = 1, \dots, n.$$

Since $1 - (s - r)\chi_{K,i}^\phi(y) > 0$ for every $i = 1, \dots, n$ by Lemma 3.3, we conclude that

$$D\nu(x)\tau_i = \chi_{K,i}^\phi(x)\tau_i, \quad \chi_{K,i}^\phi(x) = \frac{\chi_{K,i}^\phi(y)}{1 - (s - r)\chi_{K,i}^\phi(y)}$$

and

$$\frac{\chi_{K,i}^\phi(x)}{1 - r\chi_{K,i}^\phi(x)} = \frac{\chi_{K,i}^\phi(y)}{1 - s\chi_{K,i}^\phi(y)}$$

for every $i = 1, \dots, n$.

The last paragraph shows in particular that $D\nu(x)|\text{Tan}(\mathcal{W}^\phi, u)$ and $D\nu(y)|\text{Tan}(\mathcal{W}^\phi, u)$ have the same number k of distinct eigenvalues. Denoting with $N_1(x), \dots, N_k(x)$ and $N_1(y), \dots, N_k(y)$ the eigenspaces of $D\nu(x)|\text{Tan}(\mathcal{W}^\phi, u)$ and $D\nu(y)|\text{Tan}(\mathcal{W}^\phi, u)$ respectively, we can also derive from the last paragraph the inclusions $N_i(y) \subseteq N_i(x)$ for every $i = 1, \dots, k$. Since

$$N_1(y) \oplus \dots \oplus N_k(y) = \text{Tan}(\mathcal{W}^\phi, u) = N_1(x) \oplus \dots \oplus N_k(x),$$

we conclude that $N_i(y) = N_i(x)$ for every $i = 1, \dots, n$ and the proof is completed. \square

3.7 Definition. Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed. We define

$$\tilde{N}^\phi(K) = \{(a, u) \in N^\phi(K) : a + ru \in \text{dmn}(D\nu_K^\phi) \text{ for every } 0 < r < \mathbf{r}_K^\phi(a, u)\}$$

and

$$\kappa_{K,i}^\phi(a, u) = \frac{\chi_{K,i}^\phi(a + ru)}{1 - r\chi_{K,i}^\phi(a + ru)}$$

for $(a, u) \in \tilde{N}^\phi(K)$, $0 < r < \mathbf{r}_K^\phi(a, u)$ and $i = 1, \dots, n$.

3.8 Remark. Lemma 3.6 ensures that the definition of $\kappa_{K,i}^\phi(a, u)$ does not depend on the choice of r and Theorem 2.3 ensures that $\mathcal{H}^n(N^\phi(K) \setminus \tilde{N}^\phi(K)) = 0$. In particular we notice that $\tilde{N}^\phi(K)$ is an \mathcal{H}^n -measurable subset of $N^\phi(K)$. Moreover by Remark 3.5 we deduce that the functions $\kappa_{K,i}^\phi$ are $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable.

3.9 Remark. Noting (5), we conclude from Lemma 3.3 that

$$-\frac{1}{\mathbf{r}_K^\phi(a, u)} \leq \kappa_{K,i}^\phi(a, u) \leq +\infty$$

for every $(a, u) \in \tilde{N}^\phi(K)$ and $i = 1, \dots, n$.

3.10 Lemma. Let $K \subseteq \mathbf{R}^{n+1}$ be closed. Suppose $\tau_i : \tilde{N}^\phi(K) \rightarrow \mathbf{R}^{n+1}$, for $i = 1, \dots, n$, are defined so that $\tau_1(a, u), \dots, \tau_n(a, u)$ form a basis of $\text{Tan}(\mathcal{W}^\phi, u)$ with

$$D\boldsymbol{\nu}_K^\phi(a + ru)(\tau_i(a, u)) = \chi_{K,i}^\phi(a + ru)\tau_i(a, u) \quad \text{for } i = 1, \dots, n \text{ and } 0 < r < \mathbf{r}_K^\phi(a, u)$$

and $\zeta_i : \tilde{N}^\phi(K) \rightarrow \mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$, for $i = 1, \dots, n$, are defined so that

$$\zeta_i(a, u) = \begin{cases} (\tau_i(a, u), \kappa_{K,i}^\phi(a, u)\tau_i(a, u)) & \text{if } \kappa_{K,i}^\phi(a, u) < \infty \\ (0, \tau_i(a, u)) & \text{if } \kappa_{K,i}^\phi(a, u) = +\infty \end{cases}$$

For every \mathcal{H}^n -measurable subset $W \subseteq N^\phi(K)$ with $\mathcal{H}^n(W) < \infty$ and for \mathcal{H}^n almost all $(a, u) \in W$ we have that $\text{Tan}^n(\mathcal{H}^n \llcorner W, (a, u))$ is an n -dimensional plane and $\zeta_1(a, u), \dots, \zeta_n(a, u)$ form a basis of $\text{Tan}^n(\mathcal{H}^n \llcorner W, (a, u))$. Moreover,

$$\text{ap } J_n^W \mathbf{p}(a, u) = \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} \mathbf{1}_{\tilde{N}_n^\phi(K)}(a, u)$$

for \mathcal{H}^n almost all $(a, u) \in W$.

Proof. Assume $W \subseteq N^\phi(K)$ is \mathcal{H}^n -measurable and $\mathcal{H}^n(W) < \infty$ and $\lambda > 1$. For $r > 0$ we define

$$W_r = \{(a, u) \in W : \mathbf{r}_K^\phi(a, u) \geq \lambda r\},$$

that is an \mathcal{H}^n -measurable subset of W . We denote with W_r^* the set of all $(a, u) \in W_r$ such that $\text{Tan}^n(\mathcal{H}^n \llcorner W, (a, u))$ is n -dimensional plane and $\text{Tan}(\mathcal{H}^n \llcorner W, (a, u)) = \text{Tan}^n(\mathcal{H}^n \llcorner W_r, (a, u))$. It follows from [Fed69, 3.2.19] that $\mathcal{H}^n(W_r \setminus W_r^*) = 0$. Moreover we observe from coarea formula that there exists $J \subseteq \{t : t > 0\}$ with $\mathcal{H}^1(J) = 0$ such that $\mathcal{H}^n(S^\phi(K, r) \setminus \text{Unp}^\phi(K)) = 0$ for every $r \notin J$.

We fix $r > 0$, $r \notin J$, and we define

$$M_r = \{a + ru : (a, u) \in W_r\}.$$

Notice that $M_r \subseteq S^\phi(K, r) \cap \{x : \boldsymbol{\rho}_K^\phi(x) \geq \lambda\}$ and M_r is \mathcal{H}^n -measurable. By Theorem 2.2 the function $\boldsymbol{\psi}_K^\phi|_{M_r}$ is Lipschitz; moreover we notice that $\boldsymbol{\psi}_K^\phi(M_r) = W_r$ and $(\boldsymbol{\psi}_K^\phi|_{M_r})^{-1}(a, u) = a + ru$ for $(a, u) \in W_r$. We denote with M_r^* the set of all $x \in M_r$ such that $\text{Tan}(S^\phi(K, r), x)$ is an n -dimensional plane and $\text{Tan}^n(\mathcal{H}^n \llcorner M_r, x) = \text{Tan}(S^\phi(K, r), x)$. It follows from Remark 3.1 and [Fed69, 3.2.19] that $\mathcal{H}^n(M_r \setminus M_r^*) = 0$. We conclude that

$$\mathcal{H}^n(W_r \setminus (W_r^* \cap \boldsymbol{\psi}_K^\phi(M_r^*))) = 0.$$

Moreover if $(a, u) \in (W_r^* \cap \boldsymbol{\psi}_K^\phi(M_r^*)) \cap \tilde{N}^\phi(K)$ it follows from [San20a, Lemma B.2] and Remark 3.2 that $\{\tau_1(a, u), \dots, \tau_n(a, u)\}$ is a basis of $\text{Tan}(S^\phi(K, r), a + ru)$,

$$D\boldsymbol{\psi}_K^\phi(a + ru)[\text{Tan}(S^\phi(K, r), a + ru)] = \text{Tan}^n(\mathcal{H}^n \llcorner W, (a, u))$$

and

$$D\boldsymbol{\psi}_K^\phi(a + ru)(\tau_i(a, u)) = \begin{cases} \frac{1}{1+r\kappa_{K,i}^\phi(a, u)}\zeta_i(a, u) & \text{if } \kappa_{K,i}^\phi(a, u) < \infty \\ \frac{1}{r}\zeta_i(a, u) & \text{if } \kappa_{K,i}^\phi(a, u) = +\infty. \end{cases}$$

This proves that $\{\zeta_1(a, u), \dots, \zeta_n(a, u)\}$ is a basis of $\text{Tan}^n(\mathcal{H}^n \llcorner W, (a, u))$ for \mathcal{H}^n a.e. $(a, u) \in W_r$ and for every $r \notin J$.

Since $W = \bigcup_{r>0} W_r$ and $W_r \subseteq W_s$ for $0 < s < r$, there exists a sequence $r_i \searrow 0$, $r_i \notin J$, so that $W = \bigcup_{i=1}^\infty W_{r_i}$. Henceforth, the proof is complete. \square

3.11 Definition. Let $K \subseteq \mathbf{R}^{n+1}$ be closed. For every $d = 0, \dots, n$ we define

$$\tilde{N}_d^\phi(K) = \{(a, u) \in \tilde{N}^\phi(K) : \kappa_{K,d}^\phi(a, u) < \infty, \kappa_{K,d+1}^\phi(a, u) = \infty\}.$$

Moreover for every $j = 0, \dots, n$ we define $E_{K,j}^\phi : \tilde{N}^\phi(K) \rightarrow \mathbf{R}$ by

$$E_{K,j}^\phi(a, u) = \begin{cases} \sum_{1 \leq l_1 < \dots < l_j \leq d} \prod_{h=1}^j \kappa_{K,l_h}^\phi & \text{if } (a, u) \in \tilde{N}_d^\phi(K) \text{ and } d \geq j \\ 0 & \text{if } (a, u) \in \tilde{N}_d^\phi(K) \text{ and } d < j, \end{cases}$$

where for $j = 0$ this means $E_{K,0}^\phi \equiv 1$. Finally for every $r = 0, \dots, n$ we define r -th ϕ -mean curvature of K as

$$\mathbf{H}_{K,r}^\phi = \sum_{j=0}^r E_{K,j}^\phi \mathbf{1}_{\tilde{N}_{j+n-r}^\phi(K)}.$$

We can now prove a general Steiner formula for arbitrary closed sets.

3.12 Theorem. Suppose $K \subseteq \mathbf{R}^{n+1}$ is closed, $\tau_1, \dots, \tau_n, \zeta_1, \dots, \zeta_n$ are $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable functions satisfying the hypothesis in Lemma 3.10 and J is the $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable function defined on \mathcal{H}^n almost all of $N^\phi(K)$ by

$$J(a, u) = \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K).$$

Then

$$\begin{aligned} & \int_{\{x:0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} \\ &= \sum_{m=0}^n \frac{1}{n-m+1} \int_{N^\phi(K)} \inf\{\rho, \mathbf{r}_K^\phi(a, u)\}^{n+1-m} \phi(\mathbf{n}^\phi(u)) J(a, u) \mathbf{H}_{K,n-m}^\phi(a, u) \varphi(a, u) d\mathcal{H}^n(a, u) \end{aligned}$$

for every bounded Borel function $\varphi : N^\phi(K) \rightarrow \mathbf{R}$ and for every $\rho > 0$.

Proof. We define $\Omega = \{(a, u, t) : (a, u) \in N^\phi(K), 0 < t < \mathbf{r}_K^\phi(a, u)\}$ and the bijective map

$$f : \Omega \rightarrow \mathbf{R}^{n+1} \setminus (K \cup \text{Cut}^\phi(K))$$

by $f(a, u, t) = a + tu$ for $(a, u, t) \in \Omega$. We choose a countable family $\{N_i : i \geq 1\}$ of Borel subsets of $N^\phi(K)$ such that $\mathcal{H}^n(N_i) < \infty$ and $N_i \subseteq N_{i+1}$ for every $i \geq 1$ and $N^\phi(K) = \bigcup_{i=1}^\infty N_i$ (recall that $N^\phi(K)$ is a Borel and countably n -rectifiable subset of $\mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$). We define $\Omega_i = \Omega \cap (N_i \times \mathbf{R})$ for every $i \geq 1$ and we notice that Ω_i is a Borel $(n+1)$ -rectifiable subset of $\mathbf{R}^{n+1} \times \mathbf{R}^{n+1} \times \mathbf{R}$. Let $\varphi : N^\phi(K) \rightarrow \mathbf{R}$ be a bounded Borel function. Recalling that $\mathcal{L}^{n+1}(\text{Cut}^\phi(K)) = 0$, we use the dominated convergence theorem and the coarea formula [Fed69, 3.2.22] to compute

$$\begin{aligned} \int_{\{x:0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} &= \lim_{i \rightarrow \infty} \int_{f(\Omega_i) \cap \{x:0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} \\ &= \lim_{i \rightarrow \infty} \int_{f[\Omega_i \cap \{(a, u, t): 0 < t \leq \rho\}]} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} \\ &= \lim_{i \rightarrow \infty} \int_{\Omega_i \cap \{(a, u, t): 0 < t \leq \rho\}} \varphi(a, u) \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) d\mathcal{H}^{n+1}(a, u) \\ &= \lim_{i \rightarrow \infty} \int_{N_i} \int_0^{\inf\{\rho, \mathbf{r}_K^\phi(a, u)\}} \varphi(a, u) \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) dt d\mathcal{H}^n(a, u). \end{aligned}$$

Noting that

$$\text{Tan}^{n+1}(\mathcal{H}^{n+1} \llcorner \Omega_i, (a, u, t)) = \text{Tan}^n(\mathcal{H}^n \llcorner N_i, (a, u)) \times \mathbf{R} \quad \text{for } (a, u, t) \in \Omega_i,$$

we can apply Lemma 3.10 to compute

$$\begin{aligned}
& \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= \frac{|\mathbf{D}f(a, u, t)(\zeta_1(a, u), 0) \wedge \dots \wedge \mathbf{D}f(a, u, t)(\zeta_n(a, u), 0) \wedge \mathbf{D}f(a, u, t)(0, 1)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u) \wedge u|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} t^{n-d} \left(\prod_{j=1}^d (1 + t\kappa_{K,j}^\phi(a, u)) \right) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= (\mathbf{n}^\phi(u) \bullet u) \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u) \wedge \mathbf{n}^\phi(u)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} t^{n-d} \left(\prod_{j=1}^d (1 + t\kappa_{K,j}^\phi(a, u)) \right) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= \phi(\mathbf{n}^\phi(u)) J(a, u) t^{n-d} \left(\prod_{j=1}^d (1 + t\kappa_{K,j}^\phi(a, u)) \right) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= \phi(\mathbf{n}^\phi(u)) J(a, u) \sum_{j=0}^d t^{n-d+j} E_{K,j}^\phi(a, u) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u)
\end{aligned}$$

for \mathcal{H}^{n+1} a.e. $(a, u, t) \in \tilde{\Omega}_i$ and $d = 0, \dots, n$. Consequently

$$\begin{aligned}
& \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) \\
&= \sum_{d=0}^n \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) \mathbf{1}_{\tilde{N}_d^\phi(K)} \\
&= \phi(\mathbf{n}^\phi(u)) J(a, u) \sum_{d=0}^n \sum_{j=0}^d t^{n-d+j} E_{K,j}^\phi(a, u) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) \\
&= \phi(\mathbf{n}^\phi(u)) J(a, u) \sum_{m=0}^n t^{n-m} \mathbf{H}_{K,n-m}^\phi(a, u)
\end{aligned}$$

for \mathcal{H}^{n+1} a.e. $(a, u, t) \in \tilde{\Omega}_i$ and we conclude

$$\begin{aligned}
& \int_{\{x: 0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} \\
&= \sum_{m=0}^n \frac{1}{n-m+1} \int_{N^\phi(K)} \inf\{\rho, \mathbf{r}_K^\phi(a, u)\}^{n-m+1} \phi(\mathbf{n}^\phi(u)) J(a, u) \mathbf{H}_{K,n-m}^\phi(a, u) \varphi(a, u) d\mathcal{H}^n(a, u).
\end{aligned}$$

□

3.13 Remark. By Remark 3.9 we have that $1 + t\kappa_{K,i}^\phi(a, u) > 0$ for every $(a, u) \in \tilde{N}_n^\phi(K)$ and for every $0 < t < \mathbf{r}^\phi(K)$. Therefore the proof of Theorem 3.12 provides the equality

$$J(a, u) \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u) = \text{ap } J_{n+1}^{\Omega_i} f(a, u, t) \phi(\mathbf{n}^\phi)^{-1} t^{d-n} \left(\prod_{j=1}^d (1 + t\kappa_{K,j}^\phi(a, u)) \right)^{-1} \mathbf{1}_{\tilde{N}_d^\phi(K)}(a, u)$$

for every $i \geq 1$ and for \mathcal{H}^{n+1} a.e. $(a, u, t) \in \tilde{\Omega}_i$. In particular, if $\tau'_1, \dots, \tau'_n, \zeta'_1, \dots, \zeta'_n$ is another set of $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable functions satisfying the hypothesis of Lemma 3.10, then

$$\frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} = \frac{|\tau'_1(a, u) \wedge \dots \wedge \tau'_n(a, u)|}{|\zeta'_1(a, u) \wedge \dots \wedge \zeta'_n(a, u)|}$$

for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$.

In view of Remark 3.13 it is convenient to introduce the following function.

3.14 Definition. For every closed set $K \subseteq \mathbf{R}^{n+1}$, we denote with J_K^ϕ any $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable function defined on \mathcal{H}^n almost all of $N^\phi(K)$ such that

$$J_K^\phi(a, u) = \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u)|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K),$$

where $\tau_1, \dots, \tau_n, \zeta_1, \dots, \zeta_n$ are $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable functions satisfying the hypothesis in Lemma 3.10

3.15 Remark. The proof of Theorem 3.12 provides also the formula

$$\begin{aligned} & \int_{\{x: 0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} \\ &= \sum_{d=0}^n \int_{\tilde{N}_d^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \int_0^{\inf\{\rho, r_K^\phi(a, u)\}} t^{n-d} \left(\prod_{j=1}^d (1 + t \kappa_{K,j}^\phi(a, u)) \right) dt d\mathcal{H}^n(a, u) \end{aligned}$$

for every $\rho > 0$, that will be used in the proof of Theorem 5.2.

The following Corollary, besides of being of independent interest, plays a key role in the analysis of the equality case in Theorem 5.2.

3.16 Corollary. *Suppose $K \subseteq \mathbf{R}^{n+1}$ be a closed set, $s > 0$ and $r_K^\phi(a, u) \geq s$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$. Then $\{x \in \mathbf{R}^{n+1} : \delta_K^\phi(x) < s\} \subseteq \text{Unp}^\phi(K)$.*

Proof. It follows from Theorem 3.12 that

$$\int_{\{x: 0 < \delta_K^\phi(x) \leq \rho\}} (\varphi \circ \psi_K^\phi) d\mathcal{L}^{n+1} = \sum_{m=0}^{n+1} \rho^{n+1-m} I_m(\varphi) \quad \text{for } 0 < \rho < s,$$

where $I_m(\varphi) = \frac{1}{n-m+1} \int_{N^\phi(K)} \phi(\mathbf{n}^\phi) J_K^\phi \mathbf{H}_{K, n-m}^\phi \varphi d\mathcal{H}^n$ for $m = 0, \dots, n$. We conclude from [DRKS20, Theorem 5.9] that $\{x \in \mathbf{R}^{n+1} : \delta_K^\phi(x) < s\} \subseteq \text{Unp}^\phi(K)$. \square

We conclude with the following Lemma, that will be useful later.

3.17 Lemma. *For every closed set $K \subseteq \mathbf{R}^{n+1}$,*

$$\mathcal{H}^0(N^\phi(K, a)) \in \{1, 2\} \quad \text{for every } a \in \mathbf{p}(\tilde{N}_n^\phi(K))$$

and

$$\mathcal{H}^n(\mathbf{p}[N^\phi(K) \setminus \tilde{N}_n^\phi(K)]) = 0.$$

Proof. Let $(a, u) \in \tilde{N}_n^\phi(K)$ and $0 < r < r_K^\phi(a, u)$. Then $1 - r \chi_{K,i}^\phi(a + ru) > 0$ for every $i = 1, \dots, n$ and, since these numbers are the eigenvalues of $D \xi_K^\phi(a + ru) | \text{Tan}(S^\phi(K, r), a + ru)$, we conclude (noting Remark 3.2)

$$\text{Tan}(\mathcal{W}^\phi, u) = D \xi_K^\phi(a + ru) [\text{Tan}(S^\phi(K, r), a + ru)] \subseteq \text{Tan}(K, a).$$

Since $N(K, a) \subseteq \text{Nor}(K, a) \subseteq \text{Nor}(\mathcal{W}^\phi, u)$ and $\dim \text{Nor}(\mathcal{W}^\phi, u) = 1$, it follows that $\mathcal{H}^0(N^\phi(K, a)) \in \{1, 2\}$. Moreover combining Lemma 3.10 with coarea formula [Fed69, 3.2.22] we obtain

$$\begin{aligned} 0 &= \int_{W \setminus \tilde{N}_n^\phi(K)} \text{ap } J_n^W \mathbf{p}(a, u) d\mathcal{H}^n(W) = \int_{\mathbf{p}[W \setminus \tilde{N}_n^\phi(K)]} \mathcal{H}^0(N^\phi(K, x)) d\mathcal{H}^n(x) \\ & \quad \mathcal{H}^n(\mathbf{p}[W \setminus \tilde{N}_n^\phi(K)]) = 0 \end{aligned}$$

for every \mathcal{H}^n -measurable subset $W \subseteq N^\phi(K)$ with $\mathcal{H}^n(W) < \infty$. \square

3.18 Remark. In particular it follows from the previous Remark that if $K \subseteq \mathbf{R}^{n+1}$ is a convex body, then $\mathcal{H}^0(N^\phi(K, a)) = 1$ for each $a \in \mathbf{p}(\tilde{N}_n^\phi(K))$.

4 Anisotropic curvature measures for convex sets

If $K \subseteq \mathbf{R}^{n+1}$ is a closed convex set, then $r_K^\phi \equiv +\infty$ and $\kappa_{K,i}^\phi \geq 0$ for every $i = 1, \dots, n$ by (3.9). Moreover $N^\phi(K)$ is an n -dimensional compact Lipschitz manifold and $\text{Cut}^\phi(K) = \emptyset$. We introduce now the anisotropic curvature measures of a convex set.

4.1 Definition. Let $K \subseteq \mathbf{R}^{n+1}$ be a closed convex set and $m = 0, \dots, n$. The m -th support measure of K with respect to ϕ is the measure $\tilde{\mathcal{C}}_m^\phi(K, \cdot)$ over $\mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$ defined by (see [Fed69, 2.4] for the definition of the upper integral \int^*)

$$\tilde{\mathcal{C}}_m^\phi(K, B) = \frac{1}{n-m+1} \int_{B \cap N^\phi(K)}^* \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \mathbf{H}_{K, n-m}^\phi(a, u) d\mathcal{H}^n(a, u)$$

for every $B \subseteq \mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$ and the m -th curvature measure of K with respect to ϕ is the measure over \mathbf{R}^{n+1} given by (see [Fed69, 2.1.2])

$$\mathcal{C}_m^\phi(K, \cdot) = \mathbf{p}_\# \tilde{\mathcal{C}}_m^\phi(K, \cdot).$$

We also write $\mathcal{C}_m^\phi(K) = \mathcal{C}_m^\phi(K, \mathbf{R}^{n+1})$ (this is the anisotropic m -th mixed volume of K).

4.2 Remark. The measure $\tilde{\mathcal{C}}_m^\phi(K, \cdot)$ is a Radon measure over $\mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$ (see [Fed69, 2.2.5]). Therefore it follows from [Fed69, 2.2.17] that $\mathcal{C}_m^\phi(K, \cdot)$ is a Radon measure over \mathbf{R}^{n+1} .

We can now state the *local anisotropic Steiner formula for convex sets*, which readily follows from Theorem 3.12.

4.3 Corollary. Let $K \subseteq \mathbf{R}^{n+1}$ be a closed convex set. Then

$$\mathcal{L}^{n+1}(\{x \in \mathbf{R}^{n+1} : 0 < \delta_K^\phi(x) \leq \rho, \boldsymbol{\xi}_K^\phi(x) \in B\}) = \sum_{m=0}^n \rho^{n+1-m} \mathcal{C}_m^\phi(K, B)$$

for every Borel subset $B \subseteq \partial K$ and for every $\rho > 0$.

4.4 Lemma. Suppose $K \subseteq \mathbf{R}^{n+1}$ is a convex body and η is its (measure theoretic) exterior unit normal. Then

$$\mathcal{C}_n^\phi(K, B) = \int_B \phi(\eta(x)) d\mathcal{H}^n(x)$$

for every Borel set $B \subseteq \partial K$.

Proof. Noting that $\mathbf{H}_{K,0}^\phi = \mathbf{1}_{\tilde{N}_n^\phi(K)}$, Lemma 3.10 implies that

$$\text{ap } J_n^{N^\phi(K)} \mathbf{p}(a, u) = J_K^\phi(a, u) \cdot \mathbf{H}_{K,0}^\phi(a, u) \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K).$$

Therefore, noting that $N^\phi(K, x) = \{\nabla \phi(\eta(x))\}$ for \mathcal{H}^n a.e. $x \in \partial K$ by (4), we apply Coarea formula [Fed69, 3.2.22] to conclude

$$\begin{aligned} \mathcal{C}_n^\phi(K, B) &= \int_{N^\phi(K)|_B} \text{ap } J_n^{N^\phi(K)} \mathbf{p}(a, u) \phi(\mathbf{n}^\phi(u)) d\mathcal{H}^n(a, u) \\ &= \int_{B \cap \partial K} \int_{N^\phi(K, x)} \phi(\mathbf{n}^\phi(u)) d\mathcal{H}^0(u) d\mathcal{H}^n(x) \\ &= \int_{B \cap \partial K} \phi(\eta(x)) d\mathcal{H}^n(x) \end{aligned}$$

□

We now prove the anisotropic Minkowski formulae for arbitrary convex bodies.

4.5 Theorem. *If $K \subseteq \mathbf{R}^{n+1}$ be a convex body and $r = 1, \dots, n$ then*

$$\begin{aligned} & (n-r+1) \int_{N^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \mathbf{H}_{K, r-1}^\phi(a, u) d\mathcal{H}^n(a, u) \\ &= r \int_{N^\phi(K)} [a \bullet \mathbf{n}^\phi(u)] J_K^\phi(a, u) \mathbf{H}_{K, r}^\phi(a, u) d\mathcal{H}^n(a, u). \end{aligned}$$

Proof. We set $\eta(x) = \frac{\nabla \delta_K^\phi(x)}{|\nabla \delta_K^\phi(x)|}$ for $x \in \mathbf{R}^{n+1} \setminus K$ and

$$B_\rho(K) = \{x \in \mathbf{R}^{n+1} : 0 \leq \delta_K^\phi(x) \leq \rho\} \quad \text{for } 0 < \rho < \infty.$$

We notice that $B_\rho(K)$ is a convex body with $\mathcal{C}^{1,1}$ -boundary $\partial B_\rho(K) = S^\phi(K, \rho)$ and $\eta|_{\partial B_\rho(K)}$ is its exterior unit normal. Moreover for each $\rho > 0$ the map $f_\rho : N^\phi(K) \rightarrow S^\phi(K, \rho)$, defined as

$$f_\rho(a, u) = a + \rho u \quad \text{for } (a, u) \in N^\phi(K),$$

is a bi-lipschitz homeomorphism by Theorem 2.2. We observe (see proof of Theorem 3.12) that

$$J_n^{N^\phi(K)} f_\rho(a, u) = J_K^\phi(a, u) \sum_{m=0}^n \rho^{n-m} \mathbf{H}_{K, n-m}^\phi(a, u)$$

for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$. We set

$$I_m(K) = \int_{N^\phi(K)} a \bullet \mathbf{n}^\phi(u) J_K^\phi(a, u) \mathbf{H}_{K, n-m}^\phi(a, u) d\mathcal{H}^n(a, u) \quad \text{for } m = 0, \dots, n$$

and we compute

$$\begin{aligned} (n+1)\mathcal{L}^{n+1}(B_\rho(K)) &= \int_{S^\phi(K, \rho)} x \bullet \eta(x) d\mathcal{H}^n(x) \\ &= \int_{N^\phi(K)} [(a + \rho u) \bullet \eta(a + \rho u)] J_n^{N^\phi(K)} f_\rho(a, u) d\mathcal{H}^n(a, u) \\ &= \sum_{m=0}^n \rho^{n-m} I_m(K) + \sum_{m=0}^n (n-m+1) \rho^{n-m+1} \mathcal{C}_m^\phi(K). \end{aligned}$$

Employing the Steiner formula 4.3 we get

$$(n+1)\mathcal{L}^{n+1}(B_\rho(K)) = (n+1)\mathcal{L}^{n+1}(K) + (n+1) \sum_{m=0}^n \rho^{n-m+1} \mathcal{C}_m^\phi(K)$$

and we infer

$$\sum_{m=0}^{n-1} [I_m(K) - (m+1)\mathcal{C}_{m+1}^\phi(K)] \rho^{n-m} + I_n(K) - (n+1)\mathcal{L}^{n+1}(K) = 0$$

for every $\rho > 0$. It follows that $I_m(K) = (m+1)\mathcal{C}_{m+1}^\phi(K)$ for $m = 0, \dots, n-1$ and $I_n(K) = (n+1)\mathcal{L}^{n+1}(K)$. \square

4.6 Lemma. *Suppose $K \subseteq \mathbf{R}^{n+1}$ is a convex body, $r = 1, \dots, n$, $\lambda > 0$ and $\mathcal{C}_{n-r}^\phi(K, \cdot) = \lambda \mathcal{C}_n^\phi(K, \cdot)$.*

Then $\mathbf{H}_{K, r}^\phi(a, u) = 0$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K) \setminus \tilde{N}_n^\phi(K)$ and

$$\mathbf{H}_{K, r}^\phi(a, u) = (r+1)\lambda \geq \left(\frac{\mathcal{C}_n^\phi(K)}{(n+1)\mathcal{L}^{n+1}(K)} \right)^r \binom{n}{r} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in \tilde{N}_n^\phi(K).$$

Proof. We notice from Remark 3.18 that $\mathcal{H}^0(N^\phi(K, a)) = 1$ for each $a \in \mathbf{p}[\tilde{N}_n^\phi(K)]$ and

$$\mathbf{p}[N^\phi(K) \setminus \tilde{N}_n^\phi(K)] \cap \mathbf{p}[\tilde{N}_n^\phi(K)] = \emptyset.$$

From the equality $\mathcal{C}_{n-r}^\phi(K, \cdot) = \lambda \mathcal{C}_n^\phi(K, \cdot)$ we get

$$\begin{aligned} & \int_{\tilde{N}_n^\phi(K)|B}^* \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \left(\frac{1}{r+1} \mathbf{H}_{K,r}^\phi(a, u) - \lambda \right) d\mathcal{H}^n(a, u) \\ & + \frac{1}{r+1} \int_{(N^\phi(K) \setminus \tilde{N}_n^\phi(K))|B}^* \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \mathbf{H}_{K,r}^\phi(a, u) d\mathcal{H}^n(a, u) = 0 \end{aligned}$$

for every subset $B \subseteq \partial K$. Choosing $B = \mathbf{p}[N^\phi(K) \setminus \tilde{N}_n^\phi(K)]$ in the last equation, we infer that $\mathbf{H}_{K,r}^\phi(a, u) = 0$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K) \setminus \tilde{N}_n^\phi(K)$. Moreover,

$$\int_{\tilde{N}_n^\phi(K)|B}^* \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \left(\frac{1}{r+1} \mathbf{H}_{K,r}^\phi(a, u) - \lambda \right) d\mathcal{H}^n(a, u) = 0$$

for every $B \subseteq \mathbf{p}[\tilde{N}_n^\phi(K)]$ and we infer that $\frac{1}{r+1} \mathbf{H}_{K,r}^\phi(a, u) = \lambda$ for \mathcal{H}^n a.e. $(a, u) \in \tilde{N}_n^\phi(K)$.

Let η be the (measure theoretic) exterior unit normal of K . Then, using Lemma 3.10 and coarea formula, we observe

$$\begin{aligned} & \int_{N^\phi(K)} [a \bullet \mathbf{n}^\phi(u)] J_K^\phi(a, u) \mathbf{H}_{K,r}^\phi(a, u) d\mathcal{H}^n(a, u) \\ & = \int_{\tilde{N}_n^\phi(K)} [a \bullet \mathbf{n}^\phi(u)] J_K^\phi(a, u) \mathbf{H}_{K,r}^\phi(a, u) d\mathcal{H}^n(a, u) \\ & = (r+1)\lambda \int_{\tilde{N}_n^\phi(K)} [a \bullet \mathbf{n}^\phi(u)] J_K^\phi(a, u) d\mathcal{H}^n(a, u) \\ & = (r+1)\lambda \int_{N^\phi(K)} [a \bullet \mathbf{n}^\phi(u)] \text{ap } J_n^{N^\phi(K)} \mathbf{p}(a, u) d\mathcal{H}^n(a, u) \\ & = (r+1)\lambda \int_{\partial K} \int_{N^\phi(K, a)} [a \bullet \mathbf{n}^\phi(u)] d\mathcal{H}^0 u d\mathcal{H}^n a \\ & = (r+1)\lambda \int_{\partial K} [a \bullet \eta(a)] d\mathcal{H}^n a \\ & = (r+1)\lambda(n+1)\mathcal{L}^{n+1}(K). \end{aligned}$$

Moreover, employing the Newton-McLaurin inequality [Nic00, Theorem 1.1], we obtain

$$\begin{aligned} & \int_{N^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \mathbf{H}_{K,r-1}^\phi(a, u) d\mathcal{H}^n(a, u) \\ & \geq \int_{\tilde{N}_n^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \mathbf{H}_{K,r-1}^\phi(a, u) d\mathcal{H}^n(a, u) \\ & \geq [(r+1)\lambda]^{\frac{r-1}{r}} \binom{n}{r}^{\frac{1-r}{r}} \binom{n}{r-1} \int_{\tilde{N}_n^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) d\mathcal{H}^n(a, u) \\ & = [(r+1)\lambda]^{\frac{r-1}{r}} \binom{n}{r}^{\frac{1-r}{r}} \binom{n}{r-1} \mathcal{C}_n^\phi(K). \end{aligned}$$

We now use the Minkowski Formula in 4.5 and, noting that $(n-r+1) \binom{n}{r}^{-1} \binom{n}{r-1}^{\frac{1}{r}} = 1$, we conclude

$$[\lambda(r+1)]^{\frac{1}{r}} \geq \frac{\mathcal{C}_n^\phi(K)}{(n+1)\mathcal{L}^{n+1}(K)} \binom{n}{r}^{\frac{1}{r}}.$$

□

5 An optimal geometric inequality

In this section we assume that ϕ is a uniformly convex \mathcal{C}^2 norm.

5.1 Lemma. *Suppose $C \subseteq \mathbf{R}^{n+1}$ is a convex body, K is the closure of $\mathbf{R}^{n+1} \setminus C$ and*

$$\iota : \mathbf{R}^{n+1} \times \mathbf{R}^{n+1} \rightarrow \mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$$

is the linear map defined as $\iota(a, u) = (a, -u)$ for every $(a, u) \in \mathbf{R}^{n+1} \times \mathbf{R}^{n+1}$.

Then the following statements hold.

(a) $\mathcal{H}^0(N^\phi(K, a)) = 1$ and $N^\phi(K, a) = -N^\phi(C, a)$ for every $a \in \mathbf{p}(N^\phi(K))$.

(b) $\mathcal{H}^n(N^\phi(K) \setminus \tilde{N}_n^\phi(K)) = 0$.

(c) $\kappa_{K,i}^\phi(a, u) = -\kappa_{C,n+1-i}^\phi(a, -u)$ for \mathcal{H}^n a.e. $(a, u) \in \tilde{N}_n^\phi(K)$ and $i = 1, \dots, n$.

(d) $J_K^\phi(a, u) = \text{ap } J_n^{N^\phi(K)} \iota(a, u) J_C^\phi(a, -u)$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$.

Proof. The statement in (a) is contained in [San20b, Lemma 5.1] and the statement in (b) follows from [San20b, Lemma 5.1] noting that $\mathcal{H}^n(\mathbf{p}[N^\phi(K) \setminus \tilde{N}_n^\phi(K)]) = 0$ by Lemma 3.17.

We prove (c). Firstly, we prove that

$$\nu_K^\phi(y) = \{-\nu_C^\phi((1+\lambda)a - \lambda y) : a \in \xi_K^\phi(y)\} \quad \text{for every } y \in \mathbf{R}^{n+1} \setminus K \text{ and } \lambda > 0. \quad (7)$$

The equality in (7) follows noting that if $y \in \mathbf{R}^{n+1} \setminus K$, $\lambda > 0$ and $a \in \xi_K^\phi(y)$, then

$$a + \lambda \delta_K^\phi(y) \frac{a-y}{\delta_K^\phi(y)} = (1+\lambda)a - \lambda y,$$

$$N^\phi(K, a) = \left\{ \frac{y-a}{\delta_K^\phi(y)} \right\}, \quad N^\phi(C, a) = \left\{ \frac{a-y}{\delta_K^\phi(y)} \right\},$$

$$\nu_C^\phi((1+\lambda)a - \lambda y) = \frac{a-y}{\delta_K^\phi(y)}, \quad -\nu_C^\phi((1+\lambda)a - \lambda y) \in \nu_K^\phi(y).$$

Define $S = \mathbf{p}(\iota(N^\phi(K)) \setminus \tilde{N}^\phi(C))$ and notice that $\mathcal{H}^n(S) = 0$ by Remark 3.8. It follows from [San20b, Lemma 5.1] that $\mathcal{H}^n(N^\phi(K)|S) = 0$. Fix now $(a, u) \in \tilde{N}_n^\phi(K)$ with $a \notin S$, $0 < r < r_K^\phi(a, u)$ and, noting that $1 - r\chi_{K,i}^\phi(a+ru) > 0$ for every $i = 1, \dots, n$, we select $\lambda > 0$ so that

$$\chi_{K,i}^\phi(a+ru) < \frac{1}{(1+\lambda)r} \quad \text{for } i = 1, \dots, n.$$

Since $a \notin S$, then $(a, -u) \in \tilde{N}^\phi(C)$ and ν_C^ϕ is differentiable at $a - tu$ for every $t > 0$. Differentiating at $a + ru$ the equality in (7), we compute

$$D\nu_K^\phi(a+ru) = -D\nu_C^\phi(a-\lambda ru) \circ ((1+\lambda)D\xi_K^\phi(a+ru) - \mathbf{I}_{\mathbf{R}^{n+1}}).$$

If τ_1, \dots, τ_n form a basis of $\text{Tan}(\mathcal{W}^\phi, u)$ such that $D\nu_K^\phi(a+ru)(\tau_i) = \chi_{K,i}^\phi(a+ru)\tau_i$ for every $i = 1, \dots, n$, then we infer

$$D\nu_C^\phi(a-\lambda ru)(\tau_i) = \frac{\chi_{K,i}^\phi(a+ru)}{(1+\lambda)r\chi_{K,i}^\phi(a+ru) - 1} \tau_i \quad \text{for } i = 1, \dots, n$$

and, noting that $\text{Tan}(\mathcal{W}_1^\phi, -u) = \text{Tan}(\mathcal{W}_1^\phi, u)$ and $(1+\lambda)r\chi_{K,i}^\phi(a+ru) - 1 < 0$ for every $i = 1, \dots, n$, we conclude that

$$\chi_{C,n+1-i}^\phi(a-\lambda ru) = \frac{\chi_{K,i}^\phi(a+ru)}{(1+\lambda)r\chi_{K,i}^\phi(a+ru) - 1}$$

for every $i = 1, \dots, n$. Therefore,

$$\kappa_{C, n+1-i}^\phi(a, -u) = \frac{\chi_{C, n+1-i}^\phi(a - \lambda ru)}{1 - \lambda r \chi_{C, n+1-i}^\phi(a - \lambda ru)} = \frac{\chi_{K, i}^\phi(a + ru)}{r \chi_{K, i}^\phi(a + ru) - 1} = -\kappa_{K, i}^\phi(a, u)$$

for every $i = 1, \dots, n$.

Finally we prove (d). Let $\tau_1, \dots, \tau_n, \zeta_1, \dots, \zeta_n$ be $\mathcal{H}^n \llcorner N^\phi(K)$ -measurable functions satisfying the hypothesis of Lemma 3.10. The argument of the previous paragraph in combination with Lemma 3.6 shows that

$$\mathbf{D} \nu_C^\phi(a - tu)(\tau_i(a, u)) = \chi_{C, n+1-i}^\phi(a - tu) \tau_i(a, u)$$

for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$ and for every $t > 0$. Since $\kappa_{C, i}^\phi(a, -u) < \infty$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$ by (b) and (c), we infer that

$$J_C^\phi(a, -u) = \frac{|\tau_1(a, u) \wedge \dots \wedge \tau_n(a, u)|}{|\iota(\zeta_1(a, u)) \wedge \dots \wedge \iota(\zeta_n(a, u))|}$$

for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$. Since

$$\text{ap } J_n^{N^\phi(K)} \iota(a, u) = \frac{|\iota(\zeta_1(a, u)) \wedge \dots \wedge \iota(\zeta_n(a, u))|}{|\zeta_1(a, u) \wedge \dots \wedge \zeta_n(a, u)|},$$

the equation in (d) follows. \square

5.2 Theorem. *Suppose $C \subseteq \mathbf{R}^{n+1}$ is a convex body. Then*

$$(n+1)\mathcal{L}^{n+1}(C) \leq n \int_{\tilde{N}_n^\phi(C)} J_C^\phi(a, u) \frac{\phi(\mathbf{n}^\phi(u))}{\mathbf{H}_{C,1}^\phi(a, u)} d\mathcal{H}^n(a, u).$$

Moreover, if the equality holds and there exists $q < \infty$ so that $\mathbf{H}_{C,1}^\phi(a, u) \leq q$ for \mathcal{H}^n a.e. $(a, u) \in \tilde{N}_n^\phi(C)$, then $\partial C = a + s\mathcal{W}^\phi$ for some $a \in \mathbf{R}^{n+1}$ and $s > 0$.

Proof. We assume that $\mathbf{H}_{C,1}^\phi(a, u) > 0$ for \mathcal{H}^n a.e. $(a, u) \in \tilde{N}_n^\phi(C)$, otherwise the inequality is trivially true. Let K be the closure of $\mathbf{R}^{n+1} \setminus C$ and notice by Lemma 5.1 that

$$\begin{aligned} \mathbf{H}_{K,1}^\phi(a, u) &= \sum_{i=1}^n \kappa_{K,i}^\phi(a, u) \\ &= -\sum_{i=1}^n \kappa_{C, n+1-i}^\phi(a, -u) = -\mathbf{H}_{C,1}^\phi(a, -u) < 0 \end{aligned}$$

for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$. We use the inequality in Remark 3.9 to infer that

$$\mathbf{r}_K^\phi(a, u) \leq -\frac{n}{\mathbf{H}_{K,1}^\phi(a, u)} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K)$$

and $1 + t\kappa_{K,i}^\phi(a, u) > 0$ for \mathcal{H}^n a.e. $(a, u) \in N^\phi(K)$ and for every $0 < t < \mathbf{r}_K^\phi(a, u)$. Noting that $\mathbf{n}^\phi(u) = -\mathbf{n}^\phi(-u)$ for every $u \in \mathcal{W}^\phi$, we can use Lemma 5.1, the formula in Remark 3.15 and the classical arithmetic-geometric mean inequality to estimate

$$\begin{aligned} \mathcal{L}^{n+1}(C) &= \int_{N^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \int_0^{\mathbf{r}_K^\phi(a, u)} \prod_{j=1}^n (1 + t\kappa_{K,j}^\phi(a, u)) dt d\mathcal{H}^n(a, u) \\ &\leq \int_{N^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \int_0^{\mathbf{r}_K^\phi(a, u)} \left(1 + \frac{t}{n} \mathbf{H}_{K,1}^\phi(a, u)\right)^n dt d\mathcal{H}^n(a, u) \\ &\leq \int_{N^\phi(K)} \phi(\mathbf{n}^\phi(u)) J_K^\phi(a, u) \int_0^{-\frac{n}{\mathbf{H}_{K,1}^\phi(a, u)}} \left(1 + \frac{t}{n} \mathbf{H}_{K,1}^\phi(a, u)\right)^n dt d\mathcal{H}^n(a, u) \end{aligned}$$

$$\begin{aligned}
&= \int_{N^\phi(K)} \text{ap } J_n^{N^\phi(K)} \iota(a, u) J_C^\phi(a, -u) \phi(\mathbf{n}^\phi(u)) \int_0^{-\frac{n}{\mathbf{H}_{K,1}^\phi(a,u)}} \left(1 + \frac{t}{n} \mathbf{H}_{K,1}^\phi(a, u)\right)^n dt d\mathcal{H}^n(a, u) \\
&\leq \int_{\tilde{N}_n^\phi(C)} \phi(\mathbf{n}^\phi(u)) J_C^\phi(a, u) \int_0^{\frac{n}{\mathbf{H}_{C,1}^\phi(a,u)}} \left(1 - \frac{t}{n} \mathbf{H}_{C,1}^\phi(a, u)\right)^n dt d\mathcal{H}^n(a, u) \\
&= \frac{n}{n+1} \int_{\tilde{N}_n^\phi(C)} J_C^\phi(a, u) \frac{\phi(\mathbf{n}^\phi(u))}{\mathbf{H}_{C,1}^\phi(a, u)} d\mathcal{H}^n(a, u).
\end{aligned}$$

We discuss now the equality case. We assume that $\mathbf{H}_{C,1}^\phi(a, u) \leq q$ for \mathcal{H}^n a.e. $(a, u) \in \tilde{N}_n^\phi(C)$. We notice that the inequalities in the last estimate become equalities. In particular we deduce the inequality

$$\mathbf{r}_K^\phi(a, u) = -\frac{n}{\mathbf{H}_{K,1}^\phi(a, u)} \geq \frac{n}{q} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K)$$

and the umbilicity condition

$$\kappa_{K,1}^\phi(a, u) = \dots = \kappa_{K,n}^\phi(a, u) \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in N^\phi(K).$$

Consequently we infer from Corollary 3.16 that $\{x \in \mathbf{R}^{n+1} : \delta_K^\phi(x) < \frac{n}{q}\} \subseteq \text{Unp}^\phi(K)$. It follows that $S^\phi(K, r)$ is a closed $\mathcal{C}^{1,1}$ -hypersurface for every $0 < r < \frac{n}{q}$ by [DRKS20, Corollary 5.8]. Moreover the aforementioned umbilicity condition gives

$$\chi_{K,1}^\phi(x) = \dots = \chi_{K,n}^\phi(x)$$

for \mathcal{H}^n a.e. $x \in S^\phi(K, r)$ and for every $0 < r < \frac{n}{q}$. We deduce from [DRKS20, Lemma 3.2] that for every $0 < r < \frac{n}{q}$ there exists $c_r \in \mathbf{R}^{n+1}$ and $\lambda_r > 0$ so that $S^\phi(K, r) = c_r + \lambda_r \mathcal{W}^\phi$. (Notice that the last line of [DRKS20, Lemma 3.2] contains a typo: one should replace the equality $M = \partial \mathbf{B}^F(a, |\lambda|^{-1})$ with $M = \partial \mathbf{B}^{F^*}(a, |\lambda|^{-1})$, which is what the proof given there proves.) Now fix $0 < r < \frac{n}{q}$ and notice that

$$\frac{\nabla \delta_K^\phi(c_r + \lambda_r z)}{|\nabla \delta_K^\phi(c_r + \lambda_r z)|} = -\mathbf{n}^\phi(z)$$

and

$$z = \nabla \phi(\mathbf{n}^\phi(z)) = -\nabla \phi(\nabla \delta_K^\phi(c_r + \lambda_r z)) = -\boldsymbol{\nu}_K^\phi(c_r + \lambda_r z)$$

for every $z \in \mathcal{W}^\phi$, whence we infer that

$$c_r + (\lambda_r + r)\mathcal{W}^\phi = \{x - r\boldsymbol{\nu}_K^\phi(x) : x \in S^\phi(K, r)\} \subseteq \partial K.$$

This implies that $c_r + (\lambda_r + r)\mathcal{W}^\phi = \partial K$ and the proof is complete. \square

5.3 Remark. Similar inequalities have been proved for smooth domains in [HLMG09] and for certain sets of finite perimeter in [DRKS20].

5.4 Theorem. *Suppose $C \subseteq \mathbf{R}^{n+1}$ is a convex body, $r = 1, \dots, n$, $\lambda > 0$ and $\mathcal{C}_{n-r}^\phi(C, \cdot) = \lambda \mathcal{C}_n^\phi(C, \cdot)$. Then $\partial C = a + s\mathcal{W}^\phi$ for some $a \in \mathbf{R}^{n+1}$ and $s > 0$.*

Proof. The Newton-McLaurin inequality (see [Nic00, Theorem 1.1]) together with Lemma 4.6 implies

$$\frac{1}{n} \mathbf{H}_{C,1}^\phi(a, u) \geq \frac{\mathcal{C}_n^\phi(C)}{(n+1)\mathcal{L}^{n+1}(C)} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in \tilde{N}_n^\phi(C). \quad (8)$$

For every $\epsilon > 0$ we set

$$Z_\epsilon = \left\{ (a, u) \in \tilde{N}_n^\phi(C) : \frac{1}{n} \mathbf{H}_{C,1}^\phi(a, u) \geq (1 + \epsilon) \frac{\mathcal{C}_n^\phi(C)}{(n+1)\mathcal{L}^{n+1}(C)} \right\}.$$

We claim that $\mathcal{H}^n(Z_\epsilon) = 0$ for every $\epsilon > 0$. If there existed $\epsilon > 0$ so that $\mathcal{H}^n(Z_\epsilon) > 0$, then we could employ (8) to estimate

$$\begin{aligned} & n \int_{\tilde{N}_n^\phi(C)} J_C^\phi(a, u) \frac{\phi(\mathbf{n}^\phi(u))}{\mathbf{H}_{C,1}^\phi(a, u)} d\mathcal{H}^n(a, u) \\ &= n \int_{\tilde{N}_n^\phi(C) \setminus Z_\epsilon} J_C^\phi(a, u) \frac{\phi(\mathbf{n}^\phi(u))}{\mathbf{H}_{C,1}^\phi(a, u)} d\mathcal{H}^n(a, u) + n \int_{Z_\epsilon} J_C^\phi(a, u) \frac{\phi(\mathbf{n}^\phi(u))}{\mathbf{H}_{C,1}^\phi(a, u)} d\mathcal{H}^n(a, u) \\ &\leq \frac{(n+1)\mathcal{L}^{n+1}(C)}{\mathcal{C}_n^\phi(C)} \tilde{\mathcal{C}}_n^\phi(C, N^\phi(C) \setminus Z_\epsilon) + \frac{(n+1)\mathcal{L}^{n+1}(C)}{\mathcal{C}_n^\phi(C)(1+\epsilon)} \tilde{\mathcal{C}}_n^\phi(C, Z_\epsilon) \\ &< (n+1)\mathcal{L}^{n+1}(C) \end{aligned}$$

in contradiction with the inequality in Theorem 5.2.

Since $\mathcal{H}^n(Z_\epsilon) = 0$ for every $\epsilon > 0$, we infer from (8) that

$$\frac{1}{n} \mathbf{H}_{C,1}^\phi(a, u) = \frac{\mathcal{C}_n^\phi(C)}{(n+1)\mathcal{L}^{n+1}(C)} \quad \text{for } \mathcal{H}^n \text{ a.e. } (a, u) \in \tilde{N}_n^\phi(C)$$

and we obtain the conclusion employing the second part of Theorem 5.2. \square

References

- [Ale62] A. D. Alexandrov. A characteristic property of spheres. *Ann. Mat. Pura Appl. (4)*, 58:303–315, 1962.
- [AW21] Ben Andrews and Yong Wei. Volume preserving flow by powers of the k -th mean curvature. *J. Differential Geom.*, 117(2):193–222, 2021.
- [BS18] Maria Chiara Bertini and Carlo Sinestrari. Volume-preserving nonhomogeneous mean curvature flow of convex hypersurfaces. *Ann. Mat. Pura Appl. (4)*, 197(4):1295–1309, 2018.
- [DRKS20] Antonio De Rosa, Sławomir Kolasiński, and Mario Santilli. Uniqueness of critical points of the anisotropic isoperimetric problem for finite perimeter sets. *Arch. Ration. Mech. Anal.*, 238(3):1157–1198, 2020.
- [Fed59] Herbert Federer. Curvature measures. *Trans. Amer. Math. Soc.*, 93:418–491, 1959.
- [Fed69] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [HK78] Ernst Heintze and Hermann Karcher. A general comparison theorem with applications to volume estimates for submanifolds. *Ann. Sci. École Norm. Sup. (4)*, 11(4):451–470, 1978.
- [HL00] Daniel Hug and Günter Last. On support measures in Minkowski spaces and contact distributions in stochastic geometry. *Ann. Probab.*, 28(2):796–850, 2000.
- [HLMG09] Yijun He, Haizhong Li, Hui Ma, and Jianquan Ge. Compact embedded hypersurfaces with constant higher order anisotropic mean curvatures. *Indiana Univ. Math. J.*, 58(2):853–868, 2009.
- [HLW04] Daniel Hug, Günter Last, and Wolfgang Weil. A local Steiner-type formula for general closed sets and applications. *Math. Z.*, 246(1-2):237–272, 2004.
- [HM87] Gary Harris and Clyde Martin. The roots of a polynomial vary continuously as a function of the coefficients. *Proc. Amer. Math. Soc.*, 100(2):390–392, 1987.
- [Hsi54] Chuan-Chih Hsiung. Some integral formulas for closed hypersurfaces. *Math. Scand.*, 2:286–294, 1954.

- [Koh95] Peter Kohlmann. Compact convex bodies with one curvature measure near the surface measure. *J. Reine Angew. Math.*, 458:201–217, 1995.
- [Koh98] Peter Kohlmann. Characterizations via linear combinations of curvature measures. *Arch. Math. (Basel)*, 70(3):250–256, 1998.
- [KS21] Sławomir Kolasiński and Mario Santilli. Regularity of distance functions from arbitrary closed sets, 2021.
- [MR91] Sebastián Montiel and Antonio Ros. Compact hypersurfaces: the Alexandrov theorem for higher order mean curvatures. In *Differential geometry*, volume 52 of *Pitman Monogr. Surveys Pure Appl. Math.*, pages 279–296. Longman Sci. Tech., Harlow, 1991.
- [Nic00] Constantin P. Niculescu. A new look at Newton’s inequalities. *JIPAM. J. Inequal. Pure Appl. Math.*, 1(2):Article 17, 14, 2000.
- [Ros87] Antonio Ros. Compact hypersurfaces with constant higher order mean curvatures. *Rev. Mat. Iberoamericana*, 3(3-4):447–453, 1987.
- [Ros88] Antonio Ros. Compact hypersurfaces with constant scalar curvature and a congruence theorem. *J. Differential Geom.*, 27(2):215–223, 1988. With an appendix by Nicholas J. Korevaar.
- [San20a] Mario Santilli. Fine properties of the curvature of arbitrary closed sets. *Ann. Mat. Pura Appl. (4)*, 199(4):1431–1456, 2020.
- [San20b] Mario Santilli. Uniqueness of singular convex hypersurfaces with lower bounded k-th mean curvature, 2020.
- [San21] Mario Santilli. Distance functions with dense singular sets. *Comm. Partial Differential Equations*, 46(7):1319–1325, 2021.
- [Sch79] Rolf Schneider. Bestimmung konvexer Körper durch Krümmungsmasse. *Comment. Math. Helv.*, 54(1):42–60, 1979.
- [Sch14] Rolf Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, expanded edition, 2014.
- [Sin15] Carlo Sinestrari. Convex hypersurfaces evolving by volume preserving curvature flows. *Calc. Var. Partial Differential Equations*, 54(2):1985–1993, 2015.
- [Zäh86] M. Zähle. Integral and current representation of Federer’s curvature measures. *Arch. Math. (Basel)*, 46(6):557–567, 1986.

Institut für Mathematik, Universität Augsburg,
 Universitätsstr. 14, 86159, Augsburg, Germany,
 mario.santilli@math.uni-augsburg.de

New affiliation

Mario Santilli, Department of Information Engineering, Computer Science and Mathematics, Università degli Studi dell’Aquila, 67100 L’Aquila, Italy, mario.santilli@univaq.it