Boundedness of Calderón–Zygmund Operators on Special John–Nirenberg–Campanato and Hardy-Type Spaces via Congruent Cubes

Hongchao Jia, Jin Tao, Dachun Yang, Wen Yuan and Yangyang Zhang

Abstract Let $p \in [1, \infty], q \in (1, \infty), s \in \mathbb{Z}_+ := \mathbb{N} \cup \{0\}$, and $\alpha \in \mathbb{R}$. In this article, the authors introduce a reasonable version \widetilde{T} of the Calderón–Zygmund operator T on $JN^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, the special John–Nirenberg–Campanato space via congruent cubes, which coincides with the Campanato space $C_{\alpha,q,s}(\mathbb{R}^n)$ when $p = \infty$. Then the authors prove that \widetilde{T} is bounded on $JN^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}^n_+$ with $|\gamma| \leq s$, $T^*(x^\gamma) = 0$, which is a well-known assumption. To this end, the authors find an equivalent version of this assumption. Moreover, the authors show that T can be extended to a unique continuous linear operator on the Hardy-kind space $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, the predual space of $JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ with $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$, if and only if, for any $\gamma \in \mathbb{Z}^n_+$ with $|\gamma| \leq s$, $T^*(x^\gamma) = 0$. The main interesting integrands in the latter boundedness are that, to overcome the difficulty caused by that $\|\cdot\|_{HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ is no longer concave, the authors first find an equivalent norm of $\|\cdot\|_{HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$, and then establish a criterion for the boundedness of linear operators on $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ via introducing molecules of $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, using the boundedness of \widetilde{T} on $JN^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, and skillfully applying the dual relation $(HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n))^* = JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$.

1 Introduction

Let $q \in (0, \infty]$ and E be any given measurable set of \mathbb{R}^n . The *Lebesgue space* $L^q(E)$ is defined to be the set of all measurable functions f on E such that

$$||f||_{L^{q}(E)} := \begin{cases} \left[\int_{E} |f(x)|^{q} dx \right]^{\frac{1}{q}} & \text{when} \quad q \in (0, \infty), \\ \operatorname{ess \, sup} |f(x)| & \text{when} \quad q = \infty \end{cases}$$

is finite. Moreover, the *space* $L^q_{loc}(\mathbb{R}^n)$ is defined to be the set of all measurable functions f on \mathbb{R}^n such that $f\mathbf{1}_F \in L^q(\mathbb{R}^n)$ for any given bounded measurable set $F \subset \mathbb{R}^n$, here and thereafter, we use $\mathbf{1}_F$ to denote the *characteristic function* of F.

²⁰²⁰ Mathematics Subject Classification. Primary 42B20; Secondary 47A30, 42B30, 46E35, 42B35.

Key words and phrases. John-Nirenberg space, Campanato space, Hardy-kind space, Calderón-Zygmund operator, molecule, dual.

This project is partially supported by the National Natural Science Foundation of China (Grant Nos. 11971058, 12071197, 12122102 and 11871100) and the National Key Research and Development Program of China (Grant No. 2020YFA0712900).

^{*}Corresponding author, E-mail: dcyang@bnu.edu.cn/August 23, 2021/Final version.

In this article, a *cube* Q of \mathbb{R}^n always has finite side length and all its sides parallel to the coordinate axes, but Q is not necessary to be open or closed. Let Q_0 be a given cube of \mathbb{R}^n . John and Nirenberg [20] introduced the well-known space BMO (Q_0) and, in the same article, they also introduced the space $JN_p(Q_0)$ as a natural generalization of the space BMO (Q_0) , which is now called the John–Nirenberg space and can be naturally defined on \mathbb{R}^n as well (see, for instance, [42]). In what follows, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any bounded measurable set $E \subset \mathbb{R}^n$ with |E| > 0, let

$$f_E := \int_E f(x) \, dx := \frac{1}{|E|} \int_E f(x) \, dx.$$

Definition 1.1. Let $p \in (1, \infty)$ and Q_0 be a given cube of \mathbb{R}^n . The *John–Nirenberg space* $JN_p(Q_0)$ is defined to be the set of all $f \in L^1(Q_0)$ such that

$$||f||_{JN_p(Q_0)} := \sup \left\{ \sum_j |Q_j| \left[\int_{Q_j} |f(x) - f_{Q_j}| \ dx \right]^p \right\}^{\frac{1}{p}} < \infty,$$

where the supremum is taken over all collections $\{Q_j\}_j$ of interior pairwise disjoint subcubes of Q_0 .

Recently, the John–Nirenberg space has attracted more and more attention. For instance, Aalto et al. [1] introduced the John–Nirenberg space in the context of doubling metric measure spaces. Dafni et al. [6] showed the non-triviality of $JN_p(Q_0)$ and introduced a Hardy-kind space $HK_{p'}(Q_0)$ with $\frac{1}{p} + \frac{1}{p'} = 1$, which proves the predual space of the space $JN_p(Q_0)$ (see also [42] for more generalization results on Q_0 and \mathbb{R}^n). Very recently, Domínguez and Milman [7] introduced and studied sparse Brudnyi and John–Nirenberg spaces. Moreover, Tao et al. [42] studied the John–Nirenberg–Campanato space, which is a generalization of the space $JN_p(Q_0)$, and Sun et al. [41] studied the localized John–Nirenberg–Campanato space. We refer the reader to [13, 14, 16, 25, 27, 28, 44, 45] for more studies on John–Nirenberg-type spaces.

It is well known that the real-variable theory of Hardy-type and Campanato-type spaces on \mathbb{R}^n , including the boundedness of Calderón–Zygmund operators, plays an important role in harmonic analysis and partial differential equations (see, for instance, [5, 24, 35, 32, 33, 38]), but, unfortunately, it is still a challenging and open question to obtain the boundedness of some important operators, for instance, the Hardy–Littlewood maximal operator, the Calderón–Zygmund operator, and the fractional integrals, on John–Nirenberg spaces and their predual spaces, namely, some Hardy-kind spaces.

To shed some light on the boundedness of these important operators on John–Nirenberg spaces, we in [17] studied the special John–Nirenberg–Campanato space via congruent cubes, which is actually a generalization of the Campanato space, and, in this article, we study the boundedness of Calderón–Zygmund operators on these special John–Nirenberg–Campanato spaces. In what follows, for any $\ell \in (0, \infty)$, let $\Pi_{\ell}(\mathbb{R}^n)$ be the class of all collections of interior pairwise disjoint subcubes $\{Q_j\}_j$ of \mathbb{R}^n with side length ℓ ; for any $s \in \mathbb{Z}_+ := \{0, 1, 2, \ldots\}$, let $\mathcal{P}_s(\mathbb{R}^n)$ denote the set of all polynomials of degree not greater than s on \mathbb{R}^n ; moreover, for any $\gamma := (\gamma_1, \ldots, \gamma_n) \in \mathbb{Z}_+^n := (\mathbb{Z}_+)^n$ and $x := (x_1, \ldots, x_n) \in \mathbb{R}^n$, let $|\gamma| := \gamma_1 + \cdots + \gamma_n$ and $x^\gamma := x_1^{\gamma_1} \cdots x_n^{\gamma_n}$.

Definition 1.2. Let $p \in [1, \infty]$, $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. The *special John–Nirenberg–Campanato space via congruent cubes* (for short, *congruent JNC space*), $JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, is defined to be the set of all $f \in L^q_{\text{loc}}(\mathbb{R}^n)$ such that

 $||f||_{JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)}$

$$:= \begin{cases} \sup_{\ell \in (0,\infty)} \sup_{\{Q_j\}_j \in \Pi_\ell(\mathbb{R}^n)} \left[\sum_j \left| Q_j \right| \left\{ \left| Q_j \right|^{-\alpha} \left[\int_{Q_j} \left| f(x) - P_{Q_j}^{(s)}(f)(x) \right|^q dx \right]^{\frac{1}{q}} \right\}^p \right]^{\frac{1}{p}}, \quad p \in [1,\infty), \\ \sup_{\text{cube } Q \subset \mathbb{R}^n} |Q|^{-\alpha} \left[\int_{Q} \left| f(x) - P_{Q}^{(s)}(f)(x) \right|^q dx \right]^{\frac{1}{q}}, \qquad p = \infty \end{cases}$$

is finite, where, for any cube Q of \mathbb{R}^n , $P_Q^{(s)}(f)$ denotes the unique polynomial of degree not greater than s such that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$,

(1.1)
$$\int_{Q} \left[f(x) - P_{Q}^{(s)}(f)(x) \right] x^{\gamma} dx = 0.$$

Remark 1.3. (i) It is well known that, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any cube (or ball) Q of \mathbb{R}^n , $P_Q^{(0)}(f) = f_Q$ and, for any $s \in \mathbb{Z}_+$, there exists a constant $C_{(s)} \in [1, \infty)$, independent of f and Q, such that, for any $x \in Q$,

(1.2)
$$\left| P_Q^{(s)}(f)(x) \right| \le C_{(s)} \oint_Q |f(y)| \, dy;$$

see, for instance, [23, 34] for more properties of $P_Q^{(s)}(f)$.

(ii) $JN^{\mathrm{con}}_{(\infty,q,s)_{\alpha}}(\mathbb{R}^n)$ is just the well-known Campanato space $C_{\alpha,q,s}(\mathbb{R}^n)$ which was introduced by Campanato [4] and which when $\alpha = 0$ coincides with the space BMO (\mathbb{R}^n) .

Moreover, in [17], we introduced a Hardy-kind space $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ which proves the predual space of $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ (see, for instance, [17, Theorem 4.10]), here and thereafter, p' and q' denote, respectively, the conjugate indexes of p and q, namely, $\frac{1}{p} + \frac{1}{p'} = 1$ and $\frac{1}{q} + \frac{1}{q'} = 1$. In this article, we also establish the boundedness of Calderón–Zygmund operators on such a Hardy-type space (see Theorem 3.20 below).

Furthermore, inspired by the boundedness of the Calderón–Zygmund operator T from $L^{\infty}(\mathbb{R}^n)$ to BMO (\mathbb{R}^n) (see, for instance, [9, Theorem 6.6]), we also introduce the following subspace $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$ of $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$, and further study the boundedness of T on $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$; see Proposition 2.5 below.

Definition 1.4. Let $p, q \in [1, \infty]$ and $\alpha \in \mathbb{R}$. The special Riesz–Morrey space via congruent cubes, $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, is defined to be the set of all $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ such that

$$||f||_{RM^{\text{con}}_{p,q,\alpha}(\mathbb{R}^n)} := \begin{cases} \sup_{\ell \in (0,\infty)} \sup_{\{Q_j\}_j \in \Pi_{\ell}(\mathbb{R}^n)} \left\{ \sum_j \left| Q_j \right| \left[\left| Q_j \right|^{-\alpha - \frac{1}{q}} ||f||_{L^q(Q_j)} \right]^p \right\}^{\frac{1}{p}}, & p \in [1,\infty), \\ \sup_{\text{cube } Q \subset \mathbb{R}^n} |Q|^{-\alpha - \frac{1}{q}} ||f||_{L^q(Q)}, & p = \infty \end{cases}$$

is finite.

- **Remark 1.5.** (i) If $1 \le q < \alpha < p \le \infty$, then the space $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ coincides with the space $(L^q, \ell^p)^{\frac{p}{1-p\alpha}}(\mathbb{R}^n)$ which was introduced by Fofana [11]; see [8, 10, 12] for more studies on the space $(L^q, \ell^p)^{\alpha}(\mathbb{R}^n)$.
 - (ii) If we do not require that $\{Q_j\}_j$ have the same size in the definition of $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$, then such a space is just the Riesz–Morrey space $RM_{p,q,\alpha}(\mathbb{R}^n)$ introduced by Tao et al. [43] and, independently, by Fofana et al. [12]; see also, for instance, [12, 43, 46] for more studies on $RM_{p,q,\alpha}(\mathbb{R}^n)$.
- (iii) If $p = \infty$, $q \in (0, \infty)$, and $\alpha \in [-\frac{1}{q}, 0)$, then the space $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ coincides with the Morrey space introduced by Morrey [29]; see, for instance, [30, 31, 37] for more studies on Morrey-type spaces.

Let T be a Calderón–Zygmund operator as in Definition 2.9 below. In this article, we first introduce a reasonable version of the Calderón–Zygmund operator T on $JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, denoted by \widetilde{T} . Then we prove that \widetilde{T} is bounded on $JN^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}^n_+$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$, which is a well-known assumption; see Theorem 2.25 below. To this end, we find an equivalent version of this assumption; see Proposition 2.16 below. Moreover, we show that T can be extended to a unique continuous linear operator on the Hardy-kind space $HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$; see Theorem 3.20 below. To obtain the desired extension of T on $HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$, one key step is to estimate $||T(g)||_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)}$ for any $g \in \mathbb{R}$ $HK_{(p,q,s)_{\alpha}}^{\text{con-fin}}(\mathbb{R}^n)$ (see Definition 3.5 below), but the classical method used to extend T from the finite atomic Hardy space to the corresponding Hardy space is no longer feasible because $\|\cdot\|_{HK_{(p,q)}^{\text{con}}}$ (\mathbb{R}^n) is not concave with $p \in (1, \infty)$, which is essentially different from the Hardy space $H^p(\mathbb{R}^n)$ with $p \in (0,1]$. To avoid this, we first find an equivalent norm, $\|\cdot\|_{(JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*}$, of $\|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ in Proposition 3.8 below, and hence it suffices to estimate $||T(g)||_{(JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*}$. Moreover, we use the properties of molecules of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and the dual relation $(HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n))^* = JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ to reduce the estimate of $||T(g)||_{(JN_{(p',q',s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n}))^{*}}$ to the boundedness of \overline{T} on $JN_{(p',q',s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})$ obtained in Theorem 2.25; see Theorem 3.16 below for more details. To limit the length of this article, the boundedness of fractional integrals and Littlewood-Paley operators on $JN_{(p,q,s)_n}^{\text{con}}(\mathbb{R}^n)$ will be studied, respectively, in the forthcoming articles [18, 19].

The remainder of this article is organized as follows.

The main aim of Section 2 is to establish the boundedness of Calderón–Zygmund operators, respectively, on $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$ and $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$. In Subsection 2.1, we study the boundedness of Calderón–Zygmund operators on $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$; see Proposition 2.5 below. In Subsection 2.2, we first introduce a reasonable version of Calderón–Zygmund operators via borrowing some ideas from [32, Section 4]. Then we show that such a modified Calderón–Zygmund operator is bounded on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$; see Theorem 2.25 below.

In Section 3, we establish the boundedness of Calderón–Zygmund operators on the Hardy-kind space $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. In Subsection 3.1, we recall the notion of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ in Definition 3.4 below, and then prove that $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ is a Banach space in Proposition 3.9 below. In Subsection 3.2, to overcome the difficulty caused by the fact that $\|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ is no longer concave, we establish a criterion (see Theorem 3.16 below) for the boundedness of linear operators on $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ via introducing molecules of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and skillfully using the dual relation

 $(HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n))^* = JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$. In Subsection 3.3, as an application of Theorem 3.16, we prove that the Calderón–Zygmund operator T can be extended to a unique continuous linear operator on the Hardy-kind space $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$; see Theorem 3.20 below.

Finally, we make some conventions on notation. Let $\mathbb{N}:=\{1,2,\ldots\}$ and $\mathbb{Z}_+:=\mathbb{N}\cup\{0\}$. For a given $s\in\mathbb{Z}_+$ and a given ball $B\subset\mathbb{R}^n$, $\mathcal{P}_s(B)$ [resp., $\mathcal{P}_s(\mathbb{R}^n)$] denotes the set of all polynomials of degree not greater than s on B (resp., \mathbb{R}^n). We always denote by C a positive constant which is independent of the main parameters, but it may vary from line to line. We also use $C_{(\alpha,\beta,\ldots)}$ to denote a positive constant depending on the indicated parameters α,β,\ldots . The symbol $f\lesssim g$ means that $f\leq Cg$. If $f\lesssim g$ and $g\lesssim f$, we then write $f\sim g$. If $f\leq Cg$ and g=h or $g\leq h$, we then write $f\lesssim g\sim h$ or $f\lesssim g\lesssim h$, rather than $f\lesssim g=h$ or $f\lesssim g\leq h$. We use $\mathbf{0}$ to denote the origin of \mathbb{R}^n . For any $E\subset\mathbb{R}^n$, we denote by $\mathbf{1}_E$ its characteristic function. We use $\eta\to 0^+$ to denote $\eta\in(0,\infty)$ and $\eta\to 0$. For any $m\in\mathbb{N}$, let $m!:=1\times\cdots\times m$ and m!:=1. For any m:=1 in m:=1

2 Boundedness of Calderón–Zygmund Operators on $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ and $JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$

In this section, we first establish the boundedness of Calderón–Zygmund operators on the space $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$. Moreover, we give a reasonable version of Calderón–Zygmund operators on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$ via borrowing some ideas from [32, Section 4]. Then we obtain the boundedness of such Calderón–Zygmund operators on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$. To this end, we first find an equivalent version of the well-known assumption that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$; see Proposition 2.16 below.

2.1 Calderón–Zygmund Operators on $RM_{p,q,\alpha}^{con}(\mathbb{R}^n)$

This subsection is devoted to studying the boundedness of Calderón–Zygmund operators on $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$. We first recall the following notion of the (generalized) Calderón–Zygmund operator T (see, for instance, [9, Definition 5.11]).

Definition 2.1. A measurable function K on $\mathbb{R}^n \times \mathbb{R}^n \setminus \{(x, x) : x \in \mathbb{R}^n\}$ is called a *standard kernel* if there exist a $\delta \in (0, 1]$ and a positive constant C such that

(i) for any $x, y \in \mathbb{R}^n$ with $x \neq y$,

$$|K(x,y)| \le \frac{C}{|x-y|^n};$$

(ii) for any $x, y, \omega \in \mathbb{R}^n$ with $x \neq y$ and $|x - y| \ge 2|x - \omega|$,

$$(2.1) |K(x,y) - K(\omega,y)| \le C \frac{|x - \omega|^{\delta}}{|x - y|^{n + \delta}};$$

(iii) (2.1) still holds true for the second variable of K.

In what follows, for any measurable function f on \mathbb{R}^n , we define the *support* supp (f) of f by setting

$$\operatorname{supp}(f) := \overline{\{y \in \mathbb{R}^n : f(y) \neq 0\}}.$$

Definition 2.2. Let K be a standard kernel as in Definition 2.1. A linear operator T is called a *Calderón–Zygmund operator with kernel* K if T is bounded on $L^2(\mathbb{R}^n)$ and, for any given $f \in L^2(\mathbb{R}^n)$ with compact support, and for almost every $x \notin \text{supp}(f)$,

(2.2)
$$T(f)(x) = \int_{\mathbb{R}^n} K(x, y) f(y) \, dy.$$

Next, we establish the boundedness of Calderón–Zygmund operators on $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$. We give two lemmas first. The following lemma gives an equivalent characterization of $RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$, whose proof is similar to that used in [10, Proposition 4.1]; we present the details here for the convenience of the reader.

Lemma 2.3. Let $p, q \in [1, \infty)$ and $\alpha \in \mathbb{R}$. Then $f \in RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ if and only if $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ and

$$[f]_{RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} := \sup_{r \in (0,\infty)} \left[\int_{\mathbb{R}^n} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} |f(x)|^q \, dx \right]^{\frac{1}{q}} \right\}^p \, dy \right]^{\frac{1}{p}} < \infty.$$

Moreover, for any $f \in RM_{p,q,\alpha}^{con}(\mathbb{R}^n)$,

$$[f]_{RM_{p,q,\alpha}^{\operatorname{con}}(\mathbb{R}^n)} \sim ||f||_{RM_{p,q,\alpha}^{\operatorname{con}}(\mathbb{R}^n)},$$

where the positive equivalence constants are independent of f.

Proof. Let p, q, and α be as in the present lemma. We first prove that, for any $f \in L^q_{loc}(\mathbb{R}^n)$,

$$(2.3) ||f||_{RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} \gtrsim [f]_{RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}.$$

Fix $r \in (0, \infty)$. Let $2\mathbb{Z}^n := \{2k : k \in \mathbb{Z}^n\}$, $I := \{(i_1, \dots, i_n) : i_1 \dots, i_n \in \{0, 1\}\}$, and, for any $k \in \mathbb{Z}^n$, $Q_{r,k} := rk + [0, r)^n$. Observe that, for any given $k \in \mathbb{Z}^n$ and for any $y \in Q_{r,k}$, $B(y, r/2) \subset 2Q_{r,k}$. From this, $p \in (1, \infty)$, and the fact that, for any given $i \in I$, $\{2Q_{r,k+i}\}_{k \in 2\mathbb{Z}^n}$ have disjoint interiors, we deduced that, for any $f \in RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$,

$$\left[\int_{\mathbb{R}^{n}} \left\{ |B(y, r/2)|^{-\alpha} \left[\int_{B(y, r/2)} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
= \left[\sum_{i \in I} \sum_{k \in 2\mathbb{Z}^{n}} \int_{Q_{r,k+i}} \left\{ |B(y, r/2)|^{-\alpha} \left[\int_{B(y, r/2)} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
\lesssim \left[\sum_{i \in I} \sum_{k \in 2\mathbb{Z}^{n}} \int_{Q_{r,k+i}} \left\{ |2Q_{r,k+i}|^{-\alpha} \left[\int_{2Q_{r,k+i}} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}}$$

$$\leq \sum_{i \in I} \left[\sum_{k \in 2\mathbb{Z}^n} \left| 2Q_{r,k+i} \right| \left\{ \left| 2Q_{r,k+i} \right|^{-\alpha} \left[\int_{2Q_{r,k+i}} |f(x)|^q \ dx \right]^{\frac{1}{q}} \right\}^p \right]^{\frac{1}{p}}$$

$$\leq \|f\|_{RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)},$$

which implies that (2.3) holds true.

Now, to complete the proof of the present lemma, we only need to show that, for any $f \in L^q_{loc}(\mathbb{R}^n)$,

Let $\{Q_i\}_i \in \Pi_r(\mathbb{R}^n)$ with $r \in (0, \infty)$. Then it is easy to see that

$$\left[\sum_{j} |Q_{j}| \left\{ |Q_{j}|^{-\alpha} \left[\int_{Q_{j}} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} \right]^{\frac{1}{p}}$$

$$= \left[\sum_{j} \int_{Q_{j}} \left\{ |Q_{j}|^{-\alpha} \left[\int_{Q_{j}} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}}$$

$$\lesssim \left[\sum_{j} \int_{Q_{j}} \left\{ |B(y, \sqrt{n}r)|^{-\alpha} \left[\int_{B(y, \sqrt{n}r)} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}}$$

$$\lesssim \left[\int_{\mathbb{R}^{n}} \left\{ |B(y, \sqrt{n}r)|^{-\alpha} \left[\int_{B(y, \sqrt{n}r)} |f(x)|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}},$$

which implies (2.4) holds true. This finishes the proof of Proposition 2.3.

The following lemma is just [2, Proposition 8.4(a)]; see also [21, Theorem 7] for the corresponding weighted case. Recall that, for any given $r \in (0, \infty)$ and $p, q \in (1, \infty)$, the *space* $(E_q^p)_r(\mathbb{R}^n)$ is defined to be the set of all $f \in L_{loc}^q(\mathbb{R}^n)$ such that

$$||f||_{(E_q^p)_r(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}^n} \left[\int_{B(y,r)} |f(x)|^q \, dx \right]^{\frac{p}{q}} \, dy \right\}^{\frac{1}{p}} < \infty;$$

see, for instance, [2, p. 1600].

Lemma 2.4. Let $p, q \in (1, \infty)$ and T be a Calderón–Zygmund operator as in Definition 2.2. Then there exists a positive constant C such that, for any $r \in (0, \infty)$ and $f \in (E_q^p)_r(\mathbb{R}^n)$,

$$||T(f)||_{(E_q^p)_r(\mathbb{R}^n)} \le C||f||_{(E_q^p)_r(\mathbb{R}^n)}.$$

Using the above two lemmas, we immediately obtain the boundedness of Calderón–Zygmund operators on $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$; we omit the details here.

Proposition 2.5. Let $p, q \in (1, \infty)$, $\alpha \in \mathbb{R}$, and T be a Calderón–Zygmund operator as in Definition 2.2. Then T is bounded on $RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in RM_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$,

$$||T(f)||_{RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)} \le C||f||_{RM_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)}.$$

2.2 Calderón–Zygmund Singular Integral Operators on $JN^{\text{con}}_{(p,q,s)_o}(\mathbb{R}^n)$

In this subsection, we first give a reasonable version of the Calderón–Zygmund operator T on $JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, denoted by \widetilde{T} . Then we establish the boundedness of \widetilde{T} on $JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. We begin with the notion of the s-order standard kernel (see, for instance [39, Chapter III]). In what follows, for any $\gamma = (\gamma_1, \ldots, \gamma_n) \in \mathbb{Z}_+^n$, any γ -order differentiable function $F(\cdot, \cdot)$ on $\mathbb{R}^n \times \mathbb{R}^n$, and any $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$, let

$$\partial_{(1)}^{\gamma} F(x, y) := \frac{\partial^{|\gamma|}}{\partial x_1^{\gamma_1} \cdots \partial x_n^{\gamma_n}} F(x, y) \quad \text{and} \quad \partial_{(2)}^{\gamma} F(x, y) := \frac{\partial^{|\gamma|}}{\partial y_1^{\gamma_1} \cdots \partial y_n^{\gamma_n}} F(x, y).$$

Definition 2.6. Let $s \in \mathbb{Z}_+$. A measurable function K on $\mathbb{R}^n \times \mathbb{R}^n \setminus \{(x, x) : x \in \mathbb{R}^n\}$ is called an *s-order standard kernel* if there exist a positive constant C and a $\delta \in (0, 1]$ such that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, the following hold true:

(i) for any $x, y \in \mathbb{R}^n$ with $x \neq y$,

(2.5)
$$\left|\partial_{(2)}^{\gamma}K(x,y)\right| \le \frac{C}{|x-y|^{n+|\gamma|}};$$

- (ii) (2.5) still holds true for the first variable of K;
- (iii) for any $x, y, z \in \mathbb{R}^n$ with $x \neq y$ and $|x y| \ge 2|y z|$,

(2.6)
$$\left| \partial_{(2)}^{\gamma} K(x, y) - \partial_{(2)}^{\gamma} K(x, z) \right| \le C \frac{|y - z|^{\delta}}{|x - y|^{n + |\gamma| + \delta}};$$

(iv) (2.6) still holds true for the first variable of K.

In what follows, we use $\eta \to 0^+$ to denote $\eta \in (0, \infty)$ and $\eta \to 0$.

Definition 2.7. Let $s \in \mathbb{Z}_+$ and K be an s-order standard kernel as in Definition 2.6. A linear operator T is called an s-order Calderón–Zygmund singular integral operator with kernel K if T is bounded on $L^2(\mathbb{R}^n)$ and, for any given $f \in L^2(\mathbb{R}^n)$ and for almost every $x \in \mathbb{R}^n$,

(2.7)
$$T(f)(x) = \lim_{\eta \to 0^+} T_{\eta}(f)(x),$$

where

(2.8)
$$T_{\eta}(f)(x) := \int_{\mathbb{R}^n \setminus B(x,\eta)} K(x,y) f(y) \, dy.$$

Remark 2.8. Let T be as in Definition 2.7. Since (2.7) implies (2.2), we find that T is a Calderón–Zygmund operator as in Definition 2.2. Moreover, by [9, p. 102], we conclude that T is well defined on $L^q(\mathbb{R}^n)$ with $q \in (1, \infty)$ and, for any $f \in L^q(\mathbb{R}^n)$,

$$T(f) = \lim_{\eta \to 0^+} T_{\eta}(f)$$

both almost everywhere on \mathbb{R}^n and in $L^q(\mathbb{R}^n)$.

Let T be an s-order Calderón–Zygmund singular integral operator. Recall the well-known assumption on T that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, $T^*(x^\gamma) = 0$, namely, for any $a \in L^2(\mathbb{R}^n)$ having compact support and satisfying that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, $\int_{\mathbb{R}^n} a(x) x^\gamma dx = 0$, it holds true that

(2.9)
$$\int_{\mathbb{R}^n} T(a)(x)x^{\gamma} dx = 0;$$

see, for instance, [3, Definition 9.4].

Definition 2.9. Let $s \in \mathbb{Z}_+$. An *s*-order Calderón–Zygmund singular integral operator T is said to have the *vanishing moments up to order* s if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, $T^*(x^{\gamma}) = 0$.

Remark 2.10. Let K be a measurable function on $\mathbb{R}^n \setminus \{\mathbf{0}\}$ such that $K_1(x, y) := K(x - y)$, with x, $y \in \mathbb{R}^n$ and $x \neq y$, is an s-order standard kernel. Let T be an s-order Calderón–Zygmund singular integral operator with kernel K_1 . It is well known that such a T satisfies (2.9); see, for instance, [23, 26]. This indicates that (2.9) is a reasonable assumption.

Next, we give the following notion of the *s*-order modified Calderón–Zygmund operators via borrowing some ideas from [32, Section 4].

Definition 2.11. Let $s \in \mathbb{Z}_+$, K be an s-order standard kernel, and T the s-order Calderón–Zygmund operator with kernel K. For any $x, y \in \mathbb{R}^n$ with $x \neq y$, let

$$(2.10) \widetilde{K}(x, y) := K(y, x)$$

and $B_0 := B(x_0, r_0)$ be a given ball of \mathbb{R}^n with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$. The *s-order modified Calderón–Zygmund operator* \widetilde{T}_{B_0} *with kernel* \widetilde{K} is defined by setting, for any suitable function f on \mathbb{R}^n , and almost every $x \in \mathbb{R}^n$,

(2.11)
$$\widetilde{T}_{B_0}(f)(x) := \lim_{\eta \to 0^+} \widetilde{T}_{B_0,\eta}(f)(x)$$

pointwisely, where, for any $\eta \in (0, \infty)$,

$$(2.12) \qquad \widetilde{T}_{B_0,\eta}(f)(x) := \int_{\mathbb{R}^n \backslash B(x,\eta)} \left[\widetilde{K}(x,y) - \sum_{\{\gamma \in \mathbb{Z}_+^n \colon |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0,y)}{\gamma!} (x - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \backslash B_0}(y) \right] f(y) \, dy.$$

The following conclusion shows that \widetilde{T}_{B_0} coincides with the s-order Calderón–Zygmund singular integral operator with kernel \widetilde{K} on $L^q(\mathbb{R}^n)$, with $q \in (1, \infty)$, in the sense of modulo $\mathcal{P}_s(\mathbb{R}^n)$.

Proposition 2.12. Let $s \in \mathbb{Z}_+$, $q \in (1, \infty)$, K be an s-order standard kernel, and T the s-order Calderón–Zygmund singular integral operator with kernel K. Let \widetilde{K} be as in (2.10), $T_{(1)}$ an s-order Calderón–Zygmund singular integral operator with kernel \widetilde{K} as in Definition 2.7, and $B_0 := B(x_0, r_0) \subset \mathbb{R}^n$ a given ball with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$. Then, for any $f \in L^q(\mathbb{R}^n)$, $\widetilde{T}_{B_0}(f)$ in (2.11) is well defined almost everywhere on \mathbb{R}^n , and $T_{(1)}(f) - \widetilde{T}_{B_0}(f) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero.

Proof. Let $s, q, T, K, \widetilde{K}, T_{(1)}$, and $B_0 := B(x_0, r_0)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$ be as in the present proposition. We first show that, for any $f \in L^q(\mathbb{R}^n)$, $\widetilde{T}_{B_0}(f)$ exists almost everywhere on \mathbb{R}^n . Indeed, by Remark 2.8, (2.5), (2.10), the Hölder inequality, and $q \in (1, \infty)$, we find that, for any $f \in L^q(\mathbb{R}^n)$ and almost every $x \in \mathbb{R}^n$,

$$|\widetilde{T}_{B_{0}}(f)(x)| \leq \left| \lim_{\eta \to 0^{+}} \int_{\mathbb{R}^{n} \backslash B(x,\eta)} \widetilde{K}(x,y) f(y) \, dy \right|$$

$$+ \int_{\mathbb{R}^{n}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \left| \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \backslash B_{0}}(y) f(y) \right| \, dy$$

$$\leq \left| T_{(1)}(f)(x) \right| + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} |x - x_{0}|^{|\gamma|} \int_{\mathbb{R}^{n} \backslash B_{0}} \frac{|f(y)|}{|y - x_{0}|^{n+|\gamma|}} \, dy$$

$$\leq \left| T_{(1)}(f)(x) \right| + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} |x - x_{0}|^{|\gamma|} ||f||_{L^{q}(\mathbb{R}^{n})} \left[\int_{\mathbb{R}^{n} \backslash B_{0}} \frac{1}{|y - x_{0}|^{q'(n+|\gamma|)}} \, dy \right]^{\frac{1}{q'}}$$

$$< \infty,$$

This implies that $\widetilde{T}_{B_0}(f)$ exists almost everywhere on \mathbb{R}^n .

Moreover, it is easy to see that, for any $f \in L^q(\mathbb{R}^n)$ and almost every $x \in \mathbb{R}^n$,

$$T_{(1)}(f)(x) - \widetilde{T}_{B_0}(f)(x) = \int_{\mathbb{R}^n} \left[\sum_{\{y \in \mathbb{Z}^n : |y| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] f(y) \, dy.$$

From this and the estimation of (2.13), we deduce that $T_{(1)}(f) - \widetilde{T}_{B_0}(f) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero. This finishes the proof of Proposition 2.12.

Now, we show that \widetilde{T}_{B_0} in Definition 2.11 is well defined on $JN^{\mathrm{con}}_{(p,q,s)_a}(\mathbb{R}^n)$ (see Proposition 2.22 below) and changing B_0 in the definition of \widetilde{T}_{B_0} results in adding a polynomial of $\mathcal{P}_s(\mathbb{R}^n)$ (see Remark 2.23 below). To this end, we first establish some technical lemmas. The following lemma is the boundedness of Calderón–Zygmund operators on $L^q(\mathbb{R}^n)$ with $q \in (1, \infty)$; see, for instance, [9, p. 99, Theorem 5.10].

Lemma 2.13. Let $q \in (1, \infty)$ and T be a Calderón–Zygmund operator as in Definition 2.2. Then T is bounded on $L^q(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in L^q(\mathbb{R}^n)$,

$$||T(f)||_{L^q(\mathbb{R}^n)} \le C||f||_{L^q(\mathbb{R}^n)}.$$

The following conclusion might be well known but, to the best of our knowledge, we did not find a complete proof. For the convenience of the reader, we present the details here.

Lemma 2.14. Let $s \in \mathbb{Z}_+$, a ball $B \subset \mathbb{R}^n$, and $\{a_\gamma\}_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \le s\}} \subset L^1_{loc}(\mathbb{R}^n)$. If, for any $x \in \mathbb{R}^n$ (resp., $x \in B$),

(2.14)
$$P(x) := \int_{\mathbb{R}^n} \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \le s\}} a_{\gamma}(y) x^{\gamma} dy$$

is finite, then $P \in \mathcal{P}_s(\mathbb{R}^n)$ [resp., $\mathcal{P}_s(B)$].

Proof. Let s, $\{a_{\gamma}\}_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}}$, and P be as in the present lemma. Without loss of generality, we may only show the case \mathbb{R}^{n} . To this end, it suffices to prove that, for any given $\gamma_{0} := (\gamma_{0}^{(1)}, \ldots, \gamma_{0}^{(n)}) \in \mathbb{Z}_{+}^{n}$ with $|\gamma_{0}| \leq s$,

$$\left| \int_{\mathbb{R}^n} a_{\gamma_0}(y) \, dy \right| < \infty.$$

Indeed, if (2.15) holds true, using the linearity of the integral, we find that

$$P(x) = \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \le s\}} \int_{\mathbb{R}^n} a_{\gamma}(y) \, dy \, x^{\gamma}$$

and hence $P \in \mathcal{P}_s(\mathbb{R}^n)$. To prove (2.15), for any $(x_1, \dots, x_n) \in \mathbb{R}^n$ and $y \in \mathbb{R}^n$, let

(2.16)
$$F_{x_1,\dots,x_{n-1},y}(x_n) := \sum_{\{\gamma:=(\gamma_1,\dots,\gamma_n)\in\mathbb{Z}_+^n: |\gamma|\leq s\}} a_{\gamma}(y)x_1^{\gamma_1}\cdots x_n^{\gamma_n}.$$

Regard $F_{x_1,...,x_{n-1},y}$ as an element of $\mathcal{P}_s(\mathbb{R})$. Then, from the Lagrange interpolation formula, we deduce that, for any (x_1,\ldots,x_n) , $y\in\mathbb{R}^n$,

(2.17)
$$F_{x_1,\dots,x_{n-1},y}(x_n) = \sum_{i=1}^{s+1} F_{x_1,\dots,x_{n-1},y}(a_i) \prod_{j \neq i, \ 1 \le j \le s+1} \frac{x_n - a_j}{a_i - a_j},$$

where $\{a_j\}_{j=1}^{s+1} \subset \mathbb{R}$ satisfy, for any $i, j \in \{1, \ldots, s+1\}$ with $i \neq j, a_i \neq a_j$. Moreover, by (2.14) with x replaced by $(x_1, \ldots, x_{n-1}, a_i)$, and (2.16), we conclude that, for any $i \in \{1, \ldots, s+1\}$,

$$\left| \int_{\mathbb{R}^n} F_{x_1,\dots,x_{n-1},y}(a_i) \, dy \right| < \infty$$

From this and (2.17), we deduce that the coefficient of the term x_n in the right hand of (2.17) is integrable on y and, moreover, by (2.16), we find that the coefficient of the term x_n in the right hand side of (2.17) is just

$$F_{x_1,\dots,x_{n-1},y}(1) = \sum_{\{\gamma:=(\gamma_1,\dots\gamma_n)\in\mathbb{Z}_+^n: \ |\gamma|\leq s \text{ and } \gamma_n=\gamma_0^{(n)}\}} a_{\gamma}(y)x_1^{\gamma_1}\cdots x_{n-1}^{\gamma_{n-1}}$$

and hence, for any $(x_1, \ldots, x_{n-1}) \in \mathbb{R}^{n-1}$,

$$\left| \int_{\mathbb{R}^n} \sum_{\{\gamma := (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}_+^{n} : |\gamma| \le s \text{ and } \gamma_n = \gamma_0^{(n)} \}} a_{\gamma}(y) x_1^{\gamma_1} \cdots x_{n-1}^{\gamma_{n-1}} dy \right| < \infty.$$

Repeating the above procedure n-1 times, we conclude that (2.15) holds true. This finishes the proof of Lemma 2.14.

Next, we give an equivalent version of (2.9); see Proposition 2.16 below. To this end, we first establish a technical lemma. In what follows, for any $s \in \mathbb{Z}_+$, any $v \in [1, \infty]$, and any measurable subset $E \subset \mathbb{R}^n$, the *space* $L_s^v(E)$ is defined by setting

$$L_s^{\nu}(E) := \left\{ f \in L^{\nu}(E) : \int_E f(x) x^{\gamma} dx = 0 \text{ for any } \gamma \in \mathbb{Z}_+^n \text{ with } |\gamma| \le s \right\}.$$

Lemma 2.15. Let $s \in \mathbb{Z}_+$, K be an s-order standard kernel, and T the s-order Calderón–Zygmund singular integral operator with kernel K. Let \widetilde{K} be as in (2.10), $B_0 := B(x_0, r_0) \subset \mathbb{R}^n$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$ a given ball, and, for any $v := (v_1 \dots, v_n) \in \mathbb{Z}_+^n$ with $|v| \leq s$,

$$(2.18) \qquad \widetilde{T}_{B_0}(y^{\nu})(\cdot) = \lim_{\eta \to 0^+} \int_{\mathbb{R}^n \setminus B(\cdot, \eta)} \left[\widetilde{K}(\cdot, y) - \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \le s\}} \frac{\partial_x^{\gamma} \widetilde{K}(x_0, y)}{\gamma!} (\cdot - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] y^{\nu} \, dy$$

as in (2.11) with f replaced by $y^{\nu} := y_1^{\nu_1} \cdots y_n^{\nu_n}$. Then, for any $\nu \in \mathbb{Z}_+^n$ with $|\nu| \leq s$,

- (i) $\widetilde{T}_{B_0}(y^{\nu})$ is well defined almost everywhere on \mathbb{R}^n , and $\widetilde{T}_{B_0}(y^{\nu}) \in L^q_{loc}(\mathbb{R}^n)$ for any given $q \in (1, \infty)$;
- (ii) for any given ball $B_1 \subset \mathbb{R}^n$,

$$\widetilde{T}_{B_0}(y^{\nu}) - \widetilde{T}_{B_1}(y^{\nu}) \in \mathcal{P}_s(\mathbb{R}^n)$$

after changing values on a set of measure zero;

(iii) for any given ball B := B(z, r) of \mathbb{R}^n with $z \in \mathbb{R}^n$ and $r \in (0, \infty)$, and for any $a \in L_s^q(\mathbb{R}^n)$ supported in B with $q \in (1, \infty)$,

(2.19)
$$\int_{\mathbb{R}^n} T(a)(x)x^{\nu} dx = \int_{\mathbb{R}^n} a(x)\widetilde{T}_{B_0}(y^{\nu})(x) dx.$$

Proof. Let s, T, K, \widetilde{K} , and $B_0 = B(x_0, r_0)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$ be as in the present lemma. We first show (i). To this end, let \widetilde{B} be any given ball of \mathbb{R}^n and

$$R := \sup\{2|x - x_0| + 2|x_0| + 1 : x \in \widetilde{B}\}.$$

Then it is easy to see that $(B_0 \cup \widetilde{B}) \subset B(x_0, R)$ and, for any $y \notin B(x_0, R)$,

$$(2.20) |y| \ge |y - x_0| - |x_0| \ge R - |x_0| \ge 1, |y - x_0| \ge 2|x - x_0|,$$

$$|y| \ge |y - x_0| - |x_0| > |y - x_0| - \frac{R}{2} \ge \frac{1}{2}|y - x_0|,$$

and

$$|y| \le |y - x_0| + |x_0| < |y - x_0| + \frac{R}{2} \le \frac{3}{2}|y - x_0|,$$

and hence

$$(2.21) |y| \sim |y - x_0|.$$

Moreover, by (2.5), (2.10), the Taylor remainder theorem, (2.20), and (2.21), we find that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$, and $x \in B$, there exists an $\widetilde{x} \in \{\theta x + (1 - \theta)x_0 : \theta \in (0, 1)\}$ such that

$$(2.22) \quad \left| \widetilde{T}_{B_0}(y^{\nu})(x) \right| \le \left| \lim_{\eta \to 0^+} \int_{B(x_0, R) \setminus B(x, \eta)} \widetilde{K}(x, y) y^{\nu} \, dy \right|$$

$$+ \left| \lim_{\eta \to 0^{+}} \int_{B(x_{0},R) \setminus B(x,\eta)} \left[\sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] y^{\gamma} dy \right|$$

$$+ \int_{\mathbb{R}^{n} \setminus B(x_{0},R)} \left| \widetilde{K}(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x - x_{0})^{\gamma} \left| |y|^{|\nu|} dy \right|$$

$$\leq \left| T_{(1)}(y^{\nu} \mathbf{1}_{B(x_{0},R)})(x) \right| + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \int_{B(x_{0},R) \setminus B_{0}} \frac{|y|^{|\nu|} |\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)||x - x_{0}|^{|\gamma|}}{\gamma!} dy$$

$$+ \int_{\mathbb{R}^{n} \setminus B(x_{0},R)} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| = s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(\widetilde{X},y) - \partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x - x_{0})^{\gamma} \left| |y|^{|\nu|} dy \right|$$

$$\leq \left| T_{(1)}(y^{\nu} \mathbf{1}_{B(x_{0},R)})(x) \right| + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} |x - x_{0}|^{|\gamma|} \int_{B(x_{0},R) \setminus B_{0}} \frac{|y|^{|\nu|}}{|y - x_{0}|^{n+|\gamma|}} dy$$

$$+ \int_{\mathbb{R}^{n} \setminus B(x_{0},R)} \frac{|y|^{s} |x_{0} - \widetilde{X}|^{\delta} |x - x_{0}|^{s}}{|y - x_{0}|^{n+s+\delta}} dy$$

$$\leq \left| T_{(1)}(y^{\nu} \mathbf{1}_{B(x_{0},R)})(x) \right| + 1,$$

where the implicit positive constants depend on x_0 and R, $T_{(1)}$ denotes the s-order Calderón–Zygmund singular integral operator with kernel \widetilde{K} , and, in the penultimate step, we used (2.6) and (2.10) together with $|y - x_0| \ge 2|x_0 - \widetilde{x}|$. This and Remark 2.8 then finish the proof of (i).

Next, we prove (ii). Indeed, let $B_1 := B(x_1, r_1)$ be a given ball of \mathbb{R}^n with $x_1 \in \mathbb{R}^n$ and $r_1 \in (0, \infty)$. For any $x \in \mathbb{R}^n$, choose a ball $B_2 \subset \mathbb{R}^n$ such that $x \in B_2$ and $(B_0 \cup B_1) \subset B_2$. Then, from the Lebesgue dominated convergence theorem, we deduce that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$, and almost every $x \in \mathbb{R}^n$,

$$\begin{split} \widetilde{T}_{B_{0}}(y^{\nu})(x) &= \int_{\mathbb{R}^{n} \setminus B_{2}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] y^{\nu} \, dy \\ &+ \lim_{\eta \to 0^{+}} \int_{B_{2} \setminus B(x, \eta)} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] y^{\nu} \, dy \\ &= \int_{\mathbb{R}^{n}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] y^{\nu} \, dy. \end{split}$$

For any $x \in \mathbb{R}^n$, let

$$F(x) := \int_{\mathbb{R}^n} \sum_{\{y \in \mathbb{Z}^{n_+} \mid y | \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_1, y)}{\gamma!} (x - x_1)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_1}(y) - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] y^{\nu} dy.$$

Using (2.5), (2.10), and an argument similar to the estimation of (2.22), we find that, for any $x \in \mathbb{R}^n$,

$$(2.23) |F(x)| \leq \int_{\mathbb{R}^{n} \setminus B_{2}} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} - \widetilde{K}(x, y) \right| |y|^{|\gamma|} dy$$

$$+ \int_{\mathbb{R}^{n} \setminus B_{2}} \left| \widetilde{K}(x, y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \right| |y|^{|\gamma|} dy$$

$$+ \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \int_{B_{2}} \left| \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right|$$

$$- \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right| |y^{\gamma}| dy$$

$$< \infty,$$

which, combined with Lemma 2.14, further implies that, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$,

$$(2.24) \widetilde{T}_{B_0}(y^{\nu}) - \widetilde{T}_{B_1}(y^{\nu}) \in \mathcal{P}_s(\mathbb{R}^n)$$

after changing values on a set of measure zero. This finishes the proof of (ii).

Now, we prove (iii). Indeed, from the definition of T, and (2.10), we deduce that, for any $a \in L^q_s(\mathbb{R}^n)$ supported in B := B(z, r), with $q \in (1, \infty)$, $z \in \mathbb{R}^n$, and $r \in (0, \infty)$, and for any $v \in \mathbb{Z}^n_+$ with $|v| \le s$,

$$(2.25) \int_{\mathbb{R}^{n}} T(a)(x)x^{\nu} dx$$

$$= \int_{\mathbb{R}^{n}} \lim_{\eta \to 0^{+}} \int_{B \setminus B(x,\eta)} K(x,y)a(y) dy x^{\nu} dx$$

$$= \int_{\mathbb{R}^{n}} \lim_{\eta \to 0^{+}} \int_{B \setminus B(x,\eta)} \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,z)}{\gamma!} (y-z)^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus 2B}(x) \right] a(y) dy x^{\nu} dx$$

$$= \int_{2B} \lim_{\eta \to 0^{+}} \int_{B \setminus B(x,\eta)} K(x,y)a(y) dy x^{\nu} dx$$

$$+ \int_{\mathbb{R}^{n} \setminus 2B} \int_{B} \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,z)}{\gamma!} (y-z)^{\gamma} \right] a(y) dy x^{\nu} dx$$

$$=: I_{1} + I_{2}.$$

Thus, to show (2.19), we only need to calculate I_1 and I_2 , respectively. We first consider I_1 . Indeed, for any $\eta \in (0, \infty)$ and $x, y \in \mathbb{R}^n$, let

(2.26)
$$K_{\eta}(x,y) := \begin{cases} K(x,y), & |x-y| \ge \eta, \\ 0, & \text{otherwise.} \end{cases}$$

Then, for any $x \in \mathbb{R}^n$.

$$T_{\eta}(a)(x) = \int_{B} K_{\eta}(x, y)a(y) \, dy,$$

where T_{η} is as in (2.8). By this, $a \in L^{q}(\mathbb{R}^{n})$, Remark 2.8, and the fact that $x^{\nu}\mathbf{1}_{2B} \in L^{q'}(\mathbb{R}^{n})$, we find that, for any $\nu \in \mathbb{Z}_{+}^{n}$ with $|\nu| \leq s$,

$$\left| \int_{2B} T_{\eta}(a)(x) \, x^{\nu} \, dx - \int_{2B} \lim_{\eta \to 0^{+}} T_{\eta}(a)(x) x^{\nu} \, dx \right|$$

$$\leq \int_{2B} \left| T_{\eta}(a)(x) - \lim_{\eta \to 0^{+}} T_{\eta}(a)(x) \right| \left| x^{\nu} \right| \, dx$$

$$\leq \left\| T_{\eta}(a) - \lim_{\eta \to 0^{+}} T_{\eta}(a) \right\|_{L^{q}(\mathbb{R}^{n})} \left(\int_{2B} \left| x^{\nu} \right|^{q'} \, dx \right)^{\frac{1}{q'}} \to 0$$

as $\eta \to 0^+$, which implies that

(2.27)
$$\lim_{\eta \to 0^+} \int_{2R} T_{\eta}(a)(x) \, x^{\nu} \, dx = \int_{2R} \lim_{\eta \to 0^+} T_{\eta}(a)(x) x^{\nu} \, dx.$$

From this and the Fubini theorem, we deduce that, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$,

(2.28)
$$I_{1} = \lim_{\eta \to 0^{+}} \int_{2B} \int_{B} K_{\eta}(x, y) a(y) \, dy \, x^{\nu} \, dx$$
$$= \lim_{\eta \to 0^{+}} \int_{B} a(y) \int_{2B} K_{\eta}(x, y) x^{\nu} \, dx \, dy = \int_{B} a(y) \lim_{\eta \to 0^{+}} \int_{2B} K_{\eta}(x, y) x^{\nu} \, dx \, dy,$$

where the proof of the last equality is similar to (2.27). This is a desired conclusion of I_1 . Now, we consider I_2 . To this end, we first show that, for any $\nu \in \mathbb{Z}_+^n$ with $|\nu| \leq s$,

$$(2.29) \qquad \widetilde{I}_{2} := \int_{\mathbb{R}^{n} \setminus 2B} \int_{B} \left| K(x, y) - \sum_{\{ \gamma \in \mathbb{Z}_{1}^{n} : |\gamma| \le s \}} \frac{\partial_{(2)}^{\gamma} K(x, z)}{\gamma!} (y - z)^{\gamma} \right| |a(y)| |x|^{|\gamma|} \, dy \, dx < \infty.$$

Indeed, by the Tonelli theorem, the Taylor remainder theorem, we find that, for any $y \in B$, there exists a $\widetilde{y} \in B$ such that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$,

$$\begin{split} \widetilde{\mathbf{I}}_{2} &= \int_{B} \int_{\mathbb{R}^{n} \backslash 2B} \left| K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \ |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,z)}{\gamma!} (y-z)^{\gamma} \right| |x|^{|\gamma|} \, dx \, |a(y)| \, dy \\ &= \int_{B} \int_{\mathbb{R}^{n} \backslash 2B} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \ |\gamma| = s\}} \frac{\partial_{(2)}^{\gamma} K(x,\overline{y}) - \partial_{(2)}^{\gamma} K(x,z)}{\gamma!} (y-z)^{\gamma} \right| |x|^{|\gamma|} \, dx \, |a(y)| \, dy \\ &\lesssim \int_{B} \int_{\mathbb{R}^{n} \backslash 2B} \frac{|\overline{y} - z|^{\delta} |y - z|^{s}}{|x - z|^{n+s+\delta}} |x|^{|\gamma|} \, dx \, |a(y)| \, dy \\ &\lesssim r^{s+\delta} ||a||_{L^{1}(B)} \int_{\mathbb{R}^{n} \backslash 2B} \frac{|x|^{|\gamma|}}{|x - z|^{n+s+\delta}} \, dx < \infty, \end{split}$$

where, in the penultimate step, we used (2.6) and (2.10) together with $|x - z| \ge 2r > 2|\tilde{y} - z|$. This implies that (2.29) holds true. Moreover, using (2.29) and the Fubini theorem, we find that

(2.30)
$$I_{2} = \int_{B} \int_{\mathbb{R}^{n} \setminus 2B} \left[K(x, y) - \sum_{\{ \gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \le s \}} \frac{\partial_{(2)}^{\gamma} K(x, z)}{\gamma!} (y - z)^{\gamma} \right] x^{\nu} dx \, a(y) \, dy.$$

This is a desired conclusion of I_2 .

Next, we show (2.19). Indeed, from (2.25), (2.28), (2.30), (2.10), and (2.24), we deduce that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$,

$$\int_{\mathbb{R}^{n}} T(a)(x)x^{\nu} dx$$

$$= \int_{B} a(y) \lim_{\eta \to 0^{+}} \int_{\mathbb{R}^{n} \backslash B(y,\eta)} \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,z)}{\gamma!} (y-z)^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \backslash 2B}(x) \right] x^{\nu} dx dy$$

$$= \int_{B} a(y) \lim_{\eta \to 0^{+}} \int_{\mathbb{R}^{n} \backslash B(y,\eta)} \left[\widetilde{K}(y,x) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,x)}{\gamma!} (y-z)^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \backslash 2B}(x) \right] x^{\nu} dx dy$$

$$= \int_{B} a(y) \widetilde{T}_{2B}(x^{\nu})(y) dy = \int_{B} a(y) \widetilde{T}_{B_{0}}(x^{\nu})(y) dy.$$

This shows that (2.19) holds true and hence finishes the proof of Lemma 2.15.

Proposition 2.16. Let $s \in \mathbb{Z}_+$, K be an s-order standard kernel, and T an s-order Calderón–Zygmund singular integral operator with kernel K. Let \widetilde{K} be as in (2.10), $B_0 := B(x_0, r_0) \subset \mathbb{R}^n$ a given ball with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$, and, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, $\widetilde{T}_{B_0}(y^v)$ as (2.18). Then (2.9) holds true if and only if, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, $\widetilde{T}_{B_0}(y^v)(\cdot) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero.

Proof. Let s, T, K, \widetilde{K} , and $B_0 := B(x_0, r_0)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$ be as in the present proposition. Now, we show the necessity. Indeed, if (2.9) holds true, then, using (1.1), we conclude that, for any $h \in L^2(\mathbb{R}^n)$ supported in a ball $B \subset \mathbb{R}^n$, $h - P_B^{(s)}(h) \in L_s^2(B)$. By this, (2.9) with a replaced by $[h - P_B^{(s)}(h)]\mathbf{1}_B$, (2.19) with a := 2 and a replaced by $[h - P_B^{(s)}(h)]\mathbf{1}_B$, and (1.1), we conclude that, for any a := 2 and a

(2.31)
$$0 = \int_{\mathbb{R}^{n}} T\left(\left[h - P_{B}^{(s)}(h)\right] \mathbf{1}_{B}\right)(x) x^{\nu} dx$$

$$= \int_{B} \left[h(x) - P_{B}^{(s)}(h)(x)\right] \widetilde{T}_{B_{0}}(y^{\nu})(x) dx$$

$$= \int_{B} \left[h(x) - P_{B}^{(s)}(h)(x)\right] \left[\widetilde{T}_{B_{0}}(y^{\nu})(x) - P_{B}^{(s)}\left(\widetilde{T}_{B_{0}}(y^{\nu})\right)(x)\right] dx$$

$$= \int_{B} h(x) \left[\widetilde{T}_{B_{0}}(y^{\nu})(x) - P_{B}^{(s)}\left(\widetilde{T}_{B_{0}}(y^{\nu})\right)(x)\right] dx.$$

Moreover, using Lemma 2.15(i), we have $\widetilde{T}_{B_0}(y^{\nu}) - P_B^{(s)}(\widetilde{T}_{B_0}(y^{\nu})) \in L^2(B)$. From this and (2.31), we deduce that, for almost every $x \in B$,

$$\widetilde{T}_{B_0}(y^{\nu})(x) = P_B^{(s)}\left(\widetilde{T}_{B_0}(y^{\nu})\right)(x).$$

Repeating the above procedure, we find that, for any $k \in \mathbb{N}$ and almost every $x \in 2^k B$,

$$\widetilde{T}_{B_0}(y^{\nu})(x) = P_{2^k B}^{(s)} \left(\widetilde{T}_{B_0}(y^{\nu}) \right)(x)$$

and hence, for almost every $x \in \mathbb{R}^n$,

$$\widetilde{T}_{B_0}(y^{\nu})(x) = P_B^{(s)}\left(\widetilde{T}_{B_0}(y^{\nu})\right)(x).$$

This finishes the proof of the necessity.

Moreover, the sufficiency immediately follows from (2.19). This finishes the proof of Proposition 2.16.

Remark 2.17. Proposition 2.16 might be well known but, to the best of our knowledge, we did not find a complete proof.

Corollary 2.18. Let $s \in \mathbb{Z}_+$, K be an s-order standard kernel, and T an s-order Calderón–Zygmund singular integral operator with kernel K. Then (2.9) holds true if and only if, for any $q \in (1, \infty)$, $a \in L^q_s(\mathbb{R}^n)$ having compact support, and $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$,

$$\int_{\mathbb{R}^n} T(a)(x) x^{\gamma} \, dx = 0.$$

Proof. Let s, K, and T be as in the present corollary. The sufficiency obviously holds true by taking q=2. Now, we show the necessity. By (2.9) and Proposition 2.16, we conclude that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $\widetilde{T}_{B_0}(y^{\gamma}) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero, where \widetilde{T}_{B_0} is as in Definition 2.11. Using this and (2.19), we have, for any $q \in (1, \infty)$, $a \in L_s^q(\mathbb{R}^n)$ having compact support, and $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$,

$$\int_{\mathbb{R}^n} T(a)(x) x^{\gamma} dx = \int_{\mathbb{R}^n} a(x) \widetilde{T}_{B_0}(y^{\gamma})(x) dx = 0.$$

This finishes the proof of the necessity and hence of Corollary 2.18.

The following conclusion is well known; we present the details here for the convenience of the reader.

Lemma 2.19. Let $s \in \mathbb{Z}_+$. Then there exists a positive constant C such that, for any ball (or cube) $B \subset \mathbb{R}^n$, any $P \in \mathcal{P}_s(\mathbb{R}^n)$, and any $\lambda \in (1, \infty)$,

where $\deg P$ denotes the degree of P.

Proof. Let $s \in \mathbb{Z}_+$ and, for any $x \in \mathbb{R}^n$, $P(x) := \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \le \deg P\}} a_\gamma x^\gamma$ with $\deg P \le s$ and $\{a_\gamma\}_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \le \deg P\}} \subset \mathbb{C}$. We only show the case of balls because the proof of the case of cubes is similar. Indeed, by the norm-equivalence theorem (see, for instance, [40, Corollary 3.3]), we find that

where the positive equivalence constants are independent of P but depend on s and n. Moreover, for any given ball $B(x_B, r_B) \subset \mathbb{R}^n$ with $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$, and for any $x \in \mathbb{R}^n$, let

$$\widetilde{P}(x) := \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \le \deg P\}} \widetilde{a}_{\gamma} x^{\gamma} := P(r_B x + x_B),$$

where $\{\widetilde{a}_{\gamma}\}_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq \deg P\}} \subset \mathbb{C}$. Then, using (2.33), we find that

$$||P||_{L^{\infty}(\lambda B(x_B, r_B))} = ||\widetilde{P}||_{L^{\infty}(\lambda B(\mathbf{0}, 1))} \lesssim \lambda^{\deg P} \max \{ [\widetilde{a}_{\gamma} | : \gamma \in \mathbb{Z}_+^n \text{ and } |\gamma| \leq \deg P \}$$
$$\sim \lambda^{\deg P} ||\widetilde{P}||_{L^{\infty}(B(\mathbf{0}, 1))} \sim \lambda^{\deg P} ||P||_{L^{\infty}(B(x_B, r_B))},$$

which implies that (2.32) holds true. This finishes the proof of Lemma 2.19.

By Lemma 2.19, we have the following two conclusions, which play vital roles in the proofs of Propositions 2.22 and 3.11 below.

Lemma 2.20. Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\theta \in (0, 2^{-s})$. Then there exists a positive constant C such that, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any ball (or cube) $B \subset \mathbb{R}^n$,

$$\sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} \left| f(x) - P_{B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}} \leq C \sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} \left| f(x) - P_{2^{k}B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}}.$$

Proof. Let q, s, and θ be as in the present lemma. We only show the case of balls because the proof of the case of cubes is similar. Obviously, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any ball $B \subset \mathbb{R}^n$, we have

(2.34)
$$\sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} \left| f(x) - P_{B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} \left| f(x) - P_{2^{k}B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \sum_{k=1}^{\infty} \theta^{k} \sum_{j=1}^{k} 2^{(k-j+1)s} \left\| P_{2^{j-1}B}^{(s)} \left(f - P_{2^{j}B}^{(s)}(f) \right) \right\|_{L^{\infty}(2^{j-1}B)}$$

$$=: \sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} \left| f(x) - P_{2^{k}B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}} + J.$$

From $P_B^{(s)}(P) = P$ for any $P \in \mathcal{P}_s(\mathbb{R}^n)$ and any ball $B \subset \mathbb{R}^n$, Lemma 2.19, (1.2), the Hölder inequality, the Fubini theorem, and the fact that $2^s \theta \in (0, 1)$, we deduce that

$$J \lesssim \sum_{k=1}^{\infty} \theta^{k} \sum_{j=1}^{k} 2^{(k-j+1)s} \int_{2^{j-1}B} \left| f(x) - P_{2^{j}B}^{(s)}(f)(x) \right| dx$$
$$\lesssim \sum_{k=1}^{\infty} \sum_{k=j}^{\infty} \theta^{k} 2^{(k-j+1)s} \int_{2^{j-1}B} \left| f(x) - P_{2^{j}B}^{(s)}(f)(x) \right| dx$$

$$\leq \sum_{j=1}^{\infty} \theta^{j} \int_{2^{j}B} \left| f(x) - P_{2^{j}B}^{(s)}(f)(x) \right| dx$$

$$\leq \sum_{j=1}^{\infty} \theta^{j} \left[\int_{2^{j}B} \left| f(x) - P_{2^{j}B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}}.$$

This and (2.34) imply the desired conclusion and hence finish the proof of Lemma 2.20.

Lemma 2.21. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, $\beta \in (s, \infty)$, and $\alpha \in (-\infty, \frac{1}{p} + \frac{\beta}{n})$. Then there exists a positive constant C such that, for any $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and any ball B(x,r) with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$,

$$\int_{\mathbb{R}^{n}\setminus B(x,r)} \frac{|f(y) - P_{B(x,r)}^{(s)}(f)(y)|}{|x - y|^{n+\beta}} \, dy \le C r^{-\frac{n}{p} - \beta + \alpha n} ||f||_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}.$$

Proof. Let p, q, s, β , and α be as in the present lemma. By Lemma 2.20 with $\theta := 2^{-\beta}$ [Here, we need $\beta \in (s, \infty)$], the Hölder inequality, and $\alpha \in (-\infty, \frac{1}{p} + \frac{\beta}{n})$, we conclude that, for any $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and any ball B(x,r) with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$,

(2.35)
$$\int_{\mathbb{R}^{n}\setminus B(x,r)} \frac{|f(y) - P_{B(x,r)}^{(s)}(f)(y)|}{|x - y|^{n+\beta}} dy$$

$$\sim \sum_{k \in \mathbb{N}} (2^{k}r)^{-n-\beta} \int_{B(x,2^{k}r)\setminus B(x,2^{k-1}r)} \left| f(y) - P_{B(x,r)}^{(s)}(f)(y) \right| dy$$

$$\lesssim \sum_{k \in \mathbb{N}} (2^{k}r)^{-\beta} \int_{B(x,2^{k}r)} \left| f(y) - P_{B(x,r)}^{(s)}(f)(y) \right| dy$$

$$\lesssim \sum_{k \in \mathbb{N}} (2^{k}r)^{-\beta} \int_{B(x,2^{k}r)} \left| f(y) - P_{B(x,2^{k}r)}^{(s)}(f)(y) \right| dy$$

$$\lesssim \sum_{k \in \mathbb{N}} (2^{k}r)^{-\beta} \left[\int_{B(x,2^{k}r)} \left| f(y) - P_{B(x,2^{k}r)}^{(s)}(f)(y) \right|^{q} dy \right]^{\frac{1}{q}},$$

which, together with $\alpha \in (-\infty, \frac{1}{p} + \frac{\beta}{n})$ and the definition of $\|\cdot\|_{JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$, further implies that

$$\int_{\mathbb{R}^{n}\backslash B(x,r)} \frac{|f(y) - P_{B(x,r)}^{(s)}(f)(y)|}{|x - y|^{n + \beta}} dy$$

$$\sim \sum_{k \in \mathbb{N}} (2^{k}r)^{-\frac{n}{p} - \beta + \alpha n} \left| B(x, 2^{k}r) \right|^{\frac{1}{p} - \alpha} \left[\int_{B(x, 2^{k}r)} \left| f(y) - P_{B(x, 2^{k}r)}^{(s)}(f)(y) \right|^{q} dy \right]^{\frac{1}{q}}$$

$$\lesssim r^{-\frac{n}{p} - \beta + \alpha n} ||f||_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}.$$

This finishes the proof of Lemma 2.21.

Proposition 2.22. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in (-\infty, \frac{1}{p} + \frac{s+\delta}{n})$ with $\delta \in (0, 1]$ as in Definition 2.6. Let K be an s-order standard kernel as in Definition 2.6 such that there exists an s-order Calderón–Zygmund singular integral operator T with kernel K having vanishing moments up to order s. Let \widetilde{K} be as in (2.10) and $B_0 := B(x_0, r_0)$ as in (2.12) with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$. Then, for any $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, $\widetilde{T}_{B_0}(f)$ in (2.11) is well defined almost everywhere on \mathbb{R}^n .

Proof. Let $p, q, s, \alpha, \delta, K, \widetilde{K}$, and $B_0 := B(x_0, r_0)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$ be as in the present proposition. Also, let $\widetilde{T}_{B_0,\eta}$ with $\eta \in (0,\infty)$ be as in (2.12). We only prove the case $p \in [1,\infty)$ because the proof of $p = \infty$ is similar. To this end, let $f \in JN^{\text{con}}_{(p,q,s)_\alpha}(\mathbb{R}^n)$ and $\widetilde{B} := B(z,r)$ be any given ball of \mathbb{R}^n with $z \in \mathbb{R}^n$ and $r \in (0,\infty)$. Then it is easy to see that, for any given $\eta \in (0,r)$ [and hence $B(x,\eta) \subset 2\widetilde{B}$] and for any $x \in \widetilde{B}$,

$$\widetilde{T}_{B_0,\eta}(f)(x) = E_{\widetilde{B},\eta}(x) + E_{\widetilde{B}}(x) + E_{\widetilde{B},B_0,\eta}^{(1)}(x) + E_{\widetilde{B},B_0,\eta}^{(2)}(x),$$

where

$$E_{\widetilde{B},\eta}(x) := \int_{2\widetilde{B}\setminus B(x,\eta)} \widetilde{K}(x,y) \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] dy,$$

$$E_{\widetilde{B}}(x) := \int_{\mathbb{R}^n \setminus 2\widetilde{B}} \left[\widetilde{K}(x,y) - \sum_{\{y \in \mathbb{Z}_1^n : \, |y| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} \right] \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] dy,$$

$$E_{\widetilde{B},B_0,\eta}^{(1)}(x):=\int_{\mathbb{R}^n\setminus B(x,\eta)}L_{\widetilde{B},B_0}(x,y)\left[f(y)-P_{2\widetilde{B}}^{(s)}(f)(y)\right]\,dy$$

with

$$L_{\widetilde{B},B_0}(x,y) := \sum_{\{\gamma \in \mathbb{Z}_+^n: \ |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus 2\widetilde{B}}(y)$$
$$- \sum_{\{\gamma \in \mathbb{Z}_+^n: \ |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0,y)}{\gamma!} (x-x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y),$$

and

$$E_{\widetilde{B},B_0,\eta}^{(2)}(x) := \int_{\mathbb{R}^n \setminus B(x,\eta)} \left[\widetilde{K}(x,y) - \sum_{\{\gamma \in \mathbb{Z}^n : |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0,y)}{\gamma!} (x - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] P_{2\widetilde{B}}^{(s)}(f)(y) \, dy.$$

We first consider $E_{B,\eta}(x)$ for almost every $x \in \widetilde{B}$. Indeed, since $f \in JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$, from the definition of $\|\cdot\|_{JN_{(p,a,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)}$, it is easy to deduce that

$$\left|2\widetilde{B}\right|^{\frac{1}{p}-\alpha}\left[\int_{2\widetilde{B}}\left|f(y)-P_{2\widetilde{B}}^{(s)}(f)(y)\right|^{q}dy\right]^{\frac{1}{q}}\lesssim \|f\|_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}<\infty,$$

which further implies that $[f - P_{2\widetilde{B}}^{(s)}(f)]\mathbf{1}_{2\widetilde{B}} \in L^q(\mathbb{R}^n)$, and hence $E_{\widetilde{B},\eta}(x)$ is well defined. Moreover, let $T_{(1)}$ be the *s*-order Calderón–Zygmund singular integral operator with kernel \widetilde{K} . Then, by Remark 2.8, we know that

(2.36)
$$\lim_{\eta \to 0^+} E_{\widetilde{B},\eta} = T_{(1)} \left(\left[f - P_{2\widetilde{B}}^{(s)}(f) \right] \mathbf{1}_{2\widetilde{B}} \right)$$

almost everywhere on \widetilde{B} .

Next, we estimate $E_{\widetilde{B}}$. Noticing that $E_{\widetilde{B}}$ is independent of η , we only need to show that $E_{\widetilde{B}}$ is bounded on \widetilde{B} . Indeed, using the Taylor remainder theorem, we find that, for any $x \in \widetilde{B}$, there exists an $\widetilde{x} \in \widetilde{B}$ such that

$$(2.37) \left| E_{\widetilde{B}}(x) \right| \leq \int_{\mathbb{R}^{n} \setminus 2\widetilde{B}} \left| \widetilde{K}(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : \, |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} \right| \left| f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right| dy$$

$$\lesssim \int_{\mathbb{R}^{n} \setminus 2\widetilde{B}} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : \, |\gamma| = s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(\widetilde{x},y) - \partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} \right| \left| f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right| dy$$

$$\lesssim |z - \widetilde{x}|^{\delta} |x - z|^{s} \int_{\mathbb{R}^{n} \setminus 2\widetilde{B}} \frac{|f(y) - P_{2\widetilde{B}}^{(s)}(f)(y)|}{|y - z|^{n+s+\delta}} dy,$$

where, in the last inequality, we used (2.6) and (2.10) together with $|y - z| \ge 2r \ge |\widetilde{x} - z|$. From this, Lemma 2.21 with $\beta := s + \delta \in (s, \infty)$, and $\alpha \in (-\infty, \frac{1}{p} + \frac{s + \delta}{n})$, we deduce that, for any $x \in \widetilde{B}$,

$$\left| E_{\widetilde{B}}(x) \right| \lesssim r^{-\frac{n}{p} + \alpha n} ||f||_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)} < \infty.$$

This further implies that $E_{\widetilde{B}}(x)$ is well defined for any $x \in \widetilde{B}$.

Now, we estimate $E_{\widetilde{B},B_0,\eta}^{(1)}$. To this end, let $R:=3|z-x_0|+2r+r_0$. It is easy to see that

$$(2.38) (2\widetilde{B} \cup B) \subset B(z,R)$$

and, for any $y \in \mathbb{R}^n \setminus B(z, R)$,

$$(2.39) |y - x_0| \le |y - z| + |z - x_0| < 2|y - z|.$$

Moreover, if $x \in \widetilde{B}$ and $y \in B(z, R)$, then, by (2.5) and (2.10), we have

$$|L_{\widetilde{B},B_{0}}(x,y)| \leq \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus 2\widetilde{B}}(y) \right|$$

$$+ \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x-x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right|$$

$$\lesssim \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{|x-z|^{|\gamma|}}{r^{n+|\gamma|}} + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \frac{|x-x_{0}|^{|\gamma|}}{r_{0}^{n+|\gamma|}}$$

$$\lesssim \frac{1}{r^n} + \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \leq s\}} \frac{(r + |z - x_0|)^{|\gamma|}}{r_0^{n + |\gamma|}} \lesssim 1,$$

where the implicit positive constants depend on r, z, and x_0 , but are independent of η . If $x \in \widetilde{B}$ and $y \in \mathbb{R}^n \setminus B(z, R)$, then, using (2.38), the Taylor remainder theorem, and (2.6) and (2.10) together with (2.39), we find that there exist a $z_1 \in \widetilde{B}$ and a $z_2 \in B(x_0, |z - x_0| + r)$ such that

$$\begin{aligned} |L_{\widetilde{B},B_{0}}(x,y)| &= \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x-x_{0})^{\gamma} \right| \\ &\leq \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y)}{\gamma!} (x-z)^{\gamma} - \widetilde{K}(x,y) \right| \\ &+ \left| \widetilde{K}(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x-x_{0})^{\gamma} \right| \\ &\sim \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| = s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z,y) - \partial_{(1)}^{\gamma} \widetilde{K}(z_{1},y)}{\gamma!} (x-z)^{\gamma} \right| \\ &+ \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| = s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(z_{2},y) - \partial_{(1)}^{\gamma} \widetilde{K}(x_{0},y)}{\gamma!} (x-x_{0})^{\gamma} \right| \\ &\lesssim \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| = s\}} \frac{|z-z_{1}|^{\delta} |x-z|^{|\gamma|}}{|y-z|^{n+|\gamma|+\delta}} + \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| = s\}} \frac{|z_{2}-x_{0}|^{\delta} |x-x_{0}|^{|\gamma|}}{|y-z|^{n+|\gamma|+\delta}} \\ &\lesssim \frac{r^{s+\delta} + (r+|z-x_{0}|)^{s+\delta}}{|y-z|^{n+s+\delta}} \sim \frac{1}{|y-z|^{n+s+\delta}}, \end{aligned}$$

where the implicit positive constants depend on r, z, and x_0 , but are independent of η . From this, (2.40), (2.41), and Lemma 2.21 with $\beta := s + \delta \in (s, \infty)$ and $\alpha \in (-\infty, \frac{1}{p} + \frac{s+\delta}{n})$, we deduce that, for any $x \in \widetilde{B}$,

$$\begin{aligned} \left| E_{\widetilde{B},B_{0},\eta}^{(1)}(x) \right| &\leq \int_{\mathbb{R}^{n} \setminus B(x,\eta)} \left| L_{\widetilde{B},B_{0}}(x,y) \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] \right| dy \\ &\lesssim \int_{B(z,R)} \left| f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right| dy + \int_{\mathbb{R}^{n} \setminus B(z,R)} \frac{|f(y) - P_{2\widetilde{B}}^{(s)}(f)(y)|}{|y - z|^{n+s+\delta}} dy \\ &\lesssim \int_{B(z,R)} \left| f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right| dy + \int_{\mathbb{R}^{n} \setminus 2\widetilde{B}} \frac{|f(y) - P_{2\widetilde{B}}^{(s)}(f)(y)|}{|y - z|^{n+s+\delta}} dy \\ &\lesssim \left\| f - P_{2\widetilde{B}}^{(s)}(f) \right\|_{L^{1}(B(z,R))} + \|f\|_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} < \infty, \end{aligned}$$

where the implicit positive constants depend on r, z, and x_0 , but are independent of η . Using (2.42) and the Lebesgue dominated convergence theorem, we conclude that

$$P_1(\widetilde{B}, B_0; f)(\cdot) := \lim_{\eta \to 0^+} E_{\widetilde{B}, B_0, \eta}^{(1)}(\cdot) = \int_{\mathbb{R}^n} L_{\widetilde{B}, B_0}(\cdot, y) \left[f(y) - P_{\widetilde{B}}^{(s)}(f)(y) \right] dy$$

is finite everywhere on \widetilde{B} , which, combined with Lemma 2.14, further implies that

$$(2.43) P_1(\widetilde{B}, B_0; f) \in \mathcal{P}_s(B).$$

Finally, it remains to estimate $E_{\overline{B},B_0,\eta}^{(2)}$. Indeed, by Proposition 2.16, we find that there exists a polynomial $P_2(\widetilde{B},B_0;f) \in \mathcal{P}_s(\mathbb{R}^n)$ such that, for almost every $x \in \widetilde{B}$,

(2.44)
$$\lim_{n \to 0^+} E_{\overline{B}, B_0, \eta}^{(2)}(x) = P_2(\widetilde{B}, B_0; f)(x).$$

This is a desired estimate of $E_{\widetilde{B},B_0,\eta}^{(2)}$.

From (2.36), (2.37), (2.43), and (2.44), we deduce that, for any given ball $\widetilde{B} \subset \mathbb{R}^n$ and for almost every $x \in \widetilde{B}$,

$$(2.45) \widetilde{T}_{B_0}(f)(x) = \lim_{\eta \to 0^+} \widetilde{T}_{B_0,\eta}(f)(x)$$

$$= T_{(1)} \left(\left[f - P_{2\widetilde{B}}^{(s)}(f) \right] \mathbf{1}_{2\widetilde{B}} \right)(x) + E_{\widetilde{B}}(x) + P_1(\widetilde{B}, B_0; f)(x) + P_2(\widetilde{B}, B_0; f)(x).$$

Therefore, $\widetilde{T}_{B_0}(f)$ is well defined almost everywhere on \mathbb{R}^n . This finishes the proof of Proposition 2.22.

Remark 2.23. Let all the symbols be as in Proposition 2.22. We now claim that, for any given ball $B_1 \subset \mathbb{R}^n$,

$$\widetilde{T}_{B_0}(f)-\widetilde{T}_{B_1}(f)\in\mathcal{P}_s(\mathbb{R}^n)$$

after changing values on a set of measure zero. Indeed, let $B_1 := B(x_1, r_1)$ be any given ball of \mathbb{R}^n with $x_1 \in \mathbb{R}^n$ and $r_1 \in (0, \infty)$. Then, for almost every $x \in \mathbb{R}^n$, choosing a ball $\widetilde{B} \subset \mathbb{R}^n$ such that $x \in \widetilde{B}$ and $(B_0 \cup B_1) \subset \widetilde{B}$, by the Lebesgue dominated convergence theorem, we obtain

$$(2.46) \qquad \widetilde{T}_{B_{0}}(f)(x) - \widetilde{T}_{B_{1}}(f)(x)$$

$$= \int_{\mathbb{R}^{n} \setminus \widetilde{B}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right]$$

$$- \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] f(y) dy$$

$$+ \lim_{\eta \to 0^{+}} \int_{\widetilde{B} \setminus B(x, \eta)} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right]$$

$$- \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right] f(y) dy$$

$$= \int_{\mathbb{R}^{n}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right]$$

$$-\frac{\partial_{(1)}^{\gamma}\widetilde{K}(x_0,y)}{\gamma!}(x-x_0)^{\gamma}\mathbf{1}_{\mathbb{R}^n\setminus B_0}(y)\bigg]f(y)\,dy.$$

For any $x \in \mathbb{R}^n$, let

$$D(x) := \int_{\mathbb{R}^n} \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \le s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_1, y)}{\gamma!} (x - x_1)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_1}(y) - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] f(y) \, dy.$$

By (2.46), an argument similar to the estimation of (2.42), and (2.23), we conclude that, for any $x \in \mathbb{R}^n$,

$$|D(x)| \leq \int_{\mathbb{R}^{n} \setminus \widetilde{B}} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right] \right| dy$$

$$- \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right| \left| f(y) - P_{2B_{1}}^{(s)}(f)(y) \right| dy$$

$$+ \left| \int_{\mathbb{R}^{n} \setminus \widetilde{B}} \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \left[\frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right] - \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right| P_{2B_{1}}^{(s)}(f)(y) dy$$

$$+ \sum_{\{\gamma \in \mathbb{Z}_{+}^{n} : |\gamma| \leq s\}} \int_{\widetilde{B}} \left| \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{1}, y)}{\gamma!} (x - x_{1})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{1}}(y) \right|$$

$$- \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_{0}, y)}{\gamma!} (x - x_{0})^{\gamma} \mathbf{1}_{\mathbb{R}^{n} \setminus B_{0}}(y) \right| |f(y)| dy$$

$$< \infty,$$

which, together with Lemma 2.14, further implies that $\widetilde{T}_{B_0}(f) - \widetilde{T}_{B_1}(f) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero. This finishes the proof of the above claim. Based on this claim, in what follows, we write \widetilde{T} instead of \widetilde{T}_{B_0} if there exists no confusion.

The following conclusion is just [17, Proposition 2.2], which gives an equivalent characterization of $JN^{\text{con}}_{(p,q,s)_a}(\mathbb{R}^n)$.

Lemma 2.24. Let $p, q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ if and only if $f \in L^q_{\text{loc}}(\mathbb{R}^n)$ and

$$[f]_{JN_{(p,q,s)\alpha}^{\text{con}}(\mathbb{R}^n)} := \sup_{r \in (0,\infty)} \left[\int_{\mathbb{R}^n} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} \left| f(x) - P_{B(y,r)}^{(s)}(f)(x) \right|^q dx \right]^{\frac{1}{q}} \right\}^p dy \right]^{\frac{1}{p}} < \infty.$$

Moreover, for any $f \in JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$,

$$||f||_{JN_{(p,q,s)\alpha}^{\mathrm{con}}(\mathbb{R}^n)} \sim [f]_{JN_{(p,q,s)\alpha}^{\mathrm{con}}(\mathbb{R}^n)},$$

where the positive equivalence constants are independent of f.

Next, we show the boundedness of \widetilde{T} on $JN^{\text{con}}_{(p,q,s)_q}(\mathbb{R}^n)$.

Theorem 2.25. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in (-\infty, \frac{s+\delta}{n})$ with $\delta \in (0, 1]$ as in Definition 2.6. Let K be an s-order standard kernel as in Definition 2.6 and T the s-order Calderón–Zygmund singular integral operator with kernel K. Let \widetilde{K} be as in (2.10) and \widetilde{T} an s-order modified Calderón–Zygmund operator with kernel \widetilde{K} . Then \widetilde{T} is bounded on $JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$,

$$\left\|\widetilde{T}(f)\right\|_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \le C\|f\|_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}$$

if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$.

Proof. Let p, q, s, α, δ , and \widetilde{T} be as in the present theorem. We only prove the case $p \in [1, \infty)$ because the proof of the case $p = \infty$ is similar. We first show the sufficiency. Indeed, using (1.2), it is easy to see that, for any $g \in L^1_{loc}(\mathbb{R}^n)$ and any ball B of \mathbb{R}^n ,

$$\left[\int_{B} \left| g(x) - P_{B}^{(s)}(g)(x) \right|^{q} dx \right]^{\frac{1}{q}} \sim \inf_{P \in \mathcal{P}_{s}(B)} \left[\int_{B} \left| g(x) - P(x) \right|^{q} dx \right]^{\frac{1}{q}},$$

where $\mathcal{P}_s(B)$ denotes the set of all polynomials with degree not greater than s on B (see, for instance, [17, (2.12)]). From this, (2.45), the Minkowski inequality, and $P_{B(z,r)}^{(s)}(P) = P$ for any $P \in \mathcal{P}_s(\mathbb{R}^n)$, we deduce that, for any given $f \in JN_{(p,q,s)_\alpha}^{\text{con}}(\mathbb{R}^n)$ and for any $r \in (0,\infty)$,

$$(2.47) \qquad \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,r)} \left| \widetilde{T}(f)(x) - P_{B(z,r)}^{(s)}(\widetilde{T}(f))(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$\lesssim \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,r)} \left| T_{(1)} \left(\left[f - P_{2B(z,r)}^{(s)}(f) \right] \mathbf{1}_{2B(z,r)} \right)(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$+ \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,r)} \left| E_{B(z,r)}(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$\sim F_{1} + F_{2},$$

where $T_{(1)}$ and $E_{B(z,r)}$ are as in (2.45),

$$F_{1} := \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,r)} \left| T_{(1)} \left(\left[f - P_{2B(z,r)}^{(s)}(f) \right] \mathbf{1}_{2B(z,r)} \right)(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}},$$

and

$$F_{2} := \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,r)} \left| E_{B(z,r)}(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}.$$

We now estimate F_1 . Indeed, by the fact that $[f - P_{2B(z,r)}^{(s)}(f)]\mathbf{1}_{2B(z,r)} \in L^q(\mathbb{R}^n)$, and Lemmas 2.13 and 2.24, we find that

$$(2.48) F_{1} \lesssim \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{2B(z,r)} \left| f(x) - P_{2B(z,r)}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}} \lesssim ||f||_{JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}.$$

This is a desired estimate of F_1 .

Next, we estimate F_2 . Indeed, using (2.37), (2.35), the Minkowski inequality, Lemma 2.24, and $\alpha \in (-\infty, \frac{s+\delta}{n})$, we find that

$$F_{2} \lesssim \left\{ \int_{\mathbb{R}^{n}} \left[|B(z,r)|^{-\alpha} r^{s+\delta} \int_{\mathbb{R}^{n} \setminus 2B(z,r)} \frac{|f(y) - P_{2B(z,r)}^{(s)}(f)(y)|}{|y - z|^{n+s+\delta}} dy \right]^{p} dz \right\}^{\frac{1}{p}}$$

$$\lesssim \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \sum_{k \in \mathbb{N}} \frac{1}{2^{k(s+\delta)}} \left[\int_{B(z,2^{k+1}r)} \left| f(y) - P_{B(z,2^{k+1}r)}^{(s)}(f)(y) \right|^{q} dy \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$\lesssim \sum_{k \in \mathbb{N}} \frac{1}{2^{k(s+\delta)}} \left[\int_{\mathbb{R}^{n}} \left\{ |B(z,r)|^{-\alpha} \left[\int_{B(z,2^{k+1}r)} \left| f(y) - P_{B(z,2^{k+1}r)}^{(s)}(f)(y) \right|^{q} dy \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$\sim \sum_{k \in \mathbb{N}} \frac{1}{2^{k(s+\delta-n\alpha)}} \left[\int_{\mathbb{R}^{n}} \left\{ \left| B(z,2^{k+1}r) \right|^{-\alpha} \left[\int_{B(z,2^{k+1}r)} \left| f(y) - P_{B(z,2^{k+1}r)}^{(s)}(f)(y) \right|^{q} dy \right]^{\frac{1}{q}} \right\}^{p} dz \right]^{\frac{1}{p}}$$

$$\lesssim ||f||_{JN_{(p,a,s)_{p}}^{con}(\mathbb{R}^{n})}.$$

From this, (2.47), (2.48), and Lemma 2.24, we deduce that, for any $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, $\widetilde{T}(f) \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and

$$\|\widetilde{T}(f)\|_{JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})} \lesssim \|f\|_{JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})}.$$

This finishes the proof of the sufficiency.

We now show the necessity. Indeed, if \widetilde{T} is bounded on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})$, then, for any $\gamma \in \mathbb{Z}_{+}^{n}$ with $|\gamma| \leq s$,

$$\left\|\widetilde{T}(x^{\gamma})\right\|_{JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})} \lesssim \left\|x^{\gamma}\right\|_{JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})} \sim 0$$

and hence $\widetilde{T}(x^{\gamma}) \in \mathcal{P}_s(\mathbb{R}^n)$. Using this and Proposition 2.16, we find that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$, which completes the proof of the necessity and hence of Theorem 2.25. \square

Remark 2.26. (i) We should point out that Peetre [36, Remark 1.5] claimed, without giving a proof, that the Calderón–Zygmund operator on the generalized Campanato space $L_s^{q,\Phi}$, with $s \in \mathbb{N}$, which when $\Phi(r) := r^{\alpha pn+n}$ coincides with $C_{\alpha,q,s}(\mathbb{R}^n)$, is bounded under some unspecific assumptions on the kernel of the Calderón–Zygmund operator. It is doubtful whether or not this claim holds true for the Riesz transforms.

(ii) By Proposition 2.16, we further conclude that Theorem 2.25 coincides with [22, Theorem 4.21] when $p = \infty$.

3 Calderón–Zygmund Operators on Hardy-Type Spaces

This section is devoted to studying the boundedness of Calderón–Zygmund operators on the Hardy-kind space $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ which proves the predual space of $JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ (see, for instance, [17, Theorem 4.10]). To overcome the difficulty caused by the fact that $\|\cdot\|_{HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ is no longer concave, we establish a criterion for the boundedness of linear operators on $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ via using molecules of $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and the dual relation $(HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n))^* = JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$.

3.1 Hardy-Type Spaces

We first recall the notion of the Hardy-kind space $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ which was introduced in [17, Section 4].

Definition 3.1. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. A measurable function a on \mathbb{R}^n is called a $(p, q, s)_{\alpha}$ -atom supported in a cube $Q \subset \mathbb{R}^n$ if

- (i) $\{x \in \mathbb{R}^n : a(x) \neq 0\} \subset Q$;
- (ii) $||a||_{L^q(Q)} \le |Q|^{\frac{1}{q} \frac{1}{p} \alpha}$;
- (iii) $\int_O a(x)x^{\gamma} dx = 0$ for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$.

The following conclusion is just [17, Proposition 4.3].

Lemma 3.2. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, $\alpha \in \mathbb{R}$, a be a $(p, q, s)_{\alpha}$ -atom supported in the cube $Q \subset \mathbb{R}^n$, and, for any $f \in JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$,

$$\langle a, f \rangle := \int_{Q} f(x)a(x) dx,$$

where $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$. Then $\langle a, \cdot \rangle \in (JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$, where $(JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$ denotes the dual space of $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, which is defined to be the set of all continuous linear functionals on $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ equipped with the weak-* topology.

Now, we say $(p,q,s)_{\alpha}$ -atom $a \in (JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$ instead of $(a,\cdot) \in (JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$. Then we have the following notion of congruent $(p,q,s)_{\alpha}$ -polymers. Recall that, for any $\ell \in (0,\infty)$, $\Pi_{\ell}(\mathbb{R}^n)$ denotes the class of all collections of interior pairwise disjoint subcubes $\{Q_j\}_j$ of \mathbb{R}^n with side length ℓ .

Definition 3.3. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. The space of congruent $(p, q, s)_{\alpha}$ -polymers, $\widetilde{HK}_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^n)$, is defined to be the set of all $g \in (JN_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^n))^*$ satisfying that there exist an $\ell \in (0, \infty)$, $(p, q, s)_{\alpha}$ -atoms $\{a_j\}_j$ supported, respectively, in $\{Q_j\}_j \in \Pi_{\ell}(\mathbb{R}^n)$, and $\sum_j |\lambda_j|^p < \infty$ such that $g = \sum_j \lambda_j a_j$ in $(JN_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^n))^*$, where $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$. Moreover, any

 $g \in \widetilde{HK}^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ is called a *congruent* $(p,q,s)_{\alpha}$ -polymer with its norm $\|g\|_{\widetilde{HK}^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ defined by setting

$$||g||_{\widetilde{HK}_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n})} := \inf \left(\sum_{j} |\lambda_{j}|^{p} \right)^{\frac{1}{p}},$$

where the infimum is taken over all decompositions of g as above.

Definition 3.4. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. The *Hardy-type space via congruent cubes*, $HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$, is defined by setting

$$HK_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n}) := \left\{ g \in (JN_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n}))^{*} : g = \sum_{i} g_{i} \text{ in } (JN_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n}))^{*}, \right.$$

$$\left. \{g_{i}\}_{i} \subset \widetilde{HK}_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n}), \text{ and } \sum_{i} \|g_{i}\|_{\widetilde{HK}_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n})} < \infty \right\},$$

where $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$. Moreover, for any $g \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, let

$$||g||_{HK^{\operatorname{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^{n})} := \inf \sum_{i} ||g_{i}||_{\widetilde{HK}^{\operatorname{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^{n})},$$

where the infimum is taken over all decompositions of g as above.

Definition 3.5. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. The *finite atomic Hardy-type space* via congruent cubes, $HK_{(p,q,s)_{\alpha}}^{\text{con-fin}}(\mathbb{R}^n)$, is defined to be the set of all

$$g = \sum_{j=1}^{M} \lambda_j a_j$$

pointwisely, where $M \in \mathbb{N}$, $\{a_j\}_{j=1}^M$ are $(p,q,s)_{\alpha}$ -atoms supported, respectively, in cubes $\{Q_j\}_{j=1}^M$ of \mathbb{R}^n , and $\{\lambda_j\}_{j=1}^M \subset \mathbb{C}$.

The following conclusion is just [17, Propositions 4.9].

Lemma 3.6. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then $HK^{\text{con-fin}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ is dense in $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ in the sense that, for any $g \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, there exists a sequence $\{g_m\}_{m \in \mathbb{N}}$ of $HK^{\text{con-fin}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ such that

$$\lim_{m\to\infty} \|g-g_m\|_{HK^{\text{con}}_{(p,q,s)\alpha}(\mathbb{R}^n)} = 0.$$

The following conclusion is an immediate corollary of [17, Theorem 4.10] and [17, Proposition 2.7]; we omit the details here.

Lemma 3.7. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then

$$\left(HK_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n})\right)^{*} = JN_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n})$$

with equivalent norms in the following sense:

(i) any given $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ induces a linear functional \mathcal{L}_f given by setting, for any $g \in HK^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ and $\{g_i\}_i \subset \widetilde{HK}^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ with $g = \sum_i g_i \operatorname{in}(JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n))^*$,

(3.1)
$$\mathcal{L}_f(g) := \langle g, f \rangle = \sum_i \langle g_i, f \rangle.$$

Moreover, for any $g \in HK^{\text{con-fin}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, $\mathcal{L}_f(g) = \int_{\mathbb{R}^n} f(x)g(x) dx$ and there exists a positive constant C such that

$$\left\| \mathcal{L}_f \right\|_{(HK_{(p,q,s)_n}^{\text{con}}(\mathbb{R}^n))^*} \le C \|f\|_{JN_{(p,q,s)_n}^{\text{con}}(\mathbb{R}^n)};$$

(ii) conversely, for any continuous linear functional \mathcal{L} on $HK^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, there exists a unique $f \in JN^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ such that, for any $g \in HK^{\text{con-fin}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, $\mathcal{L}(g) = \int_{\mathbb{R}^n} f(x)g(x) dx$ and there exists a positive constant C such that

$$||f||_{JN_{(p,q,s)\alpha}^{\text{con}}(\mathbb{R}^n)} \le C||\mathcal{L}||_{(HK_{(p',q',s)\alpha}^{\text{con}}(\mathbb{R}^n))^*}.$$

The following conclusion can immediately be deduced from [17, Corollary 4.11], which plays a key role in the proof of Theorem 3.20 below; we omit the details here.

Proposition 3.8. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then $\|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ and $\|\cdot\|_{(JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*}$ are equivalent norms of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$.

Now, we show that $(HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n), \|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)})$ is a Banach space.

Proposition 3.9. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then

$$\left(HK_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n}), \|\cdot\|_{HK_{(p,q,s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^{n})}\right)$$

is a Banach space.

Proof. Let p, q, s, and α be as in the present proposition. Since [17, Proposition 4.8] proves that $(HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n), \|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)})$ is a linear normed space, from this, we deduce that, to prove the present proposition, it suffices to show that it is complete. To this end, let $\{g^{(m)}\}_{m\in\mathbb{N}}$ be any given Cauchy sequence of $(HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n), \|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)})$, and $g^{(0)}$ the zero element of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. Without loss of generality, we may assume that, for any $m \in \mathbb{N}$,

(3.2)
$$\|g^{(m)} - g^{(m-1)}\|_{HK_{(n,q,s)_{n}}^{\text{con}}(\mathbb{R}^{n})} < \frac{1}{2^{m}}.$$

By $HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)\subset (JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$, and Proposition 3.8, we conclude that $\{g^{(m)}\}_{m\in\mathbb{N}}$ is also a Cauchy sequence of $(HK^{\mathrm{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n),\|\cdot\|_{(JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*})$ and hence, from the completeness of $(JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$, we deduce that there exists an element $g\in (JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n))^*$ such that

(3.3)
$$\lim_{N \to \infty} \|g - g^{(N)}\|_{(JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n}))^{*}} = 0$$

and, for any $f \in JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$,

(3.4)
$$\langle g, f \rangle = \lim_{N \to \infty} \left\langle g^{(N)}, f \right\rangle.$$

Next, we show that $g \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. Indeed, using (3.2) and the definition $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, we find that, for any $m \in \mathbb{N}$, there exists a sequence $\{g_i^{(m)}\}_{i \in \mathbb{N}}$ of $\widetilde{HK}^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ such that

(3.5)
$$g^{(m)} - g^{(m-1)} = \sum_{i \in \mathbb{N}} g_i^{(m)}$$

in $(JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^n))^*$, and

Moreover, from (3.2), we deduce that, for any $f \in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ and $N \in \mathbb{N}$,

$$\left\| \sum_{m=1}^{N} \left[g^{(m)} - g^{(m-1)} \right] \right\|_{(JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n}))^{*}}$$

$$\sim \left\| \sum_{m=1}^{N} \left[g^{(m)} - g^{(m-1)} \right] \right\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \lesssim \sum_{m \in \mathbb{N}} \left\| g^{(m)} - g^{(m-1)} \right\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \lesssim 1.$$

Using this and the completeness of $(JN_{(p',q',s)_a}^{\text{con}}(\mathbb{R}^n))^*$, we find that

$$\sum_{m \in \mathbb{N}} \left[g^{(m)} - g^{(m-1)} \right] \in \left(JN_{(p',q',s)_{\alpha}}^{\operatorname{con}}(\mathbb{R}^n) \right)^*,$$

which, together with (3.4) and (3.5), implies that, for any $f \in JN_{(p',q',s)_a}^{\text{con}}(\mathbb{R}^n)$,

$$\langle g, f \rangle = \lim_{N \to \infty} \left\langle g^{(N)}, f \right\rangle = \lim_{N \to \infty} \left\langle \sum_{m=1}^{N} \left[g^{(m)} - g^{(m-1)} \right], f \right\rangle$$
$$= \left\langle \sum_{m \in \mathbb{N}} \left[g^{(m)} - g^{(m-1)} \right], f \right\rangle = \left\langle \sum_{m \in \mathbb{N}} \sum_{i \in \mathbb{N}} g_i^{(m)}, f \right\rangle.$$

Thus, $g = \sum_{m \in \mathbb{N}} \sum_{i \in \mathbb{N}} g_i^{(m)}$ in $(JN_{(p',q',s)_\alpha}^{\text{con}}(\mathbb{R}^n))^*$, which, combined with (3.6), further implies that $g \in HK_{(p,q,s)_\alpha}^{\text{con}}(\mathbb{R}^n)$. By this, (3.3), and Lemma 3.8, we find that

$$\lim_{N \to \infty} \left\| g - g^{(N)} \right\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)} = 0$$

and hence completes the proof of Proposition 3.9.

3.2 Molecules of $HK^{\text{con}}_{(p,q,s)_a}(\mathbb{R}^n)$

In this subsection, we introduce the notion of $(p, q, s, \alpha, \epsilon)$ -molecules of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, which plays an important role in the proof of the boundedness of Calderón–Zygmund operators on $HK^{\text{con}}_{(p,q,s)}(\mathbb{R}^n)$.

Via borrowing some ideas from [15, Definition 1.2], we introduce the following $(p, q, s, \alpha, \epsilon)$ molecule. In what follows, for any $z \in \mathbb{R}^n$ and $r \in (0, \infty)$, we denote by $Q_z(r)$ the cube with center z and side length r.

Definition 3.10. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, $\alpha \in (\frac{1}{q} - \frac{1}{p}, \infty)$, and $\epsilon \in (0, \infty)$. A measurable function M on \mathbb{R}^n is called a $(p, q, s, \alpha, \epsilon)$ -molecule centered at the cube $Q_z(r)$ with center $z \in \mathbb{R}^n$ and length $r \in (0, \infty)$ if

- (i) $||M\mathbf{1}_{Q_{z}(r)}||_{L^{q}(\mathbb{R}^{n})} \leq |Q_{z}(r)|^{\frac{1}{q} \frac{1}{p} \alpha};$
- (ii) for any $j \in \mathbb{N}$, $||M\mathbf{1}_{Q_{z}(2^{j}r)\setminus Q_{z}(2^{j-1}r)}||_{L^{q}(\mathbb{R}^{n})} \le 2^{\frac{jn}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)}|Q_{z}(r)|^{\frac{1}{q}-\frac{1}{p}-\alpha}$;
- (iii) $\int_{\mathbb{R}^n} M(x) x^{\gamma} dx = 0$ for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$.

Now, we show that any $(p, q, s, \alpha, \epsilon)$ -molecule induces an element of $(JN_{(p', q', s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n}))^{*}$.

Proposition 3.11. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$, $s \in \mathbb{Z}_+$, $\alpha \in (\frac{1}{q} - \frac{1}{p}, \infty)$, and $\epsilon \in (0, 1)$ be such that $\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n} < 0$. Let M be a $(p, q, s, \alpha, \epsilon)$ -molecule. Then, for any $f \in JN^{\text{con}}_{(p', q', s)_{\alpha}}(\mathbb{R}^n)$,

$$\langle M, f \rangle := \int_{\mathbb{R}^n} f(x) M(x) \, dx$$

induces a continuous linear functional on $JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$.

Proof. Let p, q, s, α , and ϵ be as in the present proposition. Let M be a $(p, q, s, \alpha, \epsilon)$ -molecule centered at the cube $Q_z(r)$ with $z \in \mathbb{R}^n$ and $r \in (0, \infty)$. Let $L_0 := Q_z(r)$ and, for any $j \in \mathbb{N}$, let $L_j := Q_z(2^j r) \setminus Q_z(2^{j-1} r)$. Then, from Definition 3.10(iii), the Hölder inequality, Definition 3.10(ii), Lemma 2.20 with

$$\theta := 2^{\frac{n}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{n}{q'}} \in (0, 2^{-s}),$$

 $\frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)+\frac{1}{q'}+\frac{s}{n}<0, \text{ and } \alpha\in(\frac{1}{q}-\frac{1}{p},\infty), \text{ we deduce that, for any } f\in JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^{n}),$

$$\begin{split} & \left| \int_{\mathbb{R}^{n}} M(x) f(x) \, dx \right| \\ & = \left| \int_{\mathbb{R}^{n}} M(x) \left[f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right] \, dx \right| \\ & \leq \sum_{j \in \mathbb{Z}_{+}} \int_{L_{j}} \left| M(x) \left[f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right] \right| \, dx \\ & \leq \sum_{j \in \mathbb{Z}_{+}} \left\| M \mathbf{1}_{L_{j}} \right\|_{L^{q}(\mathbb{R}^{n})} \left| Q_{z}(2^{j}r) \right|^{\frac{1}{q'}} \left[\int_{Q_{z}(2^{j}r)} \left| f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right|^{q'} \, dx \right]^{\frac{1}{q'}} \end{split}$$

$$\leq \sum_{j \in \mathbb{Z}_{+}} 2^{jn\left[\frac{1}{\epsilon}\left(\frac{1}{q} - \frac{1}{p} - \alpha\right) + \frac{1}{q'}\right]} |Q_{z}(r)|^{\frac{1}{p'} - \alpha} \left[\int_{Q_{z}(2^{j}r)} \left| f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right|^{q'} dx \right]^{\frac{1}{q'}}$$

$$\leq \sum_{j \in \mathbb{Z}_{+}} 2^{jn\left[\frac{1}{\epsilon}\left(\frac{1}{q} - \frac{1}{p} - \alpha\right) + \frac{1}{q'}\right]} |Q_{z}(r)|^{\frac{1}{p'} - \alpha} \left[\int_{Q_{z}(2^{j}r)} \left| f(x) - P_{Q_{z}(2^{j}r)}^{(s)}(f)(x) \right|^{q'} dx \right]^{\frac{1}{q'}}$$

$$\leq \sum_{j \in \mathbb{Z}_{+}} 2^{jn\left(\frac{1}{\epsilon} - 1\right)\left(\frac{1}{q} - \frac{1}{p} - \alpha\right)} ||f||_{JN_{(p',q',s)_{\alpha}}^{con}(\mathbb{R}^{n})} \lesssim ||f||_{JN_{(p',q',s)_{\alpha}}^{con}(\mathbb{R}^{n})},$$

which implies the desired conclusion and hence completes the proof of Proposition 3.11.

Next, we show that any $(p, q, s, \alpha, \epsilon)$ -molecule M is in $HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$ via borrowing some ideas from [23, Theorem 7.2] and [33, Lemma 6.3]. To this end, we first establish several technical lemmas. The following lemma is widely used (see, for instance, [23, p. 86]), whose proof is similar to that of [23, Lemma 4.1]; we omit the details here.

Lemma 3.12. Let $s \in \mathbb{Z}_+$. Then there exists a positive constant C such that, for any cube (or ball) Q of \mathbb{R}^n , any $f \in L^1_{loc}(\mathbb{R}^n)$, and any $x \in 2Q \setminus Q$,

$$\left| P_{2Q \setminus Q}^{(s)}(f)(x) \right| \le C \int_{2Q \setminus Q} |f(x)| \, dx,$$

where $P_{2Q\setminus Q}^{(s)}(f)$ denotes the unique polynomial $P\in \mathcal{P}_s(\mathbb{R}^n)$ such that, for any $\gamma\in \mathbb{Z}_+^n$ with $|\gamma|\leq s$,

$$\int_{2Q\setminus Q} \left[f(x) - P_{2Q\setminus Q}^{(s)}(f)(x) \right] x^{\gamma} dx = 0.$$

The following Abel transformation is well known.

Lemma 3.13. Let $\{a_j\}_{j\in\mathbb{Z}_+}$, $\{b_j\}_{j\in\mathbb{Z}_+}\subset\mathbb{C}$. Then, for any $k\in\mathbb{N}$,

$$\sum_{j=1}^{k} a_j b_j = a_k \sum_{j=1}^{k} b_j - \sum_{j=1}^{k-1} \left[\left(\sum_{i=1}^{j} b_i \right) (a_{j+1} - a_j) \right].$$

We also need the following technical lemma.

Lemma 3.14. Let $u \in (1, \infty)$, $v \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$. Then there exists a positive constant C such that, for any cube $Q_z(r) \subset \mathbb{R}^n$ with $z \in \mathbb{R}^n$ and $r \in (0, \infty)$, any $k \in \mathbb{N}$, and any $f \in JN_{(u,v,s)_\alpha}^{\text{con}}(\mathbb{R}^n)$,

$$\left[\int_{Q_{z}(2^{k}r)} \left| f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right|^{\nu} dy \right]^{\frac{1}{\nu}} \le Ck \left[2^{ks} + 2^{nk(\alpha - \frac{1}{u})} \right] r^{n(\alpha - \frac{1}{u})} ||f||_{JN_{(u,v,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}.$$

Proof. Let u, v, s, and α be as in the present lemma. By Lemma 2.19 with $\lambda := 2$, (1.2), the Hölder inequality, and the definition of $\|\cdot\|_{JN^{\mathrm{con}}_{(u,v,s)_{\alpha}}(\mathbb{R}^n)}$, we find that, for any cube $Q_z(r) \subset \mathbb{R}^n$ with $z \in \mathbb{R}^n$ and $r \in (0,\infty)$, any $k \in \mathbb{N}$, and any $f \in JN^{\mathrm{con}}_{(u,v,s)_{\alpha}}(\mathbb{R}^n)$,

$$\left[\int_{Q_{z}(2^{k}r)} \left| f(x) - P_{Q_{z}(r)}^{(s)}(f)(x) \right|^{\nu} dx \right]^{\frac{1}{\nu}}$$

$$\begin{split} &\leq \left[\int_{Q_{z}(2^{k}r)}\left|f(x)-P_{Q_{z}(2^{k}r)}^{(s)}(f)(x)\right|^{v}dx\right]^{\frac{1}{v}} + \sum_{j=1}^{k}\left\|P_{Q_{z}(2^{j}r)}^{(s)}(f)-P_{Q_{z}(2^{j}r)}^{(s)}(f)\right\|_{L^{\infty}(Q_{z}(2^{k}r))} \\ &\leq \left[\int_{Q_{z}(2^{k}r)}\left|f(x)-P_{Q_{z}(2^{k}r)}^{(s)}(f)(x)\right|^{v}dx\right]^{\frac{1}{v}} \\ &+ \sum_{j=1}^{k}2^{(k-j+1)s}\left\|P_{Q_{z}(2^{j-1}r)}^{(s)}\left(f-P_{Q_{z}(2^{j}r)}^{(s)}(f)\right)\right\|_{L^{\infty}(Q_{z}(2^{j-1}r))} \\ &\leq \left[\int_{Q_{z}(2^{k}r)}\left|f(x)-P_{Q_{z}(2^{k}r)}^{(s)}(f)(x)\right|^{v}dx\right]^{\frac{1}{v}} \\ &+ \sum_{j=1}^{k}2^{(k-j+1)s}\int_{Q_{z}(2^{j-1}r)}\left|f(x)-P_{Q_{z}(2^{j}r)}^{(s)}(f)(x)\right|dx \\ &\leq \sum_{j=1}^{k}2^{(k-j+1)s}\left[\int_{Q_{z}(2^{j}r)}\left|f(x)-P_{Q_{z}(2^{j}r)}^{(s)}(f)(x)\right|^{v}dx\right]^{\frac{1}{v}} \\ &\sim \sum_{j=1}^{k}2^{(k-j+1)s}\left|Q_{z}(2^{j}r)\right|^{\alpha-\frac{1}{u}}\left|Q_{z}(2^{j}r)\right|^{-\alpha+\frac{1}{u}}\left[\int_{Q_{z}(2^{j}r)}\left|f(x)-P_{Q_{z}(2^{j}r)}^{(s)}(f)(x)\right|^{v}dx\right]^{\frac{1}{v}} \\ &\leq k\left[2^{ks}+2^{nk(\alpha-\frac{1}{u})}\right]r^{n(\alpha-\frac{1}{u})}\|f\|_{JN_{(u,v,s)a}^{con}(\mathbb{R}^{n})}. \end{split}$$

This finishes the proof of Lemma 3.14.

Proposition 3.15. Let $p \in (1, \infty)$, $q \in (1, \infty]$, $\frac{1}{q} + \frac{1}{q'} = 1$, $s \in \mathbb{Z}_+$, $\alpha \in (\frac{1}{q} - \frac{1}{p}, \infty)$, and $\epsilon \in (0, 1)$ satisfy $\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n} < 0$. If M is a $(p, q, s, \alpha, \epsilon)$ -molecule, then $M \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and $\|M\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)} \le C$, where the positive constant C is independent of M.

Proof. Let p, q, s, α , and ϵ be as in the present proposition. Also, let M be a $(p, q, s, \alpha, \epsilon)$ molecule centered at the cube $Q_z(r)$ with $z \in \mathbb{R}^n$ and $r \in (0, \infty)$. Without loss of generality, we
may assume that $z := \mathbf{0}$ and M is a real-valued function. Let $L_0 := Q_0(r)$ and, for any $j \in \mathbb{N}$, $L_j := Q_0(2^j r) \setminus Q_0(2^{j-1} r).$ For any $j \in \mathbb{Z}_+$ and $x \in \mathbb{R}^n$, let $P_{L_j}^{(s)}(M)$ denote the unique polynomial
of $\mathcal{P}_s(\mathbb{R}^n)$ such that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, and $j \in \mathbb{Z}_+$,

(3.8)
$$\int_{L_i} \left[M(x) - P_{L_i}^{(s)}(M)(x) \right] x^{\gamma} dx = 0.$$

To prove the present proposition, we first recall some known facts on $P_j := P_{L_j}^{(s)}(M)$ with $j \in \mathbb{Z}_+$ (see, for instance, [23, pp. 85-88]). For any given $j \in \mathbb{Z}_+$, let $\{\phi_{\nu}^{(j)} : \nu \in \mathbb{Z}_+^n \text{ and } |\nu| \leq s\}$ be the *orthogonal polynomials* with weight $\frac{1}{|L_j|}$ by means of the Gram–Schmidt method from $\{x^{\nu} : \nu \in \mathbb{Z}_+^n \text{ and } |\nu| \leq s\}$ restricted on L_j , namely, $\{\phi_{\nu}^{(j)} : \nu \in \mathbb{Z}_+^n \text{ and } |\nu| \leq s\} \subset \mathcal{P}_s(\mathbb{R}^n)$ and, for any $\nu_1, \nu_2 \in \mathbb{Z}_+^n$ with $|\nu_1| \leq s$ and $|\nu_2| \leq s$,

$$\left\langle \phi_{\nu_1}^{(j)}, \phi_{\nu_2}^{(j)} \right\rangle_{L_j} := \frac{1}{|L_j|} \int_{L_j} \phi_{\nu_1}^{(j)}(x) \phi_{\nu_2}^{(j)}(x) \, dx = \begin{cases} 1, & \nu_1 = \nu_2, \\ 0, & \nu_1 \neq \nu_2. \end{cases}$$

It is easy to see that, for any $x \in L_j$ with $j \in \mathbb{Z}_+$,

(3.9)
$$P_{j}(x) = \sum_{\{v \in \mathbb{Z}_{1}^{n}: |v| \le s\}} \left\langle M, \phi_{v}^{(j)} \right\rangle_{L_{j}} \phi_{v}^{(j)}(x).$$

Moreover, for any given $j \in \mathbb{Z}_+$, let $\{\psi_{\nu}^{(j)} : \nu \in \mathbb{Z}_+^n \text{ and } |\nu| \leq s\}$ be the *dual basis* of $\{x^{\nu} : \nu \in \mathbb{Z}_+^n \text{ and } |\nu| \leq s\}$ restricted on L_j with respect to the weight $\frac{1}{|L_j|}$, namely, $\psi_{\nu}^{(j)} \in \mathcal{P}_s(\mathbb{R}^n)$ and, for any $\nu_1, \nu_2 \in \mathbb{Z}_+^n$ with $|\nu_1| \leq s$ and $|\nu_2| \leq s$,

$$\left\langle \psi_{\nu_{1}}^{(j)}, x^{\nu_{2}} \right\rangle_{L_{j}} := \frac{1}{|L_{j}|} \int_{L_{j}} \psi_{\nu_{1}}^{(j)}(x) x^{\nu_{2}} \, dx = \begin{cases} 1, & \nu_{1} = \nu_{2}, \\ 0, & \nu_{1} \neq \nu_{2}. \end{cases}$$

Using the definition of the dual basis, we find that, if, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, $j \in \mathbb{Z}_+$, and $x \in L_j$, $\phi_v^{(j)}(x) = \sum_{\{\gamma \in \mathbb{Z}_+^n: \ |\gamma| \leq s\}} m_{v,\gamma}^{(j)} x^{\gamma}$ with $\{m_{v,\gamma}^{(j)}\} \subset \mathbb{R}$, then

$$\psi_{\nu}^{(j)} = \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \le s\}} m_{\gamma,\nu}^{(j)} \phi_{\gamma}^{(j)}.$$

From this and (3.9), we deduce that, for any $j \in \mathbb{Z}_+$,

$$(3.11) P_j \mathbf{1}_{L_j} = \sum_{\{\nu \in \mathbb{Z}_+^n: |\nu| \le s\}} \langle M, x^{\nu} \rangle_{L_j} \psi_{\nu}^{(j)} \mathbf{1}_{L_j}.$$

To show $M \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, we first estimate $M\mathbf{1}_{Q_0(2^lr)}$ for any $l \in \mathbb{N}$. Indeed, for any $l \in \mathbb{N}$ and $x \in \mathbb{R}^n$,

(3.12)
$$M\mathbf{1}_{Q_{0}(2^{l}r)}(x) = \sum_{j=1}^{l} \left[M\mathbf{1}_{L_{j}}(x) - P_{j}\mathbf{1}_{L_{j}}(x) \right] + \sum_{j=0}^{l} P_{j}\mathbf{1}_{L_{j}}(x)$$
$$= \sum_{j=1}^{l} \alpha_{j}(x) + \sum_{j=0}^{l} P_{j}\mathbf{1}_{L_{j}}(x),$$

where $\alpha_j := (M - P_j) \mathbf{1}_{L_j}$. We then show that, for any $j \in \mathbb{Z}_+$, α_j is a $(p, q, s)_{\alpha}$ -atom multiplying a positive constant. Indeed, by (3.8), we conclude that, for any $j \in \mathbb{Z}_+$ and $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$,

(3.13)
$$\int_{\mathbb{R}^n} \alpha_j(x) x^{\gamma} dx = 0 \quad \text{and} \quad \operatorname{supp}(\alpha_j) \subset Q_0(2^j r).$$

Thus, for any $j \in \mathbb{Z}_+$, α_j satisfies (i) and (iii) of Definition 3.1. Moreover, we show that, for any $j \in \mathbb{Z}_+$, α_j satisfies Definition 3.1(ii) in the sense of multiplying a positive constant. Indeed, from (3.7), the Hölder inequality, and Definition 3.10(ii), we deduce that, for any $j \in \mathbb{Z}_+$,

$$\|\alpha_{j}\|_{L^{q}(\mathbb{R}^{n})} \leq (1+C) \|M\mathbf{1}_{L_{j}}\|_{L^{q}(\mathbb{R}^{n})} \leq (1+C)2^{\frac{jn}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)} |Q_{\mathbf{0}}(r)|^{\frac{1}{q}-\frac{1}{p}-\alpha}$$

$$= (1+C)2^{jn[(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha)]} |Q_{\mathbf{0}}(2^{j}r)|^{\frac{1}{q}-\frac{1}{p}-\alpha} = \lambda_{j} |Q_{\mathbf{0}}(2^{j}r)|^{\frac{1}{q}-\frac{1}{p}-\alpha},$$

where C is as in (3.7), and

(3.14)
$$\lambda_{j} := (1+C)2^{jn(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha)}.$$

This and (3.13) imply that, for any $j \in \mathbb{Z}_+$, $A_j := \frac{\alpha_j}{\lambda_j}$ is a $(p, q, s)_{\alpha}$ -atom.

To end the estimation of $M\mathbf{1}_{Q_0(2^l r)}$ for any $l \in \mathbb{N}$, it remains to consider $\sum_{j=0}^l P_j \mathbf{1}_{L_j}$ for any $l \in \mathbb{N}$. For any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, and $j \in \mathbb{Z}_+$, let

(3.15)
$$\eta_{\nu}^{(j)} := \int_{\mathbb{R}^n \setminus O_0(2^j r)} M(x) x^{\nu} dx \quad \text{and} \quad \eta_{\nu}^{(-1)} := \int_{\mathbb{R}^n} M(x) x^{\nu} dx = 0.$$

Then, by (3.11), we conclude that, for any $j \in \mathbb{Z}_+$,

$$P_{j}\mathbf{1}_{L_{j}} = \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \frac{1}{|L_{j}|} \int_{L_{j}} M(x) x^{v} dx \psi_{v}^{(j)} \mathbf{1}_{L_{j}} = \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \frac{\eta_{v}^{(j-1)} - \eta_{v}^{(j)}}{|L_{j}|} \psi_{v}^{(j)} \mathbf{1}_{L_{j}},$$

which, together with Lemma 3.13 and (3.15), further implies that, for any $l \in \mathbb{N}$,

(3.16)
$$\sum_{j=0}^{l} P_{j} \mathbf{1}_{L_{j}} = \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \sum_{j=0}^{l} \left[\eta_{v}^{(j-1)} - \eta_{v}^{(j)} \right] \frac{\psi_{v}^{(j)} \mathbf{1}_{L_{j}}}{|L_{j}|}$$

$$= \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \sum_{j=0}^{l-1} \eta_{v}^{(j)} \left[\frac{\psi_{v}^{(j+1)} \mathbf{1}_{L_{j+1}}}{|L_{j+1}|} - \frac{\psi_{v}^{(j)} \mathbf{1}_{L_{j}}}{|L_{j}|} \right] - \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \frac{\eta_{v}^{(l)} \psi_{v}^{(l)} \mathbf{1}_{L_{l}}}{|L_{l}|}$$

$$= \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \sum_{j=0}^{l-1} \widetilde{\alpha}_{v}^{(j)} - \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \frac{\eta_{v}^{(l)} \psi_{v}^{(l)} \mathbf{1}_{L_{l}}}{|L_{l}|},$$

where, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$, and $j \in \{0, \dots, l-1\}$,

$$\widetilde{\alpha}_{\nu}^{(j)} := \eta_{\nu}^{(j)} \left[\frac{\psi_{\nu}^{(j+1)} \mathbf{1}_{L_{j+1}}}{|L_{j+1}|} - \frac{\psi_{\nu}^{(j)} \mathbf{1}_{L_{j}}}{|L_{j}|} \right].$$

Now, we claim that, for any $v \in \mathbb{Z}_+$ with $|v| \le s$, and $j \in \mathbb{Z}_+$, $\widetilde{\alpha}_v^{(j)}$ is a $(p, q, s)_\alpha$ -atom multiplying a positive constant and, for any $f \in JN_{(p',q',s)_\alpha}^{\text{con}}(\mathbb{R}^n)$,

(3.17)
$$\lim_{l \to \infty} \left| \int_{\mathbb{R}^n} \frac{\eta_{\nu}^{(l)} \psi_{\nu}^{(l)}(x) \mathbf{1}_{L_l}(x)}{|L_l|} f(x) \, dx \right| = 0.$$

Indeed, using (3.10), we conclude that, for any $\nu, \gamma \in \mathbb{Z}_+^n$ with $|\nu| \leq s$ and $|\gamma| \leq s$, and $j \in \mathbb{Z}_+$,

(3.18)
$$\int_{\mathbb{R}^n} \widetilde{\alpha}_{\nu}^{(j)}(x) x^{\gamma} dx = 0 \quad \text{and} \quad \operatorname{supp}(\widetilde{\alpha}_{\nu}^{(j)}) \subset Q_{\mathbf{0}}(2^{j+1}r).$$

Moreover, from [23, (7.2)], it follows that there exists a positive constant C_0 , independent of v, j, and r, such that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$, and any $x \in L_j$ with $j \in \mathbb{Z}_+$,

(3.19)
$$\left| \psi_{\nu}^{(j)}(x) \right| \le \frac{C_0}{(2^{j-1}r)^{|\nu|}}.$$

By this, (3.15), the Hölder inequality, Definition 3.10(ii), and $\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n} < 0$, we conclude that, for any $\nu \in \mathbb{Z}_+^n$ with $|\nu| \le s$, and $j \in \mathbb{Z}_+$,

$$\begin{aligned} \left| \eta_{\nu}^{(j)} \right| &\leq \int_{\mathbb{R}^{n} \setminus Q_{\mathbf{0}}(2^{j}r)} \left| M(x) x^{\nu} \right| \, dx = \sum_{i=j+1}^{\infty} \int_{L_{i}} \left| M(x) x^{\nu} \right| \, dx \\ &\leq \sum_{i=j+1}^{\infty} \left\| M \mathbf{1}_{L_{i}} \right\|_{L^{q}(\mathbb{R}^{n})} \left| L_{i} \right|^{\frac{1}{q'}} \left(2^{i} \sqrt{n} r \right)^{|\nu|} \\ &\lesssim \sum_{i=j+1}^{\infty} 2^{\frac{in}{\epsilon} (\frac{1}{q} - \frac{1}{p} - \alpha)} |Q_{\mathbf{0}}(r)|^{\frac{1}{q} - \frac{1}{p} - \alpha} \left| Q_{\mathbf{0}}(2^{i}r) \right|^{\frac{1}{q'} + \frac{|\nu|}{n}} \\ &\sim \sum_{i=j+1}^{\infty} 2^{in \left[\frac{1}{\epsilon} (\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{|\nu|}{n} \right]} |Q_{\mathbf{0}}(r)|^{\frac{1}{p'} - \alpha + \frac{|\nu|}{n}} \\ &\sim 2^{jn \left[(\frac{1}{\epsilon} - 1)(\frac{1}{q} - \frac{1}{p} - \alpha) \right]} |Q_{\mathbf{0}}(2^{j+1}r)|^{\frac{1}{p'} - \alpha + \frac{|\nu|}{n}}. \end{aligned}$$

From (3.19) and (3.20), it follows that, for any $v \in \mathbb{Z}_+^n$ with $|v| \le s$, and $j \in \mathbb{Z}_+$,

$$\begin{split} \left\| \widetilde{\alpha}_{\nu}^{(j)} \right\|_{L^{q}(\mathbb{R}^{n})} &\leq \left\| \widetilde{\alpha}_{\nu}^{(j)} \right\|_{L^{\infty}(\mathbb{R}^{n})} \left| Q_{\mathbf{0}}(2^{j+1}r) \right|^{\frac{1}{q}} \lesssim \frac{|\eta_{\nu}^{(j)}| |Q_{\mathbf{0}}(2^{j+1}r)|^{\frac{1}{q}}}{(2^{j-1}r)^{|\nu|} |L_{j}|} \\ &\sim \frac{|\eta_{\nu}^{(j)}|}{(2^{j-1}r)^{|\nu|}} \left| Q_{\mathbf{0}}(2^{j+1}r) \right|^{-\frac{1}{q'}} \sim \left| \eta_{\nu}^{(j)} \right| \left| Q_{\mathbf{0}}(2^{j+1}r) \right|^{-\frac{1}{q'} - \frac{|\nu|}{n}} \\ &\lesssim 2^{jn[(\frac{1}{\epsilon} - 1)(\frac{1}{q} - \frac{1}{p} - \alpha)]} \left| Q_{\mathbf{0}}(2^{j+1}r) \right|^{\frac{1}{q} - \frac{1}{p} - \alpha}. \end{split}$$

This and (3.18) imply that there exists a positive constant \widetilde{C} such that, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, and $j \in \mathbb{Z}_+$, $\widetilde{A}_v^{(j)} := \frac{\widetilde{a}_v^{(j)}}{\widetilde{\lambda}_j}$ is a $(p, q, s)_{\alpha}$ -atom, where

(3.21)
$$\widetilde{\lambda}_j := \widetilde{C}(2^{jn})^{(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha)}.$$

Thus, the first claim holds true. Then we show (3.17). Indeed, by (3.19), the Hölder inequality, Lemma 3.14 with u:=p' and v:=q', (3.20), (1.2), Lemma 2.19, $\alpha\in(\frac{1}{q}-\frac{1}{p},\infty)$, $\epsilon\in(0,1)$, and $\frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)+\frac{1}{q'}+\frac{s}{n}<0$, we conclude that, for any $l\in\mathbb{N}$ and $f\in JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$,

$$\left| \int_{\mathbb{R}^{n}} \frac{\eta_{\nu}^{(l)} \psi_{\nu}^{(l)}(x) \mathbf{1}_{L_{l}}(x)}{|L_{l}|} f(x) dx \right|$$

$$\lesssim \frac{|\eta_{\nu}^{(l)}|}{2^{(l-1)|\nu|}|L_{l}|} \int_{L_{l}} |f(x)| dx \lesssim \frac{|\eta_{\nu}^{(l)}|}{2^{(l-1)|\nu|}} \int_{Q_{0}(2^{l}r)} |f(x)| dx$$

$$\lesssim \frac{|\eta_{\nu}^{(l)}|}{2^{(l-1)|\nu|}} \left[\int_{Q_{0}(2^{l}r)} |f(x)|^{q'} dx \right]^{\frac{1}{q'}}$$

$$\lesssim \frac{|\eta_{\nu}^{(l)}|}{2^{(l-1)|\nu|}} \left\{ \left[\int_{Q_{0}(2^{l}r)} |f(x) - P_{Q_{0}(r)}^{(s)}(f)(x)|^{q'} dx \right]^{\frac{1}{q'}} + \left\| P_{Q_{0}(r)}^{(s)}(f) \right\|_{L^{\infty}(Q_{0}(2^{l}r))} \right\}$$

$$\leq 2^{nl[\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'}]} l \left[2^{ls} + 2^{nl(\alpha - \frac{1}{p'})} \right] ||f||_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}$$

$$+ 2^{nl[\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n}]} ||P_{Q_{0}(r)}^{(s)}(f)||_{L^{\infty}(Q_{0}(r))}$$

$$\leq 2^{nl[\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n}]} l ||f||_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} + 2^{nl[(\frac{1}{\epsilon} - 1)(\frac{1}{q} - \frac{1}{p} - \alpha)]} l ||f||_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}$$

$$+ 2^{nl[\frac{1}{\epsilon}(\frac{1}{q} - \frac{1}{p} - \alpha) + \frac{1}{q'} + \frac{s}{n}]} \int_{Q_{0}(r)} |f(x)| dx$$

$$\to 0$$

as $l \to \infty$, where the implicit positive constants depend on r. This further implies that (3.17) holds true.

Finally, we show that $M \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. Indeed, using (3.12) and (3.16), we conclude that, for any $l \in \mathbb{N}$ and $x \in \mathbb{R}^n$,

$$(3.22) M\mathbf{1}_{Q_{0}(2^{l}r)}(x) = \sum_{j=0}^{l} \lambda_{j} A_{j}(x) + \sum_{\{\nu \in \mathbb{Z}_{+}^{n}: |\nu| \leq s\}} \sum_{j=0}^{l-1} \widetilde{\lambda}_{j} \widetilde{A}_{\nu}^{(j)}(x) - \sum_{\{\nu \in \mathbb{Z}_{+}^{n}: |\nu| \leq s\}} \frac{\eta_{\nu}^{(l)} \psi_{\nu}^{(l)}(x) \mathbf{1}_{L_{l}}(x)}{|L_{l}|}.$$

By this, Lemma 3.2, Proposition 3.11, the Lebesgue dominated convergence theorem, (3.22), and (3.17), we conclude that, for any $l \in \mathbb{N}$ and $f \in JN^{\text{con}}_{(p',q',s)_{\sigma}}(\mathbb{R}^{n})$,

$$\left| \left\langle M - \sum_{j=0}^{l} \lambda_{j} A_{j} - \sum_{j=0}^{l-1} \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \widetilde{\lambda}_{j} \widetilde{A}_{v}^{(j)}, f \right\rangle \right|$$

$$= \left| \int_{\mathbb{R}^{n}} \left[M(x) - \sum_{j=1}^{l} \lambda_{j} A_{j}(x) - \sum_{j=0}^{l-1} \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \widetilde{\lambda}_{j} \widetilde{A}_{v}^{(j)}(x) \right] f(x) dx \right|$$

$$\leq \left| \int_{\mathbb{R}^{n} \setminus Q_{0}(2^{l}r)} M(x) f(x) dx \right| + \sum_{\{v \in \mathbb{Z}_{+}^{n}: |v| \leq s\}} \left| \int_{\mathbb{R}^{n}} \frac{\eta_{v}^{(l)} \psi_{v}^{(l)}(x) \mathbf{1}_{L_{l}}(x)}{|L_{l}|} f(x) dx \right|$$

$$\to 0$$

as $l \to \infty$, and hence

$$(3.23) M = \sum_{j \in \mathbb{Z}_+} \lambda_j A_j + \sum_{j \in \mathbb{Z}_+} \sum_{\{v \in \mathbb{Z}_+^n : |v| \le s\}} \widetilde{\lambda}_j \widetilde{A}_v^{(j)}$$

in $(JN_{(p',q',s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n}))^{*}$. Moreover, since, for any $\nu \in \mathbb{Z}_{+}^{n}$ with $|\nu| \leq s$, and $j \in \mathbb{Z}_{+}$, A_{j} and $\widetilde{A}_{\nu}^{(j)}$ are $(p,q,s)_{\alpha}$ -atoms, we deduce that $\lambda_{j}A_{j}$, $\widetilde{\lambda_{j}}\widetilde{A}_{\nu}^{(j)} \in \widetilde{HK}_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})$, which, combined with (3.14), (3.21), $(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha) < 0$, and (3.23), further implies that

$$\begin{split} \|M\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^{n})} &\leq \sum_{j \in \mathbb{Z}_{+}} \left\| \lambda_{j} A_{j} \right\|_{\widetilde{HK}^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^{n})} + \sum_{j \in \mathbb{Z}_{+}} \sum_{\{\nu \in \mathbb{Z}_{+}^{n} : \ |\nu| \leq s\}} \left\| \widetilde{\lambda}_{j} \widetilde{A}_{\nu}^{(j)} \right\|_{\widetilde{HK}^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^{n})} \\ &\leq \sum_{j \in \mathbb{Z}_{+}} |\lambda_{j}| + \sum_{\{\nu \in \mathbb{Z}_{+}^{n} : \ |\nu| \leq s\}} \sum_{j \in \mathbb{Z}_{+}} \left| \widetilde{\lambda}_{j} \right| \end{split}$$

$$\lesssim \sum_{j \in \mathbb{Z}_+} 2^{jn[(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha)]} + \sum_{\{v \in \mathbb{Z}_+^n: \; |v| \leq s\}} \sum_{j \in \mathbb{Z}_+} 2^{jn[(\frac{1}{\epsilon}-1)(\frac{1}{q}-\frac{1}{p}-\alpha)]} \sim 1,$$

where the implicit constants are independent of M, and hence $M \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. This finishes the proof of Proposition 3.15.

Using Proposition 3.15, we obtain the following useful criterion for the boundedness of linear operators on $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$. In what follows, we equip $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ and $HK^{\text{con-fin}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ with the same norm $\|\cdot\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}$ unless specifically indicated.

Theorem 3.16. Let $p_1, p_2 \in (1, \infty), q_1, q_2 \in (1, \infty], \frac{1}{p_i} + \frac{1}{p_i'} = 1 = \frac{1}{q_i} + \frac{1}{q_i'}$ for any $i \in \{1, 2\}, s_1, s_2 \in \mathbb{Z}_+, \alpha_1 \in \mathbb{R}$, and $\alpha_2 \in (\frac{1}{q_2} - \frac{1}{p_2}, \infty)$. Let A be a linear operator defined on $HK^{\text{con-fin}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ and \widetilde{A} a linear operator bounded from $JN^{\text{con}}_{(p_2',q_2',s_2)_{\alpha_2}}(\mathbb{R}^n)$ to $JN^{\text{con}}_{(p_1',q_1',s_1)_{\alpha_1}}(\mathbb{R}^n)$. If the following statements hold true:

- (i) there exists a positive constant C_0 such that, for any $(p_1,q_1,s_1)_{\alpha_1}$ -atom a and some given $\epsilon \in (0,1)$ such that $\frac{1}{\epsilon}(\frac{1}{q_2}-\frac{1}{p_2}-\alpha_2)+\frac{1}{q_2'}+\frac{s_2}{n}<0, \frac{A(a)}{C_0}$ is a $(p_2,q_2,s_2,\alpha_2,\epsilon)$ -molecule;
- (ii) for any $(p_1,q_1,s_1)_{\alpha_1}$ -atom a and any $f\in JN^{\mathrm{con}}_{(p'_2,q'_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$, $\langle A(a),f\rangle=\langle a,\widetilde{A}(f)\rangle$,

then the linear operator A has a unique continuous linear extension, still denoted by A, from $HK^{\mathrm{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ to $HK^{\mathrm{con}}_{(p_2,q_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $g \in HK^{\mathrm{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$,

$$||A(g)||_{HK_{(p_2,q_2,s_2)\alpha_2}^{\text{con}}(\mathbb{R}^n)} \le C||g||_{HK_{(p_1,q_1,s_1)\alpha_1}^{\text{con}}(\mathbb{R}^n)}$$

and, moreover, for any $f \in JN^{\text{con}}_{(p'_1,q'_1,s_2)_{\alpha_2}}(\mathbb{R}^n)$,

(3.24)
$$\langle A(g), f \rangle = \langle g, \widetilde{A}(f) \rangle.$$

Proof. Let p_1 , p_2 , q_1 , q_2 , s_1 , s_2 , α_1 , α_2 , A, and A be as in the present theorem. We first prove that A has a unique continuous linear extension on $HK^{\text{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$. To this end, we claim that A is bounded from $HK^{\text{con-fin}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ to $HK^{\text{con}}_{(p_2,q_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$. Indeed, let $g \in HK^{\text{con-fin}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$. Then, from (i) and (ii) of the present theorem, Lemma 3.7, and Propositions 3.11 and 3.15, it follows that, for any $f \in JN^{\text{con}}_{(p'_2,g'_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$,

$$\begin{split} |\langle A(g),f\rangle| &= \left|\left\langle g,\widetilde{A}(f)\right\rangle\right| \lesssim \|g\|_{HK^{\mathrm{con}}_{(p_1,q_1,s_1)\alpha_1}(\mathbb{R}^n)} \left\|\widetilde{A}(f)\right\|_{JN^{\mathrm{con}}_{(p_1',q_1',s_1)\alpha_1}(\mathbb{R}^n)} \\ &\lesssim \|g\|_{HK^{\mathrm{con}}_{(p_1,q_1,s_1)\alpha_1}(\mathbb{R}^n)} \|f\|_{JN^{\mathrm{con}}_{(p_2',q_2',s_2)\alpha_2}(\mathbb{R}^n)}, \end{split}$$

which, together with Proposition 3.8, further implies that

and hence the above claim holds true.

Moreover, by (3.25), Lemma 3.6, and Proposition 3.9, we conclude that the linear operator A is bounded from the dense subspace $HK^{\operatorname{con-fin}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ of $HK^{\operatorname{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ to $HK^{\operatorname{con}}_{(p_2,q_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$. From this and [40, p. 22, Proposition 5.4], we deduce that A can be extended to a unique continuous linear operator, still denoted by A, from $HK^{\operatorname{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ to $HK^{\operatorname{con}}_{(p_2,q_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$ in the sense that, for any $g \in HK^{\operatorname{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$,

(3.26)
$$\lim_{m \to \infty} ||A(g) - A(g_m)||_{HK_{(p_2, q_2, s_2)\alpha_2}^{\text{con}}(\mathbb{R}^n)} = 0,$$

where $\{g_m\}_{m\in\mathbb{N}}$ is a sequence of $HK^{\operatorname{con-fin}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$ such that

(3.27)
$$\lim_{m \to \infty} ||g - g_m||_{HK_{(p_1, q_1, s_1)\alpha_1}^{\text{con}}(\mathbb{R}^n)} = 0,$$

A(g) is independent of the choice of $\{g_m\}_{m\in\mathbb{N}}$, and, for any $g\in HK^{\mathrm{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$,

$$||A(g)||_{HK^{\text{con}}_{(p_2,q_2,s_2)\alpha_2}(\mathbb{R}^n)} \lesssim ||g||_{HK^{\text{con}}_{(p_1,q_1,s_1)\alpha_1}(\mathbb{R}^n)}.$$

Next, we show (3.24). Indeed, for any $f \in JN^{\text{con}}_{(p'_2,q'_2,s_2)_{\alpha_2}}(\mathbb{R}^n)$ and $g \in HK^{\text{con}}_{(p_1,q_1,s_1)_{\alpha_1}}(\mathbb{R}^n)$, let $g_m := \sum_{i=1}^{M_m} \lambda_{m,i} a_{m,i}$ be as in (3.26) with $\{\lambda_{m,i}\}_{i=1}^{M_m} \subset \mathbb{C}$ and $\{a_{m,i}\}_{i=1}^{M_m}$ being $(p_1,q_1,s_1)_{\alpha_1}$ -atoms. Then, by (3.26), Lemma 3.7, (ii) of the present theorem, and (3.27), we find that

$$\langle A(g), f \rangle = \lim_{m \to \infty} \langle A(g_m), f \rangle = \lim_{m \to \infty} \sum_{i=1}^{M_m} \lambda_{m,i} \langle A(a_{m,i}), f \rangle$$
$$= \lim_{m \to \infty} \sum_{i=1}^{M_m} \lambda_{m,i} \langle a_{m,i}, \widetilde{A}(f) \rangle = \langle g, \widetilde{A}(f) \rangle,$$

which shows the above claim. This finishes the proof of Theorem 3.16.

Remark 3.17. Although we are not able to establish a molecular characterization of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ because the classical method seems no longer feasible in the present setting, the proof of Theorem 3.16 contains some thoughts of the molecular characterization of $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$.

3.3 Boundedness of Calderón–Zygmund Operators on $HK^{\text{con}}_{(p,q,s)_a}(\mathbb{R}^n)$

In this subsection, we show that the *s*-order Calderón–Zygmund singular integral operator T, having the vanishing moments up to order s, can be extended to a unique continuous linear operator on $HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$. To this end, we establish two technical lemmas. We first show that, for any $(p,q,s)_{\alpha}$ -atom a, T(a) is a (p,q,s,α,ϵ) -molecule in the sense of multiplying a positive constant independent of a.

Lemma 3.18. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $\frac{1}{q} + \frac{1}{q'} = 1$, $s \in \mathbb{Z}_+$, $\alpha \in (\frac{1}{q} - \frac{1}{p}, 1 - \frac{1}{p} + \frac{s + \delta}{n})$ with $\delta \in (0, 1]$ as in Definition 2.6, and $\epsilon \in (0, 1)$ be such that $-\frac{1}{q'} - \frac{s + \delta}{n} \leq \frac{1}{\epsilon} (\frac{1}{q} - \frac{1}{p} - \alpha)$. Let K be an s-order standard kernel and T the s-order Calderón–Zygmund singular integral operator with kernel K. Then the following two statements are equivalent:

- (i) there exists a positive constant C such that, for any $(p,q,s)_{\alpha}$ -atom a, $\frac{T(a)}{C}$ is a (p,q,s,α,ϵ) -molecule:
- (ii) for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, $T^*(x^{\gamma}) = 0$.

Proof. Let $p, q, q', \alpha, s, \alpha, \delta, \epsilon, T$, and K be as in the present lemma. We first show (i) \Rightarrow (ii). Indeed, for any given $A \in L^q_s(\mathbb{R}^n)$ with $||A||_{L^q(\mathbb{R}^n)} > 0$ supported in a ball $B(x_0, r_0)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$, it is easy to see that $\widetilde{A} := \frac{|Q_{x_0}(2r_0)|^{\frac{1}{q} - \frac{1}{p} - \alpha}A}{||A||_{L^q(\mathbb{R}^n)}}$ is a $(p, q, s)_{\alpha}$ -atom supported in $Q_{x_0}(2r_0)$. By this, (i) of the present lemma, and (2.19), we conclude that, for any $\gamma \in \mathbb{Z}^n_+$ with $|\gamma| \leq s$,

$$\int_{B(x_0,r_0)} \widetilde{A}(x)\widetilde{T}(y^{\gamma})(x) dx = \int_{\mathbb{R}^n} T(\widetilde{A})(x)x^{\gamma} dx = 0.$$

Using this and a proof similar to that of Proposition 2.16, we find that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $\widetilde{T}(y^{\gamma}) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero, which, combined with Proposition 2.16, further implies that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$. This finishes the proof that (i) \Rightarrow (ii).

Next, we show (ii) \Rightarrow (i). Let a be a $(p,q,s)_{\alpha}$ -atom supported in the cube $Q_z(r)$ with $z \in \mathbb{R}^n$ and $r \in (0,\infty)$. We show that there exists a positive constant C, independent of a, such that $\frac{T(a)}{C}$ is a (p,q,s,α,ϵ) -molecule. To this end, let $R_0 := 2\sqrt{n}r$. We first prove that there exists a positive constant C_1 , independent of a, such that

(3.28)
$$||T(a)\mathbf{1}_{Q_{z}(R_{0})}||_{L^{q}(\mathbb{R}^{n})} \leq C_{1} |Q_{z}(R_{0})|^{\frac{1}{q} - \frac{1}{p} - \alpha}.$$

Indeed, using Lemma 2.13, Definition 3.1(ii), and $r \sim R_0$, we find that

$$||T(a)\mathbf{1}_{Q_z(R_0)}||_{L^q(\mathbb{R}^n)} \lesssim ||a||_{L^q(Q_z(r))} \lesssim |Q_z(r)|^{\frac{1}{q}-\frac{1}{p}-\alpha} \sim |Q_z(R_0)|^{\frac{1}{q}-\frac{1}{p}-\alpha}.$$

This implies that (3.28) holds true.

Now, we show that there exists a positive constant C_2 , independent of a, such that, for any $j \in \mathbb{N}$,

Indeed, for any $x \notin Q_z(R_0)$ and $y \in Q_z(r)$, we have $|z - x| \ge 2|z - y|$, which, combined with $x \notin \text{supp}(a)$, Definition 3.1(iii), the Taylor remainder theorem, (2.6), the Hölder inequality, and Definition 3.1(ii), further implies that, for any $y \in Q_z(r)$, there exists a $\widetilde{y} \in Q_z(r)$ such that, for any $x \notin Q_z(R_0)$,

$$|T(a)(x)| = \left| \int_{Q_{z}(r)} K(x, y) a(y) \, dy \right|$$

$$\leq \int_{Q_{z}(r)} \left| K(x, y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x, z)}{\gamma!} (y - z)^{\gamma} \right| |a(y)| \, dy$$

$$= \int_{Q_{z}(r)} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| = s\}} \frac{\partial_{(2)}^{\gamma} K(x, \overline{y}) - \partial_{(2)}^{\gamma} K(x, z)}{\gamma!} (y - z)^{\gamma} \right| |a(y)| \, dy$$

$$\lesssim \frac{r^{s+\delta}}{|z-x|^{n+s+\delta}} \int_{Q_{z}(r)} |a(y)| \, dy \lesssim \frac{r^{\frac{n}{q'}+s+\delta}}{|z-x|^{n+s+\delta}} ||a||_{L^{q}(Q_{z}(r))}$$

$$\lesssim \left(\frac{r}{|z-x|}\right)^{\frac{n}{q'}+s+\delta} |z-x|^{-\frac{n}{q}} |Q_{z}(r)|^{\frac{1}{q}-\frac{1}{p}-\alpha}.$$

From this, $r \sim R_0$, and $-\frac{1}{q'} - \frac{s+\delta}{n} \leq \frac{1}{\epsilon} (\frac{1}{q} - \frac{1}{p} - \alpha)$, we deduce that, for any $j \in \mathbb{N}$ and $x \in Q_z(2^jR_0) \setminus Q_z(2^{j-1}R_0)$,

$$\begin{split} |T(a)(x)| & \lesssim \left(\frac{r}{2^{j}R_{0}}\right)^{\frac{n}{q'}+s+\delta} (2^{j}R_{0})^{-\frac{n}{q}} |Q_{z}(R_{0})|^{\frac{1}{q}-\frac{1}{p}-\alpha} \\ & \lesssim (2^{jn})^{\frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)} \left|Q_{z}(2^{j}R_{0}) \setminus Q_{z}(2^{j-1}R_{0})\right|^{-\frac{1}{q}} |Q_{z}(R_{0})|^{\frac{1}{q}-\frac{1}{p}-\alpha} \end{split}$$

and hence

$$\left\|T(a)\mathbf{1}_{Q_{z}(2^{j}R_{0})\backslash Q_{z}(2^{j-1}R_{0})}\right\|_{L^{q}(\mathbb{R}^{n})}\lesssim (2^{jn})^{\frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)}|Q_{z}(R_{0})|^{\frac{1}{q}-\frac{1}{p}-\alpha},$$

which implies that (3.29) holds true.

Moreover, by (ii) of the present lemma, we know that T(a) satisfies Definition 3.10(iii). From this, (3.28), and (3.29), we deduce that $\frac{T(a)}{C}$ satisfies (i), (ii), and (iii) of Definition 3.10, where $C := \max\{C_1, C_2\}$. This finishes the proof of Lemma 3.18.

Lemma 3.19. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$, $s \in \mathbb{Z}_+$, and $\alpha \in (\frac{1}{q} - \frac{1}{p}, 1 - \frac{1}{p} + \frac{s + \delta}{n})$ with $\delta \in (0, 1]$ as in Definition 2.6. Let K be an s-order standard kernel and T an s-order Calderón–Zygmund singular integral operator having the vanishing moments up to order s with kernel K. Then, for any $(p, q, s)_{\alpha}$ -atom a and any $f \in JN^{\text{con}}_{(p', q', s)_{\alpha}}(\mathbb{R}^n)$,

$$\langle T(a), f \rangle = \left\langle a, \widetilde{T}(f) \right\rangle,$$

where \widetilde{T} is as in Definition 2.11 with kernel \widetilde{K} as in (2.10).

Proof. Let $p, q, s, \alpha, \delta, T, \widetilde{T}, K$, and \widetilde{K} be as in the present lemma. Also, let a be a $(p, q, s)_{\alpha}$ -atom supported in the cube $Q_{x_0}(2r_0/\sqrt{n})$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$, and $B_0 := B(x_0, r_0)$. Then $Q_{x_0}(2r_0/\sqrt{n}) \subset B_0$. From this, Lemma 3.18, Proposition 3.11, Remark 2.8, and Definition 3.1(iii), we deduce that, for any $f \in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$,

$$(3.30) \quad \langle T(a), f \rangle = \int_{\mathbb{R}^n} T(a)(x)f(x) dx$$

$$= \int_{\mathbb{R}^n} \lim_{\eta \to 0^+} \int_{B_0 \setminus B(x,\eta)} K(x,y)a(y) dy f(x) dx$$

$$= \int_{2B_0} \lim_{\eta \to 0^+} \int_{B_0 \setminus B(x,\eta)} K(x,y)a(y) dy f(x) dx$$

$$+ \int_{\mathbb{R}^n \setminus 2B_0} \int_{B_0} \left[K(x,y) - \sum_{\{y \in \mathbb{Z}^n : |y| \le \xi\}} \frac{\partial_{(2)}^{\gamma} K(x,x_0)}{\gamma!} (y - x_0)^{\gamma} \right] a(y) dy f(x) dx$$

$$=: \Lambda_1 + \Lambda_2,$$

where T_{η} is as in (2.8).

We first consider Λ_1 . To this end, for any $\eta \in (0, \infty)$, let K_{η} be as in (2.26). Then, by $a \in L^q(\mathbb{R}^n)$, Remark 2.8, the fact that $f\mathbf{1}_{2B_0} \in L^{q'}(\mathbb{R}^n)$, and the Fubini theorem, we find that

(3.31)
$$\Lambda_{1} = \lim_{\eta \to 0^{+}} \int_{2B_{0}} \int_{B_{0}} K_{\eta}(x, y) a(y) \, dy f(x) \, dx$$
$$= \lim_{\eta \to 0^{+}} \int_{B_{0}} a(y) \int_{2B_{0}} K_{\eta}(x, y) f(x) \, dx \, dy$$
$$= \int_{B_{0}} a(y) \lim_{\eta \to 0^{+}} \int_{2B_{0}} K_{\eta}(x, y) f(x) \, dx \, dy.$$

This is a desired conclusion of Λ_1 .

Next, we consider Λ_2 . We first claim that

$$(3.32) \qquad \int_{\mathbb{R}^n \setminus 2B_0} \int_{B_0} \left| \left| K(x, y) - \sum_{\{ \gamma \in \mathbb{Z}_+^n \colon |\gamma| \le s \}} \frac{\partial_{(2)}^{\gamma} K(x, x_0)}{\gamma!} (y - x_0)^{\gamma} \right| a(y) f(x) \right| dy dx < \infty.$$

Indeed, from (2.29), we deduce that

$$(3.33) \qquad \int_{\mathbb{R}^n \setminus 2B_0} \int_{B_0} \left[\left| K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_0)}{\gamma!} (y - x_0)^{\gamma} \right| a(y) \right| \left| P_{2B_0}^{(s)}(f)(x) \right| dy dx < \infty.$$

Moreover, using the Tonelli theorem, the Taylor remainder theorem, (2.6), Lemma 2.21 with $\beta := s + \delta \in (s, \infty)$, and $\alpha \in (\frac{1}{q} - \frac{1}{p}, 1 - \frac{1}{p} + \frac{s + \delta}{n})$, we find that, for any $y \in B_0$, there exists a $\widetilde{y} \in B_0$ such that

$$\begin{split} &\int_{\mathbb{R}^{n}\backslash 2B_{0}} \int_{B_{0}} \left\| \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right] a(y) \left[f(x) - P_{2B_{0}}^{(s)}(f)(x) \right] \right| dy dx \\ &= \int_{B_{0}} \int_{\mathbb{R}^{n}\backslash 2B_{0}} \left\| \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right] a(y) \left[f(x) - P_{2B_{0}}^{(s)}(f)(x) \right] \right| dx dy \\ &= \int_{B_{0}} \int_{\mathbb{R}^{n}\backslash 2B_{0}} \left| \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: \, |\gamma| = s\}} \frac{\partial_{(2)}^{\gamma} K(x,\overline{y}) - \partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right| |a(y)| \left| f(x) - P_{2B_{0}}^{(s)}(f)(x) \right| dx dy \\ &\lesssim \int_{B_{0}} \int_{\mathbb{R}^{n}\backslash 2B_{0}} \frac{\left| \overline{y} - x_{0} \right|^{\delta} |y - x_{0}|^{s}}{|x - x_{0}|^{n+s+\delta}} |a(y)| \left| f(x) - P_{2B_{0}}^{(s)}(f)(x) \right| dx dy \\ &\lesssim r_{0}^{s+\delta} ||a||_{L^{1}(B_{0})} \int_{\mathbb{R}^{n}\backslash 2B_{0}} \frac{|f(x) - P_{2B_{0}}^{(s)}(f)(x)|}{|x - x_{0}|^{n+s+\delta}} dx \\ &\lesssim r_{0}^{(-\frac{1}{p'} + \alpha)n} ||a||_{L^{1}(B_{0})} ||f||_{JN_{(p',q',s)_{\alpha}}^{con}(\mathbb{R}^{n})} < \infty, \end{split}$$

which, together with (3.33), further implies that

$$\int_{\mathbb{R}^{n}\backslash 2B_{0}} \int_{B_{0}} \left| \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right] a(y) f(x) \right| dy dx \\
\leq \int_{\mathbb{R}^{n}\backslash 2B_{0}} \int_{B_{0}} \left| \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right] a(y) P_{B_{0}}^{(s)}(f)(x) \right| dy dx \\
+ \int_{\mathbb{R}^{n}\backslash 2B_{0}} \int_{B_{0}} \left| \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_{+}^{n}: |\gamma| \leq s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_{0})}{\gamma!} (y - x_{0})^{\gamma} \right] a(y) \left[f(x) - P_{2B_{0}}^{(s)}(f)(x) \right] dy dx \\
< \infty$$

and hence (3.32) holds true. Then, by (3.32) and the Fubini theorem, we obtain

(3.34)
$$\Lambda_2 = \int_{B_0} \int_{\mathbb{R}^n \setminus 2B_0} \left[K(x, y) - \sum_{\{y \in \mathbb{Z}^n : |y| \le \epsilon\}} \frac{\partial_{(2)}^{\gamma} K(x, x_0)}{\gamma!} (y - x_0)^{\gamma} \right] f(x) \, dx \, a(y) \, dy.$$

This is a desired conclusion of Λ_2 .

Altogether, from (3.30), (3.31), and (3.34), it follows that, for any $f \in JN_{(p,q,s)_q}^{\text{con}}(\mathbb{R}^n)$,

$$\begin{split} &\langle T(a), f \rangle \\ &= \int_{B_0} a(y) \lim_{\eta \to 0^+} \int_{2B_0 \setminus B(y,\eta)} K(x,y) f(x) \, dx \, dy \\ &+ \int_{B_0} \int_{\mathbb{R}^n \setminus 2B_0} \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_+^n : \, |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_0)}{\gamma!} (y - x_0)^{\gamma} \right] f(x) \, dx a(y) \, dy \\ &= \int_{B_0} a(y) \lim_{\eta \to 0^+} \int_{\mathbb{R}^n \setminus B(y,\eta)} \left[K(x,y) - \sum_{\{\gamma \in \mathbb{Z}_+^n : \, |\gamma| \le s\}} \frac{\partial_{(2)}^{\gamma} K(x,x_0)}{\gamma!} (y - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus 2B_0}(x) \right] f(x) \, dx \, dy \\ &= \int_{B_0} a(y) \lim_{\eta \to 0^+} \int_{\mathbb{R}^n \setminus B(y,\eta)} \left[\widetilde{K}(y,x) - \sum_{\{\gamma \in \mathbb{Z}_+^n : \, |\gamma| \le s\}} \frac{\partial_{(1)}^{\gamma} \widetilde{K}(x_0,x)}{\gamma!} (y - x_0)^{\gamma} \mathbf{1}_{\mathbb{R}^n \setminus 2B_0}(x) \right] f(x) \, dx \, dy \\ &= \int_{B_0} a(y) \widetilde{T}(f)(y) \, dy = \left\langle a, \widetilde{T}(f) \right\rangle. \end{split}$$

This finishes the proof of Lemma 3.19.

Now, we give the following main result of this subsection, which shows the boundedness of Calderón–Zygmund operators on $HK^{\text{con}}_{(p,q,s)_0}(\mathbb{R}^n)$.

Theorem 3.20. Let $p \in (1, \infty)$, $q \in (1, \infty)$, $\frac{1}{p} + \frac{1}{p'} = 1 = \frac{1}{q} + \frac{1}{q'}$, $s \in \mathbb{Z}_+$, and $\alpha \in \mathbb{R}$ satisfy $\frac{1}{q} - \frac{1}{p} < \alpha < \frac{s+\delta}{n}$ with $\delta \in (0,1]$ as in Definition 2.6. Let K be an s-order standard kernel as in Definition 2.6 and T the s-order Calderón–Zygmund singular integral operator with kernel K. Then the following two statements are equivalent:

(i) there exists a positive constant C such that, for any $(p, q, s)_{\alpha}$ -atom a,

$$(3.35) ||T(a)||_{HK^{\text{con}}_{(n,a,s)_{\alpha}}(\mathbb{R}^{n})} \le C$$

and T can be extended to a unique continuous linear operator, still denoted by T, on the space $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $g \in HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$,

$$(3.36) ||T(g)||_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \le C||g||_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}$$

and, moreover, for any $f \in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, $\widetilde{T}(f) \in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ and

(3.37)
$$\langle T(g), f \rangle = \langle g, \widetilde{T}(f) \rangle,$$

where \widetilde{T} is as in Definition 2.11 with kernel \widetilde{K} as in (2.10);

(ii) for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \le s$, $T^*(x^{\gamma}) = 0$.

Proof. Let $p, q, s, \alpha, \delta, K, T, \widetilde{K}$, and \widetilde{T} be as in the present theorem. We first show (i) \Rightarrow (ii). Indeed, using Lemma 3.7, (3.37), and (3.36), we find that, for any $f \in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$,

$$\begin{split} \left\|\widetilde{T}(f)\right\|_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} &\sim \left\|\mathcal{L}_{\widetilde{T}(f)}\right\|_{(HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n}))^{*}} \sim \sup_{\|g\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})=1}} \left|\left\langle \mathcal{L}_{\widetilde{T}(f)},g\right\rangle\right| \\ &\sim \sup_{\|g\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})=1}} \left|\left\langle g,\widetilde{T}(f)\right\rangle\right| \sim \sup_{\|g\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})=1}} \left|\left\langle T(g),f\right\rangle\right| \\ &\lesssim \sup_{\|g\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})=1}} \|T(g)\|_{HK_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \|f\|_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})} \\ &\lesssim \|f\|_{JN_{(p',q',s)_{\alpha}}^{\text{con}}(\mathbb{R}^{n})}, \end{split}$$

where $\mathcal{L}_{\widetilde{T}(f)}$ is as in (3.1) with f replaced by $\widetilde{T}(f)$. This shows that \widetilde{T} is bounded on $JN^{\mathrm{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, which, combined with Theorem 2.25, further implies that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^{\gamma}) = 0$. This finishes the proof that (i) \Rightarrow (ii).

Next, we show (ii) \Rightarrow (i). Let $\epsilon \in (0,1)$ be such that $\frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)+\frac{1}{q'}+\frac{s}{n}<0$ and $-\frac{1}{q'}-\frac{s+\delta}{n}\leq \frac{1}{\epsilon}(\frac{1}{q}-\frac{1}{p}-\alpha)$. Then, by Proposition 3.15 and Lemma 3.18, we conclude that, for any $(p,q,s)_{\alpha}$ -atom a,T(a) is a (p,q,s,α,ϵ) -molecule and $\|T(a)\|_{HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)}\leq C$, where the positive constant C is independent of a. This implies that (3.35) holds true. Moreover, from (ii) of the present theorem and Theorem 2.25, it follows that \widetilde{T} is bounded on $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ and hence, for any $f\in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$, $\widetilde{T}(f)\in JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$. Furthermore, by the boundedness of \widetilde{T} on $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$ and Lemmas 3.18 and 3.19, we conclude that (i) and (ii) of Theorem 3.16 hold true with A:=T, $\widetilde{A}:=\widetilde{T}$, $p_1:=p=:p_2,q_1:=q=:q_2,s_1:=s=:s_2,\alpha_1:=\alpha=:\alpha_2$, which further implies that (3.36) and (3.37) hold true. Then we complete the proof that (ii) \Rightarrow (i) and hence of Theorem 3.20.

- **Remark 3.21.** (i) Theorem 3.20 implies that the adjoint operator of T on $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ is just the operator \widetilde{T} defined on $JN^{\text{con}}_{(p',q',s)_{\alpha}}(\mathbb{R}^n)$; see, for instance, [24, Theorem 5] and [33, Theorem 3.1] for similar results on Campanato-type spaces.
- (ii) In Theorem 3.20, $\frac{1}{q} \frac{1}{p} < \alpha < \frac{s+\delta}{n}$ is a suitable assumption because we need $\alpha \in (\frac{1}{q} \frac{1}{p}, \infty)$ when proving that the molecule in Definition 3.10 belongs to $HK^{\text{con}}_{(p,q,s)_{\alpha}}(\mathbb{R}^n)$ (see Proposition 3.15), and $\alpha \in (-\infty, \frac{s+\delta}{n})$ when proving that \widetilde{T} is bounded on the congruent JNC space (see Theorem 2.25). It is still unknown whether or not Theorem 3.20 holds true with $\alpha \in (-\infty, \frac{1}{q} \frac{1}{p}]$.
- (iii) Let $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, K be an s-order standard kernel, and T the s-order Calderón–Zygmund singular integral operator with kernel K. An essential ingredient in this article is the boundedness of T on $L^q(\mathbb{R}^n)$, which can be deduced from the boundedness of T on $L^2(\mathbb{R}^n)$ and (i) through (iv) of Definition 2.6. However, if we directly assume that T is bounded on $L^q(\mathbb{R}^n)$, then Theorems 2.25 and 3.20 still hold true under the assumption that K only satisfies (i) and (iii) of Definition 2.6.

Acknowledgements Dachun Yang would like to thank Professor Ibrahim Fofana for him to bring their attention to references [8, 12].

References

- [1] D. Aalto, L. Berkovits, O. E. Kansanen and H. Yue, John–Nirenberg lemmas for a doubling measure, Studia Math. 204 (2011), 21-37.
- [2] P. Auscher and C. Prisuelos-Arribas, Tent space boundedness via extrapolation, Math. Z. 286 (2017), 1575-1604.
- [3] M. Bownik, Anisotropic Hardy spaces and wavelets, Mem. Amer. Math. Soc. 164 (2003), no. 781, 122 pp.
- [4] S. Campanato, Proprietà di una famiglia di spazi funzionali, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 18 (1964), 137-160.
- [5] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, Bull. Amer. Math. Soc. 83 (1977), 569-645.
- [6] G. Dafni, T. Hytönen, R. Korte and H. Yue, The space JN_p : nontriviality and duality, J. Funct. Anal. 275 (2018), 577-603.
- [7] O. Domínguez and M. Milman, Sparse Brudnyi and John–Nirenberg spaces, arXiv: 2107. 05117.
- [8] M. Dosso, I. Fofana and M. Sanogo, On some subspaces of Morrey-Sobolev spaces and boundedness of Riesz integrals, Ann. Polon. Math. 108 (2013), 133-153.
- [9] J. Duoandikoetxea, Fourier Analysis, Graduate Studies in Mathematics 29, American Mathematical Society, Providence, RI, 2001.
- [10] J. Feuto, I. Fofana and K. Koua, Integrable fractional mean functions on spaces of homogeneous type, Afr. Diaspora J. Math. (N.S.) 9 (2010), 8-30.
- [11] I. Fofana, Étude d'une classe d'espace de fonctions contenant les espaces de Lorentz, Afrika Mat. (2) 1 (1988), 29-50.

- [12] I. Fofana, F. R. Faléa and B. A. Kpata, A class of subspaces of Morrey spaces and norm inequalities on Riesz potential operators, Afr. Mat. 26 (2015), 717-739.
- [13] B. Franchi, C. Pérez and R. L. Wheeden, Self-improving properties of John–Nirenberg and Poincaré inequalities on spaces of homogeneous type, J. Funct. Anal. 153 (1998), 108-146.
- [14] N. Fusco, G. Moscariello and C. Sbordone, BMO-type seminorms and Sobolev functions, ESAIM Control Optim. Calc. Var. 24 (2018), 835-847.
- [15] G. Hu, D. Yang and Y. Zhou, Boundedness of singular integrals in Hardy spaces on spaces of homogeneous type, Taiwanese J. Math. 13 (2009), 91-135.
- [16] R. Hurri-Syrjänen, N. Marola and A. V. Vähäkangas, Aspects of local-to-global results, Bull. Lond. Math. Soc. 46 (2014), 1032-1042.
- [17] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Special John–Nirenberg–Campanato spaces via congruent cubes, Sci. China Math. (2021), https://doi.org/10.1007/s11425-021-1866-4.
- [18] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Boundedness of fractional integrals on special John–Nirenberg–Campanato and Hardy-type spaces via congruent cubes, Submitted or arXiv: 2108.01891.
- [19] H. Jia, D. Yang, W. Yuan and Y. Zhang, Estimates for Littlewood–Paley operators on special John–Nirenberg–Campanato spaces via congruent cubes, Submitted or arXiv: 2108.01559.
- [20] F. John and L. Nirenberg, On functions of bounded mean oscillation, Comm. Pure Appl. Math. 14 (1961), 415-426.
- [21] N. Kikuchi, E. Nakai, N. Tomita, K. Yabuta and T. Yoneda, Calderón–Zygmund operators on amalgam spaces and in the discrete case, J. Math. Anal. Appl. 335 (2007), 198-212.
- [22] S. Kislyakov and N. Kruglyak, Extremal Problems in Interpolation Theory, Whitney—Besicovitch Coverings, and Singular Integrals, Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) 74, Birkhäuser/Springer Basel AG, Basel, 2013.
- [23] S. Lu, Four Lectures on Real *H*^p Spaces, World Scientific Publishing Co., Inc., River Edge, NJ, 1995.
- [24] R. A. Macías and C. Segovia, Singular integrals on generalized Lipschitz and Hardy spaces, Studia Math. 65 (1979), 55-75.
- [25] P. MacManus and C. Pérez, Generalized Poincaré inequalities: sharp self-improving properties, Internat. Math. Res. Notices 1998 (1998), 101-116.
- [26] Y. Meyer and R. R. Coifman, Wavelets. Calderón–Zygmund and Multilinear Operators, Cambridge Studies in Advanced Mathematics 48, Cambridge University Press, Cambridge, 1997.
- [27] M. Milman, Marcinkiewicz spaces, Garsia–Rodemich spaces and the scale of John–Nirenberg self improving inequalities, Ann. Acad. Sci. Fenn. Math. 41 (2016), 491-501.
- [28] M. Milman, Garsia–Rodemich spaces: Bourgain–Brezis–Mironescu space, embeddings and rearrangement invariant spaces, J. Anal. Math. 139 (2019), 121-141.
- [29] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations. Trans. Amer. Math. Soc. 43 (1938), 126-166.
- [30] E. Nakai, Hardy–Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces, Math. Nachr. 166 (1994), 95-103.
- [31] E. Nakai, The Campanato, Morrey and Hölder spaces on spaces of homogeneous type, Studia Math. 176 (2006), 1-19.

- [32] E. Nakai, Singular and fractional integral operators on Campanato spaces with variable growth conditions, Rev. Mat. Complut. 23 (2010), 355-381.
- [33] E. Nakai, Singular and fractional integral operators on preduals of Campanato spaces with variable growth condition, Sci. China Math. 60 (2017), 2219-2240.
- [34] E. Nakai and Y. Sawano, Hardy spaces with variable exponents and generalized Campanato spaces, J. Funct. Anal. 262 (2012), 3665-3748.
- [35] E. Nakai and K. Yabuta, Pointwise multipliers for functions of weighted bounded mean oscillation on spaces of homogeneous type, Math. Japon. 46 (1997), 15-28.
- [36] J. Peetre, On convolution operators leaving $L^{p,\lambda}$ spaces invariant, Ann. Mat. Pura Appl. (4) 72 (1966), 295-304.
- [37] Y. Sawano and H. Tanaka, Morrey spaces for non-doubling measures, Acta Math. Sin. (Engl. Ser.) 21 (2005), 1535-1544.
- [38] E. M. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton Mathematical Series 30, Princeton University Press, Princeton, NJ, 1970.
- [39] E. M. Stein, Harmonic analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals, Princeton Mathematical Series 43, Monographs in Harmonic Analysis, III, Princeton University Press, Princeton, NJ, 1993.
- [40] E. M. Stein and R. Shakarchi, Functional Analysis. Introduction to Further Topics in Analysis, Princeton Lectures in Analysis 4, Princeton University Press, Princeton, NJ, 2011.
- [41] J. Sun, G. Xie and D. Yang, Localized John–Nirenberg–Campanato spaces, Anal. Math. Phys. 11 (2021), Paper No. 29, 49 pp.
- [42] J. Tao, D. Yang and W. Yuan, John-Nirenberg-Campanato spaces, Nonlinear Anal. 189 (2019), 111584, 36 pp.
- [43] J. Tao, D. Yang and W. Yuan, A bridge connecting Lebesgue and Morrey spaces via Riesz norms, Banach J. Math. Anal. 15 (2021), Paper No. 20, 29 pp.
- [44] J. Tao, D. Yang and W. Yuan, Vanishing John–Nirenberg spaces, Adv. Calc. Var. (2021), https://doi.org/10.1515/acv-2020-0061.
- [45] J. Tao, D. Yang and W. Yuan, A survey on function spaces of John–Nirenberg type, Submitted or arXiv: 2108.01511.
- [46] Z. Zeng, D.-C. Chang, J. Tao and D. Yang, Nontriviality of Riesz–Morrey spaces, Appl. Anal. (2021), https://doi.org/10.1080/00036811.2021.1932836.

Hongchao Jia, Jin Tao, Dachun Yang (Corresponding author), Wen Yuan and Yangyang Zhang Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, People's Republic of China

```
E-mails: hcjia@mail.bnu.edu.cn (H. Jia)
    jintao@mail.bnu.edu.cn (J. Tao)
    dcyang@bnu.edu.cn (D. Yang)
    wenyuan@bnu.edu.cn (W. Yuan)
    yangyzhang@mail.bnu.edu.cn (Y. Zhang)
```