Estimates for Littlewood–Paley Operators on Ball Campanato-Type Function Spaces

Hongchao Jia, Dachun Yang, Wen Yuan and Yangyang Zhang

Abstract Let X be a ball quasi-Banach function space on \mathbb{R}^n and assume that the Hardy–Littlewood maximal operator satisfies the Fefferman–Stein vector-valued maximal inequality on X, and let $q \in [1, \infty)$ and $d \in (0, \infty)$. In this article, the authors prove that, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ (the ball Campanato-type function space associated with X), the Littlewood–Paley g-function g(f) is either infinite everywhere or finite almost everywhere and, in the latter case, g(f) is bounded on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$. Similar results for both the Lusin-area function and the Littlewood–Paley g_{λ}^* -function are also obtained. All these results have a wide range of applications. Particularly, even when X is the weighted Lebesgue space, or the Miccolar space, all these results are new. The proofs of all these results strongly depend on several delicate estimates of Littlewood–Paley operators on the mean oscillation of the locally integrable function f on \mathbb{R}^n . Moreover, the same ideas are also used to obtain the corresponding results for the special John–Nirenberg–Campanato space via congruent cubes.

1 Introduction

It is well known that both Campanato spaces and Hardy spaces play vital roles in harmonic analysis and partial differential equations (see, for instance, [10, 61]). Recall that John and Nirenberg [39] introduced the space BMO (\mathbb{R}^n), namely, the space of functions with bounded mean oscillation, which proves the dual space of the Hardy space $H^1(\mathbb{R}^n)$ by Fefferman and Stein [16]. Later, as a generalization of the space BMO (\mathbb{R}^n), Campanato [5] introduced the Campanato space $C_{\alpha,q,s}(\mathbb{R}^n)$ which coincides with BMO (\mathbb{R}^n) when $\alpha=0$. Moreover, for any given $p\in(0,1)$, any given $q\in[1,\infty]$, and any given integer $s\in[0,\infty)\cap[n(\frac{1}{p}-1),\infty)$, $C_{\frac{1}{p}-1,q,s}(\mathbb{R}^n)$ was proved to be the dual space of the Hardy space $H^p(\mathbb{R}^n)$ by Coifman and Weiss [10].

Recall that Stein [60] introduced the Littlewood–Paley operators, namely, the Littlewood–Paley g-function, the Lusin-area function S, and the Littlewood–Paley g_{λ}^* function. Since then, estimates of Littlewood–Paley operators on Campanato-type spaces have attracted more and more attention. For instance, Wang [74] proved that, for any $f \in BMO(\mathbb{R}^n)$, g(f) is either infinite

²⁰²⁰ Mathematics Subject Classification. Primary 42B25; Secondary 42B20, 42B30, 42B35, 46E35, 47A30.

Key words and phrases. Littlewood-Paley operator, Campanato space, ball quasi-Banach function space, John-Nirenberg space.

This project is supported by the National Natural Science Foundation of China (Grant Nos. 11971058, 12071197 and 12122102) and the National Key Research and Development Program of China (Grant No. 2020YFA0712900).

^{*}Corresponding author, E-mail: dcyang@bnu.edu.cn/June 19, 2022/Final version.

everywhere or finite almost everywhere and, in the latter case, g(f) is bounded on BMO (\mathbb{R}^n). Later, Lu et al. [45] extended the main result of [74] to Campanato spaces. Moreover, Meng et al. [46] studied the boundedness of the Lusin-area function S and the Littlewood–Paley g_{λ}^* function on Campanato spaces over spaces of homogeneous type in the sense of Coifman and Weiss. Recently, Li et al. [40] studied the boundedness of multilinear Littlewood–Paley operators on amalgam-Campanato spaces. See also, for instance, [21, 42, 46, 47, 54, 66] for more studies on Littlewood–Paley operators on Campanato-type spaces.

Sawano et al. [56] introduced the ball quasi-Banach function space X and the Hardy space $H_X(\mathbb{R}^n)$ associated with X. Ball quasi-Banach function spaces contain many well-known function spaces, for instance, weighted Lebesgue spaces, Morrey spaces, mixed-norm Lebesgue spaces, Orlicz–slice spaces, and Musielak–Orlicz spaces, which implies that the results obtained in [56] have wide applications. We refer the reader to [6, 64, 65, 68, 72, 73, 76, 75, 81] for more studies on Hardy-type spaces associated with ball quasi-Banach function spaces.

As a counterpart of the dual relation $(H^p(\mathbb{R}^n))^* = C_{\frac{1}{p}-1,q,s}(\mathbb{R}^n)$, the dual space of $H_X(\mathbb{R}^n)$ has been widely studied when X is some concrete function space (see, for instance, [18, 30, 43, 52, 53, 80]). Moreover, Yan et al. [77] introduced the Campanato-type function space $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ associated with X, which proves the dual space of $H_X(\mathbb{R}^n)$ when X is concave. Very recently, Zhang et al. [78] improved the dual result in [77] via removing the assumption that X is concave. To be precise, Zhang et al. [78] introduced a new ball Campanato-type function space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ associated with X, and proved that

$$(H_X(\mathbb{R}^n))^* = \mathcal{L}_{X,q,s,d}(\mathbb{R}^n).$$

However, the mapping properties of Littlewood–Paley operators on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ are still unknown even when X is some concrete function space, for instance, the weighted Lebesgue space, the mixed-norm Lebesgue space, the variable Lebesgue space, the Orlicz space, and the Orlicz-slice space.

In this article, we study the mapping properties of Littlewood–Paley operators on the ball Campanato space $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ under the assumption that the Hardy–Littlewood maximal operator satisfies the Fefferman–Stein vector-valued maximal inequality on X (see, Assumption 2.8 below). To be precise, we prove that, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, the Littlewood–Paley g-function g(f) is either infinite everywhere or finite almost everywhere and, in the latter case, g is bounded on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ (see Theorem 3.5 below). Similar results for both the Lusin-area function and the Littlewood–Paley g^*_{λ} -function are also obtained (see Theorems 4.1 and 5.2 below). All these results have a wide range of applications. Particularly, even when $X := L^p_{\omega}(\mathbb{R}^n)$ (the weighted Lebesgue space), $X := L^p(\mathbb{R}^n)$ (the mixed-norm Lebesgue space), $X := L^p(\mathbb{R}^n)$ (the variable Lebesgue space), $X := L^p(\mathbb{R}^n)$ (the Orlicz space), and $X := (E^q_{\Phi})_t(\mathbb{R}^n)$ (the Orlicz-slice space), all these results are new. The proofs of these results strongly depend on several delicate estimates of Littlewood–Paley operators on the mean oscillation of the locally integrable function f on \mathbb{R}^n . Due to the generality and the practicability, more applications of these results are predictable. Moreover, the same ideas are also used to obtain the corresponding results for $X := JN^{\text{con}}_{p,q,\alpha}(\mathbb{R}^n)$ (the special John–Nirenberg–Campanato space via congruent cubes).

Indeed, the proofs of the boundedness of Littlewood–Paley operators on Campanato-type spaces in [44, 46, 66] strongly depend on the definition of the norm of Campanato-type spaces,

in which there exists only one cube. However, in the definition of the norm of $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, there appear countable balls which cause additional essential difficulties in the proofs of our main results. To overcome these difficulties, we establish several delicate estimates of Littlewood–Paley functions over the mean oscillation of the locally integrable function f (see Lemmas 3.10, 4.3, and 5.6 below). Moreover, we make a delicate division of \mathbb{R}^n in the proof of the boundedness of Littlewood–Paley g_{λ}^* -function on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, and then improve the existing results via widening the ranges of λ in the Littlewood–Paley g_{λ}^* -function (see Theorem 5.2 below).

The remainder of this article is organized as follows.

Section 2 is divided into two subsections to give some basic concepts. In Subsection 2.1, we first recall the concepts of quasi-Banach function spaces X and two ball Campanato-type function spaces, both $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, associated with X. Moreover, we present a mild assumption about the Fefferman–Stein vector-valued maximal inequality on X (see Assumption 2.8 below). In Subsection 2.2, we present the definitions of several general Littlewood–Paley operators, including the Littlewood–Paley g-function, the Lusin-area function S, and the Littlewood–Paley g₃-function.

Section 3 is devoted to studying the mapping properties of the Littlewood–Paley g-function on both $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$. To be precise, we first establish a technical lemma about g-function via borrowing some ideas from [74] (see Lemma 3.10 below). Then we show that, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, g(f) is either infinite everywhere or finite almost everywhere and, in the latter case, g is bounded on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ (see Theorem 3.5 below). Estimates of the Littlewood–Paley g-function on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ are also obtained in Theorem 3.5 below.

In Section 4, we study the mapping properties of the Lusin-area function S on both $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$. To be precise, we prove that, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, S(f) is either infinite everywhere or finite almost everywhere and, in the latter case, S is bounded on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ (see Theorem 4.1 below). Similar result for the Lusin-area function S on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ is also obtained.

Section 5 is devoted to studying the mapping properties of the Littlewood–Paley g_{λ}^* -function on both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,s}(\mathbb{R}^n)$. To be precise, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, g_{λ}^* is either infinite everywhere or finite almost everywhere and, in the latter case, g_{λ}^* is bounded on $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ (see Theorem 5.2 below). Similar result for $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ is also obtained in Theorem 5.1 below. Both Theorems 5.1 and 5.2 even when $X := L^p(\mathbb{R}^n)$ with $p \in (\frac{n}{n+1}, \infty)$, $\delta := 1$, and $q \in [1, \infty)$ also improve the existing results via widening the ranges of λ [see Remark 5.7(iii) below].

In Section 6, we apply the main results obtained in Sections 3, 4, and 5 to five concrete ball quasi-Banach function spaces, namely, the weighted Lebesgue space $L^p_\omega(\mathbb{R}^n)$, the mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$, the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$, the Orlicz space $L^{\Phi}(\mathbb{R}^n)$, and the Orlicz-slice space $(E^r_\Phi)_t(\mathbb{R}^n)$. Therefore, all the mapping properties of Littlewood–Paley functions on Campanato-type spaces $\mathcal{L}_{L^p_\omega(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^{\vec{p}}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^p(\cdot)(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^\Phi(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, and $\mathcal{L}_{(E^r_\Phi)_t(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$ are obtained. Moreover, using the same ideas, we also obtain the mapping properties of Littlewood–Paley operators on the special John–Nirenberg–Campanato spaces via congruent cubes, $JN^{\text{con}}_{p,q,\alpha}(\mathbb{R}^n)$, introduced in [36, Definition 1.3] (see Theorems 6.21, 6.23, and 6.25 below).

Finally, we make some conventions on notation. Let $\mathbb{N} := \{1, 2, \ldots\}$, $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$, and $\mathbb{Z}_+^n := (\mathbb{Z}_+)^n$. We always denote by C a *positive constant* which is independent of the main parameters, but it may vary from line to line. The symbol $f \leq g$ means that $f \leq Cg$. If $f \leq g$ and $g \leq f$, we then write $f \sim g$. If $f \leq Cg$ and g = h or $g \leq h$, we then write $f \leq g \sim h$ or

 $f \leq g \leq h$, rather than $f \leq g = h$ or $f \leq g \leq h$. For any $q \in [1, \infty]$, we use q' to denote the conjugate exponent of p, namely, $\frac{1}{q} + \frac{1}{q'} = 1$. For any measurable subset E of \mathbb{R}^n , we denote by $\mathbf{1}_E$ its characteristic function. We use $\mathbf{0}$ to denote the *origin* of \mathbb{R}^n . Moreover, for any $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $B(x, r) := \{y \in \mathbb{R}^n : |y - x| < r\}$. Furthermore, for any $\lambda \in (0, \infty)$ and any ball $B(x, r) \subset \mathbb{R}^n$ with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $\lambda B(x, r) := B(x, \lambda r)$.

2 Preliminaries

In this section, we give some basic concepts. In Subsection 2.1, we present the definitions of both the ball quasi-Banach function space X and its related ball Campanato-type function space. In Subsection 2.2, we recall the definition of Littlewood–Paley operators under consideration.

2.1 Campanato-Type Function Spaces Associated with X

The ball quasi-Banach function space was introduced in [56, Definition 2.2]. In what follows, we use $\mathcal{M}(\mathbb{R}^n)$ to denote the set of all measurable functions on \mathbb{R}^n . Moreover, for any $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $B(x, r) := \{y \in \mathbb{R}^n : |x - y| < r\}$ and

$$(2.1) \mathbb{B}(\mathbb{R}^n) := \{B(x, r) : x \in \mathbb{R}^n \text{ and } r \in (0, \infty)\}.$$

Definition 2.1. Let $X \subset \mathcal{M}(\mathbb{R}^n)$ be a quasi-normed linear space equipped with a quasi-norm $\|\cdot\|_X$ which makes sense for all measurable functions on \mathbb{R}^n . Then X is called a *ball quasi-Banach function space* if it satisfies:

- (i) for any $f \in \mathcal{M}(\mathbb{R}^n)$, $||f||_X = 0$ implies that f = 0 almost everywhere;
- (ii) for any $f, g \in \mathcal{M}(\mathbb{R}^n)$, $|g| \le |f|$ almost everywhere implies that $||g||_X \le ||f||_X$;
- (iii) for any $\{f_m\}_{m\in\mathbb{N}}\subset \mathcal{M}(\mathbb{R}^n)$ and $f\in \mathcal{M}(\mathbb{R}^n)$, $0\leq f_m\uparrow f$ almost everywhere as $m\to\infty$ implies that $\|f_m\|_X\uparrow\|f\|_X$ as $m\to\infty$;
- (iv) $B \in \mathbb{B}(\mathbb{R}^n)$ implies that $\mathbf{1}_B \in X$, where $\mathbb{B}(\mathbb{R}^n)$ is the same as in (2.1).
- **Remark 2.2.** (i) Let *X* be a ball quasi-Banach function space on \mathbb{R}^n . Using [75, Remark 2.6(i)], we find that, for any $f \in \mathcal{M}(\mathbb{R}^n)$, $||f||_X = 0$ if and only if f = 0 almost everywhere.
 - (ii) As was mentioned in [75, Remark 2.6(ii)], we obtain an equivalent formulation of Definition 2.1 via replacing any ball $B \in \mathbb{B}(\mathbb{R}^n)$ by any bounded measurable set E therein.
- (iii) From [14, Theorem 2], we deduce that both (ii) and (iii) of Definition (2.1) imply that *X* is complete and hence a quasi-Banach space.

Now, we recall the concepts of the convexity and the concavity of ball quasi-Banach function spaces, which is a part of [56, Definition 2.6].

Definition 2.3. Let X be a ball quasi-Banach function space and $p \in (0, \infty)$.

(i) The *p-convexification* X^p of X is defined by setting

$$X^p := \{ f \in \mathcal{M}(\mathbb{R}^n) : |f|^p \in X \}$$

equipped with the *quasi-norm* $||f||_{X^p} := |||f|^p||_X^{1/p}$ for any $f \in X^p$.

(ii) The space X is said to be *concave* if there exists a positive constant C such that, for any $\{f_k\}_{k\in\mathbb{N}}\subset \mathcal{M}(\mathbb{R}^n)$,

$$\sum_{k=1}^{\infty} ||f_k||_X \le C \left\| \sum_{k=1}^{\infty} |f_k| \right\|_X.$$

In particular, when C = 1, X is said to be *strictly concave*.

Next, we present the definition of the ball Campanato-type function space associated with X. Recall that the *Lebesgue space* $L^q(\mathbb{R}^n)$ with $q \in (0, \infty]$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{L^{q}(\mathbb{R}^{n})} := \begin{cases} \left[\int_{\mathbb{R}^{n}} |f(x)|^{q} dx \right]^{\frac{1}{q}} & \text{if } q \in (0, \infty), \\ \text{ess sup } |f(x)| & \text{if } q = \infty \end{cases}$$

is finite. Besides, for any given $q \in (0, \infty)$, we use $L^q_{loc}(\mathbb{R}^n)$ to denote the set of all the measurable functions f such that $f\mathbf{1}_E \in L^q(\mathbb{R}^n)$ for any bounded measurable set $E \subset \mathbb{R}^n$ and, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any finite measurable subset $E \subset \mathbb{R}^n$, let

$$f_E := \int_E f(x) \, dx := \frac{1}{|E|} \int_E f(x) \, dx.$$

Moreover, for any $s \in \mathbb{Z}_+$, we use $\mathcal{P}_s(\mathbb{R}^n)$ to denote the set of all the polynomials on \mathbb{R}^n with total degree not greater than s; for any ball $B \subset \mathbb{R}^n$ and any $g \in L^1_{loc}(\mathbb{R}^n)$, $P_B^{(s)}(g)$ denotes the *minimizing polynomial* of g with degree not greater than s, namely, $P_B^{(s)}(g)$ is the unique polynomial in $\mathcal{P}_s(\mathbb{R}^n)$ such that, for any $P \in \mathcal{P}_s(\mathbb{R}^n)$,

$$\int_{B} \left[g(x) - P_{B}^{(s)}(g)(x) \right] P(x) \, dx = 0.$$

Recall that Yan et al. [77, Definition 1.11] introduced the following Campanato space $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ associated with the ball quasi-Banach function space X.

Definition 2.4. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, and $s \in \mathbb{Z}_+$. Then the *Campanato space* $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$, associated with X, is defined to be the set of all the $f \in L^q_{loc}(\mathbb{R}^n)$ such that

$$||f||_{\mathcal{L}_{X,q,s}(\mathbb{R}^n)} := \sup_{B \in \mathbb{B}(\mathbb{R}^n)} \frac{|B|}{||\mathbf{1}_B||_X} \left[\int_B \left| f(x) - P_B^{(s)}(f)(x) \right|^q dx \right]^{\frac{1}{q}}$$

is finite.

In what follows, by abuse of notation, we identify $f \in \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ with $f + \mathcal{P}_s(\mathbb{R}^n)$.

Remark 2.5. (i) Let $f \in L^1_{loc}(\mathbb{R}^n)$, $s \in \mathbb{Z}_+$, and $B \in \mathbb{B}(\mathbb{R}^n)$. By [44, p. 54, Lemma 4.1], we find that there exists a positive constant C, independent of both f and B, such that

(2.2)
$$||P_B^{(s)}(f)||_{L^{\infty}(B)} \le C \int_B |f(x)| \, dx.$$

Moreover, it is well known that $P_R^{(0)}(f) = f_B$.

(ii) Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in [0, \infty)$. Recall that Campanato [5] introduced the *Campanato space* $C_{\alpha,q,s}(\mathbb{R}^n)$ which is defined to be the set of all the $f \in L^q_{loc}(\mathbb{R}^n)$ such that

(2.3)
$$||f||_{C_{\alpha,q,s}(\mathbb{R}^n)} := \sup |B|^{-\alpha} \left[\int_B \left| f(x) - P_B^{(s)}(f)(x) \right|^q \right]^{\frac{1}{q}} < \infty,$$

where the supremum is taken over all balls $B \in \mathbb{B}(\mathbb{R}^n)$. It is well known that $C_{\alpha,q,s}(\mathbb{R}^n)$ when $\alpha = 0$ coincides with the space BMO (\mathbb{R}^n) . Moreover, when $X := L^{\frac{1}{\alpha+1}}(\mathbb{R}^n)$, then $\mathcal{L}_{X,q,s}(\mathbb{R}^n) = C_{\alpha,q,s}(\mathbb{R}^n)$.

Recall that, in [78, Definition 3.2], Zhang et al. introduced the following new ball Campanato space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ associated with the ball quasi-Banach function space X.

Definition 2.6. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, $d \in (0, \infty)$, and $s \in \mathbb{Z}_+$. Then the *ball Campanato-type function space* $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, associated with X, is defined to be the set of all the $f \in L^q_{loc}(\mathbb{R}^n)$ such that

$$||f||_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)} := \sup \left\| \left\{ \sum_{i=1}^m \left(\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right)^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_X^{-1}$$

$$\times \sum_{i=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} \left| f(x) - P_{B_j}^{(s)}(f)(x) \right|^q dx \right]^{\frac{1}{q}}$$

is finite, where the supremum is taken over all $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$, and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$.

In what follows, by abuse of notation, we identify $f \in \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ with $f + \mathcal{P}_s(\mathbb{R}^n)$.

Remark 2.7. Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, $d \in (0, 1]$, and X be a concave ball quasi-Banach function space. It was proved in [78, Proposition 3.7] that $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n) = \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ with equivalent quasi-norms.

To obtain the main results of this article, we need the following mild assumption of the Fefferman–Stein vector-valued maximal inequality on X (see, for instance, [78, Assumption 2.6]).

Assumption 2.8. Let X be a ball quasi-Banach function space. Assume that there exists a $p_- \in (0, \infty)$ such that, for any given $p \in (0, p_-)$ and $u \in (1, \infty)$, there exists a positive constant C such that, for any $\{f_j\}_{j=1}^{\infty} \subset \mathcal{M}(\mathbb{R}^n)$,

(2.4)
$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^u \right\}^{\frac{1}{u}} \right\|_{X^{1/p}} \le C \left\| \left(\sum_{j \in \mathbb{N}} |f_j|^u \right)^{\frac{1}{u}} \right\|_{X^{1/p}}.$$

2.2 Littlewood-Paley Operators

In this subsection, we recall the definition of Littlewood–Paley operators. We first present the concept of Littlewood–Paley functions.

Definition 2.9. A scalar-valued function $\varphi \in L^1(\mathbb{R}^n)$ is called a *Littlewood–Paley function* if there exists a positive constant C and a $\delta \in (0, \infty)$ such that

(i)

(2.5)
$$\int_{\mathbb{R}^n} \varphi(x) \, dx = 0;$$

(ii) for any $x \in \mathbb{R}^n$,

$$|\varphi(x)| \le \frac{C}{(1+|x|)^{n+\delta}};$$

(iii) for any $x \in \mathbb{R}^n$,

$$|\nabla \varphi(x)| \le \frac{C}{(1+|x|)^{n+1+\delta}},$$

here and thereafter, $\nabla := (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})$.

Now, we recall the concept of Littlewood–Paley operators, namely, the Littlewood–Paley g-function, the Lusin-area function S, and the Littlewood–Paley g_{λ}^* -function. In what follows, for any measurable function φ on \mathbb{R}^n , and for any $t \in (0, \infty)$, let

$$\varphi_t(\cdot) := \frac{1}{t^n} \varphi\left(\frac{\cdot}{t}\right).$$

Moreover, let

$$\mathbb{R}^{n+1}_+ := \left\{ (\xi, t) \in \mathbb{R}^{n+1} : \xi \in \mathbb{R}^n, \ t \in (0, \infty) \right\}$$

and, for any $x \in \mathbb{R}^n$,

$$\Gamma(x) := \left\{ (\xi, t) \in \mathbb{R}^{n+1}_+ : |\xi - x| < t \right\}.$$

Definition 2.10. Let φ be the same as in Definition 2.9. For any $f \in L^1_{loc}(\mathbb{R}^n)$, the Littlewood–Paley g-function g(f), the Lusin-area function S(f), and the Littlewood–Paley g^*_{λ} -function $g^*_{\lambda}(f)$ with $\lambda \in (1, \infty)$ of f are defined, respectively, by setting, for any $x \in \mathbb{R}^n$,

(2.8)
$$g(f)(x) := \left[\int_0^\infty |f * \varphi_t(x)|^2 \frac{dt}{t} \right]^{\frac{1}{2}},$$

(2.9)
$$S(f)(x) := \left[\iint_{\Gamma(x)} |f * \varphi_t(\xi)|^2 \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}},$$

and

(2.10)
$$g_{\lambda}^{*}(f)(x) := \left[\iint_{\mathbb{R}^{n+1}_{+}} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} |f * \varphi_{t}(\xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}.$$

3 Littlewood–Paley g-Functions on Both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$

This section is devoted to studying the boundedness of the Littlewood–Paley g function on both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$. To this end, we first introduce the subspace $\mathcal{L}_{X,q,0}^{\mathrm{low}}(\mathbb{R}^n)$ of $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, and the subspace $\mathcal{L}_{X,q,0,d}^{\mathrm{low}}(\mathbb{R}^n)$ of $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

Definition 3.1. Let X be a ball quasi-Banach function space and $q \in [1, \infty)$. Then the *space* $\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)$ is defined to be the set of all the $f \in L^q_{\text{loc}}(\mathbb{R}^n)$ such that

$$||f||_{\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \frac{|B|}{||\mathbf{1}_B||_X} \left[\int_B \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}} < \infty,$$

where the supremum is taken over all balls $B \in \mathbb{B}(\mathbb{R}^n)$.

Definition 3.2. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, and $d \in (0, \infty)$. Then the space $\mathcal{L}^{\text{low}}_{X,q,0,d}(\mathbb{R}^n)$ is defined to be the set of all the $f \in L^q_{\text{loc}}(\mathbb{R}^n)$ such that

$$||f||_{\mathcal{L}_{X,q,0,d}^{\text{low}}(\mathbb{R}^n)} := \sup \left\| \left\{ \sum_{i=1}^m \left(\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right)^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_X^{-1}$$

$$\times \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} \left\{ f(x) - \inf_{\widetilde{x} \in B_j} f(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}}$$

is finite, where the supremum is taken over all $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$, and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$.

Remark 3.3. Let X be a concave ball quasi-Banach function space, $q \in [1, \infty)$, and $d \in (0, 1]$. In this case, by a proof similar to that of [78, Proposition 3.7], we find that $\mathcal{L}^{\text{low}}_{X,q,0,d}(\mathbb{R}^n) = \mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)$ with equivalent quasi-norms.

Next, we show that $\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n) \subset \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}^{\text{low}}_{X,q,0,d}(\mathbb{R}^n) \subset \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

Proposition 3.4. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $d \in (0, \infty)$. Then $\mathcal{L}^{low}_{X,a,0}(\mathbb{R}^n) \subset \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}^{low}_{X,a,0,d}(\mathbb{R}^n) \subset \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

Proof. Let X, q, s, and d be the same as in the present proposition. From (2.2), the fact that $P_B^{(s)}(P) = P$ for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ and any $P \in \mathcal{P}_s(\mathbb{R}^n)$, and the Hölder inequality, we deduce that, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any ball $B \in \mathbb{B}(\mathbb{R}^n)$,

$$(3.1) \qquad \left[\int_{B} \left| f(x) - P_{B}^{(s)}(f)(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} + \left[\int_{B} \left| P_{B}^{(s)}(f)(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} + \left\| P_{B}^{(s)} \left(f - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right) \right\|_{L^{\infty}(B)}$$

$$\lesssim \left[\int_{B} \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} + \int_{B} \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\} dx$$

$$\lesssim \left[\int_{B} \left\{ f(x) - \inf_{\widetilde{x} \in B} f(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}.$$

This, together with the definitions of $\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)$, $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$, $\mathcal{L}^{\text{low}}_{X,q,0,d}(\mathbb{R}^n)$, and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, then finishes the proof of Proposition 3.4.

Now, we present the main result of this section.

Theorem 3.5. Let $\delta \in (0,1]$ and g be the Littlewood–Paley g-function in (2.8). Let both X and $p_- \in (\frac{n}{n+\delta}, \infty)$ satisfy Assumption 2.8, $q \in (1, \infty)$, and $d \in (\frac{n}{n+\delta}, \infty)$. Then, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ [resp., $f \in \mathcal{L}_{X,q,0}(\mathbb{R}^n)$], g(f) is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

(3.2)
$$||g(f)||_{\mathcal{L}_{X_q,0,d}^{low}(\mathbb{R}^n)} \le C||f||_{\mathcal{L}_{X_q,0,d}(\mathbb{R}^n)}$$

$$\left[\text{resp., } \|g(f)\|_{\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{X,q,0}(\mathbb{R}^n)}\right].$$

To show Theorem 3.5, we first establish five technical lemmas. The following two lemmas are special cases, respectively, of [37, Lemmas 2.20 and 2.21].

Lemma 3.6. Let $q \in [1, \infty)$ and $\theta \in (0, 1)$. Then there exists a positive constant C such that, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any ball $B \subset \mathbb{R}^n$,

$$\sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} |f(x) - f_{B}|^{q} dx \right]^{\frac{1}{q}} \le C \sum_{k=1}^{\infty} \theta^{k} \left[\int_{2^{k}B} |f(x) - f_{2^{k}B}|^{q} dx \right]^{\frac{1}{q}}.$$

Lemma 3.7. Let $\delta \in (0, \infty)$. Then there exists a positive constant C such that, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and any ball $B(y, r) \subset \mathbb{R}^n$ with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$,

$$\int_{\mathbb{R}^n \setminus B(y,r)} \frac{r^{\delta} |f(z) - f_{B(y,r)}|}{|y - z|^{n + \delta}} \, dz \le C \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \int_{2^k B(y,r)} \left| f(z) - f_{2^k B(y,r)} \right| \, dz.$$

The following lemma on the boundedness of the Littlewood–Paley *g*-function on Lebesgue spaces is well known (see, for instance, [71, p. 312, (3.8)]).

Lemma 3.8. Let $q \in (1, \infty)$ and g be the Littlewood–Paley g-function in (2.8). Then g is bounded on $L^q(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in L^q(\mathbb{R}^n)$,

$$||g(f)||_{L^q(\mathbb{R}^n)} \le C||f||_{L^q(\mathbb{R}^n)}.$$

We also need the following lemma, which can be easily obtained by (2.4) and the fact that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$, any $\beta \in [1, \infty)$, and any $r \in (0, \infty)$, $\mathbf{1}_{\beta B} \leq (\beta + 1)^{\frac{dn}{r}} [\mathcal{M}(\mathbf{1}_B)]^{\frac{d}{r}}$ (see [7, Remark 2.11 (iii)] for more details).

Lemma 3.9. Let both X and p_- satisfy Assumption 2.8, and $d \in (0, \infty)$. Then, for any $\beta \in [1, \infty)$, any $r \in (0, \min\{d, p_-\})$, any sequence $\{B_j\}_{j \in \mathbb{N}} \subset \mathbb{B}(\mathbb{R}^n)$, and any $\{\lambda_i\}_{i \in \mathbb{N}} \subset [0, \infty)$,

$$\left\| \left(\sum_{j \in \mathbb{N}} \lambda_j^d \mathbf{1}_{\beta B_j} \right)^{\frac{1}{d}} \right\|_{X} \leq (2\beta)^{\frac{n}{r}} \left\| \left(\sum_{j \in \mathbb{N}} \lambda_j^d \mathbf{1}_{B_j} \right)^{\frac{1}{d}} \right\|_{X}.$$

Moreover, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ and any $r \in (0, p_-)$, $||\mathbf{1}_{\beta B}||_X \leq (2\beta)^{\frac{n}{r}}||\mathbf{1}_B||_X$.

Via borrowing some ideas from [66, 74], we have the following lemma which plays an important role in the proof of Theorem 3.5.

Lemma 3.10. Let $q \in (1, \infty)$, $\delta \in (0, 1]$, g be the Littlewood–Paley g-function in (2.8), and $f \in L^1_{loc}(\mathbb{R}^n)$. If $g(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then there exists a positive constant C, independent of both f and x_0 , such that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ containing x_0 ,

$$(3.3) \qquad \left[\int_{B} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \le C \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \left[\int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right|^{q} dz \right]^{\frac{1}{q}}.$$

Proof. Let q, δ , and g be the same as in the present lemma. Let $f \in L^1_{loc}(\mathbb{R}^n)$ and assume that there exists one $x_0 \in \mathbb{R}^n$ such that $g(f)(x_0) < \infty$. Also, let B := B(y, r) be any given ball of \mathbb{R}^n with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$ such that $x_0 \in B$. Moreover, for any $x \in \mathbb{R}^n$, let

$$g_r(f)(x) := \left[\int_0^{4r} |f * \varphi_t(x)|^2 \frac{dt}{t} \right]^{\frac{1}{2}}$$

and

$$g_{\infty}(f)(x) := \left[\int_{4r}^{\infty} |f * \varphi_t(x)|^2 \frac{dt}{t} \right]^{\frac{1}{2}}.$$

Then, by (2.5) and $g(f)(x_0) < \infty$ [and hence $g_\infty(f)(x_0) < \infty$], we conclude that, for any $x \in B$,

$$\begin{split} 0 &\leq g(f)(x) - \inf_{\widetilde{x} \in B} g(f)(\widetilde{x}) \\ &\leq g_r(f)(x) + g_{\infty}(f)(x) - \inf_{\widetilde{x} \in B} g_{\infty}(f)(\widetilde{x}) \\ &\leq g_r(f - f_{8B})(x) + \sup_{\{\widetilde{x} \in B: \ g_{\infty}(f)(\widetilde{x}) < \infty\}} |g_{\infty}(f)(x) - g_{\infty}(f)(\widetilde{x})| \\ &\leq g_r\left([f - f_{8B}]\mathbf{1}_{8B}\right)(x) + g_r\left([f - f_{8B}]\mathbf{1}_{\mathbb{R}^n \setminus 8B}\right)(x) \\ &+ \sup_{\{\widetilde{x} \in B: \ g_{\infty}(f)(\widetilde{x}) < \infty\}} |g_{\infty}(f)(x) - g_{\infty}(f)(\widetilde{x})| \end{split}$$

and hence

(3.4)
$$\left[\int_{B} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} |g_{r}([f - f_{8B}] \mathbf{1}_{8B})(x)|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left[\int_{B} |g_{r}([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \setminus 8B})(x)|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left\{ \int_{B} \left[\sup_{\{\widetilde{x} \in B: g_{\infty}(f)(\widetilde{x}) < \infty\}} |g_{\infty}(f)(x) - g_{\infty}(f)(\widetilde{x})| \right]^{q} dx \right\}^{\frac{1}{q}}$$

$$=: G_{1} + G_{2} + G_{3}.$$

We first estimate G_1 . Indeed, from Lemma 3.8, we deduce that

(3.5)
$$G_{1} \leq \left[\frac{1}{|B|} \int_{\mathbb{R}^{n}} |g([f - f_{8B}] \mathbf{1}_{8B})(x)|^{q} dx\right]^{\frac{1}{q}} \lesssim \left[\int_{8B} |f(x) - f_{8B}|^{q} dx \right]^{\frac{1}{q}}.$$

This is a desired estimate of G_1 .

Next, we consider G_2 . Notice that, for any $z \in \mathbb{R}^n \setminus 8B$ and $x \in B$,

$$|x-z| \sim |y-z|$$
.

Using this, (2.6), and Lemma 3.7, we conclude that, for any $x \in B$,

$$g_{r}\left(\left[f-f_{8B}\right]\mathbf{1}_{\mathbb{R}^{n}\backslash 8B}\right)(x)$$

$$\leq \left\{\int_{0}^{4r} \left[\int_{\mathbb{R}^{n}\backslash 8B} |f(z)-f_{8B}| \left|\frac{1}{t^{n}}\varphi\left(\frac{x-z}{t}\right)\right| dz\right]^{2} \frac{dt}{t}\right\}^{\frac{1}{2}}$$

$$\lesssim \left\{\int_{0}^{4r} \left[\int_{\mathbb{R}^{n}\backslash 8B} \frac{|f(z)-f_{8B}|}{(t+|x-z|)^{n+\delta}} dz\right]^{2} t^{2\delta-1} dt\right\}^{\frac{1}{2}}$$

$$\lesssim \int_{\mathbb{R}^{n}\backslash 8B} \frac{r^{\delta}|f(z)-f_{8B}|}{|y-z|^{n+\delta}} dz \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k}B} |f(z)-f_{2^{k}B}| dz$$

and hence

(3.6)
$$G_2 \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \int_{2^k B} |f(z) - f_{2^k B}| dz.$$

This is a desired estimate of G_2 .

Now, we estimate G_3 . To this end, we first consider $|g_{\infty}(f)(x) - g_{\infty}(f)(\widetilde{x})|$ for any $x \in B$ and any $\widetilde{x} \in B$ such that $g_{\infty}(f)(\widetilde{x}) < \infty$. Indeed, from the fact that $g(f) \geq 0$ everywhere on \mathbb{R}^n , we deduce that, for any $x \in B$ and any $\widetilde{x} \in B$ such that $g_{\infty}(f)(\widetilde{x}) < \infty$,

$$(3.7) |g_{\infty}(f)(x) - g_{\infty}(f)(\widetilde{x})| = \left| \left[\int_{4r}^{\infty} |f * \varphi_{t}(x)|^{2} \frac{dt}{t} \right]^{\frac{1}{2}} - \left[\int_{4r}^{\infty} |f * \varphi_{t}(\widetilde{x})|^{2} \frac{dt}{t} \right]^{\frac{1}{2}} \right|$$

$$\leq \left[\int_{4r}^{\infty} |f * \varphi_{t}(x) - f * \varphi_{t}(\widetilde{x})|^{2} \frac{dt}{t} \right]^{\frac{1}{2}}.$$

Noticing that, for any $z \in \mathbb{R}^n$ and $t \in (0, \infty)$, $|\varphi_t(z)| \leq t^{-n}$, by this, (2.5), (2.6), (2.7), the mean value theorem, and the fact that, for any $\theta \in (0, 1)$, $z \in \mathbb{R}^n \setminus BB$, and $x, \widetilde{x} \in B$,

$$|x - z + \theta(x - \widetilde{x})| \sim |y - z|,$$

we find that, for any $t \in (4r, \infty)$, $x \in B$, and $\widetilde{x} \in B$ such that $g_{\infty}(f)(\widetilde{x}) < \infty$, there exists a $\theta \in (0, 1)$ such that

$$\begin{split} |f * \varphi_{t}(x) - f * \varphi_{t}(\widetilde{x})| \\ &= |(f - f_{8B}) * \varphi_{t}(x) - (f - f_{8B}) * \varphi_{t}(\widetilde{x})| \\ &\leq \left| ([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \backslash 8B}) * \varphi_{t}(x) - ([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \backslash 8B}) * \varphi_{t}(\widetilde{x}) \right| \\ &+ |([f - f_{8B}] \mathbf{1}_{8B}) * \varphi_{t}(x)| + |([f - f_{8B}] \mathbf{1}_{8B}) * \varphi_{t}(\widetilde{x})| \\ &\leq \int_{\mathbb{R}^{n} \backslash 8B} |f(z) - f_{8B}| \left| \varphi\left(\frac{x - z}{t}\right) - \varphi\left(\frac{\widetilde{x} - z}{t}\right) \right| \frac{dz}{t^{n}} \\ &+ \int_{8B} |f(z) - f_{8B}| \left| \varphi\left(\frac{x - z}{t}\right) \right| \frac{dz}{t^{n}} + \int_{8B} |f(z) - f_{8B}| \left| \varphi\left(\frac{\widetilde{x} - z}{t}\right) \right| \frac{dz}{t^{n}} \\ &\lesssim \int_{\mathbb{R}^{n} \backslash 8B} \frac{t^{\delta} |f(z) - f_{8B}| |x - \widetilde{x}|}{(t + |x - z + \theta(x - \widetilde{x})|^{n+1+\delta}} dz + \int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} dz \\ &\sim \int_{\mathbb{R}^{n} \backslash 8B} |f(z) - f_{8B}| \frac{t^{\delta} |x - \widetilde{x}|}{(t + |y - z|)^{n+1+\delta}} dz + \int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} dz. \end{split}$$

By this, (3.7), the Minkowski inequality, and Lemma 3.7, we conclude that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\infty}(f)(\widetilde{x}) < \infty$,

$$(3.8) |g_{\infty}(f)(x) - g_{\infty}(f)(\overline{x})|$$

$$\leq \left\{ \int_{4r}^{\infty} \left[\int_{\mathbb{R}^{n} \setminus 8B} |f(z) - f_{8B}| \frac{t^{\delta} |x - \overline{x}|}{(t + |y - z|)^{n+1+\delta}} dz \right]^{2} \frac{dt}{t} \right\}^{\frac{1}{2}}$$

$$+ \left\{ \int_{4r}^{\infty} \left[\int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} dz \right]^{2} \frac{dt}{t} \right\}^{\frac{1}{2}}$$

$$\leq \int_{\mathbb{R}^{n} \setminus 8B} r |f(z) - f_{8B}| \left[\int_{4r}^{\infty} \frac{t^{2\delta - 1}}{(t + |y - z|)^{2n+2+2\delta}} dt \right]^{\frac{1}{2}} dz$$

$$+ \int_{8B} |f(z) - f_{8B}| \left(\int_{4r}^{\infty} \frac{1}{t^{2n+1}} dt \right)^{\frac{1}{2}} dz$$

$$\sim \int_{\mathbb{R}^{n} \setminus 8B} r |f(z) - f_{8B}| \left[\left(\int_{4r}^{|y - z|} + \int_{|y - z|}^{\infty} \right) \frac{t^{2\delta - 1}}{(t + |y - z|)^{2n+2+2\delta}} dt \right]^{\frac{1}{2}} dz$$

$$+ \frac{1}{r^{n}} \int_{8B} |f(z) - f_{8B}| dz$$

$$\sim \int_{\mathbb{R}^{n} \setminus 8B} \frac{r |f(z) - f_{8B}|}{|y - z|^{n+1}} dz + \int_{8B} |f(z) - f_{8B}| dz$$

$$\lesssim \sum_{k=1}^{\infty} \frac{1}{2^k} \int_{2^{k+1}B} |f(z) - f_{2^{k+1}B}| dz,$$

which, combined with (3.4), (3.5), (3.6), $\delta \in (0, 1]$, and the Hölder inequality, further implies that (3.3) holds true. This finishes the proof of Lemma 3.10.

Next, we show Theorem 3.5

Proof of Theorem 3.5. Let δ , g, X, p_- , q, and d be the same as in the present theorem. We only show the case $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ because the proof of $f \in \mathcal{L}_{X,q,0}(\mathbb{R}^n)$ is similar. We first claim that, if $g(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then g(f) is finite almost everywhere. To this end, let $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$ satisfy that $x_0 \in B_j$ for any $j \in \mathbb{N}$, and $\{\lambda_j\}_{j=1}^m \subset [0,\infty)$ that $\sum_{j\in\mathbb{N}} \lambda_j \neq 0$. Moreover, for any $i \in \{1,\ldots,m\}$ and $k \in \mathbb{N}$, let $\lambda_{i,k} := \frac{\lambda_i \|\mathbf{1}_{2^k B_i}\|_X}{\|\mathbf{1}_{B_i}\|_X}$. Then, from Lemma 3.9, we deduce that, for any given $r \in (0, \min\{d, p_-\})$ and any $k \in \mathbb{N}$,

(3.9)
$$\left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^{k}B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{2^{k}B_{i}} \right\}^{\frac{1}{d}} \right\|_{X} \lesssim 2^{\frac{kn}{r}} \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^{k}B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X} \\ \sim 2^{\frac{kn}{r}} \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}.$$

By Lemma 3.10, (3.9), the Fubini theorem, the definition of $\|\cdot\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)}$, and $\min\{d,p_-\}\in (\frac{n}{n+\delta},\infty)$, we conclude that, for any given $r\in (\frac{n}{n+\delta},\min\{d,p_-\})$,

$$(3.10) \qquad \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}^{-1} \sum_{j=1}^{m} \frac{\lambda_{j}|B_{j}|}{\|\mathbf{1}_{B_{j}}\|_{X}} \left[\int_{B_{j}} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B_{j}} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \right]^{\frac{1}{q}} \\ \lesssim \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}^{-1} \sum_{j=1}^{m} \frac{\lambda_{j}|B_{j}|}{\|\mathbf{1}_{B_{j}}\|_{X}} \sum_{k \in \mathbb{N}} \left[\frac{1}{2^{k\delta}} \int_{2^{k}B_{j}} \left| f(x) - f_{2^{k}B_{j}} \right|^{q} dx \right]^{\frac{1}{q}} \right]^{\frac{1}{q}} \\ \lesssim \sum_{k \in \mathbb{N}} 2^{k(-n-\delta+\frac{n}{r})} \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^{k}B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{2^{k}B_{j}} \right\}^{\frac{1}{q}} dx \right\}^{\frac{1}{q}} \\ \lesssim \sum_{k \in \mathbb{N}} 2^{k(-n-\delta+\frac{n}{r})} \left\| f \right\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^{n})} \lesssim \|f\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^{n})} < \infty.$$

From this, we deduce that, for any $j \in \mathbb{N}$,

$$\left[\int_{B_j} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B_j} g(f)(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}} < \infty$$

and hence $g(f)(x) < \infty$ for almost every $x \in B_j$, which, together with the arbitrariness of B_j , further implies that, for almost every $x \in \mathbb{R}^n$, $g(f)(x) < \infty$. Thus, the above claim holds true.

Now, we show that (3.2) holds true if $g(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$. Indeed, by the above claim and an argument similar to the estimation of (3.10), we conclude that, for any $m \in \mathbb{N}$, any balls $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$, and any $\{\lambda_j\}_{j=1}^m \subset [0,\infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$,

$$\left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right)^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_X^{-1} \sum_{j=1}^{m} \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} \left\{ f(x) - \inf_{\widetilde{x} \in B_j} g(f)(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}}$$

$$\lesssim \|f\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)},$$

which, combined with the definition of $\|\cdot\|_{\mathcal{L}^{low}_{X,q,0,d}(\mathbb{R}^n)}$, further implies that (3.2) holds true. This finishes the proof of Theorem 3.5.

- **Remark 3.11.** (i) Let $p \in (\frac{n}{n+1}, \infty)$, $X := L^p(\mathbb{R}^n)$, $\delta = 1$, and $q \in (1, \infty)$. In this case, $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = C_{\frac{1}{p}-1,q,0}(\mathbb{R}^n)$ and hence the corresponding result on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ of Theorem 3.5 coincides with [66, Theorem 1]. Moreover, when p = 1, we have $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = \text{BMO}(\mathbb{R}^n)$ and hence the corresponding result on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ of Theorem 3.5 in this case coincides with [47, Corollary 1.1].
 - (ii) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Littlewood–Paley g-function is bounded on either $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ or $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

4 Lusin-Area Functions on Both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$

In this section, we study the boundedness of the Lusin-area function on both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$. To be precise, we prove the following theorem.

Theorem 4.1. Let $\delta \in (0,1]$ and S be the Lusin-area function in (2.9). Let both X and $p_- \in (\frac{n}{n+\delta},\infty)$ satisfy Assumption 2.8, $q \in (1,\infty)$, and $d \in (\frac{n}{n+\delta},\infty)$. Then, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$ [resp., $f \in \mathcal{L}_{X,q,0}(\mathbb{R}^n)$], S(f) is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$||S(f)||_{\mathcal{L}_{X,q,0,d}^{\text{low}}(\mathbb{R}^n)} \le C||f||_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)}$$

$$\left[\text{resp., } \|S(f)\|_{\mathcal{L}^{\text{low}}_{X,q,0}(\mathbb{R}^n)} \le C\|f\|_{\mathcal{L}_{X,q,0}(\mathbb{R}^n)}\right].$$

To show Theorem 4.1, we establish two technique lemmas first. We begin with the boundedness of the Lusin-area function on Lebesgue spaces (see, for instance, [62, p. 46]).

Lemma 4.2. Let $q \in (1, \infty)$ and S be the Lusin-area function in (2.9). Then S is bounded on $L^q(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in L^q(\mathbb{R}^n)$,

$$||S(f)||_{L^q(\mathbb{R}^n)} \le C||f||_{L^q(\mathbb{R}^n)}.$$

We also need the following technical lemma.

Lemma 4.3. Let $q \in (1, \infty)$, $\delta \in (0, 1]$, S be the Lusin-area function in (2.8), and $f \in L^1_{loc}(\mathbb{R}^n)$. If $S(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then there exists a positive constant C, independent of both f and x_0 , such that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ containing x_0 ,

$$(4.1) \qquad \left[\int_{B} \left\{ S(f)(x) - \inf_{\widetilde{x} \in B} S(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \leq C \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \left[\int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right|^{q} dz \right]^{\frac{1}{q}}.$$

Proof. Let q, δ , and S be the same as in the present lemma. Let $f \in L^1_{loc}(\mathbb{R}^n)$ and assume that there exists one $x_0 \in \mathbb{R}^n$ such that $S(f)(x_0) < \infty$. Also, let B := B(y, r) be any given ball of \mathbb{R}^n with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$ such that $x_0 \in B$. Moreover, for any $x \in \mathbb{R}^n$, let

$$S_r(f)(x) := \left[\int_0^{4r} \int_{|\xi - x| < t} |f * \varphi_t(\xi)|^2 \frac{d\xi \, dt}{t^{n+1}} \right]^{\frac{1}{2}}$$

and

$$S_{\infty}(f)(x) := \left[\int_{4r}^{\infty} \int_{|\xi-x| < t} |f * \varphi_t(\xi)|^2 \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}.$$

Then, using (2.5) and $S(f)(x_0) < \infty$ [and hence $S_{\infty}(f)(x_0) < \infty$], we conclude that, for any $x \in B$,

$$\begin{split} 0 &\leq S(f)(x) - \inf_{\widetilde{x} \in B} S(f)(\widetilde{x}) \\ &\leq S_r(f)(x) + S_{\infty}(f)(x) - \inf_{\widetilde{x} \in B} S_{\infty}(f)(\widetilde{x}) \\ &\leq S_r\left([f - f_{8B}]\mathbf{1}_{8B}\right)(x) + S_r\left([f - f_{8B}]\mathbf{1}_{\mathbb{R}^n \setminus 8B}\right)(x) \\ &+ \sup_{\{\widetilde{x} \in B: \ S_{\infty}(f)(\widetilde{x}) < \infty\}} |S_{\infty}(f)(x) - S_{\infty}(f)(\widetilde{x})| \end{split}$$

and hence

$$(4.2) \qquad \left[\int_{B} \left\{ S(f)(x) - \inf_{\widetilde{x} \in B} S(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} |S_{r}([f - f_{8B}] \mathbf{1}_{8B})(x)|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left[\int_{B} \left| S_{r}([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \setminus 8B})(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left\{ \int_{B} \left[\sup_{(\widetilde{x} \in B: S_{\infty}(f)(\widetilde{x}) < \infty)} |S_{\infty}(f)(x) - S_{\infty}(f)(\widetilde{x})| \right]^{q} dx \right\}^{\frac{1}{q}}$$

$$=: S_{1} + S_{2} + S_{3}.$$

We first estimate S_1 . Indeed, from Lemma 4.2, we deduce that

(4.3)
$$S_{1} \leq \left[\frac{1}{|B|} \int_{\mathbb{R}^{n}} |S([f - f_{8B}] \mathbf{1}_{8B})(x)|^{q} dx\right]^{\frac{1}{q}} \lesssim \left[\int_{8B} |f(x) - f_{8B}|^{q} dx\right]^{\frac{1}{q}}.$$

This is a desired estimate of S_1 .

Next, we consider S₂. Indeed, noticing that, for any $t \in (0, 4r)$, $\xi \in B(x, t)$, and $z \in \mathbb{R}^n \setminus 8B$,

$$\begin{split} t + |z - \xi| &\geq t + |z - y| - |y - x| - |x - \xi| \\ &\geq t + \frac{|z - y|}{2} + 4r - |y - x| - |x - \xi| \gtrsim |z - y|, \end{split}$$

by this, (2.6), and Lemma 3.7, we conclude that, for any $x \in B$,

$$\begin{split} &S_{r}\left([f-f_{8B}]\mathbf{1}_{\mathbb{R}^{n}\backslash 8B}\right)(x) \\ &\leq \left\{ \int_{0}^{4r} \int_{|\xi-x|< t} \left[\int_{\mathbb{R}^{n}\backslash 8B} |f(z)-f_{8B}| \frac{1}{t^{n}} \varphi\left(\frac{\xi-z}{t}\right) dz \right]^{2} \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ &\lesssim \left\{ \int_{0}^{4r} \int_{|\xi-x|< t} \left[\int_{\mathbb{R}^{n}\backslash 8B} \frac{|f(z)-f_{8B}|}{t^{n}} \frac{t^{n+\delta}}{(t+|\xi-z|)^{n+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ &\lesssim \left\{ \int_{0}^{4r} \int_{|\xi-x|< t} \left[\int_{\mathbb{R}^{n}\backslash 8B} \frac{|f(z)-f_{8B}|}{|y-z|^{n+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1-2\delta}} \right\}^{\frac{1}{2}} \\ &\sim \left(\int_{0}^{4r} \int_{|\xi-x|< t} \frac{d\xi \, dt}{t^{n+1-2\delta}} \right)^{\frac{1}{2}} \int_{\mathbb{R}^{n}\backslash 8B} \frac{|f(z)-f_{8B}|}{|y-z|^{n+\delta}} \, dz \\ &\sim \int_{\mathbb{R}^{n}\backslash 8B} \frac{r^{\delta}|f(z)-f_{8B}|}{|y-z|^{n+\delta}} \, dz \lesssim \sum_{k=3}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k+1}B} \left|f(z)-f_{2^{k+1}B}\right| \, dz \end{split}$$

and hence

(4.4)
$$S_2 \lesssim \sum_{k=3}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k+1}B} \left| f(z) - f_{2^{k+1}B} \right| dz.$$

This is a desired estimate of S_2 .

Now, we estimate S_3 . To this end, we first estimate $|S_{\infty}(f)(x) - S_{\infty}(f)(\widetilde{x})|$ for any $x \in B$ and any $\widetilde{x} \in B$ such that $S_{\infty}(f)(\widetilde{x}) < \infty$. Indeed, from the fact that $S(f) \geq 0$ everywhere on \mathbb{R}^n , we deduce that, for any $x \in B$ and any $\widetilde{x} \in B$ such that $S_{\infty}(f)(\widetilde{x}) < \infty$,

$$(4.5) |S_{\infty}(f)(x) - S_{\infty}(f)(\widetilde{x})|$$

$$= \left| \left[\int_{4r}^{\infty} \int_{|\xi| < t} |f * \varphi_{t}(x + \xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}} - \left[\int_{4r}^{\infty} \int_{|\xi| < t} |f * \varphi_{t}(\widetilde{x} + \xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}} \right|$$

$$\leq \left[\int_{4r}^{\infty} \int_{|\xi| < t} |f * \varphi_{t}(x + \xi) - f * \varphi_{t}(\widetilde{x} + \xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}.$$

Moreover, noticing that, for any $t \in (4r, \infty)$, $\xi \in B(0, t)$, $x, \widetilde{x} \in B$, $z \in \mathbb{R}^n \setminus 8B$, and $\theta \in (0, 1)$,

$$t + |x + \xi - z + \theta(x - \widetilde{x})| \sim t + |y - z|$$

by this, (2.5), the mean value theorem, (2.6), (2.7), and the fact that $|\varphi_t(z)| \leq t^{-n}$ for any $z \in \mathbb{R}^n$ and $t \in (0, \infty)$, we conclude that, for any $t \in (4r, \infty)$, $\xi \in B(\mathbf{0}, t)$, $x \in B$, and $\widetilde{x} \in B$ such that $S_{\infty}(f)(\widetilde{x}) < \infty$, there exists a $\theta \in (0, 1)$ such that

$$\begin{split} |f * \varphi_{t}(x + \xi) - f * \varphi_{t}(\widetilde{x} + \xi)| \\ &= |(f - f_{8B}) * \varphi_{t}(x + \xi) - (f - f_{8B}) * \varphi_{t}(\widetilde{x} + \xi)| \\ &\leq \left| ([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \setminus 8B}) * \varphi_{t}(x + \xi) - ([f - f_{8B}] \mathbf{1}_{\mathbb{R}^{n} \setminus 8B}) * \varphi_{t}(x_{0} + \xi) \right| \\ &+ |([f - f_{8B}] \mathbf{1}_{8B}) * \varphi_{t}(x + \xi)| + |([f - f_{8B}] \mathbf{1}_{8B}) * \varphi_{t}(\widetilde{x} + \xi)| \\ &\leq \int_{\mathbb{R}^{n} \setminus 8B} |f(z) - f_{8B}| \left| \varphi \left(\frac{x + \xi - z}{t} \right) - \varphi \left(\frac{\widetilde{x} + \xi - z}{t} \right) \right| \frac{dz}{t^{n}} \\ &+ \int_{8B} |f(z) - f_{8B}| \left| \varphi \left(\frac{x + \xi - z}{t} \right) \right| \frac{dz}{t^{n}} \\ &+ \int_{8B} |f(z) - f_{8B}| \left| \varphi \left(\frac{\widetilde{x} + \xi - z}{t} \right) \right| \frac{dz}{t^{n}} \\ &\leq \int_{\mathbb{R}^{n} \setminus 8B} \frac{t^{\delta} |f(z) - f_{8B}| |x - \widetilde{x}|}{(t + |x + \xi - z + \theta(x - \widetilde{x})|)^{n+1+\delta}} dz + \int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} dz \\ &\sim \int_{\mathbb{R}^{n} \setminus 8B} |f(z) - f_{8B}| \frac{t^{\delta} |x - \widetilde{x}|}{(t + |y - z|)^{n+1+\delta}} dz + \int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} dz. \end{split}$$

Using this and an argument similar to that used in the estimation of (3.8), we conclude that, for any $x \in B$ and any $\widetilde{x} \in B$ such that $S_{\infty}(f)(\widetilde{x}) < \infty$,

$$\begin{split} & \left[\int_{4r}^{\infty} \int_{|\xi| < t} |f * \varphi_{t}(x + \xi) - f * \varphi_{t}(\widetilde{x} + \xi)|^{2} \frac{dt}{t^{n+1}} \right]^{\frac{1}{2}} \\ & \lesssim \left\{ \int_{4r}^{\infty} \int_{|\xi| < t} \left[\int_{\mathbb{R}^{n} \backslash 8B} |f(z) - f_{8B}| \frac{t^{\delta} |x - \widetilde{x}|}{(t + |y - z|)^{n+1+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ & + \left\{ \int_{4r}^{\infty} \int_{|\xi| < t} \left[\int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ & \sim \left\{ \int_{4r}^{\infty} \left[\int_{\mathbb{R}^{n} \backslash 8B} |f(z) - f_{8B}| \frac{t^{\delta} |x - \widetilde{x}|}{(t + |y - z|)^{n+1+\delta}} \, dz \right]^{2} \frac{dt}{t} \right\}^{\frac{1}{2}} \\ & + \left\{ \int_{4r}^{\infty} \left[\int_{8B} \frac{|f(z) - f_{8B}|}{t^{n}} \, dz \right]^{2} \frac{dt}{t} \right\}^{\frac{1}{2}} \\ & \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k}} \int_{2^{k+1}B} |f(z) - f_{2^{k+1}B}| \, dz. \end{split}$$

By this and (4.5), we conclude that

$$S_3 \lesssim \sum_{k=1}^{\infty} \frac{1}{2^k} \int_{2^k B} |f(z) - f_{2^k B}| dz,$$

which, together with (4.2), (4.3), (4.4), $\delta \in (0, 1]$, and the Hölder inequality, further implies that (4.1) holds true, and hence finishes the proof of Lemma 4.3.

Next, we show Theorem 4.1.

Proof of Theorem 4.1. Let all the symbols be the same as in the present theorem. Indeed, by repeating the proof of Theorem 3.5 with g and Lemma 3.10 therein replaced, respectively, by S and Lemma 4.3, we complete the proof of Theorem 4.1.

- **Remark 4.4.** (i) Let $p \in (\frac{n}{n+1}, \infty)$, $X := L^p(\mathbb{R}^n)$, $\delta = 1$, and $q \in (1, \infty)$. In this case, $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = C_{\frac{1}{p}-1,q,0}(\mathbb{R}^n)$ and hence the corresponding result on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ of Theorem 4.1 coincides with [66, Theorem 2]. Moreover, when p = 1, we have $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = \mathrm{BMO}(\mathbb{R}^n)$ and hence the corresponding result on $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ of Theorem 3.5 in this case coincides with [47, Corollary 1.2].
 - (ii) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Littlewood–Paley *g*-function is bounded on either $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ or $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

5 Littlewood–Paley g_{λ}^* -Functions on Both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$

The aim of this section is to estimate the Littlewood–Paley g_{λ}^* -function on both $\mathcal{L}_{X,q,0}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$. Moreover, even for the Campanato space $C_{\alpha,q,0}(\mathbb{R}^n)$, we obtain a better range of the index λ . To be precise, we prove the following two theorems.

Theorem 5.1. Let both X and p_- satisfy Assumption 2.8, $q \in [1, \infty)$, $\delta \in (0, 1]$, $\lambda \in (1, \infty)$, and g_{λ}^* be the same as in (2.10). If either of the following statements holds true:

(i)
$$p_- \in (\frac{n}{n+\frac{\delta}{2}}, \infty)$$
 and $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$;

(ii)
$$p_- \in (\frac{n}{n+\delta}, \frac{n}{n+\frac{\delta}{2}}]$$
 and $\lambda \in (\max\{1, \frac{2}{q}\} + \frac{2}{n}, \infty) \cap (\frac{2}{p_-}, \infty),$

then, for any $f \in \mathcal{L}_{X,q,0}(\mathbb{R}^n)$, $g_{\lambda}^*(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$\left\|g_{\lambda}^*(f)\right\|_{\mathcal{L}_{X,q,0}^{\text{low}}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{X,q,0}(\mathbb{R}^n)}.$$

Theorem 5.2. Let both X and $p_- \in (0, \infty)$ satisfy Assumption 2.8, $q \in [1, \infty)$, $d \in (0, \infty)$, $\delta \in (0, 1]$, $\lambda \in (1, \infty)$, and g_{λ}^* be the same as in (2.10). If either of the following statements holds true:

(i)
$$\min\{d, p_-\} \in (\frac{n}{n+\frac{\delta}{2}}, \infty)$$
 and $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$;

(ii)
$$\min\{d, p_-\} \in (\frac{n}{n+\delta}, \frac{n}{n+\frac{\delta}{2}}]$$
 and $\lambda \in (\max\{1, \frac{2}{q}\} + \frac{2}{n}, \infty) \cap (\frac{2}{\min\{d, p_-\}}, \infty),$

then, for any $f \in \mathcal{L}_{X,q,0,d}(\mathbb{R}^n)$, $g_{\lambda}^*(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

(5.1)
$$\|g_{\lambda}^{*}(f)\|_{\mathcal{L}_{X,q,0,d}^{\text{low}}(\mathbb{R}^{n})} \leq C\|f\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^{n})}.$$

To show the above two theorems, we give some technique lemmas first. The following lemma on the boundedness of the Littlewood–Paley g_{λ}^* -function on Lebesgue spaces is well known (see, for instance, [62, pp. 316-318]).

Lemma 5.3. Let $q \in (1, \infty)$, $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$, and g_{λ}^* be the same as in (2.10). Then g_{λ}^* is bounded on $L^q(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in L^q(\mathbb{R}^n)$,

$$\|g_{\lambda}^*(f)\|_{L^q(\mathbb{R}^n)} \le C\|f\|_{L^q(\mathbb{R}^n)}.$$

The following concept of fractional integrals can be found in, for instance, [61, p. 117].

Definition 5.4. Let $\beta \in (0, n)$ and $p \in [1, \frac{n}{\beta})$. The fractional integral operator I_{β} is defined by setting, for any $f \in L^p(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$I_{\beta}(f)(x) := \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n - \beta}} \, dy.$$

The following lemma is well known (see, for instance, [61, p. 119]).

Lemma 5.5. Let $\beta \in (0, n)$, $a \in (1, \frac{n}{\beta})$, and $\frac{1}{a} := \frac{1}{a} - \frac{\beta}{n}$. Then I_{β} is bounded from $L^{a}(\mathbb{R}^{n})$ into $L^{\widetilde{a}}(\mathbb{R}^{n})$, namely, there exists a positive constant C such that, for any $f \in L^{a}(\mathbb{R}^{n})$,

$$||I_{\beta}(f)||_{L^{\widetilde{a}}(\mathbb{R}^n)} \leq C||f||_{L^{a}(\mathbb{R}^n)}.$$

The following lemma plays an important role in the proofs of both Theorems 5.1 and 5.2.

Lemma 5.6. Let $q \in (1, \infty)$, $\delta \in (0, 1]$, $\lambda \in (1, \infty)$, g_{λ}^* be the Littlewood–Paley g_{λ}^* -function in (2.8), and $f \in L_{loc}^1(\mathbb{R}^n)$. Then the following two statements hold true:

(i) if $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$ and $g_{\lambda}^*(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then there exists a positive constant C, independent of both f and x_0 , such that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ containing x_0 ,

$$\left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \leq C \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k\delta}{2}}} \left[\int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right|^{q} dz \right]^{\frac{1}{q}};$$

(ii) if $\lambda \in (\max\{1, \frac{2}{q}\} + \frac{2}{n}, \infty) \cap (2, \infty)$ and $g_{\lambda}^*(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then there exists a positive constant C, independent of both f and x_0 , such that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ containing x_0 ,

$$\left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \\ \leq C \sum_{k=1}^{\infty} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \left[\int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right|^{q} dz \right]^{\frac{1}{q}}.$$

Proof. Let $q, \delta, \lambda, g_{\lambda}^*$, and f be the same as in the present lemma. We first show (i). We do this via borrowing some ideas from the proof of [47, Theorem 1.3]. Assume that there exists one $x_0 \in \mathbb{R}^n$ such that $g_{\lambda}^*(f)(x_0) < \infty$. Let B := B(y, r) with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$ be such that $x_0 \in B$ and, for any $k \in \mathbb{Z}_+$, let

(5.2)
$$J(y,k) := \left\{ (\xi,t) \in \mathbb{R}^{n+1}_+ : |\xi - y| < 2^{k+2} r \text{ and } 0 < t < 2^{k+2} r \right\}.$$

Moreover, for any $x, y \in \mathbb{R}^n$, let

(5.3)
$$g_{\lambda,y,0}^*(f)(x) := \left[\iint_{J(y,0)} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} |f * \varphi_t(\xi)|^2 \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}$$

and

$$g_{\lambda,y,\infty}^{*}(f)(x) := \left[\iint_{\mathbb{R}^{n+1}_{+} \setminus J(y,0)} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} |f * \varphi_{t}(\xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}.$$

Using (2.5) and $g_{\lambda}^*(f)(x_0) < \infty$ [and hence $g_{\lambda,\nu,\infty}^*(f)(x_0) < \infty$], we conclude that, for any $x \in B$,

$$\begin{split} 0 &\leq g_{\lambda,y,0}^*(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^*(f)(\widetilde{x}) \\ &\leq g_{\lambda,y,0}^*(f)(x) + g_{\lambda,y,\infty}^*(f)(x) - \inf_{\overline{x} \in B} g_{\lambda,y,\infty}^*(f)(\widetilde{x}) \\ &\leq g_{\lambda,y,0}^*(f - f_B)(x) + \sup_{\{\widetilde{x} \in B: \ g_{\lambda,y,\infty}^*(\widetilde{x}) < \infty\}} \left| g_{\lambda,y,\infty}^*(f)(x) - g_{\lambda,y,\infty}^*(f)(\widetilde{x}) \right| \\ &\leq g_{\lambda,y,0}^*\left([f - f_B] \mathbf{1}_B \right)(x) + g_{\lambda,y,0}^*\left([f - f_B] \mathbf{1}_{\mathbb{R}^n \setminus B} \right)(x) \\ &+ \sup_{\{\widetilde{x} \in B: \ g_{\lambda,y,\infty}^*(\widetilde{x}) < \infty\}} \left| g_{\lambda,y,\infty}^*(f)(x) - g_{\lambda,y,\infty}^*(f)(\widetilde{x}) \right| \end{split}$$

and hence

$$(5.4) \qquad \left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} \left| g_{\lambda,y,0}^{*}\left([f - f_{B}] \mathbf{1}_{B}\right)(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left[\int_{B} \left| g_{\lambda,y,0}^{*}\left([f - f_{B}] \mathbf{1}_{\mathbb{R}^{n} \setminus B}\right)(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left\{ \int_{B} \left[\sup_{(\widetilde{x} \in B: g_{\lambda,y,\infty}^{*}(f)(\widetilde{x}) < \infty\}} \left| g_{\lambda,y,\infty}^{*}(f)(x) - g_{\lambda,y,\infty}^{*}(f)(x_{0}) \right| \right]^{q} dx \right\}^{\frac{1}{q}}$$

$$=: \Lambda_{1} + \Lambda_{2} + \Lambda_{3}.$$

We first estimate Λ_1 . Indeed, by $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$ and Lemma 5.3, we find that

(5.5)
$$\Lambda_{1} \leq \left[\frac{1}{|B|} \int_{\mathbb{R}^{n}} \left| g_{\lambda}^{*} ([f - f_{B}] \mathbf{1}_{8B}) (x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$\lesssim \left[\int_{8B} |f(x) - f_B|^q dx \right]^{\frac{1}{q}} \lesssim \left[\int_{8B} |f(x) - f_{8B}|^q dx \right]^{\frac{1}{q}}.$$

This is a desired estimate of Λ_1 .

Now, we estimate Λ_2 . Indeed, it is easy to prove that, for any $(\xi, t) \in J(y, 0)$, $x \in B$, and $z \in \mathbb{R}^n \setminus 8B$, $|\xi - z| \sim |z - y|$ and $4B(y, r) \subset 8B(x, r)$. From this, (2.6), and Lemma 3.7, we deduce that, for any $x \in B$,

$$\begin{split} &g_{\lambda,y,0}^{*}\left(\left[f-f_{B}\right]\mathbf{1}_{\mathbb{R}^{n}\backslash 8B}\right)(x) \\ &= \left\{\int_{0}^{4r}\int_{|\xi-y|<4r}\left(\frac{t}{t+|x-\xi|}\right)^{\lambda n}\left[\int_{\mathbb{R}^{n}\backslash 8B}\frac{|f(z)-f_{B}|}{t^{n}}\varphi\left(\frac{\xi-z}{t}\right)dz\right]^{2}\frac{d\xi\,dt}{t^{n+1}}\right\}^{\frac{1}{2}} \\ &\lesssim \left\{\int_{0}^{4r}\int_{|\xi-y|<4r}\left(\frac{t}{t+|x-\xi|}\right)^{\lambda n}\left[\int_{\mathbb{R}^{n}\backslash 8B}\frac{|f(z)-f_{B}|}{(t+|\xi-z|)^{n+\delta}}dz\right]^{2}\frac{d\xi\,dt}{t^{n+1-2\delta}}\right\}^{\frac{1}{2}} \\ &\lesssim \left\{\int_{0}^{4r}\int_{|\xi-y|<4r}\left(\frac{t}{t+|x-\xi|}\right)^{\lambda n}\left[\int_{\mathbb{R}^{n}\backslash 8B}\frac{|f(z)-f_{B}|}{|z-y|^{n+\delta}}dz\right]^{2}\frac{d\xi\,dt}{t^{n+1-2\delta}}\right\}^{\frac{1}{2}} \\ &\lesssim \left[\sum_{k=3}^{\infty}\frac{1}{2^{k\delta}}\int_{2^{k+1}B}|f(z)-f_{B}|\,dz\right]\left[\int_{0}^{4r}\int_{|\xi-x|<8r}\left(\frac{t}{t+|x-\xi|}\right)^{\lambda n}\frac{1}{r^{2\delta}}\frac{d\xi\,dt}{t^{n+1-2\delta}}\right]^{\frac{1}{2}}, \end{split}$$

and, moreover, using the change of variable $s := \frac{\xi - x}{t}$ and the fact that $\lambda > 1$, we have

$$\begin{split} & \left[\int_{0}^{4r} \int_{|\xi-y| < 8r} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} \frac{1}{r^{2\delta}} \frac{d\xi \, dt}{t^{n+1-2\delta}} \right]^{\frac{1}{2}} \\ & = \left[\int_{0}^{4r} \int_{|s - \frac{y - x}{t}| < \frac{8r}{t}} \left(\frac{t}{t + t|s|} \right)^{\lambda n} \frac{t^{2\delta - 1}}{r^{2\delta}} \, ds \, dt \right]^{\frac{1}{2}} \\ & \leq \left[\int_{0}^{4r} \int_{\mathbb{R}^{n}} \frac{t^{2\delta - 1}}{(1 + |s|)^{\lambda n} r^{2\delta}} \, ds \, dt \right]^{\frac{1}{2}} \quad \sim \left(\int_{0}^{4r} \frac{t^{2\delta - 1}}{r^{2\delta}} \, dt \right)^{\frac{1}{2}} \sim 1. \end{split}$$

By this and Lemma 3.6 with $\theta := 2^{-\delta}$, we conclude that, for any $x \in B$,

$$g_{\lambda,y,0}^*\left([f-f_B]\mathbf{1}_{8B}\right)(x) \lesssim \sum_{k=3}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k+1}B} \left|f(z) - f_{2^{k+1}B}\right| dz$$

and hence

(5.6)
$$\Lambda_2 \lesssim \sum_{k=3}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k+1}B} \left| f(z) - f_{2^{k+1}B} \right| dz.$$

This is a desired estimate of Λ_2 .

Next, we estimate Λ_3 . To this end, we first estimate $|g_{\lambda,y,\infty}^*(f)(x) - g_{\lambda,y,\infty}^*(f)(\widetilde{x})|$ for any $x \in B(y,r)$ and $\widetilde{x} \in B$ such that $g_{\lambda,y,\infty}^*(f)(\widetilde{x}) < \infty$. Indeed, it is easy to show that, for any $x, \widetilde{x} \in B(y,r)$ and $(\xi,t) \in \mathbb{R}^n \setminus J(y,0)$,

$$t + |x - \xi| \sim t + |y - \xi| \sim t + |\widetilde{x} - \xi|$$

From this, the definition of $g_{\lambda,y,\infty}^*(f)$, the mean value theorem, (2.5), and (2.6), we deduce that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,y,\infty}^*(f)(\widetilde{x}) < \infty$, there exists a $\theta \in (0,1)$ such that

$$(5.7) |g_{\lambda,y,\infty}^{*}(f)(x) - g_{\lambda,y,\infty}^{*}(f)(\widetilde{x})|^{2}$$

$$\leq \iint_{\mathbb{R}^{n+1}_{+} \setminus J(y,0)} \left| \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} - \left(\frac{t}{t + |\widetilde{x} - \xi|} \right)^{\lambda n} \right| |f * \varphi_{t}(\xi)|^{2} \frac{d\xi \, dt}{t^{n+1}}$$

$$= \iint_{\mathbb{R}^{n+1}_{+} \setminus J(y,0)} \frac{|x - \xi| - |\widetilde{x} - \xi||}{(t + \theta|x - \xi| + (1 - \theta)|\widetilde{x} - \xi|)^{\lambda n+1}} |f * \varphi_{t}(\xi)|^{2} \frac{d\xi \, dt}{t^{n+1-\lambda n}}$$

$$\lesssim \iint_{\mathbb{R}^{n+1}_{+} \setminus J(y,0)} \frac{r}{(t + |y - \xi|)^{\lambda n+1}} \left[\iint_{\mathbb{R}^{n}} |f(z) - f_{B}| \frac{1}{t^{n}} \varphi\left(\frac{\xi - z}{t} \right) dz \right]^{2} \frac{d\xi \, dt}{t^{n+1-\lambda n}}$$

$$\lesssim r \sum_{k=1}^{\infty} \iint_{J(y,k)\setminus J(y,k-1)} \left(\frac{t}{2^{k+2}r} \right)^{\lambda n+1} \left[\iint_{\mathbb{R}^{n}\setminus 2^{k+3}B} \frac{|f(z) - f_{B}|}{(t + |\xi - z|)^{n+\delta}} dz \right]^{2} \frac{d\xi \, dt}{t^{n-2\delta+2}}$$

$$+ r \sum_{k=1}^{\infty} \iint_{J(y,k)\setminus J(y,k-1)} \left(\frac{t}{2^{k+2}r} \right)^{\lambda n+1} \left[\iint_{\mathbb{R}^{n}\setminus 2^{k+3}B} \frac{|f(z) - f_{B}|}{(t + |\xi - z|)^{n+\delta}} dz \right]^{2} \frac{d\xi \, dt}{t^{n-2\delta+2}}$$

$$=: g_{\lambda,y,\infty}^{*,1}(x) + g_{\lambda,y,\infty}^{*,2}(x).$$

Now, we estimate $g_{\lambda,y,\infty}^{*,1}(x)$ for any $x \in B$. Notice that, for any $k \in \mathbb{N}$, $y \in \mathbb{R}^n$, and $(\xi,t) \in J(y,k)$, $\frac{t}{2^{k+2}r} \in (0,1)$, which implies that, for any $\lambda_1, \lambda_2 \in (1,\infty)$ satisfying $\lambda_1 > \lambda_2$, $(\frac{t}{2^{k+2}r})^{\lambda_1 n+1} \le (\frac{t}{2^{k+2}r})^{\lambda_2 n+1}$. Thus, in the following estimation for $g_{\lambda,y,\infty}^{*,1}(x,\widetilde{x})$, we may always assume that $\lambda \in (\max\{1,\frac{2}{q}\},2)$ because $q \in (1,\infty)$. From this, the Minkowski inequality, Lemma 5.5(ii) with $\beta := \frac{n(\lambda-1)}{2} \in (0,n)$, $a := \frac{2}{\lambda} \in (1,q) \cap (1,\frac{n}{\beta})$, and $\widetilde{a} := 2$, Lemma 3.6 with $\theta := \frac{1}{2}$, and the Hölder inequality, we deduce that, for any $x \in B$,

$$(5.8) g_{\lambda,y,\infty}^{*,1}(x) \lesssim r \sum_{k=1}^{\infty} \int_{|\xi-y| < 2^{k+2}r} \int_{0}^{2^{k+2}r} \left(\frac{t}{2^{k+2}r}\right)^{\lambda n+1} \left[\int_{2^{k+3}B} \frac{|f(z) - f_{B}|}{(t+|\xi-z|)^{n+\delta}} dz \right]^{2} \frac{d\xi dt}{t^{n-2\delta+2}}$$

$$\lesssim \sum_{k=1}^{\infty} \frac{r}{(2^{k}r)^{\lambda n+1}} \int_{|\xi-y| < 2^{k+2}r} \left\{ \int_{2^{k+3}B} |f(z) - f_{B}| \right\}$$

$$\times \left[\int_{0}^{2^{k+2}r} \frac{t^{\lambda n-n-1+2\delta}}{(t+|\xi-z|)^{2n+2\delta}} dt \right]^{\frac{1}{2}} dz \right\}^{2} d\xi$$

$$\lesssim \sum_{k=1}^{\infty} \frac{r}{(2^{k}r)^{\lambda n+1}} \int_{|\xi-y| < 2^{k+2}r} \left\{ \int_{2^{k+3}B} |f(z) - f_{B}| \right\}$$

$$\times \left[\int_{0}^{|\xi-z|} \frac{t^{\lambda n-n-1+2\delta}}{(t+|\xi-z|)^{2n+2\delta}} dt + \int_{|\xi-z|}^{\infty} \frac{t^{\lambda n-n-1+2\delta}}{(t+|\xi-z|)^{2n+2\delta}} dt \right]^{\frac{1}{2}} dz \right\}^{2} d\xi$$

$$\lesssim \sum_{k=1}^{\infty} \frac{r}{(2^{k}r)^{\lambda n+1}} \int_{\mathbb{R}^{n}} \left[\int_{2^{k+3}B} \frac{|f(z) - f_{B}|}{|\xi-z|^{n-\frac{n}{2}(\lambda-1)}} dz \right]^{2} d\xi$$

$$\lesssim \sum_{k=1}^{\infty} \frac{r}{(2^{k}r)^{\lambda n+1}} \left[\int_{2^{k+3}B} |f(z) - f_{B}|^{\frac{2}{\lambda}} dz \right]^{\lambda}$$

$$\lesssim \left\{ \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k}{2}}} \left[\int_{2^{k+3}B} |f(z) - f_{2^{k+3}B}|^{\frac{2}{\lambda}} dz \right]^{\frac{1}{2}} \right\}^{2}$$

$$\lesssim \left\{ \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k}{2}}} \left[\int_{2^{k+3}B} |f(z) - f_{2^{k+3}B}|^{q} dz \right]^{\frac{1}{q}} \right\}^{2} .$$

This is a desired estimate of $g_{\lambda,\nu,\infty}^{*,1}(x)$.

Next, we estimate $g_{\lambda,y,\infty}^{*,2}(x)$ for any $x \in B$. Indeed, notice that, for any $z \in \mathbb{R}^n \setminus 2^{k+3}B$ and $(\xi,t) \in J(y,k) \setminus J(y,k-1)$,

$$|\xi - z| \sim |y - z|$$

Using this and the Fubini theorem, we conclude that, for any $x \in B$,

$$\begin{split} & g_{\lambda,y,\infty}^{*,2}(x) \\ & \sim \sum_{k=1}^{\infty} \frac{r}{(2^k r)^{\lambda n+1}} \iint_{J(y,k) \setminus J(y,k-1)} \left[\sum_{l=k+3}^{\infty} \int_{2^{l+1} B \setminus 2^l B} \frac{|f(z) - f_B|}{(t+|y-z|)^{n+\delta}} \, dz \right]^2 \frac{d\xi \, dt}{t^{n-\lambda n-2\delta+1}} \\ & \lesssim \sum_{k=1}^{\infty} \frac{r}{(2^k r)^{\lambda n+1}} \iint_{J(y,k)} \left[\sum_{l=k+3}^{\infty} \int_{2^{l+1} B} \frac{|f(z) - f_B|}{(2^l r)^{n+\delta}} \, dz \right]^2 \frac{d\xi \, dt}{t^{n-\lambda n-2\delta+1}} \\ & \sim \sum_{k=1}^{\infty} \frac{r}{(2^k r)^{\lambda n+1}} \iint_{J(y,k)} \frac{1}{r^{2\delta}} \left[\sum_{l=k+3}^{\infty} \frac{1}{2^{l\delta}} \int_{2^{l+1} B} |f(z) - f_B| \, dz \right]^2 \frac{d\xi \, dt}{t^{n-\lambda n-2\delta+1}} \\ & \lesssim \left\{ \sum_{k=1}^{\infty} \sum_{l=k+3}^{\infty} \left[\frac{1}{2^{l\delta}} \int_{2^{l+1} B} |f(z) - f_B| \, dz \right] \left[\frac{r^{1-2\delta}}{(2^k r)^{\lambda n+1}} \int_{0}^{2^{k+2} r} \int_{|\xi-y| < 2^{k+2} r} \frac{d\xi \, dt}{t^{n-\lambda n-2\delta+1}} \right]^{\frac{1}{2}} \right\}^2 \\ & \lesssim \left[\sum_{l=4}^{\infty} \sum_{k=1}^{l-3} \frac{1}{2^{l\delta}} \int_{2^{l+1} B} |f(z) - f_B| \, dz \right]^{\frac{k(2\delta-1)}{2}} \right]^2 \lesssim \left[\sum_{l=4}^{\infty} \frac{1}{2^{\frac{l\delta}{2}}} \int_{2^{l+1} B} |f(z) - f_B| \, dz \right]^2, \end{split}$$

where, in the last step, we used the fact that

$$\frac{1}{2^{l\delta}} \sum_{k=1}^{l-3} 2^{\frac{k(2\delta-1)}{2}} \lesssim \begin{cases} \frac{1}{2^{\frac{l}{2}}}, & \delta \in (\frac{1}{2}, 1), \\ \frac{l}{2^{l\delta}}, & \delta \in (0, \frac{1}{2}] \end{cases} \lesssim \frac{1}{2^{\frac{l\delta}{2}}}.$$

From this and Lemma 3.6 with $\theta := 2^{-\frac{\delta}{2}}$, we deduce that

(5.9)
$$g_{\lambda,y,\infty}^{*,2}(x) \lesssim \left\{ \sum_{l=4}^{\infty} \frac{1}{2^{\frac{l\delta}{2}}} \left[\int_{2^{l}B} \left| f(z) - f_{2^{l}B} \right|^{q} dz \right]^{\frac{1}{q}} \right\}^{2}.$$

This is a desired estimate of $g_{\lambda,v,\infty}^{*,2}(x)$.

Moreover, by (5.7), (5.8), and (5.9), we find that

$$\Lambda_3 \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k}{2}}} \left[\int_{2^{k+3}B} \left| f(z) - f_{2^{k+3}B} \right|^q dz \right]^{\frac{1}{q}},$$

which, combined with (5.4), (5.5), (5.6), and $\delta \in (0, 1]$, further implies that

$$\left[\int_{B}\left\{g_{\lambda}^{*}(f)(x)-\inf_{\widetilde{x}\in B}g_{\lambda}^{*}(f)(\widetilde{x})\right\}^{q}\,dx\right]^{\frac{1}{q}}\lesssim \sum_{k=1}^{\infty}\frac{1}{2^{\frac{k\delta}{2}}}\left[\int_{2^{k+1}B}\left|f(z)-f_{2^{k+1}B}\right|^{q}\,dz\right]^{\frac{1}{q}}.$$

This finishes the proof of (i).

Now, we prove (ii). To this end, assume that there exists one $x_0 \in \mathbb{R}^n$ such that $g_{\lambda}^*(f)(x_0) < \infty$ and a ball $B := B(y, r) \subset \mathbb{R}^n$ with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$ such that $x_0 \in B$, and let

$$\widetilde{J}(0) := \left\{ (\xi, t) \in \mathbb{R}_+^{n+1} : \xi \in \mathbb{R}^n \text{ and } t \in (0, 4r) \right\}.$$

Moreover, for any $x \in \mathbb{R}^n$, let $g_{\lambda, v, 0}^*(f)(x)$ be the same as in (5.3),

$$g_{\lambda,0}^*(f)(x) := \left[\iint_{\widetilde{J}(0) \setminus J(y,0)} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} |f * \varphi_t(\xi)|^2 \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}},$$

and

(5.10)
$$g_{\lambda,\infty}^{*}(f)(x) := \left[\iint_{\mathbb{R}^{n+1}_{+} \setminus \widetilde{J}(0)} \left(\frac{t}{t + |x - \xi|} \right)^{\lambda n} |f * \varphi_{t}(\xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}$$

$$= \left[\iint_{\mathbb{R}^{n+1}_{+} \setminus \widetilde{J}(0)} \left(\frac{t}{t + |\xi|} \right)^{\lambda n} |f * \varphi_{t}(x + \xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}},$$

where J(y,0) is the same as in (5.2) with k := 0. Then, using (2.5), we conclude that, for any $x \in B$,

$$\begin{split} 0 &\leq g_{\lambda}^*(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^*(f)(x) \\ &\leq g_{\lambda,y,0}^*(f)(x) + g_{\lambda,0}^*(f)(x) + g_{\lambda,\infty}^*(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda,\infty}^*(f)(x) \\ &\leq g_{\lambda,y,0}^*\left(f - f_B\right)(x) + g_{\lambda,0}^*\left(f - f_B\right)(x) \\ &\quad + \sup_{\left[\widetilde{x} \in B: \ g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty\right]} \left|g_{\lambda,\infty}^*(f)(x) - g_{\lambda,\infty}^*(f)(\widetilde{x})\right| \\ &\leq g_{\lambda,y,0}^*\left([f - f_B]\mathbf{1}_B\right)(x) + g_{\lambda,y,0}^*\left([f - f_B]\mathbf{1}_{\mathbb{R}^n \setminus B}\right)(x) + g_{\lambda,0}^*\left(f - f_B\right)(x) \\ &\quad + \sup_{\left[\widetilde{x} \in B: \ g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty\right]} \left|g_{\lambda,\infty}^*(f)(x) - g_{\lambda,\infty}^*(f)(\widetilde{x})\right| \end{split}$$

and hence

(5.11)
$$\left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\leq \left[\int_{B} \left| g_{\lambda,y,0}^{*} \left([f - f_{B}] \mathbf{1}_{B} \right) (x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left[\int_{B} \left| g_{\lambda,y,0}^{*} \left([f - f_{B}] \mathbf{1}_{\mathbb{R}^{n} \setminus B} \right) (x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left[\int_{B} \left| g_{\lambda,0}^{*} (f - f_{B})(x) \right|^{q} dx \right]^{\frac{1}{q}}$$

$$+ \left\{ \int_{B} \left[\sup_{\{\widetilde{x} \in B: \ g_{\lambda,\infty}^{*}(f)(\widetilde{x}) < \infty\}} \left| g_{\lambda,\infty}^{*}(f)(x) - g_{\lambda,\infty}^{*}(f)(\widetilde{x}) \right| \right]^{q} dx \right\}^{\frac{1}{q}}$$

$$=: \Lambda_{1} + \Lambda_{2} + \widetilde{\Lambda}_{3} + \widetilde{\Lambda}_{4},$$

where Λ_1 and Λ_2 are the same as in (5.4). Moreover, by (5.5) and (5.6), we find that

(5.12)
$$\Lambda_1 + \Lambda_2 \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k+1}B} \left| f(z) - f_{2^{k+1}B} \right| dz.$$

Next, we estimate $\widetilde{\Lambda}_3$. To this end, we first estimate $g_{\lambda,0}^*(f-f_B)(x)$ for any $x \in B$. Indeed, by (2.6), we find that, for any $x \in B$,

$$(5.13) g_{\lambda,0}^{*}(f-f_{B})(x)$$

$$= \left\{ \iint_{\widetilde{J}(0)\backslash J(y,0)} \left(\frac{t}{t+|x-\xi|} \right)^{\lambda n} |(f-f_{B}) * \varphi_{t}(\xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right\}^{\frac{1}{2}}$$

$$= \left\{ \int_{0}^{4r} \int_{\mathbb{R}^{n}\backslash 4B} \left(\frac{t}{t+|x-\xi|} \right)^{\lambda n} \left| \int_{\mathbb{R}^{n}} [f(z)-f_{B}] \varphi_{t}(\xi-z) dz \right|^{2} \frac{d\xi dt}{t^{n+1}} \right\}^{\frac{1}{2}}$$

$$\leq \left\{ \sum_{k=1}^{\infty} \int_{0}^{4r} \int_{2^{k+2}B\backslash 2^{k+1}B} \left(\frac{t}{2^{k}r} \right)^{\lambda n} \left[\int_{2^{k+3}B} \frac{|f(z)-f_{B}|}{(t+|\xi-z|)^{n+\delta}} dz \right]^{2} \frac{d\xi dt}{t^{n+1-2\delta}} \right\}^{\frac{1}{2}}$$

$$+ \left\{ \sum_{k=1}^{\infty} \int_{0}^{4r} \int_{2^{k+2}B\backslash 2^{k+1}B} \left(\frac{t}{2^{k}r} \right)^{\lambda n} \left[\int_{\mathbb{R}^{n}\backslash 2^{k+3}B} \frac{|f(z)-f_{B}|}{(t+|\xi-z|)^{n+\delta}} dz \right]^{2} \frac{d\xi dt}{t^{n+1-2\delta}} \right\}^{\frac{1}{2}}$$

$$=: \widetilde{\Lambda}_{3}^{(1)} + \widetilde{\Lambda}_{3}^{(2)}.$$

Now, we estimate $\widetilde{\Lambda}_3^{(1)}$. Indeed, notice that, for any $t \in (0, \infty)$,

$$\int_{\mathbb{R}^{n}} \frac{1}{(t+|\xi-z|)^{2n+2\delta}} d\xi$$

$$= \int_{\mathbb{R}^{n}} \frac{1}{(t+|\xi|)^{2n+2\delta}} d\xi = \int_{0}^{t} \frac{u^{n-1}}{(t+u)^{2n+2\delta}} du + \int_{t}^{\infty} \cdots$$

$$\leq \int_{0}^{t} \frac{u^{n-1}}{t^{2n+2\delta}} du + \int_{t}^{\infty} u^{-n-1-2\delta} du = t^{-n-2\delta}.$$

By this, the Minkowski inequality, $\lambda > 2$, and Lemma 3.6 with $\theta := 2^{-\frac{n(\lambda-2)}{2}}$, we conclude that

$$(5.14) \qquad \widetilde{\Lambda}_{3}^{(1)} \lesssim \left\{ \sum_{k=1}^{\infty} \int_{0}^{4r} \int_{\mathbb{R}^{n}} \left(\frac{t}{2^{k}r} \right)^{\lambda n} \left[\int_{2^{k+3}B} \frac{|f(z) - f_{B}|}{(t + |\xi - z|)^{n+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1-2\delta}} \right\}^{\frac{1}{2}}$$

$$\lesssim \left[\sum_{k=1}^{\infty} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \int_{0}^{4r} t^{\lambda n - n - 1 + 2\delta} \left\{ \int_{2^{k+3}B} \left[\int_{\mathbb{R}^{n}} \frac{|f(z) - f_{B}|^{2}}{(t + |\xi - z|)^{2n+2\delta}} \, d\xi \right]^{\frac{1}{2}} \, dz \right\}^{2} \, dt \right]^{\frac{1}{2}}$$

$$\lesssim \left\{ \sum_{k=1}^{\infty} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \int_{0}^{4r} t^{\lambda n - 2n - 1} \, dt \left[\int_{2^{k+3}B} |f(z) - f_{B}| \, dz \right]^{2} \, d\xi \right\}^{\frac{1}{2}}$$

$$\sim \left\{ \sum_{k=1}^{\infty} \frac{r^{-2n}}{2^{\lambda kn}} \left[\int_{2^{k+3}B} |f(z) - f_{B}| \, dz \right]^{2} \right\}^{\frac{1}{2}} \lesssim \sum_{k=1}^{\infty} \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \int_{2^{k+3}B} |f(z) - f_{B}| \, dz$$

$$\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \int_{2^{k}B} |f(z) - f_{2^{k}B}| \, dz.$$

This is a desired estimate of $\widetilde{\Lambda}_3^{(1)}$.

Next, we estimate $\widetilde{\Lambda}_3^{(2)}$. Indeed, for any $k \in \mathbb{N}$, $z \in \mathbb{R}^n \setminus 2^{k+3}B$, $t \in (0, 4r)$, and $\xi \in 2^{k+2}B$,

$$t + |\xi - z| \sim |y - z|$$
.

From this, the Fubini theorem, and $\lambda \in (1, \infty)$, we deduce that

$$\begin{split} \widetilde{\Lambda}_{3}^{(2)} &\lesssim \left\{ \sum_{k=1}^{\infty} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \int_{0}^{4r} \int_{2^{k+2}B} \left[\sum_{l=k+3}^{\infty} \int_{2^{l+1}B \setminus 2^{l}B} \frac{|f(z) - f_{B}|}{|y - z|^{n+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right\}^{\frac{1}{2}} \\ &\lesssim \left\{ \sum_{k=1}^{\infty} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \int_{0}^{4r} \int_{2^{k+2}B} \left[\sum_{l=k+3}^{\infty} \int_{2^{l+1}B} \frac{|f(z) - f_{B}|}{(2^{l}r)^{n+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right\}^{\frac{1}{2}} \\ &\sim \left\{ \sum_{k=1}^{\infty} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \int_{0}^{4r} \int_{2^{k+2}B} \frac{1}{r^{2\delta}} \left[\sum_{l=k+3}^{\infty} \frac{1}{2^{l\delta}} \int_{2^{l+1}B} |f(z) - f_{B}| \, dz \right]^{2} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right\}^{\frac{1}{2}} \\ &\lesssim \sum_{k=1}^{\infty} \sum_{l=k+3}^{\infty} \left[\frac{1}{2^{l\delta}} \int_{2^{l+1}B} |f(z) - f_{B}| \, dz \right] \left[\left(\frac{1}{2^{k}r} \right)^{\lambda n} \frac{1}{r^{2\delta}} \int_{0}^{4r} \int_{|\xi - y| < 2^{k+2}r} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right]^{\frac{1}{2}} \\ &\sim \sum_{l=4}^{\infty} \sum_{k=1}^{l-3} \frac{1}{2^{l\delta}} \int_{2^{l+1}B} |f(z) - f_{B}| \, dz 2^{\frac{kn(1-\lambda)}{2}} \sim \sum_{l=4}^{\infty} \frac{1}{2^{l\delta}} \int_{2^{l+1}B} |f(z) - f_{B}| \, dz, \end{split}$$

which, together with Lemma 3.6 with $\theta := 2^{-\delta}$, further implies that

(5.15)
$$\widetilde{\Lambda}_{3}^{(2)} \lesssim \sum_{l=1}^{\infty} \frac{1}{2^{l\delta}} \int_{2^{l}B} \left| f(z) - f_{2^{l}B} \right| dz.$$

This is a desired estimate of $\widetilde{\Lambda}_3^{(2)}$.

Moreover, by (5.13), (5.14), and (5.15), we find that, for any $x \in B$,

$$g_{\lambda,0}^* (f - f_B)(x) \lesssim \sum_{k=1}^{\infty} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \int_{2^k B} |f(z) - f_{2^k B}| dz$$

and hence

(5.16)
$$\widetilde{\Lambda}_{3} \lesssim \sum_{k=1}^{\infty} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right| dz.$$

This is a desired estimate of $\widetilde{\Lambda}_3$.

Now, we estimate Λ_4 . To this end, for any $k \in \mathbb{N}$ and $y \in \mathbb{R}^n$, let

$$\widetilde{J}(y,k) := J(y,k) \setminus \widetilde{J}(0),$$

where J(y, k) is the same as in (5.2). Then

(5.17)
$$\mathbb{R}^{n+1}_+ \setminus \widetilde{J}(0) = \bigcup_{k \in \mathbb{N}} \widetilde{J}(y, k).$$

We first estimate $|g_{\lambda,\infty}^*(f)(x) - g_{\lambda,\infty}^*(f)(\widetilde{x})|$ for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$. Indeed, from (5.10), we deduce that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$|g_{\lambda,\infty}^{*}(f)(x) - g_{\lambda,\infty}^{*}(f)(\widetilde{x})|$$

$$\leq \left[\iint_{\mathbb{R}^{n+1} \setminus \widetilde{J}(0)} \left(\frac{t}{t + |\xi|} \right)^{\lambda n} |f * \varphi_{t}(x + \xi) - f * \varphi_{t}(\widetilde{x} + \xi)|^{2} \frac{d\xi dt}{t^{n+1}} \right]^{\frac{1}{2}}.$$

Moreover, using (2.5), we find that, for any $t \in (4r, \infty)$, $\xi \in \mathbb{R}^n$, $x \in B$, and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$, and for any $k \in \mathbb{N}$,

$$\begin{split} |f*\varphi_t(x+\xi)-f*\varphi_t(\widetilde{x}+\xi)| &= |(f-f_B)*\varphi_t(x+\xi)-(f-f_B)*\varphi_t(\widetilde{x}+\xi)| \\ &\leq \left| ([f-f_B]\mathbf{1}_{2^{k+3}B})*\varphi_t(x+\xi)-([f-f_B]\mathbf{1}_{2^{k+3}B})*\varphi_t(\widetilde{x}+\xi) \right| \\ &+ \left| \left([f-f_B]\mathbf{1}_{\mathbb{R}^n \setminus 2^{k+3}B} \right)*\varphi_t(x+\xi)-\left([f-f_B]\mathbf{1}_{\mathbb{R}^n \setminus 2^{k+3}B} \right)*\varphi_t(\widetilde{x}+\xi) \right| \\ &\leq \int_{2^{k+3}B} |f(z)-f_B| \left| \varphi\left(\frac{x+\xi-z}{t}\right)-\varphi\left(\frac{\widetilde{x}+\xi-z}{t}\right) \right| \frac{dz}{t^n} \\ &+ \int_{\mathbb{R}^n \setminus 2^{k+3}B} |f(z)-f_B| \left| \varphi\left(\frac{x+\xi-z}{t}\right)-\varphi\left(\frac{\widetilde{x}+\xi-z}{t}\right) \right| \frac{dz}{t^n}. \end{split}$$

Thus, by this, (5.17), and (5.18), we find that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$\leq \left\{ \sum_{k=1}^{\infty} \iint_{\widetilde{J}(\mathbf{0},k)\backslash\widetilde{J}(\mathbf{0},k-1)} \left(\frac{t}{t+|\xi|} \right)^{\lambda n} \left[\int_{2^{k+3}B} |f(z) - f_B| \right. \\ \left. \times \left| \varphi\left(\frac{x+\xi-z}{t} \right) - \varphi\left(\frac{\widetilde{x}+\xi-z}{t} \right) \right| \frac{dz}{t^n} \right]^2 \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ \left. + \left\{ \sum_{k=1}^{\infty} \iint_{\widetilde{J}(\mathbf{0},k)\backslash\widetilde{J}(\mathbf{0},k-1)} \left(\frac{t}{t+|\xi|} \right)^{\lambda n} \left[\int_{\mathbb{R}^n \backslash 2^{k+3}B} |f(z) - f_B| \right. \\ \left. \times \left| \varphi\left(\frac{x+\xi-z}{t} \right) - \varphi\left(\frac{\widetilde{x}+\xi-z}{t} \right) \right| \frac{dz}{t^n} \right]^2 \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ =: \widetilde{\Lambda}_{A}^{(1)}(x,\widetilde{x}) + \widetilde{\Lambda}_{A}^{(2)}(x,\widetilde{x}).$$

Next, we estimate $\widetilde{\Lambda}_4^{(1)}(x, \widetilde{x})$ for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda, \infty}^*(f)(\widetilde{x}) < \infty$. Indeed, for any $(\xi, t) \in \mathbb{R}_+^{n+1}$ and $\lambda_1, \lambda_2 \in (1, \infty)$ satisfying $\lambda_1 > \lambda_2$, we conclude that

$$\left(\frac{t}{t+|\xi|}\right)^{\lambda_1 n} \le \left(\frac{t}{t+|\xi|}\right)^{\lambda_2 n}.$$

Using this and $q \in (1, \infty)$, in the following estimates for $\widetilde{\Lambda}_4^{(1)}(x, \widetilde{x})$, we may always assume that

(5.20)
$$\lambda \in \left(\max\left\{1, \frac{2}{q}\right\} + \frac{2}{n}, 2 + \frac{2}{n}\right) \cap \left(2, 2 + \frac{2}{n}\right).$$

Moreover, by the mean value theorem and (2.7), we conclude that, for any $t \in (4r, \infty)$, $x, \widetilde{x} \in B$, $z \in 2^{k+3}B$ with $y \in \mathbb{R}^n$ and $r \in (0, \infty)$, and $(\xi, t) \in \widetilde{J}(\mathbf{0}, k) \setminus \widetilde{J}(\mathbf{0}, k-1)$ with $k \in \mathbb{N}$, there exists a $\theta \in (0, 1)$ such that

$$\begin{split} & \left| \varphi \left(\frac{x + \xi - z}{t} \right) - \varphi \left(\frac{\widetilde{x} + \xi - z}{t} \right) \right| \\ & \lesssim \frac{t^{n + \delta} |x - \widetilde{x}|}{[t + |x + \xi - z + \theta(\widetilde{x} - x)|]^{n + 1 + \delta}} \lesssim \frac{t^{n + \delta} r}{(t + |x + \xi - z|)^{n + 1 + \delta}} \\ & \sim \frac{t^{n + \delta} r}{(t + |x + \xi - z|)^{n + 1 + \delta}}, \end{split}$$

where, in the last step, we used $t \in (4r, \infty)$. From this, the Minkowski inequality, and (5.20), we deduce that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$\begin{split} \widetilde{\Lambda}_{4}^{(1)}(x,\widetilde{x}) &\lesssim \left\{ \sum_{k=1}^{\infty} \iint_{\widetilde{J}(\mathbf{0},k) \setminus \widetilde{J}(\mathbf{0},k-1)} \left(\frac{t}{t+|\xi|} \right)^{\lambda n} \left[\int_{2^{k+3}B} \frac{t^{\delta} r |f(z) - f_{B}|}{(t+|x+\xi-z|)^{n+1+\delta}} \, dz \right]^{2} \, \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ &\lesssim \left\{ r^{2} \sum_{k=1}^{\infty} \int_{\mathbb{R}^{n}} \int_{0}^{2^{k+2}r} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \right. \\ &\times \left[\int_{2^{k+3}B} \frac{|f(z) - f_{B}|}{(t+|x+\xi-z|)^{n+1+\delta}} \, dz \right]^{2} \, \frac{dt \, d\xi}{t^{n-\lambda n+1-2\delta}} \right\}^{\frac{1}{2}} \end{split}$$

$$\leq \left[r^{2} \sum_{k=1}^{\infty} \left(\frac{1}{2^{k} r} \right)^{\lambda n} \int_{\mathbb{R}^{n}} \left\{ \int_{2^{k+3} B} |f(z) - f_{B}| \right. \\
\times \left[\int_{0}^{2^{k+2} r} \frac{t^{\lambda n - n - 1 + 2\delta}}{(t + |x + \xi - z|)^{2(n+1+\delta)}} dt \right]^{\frac{1}{2}} dz \right\}^{2} d\xi \right]^{\frac{1}{2}} \\
\leq \left[r^{2} \sum_{k=1}^{\infty} \left(\frac{1}{2^{k} r} \right)^{\lambda n} \int_{\mathbb{R}^{n}} \left\{ \int_{2^{k+3} B} |f(z) - f_{B}| \right. \\
\times \left[\left(\int_{0}^{|x + \xi - z|} + \int_{|x + \xi - z|}^{\infty} \right) \frac{t^{\lambda n - n - 1 + 2\delta}}{(t + |x + \xi - z|)^{2(n+1+\delta)}} dt \right]^{\frac{1}{2}} dz \right\}^{2} d\xi \right]^{\frac{1}{2}} \\
\sim \left\{ r^{2} \sum_{k=1}^{\infty} \left(\frac{1}{2^{k} r} \right)^{\lambda n} \int_{\mathbb{R}^{n}} \left[\int_{2^{k+3} B} \frac{|f(z) - f_{B}|}{|u - z|^{n - \frac{\lambda n - n - 2}{2}}} dz \right]^{2} du \right\}^{\frac{1}{2}},$$

which, combined with (5.20), Lemma 5.5(ii) with $\beta := \frac{\lambda n - n - 2}{2} \in (0, n)$, $a := \frac{1}{\frac{\lambda}{2} - \frac{1}{n}} \in (1, q) \cap (1, \frac{n}{\beta})$, and $\widetilde{a} := 2$, and Lemma 3.6 with $\theta := \frac{1}{2}$, further implies that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$\widetilde{\Lambda}_{4}^{(1)}(x,\widetilde{x}) \lesssim \left\{ r^{2} \sum_{k=1}^{\infty} \left(\frac{1}{2^{k} r} \right)^{\lambda n} \left[\int_{2^{k+3} B} |f(z) - f_{B}|^{a} dz \right]^{\frac{2}{a}} \right\}^{\frac{1}{2}} \\
\lesssim r \sum_{k=1}^{\infty} \left(\frac{1}{2^{k} r} \right)^{\frac{\lambda n}{2} - n(\frac{1}{2} - \frac{1}{n})} \left[\int_{2^{k+3} B} |f(z) - f_{B}|^{a} dz \right]^{\frac{1}{a}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k}} \left[\int_{2^{k+3} B} |f(z) - f_{2^{k} B}|^{q} dz \right]^{\frac{1}{q}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k}} \left[\int_{2^{k} B} |f(z) - f_{2^{k} B}|^{q} dz \right]^{\frac{1}{q}}.$$

This is a desired estimate of $\widetilde{\Lambda}_4^{(1)}(x, \overline{x})$.

Now, we estimate $\widetilde{\Lambda}_4^{(2)}(x,\widetilde{x})$. Indeed, by the mean value theorem and (2.7), we conclude that, for any $t \in (4r,\infty)$, $x \in B$, $\widetilde{x} \in B$ such that $g_{\lambda}^*(f)(\widetilde{x}) < \infty$, $(\xi,t) \in \widetilde{J}(\mathbf{0},k) \setminus \widetilde{J}(\mathbf{0},k-1)$ with $k \in \mathbb{N}$, and $z \in 2^{l+1}B \setminus 2^lB$ with integer $l \geq k+3$, there exists a $\theta \in (0,1)$ such that

$$\left| \varphi\left(\frac{x+\xi-z}{t}\right) - \varphi\left(\frac{\widetilde{x}+\xi-z}{t}\right) \right| \lesssim \frac{t^{n+\delta}|x-\widetilde{x}|}{[t+|x+\xi-z+\theta(\widetilde{x}-x)|]^{n+1+\delta}} \lesssim \frac{t^{n+\delta}r}{|x+\xi-z|^{n+1+\delta}} \sim \frac{t^{n+\delta}r}{(2^lr)^{n+1+\delta}},$$

where, in the last step, we used

$$|x+\xi-z| \geq |z-y| - |y-x| - |\xi| \geq |y-z| - r - 2^{k+2}r \sim |y-z| \sim 2^l r.$$

From this, the Fubini theorem, and $\lambda > 1$, we deduce that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$\begin{split} \widetilde{\Lambda}_{4}^{(2)}(x,\overline{x}) &\lesssim \left\{ \sum_{k=1}^{\infty} \iint_{\widetilde{J}(\mathbf{0},k)\backslash\widetilde{J}(\mathbf{0},k-1)} \left(\frac{t}{2^{k+2}r} \right)^{\lambda n} \left[\sum_{l=k+3}^{\infty} \int_{2^{l+1}B\backslash 2^{l}B} \frac{rt^{\delta}|f(z)-f_{B}|}{(2^{l}r)^{n+1+\delta}} \, dz \right]^{2} \frac{d\xi \, dt}{t^{n+1}} \right\}^{\frac{1}{2}} \\ &\lesssim \left\{ \sum_{k=1}^{\infty} \iint_{\widetilde{J}(\mathbf{0},k)\backslash\widetilde{J}(\mathbf{0},k-1)} \left(\frac{1}{2^{k}r} \right)^{\lambda n} \left[\sum_{l=k+3}^{\infty} \frac{r^{-\delta}}{2^{(1+\delta)l}} \, \int_{2^{l+1}B} |f(z)-f_{B}| \, dz \right]^{2} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right\}^{\frac{1}{2}} \\ &\lesssim \sum_{k=1}^{\infty} \sum_{l=k+3}^{\infty} \frac{r^{-\delta}}{2^{(1+\delta)l}} \, \int_{2^{l+1}B} |f(z)-f_{B}| \, dz \left[\left(\frac{1}{2^{k}r} \right)^{\lambda n} \, \iint_{\widetilde{J}(\mathbf{0},k)\backslash\widetilde{J}(\mathbf{0},k-1)} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right]^{\frac{1}{2}} \\ &\lesssim \sum_{k=1}^{\infty} \sum_{l=k+3}^{\infty} \frac{r^{-\delta}}{2^{(1+\delta)l}} \, \int_{2^{l+1}B} |f(z)-f_{B}| \, dz \left[\left(\frac{1}{2^{k}r} \right)^{\lambda n} \, \int_{0}^{2^{k+2}r} \int_{2^{k+2}B} \frac{d\xi \, dt}{t^{n-\lambda n+1-2\delta}} \right]^{\frac{1}{2}} \\ &\sim \sum_{l=4}^{\infty} \sum_{k=1}^{l-3} \frac{2^{k\delta}}{2^{(1+\delta)l}} \, \int_{2^{l+1}B} |f(z)-f_{B}| \, dz \sim \sum_{l=4}^{\infty} \frac{1}{2^{l}} \, \int_{2^{l+1}B} |f(z)-f_{B}| \, dz, \end{split}$$

which, together with Lemma 3.7, further implies that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

(5.22)
$$\widetilde{\Lambda}_{4}^{(2)}(x,\widetilde{x}) \lesssim \sum_{l=1}^{\infty} \frac{1}{2^{l}} \int_{2^{l}B} \left| f(z) - f_{2^{l}B} \right| dz.$$

This is a desired estimate of $\widetilde{\Lambda}_4^{(2)}(x, \widetilde{x})$.

Moreover, by (5.19), (5.21), (5.22), and the Hölder inequality, we conclude that, for any $x \in B$ and $\widetilde{x} \in B$ such that $g_{\lambda,\infty}^*(f)(\widetilde{x}) < \infty$,

$$\left|g_{\lambda,y,\infty}^*(f)(x) - g_{\lambda,y,\infty}^*(f)(\widetilde{x})\right| \lesssim \sum_{k=1}^{\infty} \frac{1}{2^k} \left[\int_{2^k B} \left| f(z) - f_{2^k B} \right|^q dz \right]^{\frac{1}{q}}.$$

Thus,

$$\widetilde{\Lambda}_4 \lesssim \sum_{k=1}^{\infty} \frac{1}{2^k} \left[\int_{2^k B} \left| f(z) - f_{2^k B} \right|^q dz \right]^{\frac{1}{q}},$$

which, combined with (5.11), (5.12), and (5.16), further implies that

$$\left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\lesssim \sum_{k=1}^{\infty} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \left[\int_{2^{k}B} \left| f(z) - f_{2^{k}B} \right|^{q} dz \right]^{\frac{1}{q}}.$$

This finishes the proof of (ii), and hence of Theorem 5.6.

Next, we prove the main results of this section. Indeed, the proof of Theorem 5.1 is similar to that of Theorem 5.2 and hence we only prove the second one.

Proof of Theorem 5.2. Let X, p_- , q, δ , r, d, λ , and g_{λ}^* be the same as in the present theorem. The proofs of both (i) and (ii) of the present theorem are strongly depend, respectively, on both (i) and (ii) of Lemma 5.6, and we only prove (ii) because the proof of (i) is similar.

We first claim that, if $g_{\lambda}^*(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then $g_{\lambda}^*(f)$ is finite almost everywhere. To this end, let $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$ be such that $x_0 \in B_j$ for any $j \in \mathbb{N}$, and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ such that $\sum_{j=1}^m \lambda_j \neq 0$. Also, for any $i \in \{1, \dots, m\}$ and $k \in \mathbb{N}$, let $\lambda_{i,k} := \frac{\lambda_i \|\mathbf{1}_{2k_B_i}\|_X}{\|\mathbf{1}_{B_i}\|_X}$. By Lemma 5.6, (3.9), the definition of $\|\cdot\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^n)}$, $\lambda \in (\frac{2}{\min\{d,p_-\}},\infty)$, and $\min\{d,p_-\} \in (\frac{n}{n+\delta},\infty)$, we conclude that, for any given $r \in (\frac{n}{n+\delta},\min\{d,p_-\}) \cap (\frac{2}{\lambda},\min\{d,p_-\})$,

$$\begin{aligned}
&\left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}^{-1} \sum_{j=1}^{m} \frac{\lambda_{j} |B_{j}|}{\|\mathbf{1}_{B_{j}}\|_{X}} \left[\int_{B_{j}} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B_{j}} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \\
&\lesssim \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}^{-1} \\
&\times \sum_{j=1}^{m} \frac{\lambda_{j} |B_{j}|}{\|\mathbf{1}_{B_{j}}\|_{X}} \sum_{k=1}^{\infty} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \left[\int_{2^{k}B_{j}} \left| f(z) - f_{2^{k}B_{j}} \right|^{q} dz \right]^{\frac{1}{q}} \\
&\lesssim \sum_{k \in \mathbb{N}} 2^{k(-n+\frac{n}{r})} \left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^{k}B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{2^{k}B_{i}} \right\}^{\frac{1}{d}} \right\|_{X}^{-1} \\
&\times \sum_{j=1}^{m} \frac{\lambda_{j,k} |2^{k}B_{j}|}{\|\mathbf{1}_{2^{k}B_{j}}\|_{X}} \left[\frac{1}{2^{k\delta}} + \frac{1}{2^{\frac{kn(\lambda-2)}{2}}} \right] \left[\int_{2^{k}B_{j}} \left| f(x) - f_{2^{k}B_{j}} \right|^{q} dx \right]^{\frac{1}{q}} \\
&\lesssim \sum_{k \in \mathbb{N}} \left[2^{k(-n-\delta+\frac{n}{r})} + 2^{kn(-\frac{\lambda}{2}+\frac{1}{r})} \right] \|f\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^{n})} \lesssim \|f\|_{\mathcal{L}_{X,q,0,d}(\mathbb{R}^{n})} < \infty.
\end{aligned}$$

From this, we deduce that, for any $j \in \mathbb{N}$,

$$\left[\int_{B_i} \left\{ g_{\lambda}^*(f)(x) - \inf_{\widetilde{x} \in B_j} g_{\lambda}^*(f)(\widetilde{x}) \right\}^q \, dx \right]^{\frac{1}{q}} < \infty$$

and hence $g_{\lambda}^*(f)(x) < \infty$ for almost every $x \in B_j$, which, together with the arbitrariness of B_j , further implies that, for almost every $x \in \mathbb{R}^n$, $g_{\lambda}^*(f)(x) < \infty$. Thus, the above claim holds true.

Now, we show that (5.1) holds true if $g_{\lambda}^*(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$. In this case, by the above claim and an argument similar to that used in the estimation of (5.23), we conclude that, for any $m \in \mathbb{N}$, any balls $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$, and any $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$,

$$\left\| \left\{ \sum_{i=1}^{m} \left(\frac{\lambda_{i}}{\|\mathbf{1}_{B_{i}}\|_{X}} \right)^{d} \mathbf{1}_{B_{i}} \right\}^{\frac{1}{d}} \right\|^{-1} \sum_{j=1}^{m} \frac{\lambda_{j} |B_{j}|}{\|\mathbf{1}_{B_{j}}\|_{X}} \left[\int_{B_{j}} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B_{j}} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \right]^{\frac{1}{q}} dx$$

$$\lesssim ||f||_{\mathcal{L}_{X,a,0,d}(\mathbb{R}^n)},$$

which, combined with the definition of $\|\cdot\|_{\mathcal{L}^{low}_{X,q,0,d}(\mathbb{R}^n)}$, further implies that (5.1) holds true. This finishes the proof of (ii), and hence of Theorem 5.2.

Remark 5.7. (i) Let X be concave and $d \in (0, 1]$. In this case, from Remark 2.5(i), we deduce that Theorem 5.2 coincides with Theorem 5.1.

- (ii) Let $X := L^1(\mathbb{R}^n)$, $\delta = 1$, and $q \in (1, \infty)$. In this case, $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = BMO(\mathbb{R}^n)$ and the case (i) of Theorem 5.1 coincides with [47, Corollary 1.3].
- (iii) Let $p \in (\frac{n}{n+1}, \infty)$, $X := L^p(\mathbb{R}^n)$, $\delta = 1$, and $q \in (1, \infty)$. In this case, $\mathcal{L}_{X,q,0}(\mathbb{R}^n) = C_{\frac{1}{p}-1,q,0}(\mathbb{R}^n)$ and Theorem 5.1 improves the range $\lambda \in (3+\frac{2}{n},\infty)$ in [66, Theorem 3] into $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $p \in (\frac{n}{n+1},\frac{n}{n+1/2}]$.
- (iv) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Littlewood–Paley g_{λ}^* -function is bounded on either $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ or $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

6 Applications

Theorems 3.5, 4.1, 5.1, and 5.2 can be applied to a plenty of concrete examples of ball quasi-Banach function spaces, for instance, weighted Lebesgue spaces, mixed-norm Lebesgue spaces, variable Lebesgue spaces, Orlicz spaces, and Orlicz–slice spaces. Therefore, we obtain the boundedness of Littlewood–Paley operators on these spaces. Moreover, to the best of our knowledge, all these results are new. These examples explicitly reveal that more applications of the main results of this article to new function spaces are possible. Furthermore, we also apply Lemmas 3.10, 4.3, and 5.6 to $JN_{p,q,\alpha}^{\rm con}(\mathbb{R}^n)$ [the special John–Nirenberg–Campanato spaces via congruent cubes (see, for instance, [36, Definition 1.3])], and then obtain the boundedness of Littlewood–Paley operators on $JN_{p,q,\alpha}^{\rm con}(\mathbb{R}^n)$ (see Subsection 6.6 below), which is also new.

6.1 Weighted Lebesgue Spaces

In this subsection, we apply Theorems 3.5, 4.1, 5.1, and 5.2 to weighted Lebesgue spaces. To this end, we first recall the definition of Muckenhoupt weights $A_p(\mathbb{R}^n)$ (see, for instance, [19, Definitions 7.1.2 and 7.1.3]).

Definition 6.1. Let $p \in [1, \infty)$ and ω be a nonnegative locally integrable function on \mathbb{R}^n . Then ω is called an $A_p(\mathbb{R}^n)$ weight, denoted by $\omega \in A_p(\mathbb{R}^n)$, if, when $p \in (1, \infty)$,

$$[\omega]_{A_p(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \frac{1}{|B|} \left[\int_B \omega(x) \, dx \right] \left\{ \frac{1}{|B|} \int_B [\omega(x)]^{-\frac{1}{p-1}} \, dx \right\}^{p-1} < \infty,$$

and

$$[\omega]_{A_1(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \frac{1}{|B|} \left[\int_B \omega(x) \, dx \right] \left\{ \underset{x \in \mathbb{R}^n}{\operatorname{ess sup}} [\omega(x)]^{-1} \right\} < \infty,$$

where the suprema are taken over all balls $B \in \mathbb{B}(\mathbb{R}^n)$. Moreover, the *class* $A_{\infty}(\mathbb{R}^n)$ is defined by setting

$$A_{\infty}(\mathbb{R}^n) := \bigcup_{p \in [1,\infty)} A_p(\mathbb{R}^n).$$

Definition 6.2. Let $p \in (0, \infty)$ and $\omega \in A_{\infty}(\mathbb{R}^n)$. The weighted Lebesgue space $L^p_{\omega}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{L^p_\omega(\mathbb{R}^n)} = \left[\int_{\mathbb{R}^n} |f(x)|^p \omega(x) \, dx \right]^{\frac{1}{p}} < \infty.$$

Remark 6.3. Let $p \in (0, \infty)$ and $\omega \in A_{\infty}(\mathbb{R}^n)$. As was pointed out in [56, Section 7.1], $L_{\omega}^p(\mathbb{R}^n)$ is a ball quasi-Banach space but it may not be a quasi-Banach function space.

In what follows, for any $\omega \in A_{\infty}(\mathbb{R}^n)$, let

$$q_{\omega} := \inf \left\{ q \in [1, \infty) : \ \omega \in A_q(\mathbb{R}^n) \right\}.$$

The following theorem is a corollary of Theorems 3.5, 4.1, 5.1, and 5.2.

Theorem 6.4. Let $\delta \in (0,1]$, $p \in (0,\infty)$, and $\omega \in A_{\infty}(\mathbb{R}^n)$ satisfy $p_- := \frac{p}{q_{\omega}} \in (\frac{n}{n+\delta},\infty)$, and let $d \in (\frac{n}{n+\delta},\infty)$. Let $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $\min\{d,p_-\} \in (\frac{n}{n+\frac{\delta}{2}},\infty)$, or $\lambda \in (\max\{1,\frac{2}{q}\}+\frac{2}{n},\infty) \cap (\frac{2}{\min\{d,p_-\}},\infty)$ when $\min\{d,p_-\} \in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}}]$. Let Λ be g, S, or g^*_{λ} in Definition 2.10. Then, for any $f \in \mathcal{L}_{L^p_{\omega}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, $\Lambda(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$\|\Lambda(f)\|_{\mathcal{L}^{\text{low}}_{L^p_{\omega}(\mathbb{R}^n),q,0,d(\mathbb{R}^n)}} \le C\|f\|_{L^p_{\omega}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n).$$

Similar results also hold true for the space $\mathcal{L}_{L^p_{\omega}(\mathbb{R}^n),q,0}(\mathbb{R}^n)$.

Proof. Let all the symbols be the same as in the present theorem. It is easy to show that $L^p_{\omega}(\mathbb{R}^n)$ is a ball quasi-Banach function space (see, for instance, [56, Section 7.1]). Moreover, by [3, Theorem 3.1(b)], we conclude that $L^p_{\omega}(\mathbb{R}^n)$ satisfies Assumption 2.8 with both $X := L^p_{\omega}(\mathbb{R}^n)$ and p_- in the present theorem. Thus, applying Theorems 3.5, 4.1, 5.1, and 5.2 with $X := L^p_{\omega}(\mathbb{R}^n)$, we then obtain the desired conclusions and hence complete the proof of Theorem 6.4.

6.2 Mixed-Norm Lebesgue Spaces

Recall that Benedek and Panzone [4] introduced the mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$ with $\vec{p} \in (0, \infty]^n$, which is a natural generalization of the Lebesgue space $L^p(\mathbb{R}^n)$. We refer the reader to [8, 29, 30, 31, 32] for more studies on mixed-norm Lebesgue spaces. To be precise, the mixed-norm Lebesgue space is defined as follows.

Definition 6.5. Let $\vec{p} := (p_1, \dots, p_n) \in (0, \infty]^n$. The mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{L^{\vec{p}}(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}} \cdots \left[\int_{\mathbb{R}} |f(x_1, \dots, x_n)|^{p_1} dx_1 \right]^{\frac{p_2}{p_1}} \cdots dx_n \right\}^{\frac{1}{p_n}} < \infty$$

with the usual modifications made when $p_i = \infty$ for some $i \in \{1, ..., n\}$. Moreover, let

(6.1)
$$p_{-} := \min\{p_{1}, \dots, p_{n}\}.$$

- **Remark 6.6.** (i) Let $\vec{p} \in (0, \infty]^n$. It is easy to show that $L^{\vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach function space (see, for instance, [81, p. 2047]). However, as was pointed in [81, Remark 7.21], $L^{\vec{p}}(\mathbb{R}^n)$ may not be a quasi-Banach function space.
 - (ii) We refer the reader to [23, 25] for more studies on the boundedness of (sub)linear operators on mixed-norm (Hardy) spaces.

The following theorem is a corollary of Theorems 3.5, 4.1, 5.1, and 5.2.

Theorem 6.7. Let $\delta \in (0,1]$, $\vec{p} \in (0,\infty)^n$, $p_- \in (\frac{n}{n+\delta},\infty)$ be the same as in (6.1), $q \in (1,\infty)$, and $d \in (\frac{n}{n+\delta},\infty)$. Let $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $\min\{d,p_-\}\in (\frac{n}{n+\frac{\delta}{2}},\infty)$, or $\lambda \in (\max\{1,\frac{2}{q}\}+\frac{2}{n},\infty)\cap (\frac{2}{\min\{d,p_-\}},\infty)$ when $\min\{d,p_-\}\in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}}]$. Let Λ be g, S, or g^*_{λ} in Definition 2.10. Then, for any $f \in \mathcal{L}_{L^{\vec{p}}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)$, $\Lambda(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$\|\Lambda(f)\|_{\mathcal{L}^{\mathrm{low}}_{L^{\vec{p}}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{L^{\vec{p}}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)}.$$

Similar results also hold true for the space $\mathcal{L}_{L^{\vec{p}}(\mathbb{R}^n),a,0}(\mathbb{R}^n)$.

Proof. Let all the symbols be the same as in the present theorem. It is easy to show that $L^{\vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach function space (see, for instance, [81, p. 2047]). Moreover, from [29, Lemma 3.5], we deduce that $L^{\vec{p}}(\mathbb{R}^n)$ satisfies Assumption 2.8 with both $X := L^{\vec{p}}(\mathbb{R}^n)$ and p_- in (6.1). Thus, applying Theorems 3.5, 4.1, 5.1, and 5.2 with $X := L^{\vec{p}}(\mathbb{R}^n)$, we then obtain the desired conclusions and hence complete the proof of Theorem 6.7.

6.3 Variable Lebesgue Spaces

Let $p(\cdot)$: $\mathbb{R}^n \to [0, \infty)$ be a measurable function. Recall that the *variable Lebesgue space* $L^{p(\cdot)}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{L^{p(\cdot)}(\mathbb{R}^n)} := \inf \left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \left[\frac{|f(x)|}{\lambda} \right]^{p(x)} dx \le 1 \right\} < \infty.$$

We refer the reader to [11, 15, 24, 33, 34, 35, 48] for more studies on variable Lebesgue spaces. In the reminder of this subsection, for any measurable function $p(\cdot)$: $\mathbb{R}^n \to (0, \infty)$, we always let

(6.2)
$$p_{-} := \underset{x \in \mathbb{R}^{n}}{\operatorname{ess inf}} \ p(x) \quad \text{and} \quad p_{+} := \underset{x \in \mathbb{R}^{n}}{\operatorname{ess sup}} \ p(x).$$

Recall that $p(\cdot)$: $\mathbb{R}^n \to (0, \infty)$ is said to be *globally log-Hölder continuous* if there exists a $p_\infty \in \mathbb{R}$ such that, for any $x, y \in \mathbb{R}^n$,

$$|p(x) - p(y)| \le \frac{1}{\log(e + 1/|x - y|)}$$
 and $|p(x) - p_{\infty}| \le \frac{1}{\log(e + |x|)}$,

where the implicit positive constants are independent of both x and y. As was pointed out in [56, Section 7.1], $L^{p(\cdot)}(\mathbb{R}^n)$ is a ball quasi-Banach space but it may not be a quasi-Banach function space.

The following theorem is a corollary of Theorems 3.5, 4.1, 5.1, and 5.2.

Theorem 6.8. Let $\delta \in (0,1]$, $p(\cdot): \mathbb{R}^n \to (0,\infty)$ be a globally log-Hölder continuous function satisfying $\frac{n}{n+\delta} < p_- \le p_+ < \infty$, $q \in (1,\infty)$, and $d \in (\frac{n}{n+\delta},\infty)$. Let $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $\min\{d,p_-\}\in (\frac{n}{n+\frac{\delta}{2}},\infty)$, or $\lambda \in (\max\{1,\frac{2}{q}\}+\frac{2}{n},\infty)\cap (\frac{2}{\min\{d,p_-\}},\infty)$ when $\min\{d,p_-\}\in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}}]$. Let Λ be g, g, or g^*_{λ} in Definition 2.10. Then, for any g in g in the latter case, there exists a positive constant g independent of g, such that

$$\|\Lambda(f)\|_{\mathcal{L}^{\mathrm{low}}_{L^{p(\cdot)}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)} \le C\|f\|_{\mathcal{L}_{L^{p(\cdot)}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)}.$$

Similar results also hold true for the space $\mathcal{L}_{L^{p(\cdot)}(\mathbb{R}^n),q,0}(\mathbb{R}^n)$.

Proof. Let all the symbols be the same as in the present theorem. It is easy to show that $L^{p(\cdot)}(\mathbb{R}^n)$ is a ball quasi-Banach function space (see, for instance, [56, Section 7.1]). Moreover, from [12, Corollary 2.1], we deduce that Assumption 2.8 with both $X := L^{p(\cdot)}(\mathbb{R}^n)$ and p_- in (6.2) is satisfied. Thus, applying Theorems 3.5, 4.1, 5.1, and 5.2 with $X := L^{p(\cdot)}(\mathbb{R}^n)$, we then obtain the desired conclusions and hence complete the proof of Theorem 6.8.

6.4 Orlicz Spaces

Recall that a function $\Phi: [0, \infty) \to [0, \infty)$ is called an *Orlicz function* if it is non-decreasing and satisfies $\Phi(0) = 0$, $\Phi(t) > 0$ whenever $t \in (0, \infty)$, and $\lim_{t \to \infty} \Phi(t) = \infty$. An Orlicz function Φ is said to be of *lower* (resp., *upper*) *type p* with $p \in (-\infty, \infty)$ if there exists a positive constant $C_{(p)}$, depending on p, such that, for any $t \in [0, \infty)$ and $s \in (0, 1)$ [resp., $s \in [1, \infty)$],

$$\Phi(st) \le C_{(p)} s^p \Phi(t).$$

An Orlicz function Φ is said to be of *positive lower* (resp., *upper*) *type* if it is of lower (resp., upper) type p for some $p \in (0, \infty)$. Then we recall the definition of Orlicz spaces as follows (see, for instance, [55, p. 58, Definition 2]).

Definition 6.9. Let Φ be an Orlicz function with positive lower type p_{Φ}^- and positive upper type p_{Φ}^+ . The *Orlicz space* $L^{\Phi}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{L^{\Phi}(\mathbb{R}^n)} := \inf \left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \Phi\left(\frac{|f(x)|}{\lambda}\right) dx \le 1 \right\} < \infty.$$

- **Remark 6.10.** (i) Let Φ be an Orlicz function on \mathbb{R}^n with positive lower type $p_{\Phi}^- \in (0, \infty)$ and positive upper type $p_{\Phi}^+ \in (0, \infty)$. As was pointed out in [56, Section 7.6], $L^{\Phi}(\mathbb{R}^n)$ is a quasi-Banach function space and hence a ball quasi-Banach function space.
 - (ii) We refer the reader to [2, 22, 20, 51, 57, 58, 59] for more studies on the boundedness of (sub)linear operators on Orlicz-type spaces.

The following theorem is a corollary of Theorems 3.5, 4.1, 5.1, and 5.2.

Theorem 6.11. Let $\delta \in (0,1]$, Φ be an Orlicz function with lower type p_{Φ}^- and upper type p_{Φ}^+ satisfying $\frac{n}{n+\delta} < p_{\Phi}^- \le p_{\Phi}^+ < \infty$, $q \in (1,\infty)$, and $d \in (\frac{n}{n+\delta},\infty)$. Let $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $\min\{d,p_{\Phi}^-\} \in (\frac{n}{n+\delta},\infty)$, or $\lambda \in (\max\{1,\frac{2}{q}\}+\frac{2}{n},\infty) \cap (\frac{2}{\min\{d,p_{\Phi}^-\}},\infty)$ when $\min\{d,p_{\Phi}^-\} \in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}}]$. Let Λ be g, g, or g^*_{λ} in Definition 2.10. Then, for any g is g, when g is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant g, independent of g, such that

$$||\Lambda(f)||_{\mathcal{L}_{L^{\Phi}(\mathbb{R}^n),q,0,d}^{\text{low}}(\mathbb{R}^n)} \leq C||f||_{\mathcal{L}_{L^{\Phi}(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)}.$$

Similar results also hold true for the space $\mathcal{L}_{L^{\Phi}(\mathbb{R}^n),q,0}(\mathbb{R}^n)$.

Proof. Let all the symbols be the same as in the present theorem. It is easy to show that $L^{\Phi}(\mathbb{R}^n)$ is a ball quasi-Banach function space (see, for instance, [56, Section 7.6]). Moreover, from [41, Theorem 2.10] (see also [56, Theorem 7.12]), we deduce that Assumption 2.8 with both $X := L^{\Phi}(\mathbb{R}^n)$ and $p_- := p_{\Phi}^-$ is satisfied. Thus, applying Theorems 3.5, 4.1, 5.1, and 5.2 with $X := L^{\Phi}(\mathbb{R}^n)$, we then obtain the desired conclusions and hence complete the proof of Theorem 6.11. \square

6.5 Orlicz-Slice Spaces

In this subsection, we apply the main results of this article to the Orlicz-slice space $(E_{\Phi}^r)_t(\mathbb{R}^n)$ which was first introduced in [80]. To this end, we first recall the following definition of Orlicz-slice spaces which is just [80, Definition 2.8].

Definition 6.12. Let $t, r \in (0, \infty)$ and Φ be an Orlicz function with positive lower type p_{Φ}^- and positive upper type p_{Φ}^+ . The *Orlicz-slice space* $(E_{\Phi}^r)_t(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$||f||_{(E_{\Phi}^r)_t(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}^n} \left[\frac{||f\mathbf{1}_{B(x,t)}||_{L^{\Phi}(\mathbb{R}^n)}}{||\mathbf{1}_{B(x,t)}||_{L^{\Phi}(\mathbb{R}^n)}} \right]^r dx \right\}^{\frac{1}{r}} < \infty.$$

- **Remark 6.13.** (i) Let $t, r \in (0, \infty)$ and Φ be an Orlicz function on \mathbb{R}^n with $0 < p_{\Phi}^- \le p_{\Phi}^+ < \infty$. By [80, Lemma 2.28], we find that $(E_{\Phi}^r)_t(\mathbb{R}^n)$ is a ball quasi-Banach function space, but it may not be a quasi-Banach function space (see, for instance, [81, Remark 7.43(i)]).
 - (ii) We refer the reader to [26, 27, 79] for more studies on the boundedness of (sub)linear operators on (local) Orlicz-slice spaces.

The following theorem is a corollary of Theorems 3.5, 4.1, 5.1, and 5.2.

Theorem 6.14. Let $\delta \in (0,1]$ and Φ be an Orlicz function with both lower type p_{Φ}^- and upper type p_{Φ}^+ satisfying $\frac{n}{n+\delta} < p_{\Phi}^- \le p_{\Phi}^+ < \infty$. Let $t \in (0,\infty)$, $r \in (\frac{n}{n+\delta},\infty)$, $q \in (1,\infty)$, and $d \in (\frac{n}{n+\delta},\infty)$. Let $\lambda \in (\max\{1,\frac{2}{q}\},\infty)$ when $\min\{d,p_{\Phi}^-,r\} \in (\frac{n}{n+\frac{\delta}{2}},\infty)$, or $\lambda \in (\max\{1,\frac{2}{q}\}+\frac{2}{n},\infty) \cap (\frac{2}{\min\{d,p_{\Phi}^-,r\}},\infty)$ when $\min\{d,p_{\Phi}^-,r\} \in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}}]$. Let Λ be g, S, or g_{λ}^* in Definition 2.10. Then, for any $f \in (\frac{n}{n+\delta},\frac{n}{n+\frac{\delta}{2}})$

 $\mathcal{L}_{(E_{\Phi}^{r})_{t}(\mathbb{R}^{n}),q,0,d}(\mathbb{R}^{n})$, $\Lambda(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$\|\Lambda(f)\|_{\mathcal{L}^{\text{low}}_{(E^r_{\Phi})_t(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{(E^r_{\Phi})_t(\mathbb{R}^n),q,0,d}(\mathbb{R}^n)}.$$

Similar results also hold true for the space $\mathcal{L}_{(E_{\Phi}^r),q,0}(\mathbb{R}^n)$.

Proof. Let all the symbols be the same as in the present theorem. By [80, Lemma 2.28], we find that $(E_{\Phi}^r)_t(\mathbb{R}^n)$ is a ball quasi-Banach function space. Moreover, from [80, Lemma 4.3], we deduce that Assumption 2.8 with both $X := (E_{\Phi}^r)_t(\mathbb{R}^n)$ and $p_- := \min\{r, p_{\Phi}^-\}$ is satisfied. Thus, applying Theorems 3.5, 4.1, 5.1, and 5.2 with $X := (E_{\Phi}^r)_t(\mathbb{R}^n)$, we then obtain the desired conclusions and hence complete the proof of Theorem 6.11.

6.6 Special John-Nirenberg-Campanato Spaces via Congruent Cubes

Recall that John and Nirenberg [39] introduced the well-known space BMO (\mathbb{R}^n) and, in the same article, they also introduced the mysterious John–Nirenberg space. Recall that Dafni et al. [13] showed the non-triviality of the John–Nirenberg space. Indeed, the John–Nirenberg space now attracts more and more attention. We refer the reader to [1, 17, 49, 50, 63, 67, 69, 70] for more studies on these spaces. However, it is still a challenging and open question to obtain the boundedness of some important operators on the John–Nirenberg space.

To shed some light on the boundedness of Littlewood–Paley operators on John–Nirenberg spaces, Jia et al. [36, Definition 1.3] introduced the special John–Nirenberg–Campanato space via congruent cubes, $JN_{(p,q,s)_{\alpha}}^{\rm con}(\mathbb{R}^n)$. In this subsection, we use Lemmas 3.10, 4.3, and 5.6 to obtain the boundedness of Littlewood–Paley operators on $JN_{(p,q,0)_{\alpha}}^{\rm con}(\mathbb{R}^n)$. We refer the reader to [38, 37], respectively, for the boundedness of Calderón–Zygmund operators and fractional integrals on $JN_{(p,q,s)_{\alpha}}^{\rm con}(\mathbb{R}^n)$.

In what follows, a *cube* always has a finite edge length and all its edges parallel to the coordinate axes, but it is not necessary to be open or closed. Moreover, for any $m \in \mathbb{Z}$, we use $\mathcal{D}_m(\mathbb{R}^n)$ to denote the set of all the subcubes of \mathbb{R}^n with side length 2^{-m} .

Definition 6.15. Let $p, q \in [1, \infty]$ and $\alpha \in \mathbb{R}$. Then the *special John–Nirenberg–Campanato space* via congruent cubes, $JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, is defined to be the set of all the $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ such that

$$f \|_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}^{\text{con}}$$

$$:= \begin{cases} \sup_{m \in \mathbb{Z}} \sup_{\{Q_j\}_j \subset \mathcal{D}_m(\mathbb{R}^n)} \left[\sum_j |Q_j| \left\{ |Q_j|^{-\alpha} \left[\int_{Q_j} |f(x) - f_{Q_j}|^q \ dx \right]^{\frac{1}{q}} \right\}^p \right]^{\frac{1}{p}} \\ \text{if } p \in [1, \infty), \\ \|f\|_{C_{\alpha,q,0}(\mathbb{R}^n)} \qquad \text{if } p = \infty \end{cases}$$

is finite, where $\|\cdot\|_{C_{\alpha,q,0}(\mathbb{R}^n)}$ is the same as in (2.3).

Remark 6.16. (i) Observe that $JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ is only a special case when s=0 of the special John–Nirenberg–Campanato space via congruent cubes, $JN_{(p,q,s)_{\alpha}}^{\text{con}}(\mathbb{R}^n)$, introduced in [36,

Definition 1.3]. If we use the same notation as there, $JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$ here is just the space $JN_{(p,q,0)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$ therein; here, for the simplicity of the presentation, we denote $JN_{(p,q,0)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$ simply by $JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$.

(ii) We point out that $JN^{\operatorname{con}}_{p,q,\alpha}(\mathbb{R}^n)$ [and hence $JN^{\operatorname{con}}_{(p,q,s)_\alpha}(\mathbb{R}^n)$] is not a ball quasi-Banach function space. Indeed, it is easy to show that, for any $f \in \mathscr{M}(\mathbb{R}^n)$, $\|f\|_{JN^{\operatorname{con}}_{p,q,\alpha}(\mathbb{R}^n)} = 0$ if and only if f is a constant almost everywhere on \mathbb{R}^n , which implies that $JN^{\operatorname{con}}_{p,q,\alpha}(\mathbb{R}^n)$ does not satisfy Definition 2.1(i), and hence $JN^{\operatorname{con}}_{p,q,\alpha}(\mathbb{R}^n)$ is not a ball quasi-Banach function space.

To obtain the main results of this section, we first introduce a proper subspace of $JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$, denoted by $LJN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)$.

Definition 6.17. Let $p, q \in [1, \infty]$ and $\alpha \in \mathbb{R}$. The *space LJN*^{con}_{p,q,α}(\mathbb{R}^n) is defined to be the set of all the $f \in L^q_{loc}(\mathbb{R}^n)$ such that

$$:= \begin{cases} \sup_{r \in (0,\infty)} \left[\int_{\mathbb{R}^n} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} \left\{ f(x) - \inf_{\widetilde{x} \in B(y,r)} f(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}} \right\}^p dy \right]^{\frac{1}{p}} \\ := \begin{cases} \sup_{r \in (0,\infty)} \left[\int_{\mathbb{R}^n} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} \left\{ f(x) - \inf_{\widetilde{x} \in B(y,r)} f(\widetilde{x}) \right\}^q dx \right]^{\frac{1}{q}} \right] \\ \text{if } p \in [1,\infty), \end{cases}$$

is finite.

Remark 6.18. Let $q \in [1, \infty)$ and $\alpha \in \mathbb{R}$. In this case, $LJN_{\infty,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ coincides with the space $C_{\alpha,q,*}(\mathbb{R}^n)$ which was introduced by Hu et al. in [28, Definition 1.4] and, moreover, $C_{0,q,*}(\mathbb{R}^n)$ coincides with the space BLO (\mathbb{R}^n) which was first introduced by Coifman and Rochberg in [9].

The following lemma gives an equivalent norm of $JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, which is just a special case of [36, Proposition 2.2].

Lemma 6.19. Let $p, q \in [1, \infty)$ and $\alpha \in \mathbb{R}$. Then $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$ if and only if $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ and

$$[f]_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} := \sup_{r \in (0,\infty)} \left[\int_{\mathbb{R}^n} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} |f(x) - f_{B(y,r)}|^q \ dx \right]^{\frac{1}{q}} \right\}^p \ dy \right]^{\frac{1}{p}}$$

is finite. Moreover, for any $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$,

$$||f||_{JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)} \sim [f]_{JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n)},$$

where the positive equivalence constants are independent of f.

By Lemma 6.19 and (3.1), we have the following conclusion; we omit the details here.

Proposition 6.20. Let $p \in [1, \infty]$, $q \in [1, \infty)$, and $\alpha \in \mathbb{R}$. Then

$$LJN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n) \subset JN_{p,q,\alpha}^{\mathrm{con}}(\mathbb{R}^n).$$

Next, we establish the boundedness of Littlewood–Paley g functions on $JN_{p,a,\alpha}^{\rm con}(\mathbb{R}^n)$.

Theorem 6.21. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $\alpha \in (-\infty, \frac{\delta}{n})$ with $\delta \in (0, 1]$, and g be the Littlewood–Paley g-function in (2.8). For any $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, g(f) is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

(6.3)
$$||g(f)||_{LJN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} \le C||f||_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}.$$

Proof. Let p, q, α, δ , and g be the same as in the present theorem. We only prove the case $p \in [1, \infty)$ because the proof of $p = \infty$ is similar. Let $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$. We first claim that, if $g(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then g(f) is finite almost everywhere. Indeed, using Lemma 3.10, $\alpha < \frac{\delta}{n} < \frac{\delta}{n} + \frac{1}{p}$, and the definition of $\|\cdot\|_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}$, we find that, for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ containing x_0 ,

$$|B|^{\frac{1}{p}-\alpha} \left[\int_{B} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k(-n\alpha + \frac{n}{p} + \delta)}} \left| 2^{k} B \right|^{\frac{1}{p} - \alpha} \left[\int_{2^{k} B} \left| f(z) - f_{2^{k} B} \right|^{q} dz \right]^{\frac{1}{q}}$$

$$\lesssim ||f||_{JN_{p,q,r}^{\text{con}}(\mathbb{R}^{n})} < \infty,$$

which further implies that g(f) is finite almost everywhere on B. By this and the arbitrariness of B, we conclude that, for almost every $x \in \mathbb{R}^n$, $g(f)(x) < \infty$ and hence the above claim holds true.

Now, we show that (6.3) holds true if $g(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$. Indeed, by Lemma 3.10, the above claim, the Minkowski inequality, the Hölder inequality, $\alpha \in (-\infty, \frac{\delta}{n})$, and Lemma 6.19, we conclude that, for any $r \in (0, \infty)$,

$$\left[\int_{\mathbb{R}^{n}} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} \left\{ g(f)(x) - \inf_{\widetilde{x} \in B(y,r)} g(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
\lesssim \left\{ \int_{\mathbb{R}^{n}} \left[|B(y,r)|^{-\alpha} \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \int_{2^{k}B(y,r)} \left| f(x) - f_{2^{k}B(y,r)} \right| dx \right]^{p} dy \right\}^{\frac{1}{p}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k\delta}} \left\{ \int_{\mathbb{R}^{n}} \left[|B(y,r)|^{-\alpha} \int_{2^{k}B(y,r)} \left| f(x) - f_{2^{k}B(y,r)} \right| dx \right]^{p} dy \right\}^{\frac{1}{p}} \\
\sim \sum_{k=1}^{\infty} \frac{1}{2^{k(\delta-\alpha n)}} \left\{ \int_{\mathbb{R}^{n}} \left[\left| 2^{k}B(y,r) \right|^{-\alpha} \int_{2^{k}B(y,r)} \left| f(x) - f_{2^{k}B(y,r)} \right| dx \right]^{p} dy \right\}^{\frac{1}{p}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k(\delta-\alpha n)}} \left\{ \int_{\mathbb{R}^{n}} \left[\left| 2^{k}B(y,r) \right|^{-\alpha} \int_{2^{k}B(y,r)} \left| f(x) - f_{2^{k}B(y,r)} \right|^{q} dx \right]^{\frac{p}{q}} dy \right\}^{\frac{1}{p}} \\
\lesssim ||f||_{JN_{p,q,\alpha}^{con}(\mathbb{R}^{n})}.$$

This further implies that

$$||g(f)||_{LJN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} \lesssim ||f||_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)},$$

and hence finishes the proof of Theorem 6.21.

- **Remark 6.22.** (i) Let $\delta = 1$ and $p = \infty$. In this case, from the conclusion of Theorem 6.21, we deduce the conclusion of [66, Theorem 1] because, by Theorem 6.21 in this case, we obtain $g(f) \in C_{\alpha,q,*}(\mathbb{R}^n)$ which, together with $C_{\alpha,q,*}(\mathbb{R}^n) \subset C_{\alpha,q}(\mathbb{R}^n)$, further implies $g(f) \in C_{\alpha,q}(\mathbb{R}^n)$, the conclusion of [66, Theorem 1]. Thus, a special case of Theorem 6.21 is also stronger than [66, Theorem 1].
 - (ii) Let $\delta = 1$, $p = \infty$, and $\alpha = 0$. In this case, Theorem 6.21 coincides with [47, Corollary 1.1].
- (iii) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Littlewood–Paley g-function is bounded on $JN_{(p,q,s)_a}^{\text{con}}(\mathbb{R}^n)$ (see [36, Definition 1.3] for the precise definition).

Next, we establish the boundedness of the Lusin-area function S on the space $JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, whose proof is just a repetition of that of Theorem 6.21 with both g and Lemma 3.10 therein replaced, respectively, by both S and Lemma 4.3 here; we omit details here.

Theorem 6.23. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $\alpha \in (-\infty, \frac{\delta}{n})$ with $\delta \in (0, 1]$, and S be the Lusin-area function in (2.9). For any $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, S(f) is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

$$||S(f)||_{LJN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} \le C||f||_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}.$$

- **Remark 6.24.** (i) Let $\delta = 1$ and $p = \infty$. In this case, from the conclusion of Theorem 6.23, we deduce the conclusion of [66, Theorem 2] because, by Theorem 6.23 in this case, we obtain $S(f) \in C_{\alpha,q,*}(\mathbb{R}^n)$ which, combined with $C_{\alpha,q,*}(\mathbb{R}^n) \subset C_{\alpha,q}(\mathbb{R}^n)$, further implies $S(f) \in C_{\alpha,q}(\mathbb{R}^n)$, the conclusion of [66, Theorem 2]. Thus, a special case of Theorem 6.23 is also stronger than [66, Theorem 2].
 - (ii) Let $\delta = 1$, $p = \infty$, and $\alpha = 0$. In this case, Theorem 6.23 coincides with [47, Corollary 1.2].
- (iii) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Lusin-area function is bounded on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^{n})$ (see [36, Definition 1.3] for the precise definition).

Now, we estimate the Littlewood–Paley g^*_{λ} function on the space $JN^{\mathrm{con}}_{p,q,\alpha}(\mathbb{R}^n)$.

Theorem 6.25. Let $p \in [1, \infty]$, $q \in (1, \infty)$, $\alpha \in \mathbb{R}$, $\delta \in (0, 1]$, $\lambda \in (1, \infty)$, and g_{λ}^* be the same as in (2.10). For any $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$, $g_{\lambda}^*(f)$ is either infinite everywhere or finite almost everywhere and, in the latter case, there exists a positive constant C, independent of f, such that

if either of the following statements holds true:

(i)
$$\alpha \in (-\infty, \frac{\delta}{2n})$$
 and $\lambda \in (\max\{1, \frac{2}{a}\}, \infty)$;

(ii)
$$\alpha \in [\frac{\delta}{2n}, \frac{\delta}{n})$$
 and $\lambda \in (\max\{1, \frac{2}{a}\} + \frac{2}{n}, \infty) \cap (2 + 2\alpha, \infty)$.

Proof. Let p, q, α, δ , and g_{λ}^* be the same as in the present theorem. The proofs of both (i) and (ii) of the present theorem strongly depend, respectively, on both (i) and (ii) of Lemma 5.6, and we only prove (i) because the proof of (ii) is similar. Also, we only show the case $p \in [1, \infty)$ because the proof of $p = \infty$ is similar. Let $f \in JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)$. We first claim that, if $g_{\lambda}^*(f)(x_0) < \infty$ for one $x_0 \in \mathbb{R}^n$, then $g_{\lambda}^*(f)$ is finite almost everywhere. Indeed, using Lemma 5.6(i), $\alpha \in (-\infty, \frac{\delta}{2n})$, and the definition of $\|\cdot\|_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}$, we find that, for any ball $B \subset \mathbb{R}^n$ containing x_0 ,

$$|B|^{\frac{1}{p}-\alpha} \left[\int_{B} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}}$$

$$\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{k(-n\alpha + \frac{n}{p} + \frac{\delta}{2})}} \left| 2^{k} B \right|^{\frac{1}{p}-\alpha} \left[\int_{2^{k} B} \left| f(z) - f_{2^{k} B} \right|^{q} dz \right]^{\frac{1}{q}}$$

$$\lesssim ||f||_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^{n})} < \infty,$$

which further implies that $g_{\lambda}^*(f)$ is finite almost everywhere on B. Using this and the arbitrariness of B, we find that, for almost every $x \in \mathbb{R}^n$, $g_{\lambda}^*(f)(x) < \infty$ and hence the above claim holds true.

Next, we show (6.4). Indeed, from the above claim, Lemma 5.6, the Minkowski inequality, $\alpha \in (-\infty, \frac{\delta}{2n}) \subset (-\infty, \frac{1}{2n})$, and Lemma 6.19, we deduce that, for any $r \in (0, \infty)$,

$$\left[\int_{\mathbb{R}^{n}} \left\{ |B(y,r)|^{-\alpha} \left[\int_{B(y,r)} \left\{ g_{\lambda}^{*}(f)(x) - \inf_{\widetilde{x} \in B(y,r)} g_{\lambda}^{*}(f)(\widetilde{x}) \right\}^{q} dx \right]^{\frac{1}{q}} \right\}^{q} dy \right]^{\frac{1}{p}} \\
\lesssim \left[\int_{\mathbb{R}^{n}} \left\{ |B(y,r)|^{-\alpha} \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k\delta}{2}}} \left[\int_{2^{k}B(y,r)} |f(z) - f_{2^{k}B(y,r)}|^{q} dz \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k\delta}{2}}} \left[\int_{\mathbb{R}^{n}} \left\{ |B(y,r)|^{-\alpha} \left[\int_{2^{k}B(y,r)} |f(z) - f_{2^{k}B(y,r)}|^{q} dz \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
\lesssim \sum_{k=1}^{\infty} \frac{1}{2^{\frac{k\delta}{2} - \alpha kn}} \left[\int_{\mathbb{R}^{n}} \left\{ |2^{k}B(y,r)|^{-\alpha} \left[\int_{2^{k}B(y,r)} |f(z) - f_{2^{k}B(y,r)}|^{q} dz \right]^{\frac{1}{q}} \right\}^{p} dy \right]^{\frac{1}{p}} \\
\lesssim ||f||_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^{n})} \end{aligned}$$

and hence

$$\|g_{\lambda}^*(f)\|_{LJN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)} \lesssim \|f\|_{JN_{p,q,\alpha}^{\text{con}}(\mathbb{R}^n)}.$$

This finishes the proof of Theorem 6.25.

Remark 6.26. (i) Let $\delta = 1$, $p = \infty$, and $\alpha = 0$. In this case, Theorem 6.25(i) coincides with [47, Corollary 1.3].

(ii) Let $\delta = 1$. In this case, we point out that the boundedness of the Littlewood–Paley g_{λ}^* function on the Campanato space $C_{\alpha,q,0}(\mathbb{R}^n)$ in Theorem 6.25 widens the range $\lambda \in (3+\frac{2}{n},\infty)$

in [66, Theorem 3] into $\lambda \in (\max\{1, \frac{2}{q}\}, \infty)$ when $\alpha \in (-\infty, \frac{1}{2n})$, or $\lambda \in (\max\{1, \frac{2}{q}\} + \frac{2}{n}, \infty) \cap (2 + 2\alpha, \infty)$ when $\alpha \in [\frac{1}{2n}, \frac{1}{n})$.

(iii) Let $s \in \mathbb{N}$. In this case, it is still unclear whether or not the Littlewood–Paley g_{λ}^* -function is bounded on $JN_{(p,q,s)_{\alpha}}^{\mathrm{con}}(\mathbb{R}^n)$ (see [36, Definition 1.3] for the precise definition).

Acknowledgements Hongchao Jia and Yangyang Zhang would like to thank Jin Tao for some useful discussions on the subject of this article.

Author contributions All the authors contributed to this article.

Data Availability Statement Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest The author declares that they have no conflict interests.

Code availability Not applicable.

References

- [1] D. Aalto, L. Berkovits, O. E. Kansanen and H. Yue, John–Nirenberg lemmas for a doubling measure, Studia Math. 204 (2011), 21-37.
- [2] R. Arai, E. Nakai and Y. Sawano, Generalized fractional integral operators on Orlicz–Hardy spaces, Math. Nachr. 294 (2021), 224-235.
- [3] K. F. Andersen and R. T. John, Weighted inequalities for vector-valued maximal functions and singular integrals, Studia Math. 69 (1980/81), 19-31.
- [4] A. Benedek and R. Panzone, The space L^p , with mixed norm, Duke Math. J. 28 (1961), 301-324.
- [5] S. Campanato, Proprietà di una famiglia di spazi funzionali, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 18 (1964), 137-160.
- [6] D.-C. Chang, S. Wang, D. Yang and Y. Zhang, Littlewood–Paley characterizations of Hardy-type spaces associated with ball quasi-Banach function spaces, Complex Anal. Oper. Theory 14 (2020), Paper No. 40, 33 pp.
- [7] Y. Chen, H. Jia and D. Yang, Boundedness of fractional integrals on Hardy spaces associated with ball quasi-Banach function spaces, Submitted or arXiv: 2206.06080.
- [8] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Anisotropic mixed-norm Hardy spaces, J. Geom. Anal. 27 (2017), 2758-2787.
- [9] R. R. Coifman and R. Rochberg, Another characterization of BMO, Proc. Amer. Math. Soc. 79 (1980), 249-254.
- [10] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, Bull. Amer. Math. Soc. 83 (1977) 569-645.
- [11] D. V. Cruz-Uribe and A. Fiorenza, Variable Lebesgue Space. Foundations and Harmonic Analysis, Appl. Number. Harmon Aanl., Birkhäuser/Springer, Heidelberg, 2013.
- [12] D. V. Cruz-Uribe, A. Fiorenza, J. M. Martell and C. Pérez, The boundedness of classical operators on variable L^p spaces, Ann. Acad. Sci. Fenn. Math. 31 (2006), 239-264.

- [13] G. Dafni, T. Hytönen, R. Korte and H. Yue, The space JN_p : nontriviality and duality, J. Funct. Anal. 275 (2018), 577-603.
- [14] R. Del Campo, A. Fernández, F. Mayoral and F. Naranjo, Orlicz spaces associated to a quasi-Banach function space: applications to vector measures and interpolation, Collect. Math. 72 (2021), 481-499.
- [15] L. Diening, P. Hästö and S. Roudenko, Function spaces of variable smoothness and integrability, J. Funct. Anal. 256 (2009), 1731-1768.
- [16] C. Fefferman and E. M. Stein, H^p spaces of several variables, Acta Math. 129 (1972), 137-193.
- [17] B. Franchi, C. Pérez and R. L. Wheeden, Self-improving properties of John–Nirenberg and Poincaré inequalities on spaces of homogeneous type, J. Funct. Anal. 153 (1998), 108-146.
- [18] J. García-Cuerva, Weighted H^p spaces, Dissertationes Math. (Rozprawy Mat.) 162 (1979), 63 pp.
- [19] L. Grafakos, Classical Fourier Analysis, Third edition, Graduate Texts in Mathematics 249, Springer, New York, 2014.
- [20] D. I. Hakim, E. Nakai and Y. Sawano, Generalized fractional maximal operators and vectorvalued inequalities on generalized Orlicz–Morrey spaces, Rev. Mat. Complut. 29 (2016), 59-90.
- [21] S. He, Q. Xue, T. Mei and K. Yabuta, Existence and boundedness of multilinear Littlewood–Paley operators on Campanato spaces, J. Math. Anal. Appl. 432 (2015), 86-102.
- [22] K.-P. Ho, Vector-valued maximal inequalities on weighted Orlicz–Morrey spaces, Tokyo J. Math. 36 (2013), 499-512.
- [23] K.-P. Ho, Strong maximal operator on mixed-norm spaces, Ann. Univ. Ferrara Sez. VII Sci. Mat. 62 (2016), 275-291.
- [24] K.-P. Ho, Mixed norm Lebesgue spaces with variable exponents and applications, Riv. Math. Univ. Parma (N.S.) 9 (2018), 21-44.
- [25] K.-P. Ho, Sublinear operators on mixed-norm Hardy spaces with variable exponents, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 31 (2020), 481-502.
- [26] K.-P. Ho, Operators on Orlicz-slice spaces and Orlicz-slice Hardy spaces, J. Math. Anal. Appl. 503 (2021), Paper No. 125279, 18 pp.
- [27] K.-P. Ho, Fractional integral operators on Orlicz slice Hardy spaces, Fract. Calc. Appl. Anal. (2022), https://doi.org/10.1007/s13540-022-00043-1.
- [28] G. Hu, Y. Meng and D. Yang, Estimates for Marcinkiewicz integrals in BMO and Campanato spaces, Glasg. Math. J. 49 (2007), 167-187.
- [29] L. Huang, J. Liu, D. Yang and W. Yuan, Atomic and Littlewood–Paley characterizations of anisotropic mixed-norm Hardy spaces and their applications, J. Geom. Anal. 29 (2019), 1991-2067.
- [30] L. Huang, J. Liu, D. Yang and W. Yuan, Dual spaces of anisotropic mixed-norm Hardy spaces, Proc. Amer. Math. Soc. 147 (2019), 1201-1215.
- [31] L. Huang, J. Liu, D. Yang and W. Yuan, Identification of anisotropic mixed-norm Hardy spaces and certain homogeneous Triebel–Lizorkin spaces, J. Approx. Theory 258 (2020), 105459, 27 pp.
- [32] L. Huang and D. Yang, On function spaces with mixed norms—a survey, J. Math. Study 54 (2021), 262-336.

- [33] M. Izuki and Y. Sawano, Variable Lebesgue norm estimates for BMO functions, Czechoslovak Math. J. 62(137) (2012), 717-727.
- [34] M. Izuki, Y. Sawano and Y. Tsutsui, Variable Lebesgue norm estimates for BMO functions. II, Anal. Math. 40 (2014), 215-230.
- [35] M. Izuki, E. Nakai, Eiichi and Y. Sawano, Wavelet characterization and modular inequalities for weighted Lebesgue spaces with variable exponent, Ann. Acad. Sci. Fenn. Math. 40 (2015), 551-571.
- [36] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Special John–Nirenberg–Campanato spaces via congruent cubes, Sci. China Math. 65 (2022), 359-420.
- [37] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Boundedness of Calderón–Zygmund operators on special John–Nirenberg–Campanato and Hardy-type spaces via congruent cubes, Anal. Math. Phys. 12 (2022), Paper No. 15, 56 pp.
- [38] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Boundedness of fractional integrals on special John–Nirenberg–Campanato and Hardy-type spaces via congruent cubes, Front. Math. China (to appear) or arXiv: 2108.01891.
- [39] F. John and L. Nirenberg, On functions of bounded mean oscillation, Comm. Pure Appl. Math. 14 (1961), 415-426.
- [40] X. Li, Q. He and D. Yan, Boundedness of multilinear Littlewood–Paley operators on amalgam-Campanato spaces, Acta Math. Sci. Ser. B (Engl. Ed.) 40 (2020), 272-292.
- [41] Y. Liang, J. Huang and D. Yang, New real-variable characterizations of Musielak–Orlicz Hardy spaces, J. Math. Anal. Appl. 395 (2012), 413-428.
- [42] Y. Liang, E. Nakai, D. Yang and J. Zhang, Boundedness of intrinsic Littlewood–Paley functions on Musielak–Orlicz Morrey and Campanato spaces, Banach J. Math. Anal. 8 (2014), 221-268.
- [43] Y. Liang and D. Yang, Musielak–Orlicz Campanato spaces and applications, J. Math. Anal. Appl. 406 (2013), 307-322.
- [44] S. Lu, Four Lectures on Real *H*^p Spaces, World Scientific Publishing Co., Inc., River Edge, NJ, 1995.
- [45] S. Lu, C. Tan and K. Yabuta, Littlewood–Paley operators on the generalized Lipschitz spaces, Georgian Math. J. 3 (1996), 69-80.
- [46] Y. Meng, E. Nakai and D. Yang, Estimates for Lusin–area and Littlewood–Paley g_{λ}^* functions over spaces of homogeneous type, Nonlinear Anal. 72 (2010), 2721-2736.
- [47] Y. Meng and D. Yang, Estimates for Littlewood–Paley operators in BMO(\mathbb{R}^n), J. Math. Anal. Appl. 346 (2008), 30-38.
- [48] A. Meskhi and Y. Sawano, Density, duality and preduality in grand variable exponent Lebesgue and Morrey spaces, Mediterr. J. Math. 15 (2018), Paper No. 100, 15 pp.
- [49] M. Milman, Marcinkiewicz spaces, Garsia–Rodemich spaces and the scale of John–Nirenberg self improving inequalities, Ann. Acad. Sci. Fenn. Math. 41 (2016), 491-501.
- [50] M. Milman, Garsia–Rodemich spaces: Bourgain–Brezis–Mironescu space, embeddings and rearrangement invariant spaces, J. Anal. Math. 139 (2019), 121-141.
- [51] E. Nakai, On generalized fractional integrals in the Orlicz spaces on spaces of homogeneous type, Sci. Math. Jpn. 54 (2001), 473-487.
- [52] E. Nakai and Y. Sawano, Hardy spaces with variable exponents and generalized Campanato spaces, J. Funct. Anal. 262 (2012), 3665-3748.

- [53] E. Nakai and Y. Sawano, Orlicz–Hardy spaces and their duals, Sci. China Math. 57 (2014), 903-962.
- [54] S. Qiu, Boundedness of Littlewood–Paley operators and Marcinkiewicz integral on $\varepsilon^{\alpha,p}$, J. Math. Res. Exposition 12 (1992), 41-50.
- [55] M. M. Rao and Z. D. Ren, Theory of Orlicz Spaces, Monographs and Textbooks in Pure and Applied Mathematics 146, Marcel Dekker, Inc., New York, 1991.
- [56] Y. Sawano, K.-P. Ho, D. Yang and S. Yang, Hardy spaces for ball quasi-Banach function spaces, Dissertationes Math. 525 (2017), 1-102.
- [57] Y. Sawano and T. Shimomura, Generalized fractional integral operators on generalized Orlicz–Morrey spaces of the second kind over non-doubling metric measure spaces, Georgian Math. J. 25 (2018), 303-311.
- [58] Y. Sawano and T. Shimomura, Maximal operator on Orlicz spaces of two variable exponents over unbounded quasi-metric measure spaces, Proc. Amer. Math. Soc. 147 (2019), 2877-2885.
- [59] Y. Sawano, S. Sugano and H. Tanaka, Orlicz–Morrey spaces and fractional operators, Potential Anal. 36 (2012), 517-556.
- [60] E. M. Stein, On the functions of Littlewood–Paley, Lusin, and Marcinkiewicz, Trans. Amer. Math. Soc. 88 (1958), 430-466.
- [61] E. M. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton Mathematical Series 30, Princeton University Press, Princeton, NJ, 1970.
- [62] E. M. Stein, Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals, Princeton Mathematical Series 43, Monographs in Harmonic Analysis, III, Princeton University Press, Princeton, NJ, 1993.
- [63] J. Sun, G. Xie and D. Yang, Localized John–Nirenberg–Campanato spaces, Anal. Math. Phys. 11 (2021), Paper No. 29, 47 pp.
- [64] J. Sun, D. Yang and W. Yuan, Weak Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: decompositions, real interpolation, and Calderón–Zygmund operators, J. Geom. Anal. 32 (2022), Paper No. 191, 85 pp.
- [65] J. Sun, D. Yang and W. Yuan, Molecular characterization of weak Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type with its application to Littelwood–Paley function characterization, Submitted.
- [66] Y. Sun, On the existence and boundedness of square function operators on Campanato spaces, Nagoya Math. J. 173 (2004), 139-151.
- [67] J. Tao, D. Yang and W. Yuan, John–Nirenberg–Campanato spaces, Nonlinear Anal. 189 (2019), 111584, 36 pp.
- [68] J. Tao, D. Yang, W. Yuan and Y. Zhang, Compactness characterizations of commutators on ball Banach function spaces, Potential Analysis (2021), https://doi.org/10.1007/s11118-021-09953-w.
- [69] J. Tao, D. Yang and W. Yuan, Vanishing John–Nirenberg spaces, Adv. Calc. Var. (2021), https://doi.org/10.1515/acv-2020-0061.
- [70] J. Tao, D. Yang and W. Yuan, A survey on several spaces of John–Nirenberg-type, Mathematics 9 (2021), Art. No. 2264, https://doi.org/10.3390/math9182264.
- [71] A. Torchinsky, Real-Variable Methods in Harmonic Analysis, Pure and Applied Mathematics 123, Academic Press, Inc., Orlando, FL, 1986.

- [72] F. Wang, D. Yang and S. Yang, Applications of Hardy spaces associated with ball quasi-Banach function spaces, Results Math. 75 (2020), Paper No. 26, 58 pp.
- [73] S. Wang, D. Yang, W. Yuan and Y. Zhang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces II: Littlewood–Paley characterizations and real interpolation, J. Geom. Anal. 31 (2021), 631-696.
- [74] S. Wang, Some properties of Littlewood–Paley *g*-function, Sci. Sinica Ser. A 28 (1985), 252-262.
- [75] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Characterizations of maximal functions, decompositions, and dual spaces, Math. Nachr. (2022), http://doi.org/10.1002/mana.202100432.
- [76] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Littlewood–Paley characterizations with applications to boundedness of Calderón–Zygmund operators, Acta Math. Sin. (Engl. Ser.) (2022), https://doi.org/10.1007/s10114-022-1573-9.
- [77] X. Yan, D. Yang and W. Yuan, Intrinsic square function characterizations of Hardy spaces associated with ball quasi-Banach function spaces, Front. Math. China 15 (2020), 769-806.
- [78] Y. Zhang, L. Huang, D. Yang and W. Yuan, New ball Campanato-type function spaces and their applications, J. Geom. Anal. 32 (2022), Paper No. 99, 42 pp.
- [79] Y. Zhang, D. Yang and W. Yuan, Real-variable characterizations of local Orlicz-slice Hardy spaces with application to bilinear decompositions, Commun. Contemp. Math. (2021), https://doi.org/10.1142/S0219199721500048.
- [80] Y. Zhang, D. Yang, W. Yuan and S. Wang, Real-variable characterizations of Orlicz-slice Hardy spaces, Anal. Appl. (Singap.) 17 (2019), 597-664.
- [81] Y. Zhang, D. Yang, W. Yuan and S. Wang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces I: Decompositions with applications to boundedness of Calderón–Zygmund operators, Sci. China Math. 64 (2021), 2007-2064.

Hongchao Jia, Dachun Yang (Corresponding author), Wen Yuan and Yangyang Zhang

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, The People Republic of China

```
E-mails: hcjia@mail.bnu.edu.cn(H. Jia)
dcyang@bnu.edu.cn(D. Yang)
wenyuan@bnu.edu.cn(W. Yuan)
yangyzhang@mail.bnu.edu.cn(Y. Zhang)
```