Sentential logics based on k-cyclic modal pseudocomplemented De Morgan algebras

Aldo Figallo-Orellano, Miguel Peréz-Gaspar and Juan Manuel Ramírez-Contreras

Abstract

The study of the theory of operators over modal pseudocomplemented De Morgan algebras was begun in the papers [15] and [16]. In this paper, we introduce and study the class of modal pseudocomplemented De Morgan algebras enriched by an automorphism k-periodic (or C_k -algebras) where k is a positive integer; for k = 2 the class coincides with the one studied in [15] where the automorphism works as a new unary operator. In the first place, we prove the class C_k -algebras is a semisimple variety and we determine the generating algebras. Afterwards, we calculate the cardinality of the free C_k -algebra with n generator. After the algebraic study, we built two sentential logics that have as algebraic counterpart the class of C_k -algebras that we denote \mathbb{L}_k^\leq and \mathbb{L}_k for every k. \mathbb{L}_k is a 1-assertional logic and \mathbb{L}_k^\leq is the degree-preserving logic both associated with the class of C_k -algebras. Working over these logics, we prove that \mathbb{L}_k^\leq is paraconsistent, which is protoalgebraic and finitely equivalential but not algebraizable. In contrast, we prove that \mathbb{L}_k is algebraizable, sharing the same theorem with \mathbb{L}_k^{\leq} , but not paraconsistent.

1 Introduction and Preliminaries

In 1978, A. Monteiro introduced tetravalent modal algebras (or TMA-algebras for short) as algebras $\langle L, \wedge, \vee, \sim, \nabla, 0, 1 \rangle$ of type (2, 2, 1, 1, 0, 0) such that $\langle L, \wedge, \vee, \sim, 0, 1 \rangle$ are De Morgan algebras which satisfy the following conditions:

$$\nabla x \lor \sim x = 1,$$

$$\nabla x \land \sim x = \sim x \land x.$$

These algebras arise as a generalization of three-valued Lukasiewicz algebras by omitting the identity $\nabla(x \wedge y) = \nabla x \wedge \nabla y$. The variety of TMAs is generated by the well-known four-element De Morgan algebra expanded with a simple modal operator ∇ (i.e., $\nabla 1 = 1$ and $\nabla x = 0$ for $x \neq 1$. Besides, $\sim 0 = 1$ and $\sim x = x$ for $x \neq 0, 1$). These algebras were studied by I. Loureiro, A. V. Figallo and A. Ziliani, see [21, 20] for instance.

Later, J. Font and M. Rius introduced some logics that have as algebraic counterpart to the class of TMA-algebras. In particular, they defined the logic TML which can be seen as the degree-preserving logic associated with the class of TMA-algebras, see [26, Proposition 3.7] and [23]. These authors also proved that the logic \mathcal{TML} is not algebraizable with Blok-Pigozzi's method. More recently, M. Figallo proved that \mathcal{TML} is in fact a paraconsistent logic and a *Logic of Formal Inconsistency* in [22]. Recall that the Logics of Formal Inconsistency (LFIs) were introduced by W. Carnielli and J. Marcos (see [8]), and the Logics of Formal Undeterminedness (LFUs) was introduced in [31]. Other paraconsistent degree-preserving logics were studied by Ertola *et al.* in [7] and [13]; besides, these kinds of logics were built over *Distributive Involutive Residuated Lattices* in [14], where the authors presented a large family of LFIs and LFUs.

Another class of algebras related with TMA-algebra are Stone Involutive Algebras (SIalgebras), these structures are De Morgan algebras enriched with a modal operator similar to ∇ considered above; they were defined by Cingoli and Gallego in [10]. Later, L. Cantú and M. Figallo studied the degree-preserving logic associated with the class of SI-algebras $(S\mathcal{I}^{\leq})$ providing a sound and complete Gentzen calculus. Besides, they proved that $S\mathcal{I}^{\leq}$ is a *strong* LFI. Later on, S. Marcelino and U. Rivieccio proved that $S\mathcal{I}^{\leq}$ is selfextensional and non-protoalgebraic logic; hence, non-algebraizable, [35].

On the other hand, A. V. Figallo considered the subvariety of pseudocomplemented De Morgan algebras which verifies: (tm) $x \vee \sim x \leq x \vee x^*$ in [17]. This author called them modal pseudocomplemented De Morgan algebras (or mpM-algebras). Recall that a pseudocomplemented De Morgan algebra A is a De Morgan algebra with a unary operator * such that every $a \in A$, the element a^* is the pseudocomplement of a; i.e. $a \wedge x = 0$ if and only if $x \leq a^*$. Later, he showed that every mpM-algebra is a TMA by defining $\nabla x = \sim (\sim x \wedge x^*)$ and $\Delta x = \sim \nabla \sim x$. In [18] (see also [19]), the authors have proven that the subdirectly irreducible mpM-algebras are three as TMAs, in fact, Hasse diagrams are the same in each case, but 3-chain-mpM-algebra is not a subalgebra of four-elements. The mentioned algebras are the following: $T_2 = \{0, 1\}$ with $0 < 1, \sim 0 = 0^* = 1, \sim 1 = 1^* = 0;$ $T_3 = \{0, a, 1\}$, with 0 < a < 1, $\sim a = a$, $a^* = 0$, $\sim 0 = 0^* = 1$, $\sim 1 = 1^* = 0$; $T_4 = \{0, a, b, 1\}$ with $a \not\leq b, b \not\leq a$ and $0 < a, b < 1, \sim b = a^* = b, \sim a = b^* = a,$ $\sim 0 = 0^* = 1, \sim 1 = 1^* = 0$. It is worth mentioning that mpM-algebras constitute a proper subvariety of the variety \mathcal{V}_0 studied by H. Sankappanavar in [38]. More recently, the theory of operators over mpM-algebras was considered in [15, 16]. In particular, the authors studied the class of mpM-algebras enriched with an automorphism of period 2, this automorphism works as a new unary operator, [16].

One of the principal objects of the present paper is to generalize the work of [16] for an automorphism of period k, where k is a positive integer. This kind of operators was studied over other algebraic structures, for instance, over Post algebras of order p by López-Martinolich in [32] (see also [33, 12]), where the variety of k-cyclic Post algebras of order p was studied. Furthermore, it was proved that there exists an equivalence between the variety generated by the simple k-cyclic Post algebra of order p and the variety generated by the finite field with p^k elements, see [33].

The paper is organized as follows: In Section 2, we introduce the class of C_k -algebras which is in fact a variety; and, we display some properties of *prime spectrum* for a given algebra. Section 3, we present a new implication in order to use A. Monteiro's techniques

to prove the variety semisimplicity. Section 4, maximal congruences and simple algebras of the class are characterized; and, as consequence, the simple C_k -algebras are determined. In Section 5, we describe the free algebra with n generators calculating their cardinality using the algebraic properties displayed before; as a by-product, we check that for k = 2and k = 1 the cardinality coincides with others obtained in the literature. Finally, in Section 6 and 7, we build a family of sentential logics that have as algebraic counterpart the class of C_k -algebras denoted \mathbb{L}_k^{\leq} and \mathbb{L}_k for every k. The logic \mathbb{L}_k is 1-assertional and \mathbb{L}_k^{\leq} is the degree-preserving logic, both associated with the class of C_k -algebras. Firstly, we prove that \mathbb{L}_k^{\leq} is an LFI (hence paraconsistent) and an LFU which is protoalgebraic and finitely equivalential but not algebraizable. Finally, we prove that \mathbb{L}_k is algebraizable which is an LFU and shares the same theorem with \mathbb{L}_k^{\leq} but it is not paraconsistent.

2 The class of k-cyclic mpM-algebras: C_k -algebras

In this section, we will introduce a new class of algebras that we call C_k -algebras which is in fact an equational one. Later, we will display algebraic properties with the purpose to determine the generating algebras of the variety. First, let us consider the following definition:

Definition 2.1 A k-cyclic mpM-algebra (for short C_k -algebra) is a pair (A, t) where A is an mpM-algebra and the function $t : A \to A$ is an automorphism such that $t^k(x) = x$ where k is an integer $k \ge 0$. Besides, we write $t^0(x) = x$ and $t^n(x) = (t^{n-1} \circ t)(x)$ if $n \ge 1$.

Sometimes we write "tx" instead of "t(x)". It is worth mentioning that the class C_k -algebra is a variety; and, as examples, we have the classes 1-cyclic mpM-algebra and 2-cyclic mpM-algebra were studied in [18] and [15], respectively. Besides, if an mpM-algebra is a 3-valued Łukasiewicz algebra then the k-cyclic version is a special class studied in [32, 33, 12]. Now, we present a Lemma that will be used in the rest of the paper.

Lemma 2.2 ([15]) In every C_k -algebra, the following conditions hold:

(T1) $\triangle 0 = 0, \ \triangle 1 = 1,$ (T2) $\triangle x < x, x < \nabla x$, (T3) if $x \leq y$ then $\triangle x < \triangle y$, (T4) $\triangle x$ is a Boolean element. (T6) $(\sim \bigtriangleup x)^* = \bigtriangleup (\sim x)^*$. (T5) $(\sim \bigtriangleup x)^* = \bigtriangleup x$, (T7) $\triangle \triangle x = \triangle x, \ \triangle \nabla x = \nabla x, \ \nabla \triangle x = \triangle x,$ (T8) $(\bigtriangleup x)^* = \sim \bigtriangleup x$. (T9) $\triangle \sim \triangle x = \sim \triangle x$, (T10) $\triangle(x \land y) = \triangle x \land \triangle y$, (T10)(bis) $\nabla(x \lor y) = \nabla x \lor \nabla y$, (T11) $x \in \triangle A$ iff $x = \triangle x$ iff $x = \nabla x$. (T13) $\sim x \land \bigtriangleup x = 0 \text{ (or } x \lor \nabla \sim x = 1),$ (T12) $\triangle A$ is an S-subalgebra of A, (T15) $\triangle(\triangle x \lor y) = \triangle x \lor \triangle y$, (T14) $x \lor \sim \bigtriangleup x = 1$, (T16) $\triangle (\sim \triangle x \lor y) = \sim \triangle x \lor \triangle y,$ (T17) $\nabla(\bigtriangleup x \land y) = \bigtriangleup x \land \nabla y$, (T18) $\triangle(\nabla x \lor y) = \nabla x \lor \triangle y$,

The relation between the congruences and certain filters is given by the following Remark.

Remark 2.3 Taking into account the studied of congruences of [15, Theorem 2.6], we can affirm that the c-filters (i.e. $t(\Delta F) = F$ where F is a filter) characterize the congruences of given C_k -algebra, moreover:

- (i) The relation $Con(A) = \{R(F) : F \in C(A)\}$ where $R(F) = \{(x, y) \in A \times A : there is f \in F \text{ such that } x \land f = y \land f\}$ is a congruence, we denote by C(A) the set of all *c*-filters of A.
- (ii) The posets Con(A) and C(A) are isomorphic where Con(A) is the poset of all congruences of A.

Now, we give a result to be used to characterize the prime spectrum for a given C_k algebra. First, recall that the notion of prime filter, ultrafilter, maximal and minimal are in the usual ones, see [1]. Also, we are going to use the well-known Birula–Rasiowa transformation φ , see, for instance, [26]. Recall that, if A is a De Morgan algebra, the map φ is defined as follows: For every prime filter P of A

$$\varphi(P) = A \setminus \sim P = A \setminus \{\sim x : x \in P\}$$

This map has the following properties:

- $\varphi(P)$ is a prime filter of A,
- $\varphi(\varphi(P)) = P$,
- if Q is a prime filter of A such that $P \subseteq Q$ then $\varphi(Q) \subseteq \varphi(P)$.

Then, we have the following Lemma:

Lemma 2.4 Let (A, t) be a C_k -algebra, $P \subseteq A$ a prime filter, and φ is the Birula–Rasiowa transformation on A. Then, the following properties hold:

- (a) $t^i(P)$ is a prime filter for $1 \le i \le k$;
- (b) P is minimal (maximal) iff $t^i(P)$ is minimal (maximal) for $1 \le i \le k$;
- (c) U is an ultrafilter iff $t^i(U)$ is an ultrafilter for $1 \le i \le k$;
- (d) $\varphi(t^i(P)) = t^i \varphi(P), \ 1 \le i \le k;$
- (e) if F is a c-filter of A and $F \subseteq P$, then $F \subseteq \varphi(P) \cap t^i(P)$;

(f) if $P \subseteq Q$ and Q is a prime filter of A, then $\varphi(P) = Q$ of P = Q.

Proof. It follows from the very definitions.

In what follows, we will study the prime spectrum of a given C_k -algebra A with an algebraic technique. First, let us observe the study could be done using topological representation through the techniques given in [16, 18].

Proposition 2.5 Let (A, t) be a \mathcal{C}_k -algebra and $U \subseteq A$ an ultrafilter. Then, $N = \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U)$ is a c-filter of A.

Proof. It is clear that N is a filter of A. Then, we only have to prove that N is closed by \triangle and t. Indeed, let $x \in N$ then by (T14) we have that $\sim x \lor \triangle x \in N$. Since $x \in \varphi(t^i U)$ for some i such that $0, \leq i \leq k-1$, we infer that $\sim x \notin t^i U$ and now from Lemma 2.4 (a) we have that $\triangle x \in t^i U$. Now, if $x \in t^i(U)$, then $\sim x \notin \varphi(t^i U)$ and so $\triangle x \in \varphi(t^i U)$. From the last assertions, we deduce that $\triangle x \in N$. On the other hand, let us suppose $x \in N$, if $x \in \bigcap_{i=0}^{k-1} t^i(U)$, then $t(x) \in \bigcap_{i=0}^{k-1} t^i(U)$. Besides, if we have that $x \in \bigcap_{i=0}^{k-1} \varphi(t^i U)$, then $t(x) \in \bigcap_{i=0}^{k-1} t^i(U)$ and from Lemma 2.4 (d) we conclude that $t(x) \in \bigcap_{i=0}^{k-1} \varphi(t^i U)$. Therefore, $t(x) \in N$ as desired.

In the following, we will show that every maximal c-filter can be expressed in terms of certain ultrafilters extending the result given [15, Theorem 5.10].

Theorem 2.6 Let (A, t) be a C_k -algebra and $N \subseteq A$. Then, N is a maximal c-filter of if only if there exists an ultrafilter U of A such that

$$N = \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U).$$

Besides, this representation of N is unique.

Proof. Let N be a maximal c-filter of A, since N is proper there is an ultrafilter U of A such that $N \subseteq U$. Then, from Lemma 2.4 we infer that $N \subseteq \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U)$ and so $N = \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U).$

Conversely, let us take U an ultrafilter of A such that $N = \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U)$. From Proposition 2.5, we have that N is a c-filter of A and since N is a proper filter, then there

is a maximal c-filter W such that $N \subseteq W$. From the necessary condition, we know that there exists an ultrafilter Y of A such that $W = \bigcap_{i=0}^{k-1} t^i Y \cap \varphi(t^i Y)$. Now, since Y is a prime filter, we infer that $t^i(U) \subseteq Y$ or $\varphi(t^i U) \subseteq Y$. Besides, from Lemma 2.4, we know that $t^i(U)$ is maximal, and therefore $t^i(U) = Y$. On the other hand, if $\varphi(t^i U) \subseteq Y$, from Lemma 2.4, we have that $t^i U = \varphi(\varphi(t^i U)) = Y$ of $\varphi(t^i U) = Y$. Therefore, N = W as desired.

Definition 2.7 Let (A, t) be a C_k -algebra, we say that a maximal c-filter N is d-periodic if d the smallest positive integer such that $t^d N = N$. In this case, we also say N is a d-filter of A.

The last Definition plays an important role in the following Lemma.

Lemma 2.8 Let (A, t) be a C_k -algebra. If N is a d-filter of A, then the following conditions hold:

- (i) d/k where / is the divide relation on integers,
- (ii) the unique ultrafilters of A such that they contain N are the form $\{t^i P, \varphi(t^i P)\}_{0 \le i \le d-1}$.

Proof.

(i): According to Definition 2.7, it is clear that $d \leq k$, then we have that k = dq + r with r < d. Thus, $t^k N = t^{dq+r} N = t^r (t^{dq} N) = t^r N$ and since (A, t) is a \mathcal{C}_k -algebra, we infer that $t^r N = N$ as desired.

(ii): It is direct consequence from Definition 2.7 and Theorem 2.6.

3 The semsimplicity of the class of C_k -algebras

In the following, we will characterize the congruences by means of certain deductive systems, for a given C_k -algebra in order to prove that the variety of C_k -algebra is semisimple. Now, let (A, t) be a C_k -algebra, then we define a new implication \neg on A as follows:

$$a \rightarrow b = \bigvee_{i=1}^{k} \nabla(\sim t^{i}a) \lor b.$$

Thus, we have the following Lemma that will be central in the rest of the section.

Lemma 3.1 In every C_k -algebra the following conditions hold:

- (i) $a \rightarrow a = 1$,
- (ii) $a \rightarrow \triangle a = 1$,
- (iii) $a \rightarrow t^s(a) = 1$, for every $s \in \mathbb{N}$,
- (iv) $a \rightarrow (a \wedge b) = a \rightarrow b$,
- (v) a < b implies $c \rightarrow a < c \rightarrow b$,

- (vi) $((a \rightarrow b) \rightarrow a) \rightarrow a) = 1$.
- (vii) $a \rightarrow (b \rightarrow a) = 1$,
- (viii) $1 \rightarrow a = 1$ implies a = 1,
- (ix) $(a \rightarrow (b \rightarrow c)) = ((a \rightarrow b) \rightarrow (a \rightarrow c))$ c)),

(x)
$$a \to 1 = 1$$
.

Proof. We only prove (iii), (iv) y (ix):

(iii): If $s \leq k$, then $1 = \nabla \sim t^s a \lor t^s(a) \leq \bigvee_{i=1}^k \nabla \sim t^i a \lor t^s(a) = a \rightarrow t^s(a)$. On the other hand, if $k \leq s$, then there are q, r positive integers such that $s = k \cdot q + r$ and r < k. Thus, $t^{s}(a) = t^{r}(a)$ and by analogous reasoning given before we have proved the property. (vi): Let us prove first: (*) $\bigvee_{n=1}^{k-1} t^n (\bigwedge_{i=1}^k t^i x) = \bigvee_{n=1}^{k-1} \bigwedge_{i=1}^k t^{n+i} x = \bigwedge_{j=1}^k t^j x$. Indeed, it is enough to show $\{t^{n_0+i}x\}_{1\leq i\leq k} = \{t^jx\}_{1\leq j\leq k}$ for every positive integer n_0 such that $1\leq n_0\leq k$.

Let $A = \{t^j x\}_{1 \le j \le k}$ and $B = \{t^{n_0+i}x\}_{1 \le i \le k}$. If j is such that $1 \le j \le k$, then we have three possible cases: (a) $j > n_0$ or (b) $j = n_0$ or (c) $j < n_0$.

We will see there is an positive integer "i" such that $t^{j}x = t^{n_{0}+i}x$ for every case. Indeed, if (a) occurs, we have that $0 < j - n_0 < k$, then taking $i = j - n_0$ the property is proved. If (b) holds, taking i = k the property is easily verified. Finally, if (c) holds, then we have that $0 < n_0 - j < k$ and so $0 < k - (n_0 - j) < k$. Hence, if we take $i = k - (n_0 - j)$, then we have that $t^j x = t^{n_0+i} x$. From the last assertions, we conclude that $A \subseteq B$.

Conversely, since $0 \le i \le k$ we have that $n_0 \le i + n_0 \le k + n_0$. Then, we have the following two cases. If (a) occurs we have $n_0 \leq i + n_0 \leq k$, then $j = i + n_0$ as desired. If (b) occurs we have $k < i + n_0 \le k + n_0$, then $i + n_0 = q \cdot k + r$ with $0 \le r < k - 1$. Thus, $t^{i+n_0}x = t^{q\cdot k+r}x = t^r x$ and if we take j = r, we have the other inclusion verified. Therefore, A = B as desired.

Now, let us show that
$$((a \to b) \to a) = a$$
. First, let us observe that $(**) a \to b = \bigvee_{j=1}^{k} \nabla \sim t^{j} a \lor b = \nabla \sim (\bigwedge_{j=1}^{k} t^{j} a) \lor b = \nabla \sim z \lor b$, where $z = \bigwedge_{j=1}^{k} t^{j} a$. Thus,
 $(a \to b) \to a = (\nabla \sim z \lor b) \to a) = \bigvee_{n=1}^{k} \nabla \sim t^{n} (\nabla \sim z \lor b) \lor a = \nabla \sim (\nabla \sim z \lor b) \lor \bigvee_{n=1}^{k-1} \nabla \sim t^{n} (\nabla \sim z \lor b) \lor a = \nabla (\sim \nabla \sim z \land \sim b) \lor \bigvee_{n=1}^{k-1} t^{n} \nabla (\sim \nabla \sim z \land \sim b) \lor a = \nabla ((\Delta z \land \sim a)) \lor a = (T17) ((\Delta z \land \nabla \sim b) \lor \bigvee_{n=1}^{k-1} t^{n} ((\Delta z \land \nabla \sim b)) \lor a) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor \bigvee_{n=1}^{k-1} t^{n} ((\Delta z \land \nabla \sim b)) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor \bigvee_{n=1}^{k-1} t^{n} ((\Delta z \land \nabla \sim b)) \lor a) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigwedge_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \sim b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor_{j=1}^{k} t^{j} a \land \nabla \to b) \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor a = (\bigvee_{j=1}^{k} t^{j} a \land \nabla \to b) \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor_{j=1}^{k} t^{j} a \lor b \lor b \lor b \lor_{j=1}^{k}$

$$\begin{split} b) \vee \bigvee_{n=1}^{k-1} t^n (\bigwedge_{j=1}^k t^j \triangle a \land \nabla \sim b) \lor a &= (\triangle a \land \bigwedge_{j=1}^{k-1} t^j \triangle a \land \nabla \sim b) \lor \bigvee_{n=1}^{k-1} t^n (\bigwedge_{j=1}^k t^j \triangle a \land \nabla \sim b) \lor a &= ((\triangle a \land \bigwedge_{j=1}^{k-1} t^j (\bigwedge_{j=1}^k t^j \triangle a \land \nabla \sim b) \lor a) \lor \bigvee_{n=1}^{k-1} t^n (\bigwedge_{j=1}^k t^j \triangle a \land \nabla \sim b) &= (T2) \ a \lor \bigvee_{n=1}^{k-1} t^n (\bigwedge_{j=1}^k t^j \triangle a \land \nabla \sim b) \\ b) &= a \lor (\bigvee_{n=1}^{k-1} t^n (\bigwedge_{j=1}^k t^j \triangle a) \land t^n \nabla \sim b). \\ \text{From the latter, (**) and (T2), we can infer that} \\ (a \to b) \to a &= a \lor (\bigwedge_{i=1}^k t^i \triangle a \land \bigvee_{n=1}^{k-1} t^n \nabla \sim b) &= a \lor ((\triangle a \land \bigwedge_{i=1}^{k-1} t^i \triangle a \land \bigvee_{n=1}^{k-1} t^n \nabla \sim b) &= a. \\ \text{Form the last assertion and (i), we have proved the property.} \\ \text{Form the last assertion and (i), we have proved the property.} \\ (ix): First, let us observe that $(a \to b) \to (a \to c) = \bigvee_{i=1}^k \nabla \sim t^i (\bigwedge_{j=1}^k \Delta t^j a \land b) \lor (\bigvee_{j=1}^k \nabla \sim t^j a \land b) \lor (\bigvee_{j=1}^k \nabla \sim t^j a \lor t^j a \lor t^j a) \lor (\bigvee_{j=1}^k \nabla \sim t^j a \lor t^j a$$$

 $a \rightarrow (b \rightarrow c)$ which completes the proof.

In the following, we will consider the notion of deductive system for the implication \neg as usual: given a \mathcal{C}_k -algebra A and $D \subset A$, we say that D is a cyclic deductive system if $(D_c 1)$ $1 \in D$ and if $(D_c 2)$ $x, x \neg y \in D$, then $y \in D$.

Lemma 3.2 Let (A,t) be a C_k -algebra and $F \subseteq A$, then the following conditions are equivalent:

- (i) F is a cyclic deductive system,
- (ii) F is a c-filter.

Proof.

(i) \Rightarrow (ii): It is clear that $1 \in F$. Now, let us suppose $a, b \in F$, from Lemma 3.1 (vii) and (iv), we have $1 = b \rightarrow (a \rightarrow b) = b \rightarrow (a \rightarrow (a \wedge b)) \in F$. Then, by (D_c2) we infer that $a \wedge b \in F$. Now, let us suppose $a \in F$ and $b \in A$ such that $a \leq b$, then form Lemma 3.1 (v) we have that $1 = a \rightarrow a \leq a \rightarrow b$. From the latter, Lemma 3.1 (viii) and (D_c2), we obtain that $b \in F$. Now, from Lemma 3.1 (ii) and (iii), we have the conditions of *c*-filter are verified.

(ii) \Rightarrow (i): Let $a, a \to b \in F$, then $\triangle t^j a \in F$ for every $1 \le j \le k$. Besides, $\triangle (a \to b) \in F$ and therefore $\bigwedge_{j=1}^k \triangle t^j a \land \triangle (a \to b) \in F$. On the other hand, we have that $\bigwedge_{j=1}^k \triangle t^j a \land \triangle (a \to b) = k$. $b) = \bigwedge_{j=1}^k \triangle t^j a \land \triangle (\bigvee_{i=1}^k \nabla \sim t^i a \lor b) = (T18) \bigwedge_{j=1}^k \triangle t^j a \land (\bigvee_{i=1}^k \nabla \sim t^i a \lor \triangle b) = (\bigwedge_{j=1}^k \triangle t^j a \land \bigvee_{i=1}^k \sim b) = (\bigwedge_{j=1}^k \triangle t^j a \land (\bigvee_{i=1}^k \nabla \sim t^i a \lor \triangle b) = H$. Since (T4) we know that $\triangle a$ is a Boolean element, then

 $(\bigwedge_{j=1}^{k} \triangle t^{j}a \land \bigvee_{i=1}^{k} \sim \triangle t^{i}a) = 0. \text{ Thus, } H = \bigwedge_{j=1}^{k} \triangle t^{j}a \land \triangle b \le \triangle b \le b. \text{ Hence, from the last assertions we conclude that } b \in F \text{ as desired.}$

From Lemma 3.2 we can say the congruences are characterized by the cyclic deductive systems. Now, according to Lemma 3.1 and results given by A. Monteiro ([34]), we have proved the following Lemma.

Lemma 3.3 For every C_k -algebra (A, t), the following conditions hold:

- (i) every cyclic deductive system of A is intersection of maximal cyclic deductive systems of A,
- (ii) the intersection of all maximal cyclic deductive systems of A is equal to $\{1\}$.
- (iii) The variety of C_k -algebras is semisimple.

4 Generating algebras of the variety

In this section, we will determine the simple algebras of the class of C_k -algebras. To this end, we first studied the maximal congruence for a given algebra. The notion of cyclic deductive generated by a set H is defined as usual and we denoted by D(H). If $H = \{a\}$ we just write D(a).

Lemma 4.1 Let (A, t) be a C_k -algebra, $a \in A$ and $H \subseteq A$, then:

(i)
$$D(a) = [\bigwedge_{j=1}^{k} t^{j} \triangle a)$$
, where [X) is the c-filter generated by X,

(ii)
$$D(H,a) = D(H \cup \{a\}) = \{x \in A : a \rightarrow x \in D(H)\},\$$

(iii)
$$D(a) = \{x \in A : a \rightarrow x = 1\}.$$

Proof. (i): Since $\bigwedge_{j=1}^{k} t^{j} \triangle a \leq \Delta a \leq a$, then $a \in [\bigwedge_{j=1}^{k} t^{j} \triangle a)$. Now, let us take $x \in [\bigwedge_{j=1}^{k} t^{j} \triangle a)$, then $\bigwedge_{j=1}^{k} t^{j} \triangle a \leq x$ and $\bigwedge_{j=1}^{k} t^{j} \triangle \Delta a \leq \Delta x$. Thus, $\Delta x \in [\bigwedge_{j=1}^{k} t^{j} \triangle a)$. Similarly, we can prove that $tx \in [\bigwedge_{j=1}^{k} t^{j} \triangle a)$. Then, from Lemma 3.2 we have that $[\bigwedge_{j=1}^{k} t^{j} \triangle a)$ is a cyclic deductive system. On the other hand, if W is a cyclic deductive system and $a \in W$, then it is simple to verify that $[\bigwedge_{j=1}^{k} t^{j} \triangle a] \subseteq W$ which completes the proof.

(ii): From Lemma 3.1 (i), we have that $a \in D(H, a)$. Besides, it is clear that $h \in D(H, a)$ for every $h \in H$. Let us see that D(H, a) is a deductive system. Indeed, from Lemma 3.1 (x) we know that $1 \in D(H, a)$. Besides, let us consider $x, x \to y \in D(H, a)$, then we have that $a \to x, a \to (x \to y) \in D(H)$. Thus, from Lemma 3.1 we infer that $1 = (a \to (x \to y)) \to ((a \to x) \to (a \to y)) \in D(H)$. From the latter and $(D_c 2)$, we have $a \to y \in D(H)$. On the other hand, if B is a cyclic deductive system such that $H \cup \{a\} \subseteq B$, then is not hard to see that $D(H \cup \{a\}) \subseteq B$ as desired.

(iii): It immediately follows from (ii) taking $H = \emptyset$.

Lemma 4.2 Let (A, t) be a C_k -algebra and let D_1 be a cyclic deductive system of A. Then,

$$D(D_1, a) = \{x \in A : \text{ there exists } d \in D_1 \text{ such that } d \land \bigwedge_{j=1}^k t^j \triangle a \le x\}.$$

Proof. Consider $B = \{x \in A : \text{ there is } d \in D_1 \text{ such that } d \land \bigwedge_{j=1}^k t^j \triangle a \leq x\}$. Then, it is clear that $D_1 \subseteq D(D_1, a)$ and that $a \in D(D_1, a)$. On the other hand, if $x, y \in B$ and since D_1 is filter then we conclude $x \land y \in B$. Now let us see that B is a *c*-filter. Indeed, let $z \in B$ then there is $w \in D_1$ such that $w \land \bigwedge_{j=1}^k t^j \triangle a \leq z$. Thus, from (T10) we have that $\triangle(w \land \bigwedge_{j=1}^k t^j \triangle a) = \triangle w \land \bigwedge_{j=1}^k t^j \triangle a \leq \Delta z$. From Lemma 3.2, D_1 is a *c*-filter we have that $\triangle w \in D_1$ and then $\triangle z \in B$. Analogously, it is possible to see that $t(z) \in B$. On the other hand, let us suppose there exists a *c*-filter *Y* such that $D_1 \cup \{a\} \subseteq Y$. If $x \in B$, then there exists $d \in D_1$ such that $d \land \bigwedge_{j=1}^k t^j \triangle a \leq x$. Now, since $\bigwedge_{j=1}^k t^j \triangle a \in Y$ and *Y* us a filter we infer that $x \in Y$, which concludes the proof.

Our next task is to determine the simple algebras of the variety of C_k -algebra. First, recall that A/θ is a simple algebra if only if θ is a maximal congruence for a given C_k -algebra A, [5, p.59]. Then, taking into account Remark 2.3 and Lemma 3.2, the maximal congruences are determined by the cyclic deductive system.

Theorem 4.3 Let (A, t) be a C_k -algebra and $M \subseteq A$ a cyclic deductive system. Then, the following conditions are equivalents:

- (1) M is maximal,
- (2) if $a \notin M$, then there exists $m \in M$ such that $\bigwedge_{j=1}^{k} t^{j}(\triangle a) \wedge m = 0$,
- (3) if $\bigwedge_{j=1}^{k} t^{j}(\bigtriangleup a) \lor b \in M$, then $a \in M$ or $b \in M$,

(4) if
$$a \notin M$$
, then $\sim \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \in M$,

(5) if $a, b \notin M$, then $a \rightarrow b, b \rightarrow a \in M$.

Proof.

(1) \Rightarrow (2): Let $a \in A$ such that $a \notin M$ and consider D = D(M, a). If $\bigwedge_{j=1}^{k} t^{j}(\triangle a) \land m \neq 0$ holds for every $m \in M$, then from Lemma 4.2 we have that $D \neq A$ and $M \subset D \subset A$, which is impossible.

 $\begin{array}{ll} (2) \Rightarrow (3): \text{ Let } b \in A \text{ such that } \bigwedge_{j=1}^{k} t^{j}(\bigtriangleup a) \lor b \in M \text{ and suppose that } a \notin M. \text{ Then,} \\ \text{by } (2) \text{ there exists } m \in M \text{ such that } \bigwedge_{j=1}^{k} t^{j}(\bigtriangleup a) \land m = 0. \text{ Thus, } (\bigwedge_{j=1}^{k} t^{j}(\bigtriangleup a) \lor b) \land m = \\ (\bigwedge_{j=1}^{k} t^{j}(\bigtriangleup a) \land m) \lor (b \land m) = b \land m \in M \text{ and therefore } b \in M. \\ (3) \Rightarrow (4): \text{ By hypothesis } a \notin M \text{ and since } \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \leq a, \text{ we have } \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \notin M. \text{ On } \\ \text{the other hand, } \sim \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \lor \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) = 1 \in M. \text{ Hence, from the last assertions and} \\ (3) \text{ we can infer } \sim \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \in M, \text{ which completes the proof.} \\ (4) \Rightarrow (5): \text{ Let } a, b \in A \text{ such that } a \notin M \text{ and } b \notin M. \text{ Thus, from } (4) \text{ we have that} \\ \sim \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) \in M. \text{ Since } \sim \bigtriangleup \bigwedge_{j=1}^{k} t^{j}(a) = \sim \bigwedge_{j=1}^{k} \bigtriangleup t^{j}(a) = \bigvee_{j=1}^{k} \sim \bigtriangleup t^{j}(a) = \bigvee_{j=1}^{k} \nabla \sim t^{j}(a) \leq a \\ a \to b, \text{ then } a \to b \in M. \text{ Analogously, it is possible to see that } b \to a \in M. \end{array}$

 $(5) \Rightarrow (1)$: Let us suppose that M is not maximal, then there is cyclic deductive system M'such that $M \subset M'$. Now, let $a \in M' \setminus M$ and $b \in A \setminus M'$, thus by (5) we have $a \to b \in M'$ but from the definition of deductive system we obtain $b \in M'$, which is a contradiction. \Box

In the following we will consider the set $K(A) = \{x \in A : tx = x = \nabla x\}$. Now, let us present the following result essential in the rest of the section.

Proposition 4.4 Let (A, t) be a C_k -algebra and $a \in A$. Then, the following conditions hold:

- (a) K(A) is an Mpm-subalgebra de A,
- (b) K(A) is a Boolean algebra,
- (c) If (S, t_s) is a \mathcal{C}_k -subalgebra of (A, t), then $K(S) = K(A) \cap S$,
- (d) [a) is a c-filter if only if $a \in K(A)$,
- (e) if $a \in K(A)$, then D(a) = [a).

Proof. The proof of (a) and (b) is analogous to the one of [15, Proposition 5.1]. Besides, (c) immediately follows from the very definitions. Moreover, (d) has an analogous proof of [15, Proposition 5.2]. Finally, if $a \in K(A)$, then by (d) we have that [a) is a cyclic deductive system. Besides, if D_1 is a cyclic deductive system such that $a \in D_1$, then it is immediately that $\bigwedge_{j=1}^k t^j \triangle \in D_1$. Therefore, from Lemma 4.1 we conclude that $D(a) \subseteq D_1$ which completes the proof.

Theorem 4.5 Let (A, t) be a C_k -algebra, then the following conditions are equivalent:

(i) (A,t) is simple algebra,

(ii) for every
$$a \in A$$
, $a \neq 1$, $\bigwedge_{j=1}^{k} t^{j}(\triangle a) = 0$

(iii) $K(A) = \{0, 1\}.$

Proof.

(i) \Leftrightarrow (iii): It is analogous to the one of [15, Theorem 5.3]. (i) \Rightarrow (ii): Since (A, t) is simple algebra, then {1} is a maximal cyclic deductive system. From the last assertions and Theorem 4.3 (2), we infer that $\bigwedge_{j=1}^{k} t^{j}(\triangle a) = 0$.

(ii) \Rightarrow (iii): Let $z \in K(A)$ and $z \neq 1$. Since $\triangle z = z = t^j z$ we have that $t^j \triangle z = t^j z$ and therefore $\bigwedge_{j=1}^k t^j(\triangle z) = z = 0$ as desired.

Definition 4.6 For a given C_k -algebra (A, t), we say that (A, t) is r-periodic if r is the smallest non-negative element such that $t^r(x) = x$ for every $x \in A$.

Let T_2 , T_3 , and T_4 be the simple algebras of the variety of Mpm-álgebras. In the following, we will consider the sets $T_{2,k}$, $T_{3,k}$, and $T_{4,k}$ of all sequences $x = (x_1, \dots, x_k)$ with $x_i \in T_{s,k}$ (s = 2, 3, 4) and with the pointwise defined operations. Indeed, these algebras are also C_k -algebras. Taking the function $t : T_{i,k} \to T_{i,k}$ defined as follows $t(x_1, x_2, \dots, x_k) = (x_k, x_1, x_2, \dots, x_{k-1})$ where $(x_1, x_2, \dots, x_k) \in T_{i,k}$ with i = 2, 3, 4. It is not hard to see that $(T_{2,k}, t)$, $(T_{3,k}, t)$ and $(T_{4,k}, t)$ are k-periodic C_k -algebras ;and, in the next, we will show they are simple algebras.

Corollary 4.7 Every C_k -algebras $(T_{i,k}, t)$ with i = 2, 3, 4 is simple one.

Proof. In the first place consider $(T_{4,k}, t)$. By construction, we have that $0, 1 \in K(T_{4,k})$. Let us show that they are the unique elements of $K(T_{4,k})$. Indeed, let $z \in K(T_{4,k})$ such that $z \neq 0$ and $z \neq 1$, then $z = (x_1, x_2, \cdots, x_k)$. Thus, we have the following three cases: (I) $x_i \in \{a, b\}$ for some $i, 1 \leq i \leq k$ or (II) $x_i \in \{0, 1\}$ for every $i, 1 \leq i \leq k$. If we have (I), since $\nabla a = \nabla b = 1$, we infer that $\nabla z \neq z$, which is a contradiction. If (II) holds, from the hypothesis there exists l such that $x_l = 0$ and $x_{l+1} = 1$ or $x_l = 1$ and $x_{l+1} = 0$. In any case, from definition of t we have that $tz \neq z$, which contradicts that $z \in K(T_{4,k})$. On the other hand, from Theorem 4.5 we have that $(T_{4,k}, t)$ is a simple \mathcal{C}_k -algebra. Analogously, it is possible to see that $(T_{2,k}, t)$ and $(T_{3,k}, t)$ are simple \mathcal{C}_k -algebras.

Corollary 4.8 The subalgebras $(T_{i,k}, t)$ with i = 3, 4 are simple algebras.

Proof. It immediately follows from Proposition 4.4.

Lemma 4.9 Every simple C_k -algebra is finite.

Proof. Let A be a simple algebra of the variety. Then, $\{1\}$ is a maximal c-filter of A and from Theorem 2.6 there is an ultrafilter U such that $\{1\} = \bigcap_{i=0}^{k-1} t^i U \cap \varphi(t^i U)$. From the latter, we can infer that the unique ultrafilters of A are $\{t^i U, \varphi(t^i U)\}_{\{0 \le i \le k-1\}}$, which is a finite set. Thus, the prime spectrum of A is finite and so A is finite.

Theorem 4.10 If (A, t) is a simple C_k -algebra, then $(A, t) \simeq (T_{i,d}, t)$ with i = 2, 3, 4 and $T_{i,d}$ is d-periodic for some positive integer d such that d/k.

Proof. According to Lemma 4.9 we have that A is finite and let us suppose that {1} is d-periodic with d/k. Then, $\{1\} = \bigcap_{i=0}^{d-1} t^i P \cap \varphi(t^i P)$ and so the prime spectrum of A is $\{t^j P, \varphi(t^j P)\}_{\{0 \le j \le d-1\}}$, see Lemma 2.8.

On the other hand, since A is a finite lattice then there a prime element $p \in \Pi(A)$ such that P = [p]. That is to say, we can consider the transformation as follows: $\psi(p) = q$

if only if $\varphi([p)) = [q)$. This allows us to identify every prime element with of the prime filter of the spectrum. Thus, p verifies the following cases: type I: $p = \psi(p)$, type II: $p < \psi(p)$ or type III: p and $\psi(p)$ are incomparable. Now, let us suppose p is type I, then the prime spectrum formed by the anti-chain $\{p, tp, \ldots, t^{d-1}p\}$. It is not hard to see that every $t^i p$ is type I. Therefore, there is a function $\alpha : A \to T_2^d$ where $\alpha(p) = (1, 0, \ldots, 0)$, $\alpha(tp) = (0, 1, 0, \ldots, 0), \ldots, \alpha(t^j p) = (0, 0, \ldots, 0, 1_j, 0, \ldots, 0)$. On the other hand, we have that $t\alpha(p) = t(1, 0, \ldots, 0) = (0, 1, 0, \ldots, 0)$ and in general, we obtain that $t^j \alpha(p) =$ $t^{j-1}t\alpha(p) = t^{j-1}t(1, 0, \ldots, 0) = t^{j-1}(0, 1, 0, \ldots, 0) = \cdots = (0, 0, \ldots, 0, 1_j, 0, \ldots, 0)$. From the latter, it is not hard to see that α is a \mathcal{C}_k -isomorphism and therefore, $(A, t) \simeq (T_{2,d}, t)$. Now, if p is type II, the prime spectrum is formed by

$$\{p, tp, \dots, t^{d-1}p, \psi(p), \psi(tp), \dots, \psi(t^{d-1}p)\}$$

where $t^i p < \psi(t^i p)$. Let us observe the prime filters of $T_{3,d}$ are of the form $(0, \ldots, 0, c, 0, \ldots, 0)$ and $(0, \ldots, 0, 1, 0, \ldots, 0)$. Thus, by analogous reasoning from the last case, we have that $(A, t) \simeq (T_{3,d}, t)$. If p is type III, the prime spectrum is the anti-chain

$$\{p, tp, \dots, t^{d-1}p, \psi(p), \psi(tp), \dots, \psi(t^{d-1}p)\}.$$

The prime elements of $T_{4,d}$ are the form of $(0, \ldots, 0, a, 0, \ldots, 0)$ and $(0, \ldots, 0, b, 0, \ldots, 0)$. Therefore, by analogous reasoning we can see that $(A, t) \simeq (T_{4,d}, t)$, which completes the proof.

It is well-known that divides relation "/" between integers is a partial order. Then, for a given positive integer k, we have the set $Div(k) = \{z : z \text{ divisor of } k\}$ can be considered as distributive latices; moreover, $k_1 \wedge k_2 = lcm(k_1, K_2)$ and $k_1 \vee k_2 = gcd(k_1, K_2)$ are the *least common multiple* and greatest common divisor, respectively. Now, it is possible to see that for every k-periodic Boolean algebra, we have \mathcal{B}_k is lattice-isomorphic to the Boolean algebra Div(k). Furthermore, for every $d \in Div(k)$ there is a unique \mathcal{B}_g associated with d which is a subalgebra of \mathcal{B}_k . Besides, \mathcal{B}_g is d-periodic characterized by $B_d = \{g \in B_k : t \text{ is an automorphism and } t^d g = g\}.$

Lemma 4.11 The subalgebras of $(T_{i,k}, t)$ are of the form $(T_{i,d}, t_d)$ with d/k and i = 2, 3, 4. Besides, $T_{2,d}$ is a C_k -subalgebra of $T_{3,d}$ and $T_{4,d}$, but $T_{3,d}$ is not subalgebra of $T_{4,d}$.

Proof. Let us take $T_{i,k}^d = \{x \in T_i^k : t^d x = x\}$, then it is clear that $(T_{i,k}^d, t_d')$ is a subalgebra of $(T_{i,k}, t)$ where the restriction $t|_{T_{i,k}^d} = t_d'$ is an automorphism over $T_{i,k}^d$. On the other hand, from Lemma 2.8 we have that the prime spectrum of $T_{i,k}^d$ has the form $\{t^i P, \varphi(t^i P)\}_{\{0 \le i \le d-1\}}$. So, $(T_{i,k}^d, t_d') \simeq (T_{i,d}, t_d)$.

On the other hand, it is clear that $(B(T_{i,k}), t_B) \simeq B_k$ for i = 2, 3 where $t|_{B(T_{i,k})} = t_B$ and B(A) is the set of Boolean elements of \mathcal{C}_k -alegbra A. Thus, there is d such that d/kand B_d is a sublagebra of B_k . So, since $(B(T_{i,d}), t_B) \simeq B_d$, we have that the sublagebras $(T_{i,k}, t)$ with i = 2, 3 are the form $(T_{i,d}, t_d)$. For the case i = 4, it is clear that $B(T_{4,k}) = T_{4,k}$. Let us suppose that d is a divisor of k, then $k = d \cdot q$. Now, we can consider the elements of $T_{4,k}$ as a sequence $x = (\overline{x}_1, \overline{x}_2, \ldots, \overline{x}_q)$ where $\overline{x}_1 = (x_1, \ldots, x_q), \ldots, \overline{x}_q = (x_{dq-q}, x_{dq-(q-1)}, \ldots, x_{dq})$. Now, let us consider the set $D = \{x \in T_{4,k} : x = (\overline{x}_1, \overline{x}_2, \ldots, \overline{x}_q), \text{ such that } \overline{x}_1 = \overline{x}_2 = \ldots = \overline{x}_d\}$. Then, it is clear that D is a subalgebra of $T_{4,k}$ and taking the automorphism $t|_D : D \to D$, we have $(D, t|_D)$ is a d-periodic. Their atoms are the form $a_1 = (a, 0, \ldots, 0), \ldots, a_d = (0, 0, \ldots, a)$ and $b_1 = (b, 0, \ldots, 0), \ldots, b_d = (0, 0, \ldots, b)$ and it is verified $t(a_i) = a_{i+1}$ and $t(b_i) = b_{i+1}$. Thus, $(D, t|_D) \simeq (T_{4,d}, t_d)$.

To end the proof, we will determine all sublagebras of $T_{4,k}$ where the elements of the sequences belong to T_2 . Let us take

$$D' = \{x \in T_{4,k} : x = (\overline{x}_1, \overline{x}_2, \dots, \overline{x}_q), \text{ such that } \overline{x_1} = \overline{x_2} = \dots = \overline{x_d} \in T_{2,q}\}$$

and

$$t' = t|_{D'} : D' \to D',$$

then it is not hard to see that $(D', t') \simeq (T_{2,d}, t_d)$. Therefore, $(T_{2,d}, t_d)$ and $(T_{4,d}, t_d)$ are the unique subalgebras of $(T_{4,k}, t)$. Since T_3 is not subalgebra of T_4 , then $T_{3,d}$ is not subalgebra of $T_{4,d}$, which completes the proof.

The following Lemma will be used in the next Section and it is an immediate consequence of Lemma 4.11.

Lemma 4.12 $T_{i,d_1} \cap T_{i,d_2} = T_{i,gcd(d_1,d_2)}, T_{2,d_1} \cap T_{i,d_2} = T_{2,gcd(d_1,d_2)}, T_{3,d_1} \cap T_{4,d_2} = T_{2,gcd(d_1,d_2)}$ with i = 2, 3, 4

As directly consequence from Corollary 4.7, Lemma 4.9, Theorem 4.10 and universal algebra results given in [5, Theorem 10.16], we conclude:

Theorem 4.13 The variety of C_k -algebras is finitely generated and locally finite. Besides, the only simple algebras are $(T_{i,k}, t)$ with i = 3, 4 and their subalgebras.

5 The cardinality of the finitely generated C_k -algebras

In this section, we will focus on the task of studying the structure of free C_k -algebras with a finite number of generators. We denote by $F_{\mathcal{C}_k}(n)$ this algebra where n is a positive integer. From the results showed in the last section, we know that for every d divisor of k, the family \mathcal{C} of the maximal c-filters of $F_{\mathcal{C}_k}(n)$ can be partitioned in the following way:

$$N_{i,d} = \{ N \in \mathcal{C} : F_{\mathcal{C}_k}(n) / N \simeq T_{i,d} \}.$$

Thus, form Theorem 4.13 we have:

$$F_{\mathcal{C}_k}(n) \simeq \prod_{d/k} T_{2,d}^{\alpha_{2,d}} \times \prod_{d/k} T_{3,d}^{\alpha_{3,d}} \times \prod_{d/k} T_{4,d}^{\alpha_{4,d}},$$

where $\alpha_{i,d} = |N_{i,d}|$ with i = 2, 3, 4.

Now, let us consider $Epi(F_{\mathcal{C}_k}(n), T_{i,d})$ the set of all epimorphisms from $F_{\mathcal{C}_k}(n)$ into $T_{i,d}$ and we denote by $Aut(T_{i,d})$ the set of all automorphisms over the algebra $T_{i,d}$ and |X| denotes the number of elements of X. It is not hard to see that $Aut(T_{l,d})$ (with l = 2, 3) has only d automorphisms and they are $t, t^2, \cdots, t^{d-1}, t^d$. Besides, it is possible to prove that $|Aut(T_{4,d})| = 2d$. Taking the function $s : Epi(F_{\mathcal{C}_k}(n), T_{i,d}) \to N_{i,d}$ defined by $s(h) = ker(h) = h^{-1}(\{1\})$ for every $h \in Epi(F_{\mathcal{C}_k}(n), T_{i,d})$, we can see that s is onto and $s^{-1}(N) = \{\alpha \circ h : \alpha \in Aut(T_{i,d})\}$. Therefore,

$$\alpha_{i,d} = \frac{|Epi(F_{\mathcal{C}_k}(n), T_{i,d})|}{|Aut(T_{l,d})|}$$

On the other hand, for every $h \in Epi(F_{\mathcal{C}_k}(n), T_{i,d})$ there is a function $f : G \to T_{i,d}$ such that $f = h|_G$. Now, if $F^*(G, T_{i,d}) = \{f : G \to T_{i,d} \text{ such that } [f(G)]_{\mathcal{C}_k} = T_{i,d}\}$, then we have:

$$|Epi(F_{\mathcal{C}_k}(n), T_{i,d})| = |F^*(G, T_{i,d})|.$$

Where $[f(G)]_{\mathcal{C}_k}$ denotes the \mathcal{C}_k -algebra generated by f(G). Let us observe that the condition $[f(G)]_{\mathcal{C}_k} = T_{i,d}$ is equivalent to asking $f(G) \subseteq T_{i,d}$ and $f(G) \not\subseteq S$ for every maximal subalgebra S of $T_{i,d}$. If we denote M(d) the set of maximal divisors of d different of d. Then, for every maximal subalgebra of $T_{i,d}$ are of the form $T_{i,x}$ with $x \in M(d)$. Thus, we can write:

$$F^*(G, T_{i,d}) = F_{i,d} \setminus \bigcup_{l \le i} \bigcup_{x \in M(d)} F_{l,x},$$

where $F_{i,d}$ is the set of all functions from G into $T_{i,d}$. Therefore,

$$F^*(G, T_{i,d})| = |F_{i,d}| - |\bigcup_{l \le i} \bigcup_{x \in M(d)} F_{l,x}| = (i^d)^n - |\bigcup_{l \le i} \bigcup_{x \in M(d)} F_{l,x}|.$$

On the other hand, it is well-known that every finite set \mathcal{J} and the family of finite sets $\{A_j\}_{j\in\mathcal{J}}$ verify:

$$|\bigcup_{j\in\mathcal{J}}A_i| = \sum_{X\subseteq\mathcal{J},\,X\neq\emptyset}(-1)^{|X|-1}|\bigcap_{j\in X}A_j|.$$

Now, let us observe that if we take $I = \{w : w \le i\}$, then:

$$|\bigcup_{l \le i} \bigcup_{x \in M(d)} F_{l,x}| = |\bigcup_{(z,t) \in I \times M(d)} F_{z,t}| = \sum_{X \subseteq I \times M(d)} (-1)^{|X|} |\bigcap_{(j_1,j_2) \in X} F_{j_1,j_2}|,$$

where

$$\bigcap_{(j_1,j_2)\in X} F_{j_1,j_2} = \{ f \in F_{i,d} : f : G \to \bigcap_{(j_1,j_2)\in X} T_{j_1,j_2} \}.$$

Calculating $\alpha_{i,d}$

If i = 2, then $I = \{2\}$ and therefore the following holds:

$$\alpha_{2,d} = \frac{(2^d)^n - |\bigcup_{x \in M(d)} F_{2,x}|}{d} = \frac{(2^d)^n - \sum_{Z \subseteq M(d), Z \neq \emptyset} (-1)^{|Z|-1} |\bigcap_{w \in Z} F_{2,w}|}{d}$$
$$= \frac{(2^d)^n - \sum_{Z \subseteq M(d), Z \neq \emptyset} (-1)^{|Z|-1} (2^{mcd(Z)})^n}{d},$$

where gcd(Z) is the set of greatest common divisors of Z.

If i = 3, then:

$$\alpha_{3,d} = \frac{(3^d)^n - |\bigcup_{(j,x)\in\{2,3\}\times M(d)} F_{j,x}\cup F_{2,d}|}{d} = \frac{(3^d)^n - |F_{2,d}\cup\bigcup_{y\in M(d)} F_{3,y}|}{d}.$$

Since

$$|F_{2,d} \cup \bigcup_{y \in M(d)} F_{3,y}| = |F_{2,d}| + |\bigcup_{y \in M(d)} F_{3,y}| - |F_{2,d} \cap \bigcup_{y \in M(d)} F_{3,y}|$$

and

$$|F_{2,d} \cap \bigcup_{y \in M(d)} F_{3,y}| = |\bigcup_{y \in M(d)} (F_{2,d} \cap F_{3,y})|,$$

we have that

$$\alpha_{3,d} = \frac{(3^d)^n - (2^d)^n - \sum_{W \subseteq M(d), W \neq \emptyset} (-1)^{|W| - 1} (3^{gcd(W)})^n + \sum_{H \subseteq M(d) \times M(d), H \neq \emptyset} (-1)^{|H| - 1} (2^{gcd(H_1 \cup H_2)})^n}{d}$$

where $H_1 = \{x : (x, y) \in H\}$ and $H_2 = \{y : (x, y) \in H\}.$

Finally, for i = 4 and by Lemma 4.11, we have:

$$\alpha_{4,d} = \frac{(4^d)^n - |\bigcup_{(j,x) \in \{2,4\} \times M(d)} F_{j,x} \cup F_{2,d}|}{2d} = \frac{(4^d)^n - |F_{2,d} \cup \bigcup_{y \in M(d)} F_{4,y}|}{2d}$$

By analogous reasoning to the last case, we have:

$$\alpha_{4,d} = \frac{(4^d)^n - (2^d)^n - \sum_{W \subseteq M(d), W \neq \emptyset} (-1)^{|W| - 1} (4^{gcd(W)})^n + \sum_{H \subseteq M(d) \times M(d), H \neq \emptyset} (-1)^{|H| - 1} (2^{gcd(H_1 \cup H_2)})^n}{2d}$$

From the last assertions, we have proved the following Theorem:

Theorem 5.1 Let $F_{\mathcal{C}_k}(n)$ be the free \mathcal{C}_k -algebra with n generators, then:

$$|F_{\mathcal{C}_k}(n)| = \prod_{d/k} 2^{\alpha_{2,d}} \cdot \prod_{d/k} 3^{\alpha_{3,d}} \cdot \prod_{d/k} 4^{\alpha_{4,d}}$$

with

$$\begin{aligned} \alpha_{2,d} &= \frac{(2^d)^n - \sum\limits_{Z \subseteq M(d), Z \neq \emptyset} (-1)^{|Z|-1} (2^{gcd(Z)})^n}{d}, \\ \alpha_{3,d} &= \frac{(3^d)^n - (2^d)^n - \sum\limits_{W \subseteq M(d), W \neq \emptyset} (-1)^{|W|-1} (3^{gcd(W)})^n + \sum\limits_{H \subseteq M(d) \times M(d), H \neq \emptyset} (-1)^{|H|-1} (2^{gcd(H_1 \cup H_2)})^n}{d}, \\ \alpha_{4,d} &= \frac{(4^d)^n - \sum\limits_{W \subseteq M(d), W \neq \emptyset} (-1)^{|W|-1} (4^{gcd(W)})^n - (2^d)^n + \sum\limits_{H \subseteq M(d) \times M(d), H \neq \emptyset} (-1)^{|H|-1} (2^{gcd(H_1 \cup H_2)})^n}{2d}. \end{aligned}$$

5.1 Particular cases

In this subsection, we will show that Theorem 5.1 has particular cases that was previously obtained in the literature in [19, 15].

\mathcal{C}_1 -algebras

If k = 1, the class of C_1 -algebras coincides with the one of mpM-algebras. According to Theorem 5.1, we have that the free C_1 -algebra with r generators is:

$$F_{\mathcal{C}_1}(r) \simeq T_{2,1}^{\alpha_{2,1}} \times T_{3,1}^{\alpha_{3,1}} \times T_{4,1}^{\alpha_{4,1}}.$$

Since $M(1) = \emptyset$ we have that $\sum_{Z \subseteq M(1)} (-1)^{|Z|} (2^{gcd(Z)})^r = 0$ and therefore $\alpha_{2,1} = 2^r$. Besides, it is possible to see that

$$\sum_{\substack{H \subseteq M(1) \times M(1), H \neq \emptyset}} (-1)^{|H|} (2^{gcd(H_1 \cup H_2)})^r = 0,$$
$$\sum_{\substack{W \subseteq M(d), W \neq \emptyset}} (-1)^{|W| - 1} (4^{gcd(W)})^r = 0 = \sum_{\substack{W \subseteq M(d), W \neq \emptyset}} (-1)^{|W| - 1} (3^{gcd(W)})^r,$$

and thus, $\alpha_{3,1} = 3^r - 2^r$ and $\alpha_{4,1} = \frac{4^r - 2^r}{2} = 2^{r-1}(2^r - 1)$. Therefore,

$$|F_{\mathcal{C}_1}(n)| = 2^{2^r} \times 3^{3^r - 2^r} \times 4^{2^{r-1}(2^r - 1)}.$$

This number was presented in [19, Theorem 4.5].

\mathcal{C}_2 -algebras

For the k = 2, we have the class of C_2 -algebras coincides with the one of S-algebras introduced in [15]. Then, the free C_2 -algebra with r generators is:

$$F_{\mathcal{C}_{2}}(r) \simeq \prod_{d=1}^{2} T_{2,d}^{\alpha_{2,d}} \times \prod_{d=1}^{2} T_{3,d}^{\alpha_{3,d}} \times \prod_{d=1}^{2} T_{4,d}^{\alpha_{4,d}}.$$
Since $|T_{i,d}| = i^{d}$, $\alpha_{2,2} = \frac{4^{r}-2^{r}}{2}$, $\alpha_{3,2} = \frac{9^{r}-3^{r}-4^{r}+2^{r}}{2}$ and $\alpha_{4,2} = \frac{4^{2r}-4^{r}-2^{2r}+2^{r}}{4}$, then
$$|F_{\mathcal{C}_{2}}(r)| = |T_{2,1}|^{2^{r}} \times |T_{2,2}|^{\frac{4^{r}-2^{r}}{2}} \times |T_{3,1}|^{3^{r}-2^{r}} \times |T_{3,2}|^{\frac{3^{2r}-3^{r}-2^{2r}+2^{r}}{2}} \times |T_{4,1}|^{\frac{4^{r}-2^{r}}{2}} \times |T_{4,2}|^{\frac{4^{2r}-4^{r}-2^{2r}+2^{r}}{4}}$$

$$= 2^{2^{r}} \times 4^{\frac{4^{r}-2^{r}}{2}} \times 3^{3^{r}-2^{r}} \times 9^{\frac{3^{2r}-3^{r}-2^{2r}+2^{r}}{2}} \times 16^{\frac{4^{r}-2^{r}}{2}} \times 16^{\frac{4^{2r}-4^{r}-2^{2r}+2^{r}}{4}}$$

$$= 2^{2^{r}} \times 4^{2\cdot\frac{4^{r}-2^{r}}{2}} \times 3^{3^{r}-2^{r}} \times 9^{\frac{9^{r}-3^{r}-4^{r}+2^{r}}{2}} \times 16^{\frac{16^{r}-4^{r}+4^{r}+2^{r}}{4}}$$

$$= 2^{2^{r}} \times 4^{2\cdot\frac{4^{r}-2^{r}}{2}} \times 3^{3^{r}-2^{r}} \times 9^{\frac{9^{r}-3^{r}-4^{r}+2^{r}}{2}} \times 16^{\frac{16^{r}-3\cdot4^{r}+2\cdot2^{r}+4^{r}-2^{r}}{4}} \times 16^{\frac{4^{r}-2^{r}}{4}}$$

$$= 2^{2^{r}} \times 4^{3\cdot\frac{4^{r}-2^{r}}{2}} \times 3^{3^{r}-2^{r}} \times 9^{\frac{9^{r}-3^{r}-4^{r}+2^{r}}{2}} \times 16^{\frac{16^{r}-3\cdot4^{r}+2\cdot2^{r}}{4}} \times 16^{\frac{4^{r}-2^{r}}{4}}$$

$$= 2^{2^{r}} \times 4^{3\cdot\frac{4^{r}-2^{r}}{2}} \times 3^{3^{r}-2^{r}} \times 9^{\frac{9^{r}-3^{r}-4^{r}+2^{r}}{2}} \times 16^{\frac{16^{r}-3\cdot4^{r}+2\cdot2^{r}}{4}} \times 16^{\frac{4^{r}-2^{r}}{4}}$$

This number was obtained in [15, Theorem 6.4].

 \mathcal{C}_k -algebras with k is prime number

$$|F_{\mathcal{C}_{k}}(r)| = \prod_{N \in N_{2,d}, d/k} |T_{2,d}|^{|N_{2,d}|} \times \prod_{N \in N_{3,d}, d/k} |T_{3,d}|^{|N_{3,d}|} \times \prod_{N \in N_{4,d}, d/k} |T_{4,d}|^{|N_{4,d}|}$$
$$= \prod_{d/k} (2^{d})^{\alpha_{2,d}} \times \prod_{d/k} (3^{d})^{\alpha_{3,d}} \times \prod_{d/k} (4^{d})^{\alpha_{4,d}}$$
$$= 2^{\alpha_{2,1}} \times (2^{k})^{\alpha_{2,k}} \times 3^{\alpha_{3,1}} \times (3^{k})^{\alpha_{3,k}} \times 4^{\alpha_{4,1}} \times (4^{k})^{\alpha_{4,k}}$$
$$= 2^{2^{r}} \times (2^{k})^{\frac{2^{kr} - 2^{r}}{k}} \times 3^{3^{r} - 2^{r}} \times (3^{k})^{\frac{3^{kr} - 3^{r} - 2^{kr} + 2^{r}}{k}} \times 4^{\frac{4^{r} - 2^{r}}{2}} \times (4^{k})^{\frac{4^{kr} - 4^{r} - 2^{kr} + 2^{r}}{2k}}$$

6 Degree-preserving logic associated with the class C_k -algebras

In this section, we show that the class of C_k -algebras can be seen as suitable algebraic semantics for a logic system. There are many ways to associate a logic system with an ordered algebraic structure. For instance, it is possible to define the *logic that preserves degrees of truth* associated with the class of C_k -algebras. This notion was introduced and studied mainly in the works [3, 27, 28, 29] with the purpose of taking advantage of the multiple values of truth, more that the two classical values of truth, in opposition to the logics that only preserve truth. These logics preserve lower bounds of truth values instead of just preserving value 1.

From now on, we shall denote by $\mathfrak{Fm} = \langle Fm, \wedge, \vee, \sim, *, t, \top, \bot \rangle$ the absolutely free algebra of type (2,2,1,1,1,0) generated by some denumerable set of variables. We call Fm the sentential formulas, and we shall refer to them by lowercase greek letters $\alpha, \beta, \gamma, \ldots$ and so on; and we shall denote the sets of formulas by uppercase greek letters Γ, Δ , etc..

Let $\mathbb{L}_k^{\leq} = \langle Fm, \models_k^{\leq} \rangle$, for every $\Gamma \cup \{\alpha\} \subseteq Fm$

(i) If
$$\Gamma = \{\alpha_1, \ldots, \alpha_n\} \neq \emptyset$$
,

$$\alpha_1, \dots, \alpha_n \models_k^{\leq} \alpha \iff \forall (A, t) \in \mathbb{C}_k, \forall v \in Hom_{\mathbb{C}_k}(Fm, (A, t)), \forall a \in A$$

if $v(\alpha_i) \geq a$, for all $i \leq n$, then $v(\alpha) \geq a$

(ii)
$$\emptyset \models_k^{\leq} \alpha \iff \forall A \in \mathbb{C}_k, \forall v \in Hom_{\mathbb{C}_k}(Fm, A), v(\alpha) = 1.$$

(iii) If $\Gamma \subseteq Fm$ is infinity,

$$\Gamma \models_k^{\leq} \alpha \iff$$
 there are $\alpha_1, \ldots, \alpha_n \in \Gamma$ such that $\alpha_1, \ldots, \alpha_n \models_k^{\leq} \alpha$.

From the above definition we have that \mathbb{L}_k^{\leq} is a sentential logic, that is, \models_k^{\leq} is a finitary consequence relation on Fm. Besides:

Proposition 6.1 Let $\alpha_1, \ldots, \alpha_n, \alpha \in Fm$. Then, the following conditions are equivalent

- (i) $\alpha_1, \ldots, \alpha_n, \models_k^{\leq} \alpha,$
- (ii) the inequality $\alpha_1 \wedge \cdots \wedge \alpha_n \preccurlyeq \alpha$ holds in the variety \mathbb{C}_k ,
- (iii) for every $(A,t) \in \mathbb{C}_k$ and every $h \in Hom_{\mathbb{C}_k}(Fm, (A,t))$, it is held that $h(\alpha_1) \wedge \cdots \wedge h(\alpha_n) \preccurlyeq h(\alpha)$.

Proof. (i) \Rightarrow (ii): Suppose that $\alpha_1, \ldots, \alpha_n \models_k^{\leq} \alpha$. Then, $\forall A \in \mathbb{C}_k, \forall v \in Hom_{\mathbb{C}_k}(Fm, A)$, $\forall a \in A$, if $v(\alpha_i) \geq a$, for all $i \leq n$, then $v(\alpha) \geq a$. Since A is a lattice, we have that $v(\alpha_i) \geq a$, for all $i \leq n$ iff $\bigwedge_{i=1}^n v(\alpha_i) \geq a$ iff $v(\bigwedge_{i=1}^n (\alpha_i)) \geq a$. So, if $v(\bigwedge_{i=1}^n (\alpha_i)) \geq a$, then $v(\alpha) \geq a$; that is, $\alpha_1 \wedge \cdots \wedge \alpha_n \preccurlyeq \alpha$.

 $(ii) \Rightarrow (iii)$ and $(iii) \Rightarrow (i)$: Immediate.

The logics of formal inconsistency (LFI) was introduced by Carnielli and Marcos in [8]. These systems were invented with the intention to generalize the da Costa's idea to have paraconsistent logics with a unary operator that allows us to recover the classical theorems. Recall that a logic $\mathcal{L} = \langle Fm, \vdash_{\mathcal{L}} \rangle$ is said to be strong LFI if the following conditions hold:

- (i) if p and q are two different propositional variables then:
 - (i.a) $p, \sim p \not\vdash_{\mathcal{L}} q$,
 - (i.b) $\circ p, p \not\vdash_{\mathcal{L}} q$,
 - (i.c) $\circ p, \sim p \not\vdash_{\mathcal{L}} q$,
- (ii) $\circ \varphi, \varphi, \sim \varphi \vdash_{\mathcal{L}} \psi$ for every φ and ψ .

The condition (i.a) for sentences is typically used to define paraconsistent logic. On the other hand, the notion of Logic Formal Undeterminedness (LFU) was introduced by Marcos in [31], they are paracomplete systems with a unary operator that permit us to recover the Excluded Middle. That is to say, if the following holds:

- (i) $\not\vdash_{\mathcal{L}} \varphi \lor \sim \varphi$, for some φ ; i.e., \mathcal{L} is paracomplete;
- (ii) there is a formula φ such that
 - (ii.a) $\circ_k \varphi \not\vdash_{\mathcal{L}} \varphi;$ (ii.b) $\circ_k \varphi \not\vdash_{\mathcal{L}} \sim \varphi;$
- (iii) $\circ_k \varphi \vdash_{\mathcal{L}} \varphi \lor \sim \varphi$, for every φ .

In the rest of the section, we will show that \mathbb{L}_k^{\leq} is LFI and LFU. Now, let us consider a unary operator \circ_k for \mathbb{L}_k^{\leq} defined as follows:

$$\circ \alpha = \bigwedge_{i=0}^{k} t^{i} ((\sim \alpha \lor \alpha) \land (\alpha \land \sim \alpha)^{*})$$

Now, let us consider the Boolean set of a given C_k -algebra (A, t), $B(A) = \{x \in A : x \land x = 0\}$. Thus, it is easy to see the following Proposition.

Proposition 6.2 Let (A, t) be a C_k -algebra, then $x \in B(A)$ iff $\circ x = 1$.

Proof. Suppose that $x \in B(A)$, then $\circ x = \bigwedge_{i=0}^{k} t^{i}((\sim x \lor x) \land (x \land \sim x)^{*}) = 1$. Conversely, let us suppose $\circ x = 1$. Then, $\circ x = \bigwedge_{i=0}^{k} t^{i}((\sim x \lor x) \land (x \land \sim x)^{*}) = 1$ and so $\sim x \lor x = 1$.

Lemma 6.3 \mathbb{L}_k^{\leq} is a genuine paraconsistent logic.

Proof. Let us consider C_k -algebra (T_4, Id) where T_4 is the simple mpM-algebra and id is the identity function. To see that (i.a) holds, it is enough to take the homomorphism $v: Fm \to (T_4, Id)$ such that $v(p) = a = v(\sim p)$ and v(q) = 0, then $v(p \land \sim p) > v(q)$. Using the same valuation, we have $\not\models_k^{\leq} \sim (p \land \sim p)$, see [2] and [36].

Lemma 6.4 The logic \mathbb{L}_{k}^{\leq} is a strong LFI with respect to \sim and \circ .

Proof. Let us consider C_k -algebra (T_4, Id) where T_4 is the simple mpM-algebra and id is the identity function. To see that (i.a) holds, it is enough to take the homomorphism $v: Fm \to (T_4, Id)$ such that $v(p) = a = v(\sim p)$ and v(q) = 0, then $v(p \land \sim p) > v(q)$. For the case (i.b), we get a function $v(\circ\top) = 1$ $v(\top) = 1$ and v(q) = 0, then $v(\circ\top \land\top) > v(q)$. For (i.c), we get a function $v(\circ\bot) = 1$ $v(\sim \bot) = 1$ and v(q) = 0, then $v(\circ\bot \land \sim \bot) > v(q)$.

Now, for the case (ii), we can observe that the identity $((\sim x \lor x) \land (x \land \sim x)^*) \land x \land \sim x = 0$ holds in mpM- algebras T_2 and T_4 . Then, since the class of mpM-algebra is a variety and the identity is in the language of mpM-algebra, we can affirm that the identity hold in every C_k -algebra (A, t), then $v(\circ p \land p \land \sim p) \leq v(q)$ for every variable p and q, which complete the proof.

Lemma 6.5 The logic \mathbb{L}_k^{\leq} is an LFU with respect to \sim and \circ .

Proof. Taking the same simple algebra (T_4, Id) of Lemma 6.4 and $v(p) = a = v(\sim p)$, we have that $\not\models_k^{\leq} p \lor \sim p$. Now, taking $\circ \perp \not\models_k^{\leq} \perp$ and $\circ \top \not\models_k^{\leq} \sim \top$. To see that $\circ \alpha \models_k^{\leq} \alpha \lor \sim \alpha$, it is enough to observe that the following algebraic property $\circ x \leq x \lor \sim x$ hold on every \mathcal{C}_k -algebra.

Next, we will see that the operator \circ enjoys the *propagation property* in the logic \mathbb{L}_k . That is to say:

Lemma 6.6 In the logic \mathbb{L}_k^{\leq} the following holds:

- (i) $\models_k^{\leq} \circ \perp$,
- (ii) $\circ \alpha \models_k^{\leq} \circ \#_1 \alpha \text{ where } \#_1 \in \{\sim, ^*, t\},$
- (iii) $\circ \alpha, \circ \beta \models_k^{\leq} \circ (\alpha \# \beta)$ where $\# \in \{\land, \lor\}$.

Proof. It is not hard to see that the algebraic equations $\circ(0) = 1$, $((x \lor \sim x) \land (x \land \sim x)^*) \land ((y \lor \sim y) \land (y \land \sim y)^*) \leq ((x \# y) \lor \sim (x \# y)) \land ((x \# y) \land \sim x(\# y))^*$ (where $\# \in \{\land,\lor\}$), $\circ x = \circ(\sim x)$ and $\circ(x) = ((x \lor \sim x) \land (x \land \sim x)^*) = ((x^* \lor \sim x^*) \land (x^* \land \sim x^*)^*) = \circ(x^*)$ hold in T_3 and T_4 . Therefore, they are held in every mpM-algebra. So, taking in mind the definition of \circ and that t is a homomorphism, we have that $\circ 1 = 1$, $\circ x \land \circ y \leq \circ(x \# y)$ (where $\# \in \{\land,\lor\}$) and $\circ x = \circ(\#_1 x)$ (with $\#_1 \in \{\sim, *, t\}$) are held in every C_k -algebra. \Box

For every positive integer k, we have that \mathbb{L}_k^{\leq} is LFI and LFU, the same phenomenon occurs in the logics studied in [9] and [14], see also [39].

6.1 \mathbb{L}_k^{\leq} as a generalized matrix logic

Recall that a *matrix logic* is a pair $\langle A, F \rangle$ where A is an algebra and F is a non-empty set of A. On the other hand, a *generalized matrix* is a pair $\langle A, C \rangle$ where A is an algebra and C is a family of subsets of A that verify the following:

- (i) $A \in \mathcal{C}$,
- (ii) If $\{X_i\}_{i \in I}$ then $\bigcup_{i \in I} X_i \in \mathcal{C}$.

Now, let us consider the matrix $M = \langle A, F \rangle$. Then the logic $\mathbb{L}_M^k = \langle Fm, \models_M \rangle$ is determined by M in the following way:

let $\Gamma \cup \{\alpha\}$ a subset of Fm

$$\Gamma \models_M \alpha \iff \forall v \in Hom(Fm, A),$$

if $v(\gamma) \in F$, for all $\gamma \in \Gamma$, then $v(\alpha) \in F$

If $\mathcal{M} = \{M_i\}_{i \in I}$ is a family of matrices, then the logic $\mathbb{L}^k_{\mathcal{M}} = \langle Fm, \models_{\mathcal{M}} \rangle$ determined by the family \mathcal{M} is defined by $\models_{\mathcal{M}} = \bigcap_{i \in I} \models_{M_i}$.

Let us consider now the logic determined by generalized matrix $\mathcal{G} = \langle A, \mathcal{C} \rangle$ that we denote $\mathbb{L}^k_{\mathcal{G}} = \langle Fm, \models_{\mathcal{G}} \rangle$ and it is defined by the following: let $\Gamma \cup \{\alpha\}$ a subset of Fm

$$\begin{array}{ll} \Gamma \models_{\mathcal{G}} \alpha & \Longleftrightarrow & \forall v \in Hom(Fm, A), \, \forall G \in \mathcal{C} \\ & \text{if } v(\gamma) \in G, \, \text{for all } \gamma \in \Gamma, \, \text{then } v(\alpha) \in G \end{array}$$

In this case, we take

 $\mathcal{M} = \{ \langle A, F \rangle : A \in \mathbb{C}_k, \text{ and } F \text{ is a lattice filter of } A \}$

and

$$\mathcal{G} = \{ \langle A, \mathcal{F}_i(A) \rangle : A \in \mathbb{C}_k \}$$

where $\mathcal{F}_i(A)$ is the family of all filters of A. Then, we have the following Lemma:

Lemma 6.7 The logic \mathbb{L}_{k}^{\leq} coincides with both logics $\mathbb{L}_{\mathcal{M}}^{k}$ and $\mathbb{L}_{\mathcal{G}}^{k}$.

Proof. The fact that the logics $\mathbb{L}^k_{\mathcal{M}}$ and $\mathbb{L}^k_{\mathcal{G}}$ coincide follows immediately from their very definition of them. On the other hand, in the variety \mathbb{C}_k the notion of filter for every algebra is elementary definable, then the class \mathcal{M} is closed under ultraproducts. Therefore, $\mathbb{L}^k_{\mathcal{M}}$ is finitary.

Now, we will see that the systems \mathbb{L}_{k}^{\leq} and $\mathbb{L}_{\mathcal{G}}^{k}$ coincide; that is to say, $\models_{k}^{\leq}=\models_{\mathcal{G}}$. Since $\models_{\mathcal{G}}$ is finitary, we make the proof using a finite set of premises. Indeed, let $\alpha_{1}, \dots, \alpha_{n}, \alpha$ be a set of formulas such that $\alpha_{1}, \dots, \alpha_{n} \models_{k}^{\leq} \alpha$. According to Proposition 6.1, for every $v \in Hom_{\mathbb{C}_{k}}(Fm, A)$ we have that $v(\alpha_{1}) \wedge \dots \wedge v(\alpha_{n}) \leq v(\alpha)$. Let us consider now $F \in \mathcal{F}_{i}(A)$ and suppose that $v(\alpha_{1}), \dots, v(\alpha_{n}) \in F$. Since F is a filter and taking in mind the definition of $\mathbb{L}_{\mathcal{G}}^{k}$, we have that $\alpha_{1}, \dots, \alpha_{n} \models_{\mathcal{G}} \alpha$.

Conversely, suppose that $\alpha_1, \dots, \alpha_n \models_{\mathcal{G}} \alpha$ and let $v \in Hom_{\mathbb{C}_k}(Fm, A)$. Now, taking a filter generated G by the set $\{v(\alpha_1), \dots, v(\alpha_n)\}$, we have that $v(\alpha_i) \in G$ with $i = 1, \dots, n$. Then, $v(\alpha) \in G$. Again taking into account Proposition 6.1, we obtain that $\alpha_1, \dots, \alpha_n \models_k^{\leq} \alpha$ as desired.

6.2 Logic \mathbb{L}_{k}^{\leq} location inside the hierarchy of Abstract algebraic logic

The theory of *algebrizable logics* was initiated by Blok and Pigozzi in [4], in that work the notion of *Leibniz operator* to characterize them was introduced. Other weak notions of algebrizable logic can be characterized by the study of Leibniz operator. Now, we will summarize the main background to develop the rest of the section.

If $\langle \mathbf{A}, F \rangle$ is an arbitrary matrix, its *Leibniz congruence* $\Omega_{\mathbf{A}}F$ is the largest of all congruences of \mathbf{A} that are compatible with F in the following sense:

$$\Omega_{\mathbf{A}}F := \{ \Theta \in \mathrm{Co}\mathbf{A} : \mathrm{if} (a, b) \in \Theta \text{ and } a \in F \mathrm{ then } b \in F \}$$

Where CoA is the poset of all congruences of A. Besides, a matrix is said to be **reduced** when $\Omega_A F$ is the identity; that is to say, the identity relation is only congruence compatible with F.

To define the notion of algebraic counterpart of a logic L, we need to consider the class of $Alg^*(L)$ of algebras reducts of the reduced models of L as follows:

$$Alg^*(L) := \{ \mathbf{A} : \exists F \subseteq A, \langle A, F \rangle \text{ is a model of } F \text{ and } \mathbf{\Omega}_{\mathbf{A}}F = Id \}$$

The logic L is algebraizable, then $Alg^*(L)$ coincides with its equivalent algebraic semantics. This is the case of Rasiowa's *implicative models* treated in [40], see also [24, Lemma 2.6].

The theory to associate generalized matrix model with a class of algebra with an arbitrary logic can be found in [25]. Indeed, if $\langle A, \mathcal{C} \rangle$ is a generalized matrix, then its Tarski congruence $\widetilde{\Omega}_{\mathbf{A}}F$ is the largest congruence of \mathbf{A} that is compatible with all members of \mathcal{C} ; it is not hard to show that:

$$\widetilde{\mathbf{\Omega}}_{\mathbf{A}}F = \bigcap_{F \in \mathcal{C}} \mathbf{\Omega}_{\mathbf{A}}F$$

A generalized matrix is *reduced* when Tarski congruence is the identity and the class of *L*-algebras Alg(L) is defined as the class of reducts of the reduced generalized matrix if *L*, i.e.:

 $Alg(L) := \{ \mathbf{A} : \exists \mathcal{C} \subseteq P(A), \langle A, \mathcal{C} \rangle \text{ is a generalized model of } F \text{ and } \widetilde{\mathbf{\Omega}}_{\mathbf{A}} \mathcal{C} = Id \}$

It is well-known that Alg(L) is the class of subdirect products of algebras in $Alg^*(L)$, and in general $Alg(L) \subseteq Alg^*(L)$ holds, see [24, Theorem 5.70]. One of the main aims of this section is to see that $Alg(\mathbb{L}_k^{\leq}) = \mathbb{C}_k$. To this end, from Proposition 6.1 we can infer that \models_k^{\leq} is a *semilattice based* with respect to \mathbb{C}_k in the Jansana's sense [30, p. 76].

Proposition 6.8 $Alg(\mathbb{L}_{k}^{\leq}) = \mathbb{C}_{k}$.

Proof. From Proposition 3.1 [30], we have that $Alg(\mathbb{L}_k^{\leq})$ is the intrinsic variety of L which is characterized by the equations $\psi \approx \psi$. Now, from Proposition 6.1 we have that $\psi \approx \psi$ iff $\phi \models_k^{\leq} \psi$ and $\psi \models_k^{\leq} \phi$ which implies the identities that characterized \mathbb{C}_k are exactly the same that the one of $Alg(\mathbb{L}_k^{\leq})$.

Recall that logic L is said to be *selfextensional* if L satisfies the following weak form of the replacement property: For any $\alpha, \beta, \psi(x, \vec{y}) \in Fm$,

if
$$\alpha \dashv \vdash_L \beta$$
 then $\psi(\alpha, \vec{y}) \dashv \vdash_L \psi(\beta, \vec{y})$.

Corollary 6.9 \mathbb{L}_k^{\leq} is selfextensional.

Proof. It is an immediate consequence from [30, Theorem 3.2] and Proposition 6.8. \Box

Now, we start with the study of the classification of \mathbb{L}_{k}^{\leq} in the Leibniz hierarchy. This hierarchy is characterized in several ways, although the majority concerns several properties of Leibniz operator given by:

$$F \mapsto \mathbf{\Omega}_{\mathbf{A}} F$$

A logic L is algebraizable ([4]) when for every matrix $\langle \mathbf{A}, F \rangle$ which is model of L for every algebra \mathbf{A} belonging the quasivariety K, the Leibniz operator $\Omega_{\mathbf{A}}F$ is an isomorphism between the lattice $\mathcal{F}_i(\mathbf{A})$ and CoA. Besides, it is possible to see that $Alg^*(L) = K$.

A logic L is a *protoalgebraic* one when Leibniz operator is monotonic; i.e., if $G \subseteq F$ then $\Omega_{\mathbf{A}}G \subseteq \Omega_{\mathbf{A}}F$. It is clear that every algebraizable logic is protoalgebraic. An intermediate class of logics between them is of the finitely equivalential ones. Recall that a logic L is said to be *equivalential* if there is a set of formulas built-up in two variables p and q, E(p,q) such that the following conditions hold:

- (E1) $\vdash_L \delta(p, p)$ for every $\delta(p, q) \in E(p, q)$,
- (E2) $E(p,q) \cup \{p\} \vdash_L q$,
- (E3) for every *n*-ary operation # and two *n*-tuples of variables $q_1, \dots, q_n, p_1, \dots, p_n$, we have: $E(p_1, q_1) \cup \dots E(p_n, q_n) \vdash_L E(\#(p_1, \dots, p_n), \#(q_1, \dots, q_n)).$

We say that L is *finitely equivalential* if the set E(p,q) is finite. To see that the logic \mathbb{L}_{k}^{\leq} is finitely equivalential, we will consider the following implication:

$$x \Rightarrow y := (x \to y) \land (\sim y \to \sim x)$$

where $x \to y := \sim ((\sim (x \lor y))^* \land (x \lor y)) \lor ((\sim (x \land y))^* \land (x \land y))$. The following Proposition presents properties of this implication to be used in the rest of the paper.

Proposition 6.10 For a given C_k -algebras (A, t) and any $x, y, z, w \in A$, the following holds:

- (i) $x \Rightarrow y = y \Rightarrow x$, (ii) $x \Rightarrow y = \sim x \Rightarrow \sim y$, (iii) $x \Rightarrow x = 1$, (iv) $(x \Rightarrow y) \land (z \Rightarrow w) \le (x \lor z) \Rightarrow (y \lor w)$, (vii) $x \Rightarrow y = x^* \Rightarrow y^*$, (viii) $(x \Rightarrow y) \land x = (x \Rightarrow y) \land y$,
- (iv) $(x \Rightarrow y) \land (y \Rightarrow z) \le (x \Rightarrow z),$ (Viii) $(x \Rightarrow y) \land x = (x \Rightarrow y) \land y$
- (v) $(x \Rightarrow y) \land (z \Rightarrow w) \le (x \land z) \Rightarrow (y \land w)$, (ix) $(1 \Rightarrow x) = 1$ iff x = 1.

Proof. Let us consider C_k -algebra (A, t). Then, it is clear that A is an mpM-algebra, see Section 1. Since the variety of mpM-algebra is generated by T_3 and T_4 , every identity that is verified in the generating algebras holds in every algebra of the class. The identities expressed in (i) to (viii) are in the language of the mpM-algebras and so it is easy to see that using the corresponding table, these identities hold in (A, t).

Now, let us suppose that $(1 \Rightarrow x) = 1$. Then, $(1 \Rightarrow x) = (1 \rightarrow x) \land (\sim x \rightarrow \sim 1) = \sim ((\sim 1)^* \land 1) \lor ((\sim x)^* \land x) \land \sim ((\sim (\sim x)^* \land (\sim x)) \lor ((\sim 0)^* \land 0) = ((\sim x)^* \land x) \land ((\sim x)^* \lor x)) = 1$. The latter identity is equivalent to $((\sim x)^* \land x) = 1$ and $((\sim x)^* \lor x)) = 1$ iff $(\sim x)^* = 1$ and x = 1, which completes the proof.

Definition 6.11 Let $\delta(p,q)$ be the following formula of \mathcal{L}_{Σ} :

$$\delta(p,q) = (p \Rightarrow q) \land (tp \Rightarrow tq) \land (t^2p \Rightarrow t^2q) \land \dots \land (t^{k-1}p \Rightarrow t^{k-1}q).$$

The equivalence sentences system for \mathbb{L}_k^{\leq} is $E(p,q) = \{\delta(p,q)\}.$

Theorem 6.12 The logic \mathbb{L}_k^{\leq} is protoalgebraic and finitely equivalential, but not algebraizable.

Proof. The fact that \models_k^{\leq} satisfies (E1) is a consequence of (iii) from Proposition 6.10. From the definition of \models_k^{\leq} and (viii) of Proposition 6.10, we have that (E2) is held. Now, the fact that \models_k^{\leq} verifies (E3) follows from (ii), (v), (vi) and (vii) of Proposition 6.10. Since the logic is finitely equivalential, then it is protoalgebraic. To see that \mathbb{L}_k^{\leq} is not algebraizable is enough to take the simple algebra (T_4, Id) where T_4 is the simple mpM-algebra and *id* is the identity function. It is not difficult to see that if we take the filters $\uparrow a = \{a, 1\}$ and $\uparrow b = \{b, 1\}$, then $\Omega_A \uparrow a = \Omega_A \uparrow b = A \times A$, then the Leibniz operator is not 1-1, by which the proof is completed.

From the proof of the last Theorem, it is clear that the logic \mathbb{L}_k^{\leq} is not weak algebraizable because the Leibniz operator is not monotonic.

Corollary 6.13 $Alg^*(\mathbb{L}_k^{\leq}) = \mathbb{C}_k$.

Proof. From Theorem 6.12 and [25, Proposition 3.2], we have that $Alg^*(\mathbb{L}_k^{\leq}) = Alg(\mathbb{L}_k^{\leq})$. From the latter and Proposition 6.8, we have completed the proof.

7 Another sentential logic associated with the class C_k algebras

In this section, we will present another sentential logic that has an algebraic counterpart the class C_k -algebras. To this end, let us consider $\mathfrak{Fm} = \langle Fm, \wedge, \vee, \sim, *, t, \top, \bot \rangle$ the absolutely free algebra of formulas of type (2, 2, 1, 1, 1, 0) generated by some denumerable set of variables. Now we will say that $\mathbb{L}_k = \langle Fm, \models_k \rangle$ is a sentential logic which is defined as follows for every $\Gamma \cup \{\alpha\} \subseteq Fm$:

 $\Gamma \models_k \alpha \quad \Longleftrightarrow \quad \forall (A,t) \in \mathbb{C}_k, \, \forall v \in Hom_{\mathbb{C}_k}(Fm, (A,t))$ if $v(\gamma) = 1$ for every $\gamma \in \Gamma$ then $v(\alpha) = 1$

In the area of algebraic logic, these kind of logics defined as above are so-called 1assertional ones. When the logic is built over a lattice, it is generally finitary. Indeed:

Proposition 7.1 The logic \mathbb{L}_k is finitary.

Proof. It immediately follows from the fact that \mathbb{C}_k is a variety and the notion of filter for every algebra is elementary definable and then the class \mathcal{C}_k -algebras is closed under ultraproducts.

Proposition 7.2 The logics \mathbb{L}_k and \mathbb{L}_k^{\leq} have the same theorems.

Proof. It immediately follows from the very definitions of \models_k and \models_k^{\leq} .

Lemma 7.3 The logic \mathbb{L}_k is paracomplete, but not paraconsistent.

Proof. Taking the simple \mathcal{C}_k -algebra (T_4, id) and v(p) = a and $v(\sim p) = a$, then we have that $\not\models_k p \lor \sim p$. Now, from the fact that the negation \sim is involutive in every \mathcal{C}_k -algebra, we have that $p, \sim p \models_k q$ holds for every variable p and q.

Lemma 7.4 The logic \mathbb{L}_k is a LFU with respect to ~ and \circ (defined in the above section).

Proof. It is an immediately consequence from Lemmas 7.3 and 6.5.

On the other hand, Blok and Pigozzi developed the theory of algebraizable logics in [4] using a generalization of the usual Lindenbaum-Tarski process. To present a characterization of algebraizable, we will consider the propositional signature, Θ , and let \mathscr{L} be a standard propositional logic defined over the language \mathcal{L}_{Θ} , with a consequence relation $\vdash_{\mathcal{L}}$. Then \mathscr{L} is algebraizable in the sense of Blok and Pigozzi if there exists a nonempty set $\Delta(p_1, p_2) \subseteq \mathcal{L}_{\Theta}$ of formulas depending on variables p_1 and p_2 , and a nonempty set $E(p_1) \subseteq \mathcal{L}_{\Theta} \times \mathcal{L}_{\Theta}$ of pairs of formulas depending on variable p_1 satisfying the following properties:

(C1) $\vdash_{\mathscr{L}} \delta(p_1, p_1)$, for every $\delta(p_1, p_2) \in \Delta(p_1, p_2)$;

- (C2) $\Delta(p_1, p_2) \vdash_{\mathscr{L}} \delta(p_2, p_1)$, for every $\delta(p_1, p_2) \in \Delta(p_1, p_2)$;
- (C3) $\Delta(p_1, p_2), \Delta(p_2, p_3) \vdash_{\mathscr{L}} \delta(p_1, p_3)$, for every $\delta(p_1, p_2) \in \Delta(p_1, p_2)$;
- (C4) $\Delta(p_1, p_{n+1}), \dots, \Delta(p_n, p_{2n}) \vdash_{\mathscr{L}} \delta(\#(p_1, \dots, p_n), \#(p_{n+1}, \dots, p_{2n}))$, for every $\delta(p_1, p_2) \in$ $\Delta(p_1, p_2)$, every n-ary connective # of Θ and every $n \ge 1$;
- (C5) $p_1 \vdash_{\mathscr{L}} \delta(\gamma(p_1), \epsilon(p_1))$, for every $\delta(p_1, p_2) \in \Delta(p_1, p_2)$ and every $\langle \gamma(p_1), \epsilon(p_1) \rangle \in E(p_1);$

(C6) $\{\delta(\gamma(p_1), \epsilon(p_1)) : \delta(p_1, p_2) \in \Delta(p_1, p_2), \langle \gamma(p_1), \epsilon(p_1) \rangle \in E(p_1)\} \vdash \varphi p_1.$

The sets $\Delta(p_1, p_2)$ and $E(p_1)$ are called system of equivalence formulas and defining equations, respectively. In order to see that the logic \mathbb{L}_k is algebraizable, we will consider the following two binary operations defined on the above section:

$$x \to y := \sim ((\sim (x \lor y))^* \land (x \lor y)) \lor ((\sim (x \land y))^* \land (x \land y))$$

$$x \leftrightarrow y := (x \to y) \land (\sim y \to \sim x)$$

Now, let us take formula $\delta(p_1, p_2)$ defined as follows:

$$\delta(p_1, p_2) = (p_1 \leftrightarrow p_2) \land (tp_1 \leftrightarrow tp_2) \land (t^2 p_1 \leftrightarrow t^2 p_2) \land \dots \land (t^{k-1} p_1 \leftrightarrow t^{k-1} p_2)$$

The system of equivalence formulas is given by $\Delta(p_1, p_2) = \{\delta(p_1, p_2)\}$ and the system of defining equations is given by $E(p_1) = \{p_1, p_1 \leftrightarrow p_1\}.$

Theorem 7.5 The logic \mathbb{L}_k is algebraizable in the sense of Blok-Pigozzi with the system of equivalence formulas $\Delta(p_1, p_2)$ and the system of defining equations $E(p_1) = \{p_1, p_1 \leftrightarrow p_1\}$.

Proof. We have to prove that \models_k satisfies (C1) to (C6). The fact that (C1) and (C2) hold follows immediately from Proposition 6.10 (iii) and (i), respectively. (C3): Let us suppose that for every $v \in Hom_{\mathbb{C}_k}(Fm, (A, t))$ we have:

$$v((p_1 \leftrightarrow p_2) \land (tp_1 \leftrightarrow tp_2) \land (t^2p_1 \leftrightarrow t^2p_2) \land \dots \land (t^{k-1}p_1 \leftrightarrow t^{k-1}p_2) = 1$$

and

$$v((p_2 \leftrightarrow p_3) \land (tp_2 \leftrightarrow tp_3) \land (t^2p_2 \leftrightarrow t^2p_3) \land \dots \land (t^{k-1}p_2 \leftrightarrow t^{k-1}p_3) = 1$$

Then, it is clear that $v(p_1 \leftrightarrow p_2) = 1$ and $v(p_2 \leftrightarrow p_3) = 1$; moreover, $v(t^i p_1 \leftrightarrow t^i p_2) = 1$ and $v(t^i p_2 \leftrightarrow t^i p_3) = 1$ for any $i = 1, \dots, k-1$. From the latter and Proposition 6.10 (iv), $v(p_1 \leftrightarrow p_3) = 1$ and $v(t^i p_1 \leftrightarrow t^i p_3) = 1$ for any $i = 1, \dots, k-1$.

(C4): Suppose for any $v \in Hom_{\mathbb{C}_k}(Fm, (A, t))$ and the following identities $v(\delta(p_1, p_2)) = 1$ and $v(\delta(p_3, p_4)) = 1$ hold. So, $v(t^i p_1 \leftrightarrow t^i p_2) = 1$ and $v(t^i p_3 \leftrightarrow t^i p_4) = 1$ for any $i = 1, \dots, k$. Now from the items (v) and (iv) of Proposition 6.10, we have that $v(t^i p_1 \# t^i p_3 \leftrightarrow t^i p_2 \# t^i p_4) = 1$ and so $v(\delta(p_1 \# p_3, p_2 \# p_4)) = 1$ where $\# \in \{\wedge, \vee\}$. Now it is not hard to see that if $v(\delta(p_1, p_2)) = 1$, then $v(\delta(\# p_1, \# p_2)) = 1$ with $\# \in \{^*, \sim, t\}$.

(C5): Suppose for any $v \in Hom_{\mathbb{C}_k}(Fm, (A, t))$ we have $v(p_1) = 1$, then it is not hard to see that $v(\delta(p_1, p_1)) = 1$ and $v(\delta(p_1, p_1 \leftrightarrow p_1)) = 1$.

(C6): Suppose for any $v \in Hom_{\mathbb{C}_k}(Fm, (A, t))$, we have $v(\delta(p_1, p_1)) = 1$ and $v(\delta(p_1, p_1 \leftrightarrow p_1)) = 1$. So, $v(p_1 \leftrightarrow p_1) = 1$ and then we have $1 \leftrightarrow v(p_1) = 1$. From Proposition 6.10 (viii), we have that $v(p_1) = 1$ as desired.

As consequence of the last Theorem and some general results of the Blok-Pigozzi theory, we have the following:

Corollary 7.6 $Alg^*(\mathbb{L}_k) = \mathbb{C}_k$.

Proof. See section 3 and Proposition 3.15 of [24].

It is important to note that \mathbb{L}_k is not Rasiowa's implicative logic using $\delta(p_1, p_2)$ as implication, see [24, Definition 2.5] because it is not verified: if $v((p_1 \leftrightarrow p_2) \land (tp_1 \leftrightarrow tp_2) \land (t^2p_1 \leftrightarrow t^2p_2) \land \cdots \land (t^{k-1}p_1 \leftrightarrow t^{k-1}p_2) = 1$, then $v(p_1) = v(p_2)$ for every \mathcal{C}_k -algebra.

8 Conclusions

In this paper, we have studied the class of k-cyclic modal pseudocomplemented De Morgan algebras proving that it is a semsisimple variety and determining the generating algebras for every positive integer k. From the algebraic studies, we have displayed a family of

sentential logics \mathbb{L}_k^{\leq} and \mathbb{L}_k that have as algebraic counterpart the class of algebras introduced previously, this fact has been settled in Corollaries 6.13 and 7.6. As the main results that emerge from our studies, it has been shown that although the logics \mathbb{L}_k^{\leq} and \mathbb{L}_k share the same theorems, \mathbb{L}_k^{\leq} is a paraconsistent non-algebraizable logic; in contrast, \mathbb{L}_k is a non-paraconsistent algebraizable logic.

References

- R. Balbes and P. Dwinger. *Distributive lattices*. University of Missouri Press, Columbia, (1974).
- [2] J.-Y Béziau, Two Genuine 3-Valued Paraconsistent Logics, pages 35–47. Springer International Publishing, Cham, 2016.
- [3] F. Bou, F. Esteva, J. M. Font, A. Gil, L. Godo, A. Torrens and V. Verdú. Logics preserving degrees of truth from varieties of residuated lattices. J. Log. Comput., 19(6):1031–1069, 2009.
- [4] W. J. Blok and D. Pigozzi. Algebraizable Logics. Memoirs of the American Mathematical Society, vol. 77, no. 396, 1989.
- [5] S. Burris and H. Sankappanavar. A course in Universal Algebra. Graduate Texts in Mathematics 78, Springer-Verlag, Berlin, (1981).
- [6] L. Cantú and M. Figallo. On the logic that preserves degrees of truth associated to involutive Stone algebras. Logic Journal of the IGPL, 28: 5, 1000–1020, 2020.
- [7] J. Castiglioni and R. Ertola-Biraben. Strict paraconsistency of truth-degree preserving intuitionistic logic with dual negation. Logic Journal of the IGPL, 22, 268–273, 2014.
- [8] W. A. Carnielli and J. Marcos. A taxonomy of C-systems. In Paraconsistency The Logical Way to the Inconsistent, vol. 228 of Lecture Notes in Pure and Applied Mathematics, pp. 1–94. Marcel Dekker, 2002.
- [9] W. Carnielli and A. Rodrigues. An epistemic approach to paraconsistency: a logic of evidence and truth. Synthese 196, 3789-3813, 2019.
- [10] R. Cignoli and M. S. de Gallego. Dualities for some De Morgan algebras with operators and Lukasiewicz algebras. J. Austral Math. Soc (Series A), 34 (1983), 377–393.
- J. Czelakowski and R. Jansana. Weakly algebraizable logics. The Journal of Symbolic Logic, 65, 641–668, 2000
- [12] J. Díaz Varela and B. López Martinolich. Resolution of Algebraic Systems of Equations in the Variety of Cyclic Post Algebras, Stud. Logica 98, 307–330 (2011).

- [13] R. Ertola, F. Esteva, T. Flaminio, L. Godo, and C. Noguera. Paraconsistency properties in degree-preserving fuzzy logics, Soft Comput. 19 (3), 531–546, 2015.
- [14] F. Esteva, A. Figallo-Orellano, T. Flaminio and L. Godo. Logics of Formal Inconsistency Based on Distributive Involutive Residuated Lattices, Journal of Logic and Computation, 2021
- [15] A. Figallo-Orellano, A. Ziliani and M. Figallo. Symmetric operators on modal pseudocomplemented De Morgan algebras. Logic journal of the IGPL, 25:4, 496–511 (2017)
- [16] A. Figallo-Orellano and I. Pascual. On monadic operators over modal pseudocomplemented De Morgan algebras and tetravalent modal algebras. Stud Logica (2019) 107: 591–611.
- [17] A. V. Figallo. Tópicos sobre álgebras modales 4-valuadas. Proceeding of the IX Simposio Latino-Americano de Lógica Matemática, (Bahía Blanca, Argentina, 1992). Notas de Lógica Matemática, Vol. 39, (1992), pp. 145–157.
- [18] A. V. Figallo, N. Oliva and A. Ziliani. Modal Pseudocomplemented De Morgan Algebras. Acta Univ. Palacki. Olomuc., Fac. rer. nat., Mathematica 53, 1 65–79, 2014
- [19] A. V. Figallo, N. Oliva and A. Ziliani. Free Modal Pseudocomplemented De Morgan Algebras. Bulletin of the Section of Logic, 47(2), 89–106, 2018
- [20] A. V. Figallo and A. Ziliani. Symmetric tetra-valued modal algebras. Notas de la Sociedad de Matemdtica de Chile, vol. 10 (1991), no. 1, pp. 133-141.
- [21] A. V. Figallo and P. Landini. On generalized I-algebras and 4-valued modal algebras. Reports on Mathematical Logic 29 (1995), 3–18.
- [22] M. Figallo. Hypersequents and Tetravalent Modal Logic TML. Ph.D. Thesis, Universidad Nacional del Sur, 2013.
- [23] M. Figallo. Cut-free sequent calculus and natural deduction for the tetravalent modal logic. Stud. Logica, 2021.
- [24] J. M. Font. Abstract Algebraic Logic An Introductory Textbook. vol. 60 of Studies in Logic - Mathematical Logic and Foundations College Publications, London, 2016. 546 pp
- [25] J. M. Font and R. Jansana. A General Algebraic Semantics for Sentential Logics. Second revised edition, Vol. 7 of Lecture Notes in Logic. Association for Symbolic Logic, 2009. First edition 1996.
- [26] J. M. Font and M. Rius. An abstract algebraic logic approach to tetravalent modal logics. J. Symbolic Logic, Vol. 65, (2000), pp. 481–518.

- [27] J. M. Font. On substructural logics preserving degrees of truth. Bulletin of the Section of Logic, 36, 117–130, 2007.
- [28] J. M. Font. Taking degrees of truth seriously. Studia Logica (Special issue on Truth Values, Part I), 91, 383–406, 2009.
- [29] J. M. Font, A. Gil, A. Torrens, and V. Verdú. On the infinite-valued Lukasiewicz logic that preserves degrees of truth. Archive for Mathematical Logic, 45, 839–868, 2006.
- [30] R. Jansana. Selfextensional logics with a conjunction. Studia Logica, 84, 63–104, 2006.
- [31] J. Marcos. Nearly every normal modal logic is paranormal. Logique et Analyse 48, 279-300, 2005.
- [32] B. F. López-Martinolich. Gröbner Bases over Cyclic Post Algebras. 2012 IEEE 42nd International Symposium on Multiple-Valued Logic, Victoria, BC, Canada, 2012, pp. 197-202
- [33] B. F. López-Martinolich. Resolución de sistemas de ecuaciones polinomiales sobre álgebras de post k-cíclicas, Ph. D. Thesis, Universidad Nacional del Sur, 2011.
- [34] A. Monteiro. Sur les algèbres de Heyting Symétriques, Portugaliae Mathematica, 39 (1980), 1–237.
- [35] S. Marcelino and U. Rivieccio. Logics of Involutive Stone Algebras. arXiv:2102.05455 [math.LO] 2021.
- [36] M. Osorio, A. Figallo-Orellano and M. Pérez-Gaspar. A family of genuine and nonalgebraisable C-systems. Journal of Applied Non-Classical Logics, 31:1, 56–84, 2021
- [37] I. Loureiro. Prime spectrum of a tetravalent modal algebra. Notre Dame Journal of Formal Logic, 24:389–394, 1983.
- [38] H. Sankappanavar. Pseudocomplemented Ockham and De Morgan algebras. Zeitschr.
 f. Math. Logik und Grundlagen Math., 32, 385–394, 1986
- [39] D. Szmuc. Defining LFIs and LFUs in extensions of infectious logics. Journal of Applied Non-Classical Logics, 26:4, 286–314, 2016
- [40] H. Rasiowa. An algebraic approach to non-clasical logics. Studies in logic and the foundations of mathematics, vol. 78. North-Holland Publishing Company, Amsterdam and London, and American Elsevier Publishing Company, Inc., New York, 1974.

A. Figallo-Orellano Departamento de Matemática Universidad Nacional del Sur (UNS) Bahía Blanca, Argentina e-mail: *aldofigallo@gmail.com*

M. Pérez-Gaspar Facultad de Ingenirería Universidad Nacional Autónoma de México (UNAM) Ciudad de México, México e-mail: *miguel.perez@fi-b.unam.mx*

J. M. Ramírez-Contreras Departamento de Actuaría, Física y Matemáticas Universidad de las Américas Puebla (UDLAP) Puebla, México e-mail: *juan.ramirez@udlap.mx*