BIASED MULTILINEAR MAPS OF ABELIAN GROUPS

SEAN EBERHARD

ABSTRACT. We adapt the theory of partition rank and analytic rank to the category of abelian groups. If A_1, \ldots, A_k are finite abelian groups and $\varphi : A_1 \times \cdots \times A_k \to \mathbf{T}$ is a multilinear map, where $\mathbf{T} = \mathbf{R}/\mathbf{Z}$, the bias of φ is defined to be the average value of $\exp(i2\pi\varphi)$. If the bias of φ is bounded away from zero we show that φ is the sum of boundedly many multilinear maps each of which factors through the standard multiplication map of $\mathbf{Z}/q\mathbf{Z}$ for some bounded prime power q. Relatedly, if $F : A_1 \times \cdots \times A_{k-1} \to B$ is a multilinear map such that $\mathbf{P}(F = 0)$ is bounded away from zero, we show that F is the sum of boundedly many multilinear functions of a particular form. These structure theorems generalize work of several authors in the elementary abelian case to the arbitrary abelian case. The set of all possible biases is also investigated.

CONTENTS

1.	Introduction	1
2.	Basic properties of bias	3
3.	The structure theorem	6
4.	Possible biases	9
References		12

1. INTRODUCTION

Suppose A_1, \ldots, A_k are finite abelian groups and $\varphi : A_1 \times \cdots \times A_k \to \mathbf{T}$ is a multilinear map, where $\mathbf{T} = \mathbf{R}/\mathbf{Z}$. Let $e(x) = \exp(i2\pi x)$ be the standard character of \mathbf{T} . The *bias* of φ is defined by

(1)
$$\operatorname{bias}(\varphi) = \mathbf{E}_{x \in A_{[k]}} e(\varphi(x))$$

Here and throughout we use the following index notation. The symbol [k] denotes the index set $\{1, \ldots, k\}$. For $I \subseteq [k]$,

$$A_I = \prod_{i \in I} A_i.$$

For $x \in A_{[k]}$,

$$x_I = (x_i)_{i \in I} \in A_I.$$

We write I^c for $[k] \setminus I$. Also, we are using the expectation symbol to denote the (finite) average over the set $A_{[k]}$.

The concept of bias often comes up in the following way. Suppose $F: A_{[k-1]} \to B$ is a multilinear map of abelian groups such that $\mathbf{P}(F=0)$ is bounded away from zero. Letting A_k be the dual group \widehat{B} , we can define a multilinear map $\varphi: A_{[k]} \to \mathbf{T}$ by

$$\varphi(x) = x_k(F(x_{\lceil k-1 \rceil})),$$

S. Eberhard has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 803711).

SEAN EBERHARD

and for this map we have $bias(\varphi) = \mathbf{P}(F = 0)$. Moreover, F is determined by φ , so if we can say something about the structure of φ then we will know something about the structure of F. Thus the study of multilinear kernels is reduced to the study of bias.

If A_1, \ldots, A_k are vector spaces over a field $\mathbf{F} = \mathbf{F}_q$ then it is usually more natural to consider a multi-**F**-linear map $\varphi : A_{[k]} \to \mathbf{F}$ and to define

$$\operatorname{bias}(\varphi) = \mathbf{E}_{x \in A_{[k]}} \chi(\varphi(x))$$

where $\chi : \mathbf{F}_q \to \mathbf{T}$ is any standardized character such as $\chi(x) = e((\operatorname{tr} x)/p)$, where $\operatorname{tr} : \mathbf{F}_q \to \mathbf{F}_p$ is the absolute trace. Bias in this context was introduced by Gowers and Wolf in [GW] and studied extensively by several authors. The term *analytic* rank is used for the quantity

$$\operatorname{AR}(\varphi) = \log_a(\operatorname{bias}(\varphi)^{-1}).$$

The *partition rank* $PR(\varphi)$ of φ is the minimal number r such that φ is the sum of r multilinear maps of the form

 $\varphi_1(x_I)\varphi_2(x_{I^c}),$

where $I \subseteq [k]$ and 0 < |I| < k. Multilinear maps of bounded partition rank are uniformly biased: in fact, $AR(\varphi) \leq PR(\varphi)$. This is one of several nice lemmas appearing in a paper by Lovett [L]. The main theorem in this field is a converse: uniformly biased multilinear maps have bounded partition rank, or in other words partition rank is bounded in terms of analytic rank.

Theorem 1.1. Suppose A_1, \ldots, A_k are finite \mathbf{F}_q -vector spaces and $\varphi : A_{[k]} \to \mathbf{F}_q$ is multilinear over \mathbf{F}_q . If $\operatorname{bias}(\varphi) \ge \epsilon > 0$ then φ has partition rank $O_{\epsilon,k,q}(1)$. In other words, $\operatorname{PR}(\varphi) \le F(k, \operatorname{AR}(\varphi), q)$ for some function F.

This result was essentially proved by Green and Tao [GT]. Subsequent research has focused on quantitative aspects (which were initially poor). The dependence on q was removed by Bhowmick and Lovett in [BL] (so $\operatorname{PR}(\varphi) \leq F(k, \operatorname{AR}(\varphi))$). In breakthrough work Janzer [J] and Milićević [M] independently proved that $\operatorname{PR}(\varphi) \leq C_k(\operatorname{AR}(\varphi)^{D_k} + 1)$ for constants C_k, D_k . Recently, Cohen and Moshkovitz [CM] proved that $\operatorname{PR}(\varphi) \leq (2^{k-1} + 1) \operatorname{AR}(\varphi) + 1$ provided that $q > q_0(k, \operatorname{AR}(\varphi))$.

In this note we consider biased multilinear maps of arbitrary abelian groups. We will use Theorem 1.1 as a block box (and only the prime field case) and we will deduce an analogous structure theorem for arbitrary abelian groups. To state this structure theorem we need a suitable notion of partition rank. For q a prime power, let $m_q : (\mathbf{Z}/q\mathbf{Z})^2 \to \mathbf{T}$ be the bilinear map defined by $m(x, y) = xy/q \mod 1$. We say that $\varphi : A_{[k]} \to \mathbf{T}$ factors through m_q if it has the form

$$\varphi(x) = m_q(\varphi_1(x_I), \varphi_2(x_{I^c}))$$

for some $I \subseteq [k]$ with 0 < |I| < k. Here $\varphi_1 : A_I \to \mathbf{Z}/q\mathbf{Z}$ and $\varphi_2 : A_{I^c} \to \mathbf{Z}/q\mathbf{Z}$ must both be multilinear.

Theorem 1.2. Suppose A_1, \ldots, A_k are finite abelian groups and $\varphi : A_{[k]} \to \mathbf{T}$ is multilinear. If $\operatorname{bias}(\varphi) \ge \epsilon > 0$ then φ is the sum of $O_{\epsilon,k}(1)$ multilinear maps each of which factors through m_q for some prime power $q \le O_{\epsilon,k}(1)$.

It is easy to see that Theorem 1.1 and Theorem 1.2 agree in the case of vector spaces over a prime finite field, but neither result is more general than the other. (A common generalization would consider *R*-modules for some commutative ring R; Theorem 1.1 would be the case $R = \mathbf{F}_q$ and Theorem 1.2 would be the case $R = \mathbf{Z}$.)

The following corollary follows from the connection mentioned already between bias and multilinear kernels. It states that a multilinear function F with $\mathbf{P}(F=0)$ bounded away from zero must be the sum of boundedly many functions each of which "crushes" a nontrivial subset of the variables before mapping to the range. Here cod(g) denotes the codomain of the function g.

Corollary 1.3. Suppose A_1, \ldots, A_{k-1}, B are finite abelian groups and $F : A_{[k-1]} \rightarrow B$ is a multilinear map such that $\mathbf{P}(F = 0) \ge \epsilon > 0$. Then there is an expression

$$F(x) = \sum_{\emptyset \neq I \subseteq [k-1]} G_I(g_I(x_I), x_{[k-1] \setminus I}),$$

where for each I the functions g_I and G_I are multilinear maps

$$g_I : A_I \to \operatorname{cod}(g_I),$$

$$G_I : \operatorname{cod}(g_I) \times A_{[k-1] \setminus I} \to B$$

and $|\operatorname{cod}(g_I)| \leq O_{\epsilon,k}(1).$

In the last section we give an application of Theorem 1.2 to the set all possible biases. Let Φ_k be the set of all multilinear maps $\varphi : A_1 \times \cdots \times A_k \to \mathbf{T}$ (for any finite abelian groups A_1, \ldots, A_k) and let

$$B_k = \{ \operatorname{bias}(\varphi) : \varphi \in \Phi_k \}.$$

We show that B_k is a small subset of [0, 1] in various senses; for example, all its limit points are algebraic.

Elsewhere we will give an application to probabilistically nilpotent finite groups.

2. Basic properties of bias

Several nice lemmas about bias over vector spaces were proved by Lovett in [L]. We need analogues in the category of abelian groups.

In this section we will sometimes consider functions $\varphi : A_{[k]} \to \mathbf{T}$ which are not multilinear, but we still define $\operatorname{bias}(\varphi)$ by (1). If $\varphi : A_{[k]} \to \mathbf{T}$ is a map we write $\varphi_{x_I} : A_{I^c} \to \mathbf{T}$ for the map obtained by restriction:

$$\varphi_{x_I}(x_{I^c}) = \varphi(x).$$

Obviously,

(2)
$$\operatorname{bias}(\varphi) = \mathbf{E}_{x_I} \operatorname{bias}(\varphi_{x_I}).$$

In particular, consider (2) in the case $I = \{i\}^c$. If $\varphi(x)$ is linear in x_i then $bias(\varphi_{x_I})$ is 1 or 0 according to whether $\varphi_{x_I} \equiv 0$, so

(3)
$$\operatorname{bias}(\varphi) = \mathbf{P}_{x_I}(\varphi_{x_I} \equiv 0).$$

This shows that, while in general $bias(\varphi)$ is complex-valued, $bias(\varphi) \in [0, 1]$ provided that φ is linear in at least one of its arguments.

Lemma 2.1. Assume $\varphi : A_{[k]} \to \mathbf{T}$ is multilinear. For each $i \in [k]$,

$$\operatorname{bias}(\varphi) \ge 1 - \prod_{j \neq i} (1 - 1/|A_j|)$$

If φ is nontrivial then

$$\operatorname{bias}(\varphi) \le 1 - \prod_{j \ne i} (1 - 1/p_j),$$

where p_j is the smallest prime divisor of $|A_j|$.

Proof. Let $I = \{i\}^c$. If $x_j = 0$ for any $j \in I$ then $\varphi_{x_I} \equiv 0$, so from (3) we have

$$1 - \operatorname{bias}(\varphi) = \mathbf{P}(\varphi_{x_I} \neq 0) \le \prod_{j \in I} \mathbf{P}(x_j \neq 0) = \prod_{j \neq i} (1 - 1/|A_j|).$$

The second inequality is proved by induction. The case k = 1 is clear, because if φ is linear and nontrivial then $bias(\varphi) = 0$, and the right-hand side is also 0. Let k > 1. By (2) with $I = \{j\}$ we have

$$1 - \operatorname{bias}(\varphi) = \mathbf{E}_{x_i}(1 - \operatorname{bias}(\varphi_{x_i})).$$

Let $B_j = \{x_j \in A_j : \varphi_{x_j} \equiv 0\}$. By induction if $x_j \notin B_j$ then

$$1 - \operatorname{bias}(\varphi_{x_j}) \ge \prod_{j' \neq i, j} (1 - 1/p_{j'})$$

Thus

$$1 - \text{bias}(\varphi) \ge (1 - |B_j|/|A_j|) \prod_{j' \neq i,j} (1 - 1/p_{j'}).$$

But B_j is a subgroup of A_j , and proper since φ is nontrivial, so $|B_j|/|A_j| \le 1/p_j$. This completes the induction.

Lemma 2.2. For $I \subseteq [k]$, let $\varphi_I : A_I \to \mathbf{T}$ be |I|-linear. Let $\varphi : A_{[k]} \to \mathbf{T}$ be the function

$$\varphi(x) = \sum_{I \subseteq [k]} \varphi_I(x_I).$$

Suppose $J \subseteq [k]$ is any subset such that $\varphi_I = 0$ for $I \supseteq J$. Then

 $|\operatorname{bias}(\varphi)| \leq \operatorname{bias}(\varphi_J).$

Proof. Let $i \in [k]$. Write

$$\varphi = \varphi_i + \varphi_{i'},$$

where

$$\varphi_i = \sum_{I \subseteq [k], i \in I} \varphi_I,$$
$$\varphi_{i'} = \sum_{I \subseteq [k], i \notin I} \varphi_I.$$

Note that φ_i is linear in x_i and $\varphi_{i'}$ is independent of x_i , so

$$\operatorname{bias}(\varphi) = \mathbf{E}_{x_{\{i\}^c}} \left(e(\varphi_{i'}) \mathbf{E}_{x_i} e(\varphi_i) \right)$$

Hence by the triangle inequality

$$\operatorname{bias}(\varphi) \leq \mathbf{E}_{x_{\{i\}^c}} |\mathbf{E}_{x_i} e(\varphi_i)|$$

But since φ_i is linear in x_i , $\mathbf{E}_{x_i} e(\varphi_i) \ge 0$, so we may drop the absolute value signs. Hence

$$|\operatorname{bias}(\varphi)| \leq \mathbf{E}_{x_{I\setminus i}} \mathbf{E}_{x_i} e(\varphi_i(x)) = \operatorname{bias}(\varphi_i).$$

Repeat for every $i \in J$.

Lemma 2.3. Let $\varphi, \psi : A_{[k]} \to \mathbf{T}$ be multilinear. Then

 $\operatorname{bias}(\varphi + \psi) \ge \operatorname{bias}(\varphi) \operatorname{bias}(\psi).$

Proof. Let $x, y \in A_{[k]}$ be independent. Then

$$\operatorname{bias}(\varphi)\operatorname{bias}(\psi) = \mathbf{E}_{x,y}e(\varphi(x) + \psi(y)) = \mathbf{E}_{x,y}e(\varphi(x) + \psi(x+y)).$$

We may expand

$$\psi(x+y) = \sum_{I \subseteq [k]} \psi_I(x_I, y_{I^c}),$$

where $\psi_I : A_I \times A_{I^c} \to \mathbf{T}$ is again k-linear. Note that $\psi_{[k]} = \psi(x)$. For each fixed $y \in A_{[k]}$, the previous lemma with J = [k] implies that

$$\mathbf{E}_{x}e(\varphi(x) + \psi(x+y))| \le \mathbf{E}_{x}e(\varphi(x) + \psi(x)) = \operatorname{bias}(\varphi + \psi).$$

Hence the lemma follows from the triangle inequality.

Suppose $\varphi: A_{[k]} \to \mathbf{T}$ is k-linear and $\psi: B_{[l]} \to \mathbf{T}$ is l-linear. We say φ factors through ψ if there is an *l*-partition

$$[k] = I_1 \cup \dots \cup I_l \qquad (I_1, \dots, I_l \neq \emptyset)$$

and for each $j \in [l]$ an $|I_j|$ -linear map

$$\varphi_j: A_{I_j} \to B_j$$

such that φ factors as

$$\varphi = \psi(\varphi_{I_1}, \ldots, \varphi_{I_l}),$$

i.e.,

$$\varphi: A_{[k]} \cong \prod_j A_{I_j} \xrightarrow{(\varphi_{I_j})_j} B_{[l]} \xrightarrow{\psi} \mathbf{T}.$$

Lemma 2.4. Suppose $\varphi : A_{[k]} \to \mathbf{T}$ and $\psi : B_{[l]} \to \mathbf{T}$ are multilinear and φ factors through ψ . Then $\operatorname{bias}(\varphi) \ge \operatorname{bias}(\psi)$.

Proof. Suppose $[k] = I_1 \cup \cdots \cup I_l$ and

$$\varphi = \psi(\varphi_{I_1}, \ldots, \varphi_{I_l}).$$

For $b \in B_{[l]}$, let

$$\varphi_b = \psi(\varphi_{I_1} + b_1, \dots, \varphi_{I_l} + b_l).$$

Clearly

 $\mathbf{E}_b \operatorname{bias}(\varphi_b) = \operatorname{bias}(\psi).$

By Lemma 2.2, for every $b \in B_{[l]}$ we have

 $|\operatorname{bias}(\varphi_b)| \leq \operatorname{bias}(\varphi_0) = \operatorname{bias}(\varphi).$

Hence $\operatorname{bias}(\psi) \leq \operatorname{bias}(\varphi)$.

Corollary 2.5. Suppose $\varphi : A_{[k]} \to \mathbf{T}$ and $\psi : B_{[l]} \to \mathbf{T}$ are multilinear and φ factors through ψ . Then $\operatorname{bias}(\varphi) \geq 1 - \prod_{j \neq i} (1 - 1/|B_j|)$ for each $i \in [l]$.

Proof. Combine the previous lemma with Lemma 2.1.

Recall that m_q denotes the map $\mathbf{Z}/q \times \mathbf{Z}/q\mathbf{Z} \to \mathbf{T}$ defined by $m(x, y) = xy/q \mod \mathbf{Z}$ 1, where q is a prime power.

Corollary 2.6. Suppose $\varphi : A_{[k]} \to \mathbf{T}$ is the sum of maps $\varphi_1, \ldots, \varphi_r$ and φ_i factors through m_{q_i} . Then

$$\operatorname{bias}(\varphi) \ge q_1^{-1} \cdots q_r^{-1}.$$

Proof. Combine the previous corollary with Lemma 2.3.

Corollary 2.7. Suppose $A'_i \leq A_i$ for each *i*. Let $\varphi : A_{[k]} \to \mathbf{T}$ be multilinear and let φ' be the restriction of φ to $A'_{[k]}$. Then $\operatorname{bias}(\varphi') \geq \operatorname{bias}(\varphi)$.

Proof. Since φ' factors through φ , we may apply Lemma 2.4 to (φ', φ) . \square

The following lemma, which gives a little more information than the second part of Lemma 2.1, illustrates the utility of the last corollary.

Lemma 2.8. Suppose $\varphi : A_{[k]} \to \mathbf{T}$ is multilinear and suppose the image of φ has an element of order $q = p^n$ for some prime p. Then $\operatorname{bias}(\varphi) \leq \operatorname{bias}(\psi)$, where $\psi : (\mathbf{Z}/q\mathbf{Z})^k \to \mathbf{T}$ is defined by

$$\psi(x_1,\ldots,x_k) = x_1\cdots x_k/q \pmod{1}$$

In particular, for $k \geq 2$,

$$\operatorname{bias}(\varphi) \le (n+1)^{k-2}/q.$$

Proof. Suppose $\varphi(a_1, \ldots, a_k)$ has order $q = p^n$. Let $A'_i = \langle a_i \rangle$ for each *i*. Then

$$\operatorname{bias}(\varphi) \leq \operatorname{bias}(\varphi|_{A'_{[\nu]}})$$

Hence without loss of generality A_i is generated by a_i for each *i*. Note that $\varphi_{qa_i} = 0$ for each *i*, so φ factors through $\prod_{i=1}^k \langle a_i \rangle / \langle qa_i \rangle$. Hence we may assume that $qa_i = 0$ for each *i*. Also note that $p^{n-1}a_i \neq 0$ for each *i*, since

$$\varphi(\dots, a_{i-1}, p^{n-1}a_i, a_{i+1}, \dots) = p^{n-1}\varphi(a_1, \dots, a_k) \neq 0.$$

Hence each a_i has order q, so φ is equivalent to the map ψ in the statement of the lemma.

For the second statement we may assume $\varphi = \psi$. Let v_p denote the *p*-adic valuation. From (3),

$$bias(\varphi) = \mathbf{P}(x_1 \cdots x_{k-1} = 0)$$

= $\mathbf{P}(v_p(x_1) + \cdots + v_p(x_{k-1}) \ge n)$
 $\le \sum_{v_1 + \cdots + v_{k-1} = n} \mathbf{P}(v_p(x_i) \ge v_i \text{ for each } i < k)$
= $\sum_{v_1 + \cdots + v_{k-1} = n} p^{-v_1 - \cdots - v_{k-1}}$
= M/a ,

where M is the number of solutions to $v_1 + \cdots + v_{k-1} = n$ in nonnegative integers. Clearly $M \leq (n+1)^{k-2}$.

3. The structure theorem

In this section we prove Theorem 1.2. Throughout we consider k to be fixed and dependence on k will not be explicit. We start with some simple reductions.

Suppose $\varphi : A_{[k]} \to \mathbf{T}$ is multilinear and $\operatorname{bias}(\varphi) \ge \epsilon$. We may decompose each A_i into its *p*-primary parts:

$$A_i = \bigoplus_p A_i^{(p)}.$$

Then φ decomposes as

$$\varphi = \sum_{p} \varphi_{p},$$

where φ_p factors as

$$A_{[k]} \to A_1^{(p)} \times \cdots \times A_k^{(p)} \to \mathbf{T}$$

The bias of φ correspondingly decomposes as

$$\operatorname{bias}(\varphi) = \prod_{p} \operatorname{bias}(\varphi_p).$$

By Lemma 2.1, $\operatorname{bias}(\varphi_p) \leq 1 - (1 - 1/p)^{k-1} \leq 1 - 1/2^{k-1}$ whenever φ_p is nontrivial, so since $\operatorname{bias}(\varphi) \geq \epsilon$ it follows that there are at most $\log \epsilon^{-1} / \log(1 - 1/2^{k-1})^{-1}$ primes such that $\varphi_p \neq 0$. Hence it suffices to consider the *p*-primary parts separately, so without loss of generality A_1, \ldots, A_k are *p*-groups.

Let $K_i = \{a_i \in A_i : \varphi_{a_i} \equiv 0\}$. By replacing A_i with A_i/K_i we may assume $K_i = 0$, so φ is nondegenerate in the sense that $\varphi_{a_i} \neq 0$ for each nonzero $a_i \in A_i$. Then if A_i has an element of order $q = p^n$, the image of φ also contains an element of order q, so by Lemma 2.8,


bias
$$(\varphi_p) \le (n+1)^{k-2}/q \le (\log_2 q + 1)^{k-2}/q.$$

This implies that q is bounded in terms of ϵ (and k). Hence we may assume that A_1, \ldots, A_k have bounded exponent (and in particular p is bounded).


We will now deduce Theorem 1.2 from Theorem 1.1 using the following elementary lemma about extending multilinear maps. For any abelian *p*-group, let A[p]denote the subgroup $\{x \in A : px = 0\}$ and let pA denote the subgroup $\{px : x \in A\}$.

Lemma 3.1. Let A_1, \ldots, A_k be finite abelian p-groups and q a power of p.


(1) (domain enlargement) Suppose $\varphi : pA_1 \times A_{[2,k]} \to \mathbf{Z}/q\mathbf{Z}$ is a multilinear map. Assume $\varphi(x) = 0$ whenever $px_i = 0$ for any i > 1. Then φ factors through $pA_1 \times \prod_{i \in [2,k]} A_i/A_i[p]$, and there is a multilinear map $\psi : A_{[k]} \to \mathbf{Z}/(pq)\mathbf{Z}$ extending φ in the sense that the following diagram commutes:



(2) (range enlargement) Suppose $\varphi : A_{[k]} \to \mathbf{Z}/q\mathbf{Z}$ is a multilinear map such that $\varphi(x) = 0$ whenever $px_i = 0$ for any $i \in [k]$. Then φ factors through $\prod_{i \in [k]} A_i/A_i[p]$, and there is a multilinear map $\psi : A_{[k]} \to \mathbf{Z}/(pq)\mathbf{Z}$ such that $\psi \mod q = \varphi$, as in the following commutative diagram:



(3) (extension of rank-one maps) Suppose $\varphi : pA_1 \times A_{[2,k]} \to \mathbf{T}$ is a multilinear map such that $\varphi(x) = 0$ whenever $px_i = 0$ for any i > 1, and factoring through m_q . Then φ extends to a multilinear map $\psi : A_{[k]} \to \mathbf{T}$ factoring through m_{pq} , as in the following commutative diagram:



Proof. By the structure theorem for finite abelian groups, we may write

 $A_i = \langle e_{i1} \rangle \oplus \dots \oplus \langle e_{id_i} \rangle \qquad (1 \le i \le k),$

where each e_{ij} has order some power of p.

(1) Write

$$\varphi_{j_1\cdots j_k} = \varphi(pe_{1j_1}, e_{2j_2}, \dots, e_{kj_k}) \in \mathbf{Z}/q\mathbf{Z}.$$

We may identify $\mathbf{Z}/q\mathbf{Z}$ with $p(\mathbf{Z}/(pq)\mathbf{Z})$. For each j_1, \ldots, j_k , let $\psi_{j_1\cdots j_k} \in \mathbf{Z}/(pq)\mathbf{Z}$ be an arbitrary solution to

$$p\psi_{j_1\cdots j_k} = \varphi_{j_1\cdots j_k}.$$

Suppose i > 1 and e_{ij_i} has order p^n . By the assumption that φ factors through $pA_1 \times \prod_{i \in [2,k]} A_i/A_i[p]$, we must have $p^{n-1}\varphi_{j_1\cdots j_k} = 0$. Hence $p^n\psi_{j_1\cdots j_k} = 0$. Similarly, if e_{1j_1} has order p^n , then

$$p^{n}\psi_{j_{1}\cdots j_{k}} = p^{n-1}\varphi_{j_{1}\cdots j_{k}} = \varphi(p^{n}e_{1j_{1}}, e_{2j_{2}}, \dots, e_{kj_{k}}) = 0.$$

Hence we may define $\psi: A_{[k]} \to \mathbf{Z}/(pq)\mathbf{Z}$ by linearly extending the definition

$$\psi(e_{1j_1},\ldots,e_{kj_k})=\psi_{j_1\cdots j_k}.$$

The maps $\psi|_{pA_1 \times A_{[2,k]}}$ and $p\varphi$ agree on generators, so they are the same. (2) Write

$$\varphi_{j_1\cdots j_k} = \varphi(e_{1j_1}, e_{2j_2}, \dots, e_{kj_k}) \in \mathbf{Z}/q\mathbf{Z}$$

For each j_1, \ldots, j_k , let $\psi_{j_1 \cdots j_k} \in \mathbf{Z}/(pq)\mathbf{Z}$ be an arbitrary solution to

$$\psi_{j_1\cdots j_k} \equiv \varphi_{j_1\cdots j_k} \pmod{q}.$$

Suppose p^n is the order of one of the generators e_{ij_i} . Then $p^{n-1}\varphi_{j_1\cdots j_k} = 0 \pmod{q}$, so $p^n\psi_{j_1\cdots j_k} = 0 \pmod{pq}$. Hence we may define $\psi: A_{[k]} \to \mathbf{Z}/(pq)\mathbf{Z}$ by linearly extending the definition

$$\psi(e_{1j_1},\ldots,e_{kj_k})=\psi_{j_1\cdots j_k}.$$

Then $\psi \mod q$ and φ agree on generators, so they are the same.

(3) By assumption $\varphi = m_q(\varphi_1, \varphi_2)$ where φ_1 is defined on $pA_1 \times A_I$ for some $I \subseteq [2,k]$ and φ_2 is defined on $A_{[2,k]\setminus I}$. By (1), φ_1 may be extended to $\psi_1 : A_1 \times A_I \to \mathbf{Z}/(pq)\mathbf{Z}$ in such a way that $\psi_1|_{pA_1 \times A_I} = p\varphi_1$. By (2), φ_2 may be extended to $\psi_2 : A_{[2,k]\setminus I} \to \mathbf{Z}/(pq)\mathbf{Z}$ in such a way that $\psi_2 \mod q = \varphi_2$. Let $\psi = m_{pq}(\psi_1, \psi_2)$. Then for $x \in pA_1 \times A_{[2,k]}$ we have

$$\psi(x) = \psi_1(x)\psi_2(x)/(pq) \mod 1 = \varphi_1(x)\varphi_2(x)/q \mod 1 = \varphi(x),$$

so ψ extends φ .

We can now prove Theorem 1.2.

Proof of Theorem 1.2. Let the exponent of A_i be p^{n_i} . We will prove the theorem by induction on $n_1 + \cdots + n_k$. If $n_i = 1$ for each *i* then each A_i is elementary abelian, hence a vector space over \mathbf{F}_p . In this case the result follows from Theorem 1.1. Hence assume $n_1 > 1$ without loss of generality. Then pA_1 and A_1/pA_1 both have smaller exponent than A_1 .

The restriction φ_1 of φ to $pA_1 \times A_{[2,k]}$ factors through a map φ'_1 defined on $pA_1 \times \prod_{i \in [2,k]} A_i/A_i[p]$, as in the following diagram:

By Corollary 2.7, $bias(\varphi'_1) = bias(\varphi_1) \ge \epsilon$, so by induction

$$\varphi_1' = \psi_1' + \dots + \psi_r'$$

for $r = O_{\epsilon}(1)$ and some

$$\psi'_1, \dots, \psi'_r : pA_1 \times \prod_{i \in [2,k]} A_i / A_i[p] \to \mathbf{T}$$

where each ψ'_i factors through m_{q_i} for some $q_i \leq O_{\epsilon}(1)$. By Lemma 3.1, each ψ'_i extends to a multilinear map $\psi_i : A_{[k]} \to \mathbf{T}$ factoring through m_{pq_i} . Let

$$\varphi_2 = \varphi - (\psi_1 + \dots + \psi_r).$$

Then φ_2 also has bias bounded away from zero (by Lemma 2.3 and Corollary 2.6), and φ_2 factors through $(A_1/pA_1) \times A_{[2,k]}$. By induction again there are multilinear maps $\psi_{r+1}, \ldots, \psi_{r_2} : (A_1/pA_1) \times A_{[2,k]} \to \mathbf{T}$, where $r_2 = O(1)$, such that each ψ_i factors through m_q for some bounded q, and such that

$$\varphi_2 = \psi_{r+1} + \cdots \psi_{r_2}.$$

Hence

$$\varphi = \psi_1 + \dots + \psi_{r_2},$$

and the induction is complete.

We end by deducing Corollary 1.3.

Proof of Corollary 1.3. Let $A_k = \widehat{B}$ and let $\varphi(x) = x_k(F(x_1, \ldots, x_{k-1}))$. Then φ is multilinear and $\operatorname{bias}(\varphi) = \mathbf{P}(f=0) \ge \epsilon > 0$, so Theorem 1.2 implies that φ is the sum of $O_{\epsilon}(1)$ multilinear functions of the form

$$\psi(x) = m_q(\varphi_1(x_I), \varphi_2(x_{I^c})),$$

where 0 < |I| < k and $q \leq O_{\epsilon}(1)$. Assume without loss of generality that $k \in I^c$. Since ψ is linear in x_k and $\widehat{A}_k \cong B$, there is some $b = b(x_{\lfloor k-1 \rfloor}) \in B$ such that

$$\psi(x) = x_k(b(x_{[k-1]}))$$

Moreover, b must be multilinear in $x_{[k-1]}$, and b can depend on x_I only through $\varphi_1(x_I)$. Hence ψ has the form

$$\psi(x) = x_k(G(g(x_I), x_{\lceil k-1 \rceil \setminus I})),$$

where $g = \varphi_1$. This implies that F is the sum of $O_{\epsilon}(1)$ maps of the form $G(g(x_I), x_{[k-1]\setminus I})$. Now fix I and suppose the terms of the form $G(g(x_I), x_{[k-1]\setminus I})$ are

$$G_j(g_j(x_I), x_{\lceil k-1 \rceil \setminus I}) \qquad (1 \le j \le n).$$

Let $g = (g_1, \ldots, g_n)$, and note that $|\operatorname{cod}(g)|$ is still bounded and $G_j(g_j(x_I), x_{[k-1]\setminus I}) = G'_j(g(x_I), x_{[k-1]\setminus I})$, where G'_j is the composite of G_j and the projection π_j . Let $G = G'_1 + \cdots + G'_n$. Then

$$\sum_{j=1}^{n} G_j(g_j(x_I), x_{[k-1]\setminus I}) = G(g(x_I), x_{[k-1]\setminus I}).$$

Hence F has the claimed form.

4. Possible biases

Let Φ_k be the set of all multilinear maps $\varphi : A_{[k]} \to \mathbf{T}$ for any finite abelian groups A_1, \ldots, A_k . In this section we probe the set of all possible biases

$$B_k = \{ bias(\varphi) : \varphi \in \Phi_k \}.$$

For example,

$$B_1 = \{0, 1\}, \qquad B_2 = \{1/n : n \ge 1\}.$$

By (3), $B_k \subseteq [0,1]$. We will show that Theorem 1.2 implies that B_k is a small subset of [0,1] in various senses. The results of this section are analogous to those

SEAN EBERHARD

proved in [E] essentially for B_3 and for the set of commuting probabilities of finite groups.

A slight generalization is useful. Call $\varphi : A_{[k]} \to \mathbf{T}$ multiaffine if $\varphi(x)$ is affinelinear in each variable x_i separately. Equivalently, φ is multiaffine if and only if it can be written

$$\varphi = \sum_{I \subseteq [k]} \varphi_I,$$

where φ_I is a multilinear map $A_I \to \mathbf{T}$. The greatest d = |I| such that φ_I is nontrivial is the *degree* of φ . Let $\Phi_{k,d}$ be the set of all multiaffine maps $\varphi : A_{[k]} \to \mathbf{T}$ (for any A_1, \ldots, A_k) of degree at most d and zero constant term, and let

$$B_{k,d} = \{ bias(\varphi) : \varphi \in \Phi_{k,d} \}$$

Thus $B_{k,d}$ is a subset of the unit disc $D = \{z \in \mathbf{C} : |z| \le 1\}.$

Example 4.1 (Gauss sums). Let $A_1 = A_2 = A_3 = \mathbf{F}_p$, where p is an odd prime, and let

$$\varphi(x,y,z) = (xy + xz + yz)/p \bmod 1 \qquad (x,y,z \in \mathbf{F}_p)$$

Thus $\varphi \in \Phi_{3,2}$. Then

bias
$$(\varphi) = \mathbf{E}_{x,y,z \in \mathbf{F}_p} e((xy + xz + yz)/p) = p^{-2}\overline{G(p)},$$

where G(p) is the Gauss sum

$$G(p) = \sum_{x \in \mathbf{F}_p} e(x^2/p).$$

It was proved by Gauss that $G(p) = p^{1/2}$ or $ip^{1/2}$ according to whether p is 1 or 3 mod 4. Hence $B_{3,2}$ includes $i^{\pm (p-1)^2/4}p^{-3/2}$ for every odd prime p.

If X is a subset of a compact metric space, let X' be the set of limit points of X. We iterate this operation transfinitely according to the following rules:

$X^{(0)} = X,$	
$X^{(\alpha+1)} = (X^{(\alpha)})'$	$(\alpha \text{ an ordinal}),$
$X^{(\lambda)} = \bigcap_{\alpha \in \mathcal{A}} X^{(\alpha)}$	(λ a nonzero limit ordinal).
$\alpha < \lambda$	

If X is countable then there is a countable ordinal β such that $X^{(\beta)} = \emptyset$. The smallest such β must be a successor ordinal, say $\alpha + 1$, and this ordinal α is called the *Cantor–Bendixson rank* of X and denoted CB(X). It is the unique ordinal such that $X^{(\alpha)}$ is finite and nonempty.¹

Theorem 4.2. The set $B_{k,d}$ has the following smallness properties.

- (1) All limit points of $B_{k,d}$ are algebraic (in fact, rational linear combinations of roots of unity).
- (2) $CB(B_{k,d}) \in \{1, \omega, \dots, \omega^{d-2}\}$ for $d \ge 2$.

In particular, Item (1) shows that $\overline{B_{k,d}}$ is countable, so in particular it is nowhere dense and measure-zero.

Proof. For $\epsilon > 0$ let

$$B_{k,d,\epsilon} = B_{k,d} \cap \{ z \in D : |z| \ge \epsilon \}.$$

¹More often β itself is called the Cantor–Bendixson rank, but the present usage is a common variant for countable sets.

Let $\varphi \in \Phi_{k,d,\epsilon}$, so $\varphi : A_{[k]} \to \mathbf{T}$ is a multiaffine map of degree d and zero constant term defined on some finite abelian groups A_1, \ldots, A_k and $|\operatorname{bias}(\varphi)| \ge \epsilon$. Write

$$\varphi = \sum_{I \subseteq [k]} \varphi_I,$$

where $\varphi_I : A_I \to \mathbf{T}$ is multilinear. Let φ_d be the sum of the terms with |I| = d. By Lemma 2.2, for each $I \subseteq [k]$ of size d we have

$$\operatorname{bias}(\varphi_I) \ge |\operatorname{bias}(\varphi)| \ge \epsilon,$$

so by Theorem 1.2 there is an expression for φ_I as the sum of $O_{\epsilon}(1)$ maps factoring through m_q for some bounded prime power q. Hence there is an expression for φ_d of the form

$$\varphi_d = \sum_{i=1}^r m_{q_i}(\varphi_i^L, \varphi_i^R),$$

where $r = O_{\epsilon}(1)$ and, for each i, φ_i^L and φ_i^R $(1 \le i \le r)$ are multilinear maps with disjoint sets of variables and codomain $\mathbf{Z}/q_i\mathbf{Z}$, where $q_i = O_{\epsilon}(1)$. Hence

$$e(\varphi_d) = F(\varphi_1^L, \dots, \varphi_r^L, \varphi_1^R, \dots, \varphi_r^R),$$

where

$$F(x_1, \ldots, x_r, y_1, \ldots, y_1) = e(x_1y_1/q_1 + \cdots + x_ry_r/q_r)$$

Let $\Gamma = (\prod_{i=1}^{r} \mathbf{Z}/q_i \mathbf{Z})^2$ and let $Q = q_1 \cdots q_r = |\Gamma|^{1/2}$. The Fourier expansion of F is

$$F(x_1, \dots, x_r, y_1, \dots, y_s) = \frac{1}{Q} \sum_{(u,v) \in \Gamma} e\left(\sum_{i=1}^r (-u_i v_i + u_i x_i + v_i y_i)/q_i\right).$$

Hence

$$e(\varphi_d) = \frac{1}{Q} \sum_{(u,v)\in\Gamma} e\left(\sum_{i=1}^r (-u_i v_i + u_i \varphi_i^L + v_i \varphi_i^R)/q_i\right)$$

and

$$\operatorname{bias}(\varphi) = \frac{1}{Q} \sum_{(u,v)\in\Gamma} e(-u_1 v_1/q_1 - \dots - u_r v_r/q_r) \operatorname{bias}(\psi_{u,v}),$$

where

$$\psi_{u,v} = \sum_{i=1}^{\prime} (u_i \varphi_i^L + v_i \varphi_i^R) / q_i + (\varphi - \varphi_d)$$

Note that $\psi_{u,v}$ is multi-affine of degree at most d-1 with zero constant term. Since $|\Gamma| \leq O_{\epsilon}(1)$, we have proved that

(4)
$$B_{k,d,\epsilon} \subseteq O_{\epsilon}(1)H_{O_{\epsilon}(1)}B_{k,d-1},$$

where H_X is the (finite) set of all complex numbers of the form ζ/m where ζ is a root of unity of order at most X and m is a positive integer of size at most X. Here we are also using the standard notation $ST = \{st : s \in S, t \in T\}$ for the product of sets S and T and $nS = S + \cdots + S$ for the sum of n copies of a set S.

We will now use induction on d to prove Items (1) and (2). As the base of the induction, note that an affine linear map with zero constant term is just a linear map, so $B_{k,1} = B_1 = \{0, 1\}$. Hence, for d = 2, (4) shows that $B_{k,2,\epsilon}$ is finite for all $\epsilon > 0$, so $B'_{k,2} \subseteq \{0\}$. Hence we may assume d > 2.

Let z be a limit point of $B_{k,d}$. Then either z = 0 or z is a limit point of $B_{k,d,\epsilon}$ for $\epsilon = |z|/2$. In the latter case (4) shows that

$$z \in O_{\epsilon}(1)H_{O_{\epsilon}(1)}\overline{B_{k,d-1}}.$$

By induction on d this shows that z is a rational linear combination of roots of unity. This proves Item (1).

Let $\alpha = CB(B_{k,d})$. Then it follows from (4) that

$$B_{k,d,\epsilon}^{(\alpha O_{\epsilon}(1))} = \emptyset.$$

Since this holds for all $\epsilon > 0$, it follows that

$$B_{k,d}^{(\alpha\omega)} \subseteq \{0\}$$

Hence

$$\operatorname{CB}(B_{k,d}) \le \operatorname{CB}(B_{k,d-1})\omega,$$

and this proves $\operatorname{CB}(B_{k,d}) \leq \omega^{d-2}$ by induction. Finally, since $B_{k,d}$ is closed under multiplication, the argument of [E, Lemma 5.1] shows that $\operatorname{CB}(B_{k,d}) = \omega^{\gamma}$ for some ordinal γ , so $\operatorname{CB}(B_{k,d}) \in \{1, \omega, \dots, \omega^{d-2}\}$.

- Since $B_k \subseteq B_{k,k}$, the theorem applies in particular to $B_k \subseteq [0, 1]$ and shows
- (1) all limit points of B_k are algebraic,
- (2) $\operatorname{CB}(B_k) \in \{1, \omega, \dots, \omega^{k-2}\}.$

It is very likely that all limit points of B_k are actually rational, but a slightly more sophisticated analysis is required to prove this. As to the Cantor–Bendixson rank, basic examples demonstrate that $CB(B_k) > 1$ for k > 2, so

$$CB(B_1) = 0, \qquad CB(B_2) = 1, \qquad CB(B_3) = \omega$$

but the value of $\operatorname{CB}(B_k)$ for k > 3 is unclear. Relatedly, it is plausible that B_k is well-ordered by the reverse of the natural order, i.e., for every $x \in B_k$ there is some $\epsilon > 0$ such that $(x - \epsilon, x) \cap B_k = \emptyset$; if this is the case it follows that B_k has order type $\omega^{\operatorname{CB}(B_k)}$. We leave these questions as open problems.

Problem 4.3.

- (1) Prove that all limit points of B_k are rational.
- (2) Determine the Cantor-Bendixson rank of B_k . In particular, determine whether $CB(B_4)$ is ω or ω^2 .
- (3) Determine whether B_k is reverse-well-ordered.

References

- [BL] A. Bhowmick and S. Lovett, Bias vs structure of polynomials in large fields, and applications in effective algebraic geometry and coding theory, 2015. arXiv:1506.02047. ↑1
- [CM] A. Cohen and G. Moshkovitz, An optimal inverse theorem, 2021. arXiv:2102.10509. ↑1
- [E] S. Eberhard, Commuting probabilities of finite groups, Bull. Lond. Math. Soc. 47 (2015), no. 5, 796–808. MR3403962 ↑4, 4
- [GT] B. Green and T. Tao, The distribution of polynomials over finite fields, with applications to the Gowers norms, Contrib. Discrete Math. 4 (2009), no. 2, 1–36. MR2592422 ↑1
- [GW] W. T. Gowers and J. Wolf, Linear forms and higher-degree uniformity for functions on Fⁿ_p, Geom. Funct. Anal. 21 (2011), no. 1, 36–69. MR2773103 ↑1
- [J] O. Janzer, Polynomial bound for the partition rank vs the analytic rank of tensors, Discrete Anal. (2020), Paper No. 7, 18. MR4107323 ↑1
- [L] S. Lovett, The analytic rank of tensors and its applications, Discrete Anal. (2019), Paper No. 7, 10. MR3964143 ↑1, 2
- [M] L. Milićević, Polynomial bound for partition rank in terms of analytic rank, Geom. Funct. Anal. 29 (2019), no. 5, 1503–1530. MR4025518 ↑1