ITERATING THE BIG-PIECES OPERATOR AND LARGER SETS

JARED KRANDEL AND RAANAN SCHUL

ABSTRACT. We show that if an Ahlfors-David regular set E of dimension k has Big Pieces of Big Pieces of Lipschitz Graphs (denoted usually by BP(BP(LG))), then $E \subset \tilde{E}$ where \tilde{E} is Ahlfors-David regular of dimension k and has Big Pieces of Lipschitz Graphs (denoted usually by BP(LG)). Our results are quantitative and, in fact, are proven in the setting of a metric space for any family of Ahlfors-David regular sets \mathcal{F} replacing LG. A simple corollary is the stability of the BP operator after 2 iterations. This was previously only known in the Euclidean setting for the case $\mathcal{F} = LG$ with substantially more complicated proofs.

1. INTRODUCTION

A closed set E (with more than one point) in a metric space X is said to be k-Ahlfors-David regular if there is a constant C > 1 such that for all $r \in (0, \operatorname{diam}(E))$ and $x \in E$ we have $C^{-1}r^k < \mathcal{H}^k(E \cap B(x, r)) < Cr^k$. For some given class \mathcal{F} of k-Ahlfors-David regular subsets (of a metric space X), we define BP(\mathcal{F}) as follows: $F \in BP(\mathcal{F})$ if F is a k-Ahlfors-David regular set for which there exists a constant $\theta > 0$ such that for any $x \in F$ and R > 0, there is a set $G_{x,R} \in \mathcal{F}$ such that

$$\mathcal{H}^{\kappa}(B(x,R)\cap F\cap G_{x,R})\geq\theta\mathcal{H}^{\kappa}(B(x,R)\cap F).$$

Conditions involving $BP(\mathcal{F})$ for various classes of sets \mathcal{F} play an important role in the theory of uniformly rectifiable sets in \mathbb{R}^n developed By David and Semmes (see e.g. [4], [6]). While the original motivation was the study of singular integral operators, the study of such conditions has taken on a life of its own.

In the context of singular integrals, the condition $BP(\mathcal{F})$ is important because it allows the uniform boundedness of a family of SIOs given by convolution with 'nice' kernels over sets in \mathcal{F} to be transported to sets in $BP(\mathcal{F})$. In particular, one can define successively weaker conditions $BP^{j}(\mathcal{F})$ for all j > 0 which all imply boundedness given that the SIOs are bounded on \mathcal{F} ; the initial case David and Semmes considered [5] used Lipschitz graphs as the base class, i.e., $E \in BP^{j}(LG)$. This raised a natural question: how do the collections $BP^{j}(LG)$ behave as j grows? It turned out that for $j \ge 2$ the collections $BP^{j}(LG)$ are all the same and their elements are called *Uniformly Rectifiable sets*. We refer the reader to [6], [5], specifically to Proposition 2.2 on page 97 of [5], and Theorem 2.29 on page 336 of [5]. For $n \in [k, 2d + 1)$ one also needs [1] to show that BPBI implies BP(BP(LG)), but this is not where most of the work goes – the proofs by David and Semmes of that stability (for $j \ge 2$) are quite sophisticated and rely on a Euclidean ambient space.

Date: July 2021.

²⁰¹⁰ Mathematics Subject Classification. 28A75, 30L99.

R. Schul was partially supported by the National Science Foundation under Grants No. DMS-1763973.

There has recently been interest in other families \mathcal{F} , in particular for the purpose of studying *Parabolic Uniform Rectifiability*. See e.g. the work in [2], where the questions about Uniform Rectifiability in the metric setting are discussed for this purpose. In fact, we refer to [2] for a great introduction on contemporary applications of the idea of Big Pieces.

An immediate corollary of the main result contained in this essay (Theorem 3.1) is that stabilization of the operator BP^{j} occurs in the setting of metric spaces for $j \ge 2$ as well. Our proof is both simple and direct.

2. ACKNOWLEDGEMENTS

The second author would like to thank Jonas Azzam for pointing out earlier work of David and Semmes.

3. A THEOREM

Theorem 3.1. Let \mathcal{F} be a class of (closed) k-Ahlfors-David regular sets in a metric space \mathbb{X} . Let $E \subseteq \mathbb{X}$ be a k-Ahlfors-David regular set with $E \in BP(BP(\mathcal{F}))$. Then there exists a set $F \subset \mathbb{X}$ such that

(i) $E \subseteq F$,

(*ii*) F is k-Ahlfors-David regular.

(*iii*) $F \in BP(\mathcal{F})$.

The constants in the conclusion are quantitative with dependance on the constants in the assumptions.

Corollary 3.2. Let \mathcal{F} be a class of closed k-Ahlfors-David regular sets in a metric space \mathbb{X} . For any j > 2, and any constants $\theta_1, ..., \theta_j > 0$ defining $BP^j(\mathcal{F})$, there are $\theta'_1, \theta'_2 > 0$ such that the family $BP(BP(\mathcal{F}))$ defined using θ'_1, θ'_2 is equal to $BP^j(\mathcal{F})$ defined using $\theta_1, ..., \theta_j$

Proof of Corollary 3.2. Let $E \in BP^3(\mathcal{F})$. Then for any $x \in E$ and $R < \operatorname{diam}(E)$ we have a set $E'_{x,R} \in BP^2(\mathcal{F})$ such that $\mathcal{H}^k(B(x,R) \cap E) \leq \mathcal{H}^k(B(x,R) \cap E \cap E'_{x,R})$. By Theorem 3.1, there is a set $F_{x,R} \in BP(\mathcal{F})$ so that $F_{x,R} \supset E'_{x,R}$. Clearly $\mathcal{H}^k(B(x,R) \cap E) \leq \mathcal{H}^k(B(x,R) \cap E \cap F_{x,R})$. We have shown $E \in BP^2(\mathcal{F})$. This gives for any $j \geq 3$ that $BP^j(\mathcal{F}) = BP^{j-1}(\mathcal{F})$, and so we are done by induction.

Proof of Theorem 3.1 for the case diam $E < \infty$. We suppose that diam $E < \infty$. In order to construct the set F, we first fix a dyadic cube decomposition of E denoted by $\Delta = \Delta(E)$ with root cube $\operatorname{root}(\Delta) = Q_0 = E$. By construction, for each cube $Q \in \Delta$ there exists a point $c(Q) \in Q$ which we call the center of Q satisfying

(3.1)
$$\operatorname{dist}(B(c(Q), c_1 \operatorname{diam} Q), E \setminus Q) \ge c_2 \operatorname{diam} Q.$$

for some constants $c_1, c_2 > 0$ (see e.g. [3]). From now on, define $B_{c(Q)} = B(c(Q), c_1 \operatorname{diam}(Q))$. We construct the set F desired in the theorem inductively. At stage 0, use the fact that $E \in \operatorname{BP}(\operatorname{BP}(\mathcal{F}))$ to find a closed set $F_{Q_0} \in \operatorname{BP}(\mathcal{F})$ such that $F_{Q_0} \subseteq B_{c(Q_0)}$ and

$$\mathcal{H}^k(B_{c(Q_0)} \cap E \cap F_{Q_0}) \gtrsim_{\theta_1} \mathcal{H}^k(B_{c(Q_0)} \cap E) \gtrsim_{c_1, c_2} \operatorname{diam}(Q_0)^k.$$

2

We define

$$F_0 = F_{Q_0}$$

We continue the construction by defining a dyadic decomposition Q_1 of the set $E \setminus F_{Q_0}$. Indeed, since F_{Q_0} is closed, $E \setminus F_{Q_0}$ is relatively open in E and for any $x \in E \setminus F_{Q_0}$, there exists some dyadic cube $Q \ni x$ of maximal diameter such that $\operatorname{dist}(Q, F_{Q_0}) > \operatorname{diam} Q$. We call the disjoint family of all such maximal cubes Q_1 , so that we have

$$E \setminus F_{Q_0} = \bigcup_{Q \in \mathcal{Q}_1} Q.$$

We now give stage 1 of the construction of F. For each $Q \in Q_1$, Again find closed a set $F_Q \in BP(\mathcal{F})$ such that

(3.2)
$$\mathcal{H}^k(B_{c(Q)} \cap E \cap F_Q) \gtrsim_{\theta_1, c_1, c_2} \mathcal{H}^k(Q).$$

We define

$$F_1 = F_{Q_0} \cup \bigcup_{Q \in \mathcal{Q}_1} F_Q.$$

Continue the construction inductively. Given the construction completed up to stage m, we define the set Q_{m+1} to be the collection of dyadic cubes with maximal diameter contained in $E \setminus F_m$ such that $Q \in Q_{m+1}$ satisfies

$$\operatorname{dist}(Q, F_m) > \operatorname{diam}(Q)$$

 \mathcal{Q}_{m+1} is a disjoint decomposition of $E \setminus F_m$ so that

$$(3.4) E \setminus F_m = \bigcup_{Q \in \mathcal{Q}_{m+1}} Q.$$

Given such a Q, let $F_Q \in BP(\mathcal{F})$ with $F_Q \subseteq B_{c(Q)}$ be such that (3.2) holds and define

(3.5)
$$F_{m+1} = F_m \cup \bigcup_{Q \in \mathcal{Q}_{m+1}} F_Q = F_{Q_0} \cup \bigcup_{Q \in \mathcal{Q}_1} F_Q \cup \cdots \cup \bigcup_{Q \in \mathcal{Q}_{m+1}} F_Q$$

Finally, set

$$F = \bigcup_{m=0}^{\infty} F_m$$

and define $Q = \bigcup_m Q_m$. Now that we have constructed the set F, we note two of its simple properties. First, given any $Q \neq Q' \in Q$, equality (3.3) implies

$$(3.7) \qquad \qquad \operatorname{dist}(F_Q, F_{Q'}) > \min\{\operatorname{diam}(Q), \operatorname{diam}(Q')\}$$

Second,

$$\lim F \subseteq E \cup \bigcup_{m=0}^{\infty} F_m$$

where $\lim F$ denotes the set of limit points of F. Indeed, suppose $x \in \lim F$ with $x_j \to x, x_j \in F_{Q_j}$. If the set $\{Q_j\}_j$ is finite, then (3.7) implies the sequence F_{Q_j} is eventually constant, say $F_{Q_j} \to F_{Q_i}$ meaning $x \in F_{Q_i}$ since F_{Q_i} is closed. If instead $\{Q_j\}_j$ is infinite, then consider a subsequence $x_{k_j} \to x$ such that $Q_{k_j} \neq Q_{k_i}$ for any i, j. The fact that x_{k_j} converges combined with (3.7) then implies diam $Q_j \to 0$. Since dist $(F_{Q_j}, E) \leq \text{diam } Q_j$, we have dist(x, E) = 0 which implies $x \in E$. (In particular, we will soon see that this implies $\mathcal{H}^k(\lim F \setminus \bigcup_m F_m) = 0$.)

We begin with proving claim (i). Notice that for any $N \in \mathbb{N}$,

$$\mathcal{H}^{k}(E \setminus F) \leq \mathcal{H}^{k}\left(E \setminus \bigcup_{m=0}^{\infty} F_{m}\right) \leq \mathcal{H}^{k}(E \setminus F_{N})$$

because the sets F_m are increasing. Letting $0 < c_0 < 1$ be the constant implicit in inequality (3.2), we can write

$$\mathcal{H}^{k}(E \setminus F_{N}) \stackrel{(3.5)}{=} \mathcal{H}^{k}\left(E \setminus F_{N-1} \setminus \bigcup_{Q \in \mathcal{Q}_{N}} F_{Q}\right) \stackrel{(3.4)}{=} \mathcal{H}^{k}\left(\bigcup_{Q \in \mathcal{Q}_{N}} Q \setminus \bigcup_{Q \in \mathcal{Q}_{N}} F_{Q}\right)$$
$$= \sum_{Q \in \mathcal{Q}_{N}} \mathcal{H}^{k}(Q \setminus F_{Q}) \leq (1 - c_{0}) \sum_{Q \in \mathcal{Q}_{N}} \mathcal{H}^{k}(Q) = (1 - c_{0})\mathcal{H}^{k}(E \setminus F_{N-1})$$

where we used the fact that $F_Q \cap F_{Q'} = \emptyset$ for $Q, Q' \in Q_N$. Since this holds for any N, we can iterate this inequality to get

$$\mathcal{H}^k(E \setminus F_N) \le (1 - c_0)^N \mathcal{H}^k(E)$$

from which we conclude $\mathcal{H}^k(E \setminus F) = 0$. To finish the proof of (i), let $x \in E$ be arbitrary. Since E is Ahlfors-David regular, for any R > 0, $\mathcal{H}^k(B(x, R)) > 0$ so that $F \cap B(x, r) \neq \emptyset$. This means x is a limit point of F, implying $x \in F$ because F is closed.

We now prove (ii). Fix any point $x \in F$ and some $R < \operatorname{diam} F$. If $x \in F \setminus \bigcup_m F_m$, then we can find a particular F_Q with $\operatorname{dist}(x, F_Q) < \frac{R}{100}$ and $\operatorname{dist}(x, F_Q) = \operatorname{dist}(x, z)$ for $z \in F_Q$. Then, we have $B(z, R/2) \subseteq B(x, R) \subseteq B(z, 2R)$, and substitute the first ball or final ball for B(x, R) in the proofs of lower and upper regularity respectively. Hence, we can assume $x \in \bigcup_m F_m$. By definition, there exists $Q_m \in Q_m$ such that $x \in F_{Q_m}$ for some $m \in \mathbb{N}$. Write

$$(3.8) \qquad \mathcal{H}^k(B(x,R)\cap F) = \sum_{\substack{F_Q\cap B(x,R)\neq\varnothing\\ \text{diam}\,Q>10R}} \mathcal{H}^k(B(x,R)\cap F_Q) + \sum_{\substack{F_Q\cap B(x,R)\neq\varnothing\\ \text{diam}\,Q\le 10R}} \mathcal{H}^k(B(x,R)\cap F_Q).$$

We will first show that F is upper regular. Let Q_I be the collection of cubes summed over in the first term of (3.8). By (3.7), we have that for any $Q, Q' \in Q_I$, $dist(F_Q, F_{Q'}) > 10R$. This means Q_I has at most one element. Given such a Q, choose $y \in B(x, R) \cap F_Q$ and write

$$\mathcal{H}^k(B(x,R)\cap F_Q) \leq \mathcal{H}^k(F_Q\cap B(y,2R)) \lesssim R^k$$

using the fact that F_Q is itself k-Ahlfors-David regular. This proves the first sum in (3.8) has the appropriate upper bound. Let Q_{II} be the collection of cubes summed over in the second term of (3.8). Since diam(Q) < 10R, any $Q \in Q_{II}$ satisfies $Q \subseteq B(x, 20R)$. We first prove a lemma

Lemma 3.3. Let $Q \in \mathcal{Q}$, and let D(Q) be the descendants of Q in \mathcal{Q} . Then

$$\mathcal{H}^k\left(\bigcup_{Q'\in D(Q)}F_{Q'}\right)=\sum_{Q'\in D(Q)}\mathcal{H}^k(F_{Q'})\lesssim_{\theta_1,c_1,c_2}\mathcal{H}^k(Q).$$

Proof of Lemma 3.3. Suppose for simple notation that $Q = Q_0$. Using the regularity of each F_Q , we have

$$\mathcal{H}^k\left(\bigcup_{Q\in D(Q_0)}F_Q\right) = \sum_{m=0}^{\infty}\sum_{Q\in D(Q_0)\cap\mathcal{Q}_m}\mathcal{H}^k(F_Q) \le C\sum_{m=0}^{\infty}\sum_{Q\in D(Q_0)\cap\mathcal{Q}_m}\mathcal{H}^k(Q).$$

In analogy to (3.4), $Q_0 \setminus F_{m-1} = \bigcup_{Q \in D(Q_0) \cap \mathcal{Q}_m} Q$ holds so that

$$\sum_{Q \in D(Q_0) \cap \mathcal{Q}_m} \mathcal{H}^k(Q) = \mathcal{H}^k(Q_0 \setminus F_{m-1}) = \mathcal{H}^k\left(Q_0 \setminus F_{m-2} \setminus \bigcup_{Q \in D(Q_0) \cap \mathcal{Q}_{m-1}} F_Q\right)$$
$$= \mathcal{H}^k\left(\bigcup_{Q \in D(Q_0) \cap \mathcal{Q}_{m-1}} Q \setminus \bigcup_{Q \in D(Q_0) \cap \mathcal{Q}_{m-1}} F_Q\right)$$
$$\leq \sum_{Q \in D(Q_0) \cap \mathcal{Q}_{m-1}} \mathcal{H}^k(Q \setminus F_Q)$$
$$\leq (1 - c_0) \sum_{Q \in D(Q_0) \cap \mathcal{Q}_{m-1}} \mathcal{H}^k(Q)$$

where c_0 was defined as the implicit constant in (3.2). Iterating this inequality, we find

$$\mathcal{H}^k\left(\bigcup_{Q\in D(Q_0)}F_Q\right)\leq C\sum_{m=0}^\infty(1-c_0)^m\mathcal{H}^k(Q_0)\lesssim_{c_0}\mathcal{H}^k(Q_0).$$

Using this lemma, we can write

1

$$\sum_{\substack{F_Q \cap B(x,R) \neq \varnothing \\ \text{diam } Q \leq 10R}} \mathcal{H}^k(B(x,R) \cap F_Q) \leq \sum_{\substack{Q \text{ maximal } \\ Q \in \mathcal{Q}_{II}}} \sum_{\substack{Q' \in D(Q) \\ Q \in \mathcal{Q}_{II}}} \mathcal{H}^k(F_{Q'}) \lesssim \sum_{\substack{Q \text{ maximal } \\ Q \in \mathcal{Q}_{II}}} \mathcal{H}^k(Q)$$
$$\leq \mathcal{H}^k(E \cap B(x, 20R)) \lesssim R^k.$$

This proves the desired bound for the second sum in (3.8), proving the upper regularity of F. Now we show that F is lower regular. If $R < 100 \operatorname{diam} Q_m$, then the claim follows immediately from the lower regularity of F_Q . If $100 \operatorname{diam} Q_m \leq R < \operatorname{diam} F$, then since $F_{Q_m} \cap Q_m \neq \emptyset$, there exists $z \in Q$ (and thus, $z \in E$) with $B(x, R) \supseteq B(z, R/2)$ and

$$\mathcal{H}^k(B(x,R)\cap F) \ge \mathcal{H}^k(B(z,R/2)\cap E) \gtrsim R^k$$

using the fact that $E \subseteq F$. This completes the proof of lower regularity, hence of (ii) as well.

١

JARED KRANDEL AND RAANAN SCHUL

Finally, we prove (iii). Fix $x \in F_{Q_m}$ and R > 0 as in the proof of (ii). Fix a constant $\alpha > 10$ to be chosen later. If $R < \alpha \operatorname{diam} Q_m$, then since $F_{Q_m} \in \operatorname{BP}(\mathcal{F})$, there exists $G_{x,R} \in \mathcal{F}$ such that

$$(3.9) \qquad \mathcal{H}^k(B(x,R) \cap F_{Q_m} \cap G_{x,R}) \ge \theta_2 \mathcal{H}^k(B(x,R) \cap F_{Q_m}) \gtrsim_{C',\alpha} R^k \gtrsim_{C''} \mathcal{H}^k(B(x,R) \cap F)$$

where C' is the regularity constant for F_{Q_m} and C'' is the regularity constant for F. Now, suppose that $\alpha \operatorname{diam} Q_m \leq R < \operatorname{diam} F$. Since $x \in F_{Q_m}$, there exists a chain of cubes $Q_i \in Q_i$, $0 \leq i \leq m$ such that

$$Q_m \subseteq Q_{m-1} \subseteq \ldots \subseteq Q_1 \subseteq Q_0.$$

Next, notice that for any choice of $\alpha > 10$, there exists a smallest cube Q_j in the above chain such that $R < \alpha \operatorname{diam} Q_j$ since for all admissible $R, R < 10 \operatorname{diam} Q_0$. Choose the constant α such that for any $y \in E \setminus F_i$, the cube $Q_{i+1} \ni y$ satisfies

(3.10)
$$\operatorname{dist}(Q_{i+1}, F_{Q_i}) < \frac{\alpha}{10} \operatorname{diam} Q_{i+1}.$$

In general, α will depend on the constants used in the construction of Δ , as it may be the case that all of the children of the cube Q_i are small relative to Q_i with bounds given in terms of these constants. With such an α chosen, let Q_j be the smallest cube in the above chain for x such that $R < \alpha \operatorname{diam} Q_j$. This means that $R \ge \alpha \operatorname{diam} Q_{j+1}$ so that (3.10) implies that there exists $y \in F_{Q_j}$ such that $B(y, R/2) \subseteq$ B(x, R). We can now repeat the argument of (3.9) with Q_j in place of Q_m to finish the proof. This completes the proof of Theorem 3.1 for the case diam $E < \infty$.

Before we turn to the case diam $E = \infty$, we need the following lemma. It says, roughly, that finite diameter subsets of E can be made regular by extending them slightly. This extension also preserves the BP(\mathcal{F}) property.

Lemma 3.4. Let $E \subseteq X$ be a k-Ahlfors-David regular set and suppose that $G \subseteq E$ satisfies diam $G = D < \infty$. For any $A \ge 1$, there exists a set $\tilde{G} \subseteq E$ such that

(i)
$$G \subseteq \tilde{G} \subseteq B(G, \frac{3D}{A}) \cap E = \{x \in E : d(x, G) < \frac{3D}{A}\},\$$

(ii) G is k-Ahlfors-David regular with constant $C(k, C_E, A)$.

Furthermore, if $E \in BP(\mathcal{F})$ with constant θ_E for some class of k-Ahlfors-David regular sets, then $\tilde{G} \in BP(\mathcal{F})$ with constant $\theta(k, \theta_E, A)$.

Proof of Lemma 3.4. We define an "interior" of the set $G \subseteq E$ by

$$I_A(G) = \left\{ x \in G : d(x, E \setminus G) \ge \frac{D}{A} \right\}.$$

The corresponding "boundary" is then

$$G \setminus I_A(G) = \left\{ x \in G : d(x, E \setminus G) < \frac{D}{A} \right\}$$

We will construct the set \tilde{G} inductively. In the first stage, we will take a maximal net of appropriate size inside $G \setminus I_A(G)$ and add in balls around each net point to G. In the second step, we consider a smaller "boundary" of this new set and repeat the above process with a finer net and smaller balls. If

we continue this process indefinitely while adding balls of exponentially decreasing radii, we get the desired set by taking a closure. We now give this construction explicitly.

Let $G_0 = G$ and let X_1 be a maximal $\frac{D}{A}$ -net for the set $G \setminus I_A(G) \subseteq E$. Define

$$G_1 = G \cup \bigcup_{x \in X_1} B\left(x, \frac{2D}{A}\right) \cap E.$$

Given the set G_n , we define X_{n+1} to be a maximal $4^{-n}\frac{D}{A}$ -net for $G_n \setminus I_{4^n \cdot A}(G_n)$ and we let

$$G_{n+1} = G_n \cup \bigcup_{x \in X_{n+1}} B\left(x, 4^{-n} \frac{2D}{A}\right) \cap E.$$

Finally, define

$$\tilde{G} = \overline{\bigcup_{n=0}^{\infty} G_n}.$$

We will now show that \tilde{G} satisfies the desired properties in the statement of the lemma.

We begin by proving (i). The maximal distance of a point $x \in \tilde{G}$ from G is just given by the sum of the radii of the balls added in each step:

$$d(x,G) \le \frac{2D}{A} \sum_{n=0}^{\infty} 4^{-n} = \frac{8}{3} \frac{D}{A} < \frac{3D}{A}.$$

We now prove (ii). First, we observe that since $\tilde{G} \subseteq E$, we immediately have, for all $x \in \tilde{G}$, R > 0,

$$\mathcal{H}^k(B(x,R)\cap \tilde{G}) \le \mathcal{H}^k(B(x,R)\cap E) \le C_E R^k.$$

Hence, \tilde{G} is upper k-Ahlfors-David regular with constant C_E . We will now show that \tilde{G} is lower regular. In order to do so, we will first prove that there exists a constant 0 < c < 1 dependent only on A such that

$$(3.11) \qquad \forall x \in \tilde{G}, \ \forall R, \ 0 < R < \operatorname{diam} \tilde{G}, \ \exists y \in E \text{ such that } B(y, cR) \cap E \subseteq B(x, R) \cap \tilde{G}$$

We note that (ii) will follow from this since for any relevant pair (x, R), we get the existence of $y \in E$ such that

$$\mathcal{H}^k(B(x,R) \cap \tilde{G}) \ge \mathcal{H}^k(B(y,cR) \cap E) \ge \frac{c^k R^k}{C_E}$$

by the lower regularity of E. We now prove (3.11). We begin by using the constant $c' = \frac{1}{10 \cdot 4^4 \cdot A}$ (we will only need to decrease it by a factor of $\frac{1}{2}$ at the end of the proof). Let $x \in \tilde{G}$ and assume $x \in G_m$ for some m. There exists some minimal n such that $x \in I_{4^n A}(G_n)$ because $x \in G_m \setminus I_{4^m A}(G_m)$ implies $x \in I_{4^{m+1}A}(G_{m+1})$ by the triangle inequality. Indeed, let $t \in X_{m+1}$ be a nearest net point to x and let $z \in E \setminus G_{m+1}$. We can calculate

$$d(x,z) \ge d(t,z) - d(t,x) \ge 4^{-m} \frac{2D}{A} - 4^{-m} \frac{D}{A} = 4^{-m} \frac{D}{A} > 4^{-m-1} \frac{D}{A}.$$

Therefore, $d(x, E \setminus G_{m+1}) > 4^{-m-1}\frac{D}{A}$ so that $x \in I_{4^{m+1}A}(G_{m+1})$. Suppose first that $n \leq 4$. In this case, we will take y = x, and we must show the inclusion of the balls given in (3.11) for any admissible value of R. For $0 < R \leq 4^{-4}\frac{D}{A}$, note that $x \in I_{4^4A}(G_4)$ implies

(3.12)
$$d(x, E \setminus \tilde{G}) \ge d(x, E \setminus G_4) > \frac{D}{4^4 A}$$

so that $B(x,R) \cap \tilde{G} = B(x,R) \cap E$. If instead $4^{-4}\frac{D}{A} < R < \operatorname{diam} \tilde{G} < D + \frac{6D}{A} < 10D$,

$$c'R = \frac{R}{10 \cdot 4^4 \cdot A} < \frac{10D}{10 \cdot 4^4 \cdot A} = 4^{-4}\frac{D}{A}$$

Which shows that

$$B(x, c'R) \cap E \subseteq B\left(x, 4^{-4}\frac{D}{A}\right) \cap E = B\left(x, 4^{-4}\frac{D}{A}\right) \cap \tilde{G}$$

by (3.12). Now, suppose n > 4. This means $x \in I_{4^nA}(G_n) \setminus I_{4^{n-1}A}(G_{n-1})$. Hence, if $R < 4^{-n}\frac{D}{A}$, then we can take y = x and note that $B(x, R) \cap \tilde{G} = B(x, R) \cap E$ in analogy to (3.12). Now, suppose $4^{-m}\frac{D}{A} \leq R < 4^{-m+1}\frac{D}{A}$ for $0 \leq m \leq n-3$. There exist net points $x_p \in X_p$ for $m+3 \leq p \leq n$ such that

$$d(x, x_n) \le 4^{-n} \frac{2D}{A},$$

$$d(x_{p+1}, x_p) \le 4^{-p} \frac{2D}{A}.$$

Hence, the triangle inequality implies

(3.13)
$$d(x, x_{m+3}) \le \frac{2D}{A} \sum_{p=m+2}^{n} 4^{-p} \le \frac{2D}{A} \left(4^{-m-2} \cdot 2 \right) = 4^{-m-1} \frac{D}{A}.$$

In this case, we choose $y = x_{m+3}$. We calculate

$$B(y,c'R) = B\left(x_{m+3}, \frac{R}{10 \cdot 4^4 \cdot A}\right) \subseteq B\left(x_{m+3}, 4^{-(m+3)}\frac{D}{10A^2}\right) \subseteq B\left(x, 4^{-m}\frac{D}{A}\right) \subseteq B(x,R)$$

using (3.13) and the fact that $4^{-m}\frac{D}{A} \leq R < 4^{-m+1}\frac{D}{A}$. In the case when $\frac{D}{A} < R < 10D$, choose $y = x_3$, the nearest net point in X_3 and observe that

$$B(y,c'R) = B\left(x_3, \frac{R}{10 \cdot 4^4 \cdot A}\right) \subseteq B\left(x_{m+3}, 4^{-4}\frac{D}{A}\right) \subseteq B\left(x, \frac{D}{A}\right) \subseteq B(x, R)$$

again using (3.13). This proves (3.11) for all $x \in G_n$ for some n. If $x \notin G_n$ for all n, then given any admissible R > 0, there is a net point $t \in X_N$ for arbitrarily large N such that $d(x,t) < \frac{R}{4}$ so that $B(t, \frac{R}{2}) \subseteq B(x, R)$ and, applying (3.11) to $B(t, \frac{R}{2})$, we get a point $y \in B(t, \frac{R}{2})$ such that $B(y, c'\frac{R}{2}) \subseteq B(t, \frac{R}{2}) \subseteq B(x, R)$. Take $c = \frac{c'}{2}$ and $B(y, cR) \subseteq B(x, R)$ so that (3.11) holds with $c = \frac{1}{20 \cdot 4^4 \cdot A}$. **Proof that** $\hat{G} \in BP(\mathcal{F})$. This follows from (3.11). Indeed, for any admissible pair (x, R), choose y as given by (3.11). Applying the $BP(\mathcal{F})$ condition for E in the ball B(y, cR) gives a set $H_{y,cR} \in \mathcal{F}$ such that

$$\mathcal{H}^{k}(B(x,R)\cap \hat{G}\cap H_{y,cR}) \geq \mathcal{H}^{k}(B(y,cR)\cap E\cap H_{y,cR}) \gtrsim_{A,\theta_{E},k} R^{k} \gtrsim_{C} \mathcal{H}^{k}(B(x,R)\cap \hat{G}).$$

This concludes the proof of the lemma.

Proof of Theorem 3.1 for the case diam $E = \infty$. Fix $x_0 \in E$. Let A > 1 and, for $n \ge 0$, set

$$B_n = B(x_0, A^n)$$

where the constant A is sufficiently large in terms of C_E , k, and θ_E , the BP constant. Let E_n be the Ahlfors-David regular extension of the set $E \cap B_n$ with constant A in Lemma 3.4 replaced with 100 so that $E_n \subseteq B(E \cap B_n, \frac{A^n}{4})$. E_n satisfies the hypotheses of the finite diameter case of the theorem, so apply the theorem to get a regular set $F_n \in BP(\mathcal{F})$ satisfying

$$E_n \subseteq F_n \subseteq B\left(x_0, \frac{5A^n}{4}\right).$$

In order to ensure bounded overlap, we then define $\tilde{F}_0 = F_0$ and \tilde{F}_n for $n \ge 1$ to be the regular extension of $F_n \setminus \frac{1}{2}B_{n-1}$ given by the lemma with constant A there replaced by 100A here. By construction, $\tilde{F}_n \subseteq B(F_n, \frac{A^{n-1}}{10})$ so that $\tilde{F}_n \cap \frac{1}{4}B_{n-1} = \emptyset$ and $\tilde{F}_n \subseteq B(x_0, 2A^n)$. We also have $\tilde{F}_n \in BP(\mathcal{F})$ with constant $\tilde{\theta}_F$ independent of n. We now define

$$F = \bigcup_{n=0}^{\infty} \tilde{F}_n$$

and claim that F satisfies conditions (i)-(iii).

Proof of (i). By definition, $E \cap (B_n \setminus \frac{1}{2}B_{n-1}) \subseteq \tilde{F}_n$ so $E = \bigcup_{n=0}^{\infty} E \cap (B_n \setminus \frac{1}{2}B_{n-1}) \subseteq F$.

Proof of (ii). For any n, \tilde{F}_n is regular with some constant $\tilde{C}_F(A, C_E, k)$ independent of n. Lower regularity of F with constant \tilde{C}_F follows immediately, so we only need to show that F is upper regular. Let $x \in \tilde{F}_n$ for some n. Observe that, for $j \ge 2$

$$d(x, \tilde{F}_{n+j}) \ge d\left(\tilde{F}_n, \frac{1}{4}B_{n+j-1}\right) \ge \frac{1}{4}A^{n+j-1} - 2A^n > A^{n+j-2}$$

provided we choose A sufficiently large. Hence, if $R \leq A^{n-2}$, then $B(x, R) \cap \tilde{F}_j = \emptyset$ for $|n-j| \geq 2$. In this case,

$$\mathcal{H}^k(B(x,R)\cap F) = \sum_{j=-1}^1 \mathcal{H}^k(B(x,R)\cap \tilde{F}_{n+j}) \lesssim_{\tilde{C}_F} R^k$$

independent of n because \tilde{F}_{n+j} is regular with constant independent of n. Now, suppose $A^j < R \leq A^{j+1}$ for $j \geq n-2$. We can write

$$\mathcal{H}^{k}(B(x,R)\cap F) = \sum_{i=0}^{j+2} \mathcal{H}^{k}(B(x,R)\cap \tilde{F}_{i}) \le \sum_{i=0}^{j+2} \mathcal{H}^{k}(\tilde{F}_{i}) \le \tilde{C}_{F} \sum_{i=0}^{j+2} \operatorname{diam}(\tilde{F}_{i})^{k} \le \tilde{C}_{F} \sum_{i=0}^{j+2} (4A)^{ik} \le 2\tilde{C}_{F} (4A)^{(j+2)k} \le (4A)^{2k+1} \tilde{C}_{F} (4R)^{k}.$$

This proves upper regularity and finishes the proof of (ii). From now on, let $C_F = C_F(C_E, A, k)$ be the regularity constant for F.

Proof of (iii) Let $x \in \tilde{F}_n$ and R > 0. Suppose first that $0 < R \le A^{n+2}$. Because $\tilde{F}_n \in BP(\mathcal{F})$ by the lemma with constant $\tilde{\theta}_F(\theta_E, A, k)$ independent of n, we get the existence of a set $G_{x,R} \in \mathcal{F}$ such that

$$\mathcal{H}^{k}(B(x,R)\cap F\cap G_{x,R}) \geq \mathcal{H}^{k}(B(x,R)\cap \dot{F}_{n}\cap G_{x,R}) \gtrsim_{\tilde{\theta}_{F},A} \mathcal{H}^{k}(B(x,R)\cap \dot{F}_{n})$$
$$\gtrsim_{C_{F}} R^{k} \gtrsim_{C_{F}} \mathcal{H}^{k}(B(x,R)\cap F).$$

using the fact that \tilde{F}_n is regular. Now, suppose $A^j < R \leq A^{j+1}$ for $j \geq n+2$. Because $x \in \tilde{F}_n$, $\frac{1}{4}A^{n-1} \leq d(x, x_0) \leq 2A^n$ so that

$$\tilde{F}_{j-2} \subseteq B\left(x_0, 2A^{j-2}\right) \subseteq B\left(x, 2A^{j-2} + 2A^n\right) \subseteq B(x, A^{j-1}) \subseteq B(x, R).$$

Using the above containment and the fact that $\tilde{F}_{j-2} \in BP(\mathcal{F})$, there exists a set $G_{x,R} \in \mathcal{F}$ with both $G_{x,R} \subseteq B(x,R)$ and

$$\mathcal{H}^k(G_{x,R} \cap \tilde{F}_{j-2}) \gtrsim_{\tilde{\theta}_F} \operatorname{diam}(\tilde{F}_{j-2})^k \gtrsim A^{(j-2)k} \gtrsim_{A,k} R^k.$$

Hence, we have $\mathcal{H}^k(B(x,R) \cap G_{x,R} \cap F) \gtrsim_{\tilde{\theta}_F,A,k} R^k \gtrsim_{C_F} \mathcal{H}^k(B(x,R) \cap F)$ as desired.

REFERENCES

- J. Azzam and R Schul. Hard Sard: quantitative implicit function and extension theorems for Lipschitz maps. *Geom. Funct. Anal.*, 22(5):1062–1123, 2012.
- [2] Simon Bortz, John Hoffman, Steve Hofmann, José Luis Luna Garcia, and Kaj Nyström. Coronizations and big pieces in metric spaces. arXiv preprint arXiv:2008.11544, 2020.
- [3] M. Christ. A *T*(*b*) theorem with remarks on analytic capacity and the Cauchy integral. *Colloq. Math.*, 60/61(2):601–628, 1990.
- [4] G. David. Wavelets and singular integrals on curves and surfaces, volume 1465 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1991.
- [5] G. David and S. Semmes. Singular integrals and rectifiable sets in \mathbb{R}^n : Beyond Lipschitz graphs. *Astérisque*, (193), 1991.
- [6] G. David and S. Semmes. Analysis of and on uniformly rectifiable sets, volume 38 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1993.

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651 *Email address*: Jared.Krandel@stonybrook.edu

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651 *Email address*: schul@math.sunysb.edu