

RECURRENCE OF MULTIPLES OF COMPOSITION OPERATORS ON WEIGHTED DIRICHLET SPACES

NOUREDDINE KARIM, OTMANE BENCHIHEB, AND MOHAMED AMOUCHE

ABSTRACT. A bounded linear operator T acting on a Hilbert space \mathcal{H} is said to be recurrent if for every non-empty open subset $U \subset \mathcal{H}$ there is an integer n such that $T^n(U) \cap U \neq \emptyset$. In this paper, we completely characterize the recurrence of scalar multiples of composition operators, induced by linear fractional self maps of the unit disk, acting on weighted Dirichlet spaces S_ν ; in particular on the Bergman space, the Hardy space, and the Dirichlet space. Consequently, we complete a previous work of Costakis et al. [19] on recurrence of linear fractional composition operators on Hardy space. In this manner, we determine the triples $(\lambda, \nu, \phi) \in \mathbb{C} \times \mathbb{R} \times LFM(\mathbb{D})$ for which the scalar multiple of composition operator λC_ϕ acting on S_ν fails to be recurrent.

1. INTRODUCTION AND PRELIMINARIES

Throughout this paper, \mathbb{C} will represent the complex plane, \mathbb{C}^* the punctured plane $\mathbb{C} \setminus \{0\}$, and $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ will be the one-point compactification of \mathbb{C} . Moreover, \mathbb{D} will stand for the open unit disk of \mathbb{C} , while \mathbb{T} will represent the unit circle of \mathbb{C} .

A bounded linear operator T acting on a Hilbert space \mathcal{H} is said to be hypercyclic if there is a vector $f \in \mathcal{H}$ whose T -orbit;

$$\mathcal{O}(f, T) = \{T^n f; n \in \mathbb{N}\},$$

is dense in \mathcal{H} . In such a case the vector f is called a hypercyclic vector. An operator T is said to be cyclic if there is a vector $f \in \mathcal{H}$ such that the space generated by $\mathcal{O}(f, T)$;

$$\text{span}(\mathcal{O}(f, T)) = \{p(T)f; p \text{ a polynomial}\},$$

is dense in \mathcal{H} . In this case the vector f is called a cyclic vector. A strong form of cyclicity and weaker than hypercyclicity is supercyclicity. An operator T is said to be supercyclic if there is a vector $f \in \mathcal{H}$ whose projective orbit;

$$\mathbb{C} \cdot \mathcal{O}(f, T) = \{\lambda T^n f; n \in \mathbb{N} \text{ and } \lambda \in \mathbb{C}\},$$

is dense in \mathcal{H} . The vector f is called a supercyclic vector.

Recall [18], that an operator T on a Hilbert space \mathcal{H} is hypercyclic if and only if it is topologically transitive; that is, for any pair of non-empty open subsets U, V of \mathcal{H} there exists some $n \in \mathbb{N}$ such that

$$T^n(U) \cap V \neq \emptyset.$$

Recently, the notions of hypercyclic and transitivity has been generalized and studied see [4, 3, 1, 2]. Another important concept in topological dynamics is that of recurrence. This notion has been initiated by Poincaré and Birkhoff, while a systematic study was given by Costakis et al. in [19]. A bounded linear operator on a Hilbert space \mathcal{H} is called recurrent if for every non-empty open subset $U \subset \mathcal{H}$ there is a positive integer n such that

$$T^n(U) \cap U \neq \emptyset.$$

2010 *Mathematics Subject Classification.* Primarily 47A16, 37B20, Secondly 46E50, 46T25.
Key words and phrases. Hypercyclicity, Recurrence, Composition operator, Dirichlet spaces.

A vector $f \in \mathcal{H}$ is said to be recurrent for T if there exists a strictly increasing sequence of positive integers $(n_k)_{k \in \mathbb{N}}$ such that

$$T^{n_k} f \rightarrow f, \text{ as } k \rightarrow \infty.$$

In that sense, every hypercyclic operator is recurrent, and every hypercyclic vector is recurrent. For more information on linear dynamics we refer to [6] and [5].

The study of linear dynamics has become a very active area of research. This work will be devoted to studying the recurrence of composition operators. Recall that if \mathcal{H} is a Hilbert space of analytic functions in the unit disk \mathbb{D} , and if ϕ is a nonconstant self map of \mathbb{D} , then the composition operator C_ϕ associated to ϕ on \mathcal{H} is defined by

$$C_\phi f = f \circ \phi \text{ for all } f \in \mathcal{H}.$$

In which case, the function ϕ called a symbol of C_ϕ . For general references on the theory of composition operators, see, e.g., Cowen and MacCluer's book [20], Shapiro's book [8] and K. H. Zhu's book [7]. The special about the composition operator is that the properties of C_ϕ depend significantly on the behaviour of the symbol ϕ . In this paper, we show that the recurrence of the composition operator is influenced by the location of the fixed points of its symbol.

For each real number ν the weighted Dirichlet space \mathcal{S}_ν is the space of functions $f(z) = \sum_{n=0}^{\infty} a_n z^n$ analytic on \mathbb{D} such that the following norm

$$(1.1) \quad \|f\|_\nu^2 = \sum_{n=0}^{\infty} |a_n|^2 (n+1)^{2\nu}$$

is finite. Endowed with the inner product

$$\left\langle \sum_{n=0}^{\infty} a_n z^n, \sum_{n=0}^{\infty} b_n z^n \right\rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n (n+1)^{2\nu},$$

the spaces \mathcal{S}_ν are Hilbert spaces, see [20, p. 16] or [13, p. 1]. For some values of ν the spaces \mathcal{S}_ν are very well known classical analytic function spaces: for instance if $\nu = 1/2$, \mathcal{S}_ν is the Dirichlet space \mathcal{D} ; for $\nu = 0$ it is the Hardy space H^2 and for $\nu = -1/2$ it is the Bergman space \mathcal{A}^2 . Observe that if $\nu_1 > \nu_2$, then the space \mathcal{S}_{ν_1} is strictly contained in \mathcal{S}_{ν_2} , and that $\|f\|_{\nu_2} \leq \|f\|_{\nu_1}$, for every $f \in \mathcal{S}_{\nu_1}$.

Also, we can define the Dirichlet space as the collection of functions analytic on \mathbb{D} whose first derivatives have square integrable modulus over \mathbb{D} . For $f \in \mathcal{D}$ the norm in \mathcal{D} has the representation

$$\|f\|_{\mathcal{D}}^2 = |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 dA(z),$$

where here $dA(z)$ is the Lebesgue area measure on \mathbb{D} normalized to have unit mass. In the Hardy space H^2 there is also an integral representation of the norm. This representation is the following

$$\|f\|_{H^2}^2 = \frac{1}{2\pi} \sup_{0 < r < 1} \int_{-\pi}^{\pi} |f(re^{i\theta})|^2 d\theta.$$

The Dirichlet space and the Hardy space will play a central role in our study.

Let $\nu \in \mathbb{R}$, we define the function on \mathbb{D} as:

$$k(z) = \sum_{n=0}^{\infty} \frac{z^n}{(n+1)^{2\nu}},$$

which is analytic on \mathbb{D} . Then for each $w \in \mathbb{D}$, the reproducing kernel is defined by

$$K_w(z) = k(\bar{w}z).$$

This is easily seen that $\|K_w\|^2 = k(|w|^2)$ and for every $f(z) = \sum_{n=0}^{\infty} a_n z^n \in S_\nu$ we have that

$$\langle f, K_w \rangle = \sum_{n=0}^{\infty} a_n w^n = f(w).$$

It is known according to a result of P.R. Hurst [9] that the composition operator C_ϕ is always bounded on S_ν when ϕ is a linear fractional map on \mathbb{D} .

Recall (see, e.g., [22, Chapter 3], [8, Chapter 0]) that linear fractional maps are those maps of the form

$$\phi(z) = \frac{az + b}{cz + d},$$

where a, b, c and d are complex numbers satisfying $ad - bc \neq 0$. They extend to the extended complex plane $\hat{\mathbb{C}}$ by defining $\phi(\infty) = a/c$, and $\phi(-d/c) = \infty$ if $c \neq 0$, while $\phi(\infty) = \infty$ if $c = 0$. The linear fractional maps can be classified according to their fixed point, which are at most two fixed points if ϕ is not the identity. Two linear fractional maps ϕ and ψ are called conjugate if there is another linear fractional map T such that

$$\phi = T^{-1}\psi T.$$

It is called parabolic if ϕ has only one fixed point α or, equivalently, it is conjugate to a translation $\psi(z) = z + \tau$, where $\tau \neq 0$. If ϕ has two distinct fixed points α and β , then ϕ is conjugate to a dilation $\psi(z) = \mu z$. In this case, the linear fractional map ϕ is called elliptic if $|\mu| = 1$; hyperbolic if $\mu > 0$ and loxodromic otherwise, see [22] for more details. The chain rule proves that the value of the derivative at the fixed point is preserved under conjugation. Therefore, the derivative of a parabolic linear fractional map at its fixed point is 1, while the derivative of a hyperbolic one is strictly less than 1 at its attractive fixed point and greater than 1 at its repulsive point. We will note $LFM(\mathbb{D})$ to refer to the set of all such maps, which are additionally self-maps of the unit disk \mathbb{D} . It is known that if $\phi \in LFM(\mathbb{D})$, then the derivative $\phi'(\eta)$ exists and is finite for every η in the unit circle \mathbb{T} . The condition $\phi(\mathbb{D}) \subset \mathbb{D}$ make some exigences on the location of the fixed points of ϕ . We have that (see [8, p. 5])

- (1) If ϕ is parabolic, then its fixed point η is in \mathbb{T} and it satisfies $\phi'(\eta) = 1$.
- (2) If ϕ is hyperbolic, the attractive fixed point is in $\bar{\mathbb{D}}$ and the other fixed point outside of \mathbb{D} and both fixed points are on \mathbb{T} if and only if ϕ is an automorphism of \mathbb{D} .
- (3) If ϕ is loxodromic or elliptic, one fixed point is in \mathbb{D} and the other fixed point outside of $\bar{\mathbb{D}}$. The elliptic ones are always automorphisms of \mathbb{D} . The fixed point in \mathbb{D} for the loxodromic ones is attractive.

Concerning the dynamics of composition operator on Dirichlet spaces, many works have been realized:

In her dissertation [14], Nina Zorboska studied composition operators induced by non-elliptic disk automorphism and proved that they are all cyclic on the Hardy space. In addition, in [15] she has obtained some results about cyclicity and hypercyclicity on the so-called smooth weighted Hardy spaces, which are the weighted Hardy spaces whose functions have continuous first derivatives on the boundary of the unit disk (basically weighted Hardy spaces strictly smaller than $\mathcal{S}_{3/2}$).

In [16, 17] Bourdon and Shapiro thoroughly characterized the cyclicity, the supercyclicity, and the hypercyclicity of linear fractional composition operators on the Hardy space $H^2(\mathbb{D})$ (see Table I). Moreover, in [17] they gave a program of transferring the cyclic and hypercyclic properties of linear fractional composition operators to general composition operators on the Hardy space.

In [13] answering some open questions posed by Zorboska [15], Gallardo-Gutiérrez and Montes-Rodríguez obtained a complete characterization of the cyclic and hypercyclic composition operators induced by linear fractional maps on weighted Dirichlet spaces \mathcal{S}_ν (see Table II).

In the present work, we give a complete characterization of recurrence of scalar multiples of composition operator whose symbols are linear fractional maps, acting on weighted Dirichlet spaces \mathcal{S}_ν ; in particular, on the Bergman space $\mathcal{S}_{-1/2}$, the Hardy space \mathcal{S}_0 and the classical Dirichlet space $\mathcal{S}_{1/2}$. To do that, spectral technics will play a significant role.

The paper is organized as follows: In the third section, we characterize the recurrence of linear fractional composition operator C_ϕ on \mathcal{S}_ν by providing a necessary and sufficient condition on ϕ and the real number ν .

The fourth section is devoted to studying the recurrence of scalar multiples of linear fractional composition operators λC_ϕ on the weighted Dirichlet spaces by providing a necessary and sufficient condition on ϕ , the real ν , and the complex scalar λ .

In tables I and II, we expose the classification of Gallardo-Gutiérrez and Montes-Rodríguez of the dynamics of composition operators and their scalar multiples on \mathcal{S}_ν . In table III, we summarize our results about the recurrence of linear fractional composition operator on \mathcal{S}_ν . Finally, in table IV, we give a summarize of the recurrence of scalar multiples of linear fractional composition operator on \mathcal{S}_ν .

Type of ϕ		Cyclic	Supercyclic	Hypercyclic
Hyperbolic	Auto-morphism	$\nu < 1/2$	$\nu < 1/2$	$\nu < 1/2$
Parabolic	Auto-morphism	$\nu < 1/2$	$\nu < 1/2$	$\nu < 1/2$
Hyperbolic	Non-Automorphism	Always	$\nu \leq 1/2$	$\nu < 1/2$
Parabolic	Non-Automorphism	$\nu \leq 3/2$	Never	Never
Interior and Exterior		Always	Never	Never
Interior and Boundary		Never	Never	Never
Elliptic rotation	Irrational	Always	Never	Never
Elliptic rotation	Rational	Never	Never	Never

TABLE I. Cyclic properties of C_ϕ on \mathcal{S}_ν , ϕ linear fractional map.

Type of ϕ	Hypercyclicity
Hyperbolic Auto-morphism	$\nu < 1/2$ and $\phi'(\eta)^{(1-2\nu)/2} < \lambda < \phi'(\eta)^{(2\nu-1)/2}$
Parabolic automorphism	$\nu < 1/2$ and $ \lambda = 1$
Hyperbolic Non-Automorphism	$\nu \leq 1/2$ and $ \lambda > \phi'(\eta)^{(1-2\nu)/2}$

TABLE II. Cyclic properties of λC_ϕ on \mathcal{S}_ν with ϕ linear fractional map and η denotes the attractive fixed point of ϕ

Our goal in the next two sections will be completing the previous tables by adding the recurrence of C_ϕ and also the recurrence of λC_ϕ on \mathcal{S}_ν . All the results about the recurrence of composition operators and its scalar multiples on \mathcal{S}_ν spaces are summarized in Table III and Table IV below.

Type of ϕ		Recurrence	Examples
Hyperbolic	Auto-morphism	$\nu < 1/2$	$\frac{3z+1}{z+3}$
Parabolic	Auto-morphism	$\nu < 1/2$	$\frac{(1+i)z-1}{z+i-1}$
Hyperbolic	Non-Automorphism	$\nu < 1/2$	$\frac{1+z}{2}$
Parabolic	Non-Automorphism	Never	$\frac{1}{2-z}$
Interior and Exterior		Never	$\frac{-z}{2+z}$
Interior and Boundary		Never	$\frac{z}{2-z}$
Elliptic rotation	Irrational	Always	$e^{i\sqrt{2}}z$
Elliptic rotation	Rational	Always	$e^{2i\pi/3}z$

TABLE III. Recurrence of C_ϕ on \mathcal{S}_ν , ϕ linear fractional map.

Type of ϕ	Recurrence
Elliptic Automorphism	$\nu \in \mathbb{R}$ and $ \lambda = 1$
Hyperbolic Automorphism	$\nu < 1/2$ and $\phi'(\eta)^{(1-2\nu)/2} < \lambda < \phi'(\eta)^{(2\nu-1)/2}$
Parabolic automorphism	$\nu < 1/2$ and $ \lambda = 1$
Hyperbolic Non-Automorphism	$\nu \leq 1/2$ and $ \lambda > \phi'(\eta)^{(1-2\nu)/2}$

TABLE IV. Recurrence of λC_ϕ on \mathcal{S}_ν , ϕ linear fractional map and η the attractive fixed point of ϕ .

For the cases not appearing in the Table IV, the operator λC_ϕ cannot be recurrent, for any $\lambda \in \mathbb{C}$.

2. RECURRENCE OF C_ϕ ON \mathcal{S}_ν

A useful tool in the study of any of the dynamic properties is the following proposition known as Comparison principle (see [8, p. 111] and [21]).

Proposition 2.1. *Let \mathcal{E} be a metric space and \mathcal{F} be a subspace that it self a linear metric space with a stronger topology. Suppose that T is a continuous linear transformation on \mathcal{E} that maps the smaller space \mathcal{F} into itself and also continuous in the topology of this space. If T is cyclic on \mathcal{F} , then it is also cyclic on \mathcal{E} and has an \mathcal{E} -cyclic vector that belongs to \mathcal{F} . Furthermore, the same is true for supercyclic and hypercyclic operators.*

Remark 2.2. Observe that the comparison principle is also true for recurrent operators. If $\nu_1 > \nu_2$, we have that \mathcal{S}_{ν_1} is a dense subspace of the space \mathcal{S}_{ν_2} . Thus we can apply the comparison principle in the sense that, if an operator T acting on \mathcal{S}_{ν_1} is recurrent, then it is recurrent on \mathcal{S}_{ν_2} .

Theorem 2.3. *Let ϕ be a linear fractional map on \mathbb{D} with an interior fixed point $p \in \mathbb{D}$. Then C_ϕ is recurrent on any of the \mathcal{S}_ν spaces if and only if ϕ is an elliptic automorphism of \mathbb{D} .*

Proof. If ϕ is an elliptic automorphism then ϕ is conjugate to a rotation (see [8, Chapter 0]). Thus, there exists a linear fractional map composition operator S and a complex number $\lambda \in \mathbb{T}$ such that $C_\phi = S^{-1}C_{\phi_\lambda}S$, where $\phi_\lambda(z) = \lambda z$, $z \in \mathbb{D}$. In order to prove that C_ϕ is recurrent, we need to prove that C_{ϕ_λ} is recurrent. Since $\lambda \in \mathbb{T}$, there exists $(n_k)_{k \in \mathbb{N}}$ such that $\lambda^{n_k} \rightarrow 1$. For any $f(z) = \sum_{m \geq 0} a_m z^m \in \mathcal{S}_\nu$ we have that

$$\|C_{\phi_\lambda}^{n_k}(f) - f\|_{\mathcal{S}_\nu} = \sum_{m \geq 0} |a_m|^2 |\lambda^{mn_k} - 1|^2 (m+1)^{2\nu} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

Thus, every $f \in \mathcal{S}_\nu$ is a recurrent vector for C_{ϕ_λ} , and so C_{ϕ_λ} is recurrent on \mathcal{S}_ν . Now, if ϕ is not an elliptic automorphism then the Denjoy-Wolff Iteration Theorem, [8, Proposition 1, Chapter 5], implies that $\phi_n(z)$ converges to p , for every $z \in \mathbb{D}$. Hence, if f is a recurrent function of C_ϕ on \mathcal{S}_ν , then there exists a sequence of positive integers $(n_k)_{k \in \mathbb{N}}$, such that $f \circ \phi_{n_k} \rightarrow f$ in \mathcal{S}_ν . Thus, by the continuity of point evaluation functional on \mathcal{S}_ν , we have that for each $z \in \mathbb{D}$:

$$f(z) = \lim_{k \rightarrow \infty} f(\phi_{n_k}(z)) = f(p).$$

We conclude that the only recurrent vectors of C_ϕ are the constant functions. \square

Theorem 2.4. *Let ϕ be a parabolic automorphism of the unit disk. Then C_ϕ is recurrent on \mathcal{S}_ν if and only if $\nu < 1/2$.*

Proof. First, if $\nu < 1/2$, then C_ϕ is hypercyclic, hence it is recurrent. Now we shall prove that this condition is necessary. Let us start by the case $\nu = 1/2$. Since ϕ is parabolic, it has a fixed point on the unit circle. Without loss of generality we may assume this fixed point is 1. Suppose that C_ϕ is recurrent, and let f be a recurrent vector for C_ϕ . Then there exists a sequence $(n_k)_{k \in \mathbb{N}}$ of positive integers such that

$$C_{\phi_{n_k}}(f) \rightarrow f,$$

in \mathcal{D} . Since ϕ has no fixed point in \mathbb{D} , by Denjoy-Wolff Iteration Theorem, the fixed point 1 is actually attractive. Thus, ϕ_{n_k} converge to 1 uniformly on compact subsets of \mathbb{D} ; in particular, $\phi_{n_k}(z)$ converge to 1, for every $z \in \mathbb{D}$. Therefore we have that

$$\|C_{\phi_{n_k}}(f) - f(1)\|_{\mathcal{D}}^2 = \|f \circ \phi_{n_k} - f(1)\|_{\mathcal{D}}^2 = |f \circ \phi_{n_k}(0) - f(1)|^2 + \int_{\mathbb{D}} |f'(\phi_{n_k}(z))|^2 |\phi'_{n_k}(z)|^2 dA(z) \rightarrow 0,$$

when $k \rightarrow \infty$. Thus, all recurrent vectors for C_ϕ are constants, and so C_ϕ cannot be recurrent in this case. Therefore, C_ϕ cannot be recurrent on \mathcal{D} . Hence, neither on any \mathcal{S}_ν with $\nu > 1/2$, by Comparison principle. \square

Theorem 2.5. *Let ϕ be a hyperbolic automorphism of the unit disk. Then C_ϕ is recurrent on \mathcal{S}_ν if and only if $\nu < 1/2$.*

Proof. If $\nu < 1/2$ then the operator C_ϕ is recurrent. For $\nu \geq 1/2$, the non-recurrence of C_ϕ follows exactly as in the case of the parabolic automorphism. \square

Now in the sequel let \mathcal{S}_ν^0 be the space of functions obtained from the space \mathcal{S}_ν by identifying functions that differ by a constant; that is $f \in \mathcal{S}_\nu^0$ if and only if there is $g \in \mathcal{S}_\nu$ such that $f - g$ is constant. Then the space \mathcal{S}_ν^0 is invariant under the operator C_ϕ , and so we can consider \tilde{C}_ϕ the restriction of C_ϕ on \mathcal{S}_ν^0 .

Lemma 2.6. *Let ϕ be an analytic self-map of the unit disk. If C_ϕ is recurrent on \mathcal{S}_ν , then so is \tilde{C}_ϕ on \mathcal{S}_ν^0 .*

Proof. We have that for every $f \in \mathcal{S}_\nu$ and $k \in \mathbb{N}$,

$$\begin{aligned} \|C_{\phi_k}f - f\|_{\mathcal{S}_\nu} &\geq \|C_{\phi_k}f - f + f(0) - f(\phi_k(0))\|_{\mathcal{S}_\nu} \\ &= \|\tilde{C}_{\phi_k}f - f\|_{\mathcal{S}_\nu^0}. \end{aligned}$$

Thus, $\text{Rec}(\tilde{C}_\phi) = \text{Rec}(C_\phi) \cap \mathcal{S}_\nu^0$. Assume that C_ϕ is recurrent, then $\text{Rec}(C_\phi)$ is dense in \mathcal{S}_ν , which implies that $\text{Rec}(\tilde{C}_\phi)$ is dense in \mathcal{S}_ν^0 . \square

Theorem 2.7. *Let ϕ be a linear fractional map on \mathbb{D} without interior fixed point. If ϕ is hyperbolic non-automorphism, then C_ϕ is recurrent on \mathcal{S}_ν if and only if $\nu < 1/2$.*

Proof. If $\nu < 1/2$ then C_ϕ is hypercyclic, thus it is recurrent. Now, suppose that $\nu > 1/2$. Since ϕ has no interior fixed point, ϕ must have an attractive fixed point $p \in \mathbb{T}$. Thus, ϕ is hyperbolic non-automorphism with attractive fixed point $p \in \mathbb{T}$. Hence, the spectrum of C_ϕ on \mathcal{S}_ν is given by:

$$\sigma(C_\phi) = \{\alpha \in \mathbb{C}; |\alpha| \leq \phi'(p)^{-\gamma}\} \cup \{\phi'(p)^k; k = 0, 1, \dots\},$$

where $\phi'(p)$ is the angular derivation of ϕ on p and $\gamma = (1 - 2\nu)/2$, see [9]. If $\nu > 1/2$, then $\gamma < 0$. In this case the singleton $\{\phi'(p)\}$ is a component of the spectrum which doesn't intersect \mathbb{T} , and so by [19, Proposition 2.11], C_ϕ cannot be recurrent. Now, suppose that $\nu = 1/2$, that is, \mathcal{S}_ν is the Dirichlet space \mathcal{D} . If C_ϕ is recurrent on the Dirichlet space, then, by Lemma 2.6, so is \tilde{C}_ϕ on $\mathcal{D}_0 := \mathcal{S}_{1/2}^0$. Since ϕ is not an automorphism, $\phi(\mathbb{D}) \subsetneq \mathbb{D}$, and the recurrence of C_ϕ implies that ϕ is injective, then there exists a non-empty open $U \subsetneq \mathbb{D}$, such that $\phi(\mathbb{D}) \cap U = \emptyset$. Hence, on \mathcal{D}_0 we have

$$\int_{\mathbb{D}} |f'(\phi(z))|^2 |\phi'(z)|^2 dA(z) = \int_{\phi(\mathbb{D})} |f'(z)|^2 dA(z) \leq \int_{\phi(\mathbb{D})} |f'(z)|^2 dA(z) + \int_U |f'(z)|^2 dA(z) < \int_{\mathbb{D}} |f'(z)|^2 dA(z).$$

Thus, $\|C_\phi\|_{\mathcal{D}_0} < 1$, and therefore, \tilde{C}_ϕ cannot be recurrent on \mathcal{D}_0 . \square

Theorem 2.8. *Let ϕ be a parabolic non-automorphism of the unit disk. Then C_ϕ is never recurrent in any of the weighted Dirichlet spaces \mathcal{S}_ν .*

Proof. Since ϕ is parabolic non-automorphism, the spectrum of C_ϕ on \mathcal{S}_ν is given by (see [23])

$$\sigma(C_\phi) = \{e^{-\alpha t}; t \geq 0\} \cup \{0\},$$

thus $\{0\}$ is a component of the spectrum which does not intersect the unit circle. Hence, by [19, Proposition 2.11], the operator C_ϕ cannot be recurrent. \square

3. RECURRENCE OF λC_ϕ ON \mathcal{S}_ν

Theorem 3.1. *Let ϕ be a holomorphic self map of \mathbb{D} with an interior fixed point. Then the operator λC_ϕ is recurrent on \mathcal{S}_ν if and only if ϕ is an elliptic automorphism and $\lambda \in \mathbb{T}$.*

Proof. Assume that ϕ is an elliptic automorphism and $|\lambda| = 1$. Then C_ϕ is recurrent by Theorem 2.3, and since $|\lambda| = 1$, then λC_ϕ is recurrent. If ϕ is not an elliptic automorphism, then C_ϕ is not recurrent, which implies that λC_ϕ is not recurrent as well. Now, if $|\lambda| \neq 1$ and ϕ is not an elliptic automorphism. Let $p \in \mathbb{D}$ be the fixed point of ϕ . Suppose that λC_ϕ is recurrent and let f be a recurrent vector. Then there exists a sequence $(n_k)_{k \in \mathbb{N}}$ of positive integers such that

$$\lambda^{n_k} f \circ \phi_{n_k} \rightarrow f,$$

in \mathcal{S}_ν . Thus,

$$f(p) = \lim_{k \rightarrow \infty} \lambda^{n_k} (C_{\phi_{n_k}} f)(p) = \lim_{k \rightarrow \infty} \lambda^{n_k} f(\phi_{n_k}(p)).$$

Now, observe that $f(\phi_{n_k}(p)) \rightarrow f(p)$ as $k \rightarrow \infty$. Therefore, if $|\lambda| < 1$, then $f(p) = 0$ for every $z \in \mathbb{D}$. If $|\lambda| > 1$, then $f(p)$ is not even defined, unless $f(p) = 0$. But $f(p)$ cannot be zero for every recurrent vector, because the set of recurrent vectors is a dense subset. \square

Theorem 3.2. *Let ϕ be a hyperbolic non-automorphism and η its boundary fixed point. Then λC_ϕ is recurrent on \mathcal{S}_ν if and only if $|\lambda| > \phi'(\eta)^{(1-2\nu)/2}$ and $\nu \leq 1/2$. In particular, we have: For the Dirichlet space ($\nu = 1/2$) the operator λC_ϕ is recurrent if and only if $|\lambda| > 1$.*

To prove Theorem 3.2, we need the following lemma and proposition:

Lemma 3.3. *Let T be a bounded operator on a separable Hilbert space \mathcal{H} . Suppose that for each non zero vector $g \in \mathcal{H}$ such that the set of complex numbers*

$$A := \{\langle Tf, g \rangle; f \in \mathcal{H}\}$$

is not dense in \mathbb{C} . Then T is not recurrent.

Proof. Every non zero vector $g \in \mathcal{H}$ defines a linear functional $\psi : f \rightarrow \langle f, g \rangle$ whose range is \mathbb{C} . Suppose that T is recurrent, then its range is dense in \mathcal{H} . Thus,

$$\mathbb{C} = \psi(\mathcal{H}) = \psi(\overline{\text{Im}(T)}) \subset \overline{\psi(\text{Im}(T))} = \overline{A},$$

which implies that the set A is dense in \mathbb{C} , a contradiction. \square

Proposition 3.4. [13, Proposition 2.16] *If $\nu < 1/2$, then for each $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathcal{S}_\nu$ there is a constant $C > 0$ such that*

$$|f'(z)| \leq \frac{C}{(1 - |z|^2)^{(3-2\nu)/2}} \quad (z \in \mathbb{D}).$$

Now we prove Theorem 3.2.

Proof of Theorem 3.2. First, if $\nu \leq 1/2$ and $|\lambda| > \phi'(\eta)^{(1-2\nu)/2}$, then C_ϕ is hypercyclic on \mathcal{S}_ν , thus it is recurrent on \mathcal{S}_ν . Now we prove that the conditions are necessary. Since the recurrence is invariant under similarity, we may suppose that the boundary fixed point is 1.

Suppose that $\nu > 1/2$. Then the reproducing kernel at 1

$$K_1(z) = \sum_{n=0}^{\infty} \frac{z^n}{(n+1)^{2\nu}}$$

is in \mathcal{S}_ν . Furthermore, for any $f \in \mathcal{S}_\nu$ we have

$$\langle \bar{\lambda} C_\phi^* K_1, f \rangle = \bar{\lambda} \langle K_1, C_\phi f \rangle = \bar{\lambda} \langle K_1, f \circ \phi \rangle = \bar{\lambda} f(\phi(1)) = \bar{\lambda} f(1) = \langle \bar{\lambda} K_1, f \rangle.$$

Then $\bar{\lambda}$ is an eigenvalue of $\bar{\lambda} C_\phi^*$. Thus, if $|\lambda| \neq 1$, the operator λC_ϕ is not recurrent, since the point spectrum of its adjoint $\bar{\lambda} C_\phi^*$, $\sigma_p(\bar{\lambda} C_\phi^*)$ must be in the unit circle by [19, Proposition 2.15]. If $|\lambda| = 1$, also in this case λC_ϕ is not recurrent, since C_ϕ is not recurrent on \mathcal{S}_ν .

Now, if $\nu = 1/2$, in this case \mathcal{S}_ν will be the Dirichlet space \mathcal{D} . Assume that λC_ϕ is recurrent on \mathcal{D} , for some $|\lambda| \leq 1$, then so is $\lambda \tilde{C}_\phi$ on \mathcal{D}_0 . But on \mathcal{D}_0 we have

$$\int_{\mathbb{D}} |\lambda f'(\phi(z))|^2 |\phi'(z)|^2 dA(z) = |\lambda|^2 \int_{\phi(\mathbb{D})} |f'(z)|^2 dA(z) < \int_{\mathbb{D}} |f'(z)|^2 dA(z).$$

Thus $\|\lambda C_\phi\|_{\mathcal{D}_0} < 1$, and therefore, $\lambda \tilde{C}_\phi$ cannot be recurrent on \mathcal{D}_0 , a contradiction. It remains to prove that if $|\lambda| \leq \mu^{(1-2\nu)/2}$ and $\nu < 1/2$, then the λC_ϕ cannot be recurrent on \mathcal{S}_ν . The growth estimate

for the derivative in Proposition 3.4 provides a constant C such that the first inequality below holds. Thus we have that

$$\begin{aligned}
 |\langle \lambda C_\phi f, z \rangle| &= 2^{2\nu} |\lambda| |(f \circ \phi)'(0)| \\
 &= 2^{2\nu} |\lambda| |f'(\phi(0))| |\phi'(0)| \\
 &= 2^{2\nu} |\lambda| |f'(1 - \mu)| \mu \\
 &\leq \frac{2^{2\nu} C |\lambda| \mu}{(1 - |1 - \mu|^2)^{(3-2\nu)/2}} \\
 &\leq \frac{2^{2\nu} C \mu^{(3-2\nu)/2}}{(2\mu - \mu^2)^{(3-2\nu)/2}} \\
 &= \frac{2^{2\nu} C}{2 - \mu}.
 \end{aligned}$$

By applying Lemma 3.3 we see that λC_ϕ is not recurrent. \square

Theorem 3.5. *Let ϕ be a parabolic automorphism of the unit disk. Then the operator λC_ϕ is recurrent on \mathcal{S}_ν if and only if $\nu < 1/2$ and $|\lambda| = 1$.*

Proof. First if $\nu < 1/2$ and $|\lambda| = 1$, then λC_ϕ is hypercyclic on \mathcal{S}_ν , which implies that it is recurrent. Now we prove that these conditions are necessary. Let us first examine the case $\nu = 1/2$. Suppose that $|\lambda| \leq 1$. Let f be a recurrent vector for λC_ϕ , then there is a sequence $(n_k)_{k \in \mathbb{N}}$ of positive integers such that $\lambda^{n_k} C_\phi^{n_k} f \rightarrow f$ in \mathcal{D} . Since ϕ is parabolic automorphism with fixed point 1. If $|\lambda| > 1$, then, by what we have just proved, $\lambda^{-1} C_{\phi^{-1}}$ is not recurrent. Now, an invertible operator is recurrent if and only if its inverse is. Consequently, λC_ϕ is not recurrent on \mathcal{D} .

Now suppose that $\nu > 1/2$, we distinguish two cases:

Case 1: If $|\lambda| \neq 1$, the reproducing kernel at 1 in \mathcal{S}_ν is an eigenvalue for the adjoint $\bar{\lambda} C_\phi^*$, a contradiction.

Case 2: If $|\lambda| = 1$, we know that C_ϕ is not recurrent on \mathcal{S}_ν , then λC_ϕ is not recurrent on \mathcal{S}_ν as well. Now suppose that $\nu < 1/2$. By Proposition the fixed point of ϕ must be on the boundary of the unit disk. We may suppose that the fixed point is 1. Let

$$\sigma(w) = \frac{i(1+w)}{1-w}, \text{ and } \psi = \sigma \circ \phi \circ \sigma^{-1}.$$

then σ is a linear fractional map of the unit disk onto the upper half plane that takes 1 to ∞ , which implies ∞ is the only fixed point of ψ , and so $\psi(w) = w + a$, where $a \neq 0$ and $\Re a = 0$. The fact that $\Re a = 0$ comes from that ϕ corresponds to an automorphism of the upper half plane. Thus ϕ satisfies the following formula

$$(3.1) \quad \phi(z) = \frac{(2-a)z + a}{-az + 2 + a} \text{ with } a \neq 0 \text{ and } \Re a = 0.$$

If $|\lambda| < 1$. By the growth estimate for the derivative's proposition, there is a constant C such that the inequality below valid

$$\begin{aligned}
|\langle \lambda C_\phi f, z \rangle| &= 2^{2\nu} |\lambda| |f'(\phi(0))| |\phi'(0)| \\
&= 2^{2\nu} |\lambda| \left| f' \left(\frac{a}{2+a} \right) \right| \frac{4}{4+|a|^2} \\
&\leq 2^{2\nu+2} C |\lambda| \left(1 - \frac{|a|^2}{4+|a|^2} \right)^{(2\nu-3)/2} \frac{1}{4+|a|^2} \\
&= \frac{2^{4\nu-1} C |\lambda|}{(4+|a|^2)^{(2\nu-1)/2}} \\
&< \frac{2^{4\nu-1} C}{(4+|a|^2)^{(2\nu-1)/2}}.
\end{aligned}$$

Thus, by Lemma 3.3, the operator λC_ϕ is not recurrent. If $|\lambda| > 1$, then $\lambda^{-1} C_{\phi_{-1}}$ is not recurrent and, therefore, neither is λC_ϕ . \square

Theorem 3.6. *Let ϕ be a hyperbolic automorphism of the unit disk and η its attractive fixed point. Then λC_ϕ is recurrent if and only if $\nu < 1/2$ and $\phi'(\eta)^{(1-2\nu)/2} < |\lambda| < \phi'(\eta)^{(2\nu-1)/2}$.*

Proof. For $\nu \geq 1/2$, the non-recurrence of λC_ϕ follows exactly as in the case of the parabolic automorphism.

Now, suppose that $\nu < 1/2$. We need an expression of ϕ . Without loss of generality, we may suppose that ϕ has -1 and 1 as its fixed points. Moreover, we may assume that 1 is the attractive fixed point. In order to compute ϕ explicitly, we use the change of variables

$$\sigma(z) = \frac{i(1-z)}{1+z}$$

that sends the unit disk onto the upper half-plane, the fixed points 1 and -1 to 0 and ∞ , respectively, and ϕ to the contraction map $\phi(w) = \mu w$, where $0 < \mu < 1$. Coming back to the unit disk we have

$$\phi(z) = \frac{(1+\mu)z + 1 - \mu}{(1-\mu)z + 1 + \mu} \text{ with } 0 < \mu < 1.$$

Observe that the derivative at the attractive fixed point is $\phi'(1) = \mu$.

Now, for any $f \in \mathcal{S}_\nu$ we have the following estimate for some constant C ,

$$\begin{aligned}
|\langle \lambda C_\phi f, z \rangle| &= 2^{2\nu} |\lambda| |f'(\phi(0))| |\phi'(0)| \\
&= 2^{2\nu} |\lambda| \left| f' \left(\frac{1-\mu}{1+\mu} \right) \right| \frac{4\mu}{(1+\mu)^2} \\
&\leq \frac{2^{2\nu} C |\lambda| \mu^{(2\nu-1)/2}}{(1+\mu)^{2\nu-1}},
\end{aligned}$$

that remain bounded for $|\lambda| \mu^{(2\nu-1)/2} \leq 1$. Therefore, if λC_ϕ is recurrent, then $|\lambda| > \mu^{(1-2\nu)/2}$. In addition, the inverse operator $\lambda^{-1} C_{\phi_{-1}}$ must also be recurrent. The attractive fixed point of ϕ_{-1} is -1 and $\phi'_{-1}(-1) = \mu$. Therefore, we must also have $|\lambda^{-1}| > \mu^{(1-2\nu)/2}$. Thus the conditions on λ are necessary for λC_ϕ to be recurrent. \square

Theorem 3.7. *Let ϕ be a parabolic non-automorphism of the unit disk. Then the operator λC_ϕ is never recurrent on any \mathcal{S}_ν .*

Proof. Since the singleton $\{0\}$ is a component of the spectrum of λC_ϕ , but every component of the spectrum of recurrent operator must intersect the unit circle, it implies that λC_ϕ is not recurrent. \square

REFERENCES

- [1] Amouch, M., Karim, N. Strong transitivity of composition operators. *Acta Math. Hungar.* 164, 458–469 (2021).
- [2] S. Shkarin, On the spectrum of frequently hypercyclic operators, *Proc. Amer. Math. Soc.*, 137 (2009), 123-134.
- [3] A. Bonilla and K.-G. Grosse-Erdmann, Upper frequent hypercyclicity and related notions, *Rev. Mat. Complut.*, 31 (2018), 673-711.
- [4] J. Bes, Q. Menet, A. Peris and Y. Puig, Recurrence properties of hypercyclic operators, *Math. Ann.*, 366 (2016), 545-5722.
- [5] F. Bayart and E. Matheron, *Dynamics of linear operators*, Cambridge University Press, Cambridge 2009.
- [6] K.-G. Grosse-Erdmann and A. Peris Manguillot, *Linear chaos*, Springer, London 2011.
- [7] K. H. Zhu, *Spaces of holomorphic functions in the unit ball*, Graduate Texts in Mathematics 226, Springer, New York, 2005.
- [8] Shapiro J. H.: *Composition operators and classical function theory*, Universitext: Tracts in Mathematics, Springer-Verlag, New York, 1993.
- [9] Hurst P. R.: Relating composition operators on different weighted Hardy spaces. *Arch. Math. (Basel)*, 68(6)1997, 503-513.
- [10] Nordgren E. A., Radjavi H. and Rosenthal P.: Composition operators and the invariant subspace problem, *C. R. Mat. Rep. Acad. Sci. Canada* 6 (1984), 279-282.
- [11] Nordgren E. A., Rosenthal P. and Wintrobe F. S.: Invertible composition operator on H^p , *J. Func. Anal.* 73 (1987), 324-344.
- [12] Shapiro J. H.: The essential norm of a composition operator, *Annals Math.* 125 (1987), 375-404.
- [13] Gallardo-Gutiérrez E. A. and Montes-Rodríguez A.: The role of the spectrum in the cyclic behavior of composition operators. *Mem. Amer. Math. Soc.*, 167(791):x+81,2004.
- [14] Zorboska N.: *Composition operators on weighted Hardy spaces*, Thesis, Univ. Toronto, 1988.
- [15] Zorboska N., Cyclic composition operators on smooth weighted Hardy spaces, *Rocky Mountain J. Math.* 29 (1999), 725-740.
- [16] P. S. Bourdon and J. H. Shapiro, Cyclic composition operators on H^2 , *Proc. Symp. Pure Math.* 51 Part 2 (1990), 43-53.
- [17] P. S. Bourdon and J. H. Shapiro, Cyclic phenomena for composition operators, *Mem. Amer. Math. Soc.* 596 (1997).
- [18] Grosse-Erdmann K.-G.: Universal families and hypercyclic operators. *Bull. Amer. Math. Soc. (N.S.)* 36 (1999), 345-381.
- [19] Costakis G., Manoussos A., Parissis I.: Recurrent Linear Operators. *Complex Anal. Oper. Theory* 8 (2014), 1601-1643.
- [20] C. Cowen and B. MacCluer, *Composition Operators on Spaces of Analytic Functions*, CRC Press, 1995.
- [21] Salas H. N.: Supercyclicity and weighted shifts, *Studia Math.* 135 (1999), 55-74.
- [22] Ahlfors L. V.: *Complex Analysis*, McGraw-Hill, New York, 1979.
- [23] Gallardo-Gutiérrez E. A. and Schroderus R.: The spectra of linear fractional composition operators on the weighted Dirichlet spaces. *Journal of Functional Analysis*, Volume 271, Issue 3, 720-745, (2016).

(Noureddine Karim, Otmane Benchiheb, and Mohamed Amouch) CHOUAIB DOUKKALI UNIVERSITY. DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE EL JADIDA, MOROCCO

Email address: noureddinekarim1894@gmail.com

Email address: otmane.benchiheb@gmail.com

Email address: amouch.m@ucd.ac.ma