A strong equivariant deformation retraction from the homeomorphism group of the projective plane to the special orthogonal group

Michael Gene Dobbins

Department of Mathematical Sciences, Binghamton University (SUNY), Binghamton, New York, USA. mdobbins@binghamton.edu

Abstract

This is the third paper in a series on oriented matroids and Grassmannians. We construct a $(O_3 \times Z_2)$ -equivariant strong deformation retraction from the homeomorphism group of the 2-sphere to O_3 , where the action of \mathbb{Z}_2 is generated by antipodal reflection acting on the right, and O_3 acts on the left by isometry. Quotienting by the antipodal map induces a $SO₃$ -equivariant strong deformation retraction from the homeomorphism group of the projective plane to $SO₃$. The same holds for subgroups of homeomorphisms that preserve the system of null sets. This confirms a conjecture of Mary-Elizabeth Hamstrom.

1 Introduction

Let us denote the multiplicative group on $\{1, -1\}$ by \mathbb{Z}_2 and denote the orthogonal and special orthogonal groups by O_n and SO_n . Let S^2 denote the 2-sphere in \mathbb{R}^3 and $\mathbf{P}^2 = \mathbf{S}^2/\mathbb{Z}_2$ denote the real projective plane. Let $hom(X)$ denote the group of homeomorphisms from a metric space X to itself with the sup-metric. That is, the distance between maps $f, g \in$ hom(X) is dist(f, g) = sup{dist(f(x), g(x)) : $x \in X$. Note that the induced metric topology is the same as the compact-open topology. Note also that if $f \in \text{hom}(S^2)$ is \mathbb{Z}_2 -equivariant, i.e. $f(-x) = -f(x)$, then there is an induced map on the projective plane, which we simply denote by f, that acts by $f(\{x, -x\}) = \{f(x), f(-x)\}\)$. Let SO₃ act on \mathbb{P}^2 in this way. Our goal is to prove the following.

Theorem 1.1. There is a strong $(O_3 \times Z_2)$ -equivariant deformation retraction from hom(S^2) to O_3 , where $(Q, s) \in O_3 \times \mathbb{Z}_2$ acts on $f \in \text{hom}(\mathbf{S}^2)$ by $(Q, s)f = Q \circ f \circ s$.

Corollary 1.2. There is a strong SO_3 -equivariant deformation retraction from hom (P^2) to SO₃, where $Q \in SO_3$ acts on $f \in \text{hom}(\mathbf{P}^2)$ by $Qf = Q \circ f$.

In 1923 Hellmuth Kneser showed that there is a strong deformation retraction from $hom(S^2)$ to O_3 [\[11\]](#page-34-0), and later Bjorn Friberg gave another more elementary proof [\[8\]](#page-34-1). Dobbins showed that a strong deformation retraction from $hom(S^2)$ to O_3 can be made to be O_3 -equivariant [\[6\]](#page-34-2). Mary-Elizabeth Hamstrom showed that $hom(P^2)$ and SO_3 have isomorphic homotopy groups. Hamstrom further conjectured that $hom(P^2)$ deformation retracts to the Lie group of SO_3 acting on \mathbf{P}^2 [\[10,](#page-34-3) p. 43], which is confirmed by Corollary [1.2.](#page-0-0)

Let hom_N(X) denote the subgroup of $f \in \text{hom}(X)$ such that for all $A \subset X$, A is a null set if and only if $f(A)$ is a null set. In other words, both f and f^{-1} have the Luzin N property. In this case we say f preserves nullity. We also prove analogous results for these groups.

Theorem 1.3. There is a strong $(O_3 \times \mathbb{Z}_2)$ -equivariant deformation retraction from $hom_N(S^2)$ to O_3 .

Corollary 1.4. There is a strong SO_3 -equivariant deformation retraction from $hom_N(P^2)$ to SO_3 .

For us, the subgroup $hom_N(S^2)$ has the advantage that for a fixed disk $D \subset S^2$, the area of $f(D)$ is continuous as a function on $f \in \text{hom}_N(S^2)$, which is not the case on hom (S^2) . See Lemma [2.8](#page-9-0) below.

1.1 Motivation from oriented matroids

Groups of automorphisms are of fundamental interest. In this case, however, the author was motivated by a conjecture from combinatorics, which in turn has applications for working with vector bundles. Some background on the conjecture that the author wrote in the previous papers in the series is repeated in this section $[5, 6]$ $[5, 6]$ $[5, 6]$. The material in this subsection is not needed to understand the rest of the paper.

An oriented matroid is a combinatorial analog to a real vector space, but where we only keep track of sign information. One way to obtain an oriented matroid is as the set of all sequences of signs of all the vectors in a vector subspace of \mathbb{R}^n . For example, the oriented matroid obtained from the space $\{(x, y) \in \mathbb{R}^2 : x + y = 0\}$ is given by the set $\{(+, -), (0, 0), (-, +)\}.$ However, oriented matroids are defined by purely combinatorial axioms, and not all oriented matroids are obtained in this way. One way the Grassmannian is defined is as the set of all k-dimensional vector subspaces of a given vector space, and this set is given a metric. Oriented matroids come equipped with analogs of dimension and subspace, and an OM-Grassmannian is a finite simplicial complex that is defined analogously from an oriented matroid. Nicolai Mnëv and Günter Ziegler conjectured that each OM-Grassmannian of a realizable oriented matroid is homotopy equivalent to the corresponding real Grassmannian [\[14,](#page-34-5) Conjecture 2.2].

Gaku Liu showed that this conjecture does not hold in general; i.e. there is an OM-Grassmannian of a realizable oriented matroid that is not homotopy equivalent to the corresponding real Grassmannian [\[13\]](#page-34-6). Although the full conjecture is false, special cases remain open, such as for MacPhersonians, which are the OM-Grassmannians of the oriented matroids corresponding to \mathbb{R}^n , and in particular for the rank 3 case, which is the analog of the Grassmannian consisting of 3-dimensional subspaces of \mathbb{R}^n . Mnëv and Ziegler had listed the rank 3 case as a theorem with the expectation that a proof would later appear in Eric Babson's Ph.D. thesis, but then it did not $[14, 2]$ $[14, 2]$ $[14, 2]$. Also, an erroneous proof that each MacPhersonian is homotopy equivalent to the corresponding Grassmannian in all ranks was published and then retracted [\[3,](#page-34-8) [4\]](#page-34-9).

Jim Lawrence showed that oriented matroids can be characterized as the combinatorial

cell decompositions of the sphere arising from essential pseudosphere arrangements [\[7\]](#page-34-10). Lawrence's theorem provides a topological model for the combinatorial axioms of oriented matroids. In the rank 3 case, these are pseudocircle arrangements, which are collections of oriented simple closed curves in the 2-sphere such that every pair of curves either coincide or intersect at exactly 2 points, in which case any third curve either separates the 2 points or passes though both points. An arrangement is said to be essential when no single point is contained in all pseudospheres.

The first paper in the series introduced spaces of weighted essential pseudosphere arrangements called pseudolinear Grassmannians, and showed that each rank 3 pseudolinear Grassmannian is homotopy equivalent to the corresponding real Grassmannian. The pseudolinear Grassmannians serve as an intermediate space between the Grassmannians and MacPhersonians, and the main results of the first paper represent a step toward showing that the conjecture holds for the rank 3 MacPhersonians by showing homotopy equivalence between this intermediate space and one side $[6]$. The second paper provided a crucial tool needed for the present paper, which will be used in Subsection $3.2 \, |5|$ $3.2 \, |5|$.

The present paper takes another step toward showing the rank 3 MacPhersonian case of the conjecture, by providing a tool for replacing pseudolinear Grassmannians with nicer spaces, namely spaces of *antipodally symmetric* weighted pseudocircle arrangements where every pseudocircle has *area 0*. The first advantage is that we can disregard the last property of pseudocircle arrangements, namely that a third curve must separate or pass though the points where two other curves intersect, since this property is already guaranteed to hold in the symmetric case. The second advantage is that the area of the cells of an arrangement vary continuously with respect to the arrangement. The present paper will not go into the precise details on how of Theorem [1.3](#page-1-0) is applied to pseudocircle arrangements.

1.2 Organization

A central idea in the proof of Theorem [1.1](#page-0-1) is to find a simple closed curve that depends continuously on the initial homeomorphism $f \in \text{hom}(\mathbf{S}^2)$ and then continuously deform that curve to a great circle. Then, each hemisphere on either side of the resulting great circle can be dealt with separately. To do so, we have to extend the deformation of the closed curve to a deformation of the homeomorphism f , which we do by using various canonical versions of the Jordan-Schoenflies theorem. For this, we use parameterizations of the sphere or of disks on the sphere that are each defined explicitly using complex analysis and the Riemann mapping theorem. In Section [2,](#page-3-0) we present these parameterizations. In Section [3,](#page-14-0) we prove Theorem [1.1.](#page-0-1) In Section [4,](#page-31-0) we modify the proof from Section [3](#page-14-0) to arrive at Theorem [1.3.](#page-1-0)

1.3 Definitions and Notation

Let **D** be the closed unit disk and S^1 be the unit circle in the complex plane. Let e_1, e_2, e_3 be the standard basis vectors in 3-space, and let $e_{-i} = -e_i$. We will use i for the imaginary unit, and we let $\overline{z} = a - bi$ be the complex conjugate of $z = a + bi$. Let R and C be the one point compactifications of the real line and the complex plane by adjoining ∞ . We denote the derivative of a function f by ∂f or by $\partial_x f(x)$ and we denote the boundary of a

topological disk C by ∂C. We will generally use square brackets around functions and round brackets around arguments passed to functions. We use this for partial function application. For example, given a function $f : W \times X \to Y$ and $w \in W$, we have $f(w) : X \to Y$ by $[f(w)](x) = f(w, x)$. We will also use square brackets when composing, inverting, and adjoining functions. For example, given $f: W \times X \to Y$, $g: Z \to Y$, and $h: Z \to W$ we write $x = [[f(w)]^{-1} \circ g](z)$ for the value x such that $f(w, x) = g(z)$, provided that $f(w) : X \to Y$ is invertible. Also, $[g, h](z) = (g(z), h(z))$. We may distinguish multiplicative inverse by using round brackets, i.e., $(x)^{-1} = \frac{1}{x}$ $\frac{1}{x}$. We let $[x \mapsto \Phi]$ denote the anonymous function that substitutes its input for the value of a variable x in the formula Φ . For example $|z \mapsto \overline{z}|$ denotes the function that returns the complex conjugate of its input. When composing f with multiplication by a constant c, we simply write $cf = [x \mapsto cf(x)]$ and $fc = [x \mapsto f(cx)]$, and similarly we write Qf and fQ for $Q \in \mathcal{O}_3$ acting on the sphere. We denote real intervals by $(a, b]_{\mathbb{R}} = \{x \in \mathbb{R} : a < x \leq b\}$ for bounds $a, b \in \mathbb{R}$ with any combination of round or square brackets for (half) open or closed intervals.

The Fréchet distance between a pair of Jordan curves is defined by

$$
dist_F(\gamma_1, \gamma_0) = \inf_{\varphi_1, \varphi_0} \sup_x \|\varphi_1(x) - \varphi_0(x)\|
$$

where the $\varphi_i : \mathbf{S}^1 \to \gamma_i$ are homeomorphisms. We will also use Hausdorff distance, which is a coarser metric than Fréchet distance. **Hausdorff distance** is defined by

$$
dist_H(\gamma_1, \gamma_0) = \inf \{ \delta : \gamma_1 \subseteq \gamma_0 \oplus \delta, \gamma_0 \subseteq \gamma_1 \oplus \delta \}.
$$

The boundary Fréchet distance between a pair of topological disks is the Fréchet distance between the boundaries of the disks. We say a curve is *rectifiable* when it has finite length and has null area when its area is 0.

We say a map is *internally conformal* when the restriction of the map to the interior of its domain is conformal. We will use the following two extensions of the Riemann mapping theorem. Carathéodory's mapping theorem says that, for every Jordan curve in \mathbb{C} , there is an internally conformal homeomorphism from the closed disk to the region bounded by the curve. Radó's theorem says that, for Jordan curves γ_k and maps h_k as in the Carathéodory's mapping theorem, if $h_k(0)$ is fixed and $\partial h_k(0) > 0$ for all k, and $\gamma_k \to \gamma_\infty$ in Fréchet distance, then $h_k \to h_\infty$ uniformly [\[16\]](#page-34-11).

2 Parameterizations

In this section we first introduce a notion of convergence called invariable convergence. We will then present several parameterizations of the sphere and disks on the sphere and their properties. These parameterizations will be used in the next section to construct a deformation retraction.

2.1 Invariable convergence

A vitally important property for us is continuity. Here we will use parameterizations of a class of topological disks, and will need these parameterizations to depend continuously on the boundary of the disk and on some specified points on the boundary. What's more, we will need the inverse map of the parameterization to depend continuously on the boundary and the specified points. In this case, uniform convergence is not applicable, since the domain of the map can change. To deal with this, we introduce a related notion of convergence called invariable convergence that is relevant to sequences of maps where the domain changes.

Let X, Y be metric spaces with metrics dist_X, dist_Y, and let $X_k \subset X$ for $k \in \{1, \ldots, \infty\}$. We say a sequence of functions $f_k : X_k \to Y$ converges *invariably* to $f_{\infty} : X_{\infty} \to Y$ when the following holds: $X_k \to X_\infty$ in Hausdorff distance, and for all $\varepsilon > 0$, there is $K(\varepsilon)$ and $\delta(\varepsilon) > 0$ such that for all $k \in \mathbb{N}$, all $x_k \in X_k$, and all $x_\infty \in X_\infty$, if $k \geq K(\varepsilon)$ and $dist_X(x_k, x_\infty) < \delta(\varepsilon)$ then $dist_Y(f_k(x_k), f_\infty(x_\infty)) < \varepsilon$. Note that K and δ cannot depend on x_k or $x_\infty.$

Lemma 2.1. Let X, Y, Z be compact metric spaces, and let $f_k : X_k \to Y_k$ and $g_k : Y_k \to Z$ be maps defined on $X_k \subset X$ and $Y_k \subset Y$ for $k \in \{1, \ldots, \infty\}.$

- 1. For $X_k = X$ fixed, $f_k \to f_\infty$ invariably if and only if $f_k \to f_\infty$ uniformly and f_∞ is continuous.
- 2. If $f_k: X_k \to Y$ is a homeomorphic embedding of X_k in Y and $f_k \to f_{\infty}$ invariably, then $f_k^{-1} \to f_{\infty}^{-1}$ invariably.
- 3. If $f_k \to f_\infty$ and $g_k \to g_\infty$ invariably, then $g_k \circ f_k \to g_\infty \circ f_\infty$ invariably.

Proof of Lemma [2.1](#page-4-0) part [1.](#page-4-1) Suppose $f_k \to f_\infty$ uniformly and f_∞ is continuous. Then, by the Heine-Cantor theorem, f_{∞} is uniformly continuous. Consider $\varepsilon > 0$. Since $f_k \to f_{\infty}$ uniformly, there is K such that for all $x \in X$ and all $k \geq K$, $dist_Y(f_k(x), f_\infty(x)) < \varepsilon/2$. Since f_{∞} is uniformly continuous, there is δ such that for all $x, x' \in X$, if $dist_X(x, x') < \delta$, then $dist_Y(f_\infty(x), f_\infty(x')) < \varepsilon/2$. Hence, for all $x_k, x_\infty \in X$, if $k \geq K$ and $dist_X(x_k, x_\infty) < \delta$, then

$$
dist_Y(f_k(x_k), f_\infty(x_\infty)) \leq dist_Y(f_k(x_k), f_\infty(x_k)) + dist_Y(f_\infty(x_k), f_\infty(x_\infty)) < \varepsilon,
$$

so $f_k \to f_\infty$ invariably.

To show the implication in the other direction, suppose $f_k \to f_\infty$ invariably. First, fix an arbitrary point $x_{\infty} \in X$ and let $x_k = x_{\infty}$ for all k. Then, the definition of invariable convergence in this case is equivariant to uniform convergence. Next, fix $\varepsilon > 0$ and $\tilde{x} \in X$ such that $dist_X(x_\infty, \tilde{x}) < \delta(\varepsilon)$ where δ is given in the definition of invariable convergence. Then, in the limit as $k \to \infty$, we have $f_k(x_k) = f_k(x_\infty) \to f_\infty(x_\infty)$, so dist $(f_\infty(x_\infty), f_\infty(\tilde{x})) < \varepsilon$, which means that f_∞ is continuous. ε , which means that f_{∞} is continuous.

Proof of Lemma [2.1](#page-4-0) part [2.](#page-4-2) Suppose for the sake of contradiction that f_k^{-1} $k⁻¹$ does not converge invariably to f_{∞}^{-1} under the hypotheses of the lemma. Then, there would be $\varepsilon > 0$ such that for all K and all $\delta > 0$, there is some $k \geq K$ and $y_k \in f(X_k)$ and $y_\infty \in f(X_\infty)$ such that $dist_Y(y_k, y_\infty) < \delta$ and $dist_X(f_k^{-1})$ $f_k^{-1}(y_k), f_\infty^{-1}(y_\infty)) \geq \varepsilon$. This means that we may assume there is some sequence x_k such that $y_k = f_k(x_k)$ converges to $y_\infty = f_\infty(x_\infty)$ and $dist_X(x_k, x_\infty) > \varepsilon$; otherwise we could restrict to a subsequence where $\delta \to 0$ and then we

could restrict to a subsequence where y_k converges, since Y is compact. We may also assume that x_k converges to a point \tilde{x} , since X is compact. Since $f_k \to f_\infty$ invariably and $x_k \to \tilde{x}$, we have $y_k = f_k(x_k) \to f_\infty(\tilde{x})$. Since $y_k \to y_\infty$, we have $f_\infty(\tilde{x}) = y_\infty$, so $\tilde{x} = f_\infty^{-1}(y_\infty) = x_\infty$, but that contradicts our choice of x_k as a sequence bounded away from x_{∞} . П

Proof of Lemma [2.1](#page-4-0) part [3.](#page-4-3) Let δ_f , δ_g and K_f , K_g be as in the definition of invariable convergence for f_k and g_k respectively. Consider $\varepsilon > 0$ and $k \ge \max(K_f(\varepsilon), K_g(\varepsilon))$ and $x \in X_k$ and $x' \in X_\infty$ such that $dist_X(x, x') < \delta_f(\delta_g(\varepsilon))$. Then, $dist_Y(f_k(x), f_\infty(x')) < \delta_g(\varepsilon)$, so $dist_Z(g_k \circ f_k(x), g_\infty \circ f_\infty(x')) < \varepsilon.$ \Box

Lemma 2.2. Let $f_k \to f_\infty : X \to Y$ be a sequence of homeomorphic embeddings of a compact metric space X in a compact metric space Y converging uniformly to a homeomorphic embedding, and let $y_k \in f_k(X)$. If $y_n \to y_\infty \in Y$, Then $f_k^{-1}(y_k) \to f_\infty^{-1}(y_\infty)$.

Proof. By Lemma [2.1](#page-4-0) part [1,](#page-4-1) $f_k \to f_\infty$ invariably, so by part [2,](#page-4-2) $f_k^{-1} \to f_\infty^{-1}$ invariably, so $f_k^{-1}(y_k) \to f_{\infty}^{-1}(y_{\infty}).$ \Box

Lemma 2.3. Let Ω , X, Y, Z be compact metric spaces spaces, and let $f : \Omega \to \text{hom}(X, Y)$ and $q : \Omega \to \text{hom}(Y, Z)$ be continuous in the sup-metric. Then, $q(\omega) \circ f(\omega)$ and $[f(\omega)]^{-1}$ depend continuously on ω in the sup-metric.

Proof. Consider $\omega_k \in \Omega$ such that $\omega_k \to \omega_\infty$. By Lemma [2.1,](#page-4-0) $f(\omega_k) \to f(\omega_\infty)$ and $g(\omega_k) \to f(\omega_\infty)$ $g(\omega_{\infty})$ converge invariably, so $g(\omega_k) \circ f(\omega_k) \to g(\omega_{\infty}) \circ f(\omega_{\infty})$ and $[f(\omega_k)]^{-1} \to [f(\omega_k)]^{-1}$ invariably, so $q(\omega) \circ f(\omega)$ and $[f(\omega)]^{-1}$ are continuous as functions of ω . \Box

2.2 A disk with four points on the boundary - F_{D4p}

Here we construct a parameterization F_{D4p} of a topological disk that depends continuously on the boundary of the disk and on 4 distinct points on the boundary.

Lemma 2.4. Let S be a simple closed curve in S^2 , and let C be one of the closed regions bounded by S, and let $p_1, p_2, p_{-1}, p_{-2} \in S$ be distinct points counter-clockwise around C as viewed from outside the sphere. Then, there is a unique internally conformal homeomorphism $f = F_{\text{D4p}}(S, p_1, p_2, p_{-1}, p_{-2}) : \mathbf{D} \to C$ such that $f(1) = p_1$, $f(-1) = p_{-1}$, and $f^{-1}(p_{-2}) =$ $-f^{-1}(p_2)$. Also, the following holds.

- 1. In the sup-metric, f depends continuously on S in Fréchet distance and p_1, p_2, p_{-1}, p_{-2} .
- 2. For $Q \in SO_3$, $F_{D4p}(Q(S, p_1, p_2, p_{-1}, p_{-2})) = Qf$.
- 3. $F_{D4p}(-(S, p_1, p_2, p_{-1}, p_{-2}); z) = -f(\overline{z}).$
- 4. $F_{\text{D4p}}(S, p_{-1}, p_{-2}, p_1, p_2; z) = f(-z)$.
- 5. $F_{D4p}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2})$ is stereographic projection through e_{-3} .

Before proving Lemma [2.4,](#page-5-0) we will define a pair of conformal automorphisms of the unit disk. Note that the conformal automorphisms of the unit disk are the linear fractional

transformations that send the disk to itself. One automorphism, $F_{3p\text{-align}}$, depends on 3 points on the boundary, and the other, $F_{4p\text{-align}}$, depends on 4 points on the boundary. We will show that these automorphisms depend continuously on the 3 or 4 points. The map F_{D4p} implied by Lemma [2.4](#page-5-0) will use the automorphism with 4 points, which will use the automorphism with 3 points.

Lemma 2.5. Let $u_1, u_1, u_{-1} \in S^1$ be distinct points in counter-clockwise order. Then, there is a unique conformal automorphism $f = F_{3p\text{-align}}(u_1, u_i, u_{-1})$ of the disk such that $f(u_n) = n$ for $n \in \{1, i, -1\}$. Moreover, f in the sup-metric depends continuously on u_1, u_i, u_{-1} .

Proof. Let

$$
f(z) = \frac{(z - u_{-1})(u_i - u_1)(i + 1) + (z - u_1)(u_i - u_{-1})(i - 1)}{(z - u_{-1})(u_i - u_1)(i + 1) - (z - u_1)(u_i - u_{-1})(i - 1)}.
$$

To verify that f has the defining properties, observe that $f(u_1) = 1$, $f(u_i) = i$, and $f(u_{-1}) =$ −1. Since linear fractional transformations send circles to circles, and 3 points determine a circle, and f respectively sends u_1, u_1, u_{-1} to $1, i, -1$, we have that f sends the unit circle $S¹$ to itself. Also, since the order of the points around the circle is the same, namely counter-clockwise, and linear fractional transformations are orientation preserving, f sends **to itself. Thus,** f **, as given by the formula above, satisfies the defining properties of** $F_{3p\text{-align}}(u_1, u_i, u_{-1})$ in the lemma.

To verify uniqueness, consider another map f_0 satisfying the hypotheses of the lemma, and let $f_1 = f_0 \circ f^{-1}$. Then, f_1 is a linear fractional transformation with fixed values 1, i, and -1 , so f_1 is the identity map, which means $f_0 = f$.

To show continuity, observe that $F_{3p\text{-align}}$ is smooth as a function of 4 variables, so the restriction of F3p-align to a compact subset of the domain is always uniformly continuous. We can always find a compact neighborhood of a point $(u_1, u_1, u_{-1}; z)$ that is a product with **D**, so $F_{3p\text{-}align}$ is continuous in the sup-metric over **D** as a function of 3 variables. \Box

Lemma 2.6. Let $u_1, u_2, u_{-1}, u_{-2} \in \mathbf{S}^1$ be distinct points in counter-clockwise order. Then, there is a conformal automorphism $f = F_{4p\text{-align}}(u_1, u_2, u_{-1}, u_{-2})$ of the disk such that $f(u_1) =$ 1, $f(u_{-1}) = -1$, and $f(u_{-2}) = -f(u_2)$. Moreover, f in the sup-metric depends continuously on u_1, u_2, u_{-1}, u_{-2} .

Proof. Let

$$
r = \frac{(u_1 - u_2)(u_{-1} - u_{-2})}{(u_1 - u_{-2})(u_{-1} - u_2)}.
$$

Note that r is the cross-ratio of $(u_1, u_{-1}; u_2, u_{-2})$. Since these 4 values appear on a common circle, r is real, and because the pair u_1, u_{-1} and u_2, u_{-2} appear in alternating order around the circle, we have $r < 0$. Let

$$
w = \frac{1 + \sqrt{r}}{1 - \sqrt{r}}
$$
 where \sqrt{r} is in the upper half-plane.

Observe that \sqrt{r} is on the upper ray of the imaginary axis by our choice of branch of the square root, so w is on the upper half of the unit circle. Let f be the linear fractional transformation that sends u_1, u_2, u_{-1} respectively to $1, w, -1$, namely

$$
f(z) = [F_{3p\text{-align}}(1, w, -1)^{-1} \circ F_{3p\text{-align}}(u_1, u_2, u_{-1})](z)
$$

=
$$
\frac{(z - u_{-1})(u_2 - u_1)(w + 1) + (z - u_1)(u_2 - u_{-1})(w - 1)}{(z - u_{-1})(u_2 - u_1)(w + 1) - (z - u_1)(u_2 - u_{-1})(w - 1)}.
$$

To verify that f has the defining properties, we can immediately observe that $f(u_1) = 1$, $f(u_2) = w$, and $f(u_{-1}) = -1$. Since u_1, u_2, u_{-1} are sent to points on the unit circle, and f is a linear fractional transformation, we have that $f(\mathbf{S}^1) = \mathbf{S}^1$. Furthermore, since u_1, u_2, u_{-1} are sent to points in the same order around the unit circle, namely counter-clockwise, we have that $f(\mathbf{D}) = \mathbf{D}$. It only remains for us to verify that $f(u_{-2}) = -w$ by computation. We have $\frac{w-1}{w+1} = r^{1/2}$ and

$$
(u_1 - u_2)(u_{-1} - u_{-2}) = r^{1/2} \sqrt{(u_1 - u_2)(u_{-1} - u_{-2})(u_1 - u_{-2})(u_{-1} - u_2)},
$$

$$
(u_1 - u_{-2})(u_{-1} - u_2) = r^{-1/2} \sqrt{(u_1 - u_2)(u_{-1} - u_{-2})(u_1 - u_{-2})(u_{-1} - u_2)},
$$

so

$$
f(u_{-2}) = \frac{(u_1 - u_2)(u_{-1} - u_{-2})(w + 1) + (u_1 - u_{-2})(u_{-1} - u_2)(w - 1)}{(u_1 - u_2)(u_{-1} - u_{-2})(w + 1) - (u_1 - u_{-2})(u_{-1} - u_2)(w - 1)}
$$

=
$$
\frac{r^{1/2}(w + 1) + r^{-1/2}(w - 1)}{r^{1/2}(w + 1) - r^{-1/2}(w - 1)} \cdot \frac{\sqrt{(u_1 - u_2)(u_{-1} - u_{-2})(u_1 - u_{-2})(u_{-1} - u_2)}}{\sqrt{(u_1 - u_2)(u_{-1} - u_{-2})(u_1 - u_{-2})(u_{-1} - u_2)}}
$$

=
$$
\frac{(w - 1) + (w + 1)}{(w - 1) - (w + 1)}
$$

=
$$
-w.
$$

Thus, f, as given by the formula above, satisfies the defining properties of $F_{4p\text{-align}}(u_1, u_2, u_{-1}, u_{-2})$ in the lemma.

To verify uniqueness, consider another map f_0 satisfying the hypotheses of the lemma, and let $f_1 = f_0 \circ f^{-1}$. Then f_1 is a linear fractional transformation that fixes 1 and -1 and sends $S¹$ to itself, so f_1 must also send $\overline{\mathbb{R}}$ to itself. Also, f_1 sends the line through $f(u_2), 0, f(u_{-2})$ to the line though $f_0(u_2), 0, f_0(u_{-2}),$ and these lines are distinct from $\overline{\mathbb{R}}$, so f_1 must send $\{0,\infty\}$ to $\{0,\infty\}$, and since f_1 sends **D** to itself, 0 must be another fixed point of f_1 , so f_1 is the identity map. Thus, $f_0 = f$.

To show continuity, observe that $F_{4p\text{-align}}$ is smooth as a function of 5 variables, so $F_{4p\text{-align}}$ is locally uniformly continuous, so $F_{4p\text{-align}}$ is continuous in the sup-metric over **D** as a function of 4 variables. \Box

Proof of Lemma [2.4.](#page-5-0) By the Riemann and Carathéodory mapping theorems, there exists an internally conformal homeomorphism g from \bf{D} to C. Fix g, and let

$$
f = g \circ (\mathbf{F}_{4p\text{-align}}(u_1, u_2, u_{-1}, u_{-2}))^{-1}
$$

where $u_n = g^{-1}(p_n)$. Observe that f satisfies the defining properties of $F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2})$ in the lemma.

To verify uniqueness, consider another map \tilde{f} satisfying the hypotheses of the lemma, and let $h = f \circ f^{-1}$. Then, $h : \mathbf{D} \to \mathbf{D}$ is an internally conformal homeomorphism such that $h(1) = 1$ and $h(-1) = -1$. By the Schwarz-Pick theorem, h is a linear fractional transformation. Also, h sends the line through $f(p_2)$, 0, $f(p_{-2})$ to the line though $f(p_2)$, 0, $f(p_{-2})$, and these lines are distinct from \mathbb{R} , so 0 must be another fixed point of h, so h is the identity map. Thus, $f=f$.

To show part [1,](#page-5-1) continuity, consider sequences S_k , $p_{1,k}$, $p_{2,k}$, $p_{-1,k}$, $p_{-2,k}$ that respectively converge to $S_{\infty}, p_{1,\infty}, p_{2,\infty}, p_{-1,\infty}, p_{-2,\infty}$ in the space of objects where the hypotheses of the lemma are satisfied, and define C_k analogously. Let $f_k = F_{\text{D4p}}(S_k, p_{1,k}, p_{2,k}, p_{-1,k}, p_{-2,k})$. Let us identify S^2 with $\overline{\mathbb{C}}$ preserving angle and orientation in such a way that $0 \in C^{\circ}_{\infty}$, $\infty \notin C_{\infty}$, $\partial f_{\infty}(0) > 0$. Then, f_k is holomorphic on the interior of **D**.

Let g_k be the internally conformal homeomorphism from **D** to C_k such that $g_k(0) = f_\infty(0)$ and $\partial g_k(0) > 0$. Since $S_k \to S_\infty$ in Fréchet distance, by Radó's theorem [\[16,](#page-34-11) Theorem 2.11], $g_k \to g_\infty$ uniformly. Let $u_{n,k} = g_k^{-1}$ $k_k^{-1}(p_{n,k})$. Then, $u_{n,k} \to u_{n,\infty}$ by Lemma [2.2,](#page-5-2) so by Lemmas [2.6](#page-6-0) and [2.1,](#page-4-0)

$$
f_k = g_k \circ (F_{4p\text{-align}}(u_{1,k}, u_{2,k}, u_{-1,k}, u_{-2,k}))^{-1} \to f_{\infty}
$$
 uniformly.

Thus, F_{D4p} is continuous in the sup-metric over **D** as a function of the 5 variables $S, p_1, p_2, p_{-1}, p_{-2}$, so part [1](#page-5-1) holds.

The map Qf is an internally conformal homeomorphism from **D** to QC that sends 1, -1 respectively to Qp_1, Qp_{-1} , and $[Qf]^{-1}(Qp_{-2}) = -[Qf]^{-1}(Qp_2)$, and since $F_{D4p}(Q(S, \ldots, p_{-2}))$ uniquely satisfies these conditions, part [2](#page-5-3) holds.

The map $h = [z \mapsto -\mathrm{F}_{\mathrm{D4p}}(S, \ldots, p_{-2}; \overline{z})] = [z \mapsto -z] \circ \mathrm{F}_{\mathrm{D4p}}(S, \ldots, p_{-2}) \circ [z \mapsto \overline{z}]$ is a homeomorphism from **D** to $-C$, and both $[z \mapsto \overline{z}]$ and $[z \mapsto -z]$ reverse orientation, and all three components of the map preserve angles on the interior, so the map h is internally conformal. Also, the map h sends 1, -1 respectively to $-p_1$ and $-p_{-1}$, and $h^{-1}(-p_{-2})$ $\overline{f^{-1}(p_{-2})} = -\overline{f^{-1}(p_2)} = -h^{-1}(-p_2)$. Since $F_{D4p}(-(S,\ldots,p_{-2}))$ uniquely satisfies these conditions, part [3](#page-5-4) holds.

The map $h = [z \mapsto f(-z)]$ is an internally conformal homeomorphism from D to C that sends 1, -1 respectively to p_{-1} and p_1 , and $h^{-1}(p_2) = -h^{-1}(p_{-2})$. Since $F_{\text{D4p}}(S, p_{-1}, p_{-2}, p_1, p_2)$ uniquely satisfies these conditions, part [4](#page-5-5) holds.

Finally, stereographic projection though the point e_{-3} from **D** to S^2 is conformal and sends $S^1, 1, i, -1, -i$ respectively to $e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2}$, so part [5](#page-5-6) holds. \Box

2.3 A disk with two points on its null area boundary - F_{D2p}

We call a curve a *null area* curve when the curve itself has area 0. Here we construct a parameterization F_{D2p} of a topological disk that depends continuously on the boundary of the disk and on 2 distinct points on the boundary, provided that that boundary is a null area curve.

Lemma 2.7. Let C be a closed region in S^2 bounded by a null area simple closed curve S, and let $p_1, p_{-1} \in S$ be distinct points on the boundary of C. Then, there is a unique internally conformal homeomorphism $f = F_{D2p}(C, p_1, p_{-1}) : D \to C$ such that $f(1) = p_1, f(-1) = p_{-1}$, and the regions $f(iH \cap D)$ and $f(-iH \cap D)$ have equal area. Also, the following hold.

- 1. In the sup-metric, f depends continuously on C in boundary Fréchet distance and $p_1, p_{-1}.$
- 2. For $Q \in SO_3$, $F_{D2p}(Q(C, p_1, p_{-1})) = Qf$.
- 3. $F_{D2p}(-(C, p_1, p_{-1}); z) = -f(\overline{z}).$
- 4. $F_{D2p}(C, p_{-1}, p_1; z) = f(-z)$.
- 5. $F_{D2p}(C_3, e_1, e_{-1})$ with $C_3 = \{x \in \mathbf{S}^2 : \langle e_3, x \rangle \geq 0\}$ is stereographic projection though e_{-3} .

When C is an open region, let $F_{D2p}(C, p_1, p_{-1}) = F_{D2p}(\overline{C}, p_1, p_{-1})$ where \overline{C} is the closure of C.

Since area is used in the definition of the map F_{D2p} , we will need the following lemma that the area of the region bounded by a null area curve varies continuously.

Lemma 2.8. For $k \in \{1, \ldots, \infty\}$, let C_k be a closed region in S^2 bounded by a simple closed curve. If $C_k \to C_\infty$ in boundary Fréchet distance, and the boundary of C_∞ has null area, then the area of C_k converges to the area of C_{∞} .

Warning 2.9. Although F_{D2p} is defined when the boundary of C has positive area, F_{D2p} is only continuous over the space of regions with null area boundary. A crucial issue here is that the conclusion of Lemma [2.8](#page-9-0) does not hold in some cases where the boundary of C_{∞} has positive area, such as for the region bounded by an Osgood curve [\[15\]](#page-34-12).

Proof of Lemma [2.8.](#page-9-0) We claim that if $dist(C_k, C_\infty) < \varepsilon$ for ε sufficiently small, then the symmetric difference $\Delta_k = (C_k \setminus C_\infty) \cup (C_\infty \setminus C_k)$ is contained in $\partial C_\infty \oplus \varepsilon$. We may assume ε is small enough that there exists points p_0, p_1 such that $(p_0 \oplus \varepsilon) \cap C_{\infty} = \emptyset$ and $(p_1 \oplus \varepsilon) \subset C_{\infty}$. Since dist(C_k, C_∞) $\lt \varepsilon$ there is a map $f : \partial C_\infty \to \partial C_k$ such that $||f(x) - x|| < \varepsilon$, so x is always closer to $f(x)$ than to p_0 or p_1 . Therefore, there is a homotopy from ∂C_k to ∂C_k that stays within $S^2 \setminus \{p_0, p_1\}$, which implies that $p_0 \notin C_k$ and $p_1 \in C_k$. Since this holds for any such pair p_0, p_1 , the claim holds.

Consider $\varepsilon_1 > 0$. Since ∂C_{∞} has area 0, there is an open cover U of ∂C_{∞} that has area at most ε_1 . Since $S^2 \setminus U$ and ∂C_{∞} are compact and disjoint, they are bounded apart by some $\varepsilon_2 > 0$ depending on ε_1 . Since $C_k \to C_{\infty}$, we have for all k sufficiently large that dist $(C_k, C_\infty) < \varepsilon_2$, so by the claim above, $\Delta_k \subseteq \partial C_\infty \oplus \varepsilon_2$, so $\Delta_k \subseteq U$, so Δ_k has area at most ε_1 . Thus, area $(\Delta_k) \to 0$ as $k \to \infty$, which implies that area $(C_k) \to \text{area}(C_{\infty})$. \Box

Proof of Lemma [2.7.](#page-9-1) To make the problem simpler, we first use a linear fractional transformation F_{swivel} that sends the unit disk D to the upper half-plane H. Such a transformation is given by

$$
F_{\text{swivel}}(z) = -i\frac{z-1}{z+1}.
$$

We also have $F_{\text{swivel}}(1) = 0$, $F_{\text{swivel}}(-1) = \infty$, and $F_{\text{swivel}}(iH \cap D) = (D \cap H)$.

By Lemma [2.4,](#page-5-0) there exists an internally conformal homeomorphism g from \bf{D} to C such that $g(1) = p_1$ and $g(-1) = p_{-1}$. Fix g, and let $R(x) = [g \circ F_{\text{swivel}}^{-1}](x \mathbf{D} \cap \mathbf{H})$, and let $a(x)$ be the area of $R(x)$, and let a_{∞} be the area of C. Then, a is a strictly increasing function from $a(0) = 0$ to $a(\infty) = a_{\infty}$, so there is a unique $r > 0$ such that $a(r) = \frac{a_{\infty}}{2}$. Let

$$
f = g \circ F_{\text{swivel}}^{-1} \circ r F_{\text{swivel}}.
$$

Observe that $F_{\text{swivel}}(iH) = D$, so $f(iH \cap D) = [g \circ F_{\text{swivel}}^{-1}](rD \cap H) = R(r)$, which occupies half the area of C. Hence, f satisfies the defining properties of $F_{D2p}(C, p_1, p_{-1})$ in the lemma.

To check uniqueness, consider another map f_0 satisfying the hypotheses of the lemma. Then, $h = F_{\text{swivel}} \circ f_0 \circ f^{-1} \circ F_{\text{swivel}}^{-1}$ is a conformal automorphism of the upper half-plane with fixed points 0, ∞ , so h acts by scaling by a positive factor s. If we had $s \neq 1$, then the sets $f_0(iH \cap D)$ [°] and $f(iH \cap D) = R(r)$ [°] would be properly nested open sets, which would contradict that these sets have the same area. Thus, h is the identity, so $f_0 = f$.

To check part [1,](#page-9-2) continuity, consider sequences C_k , $p_{1,k}$, $p_{-1,k}$ converging to C_{∞} , $p_{1,\infty}$, $p_{-1,\infty}$ under the hypotheses of the lemma. Let $f_k = F_{D2p}(C_k, p_{1,k}, p_{-1,k})$, and let $R_k = f_k(i\mathbf{H} \cap \mathbf{D})$. Suppose for the sake of contradiction that $f_k(i)$ does not converge to $f_\infty(i)$. Then, there would be some $\varepsilon > 0$ such that $|f_k(i) - f_\infty(i)| > \varepsilon$, and since S^2 is compact, we could assume $f_k(i)$ converges to a point $p_2 \neq f_\infty(i)$; otherwise restrict to a subsequence bounded away from $f_{\infty}(\mathfrak{i})$ and then restrict to a convergent subsequence.

We may also assume that $f_k(-i)$ converges to a point p_{-2} . Let $\widetilde{f} = F_{\text{D4p}}(p_{\infty,1}, p_{\infty,-1}, p_2, p_{-2})$ and $\tilde{R} = \tilde{f}(\mathbf{i}H \cap \mathbf{D})$. By uniqueness in Lemma [2.4,](#page-5-0) we have $f_k = \mathrm{F}_{\mathrm{D4p}}(p_{k,1}, p_{k,-1}, f_k(\mathbf{i}), f_k(-\mathbf{i})),$ so by continuity in Lemma [2.4,](#page-5-0) we have $f_k \to \tilde{f}$ uniformly. Hence, $R_k \to \tilde{R}$ in boundary Fréchet distance. Since the boundaries of \widetilde{R} and C_{∞} have null area, by Lemma [2.8,](#page-9-0) we have $area(R_k) \rightarrow area(R)$ and $area(C_k) = 2area(R_k) \rightarrow area(C_{\infty}) = 2area(R_{\infty})$. Hence, $area(\widetilde{R}) = area(R_{\infty})$, so by the uniqueness of F_{D2p} , we have $\widetilde{f} = f_{\infty}$, so $p_2 = f_{\infty}(i)$, which contradicts our choice of subsequence. Thus, $f_k(i)$ must converge to $f_\infty(i)$. Similarly, $f_k(-i)$ must converge to $f_{\infty}(-i)$. Therefore, by Lemma [2.4,](#page-5-0) f_k converges to f_{∞} uniformly, so part [1](#page-9-2) holds.

The map Qf is an internally conformal homeomorphism from **D** to QC that sends 1, -1 respectively to Qp_1, Qp_{-1} , and $Qf(iH\cap D)$ occupies half the area of QC . Since $F_{D2p}(Q(S, \ldots, p_{-2}))$ uniquely satisfies these conditions, part [2](#page-9-3) holds.

The map $h = [z \mapsto -F_{D2p}(C, p_1, p_{-1}; \overline{z})]$ is an internally conformal homeomorphism from D to $-C$ that sends 1, -1 respectively to $-p_1$ and $-p_{-1}$ and $h(iH ∩ D)$ occupies half the area of $-C$. Since $F_{D2p}(-(S, p_1, p_{-2}))$ uniquely satisfies these conditions, part [3](#page-9-4) holds.

The map $h = [z \mapsto F_{D2p}(C, p_1, p_{-1}; -z)]$ is an internally conformal homeomorphism from D to C that sends 1, -1 respectively to p_{-1} and p_1 and $h(i\mathbf{H} \cap \mathbf{D})$ occupies half the area of C. Since $F_{D2p}(C, p_{-1}, p_1)$ uniquely satisfies these conditions, part [4](#page-9-5) holds.

Finally, stereographic projection though the point e_{-3} from **D** to S^2 sends **D**, 1, -1 respectively to C_3, e_1, e_{-1} and sends **iH** ∩ **D** to a lune occupying half the area of C_3 , so part [5](#page-9-6) holds. \Box

$\mathbf{2.4}\quad \mathbf{The\; sphere}\text{ - } \mathrm{F_{S^24p}} \text{ and } \mathrm{F_{S^22p}}$

Here we construct two self-homeomorphisms of the sphere, F_{S^24p} and F_{S^22p} . The map F_{S^24p} depends continuously on a curve and 4 distinct points on the curve, while F_{S^22p} depends continuously on a null area curve and 2 distinct points on the curve. In each case, the equator is mapped to the given curve. These maps are constructed by parameterizing the region on either side of the given curve respectively using F_{D4p} and F_{D2p} , and then modifying the parameterizations so that they agree along their common boundary, the curve itself. Since this is done the same way in both cases, we construct a map F_{stitch} that modifies a pair parameterizations in such a way.

Given embeddings $f_i : \mathbf{D} \to \mathbf{S}^2$ with $p_1 = f_1(1) = f_2(1)$, $p_{-1} = f_1(-1) = f_2(-1)$, and $S = f_1(S^1) = f_2(S^1)$, let $f = \mathrm{F}_{\mathrm{stitch}}(f_1, f_2) : \overline{\mathbb{C}} \to \mathbf{S}^2$

be the homeomorphism defined as follows. We will assume that both f_1 and f_2 are orientation preserving; otherwise replace f_i with $f_i \circ [z \mapsto \overline{z}]$. We first define a function f_0 , which will be the restriction of f to the unit circle in the complex plane. We will define f_0 in terms of its inverse. Let

$$
f_0^{-1}: S \to \mathbf{S}^1
$$
, $f_0^{-1}(x) = \sqrt{\frac{f_1^{-1}(x)}{f_2^{-1}(x)}}$

where we choose the branch of $\sqrt{\cdot}$ so that $f_0^{-1}(p_1) = 1$, and define f_0^{-1} from $(f_0^{-1})^2$ on the rest of S by analytic continuation. Note that the arc A of S extending in the counter-clockwise direction from p_1 around the boundary of C_1 as viewed from outside the sphere is sent to the upper half-plane of $\mathbb C$. Note also that this arc A extends in the clockwise direction from p around the boundary of C_2 , since C_2 is the cell on the opposite side of the sphere from C_1 . Let

$$
\mathcal{F}_{\text{stitch}}(f_1, f_2; z) = \begin{cases} f_1(0) & z = 0\\ f_1 \circ |z| f_1^{-1} \circ f_0(z/|z|) & 0 < |z| < 1\\ f_0(z) & |z| = 1\\ f_2 \circ \frac{1}{|z|} f_2^{-1} \circ f_0(z/|z|) & 1 < |z| < \infty\\ f_2(0) & z = \infty \end{cases}
$$

Note that we use the metric on $\overline{\mathbb{C}}$ induced by stereographic projection, which makes $\overline{\mathbb{C}}$ homeomorphic to the 2-sphere.

Lemma 2.10. The map F_{stitch} satisfies the following.

- 1. In the sup-metric, $F_{\text{stitch}}(f_1, f_2)$ depends continuously on f_1, f_2 in the sup-metric.
- 2. For $Q \in SO_3$, $F_{\text{stitch}}(Q(f_1, f_2)) = Q F_{\text{stitch}}(f_1, f_2)$.
- 3. $F_{\text{stitch}}(-(f_1, f_2); z) = -F_{\text{stitch}}(f_1, f_2; \overline{z}).$
- 4. $F_{\text{stitch}}(f_2, f_1; z) = F_{\text{stitch}}(f_1, f_2; 1/z).$

Proof. Consider embeddings $f_{i,k}$ of **D** in S^2 such that $(f_{1,k}, f_{2,k})$ is in the domain of F_{stitch} for $k \in \{1, \ldots, \infty\}$, and such that $f_{i,k} \to f_{i,\infty}$ uniformly. By Lemma [2.1,](#page-4-0) $f_{i,k} \to f_{i,\infty}$ invariably, so $f_{i,k}^{-1} \to f_{i,\infty}^{-1}$ invariably.

Let $f_{0,k}$ be as in the definition of F_{stitch} for the two maps $f_{1,k}, f_{2,k}$. The branch f_{sqrt} of the multifunction $[(x_1, x_2) \mapsto \sqrt{x_1/x_2}]$ on $({\bf S}^1 \cap {\bf H}) \times ({\bf S}^1 \cap -{\bf H})$ that sends $(1, 1)$ to 1 is continuous, and so is the branch on $(\mathbf{S}^1 \cap -\mathbf{H}) \times (\mathbf{S}^1 \cap \mathbf{H})$. Also, for $x \in S_k$, $f_{1,k}^{-1}(x)$ and $f_{2,k}^{-1}(x)$ always either coincide at 1 or -1 , or are on opposite sides of the real line, so the branch f_{sqrt} is continuous on the range of $f_{1,k}^{-1} \times f_{2,k}^{-1}$ over S_k . Hence, by Lemma [2.1,](#page-4-0) $f_{0,k}^{-1} \to f_{0,\infty}^{-1}$ invariably, so $f_{0,k} \to f_{0,\infty}$, and $f_{1,k}^{-1} \circ f_{0,k} \to f_{1,\infty}^{-1} \circ f_{0,\infty}$, and $f_{2,k}^{-1} \circ f_{0,k} \to f_{2,\infty}^{-1} \circ f_{0,\infty}$ invariably as well.

Let $g_{1,k}(z) = |z| f_1^{-1} \circ f_0(z|z|)$ for $z \in \mathbf{D} \setminus 0$ and $g_{1,k}(0) = 0$, and define $g_{2,k}$ analogously. We claim that $g_{i,k} \to g_{i,\infty}$ invariably. Consider $\varepsilon > 0$, and let $K(\varepsilon) > 0$ be such that for all $k \geq K(\varepsilon)$ and all $z \in \mathbf{D}$, we have $|[f_{1,k}^{-1} \circ f_{0,k}](z) - [f_{1,\infty}^{-1} \circ f_{0,\infty}](z)| < \varepsilon$ as in the definition of uniform convergence. Then, we have $|g_{1,k}(z) - g_{1,\infty}(z)| < \frac{2|z|}{1+|z|}$ $\frac{2|z|}{1+|z|^2}\varepsilon < \varepsilon$ in the metric on $\overline{\mathbb{C}}$ induced by stereographic projection for all $k \geq K(\varepsilon)$, so $g_{1,k} \to g_{1,\infty}$ uniformly. Therefore, by Lemma [2.1,](#page-4-0) $g_{1,k} \to g_{1,\infty}$ invariably, and similarly $g_{2,k} \to g_{2,\infty}$ invariably as claimed, which implies that $F_{\text{stitch}}(f_{1,k}, f_{2,k}) \rightarrow F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ invariably. Thus, $F_{\text{stitch}}(f_{1,k}, f_{2,k}) \rightarrow F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ $F_{\text{stitch}}(f_{1,k}, f_{2,k}) \rightarrow F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ $F_{\text{stitch}}(f_{1,k}, f_{2,k}) \rightarrow F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ uniformly, so part 1 holds.

Let f_0 and \tilde{f}_0 be as in the definition of F_{stitch} for the pairs (f_1, f_2) and $Q(f_1, f_2)$. The map $[Qf_i]^{-1}$ sends Qx to f_i^{-1} $i^{-1}(x)$, so on QS, the map f_0^{-1} sends Qx to the branch of $\sqrt{f_1^{-1}(x)/f_2^{-1}(x)}$ that sends Qp_1 to 1. Hence, $\tilde{f}_0 = Qf_0$. In the case where $0 < |z| < 1$, we have $F_{\text{stitch}}(Q(f_1, f_2); z) = Qf_1|z|f_1^{-1}Q^{-1}\tilde{f}_0(z) = Qf_1|z|f_1^{-1}f_0(z) = QF_{\text{stitch}}(f_1, f_2; z)$, and similarly in the other cases. Therefore, $F_{\text{stitch}}(Q(f_1, f_2)) = Q F_{\text{stitch}}(f_1, f_2)$ $F_{\text{stitch}}(Q(f_1, f_2)) = Q F_{\text{stitch}}(f_1, f_2)$ $F_{\text{stitch}}(Q(f_1, f_2)) = Q F_{\text{stitch}}(f_1, f_2)$, so part 2 holds.

Now let f_0 and \widetilde{f}_0 be as in the definition of F_{stitch} for the pairs (f_1, f_2) and $-(f_1, f_2)$. Since the antipodal map $[x \mapsto -x]$ reverses orientation on the sphere, the map f_0^{-1} sends $-x$ to the branch of $\sqrt{\overline{f_1^{-1}(x)}/\overline{f_2^{-1}(x)}} = \overline{\sqrt{\overline{f_1^{-1}(x)}/f_2^{-1}(x)}}$ that sends $-p_1$ to 1. Hence, $f_0(z) = -f_0(\overline{z})$. In the case where $0 < |z| < 1$, we have $F_{\text{stitch}}(-(f_1, f_2); z) = [-f_1|z|f_1^{-1} \circ$ $-\widetilde{f}_0(x) = -f_1|z|f_1^{-1}f_0(\overline{z}) = -\mathrm{F}_{\mathrm{stitch}}(f_1, f_2; \overline{z}),$ and similarly in the other cases. Therefore, $F_{\text{stitch}}(-(f_1, f_2); z) = - F_{\text{stitch}}(f_1, f_2; \overline{z}),$ so part [3](#page-12-2) holds.

For $x \in S$, we have

$$
[F_{\text{stitch}}(f_2, f_1)]^{-1}(x) = \sqrt{\frac{f_2^{-1}(x)}{f_1^{-1}(x)}}
$$

=
$$
\frac{1}{\sqrt{\frac{f_1^{-1}(x)}{f_2^{-1}(x)}}}
$$

=
$$
[[z \mapsto 1/z] \circ [F_{\text{stitch}}(f_1, f_2)]^{-1}](x)
$$

=
$$
[F_{\text{stitch}}(f_1, f_2) \circ [z \mapsto 1/z]]^{-1}(x),
$$

where the branch of $\sqrt{\cdot}$ is always chosen so that p_1 is sent to 1. Thus, $F_{\text{stitch}}(f_2, f_1; z)$ $F_{\text{stitch}}(f_1, f_2; 1/z)$ on $z \in S^1$, which means part [4](#page-12-3) holds on $z \in S^1$. Consider $z \in \overline{\mathbb{C}}$, and let $x = \mathcal{F}_{\text{stitch}}(f_2, f_1; z/|z|) \in S$. In the case where $0 < |z| < 1$, we have $1 < |1/z| < \infty$, so $F_{\text{stitch}}(f_1, f_2; 1/z) = f_2 \circ |z| f_2^{-1}(x) = F_{\text{stitch}}(f_2, f_1; z)$. Also, $F_{\text{stitch}}(f_1, f_2; 1/z) = f_2(0) =$ $F_{\text{stitch}}(f_2, f_1; 0)$, so part [4](#page-12-3) holds on **D**, and similarly in the case where $1 < |z| \leq \infty$, so part [4](#page-12-3) holds on all of $\overline{\mathbb{C}}$. \Box

For a simple closed curve $S \subset \mathbf{S}^2$ and 4 distinct points p_1, p_2, p_{-1}, p_{-2} appearing in that order around the curve, let

$$
F_{S^24p}(S, p_1, p_2, p_{-1}, p_{-2}) = F_{\text{stitch}}(F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2))
$$

Lemma 2.11. The map $f = F_{S^24p}(S, p_1, p_2, p_{-1}, p_{-2})$ satisfies the following.

- 1. f in the sup-metric depends continuously on S in Fréchet distance and on p_1, \ldots, p_{-2} .
- 2. For $Q \in O_3$, $F_{S^24n}(Q(S, p_1, p_2, p_{-1}, p_{-2})) = Qf$.
- 3. $F_{S^24p}(S, p_{-1}, p_{-2}, p_1, p_2; z) = f(-z)$.
- 4. $F_{S^24p}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2})$ is stereographic projection though e_{-3} .

Proof. Since $f_1 = F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2})$ and $f_2 = F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2)$ in the sup-metric depend continuously on $S, p_1, p_2, p_{-1}, p_{-2}$ by Lemma [2.4,](#page-5-0) and f in the sup-metric depends continuously on f_1 f_1 and f_2 , part 1 holds.

By Lemmas [2.4](#page-5-0) and [2.10,](#page-11-0) we have the following. In the case where $Q \in SO_3$ we have

$$
F_{S^24p}(Q(S, p_1, p_2, p_{-1}, p_{-2})) = F_{\text{stitch}}(F_{D4p}(Q(S, p_1, p_2, p_{-1}, p_{-2})), F_{D4p}(Q(S, p_1, p_{-2}, p_{-1}, p_2)))
$$

\n
$$
= F_{\text{stitch}}(Q F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), Q F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2))
$$

\n
$$
= Q F_{\text{stitch}}(F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2))
$$

\n
$$
= Qf.
$$

In the case where $Q \in -SO_3$ we have

$$
F_{S^24p}(Q(S, p_1, p_2, p_{-1}, p_{-2}); z) = F_{\text{stitch}}(F_{D4p}(Q(S, p_1, p_2, p_{-1}, p_{-2})), F_{D4p}(Q(S, p_1, p_{-2}, p_{-1}, p_2)); z)
$$

\n
$$
= F_{\text{stitch}}(Q F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), Q F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2); \overline{z})
$$

\n
$$
= Q F_{\text{stitch}}(F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2); z)
$$

\n
$$
= Qf.
$$

Hence, part [2](#page-13-1) holds.

$$
F_{S^24p}(S, p_{-1}, p_{-2}, p_1, p_2; z) = F_{\text{stitch}}(F_{D4p}(S, p_{-1}, p_{-2}, p_1, p_2), F_{D4p}(S, p_{-1}, p_2, p_1, p_{-2}; z))
$$

= F_{\text{stitch}}(F_{D4p}(S, p_1, p_2, p_{-1}, p_{-2}), F_{D4p}(S, p_1, p_{-2}, p_{-1}, p_2; -z))
= f(-z).

Hence, part [3](#page-13-2) holds.

The map $f_1 = \mathrm{F}_{\mathrm{D4p}}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2})$ $f_1 = \mathrm{F}_{\mathrm{D4p}}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2})$ $f_1 = \mathrm{F}_{\mathrm{D4p}}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2})$ is stereographic projection though e_{-3} , and by part 2, the map $f_2 = \mathrm{F}_{\mathrm{D4p}}(e_3^{\perp}, e_1, e_{-2}, e_{-1}, e_2)$ is stereographic projection though e_3 , so the map $[z \mapsto$ $f_2(1/z)$ is the stereographic projection on $\overline{\mathbb{C}\setminus \mathbf{D}}$ though e_{-3} . Therefore, the map $\sqrt{f_1^{-1}/f_2^{-1}}$ is the stereographic projection on S^1 though e_{-3} . Thus, $F_{S^24p}(e_3^{\perp}, e_1, e_2, e_{-1}, e_{-2}) = F_{\text{stitch}}(f_1, f_2)$ is stereographic projection though e_{-3} , so part [4](#page-13-3) holds.

For a region C bounded by a null area simple closed curve, and distinct points p_1, p_{-1} on the curve, let

$$
F_{S^{2}2p}(C, p_{1}, p_{-1}) = F_{\text{stitch}}(F_{D2p}(C, p_{1}, p_{-1}), F_{D2p}(S^{2} \setminus C, p_{1}, p_{-1}))
$$

Lemma 2.12. The map $f = F_{S^22p}(C, p_1, p_{-1})$ satisfies the following.

- 1. f in the sup-metric depends continuously on C in boundary Fréchet distance and on $p_1, p_{-1}.$
- 2. For $Q \in O_3$, $F_{S^22p}(Q(C, p_1, p_{-1})) = Qf$.
- 3. $F_{S^22p}(S^2 \setminus C, p_{-1}, p_1; z) = f(-1/z).$
- 4. $F_{S^2P}(C_3, e_1, e_{-1})$ with $C_3 = \{x \in S^2 : \langle e_3, x \rangle \geq 0\}$ is stereographic projection though e_{-3} .

Proof. Since $f_1 = F_{D2p}(C, p_1, p_{-1})$ and $f_2 = F_{D2p}(S^2 \setminus C, p_1, p_{-1})$ in the sup-metric depend continuously on C, p_1, p_{-1} by Lemma [2.7,](#page-9-1) and f in the sup-metric depends continuously on f_1 f_1 and f_2 by Lemma [2.10,](#page-11-0) part 1 holds.

$$
F_{S^{2}2p}(S^{2} \setminus C, p_{-1}, p_{1}; z) = F_{\text{stitch}}(F_{D2p}(S^{2} \setminus C, p_{-1}, p_{1}), F_{D2p}(C, p_{-1}, p_{1}); z)
$$

\n
$$
= F_{\text{stitch}}(F_{D2p}(S^{2} \setminus C, p_{1}, p_{-1}), F_{D2p}(C, p_{1}, p_{-1}); -z)
$$

\n
$$
= F_{\text{stitch}}(F_{D2p}(C, p_{1}, p_{-1}), F_{D2p}(S^{2} \setminus C, p_{1}, p_{-1})); -1/z)
$$

\n
$$
= f(-1/z)
$$

Hence, part [3](#page-14-1) holds. Parts [2](#page-14-2) and [4](#page-14-3) hold by the same argument as in the proof of Lemma [2.11.](#page-13-4) \Box

3 The deformation

Our goal here is to construct the deformation retraction implied by Theorem [1.1,](#page-0-1) which we denote by ρ . We first define a collection of objects that we will refer to when we construct

Figure 1: First 4 stages of the deformation retraction starting from a map f .

Figure 3: Stages 5 and 6 of the deformation retraction. The thin lines in the disk show how evenly spaced meridians are mapped.

ρ. These objects are defined in terms of ρ or in terms of each other. In each stage of the deformation, we define the evolution of some of these objects, and ρ and the other objects will then be determined as a consequence of the defining relationships between the objects.

Recall $e_{-i} = -e_i$ and e_i denotes the *i*th standard basis vector. For $f \in \text{hom}(S^2)$ and $t \in [0,6]_{\mathbb{R}},$ let

$$
p_i(f, t) = \rho(f, t; e_i) \quad \text{for } i \in \{-3, -2, -1, 1, 2, 3\},
$$

\n
$$
S_E(f, t) = \rho(f, t; e_3^{\perp}),
$$

\n
$$
F_E(f, t) = F_{S^24p} ([S_E, p_1, p_2, p_{-1}, p_{-2}](f, t)).
$$

\n
$$
S_A(f, 0) = F_E(f, t; \overline{\mathbb{R}})
$$

\n
$$
= \bigcup_{s \in \{1, -1\}} F_{D4p} ([S_E, p_1, p_{s2}, p_{-1}, p_{-s2}](f, 0); [-1, 1]_{\mathbb{R}}),
$$

\n
$$
S_A(f, t) = [\rho(f, t) \circ f^{-1}](S_A(f, 0)),
$$

let $C_1(f, t)$ be the region bounded by $S_A(f, t)$ that contains the point p_2 , and $C_{-1}(f, t)$ be that containing p_{-2} , and let

$$
F_{A}(f,t) = F_{S^{2}2p}([C_{1}, p_{1}, p_{-1}](f,t)).
$$

The letter E in S_E and F_E is for equator, and A in S_A and F_A is for Roald Amundsen, a polar explorer. We refer to e_3^{\perp} as the equator. In some cases we may suppress the argument for cleaner notation. For example, we may simply write p_1 for $p_1(f, t)$, as long as the meaning can be understood from context.

The deformation will proceed through 6 stages: balancing, untangling, aligning, flattening, divvying, and combing. Each stage will occur over a unit interval, so that the whole deformation occurs over $[0,6]_{\mathbb{R}}$ instead of a single unit interval. The first 4 stages deform the image of the equator, S_E , to a great circle; see Figures [1](#page-15-0) and [2.](#page-16-0) The first stage, balancing, deforms the curve S_A to a bisector of the sphere. That is, S_A is deformed to make the regions on either side each have area 2π . The second stage, untangling, deforms S_A to a great circle. The third stage, aligning, moves the points p_1 and p_{-1} into antipodal position. The fourth stage, flattening, deforms the curve S_E to a great circle. Once the equator is mapped to a great circle, we deal the map in the two hemispheres on either side separately; see Figure [3.](#page-16-0) The fifth stage, divvying, deforms the map to an isometry on the equator, and the sixth stage, combing, deforms the map to an isometry on the rest of the sphere.

3.1 Balancing

In the first stage, we deform S_A so that each of the regions on either side have area 2π .

Let
$$
\omega(f,t) = F_A(f,0) \circ e^t F_A(f,0)^{-1}
$$
. Let $a(f,t)$ be the area of

$$
D(f, t) = \omega(f, t; C_1(f, 0)) = F_A(f, 0; e^t \mathbf{D}).
$$

Observe that $a(f)$ is a strictly increasing continuous function on $\overline{\mathbb{R}}$ from $a(-\infty) = 0$ to $a(\infty) = 4\pi$. Let $T(f) \in \mathbb{R}$ such that $a(f, T(f)) = 2\pi$. For $t \in [0, 1]_{\mathbb{R}}$, let

$$
\rho(f,t) = \omega(f, tT(f)) \circ f.
$$

Lemma 3.1. For all $f \in \text{hom}(S^2)$ and $t \in \mathbb{R}$, the boundary of $D(f, t)$ is rectifiable, and so has null area. Also, $C_1(f, 1) = D(f, T(f))$ and has area 2π .

Lemma 3.2. For all $t \in [0,1]_{\mathbb{R}}$ and $f \in \text{hom}(S^2)$, we have the following.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$, and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in \mathcal{O}_3$, then $\rho(f, t) = f$.
- μ , $\rho(f, 0) = f$.

Proof of Lemma [3.1.](#page-18-0) In the case where $t = 0$, the boundary of $D(f, t)$ is $S_A(f, 0)$, which is the union of two smooth curves, so the boundary is rectifiable.

Consider the case where $t < 0$. Let $f_1 = F_{D2p}([C_1, p_1, p_{-1}](f, 0))$ and $f_0 = F_A(f, 0)$ restricted to S^1 . Then,

$$
D(f, t) = F_A(f, 0; e^t \mathbf{D})
$$

= $F_{S^2 2p}([C, p_1, p_{-1}](f, 0); e^t \mathbf{D})$
= $[z \mapsto [f_1 \circ |z|f_1^{-1} \circ f_0](z/|z|)](e^t \mathbf{D} \setminus 0) \cup \{0\}$
= $f_1(e^t \mathbf{D}),$

since $f_1^{-1} \circ f_0$ sends S^1 to S^1 bijectively. Since f_1 is internally conformal, the boundary of $D(f, t)$ is smooth and therefore rectifiable. The case where $t > 0$ follows similarly. Also,

$$
C_1(f, 1) = [\rho(f, 1) \circ f^{-1}](C_1(f, 0))
$$

= $[\omega(f, T(f))](C_1(f, 0))$
= $[F_A(f, 0) \circ e^{T(f)} F_A(f, 0)^{-1}](C_1(f, 0))$
= $F_A(f, 0; e^{T(f)} \mathbf{D})$
= $D(f, T(f)),$

which by definition of T has area 2π .

Proof of Lemma [3.2](#page-18-1) part [1.](#page-18-2) Consider $f_k \in \text{hom}(\mathbf{S}^2)$ and $t_k \in [0,1]_{\mathbb{R}}$ in the domain of ρ such that $f_k \to f_\infty$ uniformly and $t_k \to t_\infty$. Then, $S_E(f_k, 0) \to S_E(f_\infty, 0)$ in Fréchet distance $p_i(f_k, 0) \to p_i(f_\infty, 0)$, so $S_A(f_k, 0) \to S_A(f_\infty, 0)$ in Fréchet distance by Lemma [2.11,](#page-13-4) which implies that $C(f_k, 0) \to C(f_{\infty}, 0)$ in boundary Fréchet distance, so $F_A(f_k, 0) \to F_A(f_{\infty}, 0)$ uniformly, which implies that $F_A(f_k, 0)^{-1} \to F_A(f_\infty, 0)^{-1}$ uniformly as well.

Suppose for the sake of contradiction that $T(f_k)$ does not converge to $T(f_{\infty})$. Since $\overline{\mathbb{R}}$ is compact, we may assume that $T(f_k) \to \tilde{T} \neq T(f_{\infty})$ for some $\tilde{T} \in \overline{\mathbb{R}}$. Let us assume for now that $\widetilde{T} < T(f_{\infty})$. Then, $e^{T(f_{\infty})} \mathbf{D} \setminus e^{T} \mathbf{D}$ has non-empty interior, so $D(f_{\infty}, T(f_{\infty})) \setminus e^{T} \mathbf{D}$ $D(f_{\infty}, \tilde{T}) = F_{\mathbf{A}}(f_{\infty}, 0; e^{T(f_{\infty})} \mathbf{D} \setminus e^{T} \mathbf{D})$ also has non-empty interior, which implies that $a(f_{\infty}, \tilde{T}) \leq a(f_{\infty}, T(f_{\infty})) = 2\pi$. Also, $D(f_k, T(f_k)) \to D(f_{\infty}, \tilde{T})$ in boundary Fréchet distance, and by Lemma [3.1,](#page-18-0) the boundary of $D(f_{\infty}, \tilde{T})$ is rectifiable, so by Lemma [2.8,](#page-9-0)

 \Box

 $a(f_k, T(f_k)) \to a(f_{\infty}, \tilde{T}) < 2\pi$, so $a(f_k, T(f_k)) < 2\pi$ for some k sufficiently large, but $a(f_k, T(f_k)) = 2\pi$ by definition of T, which is a contradiction. Similarly, we have $a(f_k, T(f_k))$ 2π for some k sufficiently large in the case where $\tilde{T} > T(f_{\infty})$, which also contradicts the definition of T. Hence, $T(f_k)$ must converge to $T(f_{\infty})$.

Thus, $\omega(f_k, t_kT(f_k)) \to \omega(f_{\infty}, t_{\infty}T(f_{\infty}))$ uniformly, and therefore $\rho(f_k, t_k) \to \rho(f_{\infty}, t_{\infty})$ uniformly. \Box

Proof of Lemma [3.2](#page-18-1) part [2.](#page-18-3)

$$
S_{E}(Qfs, 0) = Qfs(e_{3}^{\perp}) = Qf(e_{3}^{\perp}) = QS_{E}(f, 0).
$$

By Lemma [2.11,](#page-13-4) we have

$$
S_A(Qfs, 0) = F_{S^24p} ([S_E, p_1, p_2, p_{-1}, p_{-2}](Qfs, 0); \overline{\mathbb{R}})
$$

= $F_{S^24p} (Q[S_E, p_s, p_{s2}, p_{-s}, p_{-s2}](f, 0); \overline{\mathbb{R}})$
= $Q F_{S^24p} ([S_E, p_1, p_2, p_{-1}, p_{-2}](f, 0); [z \mapsto sz](\overline{\mathbb{R}}))$
= $Q S_A(f, 0).$

Since $S_A(Qfs, 0) = QS_A(f, 0)$ and $p_2(Qfs, 0) = Qp_{s2}(f, 0)$, we have $C_1(Qfs, 0) = QC_s(f, 0)$. By Lemma [2.12,](#page-14-4) we have

$$
F_A(Qfs, 0) = F_{S^22p}([C_1, p_1, p_{-1}](Qfs, 0))
$$

=
$$
F_{S^22p}(Q[C_s, p_s, p_{-s}](f, 0))
$$

=
$$
Q F_A(f, 0) \circ [z \mapsto sz^s],
$$
 (1)

$$
\omega(Qfs, t) = F_A(Qfs, 0) \circ e^t F_A(Qfs, 0)^{-1}
$$

= $Q F_A(f, 0) \circ s^2 e^{st} F_A(f, 0)^{-1} Q^{-1}$
= $Q\omega(f, st)Q^{-1}$,

so $D(Qfs, t) = QD(f, st)$, so $a(Qfs, t) = a(f, st)$, so $T(Qfs) = sT(f)$, so $\rho(Qfs,t) = \omega(Qfs,tT(Qfs))Qfs$ $= Q\omega(f, stsT(f))Q^{-1}Qfs$ $= Q \rho(f, t) s.$

Proof of Lemma [3.2](#page-18-1) parts [3](#page-18-4) and [4.](#page-18-5) For part [4,](#page-18-5) we have

$$
\rho(f,0) = \omega(f,0) \circ f = F_A(f,0) \circ e^0 F_A(f,0)^{-1} \circ f = f.
$$

Next, consider the case $f = id$. In this case, $S_E(id, 0) = e_3^{\perp}$ and $p_i(id, 0) = e_i$, so by Lemma [2.11,](#page-13-4) the map $F_E(id, 0)$ is stereographic projection though e_{-3} , so $S_A(id, 0) = e_2^{\perp}$ and $C(\mathrm{id}, 0) = \{x \in \mathbf{S}^2 : \langle e_2, x \rangle \ge 0\}$, which has area 2π , so $T(\mathrm{id}) = 0$.

In the case $f \in O_3$, we have $T(f) = T(id) = 0$, so

$$
\rho(f, t) = \omega(f, tT(f)) \circ f = \omega(f, 0) \circ f = f.
$$

 \Box

3.2 Untangling

In the second stage, we deform S_A to a great circle. To do so, S_A will evolve by level-set flow, which is a weak formulation of curvature flow for Jordan curves of area 0, an extremely mild smoothness condition. Here are the important properties of level-set flow for our purpose. Joseph Lauer showed that level-set flow is defined for all null area simple closed curves up to some positive time depending on the curve, and that the curve immediately becomes smooth and evolves by curvature flow [\[12\]](#page-34-13). Michael Gage showed that for bisectors, i.e., null area simple closed curves that divide the sphere into 2 regions that each have area 2π , the evolution by curvature flow approach a great circle as time becomes infinite [\[9\]](#page-34-14). Dobbins showed that for bisectors, level-set flow depends continuously in Fréchet distance on initial conditions [\[5\]](#page-34-4). Sigurd Angenet showed that for a pair of smooth curves evolving by curvature flow, the number of intersection points is finite and non-increasing, provided that the curves are initially distinct [\[1\]](#page-34-15). Dobbins showed that the points of intersection move along a trajectory that depends continuously on initial conditions [\[5\]](#page-34-4).

As S_A evolves, we will also have to determine how the points p_1 and p_{-1} move. For this we introduce another curve S_L , which will also evolve by curvature flow, and we will let p_1 and p_{-1} be the points where the curves S_A and S_L intersect. By Angenet's theorem this is well defines for positive time, but this might not be well defined at the limit as t becomes infinite. To deal with this we will determine a time T_1 , after which the points p_1 and p_{-1} will move according to reparameterizations of the regions on either side of S_A using the Carathéodory mapping theorem.

Let $L(f) = L_1 \cup L_{-1} \subset \overline{\mathbb{C}}$ be the lune consisting of 2 circular arcs L_1, L_{-1} from 1 to -1 such that $F_A(f, 1; L_n)$ bisects C_n . Let $\gamma_A(f)$ and $\gamma_L(f)$ be the level-set flows starting from $\gamma_A(f;0) = S_A(f,1)$ and $\gamma_L(f;0) = F_A(f,1;L(f))$. Note that since $\gamma_A(f;0)$ is a bisector, $\gamma_{A}(f;\infty) = \lim_{t\to\infty} \gamma_{A}(f;t)$ exists and is a great circle, and likewise for γ_{L} .

Let $\tau(t) = \frac{t-1}{2-t}$. Note that τ is strictly increasing on $[1,2]_{\mathbb{R}}$ and sends $[1,2]_{\mathbb{R}}$ to $[0,\infty]_{\mathbb{R}}$. For $t \in (1,2]_{\mathbb{R}},$ let $S_A(f,t) = \gamma_A(f,\tau(f,t))$ and $S_L(f,t) = \gamma_L(f,\tau(f,t))$. Let C_n be on the same side of S_A throughout the evolution.

Let $U = U(f) = \{u_1, u_{-1}\} \subset \mathbf{S}^2$ be the two points on the sphere that are perpendicular to the plane spanned by $S_A(f, 2)$ with $u_1 \in C_1(f, 2)$. Note that $S_A(f, 2)$ is a great circle, so U is well defined.

For (f, t) and n such that $u_n(f) \in (C_n(f, t))^{\circ}$, let $h_n(f, t) : C_n(f, 2) \to C_n(f, t)$ be the internally conformal homeomorphism that is fixed at $u_n = h_n(f, t; u_n)$ and $D(h_n(f, t); u_n) =$ λ id with $\lambda > 0$ where $D(h, x)$ denotes the total derivative of a function h at x. Note that h_n is uniquely determined by these conditions by the Schwarz lemma. Let

$$
r_{h}(f,t) = \sup \{ ||h_{n}(f,t;x) - x|| : x \in S_{A}(f,2), n \in \{1, -1\} \},
$$

where r_h is only defined when $h₁$ and $h₋₁$ are both defined,

$$
r_{\mathbf{u}}(f, t) = \sup \left\{ \left| (||x - u_n||) - \sqrt{2} \right| : x \in S_{\mathbf{A}}(f, t), n \in \{1, -1\} \right\},
$$

$$
r_{\mathbf{t}}(f, t) = \sup \left\{ r_{\mathbf{h}}(f, s), r_{\mathbf{u}}(f, s) : t \le s \le 2 \right\},
$$

$$
w(f, t) = (1 - r_{\mathbf{t}}(f, t))^+
$$

where $(x)^{+} = \max(x, 0)$ is the positive part of a function. We will show that $r_t(f, t)$ is positive and decreasing on $t \in [1,2]_{\mathbb{R}}$. Let $T_1(f)$ be the weighted median value of t with weight $w(f, t)$, i.e. such that

$$
\int_{t=1}^{T_1(f)} w(f, t) dt = \frac{1}{2} \int_{t=1}^2 w(f, t) dt.
$$

For a pair of bisectors that intersect at a pair of points, there is a unique trajectory starting at each intersection point that remains in the intersection of the curves as they evolve by level-set flow [\[5\]](#page-34-4). For $t \leq T_1(f)$, let $p_1(f,t)$ be the trajectory starting from $p_1(f, 1)$ that remains in $[S_A \cap S_L](f, t)$, and define p_{-1} analogously. For $t > T_1(f)$, let

$$
p_n(f, t) = [h_n(f, t) \circ h_n(f, T_1(f))^{-1}](p_n(f, T_1(f))).
$$

For $t \in (1,2]_{\mathbb{R}},$ let

$$
\rho(f,t) = \mathcal{F}_{\mathcal{A}}(f,t) \circ \mathcal{F}_{\mathcal{A}}(f,1)^{-1} \circ \rho(f,1).
$$

Lemma 3.3. For all $f \in \text{hom}(S^2)$, $S_A(f, 2)$ is a great circle.

Lemma 3.4. For all $t \in [1,2]_{\mathbb{R}}$ and $f \in \text{hom}(S^2)$, we have the following.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$, and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in \mathcal{O}_3$, then $\rho(f, t) = f$.

Proof of Lemma [3.3.](#page-21-0) Curves evolving by curvature flow approach a great circle [\[9\]](#page-34-14), and $S_A(f, 2)$ is defined as the limit as $t \to \infty$ of such a curve $\gamma_A(f, t)$, so $S_A(f, 2)$ is a great circle. \Box

Proof of Lemma [3.4](#page-21-1) part [1.](#page-21-2) Consider $f_k \to f_\infty$ in the sup-metric and $t_k \to t_\infty$. By Lemma [3.2,](#page-18-1) $S_A(f_k, 1) \to S_A(f_{\infty}, 1)$ in Fréchet distance and $p_n(f_k, 1) \to p_n(f_{\infty}, 1)$, so by Lemma [2.12](#page-14-4) $F_A(f_k, 1) \rightarrow F_A(f_\infty, 1)$ in the sup-metric.

Suppose for the sake of contradiction that $L_1(f_k)$ does not converge to $L_1(f_\infty)$ in Fréchet distance. We can parameterize the space of circular arcs through D from 1 to −1 by the point where the arc intersects the segment $[-i, i]_{\mathbb{C}}$. Moreover, the space of such arcs is compact, provided we include the two semicircles on $S¹$ between 1 and -1 . Hence, we may assume that $L_1(f_k) \to L \neq L_1(f_\infty)$; otherwise restrict to a convergent subsequence.

Let $S_{L1,k} = F_A(f_k, 1; L_1(f_k))$, and $\widetilde{S} = F_A(f_k, 1; \widetilde{L})$. Then, $S_{L1,k} \to \widetilde{S}$ in Fréchet distance, since $F_A(f_\infty, 1; L_1(f_k)) \to \widetilde{S}$ and the Fréchet distance between $F_A(f_\infty, 1; L_1(f_k))$ and $S_{L_1,k}$ converges to 0. The curves $S_A(f_\infty, 1)$ and \widetilde{S} are piecewise smooth, and therefore rectifiable, so by Lemma [2.8,](#page-9-0) the areas of the regions of $C_1(f_k)$ on either side of $S_{L1,k}$ converge to those of \widetilde{S} , but \widetilde{S} is on one side of $S_{L1,\infty}$, so \widetilde{S} does not bisect the sphere. Hence, for some k sufficiently large, $S_{L1,k}$ does not bisect the sphere, but that contradicts the definition of L_1 . Thus, $L_1(f_k)$ must converge to $L_1(f_\infty)$ in Fréchet distance. Similarly, $L_{-1}(f_k)$ converges to $L_{-1}(f_{\infty})$, so $L(f_k)$ converges to $L(f_{\infty})$ in Fréchet distance.

With this, we have that $S_{L}(f_{k}, 1) = F_{A}(f_{k}, 1; L(f_{k})) \rightarrow S_{L}(f_{\infty}, 1)$ in Fréchet distance, since $L(f_k) \to L(f_\infty)$ and $F_A(f_k, 1) \to F_A(f_\infty, 1)$, and $S_A(f_k, 1) \to S_A(f_\infty, 1)$. Since level-set flow depends continuously on initial conditions for bisectors [\[5\]](#page-34-4), $S_L(f_k, t_k) \to S_L(f_\infty, t_\infty)$ and $S_A(f_k, t_k) \to S_A(f_\infty, t_\infty)$ in Fréchet distance.

By Angenet's theorem $[S_A \cap S_L](f, t)$ is a pair of points for $t \in [1, 2)_{\mathbb{R}}$ [\[1\]](#page-34-15), and the author showed that these points move along a trajectory that depends continuously on $S_A(f, 1)$ and $S_{L}(f, 1)$ in Fréchet distance [\[5\]](#page-34-4). Let $p_{\text{AL},n}(f, t) \in [S_{A} \cap S_{L}](f, t)$ be the point on the trajectory starting from $p_{\text{AL},n}(f,1)$ for $n \in \{1,-1\}$. By [, Lemma 3.1.3], $p_{\text{AL},n}(f_k, t_k) \to p_{\text{AL},n}(f_{\infty}, t_{\infty})$ provided that $t_{\infty} < 2$.

Since $S_A(f_k, t_k) \to S_A(f_\infty, t_\infty)$ and $u_n(f_k) \to u_n(f_\infty)$, we have $r_n(f_k, t_k) \to r_n(f_\infty, t_\infty)$.

In the case where $u_n(f_\infty, t_\infty) \in (C_n(f_\infty, t_\infty))^{\circ}$, we have that $u_n(f_\infty, t_\infty)$ is bounded away from $S^2 \setminus (C_n(f_\infty, t_\infty))^\circ$, so for all k sufficiently large, $u_n(f_k, t_k) \in (C_n(f_k, t_k))^\circ$. Hence $h_n(f_k, t_k)$ is defined for all k sufficiently large, and by Rado's theorem, $h_n(f_k, t_k) \to h_n(f_\infty, t_\infty)$ uniformly, so $r_{\rm h}(f_k, t_k) \rightarrow r_{\rm h}(f_\infty, t_\infty)$.

Let T_0 be the last time that there is a point of $U(f_{\infty})$ on $S_{\mathcal{A}}(f_{\infty}, T_0)$. Consider the case where $t_{\infty} > T_0$. Consider $\varepsilon > 0$.

By the definition of r_t , there is $s \ge t_\infty$ such that $r_c(f_\infty, s) > r_t(f_\infty, t_\infty) - \varepsilon/2$ where c is either 'h' or 'u'. In either case, we have already shown that $r_c(f_k, s) \to r_c(f_\infty, s)$. In the case where $s = t_{\infty}$, we have $r_c(f_k, t_k) \to r_c(f_{\infty}, s)$, so for all k sufficiently large, $r_t(f_k, t_k) \ge r_c(f_k, t_k) >$ $r_c(f_\infty, s) - \varepsilon/2$. In the case where $s > t_\infty$, for all k sufficiently large, $r_c(f_k, s) > r_c(f_\infty, s) - \varepsilon/2$ and $t_k < s$, so again $r_t(f_k, t_k) > r_c(f_\infty, s) - \varepsilon/2$. In any case $r_t(f_k, t_k) > r_t(f_\infty, t_\infty) - \varepsilon$.

Since $r_c(f, t)$ is continuous as a function of f and t, for all k sufficiently large and all s such that $|s-t_{\infty}| \leq |t_{k}-t_{\infty}|$, we have that $r_c(f_k, s) < r_c(f_{\infty}, t_{\infty}) + \varepsilon \leq r_t(f_{\infty}, t_{\infty}) + \varepsilon$. We also have by definition of r_t that for all $s \ge t_\infty$ that $r_c(f_\infty, s) < r_t(f_\infty, t_\infty) + \varepsilon/2$ for both $c = h$ and $c = u$. For all k sufficiently large, we have $r_c(f_k, s) < r_c(f_{\infty}, s) + \varepsilon/2$, so $r_c(f_k, s) < r_t(f_\infty, t_\infty) + \varepsilon$. Together, we now have that for all k sufficiently large, all $s \ge t_k$, and $c \in \{h, u\}$ that $r_c(f_k, s) < r_t(f_\infty, t_\infty) + \varepsilon$. Therefore, $r_t(f_k, t_k) \leq r_t(f_\infty, t_\infty) + \varepsilon$.

We now have for all k sufficiently large that $|r_t(f_k, t_k) - r_t(f_{\infty}, t_{\infty})| \leq \varepsilon$ Hence, $r_t(f_k, t_k) \to$ $r_t(f_\infty, t_\infty)$, provided $t_\infty > T_0$.

In the case where $t_{\infty} = T_0$, we have $r_u(f_k, t_k) \to r_u(f_{\infty}, t_{\infty}) = \sqrt{2}$, which is the largest value r_h In the case where $t_{\infty} = t_0$, we have $r_u(f_k, t_k) \to r_u(f_{\infty}, t_{\infty}) = \sqrt{2}$, which is the largest value r_h
or r_u can attain, so $r_t(f_k, t_k) \to r_t(f_{\infty}, t_{\infty}) = \sqrt{2}$ in this case as well, and likewise in the case

where $t_{\infty} < T_0$, since for k sufficiently large, $\sqrt{2} \geq r_t(f_k, t_k) \geq r_u(f_k, T_0) \rightarrow$ √ $2 = r_{\rm t}(f_{\infty}, T_0).$ Thus, $r_t(f, t)$ is continuous as a function of f and t, and therefore w is continuous as well.

As $t \to 2$, we have $h_n(f, t) \to id_{C_n(f, 2)}$, since $S_A(f, t) \to S_A(f, 2)$, so $r_h(f, t) \to 0$, and likewise for $r(u)(f, t)$, since $S_A(f, t)$ converges to the great circle orthogonal to u_1 and u_{-1} , so $r_t(f, t)$ is decreasing and converges to 0 as $t \to 2$. Hence, $w(f, t)$ is non-negative and increasing as a function of $t \in [1,2]_{\mathbb{R}}$ and converges to 1 as $t \to 2$. Moreover, $w(f)$ is positive on its support, which is the half open interval $I(f) = (T_{w}(f), 2]_{\mathbb{R}}$ where $T_{w}(f)$ is the last time that $r_t(f, T_w(f)) = 1$, or $I(f) = [1, 2]_{\mathbb{R}}$ in the case where $r_t(f, t) < 1$ for all t.

We now have that

$$
W(f,T) = \int_{t=1}^{T} w(f,t)dt
$$

is continuous as a function of T and strictly increasing on $I(f)$, which is the support of $W(f)$. Therefore, $T_1(f)$ is well defined. By Lebesgue's dominated convergence theorem $W(f_k, T) \to W(f_{\infty}, T)$. Since $w(f) < 1$, $W(f)$ is 1-Lipschitz continuous for all f, and since $W(f_k) \to W(f_{\infty})$ pointwise, by the Arzelà-Ascoli theorem, $W(f_k)$ converges uniformly. Therefore, $[W(f_k)]^{-1}$ converges uniformly, so $T_1(f_k) = [W(f_k)]^{-1} \left(\frac{1}{2}W(f_k, 2)\right) \rightarrow T_1(f_\infty)$.

Since $C_n(f_k, t_k) \to C_n(f_\infty, t_\infty)$ and $u_n(f_k, t_k) \to u_n(f_\infty, t_\infty)$, Rado's theorem implies that $h_n(f_k, t_k) \to h_n(f_\infty, t_\infty)$ uniformly. This also implies that $[h_n(f_k, T(f_k))]^{-1} \to [h_n(f_\infty, T(f_\infty))]^{-1}$ invariably.

We now show continuity of p_n . First, consider the case where $t_k \leq T_1(f_k)$ for all k sufficiently large. Then, $p_n(f_k, t_k) = p_{\text{AL},n}(f_k, t_k)$, so $p_n(f_k, t_k) \to p_n(f_{\infty}, t_{\infty})$. This also means that $p_n(f_k, T(f_k)) \to p_n(f_{\infty}, T(f_{\infty}))$. Next, consider the case where $t_k \leq T_1(f_k)$ for all k sufficiently large. Then,

$$
p_n(f_k, t_k) = [h_n(f_k, t_k) \circ [h_n(f_k, T_1(f_k))]^{-1}](p_n(f_k, T_1(f_k))),
$$

and $p_n(f_k, T_1(f_k)) \to p_n(f_\infty, T_1(f_\infty))$ and both $h_n(f_k, t_k)$ and $[h_n(f_k, T_1(f_k))]^{-1}$ converge invariably, so again we have $p_n(f_k, t_k) \to p_n(f_\infty, t_\infty)$. If neither of the above cases hold, then we can partition the sequence $p_n(f_k, t_k)$ depending on how t_k compares to $T_1(f_k)$, and since $p_n(f_k, t_k)$ converges to $p_n(f_\infty, t_\infty)$ for each part, we have $p_n(f_k, t_k) \to p_n(f_\infty, t_\infty)$ in all cases. Thus, $F_A(f_k, t_k) \to F_A(f_\infty, t_\infty)$ by Lemma [2.12](#page-14-4) and therefore $\rho(f_k, t_k) \to \rho(f_\infty, t_\infty)$ in the sup-metric. \Box

Proof of Lemma [3.4](#page-21-1) part [2.](#page-21-3) By Lemma [3.2,](#page-18-1) we have $S_A(Qfs, 1) = Q S_A(f, 1)$ and $C_n(Qfs, 1) =$ $QC_{sn}(f, 1)$. Hence, By Lemma [2.12,](#page-14-4) we have $F_A(Qfs, 1) = Q F_A(f, 1) \circ [z \mapsto ss^s]$ as in Equa-tion [1,](#page-19-0) so $L(Qfs) = sL(f)^s$, so $\gamma_L(Qfs; 0) = Q\gamma_L(f; 0)$ and $\gamma_A(Qfs; 0) = Q\gamma_A(f; 0)$. Since the curvature $\kappa \nu$ of a curve commutes with isometries of the sphere, we have for $t \in [0,\infty]_{\mathbb{R}}$ that $\gamma_L(Qfs;t) = Q\gamma_L(f;t)$ and $\gamma_A(Qfs;t) = Q\gamma_A(f;t)$, so for $t \in [1,2]_{\mathbb{R}}$ we have that $S_A(Qfs, t) = QS_A(f, t)$ and $S_L(Qfs, t) = QS_L(f, t)$. Hence, $C_n(Qfs, t) = QC_{sn}(f, t)$ and $u_n(Qfs) = Qu_{sn}(f).$

In the case where $Q \in SO_3$, both Q and Q^{-1} are conformal, so $Qh_{sn}(f,t)Q^{-1}$ is a conformal map. In the case where $Q \in -SO_3$, both Q and Q^{-1} are angle preserving and orientation

reversing, so again $Qh_{sn}(f, t)Q^{-1}$ is a conformal map. In both cases, we have a conformal map from $C_n(Qfs, 2)$ to $C_n(Qfs, t)$ that is fixed at $u_n(Qfs)$. By the chain rule,

$$
D(Qh_{sn}(f,t)Q^{-1};u_n(Qfs)) = D(Q;h_{sn}(f,t)(u_{sn}(f))) \circ D(h_{sn}(f,t);u_{sn}(f)) \circ D(Q^{-1};Qu_{sn}(f))
$$

= $Q \circ \lambda \operatorname{id} \circ Q^{-1}$
= $\lambda \operatorname{id}$

where $\lambda > 0$ is as given in the definition of h_{sn} . Hence, for all case of $Q \in O_3$ we have $h(Qfs, t) = Qh_{sn}(f, t)Q^{-1}$. Therefore, $r_h(Qfs, t) = r_h(f, t)$ and $r_u(Qfs, t) = r_u(f, t)$, so $w(Qfs, t) = w(f, t)$, which implies that $T_1(Qfs) = T_1(f)$.

By Lemma [3.2,](#page-18-1) we also have $p_n(Qfs, 1) = Qp_{sn}(f, 1)$ for $n \in \{1, -1\}$, so $p_n(Qfs, t) =$ $Qp_{sn}(f, t)$ in the case where $t \in [1, T_1(f)]_{\mathbb{R}}$ since $S_A(Qfs, t) = QS_A(f, t)$ and $S_L(Qfs, t) =$ $Q S_{L}(f, t)$, and likewise in the case where $t \in [T_1(f), 2]_{\mathbb{R}}$ since

$$
p_n(Qfs, t) = [h_n(Qfs, t)[h_n(Qfs, T_1(Qfs))]^{-1}](p_n(Qfs, T_1(Qfs))).
$$

\n
$$
= [Qh_{sn}(f, t)Q^{-1}Q[h_{sn}(f, T_1(f))]^{-1}Q^{-1}](Qp_{sn}(f, T_1(f))).
$$

\n
$$
= Q[h_{sn}(f, t)[h_{sn}(f, T_1(f))]^{-1}](p_{sn}(f, T_1(f))).
$$

\n
$$
= Qp_{sn}(f, t)
$$

Therefore, by lemma [2.12,](#page-14-4) $F_A(Qfs, t) = Q F_A(f, t) \circ [z \mapsto ss^s]$, so

$$
\rho(Qfs,t) = F_A(Qfs,t) \circ F_A(Qfs,1)^{-1} \circ \rho(Qfs,1)
$$

= $Q F_A(f,t) \circ [z \mapsto sz^s] \circ [z \mapsto sz^s]^{-1} \circ [F_A(f,1)]^{-1}Q^{-1} \circ Q\rho(f,1)s$
= $Q F_A(f,t) \circ F_A(f,1)^{-1} \circ \rho(f,1)s$
= $Q\rho(f,t)s$

Proof of Lemma [3.4](#page-21-1) part [3.](#page-21-4) Consider $f \in O_3$. By Lemma [3.2,](#page-18-1) we have $\rho(f, 1) = f$. Hence, $S_A(f, 1) = S_A(f, 0)$ is a great circle and $p_1(f, 1) = p_1(f, 0)$ and $p_{-1}(f, -1) = p_{-1}(f, 0)$ are antipodes, so by Lemma [2.7,](#page-9-1) $F_A(f, 1)$ is isometric to a stereographic projection, so $S_L(f, 1)$ is also a great circle. Since great circles have no geodesic curvature, the evolution of a great circle by curvature flow is trivial, so $S_A(f, t) = S_A(f, 1)$ and $S_L(f, t) = S_L(f, 1)$, so $u_n(f, t) = p_{n2}(f, 1)$ and $h_n(f, t) = id$, so $r_n(f, t) = 0$ and $r_n(f, t) = 0$, so $w(f, t) = 1$, so $T_1(f) = \frac{3}{2}$. Hence, $p_n(f, t) = p_n(f, 1)$ for $n \in \{1, -1\}$; in the case of $t \leq \frac{3}{2}$ because $S_A(f, t)$ and $S_{L}(f, t)$ remain constant in t, and in the case of $t > \frac{3}{2}$ because $h_n(f, t) = id$. Therefore, $F_A(f,t) = F_A(f,1)$, which implies that $\rho(f,t) = \rho(f,1) = f$. \Box

3.3 Aligning

In the third stage, we move the points p_1 and p_{-1} to antipodal positions.

For $t \in (2,3]_{\mathbb{R}}$, we define the following. Let $S_A(f,t) = S_A(f,2)$. Let $p_1(f,t)$ move at uniform speed along the arc from $p_1(f, 2)$ to $-p_{-1}(f, 2)$ that avoids $-p_1(f, 2)$ and reach the midpoint of this arc at $t = 3$, and analogously, let $p_{-1}(f, t)$ move at uniform speed along the arc from $p_{-1}(f, 2)$ to $-p_1(f, 2)$ that avoids $-p_{-1}(f, 2)$ and reach the midpoint of this arc at $t = 3$. Let

$$
\rho(f,t) = \mathcal{F}_{\mathcal{A}}(f,t) \circ \mathcal{F}_{\mathcal{A}}(f,2)^{-1} \circ \rho(f,2).
$$

Lemma 3.5. The points $p_1(f, 3), p_{-1}(f, 3)$ are antipodes.

Lemma 3.6. On $t \in [2,3]_{\mathbb{R}}$, the following hold.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$, and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in O_3$, then $\rho(f, t) = f$.

Proof of Lemma [3.5.](#page-25-0) In the case where $p_{-1}(f, 2) = -p_1(f, 2)$, both points $p_1(f, t)$ and $p_{-1}(f, t)$ remain constant, so $p_1(f, 3), p_{-1}(f, 3)$ are antipodes. Otherwise, the signed angle θ from $p_1(f, 2)$ to $-p_{-1}(f, 2)$ is the same as the signed angle from $-p_1(f, 2)$ to $p_{-1}(f, 2)$, and the signed angle from $p_1(f, 2)$ to $p_1(f, 3)$ as well as that from $-p_1(f, 2)$ to $p_{-1}(f, 3)$ is $\theta/2$, so again the points $p_1(f, 3), p_{-1}(f, 3)$ are antipodes. \Box

Proof of Lemma [3.6](#page-25-1) part [1.](#page-25-2) Consider $f_k \in \text{hom}(\mathbf{S}^2)$ and $t_k \in [2,3]_{\mathbb{R}}$ such that $f_k \to f_{\infty}$ uniformly and $t_k \to t_\infty$. By Lemma [3.4,](#page-21-1) $S_A(f_k, t_k) = S_A(f_k, 2) \to S_A(f_\infty, t_\infty)$ and $C_1(f_k, t_k) \to$ $C_1(f_\infty, t_\infty)$. Also, the point that is a $((t-2)/2)$ -fraction of the way along the arc from p_1 to $-p_{-1}$ varies continuously as a function of $p_1, -p_{-1}$, and t. Hence, $p_1(f_k, t_k) \to p_1(f_{\infty}, t_{\infty})$ and likewise for p_{-1} , so by Lemma [2.12](#page-14-4) $F_A(f_k, t_k) \to F_A(f_{\infty}, t_{\infty})$ uniformly, so by Lemma [2.1,](#page-4-0) $[F_A(f_k, 2)]^{-1} \to [F_A(f_{\infty}, 2)]^{-1}$ uniformly, and therefore $\rho(f_k, t_k) \to \rho(f_{\infty}, t_{\infty})$. \Box

Proof of Lemma [3.6](#page-25-1) part [2.](#page-25-3) By Lemma [3.4](#page-21-1) we have $S_A(Qfs, t) = S_A(Qfs, 2) = Q S_A(f, 2)$ $QS_A(f, t)$ and $C_1(Qfs, t) = QC_s(f, t)$. Also, $p_1(Qfs, 2) = Qp_s(f, 2)$ and $-p_{-1}(Qfs, 2)$ $-Qp_{-s}(f, 2)$, so the point $p_1(Qfs, t)$ that is a $((t - 2)/2)$ -fraction of the way along the arc from $p_1(Qfs, 2)$ to $-p_{-1}(Qfs, 2)$ is also the image by Q of the point $p_s(f, t)$ that is a ($(t-2)/2$)-fraction of the way along the arc from $p_s(f, 2)$ to $-p_{-s}(f, 2)$, which means that $p_1(Qfs, t) = Qp_s(f, t)$ and likewise for p_{-1} . Therefore, by Lemma [2.12,](#page-14-4) $F_A(Qfs, t)$ $Q F_A(f, t) \circ [z \mapsto s z^s],$ which implies that $\rho(Qfs, t) = Q \rho(f, t)s$. \Box

Proof of Lemma [3.6](#page-25-1) part [3.](#page-25-4) Suppose $f \in O_3$. Then, by Lemma [3.4,](#page-21-1) $\rho(f, 2) = f \in O_3$, so $p_1(f, 2)$ and $p_{-1}(f, 2)$ are antipodes, so $p_1(f, t) = p_1(f, 2)$ and $p_{-1}(f, t) = p_{-1}(f, 2)$ are constant in t, so $F_A(f,t) = F_A(f, 2)$ is constant in t, so $\rho(f,t) = \rho(f, 2) = f$. \Box

3.4 Flattening

In the fourth stage, we deform S_E to a great circle. In each hemisphere, we map S_E to a curve S_{O} in the disk, where we deform S_{O} to a line segment, and we let S_{E} be the preimage of S_O . We deform S_O to a segment by continuously shrinking S_O to the origin and connecting the end points to the boundary of the disk by segments; see Figure [2.](#page-16-0)

For $t \in (3, 4]_{\mathbb{R}}$, we define the following. Let $p_i(f, t) = p_i(f, 3)$ for $i \in \{1, -1\}$ and $r(t) = 4-t$.

Observe r sends $[3, 4]_{\mathbb{R}}$ to $[1, 0]_{\mathbb{R}}$. Let

$$
S_O(f,3) = [F_A(f,3)]^{-1}(S_E(f,3)),
$$

\n
$$
S_O(f,t) = \bigcup \left\{ \begin{array}{c} r(t)(S_O(f,3) \cap D), & \frac{1}{r(t)}(S_O(f,3) \setminus D), \\ \left[\frac{-1}{r(t)}, -r(t)\right]_{\mathbb{R}}, & \left[r(t), \frac{1}{r(t)}\right]_{\mathbb{R}} \end{array} \right\},
$$

\n
$$
S_E(f,t) = [F_A(f,3)](S_O(f,t)),
$$

\n
$$
p_2(f,t) = [F_A(f,3) \circ r(t)[F_A(f,3)]^{-1}](p_2(f,3)),
$$

\n
$$
p_{-2}(f,t) = \left[F_A(f,3) \circ \frac{1}{r(t)}[F_A(f,3)]^{-1}\right](p_{-2}(f,3)),
$$

\n
$$
\rho(f,t) = F_E(f,t) \circ F_E(f,3)^{-1} \circ \rho(f,3).
$$

Lemma 3.7. The curve $S_E(f, 4)$ is a great circle, and $\rho(f, t)$ is well defined for $t \in (3, 4]$.

Lemma 3.8. On $t \in [3, 4]_{\mathbb{R}}$, the following hold.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$, and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in O_3$, then $\rho(f, t) = f$.

Proof of Lemma [3.7.](#page-26-0) Since $S_E(f, 3)$ and $S_A(f, 3)$ intersect in exactly the pair of points $\{p_1, p_{-1}\}(f, 3)$, the curve $S_O(f, 3)$ consists of a pair of curves from 1 to -1; one going through D and the other going through $\mathbb{C} \setminus \mathbf{D}$. Therefore, $S_{\mathcal{O}}(f, t)$ consists of a curve from $-r(t)$ to $r(t)$ through $r(t)$ **D**, and a segment from $r(t)$ to $\frac{1}{r(t)}$, and a curve from $\frac{1}{r(t)}$ to $\frac{-1}{r(t)}$ through $\overline{C \setminus \frac{1}{r(t)} \mathbf{D}}$, and a segment from $\frac{-1}{r(t)}$ back to $-r(t)$. Hence, $S_O(f, t)$ is a simple closed curve in $\overline{\mathbb{C}}$, so $S_E(f, t)$ is a simple closed curve.

By definition, we have $p_2(f, 3) \in S_E(f, 3) \cap C_1(f, 3)^\circ$, so $[F_A(f, 3)]^{-1}(p_2(f, 3)) \in S_O(f, 3) \cap D^\circ$, so $r(t)[F_A(f,3)]^{-1}(p_2(f,3)) \in r(t)(S_O(f,t)∩D°) \subset S_O(f,t)∩C_1(f,3)°$, so $p_2(f,t) ∈ S_E(f,t)∩C_2(f,3)$ $C_1(f,3)^\circ$. Similarly, $p_{-2}(f,t) \in S_E(f,t) \cap C_{-1}(f,3)^\circ$. Hence, the points p_1, p_2, p_{-1}, p_{-2} appear in that order around S_E , so $F_E(f, t)$ is well defined, and therefore $\rho(f, t)$ is well defined.

The curve $S_A(f, 4) = S_A(f, 2)$ is a great circle by Lemma [3.3,](#page-21-0) and the points $p_1(f, 4) =$ $p_1(f, 3)$ and $p_{-1}(f, 4) = p_{-1}(f, 3)$ are antipodal by Lemma [3.5.](#page-25-0) Therefore, by Lemma [2.12,](#page-14-4) $F_A(f, 4)$ is isometric to a stereographic projection, and by definition $S_O(f, 4) = \mathbb{R}$, so $S_E(f, 4) = F_A(f, 4; S_O(f, 4))$ is a great circle. \Box

Proof of Lemma [3.8](#page-26-1) part [1.](#page-26-2) Consider $f_k \in \text{hom}(\mathbf{S}^2)$ and $t_k \in [3, 4]_{\mathbb{R}}$ such that $f_k \to f_{\infty}$ uniformly and $t_k \to t_\infty$. By Lemma [3.6,](#page-25-1) $\rho(f_k, 3) \to \rho(f_\infty, 3)$ uniformly, so $p_i(f_k, 3) \to p_i(f_\infty, 3)$ and $S_E(f_k, 3) \to S_E(f_\infty, 3)$ and $S_A(f_k, 3) \to S_A(f_\infty, 3)$ in Fréchet distance, so by Lemma [2.12,](#page-14-4) $F_A(f_k, 3) \to F_A(f_\infty, 3)$ uniformly, so by Lemma [2.1,](#page-4-0) $[F_A(f_k, 3)]^{-1} \to [F_A(f_\infty, 3)]^{-1}$ uniformly, and therefore $S_O(f_k, 3) = [F_A(f_k, 3)]^{-1}(S_E(f_k, 3)) \rightarrow S_O(f_{\infty}, 3)$ in Fréchet distance.

Each part of S_O is composed of continuous functions, so $S_O(f_k, t_k) \to S_O(f_\infty, t_\infty)$ in Fréchet distance, and $F_A(f, 3)$ is uniformly continuous by the Heine-Cantor theorem, so $S_E(f_k, t_k) \to$

 $S_E(f_\infty, t_\infty)$ in Fréchet distance. Also, $p_n(f_k, t_k) \to p_n(f_\infty, t_\infty)$ for $n \in \{1, -1, 2, -2\}$, so by Lemma [2.11,](#page-13-4) $F_E(f_k, t_k) \to F_E(f_\infty, t_\infty)$ uniformly. Therefore, $\rho(f_k, t_k) \to \rho(f_\infty, t_\infty)$ by Lemma [2.1.](#page-4-0) \Box

Proof of Lemma [3.8](#page-26-1) part [2.](#page-26-3) By Lemma [3.6,](#page-25-1) $\rho(Qfs, 3) = Q\rho(f, 3)s$, so $S_{E}(Qfs, 3) = QS_{E}(f, 3)$ and $S_A(Qfs, 3) = QS_A(f, 3)$ and $p_i(Qfs, 3) = p_{si}(f, 3)$, so by Lemma [2.12,](#page-14-4) $F_A(Qfs, 3) =$ $Q F_A(f,3) \circ [z \mapsto s z^s]$. Hence, $S_O(Qfs,3) = s S_O(f,3)^s$. In the case where $s = 1$ we have $r(t)(S_O(Qfs, t) \cap D) = r(t)(S_O(f, t) \cap D)$ and analogously the other parts of S_O are unchanged, so $S_O(Qfs, t) = S_O(f, t) = sS_O(f, t)^s$. In the case where $s = -1$ we have $r(t)(S_O(Qfs, t) \cap D) = r(t)(sS_O(f, t)^s \cap D) = s(\frac{1}{r(t)})$ $\frac{1}{r(t)}(\overline{\mathrm{S}_{\mathrm{O}}(f,t)\setminus \mathbf{D}}))^{s}$, and the other parts of S_O are permuted and transformed analogously, so $S_O(Qfs, t) = s S_O(f, t)^s$. Hence, $S_{\mathcal{E}}(Qfs,t) = Q F_{\mathcal{A}}(f,3) \circ [z \mapsto ss^s](s S_{\mathcal{O}}(f,t)^s) = Q S_{\mathcal{E}}(f,t)$, and $p_2(Qfs,t) = Q p_{s2}(f,t)$ and analogously for p_{-2}, p_1, p_{-1} , so by Lemma [2.11,](#page-13-4) $F_E(Qfs, t) = Q F_E(f, t)s$. Thus, $\rho(Qfs, t) =$ $Q\rho(Q,t)s.$ \Box

Proof of Lemma [3.8](#page-26-1) part [3.](#page-26-4) Suppose $f \in O_3$. Then, by Lemma [3.6,](#page-25-1) $\rho(f, 3) = f$, so $S_E(f, 3) =$ $f(e_3^{\perp})$ is a great circle and the 4 points $p_n(f, 3) = f(e_n)$ for $n \in \{1, 2, -1, -2\}$ are spaced uniformly at right angles around $S_E(f, 3)$. By Lemma [2.11,](#page-13-4) $F_E(f, 3)$ is f composed with stereographic projection through e_{-3} , so $S_A(f, 3)$ is a great circle that is perpendicular to $S_E(f, 3)$, so by Lemma [2.12,](#page-14-4) $F_A(f, 3)$ is isometric to stereographic projection. Since the map $F_A(f,3)^{-1}$ sends $S_A(f,3)$ to $S^1 \subset \mathbb{C}$ and sends the point p_1 to 1, the map $F_A(f,3)^{-1}$ sends $S_E(f,3)$ to the line in $\overline{\mathbb{C}}$ that passes though S^1 at 1 at a right angle. That is, $F_A(f,3)^{-1}$ sends $S_{E}(f,3)$ to $\overline{\mathbb{R}}$. Hence, $S_{O}(f,t) = \overline{\mathbb{R}}$ for $t \in [3,4]_{\mathbb{R}}$, so $S_{E}(f,t) = f(e_{3}^{\perp})$. Also, $p_{2}(f,3)$ is orthogonal to $p_1(f,3)$, so $[F_A(f,3)]^{-1}(p_2(f,3)) = 0$, so $p_2(f,t) = F_A(f,3;0) = p_2(f,3)$, and similarly $p_{-2}(f, t) = p_{-2}(f, 3)$, so $F_E(f, t) = F_E(f, 3)$. Therefore, $\rho(f, t) = \rho(f, 3) = f$. \Box

3.5 Divvying

In the fifth stage, we deform the map to an isometry on the equator by linear interpolation in polar coordinates; see Figure [3.](#page-16-0)

For $t \in (4, 5]_{\mathbb{R}}$, we define the following. Let $Q = Q_{fin}(f) \in O_3$ such that $Qe_1 = p_1(f, 4)$, $Qe_2 = p_2(f, 4)$, and Q has the same orientation as f. Eventually we will deformation retract to Q.

Let

$$
\rho(f, t; x) = \left[\mathbf{F}_{\mathcal{E}}(f, 4) \right] \left(z_1^{4-t} z_0^{5-t} \right)
$$

where

$$
z_0 = \zeta_0(f; x) = [[\mathbf{F}_E(f, 4)]^{-1} \circ \rho(f, 4)](x),
$$

\n
$$
z_1 = \zeta_1(f; x) = \begin{cases} z_0 \text{ if } z_0 \in \{0, \infty\} \\ |z_0| [[\mathbf{F}_E(f, 4)]^{-1} \circ Q_{\text{fin}}(f) \circ [\rho(f, 4)]^{-1} \circ \mathbf{F}_E(f, 4)] (z_0/|z_0|) \text{ else,} \\ z_1^{4-t} z_0^{5-t} = e^{(4-t) \log(z_1) + (5-t) \log(z_0)} \end{cases}
$$

with the branch of the log with range $(-\infty,\infty)_{\mathbb{R}} + i[0,2\pi)_{\mathbb{R}}$. Note that the cases in the definition of z_1 coincide if we simply regard 0 or ∞ times an undefined unit complex number as 0 or ∞ respectively.

Lemma 3.9. For $t \in (4, 5]_{\mathbb{R}}$, $\rho(f, t) \in \text{hom}(\mathbf{S}^2)$. Also, for $x \in e_3^{\perp}$, $\rho(f, 5; x) = [Q_{\text{fin}}(f)](x)$.

Lemma 3.10. On $t \in [4, 5]_{\mathbb{R}}$, the following hold.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$ and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in O_3$, then $\rho(f,t) = f$.

Proof of Lemma [3.9.](#page-28-0) For $x \in e_3^{\perp}$, we have $\rho(f, 4; x) \in S_E(f, 4)$, so $z_0 \in S^1$, so $|z_0| = 1$. Hence, $\zeta_1(f; x)$ restricted to $x \in e_3^{\perp}$ is a composition of homeomorphisms; namely

$$
\zeta_1(f; x) = \left[\mathcal{F}_E(f, 4) \right]^{-1} \circ Q_{fin}(f) \circ [\rho(f, 4)]^{-1} \circ \mathcal{F}_E(f, 4) \right] \left(\left[\left[\mathcal{F}_E(f, 4) \right]^{-1} \circ \rho(f, 4) \right] (x) \right)
$$

=
$$
\left[\mathcal{F}_E(f, 4) \right]^{-1} \circ Q_{fin}(f) \left[(x) \right].
$$

Likewise, the map $\zeta_1(f) \circ [\zeta_0(f)]^{-1}$, which sends z_0 to z_1 , is a self-homeomorphism of S^1 , and since $Q_{fin}(f)$ and $\rho(f, 4)$ have the same orientation, this map is orientation preserving. Also,

$$
[\zeta_1(f) \circ [\zeta_0(f)]^{-1}](1) = [[\mathbf{F}_E(f, 4)]^{-1} \circ Q_{fin}(f) \circ [\rho(f, 4)]^{-1} \circ \mathbf{F}_E(f, 4)](1)
$$

\n
$$
= [[\mathbf{F}_E(f, 4)]^{-1} \circ Q_{fin}(f) \circ [\rho(f, 4)]^{-1}]](p_1(f, 4))
$$

\n
$$
= \mathbf{F}_E(f, 4; p_1(f, 4))
$$

\n
$$
= 1,
$$

so the phase of z_1 as a function of the phase of z_0 , which is a the map given by $\zeta_{\rm ph}(\theta)$ = $[\theta \mapsto -i[\log \circ \zeta_1(f) \circ [\zeta_0(f)]^{-1}](e^{i\theta})]$, is strictly increasing, and the restriction to $(0, 2\pi)_{\mathbb{R}}$ is a self-homeomorphism of $(0, 2\pi)_{\mathbb{R}}$, provided that we choose the appropriate branch of the log. Consider the map

$$
\widetilde{\rho}(f,t) = \left[\mathbf{F}_{\mathcal{E}}(f,4)\right]^{-1} \circ \rho(f,t) \circ [\rho(f,4)]^{-1} \circ \mathbf{F}_{\mathcal{E}}(f,4),
$$

which sends z_0 to $z_1^{4-t}z_0^{5-t}$. The phase of $\tilde{\rho}(f, t; z_0)$ as a function of the phase of z_0 , which
is given by β to $\tilde{\rho}(f, t; c^{i\theta}) = (4-1)\zeta_1(\theta) + (5-t)\theta$ is also a strictly increasing is given by $[\theta \mapsto -i \log(\tilde{\rho}(f, t; e^{i\theta})) = (4 - 1)\zeta_{ph}(\theta) + (5 - t)\theta$, is also a strictly increasing
solf homomorphism $(0, 2\pi)$. Hones the restriction of $\tilde{\rho}(f, t)$ to S^1 is a solf homomorphism self-homeomorphism $(0, 2\pi)_\mathbb{R}$. Hence, the restriction of $\tilde{\rho}(f, t)$ to S^1 is a self-homeomorphism of S^1 . The change in $\tilde{\rho}(f, t)$ with respect to change in magnitude is defined to be the identity. of S^1 . The change in $\tilde{\rho}(f, t)$ with respect to change in magnitude is defined to be the identity,
so $\tilde{\rho}(f, t)$ is a self-homomorphism of $\overline{\mathcal{C}}$ with the metric induced by stepsorphic projection. so $\tilde{\rho}(f, t)$ is a self-homeomorphism of $\overline{\mathbb{C}}$ with the metric induced by stereographic projection. Thus, $\rho(f, t) \in \text{hom}(\mathbf{S}^2)$, as it should be.

For the last part of the lemma, we have

$$
\rho(f, 5; x) = [F_{E}(f, 4)] (z_{1})
$$

=
$$
[F_{E}(f, 4)] ([F_{E}(f, 4)]^{-1} \circ Q_{fin}(f)](x))
$$

=
$$
[Q_{fin}(f)](x).
$$

Proof of Lemma [3.10](#page-28-1) part [1.](#page-28-2) If we extend the definition above for $\rho(f,t)$ on $t \in (4,5]_{\mathbb{R}}$ to $t = 4$, we get

$$
\begin{aligned} \left[\mathbf{F}_{\mathcal{E}}(f,4) \right] \left(z_1^{4-t} z_0^{5-t} \right) &= \left[\mathbf{F}_{\mathcal{E}}(f,4) \right] (z_0) \\ &= \left[\left[\mathbf{F}_{\mathcal{E}}(f,4) \right] \circ \left[\mathbf{F}_{\mathcal{E}}(f,4) \right]^{-1} \circ \rho(f,4) \right] (x) \\ &= \rho(f,4;x). \end{aligned}
$$

Thus, it suffices to show that $\rho(f, t)$ continuous on $t \in [4, 5]_{\mathbb{R}}$ using the definition above. Let

$$
\varphi_{Q}(f,t) = Q_{fin}(f)^{-1} \circ \rho(f,t) \circ [\rho(f,4)]^{-1} \circ Q_{fin}(f)
$$

= $[Q_{fin}(f)^{-1} \circ F_{E}(f,4)] \circ \tilde{\rho}(f,t) \circ [Q_{fin}(f)^{-1} \circ F_{E}(f,4)]^{-1}.$

On the restriction to e_3^{\perp} , $\rho_Q(f,t)$ is a composition of functions and inverses of functions that depend continuously in the sup-metric on f and t, so by Lemma [2.1,](#page-4-0) the map $\rho_{\mathcal{Q}}(f, t)$ is also continuous in the sup-metric on the restriction to e_3^{\perp} . Since change in $\tilde{\rho}(f, t)$ with respect to change in magnitude is the identity and $O_1(f)^{-1} \circ F_1(f, t)$ is the storeographic projection change in magnitude is the identity, and $Q_{fin}(f)^{-1} \circ F_E(f, 4)$ is the stereographic projection though e_3 , we have that the change in φ_Q with respect to change along a great circle though e_3 is the identity. Hence, $\rho_Q(f,t)$ is continuous in the sup-metric on all of S^2 . Thus, $\rho(f,t)$ \Box is continuous in the sup-metric

Proof of Lemma [3.10](#page-28-1) part [2.](#page-28-3) For arbitrary $Q \in O_3$ and $t \in [4, 5]$ _R we have the following.

$$
\begin{aligned} \zeta_0(Qfs) &= [\mathcal{F}_E(Qfs, 4)]^{-1} \circ \rho(Qfs, 4) \\ &= s[\mathcal{F}_E(f, 4)]^{-1} Q^{-1} \circ Q\rho(f, 4)s \\ &= s\zeta_0(f)s. \end{aligned}
$$

Since $p_i(Qfs, 4) = Qp_{si}(f, 4)$, and $QQ_{fin}(f)s$ sends e_i to $Qp_{si}(f, 4)$ for $i \in \{1, 2, -1, -2\}$ and has the same orientation as Qfs , we have $Q_{fin}(Qfs) = QQ_{fin}(f)s$.

$$
\zeta_1(Qfs;x) = |\zeta_0(Qfs;x)|[[\mathbf{F}_{\mathbf{E}}(Qfs,4)]^{-1} \circ Q_{\text{fin}}(Qfs) \circ [\rho(Qfs,4)]^{-1} \circ \mathbf{F}_{\mathbf{E}}(Qfs,4)] \left(\frac{\zeta_0(Qfs;x)}{|\zeta_0(Qfs;x)|}\right)
$$

= $|\zeta_0(f;sx)|[s[\mathbf{F}_{\mathbf{E}}(f,4)]^{-1}Q^{-1} \circ QQ_{\text{fin}}(f)s \circ s[\rho(f,4)]^{-1}Q^{-1} \circ Q \mathbf{F}_{\mathbf{E}}(f,4)s] \left(\frac{s\zeta_0(f;sx)}{|\zeta_0(f;sx)|}\right)$
= $s\zeta_1(f;sx)$

$$
\rho(Qfs, t; x) = \left[\mathcal{F}_{\mathcal{E}}(Qfs, 4) \right] \left(e^{(4-1)\log(\zeta_1(Qfs;x)) + (5-t)\log(\zeta_0(Qfs;x))} \right)
$$

\n
$$
= \left[Q \mathcal{F}_{\mathcal{E}}(f, 4)s \right] \left(e^{(4-1)\log(s\zeta_1(f;sx)) + (5-t)\log(s\zeta_0(f;sx))} \right)
$$

\n
$$
= \left[Q \mathcal{F}_{\mathcal{E}}(f, 4)s \right] \left(e^{(4-1)\log(\zeta_1(f;sx)) + (5-t)\log(\zeta_0(f;sx)) + \log(s)} \right)
$$

\n
$$
= Q\rho(f, t; sx).
$$

Proof of Lemma [3.10](#page-28-1) part [3.](#page-28-4) Suppose $f \in O_3$. Then, by Lemma [3.8,](#page-26-1) $\rho(f, 4) = f$, and since $\rho(f, 4)$ satisfies the defining properties of Q_{fin} , we have $Q_{fin}(f) = f$ as well, so $z_1 = z_0$, so $z_1^{4-t}z_0^{4-t} = z_0$, so $\rho(f,t) = \rho(f,4) = f$.

3.6 Combing

In the sixth stage, we use Alexander's trick on each hemisphere to deform the map to an isometry; see Figure [3.](#page-16-0)

For $t \in (5, 6]_{\mathbb{R}}$, we define the following. Let

$$
J(f,r;z) = \begin{cases} rg\left(f; \frac{z}{r}\right) & 0 \le |z| < r\\ z & r \le |z| \le \frac{1}{r}\\ \frac{1}{r}g(f;rz) & \frac{1}{r} < |z| \le \infty \end{cases}
$$

where $g(f) = [F_{E}(f, 4)]^{-1} \circ \rho(f, 5) \circ [Q_{fin}(f)]^{-1} \circ F_{E}(f, 4)$, and let

$$
\rho(f,t) = \mathcal{F}_{\mathcal{E}}(f,4) \circ J(f,6-t) \circ [\mathcal{F}_{\mathcal{E}}(f,4)]^{-1} \circ Q_{\text{fin}}(f).
$$

Lemma 3.11. For $t \in (5,6]_{\mathbb{R}}$, $\rho(f,t) \in \text{hom}(S^2)$.

Lemma 3.12. $\rho(f, 6) = Q_{fin}(f) \in O_3$.

Lemma 3.13. On $t \in [4, 5]_{\mathbb{R}}$, the following hold.

- 1. In the sup-metric, $\rho(f, t)$ depends continuously on f and t.
- 2. For all $Q \in \mathcal{O}_3$ and $s \in \{1, -1\}$, we have $\rho(Qfs, t) = Q\rho(f, t)s$.
- 3. If $f \in O_3$, then $\rho(f, t) = f$.

Proof of Lemma 3.11. We claim that the restriction of $g(f)$ to S^1 is the identity function. For $z \in \mathbf{S}^1$, we have $F_A(f, 4; z) \in S_A(f, 4)$, so $[[Q_{fin}(f)]^{-1} \circ F_A(f, 4)](z) \in e_3^{-1}$, so by Lemma [3.9,](#page-28-0) $[\rho(f, 5) \circ [Q_{fin}(f)]^{-1} \circ F_E(f, 4)](z) = F_E(f, 4; z),$ so $g(f; z) = [[F_E(f, 4)]^{-1} \circ F_E(f, 4)](z) = z$, so the claim holds.

Hence, if $|z| = r$ then $rg(f; \frac{z}{r})$ $(\frac{z}{r}) = z$, and if $|z| = \frac{1}{r}$ $\frac{1}{r}$ then $\frac{1}{r}g(f;rz) = z$. Therefore, $J(f,r)$ is continuous along the boundary between cases, and similarly for $[J(f,r)]^{-1}$, so $J(f,r)$ is a homeomorphism, so $\rho(f, t)$ is a homeomorphism. \Box

Proof of Lemma [3.12.](#page-30-1) At $t = 6$, we have $J(f, 0) = id_{\mathbb{C}}$, so $\rho(f, 6) = F_{\mathbb{E}}(f, 4) \circ [F_{\mathbb{E}}(f, 4)]^{-1} \circ$ $Q_{fin}(f) = Q_{fin}(f).$ \Box

Proof of Lemma [3.13](#page-30-2) part [1.](#page-30-3) By Lemmas [2.3,](#page-5-7) [3.8,](#page-26-1) and [3.10,](#page-28-1) $g(f)$ depends continuously on f, so $J(f,r)$ also depends continuously on f and r, so $\rho(f,t)$ depends continuously on f and t. \Box

Proof of Lemma [3.13](#page-30-2) part [2.](#page-30-4) For $Q \in O_3$ and $t \in (5, 6]$ _R, we have the following.

$$
g(Qfs) = [F_E(Qfs, 4)]^{-1} \circ \rho(Qfs, 5) \circ [Q_{fin}(Qfs)]^{-1} \circ F_E(Qfs, 4)
$$

= $s[F_E(f, 4)]^{-1}Q^{-1} \circ Q\rho(f, 5)s \circ s[Q_{fin}(f)]^{-1}Q^{-1} \circ QF_E(f, 4)s$
= $sg(f)s$,

so $J(Qfs, r) = sJ(f, r)s$, so

$$
\rho(Qfs,t) = \mathcal{F}_{\mathcal{E}}(Qfs,4) \circ J(Qfs,6-t) \circ [\mathcal{F}_{\mathcal{E}}(Qfs,4)]^{-1} \circ Q_{\text{fin}}(Qfs)
$$

= $Q \mathcal{F}_{\mathcal{E}}(f,4)s \circ sJ(f,6-t)s \circ s[\mathcal{F}_{\mathcal{E}}(f,4)]^{-1}Q^{-1} \circ QQ_{\text{fin}}(f)s$
= $Q\rho(f,t)s$.

Proof of Lemma [3.13](#page-30-2) part [3.](#page-30-5) Suppose $f \in O_3$. Then, $\rho(f, 5) = Q_{fin}(f) = f$, so $g(f) = id_{\mathbb{C}}$, so $J(f,r) = id_{\mathbb{C}}$, so $\rho(f,t) = f$. \Box

3.7 Putting it together

Proof of Theorem [1.1.](#page-0-1) By Lemma [3.2](#page-18-1) Part [4,](#page-18-5) we have $\rho(f, 0) = f$, so by Part [1](#page-18-2) of Lemmas [3.2,](#page-18-1) [3.4,](#page-21-1) [3.6,](#page-25-1) [3.8,](#page-26-1) [3.10,](#page-28-1) [3.13,](#page-30-2) ρ is a continuous deformation of hom(S^2). By Part [3](#page-18-4) of the above lemmas and Lemma [3.12,](#page-30-1) ρ is a strong deformation retraction to O_3 , and by Part [2](#page-18-3) of the above lemmas, ρ is $(O_3 \times Z_2)$ -equivariant. \Box

4 Nullity preserving homeomorphisms

The goal of this section is to prove Theorem [1.3.](#page-1-0) The construction will be similar, except that we will use a different definition for F_{stitch} . We will first define a function F_{radmul} that extends a homeomorphism f of the circle to a homeomorphism the disk that is smooth almost everywhere. We will extend f to the interior of the disk radially, but on each concentric circle, we mollify f by convolution with a smooth bump function that approaches the Dirac delta distribution at the boundary of the disk.

Let bump(ε) be a normalized smooth bump function with support on $[-\varepsilon, \varepsilon]_{\mathbb{R}}$. Specifically, let

$$
bump(\varepsilon; x) = \begin{cases} \frac{1}{c\varepsilon} e^{\frac{-1}{1-(\frac{x}{\varepsilon})^2}} & |x| < \varepsilon \\ 0 & |x| \ge \varepsilon \end{cases}
$$

where $c = \int_{-1}^{1} e^{\frac{-1}{1-x^2}} dx$.

Given a homeomorphism $f: \mathbf{S}^1 \to \mathbf{S}^1$ such that $f(1) = 1$, let $F_{\text{radmul}}: \mathbf{D} \to \mathbf{D}$ be defined as follows. Let $g(\theta) = -i \log (f(e^{i\theta}))$, and for $r \in (0,1)_{\mathbb{R}}$ let

$$
F_{\text{radmul}}(f; r e^{i\theta}) = r e^{i[\text{bump}(1-r)*g](\theta)}
$$

=
$$
r \exp\left(\frac{1}{c(1-r)} \int_{r-1}^{1-r} e^{\frac{-1}{1-(\frac{x}{1-r})^2}} \log\left(f\left(e^{i(\theta-x)}\right)\right) dx\right),
$$

and $F_{\text{radmul}}(f; e^{i\theta}) = f(e^{i\theta})$ and $F_{\text{radmul}}(f; 0) = 0$.

Lemma 4.1. If f is a homeomorphism of the circle, then $F_{radmul}(f)$ satisfies the following.

- 1. $F_{\text{radmul}}(f)$ preserves nullity.
- 2. $F_{radmul}(f)$ in the sup-metric depends continuously on f in the sup-metric.

3. $F_{\text{radmul}}(f \circ [z \mapsto 1/z]) = F_{\text{radmul}}(f \circ [z \mapsto \overline{z}]) = F_{\text{radmul}}(f) \circ [z \mapsto \overline{z}]$

Proof. Observe that $[\text{bump}(\varepsilon)*g](\theta) \in [\text{min}, \text{max}]_{\mathbb{R}}(g([\theta-\varepsilon,\theta+\varepsilon]_{\mathbb{R}}))$ by comparing integrals, since for any constant function c we have $[\text{bump}(\varepsilon)*c]=c$. Hence, $F_{\text{radmul}}(f)$ is continuous at the boundary of the disk.

Suppose that f is orientation preserving. Then, g is strictly increasing and quasiperiodic with $q(\theta + 2\pi) = q(\theta) + 2\pi$. Hence, bump $(\varepsilon) * q$ is also strictly increasing and quasiperiodic with $[\text{bump}(\varepsilon)*g](\theta+2\pi) = [\text{bump}(\varepsilon)*g](\theta)+2\pi$. Hence, $e^{i[\text{bump}(\varepsilon)*g](\theta)}$ is an orientation preserving homeomorphism of S^1 , so $F_{\text{radmul}}(f)$ is injective. Furthermore, ∂_{θ} bump $(\varepsilon; \theta)$ is positive on $(-\varepsilon, 0)_{\mathbb{R}}$ and negative on $(0, \varepsilon)_{\mathbb{R}}$, and since $g(\theta + \varepsilon) > g(\theta)$, this implies $\partial_{\theta}[\text{bump}(\varepsilon) * g](\theta) = [\partial_{\theta} \text{bump}(\varepsilon;\theta)] * g$ is always positive. Similarly, $F_{\text{radmul}}(f)$ is injective and ∂_{θ} [bump(ε) * g](θ) is always negative in the case where f is orientation reversing.

In either case, $\partial_{\theta} F_{\text{radmul}}(f; r e^{i\theta})$ never vanishes for $r \in (0, 1)_{\mathbb{R}}$. Also, $\partial_r |F_{\text{radmul}}(f; r e^{i\theta})|$ = $\partial_r r = 1$, so $\partial_r F_{\text{radmul}}(f; r e^{i\theta})$ never vanishes. By the dominated convergence theorem, the partial derivatives of $F_{radmul}(f)$ are continuous, so $F_{radmul}(f)$ is continuously differentiable on $\mathbf{D}^{\circ} \setminus 0$. Since $F_{\text{radmul}}(f)$ has no critical points in $\mathbf{D}^{\circ} \setminus 0$, $F_{\text{radmul}}(f)^{-1}$ is continuously differentiable on $\mathbf{D}^{\circ} \setminus 0$ by the inverse function theorem. Hence, both $F_{\text{radmul}}(f)$ and $F_{\text{radmul}}(f)^{-1}$ send null sets to null sets, so the first part holds.

For the second part, consider $\varepsilon > 0$ and f_1, f_∞ such that $|f_1 - f_\infty| < \varepsilon$, and let g_k be defined analogously. Then, $|g_1-g_\infty| < 2\varepsilon$, so for all r, we have $|[\text{bump}(1-r)*g_1] - [\text{bump}(1-r)*g_\infty]| =$ $|\text{bump}(1-r)*(g_1-g_{\infty})| < 2\varepsilon$, so $|F_{\text{radmul}}(f_1) - F_{\text{radmul}}(f_{\infty})| < 2\varepsilon$. Hence, $F_{\text{radmul}}(f)$ in the sup-metric depends continuously on f in the sup-metric.

For the third part, observe that the maps $[z \mapsto 1/z]$ and $[z \mapsto \overline{z}]$ both act on S^1 by reflecting the circle across the real line, so these are actually the same map, so the first equality holds. Also, $-i \log \left(\left[f \circ [z \mapsto 1/z] \right] (e^{i\theta}) \right) = -i \log \left(f (e^{-i\theta}) \right) = g(-\theta)$, and since bump $(1 - r)$ is symmetric about 0, we have

$$
F_{\text{radmul}}(f \circ [z \mapsto 1/z]; re^{i\theta}) = re^{i[\text{bump}(1-r)*(g\circ[\psi \mapsto -\psi])](\theta)}
$$

= $re^{i[\text{bump}(1-r)*g](-\theta)}$
= $F_{\text{radmul}}(f; re^{i(-\theta)})$
= $F_{\text{radmul}}(f; \overline{re^{i\theta}}),$

so $F_{\text{radmul}}(f \circ [z \mapsto 1/z]) = F_{\text{radmul}}(f) \circ [z \mapsto \overline{z}].$

We redefine $F_{\text{stitch}}(f_1, f_2)$ as follows. Given embeddings $f_i : \mathbf{D} \to \mathbf{S}^2$ with $p_1 = f_1(1) = f_2(1)$, $p_{-1} = f_1(-1) = f_2(-1)$, and $S = f_1(S^1) = f_2(S^1)$, let f_0 be the same as before, which is

$$
f_0^{-1}: S \to \mathbf{S}^1
$$
, $f_0^{-1}(x) = \sqrt{\frac{f_1^{-1}(x)}{f_2^{-1}(x)}}$,

 \Box

and let

$$
F_{\text{stitch}}(f_1, f_2; z) = \begin{cases} [f_1 \circ F_{\text{radmul}}(f_1^{-1} \circ f_0)](z) & |z| < 1\\ f_0(z) & |z| = 1\\ [f_2 \circ F_{\text{radmul}}(f_2^{-1} \circ f_0)](1/\bar{z}) & |z| > 1. \end{cases} \tag{2}
$$

Lemma 4.2. Lemma [2.10](#page-11-0) also holds for F_{stitch} redefined by [\(2\)](#page-33-0). Additionally:

5. If both f_1 and f_2 preserve nullity, then $F_{\text{stitch}}(f_1, f_2)$ preserves nullity.

Proof. Observe that in the case where $|z|=1$, we have

$$
f_1 \circ F_{\text{radmul}}(f_1^{-1} \circ f_0)(z) = [f_1 \circ f_1^{-1} \circ f_0](z) = f_0(z),
$$

$$
[f_2 \circ F_{\text{radmul}}(f_2^{-1} \circ f_0)](1/\overline{z}) = [f_2 \circ f_2^{-1} \circ f_0](z) = f_0(z),
$$

so the defining formulas for $F_{\text{stitch}}(f_1, f_2)$ in all 3 case coincide when $|z| = 1$. Hence, $F_{\text{stitch}}(f_1, f_2)$ is indeed a homeomorphism, and part [5](#page-33-1) holds by Lemma [4.1.](#page-31-1)

Observe that f_0 , i.e., the restriction of $F_{\text{stitch}}(f_1, f_2)$ to the unit circle, is defined in the same way as before, so Lemma [2.10](#page-11-0) still holds for f_0 . Consider embeddings $f_{i,k}$ in the domain of F_{stitch} such that $f_{i,k} \to f_{i,\infty}$ uniformly. Just as before $f_{i,k}^{-1} \circ f_{0,k} \to f_{i,\infty}^{-1} \circ f_{0,\infty}$ invariably, so $F_{\text{radmul}}(f_{i,k}^{-1} \circ f_{0,k}) \to F_{\text{radmul}}(f_{i,\infty}^{-1} \circ f_{0,\infty})$ by Lemma [4.1,](#page-31-1) so $F_{\text{stitch}}(f_{1,k}, f_{2,k}) \to$ $F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ $F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$ $F_{\text{stitch}}(f_{1,\infty}, f_{2,\infty})$, so part 1 holds, i.e., F_{stitch} is continuous. For $Q \in SO_3$, we have (Qf_i) $\mathrm{F_{radmul}}((Qf_i)^{-1} \circ (Qf_0)) \,=\, Q(f_i \circ \mathrm{F_{radmul}}(f_i^{-1}))$ $i_i^{-1} \circ f_0$), so just as before, $F_{\text{stitch}}(Q(f_1, f_2)) =$ $Q F_{\text{stitch}}(f_1, f_2)$ $Q F_{\text{stitch}}(f_1, f_2)$ $Q F_{\text{stitch}}(f_1, f_2)$, which means part 2 holds. Similarly, $F_{\text{stitch}}(-(f_1, f_2); z) = - F_{\text{stitch}}(f_1, f_2; \overline{z}),$ which means part [3](#page-12-2) holds, and $F_{\text{stitch}}(f_2, f_1; z) = F_{\text{stitch}}(f_1, f_2; 1/z)$ on $z \in S^1$, which means part [4](#page-12-3) holds on $z \in S^1$. In the case where $|z| < 1$, we have $|1/z| > 1$, so by Lemma [4.1,](#page-31-1) we have

$$
F_{\text{stitch}}(f_1, f_2; 1/z) = [f_2 \circ F_{\text{radmul}}(f_2^{-1} \circ F_{\text{stitch}}(f_1, f_2))](\overline{z})
$$

\n
$$
= [f_2 \circ F_{\text{radmul}}(f_2^{-1} \circ F_{\text{stitch}}(f_2, f_1) \circ [w \mapsto 1/w])](\overline{z})
$$

\n
$$
= [f_2 \circ F_{\text{radmul}}(f_2^{-1} \circ F_{\text{stitch}}(f_2, f_1))](z)
$$

\n
$$
= F_{\text{stitch}}(f_2, f_1; z),
$$

and likewise in the case where $|z| > 1$, so part [4](#page-12-3) holds.

Proof of Theorem [1.3.](#page-1-0) Let ρ be similar to the deformation retraction from Section [3,](#page-14-0) except with F_{stitch} redefined by equation [\(2\)](#page-33-0). By Lemma [4.2,](#page-33-2) the redefined map F_{stitch} still satisfies all relevant properties needed for the proof of Theorem [1.1,](#page-0-1) so the redefined ρ is still a strong $(O_3 \times Z_2)$ -equivariant deformation retraction from hom(S²) to O_3 . It only remains to show that if $f \in \text{hom}_N(S^2)$, then $\rho(f, t) \in \text{hom}_N(S^2)$.

For $f \in \text{hom}_N(\mathbf{S}^2)$, the curve S_E is a null set, so $F_{D4p}(S_E, p_1, p_2, p_{-1}, p_{-2})$ and $F_{D4p}(S_E, p_1, p_{-2}, p_{-1}, p_2)$ preserve nullity, so by Lemma [4.2,](#page-33-2) $F_E \in \text{hom}_N(S^2)$. Also, $F_A \in \text{hom}_N(S^2)$, so ρ at each step is a composition of maps in $hom_N(S^2)$, so $\rho(f, t) \in hom_N(S^2)$. \Box

 \Box

References

- [1] Sigurd Angenent. Parabolic equations for curves on surfaces: Part ii. intersections, blow-up and generalized solutions. Annals of Mathematics, pages 171–215, 1991.
- [2] Eric Kendall Babson. A combinatorial flag space. PhD thesis, Massachusetts Institute of Technology, 1993.
- [3] Daniel K Biss. The homotopy type of the matroid grassmannian. Annals of mathematics, 158(3):929–952, 2003.
- [4] Daniel K Biss. Erratum to "The homotopy type of the matroid Grassmannian". Annals of mathematics, 170(1):493–493, 2009.
- [5] Michael Gene Dobbins. Continuous dependence of curvature flow on initial conditions. arXiv:2106.08907, 2021.
- [6] Michael Gene Dobbins. Grassmannians and pseudosphere arrangements. Journal de l'Ecole polytechnique—Mathématiques, $8:1225-1274$, 2021.
- [7] Jon Folkman and Jim Lawrence. Oriented matroids. Journal of Combinatorial Theory, Series B, 25(2):199–236, 1978.
- [8] Bjorn Friberg. A topological proof of a theorem of Kneser. Proceedings of the American Mathematical Society, 39(2):421–426, 1973.
- [9] Michael E Gage. Curve shortening on surfaces. Annales scientifiques de l'Ecole normale $supérieure, 23(2):229-256, 1990.$
- [10] Mary-Elizabeth Hamstrom. Homotopy properties of the space of homeomorphisms on $P²$ and the Klein bottle. Transactions of the American Mathematical Society, 120(1):37– 45, 1965.
- [11] Hellmuth Kneser. Die Deformationssatze der einfach zusammenhangenden Fl¨aschen. Mathematische Zeitschrift, 25:362–372, 1926.
- [12] Joseph Lauer. The evolution of jordan curves on S^2 by curve shortening flow. $arXiv$ preprint arXiv:1601.05704, 2016.
- [13] Gaku Liu. A counterexample to the extension space conjecture for realizable oriented matroids. Journal of the London Mathematical Society, 101(1):175–193, 2020.
- [14] Nicolai E. Mnëv and Günter M. Ziegler. Combinatorial models for the finite-dimensional Grassmannians. Discrete & Computational Geometry, $10(3):241-250$, 1993.
- [15] William F Osgood. A Jordan curve of positive area. Transactions of the American Mathematical Society, 4(1):107–112, 1903.
- [16] Christian Pommerenke. Boundary Behaviour of Conformal Maps. Springer, 1992.