

Spectral extrema of $K_{s,t}$ -minor free graphs—On a conjecture of M. Tait*

Mingqing Zhai^a, Huiqiu Lin^{b†}

^a School of Mathematics and Finance, Chuzhou University, Chuzhou, Anhui 239012, China

^b Department of Mathematics, East China University of Science and Technology, Shanghai 200237, China

Abstract Minors play an important role in extremal graph theory and spectral extremal graph theory. Tait [The Colin de Verdière parameter, excluded minors, and the spectral radius, J. Combin. Theory Ser. A 166 (2019) 42–58] determined the maximum spectral radius and characterized the unique extremal graph for K_r -minor free graphs of sufficiently large order n , he also made great progress on $K_{s,t}$ -minor free graphs and posed a conjecture: Let $2 \leq s \leq t$ and $n - s + 1 = pt + q$, where n is sufficiently large and $1 \leq q \leq t$. Then $K_{s-1} \nabla (pK_t \cup K_q)$ is the unique extremal graph with the maximum spectral radius over all n -vertex $K_{s,t}$ -minor free graphs. In this paper, Tait’s conjecture is completely solved. We also determine the maximum spectral radius and its extremal graphs for n -vertex $K_{1,t}$ -minor free graphs. To prove our results, some spectral and structural tools, such as, local edge maximality, local degree sequence majorization, double eigenvectors transformation, are used to deduce structural properties of extremal graphs.

Keywords: $K_{s,t}$; minor; spectral radius; extremal graph; majorization

AMS Classification: 05C50; 05C35

1 Introduction

Given a graph H , a graph is said to be H -free if it does not contain H as a subgraph. The classic *Turán’s problem* asks what is the maximum size of an H -free graph of order n , where the maximum size is known as the *Turán number* of H and denoted by $ex(n, H)$. The study of Turán’s problem can be dated back at least to Mantel [22] in 1907, who showed that $ex(n, K_3) \leq \lfloor n^2/4 \rfloor$. Mantel’s theorem was extended by Turán’s theorem in 1941 [33]. Since then, Turán’s problem and many kinds of its variations have been paid much attention and a considerable number of influential results in extremal graph theory have been obtained (see for example, a survey, [14]). In contrast, the spectral extremal problem asks: given a graph H , what is the maximum spectral radius of an H -free graph of order n ? In the past decades much research has been done on spectral extremal graph

*Supported by the National Natural Science Foundation of China (Nos. 11971445, 11771141 and 12011530064).

†Corresponding author. E-mail addresses: mqzhai@chzu.edu.cn (M. Zhai); huiqiulin@126.com (H. Lin).

theory, see K_r [2, 36], $K_{s,t}$ [1, 29], M_k [13], $C_{k,q}$ [21], P_k [27], F_k [6], W_{2k+1} [5], $\bigcup_{i=1}^k S_{a_i}$ [7], $\bigcup_{i=1}^k P_{a_i}$ [8], C_4 [28, 40], C_6 [38], consecutive cycles [15, 19, 23, 25, 39] and a survey [26].

Given two graphs H and G , H is a *minor* of G if H can be obtained from a subgraph of G by contracting edges. A graph is said to be *H -minor free*, if it does not contain H as a minor. Let $A(G)$ be the adjacency matrix of G and $\rho(G)$ be its spectral radius. Recently, Nikiforov [24] and Tait [31, 32] studied the following spectral extremal problem.

Problem 1.1. *Given a graph H or a family \mathbb{H} , what is the maximum spectral radius of an H -minor (\mathbb{H} -minor) free graph of order n ?*

Problem 1.1 was initially paid attention in 1990. Cvetković and Rowlinson [10] conjectured that $\rho(G) \leq \rho(K_1 \nabla P_{n-1})$ for any outerplanar graph G with equality if and only if $G \cong K_1 \nabla P_{n-1}$. Boots and Royle [3] and independently Cao and Vince [4] conjectured that $\rho(G) \leq \rho(K_2 \nabla P_{n-2})$ for any planar graph G of order $n \geq 9$ with equality if and only if $G \cong K_2 \nabla P_{n-2}$. Subsequently, many scholars contributed to these two conjectures (see [4, 16, 17, 30]). Ellingham and Zha [12] showed that $\rho(G) \leq 2 + \sqrt{2n-6}$ for a planar graph G . Dvořák and Mohar [42] proved that $\rho(G) \leq \sqrt{8\Delta-16} + 3.47$ for a planar graph G with maximum degree Δ . In 2017, Tait and Tobin [31] confirmed these two old conjectures for sufficiently large n . Recently, Lin and Ning [20] confirmed Cvetković-Rowlinson conjecture completely. In 2004, Hong [18] proved that $K_3 \nabla (n-3)K_1$ uniquely attains the maximum spectral radius over all K_5 -minor free graphs. Tait [32] extended Hong's result to K_r -minor free graphs by showing the unique extremal graph is $K_{r-2} \nabla (n-r+2)K_1$. In 2017, Nikiforov [24] contributed to $K_{2,t}$ -minor free graphs, and the result was extended to $K_{s,t}$ -minor free graphs by Tait as shown in the following theorem.

Theorem 1.1. [32] *Let $2 \leq s \leq t$, n be large enough and G be an n -vertex $K_{s,t}$ -minor free graph. Then*

$$\rho(G) \leq \frac{1}{2} \left(s+t-3 + \sqrt{(s+t-3)^2 + 4(s-1)(n-s+1) - 4(s-2)(t-1)} \right),$$

with equality if and only if $t \mid n-s+1$ and $G \cong K_{s-1} \nabla \frac{n-s+1}{t} K_t$.

It should be noted that, if $t \nmid n-s+1$ then the maximum spectral radius together with its extremal graph is still unknown for $K_{s,t}$ -minor free graphs. To this end, Tait posed the following conjecture.

Conjecture 1.1. [32] *Let $2 \leq s \leq t$, n be large enough and $n-s+1 = pt+q$, where $1 \leq q \leq t$. Then, the maximum spectral radius of n -vertex $K_{s,t}$ -minor free graphs is attained by the join of K_{s-1} with p copies of K_t and a copy of K_q .*

Up to now, Conjecture 1.1 has been confirmed for $s+t=4$ [28, 40]; $s+t=5$ [24]; $s+t=6$ [35]; and $q=t$ (see Theorem 1.1). For a graph G , let \overline{G} be its complement and $S^k(G)$ be a graph obtained by subdividing k times of an edge uv with the minimum degree sum $d_G(u) + d_G(v)$. Let H^* be the Petersen graph, and $H_{s,t}$ be a star forest of order $t+1$, precisely, the disjoint union of $\lfloor \frac{t+1}{s+1} \rfloor$ stars in which all but at most one are isomorphic to $K_{1,s}$. In this paper, Conjecture 1.1 is solved.

Theorem 1.2. *Let $2 \leq s \leq t$, $n - s + 1 = pt + q$ and $\beta = \lfloor \frac{t+1}{s+1} \rfloor$, where n is large enough and $1 \leq q \leq t$. Let G^* attain the maximum spectral radius over all n -vertex $K_{s,t}$ -minor free graphs. Then*

$$G^* \cong \begin{cases} K_{s-1} \nabla ((p-1)K_t \cup \overline{H^*}) & \text{if } q = 2, t = 8 \text{ and } \beta = 1; \\ K_{s-1} \nabla ((p-1)K_t \cup S^1(\overline{H_{s,t}})) & \text{if } q = \beta = 2; \\ K_{s-1} \nabla ((p-q)K_t \cup q\overline{H_{s,t}}) & \text{if } q \leq 2(\beta-1) \text{ except } q = \beta = 2; \\ K_{s-1} \nabla (pK_t \cup K_q) & \text{otherwise.} \end{cases}$$

It remains $K_{1,t}$ -minor free graphs. Let us first consider connected case. A nice result, due to Ding, Johnson and Seymour [11], determined the maximum size and constructed its extremal graphs for connected $K_{1,t}$ -minor free graphs. However, it seems difficult to completely characterize the extremal graphs. Inspired by Ding, Johnson and Seymour, we obtain the following spectral extremal result.

Theorem 1.3. *Let $t \geq 3$, and G^* attain the maximum spectral radius over all n -vertex connected $K_{1,t}$ -minor free graphs. Then*

$$G^* \cong \begin{cases} \overline{H_{1,t}} & \text{if } n = t + 1; \\ S^{n-t}(K_t) & \text{if } n \geq t + 2. \end{cases}$$

By the connected case in Theorem 1.3, we further solve general case.

Theorem 1.4. *Let $n \geq t \geq 1$, and G be an n -vertex $K_{1,t}$ -minor free graph. Then $\rho(G) \leq t - 1$, with equality if and only if G contains a component either isomorphic to K_t , or isomorphic to K_{t+1} by deleting $\frac{t+1}{2}$ independent edges.*

Combining with above results, the spectral extremal problem on $K_{s,t}$ -minor free graphs is completely solved for large enough n . To prove our results, we use some spectral and structural tools, such as, local edge maximality (see Lemma 2.3), local degree sequence majorization (see Lemma 2.7) and double eigenvectors transformation to deduce structural properties of extremal graphs.

2 Preliminaries

As usual, $V(G)$ is the vertex set and $E(G)$ is the edge set of a graph G . The number of vertices and edges of G are called its *order* and *size*, and denoted by $|G|$ and $e(G)$, respectively. Given $u \in V(G)$ and a subgraph $H \subseteq G$ (possibly $u \notin V(H)$), $N_{V(H)}(u)$ is the set of neighbors of u in $V(H)$ and $d_{V(H)}(u)$ is its cardinality. If $u \in V(H)$, we also use $N_H(u)$ and $d_H(u)$ for convenience. If $S \subseteq V(G)$, then $G[S]$ and $G - S$ stand for the subgraphs of G induced by S and $V(G) \setminus S$, respectively. If $S \subseteq E(G)$, then $G - S$ denotes the subgraph obtained from deleting all edges in S . If $A, B \subseteq V(G)$, then $e_G(A, B)$ denotes the number of edges with one endpoint in A and the other in B , and particularly, $e_G(A, A)$ is simplified by $e_G(A)$.

Throughout this section, let s, t, n be integers with $2 \leq s \leq t$ and n sufficiently large, G^* be the extremal graph with the maximum spectral radius ρ over all n -vertex $K_{s,t}$ -minor free graphs, and $X = (x_1, x_2, \dots, x_n)^T$ be the Perron vector of G^* . Now let us introduce some important lemmas. The first is due to Chudnovsky, Reed and Seymour.

Lemma 2.1. [9, 11] Let $t \geq 3$ and $n \geq t + 2$. If G is an n -vertex connected graph with no $K_{1,t}$ -minor, then $e(G) \leq \binom{t}{2} + n - t$, and this is best possible for all n, t .

For $s = 2$, Nikiforov [24] proved that G^* contains a dominating vertex. Tait showed the following result for general s . This gives a very important information for G^* .

Lemma 2.2. [32] G^* contains a clique dominating set K with $|K| = s - 1$.

By Lemma 2.2, we can observe that G^* is $K_{s,t}$ -minor free if and only if $G^* - K$ satisfies the following property: $K_{a,b}$ -minor free for all positive integers a, b with $a + b = t + 1$ and $1 \leq a \leq \min\{s, \lfloor \frac{t+1}{2} \rfloor\}$. For convenience, we call it (s, t) -property.

Many known results indicate that an extremal graph with the maximum size usually is not an extremal graph with the maximum spectral radius. This observation also happens on $K_{s,t}$ -minor free graphs for general s and t . However, the following lemma implies that G^* has a local edge maximality.

Lemma 2.3. Let H be a disjoint union of several components of $G^* - K$ such that $|H| \leq N$ (a constant). If H' also has (s, t) -property with $V(H') = V(H)$, then $e(H') \leq e(H)$.

Proof. For convenience, let

$$X_0 = \sum_{v \in K} x_v, \quad x_1 = \max_{v \in V(H)} x_v \quad \text{and} \quad x_2 = \min_{v \in V(H)} x_v.$$

Since H has (s, t) -property, then H is $K_{1,t}$ -minor free and hence $\Delta(H) < t$. So, $\rho x_1 < X_0 + t x_1$ and $\rho x_2 \geq X_0$. It follows that

$$x_1 < \frac{X_0}{\rho - t} \quad \text{and} \quad x_2 \geq \frac{X_0}{\rho}. \quad (1)$$

We now give a claim, which will be frequently used in the subsequent proof.

Claim 2.1. Let a, b be two constants with $a > b$. Then $a x_2 > b x_1$ and $a x_2^2 > b x_1^2$.

Proof. By Lemma 2.2, $\Delta(G^*) = n - 1$ and thus $\rho \geq \rho(K_{1,n-1}) = \sqrt{n-1}$. Since n is large enough and a, b, t are constants, we can easily have

$$a x_2 - b x_1 > X_0 \left(\frac{a}{\rho} - \frac{b}{\rho - t} \right) > 0,$$

and similarly, $a x_2^2 > b x_1^2$. □

Let $G' = G^* - E(H) + E(H')$ and $\rho' = \rho(G')$. Then G' is also $K_{s,t}$ -minor free. By the way of contradiction, assume that $e(H') \geq e(H) + 1$. Then by Claim 2.1,

$$\begin{aligned} \rho' - \rho &\geq X^T(A(G') - A(G^*))X = 2 \sum_{uv \in E(H')} x_u x_v - 2 \sum_{uv \in E(H)} x_u x_v \\ &\geq 2e(H')x_2^2 - 2e(H)x_1^2 \\ &> 0, \end{aligned}$$

a contradiction with the maximality of $\rho(G^*)$. So $e(H') \leq e(H)$. □

For a graph H and two vertices $u, v \in V(H)$ (possibly, $uv \notin E(H)$), we use $d_H(uv)$ to denote $d_H(u) + d_H(v)$ for convenience.

Lemma 2.4. *Let H be a disjoint union of several components of $G^* - K$ such that $|H| \leq N$ (a constant), If H' also has (s, t) -property with $V(H') = V(H)$ and $e(H') = e(H)$, then*

$$\sum_{uv \in E(H')} d_H(uv) \leq \sum_{uv \in E(H)} d_H(uv),$$

and if equality holds, then

$$\sum_{uv \in E(H')} d_{H'}(uv) \leq \sum_{uv \in E(H)} d_H(uv).$$

Proof. Let $G' = G^* - E(H) + E(H')$ and $\rho' = \rho(G')$. Then $\rho \geq \rho'$ and

$$\frac{1}{2}\rho^2(\rho' - \rho) \geq \frac{1}{2}\rho^2 X^T (A(G') - A(G^*)) X = \sum_{uv \in E(H')} \rho x_u \rho x_v - \sum_{uv \in E(H)} \rho x_u \rho x_v. \quad (2)$$

Recall that

$$X_0 = \sum_{v \in K} x_v, \quad x_1 = \max_{v \in V(H)} x_v \quad \text{and} \quad x_2 = \min_{v \in V(H)} x_v.$$

Thus, for any $v \in V(H)$, we have

$$X_0 + d_H(v)x_2 \leq \rho x_v \leq X_0 + d_H(v)x_1. \quad (3)$$

It follows that

$$\begin{aligned} \sum_{uv \in E(H')} \rho x_u \rho x_v &\geq \sum_{uv \in E(H')} (X_0 + d_H(u)x_2)(X_0 + d_H(v)x_2), \\ \sum_{uv \in E(H)} \rho x_u \rho x_v &\leq \sum_{uv \in E(H)} (X_0 + d_H(u)x_1)(X_0 + d_H(v)x_1). \end{aligned}$$

Now let

$$a = \sum_{uv \in E(H')} d_H(uv), \quad b = \sum_{uv \in E(H)} d_H(uv), \quad c = \sum_{uv \in E(H)} d_H(u)d_H(v),$$

and suppose to the contrary that $a \geq b + 1$. Then

$$\begin{aligned} \sum_{uv \in E(H')} \rho x_u \rho x_v - \sum_{uv \in E(H)} \rho x_u \rho x_v &\geq aX_0x_2 - bX_0x_1 - cx_1^2 \\ &= X_0 \left(\left(a - \frac{1}{2} \right) x_2 - bx_1 \right) + \left(\frac{1}{2} X_0 x_2 - cx_1^2 \right). \end{aligned}$$

By Claim 2.1, $(a - \frac{1}{2})x_2 > bx_1$. And by (1),

$$\frac{1}{2}X_0x_2 - cx_1^2 > X_0^2 \left(\frac{1}{2\rho} - \frac{c}{(\rho - t)^2} \right) > 0,$$

since $\rho \geq \sqrt{n-1}$ and n is large enough. Combining with (2), we have $\rho' > \rho$, a contradiction. Hence, the first inequality holds.

We now prove the second inequality. Let $a = b$ and $Y = (y_1, y_2, \dots, y_n)^T$ be the Perron vector of G' such that y_i and x_i correspond to the same vertex i . Suppose to the contrary that $a' \geq b + 1$, and assume that

$$Y_0 = \sum_{v \in K} y_v, \quad y_1 = \max_{v \in V(H')} y_v \quad \text{and} \quad y_2 = \min_{v \in V(H')} y_v.$$

Similar with (3), we have

$$Y_0 + d_{H'}(v)y_2 \leq \rho' y_v \leq Y_0 + d_{H'}(v)y_1. \quad (4)$$

for any $v \in V(H')$. It follows that

$$y_1 < \frac{Y_0}{\rho' - t} \quad \text{and} \quad y_2 \geq \frac{Y_0}{\rho'}. \quad (5)$$

Now we can see that

$$\begin{aligned} \rho' \rho (\rho' - \rho) Y^T X &= \rho' \rho ((A(G')Y)^T X - Y^T (A(G)X)) \\ &= \sum_{uv \in E(H')} (\rho x_u \rho' y_v + \rho x_v \rho' y_u) - \sum_{uv \in E(H)} (\rho x_u \rho' y_v + \rho x_v \rho' y_u). \end{aligned}$$

Now let

$$a' = \sum_{uv \in E(H')} d_{H'}(uv), \quad b' = \sum_{uv \in E(H)} d_{H'}(uv), \quad c' = \sum_{uv \in E(H)} (d_H(u)d_{H'}(v) + d_H(v)d_{H'}(u)).$$

Note that $e(H) = e(H')$. By (3) and (4), we have

$$\begin{aligned} &\rho' \rho (\rho' - \rho) Y^T X \\ &\geq \sum_{uv \in E(H')} ((X_0 + d_H(u)x_2)(Y_0 + d_{H'}(v)y_2) + (X_0 + d_H(v)x_2)(Y_0 + d_{H'}(u)y_2)) \\ &\quad - \sum_{uv \in E(H)} ((X_0 + d_H(u)x_1)(Y_0 + d_{H'}(v)y_1) + (X_0 + d_H(v)x_1)(Y_0 + d_{H'}(u)y_1)) \\ &\geq aY_0x_2 + a'X_0y_2 - bY_0x_1 - b'X_0y_1 - c'x_1y_1. \end{aligned}$$

Recall that $d_H(uv) = d_H(u) + d_H(v)$. We can observe that

$$\sum_{uv \in E(H')} d_H(uv) = \sum_{v \in V(H')} d_H(v)d_{H'}(v) = \sum_{v \in V(H)} d_H(v)d_{H'}(v) = \sum_{uv \in E(H)} d_{H'}(uv),$$

that is, $a = b'$. Combining with $a = b$ and $a' \geq a + 1$, we have

$$\begin{aligned} \rho' \rho (\rho' - \rho) Y^T X &\geq aY_0x_2 + (a + 1)X_0y_2 - aY_0x_1 - aX_0y_1 - c'x_1y_1 \\ &= X_0 \left(\left(a + \frac{1}{2} \right) y_2 - ay_1 \right) + \left(\frac{X_0y_2}{2} + aY_0(x_2 - x_1) - c'x_1y_1 \right). \end{aligned}$$

By Claim 2.1, we have $(a + \frac{1}{2})y_2 > ay_1$. And by (1) and (5), we have

$$\frac{X_0y_2}{2} + aY_0(x_2 - x_1) - c'x_1y_1 > X_0Y_0 \left(\frac{1}{2\rho'} + \frac{a}{\rho} - \frac{a}{\rho - t} - \frac{c'}{(\rho - t)(\rho' - t)} \right) > 0,$$

since $\rho \geq \rho' \geq \rho(K_{1,n-1}) = \sqrt{n-1}$ and a, c', t are constants. It follows that $\rho' \rho (\rho' - \rho) Y^T X > 0$, that is, $\rho' > \rho$, a contradiction. This completes the proof. \square

Definition 2.1. Let $X = (x_1, x_2, \dots, x_n)^T$ and $Y = (y_1, y_2, \dots, y_n)^T$ be two decreasing real vectors. If

$$\sum_{i=1}^k x_i \leq \sum_{i=1}^k y_i, \quad k = 1, 2, \dots, n,$$

then we say that X is *weakly majorized* by Y and denote it by $X \prec_w Y$. If $X \prec_w Y$ and $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$, then we say that X is *majorized* by Y and denote it by $x \prec y$.

The following two lemmas are needed in the proof of Lemma 2.7, which is one of our main tools in this paper.

Lemma 2.5. [20] Let $X = (x_1, x_2, \dots, x_n)^T$, $Y = (y_1, y_2, \dots, y_n)^T$ be two nonnegative decreasing real vectors. If $X \prec_w Y$, then $\|X\|_p \leq \|Y\|_p$ for $p > 1$, with equality holding if and only if $X = Y$.

Lemma 2.6 (Exercises 5 (i), P74, [41]). Let $X, Y, Z \in R^n$ be three decreasing vectors. If $X \prec Y$, then $X^T \cdot Z \leq Y^T \cdot Z$.

Let $\pi(G)$ be the decreasing sequence of vertex degrees of a graph G . By using majorization, Bıykođlu and Leydold [34] showed that, for two trees T and T' with $|T| = |T'|$, if $\pi(T') \prec \pi(T)$, then $\rho(T') \leq \rho(T)$. Unfortunately, this nice tool does not work for general graphs, even for unicyclic graphs. However, the following lemma implies that G^* has a local degree sequence majorization.

Lemma 2.7. Let H and H' be defined as in Lemma 2.4. Let $\pi(H) = (d_1, \dots, d_{|H|})$ and $\pi(H') = (d'_1, \dots, d'_{|H|})$. If $\pi(H) \prec \pi(H')$, then $\pi(H) = \pi(H')$.

Proof. Let $V(H) = \{v_1, v_2, \dots, v_{|H|}\}$ with $d_H(v_i) = d_i$. Since $V(H) = V(H')$, by graph isomorphism, we may let $d_{H'}(v_i) = d'_i$ for $i = 1, 2, \dots, |H|$. Suppose to the contrary that $\pi(H) \neq \pi(H')$. Then by Lemma 2.5, we have

$$\sum_{i=1}^{|H|} d_i^2 > \sum_{i=1}^{|H|} d_i'^2. \quad (6)$$

Now let $X = Z = \pi(H)$ and $Y = \pi(H')$ in Lemma 2.6. Then we have

$$\sum_{i=1}^{|H|} d_i^2 \leq \sum_{i=1}^{|H|} d_i d'_i. \quad (7)$$

Furthermore, we see that

$$\sum_{i=1}^{|H|} d_i d'_i = \sum_{v \in V(H')} d_H(v) d_{H'}(v) = \sum_{uv \in E(H')} (d_H(u) + d_H(v)) = \sum_{uv \in E(H')} d_H(uv).$$

And similarly,

$$\sum_{i=1}^{|H|} d_i^2 = \sum_{uv \in E(H)} (d_H(u) + d_H(v)) = \sum_{uv \in E(H)} d_H(uv).$$

It follows from the first inequality of Lemma 2.4 that $\sum_{i=1}^{|H|} d_i d'_i \leq \sum_{i=1}^{|H|} d_i^2$. Combining with (7), we have $\sum_{i=1}^{|H|} d_i d'_i = \sum_{i=1}^{|H|} d_i^2$. By the second inequality of Lemma 2.4, we have

$$\sum_{uv \in E(H')} d_{H'}(uv) \leq \sum_{uv \in E(H)} d_H(uv),$$

that is, $\sum_{i=1}^{|H|} d_i'^2 \leq \sum_{i=1}^{|H|} d_i^2$, which contradicts (6). So, $\pi(H) = \pi(H')$. \square

3 Characterization of Components in $G^* - K$

Throughout this section, assume that $t \geq 4$ and $2 \leq s \leq t$. Let $G^*, K, H_{s,t}$ and related notations be defined as above. In particular, recall that each component H of $G^* - K$ satisfies (s, t) -property, that is, H is $K_{a,b}$ -minor free for all positive integers a, b with $a + b = t + 1$ and $1 \leq a \leq \gamma$, where $\gamma = \min\{s, \lfloor \frac{t+1}{2} \rfloor\}$.

Lemma 3.1. *Let G be a connected graph with $|G| = t + 1$. Then G has (s, t) -property if and only if each component of \overline{G} has at least $\gamma + 1$ vertices.*

Proof. Since $|G| = t + 1$, a $K_{a,b}$ -minor is equivalent to a copy of $K_{a,b}$ for any positive integers a, b with $a + b = t + 1$ and $1 \leq a \leq \gamma$. It is easy to see that G has (s, t) -property if and only if each component of \overline{G} has at least $\gamma + 1$ vertices. \square

Lemma 3.2. *Let $\beta = \lfloor \frac{t+1}{s+1} \rfloor$ and G be a connected graph with $|G| = t + 1$. If G has (s, t) -property, then $e(G) \leq \binom{t}{2} + \beta - 1$.*

Proof. By Lemma 3.1, each component of \overline{G} has at least $\gamma + 1$ vertices. Now assume that G is edge-maximal and \overline{G} has c components. Then each component of \overline{G} is a tree and c is maximal. This implies that $c = \lfloor \frac{t+1}{\gamma+1} \rfloor$ and $e(\overline{G}) = |G| - c$. If $s \leq \lfloor \frac{t+1}{2} \rfloor$, then $\gamma = s$ and thus $c = \beta$. It follows that

$$e(G) = \binom{t+1}{2} - e(\overline{G}) = \binom{t}{2} + c - 1 = \binom{t}{2} + \beta - 1. \quad (8)$$

If $s > \lfloor \frac{t+1}{2} \rfloor$, then $\gamma = \lfloor \frac{t+1}{2} \rfloor$ and $\beta = 1$. We also have $c = \beta$ and hence (8) holds. \square

The following lemma holds clearly.

Lemma 3.3. *Let G be a graph with $vw \in E(G)$ and $uw \notin E(G)$. If $d_G(u) \geq d_G(v)$, then $\pi(G) \prec \pi(G - \{vw\} + \{uw\})$ and $\pi(G) \neq \pi(G - \{vw\} + \{uw\})$.*

Recall that $\beta = \lfloor \frac{t+1}{s+1} \rfloor$ and $H_{s,t} = (\beta - 1)K_{1,s} \cup K_{1,\alpha}$, where $\alpha = t - (\beta - 1)(s + 1) \geq s$. The following theorem presents a clear characterization of a $(t + 1)$ -vertex component of $G^* - K$, which is a key subgraph of the extremal graphs.

Theorem 3.1. *Let H be a component of $G^* - K$. If $|H| = t + 1$, then $\beta \geq 2$, $H \cong \overline{H_{s,t}}$ and $e(H) = \binom{t}{2} + \beta - 1$.*

Proof. We first show $\beta \geq 2$. Suppose that $\beta = 1$. Then by Lemma 3.2, $e(H) \leq \binom{t}{2} = e(K_t \cup K_1)$. By Lemma 2.3, $e(H) = e(K_t \cup K_1)$, since $K_t \cup K_1$ also has (s, t) -property. Furthermore, since $\Delta(H) \leq t - 1$ and $\delta(H) \geq 1$, we can see that $\pi(H) \prec \pi(K_t \cup K_1)$ and $\pi(H) \neq \pi(K_t \cup K_1)$, which contradicts Lemma 2.7. Thus, $\beta \geq 2$ and correspondingly $s \leq \lfloor \frac{t-1}{2} \rfloor$. It follows that $\gamma = s$.

Now we characterize the structure of H . By Lemma 2.3, H is edge-maximal, and combining with Lemmas 3.1 and 3.2, we have $e(H) = \binom{t}{2} + \beta - 1$ and \overline{H} is a forest with β components, each of which has at least $\gamma + 1 (= s + 1)$ vertices. We divide the proof into two claims.

Claim 3.1. *Each component of \overline{H} is a star.*

Proof. Suppose that \overline{H} contains a component T which is not a star. Then, T contains at least two pendant edges u_1v_1 and u_2v_2 , where v_1, v_2 are leaves and $u_1 \neq u_2$. Assume without loss of generality that $d_T(u_1) \leq d_T(u_2)$. Then $d_H(u_1) \geq d_H(u_2)$. By Lemma 3.3, $\pi(H) \prec \pi(H - \{u_2v_1\} + \{u_1v_1\})$, which contradicts Lemma 2.7. \square

Claim 3.2. $\overline{H} \cong H_{s,t}$.

Proof. It suffices to show that \overline{H} has at most one component not isomorphic to $K_{1,s}$. Suppose that there exist two components T_1 and T_2 with $|T_2| \geq |T_1| \geq s + 2$. Assume that $u_i v_i \in E(T_i)$ and v_i is a leaf for $i \in \{1, 2\}$. Similar to the proof of Claim 3.1, we can see that $\pi(H) \prec \pi(H - \{u_2v_1\} + \{u_1v_1\})$, a contradiction. \square

Having Claim 3.2, it remains to show that $\overline{H_{s,t}}$ has (s, t) -property. Since $\gamma = s$ and $H_{s,t} \cong K_{1,\alpha} \cup (\beta - 1)K_{1,s}$, each component of $H_{s,t}$ has at least $\gamma + 1$ vertices. By Lemma 3.1, $\overline{H_{s,t}}$ has (s, t) -property. \square




Figure 1: The graph $H_{a,b,c}$ and its complement.

Lemma 3.4. *Let $\beta \leq 2$ and G be a connected graph with $|G| = t + 2$. If G has (s, t) -property, then $e(G) \leq \binom{t}{2} + 2$, and if equality holds, then \overline{G} is isomorphic to either some $H_{a,b,c}$ (see Fig. 1) or the Petersen graph H^* .*

Proof. Since G satisfies (s, t) -property, G contains no $K_{1,t}$ -minor. By Lemma 2.1, we have $e(G) \leq \binom{t}{2} + 2$. In the following, assume that $e(G) = \binom{t}{2} + 2$. Then $e(\overline{G}) = 2t - 1$. The proof is divided into five claims.

Claim 3.3. \overline{G} is connected and $\text{diam}(\overline{G}) = 2$, where $\text{diam}(\overline{G})$ is the diameter of \overline{G} .

Proof. If \overline{G} is not connected or $\text{diam}(\overline{G}) \geq 3$, then there exist a pair of non-adjacent vertices u, v with no common neighbor in \overline{G} . This implies that $uv \in E(G)$ and $N_G(u) \cup N_G(v) = V(G)$. Since $|G| = t + 2$, we can conclude that G contains a $K_{1,t}$ -minor, a contradiction with (s, t) -property. \square

Claim 3.4. Let $S = \{u_1, \dots, u_{|S|}\}$ be a minimum cut set of \overline{G} . Then $2 \leq |S| \leq 3$.

Proof. Let $d_G(u, v)$ be the distance of $u, v \in V(G)$. If $|S| = 1$, then $\overline{G} - \{u_1\}$ has at least two components G_1 and G_2 . Since G is connected, there exists a vertex v_1 , say $v_1 \in V(G_1)$, with $u_1 v_1 \notin E(\overline{G})$. Now, $d_{\overline{G}}(v_1, v) \geq 3$ for any $v \in V(G_2)$, contradicting Claim 3.3. Therefore, $|S| \geq 2$. Furthermore, since $\delta(\overline{G}) \geq |S|$. Thus, $2e(\overline{G}) \geq \delta(\overline{G}) \cdot |G| \geq |S|(t+2)$. Combining with $e(\overline{G}) = 2t - 1$, we have $|S| \leq 3$. \square

Now let $W = V(G) \setminus S$, $W_1 = \{w \in W \mid |N_{\overline{G}}(w) \cap S| = 1\}$ and $\overline{G}[W]$ consists of k components T_1, \dots, T_k , where $k \geq 2$.

Claim 3.5. S dominates W and each $V(T_i)$ dominates S in \overline{G} . Moreover, W_1 belongs to a single component T_i .

Proof. Firstly, suppose that $N_{\overline{G}}(w_0) \cap S = \emptyset$ for some $w_0 \in W$. Without loss of generality, assume that $w_0 \in V(T_1)$. Then $d_{\overline{G}}(w_0, w) \geq 3$ for any $w \in V(T_2)$, a contradiction. Hence, S dominates W in \overline{G} . Secondly, if a $V(T_i)$ does not dominate S in \overline{G} , say $u_1 \notin \cup_{w \in V(T_i)} N_{\overline{G}}(w)$, then $S \setminus \{u_1\}$ is a cut set, which contradicts the minimality of $|S|$.

Now we show that $W_1 \subseteq V(T_i)$ for some i . Suppose to the contrary that $w_i \in W_1 \cap V(T_i)$ for $i = 1, 2$. Since $\text{diam}(\overline{G}) = 2$, w_1 and w_2 have a unique and common neighbor, say u_1 , in S . It follows that $wu_1 \in E(\overline{G})$ for any $w \in W$ (Otherwise, either $d_{\overline{G}}(w, w_1) \geq 3$ or $d_{\overline{G}}(w, w_2) \geq 3$). Therefore, $d_{\overline{G}}(u_1) \geq |W|$. If $|S| = 2$, then

$$e(\overline{G} - \{u_1\}) \leq e(\overline{G}) - |W| = (2t - 1) - t = t - 1.$$

This implies that $\overline{G} - \{u_1\}$ is not connected, which contradicts $|S| = 2$. If $|S| = 3$, then $d_{\overline{G}}(u_2) \geq \delta(\overline{G}) \geq |S| = 3$. It follows that

$$e(\overline{G} - \{u_1, u_2\}) \leq e(\overline{G}) - (|W| + 2) = (2t - 1) - (t + 1) = t - 2.$$

Then $\{u_1, u_2\}$ is a cut set of \overline{G} , contradicting $|S| = 3$. So, the claim holds. \square

Now we may assume without loss of generality that $W_1 \subseteq V(T_1)$. Note that

$$e(\overline{G}) = e_{\overline{G}}(S) + \sum_{i=1}^k e_{\overline{G}}(V(T_i)) + \sum_{i=1}^k e_{\overline{G}}(V(T_i), S). \quad (9)$$

Claim 3.6. If $|S| = 2$, then \overline{G} is isomorphic to some $H_{a,b,c}$ (see Fig. 1).




Figure 2: The graph $H'_{a,b,c}$ and its complement.

Proof. Now, $|W| = t$. Since $W_1 \subseteq V(T_1)$, by (9) we have

$$e(\overline{G}) \geq 2|T_1| - 1 + 2(|W| - |T_1|) = 2t - 1.$$

Recall that $e(\overline{G}) = 2t - 1$. We can see that $e_{\overline{G}}(S) = 0$, T_1 is a tree with $V(T_1) = W_1$, and $W \setminus W_1$ is an independent set of \overline{G} .

We shall further show that T_1 is a star. Since $\text{diam}(\overline{G}) = 2$ and $V(T_1) = W_1$, we can observe that w_1 and w_2 have a unique and common neighbor in S for any $w_1, w_2 \in V(T_1)$ with $d_{T_1}(w_1, w_2) \geq 3$. Thus, if $\text{diam}(T_1) \geq 5$, then $V(T_1)$ dominates exactly one vertex of S in \overline{G} , contradicting Claim 3.5. If $\text{diam}(T_1) = 4$, then all vertices in $V(T_1)$ but the central vertex are adjacent to a vertex (say u_2) of S . But now, we can find a leaf w with $d_{\overline{G}}(w, u_1) = 3$, a contradiction. If $\text{diam}(T_1) = 3$, that is, T_1 is a double star, then we can similarly see that all leaves have a common neighbor (say u_2) in S , and two central vertices w_1 and w_2 are adjacent to u_1 . It follows that $\overline{G} \cong H'_{a,b,c}$ for some a, b, c with $a + b + c = t - 2$ (see Fig. 2). If $c \leq \gamma - 1$, we contract the edge $u_1 u_2$ in G and call the new vertex u in the resulting graph, then we get a complete bipartite subgraph with bipartite partition $(V(K_c) \cup \{u\}, V(K_a) \cup V(K_b) \cup \{w_1, w_2\})$. This implies that G contains a $K_{c+1, a+b+2}$ -minor, a contradiction with (s, t) -property. So, $c \geq \gamma$. By symmetry, we also have $a, b \geq \gamma$. Therefore, $t - 2 = a + b + c \geq 3\gamma$. This implies that $\gamma \leq \lfloor \frac{t-2}{3} \rfloor$ and thus $s \leq \lfloor \frac{t-2}{3} \rfloor$, which contradicts $\beta = \lfloor \frac{t+1}{s+1} \rfloor \leq 2$.

Now we conclude that T_1 is a star. Let w be the central vertex of T_1 . Without loss of generality, assume that $wu_1 \in E(\overline{G})$. Then $\overline{G} \cong H_{a,b,c}$ for some a, b, c with $a + b + c = t - 1$ (see Fig. 1), as desired. \square

Claim 3.7. *If $|S| = 3$, then \overline{G} is isomorphic to the Petersen graph H^* (see Fig. 3).*

Proof. Now $\delta(\overline{G}) \geq |S| = 3$. So, all leaves of T_1 , if exist, belong to $W \setminus W_1$. Hence,

$$e_{\overline{G}}(V(T_1)) + e_{\overline{G}}(V(T_1), S) \geq 2|T_1|, \quad (10)$$

and if equality holds, then T_1 is a cycle with $V(T_1) = W_1$.




Figure 3: The Petersen graph H^* and its complement.

On the other hand, by Claim 3.5, $V(T_1)$ dominates S in \overline{G} , which implies that each vertex of $W \setminus V(T_1)$ also dominates S in \overline{G} (Otherwise, we can find $w_1 \in V(T_1)$ and $w_2 \in W \setminus V(T_1)$ with $d_{\overline{G}}(w_1, w_2) \geq 3$). Therefore, $e_{\overline{G}}(W \setminus V(T_1), S) = 3(|W| - |T_1|)$. Combining with (9) and (10), we have

$$e(\overline{G}) \geq e_{\overline{G}}(S) + 2|T_1| + 3(|W| - |T_1|) \geq 3|W| - |T_1| \geq 2|W| + 1 = 2t - 1.$$

Since $e(\overline{G}) = 2t - 1$, we have $e_{\overline{G}}(S) = 0$, $T_1 \cong C_{|W|}$ and $|W| = |T_1| + 1$.

Note that $\delta(\overline{G}) \geq 3$. Each vertex of S has at least two neighbors in $V(T_1)$. It follows that $|T_1| \geq 6$. Furthermore, any two vertices with distance at least 3 in T_1 have a common neighbor in S . If $|T_1| \geq 7$, then all vertices in T_1 have a common neighbor in S , contradicting Claim 3.5. So, $T_1 \cong C_6$ and every pair of vertices with distance 3 in T_1 have a common neighbor in S . Thus, $\overline{G} \cong H^*$ (see Fig. 3). \square

Combining with Claim 3.6 and Claim 3.7, the proof of Lemma 3.4 is completed. \square

We now use \mathbb{H}_i , $\mathbb{H}_{>i}$ and $\mathbb{H}_{<i}$ to denote the family of components in $G^* - K$ with order i , greater than i and less than i , respectively.

Lemma 3.5. $\mathbb{H}_{>t+3} = \emptyset$.

Proof. Suppose to the contrary that $H \in \mathbb{H}_{>t+3}$. By Lemma 2.1,

$$e(H) \leq \binom{t}{2} + |H| - t. \quad (11)$$

Assume that $|H| = pt + q$, where $p \geq 1$ and $1 \leq q \leq t$, and let $H' \cong pK_t \cup K_q$ with $V(H') = V(H)$. Clearly, H' satisfies (s, t) -property. By (11), we can see that

$$e(H) \leq \binom{t}{2} + (pt + q) - t < p \binom{t}{2} + \binom{q}{2} = e(H')$$

for $|H| > t + 3$ and $t \geq 4$. If $p \leq 7$, then $|H| \leq 8t$ (a constant). By Lemma 2.3, we have $e(H') \leq e(H)$, a contradiction. Now assume that $p \geq 8$. Then by (11),

$$e(H) \leq \binom{t}{2} + pt < \frac{4p}{5} \binom{t}{2} \leq \frac{4}{5} e(H'). \quad (12)$$

Now let $G' = G^* - E(H) + E(H')$ and $\rho' = \rho(G')$. Then

$$\rho' - \rho \geq X^T(A(G') - A(G^*))X = \sum_{uv \in E(H')} 2x_u x_v - \sum_{uv \in E(H)} 2x_u x_v.$$

Recall that $x_1 = \max_{v \in V(H)} x_v$ and $x_2 = \min_{v \in V(H)} x_v$. By (12) and Claim 2.1,

$$\rho' - \rho \geq 2e(H')x_2^2 - 2e(H)x_1^2 > 2e(H')(x_2^2 - \frac{4}{5}x_1^2) > 0,$$

a contradiction. Thus we have $\mathbb{H}_{>t+3} = \emptyset$. \square

Lemma 3.6. $\mathbb{H}_t = O(\frac{n}{t})$.

Proof. By Lemma 3.5, $\mathbb{H}_{>t+3} = \emptyset$. Hence, it suffices to show $|\mathbb{H}_i| < t$ for any $i \leq t+3$ and $i \neq t$. Suppose to the contrary that $|\mathbb{H}_i| \geq t$ for some i and let \mathbb{F} be the disjoint union of any t components in \mathbb{H}_i . If $i < t$, then $e(\mathbb{F}) \leq e(tK_i) < e(iK_t)$, which contradicts Lemma 2.3.

If $i = t+1$, then by Lemma 3.2 and $\beta = \lfloor \frac{t+1}{s+1} \rfloor \leq \lfloor \frac{t+1}{3} \rfloor$, we have

$$e(\mathbb{F}) \leq t \left(\binom{t}{2} + \beta - 1 \right) < (t+1) \binom{t}{2} = e((t+1)K_t),$$

which contradicts Lemma 2.3.

If $i \in \{t+2, t+3\}$, then by Lemma 2.1,

$$e(\mathbb{F}) \leq t \left(\binom{t}{2} + i - t \right) < i \binom{t}{2} = e(iK_t),$$

also a contradiction. \square

Now we are ready to characterize a $(t+2)$ -vertex component of $G^* - K$, which is also a key subgraph of the extremal graphs.




Figure 4: The graph $S^1(\overline{H_{s,t}})$ for $\beta = \lfloor \frac{t+1}{s+1} \rfloor = 2$.

Theorem 3.2. Let H be a component of $G^* - K$. If $|H| = t+2$, then $\beta \leq 2$. Moreover, $H \cong S^1(\overline{H_{s,t}})$ for $\beta = 2$ (see Fig. 4), and $H \cong \overline{H^*}$ for $\beta = 1$.

Proof. The proof is divided into several claims.

Claim 3.8. $\beta \leq 2$.

Proof. If $\beta \geq 3$, then by Theorem 3.1, $e(\overline{H_{s,t}}) = \binom{t}{2} + \beta - 1 \geq \binom{t}{2} + 2$. By Lemma 3.6, $G^* - K$ contains a component H_0 isomorphic to K_t . Let $H' \cong \overline{H_{s,t}} \cup \overline{H_{s,t}}$ and $V(H') = V(H \cup H_0)$. By Theorem 3.1, H' has (s, t) -property. However, $e(H \cup H_0) = 2\binom{t}{2} + 2 < e(H')$, contradicting Lemma 2.3. So, $\beta \leq 2$. \square

Claim 3.9. $e(H) = \binom{t}{2} + 2$.

Proof. By Lemma 2.1, we have $e(H) \leq \binom{t}{2} + 2$. Since $K_t \cup K_2$ satisfies (s, t) -property, by Lemma 2.3, we have $e(H) \geq e(K_t \cup K_2) = \binom{t}{2} + 1$. Now suppose that $e(H) = e(K_t \cup K_2)$. Since $\Delta(H) \leq t - 1$ and $\delta(H) \geq 1$, we can see that $\pi(H) \prec \pi(K_t \cup K_2)$. By Lemma 2.7, we have

$$\pi(H) = \pi(K_t \cup K_2) = (t - 1, t - 1, \dots, t - 1, 1, 1),$$

which implies that H is obtained from K_t by deleting an edge u_1u_2 and adding two pendant edges u_1v_1 and u_2v_2 . Since $t \geq 4$, we have $N_H(u_1) \setminus \{v_1\} \neq \emptyset$ and thus

$$\rho(x_{u_1} - x_{v_1}) = (x_{v_1} - x_{u_1}) + \sum_{v \in N_H(u_1) \setminus \{v_1\}} x_v > x_{v_1} - x_{u_1},$$

that is, $x_{u_1} > x_{v_1}$. By symmetry, we have $x_{u_1} = x_{u_2}$ and $x_{v_1} = x_{v_2}$. Now, let $H' = H - \{u_1v_1, u_2v_2\} + \{u_1u_2, v_1v_2\}$ and $G' = G^* - E(H) + E(H')$. Then $H' \cong K_t \cup K_2$ and thus H' has (s, t) -property.

$$\rho(G') - \rho(G^*) \geq X^T(A(G') - A(G^*))X = 2(x_{u_1} - x_{v_2})(x_{u_2} - x_{v_1}) > 0,$$

a contradiction. Hence, $e(H) = \binom{t}{2} + 2$. \square

Now, combining Claims 3.8 and 3.9 with Lemma 3.4, H is isomorphic to either some $\overline{H_{a,b,c}}$ (see Fig. 1) or the complement of the Peterson graph H^* .

Claim 3.10. If $\beta = 2$, then $H \cong S^1(\overline{H_{s,t}})$ (see Fig. 4).

Proof. Since $\beta = \lfloor \frac{t+1}{s+1} \rfloor$, we have

$$2s + 1 \leq t \leq 3s + 1 \quad \text{and} \quad \gamma = \min\{s, \lfloor \frac{t+1}{2} \rfloor\} = s. \quad (13)$$

If $H \cong \overline{H^*}$, then $|H| = 10$, $t = 8$ and H is 6-regular. It follows that $e(H) = 30$. On the other hand, since $S^1(\overline{H_{s,8}})$ is a subdivision of $\overline{H_{s,8}}$, by Theorem 3.1 we have $e(S^1(\overline{H_{s,8}})) = e(\overline{H_{s,8}}) + 1 = \binom{t}{2} + \beta = 30$. By Fig. 4 we can see that

$$\pi(S^1(\overline{H_{s,t}})) = (t - 1, \dots, t - 1, t - s, s + 1, 2), \quad (14)$$

where $s + 1 \leq t - s \leq 6$. Since H is 6-regular, we have $\pi(H) \prec \pi(S^1(\overline{H_{s,8}}))$, contradicting Lemma 2.7. Thus, H is isomorphic to some $\overline{H_{a,b,c}}$ with $a + b + c = t - 1$.

Next we show that

$$\min\{b, c\} \geq \gamma. \quad (15)$$

If $b \leq \gamma - 1$, we contract the edge u_2w in $\overline{H_{a,b,c}}$ and call the new vertex u in the resulting graph, then we get a complete bipartite subgraph with bipartite partition $\langle V(K_b) \cup \{u\},$

$V(K_a) \cup V(K_c) \cup \{u_1\}$. This implies that H contains a $K_{b+1, a+c+1}$ -minor, a contradiction with (s, t) -property. So, $b \geq \gamma$. And by symmetry, $c \geq \gamma$.

Now by Fig. 1 we can see that

$$\pi(\overline{H_{a,b,c}}) = (t-1, \dots, t-1, a_1, a_2, a_3), \quad (16)$$

where $a_1, a_2, a_3 \in \{a+2, b+1, c+1\}$. By (13) and (15), $\min\{b, c\} \geq \gamma = s$. It follows that $a_3 \geq 2$ and $a_2 \geq s+1$. Comparing (14) with (16), we have $\pi(\overline{H_{a,b,c}}) \prec \pi(S^1(\overline{H_{s,t}}))$. By Lemma 2.7, $\pi(\overline{H_{a,b,c}}) = \pi(S^1(\overline{H_{s,t}}))$. So, $a_3 = 2$ and $a_2 = s+1$. Note that $\min\{b, c\} \geq s \geq 2$. We conclude that $a = 0$ and $\min\{b, c\} = s$, that is, $\overline{H_{a,b,c}} \cong S^1(\overline{H_{s,t}})$.

Finally, we shall prove that $S^1(\overline{H_{s,t}})$ has (s, t) -property. It suffices to show that contracting any edge, the resulting graph always has (s, t) -property. Indeed, if we contract an edge within the $(t-1)$ -clique, then the complement of the resulting graph is clearly connected. By Lemma 3.1, it has (s, t) -property. If we contract an edge out of the $(t-1)$ -clique, then the resulting graph is isomorphic to $\overline{H_{s,t}}$. By Theorem 3.1, it also has (s, t) -property. \square

Claim 3.11. *If $\beta = 1$, then $H \cong \overline{H^*}$ (see Fig. 3).*

Proof. Since $\beta = \lfloor \frac{t+1}{s+1} \rfloor = 1$, we have $t \leq 2s$ and thus $\gamma = \min\{s, \lfloor \frac{t+1}{2} \rfloor\} = \lfloor \frac{t+1}{2} \rfloor$. If H is isomorphic to some $\overline{H_{a,b,c}}$ with $a+b+c = t-1$, then by (15), we have $t-1 \geq b+c \geq 2\gamma = 2\lfloor \frac{t+1}{2} \rfloor$, a contradiction. By Lemma 3.4, H is only possibly isomorphic to $\overline{H^*}$.

It remains to show that $\overline{H^*}$ has (s, t) -property. We know that the Peterson graph H^* is 3-connected and any two non-adjacent vertices of H^* have exactly one common neighbor, which implies that contracting any edge of $\overline{H^*}$, the complement of the resulting graph is connected. By Lemma 3.1, the resulting graph has (s, t) -property, and thus $\overline{H^*}$ has (s, t) -property. \square

Combining with Claims 3.8, 3.10 and 3.11, the proof of Theorem 3.2 is completed. \square

From Lemma 3.5, we know that $\mathbb{H}_{>t+3} = \emptyset$. Now combining with Lemma 3.4, we can get a stronger result.

Theorem 3.3. $\mathbb{H}_{>t+2} = \emptyset$.

Proof. It suffices to show that $\mathbb{H}_{t+3} = \emptyset$. Suppose to the contrary that there exists a component H of $G^* - K$ with $|H| = t+3$. On one hand, $e(H) \leq \binom{t}{2} + 3$ by Lemma 2.1. On the other hand, note that $K_t \cup K_3$ also satisfies (s, t) -property. Then by Lemma 2.3, $e(H) \geq e(K_t \cup K_3) = \binom{t}{2} + 3$. Therefore, $e(H) = \binom{t}{2} + 3$. Moreover, if $\beta \geq 3$, then by Theorem 3.1, $|\overline{H_{s,t}}| = t+1$, $e(\overline{H_{s,t}}) = \binom{t}{2} + (\beta-1)$ and $\overline{H_{s,t}}$ has (s, t) -property. Selecting two copies of K_t in $G^* - K$, we have

$$e(H \cup K_t \cup K_t) = 3\binom{t}{2} + 3 < 3\binom{t}{2} + 3(\beta-1) = e(\overline{H_{s,t}} \cup \overline{H_{s,t}} \cup \overline{H_{s,t}}),$$

a contradiction with Lemma 2.3. Therefore, $\beta \leq 2$.

Claim 3.12. $\delta(H) \geq 2$.

Proof. If $\delta(H) = 1$, say $d_H(v) = 1$, then $H - \{v\}$ is connected and $e(H - \{v\}) = \binom{t}{2} + 2$. By Lemma 3.4, $H - \{v\}$ is isomorphic to either $\overline{H^*}$ or some $\overline{H_{a,b,c}}$ with $a + b + c = t - 1$. Observe that $\overline{H_{a,b,c}}$ contains a K_t -minor (by contracting edges u_1u_2 and u_2w , see Fig. 1). Hence, $H - \{v\} \cong \overline{H^*}$ (Otherwise, H contains a $K_{1,t}$ -minor).

Now let vuv be a path of length 2 in H . We know that any two non-adjacent vertices in the Petersen graph H^* have exactly one common neighbor. So, contracting uv in $H - \{v\}$, the new vertex is of degree $t - 1$ in the resulting graph. Correspondingly, contracting uv in H , the new vertex is of degree t , which contradicts that H is $K_{1,t}$ -minor free. \square

Now we have $\pi(H) \prec \pi(K_t \cup K_3) = (t - 1, \dots, t - 1, 2, 2, 2)$, since $e(H) = \binom{t}{2} + 3$ and $\delta(H) \geq 2$. By Lemma 2.7, $\pi(H) = \pi(K_t \cup K_3)$. Let $S_1 = \{v \in V(H) \mid d_H(v) = 2\}$ and $S_2 = \cup_{v \in S_1} N_H(v) \setminus S_1$. Then $|S_1| = 3$, and $1 \leq d_{S_1}(u) \leq 3$ for any $u \in S_2$.

Claim 3.13. $d_{S_1}(u) = 1$ for any $u \in S_2$.

Proof. Let R be the set of non-adjacent vertex-pairs in S_2 . Since $|V(H) \setminus S_1| = t$ and $d_H(u) = t - 1$ for any $u \in V(H) \setminus S_1$, we have $|R| = \frac{1}{2}e_H(S_1, S_2) \leq 3$. Suppose that $d_{S_1}(u_0) = c \in \{2, 3\}$ for some $u_0 \in S_2$. Then, u_0 has exactly c non-neighbors, say $\{u_1, \dots, u_c\}$, in S_2 . If $c = 3$, then $R = \{(u_0, u_i) \mid i = 1, 2, 3\}$ and there are three paths $u_0v_iu_i$ in H , where $v_i \in S_1$ and $i \in \{1, 2, 3\}$. But now we find a $K_{2,t-1}$ -minor in H by contracting two of the three paths into edges. Therefore, $c = 2$.

Since $t \geq 4$, we can find a vertex $u_3 \in N_H(u_0) \setminus S_1$. If both u_1 and u_2 are neighbors of u_3 , then H contains a double star with a non-pendant edge u_0u_3 and t leaves, and thus a $K_{1,t}$ -minor. If both u_1 and u_2 are not neighbors of u_3 , then $|R| \geq 4$, a contradiction. Hence, we may assume that $u_1u_3 \in E(H)$ and $u_2u_3 \notin E(H)$. Then, $u_1u_2 \in E(H)$ (Otherwise, we get $|R| \geq 4$ again). It follows that $P = u_0u_3u_1u_2$ is an induced path in H and $S_2 = \{u_0, u_1, u_2, u_3\}$. Furthermore, $d_{S_1}(u_0) = d_{S_1}(u_2) = 2$ and $d_{S_1}(u_1) = d_{S_1}(u_3) = 1$. Now, we can always find a double star with a non-pendant edge in $E(P)$ and t leaves, a contradiction. \square

Claim 3.14. $x_u > x_v$ for any $u \in S_2$ and $v \in S_1$.

Proof. By (3), $\rho x_u \geq X_0 + d_H(u)x_2$ and $\rho x_v \leq X_0 + d_H(v)x_1$. Combining with Claim 2.1, we have $\rho(x_u - x_v) \geq d_H(u)x_2 - d_H(v)x_1 > 0$. \square

Now by Claim 3.13, each $u_i \in S_2$ has a unique neighbor v_i in S_1 , and thus a unique non-neighbor u_j in S_2 . If $v_i = v_j$ or $v_iv_j \in E(H)$ for some $(u_i, u_j) \in R$, then u_i and u_j have $t - 1$ common neighbors after contracting the path $u_iv_iv_ju_j$ into $u_iv_iu_j$ in H . This implies that H contains a $K_{2,t-1}$ -minor. Thus, $v_i \neq v_j$ and $v_iv_j \notin E(H)$ for any $(u_i, u_j) \in R$. Let

$$H' = H - \{u_iv_i, u_jv_j \mid (u_i, u_j) \in R\} + \{u_iu_j, v_iv_j \mid (u_i, u_j) \in R\}$$

and $G' = G^* - E(H) + E(H')$. Note that $H' \cong K_t \cup K_3$. Thus, H' has (s, t) -property. Moreover, by Claim 3.14, we have

$$\rho(G') - \rho(G^*) \geq 2 \sum_{(u_i, u_j) \in R} (x_{u_i}x_{u_j} + x_{v_i}x_{v_j} - x_{u_i}x_{v_i} - x_{u_j}x_{v_j}) = 2(x_{u_i} - x_{v_j})(x_{u_j} - x_{v_i}) > 0,$$

a contradiction. So, $\mathbb{H}_{t+3} = \emptyset$. This completes the proof. \square

4 Proof of Theorem 1.2

Tait's conjecture has been confirmed for $s + t \leq 6$. This implies that Tait's conjecture holds for $t \leq 3$. In this section, we only need prove Theorem 1.2 under that $t \geq 4$.

Lemma 4.1. $|\mathbb{H}_{<t} \cup \mathbb{H}_{t+2}| \leq 1$.

Proof. By the way of contradiction assume that $H_1, H_2 \in \mathbb{H}_{<t} \cup \mathbb{H}_{t+2}$, and let $|H_1| + |H_2| = pt + q$, where $0 \leq p \leq 2$ and $1 \leq q \leq t$. For each H_i , if $|H_i| \leq t - 1$, then $e(H_i) \leq \binom{|H_i|}{2}$; and if $|H_i| = t + 2$, then $e(H_i) \leq \binom{t}{2} + 2$ by Lemma 3.2. In any case, one can check that $e(H_1 \cup H_2) < p\binom{t}{2} + \binom{q}{2} = e(pK_t \cup K_q)$, a contradiction with Lemma 2.3. \square

Theorem 4.1. Let $G^*, s, t, n, p, q, \beta$ be defined as in Theorem 1.2. If $\beta = 1$, then

$$G^* \cong \begin{cases} K_{s-1} \nabla ((p-1)K_t \cup \overline{H^*}) & \text{for } q = 2 \text{ and } t = 8; \\ K_{s-1} \nabla (pK_t \cup K_q) & \text{otherwise.} \end{cases}$$

Proof. Since $\beta = 1$, by Theorem 3.1 we have $\mathbb{H}_{t+1} = \emptyset$, and by Theorem 3.3, $\mathbb{H}_{>t+2} = \emptyset$. Furthermore, by Lemma 4.1, all but at most one component H of $G^* - K$ are isomorphic to K_t . Note that $|G^* - K| = pt + q$, where $1 \leq q \leq t$. Then, either $|H| = q$ or $|H| = t + q = t + 2$.

If $q \neq 2$, then $H \cong K_q$ and $G^* - K \cong pK_t \cup K_q$, as desired. Now assume that $q = 2$. Then either $H \cong K_2$, or $H \cong \overline{H^*}$ by Theorem 3.2 (In this case, $t = 8$). So, if $t \neq 8$, then $G^* - K \cong pK_t \cup K_2$. If $t = 8$, then $H \cong \overline{H^*}$ (Otherwise, $H \cong K_2$. Then $e(K_8 \cup K_2) = 29 < 30 = e(\overline{H^*})$, a contradiction with Lemma 2.3). \square

Lemma 4.2. $|\mathbb{H}_{t+1}| \leq 2\beta - 2$, where $\beta = \lfloor \frac{t+1}{s+1} \rfloor$.

Proof. The case $\mathbb{H}_{t+1} = \emptyset$ is trivial. Assume that $\mathbb{H}_{t+1} \neq \emptyset$. Then by Theorem 3.1, $\beta \geq 2$, $H \cong \overline{H_{s,t}}$ and $e(H) = \binom{t}{2} + \beta - 1$ for any $H \in \mathbb{H}_{t+1}$.

If $|\mathbb{H}_{t+1}| \geq 2\beta$, we select 2β copies of $\overline{H_{s,t}}$ and denote it by \mathbb{F} . Then $e(\mathbb{F}) = 2\beta \left(\binom{t}{2} + \beta - 1 \right)$. Now let $\mathbb{F}' = 2\beta K_t \cup K_{2\beta}$. Note that $2\beta < t$. Then \mathbb{F}' satisfies (s, t) -property. However,

$$e(\mathbb{F}') = 2\beta \binom{t}{2} + \binom{2\beta}{2} > 2\beta \left(\binom{t}{2} + \beta - 1 \right) = e(\mathbb{F}),$$

a contradiction. So, $|\mathbb{H}_{t+1}| \leq 2\beta - 1$.

Now assume that $|\mathbb{H}_{t+1}| = 2\beta - 1$. Let \mathbb{F} be the disjoint union of $2\beta - 1$ copies of $\overline{H_{s,t}}$, and $\mathbb{F}' = (2\beta - 1)K_t \cup K_{2\beta-1}$. Then $e(\mathbb{F}') = e(\mathbb{F})$. Recall that $H_{s,t} \cong K_{1,\alpha} \cup (\beta - 1)K_{1,s}$, where $\alpha = t - (s + 1)(\beta - 1) \geq s$. So,

$$\delta(\overline{H_{s,t}}) = (s + 1)(\beta - 1) > 2(\beta - 1) \tag{17}$$

and thus $\delta(\mathbb{F}) > 2\beta - 2$. Note that $\pi(\mathbb{F}') = (t - 1, \dots, t - 1, 2\beta - 2, \dots, 2\beta - 2)$. We can observe that $\pi(\mathbb{F}) \prec \pi(\mathbb{F}')$, a contradiction. Therefore, $|\mathbb{H}_{t+1}| \leq 2\beta - 2$. \square

Lemma 4.3. If $\mathbb{H}_{t+1} \neq \emptyset$, then $\mathbb{H}_{t+2} \cup \mathbb{H}_{<t} = \emptyset$.

Proof. Suppose to the contrary that $\mathbb{H}_{t+2} \cup \mathbb{H}_{<t} \neq \emptyset$. By Lemma 4.1, $G^* - K$ contains a unique component $H_1 \in \mathbb{H}_{t+2} \cup \mathbb{H}_{<t}$. Now let $H_2 \in \mathbb{H}_{t+1}$. Then by Theorem 3.1, $\beta \geq 2$ and $H_2 \cong \overline{H_{s,t}}$.

If $|H_1| = t + 2$, then $\beta = 2$ and $H_1 \cong S^1(\overline{H_{s,t}})$ by Theorem 3.2. Note that H_1 is a subdivision of H_2 and $e(H_2) = \binom{t}{2} + \beta - 1$. We have

$$e(H_1 \cup H_2) = 2e(H_2) + 1 = 2 \left(\binom{t}{2} + \beta - 1 \right) + 1 = 2 \binom{t}{2} + 3.$$

Now let $H' = K_t \cup K_t \cup K_3$. Then $|H'| = |H_1 \cup H_2|$ and $e(H') = e(H_1 \cup H_2)$. By (17), $\delta(H_2) > 2$. Since H_1 is a subdivision of H_2 , $\delta(H_1) = 2$ and its vertex of degree two is unique. This implies that $\pi(H_1 \cup H_2) \prec \pi(H')$ and $\pi(H_1 \cup H_2) \neq \pi(H')$, a contradiction.

If $|H_1| < t$, then $H_1 \cong K_{|H_1|}$. Then

$$|H_1| \leq \beta - 1. \quad (18)$$

Otherwise, $|H_1| > \beta - 1$, then

$$e(H_1 \cup H_2) = \binom{|H_1|}{2} + \binom{t}{2} + \beta - 1 < \binom{t}{2} + \binom{|H_1|}{2} + |H_1| = e(K_t \cup K_{|H_1|+1}),$$

a contradiction with Lemma 2.3. On the other hand, we have

$$|H_1|(\beta - 1) \leq \binom{|H_1|}{2}. \quad (19)$$

Indeed, recall that $|\mathbb{H}_t| = O(\frac{n}{t})$, thus $G^* - K$ contains a disjoint union of H_1 and $|H_1|$ copies of K_t . We denote it by \mathbb{F} . Then $e(\mathbb{F}) = \binom{|H_1|}{2} + |H_1| \binom{t}{2}$. Now let $\mathbb{F}' = |H_1|H_2$. Clearly, $|\mathbb{F}| = |\mathbb{F}'|$ and $e(\mathbb{F}') = |H_1| \left(\binom{t}{2} + \beta - 1 \right)$. By Lemma 2.3, $e(\mathbb{F}') \leq e(\mathbb{F})$. It follows that (19) holds. However, (18) and (19) contradict each other. \square

Theorem 4.2. *Let $G^*, s, t, n, p, q, \beta$ be defined as in Theorem 1.2. If $\beta \geq 2$, then*

$$G^* \cong \begin{cases} K_{s-1} \nabla ((p-1)K_t \cup S^1(\overline{H_{s,t}})) & \text{if } q = \beta = 2; \\ K_{s-1} \nabla ((p-q)K_t \cup q\overline{H_{s,t}}) & \text{if } q \leq 2(\beta-1) \text{ except } q = \beta = 2; \\ K_{s-1} \nabla (pK_t \cup K_q) & \text{if } q > 2(\beta-1). \end{cases}$$

Proof. Recall that $|G^* - K| = pt + q$, where $1 \leq q \leq t$, and $H \cong \overline{H_{s,t}}$ for any $H \in \mathbb{H}_{t+1}$. Moreover, we assert that if $q \leq 2(\beta - 1)$, then $\mathbb{H}_{<t} = \emptyset$. Indeed, if $G^* - K$ contains a component H_1 with $|H_1| < t$, then $\mathbb{H}_{>t} = \emptyset$ and $|\mathbb{H}_{<t}| = 1$ by Lemmas 4.1 and 4.3. This implies that $|H_1| = q$. Now by (19), $|H_1|(\beta - 1) \leq \binom{|H_1|}{2}$, that is, $|H_1| \geq 2(\beta - 1) + 1$, contradicts $q \leq 2(\beta - 1)$.

Now we distinguish two cases. We first assume that $q \neq 2$. Then by Lemmas 4.1 and 4.3, $G^* - K$ is isomorphic to either $pK_t \cup K_q$ or $(p - q)K_t \cup q\overline{H_{s,t}}$. If $q \leq 2(\beta - 1) (< t)$, then $\mathbb{H}_{<t} = \emptyset$, and thus $G^* - K \cong (p - q)K_t \cup q\overline{H_{s,t}}$. If $q > 2(\beta - 1)$, then by Lemma 4.2, we have $G^* - K \cong pK_t \cup K_q$.

Now assume that $q = 2$. Since $\beta \geq 2$, we have $q \leq 2(\beta - 1)$ and thus $\mathbb{H}_{<t} = \emptyset$. If $\beta > 2$, then $\mathbb{H}_{t+2} = \emptyset$ by Theorem 3.2. Thus, $G^* - K \cong (p - q)K_t \cup q\overline{H_{s,t}}$. It remains the case $q = \beta = 2$. Now if $\mathbb{H}_{t+1} \neq \emptyset$, then $\mathbb{H}_{t+2} = \emptyset$ by Lemma 4.3. This implies that

$G^* - K \cong (p-2)K_t \cup \overline{H_{s,t}} \cup \overline{H_{s,t}}$. Recall that $e(\overline{H_{s,t}}) = \binom{t}{2} + \beta - 1 = \binom{t}{2} + 1$. Now let $H' = K_t \cup S^1(\overline{H_{s,t}})$. Then $|H'| = 2|\overline{H_{s,t}}| = 2t + 2$ and

$$e(H') = e(K_t) + e(\overline{H_{s,t}}) + 1 = 2e(\overline{H_{s,t}}).$$

Since $S^1(\overline{H_{s,t}})$ is a subdivision of $\overline{H_{s,t}}$, we can easily see that $\pi(\overline{H_{s,t}} \cup \overline{H_{s,t}}) \prec \pi(H')$, a contradiction. Thus, $\mathbb{H}_{t+1} = \emptyset$. It follows that $|\mathbb{H}_{t+2}| = 1$. From Theorem 3.2, we have $G^* - K \cong (p-1)K_t \cup S^1(\overline{H_{s,t}})$. This completes the proof. \square

Combining with Theorems 4.1 and 4.2, we completes the proof of Theorem 1.2.

5 Proof of Theorems 1.3 and 1.4

Throughout this section, let G^* be the extremal graph with the maximum spectral radius over all n -vertex connected $K_{1,t}$ -minor free graphs, $\rho = \rho(G^*)$ and $X = (x_1, x_2, \dots, x_n)^T$ be the Perron vector of G^* . Furthermore, assume that $u^* \in V(G^*)$ with $x_{u^*} = \max_{u \in V(G^*)} x_u$, and let $A = N_{G^*}(u^*)$, $B = V(G^*) \setminus (A \cup \{u^*\})$, $N^2(u^*) = \{d_{G^*}(w, u^*) = 2 | w \in B\}$. Above all, we need a lemma.

Lemma 5.1. [37] *Let G be a connected graph and $X = (x_1, x_2, \dots, x_n)^T$ be its Perron vector. If $x_u \geq x_v$ and $N_G(v) \setminus (N_G(u) \cup \{u\}) = S \neq \emptyset$ for some $u, v \in V(G)$, then $\rho(G - \{vw | w \in S\} + \{uw | w \in S\}) > \rho(G)$.*

If $n \leq t$, it is clear that $G^* \cong K_n$. Moreover, there exists no connected $K_{1,2}$ -minor free graph of order $n \geq 3$. Therefore, we need only consider connected $K_{1,t}$ -minor free graphs for $t \geq 3$ and $n \geq t + 1$.

Lemma 5.2. $\rho > t - 2$ and $|A| = t - 1$.

Proof. Let $K_t - e$ be the graph obtained from K_t by deleting an edge, and recall that $S^{n-t}(K_t)$ is obtained from K_t by subdividing $n - t$ times of one edge. Clearly, $S^{n-t}(K_t)$ is $K_{1,t}$ -minor free and contains $K_t - e$ as a proper subgraph. It follows that $\rho(G^*) \geq \rho(S^{n-t}(K_t)) > \rho(K_t - e)$. Let $\rho' = \rho(K_t - e)$ and $Y = (y_1, \dots, y_1, y_2, y_2)^T$ be the Perron vector of $K_t - e$, where y_1 corresponds to the $t-2$ vertices of degree $t-1$ and y_2 corresponds to the two vertices of degree $t-2$. Then we have

$$\rho' y_1 = (t-3)y_1 + 2y_2, \quad \rho' y_2 = (t-2)y_1.$$

Solving these two equations, we have $\rho'^2 - (t-3)\rho' - 2(t-2) = 0$. Since $\rho = \rho(G^*) > \rho(K_t - e)$, we have

$$\rho^2 - (t-3)\rho > 2(t-2). \quad (20)$$

Clearly, $\rho > t+2$ for $t \geq 3$, as claimed. Furthermore, $\rho x_{u^*} = \sum_{v \in A} x_v \leq |A|x_{u^*}$. It follows that $|A| \geq \rho > t-2$. On the other hand, we see that $|A| \leq \Delta(G^*) \leq t-1$ for forbidding $K_{1,t}$ -minor. Therefore, $|A| = t-1$. \square

By Lemma 5.2, $d_A(v) \leq t-2$ for any $v \in A$. Let $A_0 = \{v \in A | d_A(v) = t-2\}$ and $A_1 = A \setminus A_0$. Clearly, $d_B(v) = 0$ for any $v \in A_0$, since $\Delta(G^*) = t-1$.

Lemma 5.3. (i) $d_B(v) \leq 1$ for any $v \in A_1$.

(ii) $d_B(w) \leq 2$ for any $w \in B$. Particularly, $d_B(w) \leq 1$ for $w \in N^2(u^*)$.

Proof. (i) Suppose that $d_B(v_0) \geq 2$ for some $v_0 \in A_1$. Then G^* contains a double star with a non-pendant edge u^*v_0 and $|A| - 1 + d_B(v_0)$ leaves. Since $|A| - 1 + d_B(v_0) \geq t$, we find a K_t -minor in G^* , a contradiction. So the claim holds.

(ii) For any vertex $w \in B$, we can find a shortest path P from w to vertices in A . Clearly, $d_{B \cap V(P)}(w) \leq 1$, and particularly, $d_{B \cap V(P)}(w) = 0$ if $w \in N^2(u^*)$. Let $v \in A$ be the other endpoint of P . Then G^* contains a tree consisting of a path $P + vu^*$ and $|A| - 1 + d_{B \setminus V(P)}(w)$ leaves. Since G^* is $K_{1,t}$ -minor free, we have $|A| - 1 + d_{B \setminus V(P)}(w) \leq t - 1$. It follows that $d_B(w) \leq d_{B \cap V(P)}(w) + 1$, and thus the claim holds. \square

Lemma 5.4. Let $v_1 \in A_1$ and $w \in B$ such that v_1w is an edge, and $\overline{N}_A(v_1)$ be the set of non-neighbors of v_1 in A . Then $x_w \leq \sum_{v \in \overline{N}_A(v_1)} x_v$.

Proof. By Lemma 5.3, w is the unique neighbor of v_1 in B . Thus we have

$$\rho x_{u^*} = \sum_{v \in A \setminus \{v_1\}} x_v + x_{v_1}, \quad \rho x_{v_1} = \sum_{v \in N_A(v_1)} x_v + x_w + x_{u^*}.$$

It follows that $\rho(x_{u^*} - x_{v_1}) = (x_{v_1} - x_{u^*}) - x_w + \sum_{v \in \overline{N}_A(v_1)} x_v$. Therefore,

$$x_w = (\rho + 1)(x_{v_1} - x_{u^*}) + \sum_{v \in \overline{N}_A(v_1)} x_v \leq \sum_{v \in \overline{N}_A(v_1)} x_v,$$

since $x_{v_1} \leq x_{u^*}$. \square

Theorem 5.1. If $n = t + 1$, then $\overline{G^*} \cong \frac{n}{2}K_{1,1}$ for even n , and $\overline{G^*} \cong K_{1,2} \cup \frac{n-3}{2}K_{1,1}$ for odd n . In other words, $\overline{G^*} \cong H_{1,t}$.

Proof. Recall that $|A| = t - 1$. Then $|B| = 1$. Say $B = \{w\}$, then $N_{G^*}(w) \subseteq A_1$. We can see that $N_{G^*}(w) = A_1$. Indeed, if there exists a vertex $v \in A_1$ with $vw \notin E(G^*)$, then $\max\{d_{G^*}(v), d_{G^*}(w)\} \leq t - 2$. So, $\Delta(G^* + vw) = \Delta(G^*) = t - 1$, and thus, $G^* + vw$ also contains no $K_{1,t}$. However, $\rho(G^* + vw) > \rho(G^*)$, a contradiction.

Recall that $A_1 = \{v \in A \mid d_A(v) \leq t - 3\}$. We now claim that $d_A(v) = t - 3$ for any $v \in A_1$. Suppose that $d_A(v_0) \leq t - 4$ for some $v_0 \in A_1$ and let $v_1 \in \overline{N}_A(v_0)$. Then $d_A(v_1) \leq t - 3$, and thus $v_1 \in A_1 = N_{G^*}(w)$. Furthermore, $d_A(v_1) = t - 3$ (Otherwise, $\Delta(G^* + v_0v_1) = \Delta(G^*)$ and $\rho(G^* + v_0v_1) > \rho(G^*)$). By Corollary ??, we have $x_w \leq x_{v_0}$. Now let $G' = G^* - \{wv_1\} + \{v_0v_1\}$. Then $\Delta(G') = \Delta(G^*)$, and since $v_0 \in A_1 = N_{G^*}(w)$, G' is still a connected graph. However, by Lemma 5.1, we get $\rho(G') > \rho(G^*)$, a contradiction. So, $d_A(v) = t - 3$ for any $v \in A_1$. Note that each vertex in A_0 dominates A . The above claim implies that $G^*[A_1]$ is isomorphic to $K_{|A_1|}$ by deleting a perfect matching. Hence, $|A_1|$ is even.

It is known that $\rho(G) \leq \Delta(G)$ for any connected graph G , with equality if and only if G is Δ -regular. Now, G^* is $t - 1$ -regular if and only if $A_1 = A$. It follows that if $t + 1$ is even, then $A_1 = A$, and thus $\overline{G^*}$ is the union of $\frac{t+1}{2}$ independent edges, that is, u^*w and others within A .

If $t + 1$ is odd, then $|A| = t - 1$ is also odd, and since $|A_1|$ is even, we have $|A_1| < |A|$ (see Fig. 5). By symmetry, $x_v = x_{u^*}$ for any $v \in A_0$, and we may let $x_v = x_1$ for any $v \in A_1$. Thus, $\rho x_w = |A_1|x_1$, $\rho x_{u^*} = |A_1|x_1 + |A_0|x_{u^*}$ and

$$\rho x_1 = (|A_1| - 2)x_1 + (|A_0| + 1)x_{u^*} + x_w.$$

Note that $|A_0| + |A_1| = |A| = t - 1$. Solving above equations, we get

$$\rho^3 - (t - 3)\rho^2 - (2t - 2)\rho + |A_0||A_1| = 0. \quad (21)$$

Since $|A_1|$ is even and $t + 1$ is odd, we have $2 \leq |A_1| \leq t - 2$ and $t \geq 4$. Thus,

$$|A_0||A_1| \geq \min\{2(t - 3), t - 2\} = t - 2,$$

with $|A_0||A_1| = t - 2$ if and only if $|A_1| = t - 2$. Since ρ is the maximum spectral radius, by (21) we can see that $|A_0||A_1|$ must attain its minimum. Therefore, $|A_1| = t - 2$ and thus $|A_0| = 1$. Now, $\overline{G^*}$ is the union of $\frac{n-3}{2}$ independent edges with all endpoints in A_1 , and a path u^*wv with $v \in A_0$. This completes the proof. \square




Figure 5: The extremal graph G^* for odd $n = t + 1$.

Now we consider the case $n \geq t + 2$. Recall that $S^k(G)$ denotes a graph obtained from G by subdividing k times of one edge.

Theorem 5.2. *If $n \geq t + 2$, then $G^* \cong S^{n-t}(K_t)$.*

Proof. Observe that a graph G is $K_{1,3}$ -minor free if and only if $\Delta(G) \leq 2$. Therefore, if $t = 3$, then $G^* \cong C_n$, in other words, $G^* \cong S^{n-3}(K_3)$. Next, let $t \geq 4$. The proof is divided into several claims.

Claim 5.1. $|A_1| \geq |N^2(u^*)| \geq 2$.

Proof. Since $|A| = t - 1$ and $n \geq t + 2$, we have $N^2(u^*) \neq \emptyset$. We first show $|N^2(u^*)| \geq 2$. Suppose to the contrary that $N^2(u^*) = \{w_0\}$. By Lemma 5.3, we have $d_B(w_0) \leq 1$, and $d_B(w) \leq 2$ for each $w \in B$. Since G^* is connected, $G^*[B]$ is a pendant path $P = w_0w_1 \cdots w_{n-t-1}$. Take an arbitrary vertex $v_0 \in N_A(w_0)$. If $\overline{N}_A(w_0) \subseteq N_{G^*}(v_0)$, then $|N_{G^*}(v_0) \cup N_{G^*}(w_0)| = |A \cup \{u^*, w_0, w_1\}| = t + 2$. Hence, G^* contains a double star

with a non-pendant edge v_0w_0 and t leaves, a contradiction. Thus, there exists a vertex $v_1 \in \overline{N}_A(w_0) \cap \overline{N}_{G^*}(v_0)$. Then $d_{G^*}(v_1) \leq t - 2$.

Now, let $G' = G^* + \{v_1w_{n-t+1}\}$ and G'' be the graph obtained from G' by contracting the path $w_0w_1w_2 \cdots w_{n-t-1}v_1$ into an edge w_0v_1 . Clearly, $|G''| = t + 1$ and $\Delta(G'') = \Delta(G') = \Delta(G^*) = t - 1$. Therefore, G'' is $K_{1,t}$ -minor free, and thus G' is too. However, $\rho(G') > \rho(G^*)$, a contradiction. So $|N^2(u^*)| \geq 2$. Furthermore, by Lemma 5.3 (i), we have $|A_1| \geq |N^2(u^*)|$. \square

Now let $A_{11} = \{v \in A_1 | d_A(v) \leq t - 4\}$. Since $n \geq t + 2$, we have $|B| \geq 2$. So we may assume that $w', w'' \in B$ with $x_{w'} = \max_{w \in B} x_w$ and $x_{w''} = \max_{w \in B \setminus \{w'\}} x_w$.

Claim 5.2. *If $d_B(v_0) = 0$ for some $v_0 \in A_1$, then $x_{w'} + x_{w''} \leq x_{u^*} + \sum_{v \in A_{11}} x_v$.*

Proof. We first show $w' \in N^2(u^*)$. Otherwise, $w' \in B \setminus N^2(u^*)$. By Lemma 5.3, $d_B(w') \leq 2$, and hence, $\rho x_{w'} = \sum_{w \in N_B(w')} x_w \leq 2x_{w'}$. It follows that $\rho \leq 2$, which contradicts Lemma 5.2 as $t \geq 4$. Since $w' \in N^2(u^*)$, we have $d_B(w') \leq 1$ and

$$\rho x_{w'} \leq x_{w''} + \sum_{v \in N_A(w')} x_v. \quad (22)$$

If $w'' \in B \setminus N^2(u^*)$, then $d_B(w'') \leq 2$ and hence $\rho x_{w''} \leq x_{w'} + x_{w''}$, which implies

$$x_{w''} \leq \frac{x_{w'}}{\rho - 1}. \quad (23)$$

Combining with (22), we have

$$x_{w'} \leq \frac{\rho - 1}{\rho^2 - \rho - 1} \sum_{v \in N_A(w')} x_v. \quad (24)$$

Furthermore, by Claim 5.1, $|N^2(u^*)| \geq 2$. Hence, there exist a vertex $w \in N^2(u^*) \setminus \{w'\}$ and a vertex v in $N_A(w) \setminus N_A(w')$. So, $d_A(w') \leq |A \setminus \{v, v_0\}| = t - 3$. Combining it with (23) and (24), we have

$$x_{w'} + x_{w''} \leq \frac{\rho}{\rho - 1} x_{w'} \leq \frac{\rho}{\rho^2 - \rho - 1} \sum_{v \in N_A(w')} x_v \leq \frac{(t - 3)\rho}{\rho^2 - \rho - 1} x_{u^*}.$$

Note that (20) implies $\rho^2 - \rho - 1 > (t - 3)\rho$. So, $x_{w'} + x_{w''} < x_{u^*}$, as desired.

Now assume that $w'' \in N^2(u^*)$. By Lemma 5.3, $d_B(w'') \leq 1$ and thus $\rho x_{w''} \leq x_{w'} + \sum_{v \in N_A(w'')} x_v$. Combining with (22), we have

$$\rho(x_{w'} + x_{w''}) \leq (x_{w'} + x_{w''}) + \sum_{v \in N_A(w') \cup N_A(w'')} x_v, \quad (25)$$

since $N_A(w') \cap N_A(w'') = \emptyset$ by Lemma 5.3. Moreover,

$$|N_A(w') \cup N_A(w'')| \leq |A \setminus \{v_0\}| = t - 2.$$

If $|N_A(w') \cup N_A(w'')| \leq t - 3$, then by (25), we have $x_{w'} + x_{w''} \leq \frac{t-3}{\rho-1} x_{u^*} < x_{u^*}$. If $d_A(v_1) \leq t - 4$ for some $v_1 \in N_A(w') \cup N_A(w'')$, then $v_1 \in A_{11}$ and

$$x_{w'} + x_{w''} \leq \frac{t-3}{\rho-1} x_{u^*} + x_{v_1} < x_{u^*} + \sum_{v \in A_{11}} x_v.$$

Next we may assume that $|N_A(w') \cup N_A(w'')| = t - 2$ and $d_A(v) = t - 3$ for any $v \in N_A(w') \cup N_A(w'')$. Now $N_A(w') \cup N_A(w'') = A \setminus \{v_0\}$. Since $v_0 \in A_1$, $d_A(v_0) \leq t - 3$ and thus v_0 has a non-neighbor $v_1 \in N_A(w') \cup N_A(w'')$. Therefore, $d_A(v_1) = t - 3$, and v_1 dominates $N_A(w') \cup N_A(w'')$. Let v_1v_2 be an edge between $N_A(w')$ and $N_A(w'')$. Then v_0 is also a non-neighbor of v_2 (Otherwise, $|N_{G^*}(v_1) \cup N_{G^*}(v_2)| = |A \cup \{u^*, w', w''\}| = t + 2$ and we get a $K_{1,t}$ -minor). Now we see that $d_A(v_0) \leq t - 4$ and thus $v_0 \in A_{11}$. On the other hand, by Lemma 5.4, we have $x_{w''} \leq x_{v_0}$. It follows that $x_{w'} + x_{w''} \leq x_{u^*} + \sum_{v \in A_{11}} x_v$. \square

Claim 5.3. $d_B(v) = 1$ for each vertex $v \in A_1$, and thus $e(A, B) = |A_1|$.

Proof. By Lemma 5.3, $d_B(v) \leq 1$ for each vertex $v \in A_1$. Suppose that $d_B(v_0) = 0$ for some $v_0 \in A_1$. We can see that $\rho x_{u^*} = \sum_{v \in A} x_v$ and

$$\rho^2 x_{u^*} = \sum_{v \in A} \sum_{u \in N_{G^*}(v)} x_u = |A|x_{u^*} + \sum_{v \in A} d_A(v)x_v + \sum_{w \in B} d_A(w)x_w. \quad (26)$$

From (26) and the definitions of A_0 and A_{11} , we have

$$\begin{aligned} (\rho^2 - (t-3)\rho)x_{u^*} &= \rho^2 x_{u^*} - (t-3) \sum_{v \in A} x_v \\ &= |A|x_{u^*} + \sum_{v \in A} (d_A(v) - t + 3)x_v + \sum_{w \in B} d_A(w)x_w \\ &\leq |A|x_{u^*} + \sum_{v \in A_0} x_v - \sum_{v \in A_{11}} x_v + \sum_{w \in B} d_A(w)x_w \\ &\leq (|A| + |A_0| + e(A, B) - 2)x_{u^*} + x_{w'} + x_{w''} - \sum_{v \in A_{11}} x_v. \end{aligned} \quad (27)$$

Combining Claim 5.2 with (27), we have

$$(\rho^2 - (t-3)\rho)x_{u^*} \leq (|A| + |A_0| + e(A, B) - 1)x_{u^*}.$$

Recall that $d_B(v) = 0$ for any $v \in A_0$, and $d_B(v) \leq 1$ for any $v \in A_1$. Therefore,

$$e(A, B) = e(A_1, B) = \sum_{v \in A_1 \setminus \{v_0\}} d_B(v) \leq |A_1 \setminus \{v_0\}| = |A_1| - 1.$$

It follows that $\rho^2 - (t-3)\rho \leq 2(|A| - 1) = 2(t-2)$, which contradicts (20). So, $d_B(v) = 1$ for any $v \in A_1$ and thus $e(A, B) = e(A_1, B) = |A_1|$. \square

Claim 5.4. $|A_{11}| \leq 2$.

Proof. Suppose to the contrary that $|A_{11}| \geq 3$. Then $|A \setminus A_{11}| \leq t - 4$, and thus

$$\rho x_{u^*} = \sum_{v \in A_{11}} x_v + \sum_{v \in A \setminus A_{11}} x_v \leq \sum_{x \in A_{11}} x_v + (t-4)x_{u^*}.$$

Therefore, by Lemma 5.2, we have

$$\sum_{v \in A_{11}} x_v \geq (\rho - t + 4)x_{u^*} \geq 2x_{u^*}. \quad (28)$$

Combining with (27) and (28), we have

$$\begin{aligned} (\rho^2 - (t-3)\rho)x_{u^*} &\leq |A|x_{u^*} + \sum_{v \in A_0} x_v - \sum_{v \in A_{11}} x_v + \sum_{w \in B} d_A(w)x_w \\ &\leq (|A| + |A_0| - 2)x_{u^*} + e(A, B)x_{u^*}. \end{aligned}$$

By Claim 5.3, we have $e(A, B) = |A_1|$. Hence,

$$\rho^2 - (t-3)\rho \leq (|A| + |A_0| + |A_1| - 2) = 2|A| - 2 = 2(t-2),$$

which contradicts (20). Thus, $|A_{11}| \leq 2$, as claimed. \square

Claim 5.5. *If $|A_1| \geq 3$, then A_{10} is a clique, where $A_{10} = \{v \in A | d_A(v) = t-3\}$.*

Proof. Recall that $A_{11} = \{v \in A | d_A(v) \leq t-4\}$. Then $A_{10} = A_1 \setminus A_{11}$. Suppose to the contrary that there exist two non-adjacent vertices $v_1, v_2 \in A_{10}$. Since $d_A(v_1) = d_A(v_2) = t-3$, both v_1 and v_2 dominate $A \setminus \{v_1, v_2\}$. Since $|A_1| \geq 3$, we can take a vertex $v_3 \in A_1 \setminus \{v_1, v_2\}$. Furthermore, by Claim 5.3, there exists a unique vertex $w_i \in N_B(v_i)$ for $i \in \{1, 2, 3\}$. If $w_3 \neq w_1$, then

$$|N_{G^*}(v_1) \cup N_{G^*}(v_3)| \geq |A \cup \{u^*, w_1, w_3\}| = t+2.$$

If $w_3 \neq w_2$, then we similarly have $|N_{G^*}(v_2) \cup N_{G^*}(v_3)| \geq t+2$. If $w_1 = w_2 = w_3$, then by Claim 5.1, $|N^2(u^*)| \geq 2$ and thus there exist a vertex $w_4 \in N^2(u^*) \setminus \{w_1\}$ and a vertex $v_4 \in N_A(w_4)$. It follows that

$$|N_{G^*}(v_1) \cup N_{G^*}(v_4)| \geq |A \cup \{u^*, w_1, w_4\}| = t+2.$$

Thus we always find a $K_{1,t}$ -minor, a contradiction. \square

Now we are ready to give the final proof of the theorem. By Claim 5.1, $|A_1| \geq |N^2(u^*)| \geq 2$. If $|A_1| = 2$, then $|N^2(u^*)| = 2$, and $G^*[A \cup \{u^*\}] \cong K_t - e$, where the unique non-edge lies in A_1 . Now we can see that $d_A(w) = 1$ and $d_B(w) \leq 1$ for each $w \in N^2(u^*)$. Moreover, $d_B(w) \leq 2$ for any $w \in B \setminus N^2(u^*)$. Since G^* is an extremal graph, we can observe that $G^*[A_1 \cup B]$ is a path with both endpoints in A_1 . This implies that $G^* \cong S^{n-t}(K_t)$, as desired.

It remains the case $|A_1| \geq 3$. We shall prove that this is impossible. By Claim 5.4, $|A_{11}| \leq 2$. Suppose that $|A_{11}| = |\{v_1, v_2\}| = 2$. For $i \in \{1, 2\}$, since $d_A(v_i) \leq t-4$, v_i has at least one non-neighbor $v'_i \in A_{10}$. By the definition of A_{10} , $d_A(v'_1) = d_A(v'_2) = t-3$, and hence $v'_1 \neq v'_2$. By Claim 5.3, there exists a unique vertex $w_i \in N_B(v'_i)$ for $i \in \{1, 2\}$ (Possibly, $w_1 = w_2$). And by Lemma 5.4, $x_{w_i} \leq x_{v_i}$ for $i \in \{1, 2\}$. Thus, $x_{w_1} + x_{w_2} \leq x_{v_1} + x_{v_2} = \sum_{v \in A_{11}} x_v$. Recall that $e(A, B) = |A_1|$. Then by (27),

$$\begin{aligned} (\rho^2 - (t-3)\rho)x_{u^*} &\leq (|A| + |A_0| + e(A, B) - 2)x_{u^*} + x_{w_1} + x_{w_2} - \sum_{v \in A_{11}} x_v \\ &\leq (|A| + |A_0| + |A_1| - 2)x_{u^*} \\ &= 2(t-2)x_{u^*}, \end{aligned}$$

a contradiction with (20). Thus $|A_{11}| \leq 1$.

According to Claim 5.5, A_{10} is a clique. Recall that $A_{10} = \{v \in A \mid d_A(v) = t - 3\}$. Therefore, each vertex $v \in A_{10}$ has a non-neighbor $v_0 \in A_{11}$. And since $|A_{11}| \leq 1$, we have that $A_{11} = \{v_0\}$ and $e(\{v_0\}, A_{10}) = 0$. Let $A_{10} = \{v_1, v_2, \dots, v_{|A_{10}|}\}$. By Claim 5.3, there exists a unique vertex $w_i \in N_B(v_i)$ for each $v_i \in A_{10}$ (Possibly, $w_i = w_j$ for some $i \neq j$). Since $|N^2(u^*)| \geq 2$ and $G^*[B]$ consists of disjoint paths, we may assume without loss of generality that

- (i) $w_{|A_{10}|} \neq w_0$;
- (ii) if $|N^2(u^*)| \geq 3$, then w_0 and $w_{|A_{10}|}$ belong to two distinct paths of $G^*[B]$.

Note that $|A_1| \geq 3$, then $|A_{10}| = |A_1| - |A_{11}| \geq 2$. Let

$$G' = G^* - \{v_i w_i \mid i = 1, 2, \dots, |A_{10}| - 1\} + \{v_i v_0 \mid i = 1, 2, \dots, |A_{10}| - 1\}.$$

By Lemma 5.4, $x_{w_i} \leq x_{v_0}$ for each $w_i \in A_{10}$. Furthermore, by Lemma 5.1 we have $\rho(G') > \rho(G^*)$. We can observe that $G'[A \cup \{u^*\}] \cong K_t - e$, where the unique non-edge is $v_0 v_{|A_{10}|}$. Moreover, $G'[B]$ is still the disjoint union of paths, and the only edges between A and B are $v_0 w_0$ and $v_{|A_{10}|} w_{|A_{10}|}$. These observations imply that G' is a subgraph of $S^{n-t}(K_t)$. It follows that $\rho(S^{n-t}(K_t)) \geq \rho(G') > \rho(G^*)$, a contradiction. This completes the proof. \square

Recall that $H_{1,t}$ is a star forest of order $t + 1$, precisely, the disjoint union of $\lfloor \frac{t+1}{2} \rfloor$ stars in which all but at most one are isomorphic to $K_{1,1}$. By Theorems 5.1 and 5.2, we complete the proof of Theorem 1.3. Furthermore, we know that for any connected graph G , $\rho(G) \leq \Delta(G)$, with equality if and only if G is a Δ -regular graph. One can see that Theorem 1.4 is a direct corollary of Theorem 1.3.

References

- [1] L. Babai, B. Guiduli, Spectral extrema for graphs: the Zarankiewicz problem, *Electron. J. Combin.* **16** (2009) no. 1, Research Paper 123, 8 pp.
- [2] B. Bollobás, V. Nikiforov, Cliques and the spectral radius, *J. Combin. Theory Ser. B* **97** (2007) 859–865.
- [3] B.N. Boots, G.F. Royle, A conjecture on the maximum value of the principal eigenvalue of a planar graph, *Geogr. Anal.* **23** (1991) 276–282.
- [4] D.S. Cao, A. Vince, The spectral radius of a planar graph, *Linear Algebra Appl.* **187** (1993) 251–257.
- [5] S. Cioabă, D.N. Desai, M. Tait, The spectral radius of graphs with no odd wheels, arXiv:2104.07729v1.
- [6] S. Cioabă, L.H. Feng, M. Tait, X.D. Zhang, The maximum spectral radius of graphs without friendship subgraphs, *Electron. J. Combin.* **27** (2020) no. 4, Research Paper 4.22, 19 pp.
- [7] M.Z. Chen, A.M. Liu, X.D. Zhang, On the spectral radius of graphs without a star forest, *Discrete Math.* **344** (2021), no. 4, 112269, 12 pp.

- [8] M.Z. Chen, A.M. Liu, X.D. Zhang, Spectral extremal results with forbidding linear forests, *Graphs Combin.* **35** (2019) 335–351.
- [9] M. Chudnovsky, B. Reed, P. Seymour, The edge-density for $K_{2,t}$ minors, *J. Combin. Theory Ser. B* **101** (2011) 18–46.
- [10] D. Cvetković, P. Rowlinson, The largest eigenvalue of a graph: A survey, *Linear Multilinear Algebra* **28** (1990) 3–33.
- [11] G. Ding, T. Johnson, P. Seymour, Spanning trees with many leaves, *J. Graph Theory* **37** (2001) 189–197.
- [12] M.N. Ellingham, X.Y. Zha, The spectral radius of graphs on surfaces, *J. Combin. Theory Ser. B* **78** (2000) 45–56.
- [13] L.H. Feng, G.H. Yu, X.D. Zhang, Spectral radius of graphs with given matching number, *Linear Algebra Appl.* **422** (2007) 133–138.
- [14] Z. Füredi, M. Simonovits, The history of degenerate (bipartite) extremal graph problems, *Bolyai Soc. Studies (The Erdős Centennial)* **25** (2013) 167–262.
- [15] J. Gao, X.M. Hou, The spectral radius of graphs without long cycles, *Linear Algebra Appl.* **566** (2019) 17–33.
- [16] Y. Hong, On the spectral radius and the genus of graphs, *J. Combin. Theory Ser. B* **65** (1995) 262–268.
- [17] Y. Hong, Sharp upper bounds of the spectral radius of graphs in terms of genus, *J. Combin. Theory Ser. B* **74** (1998) 153–159.
- [18] Y. Hong, Tree-width, clique-minors, and eigenvalues, *Discrete Math.* **274** (2004) 281–287.
- [19] B.L. Li, B. Ning, The stability method, eigenvalues and cycles of consecutive lengths, arXiv:2102.03855.
- [20] H.Q. Lin, B. Ning, A complete solution to the Cvetković-Rowlinson conjecture, *J. Graph Theory* (2021) 1–10.
- [21] Y. Li, Y. Peng, The spectral radius of graphs with no intersecting odd cycles, arXiv:2106.00587.
- [22] W. Mantel, Problem 28, soln. by H. Gouventak, W. Mantel, J. Teixeira de Mattes, F. Schuh and W.A. Wythoff, *Wiskundige Opgaven* **10** (1907) 60–61.
- [23] B. Ning, X. Peng, Extensions of the Erdős-Gallai theorem and Luo’s theorem, *Combin. Probab. Comput.* **29** (2020) 128–136.
- [24] V. Nikiforov, The spectral radius of graphs with no $K_{2,t}$ -minor, *Linear Algebra Appl.* **531** (2017) 510–515.
- [25] V. Nikiforov, A spectral condition for odd cycles in graphs, *Linear Algebra Appl.* **428** (2008) 1492–1498.
- [26] V. Nikiforov, Some new results in extremal graph theory: In surveys in Combinatorics 2011, *London Math. Society Lecture Note Ser.* **392** (2011) 141–181.
- [27] V. Nikiforov, The spectral radius of graphs without paths and cycles of specified length, *Linear Algebra Appl.* **432** (2010) 2243–2256.

- [28] V. Nikiforov, Bounds on graph eigenvalues II, *Linear Algebra Appl.* **427** (2007) 183–189.
- [29] V. Nikiforov, A contribution to the Zarankiewicz problem, *Linear Algebra Appl.* **432** (2010) 1405–1411.
- [30] J.L. Shu, Y. Hong, Upper bounds for the spectral radii of outerplanar graphs and Halin graphs, *Chinese Ann. Math. Ser. A* **21** (2000), no.6, 677–682.
- [31] M. Tait, J. Tobin, Three conjectures in extremal spectral graph theory, *J. Combin. Theory Ser. B* **126** (2017) 137–161.
- [32] M. Tait, The Colin de Verdière parameter, excluded minors, and the spectral radius, *J. Combin. Theory Ser. A* **166** (2019) 42–58.
- [33] P. Turán, On an extremal problem in graph theory, *Mat. Fiz. Lapok* **48** (1941) 436–452.
- [34] T. Bıyıkođlu, J. Leydold, Graphs with given degree sequence and maximal spectral radius, *Electron. J. Combin.* **15** (2008), no. 1, Research Paper 119, 9 pp.
- [35] B. Wang, W. Chen, L. Fang, Extremal spectral radius of $K_{3,3}/K_{2,4}$ -minor free graphs, *Linear Algebra Appl.* **628** (2021) 103–114.
- [36] H. Wilf, Spectral bounds for the clique and independence numbers of graphs, *J. Combin. Theory Ser. B* **40** (1986) 113–117.
- [37] B.F. Wu, E.L. Xiao, Y. Hong, The spectral radius of trees on k pendant vertices, *Linear Algebra Appl.* **395** (2005) 343–349.
- [38] M.Q. Zhai, H.Q. Lin, Spectral extrema of graphs: forbidden hexagon, *Discrete Math.* **343** (2020), no. 10, 112028, 6 pp.
- [39] M.Q. Zhai, H.Q. Lin, A strengthening of the spectral color critical edge theorem: books and theta graphs, arXiv:2102.04041.
- [40] M.Q. Zhai, B. Wang, Proof of a conjecture on the spectral radius of C_4 -free graphs, *Linear Algebra Appl.* **437** (2012) 1641–1647.
- [41] X.Z. Zhan, Matrix theory, Graduate Studies in Mathematics, 147. American Mathematical Society, Providence, RI, 2013. x+253 pp.
- [42] Z. Dvořák, B. Mohar, Spectral radius of finite and infinite planar graphs and of graphs of bounded genus, *J. Combin. Theory Ser. B* **100** (2010), no. 6, 729–739.