

**SOME NEW THEOREMS ON GENERATING FUNCTIONS AND THEIR
APPLICATIONS ON ODD AND EVEN CERTAIN NUMBERS ATTACHED TO p AND q
PARAMETERS**

Nabiha Saba

LMAM Laboratory and Department of Mathematics,
Mohamed Seddik Ben Yahia University, Jijel, Algeria

E-Mail: sabarnhf1994@gmail.com

Ali Boussayoud*

LMAM Laboratory and Department of Mathematics,
Mohamed Seddik Ben Yahia University, Jijel, Algeria

E-Mail: aboussayoud@yahoo.fr

***Corresponding author**

Abstract

In this study, we first provide some new theorems by using the symmetrizing operator $\delta_{e_1 e_2}^k$ for $k \in \{0, 1, 2, 3, 4\}$. After that, by using this theorems we introduce a new family of generating functions of odd and even terms of (p, q) -Fibonacci numbers, (p, q) -Lucas numbers, (p, q) -Pell numbers, (p, q) -Pell Lucas numbers, (p, q) -Jacobsthal numbers and (p, q) -Jacobsthal Lucas numbers. Then, we give the new generating functions of the products of these (p, q) -numbers with odd and even phrases of (p, q) -numbers.

2010 Mathematics Subject Classification. Primary 05E05; Secondary 11B39.

Key Words and Phrases. Symmetric functions; Generating functions; Odd (p, q) -numbers; Even (p, q) -numbers.

1. Introduction and preliminary results

The generating function can be applied to solve many problems in mathematics. This concept makes it possible to know the terms of a recurring and linear sequence with constant coefficients. This procedure demonstrates how one can find the terms of a recurrent and linear sequence without calculating the previous terms. In this work, we investigate the generating functions of odd and even (p, q) -numbers and some products of them. In this section, we first present the (p, q) -numbers and some generalization of them.

For over several years, there are many recursive sequences that have been studied in the literatures. the famous examples of these sequences are (p, q) -Fibonacci and (p, q) -Lucas numbers, (p, q) -Jacobsthal and (p, q) -Jacobsthal Lucas numbers, (p, q) -Pell and (p, q) -Pell Lucas numbers (see [8, 7, 6, 10, 22, 16, 15]), because they are extensively used in various research areas. The authors in [14] defined the generalized (p, q) -Fibonacci sequence $\{f_{p,q,n}(\alpha, \beta, \gamma)\}_{n \geq 0}$, generalized (p, q) -Pell sequence $\{l_{p,q,n}(\alpha, \beta, \gamma)\}_{n \geq 0}$ and generalized (p, q) -Jacobsthal sequence $\{C_{p,q,n}(\alpha, \beta, \gamma)\}_{n \geq 0}$ as follows:

$$f_{p,q,n} := \begin{cases} \alpha, & \text{if } n = 0 \\ \beta + \gamma p, & \text{if } n = 1 \\ pf_{p,q,n-1} + qf_{p,q,n-2}, & \text{if } n \geq 2 \end{cases}, \quad (1.1)$$

$$l_{p,q,n} := \begin{cases} \alpha, & \text{if } n = 0 \\ \beta + 2\gamma p, & \text{if } n = 1 \\ 2pl_{p,q,n-1} + ql_{p,q,n-2}, & \text{if } n \geq 2 \end{cases}, \quad (1.2)$$

and

$$C_{p,q,n} := \begin{cases} \alpha, & \text{if } n = 0 \\ \beta + \gamma p, & \text{if } n = 1 \\ pC_{p,q,n-1} + 2qC_{p,q,n-2}, & \text{if } n \geq 2 \end{cases}. \quad (1.3)$$

The particular cases of Eqs. (1.1), (1.2) and (1.3) are listed below:

Remark 1. Let $\alpha = \gamma = 0$ and $\beta = 1$ in Eqs. (1.1), (1.2) and (1.3), then we get:

- (p, q) -Fibonacci numbers $\{F_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} F_{p,q,0} = 0, F_{p,q,1} = 1 \\ F_{p,q,n} = pF_{p,q,n-1} + qF_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.4)$$

- (p, q) -Pell numbers $\{P_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} P_{p,q,0} = 0, P_{p,q,1} = 1 \\ P_{p,q,n} = 2pP_{p,q,n-1} + qP_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.5)$$

- (p, q) -Jacobsthal numbers $\{J_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} J_{p,q,0} = 0, J_{p,q,1} = 1 \\ J_{p,q,n} = pJ_{p,q,n-1} + 2qJ_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.6)$$

Remark 2. Let $\alpha = 2$, $\gamma = 1$ and $\beta = 0$ in Eqs. (1.1), (1.2) and (1.3), then we get:

- (p, q) -Lucas numbers $\{L_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} L_{p,q,0} = 2, L_{p,q,1} = p \\ L_{p,q,n} = pL_{p,q,n-1} + qL_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.7)$$

- (p, q) -Pell Lucas numbers $\{Q_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} Q_{p,q,0} = 2, Q_{p,q,1} = 2p \\ Q_{p,q,n} = 2pQ_{p,q,n-1} + qQ_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.8)$$

- (p, q) -Jacobsthal Lucas numbers $\{j_{p,q,n}\}_{n \geq 0}$, defined recursively by:

$$\begin{cases} j_{p,q,0} = 2, j_{p,q,1} = p \\ j_{p,q,n} = pj_{p,q,n-1} + 2qj_{p,q,n-2} \quad (n \geq 2) \end{cases} \quad (1.9)$$

In fact, these (p, q) -numbers are the generalization of some k -numbers, like k -Fibonacci, k -Pell, k -Jacobsthal, k -Lucas, k -Pell Lucas and k -Jacobsthal Lucas numbers, the following names, notations and recurrence relations (see Tab. 1) are used for the special cases for p and q of Eqs. (1.4)-(1.9) respectively.

p	q	k -numbers	Notation	Recurrence relation
k	1	k -Fibonacci	$F_{k,n}$	$\begin{cases} F_{k,0} = 0, F_{k,1} = 1 \\ F_{k,n} = kF_{k,n-1} + F_{k,n-2} \quad (n \geq 2) \end{cases}$
1	k	k -Pell	$P_{k,n}$	$\begin{cases} P_{k,0} = 0, P_{k,1} = 1 \\ P_{k,n} = 2P_{k,n-1} + kP_{k,n-2} \quad (n \geq 2) \end{cases}$
k	1	k -Jacobsthal	$J_{k,n}$	$\begin{cases} J_{k,0} = 0, J_{k,1} = 1 \\ J_{k,n} = kJ_{k,n-1} + 2J_{k,n-2} \quad (n \geq 2) \end{cases}$
k	1	k -Lucas	$L_{k,n}$	$\begin{cases} L_{k,0} = 2, L_{k,1} = k \\ L_{k,n} = kL_{k,n-1} + L_{k,n-2} \quad (n \geq 2) \end{cases}$
1	k	k -Pell Lucas	$Q_{k,n}$	$\begin{cases} Q_{k,0} = 2, Q_{k,1} = 2 \\ Q_{k,n} = 2Q_{k,n-1} + kQ_{k,n-2} \quad (n \geq 2) \end{cases}$
k	1	k -Jacobsthal Lucas	$j_{k,n}$	$\begin{cases} j_{k,0} = 2, j_{k,1} = k \\ j_{k,n} = kj_{k,n-1} + 2j_{k,n-2} \quad (n \geq 2) \end{cases}$

Table 1. Some k -numbers.

For the special case $k = 1$ in the Tab. 1, the sequences of Fibonacci, Pell, Jacobsthal, Lucas, Pell Lucas and Jacobsthal Lucas numbers are respectively obtained.

Next, we present some backgrounds and results about the symmetric functions.

Definition 1. [13] Let k and n be two positive integers and $\{a_1, a_2, \dots, a_n\}$ are set of given variables the k^{th} complete homogeneous symmetric function $h_k(a_1, a_2, \dots, a_n)$ is defined by:

$$h_k(a_1, a_2, \dots, a_n) = \sum_{i_1+i_2+\dots+i_n=k} a_1^{i_1} a_2^{i_2} \dots a_n^{i_n} \quad (k \geq 0),$$

with $i_1, i_2, \dots, i_n \geq 0$.

Remark 3. Set $h_0(a_1, a_2, \dots, a_n) = 1$, by usual convention. For $k < 0$, we set $h_k(a_1, a_2, \dots, a_n) = 0$.

Definition 2. [1] Let A and E be any two alphabets. We define $S_n(A - E)$ by the following form:

$$\frac{\prod_{e \in E} (1 - ez)}{\prod_{a \in A} (1 - az)} = \sum_{n=0}^{\infty} S_n(A - E) z^n, \quad (1.10)$$

with the condition $S_n(A - E) = 0$ for $n < 0$.

Equation (1.10) can be rewritten in the following form:

$$\sum_{n=0}^{\infty} S_n(A - E) z^n = \left(\sum_{n=0}^{\infty} S_n(A) z^n \right) \times \left(\sum_{n=0}^{\infty} S_n(-E) z^n \right),$$

where

$$S_n(A - E) = \sum_{j=0}^n S_{n-j}(-E) S_j(A).$$

Definition 3. [11] Given a function f on \mathbb{R}^n , the divided difference operator is defined as follows:

$$\partial_{e_i e_{i+1}}(f) = \frac{f(e_1, \dots, e_i, e_{i+1}, \dots, e_n) - f(e_1, \dots, e_{i-1}, e_{i+1}, e_i, e_{i+2}, \dots, e_n)}{e_i - e_{i+1}}.$$

Definition 4. [18] Let n be positive integer and $E = \{e_1, e_2\}$ are set of given variables. Then, the n^{th} symmetric function $S_n(e_1 + e_2)$ is defined by:

$$S_n(E) = S_n(e_1 + e_2) = \frac{e_1^{n+1} - e_2^{n+1}}{e_1 - e_2},$$

with

$$\begin{aligned} S_0(E) &= S_0(e_1 + e_2) = 1, \\ S_1(E) &= S_1(e_1 + e_2) = e_1 + e_2, \\ S_2(E) &= S_2(e_1 + e_2) = e_1^2 + e_1 e_2 + e_2^2, \\ &\vdots \end{aligned}$$

Proposition 1. [15] For $n \in \mathbb{N}$, the symmetric functions of (p, q) -Fibonacci and (p, q) -Lucas numbers, (p, q) -Jacobsthal and (p, q) -Jacobsthal Lucas numbers, (p, q) -Pell and (p, q) -Pell Lucas numbers are given by:

$$F_{p,q,n} = S_{n-1}(e_1 + [-e_2]) \text{ and } L_{p,q,n} = 2S_n(e_1 + [-e_2]) - pS_{n-1}(e_1 + [-e_2]), \text{ with } e_{1,2} = \frac{p \pm \sqrt{p^2 + 4q}}{2}.$$

$$J_{p,q,n} = S_{n-1}(e_1 + [-e_2]) \text{ and } j_{p,q,n} = 2S_n(e_1 + [-e_2]) - pS_{n-1}(e_1 + [-e_2]), \text{ with } e_{1,2} = \frac{p \pm \sqrt{p^2 + 8q}}{2}.$$

$$P_{p,q,n} = S_{n-1}(e_1 + [-e_2]) \text{ and } Q_{p,q,n} = 2S_n(e_1 + [-e_2]) - 2pS_{n-1}(e_1 + [-e_2]), \text{ with } e_{1,2} = p \pm \sqrt{p^2 + q}.$$

Definition 5. [5] Given an alphabet $E = \{e_1, e_2\}$, the symmetrizing operator $\delta_{e_1 e_2}^k$ is defined by:

$$\delta_{e_1 e_2}^k(f) = \frac{e_1^k f(e_1) - e_2^k f(e_2)}{e_1 - e_2}, \quad (k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} = \{0, 1, 2, 3, \dots\}). \quad (1.11)$$

Remark 4. If $k = 0$, the operator (1.11) gives us:

$$\begin{aligned} \delta_{e_1 e_2}^0(f) &= \frac{f(e_1) - f(e_2)}{e_1 - e_2} \\ &= \partial_{e_1 e_2}(f). \end{aligned}$$

In the present article, the sections are organized as follows: Section 2 gives new theorems by using the symmetrizing operator $\delta_{e_1 e_2}^k$. By using the theorems given in the previous section, the new generating functions for odd and even terms of (p, q) -numbers are constructed in Section 3. Section 4 gives the new generating functions for the products of (p, q) -numbers with odd and even (p, q) -numbers. Finally, some concluded remarks are discussed in Section 5.

2. New theorems on symmetric functions and their proofs

In this part, we are now in a position to provide some new theorems by using the symmetrizing operator $\delta_{e_1 e_2}^k$ for $k \in \{0, 1, 2, 3, 4\}$. We now begin with the following theorem.

Theorem 1. *Given two alphabets $A = \{a_1, a_2\}$ and $E = \{e_1, e_2\}$, we have:*

$$\sum_{n=0}^{\infty} S_n(A) S_{2n-1}(E) z^n = \frac{(a_1 + a_2)(e_1 + e_2)z - a_1 a_2(e_1 + e_2) \left((e_1 + e_2)^2 - 2e_1 e_2 \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.1)$$

Proof. By applying the divided difference operator $\partial_{e_1 e_2}$ to the series $f(e_1 z) = \sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n$, the left-hand side of the formula (2.1) can be written as:

$$\begin{aligned} \partial_{e_1 e_2} f(e_1 z) &= \frac{\sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n - \sum_{n=0}^{\infty} S_n(A) e_2^{2n} z^n}{e_1 - e_2} \\ &= \sum_{n=0}^{\infty} S_n(A) \left(\frac{e_1^{2n} - e_2^{2n}}{e_1 - e_2} \right) z^n \\ &= \sum_{n=0}^{\infty} S_n(A) S_{2n-1}(E) z^n. \end{aligned}$$

By applying the divided difference operator $\partial_{e_1 e_2}$ to the series $f(e_1 z) = \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}$, the right-hand

side of the formula (2.1) can be expressed as:

$$\begin{aligned} \partial_{e_1 e_2} f(e_1 z) &= \frac{\frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n} - \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n}}{e_1 - e_2} \\ &= \frac{\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n - \sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}{(e_1 - e_2) \left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\ &= \frac{\sum_{n=0}^{\infty} S_n(-A) \frac{e_2^{2n} - e_1^{2n}}{e_1 - e_2} z^n}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \end{aligned}$$

$$\begin{aligned}
 &= \frac{S_1(-A) \frac{e_2^2 - e_1^2}{e_1 - e_2} z + S_2(-A) \frac{e_2^4 - e_1^4}{e_1 - e_2} z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
 &= \frac{-S_1(-A) (e_1 + e_2) z - S_2(-A) \left((e_1 + e_2)^3 - 2e_1 e_2 (e_1 + e_2) \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
 &= \frac{(a_1 + a_2) (e_1 + e_2) z - a_1 a_2 (e_1 + e_2) \left((e_1 + e_2)^2 - 2e_1 e_2 \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}.
 \end{aligned}$$

Thus, this completes the proof. \square

Theorem 2. Given two alphabets $A = \{a_1, a_2\}$ and $E = \{e_1, e_2\}$, we have:

$$\sum_{n=0}^{\infty} S_n(A) S_{2n}(E) z^n = \frac{1 + e_1 e_2 (a_1 + a_2) z - e_1 e_2 a_1 a_2 \left((e_1 + e_2)^2 - e_1 e_2 \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.2)$$

Proof. By applying the operator $\delta_{e_1 e_2}$ to the series $f(e_1 z) = \sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n$, the left-hand side of the formula (2.2) can be written as:

$$\begin{aligned}
 \delta_{e_1 e_2} f(e_1 z) &= \frac{e_1 \sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n - e_2 \sum_{n=0}^{\infty} S_n(A) e_2^{2n} z^n}{e_1 - e_2} \\
 &= \sum_{n=0}^{\infty} S_n(A) \left(\frac{e_1^{2n+1} - e_2^{2n+1}}{e_1 - e_2} \right) z^n \\
 &= \sum_{n=0}^{\infty} S_n(A) S_{2n}(E) z^n.
 \end{aligned}$$

By applying the operator $\delta_{e_1 e_2}$ to the series $f(e_1 z) = \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}$, the right-hand side of the formula

(2.2) can be expressed as:

$$\begin{aligned}
\delta_{e_1 e_2} f(e_1 z) &= \frac{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n} - \frac{\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n}{\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n} \\
&= \frac{e_1 \sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n - e_2 \sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}{(e_1 - e_2) \left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
&= \frac{\sum_{n=0}^{\infty} S_n(-A) \frac{e_1 e_2^{2n} - e_2 e_1^{2n}}{e_1 - e_2} z^n}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
&= \frac{S_0(-A) - e_1 e_2 S_1(-A) z - S_2(-A) \frac{e_1^3 - e_2^3}{e_1 - e_2} z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
&= \frac{S_0(-A) - e_1 e_2 S_1(-A) z - S_2(-A) e_1 e_2 \left((e_1 + e_2)^2 - e_1 e_2 \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\
&= \frac{1 + e_1 e_2 (a_1 + a_2) z - e_1 e_2 a_1 a_2 \left((e_1 + e_2)^2 - e_1 e_2 \right) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}.
\end{aligned}$$

Thus, this completes the proof. \square

Theorem 3. Given two alphabets $A = \{a_1, a_2\}$ and $E = \{e_1, e_2\}$, we have:

$$\sum_{n=0}^{\infty} S_n(A) S_{2n+1}(E) z^n = \frac{e_1 + e_2 - a_1 a_2 e_1^2 e_2^2 (e_1 + e_2) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.3)$$

Proof. By applying the operator $\delta_{e_1 e_2}^2$ to the series $f(e_1 z) = \sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n$, the left-hand side of the formula (2.3) can be written as:

$$\begin{aligned} \delta_{e_1 e_2}^2 f(e_1 z) &= \frac{e_1^2 \sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n - e_2^2 \sum_{n=0}^{\infty} S_n(A) e_2^{2n} z^n}{e_1 - e_2} \\ &= \sum_{n=0}^{\infty} S_n(A) \left(\frac{e_1^{2n+2} - e_2^{2n+2}}{e_1 - e_2} \right) z^n \\ &= \sum_{n=0}^{\infty} S_n(A) S_{2n+1}(E) z^n. \end{aligned}$$

By applying the operator $\delta_{e_1 e_2}^2$ to the series $f(e_1 z) = \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}$, the right-hand side of the formula (2.3) can be expressed as:

$$\begin{aligned} \delta_{e_1 e_2}^2 f(e_1 z) &= \frac{\frac{e_1^2}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n} - \frac{e_2^2}{\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n}}{e_1 - e_2} \\ &= \frac{e_1^2 \sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n - e_2^2 \sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}{(e_1 - e_2) \left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\ &= \frac{\sum_{n=0}^{\infty} S_n(-A) \frac{e_1^2 e_2^{2n} - e_2^2 e_1^{2n}}{e_1 - e_2} z^n}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\ &= \frac{S_0(-A) \frac{e_1^2 - e_2^2}{e_1 - e_2} + S_2(-A) \frac{e_1^2 e_2^4 - e_2^3 e_1^4}{e_1 - e_2} z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\ &= \frac{S_0(-A) (e_1 + e_2) - S_2(-A) (e_1 e_2)^2 (e_1 + e_2) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)} \\ &= \frac{(e_1 + e_2) - a_1 a_2 e_1^2 e_2^2 (e_1 + e_2) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \end{aligned}$$

Thus, this completes the proof. \square

Based on the relationship (2.3) we get:

$$\sum_{n=0}^{\infty} S_{n-1}(A) S_{2n-1}(E) z^n = \frac{(e_1 + e_2) z - a_1 a_2 e_1^2 e_2^2 (e_1 + e_2) z^3}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.4)$$

Theorem 4. *Given two alphabets $A = \{a_1, a_2\}$ and $E = \{e_1, e_2\}$, we have:*

$$\sum_{n=0}^{\infty} S_n(A) S_{2n+2}(E) z^n = \frac{(e_1 + e_2)^2 - e_1 e_2 - e_1^2 e_2^2 (a_1 + a_2) z - a_1 a_2 e_1^3 e_2^3 z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.5)$$

Proof. The proof is similar to the proof of Theorem 3, but now using the operator $\delta_{e_1 e_2}^3$ to the series $\sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n = \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}$. \square

Note that, based on the relationship (2.5), we get:

$$\sum_{n=0}^{\infty} S_{n-1}(A) S_{2n}(E) z^n = \frac{\left((e_1 + e_2)^2 - e_1 e_2 \right) z - e_1^2 e_2^2 (a_1 + a_2) z^2 - a_1 a_2 e_1^3 e_2^3 z^3}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.6)$$

Theorem 5. *Given two alphabets $A = \{a_1, a_2\}$ and $E = \{e_1, e_2\}$, we have:*

$$\sum_{n=0}^{\infty} S_n(A) S_{2n+3}(E) z^n = \frac{(e_1 + e_2) \left((e_1 + e_2)^2 - 2e_1 e_2 \right) - e_1^2 e_2^2 (a_1 + a_2) (e_1 + e_2) z}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.7)$$

Proof. The proof is similar to the proof of Theorem 3, but now using the operator $\delta_{e_1 e_2}^4$ to the series $\sum_{n=0}^{\infty} S_n(A) e_1^{2n} z^n = \frac{1}{\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n}$. \square

From (2.7), we get:

$$\sum_{n=0}^{\infty} S_{n-1}(A) S_{2n+1}(E) z^n = \frac{(e_1 + e_2) \left((e_1 + e_2)^2 - 2e_1 e_2 \right) z - e_1^2 e_2^2 (a_1 + a_2) (e_1 + e_2) z^2}{\left(\sum_{n=0}^{\infty} S_n(-A) e_1^{2n} z^n \right) \left(\sum_{n=0}^{\infty} S_n(-A) e_2^{2n} z^n \right)}. \quad (2.8)$$

3. Generating functions of odd and even some (p, q) -numbers

In this part, we now derive the new generating functions for odd and even phrases of some special numbers with parameters p and q . By making use of the Theorems 1, 2 and 3 we investigate some special cases ($A = \{1, 0\}$ and $E = \{e_1, -e_2\}$) as follows:

Lemma 1. *Given an alphabet $E = \{e_1, -e_2\}$, then we have:*

$$\sum_{n=0}^{\infty} S_{2n-1}(e_1 + [-e_2]) z^n = \frac{(e_1 - e_2) z}{1 - \left((e_1 - e_2)^2 + 2e_1 e_2 \right) z + e_1^2 e_2^2 z^2}. \quad (3.1)$$

Lemma 2. Given an alphabet $E = \{e_1, -e_2\}$, then we have:

$$\sum_{n=0}^{\infty} S_{2n}(e_1 + [-e_2])z^n = \frac{1 - e_1e_2z}{1 - \left((e_1 - e_2)^2 + 2e_1e_2\right)z + e_1^2e_2^2z^2}. \quad (3.2)$$

Lemma 3. Given an alphabet $E = \{e_1, -e_2\}$, then we have:

$$\sum_{n=0}^{\infty} S_{2n+1}(e_1 + [-e_2])z^n = \frac{e_1 - e_2}{1 - \left((e_1 - e_2)^2 + 2e_1e_2\right)z + e_1^2e_2^2z^2}. \quad (3.3)$$

This part consists of three cases.

Case 1. The substitution of $\begin{cases} e_1 - e_2 = p \\ e_1e_2 = q \end{cases}$ in (3.1), (3.2) and (3.3), we obtain:

$$\sum_{n=0}^{\infty} S_{2n-1}(e_1 + [-e_2])z^n = \frac{pz}{1 - (p^2 + 2q)z + q^2z^2}, \quad (3.4)$$

$$\sum_{n=0}^{\infty} S_{2n}(e_1 + [-e_2])z^n = \frac{1 - qz}{1 - (p^2 + 2q)z + q^2z^2}, \quad (3.5)$$

$$\sum_{n=0}^{\infty} S_{2n+1}(e_1 + [-e_2])z^n = \frac{p}{1 - (p^2 + 2q)z + q^2z^2}, \quad (3.6)$$

respectively, and we have the following propositions and theorems.

Proposition 2. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Fibonacci numbers is given by:

$$\sum_{n=0}^{\infty} F_{p,q,2n}z^n = \frac{pz}{1 - (p^2 + 2q)z + q^2z^2}. \quad (3.7)$$

Proof. By [15], we have

$$F_{p,q,n} = S_{n-1}(e_1 + [-e_2]).$$

Writing $(2n)$ instead of (n) , we get

$$F_{p,q,2n} = S_{2n-1}(e_1 + [-e_2]).$$

Then, by relation (3.4), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} F_{p,q,2n}z^n &= \sum_{n=0}^{\infty} S_{2n-1}(e_1 + [-e_2])z^n \\ &= \frac{pz}{1 - (p^2 + 2q)z + q^2z^2}. \end{aligned}$$

Hence, we obtain the desired result. \square

Proposition 3. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Fibonacci numbers is given by:

$$\sum_{n=0}^{\infty} F_{p,q,2n+1}z^n = \frac{1 - qz}{1 - (p^2 + 2q)z + q^2z^2}. \quad (3.8)$$

Proof. By referred to [15], we have

$$F_{p,q,n} = S_{n-1}(e_1 + [-e_2]).$$

Substituting n by $(2n + 1)$, we obtain

$$F_{p,q,2n+1} = S_{2n}(e_1 + [-e_2]).$$

From the relation (3.5), we can write

$$\begin{aligned} \sum_{n=0}^{\infty} F_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n \\ &= \frac{1 - qz}{1 - (p^2 + 2q)z + q^2 z^2}. \end{aligned}$$

As required. \square

Theorem 6. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Lucas numbers is given by:

$$\sum_{n=0}^{\infty} L_{p,q,2n} z^n = \frac{2 - (p^2 + 2q)z}{1 - (p^2 + 2q)z + q^2 z^2}. \quad (3.9)$$

Proof. We have

$$L_{p,q,n} = 2S_n (e_1 + [-e_2]) - pS_{n-1} (e_1 + [-e_2]), \quad (\text{see [15]}).$$

By setting $n = 2n$, we get

$$L_{p,q,2n} = 2S_{2n} (e_1 + [-e_2]) - pS_{2n-1} (e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,2n} z^n &= \sum_{n=0}^{\infty} (2S_{2n} (e_1 + [-e_2]) - pS_{2n-1} (e_1 + [-e_2])) z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n - p \sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n. \end{aligned}$$

Multiplying the equation (3.5) by (2) and adding it to the equation obtained by (3.4) multiplying by $(-p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,2n} z^n &= \frac{2(1 - qz)}{1 - (p^2 + 2q)z + q^2 z^2} - \frac{p^2 z}{1 - (p^2 + 2q)z + q^2 z^2} \\ &= \frac{2 - (p^2 + 2q)z}{1 - (p^2 + 2q)z + q^2 z^2}. \end{aligned}$$

Which completes the proof. \square

Theorem 7. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Lucas numbers is given by:

$$\sum_{n=0}^{\infty} L_{p,q,2n+1} z^n = \frac{p + pqz}{1 - (p^2 + 2q)z + q^2 z^2}. \quad (3.10)$$

Proof. Recall that, we have $L_{p,q,n} = 2S_n (e_1 + [-e_2]) - pS_{n-1} (e_1 + [-e_2])$, (see [15]).

By putting $n = 2n + 1$, we get

$$L_{p,q,2n+1} = 2S_{2n+1} (e_1 + [-e_2]) - pS_{2n} (e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} (2S_{2n+1} (e_1 + [-e_2]) - pS_{2n} (e_1 + [-e_2])) z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n+1} (e_1 + [-e_2]) z^n - p \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n. \end{aligned}$$

Multiplying the equation (3.6) by (2) and adding it to the equation obtained by (3.5) multiplying by $(-p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,2n+1} z^n &= \frac{2p}{1 - (p^2 + 2q)z + q^2 z^2} - \frac{p^2(1 - qz)}{1 - (p^2 + 2q)z + q^2 z^2} \\ &= \frac{p + pqz}{1 - (p^2 + 2q)z + q^2 z^2}. \end{aligned}$$

Thus, this completes the proof. □

- Taking $p = k$ and $q = 1$ in the Eqs. (3.7)-(3.10), we get the new generating functions of even and odd k -Fibonacci and k -Lucas numbers. The calculation and results are listed in the Tab. 2.

Coefficient of z^n	Generating function
$F_{k,2n}$	$\frac{kz}{1 - (k^2 + 2)z + z^2}$
$F_{k,2n+1}$	$\frac{1 - z}{1 - (k^2 + 2)z + z^2}$
$L_{k,2n}$	$\frac{2 - (k^2 + 2)z}{1 - (k^2 + 2)z + z^2}$
$L_{k,2n+1}$	$\frac{k + kz}{1 - (k^2 + 2)z + z^2}$

Table 2. New generating functions for even and odd k -Fibonacci and k -Lucas numbers.

Remark 5. If we take $k = 1$ in the Tab. 2, we get the generating functions of even and odd Fibonacci and Lucas numbers given in the article of Mezo [12] as follows:

$$\sum_{n=0}^{\infty} F_{2n} z^n = \frac{z}{1 - 3z + z^2}, \quad \sum_{n=0}^{\infty} F_{2n+1} z^n = \frac{1 - z}{1 - 3z + z^2}, \quad \sum_{n=0}^{\infty} L_{2n} z^n = \frac{2 - 3z}{1 - 3z + z^2} \quad \text{and} \quad \sum_{n=0}^{\infty} L_{2n+1} z^n = \frac{1 + z}{1 - 3z + z^2}.$$

Case 2. The substitution of $\begin{cases} e_1 - e_2 = 2p \\ e_1 e_2 = q \end{cases}$ in (3.1), (3.2) and (3.3), we obtain:

$$\sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n = \frac{2pz}{1 - 2(2p^2 + q)z + q^2 z^2}, \tag{3.11}$$

$$\sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n = \frac{1 - qz}{1 - 2(2p^2 + q)z + q^2 z^2}, \tag{3.12}$$

$$\sum_{n=0}^{\infty} S_{2n+1} (e_1 + [-e_2]) z^n = \frac{2p}{1 - 2(2p^2 + q)z + q^2 z^2}, \tag{3.13}$$

respectively, and we have the following propositions and theorems.

Proposition 4. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Pell numbers is given by:

$$\sum_{n=0}^{\infty} P_{p,q,2n} z^n = \frac{2pz}{1 - 2(2p^2 + q)z + q^2 z^2}. \tag{3.14}$$

Proof. Recall that, we have

$$P_{p,q,n} = S_{n-1} (e_1 + [-e_2]), \quad (\text{see [15]}).$$

Writing $(2n)$ instead of (n) , we obtain

$$P_{p,q,2n} = S_{2n-1} (e_1 + [-e_2]).$$

Therefore, according to the relationship (3.11), we get

$$\begin{aligned} \sum_{n=0}^{\infty} P_{p,q,2n} z^n &= \sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n \\ &= \frac{2pz}{1 - 2(2p^2 + q)z + q^2 z^2}. \end{aligned}$$

Hence, we obtain the desired result. \square

Proposition 5. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Pell numbers is given by:

$$\sum_{n=0}^{\infty} P_{p,q,2n+1} z^n = \frac{1 - qz}{1 - 2(2p^2 + q)z + q^2 z^2}. \quad (3.15)$$

Proof. By referred to [15], we have

$$P_{p,q,n} = S_{n-1} (e_1 + [-e_2]).$$

Substituting n by $(2n + 1)$, we obtain

$$P_{p,q,2n+1} = S_{2n} (e_1 + [-e_2]).$$

From the relation (3.12), we can write

$$\begin{aligned} \sum_{n=0}^{\infty} P_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n \\ &= \frac{1 - qz}{1 - 2(2p^2 + q)z + q^2 z^2}. \end{aligned}$$

As required. \square

Theorem 8. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Pell Lucas numbers is given by:

$$\sum_{n=0}^{\infty} Q_{p,q,2n} z^n = \frac{2 - 2(2p^2 + q)z}{1 - 2(2p^2 + q)z + q^2 z^2}. \quad (3.16)$$

Proof. Once more, by [15] we have

$$Q_{p,q,n} = 2S_n (e_1 + [-e_2]) - 2pS_{n-1} (e_1 + [-e_2]).$$

By setting $n = 2n$, we get

$$Q_{p,q,2n} = 2S_{2n} (e_1 + [-e_2]) - 2pS_{2n-1} (e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,2n} z^n &= \sum_{n=0}^{\infty} (2S_{2n} (e_1 + [-e_2]) - 2pS_{2n-1} (e_1 + [-e_2])) z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n - 2p \sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n. \end{aligned}$$

Multiplying the equation (3.12) by (2) and adding it to the equation obtained by (3.11) multiplying by $(-2p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,2n} z^n &= \frac{2(1 - qz)}{1 - 2(2p^2 + q)z + q^2 z^2} - \frac{4p^2 z}{1 - 2(2p^2 + q)z + q^2 z^2} \\ &= \frac{2 - 2(2p^2 + q)z}{1 - 2(2p^2 + q)z + q^2 z^2}. \end{aligned}$$

Which completes the proof. \square

Theorem 9. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Pell Lucas numbers is given by:

$$\sum_{n=0}^{\infty} Q_{p,q,2n+1} z^n = \frac{2p + 2pqz}{1 - 2(2p^2 + q)z + q^2 z^2}. \quad (3.17)$$

Proof. Recall that, we have $Q_{p,q,n} = 2S_n(e_1 + [-e_2]) - 2pS_{n-1}(e_1 + [-e_2])$, (see [15]).

By putting $n = 2n + 1$, we get

$$Q_{p,q,2n+1} = 2S_{2n+1}(e_1 + [-e_2]) - 2pS_{2n}(e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,2n+1}z^n &= \sum_{n=0}^{\infty} (2S_{2n+1}(e_1 + [-e_2]) - 2pS_{2n}(e_1 + [-e_2]))z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n+1}(e_1 + [-e_2])z^n - 2p \sum_{n=0}^{\infty} S_{2n}(e_1 + [-e_2])z^n. \end{aligned}$$

Multiplying the equation (3.13) by (2) and adding it to the equation obtained by (3.12) multiplying by $(-2p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,2n+1}z^n &= \frac{4p}{1 - 2(2p^2 + q)z + q^2z^2} - \frac{2p(1 - qz)}{1 - 2(2p^2 + q)z + q^2z^2} \\ &= \frac{2p + 2pqz}{1 - 2(2p^2 + q)z + q^2z^2}. \end{aligned}$$

Thus, this completes the proof. □

- Taking $p = 1$ and $q = k$ in the Eqs. (3.14)-(3.17), we get the new generating functions of even and odd k -Pell and k -Pell Lucas numbers. The calculation and results are listed in the Tab. 3.

Coefficient of z^n	Generating function
$P_{k,2n}$	$\frac{2z}{1-2(k+2)z+k^2z^2}$
$P_{k,2n+1}$	$\frac{1-kz}{1-2(k+2)z+k^2z^2}$
$Q_{k,2n}$	$\frac{2-2(k+2)z}{1-2(k+2)z+k^2z^2}$
$Q_{k,2n+1}$	$\frac{2+2kz}{1-2(k+2)z+k^2z^2}$

Table 3. New generating functions for even and odd k -Pell and k -Pell Lucas numbers.

Remark 6. If we take $k = 1$ in the Tab. 3, we get the generating functions of even and odd Pell and Pell Lucas numbers given in the article of Mezo [12] as follows:

$$\sum_{n=0}^{\infty} P_{2n}z^n = \frac{2z}{1 - 6z + z^2}, \quad \sum_{n=0}^{\infty} P_{2n+1}z^n = \frac{1 - z}{1 - 6z + z^2}, \quad \sum_{n=0}^{\infty} Q_{2n}z^n = \frac{2 - 6z}{1 - 6z + z^2} \quad \text{and} \quad \sum_{n=0}^{\infty} Q_{2n+1}z^n = \frac{2 + 2z}{1 - 6z + z^2}.$$

Case 3. The substitution of $\begin{cases} e_1 - e_2 = p \\ e_1 e_2 = 2q \end{cases}$ in (3.1), (3.2) and (3.3), we obtain:

$$\sum_{n=0}^{\infty} S_{2n-1}(e_1 + [-e_2])z^n = \frac{pz}{1 - (p^2 + 4q)z + 4q^2z^2}, \tag{3.18}$$

$$\sum_{n=0}^{\infty} S_{2n}(e_1 + [-e_2])z^n = \frac{1 - 2qz}{1 - (p^2 + 4q)z + 4q^2z^2}, \tag{3.19}$$

$$\sum_{n=0}^{\infty} S_{2n+1}(e_1 + [-e_2])z^n = \frac{p}{1 - (p^2 + 4q)z + 4q^2z^2}, \tag{3.20}$$

respectively, and we have the following propositions and theorems.

Proposition 6. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Jacobsthal numbers is given by:

$$\sum_{n=0}^{\infty} J_{p,q,2n}z^n = \frac{pz}{1 - (p^2 + 4q)z + 4q^2z^2}. \tag{3.21}$$

Proof. By [15], we have

$$J_{p,q,n} = S_{n-1} (e_1 + [-e_2]).$$

Writing $(2n)$ instead of (n) , we get

$$J_{p,q,2n} = S_{2n-1} (e_1 + [-e_2]).$$

Then, by relation (3.18), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} J_{p,q,2n} z^n &= \sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n \\ &= \frac{pz}{1 - (p^2 + 4q)z + 4q^2 z^2}. \end{aligned}$$

Hence, we obtain the desired result. \square

Proposition 7. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Jacobsthal numbers is given by:

$$\sum_{n=0}^{\infty} J_{p,q,2n+1} z^n = \frac{1 - 2qz}{1 - (p^2 + 4q)z + 4q^2 z^2}. \quad (3.22)$$

Proof. By referred to [15], we have

$$J_{p,q,n} = S_{n-1} (e_1 + [-e_2]).$$

Substituting n by $(2n + 1)$, we obtain

$$J_{p,q,2n+1} = S_{2n} (e_1 + [-e_2]).$$

From the relation (3.19), we can write

$$\begin{aligned} \sum_{n=0}^{\infty} J_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n \\ &= \frac{1 - 2qz}{1 - (p^2 + 4q)z + 4q^2 z^2}. \end{aligned}$$

As required. \square

Theorem 10. For $n \in \mathbb{N}$, the new generating function of even (p, q) -Jacobsthal Lucas numbers is given by:

$$\sum_{n=0}^{\infty} j_{p,q,2n} z^n = \frac{2 - (p^2 + 4q)z}{1 - (p^2 + 4q)z + 4q^2 z^2}. \quad (3.23)$$

Proof. We have

$$j_{p,q,n} = 2S_n (e_1 + [-e_2]) - pS_{n-1} (e_1 + [-e_2]), \quad (\text{see [15]}).$$

By setting $n = 2n$, we get

$$j_{p,q,2n} = 2S_{2n} (e_1 + [-e_2]) - pS_{2n-1} (e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} j_{p,q,2n} z^n &= \sum_{n=0}^{\infty} (2S_{2n} (e_1 + [-e_2]) - pS_{2n-1} (e_1 + [-e_2])) z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n} (e_1 + [-e_2]) z^n - p \sum_{n=0}^{\infty} S_{2n-1} (e_1 + [-e_2]) z^n. \end{aligned}$$

Multiplying the equation (3.19) by (2) and adding it to the equation obtained by (3.18) multiplying by $(-p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} j_{p,q,2n} z^n &= \frac{2(1-2qz)}{1-(p^2+4q)z+4q^2z^2} - \frac{p^2z}{1-(p^2+4q)z+4q^2z^2} \\ &= \frac{2-(p^2+4q)z}{1-(p^2+4q)z+4q^2z^2}. \end{aligned}$$

Which completes the proof. □

Theorem 11. For $n \in \mathbb{N}$, the new generating function of odd (p, q) -Jacobsthal Lucas numbers is given by:

$$\sum_{n=0}^{\infty} j_{p,q,2n+1} z^n = \frac{p+2pqz}{1-(p^2+4q)z+4q^2z^2}. \tag{3.24}$$

Proof. By [15] we have

$$j_{p,q,n} = 2S_n(e_1 + [-e_2]) - pS_{n-1}(e_1 + [-e_2]).$$

By putting $n = 2n + 1$, we get

$$j_{p,q,2n+1} = 2S_{2n+1}(e_1 + [-e_2]) - pS_{2n}(e_1 + [-e_2]).$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} j_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} (2S_{2n+1}(e_1 + [-e_2]) - pS_{2n}(e_1 + [-e_2])) z^n \\ &= 2 \sum_{n=0}^{\infty} S_{2n+1}(e_1 + [-e_2]) z^n - p \sum_{n=0}^{\infty} S_{2n}(e_1 + [-e_2]) z^n. \end{aligned}$$

Multiplying the equation (3.20) by (2) and adding it to the equation obtained by (3.19) multiplying by $(-p)$, then we obtain the following equality

$$\begin{aligned} \sum_{n=0}^{\infty} j_{p,q,2n+1} z^n &= \frac{2p}{1-(p^2+4q)z+4q^2z^2} - \frac{p(1-2qz)}{1-(p^2+4q)z+4q^2z^2} \\ &= \frac{p+2pqz}{1-(p^2+4q)z+4q^2z^2}. \end{aligned}$$

Thus, this completes the proof. □

- Taking $p = k$ and $q = 1$ in the Eqs. (3.21)-(3.24), we get the new generating functions of even and odd k -Jacobsthal and k -Jacobsthal Lucas numbers. The calculation and results are listed in the Tab. 4.

Coefficient of z^n	Generating function
$J_{k,2n}$	$\frac{kz}{1-(k^2+4)z+4z^2}$
$J_{k,2n+1}$	$\frac{1-2z}{1-(k^2+4)z+4z^2}$
$j_{k,2n}$	$\frac{2-(k^2+4)z}{1-(k^2+4)z+4z^2}$
$j_{k,2n+1}$	$\frac{k+2kz}{1-(k^2+4)z+4z^2}$

Table 4. New generating functions for even and odd k -Jacobsthal and k -Jacobsthal Lucas numbers.

Remark 7. If we take $k = 1$ in the Tab. 4, we get the generating functions of even and odd Jacobsthal and Jacobsthal Lucas numbers given in the article of Mezo [12] as follows:

$$\sum_{n=0}^{\infty} J_{2n} z^n = \frac{z}{1-5z+4z^2}, \quad \sum_{n=0}^{\infty} J_{2n+1} z^n = \frac{1-2z}{1-5z+4z^2}, \quad \sum_{n=0}^{\infty} j_{2n} z^n = \frac{2-5z}{1-5z+4z^2} \quad \text{and} \quad \sum_{n=0}^{\infty} j_{2n+1} z^n = \frac{1+2z}{1-5z+4z^2}.$$

4. Generating functions of the products of (p, q) -numbers with odd and even (p, q) -numbers

We now consider the Theorems 1, 2, 3, 4 and 5 in order to derive a new generating functions of binary products of (p, q) -numbers with odd and even (p, q) -numbers.

We consider the following sets:

$$A = \{a_1, -a_2\} \text{ and } E = \{e_1, -e_2\}.$$

By changing a_2 to $(-a_2)$ and e_2 to $(-e_2)$ in Eqs. (2.1), (2.2), (2.3), (2.4), (2.6) and (2.8), it becomes

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{(a_1 - a_2) (e_1 - e_2) z + a_1 a_2 (e_1 - e_2) \left((e_1 - e_2)^2 + 2e_1 e_2 \right) z^2}{P(z)}, \quad (4.1)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{1 - e_1 e_2 (a_1 - a_2) z - e_1 e_2 a_1 a_2 \left((e_1 - e_2)^2 + e_1 e_2 \right) z^2}{P(z)}, \quad (4.2)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{e_1 - e_2 + a_1 a_2 e_1^2 e_2^2 (e_1 - e_2) z^2}{P(z)}, \quad (4.3)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{(e_1 - e_2) z + a_1 a_2 e_1^2 e_2^2 (e_1 - e_2) z^3}{P(z)}, \quad (4.4)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{\left((e_1 - e_2)^2 + e_1 e_2 \right) z - e_1^2 e_2^2 (a_1 - a_2) z^2 - a_1 a_2 e_1^3 e_2^3 z^3}{P(z)}, \quad (4.5)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{(e_1 - e_2) \left((e_1 - e_2)^2 + 2e_1 e_2 \right) z - e_1^2 e_2^2 (a_1 - a_2) (e_1 - e_2) z^2}{P(z)}, \quad (4.6)$$

respectively, with $P(z) = (1 - a_1 e_1^2 z)(1 - a_1 e_2^2 z)(1 + a_2 e_1^2 z)(1 + a_2 e_2^2 z)$, and we have three cases.

Case 1. Let us now consider the following conditions for Eqs. (4.1)-(4.6):

$$\begin{cases} a_1 - a_2 = p & \text{and} \\ a_1 a_2 = q \end{cases} \quad \text{and} \quad \begin{cases} e_1 - e_2 = p \\ e_1 e_2 = q \end{cases}.$$

Then it yields

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{p^2 z + pq (p^2 + 2q) z^2}{D_1}, \quad (4.7)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{1 - pqz - q^2 (p^2 + q) z^2}{D_1}, \quad (4.8)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{p + pq^3 z^2}{D_1}, \quad (4.9)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{pz + pq^3 z^3}{D_1}, \quad (4.10)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{(p^2 + q) z - pq^2 z^2 - q^4 z^3}{D_1}, \quad (4.11)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{p (p^2 + 2q) z - p^2 q^2 z^2}{D_1}, \quad (4.12)$$

with

$$D_1 = 1 - p (p^2 + 2q) z - q (p^4 + 3p^2 q + 2q^2) z^2 + pq^3 (p^2 + 2q) z^3 + q^6 z^4.$$

And we deduce the following corollaries and theorems.

Corollary 1. *Let n be a natural number. Then we have the new generating function of the product of (p, q) -Fibonacci numbers $(F_{p,q,n}F_{p,q,2n})$:*

$$\begin{aligned} \sum_{n=0}^{\infty} F_{p,q,n}F_{p,q,2n}z^n &= \frac{pz + pq^3z^3}{D_1} \\ &= \frac{pz + pq^3z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}, \end{aligned} \quad (4.13)$$

with $F_{p,q,n}F_{p,q,2n} = S_{n-1}(a_1 + [-a_2])S_{2n-1}(e_1 + [-e_2])$.

Corollary 2. *For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Fibonacci numbers $(F_{p,q,n}F_{p,q,2n+1})$ is given by:*

$$\begin{aligned} \sum_{n=0}^{\infty} F_{p,q,n}F_{p,q,2n+1}z^n &= \frac{(p^2 + q)z - pq^2z^2 - q^4z^3}{D_1} \\ &= \frac{(p^2 + q)z - pq^2z^2 - q^4z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}, \end{aligned} \quad (4.14)$$

with $F_{p,q,n}F_{p,q,2n+1} = S_{n-1}(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])$.

Theorem 12. *For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Lucas numbers $(L_{p,q,n}L_{p,q,2n})$ is given by:*

$$\sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n}z^n = \frac{4 - 3p(p^2 + 2q)z - 2q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}. \quad (4.15)$$

Proof. We have

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n}z^n &= \sum_{n=0}^{\infty} \left(\begin{array}{l} (2S_n(a_1 + [-a_2]) - pS_{n-1}(a_1 + [-a_2])) \\ \times (2S_{2n}(e_1 + [-e_2]) - pS_{2n-1}(e_1 + [-e_2])) \end{array} \right) z^n \\ &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n \\ &\quad - 2p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n-1}(e_1 + [-e_2])z^n \\ &\quad - 2p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n \\ &\quad + p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n-1}(e_1 + [-e_2])z^n. \end{aligned}$$

Using the relationships (4.7), (4.8), (4.10) and (4.11), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n}z^n &= \frac{4(1 - pqz - q^2(p^2 + q)z^2)}{D_1} - \frac{2p(p^2z + pq(p^2 + 2q)z^2)}{D_1} \\ &\quad - \frac{2p((p^2 + q)z - pq^2z^2 - q^4z^3)}{D_1} + \frac{p^2(pz + pq^3z^3)}{D_1} \\ &= \frac{4 - 3p(p^2 + 2q)z - 2q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3}{D_1} \\ &= \frac{4 - 3p(p^2 + 2q)z - 2q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}. \end{aligned}$$

So, the proof is completed. \square

Theorem 13. *Let n be a natural number. Then we have the new generating function of the product of (p, q) -Lucas numbers $(L_{p,q,n}L_{p,q,2n+1})$:*

$$\sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n+1}z^n = \frac{2p - p^2(p^2 + q)z + 3pq^2(p^2 + 2q)z^2 - p^2q^4z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}. \quad (4.16)$$

Proof. We have

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n+1}z^n &= \sum_{n=0}^{\infty} \left(\begin{array}{l} (2S_n(a_1 + [-a_2]) - pS_{n-1}(a_1 + [-a_2])) \\ \times (2S_{2n+1}(e_1 + [-e_2]) - pS_{2n}(e_1 + [-e_2])) \end{array} \right) z^n \\ &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n+1}(e_1 + [-e_2])z^n \\ &\quad - 2p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n \\ &\quad - 2p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n+1}(e_1 + [-e_2])z^n \\ &\quad + p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n. \end{aligned}$$

Using the relationships (4.8), (4.9), (4.11) and (4.12), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} L_{p,q,n}L_{p,q,2n+1}z^n &= \frac{4(p + pq^3z^2)}{D_1} - \frac{2p(1 - pqz - q^2(p^2 + q)z^2)}{D_1} \\ &\quad - \frac{2p(p(p^2 + 2q)z - p^2q^2z^2)}{D_1} + \frac{p^2((p^2 + q)z - pq^2z^2 - q^4z^3)}{D_1} \\ &= \frac{2p - p^2(p^2 + q)z + 3pq^2(p^2 + 2q)z^2 - p^2q^4z^3}{D_1} \\ &= \frac{2p - p^2(p^2 + q)z + 3pq^2(p^2 + 2q)z^2 - p^2q^4z^3}{1 - p(p^2 + 2q)z - q(p^4 + 3p^2q + 2q^2)z^2 + pq^3(p^2 + 2q)z^3 + q^6z^4}. \end{aligned}$$

So, the proof is completed. \square

Corollary 3. *Putting $p = k$ and $q = 1$ in Eqs. (4.13)-(4.16) gives the following new generating functions:*

$$\begin{aligned} \sum_{n=0}^{\infty} F_{k,n}F_{k,2n}z^n &= \frac{kz + kz^3}{1 - k(k^2 + 2)z - (k^4 + 3k^2 + 2)z^2 + k(k^2 + 2)z^3 + z^4}, \\ \sum_{n=0}^{\infty} F_{k,n}F_{k,2n+1}z^n &= \frac{(k^2 + 1)z - kz^2 - z^3}{1 - k(k^2 + 2)z - (k^4 + 3k^2 + 2)z^2 + k(k^2 + 2)z^3 + z^4}, \\ \sum_{n=0}^{\infty} L_{k,n}L_{k,2n}z^n &= \frac{4 - 3k(k^2 + 2)z - 2(k^4 + 3k^2 + 2)z^2 + k(k^2 + 2)z^3}{1 - k(k^2 + 2)z - (k^4 + 3k^2 + 2)z^2 + k(k^2 + 2)z^3 + z^4}, \\ \sum_{n=0}^{\infty} L_{k,n}L_{k,2n+1}z^n &= \frac{2k - k^2(k^2 + 1)z + 3k(k^2 + 2)z^2 - k^2z^3}{1 - k(k^2 + 2)z - (k^4 + 3k^2 + 2)z^2 + k(k^2 + 2)z^3 + z^4}. \end{aligned}$$

- Put $k = 1$ in the Corollary 3, we obtain the following table:

Coefficient of z^n	Generating function
$F_n F_{2n}$	$\frac{z+z^3}{1-3z-6z^2+3z^3+z^4}$
$F_n F_{2n+1}$	$\frac{2z-z^2-z^3}{1-3z-6z^2+3z^3+z^4}$
$L_n L_{2n}$	$\frac{4-9z-12z^2+3z^3}{1-3z-6z^2+3z^3+z^4}$
$L_n L_{2n+1}$	$\frac{2-2z+9z^2-z^3}{1-3z-6z^2+3z^3+z^4}$

Table 5. A new generating functions of the products of some numbers.

Case 2. Let us now consider the following conditions for Eqs. (4.1)-(4.6):

$$\begin{cases} a_1 - a_2 = 2p \\ a_1 a_2 = q \end{cases} \quad \text{and} \quad \begin{cases} e_1 - e_2 = 2p \\ e_1 e_2 = q \end{cases} .$$

Then it yields

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{4p^2 z + 4pq (2p^2 + q) z^2}{D_2}, \quad (4.17)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{1 - 2pqz - q^2 (4p^2 + q) z^2}{D_2}, \quad (4.18)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{2p + 2pq^3 z^2}{D_2}, \quad (4.19)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{2pz + 2pq^3 z^3}{D_2}, \quad (4.20)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{(4p^2 + q) z - 2pq^2 z^2 - q^4 z^3}{D_2}, \quad (4.21)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{4p (2p^2 + q) z - 4p^2 q^2 z^2}{D_2}, \quad (4.22)$$

with

$$D_2 = 1 - 4p (2p^2 + q) z - 2q (8p^4 + 6p^2 q + q^2) z^2 + 4pq^3 (2p^2 + q) z^3 + q^6 z^4.$$

And we deduce the following corollaries and theorems.

Corollary 4. For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Pell numbers $(P_{p,q,n} P_{p,q,2n})$ is given by:

$$\begin{aligned} \sum_{n=0}^{\infty} P_{p,q,n} P_{p,q,2n} z^n &= \frac{2pz + 2pq^3 z^3}{D_2} \\ &= \frac{2pz + 2pq^3 z^3}{1 - 4p (2p^2 + q) z - 2q (8p^4 + 6p^2 q + q^2) z^2 + 4pq^3 (2p^2 + q) z^3 + q^6 z^4}, \end{aligned} \quad (4.23)$$

with $P_{p,q,n} P_{p,q,2n} = S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2])$.

Corollary 5. Let n be a natural number. Then we have the new generating function of the product of (p, q) -Pell numbers $(P_{p,q,n} P_{p,q,2n+1})$:

$$\begin{aligned} \sum_{n=0}^{\infty} P_{p,q,n} P_{p,q,2n+1} z^n &= \frac{(4p^2 + q) z - 2pq^2 z^2 - q^4 z^3}{D_2} \\ &= \frac{(4p^2 + q) z - 2pq^2 z^2 - q^4 z^3}{1 - 4p (2p^2 + q) z - 2q (8p^4 + 6p^2 q + q^2) z^2 + 4pq^3 (2p^2 + q) z^3 + q^6 z^4} \end{aligned} \quad (4.24)$$

with $P_{p,q,n} P_{p,q,2n+1} = S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2])$.

Theorem 14. For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Pell Lucas numbers $(Q_{p,q,n}Q_{p,q,2n})$ is given by:

$$\sum_{n=0}^{\infty} Q_{p,q,n}Q_{p,q,2n}z^n = \frac{4 - 12p(2p^2 + q)z - 4q(8p^4 + 4p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3}{1 - 4p(2p^2 + q)z - 2q(8p^4 + 6p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3 + q^6z^4}. \quad (4.25)$$

Proof. We have

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,n}Q_{p,q,2n}z^n &= \sum_{n=0}^{\infty} \left(\begin{array}{l} (2S_n(a_1 + [-a_2]) - 2pS_{n-1}(a_1 + [-a_2])) \\ \times (2S_{2n}(e_1 + [-e_2]) - 2pS_{2n-1}(e_1 + [-e_2])) \end{array} \right) z^n \\ &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n \\ &\quad - 4p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2])S_{2n-1}(e_1 + [-e_2])z^n \\ &\quad - 4p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n}(e_1 + [-e_2])z^n \\ &\quad + 4p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2])S_{2n-1}(e_1 + [-e_2])z^n. \end{aligned}$$

Using the relationships (4.17), (4.18), (4.20) and (4.21), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} Q_{p,q,n}Q_{p,q,2n}z^n &= \frac{4(1 - 2pqz - q^2(4p^2 + q)z^2)}{D_2} - \frac{4p(4p^2z + 4pq(2p^2 + q)z^2)}{D_2} \\ &\quad - \frac{4p((4p^2 + q)z - 2pq^2z^2 - q^4z^3)}{D_2} + \frac{4p^2(2pz + 2pq^3z^3)}{D_2} \\ &= \frac{4 - 12p(2p^2 + q)z - 4q(8p^4 + 4p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3}{D_2} \\ &= \frac{4 - 12p(2p^2 + q)z - 4q(8p^4 + 4p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3}{1 - 4p(2p^2 + q)z - 2q(8p^4 + 6p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3 + q^6z^4}. \end{aligned}$$

So, the proof is completed. \square

Theorem 15. Let n be a natural number. Then we have the new generating function of the product of (p, q) -Pell Lucas numbers $(Q_{p,q,n}Q_{p,q,2n+1})$:

$$\sum_{n=0}^{\infty} Q_{p,q,n}Q_{p,q,2n+1}z^n = \frac{4p - 4p^2(4p^2 + q)z + 12pq^2(2p^2 + q)z^2 - 4p^2q^4z^3}{1 - 4p(2p^2 + q)z - 2q(8p^4 + 6p^2q + q^2)z^2 + 4pq^3(2p^2 + q)z^3 + q^6z^4}. \quad (4.26)$$

Proof. We have

$$\begin{aligned}
 \sum_{n=0}^{\infty} Q_{p,q,n} Q_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} \left(\begin{aligned} &(2S_n(a_1 + [-a_2]) - 2pS_{n-1}(a_1 + [-a_2])) \\ &\times (2S_{2n+1}(e_1 + [-e_2]) - 2pS_{2n}(e_1 + [-e_2])) \end{aligned} \right) z^n \\
 &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n+1}(e_1 + [-e_2]) z^n \\
 &\quad - 4p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n \\
 &\quad - 4p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n+1}(e_1 + [-e_2]) z^n \\
 &\quad + 4p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n.
 \end{aligned}$$

Using the relationships (4.18), (4.19), (4.21) and (4.22), we obtain

$$\begin{aligned}
 \sum_{n=0}^{\infty} Q_{p,q,n} Q_{p,q,2n+1} z^n &= \frac{4(2p + 2pq^3 z^2)}{D_2} - \frac{4p(1 - 2pqz - q^2(4p^2 + q)z^2)}{D_2} \\
 &\quad - \frac{4p(4p(2p^2 + q)z - 4p^2 q^2 z^2)}{D_2} + \frac{4p^2((4p^2 + q)z - 2pq^2 z^2 - q^4 z^3)}{D_2} \\
 &= \frac{4p - 4p^2(4p^2 + q)z + 12pq^2(2p^2 + q)z^2 - 4p^2 q^4 z^3}{D_2} \\
 &= \frac{4p - 4p^2(4p^2 + q)z + 12pq^2(2p^2 + q)z^2 - 4p^2 q^4 z^3}{1 - 4p(2p^2 + q)z - 2q(8p^4 + 6p^2 q + q^2)z^2 + 4pq^3(2p^2 + q)z^3 + q^6 z^4}.
 \end{aligned}$$

So, the proof is completed. □

Corollary 6. Taking $p = 1$ and $q = k$ in Eqs. (4.23)-(4.26) gives the following new generating functions:

$$\begin{aligned}
 \sum_{n=0}^{\infty} P_{k,n} P_{k,2n} z^n &= \frac{2z + 2k^3 z^3}{1 - 4(k + 2)z - 2k(k^2 + 6k + 8)z^2 + 4k^3(k + 2)z^3 + k^6 z^4}. \\
 \sum_{n=0}^{\infty} P_{k,n} P_{k,2n+1} z^n &= \frac{(k + 4)z - 2k^2 z^2 - k^4 z^3}{1 - 4(k + 2)z - 2k(k^2 + 6k + 8)z^2 + 4k^3(k + 2)z^3 + k^6 z^4}. \\
 \sum_{n=0}^{\infty} Q_{k,n} Q_{k,2n} z^n &= \frac{4 - 12(k + 2)z - 4k(k^2 + 4k + 8)z^2 + 4k^3(k + 2)z^3}{1 - 4(k + 2)z - 2k(k^2 + 6k + 8)z^2 + 4k^3(k + 2)z^3 + k^6 z^4}. \\
 \sum_{n=0}^{\infty} Q_{k,n} Q_{k,2n+1} z^n &= \frac{4 - 4(k + 4)z + 12k^2(k + 2)z^2 - 4k^4 z^3}{1 - 4(k + 2)z - 2k(k^2 + 6k + 8)z^2 + 4k^3(k + 2)z^3 + k^6 z^4}.
 \end{aligned}$$

- Put $k = 1$ in the Corollary 6, we obtain the following table:

Coefficient of z^n	Generating function
$P_n P_{2n}$	$\frac{2z + 2z^3}{1 - 12z - 30z^2 + 12z^3 + z^4}$
$P_n P_{2n+1}$	$\frac{5z - 2z^2 - z^3}{1 - 12z - 30z^2 + 12z^3 + z^4}$
$Q_n Q_{2n}$	$\frac{4 - 36z - 52z^2 + 12z^3}{1 - 12z - 30z^2 + 12z^3 + z^4}$
$Q_n Q_{2n+1}$	$\frac{4 - 20z + 36z^2 - 4z^3}{1 - 12z - 30z^2 + 12z^3 + z^4}$

Table 6. A new generating functions of the products of some numbers.

Case 3. Let us now consider the following conditions for Eqs. (4.1)-(4.6)

$$\begin{cases} a_1 - a_2 = p \\ a_1 a_2 = 2q \end{cases} \quad \text{and} \quad \begin{cases} e_1 - e_2 = p \\ e_1 e_2 = 2q \end{cases}.$$

Then it give

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{p^2 z + 2pq (p^2 + 4q) z^2}{D_3}, \quad (4.27)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{1 - 2pqz - 4q^2 (p^2 + 2q) z^2}{D_3}, \quad (4.28)$$

$$\sum_{n=0}^{\infty} S_n (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{p + 8pq^3 z^2}{D_3}, \quad (4.29)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2]) z^n = \frac{pz + 8pq^3 z^3}{D_3}, \quad (4.30)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2]) z^n = \frac{(p^2 + 2q) z - 4pq^2 z^2 - 16q^4 z^3}{D_3}, \quad (4.31)$$

$$\sum_{n=0}^{\infty} S_{n-1} (a_1 + [-a_2]) S_{2n+1} (e_1 + [-e_2]) z^n = \frac{p (p^2 + 4q) z - 4p^2 q^2 z^2}{D_3}, \quad (4.32)$$

with

$$D_3 = 1 - p (p^2 + 4q) z - 2q (p^4 + 6p^2 q + 8q^2) z^2 + 8pq^3 (p^2 + 4q) z^3 + 64q^6 z^4.$$

And we deduce the following corollaries and theorems.

Corollary 7. For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Jacobsthal numbers $(J_{p,q,n} J_{p,q,2n})$ is given by:

$$\begin{aligned} \sum_{n=0}^{\infty} J_{p,q,n} J_{p,q,2n} z^n &= \frac{pz + 8pq^3 z^3}{D_3} \\ &= \frac{pz + 8pq^3 z^3}{1 - p (p^2 + 4q) z - 2q (p^4 + 6p^2 q + 8q^2) z^2 + 8pq^3 (p^2 + 4q) z^3 + 64q^6 z^4}, \end{aligned} \quad (4.33)$$

with $J_{p,q,n} J_{p,q,2n} = S_{n-1} (a_1 + [-a_2]) S_{2n-1} (e_1 + [-e_2])$.

Corollary 8. Let n be a natural number. Then we have the new generating function of the product of (p, q) -Jacobsthal numbers $(J_{p,q,n} J_{p,q,2n+1})$:

$$\begin{aligned} \sum_{n=0}^{\infty} J_{p,q,n} J_{p,q,2n+1} z^n &= \frac{(p^2 + 2q) z - 4pq^2 z^2 - 16q^4 z^3}{D_3} \\ &= \frac{(p^2 + 2q) z - 4pq^2 z^2 - 16q^4 z^3}{1 - p (p^2 + 4q) z - 2q (p^4 + 6p^2 q + 8q^2) z^2 + 8pq^3 (p^2 + 4q) z^3 + 64q^6 z^4}. \end{aligned} \quad (4.34)$$

with $J_{p,q,n} J_{p,q,2n+1} = S_{n-1} (a_1 + [-a_2]) S_{2n} (e_1 + [-e_2])$.

Theorem 16. For $n \in \mathbb{N}$, the new generating function of the product of (p, q) -Jacobsthal Lucas numbers $(j_{p,q,n} j_{p,q,2n})$ is given by:

$$\sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n} z^n = \frac{4 - 3p (p^2 + 4q) z - 4q (p^4 + 6p^2 q + 8q^2) z^2 + 8pq^3 (p^2 + 4q) z^3}{1 - p (p^2 + 4q) z - 2q (p^4 + 6p^2 q + 8q^2) z^2 + 8pq^3 (p^2 + 4q) z^3 + 64q^6 z^4}. \quad (4.35)$$

Proof. We have

$$\begin{aligned}
 \sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n} z^n &= \sum_{n=0}^{\infty} \left(\begin{array}{c} (2S_n(a_1 + [-a_2]) - pS_{n-1}(a_1 + [-a_2])) \\ \times (2S_{2n}(e_1 + [-e_2]) - pS_{2n-1}(e_1 + [-e_2])) \end{array} \right) z^n \\
 &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n \\
 &\quad - 2p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n-1}(e_1 + [-e_2]) z^n \\
 &\quad - 2p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n \\
 &\quad + p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n-1}(e_1 + [-e_2]) z^n.
 \end{aligned}$$

Using the relationships (4.27), (4.28), (4.30) and (4.31), we obtain

$$\begin{aligned}
 \sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n} z^n &= \frac{4(1 - 2pqz - 4q^2(p^2 + 2q)z^2)}{D_3} - \frac{2p(p^2z + 2pq(p^2 + 4q)z^2)}{D_3} \\
 &\quad - \frac{2p((p^2 + 2q)z - 4pq^2z^2 - 16q^4z^3)}{D_3} + \frac{p^2(pz + 8pq^3z^3)}{D_3} \\
 &= \frac{4 - 3p(p^2 + 4q)z - 4q(p^4 + 6p^2q + 8q^2)z^2 + 8pq^3(p^2 + 4q)z^3}{D_3} \\
 &= \frac{4 - 3p(p^2 + 4q)z - 4q(p^4 + 6p^2q + 8q^2)z^2 + 8pq^3(p^2 + 4q)z^3}{1 - p(p^2 + 4q)z - 2q(p^4 + 6p^2q + 8q^2)z^2 + 8pq^3(p^2 + 4q)z^3 + 64q^6z^4}.
 \end{aligned}$$

So, the proof is completed. \square

Theorem 17. Let n be a natural number. Then we have the new generating function of the product of (p, q) -Jacobsthal Lucas numbers $(j_{p,q,n} j_{p,q,2n+1})$:

$$\sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n+1} z^n = \frac{2p - p^2(p^2 + 2q)z + 12pq^2(p^2 + 4q)z^2 - 16p^2q^4z^3}{1 - p(p^2 + 4q)z - 2q(p^4 + 6p^2q + 8q^2)z^2 + 8pq^3(p^2 + 4q)z^3 + 64q^6z^4}. \quad (4.36)$$

Proof. We have

$$\begin{aligned}
 \sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n+1} z^n &= \sum_{n=0}^{\infty} \left(\begin{array}{c} (2S_n(a_1 + [-a_2]) - pS_{n-1}(a_1 + [-a_2])) \\ \times (2S_{2n+1}(e_1 + [-e_2]) - pS_{2n}(e_1 + [-e_2])) \end{array} \right) z^n \\
 &= 4 \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n+1}(e_1 + [-e_2]) z^n \\
 &\quad - 2p \sum_{n=0}^{\infty} S_n(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n \\
 &\quad - 2p \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n+1}(e_1 + [-e_2]) z^n \\
 &\quad + p^2 \sum_{n=0}^{\infty} S_{n-1}(a_1 + [-a_2]) S_{2n}(e_1 + [-e_2]) z^n.
 \end{aligned}$$

Using the relationships (4.28), (4.29), (4.31) and (4.32), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} j_{p,q,n} j_{p,q,2n+1} z^n &= \frac{4(p+8pq^3z^2)}{D_3} - \frac{2p(1-2pqz-4q^2(p^2+2q)z^2)}{D_3} \\ &\quad - \frac{2p(p(p^2+4q)z-4p^2q^2z^2)}{D_3} + \frac{p^2((p^2+2q)z-4pq^2z^2-16q^4z^3)}{D_3} \\ &= \frac{2p-p^2(p^2+2q)z+4pq^2(p^2+4q)z^2-16p^2q^4z^3}{D_3} \\ &= \frac{2p-p^2(p^2+2q)z+12pq^2(p^2+4q)z^2-16p^2q^4z^3}{1-p(p^2+4q)z-2q(p^4+6p^2q+8q^2)z^2+8pq^3(p^2+4q)z^3+64q^6z^4}. \end{aligned}$$

So, the proof is completed. \square

Corollary 9. *Setting $p = k$ and $q = 1$ in Eqs. (4.33)-(4.36) gives the following new generating functions:*

$$\begin{aligned} \sum_{n=0}^{\infty} J_{k,n} J_{k,2n} z^n &= \frac{kz+8kz^3}{1-k(k^2+4)z-2(k^4+6k^2+8)z^2+8k(k^2+4)z^3+64z^4}. \\ \sum_{n=0}^{\infty} J_{k,n} J_{k,2n+1} z^n &= \frac{(k^2+2)z-4kz^2-16z^3}{1-k(k^2+4)z-2(k^4+6k^2+8)z^2+8k(k^2+4)z^3+64z^4}. \\ \sum_{n=0}^{\infty} j_{k,n} j_{k,2n} z^n &= \frac{4-3k(k^2+4)z-4(k^4+6k^2+8)z^2+8k(k^2+4)z^3}{1-k(k^2+4)z-2(k^4+6k^2+8)z^2+8k(k^2+4)z^3+64z^4}. \\ \sum_{n=0}^{\infty} j_{k,n} j_{k,2n+1} z^n &= \frac{2k-k^2(k^2+2)z+12k(k^2+4)z^2-16k^2z^3}{1-k(k^2+4)z-2(k^4+6k^2+8)z^2+8k(k^2+4)z^3+64z^4}. \end{aligned}$$

- Put $k = 1$ in the Corollary 9, we obtain the following table:

Coefficient of z^n	Generating function
$J_n J_{2n}$	$\frac{z+8z^3}{1-5z-30z^2+40z^3+64z^4}$
$J_n J_{2n+1}$	$\frac{3z-4z^2-16z^3}{1-5z-30z^2+40z^3+64z^4}$
$j_n j_{2n}$	$\frac{4-15z-60z^2+40z^3}{1-5z-30z^2+40z^3+64z^4}$
$j_n j_{2n+1}$	$\frac{2-3z+60z^2-16z^3}{1-5z-30z^2+40z^3+64z^4}$

Table 7. A new generating functions of the products of some numbers.

5. Conclusion

This study proposes to present new class of generating functions for some special numbers with parameters p and q by using the symmetric functions. We can summarize the sections as follows:

- **In Section 1**, we presented some backgrounds about (p, q) -numbers and some preliminary facts and results on the symmetric functions.
- **In Section 2**, we derived and proved 5 theorems (1-5) by making use of the symmetrizing operator given by Definition 5.
- **In Section 3**, By making use the theorems given in Section 2, we have derived some new generating functions of odd and even terms of (p, q) -numbers.
- **In Section 4**, By making use the symmetric functions, we investigated the new generating functions of the products of (p, q) -numbers with odd and even (p, q) -numbers.

References

- [1] A. Abderrezzak, Généralisation de la transformation d'Euler d'une série formelle. Adv. Math. 103, 180-195, 1994.
- [2] A. Boussayoud, M. Kerada, S. Araci, M. Acikgoz, Symmetric functions of the k -Fibonacci and k -Lucas numbers, Asian-European Journal of Mathematics, 14, 1-12, 2021.

- [3] A. Boussayoud, A. Abderrezzak, Complete homogeneous symmetric functions and Hadamard product, *Ars Comb.*, 144, 81-90, 2019.
- [4] A. Boussayoud, M. Kerada, N. Harrouche, On the k -Lucas numbers and Lucas polynomials, *Turkish Journal of Analysis and Number*, 5, 121-125, 2017.
- [5] A. Boussayoud, M. Kerada, Symmetric and generating functions, *Int. Electron. J. Pure Appl. Math.*, 7, 195-203, 2014.
- [6] A. Suvarnamani, M. Tatong, Some properties of the product of (p,q) -Fibonacci and (p,q) -Lucas numbers, *International Journal of GEOMATE*, 13, 16-19, 2017.
- [7] A. Suvarnamani, Some properties of (p,q) -Lucas numbers, *Kyungpook Math. J.*, 56, 367-370, 2016.
- [8] A. Suvarnamani, M. Tatong, Some properties of (p,q) -Fibonacci numbers, *Science and Technology RMUTT Journal*, 5, 17-21, 2015.
- [9] B. G. S. Doman, J. K. Williams, Fibonacci and Lucas polynomials. *Math. Proc. Camb. Philos. Soc.*, 90, 385-387, 1981.
- [10] H. H. Gulec, N. Taskara, On the (s,t) -Pell and (s,t) -Pell Lucas sequences and their matrix representations, *Appl. Math. Lett.*, 25, 1554-1559, 2012.
- [11] I. G. Macdonald, *Symmetric functions and Hall polynomials*. Oxford University Press, Oxford (1979).
- [12] I. Mező, Several generating functions for second-order recurrence sequences, *J. Integer Seq.* 12, 1-16, 2009.
- [13] M. Merca, A generalization of the symmetry between complete and elementary symmetric functions, *Indian J. Pure Appl. Math.*, 75-89, 2014.
- [14] N. Saba, A. Boussayoud, K. V. V. Kanuri, Mersenne Lucas numbers and complete homogeneous symmetric functions, *J. Math. Computer Sci.*, 24, 127-139, 2022.
- [15] N. Saba, A. Boussayoud, A. Abderrezzak, Symmetric and generating functions of generalized (p,q) -numbers, *Kuwait J. Sci.*, 48(4), 2021 (In Press).
- [16] N. Saba, A. Boussayoud, S. Araci, M. Kerada, M. Acikgoz, Construction of a new class of symmetric function of binary products of (p,q) -numbers with 2-orthogonal Chebyshev polynomials, *Bol. Soc. Math. Mex.*, 27, 1-26, 2021.
- [17] N. Saba, A. Boussayoud, A. Abderrezzak, Complete homogeneous symmetric functions of third and second-order linear recurrence sequences, *Electron. J. Math. Analysis Appl.*, 9, 221-245, 2021.
- [18] N. Saba, A. Boussayoud, M. Chelgham, Symmetric and generating functions for some generalized polynomials, *Malaya J. Mat.*, 8, 1756-1765, 2020.
- [19] N. Saba, A. Boussayoud, Complete homogeneous symmetric functions of Gauss Fibonacci polynomials and bivariate Pell polynomials, *Open J. Math. Sci.*, 4, 179-185, 2020.
- [20] N. Saba, A. Boussayoud, Ordinary generating functions of binary products of (p,q) -modified Pell numbers and k -numbers at positive and negative indices, *J. Sci. Arts.*, 3, 627-648, 2020.
- [21] N. Saba, A. Boussayoud, M. Kerada, Generating functions of even and odd Gaussian numbers and polynomials, *J. Sci. Arts.*, 1, 125-144, 2021.
- [22] S. Uygun, The (s,t) -Jacobsthal and (s,t) -Jacobsthal Lucas sequences, *Appl. Math. Sci.*, 9, 3467-3476, 2015.