

Boundary smoothness conditions for functions in $R^p(X)$

Stephen Deterding ^{*}
 Department of Mathematics & Physics
 Marshall University, Huntington, WV, USA

Abstract

Let X be a compact subset of the complex plane and let $R^p(X)$, $2 < p < \infty$, denote the closure of the rational functions with poles off X in the L^p norm. In this paper we consider three conditions that show how the functions in $R^p(X)$ can have a greater degree of smoothness at the boundary of X than might otherwise be expected. We will show that two of the conditions are equivalent and imply the third but the third does not imply the other two.

1 Introduction

Let X be a compact subset of the complex plane and let $R_0(X)$ denote the set of all rational functions with poles off X . In this paper we consider the function space $R^p(X)$, $2 < p < \infty$, the closure of $R_0(X)$ in the L^p norm, and the smoothness of the functions in this space. Many results in this area have been determined for $R(X)$, the closure of $R_0(X)$ in the uniform norm, but for $R^p(X)$ less is known. In particular, Wang proposed three conditions that demonstrate how the functions in $R(X)$ can have a greater degree of smoothness at the boundary of X than might otherwise be expected. [7] To describe these conditions, we first need to state a few definitions.

We say that $\phi(r)$ is an admissible function if it is positive and nondecreasing on $(0, \infty)$ and if the associated function $\psi(r) = \frac{r}{\phi(r)}$ is also positive and nondecreasing on $(0, \infty)$ with $\psi(0^+) = 0$. Examples of admissible functions are the power functions $\phi(r) = r^\alpha$ where $0 < \alpha < 1$ and the function $\phi(r) = \frac{r}{\log(1+r)}$. Let $\gamma(X)$ denote the analytic capacity of the set X [3, pg. 196] and let $R_x^t f(z)$ be the error at z of the t -th degree Taylor polynomial of f about x , which is defined by

^{*}Email: deterding@marshall.edu

$$R_x^t f(z) = f(z) - \sum_{j=0}^t \frac{f^{(j)}(x)}{j!} (z-x)^j.$$

Let $A_n(x)$ denote the annulus $\{z : 2^{-(n+1)} < |z-x| < 2^{-n}\}$ and let $B(x, r)$ denote the disk centered at x with radius r . Let m denote area measure. A set E is said to have full area density at x if $\frac{m(B(x, r) \setminus E)}{m(B(x, r))} \rightarrow 0$ as $r \rightarrow 0$.

Let X be a compact subset of the plane, let $x \in X$, let t be a non-negative integer, let $\phi(r)$ be an admissible function, and let $\|\cdot\|_X$ denote the uniform norm on X . Wang's three conditions are the following.

(A) For each $\epsilon > 0$ the set

$$\{y \in X : |R_x^t f(y)| \leq \epsilon \phi(|y-x|) |y-x|^t \|f\|_X \\ \text{for all } f \in R_0(X)\}$$

has full area density at x .

(B) There exists a representing measure μ for x on $R(X)$ such that $\mu(x) = 0$ and

$$\int \frac{d|\mu|(z)}{|z-x|^t \phi(|z-x|)} < \infty.$$

(C) The series

$$\sum_{n=1}^{\infty} \frac{2^{n(t+1)} \gamma(A_n(x) \setminus X)}{\phi(2^{-n})}$$

converges.

The relationship between these conditions has been studied by Wang and O'Farrell. In various papers [5-7] they showed that (B) and (C) are equivalent and imply (A), but (A) does not imply (B) or (C).

We now consider Wang's conditions applied to the space $R^p(X)$. Before we state what these conditions look like in the context of $R^p(X)$, we introduce q -capacity, which is the appropriate capacity for $R^p(X)$. For $1 < q < 2$, the q -capacity of a compact set X in the complex plane is denoted $\Gamma_q(X)$ and is defined by

$$\Gamma_q(X) = \inf \int |\nabla u|^q dm,$$

where the infimum is taken over all infinitely differentiable functions u of compact support with $u \equiv 1$ on X .

Let X be a compact subset of the plane, let $x \in X$, let t be a non-negative integer, let $2 < p < \infty$ and $q = \frac{p}{p-1}$, and let $\phi(r)$ be an admissible function. Wang's three conditions restated for $R^p(X)$ are

(A') For each $\epsilon > 0$ the set

$$\{y \in X : |R_x^t f(y)| \leq \epsilon \phi(|y-x|) |y-x|^t \|f\|_{L^p(X)} \\ \text{for all } f \in R_0(X)\}$$

has full area density at x .

(B') There exists $g \in L^q(X)$ which represents x on $R^p(X)$ such that

$$\int \frac{|g(z)|^q dm}{|z-x|^{qt} \phi(|z-x|)^q} < \infty.$$

(C') The series

$$\sum_{n=1}^{\infty} \frac{2^{n(t+1)q} \Gamma_q(A_n(x) \setminus X)}{\phi(2^{-n})^q}$$

converges.

In analogy with $R(X)$, we expect that (B') and (C') are equivalent, and imply (A'), but (A') does not imply (B') or (C'). What is known about the relationship of these conditions is the following. Wolf has shown that (B') implies (A') [8, Part 1 Theorem 4.1] and (C') [8, Part 2 Theorem 1.1] but nothing else is known. We will show that (C') implies (B') but (A') does not imply (C'), and thus (A') does not imply (B'). This shows that Wang's conditions restated for $R^p(X)$ have the same relations amongst themselves as Wang's conditions for $R(X)$.

2 The equivalence of (B') and (C')

We will first show that (C') implies (B'). In particular, we will prove this for arbitrary non-decreasing ϕ which allows us to consider the case $t = 0$ without loss of generality.

Theorem 1. *Let $\phi(r)$ be a positive non-decreasing function and let X be a compact subset of the plane and let $x \in X$. Let $2 < p < \infty$ and $q = \frac{p}{p-1}$, and suppose*

$$\sum_{n=1}^{\infty} 2^{nq} \phi(2^{-n})^{-q} \Gamma_q(A_n(x) \setminus X) < \infty.$$

Then there exists $g \in L^q(X)$ which represents x on $R^p(X)$ such that

$$\int \frac{|g(z)|^q dm}{\phi(|z-x|)^q} < \infty.$$

Proof. We may assume that X is a subset of the unit disk and $x = 0$ without loss of generality. We abbreviate $A_n(x)$ to A_n .

Let T_n be a linear functional defined by $T_n(f) = \int_{\partial A_n} \frac{f(z)}{z} dz$ and let f be a rational function with poles off X . We can assume that f is modified off X so that it is continuous, but still analytic in a neighborhood of X . Then there exist closed sets $K_n \subseteq A_n \setminus X$ with smooth boundaries such that f is analytic outside of $\cup K_n$ and

$$\int_{A_n \setminus K_n} |f|^p dm \leq 2 \int_{X \cap A_n} |f|^p dm.$$

We now bound $T_n(f)$. In [4], Hedberg constructed smooth functions P_m such that each $P_m = 1$ on K_m , has support on $A_{m-1} \cup A_m \cup A_{m+1}$, and $\int |\nabla P_m|^q dm \leq C(\Gamma_q(A_m \setminus X) + 4^{-m})$. (Note that in [4] these functions are called ϕ_m .) By modifying this construction, we can make it so that $\int |\nabla P_m|^q dm \leq C(\Gamma_q(A_m \setminus X) + 4^{-m} \phi(2^{-m})^q)$. Let $P = \sup_m P_m$ so that $P(z) = 1$ on $\cup K_m$. Then it follows from Green's Theorem and the analyticity of f that

$$\begin{aligned} \sum_{m=n}^{\infty} \int_{\partial A_m} \frac{f(z)}{z} dz &= \sum_{m=n}^{\infty} \int_{\partial K_m} \frac{f(z)}{z} dz \\ &= \sum_{m=n}^{\infty} \int_{\partial K_m} \frac{f(z)P(z)}{z} dz \\ &= \sum_{m=n}^{\infty} \int_{A_m \setminus K_m} \frac{f(z)}{z} \frac{\partial P(z)}{\partial \bar{z}} dm, \end{aligned}$$

and thus

$$T_n(f) = \sum_{m=n}^{\infty} \int_{\partial A_m} \frac{f(z)}{z} dz - \sum_{m=n+1}^{\infty} \int_{\partial A_m} \frac{f(z)}{z} dz = \int_{A_n \setminus K_n} \frac{f(z)}{z} \frac{\partial P(z)}{\partial \bar{z}} dm.$$

Hence it follows from Holder's inequality that

$$\begin{aligned} |T_n(f)| &\leq 2^{n+1} \left(\int_{A_n \setminus K_n} |f(z)|^p dm \right)^{\frac{1}{p}} \left(\int_{A_n \setminus K_n} |\nabla P|^q dm \right)^{\frac{1}{q}} \\ &\leq C 2^n \|f\|_{L^p(X \cap A_n)} [\Gamma_q(A_{n-1} \setminus X) + \Gamma_q(A_n \setminus X) + \Gamma_q(A_{n+1} \setminus X) + 4^{-n} \phi(2^{-n})^q]^{\frac{1}{q}}. \end{aligned}$$

Thus $T_n(f)$ is a bounded linear functional on $R^p(X \cap A_n)$. Thus it follows from the Hahn-Banach theorem that there exists $g_n \in L^q(X)$ with support on A_n such that $T_n(f) = \int f g_n dm$ for all f in $R^p(X \cap A_n)$ and

$$\int |g_n|^q dm \leq C 2^{nq} [\Gamma_q(A_{n-1} \setminus X) + \Gamma_q(A_n \setminus X) + \Gamma_q(A_{n+1} \setminus X) + 4^{-n} \phi(2^{-n})^q].$$

Now let $g = \frac{1}{2\pi i} \sum_{n=1}^{\infty} g_n$. Then by the Cauchy integral formula,

$$\begin{aligned} f(0) &= \frac{1}{2\pi i} \sum_{n=1}^{\infty} \int_{\partial A_n} \frac{f(z)}{z} dz \\ &= \frac{1}{2\pi i} \sum_{n=1}^{\infty} \int f(z) g_n(z) dm \\ &= \int f(z) \frac{1}{2\pi i} \sum_{n=1}^{\infty} g_n(z) dm \\ &= \int f g dm. \end{aligned}$$

Thus g represents 0 on $R^p(X)$. Finally, because g_n has support on A_n ,

$$\begin{aligned}
\int \frac{|g(z)|^q}{\phi(|z|)^q} dm &= \left(\frac{1}{2\pi}\right)^q \int \phi(|z|)^{-q} \left| \sum_{n=1}^{\infty} g_n(z) \right|^q dm \\
&= \left(\frac{1}{2\pi}\right)^q \sum_{m=1}^{\infty} \int_{A_m} \phi(|z|)^{-q} \left| \sum_{n=1}^{\infty} g_n(z) \right|^q dm \\
&\leq \sum_{m=1}^{\infty} \int_{A_m} \phi(|z|)^{-q} |g_m(z)|^q dm \\
&\leq C \sum_{m=1}^{\infty} \phi(2^{-m})^{-q} \int_{A_m} |g_m(z)|^q dm.
\end{aligned}$$

Hence

$$\begin{aligned}
\int \frac{|g(z)|^q}{|\phi(z)|^q} dm &\leq C \sum_{m=1}^{\infty} \phi(2^{-m})^{-q} 2^{mq} [\Gamma_q(A_{m-1} \setminus X) + \Gamma_q(A_m \setminus X) + \Gamma_q(A_{m+1} \setminus X) \\
&\quad + 4^{-m} \phi(2^{-m})^q] < \infty.
\end{aligned}$$

□

3 (A') does not imply (C')

Finally, we show that (A') does not imply (C'). In particular we will show the following.

Theorem 2. *Let $\phi(r)$ be an admissible function with associated function $\psi(r) = \frac{r}{\phi(r)}$ such that $\phi(0^+) = 0$, and let $2 < p < \infty$ and $q = \frac{p}{p-1}$. Then there is a compact set X containing 0 such that for each $\epsilon > 0$ the set $\{y \in X : |f(y) - f(0)| \leq \epsilon \phi(|y|) \|f\|_{L^p(X)}\}$ has full area density at 0, but*

$$\sum_{n=1}^{\infty} \phi(2^{-n})^{-q} 2^{nq} \Gamma_q(A_n(0) \setminus X) = \infty.$$

Due to the length of the proof, it will be split up into four lemmas. We first describe the construction of the set X . Again we abbreviate $A_n(0)$ to A_n . Let $a_n = \frac{3}{4} \cdot 2^{-n}$. We can choose a subsequence still denoted a_n such that $\phi(a_n) < \frac{1}{2} \phi(a_{n-1})$ and

$$\sum_{n=1}^{j-1} \psi(a_n)^{-q} < \psi(a_j)^{-q}.$$

Let $r_n = [n^{-1}a_n^q\phi(a_n)^q]^{\frac{1}{2-q}}$. Then there exists $M > 0$ such that for $n \geq M$, $\frac{r_n}{a_n} < \frac{1}{5}n^{-\frac{1}{2q}}$. For such n let D_n denote the open disk centered at a_n with radius r_n ; otherwise, let D_n be the empty set. Let Δ denote the closed unit disk and let $X = \Delta \setminus \cup_n D_n$.

Lemma 1. *Let X be the set constructed in the previous paragraph. Then*

$$\sum_{n=1}^{\infty} \phi(2^{-n})^{-q} 2^{nq} \Gamma_q(A_n \setminus X) = \infty.$$

Proof. Since the q -capacity of a disk of radius r is r^{2-q} [1],

$$\begin{aligned} \sum_{n=1}^{\infty} \phi(2^{-n})^{-q} 2^{nq} \Gamma_q(A_n \setminus X) &= \sum_{n=M}^{\infty} \phi(2^{-n})^{-q} 2^{nq} \Gamma_q(A_n \setminus X) \\ &= \sum_{n=M}^{\infty} \phi(2^{-n})^{-q} 2^{nq} n^{-1} a_n^q \phi(a_n)^q \\ &= \sum_{n=M}^{\infty} \phi(2^{-n})^{-q} 2^{nq} n^{-1} \left(\frac{3}{4}\right)^q 2^{-nq} \phi\left(\frac{3}{4} \cdot 2^{-n}\right)^q \\ &= \sum_{n=M}^{\infty} \phi(2^{-n})^{-q} n^{-1} \left(\frac{3}{4}\right)^q \phi\left(\frac{3}{4} \cdot 2^{-n}\right)^q. \end{aligned}$$

Then since $\phi(r) = \frac{r}{\psi(r)}$ and $\psi(r)$ is non-decreasing

$$\begin{aligned} \sum_{n=M}^{\infty} \phi(2^{-n})^{-q} n^{-1} \left(\frac{3}{4}\right)^q \phi\left(\frac{3}{4} \cdot 2^{-n}\right)^q &= \sum_{n=M}^{\infty} \left(\frac{2^{-n}}{\psi(2^{-n})}\right)^{-q} n^{-1} \left(\frac{3}{4}\right)^q \left(\frac{\frac{3}{4} \cdot 2^{-n}}{\psi(\frac{3}{4} \cdot 2^{-n})}\right)^q \\ &= \sum_{n=M}^{\infty} \left(\frac{3}{4}\right)^{2q} \psi(2^{-n})^q n^{-1} \psi\left(\frac{3}{4} \cdot 2^{-n}\right)^{-q} \\ &\geq \left(\frac{3}{4}\right)^{2q} \sum_{n=M}^{\infty} n^{-1} = \infty. \end{aligned}$$

Hence

$$\sum_{n=1}^{\infty} \phi(2^{-n})^{-q} 2^{nq} \Gamma_q(A_n \setminus X) = \infty.$$

□

Lemma 2. *Let $d_n(z)$ denote the distance from z to D_n and let $s_n = \frac{1}{5}n^{-\frac{1}{2q}}a_n$. Let $E = \cup_{n>M} \{z \in A_n : d_n(z) \geq r_n\}$. (See Figure 1). Then E has full area density at 0.*




Figure 1: The sets D_n , $E \cap A_n$, and F_n

Proof. Since $\frac{r_n}{a_n} < \frac{1}{5}n^{-\frac{1}{2q}}$, it follows that $r_n < s_n$. Let $B(a, r)$ denote the disk centered at a with radius r . Then it follows from the definition of E that $B(0, 2^{-j}) \setminus E$ is the union of the disks $B(a_n, r_n + s_n)$, where the union is taken over $n \geq j$. Thus

$$\frac{m(B(0, 2^{-j}) \setminus E)}{m(B(0, 2^{-j}))} = \frac{\pi \sum_{n=j}^{\infty} (r_n + s_n)^2}{2^{-2j}} = \frac{\sum_{n=j}^{\infty} (r_n + s_n)^2}{\frac{16}{9}a_j^2}.$$

However, since $r_{n+1} < \frac{1}{2}r_n$ and $s_{n+1} < \frac{1}{2}s_n$,

$$\begin{aligned} \frac{m(B(0, 2^{-j}) \setminus E)}{m(B(0, 2^{-j}))} &\leq \frac{(\sum_{n=0}^{\infty} 2^{-2n}) (r_j + s_j)^2}{\frac{16}{9}a_j^2} \\ &< \frac{3}{4} \left(\frac{2s_j}{a_j} \right)^2 \\ &= \frac{3}{25} j^{-\frac{1}{q}}. \end{aligned}$$

Thus $\frac{m(B(0, 2^{-j}) \setminus E)}{m(B(0, 2^{-j}))} \rightarrow 0$ as $j \rightarrow \infty$ and E has full area density at 0. □

Lemma 3. *There exist smooth functions P_n such that $P_n(z) = 1$ on D_n , $\int |\nabla P_n|^q dm \leq C(\Gamma_q(D_n) + 16^{-n})$ and P_n is supported on the set $F_n = \{z \in A_n : d_n(z) \leq \frac{s_n}{2}\}$.*

Proof. The proof is a modification of a construction of Hedberg [4, pg. 277]. It follows from the definition of q -capacity that there exists smooth functions ω_n such that $\omega_n = 1$ on D_n and $\int |\nabla \omega_n|^q dm \leq \Gamma_q(D_n) + 16^{-n}$. Let $\xi_n(z)$ be a piecewise linear continuous function of $|z|$ such that $\xi_n = 1$ on D_n , $\xi_n = 0$ outside F_n and $|\nabla \xi_n(z)| \leq 2s_n^{-1}$, and let $P_n(z) = \xi_n(z)\omega_n(z)$. Then

$$\int |\nabla P_n(z)|^q dm \leq \left(\int \xi_n^q |\nabla \omega_n|^q dm + \int \omega_n^q |\nabla \xi_n|^q dm \right).$$

The first integral is bounded above by $\Gamma_q(D_n) + 16^{-n}$. To bound the second integral we first note that it follows from the Gagliardo-Nirenberg-Sobolev inequality (See [2, pg.277]) that $\left(\int \omega_n^{\frac{2q}{2-q}} dm \right)^{\frac{2-q}{2}} \leq C \int |\nabla \omega_n|^q dm$. Thus by Holder's inequality

$$\begin{aligned} \int \omega_n^q |\nabla \xi_n|^q dm &\leq C \left(\int \omega_n^{\frac{2q}{2-q}} dm \right)^{\frac{2-q}{2}} \left(\int_{F_n} |\nabla \xi_n|^2 \right)^{\frac{q}{2}} \\ &\leq C \int |\nabla \omega_n|^q dm \cdot m(F_n)^{\frac{q}{2}} s_n^{-q} \\ &\leq C(\Gamma_q(D_n) + 16^{-n}) \cdot \left(r_n + \frac{s_n}{2} \right)^q s_n^{-q} \\ &\leq C(\Gamma_q(D_n) + 16^{-n}) \cdot \left(s_n + \frac{s_n}{2} \right)^q s_n^{-q} \\ &= C(\Gamma_q(D_n) + 16^{-n}) \cdot \left(\frac{3}{2} \right)^q. \end{aligned}$$

Thus $\int |\nabla P_n|^q dm \leq C(\Gamma_q(D_n) + 16^{-n})$. □

Lemma 4. *Let $\epsilon > 0$. Then for N sufficiently large,*

$$|f(y) - f(0)| \leq \epsilon \phi(|y|) \|f\|_p$$

whenever $y \in A_N \cap X^\circ \cap E$ and $f \in R_0(X)$.

Proof. Let $f \in R_0(X)$ and $y \in A_N \cap X^\circ \cap E$. Then it follows from the Cauchy integral formula that

$$\begin{aligned}
|f(y) - f(0)| &= \left| \frac{1}{2\pi i} \sum_{n=1}^{\infty} \int_{\partial D_n} \frac{f(z)dz}{z-y} - \frac{1}{2\pi i} \sum_{n=1}^{\infty} \int_{\partial D_n} \frac{f(z)dz}{z} \right| \\
&= \left| \frac{1}{2\pi i} \sum_{n=1}^{\infty} \int_{\partial D_n} \frac{yf(z)dz}{z(z-y)} \right| \\
&\leq |y| \left| \sum_{n=1}^{\infty} \int_{\partial D_n} \frac{f(z)dz}{z(z-y)} \right|.
\end{aligned}$$

Let P_n be the smooth functions constructed in Lemma 3 and let $P = \sup_n P_n$. Then $P(z) = 1$ on $\cup D_n$ and hence by Green's theorem,

$$\begin{aligned}
|f(y) - f(0)| &\leq |y| \left| \sum_{n=1}^{\infty} \int_{\partial D_n} \frac{f(z)P(z)dz}{z(z-y)} \right| \\
&= |y| \left| \sum_{n=1}^{\infty} \int_{F_n \setminus D_n} \frac{f(z) \frac{\partial P}{\partial \bar{z}} dm}{z(z-y)} \right| \\
&\leq |y| \sum_{n=1}^{\infty} \int_{F_n \setminus D_n} \left(a_n - r_n - \frac{s_n}{2} \right)^{-1} \delta_n(y)^{-1} |f(z)| (|\nabla P(z)|) dm.
\end{aligned}$$

where $\delta_n(y)$ is the distance from y to F_n . Since $\frac{r_n}{a_n} < \frac{1}{5} n^{-\frac{1}{2q}}$,

$$\left(a_n - r_n - \frac{s_n}{2} \right) \geq \left(1 - \frac{1}{5} n^{-\frac{1}{2q}} - \frac{1}{10} n^{-\frac{1}{2q}} \right) a_n \geq \frac{7}{10} a_n,$$

and hence

$$|f(y) - f(0)| \leq C|y| \sum_{n=1}^{\infty} \int_{F_n \setminus D_n} a_n^{-1} \delta_n(y)^{-1} |f(z)| (|\nabla P(z)|) dm.$$

Next, it follows from Holder's inequality that

$$\begin{aligned}
|f(y) - f(0)| &\leq C|y| \left(\sum_{n=1}^{\infty} \int_{F_n \setminus D_n} |f(z)|^p dm \right)^{\frac{1}{p}} \left(\sum_{n=1}^{\infty} a_n^{-q} \delta_n(y)^{-q} \int_{F_n \setminus D_n} |\nabla P(z)|^q dm \right)^{\frac{1}{q}} \\
&\leq C|y| \cdot \|f\|_p \left(\sum_{n=1}^{\infty} a_n^{-q} \delta_n(y)^{-q} (\Gamma_q(D_n) + 16^{-n}) \right)^{\frac{1}{q}} \\
&\leq C|y| \cdot \|f\|_p \left(\sum_{n=1}^{\infty} a_n^{-q} \delta_n(y)^{-q} (r_n^{(2-q)} + 16^{-n}) \right)^{\frac{1}{q}}.
\end{aligned}$$

We next obtain bounds for $\delta_n(y)$. Recall that $\frac{r_n}{a_n} < \frac{1}{5}n^{\frac{-1}{2q}}$ and $y \in A_N$. If $n < N$, then

$$\delta_n(y) \geq \left(a_n - r_n - \frac{s_n}{2}\right)2^{-N} \geq \frac{7}{10}a_n 2^{-N} = \frac{21}{40}2^{-n}2^{-N} \geq \frac{1}{40}2^{-n} = \frac{1}{30}a_n.$$

If $n > N$, then

$$\begin{aligned} \delta_n(y) &\geq 2^{-(N+1)} - \left(a_n + r_n + \frac{s_n}{2}\right) \geq 2^{-(N+1)} - \left(1 + \frac{3}{10}n^{\frac{-1}{2q}}\right)a_n \\ &\geq 2^{-(N+1)} - \frac{39}{40}2^{-n} \geq \frac{1}{40}2^{-(N+1)} = \frac{1}{60}a_N. \end{aligned}$$

Lastly, if $n = N$, then $\delta_N(y) \geq \frac{s_N}{2}$. Now let $t_n = r_n^{(2-q)} + 16^{-n}$. Then

$$\begin{aligned} |f(y) - f(0)| &\leq C|y| \cdot \|f\|_p \left(\sum_{n=1}^{\infty} a_n^{-q} \delta_n(y)^{-q} t_n \right)^{\frac{1}{q}} \\ &\leq C|y| \cdot \|f\|_p \left(\sum_{n=1}^{N-1} a_n^{-q} \delta_n(y)^{-q} t_n + a_N^{-q} \delta_N(y)^{-q} t_N + \sum_{n=N+1}^{\infty} a_n^{-q} \delta_n(y)^{-q} t_n \right)^{\frac{1}{q}} \\ &\leq C|y| \cdot \|f\|_p \left(\sum_{n=1}^{N-1} a_n^{-2q} t_n + a_N^{-q} s_N^{-q} t_N + \sum_{n=N+1}^{\infty} a_n^{-q} a_N^{-q} t_n \right)^{\frac{1}{q}} \\ &\leq C\phi(|y|) \cdot \|f\|_p \left(\sum_{n=1}^{N-1} a_n^{-2q} t_n \psi(a_N)^q + \phi(a_N)^{-q} s_N^{-q} t_N \right. \\ &\quad \left. + \sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} t_n \right)^{\frac{1}{q}}. \end{aligned}$$

The last line follows because $|y| \leq \frac{4}{3}\phi(|y|)\psi(a_N)$. To complete the proof, we must bound the 3 terms inside the parentheses. We begin by bounding the leftmost term

$$\sum_{n=1}^{N-1} a_n^{-2q} t_n \psi(a_N)^q = \sum_{n=1}^{N-1} a_n^{-2q} r_n^{(2-q)} \psi(a_N)^q + \sum_{n=1}^{N-1} a_n^{-2q} 16^{-n} \psi(a_N)^q.$$

We will need to bound both sums. The first sum simplifies thus.

$$\begin{aligned}
\sum_{n=1}^{N-1} a_n^{-2q} r_n^{(2-q)} \psi(a_N)^q &= \psi(a_N)^q \sum_{n=1}^{N-1} a_n^{-q} n^{-1} \phi(a_n)^q \\
&= \psi(a_N)^q \sum_{n=1}^{N-1} n^{-1} \psi(a_n)^{-q}.
\end{aligned}$$

Now we make use of the fact that $\sum_{n=1}^{j-1} \psi(a_n)^{-q} < \psi(a_j)^{-q}$ and $\sum_{n=j}^{N-1} \psi(a_n)^{-q} < \psi(a_N)^{-q}$. Thus

$$\begin{aligned}
\psi(a_N)^q \sum_{n=1}^{N-1} n^{-1} \psi(a_n)^{-q} &= \psi(a_N)^q \sum_{n=1}^{j-1} n^{-1} \psi(a_n)^{-q} + \psi(a_N)^q \sum_{n=j}^{N-1} n^{-1} \psi(a_n)^{-q} \\
&\leq \psi(a_N)^q \sum_{n=1}^{j-1} \psi(a_n)^{-q} + j^{-1} \psi(a_N)^q \sum_{n=j}^{N-1} \psi(a_n)^{-q} \\
&\leq \psi(a_N)^q \psi(a_j)^{-q} + j^{-1}.
\end{aligned}$$

To get a bound for the second sum, we note that $a_n^{-2q} = \left(\frac{3}{4}\right)^{-2q} 2^{2qn}$. Hence

$$\begin{aligned}
\sum_{n=1}^{N-1} a_n^{-2q} 16^{-n} \psi(a_N)^q &\leq \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \sum_{n=1}^{\infty} 2^{(2q-4)n} \\
&= \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \frac{2^{(2q-4)}}{1 - 2^{(2q-4)}}.
\end{aligned}$$

Thus we have the following bound for the leftmost term.

$$\sum_{n=1}^{N-1} a_n^{-2q} t_n \psi(a_N)^q \leq \psi(a_N)^q \psi(a_j)^{-q} + j^{-1} + \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \frac{2^{(2q-4)}}{1 - 2^{(2q-4)}}.$$

Then by choosing j sufficiently large, it follows that

$$\sum_{n=1}^{N-1} a_n^{-2q} t_n^{(2-q)} \psi(a_N)^q \rightarrow 0$$

as $N \rightarrow \infty$. We next bound the middle term.

$$\begin{aligned}
\phi(a_N)^{-q} s_N^{-q} t_N &= \phi(a_N)^{-q} s_N^{-q} (r_N^{2-q} + 16^{-N}) \\
&= \phi(a_N)^{-q} 5^q N^{\frac{1}{2}} a_N^{-q} (N^{-1} a_N^q \phi(a_N)^q + 16^{-N}) \\
&= 5^q N^{-\frac{1}{2}} + 5^q \phi(a_N)^{-q} N^{\frac{1}{2}} a_N^{-q} 16^{-N}.
\end{aligned}$$

Since $\phi(r) = \frac{r}{\psi(r)}$,

$$\begin{aligned}
5^q \phi(a_N)^{-q} N^{\frac{1}{2}} a_N^{-q} 16^{-N} &= 5^q a_N^{-2q} \psi(a_N)^q N^{\frac{1}{2}} 16^{-N} \\
&= \left(\frac{3}{4\sqrt{5}} \right)^{-2q} \psi(a_N)^q N^{\frac{1}{2}} 2^{(2q-4)N}.
\end{aligned}$$

Thus

$$\phi(a_N)^{-q} s_N^{-q} t_N^{2-q} = 5^q N^{-\frac{1}{2}} + \left(\frac{3}{4\sqrt{5}} \right)^{-2q} \psi(a_N)^q N^{\frac{1}{2}} 2^{(2q-4)N} \rightarrow 0$$

as $N \rightarrow \infty$. Finally, we bound the rightmost term.

$$\sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} t_n^{2-q} = \sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} r_n^{2-q} + \sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} 16^{-n}$$

We will need to bound both sums. To bound the first sum we make use of the property that $\phi(a_n) < \frac{1}{2}\phi(a_{n-1})$.

$$\begin{aligned}
\sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} r_n^{2-q} &= \phi(a_N)^{-q} \sum_{n=N+1}^{\infty} a_n^{-q} n^{-1} a_n^q \phi(a_n)^q \\
&< \phi(a_N)^{-q} (N+1)^{-1} \sum_{n=N+1}^{\infty} \phi(a_n)^q \\
&< \phi(a_N)^{-q} (N+1)^{-1} \sum_{n=1}^{\infty} \phi(a_N)^q 2^{-nq} \\
&= (N+1)^{-1} \frac{1}{2^q - 1}.
\end{aligned}$$

The second sum of the rightmost term is bounded in a similar way to the second sum of the leftmost term.

$$\begin{aligned}
\sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} 16^{-n} &= \sum_{n=N+1}^{\infty} a_n^{-q} a_N^{-q} \psi(a_N)^q 16^{-n} \\
&\leq \psi(a_N)^q \sum_{n=N+1}^{\infty} a_n^{-2q} 16^{-n} \\
&\leq \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \sum_{n=1}^{\infty} 2^{(2q-4)n} \\
&= \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \frac{2^{(2q-4)}}{1 - 2^{(2q-4)}}.
\end{aligned}$$

Therefore,

$$\sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} t_n^{(2-q)} \leq (N+1)^{-1} \frac{1}{2^q - 1} + \left(\frac{3}{4}\right)^{-2q} \psi(a_N)^q \frac{2^{(2q-4)}}{1 - 2^{(2q-4)}}$$

and hence

$$\sum_{n=N+1}^{\infty} a_n^{-q} \phi(a_N)^{-q} t_n^{(2-q)} \rightarrow 0$$

as $N \rightarrow \infty$. Thus all three sums are bounded and tend to 0 as $N \rightarrow \infty$. Hence

$$|f(y) - f(0)| \leq \epsilon \phi(|y|) \|f\|_p$$

for $y \in A_N \cap X^\circ \cap E$, provided N is chosen sufficiently large. □

References

- [1] Adams, D. and Hedberg, L.I.: Function Spaces and Potential Theory. Springer-Verlag, Berlin, (1996)
- [2] Evans, L.C.: Partial Differential Equations. Graduate Studies in Math., vol. 19, AMS, Providence, RI (1998)
- [3] Gamelin, T.: Uniform Algebras. Prentice-Hall (1969)
- [4] Hedberg, L.I.: Bounded point evaluations and capacity. J. Funct. Anal. **10**, 269-280 (1972)

- [5] O'Farrell, A.G.: Analytic capacity and equicontinuity *BLMS* **10** 276-279 (1978)
- [6] Wang, J.: An approximate Taylor's theorem for $R(X)$. *Math. Scand.*, **33**, 343-358
- [7] Wang, J.: Modulus of approximate continuity for $R(X)$. *Math. Scand.*, **34**, 219-225 (1974)
- [8] Wolf, E.: Bounded point evaluations and smoothness properties of functions in $R^p(X)$. *Trans. Amer. Math. Soc.* **238**, 71-88 (1978)