

Every $(13k - 6)$ -strong tournament with minimum out-degree at least $28k - 13$ is k -linked

Jørgen Bang-Jensen and Kasper Skov Johansen*

June 23, 2021

Abstract

A digraph D is k -linked if it satisfies that for every choice of disjoint sets $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ of vertices of D there are vertex disjoint paths P_1, \dots, P_k such that P_i is an (x_i, y_i) -path. Confirming a conjecture by Kühn et al, Pokrovskiy proved in 2015 that every $452k$ -strong tournament is k -linked and asked for a better linear bound. Very recently Meng et al proved that every $(40k - 31)$ -strong tournament is k -linked. In this note we use an important lemma from their paper to give a short proof that every $(13k - 6)$ -strong tournament of minimum out-degree at least $28k - 13$ is k -linked.

Keywords: tournament, linkage, connectivity.

1 Introduction

A digraph D is **strongly connected** if it has a directed path from x to y (an (x, y) -path) for every ordered pair of distinct vertices x, y and it is **k -strong** if it has at least $k + 1$ vertices and remains strongly connected when we delete any set of at most $k - 1$ vertices. A (di)graph D is **k -linked** if it has vertex disjoint paths P_1, \dots, P_k such that P_i is an (x_i, y_i) -path for every choice of disjoint sets $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ of vertices of D . Thomas and Wollan proved that for every natural number k every $10k$ -connected undirected graph is k -linked [11]. Hence there exists a smallest function $f(k)$ such that every $f(k)$ -connected undirected graph is k -linked.

Thomassen [13] showed that for general digraphs there is no integer ℓ so that every ℓ -strong digraph is 2-linked, thus ruling out the existence of a similar function $f(k)$ for directed k -linkage. Hence it is natural to focus on special classes of digraphs and one such important class of digraphs is tournaments, that is, digraphs in which there is exactly one arc between every pair of distinct vertices. Thomassen [12] was the first to prove that for tournaments there exists a function $h(k)$ such that every $h(k)$ -strong tournament is k -linked. Bang-Jensen [1] proved that $h(2) = 5$ but already the value of $h(3)$ is open. Thomassen's function $h(k)$ was exponential in k . This was improved to a polynomial in k by Kühn, Lapinskas, Osthus, and Patel in [8] and they conjectured that a linear function would suffice. Pokrovskiy confirmed this in [10] by showing that every $452k$ -strong tournament is k -linked.

If $X = \{x_1, \dots, x_k\}$ and $Y = \{y_1, \dots, y_k\}$ are disjoint subsets of vertices in a digraph D , we say that X **anchors** Y in D if for every permutation π of $\{1, 2, \dots, k\}$ there exists a collection of disjoint paths Q_1, \dots, Q_k in D such that Q_i is an $(x_i, y_{\pi(i)})$ -path. The proof in [10] is based on the following lemma which shows that tournaments have a rich structure.

Lemma 1. [10] *For every natural number p every tournament T on at least $11p$ vertices contains disjoint sets X, Y both of size p such that X anchors Y in T*

Very recently Meng, Rolek, Wang and Yu [9] used the following improved version of Lemma 1 together with some nice ideas to show that every $(40k - 31)$ -strong tournament is k -linked.

*Department of Mathematics and Computer Science, University of Southern Denmark, email (jbj@imada.sdu.dk and kajoh14@student.sdu.dk). Research supported by the Independent Research Fund Denmark under grant number DFF 7014-00037B. This paper was written while the second author was a masters student at Department of Mathematics and Computer Science

Lemma 2. [9] Let p be a natural number and let T be a tournament on $n \geq 9p - 6$ vertices. Then T contains a pair of disjoint vertex sets X, Y , each of size p such that X anchors Y in T .

In this paper we use Lemma 2 to obtain the following better bound.

Theorem 3. For every natural number k every $(13k - 6)$ -strong tournament of minimum out-degree at least $28k - 13$ is k -linked.

The proof in [9] that every $(40k - 31)$ -strong tournament is k -linked uses an approach similar to that used by Pokrovskiy in [10] but is based on Lemma 2 instead of Lemma 1 and some matching arguments. Our proof also uses Lemma 2 but we avoid the use of matching arguments.

Notation not already introduced or given below is consistent with [2, 3]. Let $D = (V, A)$ be a digraph and let $X \subseteq V$, then we denote by $D\langle X \rangle$ the subdigraph of D induced by X . We denote by $N_D^+(v), N_D^-(v)$ the set of out-neighbours, respectively in-neighbours of v in D , that is, the set of vertices $y \neq v$ such that $v \rightarrow y$, respectively $y \rightarrow v$ is an arc of D . We call $d_D^+(v) = |N_D^+(v)|$, $d_D^-(v) = |N_D^-(v)|$ the **out-degree**, respectively the **in-degree** of the vertex v in D . If X is a subset of the vertices of D then we use the notation $D - X$ for the digraph $D\langle V(D) \setminus X \rangle$.

2 Proof of Theorem 3

Let $T_0 = T - (X_0 \cup Y_0)$. Pick a vertex $u_1 \in V(T_0)$ of minimum out-degree in T_0 , that is $d_{T_0}^+(u_1) \leq d_{T_0}^+(v)$ for all $v \in V(T_0)$. Then we find a vertex v_1 with minimum out-degree in $T\langle N_{T_0}^+(u_1) \rangle$. Now let $A_1 = N_{T_0}^+(u_1) \cap N_{T_0}^+(v_1)$ and set $D_1 = \{u_1, v_1\}$. Also, let $T_1 = T_0 - D_1$.

For $i = 2, \dots, 9k - 6$ in that order we pick $u_i, v_i \in T_{i-1}$ such that u_i has minimum out-degree in T_{i-1} and v_i is a vertex of minimum out-degree in $T\langle N_{T_{i-1}}^+(u_i) \rangle$. Let $A_i = N_{T_{i-1}}^+(u_i) \cap N_{T_{i-1}}^+(v_i)$, $D_i = \{u_i, v_i\}$ and $T_i = T_{i-1} - D_i$. Finally set $U = \{u_1, \dots, u_{9k-6}\}$ and $V = \{v_1, \dots, v_{9k-6}\}$.

The tournament $T\langle N_{T_{i-1}}^+(u_i) \rangle$ has $d_{T_{i-1}}^+(u_i)$ vertices so it has a vertex w whose out-degree in $T\langle N_{T_{i-1}}^+(u_i) \rangle$ is at most $(d_{T_{i-1}}^+(u_i) - 1)/2$. Because v_i was picked to have smallest out-degree in $T\langle N_{T_{i-1}}^+(u_i) \rangle$ it follows that

$$|A_i| = |N_{T_{i-1}}^+(u_i) \cap N_{T_{i-1}}^+(v_i)| = d_{T\langle N_{T_{i-1}}^+(u_i) \rangle}^+(v_i) \leq (d_{T_{i-1}}^+(u_i) - 1)/2. \quad (1)$$

By (1) and the fact that u_i has minimum out-degree in T_{i-1} we get

$$|A_i| \leq (d_{T_{i-1}}^+(u_i) - 1)/2 \leq (d_{T_{i-1}}^+(z) - 1)/2, \quad \forall z \in A_i \quad (2)$$

Then since $V(T_{i-1}) \subseteq V(T)$ we get from (2) that

$$|A_i| \leq (d_T^+(z) - 1)/2 \leq (d_T^+(z) - 1)/2, \quad \forall z \in A_i. \quad (3)$$

Since V contains $9k - 6$ vertices we know, by Lemma 2, that some disjoint subsets $V' = \{v'_1, \dots, v'_k\}$ and $V'' = \{v''_1, \dots, v''_k\}$ exist in V such that V' anchors V'' within $T\langle V \rangle$. Let $V^* = V - (V' \cup V'')$. We also collect the k vertices u'_i from U , for $i = 1, \dots, k$, where v'_i was picked in the tournament induced on the out-neighbors of u'_i . Let $U' = \{u'_1, \dots, u'_k\}$ and $U^* = U - U'$. This guarantees that $u'_i \rightarrow v'_i$ is an arc and that $D_{j_i} = \{u'_i, v'_i\}$ for some $j_i \in \{1, \dots, 9k - 6\}$. Then we let $A'_i = A_{j_i}$ and we get from (3) that

$$|A'_i| \leq (d_T^+(z) - 1)/2, \quad \forall z \in A'_i. \quad (4)$$

Next we consider the subtournament $T' = T - (X_0 \cup Y_0 \cup U' \cup V)$. Because $|X_0| + |Y_0| + |U'| + |V| = 12k - 6$ it follows that each $x_i \in X_0$ has at least $16k - 7$ out-neighbors in $V(T')$ (notice that x_i is not in $V(T')$ itself). We can therefore pick a set $X' = \{x'_1, \dots, x'_k\}$ of k distinct vertices of $V(T')$ such that $x_i \rightarrow x'_i$

is an arc of T .

Now we are going to construct k disjoint paths P_1, \dots, P_k such that P_i is an (x_i, v'_i) -path and these paths are disjoint.

Let $I = \{1, 2, \dots, k\}$ and let $\hat{I} \subseteq I$ be those indexes for which x'_i has an arc to at least one of u'_i, v'_i . If $i \in \hat{I}$ then T contains either the path $x_i \rightarrow x'_i \rightarrow v'_i$ or the path $x_i \rightarrow x'_i \rightarrow u'_i \rightarrow v'_i$.

For $i \in I - \hat{I}$ we have that x'_i is in A'_i , and by (4) it follows that

$$|A'_i| \leq (d_T^+(x'_i) - 1)/2. \quad (5)$$

Consider now the out-neighbors of x'_i in T . We have

$$|X_0| + |Y_0| + |U'| + |V| + |X'| + k - 1 = 14k - 7 = (28k - 14)/2 \leq (d_T^+(x'_i) - 1)/2 \quad (6)$$

Then by adding (5) and (6) we get

$$|A'_i| + |X_0| + |Y_0| + |U'| + |V| + |X'| + k - 1 \leq d_T^+(x'_i) - 1. \quad (7)$$

That is,

$$|A'_i| + |X_0| + |Y_0| + |U'| + |V| + |X'| + k \leq d_T^+(x'_i). \quad (8)$$

In other words, we see that x'_i has at least k out-neighbors outside of $(A'_i \cup X_0 \cup Y_0 \cup U' \cup V \cup X')$. We can therefore find a set $X'' = \{x''_i | i \in I - \hat{I}\}$ of distinct vertices with $X'' \subseteq V(T') - X'$ such that $x'_i \rightarrow x''_i$ is an arc in T and such that at least one of the arcs $x''_i \rightarrow u'_i$ and $x''_i \rightarrow v'_i$ exists in T . The latter follows by the fact that for all $i \in I - \hat{I}$ the vertex x''_i is not in A'_i . Therefore we can find, for all $i = 1, \dots, k$, some path from x_i to v'_i that has one of the following forms:

- $x_i \rightarrow x'_i \rightarrow v'_i$
- $x_i \rightarrow x'_i \rightarrow u'_i \rightarrow v'_i$
- $x_i \rightarrow x'_i \rightarrow x''_i \rightarrow v'_i$
- $x_i \rightarrow x'_i \rightarrow x''_i \rightarrow u'_i \rightarrow v'_i$

Call such paths P_1, \dots, P_k and note that, by construction, they are disjoint. Then each P_i has at most two vertices in $V(T')$, namely x'_i and x''_i . All vertices of the paths P_i are contained in $(X_0 \cup U' \cup V' \cup X' \cup X'')$. Furthermore, $|X_0 \cup U' \cup V' \cup X' \cup X''| \leq |X_0 \cup U' \cup V' \cup V^* \cup X' \cup X''| \leq 12k - 6$. Hence, since T is $(13k - 6)$ -strong it follows that $T^* = T - (X_0 \cup U' \cup V' \cup V^* \cup X' \cup X'')$ is k -strong and contains both Y_0 and V'' . By Menger's Theorem there exist k disjoint paths from V'' to Y_0 in T^* . Name these R_1, \dots, R_k so that R_i ends in y_i for $i \in \{1, 2, \dots, k\}$. Now, neither of the paths R_j intersect some path P_i because the vertices of each P_i are completely contained in $(X_0 \cup U' \cup V' \cup X' \cup X'')$. Suppose now we pair up P_i and R_i for each $i \in \{1, \dots, k\}$. Let $h(P_i)$ denote the terminal (last) vertex of P_i and let $t(R_j)$ denote the initial vertex of R_j . Then $(h(P_1), \dots, h(P_k))$ is a permutation of V' and $(t(R_1), \dots, t(R_k))$ is a permutation of V'' . By the choice of V' and V'' we can find a linkage $\{M_1, \dots, M_k\}$ from $\{h(P_1), \dots, h(P_k)\}$ to $\{t(R_1), \dots, t(R_k)\}$ whose inner vertices are completely within V^* . But then since P_i and R_i are disjoint from V^* it is easy to see that Q_1, Q_2, \dots, Q_k is a linkage from X_0 to Y_0 when we set $Q_i = P_i M_i R_i$ for each $i \in \{1, 2, \dots, k\}$. □

3 Remarks

Pokrovskiy gave an infinite family of $(2k - 2)$ -strong tournaments which have arbitrary high minimum in-degree and out-degree but are not k -linked and made following conjecture.

Conjecture 4. [10] *For every natural number k there exists an integer $d(k)$ such that every $2k$ -strong tournament with minimum in-degree and minimum out-degree at least $d(k)$ is k -linked.*

The following partial result was obtained by Girão and Snyder

Theorem 5. [6] For every natural number k there exists an integer $d(k)$ such that every $4k$ -strong tournament with minimum out-degree at least $d(k)$ is k -linked.

In an unpublished paper [5] Girão, Popielarz and Snyder proved Conjecture 4 up to an additive constant of 1.

Theorem 6. [5] For every natural number k there exists an integer $d(k)$ such that every $(2k+1)$ -strong tournament with minimum out-degree at least some polynomial in k is k -linked.

The function $d(k)$ used in Theorem 6 is $d(k) = Ck^{31}$ for some appropriate constant C and hence Theorem 6 does not imply Theorem 3.

The following result shows that the condition on high minimum out-degree cannot be removed in Conjecture 4.

Theorem 7. [5] For every natural number k there exists infinitely many $(5k-1)$ -strong tournaments which are not $2k$ -linked.

Girão, Popielarz and Snyder also showed that $2k$ cannot be replaced by $2k-1$ in Conjecture 4.

Theorem 8. [5] For all integers $k \geq 2$ and $m \geq 2k$ there exists a $(2k-1)$ -strong tournament T whose minimum in-degree and out-degree is at least m such that T is not k -linked.

A digraph is **semicomplete** if it has no pair of non-adjacent vertices. Thus the class of tournaments forms a subclass of the class of semicomplete digraphs. It was pointed out in [4] that Pokrovskiy's proof in [10] also holds for semicomplete digraphs, implying that every $452k$ -strong semicomplete digraph is k -linked. The proof of Theorem 3 does not hold for semicomplete digraphs because we cannot guarantee a vertex of out-degree at most $(d_{T_{i-1}}^+(u_i) - 1)/2$ in $T\langle N_{T_{i-1}}^+(u_i) \rangle$ (just above (1)) as the digraph $T\langle N_{T_{i-1}}^+(u_i) \rangle$ could be complete. It can also be checked that the proof in [9] does not hold for semicomplete digraphs.

It is known [7] that every $(3k-2)$ -strong semicomplete digraph contains a spanning k -strong tournament. From this and Theorem 3 it follows that every $(84k-41)$ -strong semicomplete digraph is k -linked but we believe this bound is far from best possible. We also believe that Conjecture 4 holds for semicomplete digraphs.

References

- [1] J. Bang-Jensen. On the 2-linkage problem for semicomplete digraphs. In *Graph theory in memory of G. A. Dirac (Sandbjerg, 1985)*, volume 41 of *Ann. Discrete Math.*, pages 23–37. North-Holland, 1989.
- [2] J. Bang-Jensen and Gutin. G (eds.). *Classes of Directed Graphs*. Springer Monographs in Mathematics. Springer Verlag, London, 2018.
- [3] J. Bang-Jensen and G. Gutin. *Digraphs: Theory, Algorithms and Applications*. Springer-Verlag, London, 2nd edition, 2009.
- [4] J. Bang-Jensen and F. Havet. Tournaments and Semicomplete Digraphs. In J. Bang-Jensen and G. Gutin, editors, *Classes of Directed Graphs*, chapter 2, pages 35–124. Springer Verlag London, 2018.
- [5] A. Girão, K. Popielarz, and R. Snyder. $(2k+1)$ -connected tournaments with large minimum out-degree are k -linked. arXiv:1912.00710.
- [6] A. Girão and R. Snyder. Highly linked tournaments with large minimum out-degree. *J. Comb. Theory, Ser. B*, 139:251–266, 2019.
- [7] Y. Guo. Spanning local tournaments in locally semicomplete digraphs. *Discrete Appl. Math.*, 79(1-3):119–125, 1997.

- [8] D. Kühn, D. Lapinskas, J. Osthus, and V. Patel. Proof of a conjecture of Thomassen on Hamilton cycles in highly connected tournaments. *Proc. London Math. Soc.*, 109:733–762, 2014.
- [9] W. Meng, M. Rolek, Y. Wang, and G. Yu. An improved linear connectivity bound for tournaments to be highly linked. *Eur. J. Comb.*, accepted.
- [10] A. Pokrovskiy. Highly linked tournaments. *J. Combin. Theory Ser B*, 115:339–347, 2015.
- [11] W. Thomas and P. Wollan. An improved linear edge bound for graph linkages. *Eur. J. Comb.*, 26(3-4):309–324, 2005.
- [12] C. Thomassen. Connectivity in tournaments. In *Graph theory and combinatorics (Cambridge, 1983)*, pages 305–313. Academic Press, 1984.
- [13] C. Thomassen. Highly connected non-2-linked digraphs. *Combinatorica*, 11(4):393–395, 1991.